

HIGHER RANK K-THEORETIC DONALDSON–THOMAS THEORY OF POINTS

NADIR FASOLA, SERGEJ MONAVARI, AND ANDREA T. RICOLFI

ABSTRACT. We exploit the critical structure on the Quot scheme $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, in particular the associated symmetric obstruction theory, in order to study rank r *K-theoretic* Donaldson–Thomas invariants of the local Calabi–Yau 3-fold \mathbb{A}^3 . We compute the associated partition function as a plethystic exponential, proving a conjecture proposed in string theory by Awata–Kanno and Benini–Bonelli–Poggi–Tanzini. A crucial step in the proof is the fact, nontrivial if $r > 1$, that the invariants do not depend on the equivariant parameters of the framing torus $(\mathbb{C}^*)^r$. Reducing from K-theoretic to *cohomological* invariants, we compute the corresponding DT invariants, proving a conjecture of Szabo. Reducing further to *enumerative* DT invariants, we solve the higher rank DT theory of a pair (X, F) , where F is an equivariant exceptional locally free sheaf on a projective toric 3-fold X .

As a further refinement of the K-theoretic DT invariants, we formulate a mathematical definition of the chiral elliptic genus studied in physics. This allows us to define *elliptic DT invariants* of \mathbb{A}^3 in arbitrary rank, which we use to tackle a conjecture of Benini–Bonelli–Poggi–Tanzini.

CONTENTS

1. Introduction	1
2. Background material	7
3. The local Quot scheme: critical and equivariant structure	9
4. Invariants attached to the local Quot scheme	17
5. Higher rank vertex on the local Quot scheme	19
6. The higher rank K-theoretic DT partition function	23
7. The higher rank cohomological DT partition function	28
8. Elliptic Donaldson–Thomas invariants	31
9. Higher rank DT invariants of compact toric 3-folds	38
References	48

1. INTRODUCTION

1.1. Overview. Classical Donaldson–Thomas (DT in short) invariants of a smooth complex projective Calabi–Yau 3-fold Y , introduced in [64], are integers that virtually count stable torsion free sheaves on Y , with fixed Chern character γ . However, the theory is much richer than what the bare DT numbers

$$(1.1) \quad \text{DT}(Y, \gamma) \in \mathbb{Z}$$

can capture: there are extra symmetries subtly hidden in the local structure of the moduli spaces of sheaves giving rise to the classical DT invariants (1.1). This idea has been present in the physics literature for some time [31, 22].

These hidden symmetries suggest that there should exist more *refined* invariants, of which the DT numbers (1.1) are just a shadow. These branch out in two main directions:

- *motivic* Donaldson–Thomas invariants, and
- *K-theoretic* Donaldson–Thomas invariants.

For the former, which at date includes a number of interesting subbranches, the papers by Kontsevich and Soibelman [35, 36] are a good starting point, and Szendrői’s survey [63] contains an extensive bibliography on the subject. For the latter, see some recent developments after Nekrasov–Okounkov [49], such as [52, 1, 66] and [48, 50, 13] for a generalisation to Calabi–Yau 4-folds. In this paper we deal with K-theoretic DT theory. The relationship between motivic and K-theoretic, which we briefly sketch in § 6.3, will be investigated in future work.

The subtle structure of the DT moduli spaces is most evident in the *local case*, i.e. when the theory is applied to the simplest Calabi–Yau 3-fold of all, namely the affine space \mathbb{A}^3 . See [6, 20] for the rank 1 motivic DT theory of \mathbb{A}^3 , and [59] for a higher rank version. This paper solves the K-theoretic Donaldson–Thomas theory of points of \mathbb{A}^3 . In [4] it is shown that the main player in the theory, the Quot scheme

$$(1.2) \quad \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$$

of length n quotients of the free sheaf $\mathcal{O}^{\oplus r}$, is a global *critical locus*, i.e. it can be realised as $\{df=0\}$, for f a regular function on a smooth scheme. This structural result, revealing in bright light the symmetries we were talking about, is used to define the *higher rank K-theoretic DT theory of points* that is the central character in this paper. The rank 1 theory, corresponding to $\text{Hilb}^n(\mathbb{A}^3)$, was already defined, and it was solved by Okounkov [52, § 3], proving a conjecture by Nekrasov [46]. Our first main result (Theorem A) can be seen as an upgrade of his calculation, completing the study of the degree 0 K-theoretic DT theory of \mathbb{A}^3 .

In physics, remarkably, the definition of the K-theoretic DT invariants studied here already existed, and gave rise to a conjecture that our paper — again, Theorem A — proves mathematically. More precisely, our formula for the K-theoretic DT partition function DT_r^K of \mathbb{A}^3 was first conjectured by Nekrasov [46] for $r=1$ and by Awata and Kanno [3] for arbitrary r as the partition function of a quiver matrix model describing instantons of a topological $U(r)$ gauge theory on D6 branes. The reader is referred to § 1.3 of this introduction for more background on the physics picture.

We also study higher rank *cohomological DT invariants* of \mathbb{A}^3 . As we show in Corollary 7.1, these can be obtained as a suitable limit of the K-theoretic invariants. Motivated by [3, 47], a closed formula for their generating function DT_r^{coh} was conjectured by Szabo [62, Conj. 4.10] as a generalisation of the $r=1$ case established by Maulik–Nekrasov–Okounkov–Pandharipande [45, Thm. 1]. We prove this conjecture as our Theorem B. To get there, in § 5 we develop a *higher rank topological vertex* formalism based on the combinatorics of *r-colored plane partitions*,¹ generalising the classical vertex formalism of [44, 45].

We pause for a second to explain a key step in this paper. The Quot scheme (1.2), which gives rise to most of the invariants we study here, is acted on by an algebraic torus

$$\mathbf{T} = (\mathbb{C}^*)^3 \times (\mathbb{C}^*)^r,$$

and by their very definition, both the K-theoretic and the cohomological DT invariants depend, a priori, on the sets $t = (t_1, t_2, t_3)$ and $w = (w_1, \dots, w_r)$ of equivariant parameters of \mathbf{T} . A technical result of this paper, which is proved as Theorem 6.5, states that

$$(1.3) \quad \text{The K-theoretic DT invariants do not depend on the framing parameters } w.$$

¹In this paper, an r -colored plane partition is an r -tuple of classical plane partitions, see Definition 3.10.

This will allow us to take arbitrary limits to evaluate our formulae. We emphasise that this independence, automatic if $r = 1$ (see Remark 6.2), is quite surprising and highly nontrivial if $r > 1$.

1.2. Main results. We briefly outline here the main results obtained in this paper.

1.2.1. K -theoretic DT invariants. As we mentioned above, the Quot scheme (1.2) is a critical locus, thus it carries a natural symmetric (\mathbf{T} -equivariant, as we prove) perfect obstruction theory in the sense of Behrend–Fantechi [7, 8]. As we recall in §4.1.1, there is also an induced *twisted virtual structure sheaf* $\widehat{\mathcal{O}}^{\text{vir}} \in K_0^{\mathbf{T}}(\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n))$, which is a twist, by a square root of the virtual canonical bundle, of the ordinary virtual structure sheaf \mathcal{O}^{vir} . The rank r K -theoretic DT partition function of the Quot scheme of points of \mathbb{A}^3 , encoding the rank r K -theoretic DT invariants of \mathbb{A}^3 , is defined as

$$\text{DT}_r^K(\mathbb{A}^3, q, t, w) = \sum_{n \geq 0} q^n \chi(\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \widehat{\mathcal{O}}^{\text{vir}}) \in \mathbb{Z}[[t, (t_1 t_2 t_3)^{\frac{1}{2}}, w]][q],$$

where the half power is caused by the twist by the chosen square root of the virtual canonical bundle (this choice does not affect the invariants, cf. Remark 4.3).

Granting Theorem 6.5, that we stated informally in (1.3), we shall write $\text{DT}_r^K(\mathbb{A}^3, q, t) = \text{DT}_r^K(\mathbb{A}^3, q, t, w)$, ignoring the framing parameters w . In §6.2 we determine a closed formula for this series, proving the conjecture by Awata–Kanno [3]. This conjecture was checked for low number of instantons in [3, §4].

To state our first main result, we need to recall the definition of the *plethystic exponential*. Given an arbitrary power series

$$f = f(p_1, \dots, p_e; u_1, \dots, u_\ell) \in \mathbb{Q}(p_1, \dots, p_e)[[u_1, \dots, u_\ell]],$$

one sets

$$(1.4) \quad \text{Exp}(f) = \exp \left(\sum_{n > 0} \frac{1}{n} f(p_1^n, \dots, p_e^n; u_1^n, \dots, u_\ell^n) \right),$$

viewed as an element of $\mathbb{Q}(p_1, \dots, p_e)[[u_1, \dots, u_\ell]]$. Consider, for a formal variable x , the operator $[x] = x^{1/2} - x^{-1/2}$. In §6.1 we consider this operator on $K_0^{\mathbf{T}}(\text{pt})$. See §8.4 for its physical interpretation. We are now able to state our first main result.

Theorem A (Theorem 6.3). *The rank r K -theoretic DT partition function of \mathbb{A}^3 is given by*

$$(1.5) \quad \text{DT}_r^K(\mathbb{A}^3, (-1)^r q, t) = \text{Exp}(F_r(q, t_1, t_2, t_3)),$$

where, setting $t = t_1 t_2 t_3$, one defines

$$F_r(q, t_1, t_2, t_3) = \frac{[t^r]}{[t][t^{\frac{r}{2}}q][t^{\frac{r}{2}}q^{-1}]} \frac{[t_1 t_2][t_1 t_3][t_2 t_3]}{[t_1][t_2][t_3]}.$$

The case $r = 1$ of Theorem A was proved by Okounkov in [52]. The general case was proposed conjecturally in [3, 9]. Arbesfeld and Kononov informed us during the writing of this paper that they also obtained a different proof of Theorem A [2], driven by different motivations.

It is interesting to notice that Formula (1.5) is equivalent to the product decomposition

$$(1.6) \quad \text{DT}_r^K(\mathbb{A}^3, (-1)^r q, t) = \prod_{i=1}^r \text{DT}_1^K(\mathbb{A}^3, -q t^{\frac{-r-1}{2}+i}, t),$$

that we obtain in Theorem 6.9. This is precisely the product formula [50, Formula (3.14)] appearing as a limit of the (conjectural) 4-fold theory developed by Nekrasov and Piazzalunga.

Formula (1.6) is also related to its motivic cousin: as we observe in § 6.3, the motivic partition function $\mathrm{DT}_r^{\mathrm{mot}}$ of the Quot scheme of points of \mathbb{A}^3 (see [4, Prop. 2.7] and the references therein) satisfies the same product formula (1.6), after the transformation $t^{1/2} \rightarrow -\mathbb{L}^{1/2}$.

1.2.2. Cohomological DT invariants. The generating function of *cohomological DT invariants* is defined as

$$\mathrm{DT}_r^{\mathrm{coh}}(\mathbb{A}^3, q, s, v) = \sum_{n \geq 0} q^n \int_{[\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)]^{\mathrm{vir}}} 1 \in \mathbb{Q}((s, v))[[q]]$$

where $s = (s_1, s_2, s_3)$ and $v = (v_1, \dots, v_r)$, with $s_i = c_1^T(t_i)$ and $v_j = c_1^T(w_j)$ respectively, and the integral is defined in Equation (4.2) via \mathbf{T} -equivariant residues. It is a consequence of (1.3) that $\mathrm{DT}_r^{\mathrm{coh}}(\mathbb{A}^3, q, s, v)$ does not depend on v , so we will shorten it as $\mathrm{DT}_r^{\mathrm{coh}}(\mathbb{A}^3, q, s)$. In § 7.1 we explain how to recover the cohomological invariants out of the K-theoretic ones. This is the limit formula (Corollary 7.1)

$$\mathrm{DT}_r^{\mathrm{coh}}(\mathbb{A}^3, q, s) = \lim_{b \rightarrow 0} \mathrm{DT}_r^K(\mathbb{A}^3, q, e^{bs}),$$

essentially a formal consequence of our explicit expression for the K-theoretic *higher rank vertex* (cf. § 5) attached to the Quot scheme $\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$.

Theorem B (Theorem 7.2). *The rank r cohomological DT partition function of \mathbb{A}^3 is given by*

$$\mathrm{DT}_r^{\mathrm{coh}}(\mathbb{A}^3, q, s) = M((-1)^r q)^{-r} \frac{(s_1 + s_2)(s_1 + s_3)(s_2 + s_3)}{s_1 s_2 s_3},$$

where $M(t) = \prod_{m \geq 1} (1 - t^m)^{-m}$ is the MacMahon function.

The case $r = 1$ of Theorem B was proved by Maulik–Nekrasov–Okounkov–Pandharipande [45, Thm. 1]. Theorem B was conjectured by Szabo in [62] and confirmed for $r \leq 8$ and $n \leq 8$ in [9]. The specialisation $\mathrm{DT}_r^{\mathrm{coh}}(\mathbb{A}^3, q, s)|_{s_1 + s_2 + s_3 = 0} = M((-1)^r q)^r$ was already computed in physics [19].

1.2.3. Elliptic DT invariants. In § 8 we define the *virtual chiral elliptic genus* for any scheme with a perfect obstruction theory, which recovers as a special case the virtual elliptic genus defined in [23]. By means of this new invariant we introduce a refinement $\mathrm{DT}_r^{\mathrm{ell}}$ of the generating series DT_r^K , providing a mathematical definition of the *elliptic DT invariants* studied in [9]. We propose a conjecture (Conjecture 8.9) about the behaviour of $\mathrm{DT}_r^{\mathrm{ell}}$ and, granting this conjecture, we obtain a proof of a conjecture formulated by Benini–Bonelli–Poggi–Tanzini (Theorem 8.11).

1.2.4. Global geometry. So far we have only discussed results concerning *local* geometry. When X is a *projective* toric 3-fold and F is an equivariant exceptional locally free sheaf, by [59, Thm. A] there is a 0-dimensional torus equivariant (cf. Proposition 9.2) perfect obstruction theory on $\mathrm{Quot}_X(F, n)$. Therefore the higher rank Donaldson–Thomas invariants

$$\mathrm{DT}_{F,n} = \int_{[\mathrm{Quot}_X(F,n)]^{\mathrm{vir}}} 1 \in \mathbb{Z}$$

can be computed via the Graber–Pandharipande virtual localisation formula [27]. The next result confirms (in the toric case) a prediction [59, Conj. 3.5] for their generating function. Before stating it, recall that an *exceptional sheaf* on a variety X is a coherent sheaf $F \in \mathrm{Coh} X$ such that $\mathrm{Hom}(F, F) = \mathbb{C}$ (i.e. F is simple), and $\mathrm{Ext}^i(F, F) = 0$ for $i > 0$.

Theorem C (Theorem 9.7). *Let (X, F) be a pair consisting of a smooth projective toric 3-fold X along with an exceptional equivariant locally free sheaf F of rank r . Then*

$$\sum_{n \geq 0} \text{DT}_{F,n} q^n = M((-1)^r q)^r \int_X c_3(T_X \otimes K_X).$$

The corresponding formula in the Calabi–Yau case was proved in [59, § 3.2], whereas the general rank 1 case was proved in [45, Thm. 2] and [39, Thm. 0.2].

1.3. Relation to string theory. An interpretation for Donaldson–Thomas invariants is available also in the context of supersymmetric string theories. In this framework, one is interested in countings of BPS-bound states on (Calabi–Yau) 3-folds. The interest in studying the BPS sector of string theories lies in the fact that it consists of quantities which are usually protected from quantum corrections, and which can sometimes be studied non-perturbatively. On the other hand, BPS countings have been shown in many occasions to have a precise mathematical interpretation rooted in counting problems in enumerative geometry. In the case of Donaldson–Thomas theory the interpretation is that of a type IIA theory compactified on X , so that the four dimensional effective theory has an $\mathcal{N} = 2$ supersymmetry content. BPS states then preserve half of this supersymmetry and are indexed by a charge vector living on a lattice determined by the cohomology (with compact support) of the 3-fold $\Gamma \cong H^0(X, \mathbb{Z}) \oplus H^2(X, \mathbb{Z}) \oplus H^4(X, \mathbb{Z}) \oplus H^6(X, \mathbb{Z})$. These lattices can also be interpreted as the charge lattices of D-branes wrapping p -cycles on X , where

$$\text{D}p \longleftrightarrow H^{6-p}(X, \mathbb{Z}) \cong H_p(X, \mathbb{Z}),$$

for $p = 0, 2, 4, 6$. The Witten index

$$\text{Ind}_X(\gamma) = \text{Tr}_{\mathcal{H}_{\text{BPS}}^{(\gamma)}} (-1)^F$$

provides a measure of the degeneracy of the BPS states, where the trace is over the fixed charge sector of the single-particle Hilbert space $H_{\text{BPS}} = \bigoplus_{\gamma \in \Gamma} \mathcal{H}_{\text{BPS}}^{(\gamma)}$ and F is a given one-particle operator acting on the γ -component of the Hilbert space. The choice of a charge vector of the form $\gamma = (r, 0, -\beta, n)$ is then equivalent to a system of (D0–D4–D6)-branes, where r D6-branes wrap the whole 3-fold X . The index of the theory is then defined via integration over the virtual class of the moduli space of BPS states $\mathcal{M}_{\text{BPS}}^{(\gamma)}(X)$,

$$\text{Ind}_X(\gamma) = \int_{[\mathcal{M}_{\text{BPS}}^{(\gamma)}(X)]^{\text{vir}}} 1.$$

The moduli space of BPS states on $X = \mathbb{A}^3$ with charge vector $\gamma = (r, 0, 0, n)$ is then identified with the Quot scheme $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, and the partition function of the theory reproduces the generating function of degree 0 DT invariants of \mathbb{A}^3

$$\sum_{n \geq 0} \text{Ind}_{\mathbb{A}^3}(r, 0, 0, n) q^n = \text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s).$$

An equivalent interpretation can be also given in type IIB theories, where the relevant systems will be those of (D(-1)–D3–D5)-branes. The effective theory on the D(-1)-branes, which in this case are 0-dimensional objects, is a quiver matrix model encoding the critical structure of $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$. K-theoretic and elliptic versions of DT invariants can also be studied by suitably generalising the D-branes construction. One can study for instance a D0–D6 brane system with r branes on $X \times S^1$, S^1 being the worldvolume of the D0-branes. In this case the D0-branes quantum mechanics describe the K-theoretic generalisation of DT theory, so that

the supersymmetric partition function computes the equivariant Euler characteristic of the twisted virtual structure sheaf $\widehat{\mathcal{O}}^{\text{vir}}$ of the Quot scheme identified with the moduli space of BPS vacua of the theory. Analogously, the D1 string theory of a D1-D7 system on the product of X by an elliptic curve, say $X \times T^2$, provides the elliptic generalisation of higher rank DT invariants, [9], while the superconformal index realises the virtual elliptic genus of the moduli space of BPS states. In this context, a plethystic formula for the generating function of the K-theoretic DT invariants was conjectured in [47] for rank 1 and in [3] for rank r . The elliptic generalisation of DT theory was studied in [9], where a plethystic formula for the virtual elliptic genus was conjectured in the Calabi–Yau specialisation $t_1 t_2 t_3 = 1$ and its generalisation $(t_1 t_2 t_3)^r = 1$ (cf. Theorem 8.11).

1.4. Related works on virtual invariants of Quot schemes. The study of virtual invariants of Quot schemes is an active research area in enumerative geometry. See for instance the seminal work of Marian–Oprea [42], where a virtual fundamental class on the Quot scheme of the trivial bundle on a smooth projective curve was constructed. More recently, there has been a lot of activity on surfaces (partially motivated by Vafa–Witten theory), see e.g. [53, 32, 41, 43]. In connection to physics, flags of framed torsion free sheaves on \mathbb{P}^2 , as a generalisation of Quot schemes of points on \mathbb{A}^2 , were studied in [11, 12]. Many of the results mentioned above use virtual localisation to study virtual invariants of Quot schemes: our paper exploits virtual localisation as well, though in the K-theoretic setup and in the almost unexplored land of 3-folds.

Generating functions of Euler characteristics of Quot schemes on 3-folds (for quotients of sheaves of homological dimension at most 1) were computed by Gholampour–Kool in [25]. Quot schemes of *reflexive* sheaves (of rank 2) also appeared in the work of Gholampour–Kool–Young [26], as fibres of “double dual maps”, exploited to compute generating functions of Euler characteristics of more complicated moduli spaces of sheaves. Virtual invariants are also defined (again, in the rank 2 case) via localisation in [26]. In the rank 1 case, quotients of the ideal sheaf of a smooth curve C in a 3-fold Y form a closed subscheme of the Hilbert scheme of curves in Y , the key player in rank 1 DT theory. The series of virtual Euler characteristics of the associated Quot scheme was computed by the third author in [57], and later upgraded to a C -local DT/PT wall-crossing formula [56]. The motivic refinement of this DT/PT correspondence was established by Davison and the third author in [20].

Virtual classes of dimension 0 of Quot schemes (for locally free sheaves) on projective 3-folds are constructed under certain assumptions in [59]. In this paper we mainly work with the virtual class on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ coming from the critical obstruction theory found in [4].

1.5. Plan of the paper. In § 2 we recall the basics on perfect obstruction theories, virtual classes, virtual structure sheaves and how to produce virtual invariants out of these data; we review the K-theoretic virtual localisation theorem in § 2.5. Sections 3–8 are devoted to the “local Quot scheme” $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$. In § 3 we recall its critical structure and we define a \mathbf{T} -action on it, whose fixed locus is parametrised by the finitely many r -colored plane partitions (Proposition 3.12); we study the equivariant critical obstruction theory on the Quot scheme and prove that the induced virtual class on the \mathbf{T} -fixed locus is trivial (Corollary 3.15). In § 4.3 we introduce cohomological and K-theoretic DT invariants of $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$. In § 5 we develop a higher rank vertex formalism which we exploit to write down a formula (Proposition 5.2) for the virtual tangent space of a \mathbf{T} -fixed point in the Quot scheme. In § 6 we prove Theorem A as well as Formula (1.6). In § 7 we prove Theorem B and we show that DT_r^{coh} does not depend on

any choice of possibly nontrivial $(\mathbb{C}^*)^3$ -weights on $\mathcal{O}^{\oplus r}$. In §8, we give a mathematically rigorous definition of a “chiral” version of the virtual elliptic genus of [23] and use it in §8.2 to define elliptic DT invariants. In §8.3 we also give closed formulae for elliptic DT invariants in some limiting cases, based on the conjectural independence on the elliptic parameter — see Conjecture 8.9 and Remark 8.12. In particular Theorem 8.11 proves a conjecture recently appeared in the physics literature [9, Formula (3.20)]. In §9 we prove Theorem C by gluing vertex contributions from the toric charts of a projective toric 3-fold.

Conventions. We work over \mathbb{C} . A *scheme* is a separated scheme of finite type over \mathbb{C} . If Y is a scheme, we let $\mathbf{D}^{[a,b]}(Y)$ denote the derived category of coherent sheaves on Y , whose objects are complexes with vanishing cohomology sheaves outside the interval $[a, b]$. We let $K^0(Y)$ be the K-group of vector bundles on Y . When Y carries an action by an algebraic torus \mathbf{T} , we let $K_{\mathbf{T}}^0(Y)$ be the K-group of \mathbf{T} -equivariant vector bundles on Y . Similarly, we let $K_0(Y)$ denote the K-group of *coherent sheaves* on Y , and we let $K_0^{\mathbf{T}}(Y)$ be the K-group of (the abelian category of) \mathbf{T} -equivariant coherent sheaves on Y . When Y is smooth, the natural \mathbb{Z} -linear map $K^0(Y) \rightarrow K_0(Y)$, resp. $K_{\mathbf{T}}^0(\text{pt})$ -linear map $K_{\mathbf{T}}^0(Y) \rightarrow K_0^{\mathbf{T}}(Y)$, is an isomorphism. Chow groups $A^*(Y)$ and cohomology groups $H^*(Y)$ are taken with rational coefficients.

Acknowledgements. We thank Noah Arbesfeld and Yakov Kononov for generously sharing with us their progress on their paper [2], for reading a first draft of this paper and for sending us very interesting comments. Thanks to Alberto Cazzaniga for suggesting the comparison between motivic and K-theoretic factorisations. We thank Barbara Fantechi for enlightening discussions about equivariant obstruction theories. We are grateful to Martijn Kool, Giulio Bonelli and Alessandro Tanzini for the many discussions, insightful suggestions and for commenting on a first draft of this paper. Special thanks to Richard Thomas for generously sharing with us his ideas and insights on equivariant sheaves and Atiyah classes. Finally, many thanks to the anonymous referee for suggesting valuable improvements and corrections.

S.M. is supported by NWO grant TOP2.17.004. A.R. was funded by Dipartimenti di Eccellenza.

2. BACKGROUND MATERIAL

2.1. Obstruction theories and virtual classes. A *perfect obstruction theory* on a scheme X , as defined in [40, 7], is the datum of a morphism

$$\phi: \mathbb{E} \rightarrow \mathbb{L}_X$$

in $\mathbf{D}^{[-1,0]}(X)$, where \mathbb{E} is a perfect complex of perfect amplitude contained in $[-1, 0]$, such that $h^0(\phi)$ is an isomorphism and $h^{-1}(\phi)$ is surjective. Here, $\mathbb{L}_X = \tau_{\geq -1} L_X^\bullet$ is the cut-off at -1 of the full cotangent complex $L_X^\bullet \in \mathbf{D}^{[-\infty, 0]}(X)$ defined by Illusie [30]. A perfect obstruction theory is called *symmetric* (see [8]) if there exists an isomorphism $\theta: \mathbb{E} \xrightarrow{\sim} \mathbb{E}^\vee[1]$ such that $\theta = \theta^\vee[1]$.

The *virtual dimension* of X with respect to (\mathbb{E}, ϕ) is the integer $\text{vd} = \text{rk } \mathbb{E}$. This is just $\text{rk } E^0 - \text{rk } E^{-1}$ if one can write $\mathbb{E} = [E^{-1} \rightarrow E^0]$.

Remark 2.1. A symmetric obstruction theory $\mathbb{E} \rightarrow \mathbb{L}_X$ has virtual dimension 0, and moreover the *obstruction sheaf*, defined as $\text{Ob} = h^1(\mathbb{E}^\vee)$, is canonically isomorphic to the cotangent sheaf Ω_X . In particular, one has the K-theoretic identity

$$\mathbb{E} = h^0(\mathbb{E}) - h^{-1}(\mathbb{E}) = \Omega_X - T_X \in K_0(X).$$

A perfect obstruction theory determines a cone

$$\mathfrak{C} \hookrightarrow E_1 = (E^{-1})^\vee.$$

Letting $\iota: X \hookrightarrow E_1$ be the zero section of the vector bundle E_1 , the induced *virtual fundamental class* on X is the refined intersection

$$[X]^{\text{vir}} = \iota^![\mathfrak{C}] \in A_{\text{vd}}(X).$$

By a result of Siebert [61, Thm. 4.6], the virtual fundamental class depends only on the K-theory class of \mathbb{E} .

Definition 2.2 ([27, 8]). Let G be an algebraic group acting on a scheme X . A G -equivariant perfect obstruction theory on X is a choice of lift of a perfect obstruction theory $\phi: \mathbb{E} \rightarrow \mathbb{L}_X$ to the G -equivariant derived category $\mathbf{D}^{[-1,0]}(\text{Coh}_X^G)$.

2.2. From obstruction theories to virtual invariants. On a proper scheme X with a perfect obstruction theory, one can define virtual enumerative invariants by

$$\int_{[X]^{\text{vir}}} \alpha \in \mathbb{Q},$$

where $\alpha \in A^i(X)$. These intersection numbers are going to vanish if $i \neq \text{vd}$.

On the K-theoretic side, it was observed in [7, § 5.4] that a perfect obstruction theory $\mathbb{E} \rightarrow \mathbb{L}_X$ not only induces a virtual fundamental class, but also a *virtual structure sheaf*

$$\mathcal{O}_X^{\text{vir}} = [\mathbf{L}^* \mathcal{O}_{\mathfrak{C}}] \in K_0(X).$$

Its construction first appeared in [34, 18] in the context of dg-manifolds and in [7, 23] in the language of perfect obstruction theories. More recently, Thomas gave a description of $\mathcal{O}_X^{\text{vir}}$ in terms of the *K-theoretic Fulton class*, showing that it only depends on the K-theory class of \mathbb{E} [65, Cor. 4.5]. If $\pi: X \rightarrow \text{pt}$ is proper, one can use $\mathcal{O}_X^{\text{vir}}$ and K-theoretic pushforward $\pi_* = \chi(X, -)$ to define virtual invariants by

$$\chi^{\text{vir}}(X, V) = \chi(X, V \otimes \mathcal{O}_X^{\text{vir}}) \in K_0(\text{pt}) = \mathbb{Z}, \quad V \in K^0(X).$$

A virtual version [23, Cor. 3.6] of the Hirzebruch–Riemann–Roch theorem holds: one has

$$\chi^{\text{vir}}(X, V) = \int_{[X]^{\text{vir}}} \text{ch}(V) \cdot \text{td}(T_X^{\text{vir}}),$$

where, setting $E_i = (E^{-i})^\vee$, one defines the *virtual tangent bundle* of X by the formula

$$T_X^{\text{vir}} = \mathbb{E}^\vee = E_0 - E_1 \in K^0(X).$$

2.3. Torus representations and their weights. Let $\mathbf{T} = (\mathbb{C}^*)^g$ be an algebraic torus, with character lattice $\widehat{\mathbf{T}} = \text{Hom}(\mathbf{T}, \mathbb{C}^*) \cong \mathbb{Z}^g$. Let $K_0^{\mathbf{T}}(\text{pt})$ be the K-group of the category of \mathbf{T} -representations. Any finite dimensional \mathbf{T} -representation V splits as a sum of 1-dimensional representations called the *weights* of V . Each weight corresponds to a character $\mu \in \widehat{\mathbf{T}}$, and in turn each character corresponds to a monomial $t^\mu = t_1^{\mu_1} \cdots t_g^{\mu_g}$ in the coordinates of \mathbf{T} . The map

$$(2.1) \quad \text{tr}: K_0^{\mathbf{T}}(\text{pt}) \rightarrow \mathbb{Z}[t^\mu \mid \mu \in \widehat{\mathbf{T}}], \quad V \mapsto \text{tr}_V,$$

sending the class of a \mathbf{T} -module to its decomposition into weight spaces is a ring isomorphism, where tensor product on the left corresponds to the natural multiplication on the right. We will therefore sometimes identify a (virtual) \mathbf{T} -module with its character. Sometimes, to ease notation, we shall write $\text{tr}(V)$ instead of tr_V .

Example 2.3. Let $V = \sum_{\mu} t^{\mu}$ be a \mathbf{T} -module. Define $\Lambda_p^{\bullet} V = \sum_{i=0}^{\text{rk } V} p^i \Lambda^i V$ to be the *total wedge* of V . We shall write $\Lambda^{\bullet} V = \Lambda_{-1}^{\bullet} V$. As shown for instance in [52, Ex. 2.1.5], its trace satisfies

$$\text{tr}_{\Lambda_p^{\bullet} V} = \prod_{\mu} (1 - p t^{\mu}).$$

2.4. Virtual normal bundle and virtual tangent space. Let $\mathbf{T} = (\mathbb{C}^*)^g$ be an algebraic torus. If Y is a scheme carrying the trivial \mathbf{T} -action, any \mathbf{T} -equivariant coherent sheaf $B \in \text{Coh } Y$ admits a direct sum decomposition $B = \bigoplus_{\mu} B^{\mu}$ into eigensheaves, where $\mu \in \widehat{\mathbf{T}} \cong \mathbb{Z}^g$ ranges over the characters of \mathbf{T} . The \mathbf{T} -fixed part and the \mathbf{T} -moving part of B are defined as

$$B^{\text{fix}} = B^0, \quad B^{\text{mov}} = \bigoplus_{\mu \neq 0} B^{\mu}.$$

This definition extends to complexes of coherent sheaves.

If X is a scheme carrying a \mathbf{T} -action and a \mathbf{T} -equivariant perfect obstruction theory $\mathbb{E} \rightarrow \mathbb{L}_X$, and $Y \subset X^{\mathbf{T}}$ is a component of the fixed locus, then the *virtual normal bundle* of $Y \subset X$ is the complex

$$N_{Y/X}^{\text{vir}} = T_X^{\text{vir}}|_Y^{\text{mov}} = \mathbb{E}^{\vee}|_Y^{\text{mov}}.$$

If $p \in X^{\mathbf{T}}$ is a fixed point, the *virtual tangent space* of X at p is $T_p^{\text{vir}} = \mathbb{E}^{\vee}|_p \in K_0^{\mathbf{T}}(\text{pt})$.

2.5. Virtual localisation. A very useful tool to compute virtual K-theoretic invariants is the *virtual localisation theorem*. Its first version was proven in equivariant Chow theory in [27], and a K-theoretic version appeared in [23, § 7] and [54, § 3]. Suppose that X carries an action of an algebraic torus \mathbf{T} and a \mathbf{T} -equivariant perfect obstruction theory. Then $\mathcal{O}_X^{\text{vir}}$ is naturally an element of $K_0^{\mathbf{T}}(X)$ and if X is proper one can define $\chi^{\text{vir}}(X, -) = \chi(X, - \otimes \mathcal{O}_X^{\text{vir}})$ by means of the equivariant pushforward $\chi(X, -): K_0^{\mathbf{T}}(X) \rightarrow K_0^{\mathbf{T}}(\text{pt})$. The K-theoretic virtual localisation formula states that, for an arbitrary element $V \in K_{\mathbf{T}}^0(X)$ of the \mathbf{T} -equivariant K-theory, one has the identity

$$(2.2) \quad \chi^{\text{vir}}(X, V) = \chi^{\text{vir}}\left(X^{\mathbf{T}}, \frac{V|_{X^{\mathbf{T}}}}{\Lambda^{\bullet} N^{\text{vir}, \vee}}\right) \in K_0^{\mathbf{T}}(\text{pt}) \left[\frac{1}{1-t^{\mu}} \mid \mu \in \widehat{\mathbf{T}} \right],$$

where $X^{\mathbf{T}} \subset X$ is the \mathbf{T} -fixed locus, $N^{\text{vir}} \in K_{\mathbf{T}}^0(X^{\mathbf{T}})$ is K-theory class of the virtual normal bundle, i.e. of the \mathbf{T} -moving part of $T_X^{\text{vir}} = \mathbb{E}^{\vee}$ restricted to $X^{\mathbf{T}}$.

More than just being a powerful theorem, the localisation formula allows one to define invariants for *quasiprojective* \mathbf{T} -varieties, provided that they have proper \mathbf{T} -fixed locus: if this is the case, one defines $\chi^{\text{vir}}(X, V)$ to be the right hand side of (2.2). If X is proper, this definition coincides with the usual one thanks to the localisation theorem.

Remark 2.4. We thank Noah Arbesfeld for pointing out to us that for quasiprojective schemes, in contrast to the case of equivariant *cohomology*, one can directly define the equivariant Euler characteristic of a coherent sheaf via its character as a torus representation (without invoking virtual localisation), provided that the weight spaces of the sheaf are finite dimensional.

3. THE LOCAL QUOT SCHEME: CRITICAL AND EQUIVARIANT STRUCTURE

3.1. Overview. In this section we start working on the local Calabi–Yau 3-fold \mathbb{A}^3 . Fix integers $r \geq 1$ and $n \geq 0$. Our focus will be on the *local Quot scheme*

$$\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n),$$

whose points correspond to short exact sequences

$$0 \rightarrow S \rightarrow \mathcal{O}^{\oplus r} \rightarrow T \rightarrow 0$$

where T is a 0-dimensional $\mathcal{O}_{\mathbb{A}^3}$ -module with $\chi(T) = n$.

We shall use the following notation throughout.

Notation 3.1. If F is a locally free sheaf on a variety X , and $F \twoheadrightarrow T$ is a surjection onto a 0-dimensional sheaf of length n , with kernel $S \subset F$, we denote by

$$[S] \in \text{Quot}_X(F, n)$$

the corresponding point in the Quot scheme.

In this section, we will:

- recall from [4] the description of the Quot scheme as a critical locus (§ 3.2),
- describe a \mathbf{T} -action (for $\mathbf{T} = (\mathbb{C}^*)^3 \times (\mathbb{C}^*)^r$ a torus of dimension $3 + r$) on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, with isolated fixed locus consisting of direct sums of monomial ideals (§ 3.3),
- reinterpret the fixed locus $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$ in terms of colored partitions (§ 3.4),
- prove that the critical perfect obstruction theory on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ is \mathbf{T} -equivariant (Lemma 3.13), and that the induced \mathbf{T} -fixed obstruction theory on the fixed locus is trivial (Corollary 3.15).

The content of this section is the starting point for the definition (see § 4.3) of virtual invariants on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, as well as our construction (see § 5) of the *higher rank vertex formalism*.

3.2. The critical structure on the Quot scheme. Let V be an n -dimensional complex vector space. Consider the space $R_{r,n} = \text{Rep}_{(n,1)}(\tilde{\mathbb{L}}_3)$ of r -framed $(n, 1)$ -dimensional representations of the 3-loop quiver \mathbb{L}_3 , depicted in Figure 1. The notation “ $(n, 1)$ ” means that the main vertex (the one belonging to the 3-loop quiver, labelled “0” in the figure) carries a copy of V , whereas the framing vertex (labelled “ ∞ ”) carries a copy of \mathbb{C} .




FIGURE 1. The r -framed 3-loop quiver $\tilde{\mathbb{L}}_3$.

We have that $R_{r,n}$ is an affine space of dimension $3n^2 + rn$, with an explicit description as

$$\begin{aligned} R_{r,n} &= \{ (A_1, A_2, A_3, u_1, \dots, u_r) \mid A_j \in \text{End}(V), u_i \in V \} \\ &= \text{End}(V)^{\oplus 3} \oplus V^{\oplus r}. \end{aligned}$$

By [4, Prop. 2.4], there exists a stability parameter θ on the 3-loop quiver such that θ -stable framed representations $(A_1, A_2, A_3, u_1, \dots, u_r) \in R_{r,n}$ are precisely those satisfying the condition:

$$\text{the vectors } u_1, \dots, u_r \in V \text{ jointly generate } (A_1, A_2, A_3) \in \text{Rep}_n(\mathbb{L}_3).$$

Imposing this stability condition on $R_{r,n}$ we obtain an open subscheme

$$U_{r,n} \subset R_{r,n}$$

on which $\mathrm{GL}(V)$ acts freely by the rule

$$g \cdot (A_1, A_2, A_3, u_1, \dots, u_r) = (gA_1g^{-1}, gA_2g^{-1}, gA_3g^{-1}, gu_1, \dots, gu_r).$$

The quotient

$$(3.1) \quad \mathrm{Quot}_r^n = U_{r,n} / \mathrm{GL}(V)$$

is a smooth quasiprojective variety of dimension $2n^2 + rn$. In [4] the scheme Quot_r^n is referred to as the *non-commutative Quot scheme*, by analogy with the *non-commutative Hilbert scheme* [51], i.e. the moduli space of left ideals of codimension n in $\mathbb{C}\langle x_1, x_2, x_3 \rangle$ (which of course exists for an arbitrary number of free variables).

On $R_{r,n}$ one can define the function

$$h_n : R_{r,n} \rightarrow \mathbb{A}^1, \quad (A_1, A_2, A_3, u_1, \dots, u_r) \mapsto \mathrm{Tr} A_1[A_2, A_3],$$

induced by the superpotential $W = A_1[A_2, A_3]$ on the 3-loop quiver. Note that this function

- is symmetric under cyclic permutations of A_1, A_2 and A_3 , and
- does not touch the vectors u_i , which are only used to define its domain.

Moreover, $h_n|_{U_{r,n}}$ is $\mathrm{GL}(V)$ -invariant, and thus descends to a regular function

$$(3.2) \quad f_n : \mathrm{Quot}_r^n \rightarrow \mathbb{A}^1.$$

Proposition 3.2 ([4, Thm. 2.6]). *There is an identity of closed subschemes*

$$\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n) = \mathrm{crit}(f_n) \subset \mathrm{Quot}_r^n.$$

In particular, $\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ carries a symmetric perfect obstruction theory.

We use the notation $\mathrm{crit}(f)$ for the zero scheme $\{df=0\}$, for f a function on a smooth scheme. The embedding of the Quot scheme inside a non-commutative quiver model had appeared (conjecturally, and in a slightly different language) in the physics literature [19].

Every critical locus $\mathrm{crit}(f)$ has a canonical symmetric obstruction theory, determined by the Hessian complex attached to the function f . It will be referred to as the *critical obstruction theory* throughout. In the case of $Q = \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, this symmetric obstruction theory is the morphism

$$(3.3) \quad \begin{array}{ccc} \mathbb{E}_{\mathrm{crit}} = [T_{\mathrm{Quot}_r^n}|_Q & \xrightarrow{\mathrm{Hess}(f_n)} & \Omega_{\mathrm{Quot}_r^n}|_Q] \\ \phi \downarrow & (df_n)^\vee|_Q \downarrow & \downarrow \mathrm{id} \\ \mathbb{L}_Q = [\mathcal{I}/\mathcal{I}^2 & \xrightarrow{d} & \Omega_{\mathrm{Quot}_r^n}|_Q] \end{array}$$

in $\mathbf{D}^{[-1,0]}(Q)$, where we represented the truncated cotangent complex by means of the exterior derivative d constructed out of the ideal sheaf $\mathcal{I} \subset \mathcal{O}_{\mathrm{Quot}_r^n}$ of the inclusion $Q \hookrightarrow \mathrm{Quot}_r^n$.

Remark 3.3. As proved by Cazzaniga and the third author in [17], for any integer $m \geq 3$, the Quot scheme $\mathrm{Quot}_{\mathbb{A}^m}(\mathcal{O}^{\oplus r}, n)$ is canonically isomorphic to the moduli space of *framed sheaves* on \mathbb{P}^m , i.e. the moduli space of pairs (E, φ) where E is a torsion free sheaf on \mathbb{P}^m with Chern character $(r, 0, \dots, 0, -n)$ and $\varphi : E|_D \xrightarrow{\sim} \mathcal{O}_D^{\oplus r}$ is an isomorphism, for $D \subset \mathbb{P}^m$ a fixed hyperplane.

3.3. Torus actions on the local Quot scheme. In this section we define a torus action on the Quot scheme. Set

$$(3.4) \quad \mathbb{T}_1 = (\mathbb{C}^*)^3, \quad \mathbb{T}_2 = (\mathbb{C}^*)^r, \quad \mathbf{T} = \mathbb{T}_1 \times \mathbb{T}_2.$$

The torus \mathbb{T}_1 acts on \mathbb{A}^3 by the standard action

$$(3.5) \quad t \cdot x_i = t_i x_i,$$

and this action lifts to an action on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$. At the same time, the torus $\mathbb{T}_2 = (\mathbb{C}^*)^r$ acts on the Quot scheme by scaling the fibres of $\mathcal{O}^{\oplus r}$. Thus we obtain a \mathbf{T} -action on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$.

Remark 3.4. The fixed locus $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbb{T}_1}$ is proper. Indeed, a \mathbb{T}_1 -invariant surjection $\mathcal{O}^{\oplus r} \twoheadrightarrow T$ necessarily has the quotient T entirely supported at the origin $0 \in \mathbb{A}^3$. Hence

$$\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbb{T}_1} \hookrightarrow \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)_0$$

sits inside the *punctual Quot scheme* as a closed subscheme. But $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)_0$ is proper, since it is a fibre of the Quot-to-Chow morphism $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n) \rightarrow \text{Sym}^n \mathbb{A}^3$, which is a proper morphism.

We recall, verbatim from [4, Lemma 2.10], the description of the full \mathbf{T} -fixed locus induced by the product action on the local Quot scheme.

Lemma 3.5. *There is an isomorphism of schemes*

$$(3.6) \quad \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}} = \coprod_{n_1 + \dots + n_r = n} \prod_{i=1}^r \text{Hilb}^{n_i}(\mathbb{A}^3)^{\mathbb{T}_1}.$$

In particular, the \mathbf{T} -fixed locus is isolated and compact. Moreover, letting $\mathbf{T}_0 \subset \mathbf{T}$ be the subtorus defined by $t_1 t_2 t_3 = 1$, one has a scheme-theoretic identity

$$(3.7) \quad \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}_0} = \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}.$$

Proof. The main result proved by Bifet in [10] (in greater generality) implies that

$$(3.8) \quad \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}_2} = \coprod_{n_1 + \dots + n_r = n} \prod_{i=1}^r \text{Hilb}^{n_i}(\mathbb{A}^3).$$

The isomorphism (3.6) follows by taking \mathbb{T}_1 -invariants. Since $\text{Hilb}^k(\mathbb{A}^3)^{\mathbb{T}_1}$ is isolated (a disjoint union of reduced points, each corresponding to a monomial ideal of colength k), the first claim follows. Let now $\mathbb{T}_0 \subset \mathbb{T}_1$ be the subtorus defined by $t_1 t_2 t_3 = 1$, so that $\mathbf{T}_0 = \mathbb{T}_0 \times \mathbb{T}_2$. Equation (3.7) follows combining Equation (3.8) and the isomorphism $\text{Hilb}^k(\mathbb{A}^3)^{\mathbb{T}_1} \cong \text{Hilb}^k(\mathbb{A}^3)^{\mathbb{T}_0}$ proved in [8, Lemma 4.1]. \square

Remark 3.6. The \mathbf{T} -action on $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ just described can be seen as the restriction of a \mathbf{T} -action on the larger space Quot_r^n . Indeed, a torus element $\mathbf{t} = (t_1, t_2, t_3, w_1, \dots, w_r) \in \mathbf{T}$ acts on $(A_1, A_2, A_3, u_1, \dots, u_r) \in \text{Quot}_r^n$ by

$$(3.9) \quad \mathbf{t} \cdot P = (t_1 A_1, t_2 A_2, t_3 A_3, w_1 u_1, \dots, w_r u_r).$$

The critical locus $\text{crit}(f_n)$ is \mathbf{T} -invariant, and the induced action is precisely the one we described earlier in this section.

3.4. Combinatorial description of the T -fixed locus. The T -fixed locus $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^T$ is described purely in terms of r -colored plane partitions of size n , as we now explain.

We first recall the definition of a partition of arbitrary dimension.

Definition 3.7. Let $d \geq 1$ and $n \geq 0$ be integers. A $(d-1)$ -dimensional partition of n is a collection of n points $\mathcal{A} = \{\mathbf{a}_1, \dots, \mathbf{a}_n\} \subset \mathbb{Z}_{\geq 0}^d$ with the following property: if $\mathbf{a}_i = (a_{i1}, \dots, a_{id}) \in \mathcal{A}$, then whenever a point $\mathbf{y} = (y_1, \dots, y_d) \in \mathbb{Z}_{\geq 0}^d$ satisfies $0 \leq y_j \leq a_{ij}$ for all $j = 1, \dots, d$, one has $\mathbf{y} \in \mathcal{A}$. The integer $n = |\mathcal{A}|$ is called the *size* of the partition.

There is a bijective correspondence between the sets of

- $(d-1)$ -dimensional partitions of size n ,
- $(\mathbb{C}^*)^d$ -fixed points of $\text{Hilb}^n(\mathbb{A}^d)$, and
- monomial ideals $I \subset \mathbb{C}[x_1, \dots, x_d]$ of colength n .

We will be interested in the case $d = 3$, corresponding by definition to *plane partitions*. These can be visualised (cf. Figure 2) as configurations of n boxes stacked in the corner of a room (with gravity pointing in the $(-1, -1, -1)$ direction).




FIGURE 2. A plane partition of size $n = 16$.

Example 3.8. If \mathcal{A} is a $(d-1)$ -dimensional partition of size n as in Definition 3.7, the associated monomial ideal is

$$I_{\mathcal{A}} = \langle x_1^{i_1} \cdots x_d^{i_d} \mid (i_1, \dots, i_d) \in \mathbb{Z}_{\geq 0}^d \setminus \mathcal{A} \rangle \subset \mathbb{C}[x_1, \dots, x_d].$$

For instance, if $d = 3$, in the case of the plane partition pictured in Figure 2, the associated monomial ideal of colength 16 is generated by the monomials shaping the staircase of the partition, and is thus equal to

$$\langle x_3^4, x_1 x_3^2, x_1^2 x_3, x_1^5, x_1^2 x_2, x_1 x_2 x_3, x_2 x_3^2, x_1 x_2^3, x_2^3 x_3, x_2^4 \rangle \subset \mathbb{C}[x_1, x_2, x_3].$$

Here is an alternative definition of plane partitions.

Definition 3.9. A (finite) *plane partition* is a sequence $\pi = \{\pi_{ij} \mid i, j \geq 0\} \subset \mathbb{Z}_{\geq 0}$ such that $\pi_{ij} = 0$ for $i, j \gg 0$ and

$$\pi_{ij} \geq \pi_{i+1, j}, \quad \pi_{ij} \geq \pi_{i, j+1} \quad \text{for all } i, j \geq 0.$$

Definition 3.10. An r -colored *plane partition* is a tuple $\bar{\pi} = (\pi_1, \dots, \pi_r)$, where each π_i is a plane partition.

Denote by

$$|\pi| = \sum_{i, j \geq 0} \pi_{ij}$$

the *size* of a plane partition (i.e. the number n in Definition 3.7) and by $|\bar{\pi}| = \sum_{i=1}^r |\pi_i|$ the size of an r -colored plane partition.

In the light of Definition 3.9, the monomial ideal associated to a plane partition π is

$$I_\pi = \left\langle x_1^i x_2^j x_3^{\pi_{ij}} \mid i, j \geq 0 \right\rangle \subset \mathbb{C}[x_1, x_2, x_3].$$

It is clear that the colength of the ideal I_π is $|\pi|$.

Remark 3.11. A general plane partition may have infinite legs, each shaped by (i.e. asymptotic to) a standard (1-dimensional) partition, or Young diagram. We are not concerned with infinite plane partitions here, since we only deal with quotients $\mathcal{O}^{\oplus r} \rightarrow T$ with finite support.

Proposition 3.12. *There is a bijection between \mathbf{T} -fixed points $[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$ and r -colored plane partitions $\bar{\pi}$ of size n .*

Proof. For $r = 1$ this is well known: as we recalled above, monomial ideals $I \subset \mathbb{C}[x_1, x_2, x_3]$ are in bijective correspondence with plane partitions. Similarly, to each r -colored plane partition $\bar{\pi} = (\pi_1, \dots, \pi_r)$ there corresponds a subsheaf $S_{\bar{\pi}} = \bigoplus_{i=1}^r I_{\pi_i} \subset \mathcal{O}^{\oplus r}$. But these are the \mathbf{T} -fixed points of the Quot scheme by Lemma 3.5. \square

3.4.1. Computing the trace of a monomial ideal. Recall the map (2.1) sending a torus representation to its weight space decomposition. Consider the 3-dimensional torus \mathbb{T}_1 acting on the coordinate ring $R = \mathbb{C}[x_1, x_2, x_3]$ of \mathbb{A}^3 . Then we have

$$\text{tr}_R = \sum_{\square \in \mathbb{Z}_{\geq 0}^3} t^\square = \sum_{(i,j,k) \in \mathbb{Z}_{\geq 0}^3} t_1^i t_2^j t_3^k = \frac{1}{(1-t_1)(1-t_2)(1-t_3)}.$$

For a cyclic monomial ideal $\mathfrak{m}_{abc} = x_1^a x_2^b x_3^c \cdot R \subset R$, one has

$$\text{tr}_{\mathfrak{m}_{abc}} = \sum_{i \geq a} \sum_{j \geq b} \sum_{k \geq c} t_1^i t_2^j t_3^k = \frac{t_1^a t_2^b t_3^c}{(1-t_1)(1-t_2)(1-t_3)}.$$

More generally, for a monomial ideal $I_\pi \subset \mathbb{C}[x_1, x_2, x_3]$, one has

$$(3.10) \quad \text{tr}_{I_\pi} = \sum_{(i,j,k) \notin \pi} t_1^i t_2^j t_3^k.$$

These are the building blocks needed to compute tr_S for an arbitrary sheaf $S = \bigoplus_{i=1}^r I_{\pi_i}$ corresponding to a \mathbf{T} -fixed point $[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$.

3.5. The \mathbf{T} -fixed obstruction theory. Recall from [27, Prop. 1] that a torus equivariant obstruction theory on a scheme Y induces a canonical perfect obstruction theory, and hence a virtual fundamental class, on each component of the torus fixed locus. In this subsection we show that the reduced isolated locus

$$\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}} \hookrightarrow \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$$

carries the trivial \mathbf{T} -fixed perfect obstruction theory, so the induced virtual fundamental class agrees with the actual (0-dimensional) fundamental class.

We first need to check the equivariance (cf. Definition 2.2) of the critical obstruction theory \mathbb{E}_{crit} obtained in Proposition 3.2. In fact, this follows from the general fact that the critical obstruction theory on $\text{crit}(f) \subset Y$, for f a function on a smooth scheme Y , acted on by an algebraic torus \mathbf{T} , is naturally \mathbf{T} -equivariant as soon as f is \mathbf{T} -homogeneous. However, for the sake of completeness, we include a direct proof below for the case at hand.

Lemma 3.13. *The critical obstruction theory on $Q = \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ is \mathbf{T} -equivariant.*

Proof. We start with two observations:

- (1) The potential $f_n = \text{Tr } A_1[A_2, A_3]$ recalled in (3.2) is homogeneous (of degree 3) in the matrix coordinates of the non-commutative Quot scheme.
- (2) The potential f_n satisfies the relation

$$(3.11) \quad f_n(t \cdot P) = t_1 t_2 t_3 \cdot f_n(P)$$

for every $t = (t_1, t_2, t_3) \in \mathbb{T}_1$ and $P \in \text{Quot}_r^n$.

Fix a point $p \in Q = \text{crit}(f_n) \subset \text{Quot}_r^n$. Then, setting $N = 2n^2 + rn = \dim \text{Quot}_r^n$, let x_1, \dots, x_N be local holomorphic coordinates of Quot_r^n around p . Let the torus \mathbf{T} act on these coordinates as prescribed by Equation (3.9), i.e. t_1 (resp. t_2 and t_3) rescales each x_k corresponding to the entries of the first (resp. second and third) matrix, and w_l rescales the coordinates of the vector u_l , for $l = 1, \dots, r$. Formally, for a matrix coordinate x_k , we set

$$(t_1, t_2, t_3, w_1, \dots, w_r) \cdot x_k = t_{\ell(k)} x_k$$

where $\ell(k) \in \{1, 2, 3\}$ depends on whether x_k comes from an entry of A_1 , A_2 or A_3 . We also have to prescribe an action on tangent vectors and 1-forms. For a matrix coordinate x_k , we set

$$(3.12) \quad \begin{aligned} (t_1, t_2, t_3, w_1, \dots, w_r) \cdot \frac{\partial}{\partial x_k} &= \frac{t_1 t_2 t_3}{t_{\ell(k)}} \frac{\partial}{\partial x_k} \\ (t_1, t_2, t_3, w_1, \dots, w_r) \cdot dx_k &= t_{\ell(k)} dx_k. \end{aligned}$$

If x_k comes from a vector component of the l -th vector, we set

$$(3.13) \quad \begin{aligned} (t_1, t_2, t_3, w_1, \dots, w_r) \cdot \frac{\partial}{\partial x_k} &= w_l^{-1} \frac{\partial}{\partial x_k} \\ (t_1, t_2, t_3, w_1, \dots, w_r) \cdot dx_k &= w_l dx_k. \end{aligned}$$

However, the \mathbb{T}_2 -action (3.13) will be invisible in the Hessian since the function f_n does not touch the vectors.

The Hessian can be seen as a section

$$\text{Hess}(f_n) \in \Gamma\left(Q, T_{\text{Quot}_r^n}^*|_Q \otimes T_{\text{Quot}_r^n}^*|_Q\right).$$

In checking the equivariance relation

$$\mathbf{t} \cdot \text{Hess}(f_n)(\xi) = \text{Hess}(f_n)(\mathbf{t} \cdot \xi), \quad \mathbf{t} \in \mathbf{T},$$

we may ignore local coordinates x_k corresponding to vector entries, because the Hessian is automatically equivariant in these coordinates (equivariance translates into the identity $0 = 0$).

So, let us fix an x_k coming from one of the matrices. The (i, j) -component of the Hessian applied to $\partial/\partial x_k$ is given by

$$\text{Hess}_{ij}(f_n) \left(\frac{\partial}{\partial x_k} \right) = \frac{\partial^2 f_n}{\partial x_i \partial x_j}(x_1, \dots, x_N) dx_j.$$

This will vanish unless $k \in \{i, j\}$. Without loss of generality we may assume $k = i$. In this case we obtain, up to a sign convention,

$$(3.14) \quad \text{Hess}_{ij}(f_n) \left(\frac{t_1 t_2 t_3}{t_{\ell(k)}} \frac{\partial}{\partial x_k} \right) = \frac{t_1 t_2 t_3}{t_{\ell(k)}} \frac{\partial^2 f_n}{\partial x_k \partial x_j}(x_1, \dots, x_N) dx_j.$$

On the other hand, combining the observations (1) and (2) with (3.12), we obtain

$$\begin{aligned} \mathbf{t} \cdot \text{Hess}_{ij}(f_n) \left(\frac{\partial}{\partial x_k} \right) &= \frac{\partial^2 f_n}{(\partial t_{\ell(k)} x_k)(\partial t_{\ell(j)} x_j)} (t_{\ell(1)} x_1, \dots, t_{\ell(N)} x_N) t_{\ell(j)} dx_j \\ &= \frac{t_1 t_2 t_3}{t_{\ell(k)} t_{\ell(j)}} t_{\ell(j)} \frac{\partial^2 f_n}{\partial x_k \partial x_j} (x_1, \dots, x_N) dx_j, \end{aligned}$$

which agrees with the right hand side of Equation (3.14). Thus we conclude that the Hessian complex is \mathbf{T} -equivariant, as well as the morphism (3.3) to the cotangent complex. This finishes the proof. \square

The property (3.11) of f_n exhibits the differential df_n as a GL_3 -equivariant section

$$df_n \otimes \mathbf{t}^{-1} : \mathcal{O}_{\text{Quot}_r^n} \rightarrow \Omega_{\text{Quot}_r^n} \otimes \mathbf{t}^{-1},$$

where $\mathbf{t}^{-1} = (t_1 t_2 t_3)^{-1}$ is the determinant representation of $\mathbb{C}^3 = \bigoplus_{1 \leq i \leq 3} t_i^{-1} \cdot \mathbb{C}$. Therefore, explicitly, the morphism in $\mathbf{D}^{[-1,0]}(\text{Coh}_Q^{\mathbf{T}})$ lifting the critical obstruction theory (3.3) is

$$(3.15) \quad \begin{array}{ccc} [\mathbf{t} \otimes T_{\text{Quot}_r^n}|_Q & \xrightarrow{\text{Hess}(f_n)} & \Omega_{\text{Quot}_r^n}|_Q \\ \downarrow (df_n)^\vee|_Q & & \downarrow \text{id} \\ [\mathcal{I}/\mathcal{I}^2 & \xrightarrow{d} & \Omega_{\text{Quot}_r^n}|_Q \end{array}$$

so that, in particular, the equivariant K-theory class of the virtual tangent bundle attached to the (equivariant) perfect obstruction theory (3.15) is

$$(3.16) \quad T_Q^{\text{vir}} = T_{\text{Quot}_r^n}|_Q - \Omega_{\text{Quot}_r^n}|_Q \otimes \mathbf{t}^{-1} \in K_{\mathbf{T}}^0(Q).$$

This fact will be recalled and used in Propositions 5.2 and 9.3.

Lemma 3.13 implies the existence of a “ \mathbf{T} -fixed” obstruction theory

$$(3.17) \quad \mathbb{E}_{\text{crit}}|_{\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}}^{\text{fix}} \rightarrow \mathbb{L}_{\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}}$$

on the fixed locus of the Quot scheme. We proved in Lemma 3.5 that this fixed locus is 0-dimensional, isolated and reduced. The next result will imply that the virtual fundamental class induced by (3.17) on the fixed locus agrees with the actual fundamental class.

Proposition 3.14. *Let $[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$ be a torus fixed point. The deformations and obstructions of $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ at $[S]$ are entirely \mathbf{T} -movable. In particular, the virtual tangent space at $[S]$ can be written as*

$$(3.18) \quad T_S^{\text{vir}} = \mathbb{E}_{\text{crit}}^{\vee}|_{[S]}^{\text{mov}} = T_Q|_{[S]} - \Omega_Q|_{[S]} \otimes \mathbf{t}^{-1} \in K_{\mathbf{T}}^0(\text{pt}).$$

Proof. The perfect obstruction theory \mathbb{E}_{crit} on $Q = \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, made explicit in Diagram (3.15), satisfies $\mathbb{E}_{\text{crit}} \cong \mathbb{E}_{\text{crit}}^{\vee}[1] \otimes \mathbf{t}$. Its equivariant K-theory class is therefore

$$\mathbb{E}_{\text{crit}} = \Omega_Q - T_Q \otimes \mathbf{t} \in K_0^{\mathbf{T}}(Q).$$

We know by Equation (3.7) in Lemma 3.5 that no power of \mathbf{t} is a weight of $T_Q|_p$ for any fixed point $p \in Q^{\mathbf{T}}$, which implies that

$$(3.19) \quad (T_Q|_p \otimes \mathbf{t})^{\text{fix}} = 0, \quad \Omega_Q|_p^{\text{fix}} = 0.$$

The claim follows. \square

Corollary 3.15. *There is an identity*

$$[\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathrm{T}}]^{\mathrm{vir}} = [\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathrm{T}}] \in A_0^{\mathrm{T}}(\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathrm{T}}).$$

4. INVARIANTS ATTACHED TO THE LOCAL QUOT SCHEME

In this section we introduce cohomological and K-theoretic DT invariants of \mathbb{A}^3 , the main object of study of this paper, starting from the Quot scheme $\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ studied in the previous section. We first need to introduce some notation and terminology.

4.1. Some notation. Recall the tori $\mathbb{T}_1 = (\mathbb{C}^*)^3$ and $\mathbb{T}_2 = (\mathbb{C}^*)^r$ from (3.4). We let t_1, t_2, t_3 and w_1, \dots, w_r be the generators of the representation rings $K_{\mathbb{T}_1}^0(\mathrm{pt})$ and $K_{\mathbb{T}_2}^0(\mathrm{pt})$, respectively. Then one can write the equivariant cohomology rings of \mathbb{T}_1 and \mathbb{T}_2 as

$$H_{\mathbb{T}_1}^* = \mathbb{Q}[s_1, s_2, s_3], \quad H_{\mathbb{T}_2}^* = \mathbb{Q}[v_1, \dots, v_r],$$

where $s_i = c_1^{\mathbb{T}_1}(t_i)$ and $v_j = c_1^{\mathbb{T}_2}(w_j)$. For a virtual \mathbb{T} -module $V \in K_{\mathbb{T}}^0(\mathrm{pt})$, we let

$$\mathrm{tr}_V \in \mathbb{Z}((t_1, t_2, t_3, w_1, \dots, w_r))$$

denote its character, i.e. its decomposition into weight spaces. We denote by $\overline{(\cdot)}$ the involution defined on $\mathbb{Z}((t_1, t_2, t_3, w_1, \dots, w_r))$ by

$$\overline{P}(t_1, t_2, t_3, w_1, \dots, w_r) = P(t_1^{-1}, t_2^{-1}, t_3^{-1}, w_1^{-1}, \dots, w_r^{-1}).$$

4.1.1. Twisted virtual structure sheaf. For any scheme X endowed with a perfect obstruction theory $\mathbb{E} \rightarrow \mathbb{L}_X$, define as in [23, Def. 3.12], the *virtual canonical bundle*

$$\mathcal{K}_{X, \mathrm{vir}} = \det \mathbb{E} = \det(T_X^{\mathrm{vir}})^{\vee}.$$

This is just $\det E^0 \otimes (\det E^{-1})^{\vee}$ if $\mathbb{E} = E^0 - E^{-1} \in K^0(X)$. We will simply write $\mathcal{K}_{\mathrm{vir}}$ when X is clear from the context.

Lemma 4.1. *Let A be a smooth variety equipped with a regular function $f: A \rightarrow \mathbb{A}^1$, and let $X = \mathrm{crit}(f) \subset A$ be the critical locus of f , with its critical (symmetric) perfect obstruction theory $\mathbb{E}_{\mathrm{crit}} \rightarrow \mathbb{L}_X$. Then $\mathcal{K}_{X, \mathrm{vir}} \in \mathrm{Pic}(X)$ admits a square root, i.e. there exists a line bundle*

$$\mathcal{K}_{X, \mathrm{vir}}^{\frac{1}{2}} \in \mathrm{Pic}(X)$$

whose second tensor power equals $\det \mathbb{E}_{\mathrm{crit}}$.

Proof. The K-theory class of the critical perfect obstruction theory is

$$\mathbb{E}_{\mathrm{crit}} = \Omega_A|_X - T_A|_X,$$

and by definition one has

$$\mathcal{K}_{X, \mathrm{vir}} = \frac{\det \Omega_A|_X}{\det T_A|_X} = \frac{\det \Omega_A|_X}{(\det \Omega_A|_X)^{-1}} = K_A|_X \otimes K_A|_X. \quad \square$$

Let X be a scheme endowed with a perfect obstruction theory, and let $\mathcal{O}_X^{\mathrm{vir}} \in K_0(X)$ be the induced virtual structure sheaf. Assume the virtual canonical bundle admits a square root. Following [49], we define the *twisted* (or *symmetrised*) *virtual structure sheaf* as

$$\widehat{\mathcal{O}}_X^{\mathrm{vir}} = \mathcal{O}_X^{\mathrm{vir}} \otimes \mathcal{K}_{X, \mathrm{vir}}^{\frac{1}{2}}.$$

In case X carries a torus action, we will see in Remark 4.2 that $\widehat{\mathcal{O}}_X^{\mathrm{vir}}$ acquires a canonical weight.

4.2. Classical enumerative invariants. Degree 0 Donaldson–Thomas invariants of various flavours have been computed in [38, 39, 8, 28, 45, 59, 6, 15]. This paper is concerned with the general theory of *Quot schemes*, hence in the (virtual) enumeration of 0-dimensional quotients of locally free sheaves on 3-folds.

The naive (topological) Euler characteristic of the Quot scheme $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ is computed via the Gholampour–Kool formula [25, Prop. 2.3]

$$\sum_{n \geq 0} e(\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)) q^n = M(q)^r,$$

where $M(q) = \prod_{m \geq 1} (1 - q^m)^{-m}$ is the MacMahon function, the generating function for the number of plane partitions of non-negative integers. On the other hand, the Behrend weighted Euler characteristic of the Quot scheme can be computed via the formula

$$(4.1) \quad \sum_{n \geq 0} e_{\text{vir}}(\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)) q^n = M((-1)^r q)^r,$$

proved in [4, Cor. 2.8]. For a complex scheme Z of finite type over \mathbb{C} , we have set $e_{\text{vir}}(Z) = e(Z, \nu_Z)$, where ν_Z is Behrend’s constructible function [5]. See [4, Thm. A] for a proof of the analogue of (4.1) for an arbitrary pair (Y, F) consisting of a smooth 3-fold Y along with a locally free sheaf F on it. It was shown by Toda [67] that, on a projective Calabi–Yau 3-fold Y , the wall-crossing factor in the higher rank DT/PT correspondence is precisely $M((-1)^r q)^{r e(Y)}$. The relationship between Toda’s wall-crossing formula [67] and the Gholampour–Kool’s formula for Euler characteristics of Quot schemes on 3-folds [25] was clarified in [4] via a Hall algebra argument.

4.3. Virtual invariants of the Quot scheme. The scheme $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ is not proper, but carries a torus action with proper fixed locus. Thus we may define virtual invariants via equivariant residues, by setting

$$(4.2) \quad \int_{[\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)]^{\text{vir}}} 1 := \sum_{[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}} \frac{1}{e_{\mathbf{T}}(T_S^{\text{vir}})} \in \mathbb{Q}((s, v)),$$

where $s = (s_1, s_2, s_3)$ and $v = (v_1, \dots, v_r)$ are the equivariant parameters of the torus \mathbf{T} and T_S^{vir} is the virtual tangent space (3.18). The sum runs over all \mathbf{T} -fixed points $[S]$, which are isolated, reduced and with the trivial perfect obstruction theory induced from the critical obstruction theory on the Quot scheme (cf. Corollary 3.15). We refer to these invariants as (degree 0) *cohomological rank r DT invariants*, as they take value in (an extension of) the fraction field $\mathbb{Q}(s, v)$ of the \mathbf{T} -equivariant cohomology ring $H_{\mathbf{T}}^*$. We will study their generating function

$$(4.3) \quad \text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, v) = \sum_{n \geq 0} q^n \int_{[\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)]^{\text{vir}}} 1 \in \mathbb{Q}((s, v))[[q]]$$

in § 7. On the other hand, K-theoretic invariants arise as natural refinements of their cohomological counterpart. Naively, one would like to study the virtual holomorphic Euler characteristic

$$\chi(\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \mathcal{O}^{\text{vir}}) = \sum_{[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}} \text{tr} \left(\frac{1}{\Lambda^\bullet T_S^{\text{vir}, \vee}} \right) \in \mathbb{Z}((t, w)),$$

where $t = (t_1, t_2, t_3)$, $w = (w_1, \dots, w_r)$, and via the trace map tr we identify a (possibly infinite-dimensional) virtual \mathbf{T} -module with its decomposition into weight spaces. It turns out that guessing a closed formula for these invariants is incredibly difficult and, after all, not what one

should look at. Instead, Nekrasov–Okounkov [49] teach us that we should focus our attention on

$$(4.4) \quad \chi(\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \widehat{\mathcal{O}}^{\mathrm{vir}}) = \sum_{[S] \in \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathrm{T}}} \mathrm{tr} \left(\frac{\mathcal{K}_{\mathrm{vir}}^{\frac{1}{2}}}{\Lambda \bullet T_S^{\mathrm{vir}, V}} \right) \in \mathbb{Z}((t, (t_1 t_2 t_3)^{\frac{1}{2}}, w)),^2$$

where the *twisted virtual structure sheaf* $\widehat{\mathcal{O}}^{\mathrm{vir}}$ is defined in § 4.1.1 — a square root of the virtual canonical bundle exists by Lemma 4.1 and Proposition 3.2. The generating function of rank r K-theoretic DT invariants

$$(4.5) \quad \mathrm{DT}_r^K(\mathbb{A}^3, q, t, w) = \sum_{n \geq 0} q^n \chi(\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \widehat{\mathcal{O}}^{\mathrm{vir}}) \in \mathbb{Z}((t, (t_1 t_2 t_3)^{\frac{1}{2}}, w))[[q]]$$

will be studied in § 6.

Remark 4.2. To be precise, we should replace the torus \mathbf{T} with its double cover \mathbf{T}_t , the minimal cover of \mathbf{T} where the character $t^{-1/2}$ is defined, as in [49, § 7.1.2]. Then $\mathcal{K}_{\mathrm{vir}}^{1/2}$ is a \mathbf{T}_t -equivariant sheaf with character $t^{-(\dim \mathrm{Quot}_r^{\mathrm{pt}})/2}$. To ease the notation, we keep denoting the torus acting as \mathbf{T} .

Remark 4.3. As remarked in [49, 1], choices of square roots of $\mathcal{K}_{\mathrm{vir}}$ differ by a 2-torsion element in the Picard group, which implies that $\chi(\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \widehat{\mathcal{O}}^{\mathrm{vir}})$ does not depend on such choices of square roots. Thus there is no ambiguity in (4.5).

5. HIGHER RANK VERTEX ON THE LOCAL QUOT SCHEME

5.1. The virtual tangent space of the local Quot scheme. By Lemma 3.5, we can represent the sheaf corresponding to a \mathbf{T} -fixed point

$$[S] \in \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathrm{T}}$$

as a direct sum of ideal sheaves

$$S = \bigoplus_{\alpha=1}^r \mathcal{I}_{Z_\alpha} \subset \mathcal{O}^{\oplus r},$$

with $Z_\alpha \subset \mathbb{A}^3$ a finite subscheme of length n_α supported at the origin, and satisfying $n = \sum_{1 \leq \alpha \leq r} n_\alpha$. In this subsection we derive a formula for the character of

$$\chi(\mathcal{O}^{\oplus r}, \mathcal{O}^{\oplus r}) - \chi(S, S) \in K_{\mathbf{T}}^0(\mathrm{pt}),$$

where for F and G in $K_{\mathbf{T}}^0(\mathrm{pt})$, we set

$$\chi(F, G) = \mathbf{R} \mathrm{Hom}(F, G) = \sum_{i \geq 0} (-1)^i \mathrm{Ext}^i(F, G).$$

Our method follows the approach of [44, § 4.7]. Moreover, we show in Proposition 5.2 that such character agrees with the virtual tangent space T_S^{vir} induced by the critical obstruction theory.

Let Q_α be the \mathbf{T}_1 -character of the α -summand of $\mathcal{O}^{\oplus r}/S$, i.e. (cf. Equation (3.10))

$$Q_\alpha = \mathrm{tr}_{\mathcal{O}_{Z_\alpha}} = \sum_{(i,j,k) \in \pi_\alpha} t_1^i t_2^j t_3^k,$$

²In theory all equivariant weights could appear with half powers. However, by Remark 4.2, $(t_1 t_2 t_3)^{\frac{1}{2}}$ is the only one that truly appears.

where $\pi_\alpha \subset \mathbb{Z}_{\geq 0}^3$ is the plane partition corresponding to the monomial ideal $\mathcal{I}_{Z_\alpha} \subset R = \mathbb{C}[x, y, z]$. Let $P_\alpha(t_1, t_2, t_3)$ be the Poincaré polynomial of \mathcal{I}_{Z_α} . This can be computed via a \mathbb{T}_1 -equivariant free resolution

$$0 \rightarrow E_{\alpha,s} \rightarrow \cdots \rightarrow E_{\alpha,1} \rightarrow E_{\alpha,0} \rightarrow \mathcal{I}_{Z_\alpha} \rightarrow 0.$$

Writing

$$E_{\alpha,i} = \bigoplus_j R(d_{\alpha,ij}), \quad d_{\alpha,ij} \in \mathbb{Z}^3,$$

one has, independently of the chosen resolution, the formula

$$P_\alpha(t_1, t_2, t_3) = \sum_{i,j} (-1)^i t^{d_{\alpha,ij}}.$$

By [44, § 4.7] we know that there is an identity

$$(5.1) \quad Q_\alpha = \frac{1 + P_\alpha}{(1-t_1)(1-t_2)(1-t_3)}.$$

For each $1 \leq \alpha, \beta \leq r$, we can compute

$$\begin{aligned} \chi(\mathcal{I}_{Z_\alpha}, \mathcal{I}_{Z_\beta}) &= \sum_{i,j,k,l} (-1)^{i+k} \operatorname{Hom}_R(R(d_{\alpha,ij}), R(d_{\beta,kl})) \\ &= \sum_{i,j,k,l} (-1)^{i+k} R(d_{\beta,kl} - d_{\alpha,ij}), \end{aligned}$$

which immediately yields the identity

$$\operatorname{tr}_{\chi(\mathcal{I}_{Z_\alpha}, \mathcal{I}_{Z_\beta})} = \frac{P_\beta \bar{P}_\alpha}{(1-t_1)(1-t_2)(1-t_3)} \in \mathbb{Z}((t_1, t_2, t_3)).$$

It follows that, as a \mathbf{T} -representation, one has

$$\begin{aligned} \chi(S, S) &= \chi \left(\sum_\alpha w_\alpha \otimes \mathcal{I}_{Z_\alpha}, \sum_\beta w_\beta \otimes \mathcal{I}_{Z_\beta} \right) \\ &= \sum_{1 \leq \alpha, \beta \leq r} \chi(w_\alpha \otimes \mathcal{I}_{Z_\alpha}, w_\beta \otimes \mathcal{I}_{Z_\beta}), \end{aligned}$$

which yields

$$(5.2) \quad \operatorname{tr}_{\chi(S,S)} = \sum_{1 \leq \alpha, \beta \leq r} \frac{w_\alpha^{-1} w_\beta \cdot P_\beta \bar{P}_\alpha}{(1-t_1)(1-t_2)(1-t_3)}.$$

On the other hand,

$$(5.3) \quad \operatorname{tr}_{\chi(\mathcal{O}^{\oplus r}, \mathcal{O}^{\oplus r})} = \sum_{1 \leq \alpha, \beta \leq r} \frac{w_\alpha^{-1} w_\beta}{(1-t_1)(1-t_2)(1-t_3)}.$$

Combining Formulae (5.2) and (5.3) with Formula (5.1) yields the following result.

Proposition 5.1. *Let $[S] \in \operatorname{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$ be a torus fixed point. There is an identity*

$$(5.4) \quad \operatorname{tr}_{\chi(\mathcal{O}^{\oplus r}, \mathcal{O}^{\oplus r}) - \chi(S,S)} = \sum_{1 \leq \alpha, \beta \leq r} w_\alpha^{-1} w_\beta \left(Q_\beta - \frac{\bar{Q}_\alpha}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} Q_\beta \bar{Q}_\alpha \right)$$

in $\mathbb{Z}((t_1, t_2, t_3, w_1, \dots, w_r))$.

For every \mathbf{T} -fixed point $[S]$ we define associated *vertex* terms

$$(5.5) \quad V_{ij} = w_i^{-1} w_j \left(Q_j - \frac{\overline{Q}_i}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} Q_j \overline{Q}_i \right)$$

for every $i, j = 1, \dots, r$. It is immediate to see that for $r = 1$ (forcing $i = j$) we recover the vertex formalism developed in [44].

Proposition 5.1 can then be restated as

$$\mathrm{tr}_{\chi(\mathcal{O}^{\oplus r}, \mathcal{O}^{\oplus r})} - \mathrm{tr}_{\chi(S, S)} = \sum_{1 \leq i, j \leq r} V_{ij}.$$

We now relate this to the virtual tangent space (cf. § 2.4) T_S^{vir} of a point $[S] \in \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$.

Proposition 5.2. *Let $[S] \in \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$ be a \mathbf{T} -fixed point. Let $T_S^{\mathrm{vir}} = \mathbb{E}_{\mathrm{crit}|[S]}^{\vee}$ be the virtual tangent space induced by the \mathbf{T} -equivariant critical obstruction theory. Then there are identities*

$$T_S^{\mathrm{vir}} = \chi(\mathcal{O}^{\oplus r}, \mathcal{O}^{\oplus r}) - \chi(S, S) = \sum_{1 \leq i, j \leq r} V_{ij} \in K_0^{\mathbf{T}}(\mathrm{pt}).$$

Proof. Let Quot_r^n be the non-commutative Quot scheme defined in (3.1). The superpotential $f_n: \mathrm{Quot}_r^n \rightarrow \mathbb{A}^1$ defined in (3.2) is equivariant with respect to the character $(t_1, t_2, t_3) \mapsto t_1 t_2 t_3$, so it gives rise to a GL_3 -equivariant section

$$(5.6) \quad \mathcal{O}_{\mathrm{Quot}_r^n} \xrightarrow{df_n \otimes \mathfrak{t}^{-1}} \Omega_{\mathrm{Quot}_r^n} \otimes \mathfrak{t}^{-1}$$

where, starting from the representation $\mathbb{C}^3 = \bigoplus_{1 \leq i \leq 3} t_i^{-1} \cdot \mathbb{C} \in K_{\mathbb{T}_1}^0(\mathrm{pt})$, we have set

$$\mathfrak{t}^{-1} = \det \mathbb{C}^3 = (t_1 t_2 t_3)^{-1}.$$

Here, and throughout this proof, we are identifying a representation with its own character via the isomorphism (2.1). The zero locus of the section (5.6) is our Quot scheme

$$Q = \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n),$$

endowed with the \mathbf{T} -action described in § 3.3. According to Equation (3.16), the virtual tangent space computed with respect to the critical \mathbf{T} -equivariant obstruction theory on Q is

$$(5.7) \quad T_S^{\mathrm{vir}} = \left(T_{\mathrm{Quot}_r^n} - \Omega_{\mathrm{Quot}_r^n} \otimes \mathfrak{t}^{-1} \right) \Big|_{[S]}.$$

The tangent space to the smooth scheme Quot_r^n can be written, around a point $S \hookrightarrow \mathcal{O}^{\oplus r} \twoheadrightarrow V$, as

$$(5.8) \quad T_{\mathrm{Quot}_r^n} \Big|_{[S]} = (\mathbb{C}^3 - 1) \otimes (\overline{V} \otimes V) + \bigoplus_{\alpha=1}^r \mathrm{Hom}(w_{\alpha} \mathbb{C}, V),$$

where

$$\bigoplus_{\alpha=1}^r \mathrm{Hom}(w_{\alpha} \mathbb{C}, V) = \bigoplus_{\alpha=1}^r w_{\alpha}^{-1} \otimes V$$

represents the r framings on the 3-loop quiver. Let V be written as a direct sum of structure sheaves

$$V = \bigoplus_{\alpha=1}^r \mathcal{O}_{Z_{\alpha}},$$

where the α -summand has \mathbf{T} -character $w_\alpha Q_\alpha$. Substituting

$$\mathbb{C}^3 - 1 = t_1^{-1} + t_2^{-1} + t_3^{-1} - 1 = \frac{t_1 t_2 + t_1 t_3 + t_2 t_3 - t_1 t_2 t_3}{t_1 t_2 t_3}$$

$$V = \sum_{\alpha=1}^r w_\alpha Q_\alpha$$

into Formula (5.8) yields

$$T_{\text{Quot}_r^n}|_{[S]} = \frac{t_1 t_2 + t_1 t_3 + t_2 t_3 - t_1 t_2 t_3}{t_1 t_2 t_3} \sum_{1 \leq \alpha, \beta \leq r} w_\alpha^{-1} w_\beta \overline{Q}_\alpha Q_\beta + \sum_{1 \leq \alpha, \beta \leq r} w_\alpha^{-1} w_\beta Q_\beta,$$

and hence

$$\left(\Omega_{\text{Quot}_r^n} \otimes \mathfrak{t}^{-1} \right) \Big|_{[S]} = \frac{t_1 + t_2 + t_3 - 1}{t_1 t_2 t_3} \sum_{1 \leq \alpha, \beta \leq r} w_\alpha^{-1} w_\beta \overline{Q}_\alpha Q_\beta + \sum_{1 \leq \alpha, \beta \leq r} w_\alpha^{-1} w_\beta \frac{\overline{Q}_\alpha}{t_1 t_2 t_3},$$

which by Formula (5.7) yields

$$T_S^{\text{vir}} = \sum_{1 \leq \alpha, \beta \leq r} w_\alpha^{-1} w_\beta \left(Q_\beta - \frac{\overline{Q}_\alpha}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} \overline{Q}_\alpha Q_\beta \right).$$

The right hand side is shown to be equal to $\chi(\mathcal{O}^{\oplus r}, \mathcal{O}^{\oplus r}) - \chi(S, S)$ in Proposition 5.1. \square

5.2. A small variation of the vertex formalism. All locally free sheaves on \mathbb{A}^3 are trivial, but this is not true equivariantly. For example, we have $K_{\mathbb{A}^3} = \mathcal{O}_{\mathbb{A}^3} \otimes t_1 t_2 t_3 \in K_{\mathbb{T}_1}^0(\mathbb{A}^3)$, even though the ordinary canonical bundle is trivial. Consider

$$(5.9) \quad F = \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{A}^3} \otimes \lambda_i \in K_{\mathbb{T}_1}^0(\mathbb{A}^3)$$

where $\lambda = (\lambda_i)_i$ are weights of the \mathbb{T}_1 -action, i.e. monomials in the representation ring of \mathbb{T}_1 . Let $[S] \in \text{Quot}_{\mathbb{A}^3}(F, n)^{\mathbf{T}}$ be a \mathbf{T} -fixed point. It decomposes as $S = \bigoplus_{i=1}^r \mathcal{I}_{Z_i} \otimes \lambda_i \in K_{\mathbb{T}_1}^0(\mathbb{A}^3)$, where the weights λ_i are naturally inherited from F . This generalises the discussion in § 5.1, which can be recovered by setting all weights λ_i to be trivial. Just as in Proposition 5.2, we can compute

$$T_{S, \lambda}^{\text{vir}} = \chi(F, F) - \chi(S, S) \in K_{\mathbf{T}}^0(\text{pt}).$$

We find

$$T_{S, \lambda}^{\text{vir}} = \chi \left(\bigoplus_{i=1}^r \mathcal{O}_{\mathbb{A}^3} \otimes \lambda_i w_i, \bigoplus_{j=1}^r \mathcal{O}_{\mathbb{A}^3} \otimes \lambda_j w_j \right) - \chi \left(\bigoplus_{i=1}^r \mathcal{I}_{Z_i} \otimes \lambda_i w_i, \bigoplus_{i=1}^r \mathcal{I}_{Z_i} \otimes \lambda_i w_i \right).$$

Therefore we derive the same expression for the vertex formalism as before, just substituting w_i with $\lambda_i w_i$. This variation will be crucial in § 9.4 where we identify the Quot scheme of the local model with the restriction of $\text{Quot}_X(F, n)$ to an open toric chart, with X a toric projective 3-fold and F a \mathbb{T}_1 -equivariant exceptional locally free sheaf.

Define, for $\lambda = (\lambda_1, \dots, \lambda_r)$ as above and F as in (5.9), the equivariant integral

$$(5.10) \quad \int_{[\text{Quot}_{\mathbb{A}^3}(F, n)]^{\text{vir}}} 1 := \sum_{[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}} \frac{1}{e^{\mathbf{T}}(T_{S, \lambda}^{\text{vir}})} \in \mathbb{Q}((s, v)),$$

and let

$$(5.11) \quad \text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, v)_{\lambda} = \sum_{n \geq 0} q^n \int_{[\text{Quot}_{\mathbb{A}^3}(F, n)]^{\text{vir}}} 1$$

be the generating function of the invariants (5.10). We shall see (cf. Corollary 7.4) that this expression does not depend on the equivariant weights λ_i .

6. THE HIGHER RANK K-THEORETIC DT PARTITION FUNCTION

6.1. Symmetrised exterior algebras and brackets. We recall some constructions in equivariant K-theory which will be used to prove Theorem A. For a recent and more complete reference, the reader may consult [52, § 2].

Let \mathbf{T} be a torus, $V = \sum_{\mu} t^{\mu}$ a \mathbf{T} -module. Assume that $\det(V)$ is a square in $K_{\mathbf{T}}^0(\text{pt})$. Define the *symmetrised exterior algebra* of V as

$$\widehat{\Lambda}^{\bullet} V := \frac{\Lambda^{\bullet} V}{\det(V)^{\frac{1}{2}}}.$$

It satisfies the relation

$$\widehat{\Lambda}^{\bullet} V^{\vee} = (-1)^{\text{rk } V} \widehat{\Lambda}^{\bullet} V.$$

Define the operator $[\cdot]$ by

$$[t^{\mu}] = t^{\frac{\mu}{2}} - t^{-\frac{\mu}{2}}.$$

One can compute

$$\text{tr}(\widehat{\Lambda}^{\bullet} V^{\vee}) = \prod_{\mu} \frac{1 - t^{-\mu}}{t^{-\frac{\mu}{2}}} = \prod_{\mu} (t^{\frac{\mu}{2}} - t^{-\frac{\mu}{2}}) = \prod_{\mu} [t^{\mu}],$$

For a virtual \mathbf{T} -representation $V = \sum_{\mu} t^{\mu} - \sum_{\nu} t^{\nu}$, where the weight $\nu = 0$ never appears, we extend $\widehat{\Lambda}^{\bullet}$ and $[\cdot]$ by linearity and find

$$\text{tr}(\widehat{\Lambda}^{\bullet} V^{\vee}) = \frac{\prod_{\mu} [t^{\mu}]}{\prod_{\nu} [t^{\nu}]}.$$

6.2. Proof of Theorem A. By the description of the \mathbf{T} -fixed locus $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$ given in § 3.4, every colored plane partition $\overline{\pi} = (\pi_1, \dots, \pi_r)$ corresponds to a unique \mathbf{T} -fixed point $S = \bigoplus_{i=1}^r \mathcal{S}_{Z_i}$, for which we defined in Equation (5.5) the vertex terms V_{ij} by

$$V_{ij} = w_i^{-1} w_j \left(Q_j - \frac{\overline{Q}_i}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} Q_j \overline{Q}_i \right)$$

with notation as in § 5.1. The generating function (4.3) of higher rank cohomological DT invariants can be rewritten in a purely combinatorial fashion as

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, v) = \sum_{\overline{\pi}} q^{|\overline{\pi}|} \prod_{i,j=1}^r e^{\text{T}(-V_{ij})}.$$

Similarly, the generating function (4.5) of the K-theoretic invariants can be rewritten as

$$\text{DT}_r^K(\mathbb{A}^3, q, t, w) = \sum_{\overline{\pi}} q^{|\overline{\pi}|} \prod_{i,j=1}^r [-V_{ij}].$$

A closed formula for $\text{DT}_1^K(\mathbb{A}^3, q, t, w)$ was conjectured in [46] and has recently been proven by Okounkov.

Theorem 6.1 ([52, Thm. 3.3.6]). *The rank 1 K-theoretic DT partition function of \mathbb{A}^3 is given by*

$$\text{DT}_1^K(\mathbb{A}^3, -q, t, w) = \text{Exp}(F_1(q, t_1, t_2, t_3))$$

where, setting $t = t_1 t_2 t_3$, one defines

$$F_1(q, t_1, t_2, t_3) = \frac{1}{[t^{\frac{1}{2}} q][t^{\frac{1}{2}} q^{-1}]} \frac{[t_1 t_2][t_1 t_3][t_2 t_3]}{[t_1][t_2][t_3]}.$$

Remark 6.2. It is clear from the expression of the vertex in rank 1 that there is no dependence on w_1 . As pointed out to us by N. Arbesfeld, this can in fact be seen as a shadow of the fact that $\mathbb{T}_2 = \mathbb{C}^*$ acts trivially on Quot_1^n and on df_n . Surprisingly, the same phenomenon occurs in the higher rank case (cf. Theorem 6.5).

We devote the rest of this section to proving a generalisation of Theorem 6.1 to higher rank.

Theorem 6.3. *The rank r K-theoretic DT partition function of \mathbb{A}^3 is given by*

$$\text{DT}_r^K(\mathbb{A}^3, (-1)^r q, t, w) = \text{Exp}(F_r(q, t_1, t_2, t_3)),$$

where, setting $t = t_1 t_2 t_3$, one defines

$$F_r(q, t_1, t_2, t_3) = \frac{[t^r]}{[t][t^{\frac{r}{2}} q][t^{\frac{r}{2}} q^{-1}]} \frac{[t_1 t_2][t_1 t_3][t_2 t_3]}{[t_1][t_2][t_3]}.$$

Remark 6.4. This result was conjectured in [3] by Awata and Kanno, who also proved it mod q^4 , i.e. up to 3 instantons. The conjecture was confirmed numerically up to some order by Benini–Bonelli–Poggi–Tanzini [9].

The proof of Theorem 6.3 will follow essentially by taking suitable limits of the weights w_i . To perform such limits, we prove the slogan (1.3), already anticipated in the Introduction.

Theorem 6.5. *The generating function $\text{DT}_r^K(\mathbb{A}^3, q, t, w)$ is independent of the weights w_1, \dots, w_r .*

Proof. The n -th coefficient of $\text{DT}_r^K(\mathbb{A}^3, q, t, w)$ is a sum of contributions

$$[-T_{\pi}^{\text{vir}}], \quad |\pi| = n.$$

A simple manipulation shows that

$$(6.1) \quad [-T_{\pi}^{\text{vir}}] = A(t) \prod_{1 \leq i < j \leq r} \frac{\prod_{\mu_{ij}} w_i - w_j t^{\mu_{ij}}}{\prod_{\nu_{ij}} w_i - w_j t^{\nu_{ij}}} = A(t) \prod_{1 \leq i < j \leq r} \frac{\prod_{\mu_{ij}} (1 - w_i^{-1} w_j t^{\mu_{ij}})}{\prod_{\nu_{ij}} (1 - w_i^{-1} w_j t^{\nu_{ij}})},$$

where $A(t) \in \mathbb{Q}((t_1, t_2, t_3, (t_1 t_2 t_3)^{\frac{1}{2}}))$ and the number of weights μ_{ij} and ν_{ij} is the same. Thus, $\text{DT}_r^K(\mathbb{A}^3, q, t, w)$ is a homogeneous rational expression of total degree 0 with respect to the variables w_1, \dots, w_r . We aim to show that $\text{DT}_r^K(\mathbb{A}^3, q, t, w)$ has no poles of the form $1 - w_i^{-1} w_j t^{\nu_{ij}}$, implying that it is a degree 0 polynomial in the w_i , hence constant in the w_i . This generalises the strategy of [3, § 4].

Set $w = w_i^{-1} w_j t^{\nu}$ for fixed $i < j$ and $\nu \in \widehat{\mathbb{T}}_1$. To see that $1 - w$ is not a pole, we use [1, Prop. 3.2], which asserts the following: if M is a quasiprojective \mathbf{T} -scheme with a \mathbf{T} -equivariant perfect obstruction theory and proper (nonempty) fixed locus, then for any $V \in K_0^{\mathbf{T}}(M)$, the only poles of the form $(1 - w)$ that may appear in $\chi(M, V \otimes \mathcal{O}^{\text{vir}})$ arise from *noncompact weights* $w \in \widehat{\mathbf{T}}$. A weight w is called compact if the fixed locus $M^{\mathbf{T}_w} \subset M$ is proper, where \mathbf{T}_w is the maximal torus in $\ker(w) \subset \mathbf{T}$, and is called noncompact otherwise [1, Def. 3.1].

We of course want to apply [1, Prop. 3.2] to $M = Q = \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$, $\mathbf{T} = \mathbb{T}_1 \times \mathbb{T}_2$ and $V = \mathcal{K}_{\text{vir}}^{1/2}$. By Equation (6.1), our goal is to prove that $w = w_i^{-1} w_j t^{\nu}$ is a compact weight for all $i < j$ and $\nu \in \widehat{\mathbb{T}}_1$.

First of all, we observe that $\mathbf{T}_w = \ker(w)$. Indeed w is not a product of powers of weights of \mathbf{T} , hence

$$\mathcal{O}(\ker w) = \mathcal{O}(\mathbf{T}) / (w_i^{-1} w_j t^\nu - 1) \cong \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, t_3^{\pm 1}, w_1^{\pm 1}, \dots, w_{i-1}^{\pm 1}, w_{i+1}^{\pm 1}, \dots, w_r^{\pm 1}],$$

which shows that $\ker(w)$ is itself a torus (of dimension $3 + r - 1$). Next, consider the automorphism $\tau_\nu: \mathbf{T} \xrightarrow{\sim} \mathbf{T}$ defined by

$$(t_1, t_2, t_3, w_1, \dots, w_r) \mapsto (t_1, t_2, t_3, w_1, \dots, w_i t^{-\nu}, \dots, w_j, \dots, w_r).$$

It maps $\mathbf{T}_w \subset \mathbf{T}$ isomorphically onto the subtorus $\mathbb{T}_1 \times \{w_i = w_j\} \subset \mathbf{T}$. This yields an inclusion of tori

$$(6.2) \quad \mathbb{T}_1 \xrightarrow{\sim} \mathbb{T}_1 \times \{(1, \dots, 1)\} \hookrightarrow \tau(\mathbf{T}_w).$$

We consider the action $\sigma_\nu: \mathbf{T} \times Q \rightarrow Q$ where \mathbb{T}_1 translates the support of the quotient sheaf in the usual way, the i -th summand of $\mathcal{O}^{\oplus r}$ gets scaled by $w_i t^\nu$ and all other summands by w_k for $k \neq i$. In other words, in terms of the matrix-and-vectors description of Q , we set

$$\sigma_\nu(\mathbf{t}, (A_1, A_2, A_3, u_1, \dots, u_r)) = (t_1 A_1, t_2 A_2, t_3 A_3, w_1 u_1, \dots, w_i t^\nu u_i, \dots, w_r u_r),$$

just a variation of Equation (3.9) in the i -th vector component. Then, upon restricting this action to \mathbf{T}_w , we have a commutative diagram

$$\begin{array}{ccc} \mathbf{T}_w \times Q & \xrightarrow{\sigma} & Q \\ \tau_\nu \times \text{id} \downarrow & & \parallel \\ \tau_\nu(\mathbf{T}_w) \times Q & \xrightarrow{\sigma_\nu} & Q \end{array}$$

where σ is the restriction of the usual action (3.9). This diagram induces a natural isomorphism $Q^{\mathbf{T}_w} \xrightarrow{\sim} Q^{\tau_\nu(\mathbf{T}_w)}$, which combined with (6.2) yields an inclusion

$$Q^{\mathbf{T}_w} \xrightarrow{\sim} Q^{\tau_\nu(\mathbf{T}_w)} \hookrightarrow Q^{\mathbb{T}_1},$$

where $Q^{\mathbb{T}_1}$ is the fixed locus with respect to the action σ_ν . But by the same reasoning as in Remark 3.4, this fixed locus is proper (because, again, a \mathbb{T}_1 -fixed surjection $\mathcal{O}^{\oplus r} \rightarrow T$ necessarily has the quotient T entirely supported at the origin $0 \in \mathbb{A}^3$). Thus w is a compact weight, and the result follows. \square

Remark 6.6. After a first draft of this work was already finished, we were informed of an alternative way to prove Theorem 6.5, which, in a nutshell, goes as follows: one exploits the (proper) Quot-to-Chow morphism $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n) \rightarrow \text{Sym}^n \mathbb{A}^3$ to express the K-theoretic DT invariants as equivariant holomorphic Euler characteristics on $\text{Sym}^n \mathbb{A}^3$, where the framing torus \mathbb{T}_2 is acting trivially on the symmetric product. One concludes by an application of Okounkov's *rigidity principle* [52, § 2.4.1]. This strategy will be carried out in [2].

Thanks to Theorem 6.5, we may now specialise w_1, \dots, w_r to arbitrary values and take arbitrary limits. We set $w_i = L^i$ for $i = 1, \dots, r$ and compute the limit for $L \rightarrow \infty$.

Lemma 6.7. *Let $i < j$. Then we have*

$$\lim_{L \rightarrow \infty} [-V_{ij}] [-V_{ji}] \Big|_{w_i = L^i} = (-t^{\frac{1}{2}})^{|\pi_j| - |\pi_i|}.$$

Proof. Notice that all monomials in V_{ij} are of the form $w_i^{-1} w_j \lambda$ for λ a monomial in t_1, t_2, t_3 . Then

$$[w_i^{-1} w_j \lambda]_{w_i=L^i} = (L^{j-i} \lambda)^{\frac{1}{2}} (1 - L^{i-j} \lambda^{-1}).$$

Write $Q_i = \sum_{\mu} t^{\mu}$ and $Q_j = \sum_{\nu} t^{\nu}$. Taking limits, we obtain

$$\begin{aligned} \lim_{L \rightarrow \infty} [-V_{ij}]_{w_i=L^i} &= \lim_{L \rightarrow \infty} [-w_i^{-1} w_j (Q_j - \bar{Q}_i t^{-1} + \bar{Q}_i Q_j t^{-1} (1-t_1)(1-t_2)(1-t_3))]_{w_i=L^i} \\ &= \lim_{L \rightarrow \infty} L^{\frac{j-i}{2}(|\pi_i| - |\pi_j|)} \frac{\prod_{\mu} (t^{-\frac{\mu}{2}} t^{-\frac{1}{2}})}{\prod_{\nu} t^{\frac{\nu}{2}}}. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} \lim_{L \rightarrow \infty} [-V_{ji}]_{w_i=L^i} &= (-1)^{\text{rk}(-V_{ji})} \lim_{L \rightarrow \infty} [-\bar{V}_{ji}]_{w_i=L^i} \\ &= (-1)^{|\pi_i| - |\pi_j|} \lim_{L \rightarrow \infty} [-w_i^{-1} w_j (\bar{Q}_i - Q_j t - \bar{Q}_i Q_j (1-t_1)(1-t_2)(1-t_3))]_{w_i=L^i} \\ &= (-1)^{|\pi_i| - |\pi_j|} \lim_{L \rightarrow \infty} L^{\frac{j-i}{2}(|\pi_j| - |\pi_i|)} \frac{\prod_{\nu} (t^{\frac{\nu}{2}} t^{\frac{1}{2}})}{\prod_{\mu} t^{-\frac{\mu}{2}}}. \end{aligned}$$

We conclude, as required, that

$$\lim_{L \rightarrow \infty} [-V_{ij}][-\bar{V}_{ji}]_{w_i=L^i} = (-t^{\frac{1}{2}})^{|\pi_j| - |\pi_i|}. \quad \square$$

Lemma 6.8. *Let x be a variable and $c_i \in \mathbb{Z}$, for $i = 1, \dots, r$. Then we have*

$$\prod_{1 \leq i < j \leq r} x^{c_j - c_i} = \prod_{i=1}^r x^{(-r-1+2i)c_i}.$$

Proof. The assertion holds for $r = 1$ as the productory on the left hand side is empty. Assume it holds for $r - 1$. Then we have:

$$\begin{aligned} \prod_{1 \leq i < j \leq r} x^{c_j - c_i} &= \prod_{1 \leq i < j \leq r-1} x^{c_j - c_i} \prod_{i=1}^{r-1} x^{c_r - c_i} \\ &= x^{(r-1)c_r} \prod_{i=1}^{r-1} x^{(-r-1+2i)c_i} \\ &= \prod_{i=1}^r x^{(-r-1+2i)c_i}. \quad \square \end{aligned}$$

Combining Lemma 6.7 with Lemma 6.8 we can express the rank r K-theoretic DT theory of \mathbb{A}^3 as a product of r copies of the rank 1 K-theoretic DT theory. This product formula already appeared as a limit of the (conjectural) 4-fold theory developed by Nekrasov and Piazzalunga [50, Formula (3.14)].³

Theorem 6.9. *There is an identity*

$$\text{DT}_r^{\text{K}}(\mathbb{A}^3, (-1)^r q, t, w) = \prod_{i=1}^r \text{DT}_1^{\text{K}}(\mathbb{A}^3, -q t^{\frac{-r-1}{2} + i}, t).$$

³Typo warning: N. Piazzalunga kindly pointed out to us that in [50, Formula (3.14)] one should read ' $\frac{N+1}{2} - l$ ' instead of ' $N + 1 - 2l$ '.

Proof. Set $w_i = L^i$. The generating series $\mathrm{DT}_r^K(\mathbb{A}^3, q, t, w)$ can be computed in the limit $L \rightarrow \infty$:

$$\begin{aligned}
\lim_{L \rightarrow \infty} \mathrm{DT}_r^K(\mathbb{A}^3, q, t, w) &= \lim_{L \rightarrow \infty} \sum_{\bar{\pi}} q^{|\bar{\pi}|} \prod_{i,j=1}^r [-V_{ij}] \\
&= \lim_{L \rightarrow \infty} \sum_{\bar{\pi}} \prod_{i=1}^r q^{|\pi_i|} [-V_{ii}] \prod_{1 \leq i < j \leq r} [-V_{ij}] [-V_{ji}] \\
&= \sum_{\bar{\pi}} \prod_{i=1}^r q^{|\pi_i|} [-V_{ii}] \prod_{1 \leq i < j \leq r} (-t^{\frac{1}{2}})^{|\pi_j| - |\pi_i|} \\
&= \sum_{\bar{\pi}} \prod_{i=1}^r q^{|\pi_i|} [-V_{ii}] \prod_{i=1}^r (-t^{\frac{1}{2}})^{(-r-1+2i)|\pi_i|} \\
&= \sum_{\bar{\pi}} \prod_{i=1}^r [-V_{ii}] q^{|\pi_i|} (-1)^{(r+1)|\pi_i|} t^{(\frac{-r-1}{2}+i)|\pi_i|} \\
&= \sum_{\bar{\pi}} \prod_{i=1}^r [-V_{ii}] \left((-1)^{(r+1)} q t^{\frac{-r-1}{2}+i} \right)^{|\pi_i|} \\
&= \prod_{i=1}^r \mathrm{DT}_1^K(\mathbb{A}^3, (-1)^{(r+1)} q t^{\frac{-r-1}{2}+i}, t). \quad \square
\end{aligned}$$

We can now prove Theorem 6.3 (i.e. Theorem A from the Introduction).

Proof of Theorem 6.3. Define

$$G_{r,i}(q, t_1, t_2, t_3) = F_1(q t^{\frac{-r-1}{2}+i}, t_1, t_2, t_3).$$

We have

$$\begin{aligned}
\mathrm{DT}_1^K(\mathbb{A}^3, -q t^{\frac{-r-1}{2}+i}, t) &= \exp \left(\sum_{n \geq 1} \frac{1}{n} \frac{1}{[t^{\frac{n}{2}} q^n t^{n(\frac{-r-1}{2}+i)}] [t^{\frac{n}{2}} q^{-n} t^{n(\frac{r+1}{2}-i)}]} \frac{[t_1^n t_2^n] [t_1^n t_3^n] [t_2^n t_3^n]}{[t_1^n] [t_2^n] [t_3^n]} \right) \\
&= \mathrm{Exp}(G_{r,i}(q, t_1, t_2, t_3)).
\end{aligned}$$

By Theorem 6.9 and Theorem 6.1 it is enough to show that $F_r = \sum_{i=1}^r G_{r,i}$, or equivalently

$$\sum_{i=1}^r \frac{1}{[t^{\frac{1}{2}} q t^{\frac{-r-1}{2}+i}] [t^{\frac{1}{2}} q^{-1} t^{\frac{r+1}{2}-i}]} = \frac{[t^r]}{[t] [t^{\frac{r}{2}} q] [t^{\frac{r}{2}} q^{-1}]}.$$

It is easy to check this is true for $r = 1, 2$. Let now $r \geq 3$: we perform induction separately on even and odd cases. Assume the claimed identity holds for $r-2$. In both cases we have

$$\begin{aligned}
&\sum_{i=1}^r \frac{1}{[t^{\frac{1}{2}} q t^{\frac{-r-1}{2}+i}] [t^{\frac{1}{2}} q^{-1} t^{\frac{r+1}{2}-i}]} \\
&= \sum_{i=1}^{r-2} \frac{1}{[t^{\frac{1}{2}} q t^{\frac{-(r-2)-1}{2}+i}] [t^{\frac{1}{2}} q^{-1} t^{\frac{(r-2)+1}{2}-i}]} + \frac{1}{[q t^{-\frac{r}{2}+1}] [q^{-1} t^{\frac{r}{2}}]} + \frac{1}{[q t^{\frac{r}{2}}] [q^{-1} t^{-\frac{r}{2}+1}]} \\
&= \frac{[t^{r-2}]}{[t] [t^{\frac{r-2}{2}} q] [t^{\frac{r-2}{2}} q^{-1}]} - \frac{1}{[t^{\frac{r-2}{2}} q^{-1}] [t^{\frac{r}{2}} q^{-1}]} - \frac{1}{[t^{\frac{r}{2}} q] [t^{\frac{r-2}{2}} q]} \\
&= \frac{1}{[t] [t^{\frac{r}{2}} q] [t^{\frac{r}{2}} q^{-1}]} \cdot \frac{[t^{r-2}] [t^{\frac{r}{2}} q] [t^{\frac{r}{2}} q^{-1}] - [t] [t^{\frac{r}{2}} q] [t^{\frac{r-2}{2}} q] - [t] [t^{\frac{r}{2}} q^{-1}] [t^{\frac{r-2}{2}} q^{-1}]}{[t^{\frac{r-2}{2}} q] [t^{\frac{r-2}{2}} q^{-1}]} \\
&= \frac{[t^r]}{[t] [t^{\frac{r}{2}} q] [t^{\frac{r}{2}} q^{-1}]}
\end{aligned}$$

by which we conclude the proof. \square

6.3. Comparison with motivic DT invariants. Let $f: U \rightarrow \mathbb{A}^1$ be a regular function on a smooth scheme U , and let $\hat{\mu}$ be the group of all roots of unity. The critical locus $Z = \text{crit}(f) \subset U$ inherits a canonical *virtual motive* [6], i.e. a $\hat{\mu}$ -equivariant motivic class

$$[Z]_{\text{vir}} = -\mathbb{L}^{-\frac{\dim U}{2}} [\phi_f] \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}} = K_0^{\hat{\mu}}(\text{Var}_{\mathbb{C}})[\mathbb{L}^{-\frac{1}{2}}]$$

such that $e[Z]_{\text{vir}} = e_{\text{vir}}(Z)$, where $e_{\text{vir}}(-)$ is Behrend weighted Euler characteristic and the Euler number specialisation prescribes $e(\mathbb{L}^{-1/2}) = -1$. The motivic class $[\phi_f]$ is the (absolute) motivic vanishing cycle class introduced by Denef and Loeser [21].

The virtual motive of $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n) = \text{crit}(f_n)$, with respect to the critical structure of Proposition 3.2, was computed in [55, Prop. 2.3.6]. The result is as follows. Let $\text{DT}_r^{\text{mot}}(\mathbb{A}^3, q) \in \mathcal{M}_{\mathbb{C}}[[q]]$ be the generating function of the virtual motives $[\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)]_{\text{vir}}$. Then one has

$$(6.3) \quad \text{DT}_r^{\text{mot}}(\mathbb{A}^3, q) = \prod_{m \geq 1} \prod_{k=0}^{r-1} \left(1 - \mathbb{L}^{2+k-\frac{rm}{2}} q^m \right)^{-1}.$$

The case $r = 1$ was computed in [6]. The general proof of Formula (6.3) is obtained in a similar fashion in [55, 14], and via a wall-crossing technique in [15]. Moreover, it is immediate to verify that DT_r^{mot} satisfies a product formula analogous to the one proved in Theorem 6.9 for the K-theoretic invariants: we have

$$(6.4) \quad \text{DT}_r^{\text{mot}}(\mathbb{A}^3, q) = \prod_{i=1}^r \text{DT}_1^{\text{mot}}\left(\mathbb{A}^3, q \mathbb{L}^{-\frac{r-1}{2}+i}\right).$$

In particular, up to the substitution $\mathbb{L}^{\frac{1}{2}} \rightarrow -\mathbb{L}^{\frac{1}{2}}$, the factorisation (6.4) is equivalent to the K-theoretic one (Theorem 6.9). As observed in [59, § 4], the (signed) motivic partition function admits an expression in terms of the motivic exponential, namely

$$(6.5) \quad \text{DT}_r^{\text{mot}}(\mathbb{A}^3, (-1)^r q) = \text{Exp} \left(\frac{(-1)^r q \mathbb{L}^{\frac{3}{2}}}{(1 - (-1)^r q \mathbb{L}^{\frac{r}{2}})(1 - (-1)^r q \mathbb{L}^{-\frac{r}{2}})} \frac{\mathbb{L}^{\frac{r}{2}} - \mathbb{L}^{-\frac{r}{2}}}{\mathbb{L}^{\frac{1}{2}} - \mathbb{L}^{-\frac{1}{2}}} \right).$$

Given their structural similarities, we believe it is an interesting problem to compare the K-theoretic partition function with the motivic one.

It is worth noticing that Formula (6.5) can be recovered from the factorisation (6.4), just as we discovered in the K-theoretic case during the proof of Theorem 6.3. This fact follows immediately from the properties of the plethystic exponential.

Remark 6.10. A virtual motive for $\text{Quot}_Y(F, n)$ was defined in [59, § 4] for every locally free sheaf F on a 3-fold Y . Just as in the case of the naive motives of the Quot scheme [58], the resulting partition function $\text{DT}_r^{\text{mot}}(Y, q)$ only depends on the motivic class $[Y] \in K_0(\text{Var}_{\mathbb{C}})$ and on $r = \text{rk } F$. See also [16] for calculations of motivic higher rank DT and PT invariants in the presence of nonzero curve classes: the generating function $\text{DT}_r^{\text{mot}}(Y, q)$, computed easily starting with Formula (6.3), is precisely the DT/PT wall-crossing factor.

7. THE HIGHER RANK COHOMOLOGICAL DT PARTITION FUNCTION

7.1. Cohomological reduction. One should think of K-theoretic invariants as refinements of the cohomological ones, as by taking suitable limits one fully recovers $\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s)$ from $\text{DT}_r^K(\mathbb{A}^3, q, t)$. We make this precise in the remainder of this section.

Let $\mathbf{T} \cong (\mathbb{C}^*)^g$ be an algebraic torus and let t_1, \dots, t_g be its coordinates. Recall that the Chern character gives a natural transformation from (equivariant) K-theory to the (equivariant) Chow

group with rational coefficients by sending $t_i \mapsto e^{s_i}$, where $s_i = c_1^T(t_i)$. We can formally extend it to

$$\begin{array}{ccc} \mathbb{Z}[t_1^{\pm 1}, \dots, t_g^{\pm 1}] & \xrightarrow{\text{ch}} & \mathbb{Q}[[s_1, \dots, s_g]] \\ \downarrow & & \downarrow \\ \mathbb{Z}[t_1^{\pm b}, \dots, t_g^{\pm b} | b \in \mathbb{C}] & \xrightarrow{\text{ch}} & \mathbb{C}[[s_1, \dots, s_g]] \end{array}$$

by sending $t_i^b \mapsto e^{b s_i}$, where $b \in \mathbb{C}$.

In §6.1 we defined the symmetrised transformation $[t^\mu] = t^{\frac{\mu}{2}} - t^{-\frac{\mu}{2}}$. We set $[\text{ch}(t^{b\mu})] = e^{\frac{b\mu \cdot s}{2}} - e^{-\frac{b\mu \cdot s}{2}}$ as an expression in rational cohomology, which enjoys the following *linearisation* property:

$$[\text{ch}(t^{b\mu})] = e^{\frac{b\mu \cdot s}{2}} (1 - e^{-b\mu \cdot s}) = b e^T(t^\mu) + o(b^2).$$

In other words, $e^T(\cdot)$ is the first-order approximation of $[\cdot]$ in \mathbf{T} -equivariant Chow groups. For a virtual representation $V = \sum_\mu t^\mu - \sum_\nu t^\nu \in K_0^T(\text{pt})$, denote by $V^b = \sum_\mu t^{b\mu} - \sum_\nu t^{b\nu}$ the virtual representation where we formally substitute each weight t^μ with $t^{b\mu}$. We have the identity

$$[\text{ch}(V^b)] = \frac{\prod_\mu [\text{ch}(t^{b\mu})]}{\prod_\nu [\text{ch}(t^{b\nu})]} = b^{\text{rk } V} \frac{\prod_\mu (e^T(t^\mu) + o(b))}{\prod_\nu (e^T(t^\nu) + o(b))}.$$

If $\text{rk } V = 0$, by taking the limit for $b \rightarrow 0$ we conclude

$$(7.1) \quad \lim_{b \rightarrow 0} [\text{ch}(V^b)] = e^T(V).$$

It is clear from the definition of $\text{ch}(\cdot)$ and $[\cdot]$ that these two transformations commute with each other. This proves the following relation between K-theoretic invariants and cohomological invariants of the local model.

Corollary 7.1. *There is an identity*

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, \nu) = \lim_{b \rightarrow 0} \text{DT}_r^K(\mathbb{A}^3, q, e^{bs}, e^{b\nu}).$$

Proof. Follows from the description of the generating series of K-theoretic invariants as

$$\text{DT}_r^K(\mathbb{A}^3, q, t, w) = \sum_{n \geq 0} q^n \sum_{[S] \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^T} [-T_S^{\text{vir}}]$$

and by noticing that $\text{rk } T_S^{\text{vir}} = 0$. □

Thanks to the ν -independence, we can now rename

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s) = \text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, \nu).$$

We are ready to prove Theorem B from the Introduction.

Theorem 7.2. *The rank r cohomological DT partition function of \mathbb{A}^3 is given by*

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s) = M((-1)^r q)^{-r \frac{(s_1 + s_2)(s_1 + s_3)(s_2 + s_3)}{s_1 s_2 s_3}}.$$

Proof. By Corollary 7.1 and Theorem 6.3, we just need to compute the limit

$$\lim_{b \rightarrow 0} \text{DT}_r^K(\mathbb{A}^3, (-1)^r q, e^{bs}) = \lim_{b \rightarrow 0} \text{Exp}(F_r(q, t_1^b, t_2^b, t_3^b)).$$

Denote for ease of notation $s = c_1^T(t) = s_1 + s_2 + s_3$. By the definition of plethystic exponential, recalled in (1.4), we have

$$\begin{aligned} & \lim_{b \rightarrow 0} \text{Exp}(F_r(q, t_1^b, t_2^b, t_3^b)) \\ &= \exp \sum_{k \geq 1} \frac{1}{k} \left(\lim_{b \rightarrow 0} \frac{[e^{bkr s}]}{[e^{bks}][e^{\frac{bkr}{2}s} q^k][e^{\frac{bkr}{2}s} q^{-k}]} \frac{[e^{bk(s_1+s_2)}][e^{bk(s_1+s_3)}][e^{bk(s_2+s_3)}]}{[e^{bks_1}][e^{bks_2}][e^{bks_3}]} \right). \end{aligned}$$

We have

$$\lim_{b \rightarrow 0} \frac{[e^{bk(s_1+s_2)}][e^{bk(s_1+s_3)}][e^{bk(s_2+s_3)}]}{[e^{bks_1}][e^{bks_2}][e^{bks_3}]} = \frac{(s_1+s_2)(s_1+s_3)(s_2+s_3)}{s_1 s_2 s_3},$$

and

$$\lim_{b \rightarrow 0} \frac{[e^{bkr s}]}{[e^{bks}][e^{\frac{bkr}{2}s} q^k][e^{\frac{bkr}{2}s} q^{-k}]} = \frac{r}{[q^k][q^{-k}]} = -r \cdot \frac{q^k}{(1-q^k)^2}.$$

Recall the plethystic exponential form of the MacMahon function

$$M(q) = \prod_{n \geq 1} (1 - q^n)^{-n} = \text{Exp}\left(\frac{q}{(1-q)^2}\right).$$

We conclude

$$\begin{aligned} \lim_{b \rightarrow 0} \text{DT}_r^K(\mathbb{A}^3, (-1)^r q, e^{bs}) &= \exp \left(-r \cdot \frac{(s_1+s_2)(s_1+s_3)(s_2+s_3)}{s_1 s_2 s_3} \sum_{k \geq 1} \frac{1}{k} \frac{q^k}{(1-q^k)^2} \right) \\ &= M(q)^{-r \frac{(s_1+s_2)(s_1+s_3)(s_2+s_3)}{s_1 s_2 s_3}}. \end{aligned} \quad \square$$

Thus we proved Szabo's conjecture [62, Conj. 4.10].

Remark 7.3. The specialisation

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s) \Big|_{s_1+s_2+s_3=0} = M((-1)^r q)^r,$$

recovering Formula (4.1), was already known in physics, see e.g. [19].

We end this section with a small variation of Theorem B.

Corollary 7.4. Fix an r -tuple $\lambda = (\lambda_1, \dots, \lambda_r)$ of \mathbb{T}_1 -equivariant line bundles on \mathbb{A}^3 . Then there is an identity

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s) = \text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, v)_\lambda,$$

where the right hand side was defined in (5.11).

Proof. We have

$$T_{S, \lambda}^{\text{vir}} = \sum_{i,j} \lambda_i^{-1} \lambda_j V_{ij}.$$

Let $V_{ij} = \sum_{\mu} w_i^{-1} w_j t^{\mu}$ be the decomposition into weight spaces. A monomial in $T_{S, \lambda}^{\text{vir}}$ is of the form $\lambda_i^{-1} \lambda_j w_i^{-1} w_j t^{\mu}$ and its Euler class is

$$\begin{aligned} e^T(\lambda_i^{-1} \lambda_j w_i^{-1} w_j t^{\mu}) &= \mu \cdot s + v_j + c_1^T(\lambda_j) - v_i - c_1^T(\lambda_i) \\ &= \mu \cdot s + \bar{v}_j - \bar{v}_i \end{aligned}$$

where we define $\bar{v}_i = v_i + c_1^T(\lambda_i)$. We conclude that

$$\text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, v)_\lambda = \text{DT}_r^{\text{coh}}(\mathbb{A}^3, q, s, \bar{v}),$$

which does not depend on \bar{v} by Theorem 6.5. □

Example 7.5. Set $r = 2$, $n = 1$, so that the only \mathbf{T} -fixed points in $\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus 2}, 1)$ are the direct sums of ideal sheaves

$$S_1 = \mathcal{I}_{\text{pt}} \oplus \mathcal{O} \subset \mathcal{O}^{\oplus 2}, \quad S_2 = \mathcal{O} \oplus \mathcal{I}_{\text{pt}} \subset \mathcal{O}^{\oplus 2},$$

where $\text{pt} = (0, 0, 0) \in \mathbb{A}^3$ is the origin. One computes

$$\begin{aligned} T_{S_1}^{\text{vir}} &= 1 - \frac{1}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} - w_1^{-1} w_2 \frac{1}{t_1 t_2 t_3} + w_2^{-1} w_1 \\ T_{S_2}^{\text{vir}} &= 1 - \frac{1}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} - w_2^{-1} w_1 \frac{1}{t_1 t_2 t_3} + w_1^{-1} w_2. \end{aligned}$$

Therefore, the cohomological DT invariant is

$$\int_{[\text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus 2}, 1)]^{\text{vir}}} 1 = e^{\mathbf{T}(-T_{S_1}^{\text{vir}})} + e^{\mathbf{T}(-T_{S_2}^{\text{vir}})}.$$

The part that could possibly depend on the framing parameters v_1 and v_2 is, in fact, constant:

$$e^{\mathbf{T}\left(w_1^{-1} w_2 \frac{1}{t_1 t_2 t_3} - w_2^{-1} w_1\right)} + e^{\mathbf{T}\left(w_2^{-1} w_1 \frac{1}{t_1 t_2 t_3} - w_1^{-1} w_2\right)} = \frac{-v_1 + v_2 - \mathfrak{s}}{v_1 - v_2} + \frac{v_1 - v_2 - \mathfrak{s}}{v_2 - v_1} = -2.$$

Let λ_1 and λ_2 be two \mathbb{T}_1 -equivariant line bundles. After the substitutions $w_i \rightarrow w_i \lambda_i$, and setting $\bar{v}_i = v_i + c_1(\lambda_i)$, the final sum of Euler classes depending on \bar{v} becomes

$$2 \frac{\bar{v}_2 - \bar{v}_1}{\bar{v}_1 - \bar{v}_2} = -2.$$

8. ELLIPTIC DONALDSON–THOMAS INVARIANTS

8.1. Chiral elliptic genus. In [9] an elliptic (string-theoretic) generalisation of the K -theoretic DT invariants is given. In physics the invariants computed in loc. cit. are obtained as the superconformal index of a D1-D7 brane system on a type IIB $\mathcal{N} = 1$ supersymmetric background, where r D7-branes wrap the product of a 3-fold by a torus, i.e. $X_3 \times T^2$, while n D1-branes wrap T^2 . The connection with enumerative geometry is then given via BPS-bound states counting, as brane systems often provide interesting constructions of relevant moduli spaces. In the case at hand, when $X_3 = \mathbb{A}^3$, the D1-D7 brane system considered has a BPS moduli space which can be naturally identified with the moduli space parametrising quotients of length n of $\mathcal{O}_{\mathbb{A}^3}^{\oplus r}$. The superconformal index is usually identified in the physics literature with the elliptic genus of such a moduli space. This however does not coincide with the usual notion of (virtual) elliptic genus, as the coupling to the D7-branes breaks half of the chiral supersymmetry, thus leading to an effective $2d$ $\mathcal{N} = (0, 2)$ GLSM on T^2 for the D1-brane dynamics, whose Witten index generalises the K-theoretic DT invariants of \mathbb{A}^3 and provide a sort of chiral (or 1/2 BPS) version of the elliptic genus of the Quot scheme. In this section we give a mathematical definition of the elliptic invariants computed in [9], and show that our definition leads precisely to the computation of the same quantities found in [9, § 3].

Let X be a scheme carrying a perfect obstruction theory $\mathbb{E} \rightarrow \mathbb{L}_X$ of virtual dimension $\text{vd} = \text{rk } \mathbb{E}$.

Definition 8.1. If F is a rank r vector bundle on X we define

$$(8.1) \quad \mathcal{E}_{1/2}(F) = \bigotimes_{n \geq 1} \text{Sym}_{p^n}^{\bullet} (F \oplus F^{\vee}) \in 1 + p \cdot K^0(X)[[p]]$$

where the total symmetric algebra $\mathrm{Sym}_p^\bullet(F) = \sum_{i \geq 0} p^i [S^i F] \in K^0(X)[[p]]$ satisfies $\mathrm{Sym}_p^\bullet(F) = 1/\Lambda_{-p}^\bullet(F)$. Note that $\mathcal{E}_{1/2}$ defines a homomorphism from the additive group $K^0(X)$ to the multiplicative group $1 + p \cdot K^0(X)[[p]]$. Set

$$(8.2) \quad \mathcal{E}\ell\ell_{1/2}(F; p) = (-p^{-\frac{1}{12}})^{\mathrm{rk} F} \mathrm{ch}(\mathcal{E}_{1/2}(F)) \cdot \mathrm{td}(F) \in A^*(X)[[p]][p^{\pm \frac{1}{12}}],$$

where $\mathrm{td}(-)$ is the Todd class, so that $\mathcal{E}\ell\ell_{1/2}(-; p)$ extends to a group homomorphism from $K^0(X)$ to the multiplicative group of units in $A^*(X)[[p]][p^{\pm \frac{1}{12}}]$.

We can then define the virtual chiral elliptic genus as follows.

Definition 8.2. Let X be a proper scheme with a perfect obstruction theory and $V \in K^0(X)$. The *virtual chiral elliptic genus* is defined as

$$\mathrm{Ell}_{1/2}^{\mathrm{vir}}(X, V; p) = (-p^{-\frac{1}{12}})^{\mathrm{vd}} \chi^{\mathrm{vir}}(X, \mathcal{E}_{1/2}(T_X^{\mathrm{vir}}) \otimes V) \in \mathbb{Z}[[p]][p^{\pm \frac{1}{12}}].$$

By the virtual Riemann–Roch theorem of [23] we can also compute the virtual chiral elliptic genus as

$$\mathrm{Ell}_{1/2}^{\mathrm{vir}}(X, V; p) = \int_{[X]^{\mathrm{vir}}} \mathcal{E}\ell\ell_{1/2}(T_X^{\mathrm{vir}}; p) \cdot \mathrm{ch}(V).$$

Remark 8.3. One may give a more general definition by adding a “mass deformation” and defining $\mathcal{E}_{1/2}^{(y)}(F)$ for $F \in K^0(X)$ as

$$\mathcal{E}_{1/2}^{(y)}(F; p) = \bigotimes_{n \geq 1} \mathrm{Sym}_{y^{-1}p^n}^\bullet(F) \otimes \mathrm{Sym}_{yp^n}^\bullet(F^\vee) \in 1 + p \cdot K^0(X)[y, y^{-1}][[p]],$$

so we recover the standard definition of virtual elliptic genus by taking $\mathcal{E}(F) = \mathcal{E}_{1/2}^{(1)}(F; p) \otimes \mathcal{E}_{1/2}^{(y)}(-F; p)$, cf. [23, § 6].

Proposition 8.4. Let X be a proper scheme with a perfect obstruction theory and let $V \in K^0(X)$. Then the virtual chiral elliptic genus $\mathrm{Ell}_{1/2}^{\mathrm{vir}}(X, V; p)$ is deformation invariant.

Proof. The statement follows directly from Definition 8.2 and [23, Theorem 3.15]. \square

Let now $V = \sum_{\mu} t^{\mu}$ be a \mathbf{T} -module as in § 6.1. The trace of its symmetric algebra is given by

$$\mathrm{tr}(\mathrm{Sym}_p^\bullet(V)) = \mathrm{tr}\left(\frac{1}{\Lambda_{-p}^\bullet(V)}\right) = \prod_{\mu} \frac{1}{1 - p t^{\mu}}.$$

Let us now assume as in § 6.1 that $\det V$ is a square in $K_{\mathbf{T}}^0(\mathrm{pt})$ and $\mu = 0$ is not a weight of V . We can then compute the trace of the symmetric product in (8.1) as

$$\begin{aligned} \mathrm{tr}\left(\bigotimes_{n \geq 1} \mathrm{Sym}_{p^n}^\bullet(V \oplus V^\vee)\right) &= \prod_{\mu} \prod_{n \geq 1} \frac{1}{(1 - p^n t^{\mu})(1 - p^n t^{-\mu})}, \\ \mathrm{tr}\left(\frac{\bigotimes_{n \geq 1} \mathrm{Sym}_{p^n}^\bullet(V \oplus V^\vee)}{\Lambda^\bullet V^\vee}\right) &= \prod_{\mu} \frac{1}{1 - t^{-\mu}} \prod_{n \geq 1} \frac{1}{(1 - p^n t^{\mu})(1 - p^n t^{-\mu})} \\ &= (-ip^{1/8} \phi(p))^{\mathrm{rk} V} \prod_{\mu} \frac{t^{\mu/2}}{\theta(p; t^{\mu})}, \end{aligned}$$

where $\phi(p)$ is the Euler function, i.e. $\phi(p) = \prod_n (1 - p^n)$, and $\theta(p; y)$ denotes the Jacobi theta function

$$\theta(p; y) = -ip^{1/8}(y^{1/2} - y^{-1/2}) \prod_{n=1}^{\infty} (1 - p^n)(1 - yp^n)(1 - y^{-1}p^n).$$

Combining everything together we get the identity

$$\begin{aligned} (-p^{-\frac{1}{12}})^{\mathrm{rk} V} \mathrm{tr} \left(\frac{\bigotimes_{n \geq 1} \mathrm{Sym}_{p^n}^\bullet (V \oplus V^\vee) \otimes \det(V^\vee)^{1/2}}{\Lambda^\bullet V^\vee} \right) &= (-p^{-\frac{1}{12}})^{\mathrm{rk} V} (-ip^{1/8} \phi(p))^{\mathrm{rk} V} \prod_\mu \frac{1}{\theta(p; t^\mu)} \\ &= \prod_\mu i \frac{\eta(p)}{\theta(p; t^\mu)}, \end{aligned}$$

where $\eta(p)$ is the Dedekind eta function

$$\eta(p) = p^{\frac{1}{24}} \prod_{n \geq 1} (1 - p^n).$$

For a virtual \mathbf{T} -representation $V = \sum_\mu t^\mu - \sum_\nu t^\nu \in K_{\mathbf{T}}^0(\mathrm{pt})$ where $\mu = 0$ is not a weight of V , we compute

$$(-p^{-\frac{1}{12}})^{\mathrm{rk} V} \mathrm{tr} \left(\frac{\bigotimes_{n \geq 1} \mathrm{Sym}_{p^n}^\bullet (V \oplus V^\vee) \otimes \det(V^\vee)^{1/2}}{\Lambda^\bullet V^\vee} \right) = (i \cdot \eta(p))^{\mathrm{rk} V} \frac{\prod_\nu \theta(p; t^\nu)}{\prod_\mu \theta(p; t^\mu)}.$$

For the remainder of the section we set $p = e^{2\pi i \tau}$, with $\tau \in \mathbb{H} = \{\tau \in \mathbb{C} \mid \Im(\tau) > 0\}$. Denoting $\theta(\tau|z) = \theta(e^{2\pi i \tau}; e^{2\pi i z})$, θ enjoys the modular behaviour

$$\theta(\tau|z + a + b\tau) = (-1)^{a+b} e^{-2\pi i b z} e^{-i\pi b^2 \tau} \theta(\tau|z), \quad a, b \in \mathbb{Z}.$$

Analogously to the measure $[\cdot]$ for K-theoretic invariants, we define the *elliptic measure*

$$\theta[V] = (i \cdot \eta(p))^{-\mathrm{rk} V} \frac{\prod_\mu \theta(p; t^\mu)}{\prod_\nu \theta(p; t^\nu)},$$

which satisfies $\theta[\overline{V}] = (-1)^{\mathrm{rk} V} \theta[V]$. Notice that, if $\mathrm{rk} V = 0$, the elliptic measure refines both $[\cdot]$ and $e^{\mathbf{T}}(\cdot)$

$$\theta[V] \xrightarrow{p \rightarrow 0} [V] \xrightarrow{b \rightarrow 0} e^{\mathbf{T}}(V)$$

where the second limit was discussed in § 7.1.

Remark 8.5. The definition we gave for virtual chiral elliptic genus is reminiscent of what is commonly known in physics as the elliptic genus (or superconformal index) of a $2d$ $\mathcal{N} = (0, 2)$ sigma model. Our definition actually matches the one in [33] for $\mathcal{N} = (0, 2)$ Landau–Ginzburg models. Indeed, in this case we are given an n -dimensional (compact) Kähler manifold X together with a holomorphic vector bundle $E \rightarrow X$ such that $c_1(E) - c_1(T_X) = 0 \bmod 2$. If we then consider the K-theory class $[V] = [T_X] - [E]$, the superconformal index of [33] can be written as

$$\mathcal{I}(X, E; p) = (-ip^{-\frac{1}{12}})^{\mathrm{rk} V} \chi \left(X, \mathcal{E}_{1/2}(V) \otimes \det(V^\vee)^{\frac{1}{2}} \right)$$

and in terms of the Chern roots v_i, w_j of T_X and E , respectively, we also have (cf. [24])

$$\mathcal{I}(X, E; p) = \int_X \prod_{i=1}^r \frac{\theta(\tau | \frac{w_i}{2\pi i})}{\eta(p)} \prod_{j=1}^n \frac{v_j \eta(p)}{\theta(\tau | \frac{-v_j}{2\pi i})}.$$

8.2. Elliptic DT invariants.

Definition 8.6. The generating series of elliptic DT invariants $\mathrm{DT}_r^{\mathrm{ell}}(\mathbb{A}^3, q, t, w; p)$ is defined as

$$\mathrm{DT}_r^{\mathrm{ell}}(\mathbb{A}^3, q, t, w; p) = \sum_{n \geq 0} q^n \mathrm{Ell}_{1/2}^{\mathrm{vir}}(\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \mathcal{K}_{\mathrm{vir}}^{\frac{1}{2}}; p) \in \mathbb{Z}((t, t^{\frac{1}{2}}, w))[[p, q]].$$

Being that $\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)$ is not projective, but nevertheless carries the action of an algebraic torus \mathbf{T} with proper \mathbf{T} -fixed locus, we define the invariants by means of virtual localisation, as we explained in § 2.5.

At each \mathbf{T} -fixed point $[S] \in \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$, the localised contribution is

$$\mathrm{tr} \left(\frac{\bigotimes_{n \geq 1} \mathrm{Sym}_{p^n}^{\bullet} (T_S^{\mathrm{vir}} \oplus T_S^{\mathrm{vir}, \vee})}{\widehat{\Lambda}^{\bullet} T_S^{\mathrm{vir}, \vee}} \right)$$

from which we deduce that we can recover the K-theoretic invariants $\chi(\mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n), \widehat{\mathcal{O}}^{\mathrm{vir}})$ in the limit $p \rightarrow 0$. As for K-theoretic invariants, we have

$$\begin{aligned} \mathrm{DT}_r^{\mathrm{ell}}(\mathbb{A}^3, q, t, w; p) &= \sum_{n \geq 0} q^n \sum_{[S] \in \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}} \theta[-T_S^{\mathrm{vir}}] \\ &= \sum_{\bar{\pi}} q^{|\bar{\pi}|} \prod_{i,j=1}^r \theta[-V_{ij}], \end{aligned}$$

where $\bar{\pi}$ runs over all r -colored plane partitions.

Contrary to the case of K-theoretic and cohomological invariants, there exists no conjectural closed formula for elliptic DT invariants yet, even for the rank 1 case. Moreover, the generating series depends on the equivariant parameters of the framing torus, as shown in the following example.

Example 8.7. Consider $Q_1^3 = \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus 3}, 1)$, whose only \mathbf{T} -fixed points are

$$S_1 = \mathcal{I}_{\mathrm{pt}} \oplus \mathcal{O} \oplus \mathcal{O} \subset \mathcal{O}^{\oplus 3}, \quad S_2 = \mathcal{O} \oplus \mathcal{I}_{\mathrm{pt}} \oplus \mathcal{O} \subset \mathcal{O}^{\oplus 3}, \quad S_3 = \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{I}_{\mathrm{pt}} \subset \mathcal{O}^{\oplus 3},$$

with $\mathrm{pt} = (0, 0, 0) \in \mathbb{A}^3$ as in Example 7.5. We have

$$\begin{aligned} T_{S_1}^{\mathrm{vir}} &= 1 - \frac{1}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} - w_1^{-1} w_2 \frac{1}{t_1 t_2 t_3} + w_2^{-1} w_1 - w_1^{-1} w_3 \frac{1}{t_1 t_2 t_3} + w_3^{-1} w_1 \\ T_{S_2}^{\mathrm{vir}} &= 1 - \frac{1}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} - w_2^{-1} w_1 \frac{1}{t_1 t_2 t_3} + w_1^{-1} w_2 - w_2^{-1} w_3 \frac{1}{t_1 t_2 t_3} + w_3^{-1} w_2 \\ T_{S_3}^{\mathrm{vir}} &= 1 - \frac{1}{t_1 t_2 t_3} + \frac{(1-t_1)(1-t_2)(1-t_3)}{t_1 t_2 t_3} - w_3^{-1} w_1 \frac{1}{t_1 t_2 t_3} + w_1^{-1} w_3 - w_3^{-1} w_2 \frac{1}{t_1 t_2 t_3} + w_2^{-1} w_3 \end{aligned}$$

by which we may compute the corresponding elliptic invariant. Set $w_j = e^{2\pi i v_j}$ and $t_\ell = e^{2\pi i s_\ell}$, so that

$$\begin{aligned} \mathrm{Ell}_{1/2}^{\mathrm{vir}}(Q_1^3, \mathcal{K}_{\mathrm{vir}}^{\frac{1}{2}}, t, w; p) &= \diamond \cdot \left(\frac{\theta(\tau|v_2 - v_1 - s) \theta(\tau|v_3 - v_1 - s)}{\theta(\tau|v_1 - v_2) \theta(\tau|v_1 - v_3)} \right. \\ &\quad \left. + \frac{\theta(\tau|v_1 - v_2 - s) \theta(\tau|v_3 - v_2 - s)}{\theta(\tau|v_2 - v_1) \theta(\tau|v_2 - v_3)} + \frac{\theta(\tau|v_1 - v_3 - s) \theta(\tau|v_2 - v_3 - s)}{\theta(\tau|v_3 - v_1) \theta(\tau|v_3 - v_2)} \right), \end{aligned}$$

where $s = s_1 + s_2 + s_3$, with the overall factor

$$\diamond = \frac{\theta(\tau|s_1 + s_2) \theta(\tau|s_1 + s_3) \theta(\tau|s_2 + s_3)}{\theta(\tau|s_1) \theta(\tau|s_2) \theta(\tau|s_3)}.$$

Moreover, by evaluating residues in $v_i - v_j = 0$ one can realise that $\text{Ell}_{1/2}^{\text{vir}}(Q_1^3, \mathcal{K}_{\text{vir}}^{1/2}, t, w; p)$ has no poles in v_i . Indeed

$$\text{Res}_{v_1-v_2=0} \text{Ell}_{1/2}^{\text{vir}}(Q_1^3, \mathcal{K}_{\text{vir}}^{1/2}; p) = \diamond \cdot \left(\frac{\theta(\tau|s)\theta(\tau|v_3-v_2-s)}{\theta(\tau|v_2-v_3)} - \frac{\theta(\tau|s)\theta(\tau|v_3-v_2-s)}{\theta(\tau|v_2-v_3)} \right) = 0,$$

and the same occurs for any other pole involving the v_i 's. However, this does not imply the independence of the elliptic invariants from v , as we now suggest.

Set $\bar{v}_i = v_i + a_i + b_i\tau$, with $a_i, b_i \in \mathbb{Z}$, for $i = 1, 2, 3$. Applying the quasi-periodicity of theta functions, we get

$$\begin{aligned} \text{Ell}_{1/2}^{\text{vir}}(Q_1^3, \mathcal{K}_{\text{vir}}^{1/2}, t, \bar{w}; p) &= \\ &\quad \diamond \frac{\theta(\tau|\bar{v}_1-\bar{v}_2)\theta(\tau|\bar{v}_1-\bar{v}_3)\theta(\tau|\bar{v}_2-\bar{v}_3)}{\theta(\tau|v_1-v_2)\theta(\tau|v_1-v_3)\theta(\tau|v_2-v_3)} \cdot (\theta(\tau|\bar{v}_2-\bar{v}_1-s)\theta(\tau|\bar{v}_3-\bar{v}_1-s)\theta(\tau|\bar{v}_2-\bar{v}_3) \\ &\quad - \theta(\tau|\bar{v}_1-\bar{v}_2-s)\theta(\tau|\bar{v}_3-\bar{v}_2-s)\theta(\tau|\bar{v}_1-\bar{v}_3) + \theta(\tau|\bar{v}_1-\bar{v}_3-s)\theta(\tau|\bar{v}_2-\bar{v}_3-s)\theta(\tau|\bar{v}_1-\bar{v}_2)) \\ &= \diamond \frac{1}{\theta(\tau|v_1-v_2)\theta(\tau|v_1-v_3)\theta(\tau|v_2-v_3)} \cdot (e^{2\pi i s(b_2+b_3-2b_1)}\theta(\tau|v_2-v_1-s)\theta(\tau|v_3-v_1-s)\theta(\tau|v_2-v_3) \\ &\quad - e^{2\pi i s(b_1+b_3-2b_2)}\theta(\tau|v_1-v_2-s)\theta(\tau|v_3-v_2-s)\theta(\tau|v_1-v_3) \\ &\quad + e^{2\pi i s(b_1+b_2-2b_3)}\theta(\tau|v_1-v_3-s)\theta(\tau|v_2-v_3-s)\theta(\tau|v_1-v_2)). \end{aligned}$$

Notice that for general values of s the above expression is different from $\text{Ell}_{1/2}^{\text{vir}}(Q_1^3, \mathcal{K}_{\text{vir}}^{1/2}, t, w; p)$. However, if we specialise $s \in \frac{1}{3}\mathbb{Z}$, we see that in the previous example $\text{Ell}_{1/2}^{\text{vir}}(Q_1^3, \mathcal{K}_{\text{vir}}^{1/2}, t, w; p)$ becomes constant and periodic with respect to v on the lattice $\mathbb{Z} + 3\tau\mathbb{Z}$ and is holomorphic in v , from which we conclude that it is constant on v under this specialisation. Therefore, by choosing $w_j = e^{2\pi i \frac{j}{3}}$ to be third roots of unity, one can show

$$\text{Ell}_{1/2}^{\text{vir}}(Q_1^3, \mathcal{K}_{\text{vir}}^{1/2}, t, w; p) \Big|_{t=e^{2\pi i \frac{k}{3}}} = \begin{cases} (-1)^{m+1}3, & \text{if } k=3m, \quad m \in \mathbb{Z} \\ 0, & \text{if } k \notin 3\mathbb{Z}. \end{cases}$$

8.3. Limits of elliptic DT invariants. Even if a closed formula for the higher rank generating series of elliptic DT invariants is not available, we can still study its behaviour by looking at some particular limits of the variables p, t_i, w_j .

It is easy to see that, under the Calabi–Yau restriction $t = 1$, the generating series of elliptic DT invariants does not carry any more refined information than the cohomological one; in particular, we have no more dependence on the framing parameters w_j and the elliptic parameter p . We generalise this phenomenon in the following setting. Denote by $\mathbf{T}_k \subset \mathbb{T}_1$ the subtorus where $t^{\frac{1}{2}} = e^{\pi i k/r}$, $k \in \mathbb{Z}$. Define by

$$\text{DT}_{r,k}^{\text{ell}}(\mathbb{A}^3, q, t, w; p) = \text{DT}_r^{\text{ell}}(\mathbb{A}^3, q, t, w; p) \Big|_{\mathbf{T}_k}$$

the restriction of the generating series to the subtorus $\mathbf{T}_k \subset \mathbb{T}_1$, which is well-defined as no powers of the Calabi–Yau weight appear in the vertex terms (5.5) by Lemma 3.5.

Proposition 8.8. *If $k = r m \in r\mathbb{Z}$, then*

$$\text{DT}_{r,k}^{\text{ell}}(\mathbb{A}^3, q, t, w; p) = M((-1)^{r(m+1)}q)^r.$$

In particular, the dependence on t_i, w_j and p drops.

Proof. Let $S \in \text{Quot}_{\mathbb{A}^3}(\mathcal{O}^{\oplus r}, n)^{\mathbf{T}}$. Denote $T_S^{\text{vir}} = T_S^{nc} - \overline{T_S^{nc}} t^{-1}$ as in Equation (3.16), where T_S^{nc} is the tangent space of Quot_r^n at S . Denote by $T_{S,l}^{nc}$ the sub-representation of T_S^{nc} corresponding

to t^l , with $l \in \mathbb{Z}$. As there are no powers of the Calabi–Yau weight in T_S^{vir} , we have an identity $T_{S,l}^{nc} = \overline{T_{S,-l-1}^{nc}} t^{-1}$. Set

$$W = T_S^{nc} - \sum_{n \in \mathbb{Z}} T_{S,n}^{nc}.$$

We have that

$$T_S^{nc} - \overline{T_S^{nc}} t^{-1} = W - \overline{W} t^{-1}$$

and, in particular, neither W nor $\overline{W} t^{-1}$ contain constant terms and powers of the Calabi–Yau weight. Using the quasi-periodicity of the theta function $\theta(\tau|z)$, we have

$$\theta[-T_S^{\text{vir}}] = \frac{\theta[\overline{W} t^{-1}]}{\theta[W]} = (-1)^{\text{rk } W} \frac{\theta[\overline{W}]}{\theta[W]} = (-1)^{\text{rk } W(m+1)}.$$

We conclude by noticing that

$$\text{rk } W = \text{rk } T_S^{nc} = r n \pmod{2}. \quad \square$$

Motivated by Example 8.7 and Proposition 8.8, we propose the following conjecture.

Conjecture 8.9. The series $\text{DT}_{r,k}^{\text{ell}}(\mathbb{A}^3, q, t, w; p)$ does not depend on the elliptic parameter p .

Remark 8.10. Notice that the independence from the elliptic parameter p implies that we can reduce our invariants to the K-theoretic ones by setting $p = 0$, i.e.

$$\text{DT}_{r,k}^{\text{ell}}(\mathbb{A}^3, q, t, w; p) = \text{DT}_r^K(\mathbb{A}^3, q, t)|_{\mathbf{T}_k},$$

which in particular do not depend on the framing parameters.

Assuming Conjecture 8.9, we derive a closed expression for $\text{DT}_{r,k}^{\text{ell}}(\mathbb{A}^3, q, t, w; p)$, which was conjectured in [9, Equation (3.20)], motivated by string-theoretic phenomena.

Theorem 8.11. Assume Conjecture 8.9 holds and let $k \in \mathbb{Z}$. Then there is an identity

$$\text{DT}_{r,k}^{\text{ell}}(\mathbb{A}^3, q, t, w; p) = M \left((-1)^{kr} ((-1)^r q)^{\frac{r}{\gcd(k,r)}} \right)^{\gcd(k,r)}.$$

Proof. Assuming Conjecture 8.9, by Remark 8.10 we just have to prove the result for K-theoretic invariants. By Theorem 6.3,

$$\begin{aligned} & \text{DT}_r^K(\mathbb{A}^3, (-1)^r q, t) \\ &= \exp \sum_{n \geq 1} \frac{1}{n} \frac{(1 - t_1^{-n} t_2^{-n})(1 - t_1^{-n} t_3^{-n})(1 - t_2^{-n} t_3^{-n})}{(1 - t_1^{-n})(1 - t_2^{-n})(1 - t_3^{-n})} \frac{1 - t^{-rn}}{1 - t^{-n}} \frac{1}{(1 - t^{-\frac{rn}{2}} q^{-n})(1 - t^{-\frac{rn}{2}} q^n)}. \end{aligned}$$

Assume now that $t^{\frac{1}{2}} = e^{\pi i \frac{k}{r}}$, with $k \in \mathbb{Z}$; we have clearly that $t^{-\frac{rn}{2}} = (-1)^{kn}$. Moreover, we have

$$\frac{1 - t^{-rn}}{1 - t^{-n}} = \begin{cases} r, & \text{if } n \in \frac{r}{\gcd(r,k)} \mathbb{Z} \\ 0, & \text{if } n \notin \frac{r}{\gcd(r,k)} \mathbb{Z} \end{cases}$$

In particular, if $n \in \frac{r}{\gcd(r,k)} \mathbb{Z}$, we have

$$\frac{(1 - t_1^{-n} t_2^{-n})(1 - t_1^{-n} t_3^{-n})(1 - t_2^{-n} t_3^{-n})}{(1 - t_1^{-n})(1 - t_2^{-n})(1 - t_3^{-n})} = -1$$

Setting $n = \frac{r}{\gcd(r,k)} m$, with $m \in \mathbb{Z}$, we have

$$\text{DT}_r^K(\mathbb{A}^3, (-1)^r q, t) = \exp \sum_{m \geq 1} \frac{1}{m} \gcd(r, k) \cdot \frac{-1}{(1 - \overline{q}^{-m})(1 - \overline{q}^m)}$$

where to ease notation we have set $\bar{q} = ((-1)^k q)^{\frac{r}{\gcd(r,k)}}$. We conclude by using the description of the MacMahon function as a plethystic exponential

$$\mathrm{DT}_r^K(\mathbb{A}^3, (-1)^r q, t) = \mathrm{M}\left((-1)^{kr} q^{\frac{r}{\gcd(r,k)}}\right)^{\gcd(r,k)}. \quad \square$$

Remark 8.12. A key technical point in the proof of the conjecture proposed in [9, Equation (3.20)] is the assumption of the independence of $\mathrm{DT}_{r,k}^{\mathrm{ell}}(\mathbb{A}^3)$ on p , as in Conjecture 8.9. We strongly believe it should be possible to prove this assumption by exploiting modular properties of the generating series of elliptic DT invariants. One could proceed by considering the integral representation of the DT invariants given in [9]. The analysis of the K-theoretic case, which we carried out in the proof of Theorem 8.11, shows that no dependence whatsoever is present in the limit $t^{1/2} = e^{\pi i k/r}$. As elliptic DT invariants take the form of meromorphic Jacobi forms, given by quotients of theta functions, poles in the equivariant parameters are only given by shifts along the lattice $\mathbb{Z} + \tau\mathbb{Z}$ of the poles found in K-theoretic DT invariants. Then $\mathrm{DT}_{r,k}^{\mathrm{ell}}(\mathbb{A}^3)$, as a function of each of the equivariant parameters ν_i , $i = 1, \dots, r$, and s_j , $j = 1, 2, 3$, is holomorphic on the torus $\mathbb{C}/\mathbb{Z} + \tau\mathbb{Z}$, so it also carries no dependence on them. This observation may be not very surprising, if one considers the striking resemblance of the chiral virtual elliptic genus to the usual level- N elliptic genus of almost complex manifolds, which is known to be often rigid. In our case, each q -term in the elliptic generating series restricted to \mathbf{T}_k would be now invariant under modular transformations on τ , hence a constant in $p = e^{2\pi i \tau}$.

8.4. Relation to string theory. The definition for the elliptic version of Donaldson–Thomas invariants is motivated by an argument due to Witten [68], which goes as follows: let M be a $2k$ -dimensional spin manifold, and take $\mathcal{L}M = C^0(S^1, M)$ to be the free loop space on M . Then $\mathcal{L}M$ always carries a natural S^1 -action, given by the rotation of loops, so that fixed points under this action of S^1 will only be constant maps $S^1 \rightarrow M$, and $(\mathcal{L}M)^{S^1} \cong M \hookrightarrow \mathcal{L}M$. One can then study the Dirac operator on $\mathcal{L}M$ by formally computing its index using fixed point formulae. In particular, if $D : \Gamma(S_+) \rightarrow \Gamma(S_-)$ is the Dirac operator on M then

$$\mathrm{Ind}(D) = \dim \ker D - \dim \mathrm{coker} D,$$

and whenever M admits the action of a compact Lie group G one can define the G -equivariant index of D as the virtual character

$$\mathrm{Ind}_G(D)(g) = \mathrm{Tr}_{\ker D} g - \mathrm{Tr}_{\mathrm{coker} D} g, \quad g \in G,$$

which only depends on the conjugacy class of g in G . In the case of Dirac operators on loop spaces over spin manifolds, a formal computation yields

$$\mathrm{Ind}_{S^1}(D)(q) = q^{-\frac{d}{24}} \hat{A}(M) \cdot \mathrm{ch} \left(\bigotimes_{n \geq 1} \mathrm{Sym}_{q^n}^\bullet T_M \right) \cap [M],$$

where q denotes a topological generator of S^1 . The \hat{A} -genus is the characteristic class which computes the index of the Dirac complex on a spin manifold. In general, if E is any rank r complex vector bundle on M , one can define $\hat{A}(E)$ in terms of the Chern roots x_i of E as

$$\hat{A}(E) = \prod_{i=1}^r \frac{x_i/2}{\sinh x_i/2},$$

and $\hat{A}(M) = \hat{A}(TM)$, which is completely analogous to the K-theoretic invariants we have been studying so far. The previous formula can also be classically interpreted as the index of a

twisted Dirac operator over the spin structure of M . It is worth noticing that the DT partition functions coming from physics are indeed interpreted as being computing indices of twisted Dirac operators, where the twist by a vector bundle $V \rightarrow M$ makes sense only if $w_2(T_M) = w_2(V)$ so as to extend D to an operator $D: \Gamma(S_+ \otimes V) \rightarrow \Gamma(S_- \otimes V)$. In this same spirit one might also justify the definition of the half-BPS elliptic genus in terms of computations of Euler characteristics of loop spaces over (compact) almost complex manifolds. Let then X be a d -dimensional almost complex manifold, with holomorphic tangent bundle T_X , and whose corresponding free loop space will be denoted by $\mathcal{L}X$. As it was the case also in the previous situation, $\mathcal{L}X$ is naturally equipped with an S^1 action, whose fixed point will be $(\mathcal{L}X)^{S^1} \cong X \hookrightarrow \mathcal{L}X$. By formally applying the virtual localisation formula to the computation of the Euler characteristic of $\mathcal{L}X$ one gets

$$\chi_{S^1}(\mathcal{L}X) = q^{-\frac{d}{12}} \text{td}(X) \cdot \text{ch} \left(\bigotimes_{n \geq 1} \text{Sym}_{q^n}^\bullet(T_X \oplus \Omega_X) \right) \cap [X],$$

which can also be seen as the index of a twisted Spin^c -Dirac operator $\bar{\partial} + \bar{\partial}^*$. Moreover, if $c_1(T_X) = 0 \pmod{2}$, X is also spin, and it is possible to compute the index of the Dirac operator on $\mathcal{L}X$ as before.

9. HIGHER RANK DT INVARIANTS OF COMPACT TORIC 3-FOLDS

Let X be a smooth *projective* toric 3-fold, along with an exceptional locally free sheaf F of rank r . By [59, Thm. A], the Quot scheme $\text{Quot}_X(F, n)$ has a 0-dimensional perfect obstruction theory, so that the rank r Donaldson–Thomas invariant

$$\text{DT}_{F,n} = \int_{[\text{Quot}_X(F,n)]^{\text{vir}}} 1 \in \mathbb{Z}$$

is well defined. In this section we confirm the formula

$$\sum_{n \geq 0} \text{DT}_{F,n} q^n = M((-1)^r q)^r \int_X c_3(T_X \otimes K_X),$$

suggested in [59, Conj. 3.5], in the case where F is *equivariant*. This will prove Theorem C from the Introduction. The next subsection is an interlude on how to induce a torus action on the Quot scheme and on the associated universal short exact sequence starting from an equivariant structure on F . More details are given in [60], including a proof that the obstruction theory obtained in [59, Thm. A] is equivariant (but see Proposition 9.2 for a sketch).

9.1. Inducing a torus action on the Quot scheme. Let X be a quasiprojective toric variety with torus $\mathbb{T} \subset X$. Let $\sigma_X: \mathbb{T} \times X \rightarrow X$ denote the action. If F is a \mathbb{T} -equivariant coherent sheaf on X , and $Q = \text{Quot}_X(F, n)$, then σ_X has a canonical lift

$$\sigma_Q: \mathbb{T} \times Q \rightarrow Q.$$

This is proved in [37, Prop. 4.1], but we sketch here the argument for the sake of completeness⁴. Let $p_2: \mathbb{T} \times X \rightarrow X$ be the second projection and let $\vartheta: p_2^* F \xrightarrow{\sim} \sigma_X^* F$ denote the chosen

⁴We thank Martijn Kool for guiding us through the details of this construction.

\mathbb{T} -equivariant structure on F . Let $\pi_X: X \times Q \rightarrow Q$ and $\pi_Q: X \times Q \rightarrow Q$ be the projections, and set $F_Q = \pi_X^* F$. Consider the universal exact sequence

$$(9.1) \quad \begin{array}{ccc} & S \hookrightarrow F_Q \xrightarrow{u} \mathcal{T} & \\ & \downarrow & \\ & X \times Q & \\ \swarrow \pi_X & & \searrow \pi_Q \\ X & & Q \end{array}$$

and note that there is a commutative diagram

$$\begin{array}{ccc} \mathbb{T} \times X \times Q & \xrightarrow{\sigma_X \times \text{id}_Q} & X \times Q \\ p_{12} \downarrow & & \downarrow \pi_X \\ \mathbb{T} \times X & \xrightarrow{\sigma_X} & X \end{array}$$

yielding an identity $(\sigma_X \times \text{id}_Q)^* F_Q = p_{12}^* \sigma_X^* F$. The induced surjection

$$(\sigma_X \times \text{id}_Q)^* u \circ p_{12}^* \vartheta: F_{\mathbb{T} \times Q} \xrightarrow{\sim} p_{12}^* \sigma_X^* F \twoheadrightarrow (\sigma_X \times \text{id}_Q)^* \mathcal{T}$$

defines a $\mathbb{T} \times Q$ -valued point of Q , i.e. a morphism

$$\sigma_Q: \mathbb{T} \times Q \rightarrow Q.$$

It is straightforward to verify that σ_Q satisfies the axioms for a \mathbb{T} -action.

Next, we explain how to make the universal exact sequence (9.1) \mathbb{T} -equivariant. The universal property of the Quot scheme applied to σ_Q implies that there is an isomorphism of surjections

$$(\text{id}_X \times \sigma_Q)^* u \xrightarrow{\sim} (\sigma_X \times \text{id}_Q)^* u \circ p_{12}^* \vartheta.$$

This means that there is a commutative diagram

$$(9.2) \quad \begin{array}{ccccc} (\text{id}_X \times \sigma_Q)^* F_Q & \xrightarrow{(\text{id}_X \times \sigma_Q)^* u} & & (\text{id}_X \times \sigma_Q)^* \mathcal{T} & \\ \parallel & & & \downarrow \wr & \\ p_{12}^* p_2^* F & \xrightarrow{p_{12}^* \vartheta} & p_{12}^* \sigma_X^* F & \xrightarrow{(\sigma_X \times \text{id}_Q)^* u} & (\sigma_X \times \text{id}_Q)^* \mathcal{T} \end{array}$$

where we used the identity $(\sigma_X \times \text{id}_Q)^* F_Q = p_{12}^* \sigma_X^* F$ in the bottom row.

Consider the morphism

$$\varphi: X \times Q \times \mathbb{T} \rightarrow X \times Q, \quad (x, f, t) \mapsto (\sigma_X(t, x), \sigma_Q(t^{-1}, f)).$$

We view this as a \mathbb{T} -action on $X \times Q$. Note that $\pi_X \circ \varphi = \sigma_X \circ p_{12}$. The moduli map $Q \times \mathbb{T} \rightarrow Q$ corresponding to the family of quotients

$$\varphi^* u \circ p_{12}^* \vartheta: F_{Q \times \mathbb{T}} = p_{12}^* p_2^* F \xrightarrow{\sim} p_{12}^* \sigma_X^* F = \varphi^* F_Q \xrightarrow{\varphi^* u} \varphi^* \mathcal{T}$$

is easily seen to agree with the first projection $p_1: Q \times \mathbb{T} \rightarrow Q$. Indeed, if \mathcal{T}_f denotes the quotient of F corresponding to a point $f = [F \twoheadrightarrow \mathcal{T}_f] \in Q$, it is immediate to see that

$$\varphi^* \mathcal{T} \big|_{X \times f \times t} = \mathcal{T}_{\sigma_Q(\sigma_Q(t, f), t^{-1})} = \mathcal{T}_{\sigma_Q(t t^{-1}, f)} = \mathcal{T}_f$$

for all $t \in \mathbb{T}$.

Let then $q = \text{id}_X \times p_1: X \times Q \times \mathbb{T} \rightarrow X \times Q$ be the projection. Since $\varphi^* u \circ p_{12}^* \vartheta$ corresponds to the projection $p_1: Q \times \mathbb{T} \rightarrow Q$, by the universal property of (Q, u) we obtain an isomorphism

of surjections $q^*u \xrightarrow{\sim} \varphi^*u$ that, after setting $\mathcal{S} = \ker(u: F_Q \rightarrow \mathcal{T})$, we can extend to an isomorphism of short exact sequences

$$\begin{array}{ccccc} q^*\mathcal{S} & \hookrightarrow & F_{Q \times \mathbb{T}} & \xrightarrow{q^*u} & q^*\mathcal{T} \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \varphi^*\mathcal{S} & \hookrightarrow & \varphi^*F_Q & \xrightarrow{\varphi^*u} & \varphi^*\mathcal{T} \end{array}$$

on $X \times Q \times \mathbb{T}$, where the middle vertical isomorphism is $p_{12}^*\vartheta$ and is a \mathbb{T} -equivariant structure on F_Q because ϑ is. The diagram allows us to conclude that a \mathbb{T} -equivariant structure on F induces a canonical \mathbb{T} -equivariant structure on the universal short exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow F_Q \rightarrow \mathcal{T} \rightarrow 0.$$

9.2. The (equivariant) obstruction theory. Throughout this subsection, F denotes an exceptional locally free sheaf of rank r on a smooth projective toric 3-fold X . In other words, F is simple, i.e. $\mathrm{Hom}(F, F) = \mathbb{C}$, and $\mathrm{Ext}^i(F, F) = 0$ for $i > 0$.

By [59, Thm. A], there is a 0-dimensional perfect obstruction theory

$$(9.3) \quad \mathbb{E} \rightarrow \mathbb{L}_{\mathrm{Quot}_X(F, n)},$$

governed by

$$\mathrm{Def}|_{[S]} = \mathrm{Ext}^1(\mathcal{S}, \mathcal{S}), \quad \mathrm{Obs}|_{[S]} = \mathrm{Ext}^2(\mathcal{S}, \mathcal{S})$$

around a point $[S] \in \mathrm{Quot}_X(F, n)$. We set $Q = \mathrm{Quot}_X(F, n)$ for brevity, and we denote by π_Q and π_X the projections from $X \times Q$, as in (9.1). Note that $\omega_{\pi_Q} = \pi_X^*\omega_X$.

As we recall during the (sketch of) proof of Proposition 9.2 below, we have

$$\mathbb{E} = \mathbf{R}\pi_{Q*}(\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0 \otimes \omega_{\pi_Q})[2]$$

where $\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0$ is the shifted cone of the trace map $\mathrm{tr}: \mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S}) \rightarrow \mathcal{O}_{X \times Q}$.

Proposition 9.1. *There is an identity*

$$\mathbb{E}^\vee = \mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(F_Q, F_Q) - \mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S}) \in K_0(\mathrm{Quot}_X(F, n)).$$

Proof. As in the proof of [59, Theorem 2.5] we have

$$\begin{aligned} \mathbb{E} &= (\mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0)^\vee[-1] \\ &= (\mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S}))^\vee[-1] - (\mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(\mathcal{O}, \mathcal{O}))^\vee[-1] \\ &= (\mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S}))^\vee[-1] - (\mathbf{R}\pi_{Q*}\mathbf{R}\mathcal{H}om(F_Q, F_Q))^\vee[-1], \end{aligned}$$

where the last identity uses that F is an exceptional sheaf. \square

The following result is proved in [60, Thm. B] in greater generality, but we sketch a proof here for the reader's convenience. Denote by $\mathbb{T} = (\mathbb{C}^*)^3$ the torus of X .

Proposition 9.2. *Let (X, F) be a pair consisting of a smooth projective toric 3-fold X along with an exceptional locally free \mathbb{T} -equivariant sheaf F . Then the perfect obstruction theory (9.3) on $\mathrm{Quot}_X(F, n)$ is \mathbb{T} -equivariant.*

The definition of equivariant obstruction theory was recalled in Definition 2.2.

Proof. As we explained in § 9.1, by equivariance of F the Quot scheme $Q = \text{Quot}_X(F, n)$ inherits a canonical \mathbb{T} -action with respect to which the universal short exact sequence $\mathcal{S} \hookrightarrow F_Q \twoheadrightarrow \mathcal{T}$ can be made \mathbb{T} -equivariant. On the other hand, the perfect obstruction theory (9.3) is obtained by projecting the truncated Atiyah class

$$\text{At}_{\mathcal{S}} \in \text{Ext}_{X \times Q}^1(\mathcal{S}, \mathcal{S} \otimes \mathbb{L}_{X \times Q})$$

onto the Ext group

$$\begin{aligned} \text{Ext}_{X \times Q}^1(\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0, \pi_Q^* \mathbb{L}_Q) &= \text{Ext}_{X \times Q}^{-2}(\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0 \otimes \omega_{\pi_Q}, \pi_Q^* \mathbb{L}_Q \otimes \omega_{\pi_Q}[3]) \\ (9.4) \quad &= \text{Ext}_{X \times Q}^{-2}(\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0 \otimes \omega_{\pi_Q}, \pi_Q^! \mathbb{L}_Q) \\ &\cong \text{Ext}_Q^{-2}(\mathbf{R}\pi_{Q*}(\mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S})_0 \otimes \omega_{\pi_Q}), \mathbb{L}_Q). \end{aligned}$$

The last isomorphism is Grothendieck duality along the smooth projective morphism π_Q . Now we need three ingredients to finish the proof:

- The Atiyah class $\text{At}_{\mathcal{S}}$ is a \mathbb{T} -invariant extension,
- Grothendieck duality preserves \mathbb{T} -invariant extensions, and
- \mathbb{T} -invariant extensions correspond to morphisms in the equivariant derived category.

These assertions are proved in [60]. \square

We let $\Delta(X)$ denote the set of vertices in the Newton polytope of the toric 3-fold X . Then

$$X^{\mathbb{T}} = \{ p_{\alpha} \mid \alpha \in \Delta(X) \} \subset X$$

will denote the fixed locus of X . For a given vertex α , let $U_{\alpha} \cong \mathbb{A}^3$ be the canonical chart containing the fixed point p_{α} . The \mathbb{T} -action on this chart can be taken to be the standard action (3.5). For every α , there is a \mathbb{T} -equivariant open immersion

$$\iota_{n,\alpha}: \text{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n) \hookrightarrow Q = \text{Quot}_X(F, n)$$

parametrising quotients whose support is contained in U_{α} . We think of $F|_{U_{\alpha}}$ as an equivariant sheaf on \mathbb{A}^3 , hence of the form described in (5.9). We denote by $\mathbb{E}_{n,\alpha}^{\text{crit}}$ the critical obstruction theory on $\text{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n)$ from Proposition 3.2.

It is natural to ask whether the restriction of the global perfect obstruction theory (9.3) along $\iota_{n,\alpha}$ agrees with the critical symmetric perfect obstruction theory described in § 3.2 (see Conjecture 9.8). However, what we really need is the following weaker result.

Proposition 9.3. *Let $\mathbb{E} \in K_0(Q)$ be the class of the global perfect obstruction theory (9.3). Then*

$$\mathbb{E}_{n,\alpha}^{\text{crit}} = \iota_{n,\alpha}^* \mathbb{E} \in K_0(\text{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n)).$$

Considering the two obstruction theories as \mathbb{T} -equivariant, the same identity holds in equivariant K-theory:

$$\mathbb{E}_{n,\alpha}^{\text{crit}} = \iota_{n,\alpha}^* \mathbb{E} \in K_0^{\mathbb{T}}(\text{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n)).$$

Proof. The chart U_{α} is Calabi–Yau, so by [59, Prop. 2.9] the induced perfect obstruction theory $\iota_{n,\alpha}^* \mathbb{E}$ is symmetric. Since by Remark 2.1 all symmetric perfect obstruction theories share the same class in K-theory, the first statement follows.

To prove the \mathbb{T} -equivariant equality, we need a slightly more refined analysis. Just for this proof, let us shorten

$$\mathbb{E}_{\text{cr}} = \mathbb{E}_{n,\alpha}^{\text{crit}} \quad \text{and} \quad \mathbb{E} = \iota_{n,\alpha}^* \mathbb{E},$$

to ease notation. We know by Diagram (3.15) that we can write

$$(9.5) \quad \mathbb{E}_{\text{cr}} = [\mathfrak{t} \otimes T_{\text{Quot}_r^n}|_Q \rightarrow \Omega_{\text{Quot}_r^n}|_Q] = \Omega - \mathfrak{t} \otimes T \in K_0^{\mathbb{T}}(\text{Quot}_{U_a}(F|_{U_a}, n)),$$

where Ω (resp. T) denotes the cotangent sheaf (resp. the tangent sheaf) of $\text{Quot}_{U_a}(F|_{U_a}, n)$, equipped with its natural equivariant structure.

Let $\pi: U_a \times \text{Quot}_{U_a}(F|_{U_a}, n) \rightarrow \text{Quot}_{U_a}(F|_{U_a}, n)$ be the projection, let S be the universal kernel living on the product and set $\mathfrak{t}_\pi = \pi^* \mathfrak{t}^{-1}$. By definition,

$$\mathbb{E} = \mathbf{R}\pi_*(\mathbf{R}\mathcal{H}om(S, S)_0 \otimes \omega_\pi)[2].$$

The equivariant isomorphism $\omega_\pi \xrightarrow{\sim} \mathcal{O} \otimes \mathfrak{t}_\pi^{-1}$ along with the projection formula yield

$$(9.6) \quad \mathfrak{t}^{-1} \otimes \mathbb{E} \xrightarrow{\sim} \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(S, S)_0[2].$$

We next show the right hand side is canonically isomorphic to $\mathbb{E}^\vee[1]$. We have

$$\begin{aligned} \mathbb{E}^\vee[1] &= \mathbf{R}\mathcal{H}om(\mathbf{R}\pi_*(\mathbf{R}\mathcal{H}om(S, S)_0 \otimes \omega_\pi), \mathcal{O})[-1] && \text{definition of } (-)^\vee \\ &= \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathbf{R}\mathcal{H}om(S, S)_0 \otimes \omega_\pi, \omega_\pi[3])[-1] && \text{Grothendieck duality} \\ &= \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathbf{R}\mathcal{H}om(S, S)_0, \mathcal{O})[2] && \text{shift} \\ &= \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(S, S)_0^\vee[2] && \text{definition of } (-)^\vee \\ &= \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(S, S)_0[2] && \mathbf{R}\mathcal{H}om(S, S)_0 \text{ is self-dual} \end{aligned}$$

in the derived category of \mathbb{T} -equivariant coherent sheaves on $\text{Quot}_{U_a}(F|_{U_a}, n)$, which by (9.6) proves that

$$\mathfrak{t}^{-1} \otimes \mathbb{E} \cong \mathbb{E}^\vee[1].$$

We thus have

$$\Omega \cong h^0(\mathbb{E}) \cong \mathfrak{t} \otimes h^0(\mathbb{E}^\vee[1]) \cong \mathfrak{t} \otimes \mathcal{E}xt_\pi^2(S, S),$$

where we use the standard notation $\mathcal{E}xt_\pi^i(-, -)$ for the i th derived functor of $\pi_* \circ \mathcal{H}om(-, -)$. We conclude

$$\begin{aligned} \mathbb{E} &= h^0(\mathbb{E}) - h^{-1}(\mathbb{E}) \\ &= \Omega - h^1(\mathbb{E}^\vee)^\vee \\ &= \Omega - h^0(\mathbb{E}^\vee[1])^\vee \\ &= \Omega - \mathcal{E}xt_\pi^2(S, S)^\vee \\ &= \Omega - (\mathfrak{t}^{-1} \otimes \Omega)^\vee \\ &= \Omega - \mathfrak{t} \otimes T \\ &= \mathbb{E}_{\text{cr}}. \end{aligned} \quad \square$$

9.3. The fixed locus of the Quot scheme and its virtual class. In this subsection we describe $\text{Quot}_X(F, n)^\mathbb{T}$ and we compute its virtual fundamental class, obtained via Proposition 9.2.

If \mathbf{n} denotes a generic tuple $\{n_\alpha \mid \alpha \in \Delta(X)\}$ of non-negative integers, we set $|\mathbf{n}| = \sum_{\alpha \in \Delta(X)} n_\alpha$.

Lemma 9.4. *There is a scheme-theoretic identity*

$$\text{Quot}_X(F, n)^\mathbb{T} = \coprod_{|\mathbf{n}|=n} \prod_{\alpha \in \Delta(X)} \text{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T}.$$

Proof. Let B be a (connected) scheme over \mathbb{C} . Let F_B be the pullback of F along the first projection $X \times B \rightarrow X$, and fix a B -flat family of quotients

$$\rho: F_B \twoheadrightarrow \mathcal{T}$$

defining a B -valued point $B \rightarrow \mathrm{Quot}_X(F, n)^\mathbb{T}$. Then, by restriction, we obtain, for each $\alpha \in \Delta(X)$, a B -flat family of quotients

$$(9.7) \quad \rho_\alpha: F_B|_{U_\alpha \times B} \twoheadrightarrow \mathcal{T}_\alpha = \mathcal{T}|_{U_\alpha \times B},$$

and we let n_α be the length of the fibres of \mathcal{T}_α . Each ρ_α corresponds to a B -valued point $g_\alpha: B \rightarrow \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T}$, thus we obtain a B -valued point

$$(g_\alpha)_\alpha: B \rightarrow \prod_{\alpha \in \Delta(X)} \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T}.$$

Note that the original family \mathcal{T} is recovered as the direct sum $\bigoplus_\alpha \mathcal{T}_\alpha$, in particular $n = \sum_\alpha n_\alpha$. Conversely, suppose given a tuple of B -families of \mathbb{T} -fixed quotients

$$\left((F|_{U_\alpha})_B \twoheadrightarrow \mathcal{T}_\alpha \right)_\alpha.$$

We obtain B -valued points

$$B \rightarrow \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T} \subset \mathrm{Quot}_X(F, n_\alpha)^\mathbb{T}.$$

Since the support of these families is disjoint, we can form the direct sum

$$\mathcal{T} = \bigoplus_\alpha \mathcal{T}_\alpha$$

to obtain a new B -flat family, representing a B -valued point of $\mathrm{Quot}_X(F, n)^\mathbb{T}$, as required. \square

Our next goal is to show that, under the identification of Lemma 9.4, the induced virtual fundamental class of the \mathbf{n} -th connected component of $\mathrm{Quot}_X(F, n)^\mathbb{T}$ is the box product of the virtual fundamental classes of $\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T}$, whose perfect obstruction theory is the \mathbb{T} -fixed part of the critical one, studied in § 3.2. For the rest of the section we restrict our attention to each connected component

$$(9.8) \quad \mathrm{Quot}_X(F, n)^\mathbb{T}_\mathbf{n} = \prod_{\alpha \in \Delta(X)} \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T} \subset \mathrm{Quot}_X(F, n)^\mathbb{T},$$

and we denote by

$$(9.9) \quad \begin{array}{ccc} \mathcal{S} \hookrightarrow \mathcal{F} \twoheadrightarrow \mathcal{T} & & \mathcal{S}_\alpha \hookrightarrow \mathcal{F}_\alpha \twoheadrightarrow \mathcal{T}_\alpha \\ \downarrow & & \downarrow \\ X \times \mathrm{Quot}_X(F, n)^\mathbb{T}_\mathbf{n} & & U_\alpha \times \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T} \\ \downarrow \pi & & \downarrow \pi_\alpha \\ \mathrm{Quot}_X(F, n)^\mathbb{T}_\mathbf{n} & \xrightarrow{p_\alpha} & \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T} \end{array}$$

the various universal structures and projection maps between these moduli spaces. For instance, \mathcal{F}_α is the pullback of $F|_{U_\alpha}$ along the projection $U_\alpha \times \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^\mathbb{T} \rightarrow U_\alpha$. Let $\mathbb{E}_\mathbf{n}$ be the restriction of $\mathbb{E} \in \mathbf{D}(\mathrm{Quot}_X(F, n))$ to the closed subscheme $\mathrm{Quot}_X(F, n)^\mathbb{T}_\mathbf{n} \subset \mathrm{Quot}_X(F, n)$.

Proposition 9.5. *There are identities in $K_0^{\mathbb{T}}(\mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}})$*

$$\begin{aligned} \mathbb{E}_{\mathbf{n}}^{\vee} &= \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{F}, \mathcal{F}) - \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S}) \\ &= \sum_{\alpha \in \Delta(X)} p_{\alpha}^* (\mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{F}_{\alpha}, \mathcal{F}_{\alpha}) - \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{S}_{\alpha}, \mathcal{S}_{\alpha})). \end{aligned}$$

Proof. Exploiting the universal short exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow \mathcal{F} \rightarrow \mathcal{T} \rightarrow 0$$

on $X \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}} \subset X \times \mathrm{Quot}_X(F, n)$, and Proposition 9.1, we may write

$$\begin{aligned} \mathbb{E}_{\mathbf{n}}^{\vee} &= \mathbb{E}^{\vee}|_{\mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}} = \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{F}, \mathcal{F}) - \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{S}) \\ &= \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{S}, \mathcal{T}) + \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{T}, \mathcal{S}) + \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(\mathcal{T}, \mathcal{T}). \end{aligned}$$

Similarly, we have

$$\begin{aligned} (9.10) \quad \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{F}_{\alpha}, \mathcal{F}_{\alpha}) - \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{S}_{\alpha}, \mathcal{S}_{\alpha}) \\ = \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{S}_{\alpha}, \mathcal{T}_{\alpha}) + \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{T}_{\alpha}, \mathcal{S}_{\alpha}) + \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(\mathcal{T}_{\alpha}, \mathcal{T}_{\alpha}). \end{aligned}$$

In the following, we write (G_1, G_2) for any of the three pairs $(\mathcal{S}, \mathcal{T})$, $(\mathcal{T}, \mathcal{S})$ or $(\mathcal{T}, \mathcal{T})$. Applying the Grothendieck spectral sequence yields

$$\begin{aligned} \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(G_1, G_2) &= \sum_{i,j} (-1)^{i+j} \mathbf{R}^i \pi_* \mathcal{E}xt^j(G_1, G_2) \\ &= \sum_j (-1)^j \pi_* \mathcal{E}xt^j(G_1, G_2), \end{aligned}$$

where we used cohomology and base change along with the fact that $\mathbf{R}^i \pi_*$ of a 0-dimensional sheaf vanishes for $i > 0$. The standard Čech cover $\{U_{\alpha}\}_{\alpha \in \Delta(X)}$ of X pulls back to a Čech cover $\{V_{\alpha}\}_{\alpha \in \Delta(X)}$ of $X \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}$, where $V_{\alpha} = U_{\alpha} \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}$. For a finite family of indices $I \subset \mathbb{N}$, set $V_I = \bigcap_{\alpha \in I} V_{\alpha}$ and let $j_I: V_I \rightarrow X \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}$ be the natural open immersion. We have a Čech resolution $\mathcal{E}xt^j(G_1, G_2) \rightarrow \mathfrak{C}^{\bullet}$, where \mathfrak{C}^{\bullet} is defined degree-wise (see e.g. [29, Lemma III.4.2]) by

$$\mathfrak{C}^k = \bigoplus_{|I|=k+1} j_{I*} j_I^* \mathcal{E}xt^j(G_1, G_2).$$

Notice that \mathcal{T} vanishes on the restriction to any double intersection $U_{\alpha\beta} \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}$, where $U_{\alpha\beta} = U_{\alpha} \cap U_{\beta}$. This implies that the only contribution of the Čech cover is given by \mathfrak{C}^0 , thus

$$\begin{aligned} \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(G_1, G_2) &= \sum_j (-1)^j \pi_* \sum_{\alpha \in \Delta(X)} j_{\alpha*} j_{\alpha}^* \mathcal{E}xt^j(G_1, G_2) \\ &= \sum_{\alpha \in \Delta(X)} \sum_j (-1)^j (\pi \circ j_{\alpha})_* j_{\alpha}^* \mathcal{E}xt^j(G_1, G_2). \end{aligned}$$

Consider the following cartesian diagram

$$\begin{array}{ccc} U_{\alpha} \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}} & \xrightarrow{\tilde{p}_{\alpha}} & U_{\alpha} \times \mathrm{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n_{\alpha})^{\mathbb{T}} \\ \downarrow j_{\alpha} & & \downarrow \\ X \times \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}} & \longrightarrow & X \times \mathrm{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n_{\alpha})^{\mathbb{T}} \\ \downarrow \pi & & \downarrow \pi_{\alpha} \\ \mathrm{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}} & \xrightarrow{p_{\alpha}} & \mathrm{Quot}_{U_{\alpha}}(F|_{U_{\alpha}}, n_{\alpha})^{\mathbb{T}} \end{array}$$

As it was already clear from the proof of Lemma 9.4, the universal short exact sequences in Diagram (9.9) satisfy $j_\alpha^*(\mathcal{S} \hookrightarrow \mathcal{F} \twoheadrightarrow \mathcal{T}) = \tilde{p}_\alpha^*(\mathcal{S}_\alpha \hookrightarrow \mathcal{F}_\alpha \twoheadrightarrow \mathcal{T}_\alpha)$. If $(G_{1\alpha}, G_{2\alpha})$ denotes any of the pairs belonging to the set $\{(\mathcal{S}_\alpha, \mathcal{T}_\alpha), (\mathcal{T}_\alpha, \mathcal{S}_\alpha), (\mathcal{T}_\alpha, \mathcal{T}_\alpha)\}$, we can write

$$\begin{aligned} j_\alpha^* \mathcal{E}xt^j(G_1, G_2) &= \mathbf{L}j_\alpha^* \mathcal{E}xt^j(G_1, G_2) \\ &= \mathcal{E}xt^j(\mathbf{L}j_\alpha^* G_1, \mathbf{L}j_\alpha^* G_2) \\ &= \mathcal{E}xt^j(\tilde{p}_\alpha^* G_{1,\alpha}, \tilde{p}_\alpha^* G_{2,\alpha}) \\ &= \tilde{p}_\alpha^* \mathcal{E}xt^j(G_{1\alpha}, G_{2\alpha}). \end{aligned}$$

We deduce, by flat base change,

$$(\pi \circ j_\alpha)_* j_\alpha^* \mathcal{E}xt^j(G_1, G_2) = (\pi \circ j_\alpha)_* \tilde{p}_\alpha^* \mathcal{E}xt^j(G_{1\alpha}, G_{2\alpha}) = p_\alpha^* \pi_{\alpha*} \mathcal{E}xt^j(G_{1\alpha}, G_{2\alpha}).$$

Combining again the Grothendieck spectral sequence, cohomology and base change and the vanishing of higher derived pushforwards on 0-dimensional sheaves, we conclude that

$$\begin{aligned} \mathbf{R}\pi_* \mathbf{R}\mathcal{H}om(G_1, G_2) &= \sum_{\alpha \in \Delta(X)} p_\alpha^* \sum_j (-1)^j \pi_{\alpha*} \mathcal{E}xt^j(G_{1\alpha}, G_{2\alpha}) \\ &= \sum_{\alpha \in \Delta(X)} p_\alpha^* \mathbf{R}\pi_{\alpha*} \mathbf{R}\mathcal{H}om(G_{1\alpha}, G_{2\alpha}). \end{aligned}$$

Now the result follows from Equation (9.10). \square

Corollary 9.6. *The virtual fundamental class of $\text{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}$ is expressed as the product of the virtual fundamental classes*

$$[\text{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}]^{\text{vir}} = \prod_{\alpha \in \Delta(X)} p_\alpha^* [\text{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}}]^{\text{vir}}.$$

Before we prove the corollary, let us explain what virtual classes are involved. The left hand side is the virtual class induced by the \mathbb{T} -fixed obstruction theory

$$\mathbb{E}_{\mathbf{n}}^{\mathbb{T}\text{-fix}} \rightarrow \mathbb{L}_{\text{Quot}_X(F, n)_{\mathbf{n}}^{\mathbb{T}}},$$

whereas $[\text{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}}]^{\text{vir}}$ is the virtual class induced by the obstruction theory

$$\iota_{n_\alpha, \alpha}^* \mathbb{E} \rightarrow \mathbb{L}_{\text{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)}$$

by restricting to the \mathbb{T} -fixed locus and taking the \mathbb{T} -fixed part. Note that by Proposition 9.3, the perfect obstruction theory

$$\mathbb{E}_{n_\alpha, \alpha}^{\text{crit}} \Big|_{\text{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}}}^{\mathbb{T}\text{-fix}} \rightarrow \mathbb{L}_{\text{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}}}$$

induces the same virtual class. This follows from the general fact that the (equivariant) virtual fundamental class depends only on the class in (equivariant) K-theory of the perfect obstruction theory — cf. [61, Theorem 4.6], where all the ingredients are naturally equivariant.

Proof. The statement follows by taking \mathbb{T} -fixed parts in Proposition 9.5 and by Siebert's result [61, Theorem 4.6] mentioned above. \square

9.4. Higher rank Donaldson–Thomas invariants of compact 3-folds. For a pair (X, F) consisting of a smooth projective toric 3-fold X and an exceptional locally free sheaf F , the perfect obstruction theory (9.3) gives rise to a 0-dimensional virtual fundamental class

$$[\mathrm{Quot}_X(F, n)]^{\mathrm{vir}} \in A_0(\mathrm{Quot}_X(F, n)),$$

allowing one to define higher rank Donaldson–Thomas invariants

$$\mathrm{DT}_{F,n} = \int_{[\mathrm{Quot}_X(F,n)]^{\mathrm{vir}}} 1 \in \mathbb{Z}.$$

Define the generating function

$$\mathrm{DT}_F(q) = \sum_{n \geq 0} \mathrm{DT}_{F,n} q^n.$$

We next compute this series in the case of a \mathbb{T} -equivariant exceptional locally free sheaf, thus proving Theorem C from the Introduction.

Theorem 9.7. *Let (X, F) be a pair consisting of a smooth projective toric 3-fold X along with an exceptional \mathbb{T} -equivariant locally free sheaf F . Then*

$$\mathrm{DT}_F(q) = M((-1)^r q)^r \int_X c_3(T_X \otimes K_X).$$

Proof. Set $Q = \mathrm{Quot}_X(F, n)$ and $Q_\alpha = \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)$. Since by Proposition 9.2 the perfect obstruction theory on Q is \mathbb{T} -equivariant, we can apply the virtual localisation formula

$$\mathrm{DT}_{F,n} = \int_{[Q^\mathbb{T}]^{\mathrm{vir}}} e^\mathbb{T}(-N_{Q^\mathbb{T}/Q}^{\mathrm{vir}}),$$

where $N_{Q^\mathbb{T}/Q}^{\mathrm{vir}}$ is the virtual normal bundle on the \mathbb{T} -fixed locus computed in Lemma 9.4. By taking \mathbb{T} -moving parts in Proposition 9.5, we obtain the K-theoretic identity

$$N_{Q^\mathbb{T}/Q}^{\mathrm{vir}} = \sum_{\alpha \in \Delta(X)} p_\alpha^* N_{Q_\alpha^\mathbb{T}/Q_\alpha}^{\mathrm{vir}}$$

of virtual normal bundles. Thus by Corollary 9.6 we have

$$\int_{[Q^\mathbb{T}]^{\mathrm{vir}}} e^\mathbb{T}(-N_{Q^\mathbb{T}/Q}^{\mathrm{vir}}) = \sum_{|\mathbf{n}|=n} \prod_{\alpha \in \Delta(X)} \int_{[\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}_1}]^{\mathrm{vir}}} e^\mathbb{T}(-N_{Q_\alpha^\mathbb{T}/Q_\alpha}^{\mathrm{vir}}).$$

In particular, the virtual fundamental class $[\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}}]^{\mathrm{vir}}$ agrees with the one coming from the critical structure. Moreover, by the virtual localisation formula applied with respect to $(\mathbb{C}^*)^r$, we have

$$\int_{[\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)^{\mathbb{T}}]^{\mathrm{vir}}} e^\mathbb{T}(-N_{Q_\alpha^\mathbb{T}/Q_\alpha}^{\mathrm{vir}}) = \int_{[\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)]^{\mathrm{vir}}} 1,$$

where the right hand side is defined equivariantly in § 5.2. Finally, by Corollary 7.4, we have an identity

$$\int_{[\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n_\alpha)]^{\mathrm{vir}}} 1 = \int_{[\mathrm{Quot}_{U_\alpha}(\mathcal{O}_{U_\alpha}^{\oplus r}, n_\alpha)]^{\mathrm{vir}}} 1$$

of equivariant integrals, where in the right hand side we take $\mathcal{O}_{U_\alpha}^{\oplus r}$ with the trivial \mathbb{T} -equivariant weights. Therefore we conclude

$$\begin{aligned} \mathrm{DT}_F(q) &= \sum_{n \geq 0} q^n \sum_{|\mathbf{n}|=n} \prod_{\alpha \in \Delta(X)} \int_{[\mathrm{Quot}_{U_\alpha}(\mathcal{O}_{U_\alpha}^{\oplus r}, n_\alpha)]^{\mathrm{vir}}} 1 \\ &= \prod_{\alpha \in \Delta(X)} \sum_{n_\alpha \geq 0} q^{n_\alpha} \int_{[\mathrm{Quot}_{U_\alpha}(\mathcal{O}_{U_\alpha}^{\oplus r}, n_\alpha)]^{\mathrm{vir}}} 1 \\ &= \prod_{\alpha \in \Delta(X)} M((-1)^r q)^{-r \frac{(s_1^\alpha + s_2^\alpha)(s_1^\alpha + s_3^\alpha)(s_2^\alpha + s_3^\alpha)}{s_1^\alpha s_2^\alpha s_3^\alpha}}. \end{aligned}$$

We have used Theorem 7.2 to obtain the last identity, in which we have denoted $s_1^\alpha, s_2^\alpha, s_3^\alpha$ the tangent weights at p_α . We conclude taking logarithms:

$$\begin{aligned} \log \mathrm{DT}_F(q) &= \sum_{\alpha \in \Delta(X)} -r \frac{(s_1^\alpha + s_2^\alpha)(s_1^\alpha + s_3^\alpha)(s_2^\alpha + s_3^\alpha)}{s_1^\alpha s_2^\alpha s_3^\alpha} \log M((-1)^r q) \\ &= r \int_X c_3(T_X \otimes K_X) \cdot \log M((-1)^r q) \end{aligned}$$

where the prefactor is computed through ordinary Atiyah–Bott localisation. \square

We have thus proved Conjecture 3.5 in [59] in the toric case. The general case is still open and will be investigated in future work.

9.5. Conjecture: two obstruction theories are the same. We close this subsection with a couple of conjectures relating the different obstruction theories appeared in the previous section.

Conjecture 9.8. Let \mathbb{E} be the perfect obstruction theory (9.3). Then its restriction along the open subscheme $\iota_{n,\alpha}: \mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n) \hookrightarrow \mathrm{Quot}_X(F, n)$ agrees, as a symmetric perfect obstruction theory, with the critical obstruction theory $\mathbb{E}_{\mathrm{crit}}$ of Proposition 3.2.

One can also ask whether $\iota_{n,\alpha}^* \mathbb{E}$ and $\mathbb{E}_{\mathrm{crit}}$ are \mathbb{T} -equivariantly isomorphic over the cotangent complex of $\mathrm{Quot}_{U_\alpha}(F|_{U_\alpha}, n)$. This is of course stronger than the statement of Proposition 9.3.

A similar conjecture (essentially the rank 1 specialisation of Conjecture 9.8) can be stated for the moduli space $\mathrm{Hilb}^n(\mathbb{A}^3) = \mathrm{Quot}_{\mathbb{A}^3}(\mathcal{O}, n)$, without reference to a compactification $\mathbb{A}^3 \subset X$. The Hilbert scheme of points has two symmetric perfect obstruction theories: the critical obstruction theory $\mathbb{E}_{\mathrm{crit}}$ (Proposition 3.2) and the one coming from moduli of ideal sheaves: if $p: \mathbb{A}^3 \times \mathrm{Hilb}^n(\mathbb{A}^3) \rightarrow \mathrm{Hilb}^n(\mathbb{A}^3)$ is the projection and \mathcal{I} is the universal ideal sheaf, one has the obstruction theory

$$\mathrm{Rp}_* \mathbf{R}\mathcal{H}om(\mathcal{I}, \mathcal{I})_0[2] \rightarrow \mathbb{L}_{\mathrm{Hilb}^n(\mathbb{A}^3)}$$

obtained from the Atiyah class $\mathrm{At}_{\mathcal{I}}$ the way we sketched in (9.4).

Conjecture 9.9. There is an isomorphism of perfect obstruction theories

$$\begin{array}{ccc} \mathbb{E}_{\mathrm{crit}} & \xrightarrow{\sim} & \mathrm{Rp}_* \mathbf{R}\mathcal{H}om(\mathcal{I}, \mathcal{I})_0[2] \\ & \searrow & \swarrow \\ & \mathbb{L}_{\mathrm{Hilb}^n(\mathbb{A}^3)} & \end{array}$$

on the Hilbert scheme of points $\mathrm{Hilb}^n(\mathbb{A}^3)$.

REFERENCES

1. Noah Arbesfeld, *K-theoretic Donaldson–Thomas theory and the Hilbert scheme of points on a surface*, [ArXiv:1905.04567](#), 2019.
2. Noah Arbesfeld and Yakov Kononov, *Boxcounting and Quot schemes*, To appear.
3. Hidetoshi Awata and Hiroaki Kanno, *Quiver matrix model and topological partition function in six dimensions*, J. High Energy Phys. **2009** (2009), no. 7, 076, 23.
4. Sjoerd Beentjes and Andrea T. Ricolfi, *Virtual counts on Quot schemes and the higher rank local DT/PT correspondence*, To appear in Math. Res. Lett., 2018.
5. Kai Behrend, *Donaldson–Thomas type invariants via microlocal geometry*, Ann. of Math. **2** (2009), no. 170, 1307–1338.
6. Kai Behrend, Jim Bryan, and Balázs Szendrői, *Motivic degree zero Donaldson–Thomas invariants*, Invent. Math. **192** (2013), no. 1, 111–160.
7. Kai Behrend and Barbara Fantechi, *The intrinsic normal cone*, Invent. Math. **128** (1997), no. 1, 45–88.
8. Kai Behrend and Barbara Fantechi, *Symmetric obstruction theories and Hilbert schemes of points on threefolds*, Algebra Number Theory **2** (2008), 313–345.
9. Francesco Benini, Giulio Bonelli, Matteo Poggi, and Alessandro Tanzini, *Elliptic non-Abelian Donaldson–Thomas invariants of \mathbb{C}^3* , J. High Energy Phys. **2019** (2019), no. 7, 068, 41.
10. Emili Bifet, *Sur les points fixes du schéma $\text{Quot}_{\mathcal{O}_X^r/X/k}$ sous l’action du tore $\mathbf{G}_{m,k}^r$* , C. R. Acad. Sci. Paris Sér. I Math. **309** (1989), no. 9, 609–612.
11. Giulio Bonelli, Nadir Fasola, and Alessandro Tanzini, *Defects, nested instantons and comet shaped quivers*, [ArXiv:1907.02771](#), 2019.
12. Giulio Bonelli, Nadir Fasola, and Alessandro Tanzini, *Flags of sheaves, quivers and symmetric polynomials*, [ArXiv:1911.12787](#), 2019.
13. Yalong Cao, Martijn Kool, and Sergej Monavari, *K -theoretic DT/PT correspondence for toric Calabi–Yau 4-folds*, [ArXiv:1906.07856](#), 2019.
14. Alberto Cazzaniga, *On some computations of refined Donaldson–Thomas invariants*, PhD Thesis, University of Oxford, 2015.
15. Alberto Cazzaniga, Dimbinaina Ralaivaosaona, and Andrea T. Ricolfi, *Higher rank motivic Donaldson–Thomas invariants of \mathbb{A}^3 via wall-crossing, and asymptotics*, [ArXiv:2004.07020](#), 2020.
16. Alberto Cazzaniga and Andrea T. Ricolfi, *Framed motivic Donaldson–Thomas invariants of small crepant resolutions*, [ArXiv:2004.07837](#), 2020.
17. Alberto Cazzaniga and Andrea T. Ricolfi, *Framed sheaves on projective space and Quot schemes*, [ArXiv:2004.13633](#), 2020.
18. Ionuț Ciocan-Fontanine and Mikhail Kapranov, *Virtual fundamental classes via dg-manifolds*, Geom. Topol. **13** (2009), no. 3, 1779–1804.
19. Michele Cirafici, Annamaria Sinkovics, and Richard J. Szabo, *Cohomological gauge theory, quiver matrix models and Donaldson–Thomas theory*, Nuclear Phys. B **809** (2009), no. 3, 452–518.
20. Ben Davison and Andrea T. Ricolfi, *The local motivic DT/PT correspondence*, [ArXiv:1905.12458](#), 2019.
21. Jan Denef and François Loeser, *Geometry on arc spaces of algebraic varieties*, 3rd European congress of mathematics (ECM), Barcelona, Spain, July 10–14, 2000. Volume I, Basel: Birkhäuser, 2001, pp. 327–348.
22. Tudor Dimofte and Sergei Gukov, *Refined, motivic, and quantum*, Lett. Math. Phys. **91** (2010), no. 1, 1–27.
23. Barbara Fantechi and Lothar Göttsche, *Riemann–Roch theorems and elliptic genus for virtually smooth schemes*, Geom. Topol. **14** (2010), no. 1, 83–115.
24. Sebastian Franco, Dongwook Ghim, Sangmin Lee, and Rak-Kyeong Seong, *Elliptic genera of 2d (0,2) gauge theories from brane brick models*, J. High Energy Phys. (2017), no. 6, 068, front matter+46.
25. Amin Gholampour and Martijn Kool, *Higher rank sheaves on threefolds and functional equations*, [Épiga](#), 2019.
26. Amin Gholampour, Martijn Kool, and Benjamin Young, *Rank 2 Sheaves on Toric 3-Folds: Classical and Virtual Counts*, Int. Math. Res. Not. **2018** (2018), no. 10, 2981–3069.
27. Tom Graber and Rahul Pandharipande, *Localization of virtual classes*, Invent. Math. **135** (1999), no. 2, 487–518.
28. Martin G. Gulbrandsen and Andrea T. Ricolfi, *The Euler characteristic of the generalized Kummer scheme of an abelian threefold*, Geom. Dedicata **182** (2016), 73–79.
29. Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977, Graduate Texts in Mathematics, No. 52.

30. Luc Illusie, *Complexe cotangent et déformations. I*, Lecture Notes in Mathematics, vol. 239, Springer-Verlag, Berlin-New York, 1971.
31. Amer Iqbal, Can Kozçaz, and Cumrun Vafa, *The refined topological vertex*, J. High Energy Phys. (2009), no. 10, 069.
32. Drew Johnson, Dragos Oprea, and Rahul Pandharipande, *Rationality of descendent series for Hilbert and Quot schemes of surfaces*, [ArXiv:2002.05861](#), 2020.
33. Toshiya Kawai and Kenji Mohri, *Geometry of $(0, 2)$ Landau-Ginzburg orbifolds*, Nuclear Phys. B **425** (1994), no. 1-2, 191–216.
34. Maxim Kontsevich, *Enumeration of rational curves via torus actions*, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. 129, Birkhäuser Boston, Boston, MA, 1995, pp. 335–368.
35. Maxim Kontsevich and Yan Soibelman, *Stability structures, motivic Donaldson–Thomas invariants and cluster transformations*, [ArXiv:0811.2435](#), 2008.
36. Maxim Kontsevich and Yan Soibelman, *Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson–Thomas invariants*, Commun. Number Theory Phys. **5** (2011), no. 2, 231–352.
37. Martijn Kool, *Fixed point loci of moduli spaces of sheaves on toric varieties*, Adv. Math. **227** (2011), no. 4, 1700–1755.
38. Marc Levine and Rahul Pandharipande, *Algebraic cobordism revisited*, Invent. Math. **176** (2009), no. 1, 63–130.
39. Jun Li, *Zero dimensional Donaldson–Thomas invariants of threefolds*, Geom. Topol. **10** (2006), 2117–2171.
40. Jun Li and Gang Tian, *Virtual moduli cycles and Gromov–Witten invariants of algebraic varieties*, J. Amer. Math. Soc. **11** (1998), no. 1, 119–174.
41. Woonam Lim, *Virtual χ_{-y} -genera of Quot schemes on surfaces*, [ArXiv:2003.04429](#), 03 2020.
42. Alina Marian and Dragos Oprea, *Virtual intersections on the Quot scheme and Vafa–Intriligator formulas*, Duke Math. J. **136** (2007), no. 1, 81–113.
43. Alina Marian, Dragos Oprea, and Rahul Pandharipande, *Segre classes and Hilbert schemes of points*, Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), no. 1, 239–267.
44. Daves Maulik, Nikita Nekrasov, Andrei Okounkov, and Rahul Pandharipande, *Gromov–Witten theory and Donaldson–Thomas theory, I*, Compos. Math. **142** (2006), no. 5, 1263–1285.
45. Daves Maulik, Nikita Nekrasov, Andrei Okounkov, and Rahul Pandharipande, *Gromov–Witten theory and Donaldson–Thomas theory, II*, Compos. Math. **142** (2006), no. 5, 1286–1304.
46. Nikita Nekrasov, *Z-theory: chasing m/f theory*, Comptes Rendus Physique **6** (2005), no. 2, 261–269, Strings, gravity, and the quest for unification - Strings 04, part II.
47. Nikita Nekrasov, *Instanton partition functions and M-theory*, Jpn. J. Math. **4** (2009), no. 1, 63–93.
48. Nikita Nekrasov, *Magnificent four*, Ann. Inst. Henri Poincaré D **7** (2020), no. 4, 505–534.
49. Nikita Nekrasov and Andrei Okounkov, *Membranes and sheaves*, Algebr. Geom. **3** (2016), no. 3, 320–369.
50. Nikita Nekrasov and Nicolò Piazzalunga, *Magnificent four with colors*, Comm. Math. Phys. **372** (2019), no. 2, 573–597.
51. Madhav Nori, *Appendix to the paper by C. S. Seshadri: Desingularisation of the moduli varieties of vector bundles over curves*, Proceedings of the International Symposium on Algebraic Geometry, 1977.
52. Andrei Okounkov, *Lectures on K-theoretic computations in enumerative geometry*, Geometry of moduli spaces and representation theory, IAS/Park City Math. Ser., vol. 24, Amer. Math. Soc., Providence, RI, 2017, pp. 251–380.
53. Dragos Oprea and Rahul Pandharipande, *Quot schemes of curves and surfaces: virtual classes, integrals, Euler characteristics*, [ArXiv:1903.08787](#), 2019.
54. Feng Qu, *Virtual pullbacks in K-theory*, Ann. Inst. Fourier (Grenoble) **68** (2018), no. 4, 1609–1641.
55. Andrea T. Ricolfi, *Local Donaldson–Thomas invariants and their refinements*, Ph.D. thesis, University of Stavanger, 2017.
56. Andrea T. Ricolfi, *The DT/PT correspondence for smooth curves*, Math. Z. **290** (2018), no. 1-2, 699–710.
57. Andrea T. Ricolfi, *Local contributions to Donaldson–Thomas invariants*, Int. Math. Res. Not. IMRN **2018** (2018), no. 19, 5995–6025.
58. Andrea T. Ricolfi, *On the motive of the Quot scheme of finite quotients of a locally free sheaf*, J. Math. Pures Appl. **144** (2020), 50–68.
59. Andrea T. Ricolfi, *Virtual classes and virtual motives of Quot schemes on threefolds*, Adv. Math. **369** (2020), 107182.

60. Andrea T. Ricolfi, *The equivariant Atiyah class*, To appear in C. R. Math. Acad. Sci. Paris, 2021.
61. Bernd Siebert, *Virtual fundamental classes, global normal cones and Fulton's canonical classes*, Frobenius manifolds, Aspects Math., vol. 36, Friedr. Vieweg, Wiesbaden, 2004, pp. 341–358.
62. Richard J. Szabo, $\mathcal{N} = 2$ gauge theories, instanton moduli spaces and geometric representation theory, J. Geom. Phys. **109** (2016), 83–121.
63. Balázs Szendrői, *Cohomological Donaldson–Thomas theory*, String-Math 2014, Proc. Sympos. Pure Math., vol. 93, Amer. Math. Soc., Providence, RI, 2016, pp. 363–396.
64. Richard P. Thomas, *A holomorphic Casson invariant for Calabi–Yau 3-folds, and bundles on K3 fibrations*, J. Diff. Geo. **54** (2000), no. 2, 367–438.
65. Richard P. Thomas, *A K-theoretic Fulton class*, Facets of Algebraic Geometry: A Volume in Honour of William Fulton's 80th Birthday, Cambridge University Press, 2020.
66. Richard P. Thomas, *Equivariant K-Theory and Refined Vafa–Witten Invariants*, Comm. Math. Phys. **378** (2020), no. 2, 1451–1500.
67. Yukinobu Toda, *Hall algebras in the derived category and higher-rank DT invariants*, Algebr. Geom. **7** (2020), no. 3, 240–262.
68. Edward Witten, *The index of the Dirac operator in loop space*, Elliptic curves and modular forms in algebraic topology (Princeton, NJ, 1986), Lecture Notes in Math., vol. 1326, Springer, Berlin, 1988, pp. 161–181.

SISSA TRIESTE, VIA BONOMEA 265, 34136 TRIESTE, ITALY

INSTITUTE FOR GEOMETRY AND PHYSICS, VIA BEIRUT 4, 34100 TRIESTE, ITALY

ISTITUTO NAZIONALE DI FISICA NUCLEARE, SEZIONE DI TRIESTE, VIA VALERIO 2, 34127 TRIESTE, ITALY

Email address: nfasola@sissa.it

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY, P.O. BOX 80010 3508 TA UTRECHT, THE NETHERLANDS

Email address: s.monavari@uu.nl

SISSA TRIESTE, VIA BONOMEA 265, 34136 TRIESTE, ITALY

INSTITUTE FOR GEOMETRY AND PHYSICS, VIA BEIRUT 4, 34100 TRIESTE, ITALY

Email address: aricolfi@sissa.it