

TWO-PHASE STOKES FLOW BY CAPILLARITY IN FULL 2D SPACE: AN APPROACH VIA HYDRODYNAMIC POTENTIALS

BOGDAN-VASILE MATIOC AND GEORG PROKERT

ABSTRACT. We study the two-phase Stokes flow driven by surface tension with two fluids of equal viscosity, separated by an asymptotically flat interface with graph geometry. The flow is assumed to be two-dimensional with the fluids filling the entire space. We prove well-posedness and parabolic smoothing in Sobolev spaces up to critical regularity. The main technical tools are an analysis of nonlinear singular integral operators arising from the hydrodynamic single-layer potential and abstract results on nonlinear parabolic evolution equations.

1. INTRODUCTION

One of the standard methods in the analysis of moving boundary problems is the reformulation of these problems as evolution equations in function spaces to which methods of Functional Analysis can be applied, depending on the character of the problem under investigation. The difficulty of this typically consists in the fact that the resulting evolution equations are nonlocal and strongly nonlinear. For moving boundary problems with domains of general shape this approach typically involves the transformation of the moving domain to a fixed reference domain by an unknown, time dependent diffeomorphism, and the (explicit or implicit) use of solution operators for boundary value problems with variable coefficients on this reference domain. This approach often implies restrictions to results of perturbation type, i.e. either short-time solutions, or solutions for (in some sense) small data. However, this can be avoided in special situations where

- the geometry is simpler (e.g. full space, with the moving boundary being a graph), and
- the underlying PDE is elliptic and has constant coefficients.

In such situations, one can use the classical methods of potential theory to solve the PDEs directly, i.e. without transformations of the domain, and reformulate the moving boundary problem as an evolution equation that involves nonlinear, singular integral operators.

This strategy proved to be successful for various versions of the Muskat (or two-phase Hele-Shaw) problem, see e.g. the survey articles [6, 7]. While the constant coefficient elliptic operator underlying the Muskat problem is simply the Laplacian, the related moving boundary problems of quasistationary Stokes flow are based on the Stokes operator (which is also elliptic in a sense that can be made precise). Concretely, in this paper we are interested in the following moving boundary problem of Stokes flow driven by the capillarity of the

2010 *Mathematics Subject Classification.* 35R37; 76D07; 35K55.

Key words and phrases. Stokes problem; Two-phase; Singular integrals; Contour integral formulation.

moving interface $t \mapsto \Gamma(t)$ between two fluid phases $\Omega^\pm(t)$ in \mathbb{R}^2 :

$$\left. \begin{aligned} \mu \Delta v^\pm - \nabla q^\pm &= 0 && \text{in } \Omega^\pm(t), \\ \operatorname{div} v^\pm &= 0 && \text{in } \Omega^\pm(t), \\ v^+ &= v^- && \text{on } \Gamma(t), \\ [T(v, q)] \tilde{\nu} &= -\sigma \tilde{\kappa} \tilde{\nu} && \text{on } \Gamma(t), \\ (v^\pm, q^\pm) &\rightarrow 0 && \text{for } |x| \rightarrow \infty, \\ V_n &= v \cdot \tilde{\nu} && \text{on } \Gamma(t). \end{aligned} \right\} \quad (1.1)$$

Here, $v^\pm : \Omega^\pm(t) \rightarrow \mathbb{R}^2$ is a vector field representing the velocity of the liquid located in $\Omega^\pm(t)$ and $q^\pm : \Omega^\pm(t) \rightarrow \mathbb{R}$ its pressure. Furthermore, $\tilde{\nu}$ is the unit exterior normal to $\partial\Omega^-(t)$ and $\tilde{\kappa}$ denotes the curvature of the interface. Moreover, $[T(v, q)]$ denotes the jump of the stress tensor across $\Gamma(t)$, see (2.4), (2.6) below. The positive constants μ and σ denote the viscosity of the liquids and the surface tension coefficient of the interface, respectively. We assume that

$$\Gamma(t) = \partial\Omega^\pm(t), \quad \Omega^+(t) \cup \Omega^-(t) \cup \Gamma(t) = \mathbb{R}^2,$$

and that $\Gamma(t)$ is a graph over a suitable straight line. Equation (1.1)₆ determines the motion of the interface by prescribing its normal velocity as coinciding with the normal component of the velocity at $\Gamma(t)$, i.e. the interface is transported by the liquid flow. The interface $\Gamma(t)$ is assumed to be known at time $t = 0$.

For the Stokes operator, it is possible to set up a treatment of boundary value problems based on so-called hydrodynamic potentials [8] in strict analogy to the potentials for the Laplacian. It is this analogy that enables us to study the moving boundary problem of two-phase Stokes flow driven by capillarity (at least in 2D and with equal viscosity in both phases) along the same lines as for the Muskat problem. This has first been exploited in [4] to obtain an existence result for all positive times, with initial data that are small in a space of Fourier transforms of bounded measures. To the best of our knowledge, this is the only result available on two-phase Stokes flow in the unbounded geometry considered here.

It is the aim of the present paper to analyze Problem (1.1) in Sobolev spaces (up to critical regularity) in an L_2 -based setting. The use of these spaces also implies that the interface is asymptotically flat.

We will establish

- existence and uniqueness of maximal solutions with initial data that are arbitrary within our phase space;
- a corresponding semiflow property;
- parabolic smoothing up to C^∞ of solutions in time and space (away from the initial time);
- a criterion for global existence of solutions, or equivalently, a necessary condition for blow-up.

Essentially, these results are obtained by applying the theory of maximal regularity for nonlinear parabolic equations in weighted Hölder spaces of vector-valued functions presented in [9]. Since the Stokes flow (1.1) is driven by capillarity, it turns out that the problem is parabolic “everywhere”, i.e. the parabolicity condition is just positivity of the surface energy. We emphasize that while for the discussion of the boundary value problem (1.1)_{1–5} at a fixed time t we need to assume H^3 -smoothness of the interface, see Section 2, the corresponding nonlinear evolution equation (3.4) is shown to be well-posed in all subcritical spaces H^s

with $s > 3/2$. Hence, we obtain a “weak” solution concept allowing for less regular initial data. Nevertheless, for positive times all solutions are classical due to parabolic smoothing.

The structure of the paper is as follows: In Section 2 we consider the underlying two-phase boundary value problem for the Stokes equations (1.1)_{1–5} with fixed interface, and show that it is solved by the so-called hydrodynamic single-layer potential. We prove this by investigating its behavior near and on the interface (recovering results from [8] in our slightly different setting) and show that it vanishes in the far-field limit. This asymptotic result can be interpreted as nonoccurrence of the 2D Stokes paradoxon in our setting, which is essentially due to the fact that the curvature vector of the interface is a derivative (by arclength) of a vector that approaches a constant at infinity (see Eqn. (2.1)). In Section 3 we first rewrite our moving boundary problem as an evolution equation for the function that parametrizes the interface between the fluids and announce our main result. The remainder of the section is dedicated to its proof. We linearize the evolution equation, and then establish its parabolic character (see Theorem 3.7). The localization procedure by which this is accomplished demands the main technical effort. Once parabolicity is established, the results follow from general facts on (fully) nonlinear problems of this type as given in [9].

Throughout the paper, some longer proofs are deferred to appendices.

2. THE FIXED TIME PROBLEM

In this section we study the two-phase boundary value problem for the Stokes equations (1.1)_{1–5} with fixed domains Ω^\pm and boundary $\partial\Omega^\pm := \Gamma$ as defined by

$$\Omega^\pm := \Omega_f^\pm := \{(x_1, x_2) \in \mathbb{R}^2 \mid x_2 \gtrless f(x_1)\}, \quad \Gamma := \Gamma_f := \partial\Omega^\pm = \{(x_1, f(x_1)) \mid x_1 \in \mathbb{R}\}.$$

The function $f \in H^3(\mathbb{R})$ is fixed. Note that Γ is the image of the x_1 -axis under the diffeomorphism $\Xi := \Xi_f := (\text{id}_{\mathbb{R}}, f)$. Further, let ν be the componentwise pull-back under Ξ of the unit normal on Γ exterior to Ω^- , i.e.

$$\nu := \frac{1}{\omega}(-f', 1)^\top, \quad \omega := \omega_f := (1 + f'^2)^{1/2}.$$

Let $\kappa := \omega^{-3}f'' \in H^1(\mathbb{R})$ be the pull-back under Ξ of the curvature of Γ . In view of

$$\kappa\nu = \omega^{-1}(\omega^{-1}(1, f')^\top)' \tag{2.1}$$

we will use the relation

$$\kappa\nu = \omega^{-1}g', \quad g := g_f := (g_1, g_2)^\top := (\omega^{-1} - 1, \omega^{-1}f')^\top =: (-\phi_1(f), \phi_2(f))^\top, \tag{2.2}$$

where

$$\phi_1(f) = \frac{f'^2}{\omega + \omega^2} \quad \text{and} \quad \phi_2(f) = \frac{f'}{\omega} \tag{2.3}$$

belong to $H^2(\mathbb{R})$.

For any functions z^\pm defined on Ω^\pm , respectively, and having limits at some $\xi \in \Gamma$ we will write

$$[z](\xi) := \lim_{\Omega^+ \ni x \rightarrow \xi} z^+(x) - \lim_{\Omega^- \ni x \rightarrow \xi} z^-(x). \tag{2.4}$$

We fix a common viscosity $\mu > 0$ as well as a surface tension coefficient $\sigma > 0$ and seek solutions

$$(v^\pm, q^\pm) \in (C^2(\Omega^\pm, \mathbb{R}^2) \cap C^1(\overline{\Omega^\pm}, \mathbb{R}^2)) \times (C^1(\Omega^\pm) \cap C(\overline{\Omega^\pm}))$$

to the two-phase boundary value problem

$$\left. \begin{aligned} \mu \Delta v^\pm - \nabla q^\pm &= 0 && \text{in } \Omega^\pm, \\ \operatorname{div} v^\pm &= 0 && \text{in } \Omega^\pm, \\ v^+ &= v^- && \text{on } \Gamma, \\ [T(v, q)](\nu \circ \Xi^{-1}) &= -\sigma(\kappa\nu) \circ \Xi^{-1} && \text{on } \Gamma, \\ (v^\pm, q^\pm) &\rightarrow 0 && \text{for } |x| \rightarrow \infty, \end{aligned} \right\} \quad (2.5)$$

with Ω^\pm and Γ as defined above. Here $T(v, q) = (T_{ij}(v, q))_{1 \leq i, j \leq 2}$ denotes the stress tensor that is given by

$$T_{ij}(v, q) := -q\delta_{ij} + \mu(\partial_i v_j + \partial_j v_i). \quad (2.6)$$

The structure of the problem allows us to represent the solution as a hydrodynamic single-layer potential [8]. For this, we introduce for $x \in \mathbb{R}^2$ and $k = 1, 2$ the fundamental solutions

$$(\mathcal{U}_x^k, \mathcal{P}_x^k) : \mathbb{R}^2 \setminus \{x\} \longrightarrow \mathbb{R}^2 \times \mathbb{R}$$

to the Stokes equations in \mathbb{R}^2 by the formulas $\mathcal{U}_x^k = (\mathcal{U}_{x,1}^k, \mathcal{U}_{x,2}^k)^\top$ and

$$\begin{aligned} \mathcal{U}_{x,j}^k(y) &= -\frac{1}{4\pi\mu} \left(\delta_{jk} \ln \frac{1}{|x-y|} + \frac{(x_j - y_j)(x_k - y_k)}{|x-y|^2} \right), \quad j = 1, 2, \\ \mathcal{P}_x^k(y) &= \frac{1}{2\pi} \frac{x_k - y_k}{|x-y|^2}. \end{aligned}$$

These functions solve (in distributional sense) the Stokes equations

$$\begin{aligned} \mu \Delta \mathcal{U}_x^k - \nabla \mathcal{P}_x^k &= \delta_x e^k \quad \text{in } \mathcal{D}'(\mathbb{R}^2), \\ \operatorname{div} \mathcal{U}_x^k &= 0 \quad \text{in } \mathcal{D}'(\mathbb{R}^2), \end{aligned}$$

with $e^1 = (1, 0)$ and $e^2 = (0, 1)$. Moreover, fixing $x = 0$, setting $y = (\xi, \eta)$ and $r := \sqrt{\xi^2 + \eta^2}$, and differentiating the solutions with respect to ξ and η we get the following solutions to the homogeneous Stokes system in $\mathbb{R}^2 \setminus \{0\}$, with corresponding pressures:

$$\begin{aligned} 4\pi \partial_\xi \mathcal{U}_0^1(\xi, \eta) &= \frac{1}{\mu r^4} \begin{pmatrix} \xi(\xi^2 - \eta^2) \\ \eta(\xi^2 - \eta^2) \end{pmatrix}, & 2\pi \partial_\xi \mathcal{P}_0^1(\xi, \eta) &= \frac{\xi^2 - \eta^2}{r^4}, \\ 4\pi \partial_\eta \mathcal{U}_0^1(\xi, \eta) &= \frac{1}{\mu r^4} \begin{pmatrix} \eta(\eta^2 + 3\xi^2) \\ \xi(\eta^2 - \xi^2) \end{pmatrix}, & 2\pi \partial_\eta \mathcal{P}_0^1(\xi, \eta) &= \frac{2\xi\eta}{r^4}, \\ 4\pi \partial_\xi \mathcal{U}_0^2(\xi, \eta) &= \frac{1}{\mu r^4} \begin{pmatrix} \eta(\xi^2 - \eta^2) \\ \xi(\xi^2 + 3\eta^2) \end{pmatrix}, & 2\pi \partial_\xi \mathcal{P}_0^2(\xi, \eta) &= \frac{2\xi\eta}{r^4}, \\ 2\pi \partial_\eta \mathcal{U}_0^2(\xi, \eta) &= \frac{1}{\mu r^4} \begin{pmatrix} \xi(\eta^2 - \xi^2) \\ \eta(\eta^2 - \xi^2) \end{pmatrix}, & 2\pi \partial_\eta \mathcal{P}_0^2(\xi, \eta) &= \frac{\eta^2 - \xi^2}{r^4}. \end{aligned} \quad (2.7)$$

We are going to prove that the functions $(v^\pm, q^\pm) := (v, q)|_{\Omega^\pm}$ with

$$\begin{aligned} v(x, y) &:= \sigma \int_{\mathbb{R}} \partial_s \left(\mathcal{U}_0^k((x, y) - (s, f(s))) \right) g_k(s) ds, \\ q(x, y) &:= -\sigma \int_{\mathbb{R}} \mathcal{P}_0^k((x, y) - (s, f(s))) g'_k(s) ds \end{aligned} \quad (2.8)$$

and g_1, g_2 as defined in (2.2) constitute the unique solution to (2.5). These functions are defined first for $(x, y) \in \mathbb{R}^2 \setminus \Gamma$, their behavior near Γ will be investigated later.

Observe first that the kernels of the integral operators in (2.8) are smooth with respect to $(x, y) \in \mathbb{R}^2 \setminus \Gamma$. Moreover, $g'_k \in H^1(\mathbb{R})$ and

$$\mathcal{P}_0^k((x, y) - (s, f(s))) = O(s^{-1}) \quad \text{for } |s| \rightarrow \infty,$$

so that the integrand in (2.8)₂ belongs to $L_1(\mathbb{R})$. Furthermore, $g_k \rightarrow 0$ for $|s| \rightarrow \infty$, so we can use integration by parts to obtain

$$q(x, y) = \sigma \int_{\mathbb{R}} \partial_s (\mathcal{P}_0^k((x, y) - (s, f(s)))) g_k(s) ds.$$

Besides, recalling (2.7), we get

$$\partial_s (\mathcal{U}_0^k((x, y) - (s, f(s)))) = O(s^{-1}) \quad \text{for } |s| \rightarrow \infty,$$

and since $g_k \in H^2(\mathbb{R})$, it follows that also v is well-defined. Altogether, we obtain the following representation for the velocity field and the pressure:

$$(v, q)^\top(x, y) = - \int_{\mathbb{R}} \partial_s M(\xi, \eta) g(s) ds = \int_{\mathbb{R}} (\partial_\xi M(\xi, \eta) + f'(s) \partial_\eta M(\xi, \eta)) g(s) ds \quad (2.9)$$

for $(x, y) \in \mathbb{R}^2 \setminus \Gamma$, where

$$\xi := x - s, \quad \eta := y - f(s), \quad r := \sqrt{\xi^2 + \eta^2}.$$

$$M(\xi, \eta) := -\sigma \begin{pmatrix} \mathcal{U}_0^1 & \mathcal{U}_0^2 \\ \mathcal{P}_0^1 & \mathcal{P}_0^2 \end{pmatrix} (\xi, \eta).$$

Theorem 2.1. *Given $f \in H^3(\mathbb{R})$, Problem (2.5) has the unique solution (v^\pm, q^\pm) given by (2.8) or equivalently (2.9).*

Proof. 1. (v^\pm, q^\pm) solves the Stokes equations:

Denote the integrand in (2.9) by

$$I : (\mathbb{R}^2 \setminus \Gamma) \times \mathbb{R} \longrightarrow \mathbb{R}^2 \times \mathbb{R}.$$

Any partial derivative $\partial_x^\alpha \partial_y^\beta I$ can be dominated by an absolutely integrable function, locally uniformly in (x, y) , so that differentiation with respect to x and y and integration with respect to s can be interchanged. In particular, it follows that

$$(v^\pm, q^\pm) \in C^\infty(\Omega^\pm, \mathbb{R}^2) \times C^\infty(\Omega^\pm).$$

As the columns of $\partial_\xi M, \partial_\eta M$ represent (as functions of (x, y)) solutions to the homogeneous Stokes equations (cf. (2.7)), (v^\pm, q^\pm) is also a solution to these equations on $\mathbb{R}^2 \setminus \Gamma$.

2. Uniqueness:

We have to show that any solution

$$(u^\pm, p^\pm) \in (C^2(\Omega^\pm, \mathbb{R}^2) \cap C^1(\overline{\Omega^\pm}, \mathbb{R}^2)) \times (C^1(\Omega^\pm) \cap C(\overline{\Omega^\pm}))$$

to

$$\left. \begin{aligned} \mu \Delta u^\pm - \nabla p^\pm &= 0 && \text{in } \Omega^\pm, \\ \operatorname{div} u^\pm &= 0 && \text{in } \Omega^\pm, \\ u^+ &= u^- && \text{on } \Gamma, \\ [T(u, p)] \tilde{\nu} &= 0 && \text{on } \Gamma, \\ (u, p) &\rightarrow 0 && \text{for } |x| \rightarrow \infty, \end{aligned} \right\}$$

where $\tilde{\nu} := \nu \circ \Xi^{-1}$, is identically zero. Let $\tilde{\tau} := (\omega^{-1}(1, f')^\top) \circ \Xi^{-1}$ be the unit tangential vector field along Γ , oriented to the right. Observe first that

$$T(u^\pm, p^\pm) \tilde{\nu} = \mu(\partial_{\tilde{\nu}} u^\pm + \partial_{\tilde{\tau}}(u_2, -u_1)^\top + \tilde{\nu} \operatorname{div} u^\pm) - p^\pm \tilde{\nu},$$

so under our assumptions

$$[T(u, p)] \tilde{\nu} = \mu[\partial_{\tilde{\nu}} u] - [p] \tilde{\nu} = 0. \quad (2.10)$$

We now define

$$(V, Q) := \mathbf{1}_{\Omega^+}(u^+, p^+) + \mathbf{1}_{\Omega^-}(u^-, p^-) \in L_{1,\text{loc}}(\mathbb{R}^2, \mathbb{R}^2 \times \mathbb{R}).$$

Taking distributional derivatives and using the continuity of u across Γ yields

$$\begin{aligned} \Delta V_i &= \mathbf{1}_{\Omega^+} \Delta u_i^+ + \mathbf{1}_{\Omega^-} \Delta u_i^- + [\partial_{\tilde{\nu}} u_i] \delta_\Gamma, \\ \partial_i Q &= \mathbf{1}_{\Omega^+} \partial_i p^+ + \mathbf{1}_{\Omega^-} \partial_i p^- + [p] \tilde{\nu}_i \delta_\Gamma, \\ \operatorname{div} V &= 0, \end{aligned}$$

where, given $a \in L_{1,\text{loc}}(\Gamma)$, the distribution $a\delta_\Gamma$ is defined by

$$\langle a\delta_\Gamma, \phi \rangle := \int_\Gamma a\phi \, d\Gamma, \quad \phi \in C_0^\infty(\mathbb{R}^2).$$

So, from this and (2.10) we get

$$\mu \Delta V - \nabla Q = 0 \quad \text{in } \mathcal{D}'(\mathbb{R}^2).$$

In particular, taking the divergence of this equation yields $\Delta Q = 0$, i.e. Q is a harmonic function on the full space \mathbb{R}^2 , and the asymptotic condition implies $Q = 0$ via Liouville's theorem. This implies in turn that V_1 and V_2 are harmonic, and are therefore zero by the same argument.

3. The behavior of (v^\pm, q^\pm) near Γ is addressed in Appendix A. In particular, it is shown that $(v^\pm, q^\pm) \in C^1(\overline{\Omega^\pm}, \mathbb{R}^2) \times C(\overline{\Omega^\pm})$ satisfies Eqns. (2.5)₃₋₄.

4. The far-field boundary condition (2.5)₅ is established in Appendix B. \square

3. THE EVOLUTION PROBLEM

In the first part of this section we introduce some notation which is then used to recast the Stokes problem (1.1) as an evolution problem for f only, see (3.4) below. In the second part we establish our main result stated in Theorem 3.2.

3.1. A class of singular integral operators. We first introduce a class of multilinear singular integral operators which are needed in the second part of this section. Given $n, m \in \mathbb{N}$ and Lipschitz continuous functions $a_1, \dots, a_m, b_1, \dots, b_n : \mathbb{R} \rightarrow \mathbb{R}$, denote by $B_{n,m}$ the singular integral operator

$$B_{n,m}(a_1, \dots, a_m)[b_1, \dots, b_n, h](x) := \text{PV} \int_{\mathbb{R}} \frac{h(x-y)}{y} \frac{\prod_{i=1}^n (\delta_{[x,y]} b_i/y)}{\prod_{i=1}^m [1 + (\delta_{[x,y]} a_i/y)^2]} dy, \quad (3.1)$$

where $\delta_{[x,y]} u := u(x) - u(x-y)$, and for brevity

$$B_{n,m}^0(f)[h] := B_{n,m}(f, \dots, f)[f, \dots, f, h] \quad (3.2)$$

(with the appropriate number of identical arguments f filled in). Here PV denotes the principle value. Below we write $C^{1-}(X, Y)$ for the space of locally Lipschitz maps from X to Y . The following properties are extensively used in our analysis.

Lemma 3.1.

- (i) *Given Lipschitz continuous functions $a_1, \dots, a_m, b_1, \dots, b_n : \mathbb{R} \rightarrow \mathbb{R}$, there exists a constant C depending only on n, m and $\max_{i=1, \dots, m} \|a'_i\|_{\infty}$, such that*

$$\|B_{n,m}(a_1, \dots, a_m)[b_1, \dots, b_n, \cdot]\|_{\mathcal{L}(L_2(\mathbb{R}))} \leq C \prod_{i=1}^n \|b'_i\|_{\infty}.$$

Moreover, $B_{n,m} \in C^{1-}((W_{\infty}^1(\mathbb{R}))^m, \mathcal{L}_{n+1}((W_{\infty}^1(\mathbb{R}))^n \times L_2(\mathbb{R}), L_2(\mathbb{R})))$.

- (ii) *Given $s \in (3/2, 2)$, there exists a constant C , depending only on n, m, s , and $\max_{1 \leq i \leq m} \|a_i\|_{H^s}$, such that*

$$\|B_{n,m}(a_1, \dots, a_m)[b_1, \dots, b_n, h]\|_{H^{s-1}} \leq C \|h\|_{H^{s-1}} \prod_{i=1}^n \|b_i\|_{H^s}$$

for all $a_1, \dots, a_m, b_1, \dots, b_n \in H^s(\mathbb{R})$ and $h \in H^{s-1}(\mathbb{R})$.

Moreover, $B_{n,m} \in C^{1-}((H^s(\mathbb{R}))^m, \mathcal{L}_{n+1}((H^s(\mathbb{R}))^n \times H^{s-1}(\mathbb{R}), H^{s-1}(\mathbb{R})))$.

- (iii) *Let $n, m \in \mathbb{N}$, $n \geq 1$, and $3/2 < s' < s < 2$ be given. There exists a constant C , depending only on n, m, s, s' , and $\max_{1 \leq i \leq m} \|a_i\|_{H^s}$, such that*

$$\begin{aligned} & \|B_{n,m}(a_1, \dots, a_m)[b_1, \dots, b_n, h] - h B_{n-1,m}(a_1, \dots, a_m)[b_2, \dots, b_n, b'_1]\|_{H^{s-1}} \\ & \leq C \|b_1\|_{H^{s'}} \|h\|_{H^{s-1}} \prod_{i=2}^n \|b_i\|_{H^s} \end{aligned}$$

for all $a_1, \dots, a_m, b_1, \dots, b_n \in H^s(\mathbb{R})$ and $h \in H^{s-1}(\mathbb{R})$.

Proof. The claim (i) is established in [10, Lemma 3.1], while the properties (ii) and (iii) are proven in [1, Lemmas 2.5 and 2.6]. \square

3.2. Formulation of the evolution equation and the main result. In view of Theorem 2.1 we may recast the two-phase Stokes moving boundary problem (1.1) as a nonlinear and nonlocal evolution problem of the form

$$\frac{df}{dt}(t) = -f'(t)v_1|_{\Gamma \circ \Xi} + v_2|_{\Gamma \circ \Xi} \quad (3.3)$$

with $v = (v_1, v_2)$ given by (2.8). Recalling (A.11) and (2.2), we conclude that

$$\frac{df}{dt}(t) = \Psi(f(t)), \quad t \geq 0, \quad f(0) = f_0, \quad (3.4)$$

where we have set

$$\Psi(f) := -\frac{\sigma}{4\pi\mu}f'\Psi_1(f) + \frac{\sigma}{4\pi\mu}\Psi_2(f), \quad (3.5)$$

with

$$\Psi_1(f) := (B_{0,2}^0(f) - B_{2,2}^0(f))[\phi_1 + f'\phi_2] + B_{1,2}^0(f)[3f'\phi_1 - \phi_2] + B_{3,2}^0(f)[f'\phi_1 + \phi_2],$$

$$\Psi_2(f) := -B_{0,2}^0(f)[f'\phi_1 + \phi_2] + (B_{1,2}^0(f) - B_{3,2}^0(f))[\phi_1 + f'\phi_2] + B_{2,2}^0(f)[f'\phi_1 - 3\phi_2].$$

We recall from (2.2) the shorthand notations

$$\phi_1(f) = \frac{f'^2}{\sqrt{1+f'^2} + 1 + f'^2} \quad \text{and} \quad \phi_2(f) = \frac{f'}{\sqrt{1+f'^2}}.$$

The following theorem contains the main results of this paper.

Theorem 3.2. *Let $s \in (3/2, 2)$ be given. Then, the following statements hold true:*

(i) (Well-posedness) *Given $f_0 \in H^s(\mathbb{R})$, there exists a unique maximal solution*

$$f = f(\cdot; f_0) \in C([0, T_+), H^s(\mathbb{R})) \cap C^1([0, T_+), H^{s-1}(\mathbb{R})),$$

where $T_+ = T_+(f_0) \in (0, \infty]$, to (3.4). Moreover, $[(t, f_0) f(t; f_0)]$ defines a semiflow on $H^s(\mathbb{R})$.

(ii) (Parabolic smoothing)

(iia) *The map $[(t, x) \mapsto f(t, x)] : (0, T_+) \times \mathbb{R} \rightarrow \mathbb{R}$ is a C^∞ -function.*

(iib) *For any $k \in \mathbb{N}$, we have $f \in C^\infty((0, T_+), H^k(\mathbb{R}))$.*

(iii) (Global existence) *If*

$$\sup_{[0, T] \cap [0, T_+(f_0))} \|f(t)\|_{H^s} < \infty$$

for each $T > 0$, then $T_+(f_0) = +\infty$.

Remark 3.3. If f is a solution to (3.4), then, given $\lambda > 0$, also

$$f_\lambda(t, x) := \lambda^{-1}f(\lambda t, \lambda x),$$

is a solution to (3.4) (with initial datum $\lambda^{-1}f(0, \cdot)$). This property identifies $H^{3/2}(\mathbb{R})$ as a critical space for the evolution problem (3.4). Therefore, our result in Theorem 3.2 covers all subcritical spaces.

Remark 3.4. We expect the solutions to be even analytic in space and time away from $t = 0$. However, we prefer to formulate and prove our result in the C^∞ -class, refraining from the considerable technicalities needed for a proof of the analytic counterpart of Lemma 3.6 below (see [11, Proposition 5.1] for a related analyticity result).

In order to study the mapping properties of the operator Ψ we need the following lemmas.

Lemma 3.5. *Given $s \in (3/2, 2)$, we have $\phi_i \in C^\infty(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))$, $i = 1, 2$. Moreover, given $f_0 \in H^s(\mathbb{R})$, the Fréchet derivative $\partial\phi_i(f_0)$ is given by*

$$\partial\phi_i(f_0) = a_i(f_0) \frac{d}{dx},$$

with a_i defined by

$$a_1(f_0) = \frac{f_0'(2 + f_0'^2 + 2\sqrt{1 + f_0'^2})}{\sqrt{1 + f_0'^2}(\sqrt{1 + f_0'^2} + 1 + f_0'^2)^2} \quad \text{and} \quad a_2(f_0) = \frac{1}{(1 + f_0'^2)^{3/2}}.$$

Proof. For the smoothness result we refer to Lemma C.3 in Appendix C. The representations for the derivatives $\partial\phi_i(f_0)$ follow from straightforward calculations. \square

Lemma 3.6. *Given $s \in (3/2, 2)$, we have*

$$\Psi \in C^\infty(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})).$$

Proof. The claim follows from Lemma 3.5 and Corollary C.5. \square

In order to establish our main result in Theorem 3.2 we next prove that, given $f_0 \in H^s(\mathbb{R})$, the Fréchet derivative $\partial\Psi(f_0) \in \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))$ generates a strongly continuous and analytic semigroup in $\mathcal{L}(H^{s-1}(\mathbb{R}))$, that is

$$-\partial\Psi(f_0) \in \mathcal{H}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})).$$

This property, which identifies (3.4) as a nonlinear evolution problem of parabolic type, is established in Theorem 3.7 below.

Theorem 3.7. *Given $f_0 \in H^s(\mathbb{R})$, we have*

$$-\partial\Psi(f_0) \in \mathcal{H}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})). \quad (3.6)$$

The subsequent analysis is devoted to the proof of Theorem 3.7. To start, we fix $f_0 \in H^s(\mathbb{R})$ and $s' \in (3/2, s)$, and we note that

$$\partial\Psi(f_0)[f] = -\frac{\sigma}{4\pi\mu} f' \Psi_1(f_0) - \frac{\sigma}{4\pi\mu} f_0' \partial\Psi_1(f_0)[f] + \frac{\sigma}{4\pi\mu} \partial\Psi_2(f_0)[f], \quad f \in H^s(\mathbb{R}).$$

To calculate the derivatives of Ψ_i we use Lemma C.4 to get

$$\partial B_{n,2}^0(f_0)[f][h] = nB_{n,2}(f_0, f_0)[f, f_0, \dots, f_0, h] - 4B_{n+2,3}(f_0, f_0, f_0)[f, f_0, \dots, f_0, h]$$

and Lemma 3.1 (iii) to rewrite this for $n \geq 0$ as

$$\begin{aligned} \partial B_{n,2}^0(f_0)[f][h] &= h(nB_{n-1,2}^0(f_0)[f'] - 4B_{n+1,3}^0(f_0)[f']) + R_n[f] \\ &= h(nB_{n-1,3}^0(f_0)[f'] + (n-4)B_{n+1,3}^0(f_0)[f']) + R_n[f], \end{aligned}$$

where $nB_{n-1,3}^0(f_0) := 0$ for $n = 0$ and

$$\|R_n f\|_{H^{s-1}} \leq C \|h\|_{H^{s-1}} \|f\|_{H^{s'}}.$$

The constant C is independent of $f \in H^s(\mathbb{R})$ and $h \in H^{s-1}(\mathbb{R})$. From this and the definition of Ψ_i , $i = 1, 2$, we get

$$\partial\Psi_i(f_0)[f] = T_{i,1}(f_0)[f] + T_{i,2}(f_0)[f] + T_{i,\text{lot}}(f_0)[f], \quad i = 1, 2, \quad (3.7)$$

where

$$\begin{aligned}
T_{1,1}(f_0)[f] &:= (B_{0,2}^0 - B_{2,2}^0)[(a_1 + \phi_2 + f'_0 a_2)f'] + B_{1,2}^0[(3(\phi_1 + f'_0 a_1) - a_2)f'] \\
&\quad + B_{3,2}^0[(\phi_1 + f'_0 a_1 + a_2)f'], \\
T_{1,2}(f_0)[f] &:= \phi_1(3f'_0 B_{0,3}^0 - 6B_{1,3}^0 - 6f'_0 B_{2,3}^0 + 2B_{3,3}^0 - f'_0 B_{4,3}^0)[f'] \\
&\quad + \phi_2(-B_{0,3}^0 - 6f'_0 B_{1,3}^0 + 6B_{2,3}^0 + 2f'_0 B_{3,3}^0 - B_{4,3}^0)[f'], \\
T_{2,1}(f_0)[f] &:= -B_{0,2}^0[(\phi_1 + f'_0 a_1 + a_2)f'] + (B_{1,2}^0 - B_{3,2}^0)[(a_1 + \phi_2 + f'_0 a_2)f'] \\
&\quad + B_{2,2}^0[(\phi_1 + f'_0 a_1 - 3a_2)f'], \\
T_{2,2}(f_0)[f] &:= \phi_1(B_{0,3}^0 + 6f'_0 B_{1,3}^0 - 6B_{2,3}^0 - 2f'_0 B_{3,3}^0 + B_{4,3}^0)[f'] \\
&\quad + \phi_2(f'_0 B_{0,3}^0 - 2B_{1,3}^0 - 6f'_0 B_{2,3}^0 + 6B_{3,3}^0 + f'_0 B_{4,3}^0)[f'],
\end{aligned} \tag{3.8}$$

with shortened notation $a_i = a_i(f_0)$, $\phi_i = \phi_i(f_0)$, $B_{n,m}^0 = B_{n,m}^0(f_0)$, and

$$\|T_{i,\text{lot}}(f_0)[f]\|_{H^{s-1}} \leq C \max\{\|\phi_1\|_{H^{s-1}}, \|\phi_2\|_{H^{s-1}}\} \|f\|_{H^{s'}} \leq C \|f\|_{H^{s'}}, \quad f \in H^s(\mathbb{R}). \tag{3.9}$$

Having computed the derivative $\partial\Psi(f_0)$, it remains to establish (3.6), which is achieved via an intricate localization type procedure. To proceed, we fix for each $\varepsilon \in (0, 1)$ a so-called finite ε -localization family, that is a set

$$\{(\pi_j^\varepsilon, x_j^\varepsilon) : -N + 1 \leq j \leq N\} \subset C_0^\infty(\mathbb{R}, [0, 1]) \times \mathbb{R}$$

such that

- $\pi_j^\varepsilon \in C_0^\infty(\mathbb{R}, [0, 1])$,
- $\text{supp } \pi_j^\varepsilon$ is an interval of length ε for all $|j| \leq N - 1$;
- $\text{supp } \pi_N^\varepsilon \subset (-\infty, -1/\varepsilon] \cup [1/\varepsilon, \infty)$;
- $\pi_j^\varepsilon \cdot \pi_l^\varepsilon = 0$ if $[|j - l| \geq 2, \max\{|j|, |l|\} \leq N - 1]$ or $[|l| \leq N - 2, j = N]$;
- $\sum_{j=-N+1}^N (\pi_j^\varepsilon)^2 = 1$;
- $\|(\pi_j^\varepsilon)^{(k)}\|_\infty \leq C\varepsilon^{-k}$ for all $k \in \mathbb{N}$, $-N + 1 \leq j \leq N$;
- $x_j^\varepsilon \in \text{supp } \pi_j^\varepsilon$, $|j| \leq N - 1$.

The real number x_N^ε plays no role in the analysis below. To each finite ε -localization family we associate a second family

$$\{\chi_j^\varepsilon : -N + 1 \leq j \leq N\} \subset C_0^\infty(\mathbb{R}, [0, 1])$$

with the following properties

- $\chi_j^\varepsilon = 1$ on $\text{supp } \pi_j^\varepsilon$;
- $\text{supp } \chi_j^\varepsilon$ is an interval of length 3ε and with the same midpoint as $\text{supp } \pi_j^\varepsilon$, $|j| \leq N - 1$;
- $\text{supp } \chi_N^\varepsilon \subset [|x| \geq 1/\varepsilon - \varepsilon]$, $x_N^\varepsilon := 0$.

Each finite ε -localization family induces norms on $H^r(\mathbb{R})$, $r \geq 0$, that are equivalent to the standard norm. Indeed, it is not difficult to prove that, given $\varepsilon \in (0, 1)$ and $r \geq 0$, there exists a constant $c = c(\varepsilon, r) \in (0, 1)$ such that

$$c\|f\|_{H^r} \leq \sum_{j=-N+1}^N \|\pi_j^\varepsilon f\|_{H^r} \leq c^{-1}\|f\|_{H^r}, \quad f \in H^r(\mathbb{R}). \quad (3.10)$$

It is useful to consider the continuous path $\Phi : [0, 1] \rightarrow \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))$ defined by

$$\Phi(\tau) := -\frac{\tau\sigma}{4\pi\mu}\Psi_1(f_0)\frac{d}{dx} - \frac{\sigma\tau}{4\pi\mu}f_0'\partial\Psi_1(\tau f_0) + \frac{\sigma}{4\pi\mu}\partial\Psi_2(\tau f_0), \quad \tau \in [0, 1].$$

We next locally approximate the operator $\Phi(\tau)$, $\tau \in [0, 1]$, by certain Fourier multipliers $\mathbb{A}_{j,\tau}$. It is worth emphasizing that $\Phi(1) = \partial\Psi(f_0)$, while $\Phi(0)$ is the Fourier multiplier given by

$$\Phi(0) = -\frac{\sigma}{4\mu}H \circ \frac{d}{dx} = -\frac{\sigma}{4\mu}\left(-\frac{d^2}{dx^2}\right)^{1/2},$$

with H denoting the Hilbert transform. The homotopy Φ is used to conclude invertibility of $\lambda - \Phi(1)$ from $\lambda - \Phi(0)$ for sufficiently large λ . We also point out the estimate

$$\|gh\|_{H^{s-1}} \leq 2(\|g\|_\infty\|h\|_{H^{s-1}} + \|h\|_\infty\|g\|_{H^{s-1}}) \quad \text{for } g, h \in H^{s-1}(\mathbb{R}), \quad s \in (3/2, 2), \quad (3.11)$$

which is used several times in the arguments that follow.

Theorem 3.8. *Let $\gamma > 0$ be given and fix $s' \in (3/2, s)$. Then, there exist $\varepsilon \in (0, 1)$, a constant $K = K(\varepsilon)$, and bounded operators*

$$\mathbb{A}_{j,\tau} \in \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})), \quad j \in \{-N+1, \dots, N\}, \quad \tau \in [0, 1],$$

such that

$$\|\pi_j^\varepsilon\Phi(\tau)[f] - \mathbb{A}_{j,\tau}[\pi_j^\varepsilon f]\|_{H^{s-1}} \leq \gamma\|\pi_j^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}} \quad (3.12)$$

for all $j \in \{-N+1, \dots, N\}$, $\tau \in [0, 1]$, and $f \in H^s(\mathbb{R})$. The operators $\mathbb{A}_{j,\tau}$ are defined by

$$\mathbb{A}_{j,\tau} := -\alpha_\tau(x_j^\varepsilon)\left(-\frac{d^2}{dx^2}\right)^{1/2} + \beta_\tau(x_j^\varepsilon)\frac{d}{dx}, \quad |j| \leq N-1, \quad \mathbb{A}_{N,\tau} := -\frac{\sigma}{4\mu}\left(-\frac{d^2}{dx^2}\right)^{1/2},$$

with functions α_τ, β_τ given by

$$\alpha_\tau := \frac{\sigma}{4\mu}\left[a_2(\tau f_0) + \tau f_0' a_1(\tau f_0)\right], \quad \beta_\tau := -\frac{\sigma\tau}{4\pi\mu}\Psi_1(f_0).$$

Proof. Let $\varepsilon \in (0, 1)$. In the following we denote by K constants that may depend on ε .

The estimate (3.9) implies

$$\|\pi_j^\varepsilon f_0' T_{1,\text{lot}}(\tau f_0)[f]\|_{H^{s-1}} + \|\pi_j^\varepsilon T_{2,\text{lot}}(\tau f_0)[f]\|_{H^{s-1}} \leq K\|f\|_{H^{s'}}, \quad -N+1 \leq j \leq N. \quad (3.13)$$

Next we consider the operators $\Psi_1(f_0)(d/dx)$, $f_0' T_{1,j}(\tau f_0)$, and $T_{2,j}(\tau f_0)$, $j = 1, 2$, which we approximate successively.

Step 1. We first consider the term $f'\Psi_1(f_0)$. Using $\Psi_1(f_0) \in H^{s-1}(\mathbb{R}) \hookrightarrow C^{s-3/2}(\mathbb{R})$ and the property $\chi_j^\varepsilon \pi_j^\varepsilon = \pi_j^\varepsilon$ together with (3.11), we get

$$\begin{aligned} & \|\pi_j^\varepsilon f'\Psi_1(f_0) - \Psi_1(f_0)(x_j^\varepsilon)(\pi_j^\varepsilon f)'\|_{H^{s-1}} \\ & \leq 2\|\chi_j^\varepsilon(\Psi_1(f_0) - \Psi_1(f_0)(x_j^\varepsilon))\|_\infty\|(\pi_j^\varepsilon f)'\|_{H^{s-1}} + K\|f\|_{H^{s'}} \\ & \leq \frac{\gamma}{3}\|\pi_j^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}} \end{aligned} \quad (3.14)$$

for $|j| \leq N - 1$, provided that ε is sufficiently small. Furthermore, since $\Psi_1(f_0)$ vanishes at infinity, we have

$$\begin{aligned} \|\pi_N^\varepsilon f' \Psi_1(f_0)\|_{H^{s-1}} &\leq 2\|\chi_N^\varepsilon \Psi_1(f_0)\|_\infty \|(\pi_N^\varepsilon f)'\|_{H^{s-1}} + K\|f\|_{H^{s'}} \\ &\leq \frac{\gamma}{3}\|\pi_N^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}}, \end{aligned} \quad (3.15)$$

provided that ε is sufficiently small.

Step 2. We now consider the operators $f_0' T_{1,2}(\tau f_0)$ and $T_{2,2}(\tau f_0)$. Repeated use of Lemma D.2 and Lemma D.3 yields

$$\begin{aligned} \|\tau \pi_j^\varepsilon f_0' T_{1,2}(\tau f_0)[f] - a_{1,j,\tau}(x_j^\varepsilon) B_{0,0}[(\pi_j^\varepsilon f)']\|_{H^{s-1}} &\leq \frac{\gamma}{6}\|\pi_j^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}}, \\ \|\pi_j^\varepsilon T_{2,2}(\tau f_0)[f] - a_{2,j,\tau}(x_j^\varepsilon) B_{0,0}[(\pi_j^\varepsilon f)']\|_{H^{s-1}} &\leq \frac{\gamma}{6}\|\pi_j^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}} \end{aligned} \quad (3.16)$$

for $|j| \leq N - 1$, and

$$\|\tau \pi_N^\varepsilon f_0' T_{1,2}(\tau f_0)[f]\|_{H^{s-1}} + \|\pi_N^\varepsilon T_{2,2}(\tau f_0)[f]\|_{H^{s-1}} \leq \frac{\gamma}{3}\|\pi_N^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}}, \quad (3.17)$$

provided that ε is sufficiently small. The functions $a_{i,j,\tau}$, $i = 1, 2$, $|j| \leq N - 1$, are given by

$$\begin{aligned} a_{1,j,\tau} &= -\phi_1(\tau f_0) \frac{(\tau f_0')^4 + 3(\tau f_0')^2}{(1 + (\tau f_0')^2)^2} + \phi_2(\tau f_0) \frac{(\tau f_0')^3 - \tau f_0'}{(1 + (\tau f_0')^2)^2}, \\ a_{2,j,\tau} &= -\phi_1(\tau f_0) \frac{(\tau f_0')^2 - 1}{(1 + (\tau f_0')^2)^2} + \phi_2(\tau f_0) \frac{(\tau f_0')^3 - \tau f_0'}{(1 + (\tau f_0')^2)^2}. \end{aligned}$$

Step 3. We now consider the operators $f_0' T_{1,1}(\tau f_0)$ and $T_{2,1}(\tau f_0)$. We first observe that $a_1(f_0)$, $a_2(f_0) - 1$, $\phi_i(f_0) \in H^{s-1}(\mathbb{R})$, $i = 1, 2$. Repeated use of Lemmas D.2–D.6 leads to

$$\begin{aligned} \|\tau \pi_j^\varepsilon f_0' T_{1,1}(\tau f_0)[f] - a_{3,j,\tau}(x_j^\varepsilon) B_{0,0}[(\pi_j^\varepsilon f)']\|_{H^{s-1}} &\leq \frac{\gamma}{6}\|\pi_j^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}}, \\ \|\pi_j^\varepsilon T_{2,1}(\tau f_0)[f] - a_{4,j,\tau}(x_j^\varepsilon) B_{0,0}[(\pi_j^\varepsilon f)']\|_{H^{s-1}} &\leq \frac{\gamma}{6}\|\pi_j^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}} \end{aligned} \quad (3.18)$$

for $|j| \leq N - 1$ and

$$\begin{aligned} \|\tau \pi_N^\varepsilon f_0' T_{1,1}(\tau f_0)[f]\|_{H^{s-1}} + \|\pi_N^\varepsilon T_{2,1}(\tau f_0)[f] - B_{0,0}[(\pi_N^\varepsilon f)']\|_{H^{s-1}} \\ \leq \frac{\gamma}{3}\|\pi_N^\varepsilon f\|_{H^s} + K\|f\|_{H^{s'}}, \end{aligned} \quad (3.19)$$

provided that ε is sufficiently small. The functions $a_{i,j,\tau}$, $i = 3, 4$, $|j| \leq N - 1$, are given by

$$\begin{aligned} a_{3,j,\tau} &= a_1(\tau f_0) \tau f_0' + \phi_1(\tau f_0) \frac{(\tau f_0')^4 + 3(\tau f_0')^2}{(1 + (\tau f_0')^2)^2} - \phi_2(\tau f_0) \frac{(\tau f_0')^3 - \tau f_0'}{(1 + (\tau f_0')^2)^2}, \\ a_{4,j,\tau} &= -a_2(\tau f_0) + \phi_1(\tau f_0) \frac{(\tau f_0')^2 - 1}{(1 + (\tau f_0')^2)^2} - \phi_2(\tau f_0) \frac{(\tau f_0')^3 - \tau f_0'}{(1 + (\tau f_0')^2)^2}. \end{aligned}$$

Gathering (3.13)–(3.19), we conclude that (3.12) holds true and the proof is complete. \square

We now consider the symbol of the Fourier multipliers $\mathbb{A}_{j,\tau}$ found in Theorem 3.8. Making use of $\Psi_1(f_0) \in H^{s-1}(\mathbb{R})$ and recalling the definition of the functions a_i in Lemma 3.5, we conclude there exists $\eta \in (0, 1)$ such that

$$\eta \leq \alpha_\tau \leq \frac{1}{\eta} \quad \text{and} \quad |\beta_\tau| \leq \frac{1}{\eta}, \quad \tau \in [0, 1].$$

Given $\alpha \in [\eta, 1/\eta]$ and $|\beta| \leq 1/\eta$, we now introduce the Fourier multipliers

$$\mathbb{A}_{\alpha,\beta} := -\alpha \left(-\frac{d^2}{dx^2} \right)^{1/2} + \beta \frac{d}{dx} \in \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})).$$

It is a matter of direct computations, see e.g. [11, Proposition 4.3], to find a constant $\kappa_0 \geq 1$ such that

$$\bullet \quad \lambda - \mathbb{A}_{\alpha,\beta} \in \text{Isom}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})), \quad \text{Re } \lambda \geq 1, \quad (3.20)$$

$$\bullet \quad \kappa_0 \|(\lambda - \mathbb{A}_{\alpha,\beta})[f]\|_{H^{s-1}} \geq |\lambda| \cdot \|f\|_{H^{s-1}} + \|f\|_{H^s}, \quad f \in H^s(\mathbb{R}), \text{Re } \lambda \geq 1. \quad (3.21)$$

Theorem 3.8 and the relations (3.20) and (3.21) combined enable us to establish the result announced in Theorem 3.7.

Proof of Theorem 3.7. Let $s' \in (3/2, s)$ and let $\kappa_0 \geq 1$ be the constant in (3.21). Theorem 3.8 with $\gamma := 1/2\kappa_0$ implies that there are $\varepsilon \in (0, 1)$, a constant $K = K(\varepsilon) > 0$ and bounded operators $\mathbb{A}_{j,\tau} \in \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))$, $-N + 1 \leq j \leq N$ and $\tau \in [0, 1]$, satisfying

$$2\kappa_0 \|\pi_j^\varepsilon \Phi(\tau)[f] - \mathbb{A}_{j,\tau}[\pi_j^\varepsilon f]\|_{H^{s-1}} \leq \|\pi_j^\varepsilon f\|_{H^s} + 2\kappa_0 K \|f\|_{H^{s'}}, \quad f \in H^s(\mathbb{R}).$$

Moreover, (3.21) yields

$$2\kappa_0 \|(\lambda - \mathbb{A}_{j,\tau})[\pi_j^\varepsilon f]\|_{H^{s-1}} \geq 2|\lambda| \cdot \|\pi_j^\varepsilon f\|_{H^{s-1}} + 2\|\pi_j^\varepsilon f\|_{H^s}$$

for all $-N + 1 \leq j \leq N$, $\tau \in [0, 1]$, $\text{Re } \lambda \geq 1$, and $f \in H^s(\mathbb{R})$. Combining these inequalities, we conclude that

$$\begin{aligned} 2\kappa_0 \|\pi_j^\varepsilon (\lambda - \Phi(\tau))[f]\|_{H^{s-1}} &\geq 2\kappa_0 \|(\lambda - \mathbb{A}_{j,\tau})[\pi_j^\varepsilon f]\|_{H^{s-1}} - 2\kappa_0 \|\pi_j^\varepsilon \Phi(\tau)[f] - \mathbb{A}_{j,\tau}[\pi_j^\varepsilon f]\|_{H^{s-1}} \\ &\geq 2|\lambda| \cdot \|\pi_j^\varepsilon f\|_{H^{s-1}} + \|\pi_j^\varepsilon f\|_{H^s} - 2\kappa_0 K \|f\|_{H^{s'}}. \end{aligned}$$

We now sum up over j to deduce from (3.10), Young's inequality, and the interpolation property

$$[H^{s_0}(\mathbb{R}), H^{s_1}(\mathbb{R})]_\theta = H^{(1-\theta)s_0 + \theta s_1}(\mathbb{R}), \quad \theta \in (0, 1), \quad -\infty < s_0 \leq s_1 < \infty, \quad (3.22)$$

where $[\cdot, \cdot]_\theta$ denotes the complex interpolation functor, that there exist constants $\kappa \geq 1$ and $\omega > 1$ such that

$$\kappa \|(\lambda - \Phi(\tau))[f]\|_{H^{s-1}} \geq |\lambda| \cdot \|f\|_{H^{s-1}} + \|f\|_{H^s} \quad (3.23)$$

for all $\tau \in [0, 1]$, $\text{Re } \lambda \geq \omega$, and $f \in H^s(\mathbb{R})$.

Additionally, (3.20) implies that $\omega - \Phi(0) \in \text{Isom}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))$. The method of continuity [2, Proposition I.1.1.1] and (3.23) imply that also

$$\omega - \Phi(1) = \omega - \partial\Psi(f_0) \in \text{Isom}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})). \quad (3.24)$$

Combining (3.23) (with $\tau = 1$) and (3.24), we conclude that (3.6) holds true, cf. [2, Chapter I]. \square

We are now in a position to prove the main result, for which we can exploit abstract theory for fully nonlinear parabolic problems from [9].

Proof of Theorem 3.2. Well-posedness: Lemma 3.6 and Theorem 3.7 show that the assumptions of [9, Theorem 8.1.1] are satisfied for the evolution problem (3.4). This theorem ensures that for each $f_0 \in H^s(\mathbb{R})$ there exists a positive time $T > 0$ and a solution $f(\cdot; f_0)$ to (3.4) such that ¹

$$f \in C([0, T], H^s(\mathbb{R})) \cap C^1([0, T], H^{s-1}(\mathbb{R})) \cap C_\alpha^\alpha((0, T], H^s(\mathbb{R}))$$

for some $\alpha \in (0, 1)$. Furthermore, it states that the solution is unique within the set

$$\bigcup_{\alpha \in (0, 1)} C_\alpha^\alpha((0, T], H^s(\mathbb{R})) \cap C([0, T], H^s(\mathbb{R})) \cap C^1([0, T], H^{s-1}(\mathbb{R})).$$

We improve this statement by showing that the solution is actually unique within

$$C([0, T], H^s(\mathbb{R})) \cap C^1([0, T], H^{s-1}(\mathbb{R})).$$

Indeed, suppose $\tilde{f} : [0, T] \rightarrow H^s(\mathbb{R})$ is another solution to (3.4) satisfying the same initial condition f_0 . Since (3.4) is an autonomous problem, we may assume $f(t) \neq \tilde{f}(t)$ for $t \in (0, T]$. Let now $s' \in (3/2, s)$ and set $\alpha := s - s' \in (0, 1)$. Using (3.22), we find a constant $C > 0$ such that

$$\|f(t_1) - f(t_2)\|_{H^{s'}} + \|\tilde{f}(t_1) - \tilde{f}(t_2)\|_{H^{s'}} \leq C|t_1 - t_2|^\alpha, \quad t_1, t_2 \in [0, T], \quad (3.25)$$

which shows that $f, \tilde{f} \in C_\alpha^\alpha((0, T], H^{s'}(\mathbb{R}))$. Applying the uniqueness statement from [9, Theorem 8.1.1] to (3.4) with $\Psi \in C^\omega(H^{s'}(\mathbb{R}), H^{s'-1}(\mathbb{R}))$ shows now that $f = \tilde{f}$ on $[0, T]$. This unique solution can be extended up to a maximal existence time $T_+(f_0)$, see [9, Section 8.2]. Finally, [9, Proposition 8.2.3] shows that the solution map defines a semiflow on $H^s(\mathbb{R})$. This proves (i).

Parabolic smoothing: The uniqueness statement in (i) enables us to use a parameter trick which was successfully applied also to other problems, cf., e.g., [3, 5, 11, 12], in order to establish (ia) and (ib). In our setting the proof details are similar to those in [10, Theorem 1.2 (v)] or [1, Theorem 1.2 (ii)] and therefore we omit them.

Global existence: We prove the statement by contradiction. Assume there exists a maximal solution $f \in C([0, T_+), H^s(\mathbb{R})) \cap C^1([0, T_+), H^{s-1}(\mathbb{R}))$ to (3.4) with $T_+ < \infty$ and such that

$$\sup_{[0, T_+)} \|f(t)\|_{H^s} < \infty. \quad (3.26)$$

The bound (3.26) together with Lemma 3.1 (ii) implies that

$$\sup_{[0, T_+)} \left\| \frac{df}{dt}(t) \right\|_{H^{s-1}} = \sup_{[0, T_+)} \|\Psi(f(t))\|_{H^{s-1}} < \infty. \quad (3.27)$$

Choosing $s' \in (3/2, s)$, we may argue as above, see (3.25), to conclude from (3.26) and (3.27) that $f : [0, T_+) \rightarrow H^{s'}(\mathbb{R})$ is uniformly continuous. Applying [9, Proposition 8.2.1], to (3.4) with $\Psi \in C^\omega(H^{s'}(\mathbb{R}), H^{s'-1}(\mathbb{R}))$ and the arguments used in the proof of (i), we may extend the solution f to an interval $[0, T'_+)$ with $T_+ < T'_+$ and such that

$$f \in C([0, T'_+), H^{s'}(\mathbb{R})) \cap C^1([0, T'_+), H^{s'-1}(\mathbb{R})).$$

¹Given $\alpha \in (0, 1)$ and $T > 0$, we set

$$C_\alpha^\alpha((0, T], H^s(\mathbb{R})) := \left\{ f : (0, T] \rightarrow X : f \text{ is bounded and } [f]_{C_\alpha^\alpha} := \sup_{s \neq t} \frac{|t^\alpha f(t) - s^\alpha f(s)|}{|t - s|^\alpha} < \infty \right\}.$$

The parabolic smoothing property established in (iib) (with s replaced by s') implies in particular that $f \in C([0, T'_+), H^s(\mathbb{R}))$, in contradiction to the maximality of f . This completes our arguments. \square

APPENDIX A. THE HYDRODYNAMIC POTENTIAL NEAR Γ

This appendix is devoted to the study of the properties of the functions (v^\pm, q^\pm) defined in (2.8) near the boundary Γ . Lemma A.1 below establishes the corresponding part of Theorem 2.1.

Lemma A.1. *Given $f \in H^3(\mathbb{R})$, the functions (v^\pm, q^\pm) given by (2.8) satisfy*

$$(v^\pm, q^\pm) \in C^1(\overline{\Omega^\pm}, \mathbb{R}^2) \times C(\overline{\Omega^\pm})$$

and solve the equations

$$\left. \begin{aligned} [v] &= 0 && \text{on } \Gamma, \\ [T(v, q)](\nu \circ \Xi^{-1}) &= -\sigma(\kappa\nu) \circ \Xi^{-1} && \text{on } \Gamma. \end{aligned} \right\} \quad (\text{A.1})$$

Before establishing Lemma A.1 we make the following observation.

Remark A.2. It is shown in the proof of Lemma A.1 that not only v is continuous in \mathbb{R}^2 , but also the first order partial derivatives of v , that is

$$v \in C^1(\mathbb{R}^2).$$

In particular, we get

$$[T(v, q)]\nu = -[q]\nu \quad \text{on } \Gamma.$$

Proof of Lemma A.1. From [10, Lemma 2.1 and Eq. (2.7)] we infer that, given $\psi \in H^1(\mathbb{R})$, the functions $A^\pm : \Omega^\pm \rightarrow \mathbb{R}$ defined by

$$A^\pm(x, y) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{(\xi, \eta)^\top}{r^2} \psi(s) ds = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{(\xi^3 + \xi\eta^2, \xi^2\eta + \eta^3)^\top}{r^4} \psi(s) ds, \quad (\text{A.2})$$

belong to $C(\overline{\Omega^\pm})$ and, given $x \in \mathbb{R}$, we have

$$A^\pm(x, f(x)) = \frac{1}{2\pi} \text{PV} \int_{\mathbb{R}} \frac{(x-s, f(x)-f(s))^\top}{(x-s)^2 + (f(x)-f(s))^2} \psi(s) ds \pm \frac{1}{2} \frac{\nu\psi}{\omega}(x), \quad (\text{A.3})$$

where PV denotes the principal value. Recalling (2.2) and (2.8), it follows that $q^\pm \in C(\overline{\Omega^\pm})$ and

$$[q] \circ \Xi = \sigma\kappa. \quad (\text{A.4})$$

We next consider the behavior of the velocity v^\pm near Γ . To this end, we first introduce the integral operators $[\phi \mapsto Z_n[\phi]]$, $n = 0, \dots, 3$, by

$$Z_n[\phi](x, y) := \int_{\mathbb{R}} \frac{\xi^{3-n}\eta^n}{r^4} \phi(s) ds, \quad (x, y) \in \mathbb{R}^2 \setminus \Gamma, \quad \phi \in H^1(\mathbb{R}),$$

and let

$$\{w\}^\pm(x^*) := \lim_{\Omega^\pm \ni (x, y) \rightarrow (x^*, f(x^*))} w(x, y), \quad x^* \in \mathbb{R},$$

be the one-sided limits of any function $w : \mathbb{R}^2 \setminus \Gamma \rightarrow \mathbb{R}$ at Γ , whenever these limits exist.

Using this notation and the operators $B_{n,2}^0$ defined in (3.2), we have from (A.3)

$$\left. \begin{aligned} \{Z_0[\phi] + Z_2[\phi]\}^\pm &= B_{0,2}^0(f)[\phi] + B_{2,2}^0(f)[\phi] \mp \frac{\pi f'}{\omega^2}, \\ \{Z_1[\phi] + Z_3[\phi]\}^\pm &= B_{1,2}^0(f)[\phi] + B_{3,2}^0(f)[\phi] \pm \frac{\pi}{\omega^2}. \end{aligned} \right\} \quad (\text{A.5})$$

For $(x, y) \in \mathbb{R}^2 \setminus \Gamma$, $\phi \in H^1(\mathbb{R})$, consider further the integral

$$I[\phi](x, y) := \int_{\mathbb{R}} \frac{(\xi\eta, \eta^2)^\top}{r^2} \phi'(s) ds. \quad (\text{A.6})$$

A straightforward application of Lebesgue's dominated convergence theorem shows that I extends continuously to Γ , that is $I \in C(\mathbb{R}^2)$. Integration by parts yields

$$I[\phi] = ((Z_0 - Z_2)[f'\phi] - (Z_1 - Z_3), 2Z_1[f'\phi] - 2Z_2[\phi])^\top \quad \text{on } \mathbb{R}^2 \setminus \Gamma, \quad (\text{A.7})$$

and

$$I[\phi] \circ \Xi = ((B_{0,2}^0(f) - B_{2,2}^0(f))[f'\phi] - (B_{1,2}^0(f) - B_{3,2}^0(f))[\phi], 2B_{1,2}^0(f)[f'\phi] - 2B_{2,2}(f)[\phi])^\top. \quad (\text{A.8})$$

So, by continuity of I ,

$$\left. \begin{aligned} \{(Z_0 - Z_2)[f'\phi] - (Z_1 - Z_3)\}^\pm &= (B_{0,2}^0(f) - B_{2,2}^0(f))[f'\phi] - (B_{1,2}^0(f) - B_{3,2}^0(f))[\phi], \\ \{2Z_1[f'\phi] - 2Z_2[\phi]\}^\pm &= 2B_{1,2}^0(f)[f'\phi] - 2B_{2,2}(f)[\phi]. \end{aligned} \right\} \quad (\text{A.9})$$

Observe that Eqns. (A.5), (A.9) constitute a linear system of the form

$$\left\{ \sum_{n=0}^3 Z_n[a_{in}\phi] \right\}^\pm = \sum_{n=0}^3 B_{n,2}^0(f)[a_{in}\phi] \pm J_i\phi, \quad i = 1, \dots, 4, \quad \phi \in H^1(\mathbb{R}), \quad (\text{A.10})$$

with coefficient matrix

$$\mathcal{A} := (a_{in}) := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ f' & -1 & -f' & 1 \\ 0 & 2f' & -2 & 0 \end{pmatrix}$$

and with $(J_1, \dots, J_4) := \frac{\pi}{\omega^2}(-f', 1, 0, 0)^\top$. The matrix \mathcal{A} is regular and has inverse

$$\mathcal{A}^{-1} := (a^{ki}) := \frac{1}{2\omega^2} \begin{pmatrix} 2 + f'^2 & -f' & -f' & 1 \\ f' & 1 & -1 & f' \\ f'^2 & f' & -f' & -1 \\ -f' & 1 + 2f'^2 & 1 & -f' \end{pmatrix}.$$

Observe that for $k = 0, \dots, 3$, $i = 1, \dots, 4$, and $\phi \in H^1(\mathbb{R})$ we have $a^{ki}\phi \in H^1(\mathbb{R})$.

For $k = 0, \dots, 3$, replace ϕ by $a^{ki}\phi$ in the i -th equation of (A.10) and sum over i . This gives

$$\{Z_k[\phi]\}^\pm = B_{k,2}^0(f)[\phi] \pm \sum_{i=1}^4 a^{ki} J_i \phi,$$

or equivalently

$$\begin{aligned} \frac{1}{2\pi} \left\{ \int_{\mathbb{R}} \frac{(\xi^3, \xi^2\eta, \xi\eta^2, \eta^3)^\top}{r^4} \phi ds \right\}^\pm &= \frac{(B_{0,2}^0(f), B_{1,2}^0(f), B_{2,2}^0(f), B_{0,2}^0(f))^\top [\phi]}{2\pi} \\ &\mp \left(\frac{f^3 + 3f'}{4\omega^4}, \frac{f'^2 - 1}{4\omega^4}, \frac{f'^3 - f'}{4\omega^4}, -\frac{3f'^2 + 1}{4\omega^4} \right)^\top \phi. \end{aligned}$$

In view of this, we get directly from (2.7) and (2.8)₁ that $[v] = 0$ and

$$\begin{aligned} v_1|_{\Gamma \circ \Xi} &= \frac{(B_{2,2}^0(f) - B_{0,2}^0(f))[g_1 - f'g_2] - B_{1,2}^0(f)[3f'g_1 + g_2] - B_{3,2}^0(f)[f'g_1 - g_2]}{4\pi\mu}, \\ v_2|_{\Gamma \circ \Xi} &= \frac{B_{0,2}^0(f)[f'g_1 - g_2] + (B_{3,2}^0(f) - B_{1,2}^0(f))[g_1 - f'g_2] - B_{2,2}^0(f)[f'g_1 + 3g_2]}{4\pi\mu}. \end{aligned} \tag{A.11}$$

Moreover, first differentiating with respect to x and y under the integral in (2.8)₁ and then using integration by parts and (2.2) we find

$$\begin{aligned} \partial_x v(x, y) &= \sigma \int_{\mathbb{R}} \partial_\xi \mathcal{U}_0^1(\xi, \eta)(f'\kappa)(s) - \partial_\xi \mathcal{U}_0^2(\xi, \eta)\kappa(s) ds, \\ \partial_y v(x, y) &= \sigma \int_{\mathbb{R}} \partial_\eta \mathcal{U}_0^1(\xi, \eta)(f'\kappa)(s) - \partial_\eta \mathcal{U}_0^2(\xi, \eta)\kappa(s) ds. \end{aligned}$$

It is now straightforward to check that $v^\pm \in C^1(\overline{\Omega^\pm})$ and $[\nabla v] = 0$. Together with (A.4), this implies (A.1)₂, and the proof is complete. \square

APPENDIX B. THE HYDRODYNAMIC POTENTIAL IN THE FAR-FIELD LIMIT

In this appendix we prove that the functions (v, q) defined in (2.8) satisfy the far-field boundary condition (2.5)₅. While the claim for q follows directly from [10, Lemma 2.1], proving that the velocity vanishes at infinity is more elaborate and necessitates some preparation.

Recall that $f \in H^3(\mathbb{R})$ is fixed. We define functions $[\phi \mapsto (F, G)[\phi]]$ according to (cf. (A.6))

$$(F, G)[\phi](x, y) := I[\phi](x, y) = \int_{\mathbb{R}} \frac{(\xi\eta, \eta^2)}{r^2} \phi'(s) ds, \quad \phi \in H^1(\mathbb{R}). \tag{B.1}$$

We recall from Appendix A that $F, G \in C(\mathbb{R}^2)$. Moreover, from (A.8) and Lemma 3.1 (ii), we get

$$(F, G)|_{\Gamma \circ \Xi} \in C^\beta(\mathbb{R}) \tag{B.2}$$

for some $\beta \in (0, 1)$, and

$$(F, G)|_{\Gamma \circ \Xi} \rightarrow 0 \quad \text{for } |x| \rightarrow \infty. \tag{B.3}$$

We first prove a bound for F and G at moderate distances from the interface, in terms of their values at the interface.

Lemma B.1 (Vertical differences). *Given $f \in H^3(\mathbb{R})$ and $\phi \in H^2(\mathbb{R})$, there exist constants $\alpha \in (0, 1)$ and $C_0 > 0$ such that for*

$$(x, y) \in S := \mathbb{R} \times [-\|f\|_\infty - 1, \|f\|_\infty + 1]$$

we have

$$|(F, G)(x, y) - (F, G)(x, f(x))| \leq C_0 |y - f(x)|^\alpha.$$

Proof. We show the estimate for F only, the arguments for G are analogous with some obvious modifications. Given $z = (x, y) \in S$, we choose $\bar{z} := (x_0, f(x_0))$ such that

$$|z - \bar{z}| = \min\{|z - (s, f(s))| \mid s \in \mathbb{R}\}.$$

After a change of variables we split

$$\begin{aligned} & |F(x, y) - F(x_0, f(x_0))| \\ & \leq \int_{\mathbb{R}} |\phi'(x-s) - \phi'(x_0-s)| \frac{|s||y-f(x-s)|}{s^2 + (y-f(x-s))^2} ds \\ & \quad + \int_{\mathbb{R}} |\phi'(x_0-s)| \left| \frac{s(y-f(x-s))}{s^2 + (y-f(x-s))^2} - \frac{s(f(x_0)-f(x_0-s))}{s^2 + (f(x_0)-f(x_0-s))^2} \right| ds. \end{aligned}$$

We estimate the terms on the right separately. For the first one we use $\phi' \in C^{2\alpha}(\mathbb{R})$ for some $\alpha \in (0, \min\{\beta, 1/2\}]$ to obtain

$$\begin{aligned} & \int_{\mathbb{R}} |\phi'(x-s) - \phi'(x_0-s)| \frac{|s||y-f(x-s)|}{s^2 + (y-f(x-s))^2} ds \\ & \leq C|x-x_0|^\alpha \int_{\mathbb{R}} |\phi'(x-s) - \phi'(x_0-s)|^{1/2} \frac{|s||y-f(x-s)|}{s^2 + (y-f(x-s))^2} ds \\ & \leq C|x-x_0|^\alpha \|\phi'\|_2^{1/2} \left(\int_{\mathbb{R}} \left(\frac{|s||y-f(x-s)|}{s^2 + (y-f(x-s))^2} \right)^{4/3} ds \right)^{3/4} \\ & \leq C|x-x_0|^\alpha \left(\int_{\mathbb{R}} \min\{1, s^{-4/3}\} ds \right)^{3/4} \leq C|z - \bar{z}|^\alpha. \end{aligned}$$

To estimate the second term we write for brevity $\zeta := y - f(x-s)$, $\zeta_0 := f(x_0) - f(x_0-s)$ and observe

$$\begin{aligned} \left| \frac{s\zeta}{s^2 + \zeta^2} - \frac{s\zeta_0}{s^2 + \zeta_0^2} \right| &= \left| \frac{s^3 - s\zeta\zeta_0}{(s^2 + \zeta^2)(s^2 + \zeta_0^2)} \right| |\zeta - \zeta_0| \leq \left(\frac{|s|}{s^2 + \zeta^2} + \frac{|\zeta|}{s^2 + \zeta^2} \right) |\zeta - \zeta_0| \\ &\leq \frac{2}{(s^2 + \zeta^2)^{1/2}} (|y - f(x_0)| + C|x - x_0|) \leq \frac{C}{(s^2 + \zeta^2)^{1/2}} |z - \bar{z}|, \end{aligned}$$

to obtain

$$\int_{\mathbb{R}} |\phi'(x_0-s)| \left| \frac{s\zeta}{s^2 + \zeta^2} - \frac{s\zeta_0}{s^2 + \zeta_0^2} \right| ds \leq C \int_{\mathbb{R}} |\phi'(x_0-s)| \frac{|z - \bar{z}|}{(s^2 + \zeta^2)^{1/2}} ds.$$

We split the integral on the right. For $|s| < 1$ we use the minimality property of \bar{z} to obtain

$$\frac{|z - \bar{z}|}{(s^2 + \zeta^2)^{1/2}} = \frac{|z - \bar{z}|^{1/2}}{(s^2 + \zeta^2)^{1/4}} \frac{|z - \bar{z}|^{1/2}}{|(x, y) - (x-s, f(x-s))|^{1/2}} \leq \frac{|z - \bar{z}|^{1/2}}{s^{1/2}}$$

and thus

$$\int_{\{|s| < 1\}} |\phi'(x_0-s)| \frac{|z - \bar{z}|}{(s^2 + \zeta^2)^{1/2}} ds \leq C \|\phi'\|_\infty |z - \bar{z}|^{1/2} \int_{\{|s| < 1\}} s^{-1/2} ds \leq C|z - \bar{z}|^{1/2}.$$

For $|s| > 1$ we estimate directly

$$\int_{\{|s|>1\}} |\phi'(x_0 - s)| \frac{|z - \bar{z}|}{(s^2 + \zeta^2)^{1/2}} ds \leq |z - \bar{z}| \|\phi'\|_2 \left(\int_{\{|s|>1\}} s^{-2} ds \right)^{1/2} \leq C|z - \bar{z}|.$$

Summarizing and using the boundedness of F on S (which follows by applying Hölder's inequality to (B.1)) we get

$$|F(z) - F(\bar{z})| \leq C|z - \bar{z}|^\alpha,$$

and consequently, using (B.2),

$$\begin{aligned} |F(x, y) - F(x, f(x))| &\leq |F(x, y) - F(x_0, f(x_0))| + |F(x_0, f(x_0)) - F(x, f(x))| \\ &\leq C(|z - \bar{z}|^\alpha + |x - x_0|^\alpha) \leq C|z - \bar{z}|^\alpha \\ &\leq C|y - f(x)|^\alpha, \end{aligned}$$

where the minimality property of \bar{z} has been used again in the last step. \square

We next prove that the functions F, G defined in (B.1) vanish at infinity.

Lemma B.2. *Given $f \in H^3(\mathbb{R})$ and $\phi \in H^2(\mathbb{R})$, we have*

$$(F, G) \rightarrow 0 \quad \text{for } |(x, y)| \rightarrow \infty.$$

Proof. We will show the result for F , the proof for G is essentially analogous. The result is proved in the following three steps:

- (i) For any $\varepsilon > 0$ there are $x_0 > 0, \delta \in (0, 1)$ such that $|y| < \delta$ and $|x| > x_0$ imply $|F(x, y)| < \varepsilon$,
- (ii) For any $\varepsilon > 0$ there is a $\delta \in (0, 1)$ such that $x \in \mathbb{R}$ and $|y| > \delta^{-1}$ imply $|F(x, y)| < \varepsilon$,
- (iii) For all $\varepsilon > 0, \delta \in (0, 1)$ there is an $x_1 > 0$ such that $|x| > x_1$ and $\delta \leq |y| \leq \delta^{-1}$ imply $|F(x, y)| < \varepsilon$.

To show (i), fix $\varepsilon > 0$, choose first $\delta > 0$ small enough to ensure $C_0(2\delta)^\alpha < \varepsilon/2$ with C_0 and α from Lemma B.1, and then x_0 large enough to guarantee that $|f(x)| < \delta$ and $|F(x, f(x))| < \varepsilon/2$ whenever $|x| > x_0$, which is possible by (B.3). Now it follows from Lemma B.1 that $|F(x, y)| < \varepsilon$ for (x, y) with $|x| > x_0, |y| < \delta$.

For (ii) we have to prove that $F(x, y) \rightarrow 0$ for $|y| \rightarrow \infty$, uniformly in $x \in \mathbb{R}$. From (A.7) we immediately have

$$|F(x, y)| \leq C \int_{\mathbb{R}} \frac{|\phi(s)|(|\xi| + |\eta|)}{r^2} ds.$$

Choosing $|y| > 2\|f\|_\infty$, we get $|y|/2 \leq |\eta| \leq 3|y|/2$ and using the Cauchy-Schwarz inequality we get

$$\begin{aligned} |F(x, y)| &\leq C \int_{\mathbb{R}} \frac{|\phi(s)|(|\xi| + |y|)}{4\xi^2 + |y|^2} ds \leq C\|\phi\|_2 \left(\int_{\mathbb{R}} \left(\frac{|\xi| + |y|}{4\xi^2 + |y|^2} \right)^2 ds \right)^{1/2} \\ &\leq \frac{C}{|y|^{1/2}} \left(\int_{\mathbb{R}} \frac{(t+1)^2}{(4t^2+1)^2} dt \right)^{1/2} \leq \frac{C}{|y|^{1/2}}, \end{aligned}$$

where we changed variables according to $t := (x - s)/|y|$ in the last step. This proves (ii).

To show (iii), let $\varepsilon > 0$ and $\delta \in (0, 1)$ be fixed. Given $y \in \mathbb{R}$ with $\delta \leq |y| \leq \delta^{-1}$, it follows that $|\eta| \leq \delta^{-1} + \|f\|_\infty$. Choose $s_0 > 0$ large enough to ensure

$$\bullet \quad (\delta^{-1} + \|f\|_\infty) \sqrt{2\pi/\delta} \left(\int_{|s|>s_0} \phi'^2(s) ds \right)^{1/2} < \varepsilon/2 \quad \text{and} \quad (\text{B.4})$$

$$\bullet \quad |f(s)| < \delta/2 \quad \text{for } |s| > s_0. \quad (\text{B.5})$$

Set $x_0 := 2s_0$. Given $x \in \mathbb{R}$ with $|x| \geq x_0$, we obtain in view of (B.1) the estimate

$$|F(x, y)| \leq \int_{\mathbb{R}} \frac{|\xi\eta|}{r^2} |\phi'(s)| ds.$$

We split the integral on the right as follows. If $|s| < s_0$, then $|\xi| > |x|/2$ and

$$\int_{\{|s|<s_0\}} \frac{|\xi\eta|}{r^2} |\phi'(s)| ds \leq \frac{4s_0(\delta^{-1} + \|f\|_\infty) \|\phi'\|_\infty}{|x|} = \frac{C}{|x|} < \frac{\varepsilon}{2}$$

provided that $|x| > x_1$ with $x_1 > x_0$ chosen sufficiently large.

If $|s| > s_0$, then $|\eta| > \delta/2$ by (B.5) and, using (B.4) and the Cauchy-Schwarz inequality, we get

$$\begin{aligned} \int_{\{|s|>s_0\}} \frac{|\xi\eta|}{r^2} |\phi'(s)| ds &\leq (\delta^{-1} + \|f\|_\infty) \int_{\{|s|>s_0\}} \frac{|\xi|}{\xi^2 + (\delta/2)^2} |\phi'(s)| ds \\ &\leq (\delta^{-1} + \|f\|_\infty) \left(\int_{\mathbb{R}} \frac{1}{t^2 + (\delta/2)^2} dt \right)^{1/2} \left(\int_{|s|>s_0} \phi'^2(s) ds \right)^{1/2} \\ &\leq (\delta^{-1} + \|f\|_\infty) \sqrt{2\pi/\delta} \left(\int_{|s|>s_0} \phi'^2(s) ds \right)^{1/2} < \frac{\varepsilon}{2}. \end{aligned}$$

This completes the proof. \square

We are now in a position to prove the desired decay behavior.

Lemma B.3. *Given $f \in H^3(\mathbb{R})$, the functions (v^\pm, q^\pm) given by (2.8) satisfy*

$$(v^\pm, q^\pm) \rightarrow 0 \quad \text{for } |(x, y)| \rightarrow \infty.$$

Proof. Since $f \in H^3(\mathbb{R})$, for $\psi \in H^1(\mathbb{R})$ the functions A_\pm defined in (A.2) satisfy $A_\pm \rightarrow 0$ for $|(x, y)| \rightarrow \infty$, cf. [10, Lemma 2.1]. Recalling the definition (2.8)₂ of q , it immediately follows (since $g'_k \in H^1(\mathbb{R})$, $k = 1, 2$) that

$$q \rightarrow 0 \quad \text{for } |(x, y)| \rightarrow \infty.$$

Moreover, the equation (2.8)₁ together with the relation $\partial_s(\xi^2/r^2) = -\partial_s(\eta^2/r^2)$ enable us to write

$$v(x, y) = \frac{\sigma}{4\pi\mu} \int_{\mathbb{R}} \frac{1}{r^2} \begin{pmatrix} -\eta^2 & \xi\eta \\ \xi\eta & \eta^2 \end{pmatrix} g'(s) ds - \frac{\sigma}{4\pi\mu} \int_{\mathbb{R}} \frac{\xi + f'(s)\eta}{r^2} g(s) ds, \quad (x, y) \in \mathbb{R}^2 \setminus \Gamma.$$

Since $g_k \in H^2(\mathbb{R})$, $k = 1, 2$, Lemma B.2 implies that the first integral vanishes at infinity. Due to [10, Lemma 2.1], also the second integral vanishes in the far field limit, hence

$$v^\pm \rightarrow 0 \quad \text{for } |(x, y)| \rightarrow \infty. \quad \square$$

APPENDIX C. SMOOTHNESS OF SOME NONLINEAR OPERATORS

In this appendix we establish the smoothness of certain nonlinear operators we are confronted with in Section 3. In the following $r \in (1/2, 1)$ is fixed.

Lemma C.1. *Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ be locally Lipschitz continuous, i.e. for any $K > 0$ there is a constant $L_K > 0$ such that*

$$|\psi(x) - \psi(x')| \leq L_K |x - x'| \quad \text{for all } x, x' \in [-K, K].$$

Let $z \in H^r(\mathbb{R})$. Then, the map $[h \mapsto (\psi \circ z)h]$ is in $\mathcal{L}(H^r(\mathbb{R}))$.

Corollary C.2. *If, in addition to the assumptions of Lemma C.1, ψ is strictly positive, then the linear operator $[h \mapsto (\psi \circ z)h]$ is an isomorphism of $H^r(\mathbb{R})$.*

Proof of Lemma C.1. Recall that a norm in $H^r(\mathbb{R})$, which is equivalent to the standard norm, is given by

$$\|z\|_{H^r}^2 = \|z\|_2^2 + [z]_{H^r}^2 \quad \text{with} \quad [z]_{H^r}^2 = \int_{\mathbb{R}} \frac{\|\tau_\xi z - z\|_2^2}{|\xi|^{1+2r}} d\xi,$$

where $\tau_\xi := [z \mapsto z(\cdot - \xi)]$ denotes the right shift operator. Let $K = \|z\|_\infty$ and observe

$$\|\psi \circ z\|_\infty \leq |\psi(0)| + KL_K,$$

$$\|(\psi \circ z)h\|_2 \leq \|\psi \circ z\|_\infty \|h\|_2 \leq C \|h\|_2,$$

$$[(\psi \circ z)h]_{H^r}^2 = \int_{\mathbb{R}} \frac{\|\tau_\xi((\psi \circ z)h) - (\psi \circ z)h\|_2^2}{|\xi|^{1+2r}} d\xi,$$

$$\|\tau_\xi((\psi \circ z)h) - (\psi \circ z)h\|_2 \leq \|\psi \circ z\|_\infty \|\tau_\xi h - h\|_2 + \|h\|_\infty L_K \|\tau_\xi z - z\|_2$$

to find

$$\|(\psi \circ z)h\|_{H^r} \leq C \|h\|_{H^r}, \quad h \in H^r(\mathbb{R}).$$

□

We now use Corollary C.2 to establish in Lemma C.3 the smoothness of three nonlinear operators. Lemma C.3 is the main argument in the proof of Lemma 3.5.

Lemma C.3. *The maps*

$$(i) \quad z \mapsto \sqrt{z^2 + 1} - 1, \quad (ii) \quad z \mapsto \frac{z^2}{\sqrt{z^2 + 1} + z^2 + 1}, \quad (iii) \quad z \mapsto \frac{z}{\sqrt{z^2 + 1}}$$

are (C^∞) -smooth from $H^r(\mathbb{R})$ to $H^r(\mathbb{R})$.

Proof. Fix any $z_0 \in H^r(\mathbb{R})$. Note first that

$$u_0 := \sqrt{z_0^2 + 1} - 1 = \frac{z_0}{\sqrt{z_0^2 + 1} + 1} z_0 \in H^r(\mathbb{R})$$

by Lemma C.1. Furthermore, a pair $(u, z) \in (H^r(\mathbb{R}))^2$ close to (u_0, z_0) is a solution to the equation

$$F(z, u) = u^2 + 2u - z^2 = 0$$

if and only if $u = U(z) := \sqrt{z^2 + 1} - 1$. The mapping F is smooth from $(H^r(\mathbb{R}))^2$ to $H^r(\mathbb{R})$, and its Fréchet derivative $\partial_u F(z_0, u_0)$ is given by

$$\partial_u F(z_0, u_0)h = 2h\sqrt{z_0^2 + 1},$$

which is an isomorphism of $H^r(\mathbb{R})$ by Corollary C.2. Now (i) follows from (the C^∞ -version of) the Implicit Function theorem.

The proof of (ii) is similar. First we note that

$$v_0 := \frac{z_0^2}{\sqrt{z_0^2 + 1} + z_0^2 + 1} \in H^r(\mathbb{R})$$

by Lemma C.1, and then we apply the Implicit Function theorem near (z_0, v_0) to the equation

$$G(z, v) = z^2v + 2v + (\sqrt{z^2 + 1} - 1)v - z^2 = 0,$$

where G is smooth from $(H^r(\mathbb{R}))^2$ to $H^r(\mathbb{R})$ due to (i). Furthermore

$$\partial_v G(z_0, v_0) = [h \mapsto (\sqrt{z_0^2 + 1} + z_0^2 + 1)h]$$

is an isomorphism due to Corollary C.2.

Finally, (iii) follows from (ii) and the identity

$$\frac{z}{\sqrt{z^2 + 1}} = z - z \frac{z^2}{\sqrt{z^2 + 1} + z^2 + 1}.$$

□

In the final part of this appendix we establish the smoothness of certain multilinear singular operators $B_{n,m}^k$. Corollary C.5 below implies in particular that, given $n, m \in \mathbb{N}$, the mapping

$$[f \mapsto B_{n,m}^0(f)[\cdot]] : H^s(\mathbb{R}) \rightarrow \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})),$$

where $s \in (3/2, 2)$ and with $B_{n,m}^0$ as defined in (3.2), is smooth. This property is essential when proving Lemma 3.6.

We start by introducing, for given $k, n, m \in \mathbb{N}$, operators

$$B_{n,m}^k : H^s(\mathbb{R}) \longrightarrow \mathcal{L}_{\text{sym}}^k(H^s(\mathbb{R}), \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R})))$$

by

$$B_{n,m}^k(f)[f_1, \dots, f_k][h] = B_{n+k,m}(f, \dots, f)[f, \dots, f, f_1, \dots, f_k, h], \quad h \in H^s(\mathbb{R}),$$

where $B_{n+k,m}$ are defined in (3.1). Observe that for $k = 0$ this definition agrees with (3.2).

Lemma C.4. *Given $k, n, m \in \mathbb{N}$ and $f \in H^s(\mathbb{R})$, the map $B_{n,m}^k$ is Fréchet differentiable at f , and the Fréchet derivative $\partial B_{n,m}^k(f)$ is given by*

$$\partial B_{n,m}^k(f)[g][f_1, \dots, f_k] = nB_{n-1,m}^{k+1}(f)[f_1, \dots, f_k, g] - 2mB_{n+1,m+1}^{k+1}(f)[f_1, \dots, f_k, g]$$

for $g \in H^s(\mathbb{R})$.

A straightforward consequence of Lemma C.4 is the following corollary.

Corollary C.5. *Given $k, n, m \in \mathbb{N}$, we have*

$$B_{n,m}^k \in C^\infty(H^s(\mathbb{R}), \mathcal{L}_{\text{sym}}^k(H^s(\mathbb{R}), \mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))))).$$

Proof of Lemma C.4. Given $f, g, f_1, \dots, f_k \in H^s(\mathbb{R})$, we consider the remainder

$$\begin{aligned} R(f, g)[f_1, \dots, f_k] &:= (B_{n,m}^k(f + g) - B_{n,m}^k(f))[f_1, \dots, f_k] \\ &\quad - nB_{n-1,m}^{k+1}(f)[f_1, \dots, f_k, g] + 2mB_{n+1,m+1}^{k+1}(f)[f_1, \dots, f_k, g] \end{aligned}$$

and represent it (by elementary algebraic operations) as

$$\begin{aligned}
& R(f, g)[f_1, \dots, f_k] \\
&= \sum_{j=0}^{n-2} (n-j-1) B_{n+k, m}(f+g, \dots, f+g) \underbrace{[f+g, \dots, f+g]}_j \underbrace{[f, \dots, f]}_{n-j-2} [f_1, \dots, f_k, g, g, \cdot] \\
&\quad - \sum_{l=0}^{m-1} B_{n+k+2, m+1} \underbrace{(f+g, \dots, f+g)}_{m-l} \underbrace{[f, \dots, f]}_{l+1} [f, \dots, f, n(2f+g) + f, f_1, \dots, f_k, g, g, \cdot] \\
&\quad + 2 \sum_{l=0}^{m-1} \sum_{j=0}^{m-l-1} B_{n+k+4, m+2} \underbrace{(f+g, \dots, f+g)}_{m-l-j} \underbrace{[f, \dots, f]}_{l+j+2} [f, \dots, f, 2f+g, f_1, \dots, f_k, g, g, \cdot].
\end{aligned}$$

Sums with negative upper summation limit are to be neglected. For $\|g\|_{H^s} \leq 1$, we obtain from Lemma 3.1 (ii)

$$\|R(f, g)[f_1, \dots, f_k]\|_{\mathcal{L}(H^s(\mathbb{R}), H^{s-1}(\mathbb{R}))} \leq C \prod_{i=1}^k \|f_i\|_{H^s} \|g\|_{H^s}^2.$$

This implies the result. \square

APPENDIX D. SOME AUXILIARY RESULTS RELATED TO LOCALIZATION

The following commutator estimate is used in the proof of Lemmas D.5 and D.6 below.

Lemma D.1. *Let $n, m \in \mathbb{N}$, $s \in (3/2, 2)$, $f \in H^s(\mathbb{R})$, and $\varphi \in C^1(\mathbb{R})$ with uniformly continuous derivative φ' be given. Then, there exist a constant K that depends only on $n, m, \|\varphi'\|_\infty$, and $\|f\|_{H^s}$ such that*

$$\|\varphi B_{n, m}^0(f)[h] - B_{n, m}^0(f)[\varphi h]\|_{H^1} \leq K \|h\|_2 \quad (\text{D.1})$$

for all $h \in L_2(\mathbb{R})$.

Proof. See [1, Lemma A.1]. \square

The next four lemmas are used in the proof of Theorem 3.8. We recall the definition of an ε -localization family from Section 3.

Lemma D.2. *Let $n, m \in \mathbb{N}$, $3/2 < s' < s < 2$, and $\nu \in (0, \infty)$ be given. Let further $f \in H^s(\mathbb{R})$ and $\bar{\omega} \in \{1\} \cup H^{s-1}(\mathbb{R})$. For any sufficiently small $\varepsilon \in (0, 1)$, there is a constant $K = K(\varepsilon, n, m, \|f\|_{H^s}, \|\bar{\omega}\|_{H^{s-1}})$ such that*

$$\left\| \pi_j^\varepsilon \bar{\omega} B_{n, m}^0(f)[h] - \frac{\bar{\omega}(x_j^\varepsilon) (f'(x_j^\varepsilon))^n}{[1 + (f'(x_j^\varepsilon))^2]^m} B_{0, 0}[\pi_j^\varepsilon h] \right\|_{H^{s-1}} \leq \nu \|\pi_j^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}}$$

for all $|j| \leq N-1$ and $h \in H^{s-1}(\mathbb{R})$.

Proof. See [1, Lemma A.2]. \square

The next two lemmas are the analogues of Lemma D.2 corresponding to the case $j = N$.

Lemma D.3. *Let $n, m \in \mathbb{N}$, $3/2 < s' < s < 2$, and $\nu \in (0, \infty)$ be given. Let further $f \in H^s(\mathbb{R})$ and $\bar{\omega} \in H^{s-1}(\mathbb{R})$. For any sufficiently small $\varepsilon \in (0, 1)$, there is a constant $K = K(\varepsilon, n, m, \|f\|_{H^s}, \|\bar{\omega}\|_{H^{s-1}})$ such that*

$$\|\pi_N^\varepsilon \bar{\omega} B_{n,m}^0(f)[h]\|_{H^{s-1}} \leq \nu \|\pi_N^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}}$$

for $h \in H^{s-1}(\mathbb{R})$.

Proof. See [1, Lemma A.3]. □

Lemma D.4 is the counterpart of Lemma D.3 in the case when $\bar{\omega} = 1$.

Lemma D.4. *Let $n, m \in \mathbb{N}$, $3/2 < s' < s < 2$, and $\nu \in (0, \infty)$ be given. Let further $f \in H^s(\mathbb{R})$. For any sufficiently small $\varepsilon \in (0, 1)$, there is a constant $K = K(\varepsilon, n, m, \|f\|_{H^s})$ such that*

$$\|\pi_N^\varepsilon B_{0,m}^0(f)[h] - B_{0,0}[\pi_N^\varepsilon h]\|_{H^{s-1}} \leq \nu \|\pi_N^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}}$$

and

$$\|\pi_N^\varepsilon B_{n,m}^0(f)[h]\|_{H^{s-1}} \leq \nu \|\pi_N^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}}, \quad n \geq 1,$$

for all $h \in H^{s-1}(\mathbb{R})$.

Proof. See [1, Lemma A.4]. □

We now prove a lemma which deals with a similar situation as in Lemma D.2.

Lemma D.5. *Let $n, m \in \mathbb{N}$, $3/2 < s' < s < 2$, and $\nu \in (0, \infty)$ be given. Let further $f \in H^s(\mathbb{R})$, $a \in H^{s-1}(\mathbb{R})$, and $\bar{\omega} \in \{1\} \cup H^{s-1}(\mathbb{R})$. For any sufficiently small $\varepsilon \in (0, 1)$, there is a constant K depending on $\varepsilon, n, m, \|f\|_{H^s}, \|a\|_{H^{s-1}}$, and $\|\bar{\omega}\|_{H^{s-1}}$ (if $\bar{\omega} \neq 1$) such that*

$$\left\| \pi_j^\varepsilon \bar{\omega} B_{n,m}^0(f)[ah] - \frac{a(x_j^\varepsilon) \bar{\omega}(x_j^\varepsilon) (f'(x_j^\varepsilon))^n}{[1 + (f'(x_j^\varepsilon))^2]^m} B_{0,0}[\pi_j^\varepsilon h] \right\|_{H^{s-1}} \leq \nu \|\pi_j^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}}$$

for all $|j| \leq N-1$ and $h \in H^{s-1}(\mathbb{R})$.

Proof. Let first $\bar{\omega} \in H^{s-1}(\mathbb{R})$ (the case $\bar{\omega} = 1$ is similar). It is suitable to decompose

$$\pi_j^\varepsilon \bar{\omega} B_{n,m}^0(f)[ah] - \frac{a(x_j^\varepsilon) \bar{\omega}(x_j^\varepsilon) (f'(x_j^\varepsilon))^n}{[1 + (f'(x_j^\varepsilon))^2]^m} B_{0,0}[\pi_j^\varepsilon h] = \bar{\omega}(T_1 + T_2) + a(x_j^\varepsilon) T_3,$$

where

$$T_1 := \pi_j^\varepsilon B_{n,m}^0(f)[(a - a(x_j^\varepsilon))h] - B_{n,m}^0(f)[\pi_j^\varepsilon (a - a(x_j^\varepsilon))h],$$

$$T_2 := B_{n,m}^0(f)[\pi_j^\varepsilon (a - a(x_j^\varepsilon))h],$$

$$T_3 := \pi_j^\varepsilon \bar{\omega} B_{n,m}^0(f)[h] - \frac{\bar{\omega}(x_j^\varepsilon) (f'(x_j^\varepsilon))^n}{[1 + (f'(x_j^\varepsilon))^2]^m} B_{0,0}[\pi_j^\varepsilon h].$$

Lemma D.1 yields

$$\|\bar{\omega} T_1\|_{H^{s-1}} \leq K \|(a - a(x_j^\varepsilon))h\|_2 \leq K \|h\|_2.$$

Besides, using Lemma 3.1 (ii), (3.11), the identity $\chi_j^\varepsilon \pi_j^\varepsilon = \pi_j^\varepsilon$, and the fact that $a \in C^{s-3/2}(\mathbb{R})$, we see that

$$\begin{aligned} \|\bar{\omega} T_2\|_{H^{s-1}} &\leq C \|\pi_j^\varepsilon (a - a(x_j^\varepsilon)) h\|_{H^{s-1}} \leq C \|\chi_j^\varepsilon (a - a(x_j^\varepsilon))\|_\infty \|\pi_j^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}} \\ &\leq \frac{\nu}{2} \|\pi_j^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}} \end{aligned}$$

provided that ε is sufficiently small (where $C = C(n, m, \|\bar{\omega}\|_{H^{s-1}}, \|f\|_{H^s})$). Since T_3 can be estimated by using Lemma D.2, we have established the desired claim. \square

Lemma D.6 below can be seen as the analogue of Lemma D.5 corresponding to the case $j = N$.

Lemma D.6. *Let $n, m \in \mathbb{N}$, $3/2 < s' < s < 2$, and $\nu \in (0, \infty)$ be given. Let further $f \in H^s(\mathbb{R})$, $a \in H^{s-1}(\mathbb{R})$, and $\bar{\omega} \in \{1\} \cup H^{s-1}(\mathbb{R})$. For any sufficiently small $\varepsilon \in (0, 1)$, there is a constant K depending on ε , n , m , $\|f\|_{H^s}$, $\|a\|_{H^{s-1}}$, and $\|\bar{\omega}\|_{H^{s-1}}$ (if $\bar{\omega} \neq 1$) such that*

$$\|\pi_N^\varepsilon \bar{\omega} B_{n,m}^0(f)[ah]\|_{H^{s-1}} \leq \nu \|\pi_N^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}}$$

for all $h \in H^{s-1}(\mathbb{R})$.

Proof. It is suitable to decompose

$$\pi_j^\varepsilon \bar{\omega} B_{n,m}^0(f)[ah] = \bar{\omega}(T_1 + T_2),$$

where

$$\begin{aligned} T_1 &:= \pi_j^\varepsilon B_{n,m}^0(f)[ah] - B_{n,m}^0(f)[\pi_j^\varepsilon ah], \\ T_2 &:= B_{n,m}^0(f)[\pi_j^\varepsilon ah]. \end{aligned}$$

Lemma D.1 yields

$$\|\bar{\omega} T_1\|_{H^{s-1}} \leq K \|ah\|_2 \leq K \|h\|_2.$$

Moreover, appealing to Lemma 3.1 (ii), (3.11), the identity $\chi_j^\varepsilon \pi_j^\varepsilon = \pi_j^\varepsilon$, and the fact that a vanishes at infinity, we find

$$\begin{aligned} \|\bar{\omega} T_2\|_{H^{s-1}} &\leq C \|\pi_j^\varepsilon ah\|_{H^{s-1}} \leq C \|\chi_j^\varepsilon a\|_\infty \|\pi_j^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}} \\ &\leq \nu \|\pi_j^\varepsilon h\|_{H^{s-1}} + K \|h\|_{H^{s'-1}} \end{aligned}$$

provided that ε is sufficiently small. This completes the proof. \square

Acknowledgements: The research leading to this paper was carried out while the second author enjoyed the hospitality of DFG Research Training Group 2339 “Interfaces, Complex Structures, and Singular Limits in Continuum Mechanics - Analysis and Numerics ” at the Faculty of Mathematics of Regensburg University.

REFERENCES

- [1] H. Abels and B.-V. Matioc. Well-posedness of the Muskat problem in subcritical L_p -Sobolev spaces. 2020. arXiv:2003.07656.
- [2] H. Amann. *Linear and Quasilinear Parabolic Problems. Vol. I*, volume 89 of *Monographs in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1995. Abstract linear theory.
- [3] S. B. Angenent. Nonlinear analytic semiflows. *Proc. Roy. Soc. Edinburgh Sect. A*, 115(1-2):91107, 1990.
- [4] A. Badaea and J. Duchon. Capillary driven evolution of an interface between viscous fluids. *Nonlinear Anal.*, 31(3-4):385–403, 1998.

- [5] J. Escher and G. Simonett. Analyticity of the interface in a free boundary problem. *Math. Ann.*, 305(3):439–459, 1996.
- [6] F. Gancedo. A survey for the Muskat problem and a new estimate. *SeMA J.*, 74(1):21–35, 2017.
- [7] Granero-Belinchón, Rafael and Lazar, Omar. Growth in the muskat problem. *Math. Model. Nat. Phenom.*, 15:7, 2020.
- [8] O. A. Ladyzhenskaya. *The mathematical theory of viscous incompressible flow*. Revised English edition. Translated from the Russian by Richard A. Silverman. Gordon and Breach Science Publishers, New York-London, 1963.
- [9] A. Lunardi. *Analytic Semigroups and Optimal Regularity in Parabolic Problems*. Progress in Nonlinear Differential Equations and their Applications, 16. Birkhäuser Verlag, Basel, 1995.
- [10] B.-V. Matioc. Viscous displacement in porous media: the Muskat problem in 2D. *Trans. Amer. Math. Soc.*, 370(10):7511–7556, 2018.
- [11] B.-V. Matioc. The Muskat problem in two dimensions: equivalence of formulations, well-posedness, and regularity results. *Anal. PDE*, 12(2):281–332, 2019.
- [12] J. Prüss, Y. Shao, and G. Simonett. On the regularity of the interface of a thermodynamically consistent two-phase Stefan problem with surface tension. *Interfaces Free Bound.*, 17(4):555–600, 2015.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, DEUTSCHLAND.
E-mail address: `bogdan.matioc@ur.de`

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, TECHNICAL UNIVERSITY EINDHOVEN, THE NETHERLANDS.
E-mail address: `g.prokert@tue.nl`