

Heteroclinic and homoclinic solutions for nonlinear second-order coupled systems with ϕ -Laplacians

Robert de Sousa^(‡) and Feliz Minhós^(†)

^(‡)Faculdade de Ciências e Tecnologia,
Núcleo de Matemática e Aplicações (NUMAT),
Universidade de Cabo Verde. Campus de Palmarejo,
279 Praia, Cabo Verde
robert.sousa@docente.unicv.edu.cv

^(†)Departamento de Matemática, Escola de Ciências e Tecnologia,
Centro de Investigação em Matemática e Aplicações (CIMA),
Instituto de Investigação e Formação Avançada,
Universidade de Évora. Rua Romão Ramalho, 59,
7000-671 Évora, Portugal
fminhos@uevora.pt

Abstract

In this paper we present sufficient conditions for the existence of heteroclinic or homoclinic solutions for second order coupled systems of differential equations on the real line.

We point out that it is required only conditions on the homeomorphisms and no growth or asymptotic conditions are assumed on the nonlinearities.

The arguments make use of the fixed point theory, L^1 -Carathéodory functions and Schauder's fixed point theorem.

An application to a family of second order nonlinear coupled systems of two degrees of freedom, shows the applicability of the main theorem.

2010 Mathematics Subject Classification: 47H10, 34K25, 34B27, 34L30

Keywords: Heteroclinic and homoclinic solutions, coupled systems, L^1 -Carathéodory functions, homeomorphism, Schauder's fixed-point theorem, operator theory.

1 Introduction

In this paper, we consider the second order coupled system on the real line

$$\begin{cases} (a(t)\phi(u'(t)))' = f(t, u(t), v(t), u'(t), v'(t)), \\ (b(t)\psi(v'(t)))' = h(t, u(t), v(t), u'(t), v'(t)), \quad t \in \mathbb{R}, \end{cases} \quad (1)$$

with ϕ and ψ increasing homeomorphisms verifying some adequate relations on their inverses, $a, b : \mathbb{R} \rightarrow (0, +\infty[$ are continuous functions, $f, h : \mathbb{R}^5 \rightarrow \mathbb{R}$ are L^1 -Carathéodory functions, together with asymptotic conditions

$$u(-\infty) = A, \quad u'(+\infty) = 0, \quad v(-\infty) = B, \quad v'(+\infty) = 0 \quad (2)$$

for $A, B \in \mathbb{R}$.

Heteroclinic trajectories play an important role in geometrical analysis of dynamical systems, connecting unstable and stable equilibria having two or more equilibrium points, [12]. In fact, the homoclinic or heteroclinic orbits are a kind of spiral structures, which are general phenomena in nature, [25]. Graphical illustrations and a very complete explanation on homoclinics and heteroclinics bifurcations can be seen in [11]. A planar homoclinic theorem and heteroclinic orbits, to analyze fluid models, is studied in [6]. Applications of dynamic systems techniques to the problem of heteroclinic connections and resonance transitions, are treated in [15], on planar circular domains. To prove the existence of heteroclinic solutions, for a class of non-autonomous second-order equations, see [2, 9, 18]. Topological, variational and minimization methods to find heteroclinic connections can be found in [24].

On heteroclinic coupled systems, among many published works, we highlight some of them:

In [1], Aguiar et al. consider the dynamics of small networks of coupled cells, with one of the points, analyzed as invariant subsets, can support robust heteroclinic attractors;

In [3], Ashwin and Karabacak study coupled phase oscillators and discuss heteroclinic cycles and networks between partially synchronized states and in [14], they analyze coupled phase oscillators, highlighting a dynamic mechanism, nothing more than a heteroclinic network;

In [5], the authors investigate such heteroclinic network between partially synchronized states, where the phases cluster are divided into three groups;

Moreover, in [10], the authors present some applications, results, methods and problems that have been recently reported and, in addition, they

suggest some possible research directions, and some problems for further studies on homoclinics and heteroclinics.

Cabada and Cid, in [7], study the following boundary value problem on the real line

$$\begin{cases} (\phi(u'(t)))' = f(t, u(t), u'(t)), & \text{on } \mathbb{R}, \\ u(-\infty) = -1, \quad u(+\infty) = 1, \end{cases}$$

with a singular ϕ -Laplacian operator where f is a continuous function that satisfies suitable symmetric conditions.

In [8], Calamai discusses the solvability of the following strongly nonlinear problem:

$$\begin{cases} (a(x(t))\phi(x'(t)))' = f(t, x(t), x'(t)), & t \in \mathbb{R}, \\ x(-\infty) = \alpha, \quad x(+\infty) = \beta, \end{cases}$$

where $\alpha < \beta$, $\phi : (-r, r) \rightarrow \mathbb{R}$ is a general increasing homeomorphism with bounded domain (singular ϕ -Laplacian), a is a positive, continuous function and f is a Carathéodory nonlinear function.

Recently, in [13], Kajiwara proved the existence of a heteroclinic solution of the FitzHugh-Nagumo type reaction-diffusion system, under certain conditions on the heterogeneity.

Motivated by these works and applying the techniques suggested in [16, 20, 21, 22], we apply the fixed point theory, to obtain sufficient conditions for the existence of heteroclinic solutions of the coupled system (1), (2), assuming some adequate conditions on ϕ^{-1} , ψ^{-1} .

We emphasize that it is the first time where heteroclinic solutions for second order coupled differential systems are considered for systems with full nonlinearities depending on both unknown functions and their first derivatives. An example illustrates the potentialities of our main result, and an application to coupled nonlinear systems of two degrees of freedom (2-DOF), shows the applicability of the main theorem.

This paper is organized as it follows: Section 2 contains some preliminary results. In section 3 we present the main theorem: an existence result of, at least, a pair of heteroclinic solutions. An application to a family of coupled 2-DOF nonlinear systems is presented in last section.

2 Notations and preliminary results

Consider the following spaces

$$X := \left\{ x \in C^1(\mathbb{R}) : \lim_{|t| \rightarrow \infty} x^{(i)}(t) \in \mathbb{R}, i = 0, 1 \right\},$$

equipped with the norm

$$\|x\|_X = \max \{ \|x\|_\infty, \|x'\|_\infty \},$$

where

$$\|x\|_\infty := \sup_{t \in \mathbb{R}} |x(t)|,$$

and $X^2 := X \times X$ with

$$\|(u, v)\|_{X^2} = \max \{ \|u\|_X, \|v\|_X \}.$$

It can be proved that $(X, \|\cdot\|_X)$ and $(X^2, \|\cdot\|_{X^2})$ are Banach spaces.

Remark 1 *If $w \in X$ then $w'(\pm\infty) = 0$.*

By solution of problem (1), (2) we mean a pair $(u, v) \in X^2$ such that

$$a(t)\phi(u'(t)) \in W^{1,1}(\mathbb{R}) \text{ and } b(t)\psi(v'(t)) \in W^{1,1}(\mathbb{R}),$$

verifying (1), (2).

For the reader's convenience we consider the definition of L^1 -Carathéodory functions:

Definition 2 *A function $g : \mathbb{R}^5 \rightarrow \mathbb{R}$ is L^1 -Carathéodory if*

- i) for each $(x, y, z, w) \in \mathbb{R}^4$, $t \mapsto g(t, x, y, z, w)$ is measurable on \mathbb{R} ;*
- ii) for a.e. $t \in \mathbb{R}$, $(x, y, z, w) \mapsto g(t, x, y, z, w)$ is continuous on \mathbb{R}^4 ;*
- iii) for each $\rho > 0$, there exists a positive function $\vartheta_\rho \in L^1(\mathbb{R})$ such that, whenever $x, y, z, w \in [-\rho, \rho]$, then*

$$|g(t, x, y, z, w)| \leq \vartheta_\rho(t), \text{ a.e. } t \in \mathbb{R}. \quad (3)$$

Along this chapter we assume that

(H1) $\phi, \psi : \mathbb{R} \rightarrow \mathbb{R}$ are increasing homeomorphisms such that

- a)** $\phi(\mathbb{R}) = \mathbb{R}$, $\phi(0) = 0$, $\psi(\mathbb{R}) = \mathbb{R}$, $\psi(0) = 0$;
- b)** $|\phi^{-1}(x)| \leq \phi^{-1}(|x|)$, $|\psi^{-1}(x)| \leq \psi^{-1}(|x|)$.

(H2) $a, b : \mathbb{R} \rightarrow (0, +\infty[$ are positive continuous functions such that

$$\lim_{t \rightarrow \pm\infty} \frac{1}{a(t)} \in \mathbb{R} \text{ and } \lim_{t \rightarrow \pm\infty} \frac{1}{b(t)} \in \mathbb{R}.$$

A convenient criterion for the compacity of the operators is given by next theorem:

Theorem 3 ([20], Theorem 2.3) *A set $M \subset X$ is relatively compact if the following conditions hold:*

- i) *both $\{t \rightarrow x(t) : x \in M\}$ and $\{t \rightarrow x'(t) : x \in M\}$ are uniformly bounded;*
- ii) *both $\{t \rightarrow x(t) : x \in M\}$ and $\{t \rightarrow x'(t) : x \in M\}$ are equicontinuous on any compact interval of \mathbb{R} ;*
- iii) *both $\{t \rightarrow x(t) : x \in M\}$ and $\{t \rightarrow x'(t) : x \in M\}$ are equiconvergent at $\pm\infty$, that is, for any given $\epsilon > 0$, there exists $t_\epsilon > 0$ such that*

$$|f(t) - f(\pm\infty)| < \epsilon, \quad |f'(t) - f'(\pm\infty)| < \epsilon, \quad \forall |t| > t_\epsilon, f \in M.$$

The existence tool will be given by Schauder's fixed point theorem:

Theorem 4 ([23]) *Let Y be a nonempty, closed, bounded and convex subset of a Banach space X , and suppose that $P : Y \rightarrow Y$ is a compact operator. Then P has at least one fixed point in Y .*

3 Existence of heteroclinics

In this section we prove the existence for a pair of heteroclinic solutions to the coupled system (1), (2), for some constants $A, B \in \mathbb{R}$.

Theorem 5 *Let $\phi, \psi : \mathbb{R} \rightarrow \mathbb{R}$ be increasing homeomorphisms and $a, b : \mathbb{R} \rightarrow (0, +\infty[$ continuous functions satisfying (H1) and (H2). Assume that $f, h : \mathbb{R}^5 \rightarrow \mathbb{R}$ are L^1 -Carathéodory functions and there is $R > 0$ and $\vartheta_R, \theta_R \in L^1(\mathbb{R})$ such that*

$$\int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_R(r) dr}{a(s)} \right) ds < +\infty, \quad (4)$$

$$\int_{-\infty}^{+\infty} \psi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \theta_R(r) dr}{b(s)} \right) ds < +\infty, \quad (5)$$

with

$$\sup_{t \in \mathbb{R}} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_R(r) dr}{a(t)} \right) ds < +\infty, \quad \sup_{t \in \mathbb{R}} \psi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \theta_R(r) dr}{b(t)} \right) ds < +\infty,$$

$$|f(t, x, y, z, w)| \leq \vartheta_R(t), \quad (6)$$

$$|h(t, x, y, z, w)| \leq \theta_R(t), \quad (7)$$

whenever $x, y, z, w \in [-R, R]$.

Then for given $A, B \in \mathbb{R}$ the problem (1), (2) has, at least, a pair of heteroclinic solutions $(u, v) \in X^2$.

Proof. Define the operators $T_1 : X^2 \rightarrow X$, $T_2 : X^2 \rightarrow X$ and $T : X^2 \rightarrow X^2$ by

$$T(u, v) = (T_1(u, v), T_2(u, v)), \quad (8)$$

with

$$\begin{aligned} (T_1(u, v))(t) &= \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds + A, \\ (T_2(u, v))(t) &= \int_{-\infty}^t \psi^{-1} \left(\frac{\int_{-\infty}^s h(r, u(r), v(r), u'(r), v'(r)) dr}{b(s)} \right) ds + B, \end{aligned}$$

with A and B given by (2).

In order to apply Theorem 4, we shall prove that T is compact and has a fixed point.

To simplify the proof, we detail the arguments only for $T_1(u, v)$, as for the operator $T_2(u, v)$ the technique is similar.

To be clear, we divide the proof into claims **(i)-(v)**.

(i) T is well defined and continuous in X^2 .

Let $(u, v) \in X^2$ and take $\rho > 0$ such that $\|(u, v)\|_{X^2} < \rho$. As f is a L^1 -Carathéodory function, there exists a positive function $\vartheta_\rho \in L^1(\mathbb{R})$ verifying (6). So,

$$\begin{aligned} & \int_{-\infty}^t |f(r, u(r), v(r), u'(r), v'(r))| dr \\ & \leq \int_{-\infty}^{+\infty} |f(r, u(r), v(r), u'(r), v'(r))| dr \leq \int_{-\infty}^{+\infty} \vartheta_\rho(t) dt < +\infty. \end{aligned}$$

So, T_1 is continuous on X . Furthermore,

$$(T_1(u, v))'(t) = \phi^{-1} \left(\frac{\int_{-\infty}^t f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right)$$

is also continuous on X and, therefore, $T_1(u, v) \in C^1(\mathbb{R})$.

By (2), (4), (6) and (H2),

$$\begin{aligned} & \lim_{t \rightarrow -\infty} T_1(u, v)(t) \\ &= \lim_{t \rightarrow -\infty} \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds + A = A, \end{aligned}$$

$$\begin{aligned} & \lim_{t \rightarrow +\infty} T_1(u, v)(t) \\ &= \int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds + A < +\infty, \end{aligned}$$

and

$$\begin{aligned} \lim_{t \rightarrow \pm\infty} (T_1(u, v)(t))' &= \lim_{t \rightarrow \pm\infty} \phi^{-1} \left(\frac{\int_{-\infty}^t f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right) \\ &\leq \lim_{t \rightarrow \pm\infty} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_\rho(r) dr}{a(t)} \right) < +\infty. \end{aligned}$$

Therefore, $T_1(u, v) \in X$, and, by the same arguments, $T_2(u, v) \in X$. So, $T(u, v) \in X^2$.

(ii) TM is uniformly bounded on $M \subseteq X^2$, for some bounded M .

Let M be a bounded set of X^2 , defined by

$$M := \{(u, v) \in X^2 : \max \{\|u\|_\infty, \|u'\|_\infty, \|v\|_\infty, \|v'\|_\infty\} \leq \rho_1\}, \quad (9)$$

for some $\rho_1 > 0$.

By (4), (6), (H1) and (H2), we have

$$\begin{aligned} & \|T_1(u, v)(t)\|_\infty \\ &= \sup_{t \in \mathbb{R}} \left| \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds + A \right| \\ &\leq \sup_{t \in \mathbb{R}} \int_{-\infty}^t \left| \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) \right| ds + |A| \\ &\leq \sup_{t \in \mathbb{R}} \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^s |f(r, u(r), v(r), u'(r), v'(r))| dr}{a(s)} \right) ds + |A| \\ &\leq \int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_1}(r) dr}{a(s)} \right) ds + |A| < +\infty, \end{aligned}$$

and

$$\begin{aligned}
& \| (T_1(u, v))' (t) \|_\infty \\
&= \sup_{t \in \mathbb{R}} \left| \phi^{-1} \left(\frac{\int_{-\infty}^t f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right) \right| \\
&\leq \sup_{t \in \mathbb{R}} \phi^{-1} \left(\frac{\int_{-\infty}^t |f(r, u(r), v(r), u'(r), v'(r))| dr}{a(t)} \right) \\
&\leq \sup_{t \in \mathbb{R}} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_1}(r) dr}{a(t)} \right) < +\infty.
\end{aligned}$$

So, $\|T_1(u, v)(t)\|_X < +\infty$, that is, T_1M is uniformly bounded on X .

By similar arguments, T_2 is uniformly bounded on X . Therefore TM is uniformly bounded on X^2 .

(iii) TM is equicontinuous on X^2 .

Let $t_1, t_2 \in [-K, K] \subseteq \mathbb{R}$ for some $K > 0$, and suppose, without loss of generality, that $t_1 \leq t_2$. Thus, by (4), (6) and (H1),

$$\begin{aligned}
& |T_1(u, v)(t_1) - T_1(u, v)(t_2)| \\
&= \left| \int_{-\infty}^{t_1} \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds \right. \\
&\quad \left. - \int_{-\infty}^{t_2} \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds \right| \\
&\leq \int_{t_1}^{t_2} \phi^{-1} \left(\frac{\int_{-\infty}^s |f(r, u(r), v(r), u'(r), v'(r))| dr}{a(s)} \right) ds \\
&\leq \int_{t_1}^{t_2} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_1}(r) dr}{a(s)} \right) ds \rightarrow 0,
\end{aligned}$$

uniformly for $(u, v) \in M$, as $t_1 \rightarrow t_2$, and

$$\begin{aligned}
& |(T_1(u, v))'(t_1) - (T_1(u, v))'(t_2)| \\
&= \left| \phi^{-1} \left(\frac{\int_{-\infty}^{t_1} f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t_1)} \right) \right. \\
&\quad \left. - \phi^{-1} \left(\frac{\int_{-\infty}^{t_2} f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t_2)} \right) \right| \rightarrow 0,
\end{aligned}$$

uniformly for $(u, v) \in M$, as $t_1 \rightarrow t_2$.

Therefore, $T_1 M$ is equicontinuous on X . Analogously, it can be proved that $T_2 M$ is equicontinuous on X . So, TM is equicontinuous on X^2 .

(iv) TM is equiconvergent at $t = \pm\infty$.

Let $(u, v) \in M$. For the operator T_1 , we have, by (4), (6) and (H1),

$$\begin{aligned} & \left| T_1(u, v)(t) - \lim_{t \rightarrow -\infty} T_1(u, v)(t) \right| \\ &= \left| \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds \right| \\ &\leq \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_1}(r) dr}{a(s)} \right) ds \rightarrow 0, \end{aligned}$$

uniformly in $(u, v) \in M$, as $t \rightarrow -\infty$, and,

$$\begin{aligned} & \left| T_1(u, v)(t) - \lim_{t \rightarrow +\infty} T_1(u, v)(t) \right| \\ &= \left| \int_{-\infty}^t \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds \right. \\ &\quad \left. - \int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds \right| \\ &= \left| \int_t^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds \right| \\ &\leq \int_t^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_1}(r) dr}{a(s)} \right) ds \rightarrow 0, \end{aligned}$$

uniformly in $(u, v) \in M$, as $t \rightarrow +\infty$.

For the derivative it follows that,

$$\begin{aligned} & \left| (T_1(u, v))'(t) - \lim_{t \rightarrow +\infty} (T_1(u, v))'(t) \right| \\ &= \left| \phi^{-1} \left(\frac{\int_{-\infty}^t f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right) \right. \\ &\quad \left. - \phi^{-1} \left(\lim_{t \rightarrow +\infty} \frac{\int_{-\infty}^{+\infty} f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right) \right| \rightarrow 0, \end{aligned}$$

uniformly in $(u, v) \in M$, as $t \rightarrow +\infty$, and

$$\begin{aligned}
& \left| (T_1(u, v))'(t) - \lim_{t \rightarrow -\infty} (T_1(u, v))'(t) \right| \\
&= \left| \phi^{-1} \left(\frac{\int_{-\infty}^t f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right) \right| \\
&\leq \phi^{-1} \left(\left| \frac{\int_{-\infty}^t f(r, u(r), v(r), u'(r), v'(r)) dr}{a(t)} \right| \right) \\
&\leq \phi^{-1} \left(\frac{\int_{-\infty}^t \vartheta_{\rho_1}(r) dr}{|a(t)|} \right) \rightarrow 0
\end{aligned}$$

uniformly in $(u, v) \in M$, as $t \rightarrow -\infty$.

Therefore, $T_1 M$ is equiconvergent at $\pm\infty$ and, following a similar technique, we can prove that $T_2 M$ is equiconvergent at $\pm\infty$, too. So, TM is equiconvergent at $\pm\infty$.

By Theorem 3, TM is relatively compact.

(v) $T : X \rightarrow X$ has a fixed point.

In order to apply Schauder's fixed point theorem for operator $T(u, v)$, we need to prove that $TD \subset D$, for some closed, bounded and convex $D \subset X^2$.

Consider

$$D := \{(u, v) \in X^2 : \|(u, v)\|_{X^2} \leq \rho_2\},$$

with $\rho_2 > 0$ such that

$$\rho_2 := \max \left\{ \begin{array}{l} \rho_1, \int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_2}(r) dr}{a(s)} \right) ds + |A|, \\ \int_{-\infty}^{+\infty} \psi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \theta_{\rho_2}(r) dr}{b(s)} \right) ds + |B|, \\ \sup_{t \in \mathbb{R}} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \vartheta_{\rho_2}(r) dr}{a(t)} \right), \sup_{t \in \mathbb{R}} \psi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \theta_{\rho_2}(r) dr}{b(t)} \right) \end{array} \right\}$$

with ρ_1 given by (9).

Following similar arguments as in (ii), we have, for $(u, v) \in D$,

$$\begin{aligned}
\|T(u, v)\|_{X^2} &= \|(T_1(u, v), T_2(u, v))\|_{X^2} \\
&= \max \{ \|T_1(u, v)\|_X, \|T_2(u, v)\|_X \} \\
&= \max \left\{ \|T_1(u, v)\|_\infty, \|(T_1(u, v))'\|_\infty, \|T_2(u, v)\|_\infty, \|(T_2(u, v))'\|_\infty \right\} \leq \rho_2,
\end{aligned}$$

and $TD \subset D$.

By Theorem 4, the operator $T(u, v) = (T_1(u, v), T_2(u, v))$ has a fixed point $(u, v) \in X^2$.

By standard arguments, it can be proved that this fixed point defines a pair of heteroclinic or homoclinic solutions of problem (1), (2). ■

Remark 6 *If*

$$\int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^s f(r, u(r), v(r), u'(r), v'(r)) dr}{a(s)} \right) ds = 0$$

and

$$\int_{-\infty}^{+\infty} \psi^{-1} \left(\frac{\int_{-\infty}^s h(r, u(r), v(r), u'(r), v'(r)) dr}{b(s)} \right) ds = 0.$$

the solutions $(u, v) \in X^2$ of problem (1), (2), will be a pair of homoclinic solutions.

4 Application to coupled systems of nonlinear 2-DOF model

Generic nonlinear coupled systems of two degrees of freedom (2-DOF), are especially important in Physics and Mechanics. For example in [19], the authors use this type of system to investigate the transient in a system containing a linear oscillator, linearly coupled to an essentially nonlinear attachment with a comparatively small mass. The family of coupled nonlinear systems of 2-DOF is used to study the global bifurcations in the motion of an externally forced coupled nonlinear oscillatory system or for the nonlinear vibration absorber subjected to periodic excitation, see [17]. Moreover, in [4], the authors deal with the stochastic moment stability of such systems.

Motivated by these works, in this section we consider an application of system (1), (2), to a family of coupled non-linear systems of 2-DOF model, given by the nonlinear coupled system (see [17])

$$\left\{ \begin{array}{l} \left((1+t^4) (q_1'(t))^3 \right)' = \frac{t^4}{(1+t^6)^2} \left[2\zeta\omega_0 (q_1'(t))^3 + \omega_0^2 q_1(t) + \gamma((q_1(t))^3 - 3d^2 q_1(t)q_2(t)) + \cos(t) \right], \\ \tau^2 \left((1+t^4) (q_2'(t))^3 \right)' = \frac{t^4}{(1+t^6)^2} \left[2\zeta\omega_0 (q_2'(t))^3 + \omega_0^2 q_2(t) + \gamma(d^2 (q_2(t))^3 - 3(q_1(t))^2 q_2(t)) \right], \end{array} \right. t \in \mathbb{R}, \quad (10)$$

where

- $q_1(t)$ and $q_2(t)$ represent the generalized coordinates;
- d, τ, γ are positive constant coefficients which depend on the characteristics of the physical or mechanical system under consideration;
- $\cos(t)$ is related to the type of excitation of the system under consideration;
- ζ, ω_0 , are the damping coefficient and the frequency, respectively.

As the asymptotic conditions we consider

$$q_1(-\infty) = A, \quad q_1'(+\infty) = 0, \quad q_2(-\infty) = B, \quad q_2'(+\infty) = 0, \quad (11)$$

with $A, B \in \mathbb{R}$, and, moreover, assume that the real coefficients $\zeta, \omega_0, \gamma, d, r$ are such that the integrals

$$\int_{-\infty}^{+\infty} \left(\sqrt[3]{\frac{\int_{-\infty}^s \frac{r^4}{(1+r^6)^2} \left[\frac{2\zeta\omega_0 (q_1'(r))^3 + \omega_0^2 q_1(r) + \gamma((q_1(r))^3 - 3d^2 q_1(r)q_2(r)) + \cos(r)}{1+s^4} \right] dr}{1+s^4}} \right) ds \quad (12)$$

and

$$\int_{-\infty}^{+\infty} \left(\sqrt[3]{\frac{\int_{-\infty}^s \frac{r^4}{\tau^2(1+r^6)^2} \left[\frac{2\zeta\omega_0 (q_2'(r))^3 + \omega_0^2 q_2(r) + \gamma(d^2 (q_2(r))^3 - 3(q_1(r))^2 q_2(r))}{1+s^4} \right] dr}{1+s^4}} \right) ds. \quad (13)$$

are finite.

It is clear that (10) is a particular case of (1) with:

$$\phi(z) = \psi(z) = z^3, \quad a(t) = b(t) = 1 + t^4,$$

$f, h : \mathbb{R}^5 \rightarrow \mathbb{R}$ are L^1 -Carathéodory functions where

$$\begin{aligned} f(t, x, y, z, w) &= \frac{t^4}{(1+t^6)^2} (2\zeta\omega_0 z^3 + \omega_0^2 x + \gamma x^3 - 3d^2 xy + \cos(t)) \\ &\leq \frac{t^4}{(t^6+1)^2} (2|\zeta\omega_0|\rho^3 + \omega_0^2 \rho + \gamma \rho^3 + 3d^2 \rho^2 + 1) \\ &:= \delta_\rho(t) \end{aligned}$$

$$\begin{aligned} h(t, x, y, z, w) &= \frac{t^4}{\tau^2(1+t^6)^2} (2\zeta\omega_0 w^3 + \omega_0^2 y + \gamma d^2 y^3 - 3x^2 y) \\ &\leq \frac{t^4}{\tau^2(t^6+1)^2} (2|\zeta\omega_0|\rho^3 + \omega_0^2 \rho + \gamma d^2 \rho^3 + 3\rho^3) \\ &:= \varepsilon_\rho(t) \end{aligned}$$

where $\delta_\rho(t)$ and $\varepsilon_\rho(t)$ are functions in $L^1(\mathbb{R})$, for $\rho > 0$ such that

$$\rho := \max \{|x|, |y|, |z|, |w|\}. \quad (14)$$

Moreover, conditions (H1) and (H2) hold as,

- $\phi(\mathbb{R}) = \psi(w) = \mathbb{R}$ and $\phi(0) = \psi(0) = 0$;
- $|\phi^{-1}(z)| = |\sqrt[3]{z}| = \phi^{-1}(|z|) = \sqrt[3]{|z|}$ and $|\psi^{-1}(w)| = |\sqrt[3]{w}| = \psi^{-1}(|w|) = \sqrt[3]{|w|}$;
- $\lim_{t \rightarrow \pm\infty} \frac{1}{a(t)} = \lim_{t \rightarrow \pm\infty} \frac{1}{1+t^4} = \lim_{t \rightarrow \pm\infty} \frac{1}{b(t)} = 0$.

For $\rho > 0$ such that

$$\begin{aligned} &\int_{-\infty}^{+\infty} \phi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \delta_\rho(r) dr}{a(s)} \right) ds \\ &= \int_{-\infty}^{+\infty} \left(\sqrt[3]{\frac{\int_{-\infty}^{+\infty} r^4 \frac{(2|\zeta\omega_0|\rho^3 + \omega_0^2 \rho + \gamma \rho^3 + 3d^2 \rho^2 + 1)}{(1+r^6)^2} dr}{1+s^4}} \right) ds < \rho \end{aligned} \quad (15)$$

and

$$\begin{aligned}
& \int_{-\infty}^{+\infty} \psi^{-1} \left(\frac{\int_{-\infty}^{+\infty} \varepsilon_{\rho}(r) dr}{b(s)} \right) ds \\
&= \int_{-\infty}^{+\infty} \left(\sqrt[3]{\frac{\int_{-\infty}^{+\infty} r^4 \frac{(2|\zeta\omega_0|\rho^3 + \omega_0^2\rho + \gamma d^2\rho^3 + 3\rho^3)}{\tau^2(1+r^6)^2} dr}{1+s^4}} \right) ds < \rho, \quad (16)
\end{aligned}$$

by Theorem 5, the system (10) together with the asymptotic conditions (11), has at least a pair $(q_1, q_2) \in X^2$ of heteroclinic solutions, since the integrals (12) and (13) are finite. As example, in particular, for

$$|\zeta| = \frac{1}{2\sqrt{1000}}, \quad |\omega_0| = \frac{1}{\sqrt{1000}}, \quad \gamma = \frac{1}{1000}, \quad d^2 = \frac{1}{3000}, \quad \tau = 23$$

the conditions (15) and (16) hold for $\rho > 6.3542$.

For the values of the above parameters, $A = 10$ and $B = 8$, the heteroclinic solutions q_1 and q_2 have the graphs given in Figure 1. In Figure 2 we present the real shape of the q_1 trajectory, which is not detailed in Figure 1 due to the scale range.




Figure 1: $A=10$, $B=8$

Remark that, if the integrals (12) and (13) are null, then the system (10) has a pair of homoclinic solutions $(q_1, q_2) \in X^2$.




Figure 2: $A=10$

References

- [1] M. Aguiar, P. Ashwin, A. Dias, M. Field, *Dynamics of Coupled Cell Networks: Synchrony, Heteroclinic Cycles and Inflation*, Journal of Nonlinear Science 21 (2), (2011) pp. 271–323.
- [2] C. Alves, *Advances in Nonlinear Dynamics: Methods and Applications*, Nonlinear Differ. Equ. Appl. 22 (5) (2015) 1195–1212.
- [3] P. Ashwin, Ö. Karabacak, *Robust Heteroclinic Behaviour, Synchronization, and Ratcheting of Coupled Oscillators*, Dynamics, Games and Science II, (2011) pp 125–140.
- [4] S. Ariaratnam, D. Tam and W-C. Xie, *Lyapunov Exponents of Two-Degrees-of-Freedom Linear Stochastic Systems*, Stochastic Structural Dynamics 1, New Theoretical Developments, Springer-Verlag Berlin Heidelberg 1991.
- [5] P. Ashwin, G. Orosz, J. Wordsworth, S. Townley, *Dynamics on Networks of Cluster States for Globally Coupled Phase Oscillators*, SIAM J. Applied Dynamical Systems, Vol. 6, No. 4, pp. 728–758.
- [6] A. Bertozzi, *Heteroclinic orbits and chaotic dynamics in planar fluid flows*, Siam J. Math. Anal. Vol. 19, No. 6, November 1988.

- [7] A. Cabada, J. Cid, *Heteroclinic solutions for non-autonomous boundary value problems with singular Φ -Laplacian operators*, Discret and Continuous Dynamical Systems Supplement 2009, 118–122.
- [8] A. Calamai, *Heteroclinic solutions of boundary value problems on the real line involving singular Φ -Laplacian operators*, J. Math. Anal. Appl. 378 (2011) 667–679.
- [9] E. Ellero, F. Zanolin, *Homoclinic and heteroclinic solutions for a class of second-order non-autonomous ordinary differential equations: multiplicity results for stepwise potentials*, Boundary Value Problems (2013) 2013: 167. <https://doi.org/10.1186/1687-2770-2013-167>.
- [10] B. Feng, R. Hu, *A Survey On Homoclinic And Heteroclinic Orbits*, Applied Mathematics E-Notes, 3(2003), 16–37.
- [11] A. Homburg, B. Sandstede, *Homoclinic and Heteroclinic Bifurcations in Vector Fields*, Handbook of Dynamical Systems Volume 3, 2010, Pages 379–524, doi: 10.1016/S1874-575X(10)00316-4.
- [12] E. Izhikevich, *Dynamical Systems in Neuroscience: The Geometry of Excitability and Bursting*, The MIT Press, Computational Neuroscience, Cambridge (2007).
- [13] T. Kajiwara, *A Heteroclinic Solution to a Variational Problem Corresponding to FitzHugh-Nagumo type Reaction-Diffusion System with Heterogeneity*, Communications on Pure and Applied Analysis Volume 16, Number 6, November (2017) 2133–2156.
- [14] Ö. Karabacak, P. Ashwin, *Heteroclinic Ratchets in Networks of Coupled Oscillators*, J. Nonlinear Science, 20 (2010) 105–129.
- [15] W. Koon, M. Lo, J. Marsden, S. Ross, *Heteroclinic orbits and chaotic dynamics in planar fluid flows*, Chaos 10(2) (2000) 427–469.
- [16] Y. Liu, S. Chen *Existence of bounded solutions of integral boundary value problems for singular differential equations on whole lines*, International Journal of Mathematics, 25(8) (2014) 1450078 (28 pages).
- [17] N. Malhotra, N. Sri Namachchivaya, *Global Bifurcations in Externally Excited Two-Degree-of-Freedom Nonlinear Systems*, Nonlinear Dynamics, Volume 8, No. 1, (1995) 85–109.

- [18] C. Marcelli, F. Papalini, *Heteroclinic connections for fully non-linear non-autonomous second-order differential equations*, J. Differential Equations 241 (2007) 160–183.
- [19] Y. Mikhlin, T. Bunakova, G. Rudneva, N. Perepelkin, *Transient in 2-DOF Nonlinear Systems*, ENOC 2008, Saint Petersburg, Russia, June, 30-July, 4 2008.
- [20] F. Minhós, H. Carrasco, *Existence of homoclinic solutions for nonlinear second-order problems*, Mediterr. J. Math. 13 (2016) 3849–3861, doi:10.1007/s00009-016-0718-4.
- [21] F. Minhós, *Sufficient conditions for the existence of heteroclinic solutions for φ -Laplacian differential equations*, Complex Variables and Elliptic Equations, volume 62, Issue 1, (2017) 123–134.
- [22] F. Minhós, R. de Sousa, *Existence of homoclinic solutions for nonlinear second-order coupled systems*, J. Differential Equations (2018), <https://doi.org/10.1016/j.jde.2018.07.072>.
- [23] E. Zeidler, *Nonlinear Functional Analysis and Its Applications: Fixed-Point Theorems*, Springer, New York (1986).
- [24] V. Zelati, P. Rabinowitz, *Heteroclinic solutions between stationary points at different energy levels*, Topological Methods in Nonlinear Analysis Journal of the Juliusz Schauder Center Volume 17, 2001, 1–21.
- [25] F. Zun-Tao, L. Shi-Da, L. Shi-Kuo, L. Fu-Ming, X. Guo-Jun, *Homoclinic (Heteroclinic) Orbit of Complex Dynamical System and Spiral Structure*, Commun. Theor. Phys. (Beijing, China) 43 (4), (2005) pp. 601–603.