

# Local Langlands correspondences in $\ell$ -adic coefficients

Naoki Imai

## Abstract

Let  $\ell$  be a prime number different from the residue characteristic of a non-archimedean local field  $F$ . We give formulations of  $\ell$ -adic local Langlands correspondences for connected reductive algebraic groups over  $F$ , which should be independent of a choice of an isomorphism between the  $\ell$ -adic coefficient field and the complex number field.

## Introduction

The local Langlands correspondence for a connected reductive algebraic group over a non-archimedean local field  $F$  is usually formulated with coefficients in  $\mathbb{C}$  because of its relation with automorphic representations. On the other hand, when we discuss a realization of the local Langlands correspondence in  $\ell$ -adic cohomology, we need a correspondence over  $\overline{\mathbb{Q}}_\ell$ , where  $\ell$  is a prime number different from the residue characteristic of  $F$ . We can take an isomorphism  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$  and use it to transfer the local Langlands correspondence over  $\mathbb{C}$  to a local Langlands correspondence over  $\overline{\mathbb{Q}}_\ell$ . However, the obtained correspondence over  $\overline{\mathbb{Q}}_\ell$  depends on the choice of  $\iota$ . In [BH06], Bushnell–Henniart formulated an  $\ell$ -adic local Langlands correspondence for  $GL_2$ , which is independent of a choice of an isomorphism  $\mathbb{C} \rightarrow \overline{\mathbb{Q}}_\ell$  by making some twists of L-parameters. The  $\ell$ -adic local Langlands correspondence is suitable to describe the non-abelian Lubin–Tate theory in the sense that it is canonical (*cf.* [IT, 5]).

In this paper, we discuss formulations of  $\ell$ -adic local Langlands correspondences for general connected reductive groups. A natural idea is to make similar twists as  $GL_2$ -case for L-parameters. However, we can not make such twists in general as explained in Example 2.1. A problem is that we do not have enough space inside the Langlands dual group to make an appropriate twist. To overcome this problem, we have two approaches. One is to introduce  $\ell$ -adic C-parameters using C-groups, which make spaces for twists. Another is to introduce Tannakian  $\ell$ -adic L-parameters that incorporate necessary twists in the realizations. Using these parameters, we formulate  $\ell$ -adic local Langlands correspondences, which we conjecture to be independent of a choice of an isomorphism  $\mathbb{C} \rightarrow \overline{\mathbb{Q}}_\ell$ . We show that two conjectures formulated by using  $\ell$ -adic C-parameters and Tannakian  $\ell$ -adic L-parameters are equivalent. Further, we confirm that the conjectures are true in  $GL_n$ -case. The formulation of the  $\ell$ -adic local Langlands correspondence using Tannakian  $\ell$ -adic L-parameters is motivated by the Kottwitz conjecture for local Shimura varieties in [RV14, Conjecture 7.4]. In number field cases, a relation between C-groups and the Kottwitz conjecture for Shimura varieties is discussed in [Joh13].

---

2010 *Mathematics Subject Classification*. Primary: 11F70; Secondary: 11F80.

In Section 1, we recall various versions of L-parameters and explain their relations. In Section 2, after explaining the problem, we give formulations of the  $\ell$ -adic local Langlands correspondences introducing the  $\ell$ -adic C-parameter and the Tannakian  $\ell$ -adic L-parameter.

## Acknowledgements

The author would like to thank Teruhisa Koshikawa for his helpful comment.

## Notation

For a field  $F$ , let  $\Gamma_F$  denote the absolute Galois group of  $F$ . For a non-archimedean local field  $F$ , let  $W_F$  and  $I_F$  denote the Weil group of  $F$  and its inertia subgroup. For a homomorphism  $\phi: G \rightarrow H$  of groups and  $g \in G$ , let  $\phi^g$  denote the homomorphism defined by  $\phi^g(g') = \phi(gg'g^{-1})$  for  $g' \in G$ . For a group  $G$ , let  $Z(G)$  denote the center of  $G$ . For a group  $G$ , a subgroup  $H \subset G$  and a subset  $S \subset G$ , let  $Z_H(S)$  denote the centralizer of  $S$  in  $H$ .

# 1 Langlands parameter

## 1.1 Over algebraically closed field

Let  $F$  be a non-archimedean local field of residue characteristic  $p$ . Let  $q$  be the number of elements of the residue field of  $F$ . Let  $v_F: F^\times \rightarrow \mathbb{Z}$  be the normalized valuation of  $F$ . Let

$$\text{Art}_F: F^\times \xrightarrow{\sim} W_F^{\text{ab}}$$

be the Artin reciprocity isomorphism normalized so that a uniformizer is sent to a lift of the geometric Frobenius element. For  $w \in W_F$ , we put

$$d_F(w) = v_F(\text{Art}_F^{-1}(\bar{w})), \quad |w| = q^{-d_F(w)},$$

where  $\bar{w}$  denotes the image of  $w$  in  $W_F^{\text{ab}}$ .

Let  $G$  be a connected reductive algebraic group over  $F$ . Let  $C$  be an algebraically closed field of characteristic 0. Let  $\widehat{G}$  be the dual group of  $G$  over  $C$ . Let  ${}^L G = \widehat{G} \rtimes W_F$  be the L-group of  $G$ . Let  $p_{\widehat{G}}: {}^L G \rightarrow \widehat{G}$  denote the projection map.

**Definition 1.1.** *Let  $H$  be a group with action of  $W_F$ . Let  $\phi: H \rtimes W_F \rightarrow {}^L G(C)$  be a homomorphism of groups over  $W_F$ .*

- (1) *We say that  $\phi$  is semisimple if all the elements of  $\phi(W_F)$  are semisimple.*
- (2) *We say that  $\phi$  is relevant if any parabolic subgroup  $P$  of  ${}^L G$  containing  $\text{Im } \phi$  is relevant in the sense of [Bor79, 3.3].*

*We say that two homomorphisms  $H \rtimes W_F \rightarrow {}^L G(C)$  of groups over  $W_F$  are equivalent if they are conjugate by an element of  $\widehat{G}(C)$ .*

Let  $WD_F = \mathbb{G}_a \rtimes W_F$  be the Weil–Deligne group scheme over  $\mathbb{Q}$  for  $F$  defined in [Del73, 8.3.6].

**Definition 1.2** (cf. [Bor79, 8.2]). (1) An  $L$ -homomorphism of Weil–Deligne type for  $G$  over  $C$  is a homomorphism

$$\xi: WD_F(C) \rightarrow {}^L G(C)$$

of groups over  $W_F$  such that  $\xi|_{\mathbb{G}_a(C)}$  is algebraic and  $(p_{\widehat{G}} \circ \xi)|_{W_F}$  is smooth.

(2) An  $L$ -parameter of Weil–Deligne type for  $G$  over  $C$  is a semisimple, relevant  $L$ -homomorphism of Weil–Deligne type for  $G$  over  $C$ .

Let  $\mathcal{L}_C^{\text{WD}}(G)$  denote the set of equivalence classes of  $L$ -homomorphisms of Weil–Deligne type for  $G$  over  $C$ . Let  $\Phi_C^{\text{WD}}(G)$  denote the set of equivalence classes of  $L$ -parameters of Weil–Deligne type for  $G$  over  $C$ .

**Definition 1.3.** (1) A Weil–Deligne  $L$ -homomorphism for  $G$  over  $C$  is a pair  $(\tau, N)$  of a homomorphism  $\tau: W_F \rightarrow {}^L G(C)$  of groups over  $W_F$  and  $N \in \text{Lie}(\widehat{G})(C)$  such that  $p_{\widehat{G}} \circ \tau$  is smooth and  $\text{Ad}(\tau(w))N = |w|N$  for  $w \in W_F$ . The second component  $N$  of a Weil–Deligne  $L$ -homomorphism  $(\tau, N)$  is called a monodromy operator. We say that two Weil–Deligne  $L$ -homomorphisms for  $G$  over  $C$  are equivalent if they are conjugate by an element of  $\widehat{G}(C)$ .

(2) A Weil–Deligne  $L$ -parameter for  $G$  over  $C$  is a Weil–Deligne  $L$ -homomorphism  $(\tau, N)$  for  $G$  over  $C$  such that  $\tau$  is semisimple and any parabolic subgroup  $P$  of  ${}^L G$  containing  $\tau(W_F)$  and satisfying  $N \in \text{Lie}(P \cap \widehat{G})(C)$  is relevant.

Let  $\mathcal{L}_C^{\text{M}}(G)$  denote the set of equivalence classes of Weil–Deligne  $L$ -homomorphisms for  $G$  over  $C$ . Let  $\Phi_C^{\text{M}}(G)$  denote the set of equivalence classes of Weil–Deligne  $L$ -parameters for  $G$  over  $C$ .

**Remark 1.4.** Let  $\iota: C \simeq C'$  be an isomorphism of algebraically closed fields of characteristic 0. Then  $\iota$  induces bijections  $\mathcal{L}_C^{\text{WD}}(G) \simeq \mathcal{L}_{C'}^{\text{WD}}(G)$ ,  $\Phi_C^{\text{WD}}(G) \simeq \Phi_{C'}^{\text{WD}}(G)$ ,  $\mathcal{L}_C^{\text{M}}(G) \simeq \mathcal{L}_{C'}^{\text{M}}(G)$  and  $\Phi_C^{\text{M}}(G) \simeq \Phi_{C'}^{\text{M}}(G)$ .

**Lemma 1.5.** Let  $(\tau, N)$  be a Weil–Deligne  $L$ -homomorphism for  $G$  over  $C$ . Then  $N$  is a nilpotent element of  $\text{Lie}(\widehat{G}^{\text{der}}(C))$ .

*Proof.* We take a finite separable extension  $F'$  of  $F$  that splits  $G$ . By the condition  $\text{Ad}(\tau(w))N = |w|N$  for  $w \in W_{F'}$ , we have  $N \in \text{Lie}(\widehat{G}^{\text{der}}(C))$ . Further, the adjoint endomorphism  $\text{ad}(N) \in \text{End}(\text{Lie}(\widehat{G}^{\text{der}}(C)))$  satisfies

$$\text{Ad}(\tau(w)) \circ \text{ad}(N) = |w| \text{ad}(N) \circ \text{Ad}(\tau(w))$$

for  $w \in W_{F'}$ . Hence  $N$  is a nilpotent element.  $\square$

**Proposition 1.6.** For an  $L$ -homomorphism of Weil–Deligne type  $\xi$  for  $G$  over  $C$ , we put  $\tau_\xi = \xi|_{W_F}$  and  $N_\xi = \text{Lie}(\xi|_{\mathbb{G}_a(C)})(1)$ . Then  $\xi \mapsto (\tau_\xi, N_\xi)$  induces bijections  $\mathcal{L}_C^{\text{WD}}(G) \simeq \mathcal{L}_C^{\text{M}}(G)$  and  $\Phi_C^{\text{WD}}(G) \simeq \Phi_C^{\text{M}}(G)$ .

*Proof.* Let  $(\tau, N)$  be a Weil–Deligne  $L$ -homomorphism for  $G$  over  $C$ . By Lemma 1.5, there is a unipotent radical  $U$  of a Borel subgroup of  $\widehat{G}$  such that  $N \in \text{Lie}(U(C))$ . Then there is an exponential map

$$\exp: \text{Lie}(U(C)) \rightarrow U(C)$$

as in [Ser96, 2.2]. We define

$$\xi_{(\tau, N)}: WD_F(C) \rightarrow {}^L G(C)$$

by  $\xi_{(\tau, N)}(a, w) = \exp(aN)\tau(w)$ . Then  $(\tau, N) \mapsto \xi_{(\tau, N)}$  induces the inverses.  $\square$

## 1.2 Over $\mathbb{C}$

Assume that  $C = \mathbb{C}$  in this subsection.

**Definition 1.7** (cf. [AG91, I.2]). *An L-parameter for  $G$  is a semisimple, relevant continuous homomorphism*

$$\phi: SU_2(\mathbb{R}) \times W_F \rightarrow {}^L G(\mathbb{C})$$

of groups over  $W_F$ .

Let  $\Phi(G)$  denote the set of equivalence classes of L-parameters for  $G$ .

**Definition 1.8** (cf. [Lan83, IV.2]). *An L-parameter of  $SL_2$ -type for  $G$  is a semisimple, relevant continuous homomorphism*

$$\phi: SL_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$$

of groups over  $W_F$  such that  $\phi|_{SL_2(\mathbb{C})}$  is algebraic.

Let  $\Phi^{\text{SL}}(G)$  denote the set of equivalence classes of L-parameters of  $SL_2$ -type for  $G$ . By the restriction with respect to  $SU_2(\mathbb{R}) \subset SL_2(\mathbb{C})$ , we have a bijection  $\Phi^{\text{SL}}(G) \simeq \Phi(G)$ .

**Lemma 1.9.** *Let  $H$  be a compact topological group. Let  $\phi: H \rightarrow {}^L G(\mathbb{C})$  be a continuous homomorphism. Then the centralizer of  $\phi(H)$  in  $\widehat{G}$  is reductive.*

*Proof.* Let  $A$  be the Zariski closure of the image of  $\phi(H)$  under  $\text{Ad}: {}^L G \rightarrow \text{Aut}(\widehat{G})$ . Then  $A$  is reductive since  $H$  is compact. Hence the claim follows from [Kot84, 10.1.2 Lemma].  $\square$

The following is a slight generalization of [Hei06, Proposition 3.5] and [KL87, 2.4].

**Lemma 1.10.** *Let  $\mathcal{G}$  be a reductive group over  $\mathbb{C}$ . Let  $s$  be a semisimple element of  $\mathcal{G}(\mathbb{C})$  and  $u$  be a unipotent element of  $\mathcal{G}(\mathbb{C})$  such that  $sus^{-1} = u^q$ . Then there is an algebraic homomorphism  $\theta: SL_2(\mathbb{C}) \rightarrow \mathcal{G}(\mathbb{C})$  such that*

$$\theta \left( \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) = u, \quad \theta \left( \begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right) s^{-1} \in Z_{\mathcal{G}(\mathbb{C})}(\text{Im } \theta).$$

Further, such  $\theta$  is unique up to conjugation by  $Z_{\mathcal{G}(\mathbb{C})}(\{s, u\})$ .

*Proof.* This is proved in the same way as [Hei06, Proposition 3.5] and [KL87, 2.4 (h)].  $\square$

Let  $\phi: SL_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$  be an L-parameter of  $SL_2$ -type for  $G$ . We define  $\xi_\phi: WD_F(\mathbb{C}) \rightarrow {}^L G(\mathbb{C})$  by

$$\xi_\phi(a, w) = \phi \left( \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{-\frac{1}{2}} \end{pmatrix}, w \right).$$

Define  $\Xi: \Phi^{\text{SL}}(G) \rightarrow \Phi_{\mathbb{C}}^{\text{WD}}(G)$  by  $\Xi([\phi]) = [\xi_\phi]$ .

**Proposition 1.11.** *The map  $\Xi$  is a bijection.*

*Proof.* Let  $\xi$  be a Weil–Deligne L-parameter for  $G$  over  $\mathbb{C}$ . Let  $\mathcal{H}$  be the centralizer of  $\xi(I_F)$  in  $\widehat{G}$ . Then  $\mathcal{H}$  is reductive by Lemma 1.9. Take a lift  $\sigma_q \in W_F$  of the  $q$ -th power Frobenius element. Then  $\mathcal{H}$  is stable under conjugation by  $\xi(\sigma_q)$ , since  $I_F$  is a normal subgroup of  $W_F$ . Take a positive integer  $m$  such that  $\xi(\sigma_q)^m$  commutes with  $\xi(I_F)$  and  $\sigma_q^m$  commutes with  $\widehat{G}$  in  ${}^L G$ . Then  $(1, \sigma_q^m)^{\mathbb{Z}}$  is a normal subgroup of  $\mathcal{H} \cdot \xi(\sigma_q)^{\mathbb{Z}} \subset {}^L G$ . We put

$$\mathcal{G} = (\mathcal{H} \cdot \xi(\sigma_q)^{\mathbb{Z}}) / (1, \sigma_q^m)^{\mathbb{Z}}.$$

Then  $\mathcal{G}$  is a reductive group. We view  $\mathcal{H}$  as an algebraic subgroup of  $\mathcal{G}(\mathbb{C})$ . Let  $s$  be the image of  $\xi(\sigma_q)$  in  $\mathcal{G}$ . Since  $\xi$  is semisimple,  $s$  is semisimple. Let  $u \in \widehat{G}(\mathbb{C})$  be the image of  $1 \in \mathbb{G}_a(\mathbb{C})$  under  $\xi$ . Then  $u$  belongs to  $\mathcal{H}(\mathbb{C})$ . Hence we can view  $u$  as an element of  $\mathcal{G}(\mathbb{C})$ . By Lemma 1.10, there is a morphism  $\theta: SL_2(\mathbb{C}) \rightarrow \mathcal{G}(\mathbb{C})$  such that

$$\theta \left( \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) = u, \quad \theta \left( \begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right) s^{-1} \in Z_{\mathcal{G}(\mathbb{C})}(\text{Im } \theta).$$

Since  $\theta$  factors through the identity connected component of  $\mathcal{G}(\mathbb{C})$ , it factors through  $\mathcal{H}(\mathbb{C})$ . Hence  $\theta$  determines a morphism  $\theta_{\widehat{G}}: SL_2(\mathbb{C}) \rightarrow \widehat{G}(\mathbb{C})$ . We define  $\phi_{\xi}: SL_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$  by  $\phi_{\xi}|_{SL_2(\mathbb{C})} = \theta_{\widehat{G}}$  and

$$\phi_{\xi}(1, w) = \theta_{\widehat{G}} \left( \begin{pmatrix} |w|^{-\frac{1}{2}} & 0 \\ 0 & |w|^{\frac{1}{2}} \end{pmatrix} \right) \xi(0, w).$$

Then  $\xi \mapsto \phi_{\xi}$  induces the inverse of  $\Xi$ . □

### 1.3 Over $\overline{\mathbb{Q}}_{\ell}$

Let  $\ell$  be a prime number different from  $p$ . Assume that  $C = \overline{\mathbb{Q}}_{\ell}$  in this subsection.

**Definition 1.12.** (1) An  $\ell$ -adic L-homomorphism for  $G$  is a continuous homomorphism

$$\varphi: W_F \rightarrow {}^L G(\overline{\mathbb{Q}}_{\ell})$$

of groups over  $W_F$ . We say that an  $\ell$ -adic L-homomorphism  $\varphi$  for  $G$  is Frobenius-semisimple if  $\varphi(\sigma_q)$  is semisimple for any lift  $\sigma_q \in W_F$  of the  $q$ -th power Frobenius element.

(2) An  $\ell$ -adic L-parameter for  $G$  is an Frobenius-semisimple, relevant L-homomorphisms for  $G$ .

Let  $\mathcal{L}_{\ell}(G)$  denote the set of the equivalence class of  $\ell$ -adic L-homomorphisms. Let  $\Phi_{\ell}(G)$  denote the set of the equivalence class of  $\ell$ -adic L-parameters of  $G$ .

Let  $t_{\ell}: I_F \rightarrow \mathbb{Z}_{\ell}(1)$  be the  $\ell$ -adic tame character. We take an isomorphism  $\mathbb{Z}_{\ell}(1) \simeq \mathbb{Z}_{\ell}$  and let

$$t'_{\ell}: I_F \xrightarrow{t_{\ell}} \mathbb{Z}_{\ell}(1) \simeq \mathbb{Z}_{\ell}$$

Let  $\xi$  be an L-homomorphism of Weil–Deligne type for  $G$  over  $\overline{\mathbb{Q}}_{\ell}$ . We take a lift  $\sigma_q \in W_F$  of the  $q$ -th power Frobenius element and define  $\varphi_{\xi}: W_F \rightarrow {}^L G(\overline{\mathbb{Q}}_{\ell})$  by

$$\varphi_{\xi}(\sigma_q^m \sigma) = \xi(t'_{\ell}(\sigma), \sigma_q^m \sigma)$$

for  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ .

Define  $\Theta: \mathcal{L}_{\overline{\mathbb{Q}}_{\ell}}^{\text{WD}}(G) \rightarrow \mathcal{L}_{\ell}(G)$  by  $\Theta([\xi]) = [\varphi_{\xi}]$ .

**Proposition 1.13.** *The map  $\Theta$  is a bijection. Further it induces a bijection  $\Phi_{\overline{\mathbb{Q}}_\ell}^{\text{WD}}(G) \rightarrow \Phi_\ell(G)$*

*Proof.* Let  $\varphi$  be an  $\ell$ -adic L-homomorphism for  $G$ . Take a finite Galois extension  $F'$  of  $F$  such that  $G$  splits over  $F'$ . Take a representation

$$\eta: {}^L G(\overline{\mathbb{Q}}_\ell) \rightarrow GL_n(\overline{\mathbb{Q}}_\ell)$$

which factors through a faithful algebraic representation

$$\overline{\eta}: \widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \text{Gal}(F'/F) \rightarrow GL_n(\overline{\mathbb{Q}}_\ell).$$

Applying Grothendieck's monodromy theorem (*cf.* [ST68, Appendix, Proposition]) to  $\eta \circ \varphi$ , we obtain a homomorphism

$$\xi_{GL_n}: WD_F(\overline{\mathbb{Q}}_\ell) \rightarrow GL_n(\overline{\mathbb{Q}}_\ell)$$

such that  $\xi_{GL_n}|_{\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)}$  is algebraic,  $\xi_{GL_n}|_{W_F}$  is smooth and

$$\xi_{GL_n}(t_\ell(\sigma), \sigma_q^m \sigma) = (\eta \circ \varphi)(\sigma_q^m \sigma)$$

for  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ . Take a finite separable extension  $F''$  of  $F'$  such that  $\xi_{GL_n}|_{I_{F''}}$  is trivial. Since  $t'_\ell(I_{F''})$  is Zariski dense in  $\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)$ , the algebraic morphism  $\xi_{GL_n}|_{\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)}$  factors through the inclusion

$$\widehat{G}(\overline{\mathbb{Q}}_\ell) \hookrightarrow \widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \text{Gal}(F'/F) \xrightarrow{\overline{\eta}} GL_n(\overline{\mathbb{Q}}_\ell)$$

via an algebraic morphism  $\alpha: \mathbb{G}_a(\overline{\mathbb{Q}}_\ell) \rightarrow \widehat{G}(\overline{\mathbb{Q}}_\ell)$ . We define a homomorphism

$$\xi_\varphi: WD_F(\overline{\mathbb{Q}}_\ell) \rightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$$

by

$$\xi_\varphi(a, \sigma_q^m \sigma) = \alpha(a - t'_\ell(\sigma))\varphi(\sigma_q^m \sigma)$$

for  $a \in \mathbb{G}_a(\overline{\mathbb{Q}}_\ell)$ ,  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ . Then  $\varphi \mapsto \xi_\varphi$  induces the inverses.  $\square$

## 2 Local Langlands correspondence

### 2.1 Problem

Let  $\text{Irr}(G(F))$  denote the set of isomorphism classes of irreducible smooth representations of  $G(F)$  over  $\mathbb{C}$ . The Langlands conjecture says that there is a surjective map

$$\text{LL}_G: \text{Irr}(G(F)) \rightarrow \Phi(G)$$

with finite fibers satisfying various properties (*cf.* [Bor79, 10], [Kal16, Conjecture G]). Assume the above conjecture in the sequel.

Let  $\text{Irr}_\ell(G(F))$  denote the set of isomorphism classes of irreducible smooth representations of  $G(F)$  over  $\overline{\mathbb{Q}}_\ell$ . If we fix an isomorphism  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$ , we have a surjection

$$\text{LL}_{G,\ell}^\iota: \text{Irr}_\ell(G(F)) \rightarrow \Phi_\ell(G)$$

sending  $[\pi] \in \text{Irr}_\ell(G(F))$  to the image of  $\text{LL}_G([\pi \otimes_{\overline{\mathbb{Q}}_\ell, \iota^{-1}} \mathbb{C}])$  under the bijection

$$\Phi(G) \simeq \Phi^{\text{SL}}(G) \simeq \Phi_{\mathbb{C}}^{\text{WD}}(G) \xrightarrow{\Phi_\ell^{\text{WD}}(G)} \Phi_{\overline{\mathbb{Q}}_\ell}^{\text{WD}}(G) \simeq \Phi_\ell(G),$$

where  $\Phi_\ell^{\text{WD}}(G)$  is a bijection induced by  $\iota$  as in Remark 1.4. However,  $\text{LL}_{G,\ell}^\iota$  depends on the choice of an isomorphism  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$ .

In [BH06, 35.1], an  $\ell$ -adic local Langlands correspondence for  $GL_2$  is constructed. We recall the construction here. For  $[\phi] \in \Phi(GL_2)$ , we define  $[\tilde{\phi}] \in \Phi(GL_2)$  by

$$\tilde{\phi}(g, w) = \left( \left( \begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{\frac{1}{2}} \end{pmatrix}, 1 \right), 1 \right) \phi(g, w).$$

We define a bijection  $\gamma: \Phi(GL_2) \rightarrow \Phi(GL_2)$  by  $[\phi] \mapsto [\tilde{\phi}]$ . We define

$$\text{LL}_{GL_2,\ell}: \text{Irr}_\ell(GL_2(F)) \rightarrow \Phi_\ell(GL_2)$$

by sending  $[\pi] \in \text{Irr}_\ell(GL_2(F))$  to the image of  $\text{LL}_{GL_2}([\pi \otimes_{\overline{\mathbb{Q}}_\ell, \iota^{-1}} \mathbb{C}])$  under the bijection

$$\Phi(GL_2) \xrightarrow{\tilde{\gamma}} \Phi(GL_2) \simeq \Phi^{\text{SL}}(GL_2) \simeq \Phi_{\mathbb{C}}^{\text{WD}}(GL_2) \xrightarrow{\Phi_\ell^{\text{WD}}(G)} \Phi_{\overline{\mathbb{Q}}_\ell}^{\text{WD}}(GL_2) \simeq \Phi_\ell(GL_2)$$

using an isomorphism  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$ . Then  $\text{LL}_{GL_2,\ell}$  is independent of the choice of  $\iota$ .

We can not make a similar twist for an L-parameter of a general connected reductive group  $G$  as the following example suggests.

**Example 2.1.** *We have a commutative diagram*

$$\begin{array}{ccc} \text{Irr}(PGL_2(F)) & \xrightarrow{\text{LL}_{PGL_2}} & \Phi(PGL_2) \\ \downarrow & & \downarrow \\ \text{Irr}(GL_2(F)) & \xrightarrow{\text{LL}_{GL_2}} & \Phi(GL_2) \end{array}$$

by the functoriality. On the other hand, there does not exist a map

$$\text{LL}_{PGL_2,\ell}: \text{Irr}_\ell(PGL_2(F)) \rightarrow \Phi_\ell(PGL_2)$$

which makes the commutative diagram

$$\begin{array}{ccc} \text{Irr}_\ell(PGL_2(F)) & \xrightarrow{\text{LL}_{PGL_2,\ell}} & \Phi_\ell(PGL_2) \\ \downarrow & & \downarrow \\ \text{Irr}_\ell(GL_2(F)) & \xrightarrow{\text{LL}_{GL_2,\ell}} & \Phi_\ell(GL_2), \end{array}$$

because  $(\det \circ p_{\widehat{GL_2}} \circ \varphi)(w) = |w|$  for  $w \in W_F$  if  $[\varphi] \in \Phi_\ell(GL_2)$  is the image under  $\text{LL}_{GL_2,\ell}$  of an element of  $\text{Irr}_\ell(GL_2(F))$  coming from  $\text{Irr}_\ell(PGL_2(F))$ .

## 2.2 $\ell$ -adic C-parameter

Let  ${}^C G = {}^L \tilde{G}$  be the C-group defined in [BG14, Definition 5.38]. We take a Borel subgroup  $B \subset G_{\overline{F}}$  and a maximal torus  $T \subset B$  defined over  $\overline{F}$ . Let  $\rho_G$  denote the half sum of positive roots of  $G$  with respect to  $T$  and  $B$ . Then  $2\rho_G$  defines a character  $\chi_G: \mathbb{G}_m \rightarrow \widehat{T}$ . We put  $z_G = \chi_G(-1)$ . Then  $z_G$  is central in  $\widehat{G}$  and independent of choices of  $B$  and  $T$  by [BG14, Proposition 5.39]. By the independent of choices, we see that  $z_G \in Z(\widehat{G})^{\Gamma_F} \subset {}^L G$ .

We have

$${}^C G \simeq (\widehat{G} \times \mathbb{G}_m / \langle (z_G, -1) \rangle) \rtimes W_F$$

by [BG14, Proposition 5.39]. We sometimes express a point of  ${}^C G$  as  $[(g, z, w)]$  using the above isomorphism. We define  $t_{\mathbb{G}_m}: {}^C G \rightarrow \mathbb{G}_m$  by  $t_{\mathbb{G}_m}([(g, z, w)]) = z^2$ . We have an exact sequence

$$1 \rightarrow {}^L G \rightarrow {}^C G \xrightarrow{t_{\mathbb{G}_m}} \mathbb{G}_m \rightarrow 1.$$

**Definition 2.2.** An  $\ell$ -adic C-parameter for  $G$  is an  $\ell$ -adic L-parameter  $\varphi$  for  $\tilde{G}$  such that  $(t_{\mathbb{G}_m} \circ \varphi)(w) = q^{d_F(w)}$ .

Let  $\Phi_\ell^C(G)$  denote the equivalence classes of  $\ell$ -adic C-parameter for  $G$ .

We take  $c \in \overline{\mathbb{Q}_\ell}$  such that  $c^2 = q$ . Define  $i_c: {}^L G \rightarrow {}^C G$  by  $i_c(g, w) = [(g, c^{d_F(w)}, w)]$ . For an  $\ell$ -adic L-parameter  $\varphi$  for  $G$ , we put  $\varphi_c = i_c \circ \varphi$ .

**Lemma 2.3.** We have a bijection between the set of the  $\ell$ -adic L-parameters for  $G$  and the set of the  $\ell$ -adic C-parameters for  $G$  given by sending  $\varphi$  to  $\varphi_c$ . Further, this induces a bijection  $\Phi_\ell(G) \simeq \Phi_\ell^C(G)$ .

*Proof.* The first claim follows from the definitions. If two  $\ell$ -adic L-parameters for  $\tilde{G}$  are conjugate by an element of  $\tilde{G}(\overline{\mathbb{Q}_\ell})$ , then they are conjugate by an element of  $G(\overline{\mathbb{Q}_\ell})$ , since  $\tilde{G}(\overline{\mathbb{Q}_\ell})$  is generated by  $G(\overline{\mathbb{Q}_\ell})$  and  $Z_{\tilde{G}(\overline{\mathbb{Q}_\ell})}({}^C G(\overline{\mathbb{Q}_\ell}))$ . Hence the second claim follows from the first one.  $\square$

We take an isomorphism  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}_\ell}$ . We define a map

$$\mathcal{C}_{G,\ell}^\iota: \Phi_\ell(G) \rightarrow \Phi_\ell^C(G)$$

by sending  $[\varphi]$  to  $[\varphi_c]$  where  $c = \iota(q^{\frac{1}{2}})$ . Then  $\mathcal{C}_{G,\ell}^\iota$  is a bijection by Lemma 2.3.

We define

$$\text{LL}_{G,\ell}^{\mathcal{C},\iota}: \text{Irr}_\ell(G(F)) \rightarrow \Phi_\ell^C(G)$$

as  $\mathcal{C}_{G,\ell}^\iota \circ \text{LL}_{G,\ell}^\iota$ .

**Conjecture 2.4.** The map  $\text{LL}_{G,\ell}^{\mathcal{C},\iota}$  is independent of a choice of  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}_\ell}$ .

## 2.3 Tannakian $\ell$ -adic L-parameter

For an algebraic group  $H$  over a field, a character  $\chi$  of  $H$  and a cocharacter  $\mu$  of  $H$ , we define  $\langle \chi, \mu \rangle_H \in \mathbb{Z}$  by

$$(\chi \circ \mu)(z) = z^{\langle \chi, \mu \rangle_H}.$$

Let  $\mathcal{M}_G$  be the conjugacy classes of cocharacters  $\mathbb{G}_m \rightarrow G_{\overline{F}}$ . Let  $[\mu] \in \mathcal{M}_G$ . We put

$$d_G([\mu]) = \langle 2\rho_G, \mu \rangle_T,$$



where we take a Borel subgroup  $B \subset G_{\overline{F}}$ , a maximal torus  $T \subset B$  defined over  $\overline{F}$  and a dominant representative  $\mu \in X_*(T)$ . Let  $E_{[\mu]}$  be the field of definition of  $[\mu]$ . Let  $r_{\widehat{G},[\mu]}$  be the irreducible representation of  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$  of highest weight  $\mu$  viewed as a dominant character of a maximal torus of  $\widehat{G}$ .

We take  $c \in \overline{\mathbb{Q}}_\ell$  such that  $c^2 = q$ . For an integer  $m$ , let

$$\left(\frac{m}{2}\right)_c$$

denote the twist by the character  $W_F \rightarrow \overline{\mathbb{Q}}_\ell$  sending  $w$  to  $c^{md_F(w)}$ .

Let  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G)$  denote the category of continuous finite dimensional representations of  ${}^L G(\overline{\mathbb{Q}}_\ell)$  over  $\overline{\mathbb{Q}}_\ell$  whose restrictions to  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$  is algebraic. Let  $r: {}^L G(\overline{\mathbb{Q}}_\ell) \rightarrow \text{Aut}(V)$  be an object in  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G)$ . Then we have a decomposition

$$V = \bigoplus_{[\mu] \in \mathcal{M}_G} V_{[\mu]}$$

as representations of  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$  where  $V_{[\mu]}$  is the  $r_{\widehat{G},[\mu]}$ -typic part of  $V$ . For an  $\ell$ -adic L-parameter  $\varphi$ , we define  $(r \circ \varphi)_c: W_F \rightarrow \text{Aut}(V)$  by

$$\bigoplus_{[\mu] \in \mathcal{M}_G} V_{[\mu]} \left( \frac{d_G([\mu])}{2} \right)_c,$$

which is well-defined since  $d_G(w[\mu]) = d_G([\mu])$  for  $w \in W_F$ .

For an  $\ell$ -adic L-parameter  $\varphi$  for  $G$ , we define a Tannakian functor

$$\mathcal{F}_{\varphi,c}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$$

by

$$\mathcal{F}_{\varphi,c}(r) = (r \circ \varphi)_c.$$

**Definition 2.5.** A Tannakian  $\ell$ -adic L-parameter for  $G$  is a functor

$$\mathcal{F}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$$

which is equal to  $\mathcal{F}_{\varphi,c}$  for an  $\ell$ -adic L-parameter  $\varphi$  for  $G$ . We say that two Tannakian  $\ell$ -adic L-parameters  $\mathcal{F}$  and  $\mathcal{F}'$  are equivalent if there is  $g \in \widehat{G}(\overline{\mathbb{Q}}_\ell)$  such that, for all  $r \in \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G)$ , we have  $\mathcal{F}(r) = \mathcal{F}'(r)^{r(g)}$ .

**Lemma 2.6.** The set of the Tannakian  $\ell$ -adic L-parameters for  $G$  is independent of a choice of  $c \in \overline{\mathbb{Q}}_\ell$  such that  $c^2 = q$ .

*Proof.* We have

$$\mu(z_G) = (-1)^{\langle \mu, \chi \rangle_{\widehat{T}}} = (-1)^{d_G([\mu])}$$

for  $\mu \in X^*(\widehat{T}) \simeq X_*(T)$ . Let

$$\chi_{z_G}: W_F \rightarrow Z(\widehat{G})^{\Gamma_F}(\overline{\mathbb{Q}}_\ell) \hookrightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$$

be the character sending  $w$  to  $z_G^{d_F(w)}$ . Then we have

$$(r \circ \varphi)_{-c} = (r \circ (\chi_{z_G} \varphi))_c \tag{2.1}$$

for an  $\ell$ -adic L-parameter  $\varphi$  and  $r \in \text{Rep}({}^L G)$ . Since  $\chi_{z_G} \varphi$  is also an  $\ell$ -adic L-parameter, the claim follows.  $\square$

Let  $\Phi_\ell^T(G)$  be the set of equivalence classes of Tannakian  $\ell$ -adic L-parameters for  $G$ . This set is independent of a choice of  $c$  by Lemma 2.6.

**Lemma 2.7.** *We have a bijection between the set of the  $\ell$ -adic L-parameters for  $G$  and the set of the Tannakian  $\ell$ -adic L-parameters for  $G$  given by sending  $\varphi$  to  $F_{\varphi,c}$ . Further, this induces a bijection  $\Phi_\ell(G) \simeq \Phi_\ell^T(G)$ .*

*Proof.* We show the first claim. The map is surjective by Definition 2.5. Assume that  $\varphi$  and  $\varphi'$  are different  $\ell$ -adic L-parameters for  $G$  and  $\mathcal{F}_{\varphi,c} = \mathcal{F}_{\varphi',c}$ . We take  $w \in W_F$  such that  $\varphi(w) \neq \varphi'(w)$ . Further, we take a finite Galois extension  $F'$  of  $F$  such that  $G$  splits over  $F'$  and the images of  $\varphi(w)$  and  $\varphi'(w)$  in  $\widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \text{Gal}(F'/F)$  are different. By considering a representation of  ${}^L G(\overline{\mathbb{Q}}_\ell)$  which factors through a faithful algebraic representation  $\widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \text{Gal}(F'/F)$ , we have a contradiction to  $\mathcal{F}_{\varphi,c} = \mathcal{F}_{\varphi',c}$ . Hence the map is injective.

The second claim follows from the first one.  $\square$

For a Tannakian  $\ell$ -adic L-parameter  $\mathcal{F}$  for  $G$ , we take an  $\ell$ -adic L-parameter  $\varphi$  for  $G$  such that  $\mathcal{F} = \mathcal{F}_{\varphi,c}$ . Then the centralizer  $S_\varphi = Z_{\widehat{G}(\overline{\mathbb{Q}}_\ell)}(\text{Im } \varphi)$  is independent of a choice of  $c$  by (2.1). We write  $S_{\mathcal{F}}$  for  $S_\varphi$ . Then  $\mathcal{F}$  naturally factors through

$$\mathcal{F}_S: \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_{\mathcal{F}} \times W_F).$$

For a finite separable extension  $F'$  of  $F$ , we define the restriction

$$\mathcal{F}|_{F'}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G_{F'}) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_{F'})$$

of  $\mathcal{F}$  to  $F'$  by the usual restriction to  $W_{F'}$  of an  $\ell$ -adic L-parameter for  $G$  and bijections given by Lemma 2.7. Then  $\mathcal{F}|_{F'}$  naturally factors through

$$\mathcal{F}_S|_{F'}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G_{F'}) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_{\mathcal{F}} \times W_{F'}).$$

We take an isomorphism  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$ . We define a map

$$\mathcal{T}_{G,\ell}^\iota: \Phi_\ell(G) \rightarrow \Phi_\ell^T(G)$$

by sending  $[\varphi]$  to  $[F_{\varphi,c}]$  where  $c = \iota(q^{\frac{1}{2}})$ . Then  $\mathcal{T}_{G,\ell}^\iota$  is a bijection by Lemma 2.7.

We define

$$\text{LL}_{G,\ell}^{\text{T},\iota}: \text{Irr}_\ell(G(F)) \rightarrow \Phi_\ell^T(G)$$

as  $\mathcal{T}_{G,\ell}^\iota \circ \text{LL}_{G,\ell}^\iota$ .

**Conjecture 2.8.** *The map  $\text{LL}_{G,\ell}^{\text{T},\iota}$  is independent of a choice of  $\iota: \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$ .*

**Remark 2.9.** *Conjecture 2.8 is motivated by the Kottwitz conjecture for local Shimura varieties in [RV14, Conjecture 7.4]. We use notations in [RV14, Conjecture 7.4]. Let  $[\pi] \in \text{Irr}_\ell(G)$  and  $[\rho] \in \text{Irr}_\ell(J)$ . Assume that*

$$[\mathcal{F}^\iota] = \text{LL}_{G,\ell}^{\text{T},\iota}([\pi]) = \text{LL}_{J,\ell}^{\text{T},\iota}([\rho])$$

for a Tannakian  $\ell$ -adic L-parameter  $\mathcal{F}^\iota$ . Let  $\delta_{\pi,\rho}^\iota$  be the representation of  $S_{\mathcal{F}^\iota}$  over  $\overline{\mathbb{Q}}_\ell$  determined by  $\iota$  and the construction of  $\check{\tau}_\pi \otimes \tau_\rho$  in [RV14, p. 312] (cf. [KW17, 2.3] for a construction in a more general case). If the Kottwitz conjecture is true, then the isomorphism class of the  $W_{E_{[\mu]}}$ -representation

$$\text{Hom}_{S_{\mathcal{F}^\iota}}(\delta_{\pi,\rho}^\iota, \mathcal{F}_S|_{E_{[\mu]}}(r_{[\mu]}))$$

is independent of  $\iota$ , since the  $\ell$ -adic cohomology of local Shimura varieties and their group actions are independent of  $\iota$ .

## 2.4 Comparison

**Proposition 2.10.** *Conjecture 2.4 and Conjecture 2.8 are equivalent.*

*Proof.* Let  $\iota': \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$  be another choice. We put  $c' = \iota'(q^{\frac{1}{2}})$ . Let  $[\pi] \in \text{Irr}_\ell(G(F))$ . Assume that  $\text{LL}'_{G,\ell}([\pi]) = [\varphi]$  and  $\text{LL}'_{G,\ell}([\pi]) = [\varphi']$ .

Assume that Conjecture 2.4 is true. Let  $r: {}^L G(\overline{\mathbb{Q}}_\ell) \rightarrow \text{Aut}(V)$  be an object in  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G)$ . Then we have a decomposition

$$V = \bigoplus_{[\mu] \in \mathcal{M}_G} V_{[\mu]}$$

as representations of  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$  where  $V_{[\mu]}$  is the  $r_{\widehat{G},[\mu]}$ -typic part of  $V$ . We extend  $r$  to  $\tilde{r} \in \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^C G)$  by letting  $\mathbb{G}_m$  act on  $V_{[\mu]}$  by  $z \mapsto z^{d_G([\mu])}$ . Then we have

$$\tilde{r} \circ \varphi_c = \tilde{r} \circ i_c \circ \varphi = \mathcal{F}_{\varphi,c}(r). \quad (2.2)$$

By Conjecture 2.4,  $\varphi_c$  and  $\varphi'_{c'}$  are conjugate by an element of  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$ . It implies that they are conjugate by an element of  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$ . Hence, Conjecture 2.8 is also true by (2.2).

Assume that Conjecture 2.8 is true. Then there is  $g \in \widehat{G}(\overline{\mathbb{Q}}_\ell)$  such that  $\mathcal{F}_{\varphi,c}(r) = \mathcal{F}_{\varphi',c'}(r)^{r(g)}$  for all  $r \in \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G)$ . Replacing  $\varphi'$  by  $\varphi'^g$ , we may assume that  $\mathcal{F}_{\varphi,c}(r) = \mathcal{F}_{\varphi',c'}(r)$  for all  $r \in \text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^L G)$ . Assume that  $\varphi_c \neq \varphi'_{c'}$ . We take an object  $\tilde{r}: {}^C G(\overline{\mathbb{Q}}_\ell) \rightarrow \text{Aut}(V)$  in  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}({}^C G)$  such that  $\tilde{r} \circ \varphi_c \neq \tilde{r} \circ \varphi'_{c'}$ .

Then we have a decomposition

$$V = \bigoplus_{[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}} V_{[\tilde{\mu}]}$$

as representations of  $\widehat{G}(\overline{\mathbb{Q}}_\ell)$  where  $V_{[\tilde{\mu}]}$  is the  $r_{\widehat{G},[\tilde{\mu}]}$ -typic part of  $V$ . For  $[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}$ , let  $n_{[\tilde{\mu}]}$  be an integer such that  $\tilde{\mu}|_{\mathbb{G}_m}(z) = z^{n_{[\tilde{\mu}]}}$ , where we view  $\tilde{\mu}$  as a character of a maximal torus of  $\widehat{G}$ . Then  $\tilde{r} \circ \varphi_c = \tilde{r} \circ i_c \circ \varphi$  is given by

$$V = \bigoplus_{[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}} V_{[\tilde{\mu}]} \left( \frac{n_{[\tilde{\mu}]}}{2} \right)_c. \quad (2.3)$$

Let  $r$  be the restriction of  $\tilde{r}$  to  ${}^L G$ . Then,  $\mathcal{F}_{\varphi,c}(r)$  is obtained by

$$V = \bigoplus_{[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}} V_{[\tilde{\mu}]} \left( \frac{d_G([\mu])}{2} \right)_c, \quad (2.4)$$

where  $[\mu]$  is the image in  $\mathcal{M}_G$  of  $[\tilde{\mu}]$ . We have

$$n_{[\tilde{\mu}]} \equiv d_G([\mu]) \pmod{2}, \quad (2.5)$$

since

$$(-1)^{n_{[\tilde{\mu}]}} = \mu(z_G) = (-1)^{d_G([\mu])}.$$

If we replace  $\iota$  by  $\iota'$ , then  $\mathcal{F}_{\varphi,c}(r)$  does not change, hence  $\tilde{r} \circ \varphi_c$  also does not change by (2.3), (2.4) and (2.5). This contradicts  $\tilde{r} \circ \varphi_c \neq \tilde{r} \circ \varphi'_{c'}$ . Hence Conjecture 2.4 is true.  $\square$

**Theorem 2.11.** *Conjecture 2.4 and Conjecture 2.8 are true for  $GL_n$ .*

*Proof.* Recall that the local Langlands correspondence for  $GL_n$  is known by [LRS93] and [HT01]. Assume that  $\text{LL}_{GL_n, \ell}^{C, \iota}([\pi]) = [\varphi']$ . Note that  $z_{GL_n} = (-1)^{n-1}$ . Let

$$\tilde{r}: {}^C GL_n \rightarrow \widehat{GL}_n; [(g, z, w)] \mapsto gz^{n-1}.$$

By [BH00, 3.2 Theorem], we can show that  $[\tilde{r} \circ \varphi']$  is independent of  $\iota$  similarly as the proof of [BH06, 35.1 Theorem].

Let  $\iota': \mathbb{C} \simeq \overline{\mathbb{Q}}_\ell$  be another choice. Taking a conjugation of  $\varphi'$  by an element of  $\widehat{GL}_n(\overline{\mathbb{Q}}_\ell)$ , we may assume that  $\tilde{r} \circ \varphi' = \tilde{r} \circ \varphi'^{\iota'}$ . Hence there is  $\chi: W_F \rightarrow \overline{\mathbb{Q}}_\ell^\times$  such that

$$\varphi'^{\iota'}(w) = \varphi'(w)[(\chi(w)^{1-n}, \chi(w), 1)]$$

for  $w \in W_F$ . By Definition 2.2, we have  $t_{\mathbb{G}_m} \circ \varphi' = t_{\mathbb{G}_m} \circ \varphi'^{\iota'}$ . Hence we have  $\chi^2 = 1$ . This implies that  $\varphi' = \varphi'^{\iota'}$ . Hence Conjecture 2.4 is true. Then Conjecture 2.8 is also true by Proposition 2.10.  $\square$

## References

- [AG91] J. Arthur and S. Gelbart, Lectures on automorphic  $L$ -functions, in  $L$ -functions and arithmetic (Durham, 1989), vol. 153 of *London Math. Soc. Lecture Note Ser.*, pp. 1–59, Cambridge Univ. Press, Cambridge, 1991.
- [BG14] K. Buzzard and T. Gee, The conjectural connections between automorphic representations and Galois representations, in Automorphic forms and Galois representations. Vol. 1, vol. 414 of *London Math. Soc. Lecture Note Ser.*, pp. 135–187, Cambridge Univ. Press, Cambridge, 2014.
- [BH00] C. J. Bushnell and G. Henniart, Davenport-Hasse relations and an explicit Langlands correspondence. II. Twisting conjectures, vol. 12, pp. 309–347, 2000, colloque International de Théorie des Nombres (Talence, 1999).
- [BH06] C. J. Bushnell and G. Henniart, The local Langlands conjecture for  $GL(2)$ , vol. 335 of *Grundlehren der Mathematischen Wissenschaften*, Springer-Verlag, Berlin, 2006.
- [Bor79] A. Borel, Automorphic  $L$ -functions, in Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979 pp. 27–61.
- [Del73] P. Deligne, Les constantes des équations fonctionnelles des fonctions  $L$ , in Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), pp. 501–597. Lecture Notes in Math., Vol. 349, Springer, Berlin, 1973.
- [Hei06] V. Heiermann, Orbites unipotentes et pôles d'ordre maximal de la fonction  $\mu$  de Harish-Chandra, *Canad. J. Math.* 58 (2006), no. 6, 1203–1228.

- [HT01] M. Harris and R. Taylor, The geometry and cohomology of some simple Shimura varieties, vol. 151 of *Annals of Mathematics Studies*, Princeton University Press, Princeton, NJ, 2001, with an appendix by Vladimir G. Berkovich.
- [IT] N. Imai and T. Tsushima, Geometric realization of the local Langlands correspondence for representations of conductor three, to appear in *Publ. Res. Inst. Math. Sci.*
- [Joh13] C. Johansson, A remark on a conjecture of Buzzard-Gee and the cohomology of Shimura varieties, *Math. Res. Lett.* 20 (2013), no. 2, 279–288.
- [Kal16] T. Kaletha, The local Langlands conjectures for non-quasi-split groups, in *Families of automorphic forms and the trace formula*, Simons Symp., pp. 217–257, Springer, [Cham], 2016.
- [KL87] D. Kazhdan and G. Lusztig, Proof of the Deligne-Langlands conjecture for Hecke algebras, *Invent. Math.* 87 (1987), no. 1, 153–215.
- [Kot84] R. E. Kottwitz, Stable trace formula: cuspidal tempered terms, *Duke Math. J.* 51 (1984), no. 3, 611–650.
- [KW17] T. Kaletha and J. Weinstein, On the Kottwitz conjecture for moduli spaces of local shtukas, 2017, arXiv:1709.06651.
- [Lan83] R. P. Langlands, Les débuts d’une formule des traces stable, vol. 13 of *Publications Mathématiques de l’Université Paris VII [Mathematical Publications of the University of Paris VII]*, Université de Paris VII, U.E.R. de Mathématiques, Paris, 1983.
- [LRS93] G. Laumon, M. Rapoport and U. Stuhler,  $\mathcal{D}$ -elliptic sheaves and the Langlands correspondence, *Invent. Math.* 113 (1993), no. 2, 217–338.
- [RV14] M. Rapoport and E. Viehmann, Towards a theory of local Shimura varieties, *Münster J. Math.* 7 (2014), no. 1, 273–326.
- [Ser96] J.-P. Serre, Exemples de plongements des groupes  $\mathrm{PSL}_2(\mathbf{F}_p)$  dans des groupes de Lie simples, *Invent. Math.* 124 (1996), no. 1-3, 525–562.
- [ST68] J.-P. Serre and J. Tate, Good reduction of abelian varieties, *Ann. of Math. (2)* 88 (1968), 492–517.

Naoki Imai

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan

naoki@ms.u-tokyo.ac.jp