

# Local Langlands correspondences in $\ell$ -adic coefficients

Naoki Imai

## Abstract

Let  $\ell$  be a prime number different from the residue characteristic of a non-archimedean local field  $F$ . We give formulations of  $\ell$ -adic local Langlands correspondences for connected reductive algebraic groups over  $F$ , which we conjecture to be independent of a choice of an isomorphism between the  $\ell$ -adic coefficient field and the complex number field.

## Introduction

The local Langlands correspondence for a connected reductive algebraic group over a non-archimedean local field  $F$  is usually formulated with coefficients in  $\mathbb{C}$  because of its relation with automorphic representations. On the other hand, when we discuss a realization of the local Langlands correspondence in  $\ell$ -adic cohomology, we need a correspondence over  $\overline{\mathbb{Q}}_\ell$ , where  $\ell$  is a prime number different from the residue characteristic of  $F$ . We can take an isomorphism  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$  and use it to transfer the local Langlands correspondence over  $\mathbb{C}$  to a local Langlands correspondence over  $\overline{\mathbb{Q}}_\ell$ . However, the obtained correspondence over  $\overline{\mathbb{Q}}_\ell$  depends on the choice of  $\iota$ . In [BH06], Bushnell–Henniart formulated an  $\ell$ -adic local Langlands correspondence for  $\mathrm{GL}_2$ , which is independent of a choice of an isomorphism  $\mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$  by making some twists of L-parameters. The  $\ell$ -adic local Langlands correspondence is suitable to describe the non-abelian Lubin–Tate theory in the sense that it is canonical (*cf.* [IT12, 5]).

In this paper, we discuss formulations of  $\ell$ -adic local Langlands correspondences for general connected reductive groups. A natural idea is to make similar twists as  $\mathrm{GL}_2$ -case for L-parameters. However, we can not make such twists in general as explained in Example 2.1. A problem is that we do not have enough space inside the Langlands dual group to make an appropriate twist. To overcome this problem, we have two approaches. One is to introduce  $\ell$ -adic C-parameters using C-groups, which make spaces for twists. Another is to introduce Tannakian  $\ell$ -adic L-parameters that incorporate necessary twists in the realizations. Using these parameters, we formulate  $\ell$ -adic local Langlands correspondences, which we conjecture to be independent of a choice of an isomorphism  $\mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$ . We show that two conjectures formulated by using  $\ell$ -adic C-parameters and Tannakian  $\ell$ -adic L-parameters are equivalent. Further, we confirm that the conjectures are true in  $\mathrm{GL}_n$ -case. The formulation of the  $\ell$ -adic

---

2010 *Mathematics Subject Classification.* Primary: 11F70; Secondary: 11F80.

local Langlands correspondence using Tannakian  $\ell$ -adic L-parameters is motivated by the Kottwitz conjecture for local Shimura varieties in [RV14, Conjecture 7.4]. In the number field case, a relation between C-groups and the Kottwitz conjecture for Shimura varieties is discussed in [Joh13].

In Section 1, we recall various versions of L-parameters and explain their relations. In Section 2, after explaining the problem, we give formulations of the  $\ell$ -adic local Langlands correspondences introducing the  $\ell$ -adic C-parameter and the Tannakian  $\ell$ -adic L-parameter.

After we put a former version of this paper on arXiv, related papers [Ber20] and [Zhu20] appeared on arXiv. [Ber20] also explains a relation between C-groups and the local Langlands correspondence. In a similar philosophy as this paper, formulations of Satake isomorphisms using C-groups are explained in [Ber20] and [Zhu20].

## Acknowledgements

The author would like to thank Teruhisa Koshikawa for his helpful comment. This work was supported by JSPS KAKENHI Grant Number 18H01109.

## Notation

For a field  $F$ , let  $\Gamma_F$  denote the absolute Galois group of  $F$ . For a non-archimedean local field  $F$ , let  $W_F$  and  $I_F$  denote the Weil group of  $F$  and its inertia subgroup. For a homomorphism  $\phi: G \rightarrow H$  of groups and  $g \in G$ , let  $\phi^g$  denote the homomorphism defined by  $\phi^g(g') = \phi(gg'g^{-1})$  for  $g' \in G$ . For a group  $G$ , let  $Z(G)$  denote the center of  $G$ . For a group  $G$ , a subgroup  $H \subset G$  and a subset  $S \subset G$ , let  $Z_H(S)$  denote the centralizer of  $S$  in  $H$ . Let  $\text{Ad}$  denote the adjoint action of a group, and  $\text{ad}$  denote the adjoint action of a Lie algebra.

# 1 Langlands parameters

## 1.1 Over algebraically closed fields

Let  $F$  be a non-archimedean local field of residue characteristic  $p$ . Let  $q$  be the number of elements of the residue field of  $F$ . Let  $v_F: F^\times \rightarrow \mathbb{Z}$  be the normalized valuation of  $F$ . Let

$$\text{Art}_F: F^\times \xrightarrow{\sim} W_F^{\text{ab}}$$

be the Artin reciprocity isomorphism normalized so that a uniformizer is sent to a lift of the geometric Frobenius element. For  $w \in W_F$ , we put

$$d_F(w) = v_F(\text{Art}_F^{-1}(\overline{w})), \quad |w| = q^{-d_F(w)},$$

where  $\overline{w}$  denotes the image of  $w$  in  $W_F^{\text{ab}}$ .

Let  $G$  be a connected reductive algebraic group over  $F$ . Let  $C$  be an algebraically closed field of characteristic 0. Let  $\widehat{G}$  be the dual group of  $G$  over  $C$ . Let  ${}^L G = \widehat{G} \rtimes W_F$  be the L-group of  $G$  defined in [Bor79, 2.3]. We say that an element of  ${}^L G(C)$  is semisimple if its image in  $\widehat{G}(C) \rtimes \text{Gal}(F'/F)$  is semisimple for any finite Galois extension  $F'$  over  $F$  that splits  $G$ . Let  $p_{\widehat{G}}: {}^L G(C) \rightarrow \widehat{G}(C)$  denote the projection map.

**Definition 1.1.** Let  $H$  be a group with action of  $W_F$ . Let  $\phi: H \rtimes W_F \rightarrow {}^L G(C)$  be a homomorphism of groups over  $W_F$ .

- (1) We say that  $\phi$  is semisimple if all the elements of  $\phi(W_F)$  are semisimple.
- (2) We say that  $\phi$  is relevant if any parabolic subgroup  $P$  of  ${}^L G$  containing  $\text{Im } \phi$  is relevant in the sense of [Bor79, 3.3].

We say that two homomorphisms  $H \rtimes W_F \rightarrow {}^L G(C)$  of groups over  $W_F$  are equivalent if they are conjugate by an element of  $\widehat{G}(C)$ .

**Lemma 1.2.** Let  $\phi: W_F \rightarrow {}^L G(C)$  be a homomorphism of groups over  $W_F$  such that  $p_{\widehat{G}} \circ \phi$  is smooth. Let  $\sigma_q \in W_F$  be a lift of the  $q$ -th power Frobenius element. Assume that  $\phi(\sigma_q)$  is semisimple. Then  $\phi$  is semisimple.

*Proof.* We write  $w \in W_F$  as  $\sigma_q^m \sigma$  for  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ . Since  $(p_{\widehat{G}} \circ \phi)(I_F)$  is finite, there is a positive integer  $d$  such that  $\phi((\sigma_q^m \sigma)^d) = \phi(\sigma_q^{dm})$ . Then the claim follows, because  $\phi(w)$  is semisimple if and only if  $\phi(w)^d$  is semisimple.  $\square$

Let  $WD_F = \mathbb{G}_a \rtimes W_F$  be the Weil–Deligne group scheme over  $\mathbb{Q}$  for  $F$  defined in [Del73, 8.3.6].

**Definition 1.3** (cf. [Bor79, 8.2]). (1) An  $L$ -homomorphism of Weil–Deligne type for  $G$  over  $C$  is a homomorphism

$$\xi: WD_F(C) \rightarrow {}^L G(C)$$

of groups over  $W_F$  such that  $\xi|_{\mathbb{G}_a(C)}$  is algebraic and  $(p_{\widehat{G}} \circ \xi)|_{W_F}$  is smooth.

- (2) An  $L$ -parameter of Weil–Deligne type for  $G$  over  $C$  is a semisimple relevant  $L$ -homomorphism of Weil–Deligne type for  $G$  over  $C$ .

Let  $\mathcal{L}_C^{\text{WD}}(G)$  denote the set of equivalence classes of  $L$ -homomorphisms of Weil–Deligne type for  $G$  over  $C$ . Let  $\Phi_C^{\text{WD}}(G)$  denote the set of equivalence classes of  $L$ -parameters of Weil–Deligne type for  $G$  over  $C$ .

**Definition 1.4.** (1) A Weil–Deligne  $L$ -homomorphism for  $G$  over  $C$  is a pair  $(\tau, N)$  of a homomorphism  $\tau: W_F \rightarrow {}^L G(C)$  of groups over  $W_F$  and  $N \in \text{Lie}(\widehat{G})(C)$  such that  $p_{\widehat{G}} \circ \tau$  is smooth and  $\text{Ad}(\tau(w))N = |w|N$  for  $w \in W_F$ . The second component  $N$  of a Weil–Deligne  $L$ -homomorphism  $(\tau, N)$  is called a monodromy operator. We say that two Weil–Deligne  $L$ -homomorphisms for  $G$  over  $C$  are equivalent if they are conjugate by an element of  $\widehat{G}(C)$ .

- (2) A Weil–Deligne  $L$ -parameter for  $G$  over  $C$  is a Weil–Deligne  $L$ -homomorphism  $(\tau, N)$  for  $G$  over  $C$  such that  $\tau$  is semisimple and any parabolic subgroup  $P$  of  ${}^L G$  containing  $\tau(W_F)$  and satisfying  $N \in \text{Lie}(P \cap \widehat{G})(C)$  is relevant.

Let  $\mathcal{L}_C^{\text{M}}(G)$  denote the set of equivalence classes of Weil–Deligne  $L$ -homomorphisms for  $G$  over  $C$ . Let  $\Phi_C^{\text{M}}(G)$  denote the set of equivalence classes of Weil–Deligne  $L$ -parameters for  $G$  over  $C$ .

**Remark 1.5.** Let  $\iota: C \xrightarrow{\sim} C'$  be an isomorphism of algebraically closed fields of characteristic 0. Then  $\iota$  induces bijections  $\mathcal{L}_C^{\text{WD}}(G) \simeq \mathcal{L}_{C'}^{\text{WD}}(G)$ ,  $\Phi_C^{\text{WD}}(G) \simeq \Phi_{C'}^{\text{WD}}(G)$ ,  $\mathcal{L}_C^{\text{M}}(G) \simeq \mathcal{L}_{C'}^{\text{M}}(G)$  and  $\Phi_C^{\text{M}}(G) \simeq \Phi_{C'}^{\text{M}}(G)$ .

**Lemma 1.6.** Let  $(\tau, N)$  be a Weil–Deligne L-homomorphism for  $G$  over  $C$ . Then  $N$  is an nilpotent element of  $\text{Lie}(\widehat{G}^{\text{der}}(C))$ .

*Proof.* We take a finite separable extension  $F'$  of  $F$  that splits  $G$ . By the condition  $\text{Ad}(\tau(w))N = |w|N$  for  $w \in W_{F'}$ , we have  $N \in \text{Lie}(\widehat{G}^{\text{der}}(C))$ . Further, the adjoint endomorphism  $\text{ad}(N) \in \text{End}(\text{Lie}(\widehat{G}^{\text{der}}(C)))$  satisfies

$$\text{Ad}(\tau(w)) \circ \text{ad}(N) = |w| \text{ad}(N) \circ \text{Ad}(\tau(w))$$

for  $w \in W_F$ . Hence  $N$  is an nilpotent element.  $\square$

**Proposition 1.7.** For an L-homomorphism of Weil–Deligne type  $\xi$  for  $G$  over  $C$ , we put  $\tau_\xi = \xi|_{W_F}$  and  $N_\xi = \text{Lie}(\xi|_{\text{Ga}(C)})(1)$ . Then  $\xi \mapsto (\tau_\xi, N_\xi)$  induces bijections  $\mathcal{L}_C^{\text{WD}}(G) \simeq \mathcal{L}_C^{\text{M}}(G)$  and  $\Phi_C^{\text{WD}}(G) \simeq \Phi_C^{\text{M}}(G)$ .

*Proof.* Let  $(\tau, N)$  be a Weil–Deligne L-homomorphism for  $G$  over  $C$ . By Lemma 1.6, there is a unipotent radical  $U$  of a Borel subgroup of  $\widehat{G}$  such that  $N \in \text{Lie}(U(C))$ . Then there is an exponential map

$$\exp: \text{Lie}(U(C)) \rightarrow U(C)$$

as in [Ser96, 2.2]. We define

$$\xi_{(\tau, N)}: WD_F(C) \rightarrow {}^L G(C)$$

by  $\xi_{(\tau, N)}(a, w) = \exp(aN)\tau(w)$ . Then  $(\tau, N) \mapsto \xi_{(\tau, N)}$  induces the inverses.  $\square$

## 1.2 Over $\mathbb{C}$

Assume that  $C = \mathbb{C}$  in this subsection.

**Definition 1.8** (cf. [AG91, I.2]). An L-parameter for  $G$  is a semisimple relevant continuous homomorphism

$$\phi: \text{SU}_2(\mathbb{R}) \times W_F \rightarrow {}^L G(\mathbb{C})$$

of groups over  $W_F$ .

Let  $\Phi(G)$  denote the set of equivalence classes of L-parameters for  $G$ .

**Definition 1.9** (cf. [Lan83, IV.2]). An L-parameter of  $\text{SL}_2$ -type for  $G$  is a semisimple relevant continuous homomorphism

$$\phi: \text{SL}_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$$

of groups over  $W_F$  such that  $\phi|_{\text{SL}_2(\mathbb{C})}$  is algebraic.

Let  $\Phi^{\text{SL}}(G)$  denote the set of equivalence classes of L-parameters of  $\text{SL}_2$ -type for  $G$ .

**Proposition 1.10.** *The map  $\Phi^{\text{SL}}(G) \rightarrow \Phi(G)$  induced by the restriction with respect to  $\text{SU}_2(\mathbb{R}) \subset \text{SL}_2(\mathbb{C})$  is a bijection.*

*Proof.* Let  $\phi: \text{SU}_2(\mathbb{R}) \times W_F \rightarrow {}^L G(\mathbb{C})$  be an L-parameter for  $G$ . Let  $\mathcal{H}$  be the centralizer of  $\phi(W_F)$  in  $\widehat{G}$ . Then  $\mathcal{H}$  is reductive by [Kot84b, 10.1.1 Lemma]. We take a compact real form  $\mathcal{H}_c$  of  $\mathcal{H}$  such that  $\phi(\text{SU}_2(\mathbb{R})) \subset \mathcal{H}_c(\mathbb{R})$ . Let  $\phi_{\text{SU}_2}: \text{SU}_2(\mathbb{R}) \rightarrow \mathcal{H}_c(\mathbb{R})$  be the restriction of  $\phi$  to  $\text{SU}_2(\mathbb{R})$ . The continuous homomorphism  $\phi_{\text{SU}_2}$  extends to an algebraic morphism  $\phi_{\text{SL}_2}: \text{SL}_2(\mathbb{C}) \rightarrow \mathcal{H}(\mathbb{C})$  uniquely by [OV90, 5.2.5 Theorem 11], since any continuous homomorphism between real Lie groups are differentiable. Let  $\phi^{\text{SL}}: \text{SL}_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$  be a homomorphism defined by  $\phi_{\text{SL}_2}$  and  $\phi|_{W_F}$ . Then  $\phi \mapsto \phi^{\text{SL}}$  induces the inverse of the map  $\Phi^{\text{SL}}(G) \rightarrow \Phi(G)$ .  $\square$

**Lemma 1.11.** *Let  $H$  be a compact topological group. Let  $\phi: H \rightarrow {}^L G(\mathbb{C})$  be a continuous homomorphism. Then the centralizer of  $\phi(H)$  in  $\widehat{G}$  is reductive.*

*Proof.* Let  $K$  be the image of  $\phi(H)$  under  $\text{Ad}: {}^L G \rightarrow \text{Aut}(\widehat{G})$ . Then  $\widehat{G}^K$  is reductive as in the proof of [Kot84b, 10.1.2 Lemma]. Hence we obtain the claim.  $\square$

The following is a slight generalization of [Hei06, Proposition 3.5] and [KL87, 2.4].

**Lemma 1.12.** *Let  $\mathcal{G}$  be a reductive group over  $\mathbb{C}$ . Let  $s$  be a semisimple element of  $\mathcal{G}(\mathbb{C})$  and  $u$  be a unipotent element of  $\mathcal{G}(\mathbb{C})$  such that  $sus^{-1} = u^q$ . Then there is an algebraic homomorphism  $\theta: \text{SL}_2(\mathbb{C}) \rightarrow \mathcal{G}(\mathbb{C})$  such that*

$$\theta \left( \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) = u, \quad \theta \left( \begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right) s^{-1} \in Z_{\mathcal{G}(\mathbb{C})}(\text{Im } \theta).$$

*Further, such  $\theta$  is unique up to conjugation by  $Z_{\mathcal{G}(\mathbb{C})}(\{s, u\})$ .*

*Proof.* The first claim is proved in the same way as [Hei06, Proposition 3.5]. We recall the argument briefly. We take an algebraic homomorphism  $\theta: \text{SL}_2(\mathbb{C}) \rightarrow \mathcal{G}(\mathbb{C})$  such that

$$\theta \left( \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) = u.$$

We define  $\mathcal{G}'$  to be the semi-direct product  $\mathcal{G} \rtimes \mathbb{G}_m$ , where the action of  $\mathbb{G}_m$  on  $\mathcal{G}$  is defined by

$$z \cdot g = \text{Ad} \left( \theta \left( \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} \right) \right) g$$

Then  $Z_{\mathcal{G}'(\mathbb{C})}(\text{Im } \theta)$  is a reductive group by [Kot84b, 10.1.2 Lemma] applied to the identity component of  $\mathcal{G}'$ . Further, we can see that  $Z_{\mathcal{G}'(\mathbb{C})}(\text{Im } \theta)$  is a maximal reductive subgroup of  $Z_{\mathcal{G}'(\mathbb{C})}(u)$  as in the proof of [BV85, Proposition 2.4]. Then any maximal reductive subgroup of  $Z_{\mathcal{G}'(\mathbb{C})}(u)$  is conjugate to  $Z_{\mathcal{G}'(\mathbb{C})}(\text{Im } \theta)$  by [Hoc81, VIII. Theorem 4.3] (cf. [Mos56, Theorem 7.1]). Hence, by replacing  $\theta$  by its conjugate under  $Z_{\mathcal{G}'(\mathbb{C})}(u)$ , we may assume that  $(s, q^{-\frac{1}{2}}) \in Z_{\mathcal{G}'(\mathbb{C})}(\text{Im } \theta)$ . Then  $\theta$  satisfies the conditions in the first claim.

The second claim is proved in the same way as [KL87, 2.4 (h)].  $\square$

Let  $\phi: \mathrm{SL}_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$  be an L-parameter of  $\mathrm{SL}_2$ -type for  $G$ . We define  $\xi_\phi: \mathrm{WD}_F(\mathbb{C}) \rightarrow {}^L G(\mathbb{C})$  by

$$\xi_\phi(a, w) = \phi \left( \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{-\frac{1}{2}} \end{pmatrix}, w \right).$$

We define  $\Xi: \Phi^{\mathrm{SL}}(G) \rightarrow \Phi_{\mathbb{C}}^{\mathrm{WD}}(G)$  by  $\Xi([\phi]) = [\xi_\phi]$ .

**Proposition 1.13.** *The map  $\Xi$  is a bijection.*

*Proof.* Let  $\xi$  be a Weil–Deligne L-parameter for  $G$  over  $\mathbb{C}$ . Let  $\mathcal{H}$  be the centralizer of  $\xi(I_F)$  in  $\widehat{G}$ . Then  $\mathcal{H}$  is reductive by Lemma 1.11. Take a lift  $\sigma_q \in W_F$  of the  $q$ -th power Frobenius element. Then  $\mathcal{H}$  is stable under conjugation by  $\xi(\sigma_q)$ , since  $I_F$  is a normal subgroup of  $W_F$ . Take a positive integer  $m_0$  such that  $\sigma_q^{m_0}$  commutes with  $\widehat{G}$  in  ${}^L G$  and

$$(p_{\widehat{G}} \circ \xi)(0, \sigma_q^{m_0} \sigma \sigma_q^{-m_0}) = (p_{\widehat{G}} \circ \xi)(0, \sigma)$$

for any  $\sigma \in I_F$ . Then  $(1, \sigma_q^{m_0})^{\mathbb{Z}}$  is a normal subgroup of  $\mathcal{H} \cdot \xi(\sigma_q)^{\mathbb{Z}} \subset {}^L G$ . We put

$$\mathcal{G} = (\mathcal{H} \cdot \xi(\sigma_q)^{\mathbb{Z}}) / (1, \sigma_q^{m_0})^{\mathbb{Z}}.$$

Then  $\mathcal{G}$  is a reductive group, because the identity component of  $\mathcal{G}$  is equal to the identity component of  $\mathcal{H}$ . We view  $\mathcal{H}$  as an algebraic subgroup of  $\mathcal{G}$ . Let  $s$  be the image of  $\xi(0, \sigma_q)$  in  $\mathcal{G}(\mathbb{C})$ . Since  $\xi$  is semisimple, the element  $s$  is semisimple. Let  $u \in \widehat{G}(\mathbb{C})$  be the image of  $1 \in \mathbb{G}_a(\mathbb{C})$  under  $\xi$ . Then  $u$  belongs to  $\mathcal{H}(\mathbb{C})$ . Hence we can view  $u$  as an element of  $\mathcal{G}(\mathbb{C})$ . By Lemma 1.12, there is a morphism  $\theta: \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathcal{G}(\mathbb{C})$  such that

$$\theta \left( \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) = u, \quad \theta \left( \begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right) s^{-1} \in Z_{\mathcal{G}(\mathbb{C})}(\mathrm{Im} \theta). \quad (1.1)$$

Since  $\theta$  factors through the identity component of  $\mathcal{G}(\mathbb{C})$ , it factors through  $\mathcal{H}(\mathbb{C})$ . Hence  $\theta$  determines a morphism  $\theta_{\widehat{G}}: \mathrm{SL}_2(\mathbb{C}) \rightarrow \widehat{G}(\mathbb{C})$ . We note that

$$\theta \left( \begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right)$$

commutes with  $\xi(W_F)$ , because it commutes with  $s$  by the latter condition in (1.1) and we have  $\mathrm{Im} \theta_{\widehat{G}} \subset \mathcal{H}(\mathbb{C})$ . We define  $\phi_\xi: \mathrm{SL}_2(\mathbb{C}) \times W_F \rightarrow {}^L G(\mathbb{C})$  by  $\phi_\xi|_{\mathrm{SL}_2(\mathbb{C})} = \theta_{\widehat{G}}$  and

$$\phi_\xi(1, w) = \theta_{\widehat{G}} \left( \begin{pmatrix} |w|^{-\frac{1}{2}} & 0 \\ 0 & |w|^{\frac{1}{2}} \end{pmatrix} \right) \xi(0, w).$$

Let  $\theta'$  be another choice of  $\theta$ . Then  $\theta' = \mathrm{Ad}(g')\theta$  for some  $g' \in Z_{\mathcal{G}(\mathbb{C})}(\{s, u\})$  by Lemma 1.12. Since we have

$$\theta \left( \begin{pmatrix} q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} \end{pmatrix} \right) s^{-1} \in Z_{\mathcal{G}(\mathbb{C})}(\mathrm{Im} \theta) \cap Z_{\mathcal{G}(\mathbb{C})}(\{s, u\})$$

by (1.1), we may replace  $g'$  and assume that  $g' \in Z_{\mathcal{H}(\mathbb{C})}(\{s, u\})$ . Then  $g'$  commutes with  $\xi(W_F)$ . Hence  $[\phi_\xi]$  is independent of the choice of  $\theta$ . We can see that  $\xi \mapsto [\phi_\xi]$  induces the inverse of  $\Xi$ .  $\square$

### 1.3 Over $\overline{\mathbb{Q}_\ell}$

Let  $\ell$  be a prime number different from  $p$ . Assume that  $C = \overline{\mathbb{Q}_\ell}$  in this subsection.

**Definition 1.14.** (1) An  $\ell$ -adic  $L$ -homomorphism for  $G$  is a continuous homomorphism

$$\varphi: W_F \rightarrow {}^L G(\overline{\mathbb{Q}_\ell})$$

of groups over  $W_F$ . We say that an  $\ell$ -adic  $L$ -homomorphism  $\varphi$  for  $G$  is Frobenius-semisimple if  $\varphi(\sigma_q)$  is semisimple for any lift  $\sigma_q \in W_F$  of the  $q$ -th power Frobenius element.

(2) An  $\ell$ -adic  $L$ -parameter for  $G$  is an Frobenius-semisimple relevant  $\ell$ -adic  $L$ -homomorphism for  $G$ .

Let  $\mathcal{L}_\ell(G)$  denote the set of the equivalence classes of  $\ell$ -adic  $L$ -homomorphisms. Let  $\Phi_\ell(G)$  denote the set of the equivalence classes of  $\ell$ -adic  $L$ -parameters of  $G$ .

Let  $t_\ell: I_F \rightarrow \mathbb{Z}_\ell(1)$  be the  $\ell$ -adic tame character. We take an isomorphism  $\mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$  and let

$$t'_\ell: I_F \xrightarrow{t_\ell} \mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$$

Let  $\xi$  be an  $L$ -homomorphism of Weil–Deligne type for  $G$  over  $\overline{\mathbb{Q}_\ell}$ . We take a lift  $\sigma_q \in W_F$  of the  $q$ -th power Frobenius element and define  $\varphi_\xi: W_F \rightarrow {}^L G(\overline{\mathbb{Q}_\ell})$  by

$$\varphi_\xi(\sigma_q^m \sigma) = \xi(t'_\ell(\sigma), \sigma_q^m \sigma)$$

for  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ .

**Lemma 1.15.** The equivalence class  $[\varphi_\xi] \in \mathcal{L}_\ell(G)$  of  $\varphi_\xi$  constructed above is independent of choices of  $\sigma_q$  and an isomorphism  $\mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$ .

*Proof.* Let  $\sigma'_q$  be another choice of a lift of the  $q$ -th power Frobenius element. Then  $\sigma'_q = \sigma_q \sigma'$  for some  $\sigma' \in I_F$ . We define  $\varphi'_\xi$  similarly as  $\varphi_\xi$  using  $\sigma'_q$  instead of  $\sigma_q$ . We put

$$g = \xi\left(\frac{t'_\ell(\sigma')}{q-1}, 1\right).$$

Then we have

$$\text{Ad}(g)(\varphi_\xi(\sigma)) = \xi(t'_\ell(\sigma), \sigma) = \varphi'_\xi(\sigma)$$

for  $\sigma \in I_F$ , and

$$\text{Ad}(g)(\varphi_\xi(\sigma_q)) = g(\text{Ad}(\varphi_\xi(\sigma_q))(g^{-1})) \varphi_\xi(\sigma_q) = \xi(-t'_\ell(\sigma'), \sigma_q) = \varphi'_\xi(\sigma_q).$$

Hence we have  $[\varphi_\xi] = [\varphi'_\xi]$ .

Let

$$t''_\ell: I_F \xrightarrow{t_\ell} \mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$$

be a homomorphism obtained from another choice of an isomorphism  $\mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$ . Then we have  $t''_\ell = ut'_\ell$  for some  $u \in \mathbb{Z}_\ell^\times$ . Take a positive integer  $m_0$  such that  $\sigma_q^{m_0}$  commutes with  $\widehat{G}$  in  ${}^L G$  and

$$(p_{\widehat{G}} \circ \xi)(0, \sigma_q^{m_0} \sigma \sigma_q^{-m_0}) = (p_{\widehat{G}} \circ \xi)(0, \sigma) \quad (1.2)$$

for any  $\sigma \in I_F$ . We put  $h_0 = (p_{\widehat{G}} \circ \xi)(0, \sigma_q^{m_0})$ . If  $\xi|_{\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)}$  is trivial, there is nothing to prove. Hence we assume that  $\xi|_{\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)}$  is non-trivial. Let  $U_\xi$  be the algebraic subgroup of  $\widehat{G}$  defined by  $\xi(\mathbb{G}_a(\overline{\mathbb{Q}}_\ell))$ . Let  $\mathcal{H}$  be the intersection of the normalizer of  $U_\xi$  in  $\widehat{G}$  and the centralizer of  $\xi(W_F)$  in  $\widehat{G}$ . We have

$$\mathrm{Ad}(h_0)(\xi(a, 1)) = \mathrm{Ad}(\xi(\sigma_q^{m_0}))(\xi(a, 1)) = \xi(q^{m_0}a, 1)$$

for  $a \in \mathbb{G}_a(\overline{\mathbb{Q}}_\ell)$  and

$$\begin{aligned} \mathrm{Ad}(h_0)(\xi(0, \sigma_q)) &= \mathrm{Ad}((1, \sigma_q^{-m_0})) \mathrm{Ad}(\xi(0, \sigma_q^{m_0}))(\xi(0, \sigma_q)) = \xi(0, \sigma_q), \\ \mathrm{Ad}(h_0)(\xi(0, \sigma)) &= \mathrm{Ad}((1, \sigma_q^{-m_0}))(\xi(0, \sigma_q^{m_0} \sigma \sigma_q^{-m_0})) = \xi(0, \sigma) \end{aligned}$$

for  $\sigma \in I_F$  using (1.2). Hence we have  $h_0 \in \mathcal{H}(\overline{\mathbb{Q}}_\ell)$ . The morphism

$$f: \mathcal{H} \rightarrow \mathrm{Aut}(U_\xi) \simeq \mathbb{G}_m$$

induced by the adjoint action is surjective, because  $f(h_0) = q^{m_0}$  is not of finite order. Hence we can take  $h \in \mathcal{H}(\overline{\mathbb{Q}}_\ell)$  such that  $f(h) = u$ . Then we have

$$\mathrm{Ad}(h) (\xi(t'_\ell(\sigma), \sigma_q^m \sigma)) = (t''_\ell(\sigma), \sigma_q^m \sigma)$$

for  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ . Therefore  $[\varphi_\xi]$  is independent of a choice an isomorphism  $\mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$ .  $\square$

We define  $\Theta: \mathcal{L}_{\overline{\mathbb{Q}}_\ell}^{\mathrm{WD}}(G) \rightarrow \mathcal{L}_\ell(G)$  by  $\Theta([\xi]) = [\varphi_\xi]$ .

**Proposition 1.16.** *The map  $\Theta$  is a bijection. Further it induces a bijection  $\Phi_{\overline{\mathbb{Q}}_\ell}^{\mathrm{WD}}(G) \rightarrow \Phi_\ell(G)$ .*

*Proof.* Let  $\varphi$  be an  $\ell$ -adic L-homomorphism for  $G$ . Take a finite Galois extension  $F'$  of  $F$  such that  $G$  splits over  $F'$ . Take a representation

$$\eta: {}^L G(\overline{\mathbb{Q}}_\ell) \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_\ell)$$

which factors through a faithful algebraic representation

$$\overline{\eta}: \widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \mathrm{Gal}(F'/F) \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_\ell).$$

Applying Grothendieck's monodromy theorem (cf. [ST68, Appendix, Proposition]) to  $\eta \circ \varphi$ , we obtain a homomorphism

$$\xi_{\mathrm{GL}_n}: WD_F(\overline{\mathbb{Q}}_\ell) \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_\ell)$$

such that  $\xi_{\mathrm{GL}_n}|_{\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)}$  is algebraic,  $\xi_{\mathrm{GL}_n}|_{W_F}$  is smooth and

$$\xi_{\mathrm{GL}_n}(t'_\ell(\sigma), \sigma_q^m \sigma) = (\eta \circ \varphi)(\sigma_q^m \sigma)$$

for  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ . Take a finite separable extension  $F''$  of  $F'$  such that  $\xi_{\mathrm{GL}_n}|_{I_{F''}}$  is trivial. Since  $t'_\ell(I_{F''})$  is Zariski dense in  $\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)$ , the algebraic morphism  $\xi_{\mathrm{GL}_n}|_{\mathbb{G}_a(\overline{\mathbb{Q}}_\ell)}$  factors through the inclusion

$$\widehat{G}(\overline{\mathbb{Q}}_\ell) \hookrightarrow \widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \mathrm{Gal}(F'/F) \xrightarrow{\overline{\eta}} \mathrm{GL}_n(\overline{\mathbb{Q}}_\ell)$$



via an algebraic morphism  $\alpha: \mathbb{G}_a(\overline{\mathbb{Q}}_\ell) \rightarrow \widehat{G}(\overline{\mathbb{Q}}_\ell)$ . We define a homomorphism

$$\xi_\varphi: WD_F(\overline{\mathbb{Q}}_\ell) \rightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$$

by

$$\xi_\varphi(a, \sigma_q^m \sigma) = \alpha(a - t'_\ell(\sigma)) \varphi(\sigma_q^m \sigma)$$

for  $a \in \mathbb{G}_a(\overline{\mathbb{Q}}_\ell)$ ,  $m \in \mathbb{Z}$  and  $\sigma \in I_F$ . Then  $\varphi \mapsto \xi_\varphi$  induces the inverse of  $\Theta$ . The bijection  $\Theta$  induces a bijection  $\Phi_{\overline{\mathbb{Q}}_\ell}^{\text{WD}}(G) \rightarrow \Phi_\ell(G)$  by Lemma 1.2 and Lemma 1.15.  $\square$

**Corollary 1.17.** *Let  $\sigma_q \in W_F$  be a lift of the  $q$ -th power Frobenius element. Then an  $\ell$ -adic  $L$ -homomorphism  $\varphi$  for  $G$  is Frobenius-semisimple if  $\varphi(\sigma_q)$  is semisimple.*

*Proof.* By Proposition 1.16, we take an  $L$ -homomorphism  $\xi$  of Weil–Deligne type for  $G$  over  $\overline{\mathbb{Q}}_\ell$  such that  $[\varphi] = [\varphi_\xi]$ , where  $\varphi_\xi$  is defined using  $\sigma_q$ . Then  $\xi|_{W_F}$  is semisimple by Lemma 1.2, because  $\xi(0, \sigma_q) = \varphi_\xi(\sigma_q)$  is semisimple. Let  $\sigma'_q \in W_F$  be another lift of the  $q$ -th power Frobenius element. We define  $\varphi'_\xi$  using  $\sigma'_q$ . Then  $[\varphi] = [\varphi'_\xi]$  by Lemma 1.15. Hence  $\varphi(\sigma'_q)$  is semisimple, because  $\varphi'_\xi(\sigma'_q) = \xi(0, \sigma'_q)$  is semisimple.  $\square$

## 2 Local Langlands correspondence

### 2.1 Problem

Let  $\text{Irr}(G(F))$  denote the set of isomorphism classes of irreducible smooth representations of  $G(F)$  over  $\mathbb{C}$ . The local Langlands correspondence is a surjective map

$$\text{LL}_G: \text{Irr}(G(F)) \rightarrow \Phi(G)$$

with finite fibers satisfying various properties (*cf.* [Bor79, 10], [Kal16, Conjecture G]). We assume the existence of the local Langlands correspondence in the sequel.

Let  $\text{Irr}_\ell(G(F))$  denote the set of isomorphism classes of irreducible smooth representations of  $G(F)$  over  $\overline{\mathbb{Q}}_\ell$ . If we fix an isomorphism  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$ , we have a surjection

$$\text{LL}_{G,\ell}^\iota: \text{Irr}_\ell(G(F)) \rightarrow \Phi_\ell(G)$$

sending  $[\pi] \in \text{Irr}_\ell(G(F))$  to the image of  $\text{LL}_G([\pi \otimes_{\overline{\mathbb{Q}}_\ell, \iota^{-1}} \mathbb{C}])$  under the bijection

$$\Phi(G) \simeq \Phi^{\text{SL}}(G) \simeq \Phi_{\mathbb{C}}^{\text{WD}}(G) \xrightarrow{\Phi_\iota^{\text{WD}}(G)} \Phi_{\overline{\mathbb{Q}}_\ell}^{\text{WD}}(G) \simeq \Phi_\ell(G),$$

where  $\Phi_\iota^{\text{WD}}(G)$  is a bijection induced by  $\iota$  as in Remark 1.5. However,  $\text{LL}_{G,\ell}^\iota$  depends on the choice of an isomorphism  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$ .

In [BH06, 35.1], an  $\ell$ -adic local Langlands correspondence for  $\text{GL}_2$  is constructed. We recall the construction here. For  $[\phi] \in \Phi(\text{GL}_2)$ , we define  $[\widetilde{\phi}] \in \Phi(\text{GL}_2)$  by

$$\widetilde{\phi}(g, w) = \left( \begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{\frac{1}{2}} \end{pmatrix}, 1 \right) \phi(g, w).$$

We define a bijection  $\Lambda: \Phi(\text{GL}_2) \rightarrow \Phi(\text{GL}_2)$  by  $[\phi] \mapsto [\widetilde{\phi}]$ . We define

$$\text{LL}_{\text{GL}_2,\ell}: \text{Irr}_\ell(\text{GL}_2(F)) \rightarrow \Phi_\ell(\text{GL}_2)$$

by sending  $[\pi] \in \text{Irr}_\ell(\text{GL}_2(F))$  to the image of  $\text{LL}_{\text{GL}_2}([\pi \otimes_{\overline{\mathbb{Q}_\ell, \iota^{-1}} \mathbb{C}}])$  under the bijection

$$\Phi(\text{GL}_2) \xrightarrow{\Lambda} \Phi(\text{GL}_2) \simeq \Phi^{\text{SL}}(\text{GL}_2) \simeq \Phi_{\mathbb{C}}^{\text{WD}}(\text{GL}_2) \xrightarrow{\Phi_{\iota}^{\text{WD}}(G)} \Phi_{\overline{\mathbb{Q}_\ell}}^{\text{WD}}(\text{GL}_2) \simeq \Phi_\ell(\text{GL}_2)$$

using an isomorphism  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}_\ell}$ . Then  $\text{LL}_{\text{GL}_2, \ell}$  is independent of the choice of  $\iota$ .

We can not make a similar twist for an L-parameter of a general connected reductive group  $G$  as the following example suggests.

**Example 2.1.** *We have a commutative diagram*

$$\begin{array}{ccc} \text{Irr}(\text{PGL}_2(F)) & \xrightarrow{\text{LL}_{\text{PGL}_2}} & \Phi(\text{PGL}_2) \\ \downarrow & & \downarrow \\ \text{Irr}(\text{GL}_2(F)) & \xrightarrow{\text{LL}_{\text{GL}_2}} & \Phi(\text{GL}_2) \end{array}$$

by functoriality. On the other hand, there does not exist a map

$$\text{LL}_{\text{PGL}_2, \ell}: \text{Irr}_\ell(\text{PGL}_2(F)) \rightarrow \Phi_\ell(\text{PGL}_2)$$

which makes the commutative diagram

$$\begin{array}{ccc} \text{Irr}_\ell(\text{PGL}_2(F)) & \xrightarrow{\text{LL}_{\text{PGL}_2, \ell}} & \Phi_\ell(\text{PGL}_2) \\ \downarrow & & \downarrow \\ \text{Irr}_\ell(\text{GL}_2(F)) & \xrightarrow{\text{LL}_{\text{GL}_2, \ell}} & \Phi_\ell(\text{GL}_2), \end{array}$$

because  $(\det \circ \widehat{\text{op}}_{\text{GL}_2} \circ \varphi)(w) = |w|$  for  $w \in W_F$  if  $[\varphi] \in \Phi_\ell(\text{GL}_2)$  is the image under  $\text{LL}_{\text{GL}_2, \ell}$  of an element of  $\text{Irr}_\ell(\text{GL}_2(F))$  coming from  $\text{Irr}_\ell(\text{PGL}_2(F))$ .

## 2.2 $\ell$ -adic C-parameter

A C-group is constructed in [BG14, Definition 5.38] for a connected reductive group over a number field. We recall the construction here in our setting. Let  $G^{\text{ad}}$  be the adjoint quotient of  $G$ , and  $G^{\text{sc}}$  be the simply-connected cover of  $G^{\text{ad}}$ . Let  $\gamma: Z(G^{\text{sc}}) \rightarrow \mathbb{G}_m$  be the restriction to  $Z(G^{\text{sc}})$  of the half sum of the positive roots of  $G^{\text{sc}}$ , where we take a maximal torus  $T^{\text{sc}}$  and a Borel subgroup  $B^{\text{sc}}$  of  $G_{\overline{F}}^{\text{sc}}$  such that  $T^{\text{sc}} \subset B^{\text{sc}}$  to define the the positive roots, but  $\gamma$  is independent of the choice. By pushing forward the exact sequence

$$1 \rightarrow Z(G^{\text{sc}}) \rightarrow G^{\text{sc}} \rightarrow G^{\text{ad}} \rightarrow 1$$

by  $\gamma$ , we obtain an extension

$$1 \rightarrow \mathbb{G}_m \rightarrow G^1 \rightarrow G^{\text{ad}} \rightarrow 1.$$

By taking the pullback of this extension along the natural morphism  $G \rightarrow G^{\text{ad}}$ , we obtain an extension

$$1 \rightarrow \mathbb{G}_m \rightarrow \tilde{G} \rightarrow G \rightarrow 1.$$

We define the C-group  ${}^C G$  of  $G$  as the L-group  ${}^L \tilde{G}$  of  $\tilde{G}$ .

The character

$$G^{\text{sc}} \times \mathbb{G}_m \rightarrow \mathbb{G}_m; (g, z) \mapsto z^2$$

induces a character  $G^1 \rightarrow \mathbb{G}_m$ , since  $\gamma^2 = 1$  by the construction of  $\gamma$ . It further induces a character  $\tilde{G} \rightarrow \mathbb{G}_m$  by taking composition with the natural morphism  $\tilde{G} \rightarrow G^1$ . Hence we obtain a morphism

$$\tilde{G} \rightarrow G \times \mathbb{G}_m. \quad (2.1)$$

We take a Borel subgroup  $B \subset G_{\overline{F}}$  and a maximal torus  $T \subset B$  over  $\overline{F}$ . Let  $\rho_G$  denote the half sum of positive roots of  $G$  with respect to  $T$  and  $B$ . Then  $2\rho_G$  defines a cocharacter  $\delta_G: \mathbb{G}_m \rightarrow \hat{T}$ . We put

$$z_G = \delta_G(-1).$$

Then  $z_G$  is central in  $\hat{G}$  and independent of choices of  $B$  and  $T$  as in [BG14, Proposition 5.39]. By the independent of choices, we see that  $z_G \in Z(\hat{G})^{\Gamma_F} \subset {}^L G$ .

Then the morphism (2.1) induces the isomorphism

$$(\hat{G} \times \mathbb{G}_m / \langle (z_G, -1) \rangle) \rtimes W_F \simeq {}^C G \quad (2.2)$$

as in [BG14, Proposition 5.39]. We sometimes express a point of  ${}^C G$  as  $[(g, z, w)]$  using the above isomorphism. We define  $t_{\mathbb{G}_m}: {}^C G \rightarrow \mathbb{G}_m$  by  $t_{\mathbb{G}_m}([(g, z, w)]) = z^2$ . We have an exact sequence

$$1 \rightarrow {}^L G \rightarrow {}^C G \xrightarrow{t_{\mathbb{G}_m}} \mathbb{G}_m \rightarrow 1.$$

**Definition 2.2.** An  $\ell$ -adic  $C$ -parameter for  $G$  is an  $\ell$ -adic  $L$ -parameter  $\varphi$  for  $\tilde{G}$  such that  $(t_{\mathbb{G}_m} \circ \varphi)(w) = q^{d_F(w)}$ .

Let  $\Phi_\ell^C(G)$  denote the equivalence classes of  $\ell$ -adic  $C$ -parameters for  $G$ . We take  $c \in \overline{\mathbb{Q}_\ell}$  such that  $c^2 = q$ . We define

$$i_c: {}^L G \rightarrow {}^C G$$

by  $i_c(g, w) = [(g, c^{d_F(w)}, w)]$ . For an  $\ell$ -adic  $L$ -parameter  $\varphi$  for  $G$ , we put

$$\varphi_c = i_c \circ \varphi.$$

**Lemma 2.3.** We have a bijection between the set of the  $\ell$ -adic  $L$ -parameters for  $G$  and the set of the  $\ell$ -adic  $C$ -parameters for  $G$  given by sending  $\varphi$  to  $\varphi_c$ . Further, this induces a bijection  $\Phi_\ell(G) \simeq \Phi_\ell^C(G)$ .

*Proof.* The first claim follows from the definitions. If two  $\ell$ -adic  $L$ -parameters for  $\tilde{G}$  are conjugate by an element of  $\hat{\tilde{G}}(\overline{\mathbb{Q}_\ell})$ , then they are conjugate by an element of  $\hat{G}(\overline{\mathbb{Q}_\ell})$ , since  $\hat{\tilde{G}}(\overline{\mathbb{Q}_\ell})$  is generated by  $\hat{G}(\overline{\mathbb{Q}_\ell})$  and  $Z_{\hat{\tilde{G}}(\overline{\mathbb{Q}_\ell})}({}^C G(\overline{\mathbb{Q}_\ell}))$ . Hence the second claim follows from the first one.  $\square$

We take an isomorphism  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}_\ell}$ . We define a map

$$\mathcal{C}_{G,\ell}^\iota: \Phi_\ell(G) \rightarrow \Phi_\ell^C(G)$$

by sending  $[\varphi]$  to  $[\varphi_c]$  where  $c = \iota(q^{\frac{1}{2}})$ . Then  $\mathcal{C}_{G,\ell}^\iota$  is a bijection by Lemma 2.3.

We define

$$\text{LL}_{G,\ell}^{C,\iota}: \text{Irr}_\ell(G(F)) \rightarrow \Phi_\ell^C(G)$$

as  $\mathcal{C}_{G,\ell}^\iota \circ \text{LL}_{G,\ell}^\iota$ .

**Conjecture 2.4.** The map  $\text{LL}_{G,\ell}^{C,\iota}$  is independent of a choice of  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}_\ell}$ .

### 2.3 Tannakian $\ell$ -adic L-parameter

Assume that  $C = \overline{\mathbb{Q}}_\ell$ . For a topological group  $H$ , let  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}(H)$  be the category of continuous finite dimensional representations of  $H$  over  $\overline{\mathbb{Q}}_\ell$ . For an algebraic group  $H$  over a field, a character  $\chi$  of  $H$  and a cocharacter  $\mu$  of  $H$ , we define  $\langle \chi, \mu \rangle_H \in \mathbb{Z}$  by

$$(\chi \circ \mu)(z) = z^{\langle \chi, \mu \rangle_H}.$$

For a cocharacter  $\mu \in X_*(T)$  of a torus  $T$  over  $\overline{F}$ , let  $\hat{\mu} \in X^*(\hat{T})$  denote the corresponding character of the dual torus  $\hat{T}$ .

Let  $\mathcal{M}_G$  be the conjugacy classes of cocharacters  $\mathbb{G}_m \rightarrow G_{\overline{F}}$ . Let  $[\mu] \in \mathcal{M}_G$ . We put

$$d_G([\mu]) = \langle 2\rho_G, \mu \rangle_T,$$

where we take a Borel subgroup  $B \subset G_{\overline{F}}$ , a maximal torus  $T \subset B$  defined over  $\overline{F}$  and a dominant representative  $\mu \in X_*(T)$ . Let  $E_{[\mu]}$  be the field of definition of  $[\mu]$ . Let  $r_{\hat{G}, [\mu]}$  be the irreducible representation of  $\hat{G}(\overline{\mathbb{Q}}_\ell)$  of highest weight  $\hat{\mu}$  viewed as a dominant character of a maximal torus of  $\hat{G}$ .

We take  $c \in \overline{\mathbb{Q}}_\ell$  such that  $c^2 = q$ . For an integer  $m$ , let

$$\left(\frac{m}{2}\right)_c$$

denote the twist by the character  $W_F \rightarrow \overline{\mathbb{Q}}_\ell$  sending  $w$  to  $c^{md_F(w)}$ .

Let  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G)$  denote the category of continuous finite dimensional representations of  ${}^L G(\overline{\mathbb{Q}}_\ell)$  over  $\overline{\mathbb{Q}}_\ell$  whose restrictions to  $\hat{G}(\overline{\mathbb{Q}}_\ell)$  are algebraic. Let  $r: {}^L G(\overline{\mathbb{Q}}_\ell) \rightarrow \text{Aut}(V)$  be an object in  $\text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G)$ . Then we have a decomposition

$$V = \bigoplus_{[\mu] \in \mathcal{M}_G} V_{[\mu]}$$

as representations of  $\hat{G}(\overline{\mathbb{Q}}_\ell)$  where  $V_{[\mu]}$  is the  $r_{\hat{G}, [\mu]}$ -typic part of  $V$ . For an  $\ell$ -adic L-parameter  $\varphi$ , we define  $(r \circ \varphi)_c: W_F \rightarrow \text{Aut}(V)$  by

$$V = \bigoplus_{[\mu] \in \mathcal{M}_G} V_{[\mu]} \left( \frac{d_G([\mu])}{2} \right)_c,$$

which means that we twist  $r \circ \varphi: W_F \rightarrow \text{Aut}(V)$  by

$$\left( \frac{d_G([\mu])}{2} \right)_c$$

on each direct summand  $V_{[\mu]}$ . This is well-defined, because  $d_G(w[\mu]) = d_G([\mu])$  for  $w \in W_F$ .

For an  $\ell$ -adic L-parameter  $\varphi$  for  $G$ , we define a tensor functor

$$\mathcal{F}_{\varphi, c}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$$

by

$$\mathcal{F}_{\varphi, c}(r) = (r \circ \varphi)_c.$$

**Definition 2.5.** A Tannakian  $\ell$ -adic L-parameter for  $G$  is a functor

$$\mathcal{F}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_F)$$

which is equal to  $\mathcal{F}_{\varphi,c}$  for an  $\ell$ -adic L-parameter  $\varphi$  for  $G$ . We say that two Tannakian  $\ell$ -adic L-parameters  $\mathcal{F}$  and  $\mathcal{F}'$  are equivalent if there is  $g \in \widehat{G}(\overline{\mathbb{Q}}_\ell)$  such that, for all  $r \in \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G)$ , we have  $\mathcal{F}(r) = \mathcal{F}'(r)^{r(g)}$ .

**Lemma 2.6.** The set of the Tannakian  $\ell$ -adic L-parameters for  $G$  is independent of a choice of  $c \in \overline{\mathbb{Q}}_\ell$  such that  $c^2 = q$ .

*Proof.* We have

$$\widehat{\mu}(z_G) = (-1)^{\langle \widehat{\mu}, \delta_G \rangle_{\widehat{T}}} = (-1)^{d_G([\mu])} \quad (2.3)$$

for  $\mu \in X_*(T)$ . Let

$$\omega_{z_G}: W_F \rightarrow Z(\widehat{G})^{\Gamma_F}(\overline{\mathbb{Q}}_\ell) \hookrightarrow {}^L G(\overline{\mathbb{Q}}_\ell)$$

be the character sending  $w$  to  $z_G^{d_F(w)}$ . By (2.3), we have

$$(r \circ \varphi)_{-c} = (r \circ (\omega_{z_G} \varphi))_c \quad (2.4)$$

for an  $\ell$ -adic L-parameter  $\varphi$  and  $r \in \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G)$ . Since  $\omega_{z_G} \varphi$  is also an  $\ell$ -adic L-parameter, the claim follows.  $\square$

Let  $\Phi_\ell^T(G)$  be the set of equivalence classes of Tannakian  $\ell$ -adic L-parameters for  $G$ . This set is independent of a choice of  $c$  by Lemma 2.6.

**Lemma 2.7.** We have a bijection between the set of the  $\ell$ -adic L-parameters for  $G$  and the set of the Tannakian  $\ell$ -adic L-parameters for  $G$  given by sending  $\varphi$  to  $\mathcal{F}_{\varphi,c}$ . Further, this induces a bijection  $\Phi_\ell(G) \simeq \Phi_\ell^T(G)$ .

*Proof.* We show the first claim. The map is surjective by Definition 2.5. Assume that  $\varphi$  and  $\varphi'$  are different  $\ell$ -adic L-parameters for  $G$  and  $\mathcal{F}_{\varphi,c} = \mathcal{F}_{\varphi',c}$ . We take  $w \in W_F$  such that  $\varphi(w) \neq \varphi'(w)$ . Further, we take a finite Galois extension  $F'$  of  $F$  such that  $G$  splits over  $F'$  and the images of  $\varphi(w)$  and  $\varphi'(w)$  in  $\widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \text{Gal}(F'/F)$  are different. By considering a representation of  ${}^L G(\overline{\mathbb{Q}}_\ell)$  which factors through a faithful algebraic representation of  $\widehat{G}(\overline{\mathbb{Q}}_\ell) \rtimes \text{Gal}(F'/F)$ , we have a contradiction to  $\mathcal{F}_{\varphi,c} = \mathcal{F}_{\varphi',c}$ . Hence the map is injective.

The second claim follows from the first one.  $\square$

For a Tannakian  $\ell$ -adic L-parameter  $\mathcal{F}$  for  $G$ , we take an  $\ell$ -adic L-parameter  $\varphi$  for  $G$  such that  $\mathcal{F} = \mathcal{F}_{\varphi,c}$ . Then the centralizer  $S_\varphi = Z_{\widehat{G}(\overline{\mathbb{Q}}_\ell)}(\text{Im } \varphi)$  is independent of a choice of  $c$  by (2.4). We write  $S_\mathcal{F}$  for  $S_\varphi$ . Then  $\mathcal{F}$  naturally factors through

$$\mathcal{F}_S: \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_\mathcal{F} \times W_F).$$

For a finite separable extension  $F'$  of  $F$ , we define the restriction

$$\mathcal{F}|_{F'}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G_{F'}) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(W_{F'})$$

of  $\mathcal{F}$  to  $F'$  by the usual restriction to  $W_{F'}$  of an  $\ell$ -adic L-parameter for  $G$  and bijections given by Lemma 2.7. Let

$$\mathcal{F}_S|_{F'}: \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G_{F'}) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_{\mathcal{F}} \times W_{F'})$$

be the composition of

$$(\mathcal{F}|_{F'})_S: \text{Rep}_{\overline{\mathbb{Q}}_\ell}^{\text{alg}}({}^L G_{F'}) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_{\mathcal{F}|_{F'}} \times W_{F'})$$

and the natural functor

$$\text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_{\mathcal{F}|_{F'}} \times W_{F'}) \rightarrow \text{Rep}_{\overline{\mathbb{Q}}_\ell}(S_{\mathcal{F}} \times W_{F'})$$

induced by the restriction with respect to  $S_{\mathcal{F}} \subset S_{\mathcal{F}|_{F'}}$ .

We take an isomorphism  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$ . We define a map

$$\mathcal{T}_{G,\ell}^\iota: \Phi_\ell(G) \rightarrow \Phi_\ell^\text{T}(G)$$

by sending  $[\varphi]$  to  $[F_{\varphi,c}]$  where  $c = \iota(q^{\frac{1}{2}})$ . Then  $\mathcal{T}_{G,\ell}^\iota$  is a bijection by Lemma 2.7.

We define

$$\text{LL}_{G,\ell}^{\text{T},\iota}: \text{Irr}_\ell(G(F)) \rightarrow \Phi_\ell^\text{T}(G)$$

as  $\mathcal{T}_{G,\ell}^\iota \circ \text{LL}_{G,\ell}^\iota$ .

**Conjecture 2.8.** *The map  $\text{LL}_{G,\ell}^{\text{T},\iota}$  is independent of a choice of  $\iota: \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$ .*

**Remark 2.9.** *Conjecture 2.8 is motivated by the Kottwitz conjecture for local Shimura varieties in [RV14, Conjecture 7.4]. We use notations in [RV14, Conjecture 7.4]. Let  $[\pi] \in \text{Irr}_\ell(G)$  and  $[\rho] \in \text{Irr}_\ell(J)$ . Assume that*

$$[\mathcal{F}^\iota] = \text{LL}_{G,\ell}^{\text{T},\iota}([\pi]) = \text{LL}_{J,\ell}^{\text{T},\iota}([\rho])$$

for a Tannakian  $\ell$ -adic L-parameter  $\mathcal{F}^\iota$ . Let  $\delta_{\pi,\rho}^\iota$  be the representation of  $S_{\mathcal{F}^\iota}$  over  $\overline{\mathbb{Q}}_\ell$  determined by  $\iota$  and  $\check{\tau}_{\pi'} \otimes \tau_{\rho'}$  constructed in [RV14, p. 312] (cf. [KW17, 2.3] for a construction in a more general case), where  $\pi' = \pi \otimes_{\overline{\mathbb{Q}}_\ell, \iota^{-1}} \mathbb{C}$  and  $\rho' = \rho \otimes_{\overline{\mathbb{Q}}_\ell, \iota^{-1}} \mathbb{C}$ . Let  $r_{[\mu]}$  be an extension of  $r_{\widehat{G},[\mu]}$  to  ${}^L G_{E_{[\mu]}}$  constructed by [Kot84a, (1.1.3), (2.1.2)] using [Bor79, 2.4 Remark (3)]. If the Kottwitz conjecture is true, then the isomorphism class of the  $W_{E_{[\mu]}}$ -representation

$$\text{Hom}_{S_{\mathcal{F}^\iota}}(\delta_{\pi,\rho}^\iota, \mathcal{F}_S|_{E_{[\mu]}}(r_{[\mu]}))$$

is independent of  $\iota$ , since the  $\ell$ -adic cohomology of local Shimura varieties and their group actions are independent of  $\iota$ .

## 2.4 Comparison

**Theorem 2.10.** *Conjecture 2.4 and Conjecture 2.8 are equivalent.*

*Proof.* Let  $\iota': \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}_\ell}$  be another choice. We put  $c' = \iota'(q^{\frac{1}{2}})$ . Let  $[\pi] \in \text{Irr}_\ell(G(F))$ . Assume that  $\text{LL}_{G,\ell}^\iota([\pi]) = [\varphi]$  and  $\text{LL}_{G,\ell}^{\iota'}([\pi]) = [\varphi']$ .

Assume that Conjecture 2.4 is true. Let  $r: {}^L G(\overline{\mathbb{Q}_\ell}) \rightarrow \text{Aut}(V)$  be an object in  $\text{Rep}_{\overline{\mathbb{Q}_\ell}}^{\text{alg}}({}^L G)$ . Then we have a decomposition

$$V = \bigoplus_{[\mu] \in \mathcal{M}_G} V_{[\mu]}$$

as representations of  $\widehat{G}(\overline{\mathbb{Q}_\ell})$  where  $V_{[\mu]}$  is the  $r_{\widehat{G},[\mu]}$ -typic part of  $V$ . We extend  $r$  to  $\tilde{r} \in \text{Rep}_{\overline{\mathbb{Q}_\ell}}^{\text{alg}}({}^C G)$  by letting  $\mathbb{G}_m$  act on  $V_{[\mu]}$  by  $z \mapsto z^{d_G([\mu])}$  using (2.2), where the extension is well-defined by (2.3). Then we have

$$\tilde{r} \circ \varphi_c = \tilde{r} \circ i_c \circ \varphi = \mathcal{F}_{\varphi,c}(r). \quad (2.5)$$

By Conjecture 2.4,  $\varphi_c$  and  $\varphi'_{c'}$  are conjugate by an element of  $\widehat{G}(\overline{\mathbb{Q}_\ell})$ . It implies that they are conjugate by an element of  $\widehat{G}(\overline{\mathbb{Q}_\ell})$ . Hence, Conjecture 2.8 is also true by (2.5).

Assume that Conjecture 2.8 is true. Then there is  $g \in \widehat{G}(\overline{\mathbb{Q}_\ell})$  such that  $\mathcal{F}_{\varphi,c}(r) = \mathcal{F}_{\varphi',c'}(r)^{r(g)}$  for all  $r \in \text{Rep}_{\overline{\mathbb{Q}_\ell}}^{\text{alg}}({}^L G)$ . Replacing  $\varphi'$  by  $\varphi'^g$ , we may assume that  $\mathcal{F}_{\varphi,c}(r) = \mathcal{F}_{\varphi',c'}(r)$  for all  $r \in \text{Rep}_{\overline{\mathbb{Q}_\ell}}^{\text{alg}}({}^L G)$ . Assume that  $\varphi_c \neq \varphi'_{c'}$ . We take an object  $\tilde{r}: {}^C G(\overline{\mathbb{Q}_\ell}) \rightarrow \text{Aut}(V)$  in  $\text{Rep}_{\overline{\mathbb{Q}_\ell}}^{\text{alg}}({}^C G)$  such that  $\tilde{r} \circ \varphi_c \neq \tilde{r} \circ \varphi'_{c'}$ .

Then we have a decomposition

$$V = \bigoplus_{[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}} V_{[\tilde{\mu}]}$$

as representations of  $\widehat{G}(\overline{\mathbb{Q}_\ell})$  where  $V_{[\tilde{\mu}]}$  is the  $r_{\widehat{G},[\tilde{\mu}]}$ -typic part of  $V$ . For  $[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}$ , let  $n_{[\tilde{\mu}]}$  be an integer such that

$$\widehat{\mu}([(1, z, 1)]) = z^{n_{[\tilde{\mu}]}},$$

where we view  $\widehat{\mu}$  as a character of a maximal torus of  $\widehat{G}$ . Then  $\tilde{r} \circ \varphi_c = \tilde{r} \circ i_c \circ \varphi$  is given by

$$V|_{{}^L G} = \bigoplus_{[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}} (V_{[\tilde{\mu}]}|_{{}^L G}) \left( \frac{n_{[\tilde{\mu}]}}{2} \right)_c. \quad (2.6)$$

Let  $r$  be the restriction of  $\tilde{r}$  to  ${}^L G$ . Then  $\mathcal{F}_{\varphi,c}(r)$  is given by

$$V|_{{}^L G} = \bigoplus_{[\tilde{\mu}] \in \mathcal{M}_{\widehat{G}}} (V_{[\tilde{\mu}]}|_{{}^L G}) \left( \frac{d_G([\mu])}{2} \right)_c, \quad (2.7)$$

where  $[\mu]$  is the image in  $\mathcal{M}_G$  of  $[\tilde{\mu}]$ . We have

$$n_{[\tilde{\mu}]} \equiv d_G([\mu]) \pmod{2}, \quad (2.8)$$

since

$$(-1)^{n_{[\tilde{\mu}]}} = \widehat{\mu}([(1, -1, 1)]) = \widehat{\mu}(z_G) = (-1)^{d_G([\mu])}$$

by (2.3). If we replace  $\iota$  by  $\iota'$ , then  $\mathcal{F}_{\varphi,c}(r)$  does not change, hence  $\tilde{r} \circ \varphi_c$  also does not change by (2.6), (2.7) and (2.8). This contradicts  $\tilde{r} \circ \varphi_c \neq \tilde{r} \circ \varphi'_{c'}$ . Hence Conjecture 2.4 is true.  $\square$

**Corollary 2.11.** *Conjecture 2.4 and Conjecture 2.8 are true for  $\mathrm{GL}_n$ .*

*Proof.* Recall that the local Langlands correspondence for  $\mathrm{GL}_n$  is known by [LRS93] and [HT01]. Assume that  $\mathrm{LL}_{\mathrm{GL}_n, \ell}^{\mathbb{C}, \iota}([\pi]) = [\varphi']$ . Note that  $z_{\mathrm{GL}_n} = (-1)^{n-1}$ , because

$$\delta_{\mathrm{GL}_n} : \mathbb{G}_m \rightarrow \widehat{T}_n \subset \widehat{\mathrm{GL}}_n; \quad z \mapsto \begin{pmatrix} z^{n-1} & 0 & \dots & 0 \\ 0 & z^{n-3} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & z^{1-n} \end{pmatrix},$$

where we define  $\delta_{\mathrm{GL}_n}$  using the diagonal maximal torus  $T_n$  of  $\mathrm{GL}_n$  and the Borel subgroup of the upper triangular matrices in  $\mathrm{GL}_n$ . Let

$$\tilde{r} : {}^C \mathrm{GL}_n \rightarrow \widehat{\mathrm{GL}}_n; \quad [(g, z, w)] \mapsto gz^{n-1}.$$

We can show that  $[\tilde{r} \circ \varphi']$  is independent of  $\iota$  using [Hen02, 1.8 Corollaire] and [BH00, 3.2 Theorem] similarly as the proof of [BH06, 35.1 Theorem].

Let  $\iota' : \mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_\ell$  be another choice. Taking a conjugation of  $\varphi'$  by an element of  $\widehat{\mathrm{GL}}_n(\overline{\mathbb{Q}}_\ell)$ , we may assume that  $\tilde{r} \circ \varphi' = \tilde{r} \circ \varphi'^{'}.$  Hence there is  $\chi : W_F \rightarrow \overline{\mathbb{Q}}_\ell^\times$  such that

$$\varphi'(w) = \varphi'^{(w)}[(\chi(w)^{1-n}, \chi(w), 1)]$$

for  $w \in W_F$ . By Definition 2.2, we have  $t_{\mathbb{G}_m} \circ \varphi' = t_{\mathbb{G}_m} \circ \varphi'^{'}.$  Hence we have  $\chi^2 = 1$ . This implies that  $\varphi' = \varphi'^{'}.$  since  $z_{\mathrm{GL}_n} = (-1)^{n-1}$ . Hence Conjecture 2.4 is true. Then Conjecture 2.8 is also true by Theorem 2.10.  $\square$

## References

- [AG91] J. Arthur and S. Gelbart, Lectures on automorphic  $L$ -functions, in  $L$ -functions and arithmetic (Durham, 1989), vol. 153 of London Math. Soc. Lecture Note Ser., pp. 1–59, Cambridge Univ. Press, Cambridge, 1991.
- [Ber20] J. Bernstein, Hidden sign in Langlands' correspondence, 2020, arXiv:2004.10487.
- [BG14] K. Buzzard and T. Gee, The conjectural connections between automorphic representations and Galois representations, in Automorphic forms and Galois representations. Vol. 1, vol. 414 of London Math. Soc. Lecture Note Ser., pp. 135–187, Cambridge Univ. Press, Cambridge, 2014.
- [BH00] C. J. Bushnell and G. Henniart, Davenport-Hasse relations and an explicit Langlands correspondence. II. Twisting conjectures, vol. 12, pp. 309–347, 2000, colloque International de Théorie des Nombres (Talice, 1999).
- [BH06] C. J. Bushnell and G. Henniart, The local Langlands conjecture for  $\mathrm{GL}(2)$ , vol. 335 of Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, Berlin, 2006.



- [Bor79] A. Borel, Automorphic  $L$ -functions, in Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979 pp. 27–61.
- [BV85] D. Barbasch and D. A. Vogan, Jr., Unipotent representations of complex semisimple groups, *Ann. of Math. (2)* 121 (1985), no. 1, 41–110.
- [Del73] P. Deligne, Les constantes des équations fonctionnelles des fonctions  $L$ , in Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), pp. 501–597. *Lecture Notes in Math.*, Vol. 349, Springer, Berlin, 1973.
- [Hei06] V. Heiermann, Orbites unipotentes et pôles d’ordre maximal de la fonction  $\mu$  de Harish-Chandra, *Canad. J. Math.* 58 (2006), no. 6, 1203–1228.
- [Hen02] G. Henniart, Une caractérisation de la correspondance de Langlands locale pour  $GL(n)$ , *Bull. Soc. Math. France* 130 (2002), no. 4, 587–602.
- [Hoc81] G. P. Hochschild, Basic theory of algebraic groups and Lie algebras, vol. 75 of *Graduate Texts in Mathematics*, Springer-Verlag, New York-Berlin, 1981.
- [HT01] M. Harris and R. Taylor, The geometry and cohomology of some simple Shimura varieties, vol. 151 of *Annals of Mathematics Studies*, Princeton University Press, Princeton, NJ, 2001, with an appendix by Vladimir G. Berkovich.
- [IT12] N. Imai and T. Tsushima, Geometric realization of the local Langlands correspondence for representations of conductor three, 2012, arXiv:1205.0734, to appear in *Publ. Res. Inst. Math. Sci.*
- [Joh13] C. Johansson, A remark on a conjecture of Buzzard-Gee and the cohomology of Shimura varieties, *Math. Res. Lett.* 20 (2013), no. 2, 279–288.
- [Kal16] T. Kaletha, The local Langlands conjectures for non-quasi-split groups, in *Families of automorphic forms and the trace formula*, *Simons Symp.*, pp. 217–257, Springer, [Cham], 2016.
- [KL87] D. Kazhdan and G. Lusztig, Proof of the Deligne-Langlands conjecture for Hecke algebras, *Invent. Math.* 87 (1987), no. 1, 153–215.
- [Kot84a] R. E. Kottwitz, Shimura varieties and twisted orbital integrals, *Math. Ann.* 269 (1984), no. 3, 287–300.
- [Kot84b] R. E. Kottwitz, Stable trace formula: cuspidal tempered terms, *Duke Math. J.* 51 (1984), no. 3, 611–650.
- [KW17] T. Kaletha and J. Weinstein, On the Kottwitz conjecture for moduli spaces of local shtukas, 2017, arXiv:1709.06651.

- [Lan83] R. P. Langlands, Les débuts d'une formule des traces stable, vol. 13 of Publications Mathématiques de l'Université Paris VII [Mathematical Publications of the University of Paris VII], Université de Paris VII, U.E.R. de Mathématiques, Paris, 1983.
- [LRS93] G. Laumon, M. Rapoport and U. Stuhler,  $\mathscr{D}$ -elliptic sheaves and the Langlands correspondence, *Invent. Math.* 113 (1993), no. 2, 217–338.
- [Mos56] G. D. Mostow, Fully reducible subgroups of algebraic groups, *Amer. J. Math.* 78 (1956), 200–221.
- [OV90] A. L. Onishchik and E. B. Vinberg, Lie groups and algebraic groups, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990, translated from the Russian and with a preface by D. A. Leites.
- [RV14] M. Rapoport and E. Viehmann, Towards a theory of local Shimura varieties, *Münster J. Math.* 7 (2014), no. 1, 273–326.
- [Ser96] J.-P. Serre, Exemples de plongements des groupes  $\mathrm{PSL}_2(\mathbf{F}_p)$  dans des groupes de Lie simples, *Invent. Math.* 124 (1996), no. 1-3, 525–562.
- [ST68] J.-P. Serre and J. Tate, Good reduction of abelian varieties, *Ann. of Math.* (2) 88 (1968), 492–517.
- [Zhu20] X. Zhu, A note on Integral Satake isomorphisms, 2020, arXiv:2005.13056.

Naoki Imai

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba,  
Meguro-ku, Tokyo, 153-8914, Japan  
naoki@ms.u-tokyo.ac.jp