

GLOBAL SOLUTION OF THE 3D RELATIVISTIC VLASOV-POISSON SYSTEM FOR A CLASS OF LARGE DATA

XUECHENG WANG

Abstract

Given smooth localized arbitrary large radial initial data, we prove global existence of smooth solution for the 3D relativistic Vlasov-Poisson system for the repulsive case. The compact support assumption on initial data is not needed.

1. INTRODUCTION

We consider the following relativistic Vlasov-Poisson system,

$$(RVP) \quad \begin{cases} \partial_t f + \hat{v} \cdot \nabla_x f + \gamma \nabla_x \phi \cdot \nabla_v f = 0 \\ \Delta \phi = \rho(t), \quad \rho(t) := \int_{\mathbb{R}^3} f(t, x, v) dv, \quad f(0, x, v) = f_0(x, v), \end{cases} \quad (1.1)$$

where $\hat{v} := v/\sqrt{1+|v|^2}$, $\gamma \in \{1, -1\}$, $f(t, x, v) \geq 0$ denotes the distribution of particles and $\rho(t, x)$ denotes the density of particles. The case $\gamma = 1$ is called the repulsive case, which is also called the plasma physics case, and the case $\gamma = -1$ is called the attractive case, which is also called the stellar dynamics case.

For the relativistic Vlasov-Poisson system (1.1), the following conservation laws hold,

$$E(t) := \int_{\mathbb{R}^n} |\nabla_x \phi(t)|^2 + \gamma \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v| f(t, x, v) dx dv = E(0), \quad \|f(t, x, v)\|_{L_{x,v}^p} = \|f(0, x, v)\|_{L_{x,v}^p}, \quad (1.2)$$

where $p \in [1, \infty]$. If the initial data is smooth and its high moment is small, then the system (1.1) admits global solution regardless the sign of γ , the regularity of initial data can be propagated, and the density and its derivatives decay sharply over time, see [10].

As pointed out by Glassey-Schaeffer in [2], RVP is worse behaved than the non relativistic case, in which there is a large literature and we do not try to elaborate it here but refer readers to Anderson [1] and Mouhot [8] and references therein for more detailed introduction. A remarkable result by Lions-Perthame [6] showed that the nonrelativistic Vlasov-Poisson system in repulsive case admits global classic solution for general initial data, see also [9].

To the best knowledge of the author, we don't have satisfactory picture of the system (1.1) for "general" initial data. There are some results if we assume that the initial data is radial in the following sense,

$$f_0(Rx, Rv) = f_0(x, v), \quad \forall R \in SO(3). \quad (1.3)$$

A well-known result by Glassey-Schaeffer [2] says that the sign of γ matters in the large data theory. Roughly speaking, there exists a class of compact support radial data for which the solution of the repulsive case blows up in finite time while the solution of the attractive case globally exists over time. More precisely, the system (1.1) for the attractive case ($\gamma = 1$) admits global classical solution if the initial data poses radial symmetry and has compact support in v . The solution of the repulsive case ($\gamma = -1$) blows up in finite time for radial initial data which has compact support in " v " and negative energy. Kiessling-Tahvildar-Zadeh [5] established sharp constants C_β for the L^β -norm of initial data for the negative energy condition.

In this paper, we mainly study the attractive case, i.e., $\gamma = 1$ and also restrict ourself to the radial symmetry case, which enable us to use the radial symmetry and the fact that the first moment is

bounded from the first conservation law in (1.2). However, we do not impose compact support assumption and do not restrict the size of initial data.

We aim to prove the following result.

Theorem 1.1. *Assume that the radial initial data (in the sense of (1.3)) $f_0(x, v) \in H^s(\mathbb{R}_x^3 \times \mathbb{R}_v^3)$, $s \in \mathbb{Z}_+$, $s \geq 6$ decays polynomially as $(x, v) \rightarrow \infty$. More precisely, for $N_0 := 100$, we assume the following estimate holds for the initial data,*

$$\sum_{\alpha \in \mathbb{Z}_+^6, |\alpha| \leq s} \|(1 + |x| + |v|)^{N_0} \nabla_{x,v}^\alpha f_0(x, v)\|_{L_{x,v}^2} < +\infty. \quad (1.4)$$

Then the relativistic Vlasov-Poisson system (1.1) for the attractive case, i.e., $\gamma = 1$, admits global solution in $H^s(\mathbb{R}_x^3 \times \mathbb{R}_v^3)$.

Remark 1.1. The primary goal of this paper is to obtain global solution for a class of large data without the compact support assumption. This improves the previous result by Glassey-Schaeffer [2], in which the initial data has compact support. The goal of optimizing s and N_0 is not pursued here. To remove the radial assumption, which is a very interesting problem, new ingredients are needed.

Acknowledgment The author is supported by NSFC-11801299.

2. PROOF OF THEOREM 1.1

2.1. Notation. For any two numbers A and B , we use $A \lesssim B$ and $B \gtrsim A$ to denote $A \leq CB$, where C is an absolute constant.

We use the convention that all constants which only depend on the initial data, e.g., the conserved quantities, will be treated as absolute constants.

2.2. Local theory and the reduction of the proof. Because our assumption on the initial data is stronger than the assumption imposed on the distribution function in [7], by using the same argument used by Luk-Strain [7] for the relativistic Vlasov-Maxwell system, which is more difficult, we can reduce the proof of global existence to the L_x^∞ -estimate of the scalar field $\nabla_x \phi$, which corresponds to the acceleration of the speed of particles.

More precisely, assume that T^* is the maximal time of existence of solution. If we can show that $\nabla_x \phi \in L^\infty([0, T^*) \times \mathbb{R}_x^3)$, then the lifespan of the solution can be extended to $[0, T^* + \epsilon]$ for some positive number ϵ . That is to say, the solution of RVP (1.1) exists globally in time.

To control the L_x^∞ -norm of the acceleration term $\nabla_x \phi$, now it's a standard argument to show that it is controlled by a high moment of the distribution function, see also Lemma 2.2.

2.3. Propagation of moments. We define

$$M_n(t, x) := \int_{\mathbb{R}^3} (1 + |v|)^n f(t, x, v) dv, \quad M_n(t) := \int_{\mathbb{R}^3} M_n(t, x) dx, \quad n := \lceil N_0/10 \rceil. \quad (2.1)$$

From the conservation laws (1.2), we know that $M_1(t)$ is always bounded from the above. Moreover, we define

$$\tilde{M}_n(t) := (1 + t)^{2n} + \sup_{s \in [0, t]} M_n(s). \quad (2.2)$$

We have two basic estimates for the L_x^∞ -norm of the acceleration term $\nabla_x \phi$, which will be elaborated in the next two Lemmas. The first estimate (2.3) is available mainly because of the radial symmetry and the conservation law. The second estimate (2.5) is standard.

Lemma 2.1. *For any $t \in [0, T^*)$, $x \in \mathbb{R}^3$, the following point-wise estimate holds,*

$$|\nabla_x \phi(t, x)| \lesssim \frac{1}{|x|^2}. \quad (2.3)$$

Proof. Recall that the distribution f is radial, see (1.3). As a result, we know that the density function $\rho(t, x)$ is radial,

$$\forall x \in \mathbb{R}^3, \quad \rho(t, Rx) = \int_{\mathbb{R}^3} f(t, Rx, v) dv = \int_{\mathbb{R}^3} f(t, Rx, R\omega) |\det(R)| d\omega = \int_{\mathbb{R}^3} f(t, x, \omega) d\omega = \rho(t, x).$$

Hence $\phi(t, x)$ is also radial. Define

$$\tilde{\phi}(t, r) := \phi(t, r, 0, 0), \quad \tilde{\rho}(t, r) := \rho(t, r, 0, 0), \quad \implies \phi(t, x) = \tilde{\phi}(t, |x|), \rho(t, x) = \tilde{\rho}(t, |x|).$$

The Poisson equation for ϕ in (1.1) is reduced as follows,

$$\begin{aligned} (\partial_r^2 + \frac{2}{r}\partial_r)\tilde{\phi}(t, r) &= \tilde{\rho}(t, r), \quad \partial_r^2(r\tilde{\phi})(t, r) = r\tilde{\rho}(t, r), \quad \implies \partial_r(r\tilde{\phi}) = \int_0^r s\tilde{\rho}(t, s) ds + c \\ \implies r\tilde{\phi} &= \int_0^r \int_0^\tau s\tilde{\rho}(t, s) ds d\tau + cr = \int_0^r s(r-s)\tilde{\rho}(t, s) ds + cr \\ \implies \tilde{\phi}(t, r) &= \int_0^r s\tilde{\rho}(t, s) ds - \frac{1}{r} \int_0^r s^2\tilde{\rho}(t, s) ds + c, \quad \implies \partial_r\tilde{\phi}(t, r) = \frac{1}{r^2} \int_0^r s^2\tilde{\rho}(t, s) ds \end{aligned}$$

Hence, from the above equality and the conservation law in (1.2), the following estimate holds point-wisely,

$$\implies \nabla_x \phi = \frac{x}{|x|} \partial_r \tilde{\phi}(t, |x|) = \frac{x}{|x|^3} \int_0^{|x|} s^2 \tilde{\rho}(t, s) ds, \quad \implies \nabla_x \phi = \frac{x}{|x|} |\nabla_x \phi|, \quad |\nabla_x \phi| \lesssim \frac{1}{|x|^2}. \quad (2.4)$$

□

Lemma 2.2. *Let $\epsilon \in (0, 10^{-10})$ be some fixed sufficiently small constant, then the following estimate holds for any $t \in [0, T^*]$,*

$$\|\nabla_x \phi(t, x)\|_{L_x^\infty} \lesssim 1 + (M_n(t))^{(5+\epsilon)/((3-\epsilon)(n-1))}. \quad (2.5)$$

Proof. Note that

$$\nabla_x \phi(x) = \int_{\mathbb{R}^3} \frac{\rho(t, y)(y-x)}{|x-y|^3} dy = \int_{|y-x| \leq \delta} \frac{\rho(t, y)(y-x)}{|x-y|^3} dy + \int_{|y-x| \geq \delta} \frac{\rho(t, y)(y-x)}{|x-y|^3} dy. \quad (2.6)$$

From the conservation law in (1.2), the first part of the above equation is controlled as follows,

$$\left| \int_{|y-x| \geq \delta} \frac{\rho(t, y)(y-x)}{|x-y|^3} dy \right| \lesssim \frac{1}{\delta^2}. \quad (2.7)$$

For the second part, we use the Hölder inequality by choosing $p = (3-\epsilon)/2$ and $q = (3-\epsilon)/(1-\epsilon)$. As a result, we have

$$\left| \int_{|y-x| \leq \delta} \frac{\rho(t, y)(y-x)}{|x-y|^3} dy \right| \lesssim \left(\int_{|y-x| \leq \delta} \frac{1}{|y-x|^{2p}} dy \right)^{1/p} \left(\int_{\mathbb{R}^n} (\rho(t, y))^q dy \right)^{1/q} \lesssim \delta^{2\epsilon/(3-\epsilon)} \|\rho(t, x)\|_{L^q}. \quad (2.8)$$

Since $\|f(t, x, v)\|_{L_{x,v}^\infty}$ and $M_1(t)$ is bounded all time from the conservation law in (1.2), the following two estimates hold,

$$\rho(t, x) = \int_{\mathbb{R}^3} f(t, x, v) dv \lesssim R^3 + R^{-6/(1-\epsilon)} M_{6/(1-\epsilon)}(t, x) \lesssim (M_{6/(1-\epsilon)}(t, x))^{(1-\epsilon)/(3-\epsilon)},$$

Note that, for $m < n$, we have

$$\begin{aligned} M_m(t) &= \int_{|v| \leq R} (1+|v|)^m f(t, x, v) dx dv + \int_{|v| \geq R} (1+|v|)^m f(t, x, v) dx dv \\ &\lesssim R^{m-1} + R^{m-n} M_n(t) \lesssim (M_n(t))^{(m-1)/(n-1)}. \end{aligned}$$

From the above two estimates, we have

$$\|\rho(t, x)\|_{L^q} \lesssim (M_n(t))^{(5+\epsilon)/((3-\epsilon)(n-1))} \quad (2.9)$$

Hence, from the estimates (2.6–2.9), after letting $\delta = 1$, we have

$$\|\nabla_x \phi(x)\|_{L_x^\infty} \lesssim 1 + (M_n(t))^{(5+\epsilon)/((3-\epsilon)(n-1))}. \quad (2.10)$$

Hence finishing the proof of the desired estimate (2.5). \square

From the first estimate (2.3), we know that the acceleration force of particles is weak if the particles are far away from the origin. Meanwhile, from the second estimate (2.5), we know that the acceleration force of particles is not too strong even if the particles are very close to the origin.

We will show a key observation that the majority of localized particles will travel toward infinity after the speed of particles reaches a threshold. Hence, because of the first estimate (2.3), we know that the majority of localized particles will not be accelerated very much in later time.

The result of this observation is summarized in Lemma 2.3. Before proceeding to detailed analysis, we need some preparation. From (1.1) and (2.4), the backward characteristics associated with the Vlasov-Poisson equation read as follow,

$$\begin{cases} \frac{d}{ds} X(s; t, x, v) = \widehat{V}(s; t, x, v) \\ \frac{d}{ds} V(s; t, x, v) = \nabla_x \phi(X(s; t, x, v)) = \frac{X(s; t, x, v)}{|X(s; t, x, v)|} |\nabla_x \phi(X(s; t, x, v))| \\ X(t; t, x, v) = x, \quad V(t; t, x, v) = v \end{cases} \quad (2.11)$$

Hence, the dynamics of the lengths of the characteristics read as follows,

$$\frac{d}{ds} |X(s; t, x, v)| = \frac{X(s; t, x, v)}{|X(s; t, x, v)|} \cdot \widehat{V}(s; t, x, v), \quad (2.12)$$

$$\frac{d}{ds} |V(s; t, x, v)| = \frac{V(s; t, x, v) \cdot X(s; t, x, v)}{|V(s; t, x, v)| |X(s; t, x, v)|} |\nabla_x \phi(X(s; t, x, v))|. \quad (2.13)$$

From the above two equations, we can see that the quantity $V(t, s, x, v) \cdot X(t, s, x, v)$ plays an essential role. We analyze its dynamics over time as follows,

$$\begin{aligned} \frac{d}{ds} \frac{V(s; t, x, v) \cdot X(s; t, x, v)}{|V(s; t, x, v)|} &= |\widehat{V}(s; t, x, v)| + \frac{\partial_s V(s; t, x, v) \cdot X(s; t, x, v)}{|V(s; t, x, v)|} \\ &\quad - \frac{V(t, s, x, v) \cdot X(t, s, x, v)}{|V(t, s, x, v)|^2} \frac{V(t, s, x, v) \cdot \partial_s V(t, s, x, v)}{|V(t, s, x, v)|} \\ &= |\widehat{V}(s; t, x, v)| + \frac{|X(s; t, x, v)|}{|V(s; t, x, v)|} \left(1 - \frac{(V(s; t, x, v) \cdot X(s; t, x, v))^2}{|V(s; t, x, v)|^2 |X(s; t, x, v)|^2}\right) |\nabla_x \phi(X(s; t, x, v))| \geq 0, \end{aligned} \quad (2.14)$$

From the above equation, we know that the quantity $V(s; t, x, v) \cdot X(s; t, x, v)$ is an increasing function with respect to time “ s ”.

We define a set of majorities of particles, which initially localize around zero, at time s as follows,

$$R(t, s) := \{(X(s; t, x, v), V(s; t, x, v)) : |X(0; t, x, v)| + |V(0; t, x, v)| \leq (\tilde{M}_n(t))^{1/(2n)}\}.$$

Lemma 2.3. *For any $t \in [0, T^*)$, the following relation holds for some sufficiently large absolute constant C ,*

$$R(t, t) \subset B(0, C(\tilde{M}_n(t))^{1/(2n)}) \times B(0, C(\tilde{M}_n(t))^{(5+2\epsilon)/((6-2\epsilon)(n-1))}) \quad (2.15)$$

Proof. Let $t \in [0, T^*)$ be fixed. Note that, from the equation (2.12), the following rough estimate holds for the length of $X(s; t, x, v)$,

$$|X(s; t, x, v)| \leq |X(0; t, x, v)| + |s| \leq 2(\tilde{M}_n(t))^{1/(2n)}, \quad s \in [0, t]. \quad (2.16)$$

We define the maximal time such that the velocity characteristic doesn't exceed the threshold as follows,

$$\tau := \sup\{s : \forall \kappa \in [0, s], |V(\kappa; t, x, v)| \leq (\tilde{M}_n(t))^{(5+2\epsilon)/((6-2\epsilon)(n-1))}\}.$$

From the continuity of characteristics, we know that $\tau > 0$. If $\tau = t$, then we are done. It remains to consider the case when $0 < \tau < t$.

Note that, from the equation (2.13), we know that $|V(s; t, x, v)|$ is decreasing if $V(s; t, x, v) \cdot X(s; t, x, v) < 0$. Hence, at the time τ , we have $V(\tau; t, x, v) \cdot X(\tau; t, x, v) \geq 0$. Otherwise, it contradicts the definition of the maximal time. From the monotonicity of $V(s; t, x, v) \cdot X(s; t, x, v)$, see the equation (2.14), we know that $V(s; t, x, v) \cdot X(s; t, x, v) \geq 0$ for $s \in [\tau, t]$, which implies that $|V(t, s, x, v)| \geq |V(t, \tau, x, v)|$ for all $s \in [\tau, t]$ from the equation (2.13). To sum up, $\forall s \in [\tau, t]$, we have

$$|V(s; t, x, v)| \geq |V(\tau; t, x, v)| = (\tilde{M}_n(t))^{(5+2\epsilon)/((6-2\epsilon)(n-1))}, \quad V(s; t, x, v) \cdot X(s; t, x, v) \geq 0. \quad (2.17)$$

Starting from the time τ , from the above estimate and the equation (2.14), we have the following estimate for any $s \in [\tau, t]$,

$$\frac{V(s; t, x, v) \cdot X(s; t, x, v)}{|V(s; t, x, v)|} = \frac{V(\tau; t, x, v) \cdot X(\tau; t, x, v)}{|V(\tau; t, x, v)|} + \int_{\tau}^s \frac{d}{ds} \frac{V(\kappa; t, x, v) \cdot X(\kappa; t, x, v)}{|V(\kappa; t, x, v)|} d\kappa \gtrsim s - \tau. \quad (2.18)$$

From the equation (2.12) and the estimates (2.17) and (2.18), we have

$$|X(s; t, x, v)|^2 - |X(\tau; t, x, v)|^2 = \int_{\tau}^s \frac{d}{ds} |X(\kappa; t, x, v)|^2 d\kappa \gtrsim \int_{\tau}^s (\kappa - \tau) d\kappa \gtrsim (s - \tau)^2.$$

From the above estimate, the estimate (2.10), and the equation in (2.11), the following estimate holds for any $s \in [\tau, t]$,

$$\begin{aligned} |V(s; t, x, v)| &\leq |V(\tau; t, x, v)| + \int_{\tau}^s \left| \frac{d}{ds} V(\kappa; t, x, v) \right| d\kappa \\ &\lesssim (\tilde{M}_n(t))^{(5+2\epsilon)/((6-2\epsilon)(n-1))} + \int_{\tau}^s |\nabla_x \phi(X(t, \kappa, x, v))| d\kappa \\ &\lesssim (\tilde{M}_n(t))^{(5+2\epsilon)/((6-2\epsilon)(n-1))} + \int_{\tau}^{\tau+\delta} (\tilde{M}_n(t))^{(5+\epsilon)/((3-\epsilon)(n-1))} d\kappa + \int_{\tau+\delta}^s \frac{1}{(\kappa - \tau)^2} d\kappa \\ &\lesssim (\tilde{M}_n(t))^{(5+2\epsilon)/((6-2\epsilon)(n-1))}, \quad \text{by letting } \delta = (\tilde{M}_n(t))^{-(5+\epsilon)/((6-2\epsilon)(n-1))}. \end{aligned} \quad (2.19)$$

To sum up, our desired conclusion (2.15) holds from (2.16) and (2.19). \square

The particles that don't belong to the majority set $R(t, t)$ can be tracked roughly back to their positions at the initial time. It allows us to control their magnitude from the assumption of the initial data in (1.4).

Lemma 2.4. *For any $t \in [0, T^*)$, $x, v \in \mathbb{R}^3$, s.t., $|v| \gtrsim (\tilde{M}_n(t))^{(5+3\epsilon)/((6-2\epsilon)(n-1))}$, we have*

$$|f(t, x, v)| \lesssim (\tilde{M}_n(t))^{-4} (1 + |x|)^{-4}. \quad (2.20)$$

Moreover, if $|v| \gtrsim (\tilde{M}_n(t))^{5/(2n)}$, then we have

$$|f(t, x, v)| \lesssim (1 + |x|)^{-4} (1 + |v|)^{-n-4}. \quad (2.21)$$

Proof. Note that, from the relation (2.15), we have $|X(0; t, x, v)| + |V(0; t, x, v)| \gtrsim (\tilde{M}_n(t))^{1/(2n)}$ if $|x| \gtrsim (\tilde{M}_n(t))^{(1+\epsilon)/(2n)}$ or $|v| \gtrsim (\tilde{M}_n(t))^{(5+3\epsilon)/((6-2\epsilon)(n-1))}$. If $|x| \gtrsim (\tilde{M}_n(t))^{(1+\epsilon)/(2n)}$, then we have

$$|X(0; t, x, v)| \geq |x| - t \geq |x| - (\tilde{M}_n(t))^{1/(2n)} \gtrsim (1 + |x|),$$

Therefore, from the above estimate and the assumption on the initial data in (1.4), the following estimate holds if $|v| \gtrsim (\tilde{M}_n(t))^{(5+3\epsilon)/((6-2\epsilon)(n-1))}$ regardless the size of $|x|$,

$$|f(t, x, v)| = |f_0(X(0; t, x, v), V(0; t, x, v))| \leq (\tilde{M}_n(t))^{-4}(1 + |x|)^{-4}. \quad (2.22)$$

Moreover, if $|v| \gtrsim (\tilde{M}_n(t))^{5/(2n)}$, then the following estimate holds from the equation (2.12) and the estimate (2.5),

$$|V(0; t, x, v)| \gtrsim |v| - \int_0^t (1 + (M_n(s))^{(5+\epsilon)/((3-\epsilon)(n-1))}) ds \gtrsim (1 + |v|).$$

From the above two estimates and the assumption on the initial data in (1.4), the following estimate holds if $|v| \gtrsim (\tilde{M}_n(t))^{5/(2n)}$ regardless the size of $|x|$,

$$|f(t, x, v)| = |f_0(X(0; t, x, v), V(0; t, x, v))| \lesssim (1 + |x|)^{-4}(1 + |v|)^{-n-4}.$$

Hence finishing the desired estimates (2.20) and (2.21). \square

Proof of Theorem 1.1: Based on the possible size of “ $|v|$ ”, we decompose $M_n(t)$ into three parts as follows,

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} (1 + |v|)^n |f(t, x, v)| dx dv = I + II + III,$$

where

$$\begin{aligned} I &:= \int_{\mathbb{R}^3} \int_{|v| \geq (\tilde{M}_n(t))^{3/n}} (1 + |v|)^n |f(t, x, v)| dx dv, \\ II &:= \int_{\mathbb{R}^3} \int_{(\tilde{M}_n(t))^{(5+4\epsilon)/((6-2\epsilon)(n-1))} \leq |v| \leq (\tilde{M}_n(t))^{3/n}} (1 + |v|)^n |f(t, x, v)| dx dv, \\ III &:= \int_{\mathbb{R}^3} \int_{|v| \leq (\tilde{M}_n(t))^{(5+4\epsilon)/((6-2\epsilon)(n-1))}} (1 + |v|)^n |f(t, x, v)| dx dv. \end{aligned}$$

Therefore, from the estimates (2.20) and (2.21) in Lemma 2.4, we have

$$|I| + |II| \lesssim 1.$$

From the conservation law (1.2), we have

$$|III| \lesssim (\tilde{M}_n(t))^{(5+4\epsilon)n/((6-2\epsilon)(n-1))}.$$

To sum up, we have

$$M_n(t) \lesssim (\tilde{M}_n(t))^{(5+4\epsilon)n/((6-2\epsilon)(n-1))}.$$

Since the above estimate holds for any $t \in [0, T^*]$ and $\tilde{M}_n(t)$ is an increasing function with respect to t , the following estimate holds for any $s \in [0, t]$,

$$M_n(s) \lesssim (\tilde{M}_n(s))^{(5+4\epsilon)n/((6-2\epsilon)(n-1))} \leq (\tilde{M}_n(t))^{(5+4\epsilon)n/((6-2\epsilon)(n-1))},$$

Hence

$$\tilde{M}_n(t) = \sup_{s \in [0, t]} M_n(s) + (1+t)^{2n} \lesssim (\tilde{M}_n(t))^{(5+4\epsilon)n/((6-2\epsilon)(n-1))} + (1+t)^{2n}, \implies \tilde{M}_n(t) \lesssim (1+t)^{2n}.$$

Therefore, from the estimate (2.5) in Lemma 2.2, we have

$$\|\nabla_x \phi(t, x)\|_{L_x^\infty} \lesssim (1+t)^{2n}.$$

We have shown the desired fact that $\nabla_x \phi \in L^\infty([0, T^*) \times \mathbb{R}_x^3)$. Hence finishing the proof of Theorem 1.1. □

REFERENCES

- [1] H. Andréasson. The Einstein-Vlasov system/kinetic theory, *Living Rev. Relativ.*, (2011), 14:4.
- [2] R.T. Glassey and J. Schaeffer. On symmetric solutions of the relativistic Vlasov-Poisson system. *Commun. Math. Phys.* 101, 459-473 (1985).
- [3] M. Hadzic and G. Rein. Global existence and nonlinear stability for the relativistic Vlasov-Poisson system in the gravitational case, *Indiana Univ. Math. J.* 56 (2007), 2453–2488.
- [4] E. Horst. Global strong solutions of Vlasov’s Equation. Necessary and sufficient conditions for their existence. Partial differential equations. *Banach Cent. Publ.* 19, 143–153 (1987).
- [5] M. K.-H. Kiessling and A. S. Tahvildar-Zadeh. On the relativistic Vlasov-Poisson system. *Indiana Univ. Math. J.* 57 (2008), no. 7, 3177–3207.
- [6] P.L. Lions and B. Perthame. Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system. *Invent. Math.*, 105 (1991), 415–430.
- [7] J. Luk and R. Strain. Strichartz estimates and the moment bounds for the relativistic Vlasov-Maxwell system. *Arch. Rational Mech. Anal.* 219(2016), 445–552.
- [8] C. Mouhot. Stabilité orbitale pour le système de Vlasov-Poisson gravitationnel (d’après Lemou-Méhats-Raphaël, Guo, Lin, Rein et al.). *Astérisque*, 352(2013), Exp. No. 1044, vii, 35–82. Séminaire Bourbaki, vol. 2011/2012. Exposés 1043–1058.
- [9] K. Pfaffelmoser. Global classical solutions for the Vlasov-Poisson system in three dimensions for general initial data, *J. Diff. Eqns.*, 95 (1992), 281–303.
- [10] X. Wang. Decay estimates for the 3D relativistic and non-relativistic Vlasov-Poisson systems, arXiv:1805.10837.

YMSC, TSINGHUA UNIVERSITY, BEIJING, CHINA, 100084

E-mail address: xuecheng@tsinghua.edu.cn, xuecheng.wang.work@gmail.com