

Parametric analysis of COVID-19 expansion in European countries in the period of February to June 2020

Martin Spousta^a

^a*Institute of Particle and Nuclear Physics, Charles University, Prague, Czech Republic*

Abstract

The data on number of registered cases of COVID-19 disease in twenty European countries is analyzed by the least-squares fitting procedure with generic analytic functions. Three regimes of the expansion of the disease are identified and quantified – early exponential expansion, damped exponential, and linear expansion. Differences among countries in the early expansion period are quantified. The velocity of the expansion in the exponential regime lies within one standard deviation from the average value for 11 countries. The number of infected individuals at the initial time is excessively high for Italy, 7 standard deviations from the average value.

Method for predicting the expansion based on extrapolation in the parametric space is presented. One week predictions based on extrapolations have average precision of 18% and 29% during the later period of the damped exponential expansion for the case of Italy and Czechia, respectively. The method based on extrapolations in the parametric space may provide an elementary method to quantify the impact of restrictive measures on the spreading of the disease.

Keywords: Outbreak, Infection, SARS-CoV-2, COVID-19, Parametric analysis, European countries

1. Introduction

The outbreak of new coronavirus SARS-CoV-2, causing severe respiratory tract infection in humans, known as COVID-19, is a global health concern. Restrictive measures were adopted in many countries to mitigate the impact of the spread of the disease on the public health system [1].

The spread of an infectious disease is usually modeled using compartmental epidemiological models, see e.g. review in Ref. [2] or examples of their recent implementation in Refs. [3–5]. These models are often based on the numerical solution of a system of coupled linear differential equations for the number of susceptible (S), infectious (I), and recovered (R) cases (SIR models). For the simplest SIR model, a set of three differential equation characterizes the transitions among the three classes.

While it is not possible to obtain an analytic solution for a complex system of the SIR model equations, it is possible to construct generic analytic functions describing the data and quantify the robustness of such a description based on statistical tests. This approach is the backbone of this analysis. The main goal of this analysis is to identify and quantify robust features in the dynamics of spread of the disease and present a method that may help quantify the impact of restrictive measures on the spreading of the disease.

2. Method and dataset

In the simple SIR model, the equation which describes the change in the number of infectious cases reads $dI/dt = (c_1 S/N - c_2)I$ where $N = S + I + R$ and c_1, c_2 are constants (see e.g. [6] for details). At the beginning of the outbreak when no restrictive measures are applied and the susceptibility of the population to the virus is 100%, one has $S \gg I$ and $R \rightarrow 0$. This leads to $dI/dt \propto I$ implying an exponential increase of the number of infected individuals with time,

$$I = I_0 \exp[at], \quad (1)$$

where I_0 and a are constants, combinations of c_1, N , and S . When restrictive measures are imposed, c_1 is not a constant but a function $c_1 = c_1(t, \dots)$ introducing more complexity to the set of SIR model equations. Alternatively, the evolution under restrictive measures can be described by exponential dependence with a time-dependent parameter, that is, by the function

$$F(t) = N_0 \exp[a(t)t] = N_0 \exp[b_1 t + b_2 t^2 + b_3 t^3 + \dots], \quad (2)$$

where the *r.h.s.* of (2) represents a Taylor expansion of the time dependent parameter a and N_0 reflects number of infected individuals at $t = 0$. Full description of the data can be achieved with n b_n parameters where n is a large number. The *r.h.s.* of (2) characterizes the basic exponential dependence plus any smooth deviation from this dependence.

A large number of free parameters in (2) may be reduced and interpreted. At the beginning of the outbreak, the number of cases is described by a pure

exponential function ($b_1 \neq 0, b_{n>1} = 0$) as argued in Eq. (1). This hypothesis is tested in Section 3.1 and the parameters of the exponential function are compared among European countries. Once the non-zero b_2 is needed to describe the data, the expansion starts to slow down with respect to the purely exponential propagation. This is quantified in Section 3.2. In an idealized case, only one b_2 coefficient along with previously fitted b_1 coefficient would be needed to describe the data. However, such a simple two-component characterization is shown to be invalid in Section 3.2, and higher-order terms are needed or alternatively time-dependent b_2 and N_0 coefficients are needed to describe the data. The latter approach is used in this analysis. The evolution of parameters with time is studied, and predictability of the spread based on extrapolations is quantified. After a certain period, the description by a linear function outperforms the description by (2). The transition to a linear spread is quantified in Section 3.3.

The cumulative number of registered cases of COVID-19 disease in European countries is used as input data to this analysis. Specifically, the analysis uses data from 20 European countries, 17 countries of the European Union exceeding 5 million of inhabitants and data from Norway, Switzerland, and the United Kingdom. Data are taken from information system [7, 8] for a period of 15th of February to 1st of June 2020. Analysis uses statistical tools implemented in *ROOT 6.1* [9].

3. Results

3.1. Period of exponential expansion

During the first period, when there is no implementation of restrictive measures and the susceptibility of the population to the virus is 100%, the number of all registered cases exponentially grows. This is described by Equation (2) with $b_{n>1} = 0$ and b_1 quantifying the velocity of the spread of the disease, here in units day^{-1} . To quantify the expansion during this period we perform the least-squares fitting of the number of all registered cases by an exponential function in the time interval of $t_{min} - t_{max}$. The default lower bound t_{min} was selected by a requirement of at least 50 cases to reduce fluctuations due to too small statistical sample. The upper bound t_{max} was selected individually for each country based on comparing with an alternative description allowing for non-zero b_2 coefficient for each day when the number of cases exceeds 500. The comparison is based on a χ^2 test. The value of t_{max} represents a first day when fit with non-zero b_2 provides a better description than a fit with zero b_2 . The resulting b_1 and N_0 coefficients are summarized in Table 1. The boundary condition leading to the definition of t_{min} (50 cases) was varied by 20% both up and down, which leads to alternative results. These alternative results were found to correspond with the nominal results within one standard deviation for 90% of countries.

Table 1 contains also results of Kolmogorov-Smirnov (KS) test. Based on p -values, one can accept the hypothesis that the exponential function provides

Country	N_0	b_1 [day $^{-1}$]	p_{KS}
Italy	6.61 ± 1.02	0.39 ± 0.01	1.00
Germany	0.57 ± 0.13	0.35 ± 0.01	1.00
Portugal	0.01 ± 0.00	0.34 ± 0.01	1.00
Sweden	0.23 ± 0.06	0.31 ± 0.01	1.00
Spain	0.92 ± 0.20	0.30 ± 0.01	1.00
France	1.38 ± 0.39	0.30 ± 0.02	1.00
Austria	0.13 ± 0.03	0.30 ± 0.01	1.00
Norway	0.37 ± 0.14	0.29 ± 0.02	1.00
Denmark	0.27 ± 0.05	0.29 ± 0.01	0.94
Czechia	0.10 ± 0.03	0.27 ± 0.01	1.00
Switzerland	0.80 ± 0.11	0.27 ± 0.01	1.00
Netherlands	0.83 ± 0.15	0.26 ± 0.01	1.00
Poland	0.08 ± 0.03	0.25 ± 0.01	1.00
Belgium	0.94 ± 0.12	0.24 ± 0.00	1.00
United Kingdom	1.85 ± 0.37	0.22 ± 0.01	1.00
Greece	0.97 ± 0.19	0.19 ± 0.01	1.00
Romania	0.84 ± 0.16	0.18 ± 0.01	1.00
Hungary	0.26 ± 0.06	0.17 ± 0.01	1.00
Bulgaria	2.96 ± 0.37	0.11 ± 0.00	1.00
Slovakia	4.28 ± 0.07	0.10 ± 0.00	0.96

Table 1: Parameters of fit (N_0 , b_1) of the total number of cases in European countries during the first period of the expansion (for the definition see the text). Fit function is given by Equation (2) with $b_i = 0$ for $i > 1$. The p -values for Kolmogorov-Smirnov test. Countries are ordered by a decreasing value of b_1 coefficient.

a suitable description of the spread of the disease in the early stage for all the countries.

The average value of b_1 is 0.25 ± 0.04 day $^{-1}$. The value of b_1 for 11 out of 20 countries lies within one standard deviation from the average value. The value of N_0 (which reflects the number of infected individuals at $t = 0$) oscillates between zero and one for the majority of countries, but it is excessively high for the case of Italy (7 standard deviations from the average value calculated excluding Italy). This may be interpreted as evidence for the validity of a hypothesis that the spread in Italy has started sooner than when the first cases were reported and systematically traced (see e.g. Ref. [10]).

To further check the universality of the initial velocity of the expansion quantified by b_1 coefficient, we studied magnitude of Pearson's correlation coefficient quantifying the correlation between b_1 and other quantity x , $\rho(b_1, x)$. For x we tested: 1) N_0 ; 2) number of days between 15th of February (first day when cases start to be collected European-wide [7]) and the day when the number of identified cases exceeds 50 in a given country (d_{50}); 3) the number of inhabitants in a given country n_{inhab} . The absolute value of the correlation coefficient




Figure 1: The N_0 (left), b_2 (middle), and number of all cases (right) as a function of time for Italy (green) and Czechia (red). The published data on the number of registered cases (full markers) are shown together with one-week predictions (open markers) (for details see the text).

was found to be smaller than 0.2 for all the quantities, implying that b_1 is not correlated with any of N_0 , d_{50} , and n_{inhab} .

The value of b_1 can be further used as an input to specific epidemiological studies. In particular, it can be used to estimate the basic reproduction number, R_0 , [11] using Laplace transform as $R_0 = 1/[\int_0^\infty \exp(-b_1 \kappa) h(\kappa) d\kappa]$ [12, 13]. Here κ and $h(\kappa)$ represent the generation interval (i.e. the average time interval from onset of one individual to the onset of another individual) and its probability distribution, respectively.

3.2. Period of exponential expansion with damping

As shown in the previous section, the growth of the number of cases during the first period of spread of the disease can be characterized by an exponential function. The parameter b_1 can be interpreted as a parameter quantifying the uncontrolled spread of the disease. This first period ends when the fit starts to prefer a non-zero b_2 . Function from *r.h.s.* of Equation (2) with $b_i = 0$ for $i > 2$ is fitted to the number of all registered cases in a given country in a time window of 7 days. This time window moves in steps of one day to quantify the evolution of N_0 and b_2 with time.

While the complete description of the spread can be done by the function $a(t)$ from Equation (2), here we take only the first two coefficients of the Taylor expansion of that function. We have tested that this provides a good description of the data ($p > 0.995$ from KS tests) in 7-day time window during the full second period. Allowing for non-zero b_3 or higher-order terms does not bring an improvement in the description. Therefore, one parameter, b_2 , quantifies the degree of deviation from a purely exponential expansion in a given time. The more negative b_2 implies more significant “damping” of the original exponential growth. The units of b_2 used here are day^{-2} .

The example of time evolution of N_0 and b_2 for one large country (Italy) and one small country (Czechia) is shown in the left and middle panel of Figure 1. The parameter b_2 starts at zero (purely exponential expansion) and then rapidly decreases. The largest negative values are achieved for Italy. The b_2 distributions are qualitatively similar: first b_2 decreases fast, later the decrease is slower, ending at a minimum followed by a shallow increase where the b_2 achieves similar values for both countries. While the interpretation of b_2 parameter is straightforward, one can call it a “push-back” parameter, the interpretation of time-dependent N_0 is not straightforward anymore and it is just a nuisance parameter.

In the case that b_2 turns to a constant in time, the evolution is predictable by simple analytic calculation. In particular the time t' when no new cases should be registered, is given by a derivative of Equation (2) with respect to time, leading to $t' = -b_1/(2b_2)$. Further, if N_0 is constant in time, number of all cases can be calculated by evaluating (2) at time t' . While b_2 and N_0 exhibit only a mild evolution during certain periods, these parameters do not converge to constant values. Prediction of the evolution of the value of registered cases cannot therefore be done analytically in general. However, it can be done by extrapolating the evolution of parameters to the future. Extrapolations based on constant, linear, polynomial, and exponential functions were tested. Prediction from extrapolation was calculated for a value of registered cases in 7 days after a given time. These one-week predictions are shown along with the data in the right panel of Figure 1. For the period when b_2 is decreasing, it was found that an extrapolation using constant values of parameters provides the best matching between estimates of the number of registered cases based on these extrapolations and the actual data. For the period when b_2 starts to increase, a linear extrapolation and exponential extrapolation of b_2 and N_0 , respectively, provide predictions that are closest to the data.

For the case of Italy, the predictions differ from the data on average by $265\% \pm 160\%$ during the first 14 days after the time when data start to prefer the parameterization with non-zero b_2 . This substantial disagreement is given by a fast decrease of b_2 during the first two weeks which we did not succeed to describe universally by the extrapolation. For the next 50 days, the predictions differ from the data on average by $29\% \pm 19\%$ and $18\% \pm 10\%$ in the case of Italy and Czechia, respectively. These findings imply that the extrapolation of trends in the parameter space used in this study does not allow to make realistic predictions in the period of two weeks after the time when b_2 starts to be non-zero. However, for the later times, the method allows to make estimates with reasonable precision.

The method based on extrapolating the trends in the parametric space may provide an elementary method to quantify the impact of restrictive measures on the number of registered cases. This may be done by evaluating a difference between the extrapolation and the data for a time period when a given restrictive measure is applied.

The fact that the method based on extrapolations provide a reasonable estimate of the future evolution for a period of one week is simple evidence for a

significant inertia in the evolution.

3.3. Period of linear evolution

The functional form of Equation (2) stops describing the data well at some point. This reflects the fact that the spread of the disease turns from the damped exponential expansion to a linear evolution. This is quantified in Figure 2 for the case of Italy in terms of the sum of residuals for the damped exponential fit and the linear fit. At around day 55, the linear function starts to characterize the data significantly better. The transition period between the damped exponential and linear growth takes about two weeks.

It is interesting to compare the value of b_2 for which the transition to linear evolution appears with the value of b_1 . The transition occurs for $\sqrt{|b_2|}$ of $0.047 - 0.051 \text{ day}^{-1}$ and $0.047 - 0.049 \text{ day}^{-1}$ for Italy and Czechia, respectively. These values are roughly five times smaller than the average value of b_1 determined in Section 3.1, $b_1 = 0.25 \pm 0.04 \text{ day}^{-1}$.

It might be interesting to further explore if the transition to the linear evolution can be connected with some specific values of b_2 universally. If so, then the transition point may be predictable within the proposed parametric description of the spread. This is left open for further studies.




Figure 2: Sum of fit residuals, $\sigma|\hat{r}|$, for the damped exponential and a linear fit of the number of all registered case evaluated as a function of time for Italy.

4. Discussion

Compartmental models represent an established way to model the spread of an infectious disease. As the COVID-19 pandemic evolves, they are being tuned and improved. In general, adding complexity into the description of the problem and adding more free parameters do not automatically guarantee a better precision of the description – see e.g. discussion in Refs. [14–16]. Work presented in this paper represents an alternative to complex compartmental

modeling, which may provide new insight into the dynamics of the spread by its parametric analysis.

The spread of the disease is also often described by growth-law models, see e.g. Refs. [12, 17–19]. Compared to the growth-law models, the parametric analysis presented here starts from generic analytic functions, a form of which is constrained by the data via statistical tests. Parameters are then let free to evolve. Trends in this evolution can be studied and extrapolated to future. This allows to make predictions and compare with the data which may provide an elementary method to judge the impact of restrictive measures. For example, if the data show a larger decrease compared to the extrapolation, then one gets an evidence that the restrictive measures applied earlier are efficient. This allows to attempt to compare the impact of different restrictive measures in a uniform way. This approach may be useful since the restrictive measures are complex and therefore hard to model. The undemanding approach used here may also allow to access larger datasets.

The cumulative number of registered COVID-19 cases from datasets used in this analysis represents a complex quantity that may be influenced by various external factors. In particular, these are the number of performed diagnostic tests and their sensitivity, number of asymptomatic cases, or simply a lack of accuracy (see e.g. discussion in Refs. [3, 15, 20]). This needs to be taken into account when interpreting results such as those presented in Table 1.

5. Conclusions

The analysis presented in this paper aims to add new information helping to understand the behaviour of the COVID-19 expansion in European countries. Data collected from 20 European countries on number of registered COVID-19 cases was analyzed for a period of 15th of February to 1st of June 2020. Three regimes of the expansion are identified and quantified – early exponential expansion, damped exponential, and linear expansion. Coefficient quantifying the velocity of the expansion in the early period (b_1) and coefficient quantifying the number of cases at time zero (N_0) were evaluated and compared for 20 European countries. The b_1 coefficient was found to lie within one standard deviation from the average value for 11 out of 20 countries. The N_0 for Italy was found to be excessively high, 7 standard deviations from the average value. The transition from the damped exponential regime to the linear regime is discussed and quantified in terms of the “push-back” parameter (b_2).

A method to predict the evolution of the value of registered cases in the period of damped exponential expansion which is based on extrapolations in parametric space was put forward. It was shown that it has an average precision of 18% and 29% in the period of 15 – 55 days after the start of the damped exponential expansion for the case of Italy and Czechia, respectively. This method may provide an elementary method to quantify the impact of restrictive measures on the number of registered cases.

Acknowledgment

The author would like to thank Jiří Dolejší, Dalibor Nosek, and Alfredo Iorio for useful discussions and careful reading of the manuscript.

Disclosure statement

The author declares no conflict of interest.

Data availability statement

The data that support the findings of this study are openly available in Ref. [\[7\]](#).

References

- [1] World Health Organization - Coronavirus disease (COVID-2019) situation reports.
URL <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
- [2] C. T. Bauch, J. O. Lloyd-Smith, M. P. Coffee, A. P. Galvani, [Dynamically modeling SARS and other newly emerging respiratory illnesses](#), *Epidemiology* 16 (6) (2005) 791–801. doi:10.1097/01.ede.0000181633.80269.4c.
URL <https://doi.org/10.1097/01.ede.0000181633.80269.4c>
- [3] E. Bertuzzo, L. Mari, D. Pasetto, S. Miccoli, R. Casagrandi, M. Gatto, A. Rinaldo, [The geography of COVID-19 spread in Italy and implications for the relaxation of confinement measures](#), *Nature Communications* 11 (1). doi:10.1038/s41467-020-18050-2.
URL <https://doi.org/10.1038/s41467-020-18050-2>
- [4] C. Hou, J. Chen, Y. Zhou, L. Hua, J. Yuan, S. He, Y. Guo, S. Zhang, Q. Jia, C. Zhao, J. Zhang, G. Xu, E. Jia, [The effectiveness of quarantine of Wuhan city against the corona virus disease 2019 \(COVID-19\): A well-mixed SEIR model analysis](#), *Journal of Medical Virology* 92 (7) (2020) 841–848. doi:10.1002/jmv.25827.
URL <https://doi.org/10.1002/jmv.25827>
- [5] K. Prem, Y. Liu, T. W. Russell, A. J. Kucharski, R. M. Eggo, N. Davies, M. Jit, P. Klepac, S. Flasche, S. Clifford, C. A. B. Pearson, J. D. Munday, S. Abbott, H. Gibbs, A. Rosello, B. J. Quilty, T. Jombart, F. Sun, C. Diamond, A. Gimma, K. van Zandvoort, S. Funk, C. I. Jarvis, W. J. Edmunds, N. I. Bosse, J. Hellewell, [The effect of control strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a modelling study](#), *The Lancet Public Health* 5 (5) (2020) e261–e270. doi:10.1016/s2468-2667(20)30073-6.
URL [https://doi.org/10.1016/s2468-2667\(20\)30073-6](https://doi.org/10.1016/s2468-2667(20)30073-6)
- [6] H. W. Hethcote, [The mathematics of infectious diseases](#), *SIAM Review* 42 (4) (2000) 599–653. doi:10.1137/s0036144500371907.
URL <https://doi.org/10.1137/s0036144500371907>
- [7] [Worldometer - collection of online world statistics](#).
URL <https://www.worldometers.info/coronavirus/>
- [8] E. Dong, H. Du, L. Gardner, [An interactive web-based dashboard to track COVID-19 in real time](#), *The Lancet Infectious Diseases* doi:10.1016/s1473-3099(20)30120-1.
URL [https://doi.org/10.1016/s1473-3099\(20\)30120-1](https://doi.org/10.1016/s1473-3099(20)30120-1)
- [9] R. Brun, F. Rademakers, ROOT: An object oriented data analysis framework, *Nucl. Instrum. Meth. A* 389 (1997) 81–86. doi:10.1016/S0168-9002(97)00048-X.

- [10] A. R. Tuite, V. Ng, E. Rees, D. Fisman, [Estimation of COVID-19 outbreak size in Italy](#), The Lancet Infectious Diseases 20 (5) (2020) 537. doi:10.1016/s1473-3099(20)30227-9.
URL [https://doi.org/10.1016/s1473-3099\(20\)30227-9](https://doi.org/10.1016/s1473-3099(20)30227-9)
- [11] P. van den Driessche, J. Watmough, [Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission](#), Mathematical Biosciences 180 (1-2) (2002) 29–48. doi:10.1016/s0025-5564(02)00108-6.
URL [https://doi.org/10.1016/s0025-5564\(02\)00108-6](https://doi.org/10.1016/s0025-5564(02)00108-6)
- [12] S. Zhao, S. S. Musa, H. Fu, D. He, J. Qin, [Simple framework for real-time forecast in a data-limited situation: the Zika virus \(ZIKV\) outbreaks in Brazil from 2015 to 2016 as an example](#), Parasites & Vectors 12 (1). doi:10.1186/s13071-019-3602-9.
URL <https://doi.org/10.1186/s13071-019-3602-9>
- [13] S. Zhao, Q. Lin, J. Ran, S. S. Musa, G. Yang, W. Wang, Y. Lou, D. Gao, L. Yang, D. He, M. H. Wang, [Preliminary estimation of the basic reproduction number of novel coronavirus \(2019-nCoV\) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak](#), International Journal of Infectious Diseases 92 (2020) 214–217. doi:10.1016/j.ijid.2020.01.050.
URL <https://doi.org/10.1016/j.ijid.2020.01.050>
- [14] N. P. Jewell, J. A. Lewnard, B. L. Jewell, [Predictive mathematical models of the COVID-19 pandemic](#), JAMA 323 (19) (2020) 1893. doi:10.1001/jama.2020.6585.
URL <https://doi.org/10.1001/jama.2020.6585>
- [15] D. Adam, [Special report: The simulations driving the world’s response to COVID-19](#), Nature 580 (7803) (2020) 316–318. doi:10.1038/d41586-020-01003-6.
URL <https://doi.org/10.1038/d41586-020-01003-6>
- [16] W. C. Roda, M. B. Varughese, D. Han, M. Y. Li, [Why is it difficult to accurately predict the COVID-19 epidemic?](#), Infectious Disease Modelling 5 (2020) 271–281. doi:10.1016/j.idm.2020.03.001.
URL <https://doi.org/10.1016/j.idm.2020.03.001>
- [17] Y.-H. Hsieh, S. Ma, [Intervention measures, turning point, and reproduction number for dengue, Singapore, 2005](#), The American Journal of Tropical Medicine and Hygiene 80 (1) (2009) 66–71. doi:10.4269/ajtmh.2009.80.66.
URL <https://doi.org/10.4269/ajtmh.2009.80.66>
- [18] Y.-H. Hsieh, [Temporal patterns and geographic heterogeneity of Zika virus \(ZIKV\) outbreaks in French Polynesia and Central America](#), PeerJ 5 (2017)

e3015. [doi:10.7717/peerj.3015](https://doi.org/10.7717/peerj.3015).
URL <https://doi.org/10.7717/peerj.3015>

- [19] P. Castorina, A. Iorio, D. Lanteri, [Data analysis on coronavirus spreading by macroscopic growth laws](#), International Journal of Modern Physics C 31 (07) (2020) 2050103. [doi:10.1142/s012918312050103x](https://doi.org/10.1142/s012918312050103x).
URL <https://doi.org/10.1142/s012918312050103x>
- [20] Y. Bai, L. Yao, T. Wei, F. Tian, D.-Y. Jin, L. Chen, M. Wang, [Presumed asymptomatic carrier transmission of COVID-19](#), JAMA 323 (14) (2020) 1406. [doi:10.1001/jama.2020.2565](https://doi.org/10.1001/jama.2020.2565).
URL <https://doi.org/10.1001/jama.2020.2565>