

(Θ_n, sl_n) - GRADED LIE ALGEBRAS ($n = 3, 4$)

HOGIR MOHAMMED YASEEN

ABSTRACT. Let \mathbb{F} be a field of characteristic zero and let \mathfrak{g} be a non-zero finite-dimensional split semisimple Lie algebra with root system Δ . Let Γ be a finite set of integral weights of \mathfrak{g} containing Δ and $\{0\}$. Following [2, 10], we say that a Lie algebra L over \mathbb{F} is *generalized root graded*, or more exactly (Γ, \mathfrak{g}) -*graded*, if L contains a semisimple subalgebra isomorphic to \mathfrak{g} , the \mathfrak{g} -module L is the direct sum of its weight subspaces L_α ($\alpha \in \Gamma$) and L is generated by all L_α with $\alpha \neq 0$ as a Lie algebra. Let $\mathfrak{g} \cong sl_n$ and

$$\Theta_n = \{0, \pm\varepsilon_i \pm \varepsilon_j, \pm\varepsilon_i, \pm 2\varepsilon_i \mid 1 \leq i \neq j \leq n\}$$

where $\{\varepsilon_1, \dots, \varepsilon_n\}$ is the set of weights of the natural sl_n -module. In [9], we classify (Θ_n, sl_n) -graded Lie algebras for $n > 4$. In this paper we describe the multiplicative structures and the coordinate algebras of (Θ_n, sl_n) -graded Lie algebras ($n = 3, 4$). In $n = 3$, we assume that

$$[V(2\omega_1) \otimes C, V(2\omega_1) \otimes C] = [V(2\omega_2) \otimes C', V(2\omega_2) \otimes C'] = 0$$

where $V(\omega)$ is the simple \mathfrak{g} -module of highest weight ω , $C = \text{Hom}_{\mathfrak{g}}(V(2\omega_1), L)$ and $C' = \text{Hom}_{\mathfrak{g}}(V(2\omega_2), L)$.

CONTENTS

1. Introduction	1
2. tensor product decompositions for the modules in Θ_n^+ ($n = 3, 4$).	2
3. Multiplication in (Θ_n, sl_n) -graded Lie algebras ($n = 3, 4$)	5
3.1. Multiplication in Θ_3 -graded Lie algebras	5
3.2. Multiplication in Θ_4 -graded Lie algebras	8
4. Coordinate algebra of (Θ_n, sl_n) -graded Lie algebras ($n = 3, 4$)	10
4.1. Unital algebra \mathfrak{a}	10
4.2. Coordinate algebra \mathfrak{b}	13
References	15

1. INTRODUCTION

In 1992, Berman and Moody introduced root-graded Lie algebras to study toroidal Lie algebras and Slodowy matrix algebras intersection. Nonetheless, this concept previously appeared in the study of simple Lie algebras by Seligman [9]. Root graded Lie algebras of simply-laced finite root systems were classified up to central isogeny by Berman and Moody in [7]. The case of double-laced finite root systems was settled by Benkart and Zelmanov [6]. Non-reduced systems BC_n were considered by Allison, Benkart and Gao [1] (for $n \geq 2$) and by Benkart and Smirnov [5] (for $n = 1$). The aim of this paper is to

describe the multiplicative structures and the coordinate algebras of (Θ_n, sl_n) -graded Lie algebras where $n = 3, 4$ and

$$\Theta_n = \{0, \pm\varepsilon_i \pm \varepsilon_j, \pm\varepsilon_i, \pm 2\varepsilon_i \mid 1 \leq i \neq j \leq n\}$$

and $\{\varepsilon_1, \dots, \varepsilon_n\}$ is the set of weights of the natural sl_n -module. Throughout the paper, the ground field \mathbb{F} is of characteristic zero and the grading subalgebra $\mathfrak{g} \cong sl_n$. We denote by Θ_n^+ the set of dominant weights in Θ_n and the corresponding simple sl_n -modules. Thus,

$$\Theta_4^+ = \{\omega_1 + \omega_3 = \varepsilon_1 - \varepsilon_4, \omega_1 = \varepsilon_1, \omega_3 = -\varepsilon_4, 2\omega_1 = 2\varepsilon_1, 2\omega_3 = -2\varepsilon_4, \omega_2 = \varepsilon_1 + \varepsilon_2, 0\},$$

$$\Theta_3^+ = \{\omega_1 + \omega_3 = \varepsilon_1 - \varepsilon_4, \omega_1 = \varepsilon_1, \omega_3 = -\varepsilon_4, 2\omega_1 = 2\varepsilon_1, 2\omega_3 = -2\varepsilon_4, 0\}.$$

These are the highest weights of the irreducible \mathfrak{g} -modules. First we compute tensor product decompositions for the modules in Θ_n^+ . Then we describe the multiplicative structures of (Θ_n, sl_n) -graded Lie algebras. The coordinate algebra of these Lie algebras are described in Section 4.

2. TENSOR PRODUCT DECOMPOSITIONS FOR THE MODULES IN Θ_n^+ ($n = 3, 4$).

We begin with the general definition of Lie algebras graded by finite weight systems.

Definition 2.1. [2] Let Δ be a root system and let Γ be a finite set of integral weights of Δ containing Δ and $\{0\}$. A Lie algebra L is called (Γ, \mathfrak{g}) -graded (or simply Γ -graded) if

($\Gamma 1$) L contains as a subalgebra a non-zero finite-dimensional split semisimple Lie algebra

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_\alpha,$$

whose root system is Δ relative to a split Cartan subalgebra $\mathfrak{h} = \mathfrak{g}_0$;

($\Gamma 2$) $L = \bigoplus_{\alpha \in \Gamma} L_\alpha$ where $L_\alpha = \{x \in L \mid [h, x] = \alpha(h)x \text{ for all } h \in \mathfrak{h}\}$;

($\Gamma 3$) $L_0 = \sum_{\alpha, -\alpha \in \Gamma \setminus \{0\}} [L_\alpha, L_{-\alpha}]$.

The subalgebra \mathfrak{g} is called the *grading subalgebra* of L . If \mathfrak{g} is the split simple Lie algebra and $\Gamma = \Delta \cup \{0\}$ then L is said to be *root-graded*.

We fix a base $\Pi = \{\varepsilon_i - \varepsilon_{i+1} \text{ for } i = 1, 2, \dots, n-1\}$ of simple roots for the root system $A_{n-1} = \{\varepsilon_i - \varepsilon_j \mid 1 \leq i \neq j \leq n\}$. Let L be a Lie algebra containing a non-zero split simple subalgebra \mathfrak{g} . Using the same arguments of [10, Lemma 3.1.2 and Proposition 3.2.2] we get the following:

(1) A Lie algebra L is (Θ_3, \mathfrak{g}) -graded if and only if L is generated by \mathfrak{g} as an ideal and the \mathfrak{g} -module L decomposes into copies of the adjoint module, its symmetric S^2V , its exterior squares \wedge^2V , their duals and the one dimensional trivial \mathfrak{g} -module.

(2) A Lie algebra L is (Θ_4, \mathfrak{g}) -graded if and only if L is generated by \mathfrak{g} as an ideal and the \mathfrak{g} -module L decomposes into copies of the adjoint module (we will denote it by the same letter \mathfrak{g}), the natural module V , its exterior squares \wedge^2V , its symmetric S^2V , their duals and the one dimensional trivial \mathfrak{g} -module.

Thus, by collecting isotypic components, we get the following decomposition of the \mathfrak{g} -module L :

(2.1)

$$L = (\mathfrak{g} \otimes A) \oplus (S \otimes C) \oplus (S' \otimes C') \oplus (\Lambda \otimes E) \oplus (\Lambda' \otimes E') \oplus D \text{ for } n = 3;$$

$$L = (\mathfrak{g} \otimes A) \oplus (V \otimes B) \oplus (V' \otimes B') \oplus (S \otimes C) \oplus (S' \otimes C') \oplus (\Lambda \otimes E) \oplus D \text{ for } n = 4;$$

where A, B, B', C, C', E are vector spaces,

$$\begin{aligned} \mathfrak{g} &:= V(\omega_1 + \omega_{n-1}), & V &:= V(\omega_1), & V' &:= V(\omega_{n-1}), \\ S &:= V(2\omega_1), & S' &:= V(2\omega_{n-1}), & \Lambda &:= V(\omega_2), & \Lambda' &:= V(\omega_{n-2}) \end{aligned}$$

and D is the sum of the trivial \mathfrak{g} -modules. Note that $\Lambda \cong \Lambda'$ for $n = 4$.

Alternatively, these spaces can also be viewed as the corresponding \mathfrak{g} -mod Hom-spaces: $A = \text{Hom}_{\mathfrak{g}}(\mathfrak{g}, L)$, $B = \text{Hom}_{\mathfrak{g}}(V, L)$, etc, so for each simple \mathfrak{g} -module M , the space $M \otimes \text{Hom}_{\mathfrak{g}}(M, L)$ is canonically identified with the M -isotypic component of L via the evaluation map

$$(2.2) \quad M \otimes \text{Hom}_{\mathfrak{g}}(M, L) \rightarrow L, \quad m \otimes \varphi \mapsto \varphi(m),$$

see [8, Proposition 4.1.15].

We can write the following examples of Θ_n -graded Lie algebras $(n = 3, 4)$:

1) Let $L = sl_{n+k}$ and let \mathfrak{g} be the copy of sl_n in the northwest corner. We consider the adjoint action of \mathfrak{g} on L . Then the \mathfrak{g} -module L decomposes into k copies of the natural module $V = \mathbb{F}^n$, k copies of the dual module $V' = \text{Hom}(V, \mathbb{F})$, an adjoint module \mathfrak{g} and one dimensional trivial \mathfrak{g} -modules in its southeast corner. Then

$$L = \mathfrak{g} \oplus V^{\oplus k} \oplus V'^{\oplus k} \oplus D$$

where D is the sum of the trivial sl_n -modules. As a result, we may write

$$L = \mathfrak{g} \oplus (V \otimes B) \oplus (V' \otimes B') \oplus D$$

where $B \cong B' \cong \mathbb{F}^k$. Then L is (A_{n-1}, \mathfrak{g}) -graded. Bahturin and Benkart [3] (for $n > 3$) and Benkart and Elduque [4] (for $n = 3$) described the multiplicative structure of this type of Lie algebras.

2) Any Lie algebra which is (A_{n-1}, sl_n) -graded is also Θ_n -graded. For such a Lie algebra, the space $L_\alpha = \{0\}$ for all α not in A_{n-1} .

3) Let $L = sl_{2n+1}$ and $\mathfrak{g} = \left\{ \left[\begin{array}{ccc} x & 0 & 0 \\ 0 & -x^t & 0 \\ 0 & 0 & 0 \end{array} \right] \mid x \in sl_n \right\} \subset L$ where $n = 3, 4$. We

consider the adjoint action of \mathfrak{g} on L . Then L is (Θ_n, \mathfrak{g}) -graded .

Definition 2.2. (1) We identify the \mathfrak{g} -modules V and V' with the space $\mathbb{F}^n (n = 3, 4)$ of column vectors with the following actions:

$$\begin{aligned} x.v &= xv \text{ for } x \in sl_n, v \in V, \\ x.v' &= -x^t v' \text{ for } x \in sl_n, v' \in V'. \end{aligned}$$

(2) We identify S and S' (resp. Λ and Λ') with symmetric (resp. skew-symmetric) $n \times n$ matrices over \mathbb{F} ($n = 3, 4$). Then S, S', Λ and Λ' are \mathfrak{g} -modules under the actions:

$$\begin{aligned} x.s &= xs + sx^t \text{ for } x \in sl_n, s \in S, \\ x.\lambda &= x\lambda + \lambda x^t \text{ for } x \in sl_n, \lambda \in \Lambda, \\ x.s' &= -s'x - x^t s' \text{ for } x \in sl_n, s' \in S, \\ x.\lambda' &= -\lambda'x - x^t \lambda' \text{ for } x \in sl_n, \lambda' \in \Lambda'. \end{aligned}$$

Since the subalgebra \mathfrak{g} of L is a \mathfrak{g} -submodule, there exists a distinguished element 1 of A such that $\mathfrak{g} = \mathfrak{g} \otimes 1$. In particular,

$$(2.3) \quad [x \otimes 1, y \otimes b] = x.y \otimes b.$$

where $x \otimes 1$ is in $\mathfrak{g} \otimes 1$, $y \otimes b$ belongs to one of the components in (2.1), and $x.y$ is as in Definition 2.2.

From equation 2.1 and Definition 2.2 and the properties that $V \cong \Lambda'$ and $V' \cong \Lambda$ for sl_3 , we get the following decomposition of the \mathfrak{g} -module L :

$$(2.4) \quad L = (\mathfrak{g} \otimes A) \oplus (S \otimes C) \oplus (S' \otimes C') \oplus (\Lambda \otimes E) \oplus (\Lambda' \otimes E') \oplus D$$

From equation 2.1 and Definition 2.2 and the property that $\Lambda \cong \Lambda'$ for sl_4 , we get the following decomposition of the \mathfrak{g} -module L :

$$(2.5) \quad L = (\mathfrak{g} \otimes A) \oplus (V \otimes B) \oplus (V' \otimes B') \oplus (S \otimes C) \oplus (S' \otimes C') \oplus (\Lambda \otimes E) \oplus D.$$

Using [10, Section 3.4], in Tables 1-2 we get the following Θ -components of all tensor product decompositions for the modules in Θ_n^+ ($n = 3, 4$). If the cell in row X and column Y contains Z this means that $\Theta(X \otimes Y) \cong Z$.

\otimes	\mathfrak{g}	S	S'	$V \cong \Lambda'$	$V' \cong \Lambda$
\mathfrak{g}	$\mathfrak{g} + \mathfrak{g} + T$	$S + \Lambda$	$S' + \Lambda'$	$S' + \Lambda'$	$S + \Lambda$
S	$S + \Lambda$	S'	$\mathfrak{g} + T$	\mathfrak{g}	Λ'
S'	$S' + \Lambda'$	$\mathfrak{g} + T$	S	Λ	\mathfrak{g}
V	$S' + \Lambda'$	\mathfrak{g}	Λ	$S + \Lambda$	$\mathfrak{g} + T$
V'	$S + \Lambda$	Λ'	\mathfrak{g}	$\mathfrak{g} + T$	$S' + \Lambda'$

TABLE 1. Θ -component of tensor product decompositions for sl_3

\otimes	\mathfrak{g}	S	$\Lambda \cong \Lambda'$	S'	V	V'
\mathfrak{g}	$\mathfrak{g} + \mathfrak{g} + T$	$S + \Lambda$	$S + \Lambda + S'$	$S' + \Lambda$	V	V'
S	$S + \Lambda$	0	\mathfrak{g}	$\mathfrak{g} + T$	0	V
Λ	$S + \Lambda + S'$	\mathfrak{g}	$\mathfrak{g} + T$	\mathfrak{g}	V'	V
S'	$S' + \Lambda$	$\mathfrak{g} + T$	\mathfrak{g}	0	V'	0
V	V	0	V'	V'	$S + \Lambda$	$\mathfrak{g} + T$
V'	V'	V	V	0	$\mathfrak{g} + T$	$S' + \Lambda$

TABLE 2. Θ -component of tensor product decompositions for sl_4

In Tables and 4 below , if the cell in row X and column Y contains Z , this means that there is a bilinear map $X \otimes Y \rightarrow Z$ given by $x \otimes y \mapsto (x, y)_Z$. For simplicity of notation, we will write dy instead of $(d, y)_D$ if $X = Z = D$ and we will write $\langle x, y \rangle$ instead of $(x, y)_D$ if $X, Y \neq D$ and $Z = D$. In the case $X = Y = Z = A$, we have two bilinear products $a_1 \otimes a_2 \mapsto a_1 \circ a_2$ and $a_1 \otimes a_2 \mapsto [a_1, a_2]$ for $a_1, a_2 \in A$. Note that some of the cells are empty. The corresponding products $X \otimes Y \rightarrow Z$ will be defined later by extending the existing maps $Y \otimes X \rightarrow Z$. This will make the table symmetric.

.	A	C	C'	E	E'	D
A	$(A, \circ, []), D$	C, E		C, E		
C		0	A, D	E'	A	
C'	C', E'		0	A	E	
E		E'		C', E'	A, D	
E'	C', E'		E		C, E	
D	A	C	C'	E	E'	D

TABLE 3. Bilinear products for $n = 3$

.	A	B	B'	C	C'	E	D
A	$(A, \circ, []), D$	B		C, E		C, E, C'	
B		C, E	A, D	0		0	
B'	A		C', E	B	0	B	
C		0		0	A, D	0	
C'	C', E	B'	0		0	A	
E		0		0		0	
D	A	B	B'	C	C'	E	D

TABLE 4. Bilinear products for $n = 4$

.Let x and y be $n \times n$ matrices. We will use the following products:

$$\begin{aligned}
 [x, y] &= xy - yx, \\
 x \circ y &= xy + yx - \frac{2}{n} \text{tr}(xy)I, \\
 x \diamond y &= xy + yx, \\
 (x | y) &= \frac{1}{n} \text{tr}(xy).
 \end{aligned}$$

3. MULTIPLICATION IN (Θ_n, sl_n) -GRADED LIE ALGEBRAS ($n = 3, 4$)

3.1. Multiplication in Θ_3 -graded Lie algebras. Let L be a Θ_3 -graded Lie algebra with the grading subalgebra $\mathfrak{g} \cong sl_3$ and the properties that

$$(3.1) \quad [V(2\omega_1) \otimes C, V(2\omega_1) \otimes C] = [V(2\omega_2) \otimes C', V(2\omega_2) \otimes C'] = 0.$$

where $V(\omega)$ is the simple \mathfrak{g} -module of highest weight ω , $C = \text{Hom}_{\mathfrak{g}}(V(2\omega_1), L)$ and $C' = \text{Hom}_{\mathfrak{g}}(V(2\omega_2), L)$. In 2.4, we show that L is the direct sum of finite-dimensional irreducible \mathfrak{g} -modules whose highest weights are in Θ_3^+ , i.e. as a \mathfrak{g} -module,

$$L = (\mathfrak{g} \otimes A) \oplus (S \otimes C) \oplus (S' \otimes C') \oplus (\Lambda \otimes E) \oplus (\Lambda' \otimes E') \oplus D.$$

Note that $V' = U(sl_3)e_3$ and $\Lambda = U(sl_3)(E_{1,2} - E_{2,1})$ are highest weight modules with highest weight ω_2 where $E_{i,j}$ denote the matrix units and $e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ (resp. $V = U(sl_4)e_1$ and $\Lambda' = U(sl_4)(E_{3,4} - E_{4,3})$ are highest weight module with highest weight ω_1 where $e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$). Define $f : \Lambda' \rightarrow V$ by $f(x.(E_{3,4} - E_{4,3})) = x.e_1$ and $g : \Lambda \rightarrow V'$ by $g(x.(E_{1,2} - E_{2,1})) = x.e_3$ for all $x \in U(sl_4)$. This allows us to identify Λ with V' and Λ' with V .

In (3.2) we list bases for all non-zero \mathfrak{g} -module homomorphism spaces $\text{Hom}_{\mathfrak{g}}(X \otimes Y, Z)$ where $X, Y, Z \in \{\mathfrak{g}, V, V', S, S', T\}$ and X and Y are both non-trivial. Note that all of them are 1-dimensional except the first one (which is 2-dimensional).

$$(3.2) \quad \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{g}) = \text{span}\{x \otimes y \mapsto xy - yx, x \otimes y \mapsto xy + yx - \frac{2}{3} \text{tr}(xy)I\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda', \mathfrak{g}) = \text{span}\{\lambda \otimes \lambda' \mapsto \lambda\lambda' - \frac{\text{tr}(\lambda\lambda')}{3}I\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda, \Lambda') = \text{span}\{\lambda_1 \otimes \lambda_2 \mapsto f(\lambda'_1)(f(\lambda'_2))^t - f(\lambda'_2)(f(\lambda'_1))^t\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda, S') = \text{span}\{\lambda_1 \otimes \lambda_2 \mapsto f(\lambda'_1)(f(\lambda'_2))^t + f(\lambda'_2)(f(\lambda'_1))^t\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda, \Lambda) = \text{span}\{\lambda_1 \otimes \lambda_2 \mapsto g(\lambda_1)(g(\lambda_2))^t - g(\lambda_2)(g(\lambda_1))^t\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda, S) = \text{span}\{\lambda_1 \otimes \lambda_2 \mapsto g(\lambda_1)(g(\lambda_2))^t + g(\lambda_2)(g(\lambda_1))^t\},$$

$$\text{Hom}_{\mathfrak{g}}(S \otimes \Lambda', \mathfrak{g}) = \text{span}\{s \otimes \lambda' \mapsto s\lambda'\},$$

$$\text{Hom}_{\mathfrak{g}}(S' \otimes \Lambda, \mathfrak{g}) = \text{span}\{s' \otimes \lambda \mapsto s'\lambda\},$$

$$\text{Hom}_{\mathfrak{g}}(S \otimes S', \mathfrak{g}) = \text{span}\{s \otimes s' \mapsto ss' - \frac{\text{tr}(ss')}{3}I\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda' \otimes \mathfrak{g}, \Lambda') = \text{span}\{\lambda' \otimes x \mapsto \lambda'x + x^t\lambda'\},$$

$$\text{Hom}_{\mathfrak{g}}(S \otimes \Lambda, \Lambda') = \text{span}\{s \otimes \lambda \mapsto sg(\lambda)\},$$

$$\text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \Lambda, S) = \text{span}\{x \otimes \lambda \mapsto x\lambda - \lambda x^t\},$$

$$\text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \Lambda, \Lambda) = \text{span}\{x \otimes \lambda \mapsto x\lambda + \lambda x^t\},$$

$$\text{Hom}_{\mathfrak{g}}(S' \otimes \Lambda', \Lambda) = \text{span}\{s' \otimes \lambda' \mapsto s'f(\lambda')\},$$

$$\text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes S, S) = \text{span}\{x \otimes s \mapsto xs + sx^t\},$$

$$\text{Hom}_{\mathfrak{g}}(S' \otimes \mathfrak{g}, S') = \text{span}\{s' \otimes x \mapsto s'x + x^t s'\},$$

$$\text{Hom}_{\mathfrak{g}}(\Lambda' \otimes \mathfrak{g}, S') = \text{span}\{\lambda' \otimes x \mapsto \lambda'x - x^t\lambda'\},$$

$$\begin{aligned} \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes S, \Lambda) &= \text{span}\{x \otimes s \mapsto xs - sx^t\}, \\ \text{Hom}_{\mathfrak{g}}(S' \otimes \mathfrak{g}, \Lambda') &= \text{span}\{s' \otimes x \mapsto s'x - x^t s'\}, \\ \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \mathfrak{g}, T) &= \text{span}\{x_1 \otimes x_2 \mapsto \frac{1}{3} \text{tr}(x_1 x_2)\}, \\ \text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda', T) &= \text{span}\{\lambda \otimes \lambda' \mapsto \frac{1}{3} \text{tr}(\lambda \lambda')\}. \\ \text{Hom}_{\mathfrak{g}}(S \otimes S', T) &= \text{span}\{s \otimes s' \mapsto \frac{1}{3} \text{tr}(s s')\}, \end{aligned}$$

Following the methods in [1, 9, 7, 6], using the results of (3.2) and Table 1, we may suppose that the multiplication in L is given as follows. For all $x, y \in sl_3$, $s \in S$, $\lambda_1, \lambda_2 \in \Lambda$, $s' \in S'$, $\lambda'_1, \lambda'_2 \in \Lambda'$ and for all $a, a_1, a_2 \in A$, $c \in C$, $c' \in C'$, $e \in E$, $e' \in E'$ and $d, d_1, d_2 \in D$,

(3.3)

$$\begin{aligned} [x \otimes a_1, y \otimes a_2] &= (x \circ y) \otimes \frac{[a_1, a_2]}{2} + [x, y] \otimes \frac{a_1 \circ a_2}{2} + (x | y) \langle a_1, a_2 \rangle, \\ [\lambda \otimes e, \lambda' \otimes e'] &= (\lambda \lambda' - (\lambda | \lambda') I) \otimes (e, e')_A + (\lambda | \lambda') \langle e, e' \rangle = -[\lambda' \otimes e', \lambda \otimes e], \\ [\lambda_1 \otimes e_1, \lambda_2 \otimes e_2] &= (g(\lambda_1)(g(\lambda_2))^t + g(\lambda_2)(g(\lambda_1))^t) \otimes \frac{(e_1, e_2)_C}{2} \\ &\quad + (g(\lambda_1)(g(\lambda_2))^t - g(\lambda_2)(g(\lambda_1))^t) \otimes \frac{(e_1, e_2)_E}{2}, \\ [\lambda'_1 \otimes e'_1, \lambda'_2 \otimes e'_2] &= (f(\lambda'_1)(f(\lambda'_2))^t + f(\lambda'_2)(f(\lambda'_1))^t) \otimes \frac{(e'_1, e'_2)_{C'}}{2} \\ &\quad + (f(\lambda'_1)(f(\lambda'_2))^t - f(\lambda'_2)(f(\lambda'_1))^t) \otimes \frac{(e'_1, e'_2)_{E'}}{2}, \\ [s \otimes c, s' \otimes c'] &= (s s' - (s | s') I) \otimes (c, c')_A + (s | s') \langle c, c' \rangle = -[s' \otimes c', s \otimes c], \\ [x \otimes a, s \otimes c] &= (x s + s x^t) \otimes \frac{(a, c)_C}{2} + (x s - s x^t) \otimes \frac{(a, c)_E}{2} = -[s \otimes c, x \otimes a], \\ [x \otimes a, \lambda \otimes e] &= (x \lambda + \lambda x^t) \otimes \frac{(a, e)_E}{2} + (x \lambda - \lambda x^t) \otimes \frac{(a, e)_C}{2} = -[\lambda \otimes e, x \otimes a], \\ [s' \otimes c', x \otimes a] &= (s' x + x^t s') \otimes \frac{(c', a)_{C'}}{2} + (s' x - x^t s') \otimes \frac{(c', a)_{E'}}{2} = -[x \otimes a, s' \otimes c'], \\ [\lambda' \otimes e', x \otimes a] &= (\lambda' x + x^t \lambda') \otimes \frac{(e', a)_{E'}}{2} + (\lambda' x - x^t \lambda') \otimes \frac{(e', a)_{C'}}{2} = -[x \otimes a, \lambda' \otimes e'], \\ [s \otimes c, \lambda' \otimes e'] &= s \lambda' \otimes (c, e')_A = -[\lambda' \otimes e', s \otimes c], \\ [s' \otimes c', \lambda \otimes e] &= s' \lambda \otimes (c', e)_A = -[\lambda \otimes e, s' \otimes c'], \\ [s' \otimes c', \lambda' \otimes e'] &= s' f(\lambda') \otimes (c', e')_E = -[\lambda' \otimes e', s' \otimes c'], \\ [\lambda \otimes e, s \otimes c] &= s g(\lambda) \otimes (e, c)_{E'} = -[s \otimes c, \lambda \otimes e], \\ [d, x \otimes a] &= x \otimes da = -[x \otimes a, d], \\ [d, s \otimes c] &= s \otimes dc = -[s \otimes c, d], \end{aligned}$$

$$\begin{aligned} [d, \lambda \otimes e] &= \lambda \otimes de = -[\lambda \otimes e, d], \\ [d, s' \otimes c'] &= s' \otimes dc' = -[s' \otimes c', d], \\ [d, \lambda' \otimes e'] &= \lambda' \otimes de' = -[\lambda' \otimes e', d], \\ [d_1, d_2] &\in D, \end{aligned}$$

3.2. Multiplication in Θ_4 -graded Lie algebras. Let L be a Θ_4 -graded Lie algebra with the grading subalgebra $\mathfrak{g} \cong sl_4$. In 2.4, we show that L is the direct sum of finite-dimensional irreducible \mathfrak{g} -modules whose highest weights are in Θ_3^+ , i.e. as a \mathfrak{g} -module,

$$L = (\mathfrak{g} \otimes A) \oplus (V \otimes B) \oplus (V' \otimes B') \oplus (S \otimes C) \oplus (S' \otimes C') \oplus (\Lambda \otimes E) \oplus D.$$

Note that $\Lambda = U(sl_4)(E_{1,2} - E_{2,1})$ and $\Lambda' = U(sl_4)(E_{3,4} - E_{4,3})$ are highest weight modules with highest weight ω_2 . Define $f : \Lambda' \rightarrow \Lambda$ by $f(x.(E_{3,4} - E_{4,3})) = x.(E_{1,2} - E_{2,1})$ for all $x \in U(sl_4)$. This allows us to identify Λ with Λ' .

In (3.4) we list bases for all non-zero \mathfrak{g} -module homomorphism spaces $\text{Hom}_{\mathfrak{g}}(X \otimes Y, Z)$ where $X, Y, Z \in \{\mathfrak{g}, V, V', S, \Lambda, S', T\}$ and X and Y are both non-trivial. Note that all of them are 1-dimensional except the first one (which is 2-dimensional).

$$\begin{aligned} (3.4) \quad \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \mathfrak{g}, \mathfrak{g}) &= \text{span}\{x \otimes y \mapsto xy - yx, x \otimes y \mapsto xy + yx - \frac{2}{4} \text{tr}(xy)I\}, \\ \text{Hom}_{\mathfrak{g}}(V \otimes V', \mathfrak{g}) &= \text{span}\{u \otimes v' \mapsto uv'^t - \frac{\text{tr}(uv'^t)}{4}I\}, \\ \text{Hom}_{\mathfrak{g}}(S \otimes \Lambda, \mathfrak{g}) &= \text{span}\{s \otimes \lambda \mapsto sf^{-1}(\lambda)\}, \\ \text{Hom}_{\mathfrak{g}}(S' \otimes \Lambda, \mathfrak{g}) &= \text{span}\{s' \otimes \lambda \mapsto s'\lambda\}, \\ \text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda, \mathfrak{g}) &= \text{span}\{\lambda \otimes \lambda' \mapsto \lambda f^{-1}(\lambda') - \frac{\text{tr}(\lambda f^{-1}(\lambda'))}{4}I\}, \\ \text{Hom}_{\mathfrak{g}}(S \otimes S', \mathfrak{g}) &= \text{span}\{s \otimes s' \mapsto ss' - \frac{\text{tr}(ss')}{4}I\}, \\ \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes V, V) &= \text{span}\{x \otimes v \mapsto xv\}, \\ \text{Hom}_{\mathfrak{g}}(\Lambda \otimes V', V) &= \text{span}\{\lambda \otimes v' \mapsto \lambda v'\}, \\ \text{Hom}_{\mathfrak{g}}(S \otimes V', V) &= \text{span}\{s \otimes v' \mapsto sv'\}, \\ \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes V', V') &= \text{span}\{x \otimes v' \mapsto x^t v'\}, \\ \text{Hom}_{\mathfrak{g}}(S' \otimes V, V') &= \text{span}\{s' \otimes v \mapsto s'v\}, \\ \text{Hom}_{\mathfrak{g}}(\Lambda \otimes V, V') &= \text{span}\{\lambda \otimes v \mapsto f^{-1}(\lambda)v\}, \\ \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes S, S) &= \text{span}\{x \otimes s \mapsto xs + sx^t\}, \\ \text{Hom}_{\mathfrak{g}}(V \otimes V, S) &= \text{span}\{u \otimes v \mapsto uv^t + vu^t\}, \\ \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \Lambda, S) &= \text{span}\{x \otimes \lambda \mapsto x\lambda - \lambda x^t\}, \\ \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \Lambda, S') &= \text{span}\{x \otimes \lambda \mapsto f^{-1}(\lambda)x - x^t f^{-1}(\lambda)\}, \\ \text{Hom}_{\mathfrak{g}}(S' \otimes \mathfrak{g}, S') &= \text{span}\{s' \otimes x \mapsto s'x + x^t s'\}, \end{aligned}$$

$$\begin{aligned}
 \text{Hom}_{\mathfrak{g}}(V' \otimes V', S') &= \text{span}\{u' \otimes v' \mapsto u'v'^t + v'u'^t\}, \\
 \text{Hom}_{\mathfrak{g}}(\Lambda' \otimes \mathfrak{g}, S') &= \text{span}\{\lambda' \otimes x \mapsto \lambda'x - x^t\lambda'\}, \\
 \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \Lambda, \Lambda) &= \text{span}\{x \otimes \lambda \mapsto x\lambda + \lambda x^t\}, \\
 \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes S, \Lambda) &= \text{span}\{x \otimes s \mapsto xs - sx^t\}, \\
 \text{Hom}_{\mathfrak{g}}(V \otimes V, \Lambda) &= \text{span}\{u \otimes v \mapsto uv^t - vu^t\}, \\
 \text{Hom}_{\mathfrak{g}}(S' \otimes \mathfrak{g}, \Lambda) &= \text{span}\{s' \otimes x \mapsto f(s'x - x^t s')\}, \\
 \text{Hom}_{\mathfrak{g}}(V' \otimes V', \Lambda) &= \text{span}\{u' \otimes v' \mapsto f(u'v'^t - v'u'^t)\}, \\
 \text{Hom}_{\mathfrak{g}}(\mathfrak{g} \otimes \mathfrak{g}, T) &= \text{span}\{x_1 \otimes x_2 \mapsto \frac{1}{4} \text{tr}(x_1 x_2)\}, \\
 \text{Hom}_{\mathfrak{g}}(V' \otimes V, T) &= \text{span}\{v^t \otimes u \mapsto \frac{1}{4} \text{tr}(uv^t)\}, \\
 \text{Hom}_{\mathfrak{g}}(S \otimes S', T) &= \text{span}\{s \otimes s' \mapsto \frac{1}{4} \text{tr}(ss')\}, \\
 \text{Hom}_{\mathfrak{g}}(\Lambda \otimes \Lambda, T) &= \text{span}\{\lambda \otimes \lambda' \mapsto \frac{1}{4} \text{tr}(\lambda f^{-1}(\lambda'))\}.
 \end{aligned}$$

Following the methods in [1, 9, 7, 6], using the results of (3.4) and Table 2, we may suppose that the multiplication in L is given as follows. For all $x, y \in sl_4$, $u, v \in V$, $u', v' \in V'$, $s \in S$, $\lambda \in \Lambda$, $s' \in S'$, $\lambda' \in \Lambda'$ and for all $a, a_1, a_2 \in A$, $b, b_1, b_2 \in B$, $b', b'_1, b'_2 \in B'$, $c \in C$, $c' \in C'$, $e \in E$ and $d, d_1, d_2 \in D$,

(3.5)

$$\begin{aligned}
 [x \otimes a_1, y \otimes a_2] &= (x \circ y) \otimes \frac{[a_1, a_2]}{2} + [x, y] \otimes \frac{a_1 \circ a_2}{2} + (x | y) \langle a_1, a_2 \rangle, \\
 [u \otimes b, v' \otimes b'] &= (uv'^t - \frac{\text{tr}(uv'^t)}{n} I) \otimes (b, b')_A + \frac{2}{n} \text{tr}(uv'^t) \langle b, b' \rangle = -[v' \otimes b', u \otimes b], \\
 [s \otimes c, s' \otimes c'] &= (ss' - (s | s') I) \otimes (c, c')_A + (s | s') \langle c, c' \rangle = -[s' \otimes c', s \otimes c], \\
 [\lambda_1 \otimes e_1, \lambda_2 \otimes e_2] &= (\lambda_1 f^{-1}(\lambda_2) - (\lambda_1 | f^{-1}(\lambda_2)) I) \otimes (e_1, e_2)_A + (\lambda_1 | f^{-1}(\lambda_2)) \langle e_1, e_2 \rangle, \\
 [u \otimes b_1, v \otimes b_2] &= (uv^t + vu^t) \otimes \frac{(b_1, b_2)_C}{2} + (uv^t - vu^t) \otimes \frac{(b_1, b_2)_E}{2}, \\
 [u' \otimes b'_1, v' \otimes b'_2] &= (u'v'^t + v'u'^t) \otimes \frac{(b'_1, b'_2)_{C'}}{2} + f(u'v'^t - v'u'^t) \otimes \frac{(b'_1, b'_2)_E}{2}, \\
 [x \otimes a, s \otimes c] &= (xs + sx^t) \otimes \frac{(a, c)_C}{2} + (xs - sx^t) \otimes \frac{(a, c)_E}{2} = -[s \otimes c, x \otimes a], \\
 [x \otimes a, \lambda \otimes e] &= (x\lambda + \lambda x^t) \otimes \frac{(a, e)_E}{2} + (x\lambda - \lambda x^t) \otimes \frac{(a, e)_C}{2}, \\
 &\quad + (f^{-1}(\lambda)x - x^t f^{-1}(\lambda)) \otimes \frac{(a, e)_{C'}}{2} = -[\lambda \otimes e, x \otimes a] \\
 [s' \otimes c', x \otimes a] &= (s'x + x^t s') \otimes \frac{(c', a)_{C'}}{2} + f(s'x - x^t s') \otimes \frac{(c', a)_E}{2} = -[x \otimes a, s' \otimes c'], \\
 [s \otimes c, \lambda \otimes e] &= s f^{-1}(\lambda) \otimes (c, e')_A = -[\lambda \otimes e, s \otimes c],
 \end{aligned}$$

$$\begin{aligned}
 [s' \otimes c', \lambda \otimes e] &= s' \lambda \otimes (c', e)_A = -[\lambda \otimes e, s' \otimes c'], \\
 [x \otimes a, u \otimes b] &= xu \otimes (a, b)_B = -[u \otimes b, x \otimes a], \\
 [s' \otimes c', u \otimes b] &= s' u \otimes (c', b)_{B'} = -[u \otimes b, s' \otimes c'], \\
 [\lambda \otimes e, u \otimes b] &= f^{-1}(\lambda)u \otimes (e, b)_{B'} = -[u \otimes b, \lambda \otimes e], \\
 [u' \otimes b', x \otimes a] &= x^t u' \otimes (b', a)_{B'} = -[x \otimes a, u' \otimes b'], \\
 [u' \otimes b', s \otimes c] &= s u' \otimes (b', c)_B = -[s \otimes c, u' \otimes b'], \\
 [u' \otimes b', \lambda \otimes e] &= -\lambda u' \otimes (b', e)_B = -[\lambda \otimes e, u' \otimes b'], \\
 [d, x \otimes a] &= x \otimes da = -[x \otimes a, d], \\
 [d, u \otimes b] &= u \otimes db = -[u \otimes b, d], \\
 [d, s \otimes c] &= s \otimes dc = -[s \otimes c, d], \\
 [d, \lambda \otimes e] &= \lambda \otimes de = -[\lambda \otimes e, d], \\
 [d, s' \otimes c'] &= s' \otimes dc' = -[s' \otimes c', d], \\
 [d, u' \otimes b'] &= u' \otimes db' = -[u' \otimes b', d], \\
 [d_1, d_2] &\in D,
 \end{aligned}$$

All other products of the homogeneous components of the decomposition (2.1) are zero.

4. COORDINATE ALGEBRA OF (Θ_n, sl_n) -GRADED LIE ALGEBRAS ($n = 3, 4$)

Let L be a Θ_n -graded Lie algebra with the grading subalgebra $\mathfrak{g} \cong sl_n$. Throughout this section we assume that $n = 4$ or $n = 3$ and the conditions (3.1) hold. Let $\mathfrak{g}^\pm = \{x \in sl_n \mid x^t = \pm x\}$. Then

$$(4.1) \quad \mathfrak{g} \otimes A = (\mathfrak{g}^+ \oplus \mathfrak{g}^-) \otimes A = (\mathfrak{g}^+ \otimes A) \oplus (\mathfrak{g}^- \otimes A) = (\mathfrak{g}^+ \otimes A^-) \oplus (\mathfrak{g}^- \otimes A^+)$$

where A^\pm is a copy of the vector space A . Recall that we identify \mathfrak{g} with $\mathfrak{g} \otimes 1$ where 1 is a distinguished element of A . We denote by a^\pm the image of $a \in A$ in the space A^\pm . Denote

$$\mathfrak{a} := A^+ \oplus A^- \oplus C \oplus C' \oplus E \oplus E' \text{ for } n = 3.$$

$$\mathfrak{a} := A^+ \oplus A^- \oplus C \oplus E \oplus C' \quad \text{and} \quad \mathfrak{b} := \mathfrak{a} \oplus B \oplus B' \text{ for } n = 4.$$

In Section 3 we described the multiplicative structures of these Lie algebras. In this section we describe the coordinate algebras of them. and we show that the product in L induces an algebra structure on both \mathfrak{a} and \mathfrak{b} .

4.1. Unital algebra \mathfrak{a} . We are going to define Lie and Jordan multiplication on \mathfrak{a} by extending the bilinear products given in Tables 5 and 6 in a natural way. It can be shown that all products $(\alpha_1, \alpha_2)_Z$ with $\alpha_1, \alpha_2 \in \mathfrak{a}$ are either symmetric or skew-symmetric, except in the cases $(\alpha_1 = c \text{ and } \alpha_2 = e \text{ or } \alpha_1 = c' \text{ and } \alpha_2 = e')$ for $n = 3$. This is why we will write $(\alpha_1 \circ \alpha_2)_Z$ or $[\alpha_1, \alpha_2]_Z$, respectively, instead of $(\alpha_1, \alpha_2)_Z$. The aim of this subsection is to show that \mathfrak{a} is a unital algebra with respect to the new multiplication given by $\alpha_1 \alpha_2 := \frac{[\alpha_1, \alpha_2]}{2} + \frac{\alpha_1 \circ \alpha_2}{2}$ for all homogeneous $\alpha_1, \alpha_2 \in \mathfrak{a}$ with the

products $[\]$ and \circ given by Table 5, except in the cases $(\alpha_1 = c$ and $\alpha_2 = e$ or $\alpha_1 = c'$ and $\alpha_2 = e')$ for $n = 3$ and $a^\pm \lambda$ for $n = 4$, see (4.2).

Remark 4.1. By using the same arguments of [10, Remark 4.1.1]) some of the products in (3.5) and (3.2) can be rewritten in terms of symmetric and skew-symmetric elements. For example for $n = 4$ these products can be written as follows: For all $x^\pm, x_1^\pm, x_2^\pm \in \mathfrak{g}^\pm$, $u, v \in V$, $u', v' \in V'$, $s \in S$, $\lambda, \lambda_1, \lambda_2 \in \Lambda$, $s' \in S'$ and for all $a^\pm, a_1^\pm, a_2^\pm \in A$, $b, b_1, b_2 \in B$, $b', b'_1, b'_2 \in B'$, $c \in C$, $c' \in C'$, $e \in E$, $d \in D$,

$$[x_1^+ \otimes a_1^-, x_2^+ \otimes a_2^-] = x_1^+ \circ x_2^+ \otimes \frac{[a_1^-, a_2^-]_{A^-}}{2} + [x_1^+, x_2^+] \otimes \frac{(a_1^- \circ a_2^-)_{A^+}}{2} + (x_1^+ | x_2^+) \langle a_1^-, a_2^- \rangle,$$

$$[x_1^- \otimes a_1^+, x_2^- \otimes a_2^+] = x_1^- \circ x_2^- \otimes \frac{[a_1^+, a_2^+]_{A^-}}{2} + [x_1^-, x_2^-] \otimes \frac{(a_1^+ \circ a_2^+)_{A^+}}{2} + (x_1^- | x_2^-) \langle a_1^+, a_2^+ \rangle,$$

$$[x_1^+ \otimes a_1^-, x_1^- \otimes a_1^+] = x_1^+ \diamond x_1^- \otimes \frac{[a_1^-, a_1^+]_{A^+}}{2} + [x_1^+, x_1^-] \otimes \frac{(a_1^- \circ a_1^+)_{A^-}}{2}.$$

$$[x^+ \otimes a^-, s \otimes c] = x^+ \diamond s \otimes \frac{[a^-, c]_C}{2} + [x^+, s] \otimes \frac{(a^- \circ c)_E}{2},$$

$$[x^- \otimes a^+, s \otimes c] = x^- \diamond s \otimes \frac{[a^+, c]_E}{2} + [x^-, s] \otimes \frac{(a^+ \circ c)_C}{2}.$$

$$[x^+ \otimes a^-, \lambda \otimes e] = x^+ \diamond \lambda \otimes \frac{[a^-, e]_E}{2} + [x^+, \lambda] \otimes \frac{(a^- \circ e)_C}{2} + [x^+, f^{-1}(\lambda)] \otimes \frac{(a^- \circ e)_{C'}}{2}.$$

$$[x^- \otimes a^+, \lambda \otimes e] = x^- \diamond \lambda \otimes \frac{[a^+, e]_C}{2} + [x^-, \lambda] \otimes \frac{(a^+ \circ e)_E}{2} + x^- \diamond f^{-1}(\lambda) \otimes \frac{[a^+, e]_{C'}}{2}.$$

$$[s' \otimes c', x^+ \otimes a^-] = s' \diamond x^+ \otimes \frac{[c', a^-]_{C'}}{2} + f([s', x^+]) \otimes \frac{(c' \circ a^-)_E}{2},$$

$$[s' \otimes c', x^- \otimes a^+] = f(s' \diamond x^-) \otimes \frac{[c', a^+]_E}{2} + [s', x^-] \otimes \frac{(c' \circ a^+)_{C'}}{2}.$$

$$[s \otimes c, s' \otimes c'] = s \circ s' \otimes \frac{[c, c']_{A^-}}{2} + [s, s'] \otimes \frac{(c \circ c')_{A^+}}{2} + (s | s') \langle c, c' \rangle.$$

$$[\lambda_1 \otimes e_1, \lambda_2 \otimes e_2] = \lambda_1 \circ f^{-1}(\lambda_2) \otimes \frac{[e_1, e_2]_{A^-}}{2} + [\lambda_1, f^{-1}(\lambda_2)] \otimes \frac{(e_1 \circ e_2)_{A^+}}{2} + (\lambda_1 | f^{-1}(\lambda_2)) \langle e_1, e_2 \rangle.$$

$$[s \otimes c, \lambda \otimes e] = s \diamond \lambda \otimes \frac{[c, e]_{A^+}}{2} + [s, \lambda] \otimes \frac{(c \circ e)_{A^-}}{2}.$$

$$[s' \otimes c', \lambda \otimes e] = s' \diamond \lambda \otimes \frac{[c', e]_{A^+}}{2} + [s', \lambda] \otimes \frac{(c' \circ e)_{A^-}}{2}.$$

The mappings $\alpha \otimes \beta \mapsto (\alpha \circ \beta)_{Z_1}$ and $\alpha \otimes \beta \mapsto [\alpha, \beta]_{Z_2}$ can be extended to $Y \otimes X$ in a consistent way by defining $(\beta \circ \alpha)_{Z_1} = (\alpha \circ \beta)_{Z_1}$ and $[\beta, \alpha]_{Z_2} = -[\alpha, \beta]_{Z_2}$. For $n = 3$, $\alpha \in C$ and $\beta \in E$ we will write $\alpha\beta$ (resp. $\beta\alpha$) instead of $(\alpha, \beta)_Z$ (resp. $(\beta, \alpha)_Z$). and the map can be extended by by defining $(\beta\alpha)_{E'} = -(\alpha\beta)_{E'}$ (resp. $(\beta\alpha)_E = -(\alpha\beta)_E$). In Tables 5 and 6 below, if the cell in row X and column Y contains (Z_1, \circ) , and $(Z_2, [\])$ this means that there is a symmetric bilinear map $X \times Y \rightarrow Z_1$, given by $\alpha \otimes \beta \mapsto (\alpha \circ \beta)_{Z_1}$ and a skew symmetric bilinear map $X \times Y \rightarrow Z_2$, given by $\alpha \otimes \beta \mapsto [\alpha, \beta]_{Z_2}$ ($\alpha \in X, \beta \in Y$).

.	A^+	A^-	C	C'	E	E'
A^+	(A^+, \circ) $(A^-, [])$	(A^-, \circ) $(A^+, [])$	(C, \circ)	(C', \circ)	(C, \circ) $(E, [])$	(E', \circ) $(C', [])$
A^-	(A^-, \circ) $(A^+, [])$	(A^+, \circ) $(A^-, [])$	$(C, [])$	$(C', [])$	(E, \circ) $(C, [])$	(C', \circ) $(E', [])$
C	(C, \circ)	$(C, [])$	0	(A^+, \circ) $(A^-, [])$	E'	(A^-, \circ) $(A^+, [])$
C'	(C', \circ)	$(C', [])$	(A^+, \circ) $(A^-, [])$	0	(A^-, \circ) $(A^+, [])$	E
E	(C, \circ) $(E, [])$	(E, \circ) $(C, [])$	E'	(A^-, \circ) $(A^+, [])$	(E', \circ) $(C', [])$	(A^+, \circ) $(A^-, [])$
E''	(E', \circ) $(C', [])$	(C', \circ) $(E', [])$	(A^-, \circ) $(A^+, [])$	E	(A^+, \circ) $(A^-, [])$	(E, \circ) $(C, [])$

TABLE 5. Products of homogeneous components of \mathfrak{a} for $n = 3$

.	A^+	A^-	C	E	C'
A^+	(A^+, \circ) $(A^-, [])$	(A^-, \circ) $(A^+, [])$	(C, \circ) $(E, [])$	(E, \circ) $(C, [])$ (C', \circ)	(C', \circ) $(E, [])$
A^-	(A^-, \circ) $(A^+, [])$	(A^+, \circ) $(A^-, [])$	(E, \circ) $(C, [])$	(C, \circ) $(E, [])$ (C', \circ)	(E, \circ) $(C', [])$
C	(C, \circ) $(E, [])$	(E, \circ) $(C, [])$	0	(A^-, \circ) $(A^+, [])$	(A^+, \circ) $(A^-, [])$
E	(E, \circ) $(C, [])$ (C', \circ)	(C, \circ) $(E, [])$ (C', \circ)	(A^-, \circ) $(A^+, [])$	(A^+, \circ) $(A^-, [])$	(A^-, \circ) $(A^+, [])$
C'	(C', \circ) $(E, [])$	(E, \circ) $(C', [])$	(A^+, \circ) $(A^-, [])$	(A^-, \circ) $(A^+, [])$	0

TABLE 6. Products of homogeneous components of \mathfrak{a} for $n = 4$

We are going to show that $\mathfrak{a} = A^+ \oplus A^- \oplus C \oplus E \oplus C'$ is an associative algebra with respect to multiplication defined as follows:

$$\alpha_1 \alpha_2 := \frac{[\alpha_1, \alpha_2]}{2} + \frac{\alpha_1 \circ \alpha_2}{2}$$

for all homogeneous $\alpha_1, \alpha_2 \in \mathfrak{a}$ with the products $[]$ and \circ given by Table 5, except in the case $a^\pm \lambda$ which is equal to

$$(4.2) \quad \begin{aligned} a^- \lambda &= \frac{[a^-, e]_E}{2} + \frac{(a^- \circ e)_C}{2} + \frac{(a^- \circ e)_{C'}}{2}, \\ a^+ \lambda &= \frac{[a^+, e]_C}{2} + \frac{(a^+ \circ e)_E}{2} + \frac{[a^+, e]_{C'}}{2}. \end{aligned}$$

Note that $[\alpha_1, \alpha_2] = \alpha_1\alpha_2 - \alpha_2\alpha_1$ and $\alpha_1 \circ \alpha_2 = \alpha_1\alpha_2 + \alpha_2\alpha_1$.

Proposition 4.2. 1) $\mathcal{A} = A^- \oplus A^+$ is an associative subalgebra of \mathfrak{a} with identity element 1^+ .

2) $C \oplus E$ and $C' \oplus E'$ are \mathcal{A} -bimodules for $n = 3$.

3) $C \oplus E \oplus C'$ are \mathcal{A} -bimodule for $n = 4$.

Proof. 1) This is similar to the proof of [10, Theorem 4.1.3].

(2) and (3) can be deduce from tensor product decompositions for sl_n ($n = 3, 4$), see Tables 1 and 2. \square

Theorem 4.3. The linear transformation $\gamma : \mathfrak{a} \rightarrow \mathfrak{a}$ defined by

$$\gamma(a^-) = -a^-, \gamma(a^+) = a^+, \gamma(c) = -c, \gamma(e) = e, \gamma(c') = -c',$$

is an antiautomorphism of order 2 of the algebra \mathfrak{a} .

Proof. See [9, Theorem 4.1.6]. \square

4.2. Coordinate algebra \mathfrak{b} . Let L be an Θ_4 -graded Lie algebra with the grading subalgebra $\mathfrak{g} \cong sl_4$. Recall that we denote

$$\mathfrak{a} := A^+ \oplus A^- \oplus C \oplus E \oplus C' \quad \text{and} \quad \mathfrak{b} := \mathfrak{a} \oplus B \oplus B'.$$

The aim of this subsection is to show that \mathfrak{b} is an algebra with identity 1^+ with respect to the multiplication extending that on \mathfrak{a} given in Table 7. It can be shown that all products $(\beta_1, \beta_2)_Z$ with $\beta_1, \beta_2 \in B \oplus B'$ are either symmetric or skew-symmetric. This is why we will write $(\beta_1 \circ \beta_2)_Z$ or $[\beta_1, \beta_2]_Z$, respectively, instead of $(\beta_1, \beta_2)_Z$. For $\alpha \in \mathfrak{a}$ and $\beta \in B \oplus B'$ we will write $\alpha\beta$ (resp. $\beta\alpha$) instead of $(\alpha, \beta)_Z$ (resp. $(\beta, \alpha)_Z$). Let $b \in B$ and $b' \in B'$. We define $b\alpha := \gamma(\alpha)b$ and $\alpha b' := b'\gamma(\alpha)$. We will show that $B \oplus B'$ is an \mathfrak{a} -bimodule.

Recall that

$$x \otimes a = \frac{(x + x^t)}{2} \otimes a + \frac{(x - x^t)}{2} \otimes a \in \mathfrak{g}^+ \otimes A + \mathfrak{g}^- \otimes A.$$

Let $u \otimes b \in V \otimes B$ and $v' \otimes b' \in V' \otimes B'$. We need the following formula from (3.5):

$$[u \otimes b, v' \otimes b'] = (uv^t - \frac{\text{tr}(uv^t)}{4}I) \otimes (b, b')_A + \frac{2 \text{tr}(uv^t)}{4} \langle b, b' \rangle.$$

By splitting $(b, b')_A$ into symmetric and skew-symmetric parts and using the equations

$$\begin{aligned} (uv^t - \frac{\text{tr}(uv^t)}{4}I) + (uv^t - \frac{\text{tr}(uv^t)}{4}I)^t &= uv^t + v'u^t - \frac{2 \text{tr}(uv^t)}{4}I, \\ (uv^t - \frac{\text{tr}(uv^t)}{4}I) - (uv^t - \frac{\text{tr}(uv^t)}{4}I)^t &= uv^t - v'u^t, \end{aligned}$$

we get

$$\begin{aligned} [u \otimes b, v' \otimes b'] &= (uv^t + v'u^t - \frac{2 \text{tr}(uv^t)}{4}I) \otimes \frac{[b, b']_{A^-}}{2} + \\ (4.3) \quad & (uv^t - v'u^t) \otimes \frac{(b \circ b')_{A^+}}{2} + \frac{2 \text{tr}(uv^t)}{4} \langle b, b' \rangle. \end{aligned}$$

Let $b, b_1, b_2 \in B$ and $b', b'_1, b'_2 \in B'$. Using (3.5) and (4.3) we get

$$\begin{aligned}
 [u \otimes b_1, v \otimes b_2] &= (uv^t + vu^t) \otimes \frac{[b_1, b_2]_C}{2} + (uv^t - vu^t) \otimes \frac{(b_1 \circ b_2)_E}{2}, \\
 [u' \otimes b'_1, v' \otimes b'_2] &= (u'v'^t + v'u'^t) \otimes \frac{[b'_1, b'_2]_{C'}}{2} + f(u'v'^t - v'u'^t) \otimes \frac{(b'_1 \circ b'_2)_E}{2}, \\
 [u \otimes b, v' \otimes b'] &= (uv'^t + v'u^t - \frac{2 \operatorname{tr}(uv'^t)}{4} I) \otimes \frac{[b, b']_{A^-}}{2} + \\
 (4.4) \quad & (uv'^t - v'u^t) \otimes \frac{(b \circ b')_{A^+}}{2} + \frac{2 \operatorname{tr}(uv'^t)}{4} \langle b, b' \rangle.
 \end{aligned}$$

We define

$$\begin{aligned}
 b_1 b_2 &:= \frac{[b_1, b_2]_C}{2} + \frac{(b_1 \circ b_2)_E}{2}, & b'_1 b'_2 &:= \frac{[b'_1, b'_2]_{C'}}{2} + \frac{(b'_1 \circ b'_2)_E}{2}, \\
 bb' &:= \frac{[b, b']_{A^-}}{2} + \frac{(b \circ b')_{A^+}}{2}, & b'b &:= -\frac{[b, b']_{A^-}}{2} + \frac{(b \circ b')_{A^+}}{2}.
 \end{aligned}$$

Then $\mathfrak{b} = \mathfrak{a} \oplus B \oplus B'$ is an algebra with multiplication extending that on \mathfrak{a} . The following table describes the products of homogeneous elements of \mathfrak{b} (use Table 5 for the products on \mathfrak{a}).

.	$A^+ + A^-$	$C + E$	C'	B	B'
$A^+ + A^-$	$A^+ + A^-$	$C + E + C'$	$C' + E$	B	B'
$C + E$	$C + E + C'$	0	$A^+ + A^-$	0	B
C'	$C' + E$	$A^+ + A^-$	0	B'	0
B	B	0	B'	$C + E$	$A^+ + A^-$
B'	B'	B	0	$A^+ + A^-$	$C' + E$

TABLE 7. Products in \mathfrak{b}

Using the same arguments of [10, Section 4.2] we get the following properties:

Theorem 4.4. (1) The linear transformation $\eta : \mathfrak{b} \rightarrow \mathfrak{b}$ defined by $\eta(\alpha) = \gamma(\alpha)$, $\eta(b) = b$ and $\eta(b') = b'$ for all $\alpha \in \mathfrak{a}$, $b \in B$ and $b' \in B'$ is an antiautomorphism of order 2 of the algebra \mathfrak{b} .

- (2) 1^+ is the identity element of \mathfrak{b} .
(3) Let $b \in B$, $b' \in B'$ and $\alpha \in \mathfrak{a}$. Then

$$\begin{aligned}
 [z \otimes \alpha, u \otimes b] &= zu \otimes ab = -[u \otimes b, z \otimes \alpha], \\
 [u' \otimes b', z \otimes \alpha] &= z^t u' \otimes b' \alpha = -[z \otimes \alpha, u' \otimes b'].
 \end{aligned}$$

- (4) $B \oplus B'$ is an \mathfrak{a} -bimodule.
(5) B and B' are \mathcal{A} -bimodules.

Proof. (1) This is similar to [10, Theorem 4.2.1].

(2) This is similar to [10, Theorem 4.2.2].

(3) We deduce this by using (3.5) and Table 7.

(4) See [10, Proposition 4.2.4].

(5) We deduce this from the properties that B and B' are invariant under multiplication by \mathcal{A} , see Table 7. \square

The mapping $\langle, \rangle : X \otimes X' \rightarrow D$ with $X = B, C$ can be extended to $X' \otimes X$ in a consistent way by defining $\langle x', x \rangle := -\langle x, x' \rangle$. Let $X, Y \in \{A^+, A^-, B, B', C, C', E\}$. Recall also the maps $\langle, \rangle : A^\pm \otimes A^\pm \rightarrow D$ described previously (see Remark 4.1(a)). For the convenience, we extend the mappings to the whole space \mathfrak{b} by defining the remaining $\langle X, Y \rangle$ to be zero. Hence

$$\langle \mathfrak{b}, \mathfrak{b} \rangle = \langle A^+, A^+ \rangle + \langle A^-, A^- \rangle + \langle B, B' \rangle + \langle C, C' \rangle + \langle E, E \rangle.$$

It follows from condition $(\Gamma 3)$ in Definition 2.1 that

$$(4.5) \quad D = \langle \mathfrak{b}, \mathfrak{b} \rangle.$$

Proposition 4.5. (1) $[d, \langle \alpha, \beta \rangle] = \langle d\alpha, \beta \rangle + \langle \alpha, d\beta \rangle$ for all $\alpha, \beta \in \mathfrak{b}$ and $d \in D$.

(2) $\langle A^+, A^+ \rangle, \langle A^-, A^- \rangle, \langle B, B' \rangle, \langle C, C' \rangle$ and $\langle E, E \rangle$ are ideals of the Lie algebra D .

(3) D acts by derivations on \mathfrak{b} and leaves all subspaces A^+, A^-, B, \dots, E invariant.

Proof. This is similar to [10, Proposition 4.2.8]. \square

REFERENCES

- [1] Bruce Allison, Georgia Benkart, and Yun Gao, Lie algebras graded by the root systems $BC_r, r > 1$, no. 751, American Mathematical Soc., 2002.
- [2] Yuri Bahturin, Alexander Baranov, and Alexander Zalesski, Simple Lie subalgebras of locally nite associative algebras, Journal of Algebra 281 (2004), no. 1, 225-246.
- [3] Yuri Bahturin and Georgia Benkart, Some constructions in the theory of locally nite simple Lie algebras, J. Lie Theory 14 (2004), no. 1, 243-270.
- [4] Georgia Benkart and Alberto Elduque, Lie algebras with prescribed sl_3 decomposition, Proceedings of the American Mathematical Society 140 (2012), no. 8, 2627-2638.
- [5] Georgia Benkart and Oleg Smirnov, Lie algebras graded by the root system BC_1 , J. Lie theory 13 (2003), no. 1, 91-132.
- [6] Georgia Benkart and Em Zelmanov, Lie algebras graded by nite root systems and intersection matrix algebras, Inventiones mathematicae 126 (1996), no. 1, 1-45.
- [7] Stephen Berman and Robert Moody, Lie algebras graded by nite root systems and the intersection matrix algebras of Slodowy, Inventiones mathematicae 108 (1992), no. 1, 323-347.
- [8] Roe Goodman and Nolan R Wallach, Symmetry, representations, and invariants, vol. 255, Springer, 2009.
- [9] George Seligman, Rational methods in Lie algebras, vol. 17, Marcel Dekker Inc, 1976.
- [10] Hogar M Yaseen, Generalized root graded lie algebras, Ph.D. thesis, Department of Mathematics; University of Leicester, 2018.

DEPARTMENT OF MATHEMATICS, SALAHADDIN UNIVERSITY-ERBIL, IRAQ.

E-mail address: hogr.yaseen@su.edu.krd, hogrmyaseen@gmail.com