

Clustering of Arrivals in Queueing Systems: Autoregressive Conditional Duration Approach¹

Petra Tomanová

University of Economics, Prague
Winston Churchill Square 1938/4, 130 67 Prague 3, Czechia
petra.tomanova@vse.cz

Vladimír Holý

University of Economics, Prague
Winston Churchill Square 1938/4, 130 67 Prague 3, Czechia
vladimir.holy@vse.cz

April 1, 2020

Abstract: Arrivals in queueing systems are typically assumed to be independent and exponentially distributed. Our analysis of an online bookshop, however, shows that there is autocorrelation structure present. First, we adjust the inter-arrivals times (i.e. durations) for diurnal and seasonal patterns. Second, we model adjusted durations by the autoregressive conditional duration (ACD) model based on the generalized gamma distribution with the generalized autoregressive score (GAS) dynamics. Third, in a simulation study, we investigate the effects of the dynamic arrival model on the number of customers, the busy period and the response time in queueing systems with single and multiple servers. We find that ignoring the autocorrelation structure leads to significantly underestimated performance measures and consequently suboptimal decisions. The proposed approach serves as a general methodology for treatment of arrivals clustering in practice.

Keywords: Inter-Arrival Times, Queueing Theory, Autoregressive Conditional Duration Model, Generalized Autoregressive Score Model, Retail Business.

JEL Codes: C41, C44, M11.

1 Introduction

In various applications of the operations research, it is undeniable that the characteristics of models evolve over time. The parameters of interest can depend on the time of day and season (see e.g. Kayacı Çodur and Yılmaz, 2020) as well as on their past values and other past indicators (see e.g. Bruzda, 2020). In the paper, we focus on the latter dependency in arrivals to queueing systems from the perspective of the autoregressive conditional duration models with the generalized autoregressive score dynamics.

Many standard queueing systems consider inter-arrival times to be independent due to analytical tractability. Some studies, however, explicitly consider autocorrelation and model arrivals using the *Markovian arrival process (MAP)* (see e.g. Adan and Kulkarni, 2003; Buchholz and Kriege, 2017; Manafzadeh Dizbin and Tan, 2019), the *Markov renewal process* (see e.g. Tin, 1985; Patuwo et al., 1993; Szekli et al., 1994), the *moving average process* (see e.g. Finch, 1963; Finch and Pearce, 1965; Pearce, 1967) or the *discrete autoregressive process* (see e.g. Hwang and Sohraby, 2003; Kamoun, 2006; Miao and Lee, 2013). Hwang and Sohraby (2003) argue that the time series models with few parameters are more suitable in practice than the MAP models which might be overparameterized. Simulation studies investigating the autocorrelation in arrivals include Livny et al. (1993), Resnick and Samorodnitsky (1997), Altiok and Melamed (2001), Nielsen (2007) and Civelek et al. (2009). Overall, these studies show that ignoring the autocorrelation structure, if present, leads to biased performance measures in queueing systems.

¹Preliminary results were presented in Tomanová (2018) and Tomanová (2019).

The arrival processes are also extensively studied in the financial high-frequency literature. In this field, the duration analysis deals with modeling of times between successive transactions, times until the price reaches a certain level and times until a certain volume is traded. Typically, the *autoregressive conditional duration (ACD) model* of Engle and Russell (1998) is utilized. Its dynamics is analogous to the *generalized autoregressive conditional heteroskedasticity (GARCH) model* of Bollerslev (1986). In its basic version, the ACD model is based on the exponential distribution but many other distributions are considered in the literature as well. Notably, Lunde (1999) introduces the generalized gamma distribution to the ACD model. Bauwens et al. (2004) and Fernandes and Grammig (2005) find that the generalized gamma distribution is more adequate than the exponential, Weibull and Burr distributions in financial applications. Hautsch (2003) further finds that the four-parameter generalized F distribution reduces to the three-parameter generalized gamma distribution in most cases of financial durations. For a survey of financial duration analysis, see Pacurar (2008) and Saranjeet and Ramanathan (2018).

A modern approach to time series modeling is the *general autoregressive score (GAS) model* of Creal et al. (2013), also known as the *dynamic conditional score (DCS) model* by Harvey (2013). The GAS model is an observation-driven model providing a general framework for modeling of time-varying parameters of any underlying probability distribution. It captures dynamics of time-varying parameters by the autoregressive term and the score of the conditional density function utilizing the shape of the density function. The theoretical properties of the GAS models together with their estimation by the maximum likelihood method are investigated e.g. by Blasques et al. (2014) and Blasques et al. (2018). The empirical performance of the GAS models is studied e.g. by Koopman et al. (2016) and Blazsek and Licht (2020). The GARCH model based on the normal distribution and the ACD model based on the exponential distribution are both special cases of the GAS class. The ACD model based on the zero-inflated negative binomial distribution with the GAS dynamics is proposed by Blasques et al. (2020). Many more models belonging to the GAS family with various applications are proposed in the literature – see Lucas, 2020 for a comprehensive list of papers.

In the paper, we put together these three cornerstones – the queueing theory, the ACD models and the GAS models – and demonstrate that they fit together perfectly. We analyze inter-arrival times between orders from an online Czech bookshop. First, we adjust arrivals for diurnal and seasonal patterns using the cubic spline. Next, we find that the adjusted inter-arrival times exhibit strong clustering behavior – short inter-arrival times are usually followed by short times. To capture this autocorrelation, we utilize the dynamic model based on the generalized gamma distribution with the GAS dynamics in the spirit of the ACD models. We confirm that the proposed specification is quite suitable for the observed data. Finally, we investigate the effects of the proposed arrivals model on queueing systems with single and multiple servers and exponential services. In a simulation study, we show that various performance measures – the number of customers in the system, the busy period of servers and the response time – have higher mean and variance as well as heavier tails for the proposed dynamic arrivals model than for the standard static model. Furthermore, we illustrate how misspecification of the arrivals model can lead to suboptimal decisions.

The rest of the paper is structured as follows. In Section 2, we present the model based on the generalized gamma distribution with the GAS dynamics for diurnally adjusted inter-arrival times. In Section 3, we show that real data of a retail store exhibit the autocorrelation structure that is well captured by our model. In Section 4, we investigate the impact of the proposed arrivals model on the performance measures using simulations. We conclude the paper in Section 5.

2 Dynamic Model for Arrivals

2.1 Diurnal and Seasonal Adjustment

Despite that we focus on time dependence in terms of the autoregressive behavior, we deal with diurnal, weekly and monthly seasonality patterns as well. To capture the non-linear behavior of the diurnal and seasonal patterns and to properly adjust the durations, the cubic spline method is utilized. The *cubic spline* is a piecewise cubic polynomial with continuous derivatives up to order

two at each k -th fixed point called knot, $k = 1, \dots, K$. Bruce and Bruce (2017) point out that the cubic spline method is often a superior approach to the polynomial regression since the polynomial regression often leads to undesirable "wiggleness" in the regression equation.

To take into account the specifics of raw durations $\{\tilde{y}_i\}_{i=1}^n$, we define the cubic spline with knots at $\{\xi_k\}_{k=1}^K$ as

$$\log \tilde{y}_i = \beta_0 + \beta_1 b_1(x_i) + \beta_2 b_2(x_i) + \dots + \beta_{K+3} b_{K+3}(x_i) + \gamma t_i + \varepsilon_i, \quad (1)$$

where t_i is the trend variable and $\{b_j\}_{j=1}^{K+3}$ are the basis functions. The basis functions are equal to (i) the variable x_i , $b_1(x_i) = x_i$; (ii) its square, $b_2(x_i) = x_i^2$; (iii) its cube, $b_3(x_i) = x_i^3$; and (iv) truncated power functions, $b_{k+3}(x_i) = \max\{0, (x_i - \xi_k)^3\}$, $k = 1, \dots, K$. The trend variable t_i is linear in time (not linear in observations), $t_1 = 0$ and $t_i = \sum_{j=1}^{i-1} \tilde{y}_j$ for $i = 2, \dots, n$, to take into account the irregularly spaced observations. Moreover, the logarithmic transformation of \tilde{y} ensures the non-negativity of adjusted durations. Equidistant intervals are used for knots identification since intervals based on quantiles might lead to a too-small number of knots allocated to off-peak hours.

Regression parameters in (1) are estimated by the *weighted least squares (WLS)* method with weights being the durations. WLS naturally compensates for the possibility that during a particular time interval either a small number of long durations or a higher number of shorter durations is observed, i.e. the number of observed durations within a time interval depends on the duration values themselves. Unlike the ordinary least squares, this approach properly weights the durations during hours that exhibit a small median but a huge dispersion. Once the parameters are estimated, the diurnally and seasonally adjusted and detrended durations y_i are set to exponentiated residuals from regression (1).

2.2 Generalized Gamma Distribution

Next, we consider the adjusted durations y_i to follow the generalized gamma distribution. The *generalized gamma distribution* is a continuous probability distribution for non-negative variables proposed by Stacy (1962). It is a three-parameter generalization of the two-parameter gamma distribution and contains the exponential distribution and the Weibull distribution as well. The distribution has the scale parameter α and the shape parameters $\psi > 0$ and $\varphi > 0$. We use the parametrization allowing for arbitrary values of α which is quite suitable for modeling of its dynamics. The probability density function is

$$f(y|\alpha, \psi, \varphi) = \frac{1}{\Gamma(\psi)} \frac{\varphi}{e^\alpha} \left(\frac{y}{e^\alpha}\right)^{\psi\varphi-1} e^{-\left(\frac{y}{e^\alpha}\right)^\varphi} \quad \text{for } y \in (0, \infty), \quad (2)$$

where $\Gamma(\cdot)$ is the gamma function. The expected value and variance is

$$\begin{aligned} \mathbb{E}[Y] &= e^\alpha \frac{\Gamma(\psi + \varphi^{-1})}{\Gamma(\psi)}, \\ \text{var}[Y] &= e^{2\alpha} \frac{\Gamma(\psi + 2\varphi^{-1})}{\Gamma(\psi)} - \left(e^\alpha \frac{\Gamma(\psi + \varphi^{-1})}{\Gamma(\psi)} \right)^2. \end{aligned} \quad (3)$$

The score for the parameter α is

$$\nabla_\alpha(y, \alpha, \psi, \varphi) = \frac{\partial \log f(y|\alpha, \psi, \varphi)}{\partial \alpha} = \varphi (y^\varphi e^{-\varphi\alpha} - \psi) \quad \text{for } y \in (0, \infty). \quad (4)$$

The Fisher information for the parameter α is

$$\mathcal{I}_\alpha(\alpha, \psi, \varphi) = \mathbb{E} [\nabla_\alpha(y, \alpha, \psi, \varphi)^2 | \alpha, \psi, \varphi] = \psi\varphi^2. \quad (5)$$

Note that the Fisher information for α is not dependent on α itself. Special cases of the generalized gamma distribution include the gamma distribution for $\varphi = 1$, the Weibull distribution for $\psi = 1$ and the exponential distribution for $\psi = 1$ and $\varphi = 1$. The generalized gamma distribution itself is contained in a larger family – the generalized F distribution with four parameters.

2.3 Generalized Autoregressive Score Dynamics

Finally, we consider the scale parameter to be time-varying. In the *generalized autoregressive score (GAS)* framework of Creal et al. (2013), the time-varying parameters are linearly dependent on their lagged values and the lagged values of the score of the conditional density. Typically, only first lag is utilized. In our case, parameter α_i follows recursion

$$\begin{aligned}\alpha_{i+1} &= c + b\alpha_i + a\nabla_{\alpha}(y_i, \alpha_i, \psi, \varphi) \\ &= c + b\alpha_i + a\varphi(y_i^{\varphi}e^{-\varphi\alpha_i} - \psi),\end{aligned}\tag{6}$$

where c is the constant parameter, b is the autoregressive parameter, a is the score parameter and $\nabla_{\alpha}(y_i, \alpha_i, \psi, \varphi)$ is the score defined in (4). In the GAS framework, the score can be scaled by the inverse of the Fisher information or the square of the inverse of the Fisher information. In our case, however, both scaling functions based on the Fisher information and the unit scaling as well lead to the same model as the Fisher information does not depend on α_i . The score for time-varying parameter α_i is the gradient of the log-likelihood with respect to α_i and indicates how sensitive the log-likelihood is to parameter α_i . In the GAS model, the score drives the time variation in parameter α_i based on the shape of the generalized gamma density function.

Let $\theta = (c, b, a, \psi, \varphi)$ denote the vector of parameters in the model. We can estimate θ straightforwardly by the maximum likelihood method. The log-likelihood function is given by

$$\ell(\theta) = \ln f(y_0|\alpha_0, \psi, \varphi) + \sum_{i=1}^n \ln f(y_i|\alpha_i, \psi, \varphi),\tag{7}$$

where $f(\cdot)$ is the generalized gamma density function given by (2). We deliberately set aside the first term as the time-varying parameter α_i needs to be initialized at $i = 0$. We set the value of α_0 to the long-term mean value $c/(1 - b)$. Subsequent values of α_i , $i = 1, \dots, n$, then follow recursion (6). Finally, the parameter estimates $\hat{\theta}$ are obtained by non-linear optimization problem

$$\hat{\theta} \in \max_{\theta} \ell(\theta).\tag{8}$$

3 Empirical Evidence

3.1 Data Overview and Preparation

The data sample is obtained from the database of an online bookshop with one brick-and-mortar location in Prague, Czechia. The data cover the period of June 8 – December 20, 2018, resulting in 28 full weeks and 5 753 observations. The precision of the timestamp is one minute. Thus, zero durations might occur in the data due to two or more orders that arrive within one minute. Since the generalized gamma distribution has a strictly positive support, the zero durations are set to a small positive number. Bauwens (2006) replaces the zero durations with a value equal to the half of the minimal positive duration and argued that this is a more correct approach than their discarding. Hence, all 81 zero durations are set to 0.5 minutes accordingly.

3.2 Diurnal and Seasonal Patterns

The raw duration median is 24 minutes and the mean is 49 minutes – more than double due to long durations during nights (specifically hours between midnight and 9 AM, see Figure 1). Hours between 9 and 11 AM exhibit many short durations and several very long durations resulting in high dispersion (SD = 111.39). The rest of the rush hours (until 5 PM) shows a similar duration median but much lower dispersion (SD = 35.98). Moreover, strong weekly and monthly seasonal patterns are observed. The highest order counts (and consequently lower duration values) occur at the beginning of a week and decrease until Saturday. On Sundays, order counts increase again and exhibit the highest dispersion. During the summer months, the order counts are rather low (resulting in higher duration values) and linearly increase until December.




Figure 1: Raw durations and their fitted diurnal/seasonal pattern.

To obtain the diurnally and seasonally adjusted and detrended durations, the regression equation (1) is estimated with one knot for every 90 minutes and weekly aggregation resulting in the same daily seasonal component for Mondays, Tuesdays, etc. To ensure continuity between Sundays and Mondays, the sample is stacked three times consecutively and the adjusted durations are computed based on the second sub-sample. Parameters are estimated by the WLS.

The fitted values are shown in Figure 1. Note that they do not coincide with the smooth cubic spline function due to a linear trend which makes the corresponding fitted line "saw-toothed". The diurnally and seasonally adjusted and detrended durations are computed as the exponentiated residuals and for convenience, they are standardized to have unit mean. Their values range from 0.002 to 11.23 minutes.

3.3 Fit of the Dynamic Model

Even after the seasonal and diurnal adjustment, the durations still tend to cluster over time – long (short) durations are likely to be followed by long (short) durations. This dependence is not particularly strong but is statistically significant nevertheless as illustrated in Figure 2. To capture the autocorrelation, we utilize the dynamic model based on the generalized gamma distribution with the GAS dynamics presented in Section 2.3. For comparison, we also report the results for static and dynamic models based on special cases of the generalized gamma distribution (G.G.), namely for the exponential (Exp.), Weibull and gamma distributions.

Parameter estimates and the performance evaluation in terms of the *Akaike information criterion* (*AIC*) of both static and dynamic duration models are shown in Table 1. The *AIC* values are at least by 43.59 higher for dynamic models than for their static counterparts. However, the differences among dynamic models are not so striking – the highest difference is between exponential and generalized gamma distributions (by 5.94). The best performing model is the most general one – the dynamic GAS model utilizing the generalized gamma distribution. The dynamic models based on either the exponential or generalized gamma distributions in comparison with their static counterparts are further analyzed in the simulation study of queueing systems.




Figure 2: The autocorrelation function (ACF) and the partial autocorrelation function (PACF) of adjusted durations.

Model		Estimate					Model Fit	
Spec.	Dist.	c	b	a	ψ	φ	Lik.	AIC
Static	Exp.	0.00	0.00	0.00	1.00	1.00	-5 753.00	11 508.00
Static	Weibull	-0.01	0.00	0.00	1.00	0.97	-5 748.93	11 501.86
Static	Gamma	0.04	0.00	0.00	0.96	1.00	-5 749.77	11 503.54
Static	G. G.	-0.12	0.00	0.00	1.08	0.93	-5 748.37	11 502.75
Dyn.	Exp.	0.00	0.76	0.06	1.00	1.00	-5 728.28	11 462.56
Dyn.	Weibull	0.00	0.75	0.06	1.00	0.97	-5 724.89	11 457.79
Dyn.	Gamma	0.01	0.76	0.06	0.97	1.00	-5 725.97	11 459.95
Dyn.	G. G.	-0.06	0.72	0.07	1.15	0.90	-5 723.31	11 456.62

Table 1: The estimated parameters of the duration models with the log-likelihood value (Lik.) and the Akaike information criterion (AIC).




Figure 3: The probability mass functions of the number of customers in the system and density functions of the response time for the static and dynamic arrival models based on the generalized gamma distribution in the queueing system with single server and $\mu = 1.10$.

4 Impact on Queueing Systems

4.1 System with Single Server

We investigate the effects of various arrival models on performance measures in queueing systems using simulations. We consider models based on the exponential and generalized gamma distributions with the static and dynamic specifications. The coefficients of the models are taken from Table 1. In all models, the rate of arrivals is $\lambda = 1$. First, we focus on the queueing system with single server only. We consider the service times to be independent and exponentially distributed with the rate μ ranging from 1.1 to 1.5. We simulate the arrival and service processes and measure the number of customers in the system, the busy period of the server and the response time. The number of simulations we perform ranges from $2 \cdot 10^7$ to $5 \cdot 10^7$ as systems with lower μ require higher number of simulations.

The results are reported in Table 2. For all values of μ , the systems based on the generalized gamma distribution have higher values of performance measures than the systems based on the exponential distribution in terms of the mean, standard deviation and 95 percent quantile. Similarly, systems with the dynamic specification have higher values of performance measures than the systems with the static specification. The left plot of Figure 3 shows how the probability mass function of the number of customers differs for the static and dynamic models. The dynamic model has higher probability of the empty system as there tend to be longer periods of low activity. It has also higher probabilities of large numbers of customers in the system as arrivals tend to cluster. The right plot of Figure 3 shows how the density function of the response time differs for the static and dynamic models. In the dynamic model, customers simply have to wait longer. The differences between the static and dynamic models are naturally weaker for larger μ .

These results carry a warning for practice. When the standard M/M/1 system is assumed but the arrivals actually follow the GAS model based on the generalized gamma distribution, the performance measures are significantly underestimated. For example, the mean number of customers and the mean response time are 22 percent lower than the actual value for $\mu = 1.1$. It is therefore crucial to correctly specify the model for arrivals.

μ	Queueing System		No. of Customers			Busy Period			Response Time		
	Spec.	Dist.	M	SD	95%	M	SD	95%	M	SD	95%
1.10	Static	Exp.	10.00	10.49	31.00	10.00	45.83	39.81	10.00	10.00	29.96
1.10	Static	G. G.	10.38	10.94	32.00	10.36	47.49	41.31	10.38	10.42	31.13
1.10	Dyn.	Exp.	12.38	13.30	39.00	10.88	54.47	41.43	12.38	12.58	37.53
1.10	Dyn.	G. G.	12.83	13.84	40.00	11.24	56.07	43.05	12.83	13.07	38.92
1.20	Static	Exp.	5.00	5.48	16.00	5.00	16.58	22.07	5.00	5.00	14.98
1.20	Static	G. G.	5.18	5.71	17.00	5.17	17.20	22.85	5.18	5.20	15.56
1.20	Dyn.	Exp.	5.97	6.81	20.00	5.36	19.47	23.35	5.97	6.14	18.27
1.20	Dyn.	G. G.	6.21	7.09	20.00	5.56	20.22	24.39	6.21	6.37	19.00
1.30	Static	Exp.	3.33	3.80	11.00	3.33	9.23	14.81	3.33	3.33	9.99
1.30	Static	G. G.	3.45	3.96	11.00	3.44	9.58	15.31	3.45	3.46	10.35
1.30	Dyn.	Exp.	3.87	4.62	13.00	3.54	10.73	15.72	3.87	3.99	11.88
1.30	Dyn.	G. G.	4.04	4.85	14.00	3.67	11.18	16.40	4.04	4.18	12.40
1.40	Static	Exp.	2.50	2.96	8.00	2.50	6.12	10.97	2.50	2.50	7.49
1.40	Static	G. G.	2.58	3.08	9.00	2.58	6.35	11.32	2.58	2.59	7.75
1.40	Dyn.	Exp.	2.84	3.52	10.00	2.63	7.04	11.56	2.84	2.93	8.71
1.40	Dyn.	G. G.	2.96	3.70	10.00	2.73	7.35	12.07	2.96	3.06	9.08
1.50	Static	Exp.	2.00	2.45	7.00	2.00	4.47	8.61	2.00	2.00	5.99
1.50	Static	G. G.	2.06	2.54	7.00	2.06	4.63	8.87	2.06	2.07	6.18
1.50	Dyn.	Exp.	2.23	2.87	8.00	2.09	5.08	9.04	2.23	2.30	6.84
1.50	Dyn.	G. G.	2.32	3.00	8.00	2.16	5.31	9.41	2.32	2.40	7.12

Table 2: Mean values (M), standard deviations (SD) and 95%-quantiles (95%) of the number of customers in the system, the busy period of the server and the response time in various queueing systems with single server.




Figure 4: Costs related to the number of servers and long queue for the static and dynamic arrival models based on the generalized gamma distribution in the queueing systems with multiple servers and $\mu = 0.10$.

4.2 System with Multiple Servers

Next, we consider the queueing systems with multiple servers. We base the simulations on the same setting as in the previous section. The only difference lies in the service structure. We consider the number of servers c ranging from 11 to 15 with the individual service rate $\mu = 0.1$. Such values result in the same server utilizations $\rho = \lambda/(c\mu)$ as in the previous section. Again, we measure the number of customers in the system, the busy period of the servers and the response time. By the busy period, we mean the full busy period, i.e. the duration of the state in which all servers are busy.

The results are reported in Table 3. The findings are very similar to the system with single server – the generalized gamma distribution and the dynamic specification increase all performance measures. When incorrectly assuming the M/M/c system, the specification error is distinct but not as high as in the case of single server. For example, the mean number of customers and the mean response time are 14 percent lower when assuming the M/M/11 system than the actual value for arrivals based on the generalized gamma distribution with the dynamic specification.

We illustrate how misspecification of the arrival model can affect decision making in the following toy example. Let us consider that there are two types of costs associated with the operation of the system – the cost of running one server per unit of time $C_1 = 10$ and the cost of the queue longer than 30 customers per unit of time $C_2 = 3000$. The analytic department is faced with the question of how many servers to operate. The composition of costs for different number of servers is shown in Figure 4. The optimal number of servers according to the static model is 12 while it is 13 for the dynamic model. An analyst assuming the static model believes that the total optimal costs per unit of time are 127.43 while they actually are 142.62 for suboptimal choice of 12 servers. An analyst correctly specifying the dynamic model finds out that the lowest possible costs are 132.53 for the optimal choice of 13 servers. The decision based on the misspecified arrival model therefore results in total cost increase of 8 percent.

5 Conclusion

We analyze the dependence of inter-arrival times in queueing systems and demonstrate the negative impact of arrival model misspecification on decision making. To capture the autocorrelation structure of inter-arrival times, we propose to utilize the dynamic model based on the generalized gamma

c	Queueing System		No. of Customers			Busy Period			Response Time		
	Spec.	Dist.	M	SD	95%	M	SD	95%	M	SD	95%
11	Static	Exp.	16.82	10.66	38.00	10.01	46.01	39.81	16.82	13.80	43.86
11	Static	G. G.	17.15	11.12	39.00	10.36	47.51	41.28	17.15	14.08	44.68
11	Dyn.	Exp.	19.08	13.49	46.00	12.37	58.51	49.89	19.08	15.70	49.83
11	Dyn.	G. G.	19.45	14.01	47.00	12.81	60.38	51.71	19.45	16.07	50.90
12	Static	Exp.	12.25	5.84	24.00	5.01	16.63	22.12	12.25	10.84	33.57
12	Static	G. G.	12.38	6.07	24.00	5.17	17.22	22.85	12.38	10.91	33.85
12	Dyn.	Exp.	13.09	7.19	27.00	6.08	21.06	27.61	13.09	11.34	35.38
12	Dyn.	G. G.	13.26	7.46	28.00	6.33	21.80	28.78	13.26	11.46	35.78
13	Static	Exp.	10.95	4.39	19.00	3.34	9.28	14.85	10.95	10.27	31.30
13	Static	G. G.	11.02	4.53	19.00	3.44	9.60	15.31	11.02	10.30	31.40
13	Dyn.	Exp.	11.36	5.23	21.00	3.99	11.63	18.33	11.36	10.45	32.01
13	Dyn.	G. G.	11.46	5.46	22.00	4.16	12.14	19.11	11.46	10.51	32.24
14	Static	Exp.	10.44	3.76	17.00	2.51	6.14	10.99	10.44	10.10	30.53
14	Static	G. G.	10.48	3.88	18.00	2.59	6.39	11.35	10.48	10.11	30.59
14	Dyn.	Exp.	10.66	4.36	19.00	2.95	7.64	13.43	10.66	10.17	30.85
14	Dyn.	G. G.	10.71	4.54	19.00	3.08	8.01	14.03	10.71	10.20	30.97
15	Static	Exp.	10.20	3.46	16.00	2.01	4.48	8.63	10.20	10.04	30.22
15	Static	G. G.	10.23	3.56	16.00	2.07	4.69	8.95	10.23	10.04	30.25
15	Dyn.	Exp.	10.33	3.93	17.00	2.34	5.53	10.46	10.33	10.07	30.38
15	Dyn.	G. G.	10.35	4.07	18.00	2.43	5.81	10.93	10.35	10.08	30.44

Table 3: Mean values (M), standard deviations (SD) and 95%-quantiles (95%) of the number of customers in the system, the full busy period of servers and the response time in various queueing systems with multiple servers and $\mu = 0.10$.

distribution with the GAS dynamics. We argue that this approach is superior to the standard model assuming the exponential distribution with a constant rate since it leads to a more faithful representation of the mean and extreme values of the arrival process. Our approach consists of three steps.

1. We construct a suitable model for capturing the diurnal and seasonal dependencies which takes into account a specific time-structure of durations. We utilize a cubic spline approach and propose to estimate the parameters by the weighted ordinary least square method to properly adjust durations during hours that exhibit a small median but a huge dispersion.
2. We argue that the GAS models based on the generalized gamma distribution and its special cases fit the data better than their static counterparts. This is due to the fact that the static models ignore the autocorrelation structure which is still present even after the proper diurnal and seasonal adjustment.
3. We compare both static and dynamic models in the simulation study of queueing systems with single and multiple servers and exponential services. We show that ignoring the autocorrelation structure leads to biased performance measures. The number of customers in the system, the busy period of servers and the response time have higher mean and variance as well as heavier tails for the proposed dynamic arrivals model than for the standard static model. Most importantly, we show how the trust in the standard static model for inter-arrival times leads to suboptimal decisions and consequently to a profit loss.

A proper treatment of arrival dependence is of a great importance since its ignorance generates extra costs. Our approach is useful for process simulations and consequently for process optimization and process quality assessment.

Acknowledgements

We would like to thank organizers and participants of the 7th International Conference on Management (Nový Smokovec, September 26–29, 2018), 30th European Conference on Operational Research (Dublin, June 23–26, 2019), 15th International Symposium on Operations Research in Slovenia (Bled, September 25–27, 2019) and 3rd International Conference on Advances in Business and Law (Dubai, November 23–24, 2019) for fruitful discussions.

Funding

The work on this paper was supported by the Internal Grant Agency of the University of Economics, Prague under project F4/27/2020 and the Czech Science Foundation under project 19-08985S.

References

- ADAN, I. J. B. F., KULKARNI, V. G. 2003. Single-Server Queue with Markov-Dependent Inter-Arrival and Service Times. *Queueing Systems*. Volume 45. Issue 2. Pages 113–134. ISSN 0257-0130. <https://doi.org/10.1023/a:1026093622185>.
- ALTIOK, T., MELAMED, B. 2001. The Case for Modeling Correlation in Manufacturing Systems. *IIE Transactions*. Volume 33. Issue 9. Pages 779–791. ISSN 0740-817X. <https://doi.org/10.1080/07408170108936872>.
- BAUWENS, L. 2006. Econometric Analysis of Intra-Daily Trading Activity on the Tokyo Stock Exchange. *Monetary and Economic Studies*. Volume 24. Issue 1. Pages 1–24. ISSN 0288-8432. <http://www.imes.boj.or.jp/research/abstracts/english/me24-1-1.html>.

- BAUWENS, L., GIOT, P., GRAMMIG, J., VEREDAS, D. 2004. A Comparison of Financial Duration Models via Density Forecasts. *International Journal of Forecasting*. Volume 20. Issue 4. Pages 589–609. ISSN 0169-2070. <https://doi.org/10.1016/j.ijforecast.2003.09.014>.
- BLASQUES, F., KOOPMAN, S. J., LUCAS, A. 2014. Stationarity and Ergodicity of Univariate Generalized Autoregressive Score Processes. *Electronic Journal of Statistics*. Volume 8. Issue 1. Pages 1088–1112. ISSN 1935-7524. <https://doi.org/10.1214/14-ejs924>.
- BLASQUES, F., GORGI, P., KOOPMAN, S. J., WINTENBERGER, O. 2018. Feasible Invertibility Conditions and Maximum Likelihood Estimation for Observation-Driven Models. *Electronic Journal of Statistics*. Volume 12. Issue 1. Pages 1019–1052. ISSN 1935-7524. <https://doi.org/10.1214/18-ejs1416>.
- BLASQUES, F., HOLÝ, V., TOMANOVÁ, P. 2020. *Zero-Inflated Autoregressive Conditional Duration Model for Discrete Trade Durations with Excessive Zeros*. Working Paper. <https://arxiv.org/abs/1812.07318>.
- BLAZSEK, S., LICHT, A. 2020. Dynamic Conditional Score Models: A Review of Their Applications. *Applied Economics*. Volume 52. Issue 11. Pages 1181–1199. ISSN 0003-6846. <https://doi.org/10.1080/00036846.2019.1659498>.
- BOLLERSLEV, T. 1986. Generalized Autoregressive Conditional Heteroskedasticity. *Journal of Econometrics*. Volume 31. Issue 3. Pages 307–327. ISSN 0304-4076. [https://doi.org/10.1016/0304-4076\(86\)90063-1](https://doi.org/10.1016/0304-4076(86)90063-1).
- BRUCE, P., BRUCE, A. 2017. *Practical Statistics for Data Scientists: 50 Essential Concepts*. Sebastopol. O'Reilly Media. ISBN 978-1-4919-5295-5. <https://www.oreilly.com/library/view/practical-statistics-for/9781491952955/>.
- BRUZDA, J. 2020. Multistep Quantile Forecasts for Supply Chain and Logistics Operations: Bootstrapping, the GARCH Model and Quantile Regression Based Approaches. *Central European Journal of Operations Research*. Volume 28. Issue 1. Pages 309–336. ISSN 1435-246X. <https://doi.org/10.1007/s10100-018-0591-2>.
- BUCHHOLZ, P., KRIEGE, J. 2017. Fitting Correlated Arrival and Service Times and Related Queueing Performance. *Queueing Systems*. Volume 85. Issue 3-4. Pages 337–359. ISSN 0257-0130. <https://doi.org/10.1007/s11134-017-9514-5>.
- CIVELEK, I., BILLER, B., SCHELLER-WOLF, A. 2009. *The Impact of Dependence on Queueing Systems*. Working Paper. <https://www.researchgate.net/publication/228814043>.
- CREAL, D., KOOPMAN, S. J., LUCAS, A. 2013. Generalized Autoregressive Score Models with Applications. *Journal of Applied Econometrics*. Volume 28. Issue 5. Pages 777–795. ISSN 0883-7252. <https://doi.org/10.1002/jae.1279>.
- ENGLE, R. F., RUSSELL, J. R. 1998. Autoregressive Conditional Duration: A New Model for Irregularly Spaced Transaction Data. *Econometrica*. Volume 66. Issue 5. Pages 1127–1162. ISSN 0012-9682. <https://doi.org/10.2307/2999632>.
- FERNANDES, M., GRAMMIG, J. 2005. Nonparametric Specification Tests for Conditional Duration Models. *Journal of Econometrics*. Volume 127. Issue 1. Pages 35–68. ISSN 0304-4076. <https://doi.org/10.1016/j.jeconom.2004.06.003>.
- FINCH, P. D. 1963. The Single Server Queueing System with Non-Recurrent Input-Process and Erlang Service Time. *Journal of the Australian Mathematical Society*. Volume 3. Issue 2. Pages 220–236. ISSN 1446-8107. <https://doi.org/10.1017/s1446788700027968>.

- FINCH, P. D., PEARCE, C. 1965. A Second Look at a Queueing System with Moving Average Input Process. *Journal of the Australian Mathematical Society*. Volume 5. Issue 1. Pages 100–106. ISSN 1446-8107. <https://doi.org/10.1017/s144678870002591x>.
- HARVEY, A. C. 2013. *Dynamic Models for Volatility and Heavy Tails: With Applications to Financial and Economic Time Series*. First Edition. New York. Cambridge University Press. ISBN 978-1-107-63002-4. <https://doi.org/10.1017/cbo9781139540933>.
- HAUTSCH, N. 2003. Assessing the Risk of Liquidity Suppliers on the Basis of Excess Demand Intensities. *Journal of Financial Econometrics*. Volume 1. Issue 2. Pages 189–215. ISSN 1479-8409. <https://doi.org/10.1093/jjfinec/nbg010>.
- HWANG, G. U., SOHRABY, K. 2003. On the Exact Analysis of a Discrete-Time Queueing System with Autoregressive Inputs. *Queueing Systems*. Volume 43. Issue 1-2. Pages 29–41. ISSN 0257-0130. <https://doi.org/10.1023/a:1021848330183>.
- KAMOUN, F. 2006. The Discrete-Time Queue with Autoregressive Inputs Revisited. *Queueing Systems*. Volume 54. Issue 3. Pages 185–192. ISSN 0257-0130. <https://doi.org/10.1007/s11134-006-9591-3>.
- KAYACI ÇODUR, M., YILMAZ, M. 2020. A Time-Dependent Hierarchical Chinese Postman Problem. *Central European Journal of Operations Research*. Volume 28. Issue 1. Pages 337–366. ISSN 1435-246X. <https://doi.org/10.1007/s10100-018-0598-8>.
- KOOPMAN, S. J., LUCAS, A., SCHARTH, M. 2016. Predicting Time-Varying Parameters with Parameter-Driven and Observation-Driven Models. *Review of Economics and Statistics*. Volume 98. Issue 1. Pages 97–110. ISSN 0034-6535. https://doi.org/10.1162/rest_a_00533.
- LIVNY, M., MELAMED, B., TSIOLIS, A. K. 1993. The Impact of Autocorrelation on Queueing Systems. *Management Science*. Volume 39. Issue 3. Pages 322–339. ISSN 0025-1909. <https://doi.org/10.2307/2632647>.
- LUCAS, A. 2020. *Generalized Autoregressive Score Models*. Online. <http://www.gasmodel.com>.
- LUNDE, A. 1999. *A Generalized Gamma Autoregressive Conditional Duration Model*. Working Paper. <https://www.researchgate.net/publication/228464216>.
- MANAFZADEH DIZBIN, N., TAN, B. 2019. Modelling and Analysis of the Impact of Correlated Inter-Event Data on Production Control Using Markovian Arrival Processes. *Flexible Services and Manufacturing Journal*. Volume 31. Issue 4. Pages 1042–1076. ISSN 1936-6582. <https://doi.org/10.1007/s10696-018-9329-7>.
- MIAO, D. W. C., LEE, H. C. 2013. Second-Order Performance Analysis of Discrete-Time Queues Fed by DAR(2) Sources with a Focus on the Marginal Effect of the Additional Traffic Parameter. *Applied Stochastic Models in Business and Industry*. Volume 29. Issue 1. Pages 45–60. ISSN 1524-1904. <https://doi.org/10.1002/asmb.939>.
- NIELSEN, E. H. 2007. Autocorrelation in Queuing Network-Type Production Systems - Revisited. *International Journal of Production Economics*. Volume 110. Issue 1-2. Pages 138–146. ISSN 0925-5273. <https://doi.org/10.1016/j.ijpe.2007.02.014>.
- PACURAR, M. 2008. Autoregressive Conditional Duration Models in Finance: A Survey of the Theoretical and Empirical Literature. *Journal of Economic Surveys*. Volume 22. Issue 4. Pages 711–751. ISSN 0950-0804. <https://doi.org/10.1111/j.1467-6419.2007.00547.x>.
- PATUWO, B. E., DISNEY, R. L., MCNICKLE, D. C. 1993. The Effect of Correlated Arrivals on Queues. *IIE Transactions*. Volume 25. Issue 3. Pages 105–110. ISSN 0740-817X. <https://doi.org/10.1080/07408179308964296>.

- PEARCE, C. 1967. An Imbedded Chain Approach to a Queue with Moving Average Input. *Operations Research*. Volume 15. Issue 6. Pages 1117–1130. ISSN 0030-364X. <https://doi.org/10.1287/opre.15.6.1117>.
- RESNICK, S., SAMORODNITSKY, G. 1997. Performance Decay in a Single Server Exponential Queueing Model with Long Range Dependence. *Operations Research*. Volume 45. Issue 2. Pages 235–243. ISSN 0030-364X. <https://doi.org/10.1287/opre.45.2.235>.
- SARANJEET, K. B., RAMANATHAN, T. V. 2018. Conditional Duration Models for High-Frequency Data: A Review on Recent Developments. *Journal of Economic Surveys*. Volume 33. Issue 1. Pages 252–273. ISSN 0950-0804. <https://doi.org/10.1111/joes.12261>.
- STACY, E. W. 1962. A Generalization of the Gamma Distribution. *The Annals of Mathematical Statistics*. Volume 33. Issue 3. Pages 1187–1192. ISSN 0003-4851. <https://doi.org/10.2307/2237889>.
- SZEKLI, R., DISNEY, R. L., HUR, S. 1994. MR/GI/1 Queues by Positively Correlated Arrival Stream. *Journal of Applied Probability*. Volume 31. Issue 2. Pages 497–514. ISSN 0021-9002. <https://doi.org/10.1017/s0021900200045009>.
- TIN, P. 1985. A Queueing System with Markov-Dependent Arrivals. *Journal of Applied Probability*. Volume 22. Issue 3. Pages 668–677. ISSN 0021-9002. <https://doi.org/10.1017/s0021900200029417>.
- TOMANOVÁ, P. 2018. Measuring Intensity of Order Arrivals and Process Quality Assessment of an Online Bookshop: A Case Study from the Czech Republic. In *Proceedings of the 7th International Conference on Management*. Prešov. Bookman s.r.o. Pages 768–773. ISBN 978-80-8165-301-8. <http://www.managerconf.com/>.
- TOMANOVÁ, P. 2019. Clustering of Arrivals and Its Impact on Process Simulation. In *Proceedings of the 15th International Symposium on Operations Research in Slovenia*. Bled. Slovenian Society Informatika. Pages 314–319. ISBN 978-961-6165-55-6. <http://fgg-web.fgg.uni-lj.si/{~}/sdrobne/sor/SOR'19-Proceedings.pdf>.