

Detection of genuine tripartite entanglement by two bipartite entangled states

Yize Sun^{1,*} and Lin Chen^{1,2,†}

¹*School of Mathematical Sciences, Beihang University, Beijing 100191, China*

²*International Research Institute for Multidisciplinary Science, Beihang University, Beijing 100191, China*

(Dated: April 2, 2020)

It is an interesting problem to construct genuine tripartite entangled states based on the collective use of two bipartite entangled states. We consider the case that the states are two-qubit Werner states, we construct the interval of parameter of Werner states such that the tripartite state is genuine entangled. Further, we present the way of detecting the tripartite genuine entanglement using current techniques in experiments. We also investigate the lower bound of genuine multipartite entanglement concurrence.

PACS numbers: 03.65.Ud, 03.67.Mn

I. INTRODUCTION

Genuine multipartite entanglement offers significant advantages compared with bipartite entanglement in quantum tasks [1–3]. Multipartite private states from which secret keys are directly accessible to trusted partners are genuinely multipartite entangled states [4]. Furthermore, genuine entanglement (GE) is the basic ingredient in measurement-based quantum computation [5, 6], and is beneficial in various quantum communication protocols [7]. Many efforts have been devoted towards the detection of genuine entanglement. For instance, GE can be computed efficient by the generalized geometric measure [8], a series of linear and nonlinear entanglement witnesses [9–17], generalized concurrence [18–21] and Bell-like inequalities [22]. Although these methods were derived and a characterization in terms of semidefinite programs was developed [23], the problem of detecting GE remains far from being satisfactorily solved.

In the one-dimensional cluster-Ising model [24], the genuine tripartite entanglement between three adjacent spins is the only source of genuine multipartite entanglement. The tripartite state is either a GE state or a biseparable state. The determination of bipartite entanglement has much more tools in theory and experiments [25, 26] than that of tripartite states [10]. It is known that all positive partial transpose (PPT) state are never distillable, so distillable state must be non-positive partial transpose (NPT) state [52]. The distillability problem is a main open problem of quantum information. It asks whether bipartite NPT states can be asymptotically converted into pure entangled states under local operations and classical communications (LOCC) [27, 28]. To solve the problem, there are several attempts by converting states into Werner states by LOCC [29–31]. On the other hand, progress towards distilling entangled states of given dimensions or deficient rank has been made steadily. For example, in [32], it shows that a bipartite NPT quantum state of rank four is distillable. Besides, any biseparable state is a PPT mixture [16]. As far as we know, the relation between genuine tripartite entangled state and NPT or PPT states is not well studied.

In this paper, we investigate the tripartite GE of tensor product of two NPT Werner states $\rho_w(p_1, d)$, $\rho_w(p_2, d)$. In [27], if Werner states form $\rho_w(p, d)$ were distillable then it equals to NPT states would be distillable through the reductions. Then we present the region of detecting GE of constructing tripartite state ρ_{ABC} containing $\rho_w(p_1, 2)_{AC_1} \otimes \rho_w(p_2, 2)_{BC_2}$ of systems A, B and (C_1C_2) in Theorem 5. Besides, we present the region of parameter p for detecting GE of tensor product of two Werner states in Theorem 5. There exist a neighborhood $h = [0, 0.05934)$ such that $\rho_w(-1 + \epsilon, 2)_{AC_1} \otimes \rho_w(-1 + \epsilon, 2)_{BC_2}$ is a tripartite genuine entangled state for all $\epsilon \in [0, h]$. Then we discuss the realization of Theorem 5 in experientment, and investigate Conjecture 3 when one of $\rho_w(p_1, 2)_{AC_1}$ and $\rho_w(p_2, 2)_{BC_2}$ is a PPT entangled state. We also correct the lower bound for genuine multipartite entanglement concurrence in Theorem 6 and apply the method in Example 7 and Example 8. Then we use it to detect the GE of $\rho_w(p_1, 2)_{AC_1} \otimes \rho_w(p_2, 2)_{BC_2}$ in Appendix A. If the conjecture is true, then we can construct genuine tripartite entanglement state from two Werner states. Furthermore, it is a special case of constructing an $(n + 2)$ -partite genuine entanglement state from two $(n + 1)$ -partite states.

In quantum information theory, the strong subadditivity plays a crucial role in nearly every nontrivial insight [33–35]. For a tripartite state ρ_{ABC} , the inequality says $S(\rho_{AC}) + S(\rho_{BC}) \geq S(\rho_{ABC}) + S(\rho_C)$, where $S(\alpha)$ is the von

*sunyize@buaa.edu.cn

†linchen@buaa.edu.cn (corresponding author)




FIG. 1: Two bipartite states of systems A, C_1 and B, C_2 can construct a tripartite state ρ_{ABC} of systems A, B and C , where $C = C_1C_2$. If α_{AC_1} and β_{BC_2} are entangled, then we conjecture that ρ_{ABC} is a tripartite genuine entangled state. We partially prove the conjecture in Theorem 5 by showing that ρ_{ABC} is a tripartite genuine entangled state for the interval $[-1, -0.94066]$ of parameter of Werner states.

Neumann entropy of quantum state α . The inequality is saturated when there exists a decomposition of system C as $\mathcal{H}_C = \mathcal{H}_{C_1} \otimes \mathcal{H}_{C_2}$ into a tensor product $\rho_{ABC} = \rho_{AC_1} \otimes \rho_{BC_2}$, where $\rho_{AC_1} \in \mathcal{B}(\mathcal{H}_A \otimes \mathcal{H}_{C_1})$, $\rho_{BC_2} \in \mathcal{B}(\mathcal{H}_B \otimes \mathcal{H}_{C_2})$. Indeed it is a special case of the necessary and sufficient condition constructed in [34]. Studying the conjecture in Fig. 1 thus may help understand the relation between the genuine entanglement and strong subadditivity.

The rest of this paper is organized as follows. In Sec. II we introduce the preliminary knowledge used in this paper. In Sec. III we present our results on the detection of genuine entanglement. In Sec. IV we investigate the lower bound of GE concurrence. This paper ends up with a conclusion in Sec. V.

II. PRELIMINARIES

In this section we introduce the preliminary knowledge used in this paper. First, genuinely multiparty entangled state is a particularly useful notion in the theory of entanglement but also have found an application, for example, in quantum error correction and cryptography. Besides, genuinely multiparty entangled states can construct genuinely entangled subspace [36]. Next, we define the notations of generators of special unitary group $SU(d)$ and the k norm for $m \times n$ matrix. Then we present two methods to detect GE of quantum state in Theorem 1 and Theorem 2.

We review the definition of genuinely multiparty entangled state. A multipartite quantum state that is not separable with respect to any bipartition is said to be genuine multipartite entangled [37]. Denote \mathcal{H}_i^d as d -dimensional Hilbert spaces, $i = 1, 2, \dots, n$. An n -partite pure state $|\psi\rangle \in \mathcal{B}(\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_n)$ is called biseparable if it can be written as

$$|\psi\rangle = |\psi_{S_1}\rangle \otimes |\psi_{S_2}\rangle, \quad (1)$$

where $|\psi_{S_1}\rangle \in \mathcal{B}(\mathcal{H}_{S_1}) = \mathcal{B}(\mathcal{H}_{s_1} \otimes \mathcal{H}_{s_2} \otimes \dots \otimes \mathcal{H}_{s_k})$, $|\psi_{S_2}\rangle \in \mathcal{B}(\mathcal{H}_{S_2}) = \mathcal{B}(\mathcal{H}_{s_{k+1}} \otimes \mathcal{H}_{s_{k+2}} \otimes \dots \otimes \mathcal{H}_{s_n})$, and $\{s_1, \dots, s_k | s_{k+1}, \dots, s_n\}$ is a particular order of $\{1, 2, \dots, n\}$. An n -partite mixed state ρ is biseparable if it can be written as a convex combination of biseparable pure states

$$\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|, \quad (2)$$

where $0 < p_i \leq 1$, $\sum p_i = 1$, and $|\psi_i\rangle$ is biseparable with respect to different bipartitions. Otherwise, it is called genuinely n -partite entangled.

Next, let $\lambda_i, i = 1, \dots, d^2 - 1$, denote the generator of the special unitary group $SU(d)$ [38]. For example, when $d = 3$, the generators λ -matrices of the unimodular unitary group $SU(3)$ are as follows,

$$\lambda_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \lambda_2 = \begin{bmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \lambda_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

$$\lambda_4 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad \lambda_5 = \begin{bmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{bmatrix}, \quad \lambda_6 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix},$$

$$\lambda_7 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{bmatrix}, \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{bmatrix}, \quad (3)$$

and $\text{Tr}(\lambda_k \lambda_l) = 2\delta_{kl}$, $k, l = 1, 2, \dots, 8$. Besides, the generators of $\text{SU}(d)$ are the elements of Bloch vector, which gives the desirable description of the states for N -level systems. Let I be the $d \times d$ identity matrix. Any $\rho \in \mathcal{B}(\mathcal{H}_1^d \otimes \mathcal{H}_2^d \otimes \mathcal{H}_3^d)$ can be represented as follows,

$$\begin{aligned} \rho &= \frac{1}{d^3} I \otimes I \otimes I + \frac{1}{2d^2} \left(\sum t_i^1 \lambda_i \otimes I \otimes I \right. \\ &+ \sum t_j^2 I \otimes \lambda_j \otimes I + \sum t_k^3 I \otimes I \otimes \lambda_k \left. \right) \\ &+ \frac{1}{4d} \left(\sum t_{ij}^{12} \lambda_i \otimes \lambda_j \otimes I + \sum t_{ik}^{13} \lambda_i \otimes I \otimes \lambda_k \right. \\ &+ \left. \sum t_{jk}^{23} I \otimes \lambda_j \otimes \lambda_k \right) + \frac{1}{8} \sum t_{ijk}^{123} \lambda_i \otimes \lambda_j \otimes \lambda_k. \end{aligned} \quad (4)$$

We know that $\text{Tr}(\lambda_k \lambda_l) = 2\delta_{kl}$ from the second section of [39]. Then according to (4), we obtain that $t_i^1 = \text{Tr}(\rho \lambda_i \otimes I \otimes I)$. Similarly, $t_j^2 = \text{Tr}(\rho I \otimes \lambda_j \otimes I)$, $t_k^3 = \text{Tr}(\rho I \otimes I \otimes \lambda_k)$, $t_{ij}^{12} = \text{Tr}(\rho \lambda_i \otimes \lambda_j \otimes I)$, $t_{ik}^{13} = \text{Tr}(\rho \lambda_i \otimes I \otimes \lambda_k)$, $t_{jk}^{23} = \text{Tr}(\rho I \otimes \lambda_j \otimes \lambda_k)$, and $t_{ijk}^{123} = \text{Tr}(\rho \lambda_i \otimes \lambda_j \otimes \lambda_k)$. Set $T^{(1)}, T^{(2)}, T^{(3)}, T^{(12)}, T^{(13)}, T^{(23)}$, and $T^{(123)}$ to be the vectors with the entries $t_i^1, t_j^2, t_k^3, t_{ij}^{12}, t_{ik}^{13}, t_{jk}^{23}$ and t_{ijk}^{123} , $i, j, k = 1, 2, \dots, d^2 - 1$. Let T_{123}, T_{213} and T_{312} be the matrices with entries $t_{i,(d^2-1)(j-1)+k} = t_{ijk}$, $t_{j,(d^2-1)(i-1)+k} = t_{ijk}$ and $t_{k,(d^2-1)(i-1)+j} = t_{ijk}$, respectively. Then we show three norms for an $m \times n$ matrix M as follows,

$$\|M\| := \sqrt{\sum_{i,j} M_{i,j}^2} = \sqrt{\sum_i \sigma_i^2}, \quad (5)$$

$$\|M\|_{tr} := \text{Tr} \sqrt{M^T M} = \sum_i \sigma_i, \quad (6)$$

$$\|M\|_k := \sum_{i=1}^k \sigma_i, \quad (7)$$

where $\{\sigma_i\}$ ($i = 1, \dots, \min(m, n)$) denote the singular values of the matrix, which are in non-increasing order. Notice that the last Ky Fan norm is the trace norm.

Let $\|M\|_k = \sum_{i=1}^k \sigma_i$ denote the k norm for an $n \times n$ matrix M , where $\sigma_i, i = 1, \dots, n$, are the singular values of M in decreasing order. After by presenting the definition of $M_k(\rho)$, then we obtain Theorem 1 in the following.

Theorem 1 Consider the average matricization norm

$$M_k(\rho) = \frac{1}{3} (\|T_{123}\|_k + \|T_{213}\|_k + \|T_{312}\|_k) \quad (8)$$

for a tripartite qudit state ρ . If it holds that

$$M_k(\rho) > \frac{2\sqrt{2}}{3} (2\sqrt{k} + 1) \frac{d-1}{d} \sqrt{\frac{d+1}{d}} \quad (9)$$

for any $k = 1, 2, \dots, d^2 - 1$, then ρ is a genuine multipartite entangled.

In the following, by using Theorem 1, we shall detect the genuine entanglement of another type of tripartite states. This state is tensor product of two Werner states, which are invariant under all unitaries of the form $U \otimes U$ [40]. Theorem 1 can detect GE not only for tripartite qubit systems but for any tripartite qudit system.

Next, denote $\|\cdot\|$ the Frobenius norm of a vector or a matrix $\|M\| = \|M\|_{KF}$ is just the Ky-Fan norm. From Vicente criterion in [41], then we have Theorem 2.

Theorem 2 For an arbitrary tripartite qudit state it holds that

$$\|T_{jlm}\| > \sqrt{\frac{8(d-1)(d^2-1)}{d^3}}, \quad (10)$$

then the state is genuine multipartite entangled.

The result shows that a high value of this measure can imply not only some entanglement but even genuine multipartite entanglement. The power of this condition increases with the subsystem dimension improving remarkably on [10]. By applying the condition of Theorem 2, all the quantities are invariant under local unitary transformations on the density matrix, then we can detect GE of the states in Example 8. Hence, if the lower bound is already large enough to violate the inequality given by Theorem 2, then we conclude with certainty the presence of genuine multipartite entanglement. On the analogy of Theorem 2, we also hope to improve it for larger d .

III. GENUINE ENTANGLEMENT DETECTION OF TENSOR PRODUCT OF TWO BIPARTITE ENTANGLED STATES

In this section, we consider how to construct a tripartite GE state for two bipartite entangled states α_{AC_1} and β_{BC_2} by involving the tensor product and the Kronecker product, it provides a systematical method to construct GE states. We mainly investigate the following conjecture 3 for bipartite entangled states. It's known that each NPT bipartite state can be convert into an NPT Werner state by using LOCC, then we introduce the definition of Werner state at first. The Werner state on $\mathbb{C}^d \otimes \mathbb{C}^d$ is defined as

$$\rho_w(p, d) := \frac{1}{d^2 + pd} (I_d \otimes I_d + p \sum_{i,j=0}^{d-1} |i, j\rangle\langle j, i|), \quad (11)$$

where the parameter $p \in [-1, 1]$. It has been proved [27] that $\rho_w(p, d)$ is (i) separable when $p \in [-\frac{1}{d}, 1]$; (ii) NPT and one-copy undistillable when $p \in [-\frac{1}{2}, -\frac{1}{d})$; and (iii) NPT and one-copy distillable when $p \in [-1, -\frac{1}{2})$. Hence studying the Werner state with $p = -1/2$ and 1 would characterize the behavior of Werner states over the whole interval of p . This is one of the motivations why we propose the following conjecture.

Conjecture 3 (i) Suppose α_{AC_1} and β_{BC_2} are two bipartite entangled states of systems A, C_1 and systems B, C_2 , respectively. Then $\alpha_{AC_1} \otimes \beta_{BC_2}$ is a GE state of systems A, B and C , where $C = (C_1 C_2)$.

(ii) Suppose α_{AC_1} and β_{BC_2} are two-qubit entangled states. Then we have $\alpha_{AC_1} \otimes \beta_{BC_2}$ is a tripartite state if and only if there is a neighborhood $[0, h)$, for all $\epsilon \in [0, h)$, the state $\rho_w(2, -1 + \epsilon)_{AC_1} \otimes \rho_w(2, -1 + \epsilon)_{BC_2}$ is a tripartite GE state.

One may show that Conjecture 3 (ii) is a special case of (i). We first consider to attack the generic one in Theorem 4, by introducing a known result from [42].

Theorem 4 Conjecture 3 (i) holds if the range of α or β is not spanned by product states.

Next, to detect the GE of tripartite state $\rho_w(2, -1 + \epsilon)_{AC_1} \otimes \rho_w(2, -1 + \epsilon)_{BC_2}$ in Conjecture 3, we construct a special state ρ in Theorem 5.

Theorem 5 Consider a mixed state $\rho = \frac{1-x}{64}I + x(\alpha_{AC_1} \otimes \beta_{BC_2})$, where α_{AC_1} and β_{BC_2} are two-qubit NPT Werner state, then $\alpha_{AC_1} \otimes \beta_{BC_2} \in \mathcal{H}_{ABC}$ is a tripartite state with $C = C_1 C_2$.

(i) When $p_1 = p_2 = -1$, the region of the GE detection of state ρ is maximum for $0.902646 < x \leq 1$.

(ii) When $x = 1$, then the GE of state ρ can be detected for $-1 \leq p_1 \leq -0.940198$, $-1 \leq p_2 \leq -0.94066$.

Proof. (i) Suppose $\alpha_{AC_1} = \rho_w(p_1, 2) = \frac{1}{(4+2p_1)}(I_2 \otimes I_2 + p_1 \sum_{i,j=0}^1 |i, j\rangle\langle j, i|)$ and $\beta_{BC_2} = \rho_w(p_2, 2) = \frac{1}{(4+2p_2)}(I_2 \otimes I_2 + p_2 \sum_{i,j=0}^1 |i, j\rangle\langle j, i|)$, where $p_1, p_2 \in [-1, -\frac{1}{2})$. It shows that $\rho \in \mathcal{B}(\mathcal{H}^2 \otimes \mathcal{H}^2 \otimes \mathcal{H}^4)$, then we have $\rho \in \mathcal{B}(\mathcal{H}^4 \otimes \mathcal{H}^4 \otimes \mathcal{H}^4)$. The generators λ -matrices of SU(4) are as follows,

$$\lambda_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \lambda_2 = \begin{bmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \lambda_3 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \lambda_4 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (12)$$

$$\lambda_5 = \frac{1}{\sqrt{6}} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -3 \end{bmatrix}, \lambda_6 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \lambda_7 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \lambda_8 = \begin{bmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (13)$$

$$\lambda_9 = \begin{bmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \end{bmatrix}, \lambda_{10} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{bmatrix}, \lambda_{11} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \end{bmatrix}, \lambda_{12} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (14)$$

$$\lambda_{13} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}, \lambda_{14} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \lambda_{15} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}. \quad (15)$$

In the following, we detect GE of ρ by Theorem 1. First, we assume that $P = |0\rangle\langle 0| + |1\rangle\langle 1|$, where $|0\rangle, |1\rangle \in \mathbb{C}^4$. Then we have $(P \otimes P \otimes I_4)\rho(P \otimes P \otimes I_4) = \rho$. From the definition of t_{ijk}^{123} , we obtain that

$$\begin{aligned} t_{ijk}^{123} &= \text{Tr}(\rho(\lambda_i \otimes \lambda_j \otimes \lambda_k)) \\ &= \text{Tr}(\rho(P\lambda_i P \otimes P\lambda_j P \otimes \lambda_k)). \end{aligned} \quad (16)$$

According to (16) and the form of generators λ -matrices, we have t_{ijk}^{123} is nonzero when $i, j \in \{1, 2, 3, 4, 5\}$. By respectively computing $\|T_{\underline{1}23}\|_k$, $\|T_{\underline{2}13}\|_k$ and $\|T_{\underline{3}12}\|_k$, $k = 4$, we obtain that

$$\begin{aligned} \max M_k(\rho) &= \max \frac{1}{3} (\|T_{\underline{1}23}\|_k + \|T_{\underline{2}13}\|_k + \|T_{\underline{3}12}\|_k) \\ &= 4.37918x \end{aligned} \quad (17)$$

when $p_1 = p_2 = -1$ in Fig. 2. Suppose that $k = 4$, then we have

$$\max M_k(\rho) - \frac{5}{2}\sqrt{\frac{5}{2}} = 4.37918x - 3.95285 > 0. \quad (18)$$

By analysing the maximum of the coefficient of x in Fig. 2, then Theorem 1 can detect the GE for $0.902646 < x \leq 1$.

(ii) Consider the boundary condition of p_1 and p_2 in the region of x that GE can be detected. Then in Fig. 3, we know that the boundary condition of p_1, p_2 is

$$\frac{1}{3} (\|T_{\underline{1}23}\|_k + \|T_{\underline{2}13}\|_k + \|T_{\underline{3}12}\|_k) = \frac{5}{2}\sqrt{\frac{5}{2}}, \quad (19)$$

where $x = 1$.

□

To compare the region of x that GE of state ρ can be detected, then we introduce another method in the next section. In Appendix A, by using other two different ways to detect GE of state ρ . Thus, we obtain that the maximum region of x that can detect ρ by using Theorem 1. Let $Q = |1\rangle\langle 1| + |2\rangle\langle 2|$, then we have

$$(Q \otimes Q)\rho_w(p, d)(Q \otimes Q) \propto \rho_w(p, 2), 2 \leq d. \quad (20)$$

From (20), if $\rho_w(p_1, 2) \otimes \rho_w(p_2, 2)$ is a tripartite GE state, $\rho_w(p_1, d_1) \otimes \rho_w(p_2, d_2)$ is also a tripartite GE state. According to the results of Theorem 5, we have $\rho_w(p_1, d_1) \otimes \rho_w(p_2, d_2), 2 \leq d_1, d_2$ is also a tripartite GE state for the region of p_1, p_2 in Fig. 3. Because NPT states can be convert into NPT Werner states by LOCC, we have obtained a large set of distillable NPT states whose tensor product is a tripartite GE state.

In the following we discuss the realization of Theorem 5 experimentally. The realization of two-qubit Werner states has been extensively studied in experiment over the past several years, using techniques of photon polarization, spontaneous parametric down conversion (SPDC), and semiconductor quantum dot [43–46]. So it is feasible to practically detect the genuine entanglement of the state $\rho_{ABC} = \alpha_{AC_1} \otimes \beta_{BC_2}$ with the two bipartite NPT entangled




FIG. 2: One finds from the figure that when $p_1 \rightarrow -1, p_2 \rightarrow -1$, the region of x that GE of state ρ can be detected is increasing. Specially, when $p_1 = p_2 = -1$, the $\max M_k(\rho) = 4.37918x$ for $0 \leq x \leq 1$.




FIG. 3: One finds from the figure that $-1 \leq p_1 \leq -0.940198$ and $-1 \leq p_2 \leq -0.94066$ in the region of x that GE can be detected.

states α and β . First of all in theory, we respectively convert them into two Werner states under twirling operations $U \otimes U$ [40]. This is a two-qubit local unitary operation and may be implemented effectively. Next we experimentally prepare the two Werner states in system AC_1 and BC_2 , respectively, using the techniques in [43–46]. Note that their parameters p_1, p_2 in (11) may be influenced due to white noise. By measuring the Werner states using quantum tomography, we may determine whether the parameters p_1, p_2 are restricted in the interval in Theorem 5 (ii). If this is true then we may detect their genuine entanglement using Theorem 5 (ii). It will support Conjecture 3 from a practical point of view.

So far we have investigated Conjecture 3 for NPT entangled states α and β . Finally we discuss Conjecture 3 when one of α and β is a PPT entangled state. It follows from Theorem 4 that Conjecture 3 holds when one of α, β satisfies that its range is not spanned by product vectors. For example, the PPT entangled states constructed from the unextendible product bases (UPBs) are such states [47]. Another example is the so-called completely symmetric state $\alpha = \sum_{i=0}^6 \lambda_i |\phi_i\rangle\langle\phi_i| - \lambda |\phi_7\rangle\langle\phi_7|$ [48], where $\lambda_i, \lambda > 0$, $|\phi_i\rangle = |x_i, x_i\rangle$, for $i = 0, 1, \dots, 6$, and

$$\begin{aligned}
 |x_i\rangle &= |i\rangle, i = 0, 1, 2, 3 \\
 |x_4\rangle &= |0\rangle + |1\rangle + |2\rangle + |3\rangle, \\
 |x_5\rangle &= |0\rangle + 2|1\rangle + 3|2\rangle + 4|3\rangle, \\
 |x_6\rangle &= |0\rangle - 2|1\rangle + 3|2\rangle - 4|3\rangle.
 \end{aligned} \tag{21}$$

It has been proven that [48] by choosing a positive constant λ we obtain that α is a 4×4 PPT entangled state of rank six, and the range of α has no product vectors. Since the coefficients λ_i 's of σ are arbitrary positive numbers, we have constructed a family of PPT entangled states α satisfying Conjecture 3.

The third example is the state

$$\begin{aligned}
\beta = & (|00\rangle + |11\rangle + |22\rangle)(\langle 00| + \langle 11| + \langle 22|) \\
& + (|01\rangle + |10\rangle + |33\rangle)(\langle 01| + \langle 10| + \langle 33|) \\
& + |12\rangle\langle 12| + |13\rangle\langle 13| + |30\rangle\langle 30| + |21\rangle\langle 21| \\
& + |02\rangle\langle 02| + |20\rangle\langle 20| + |03\rangle\langle 03| + |31\rangle\langle 31|.
\end{aligned} \tag{22}$$

It has been used for the separability criteria using symmetric extension [49], and the well-known PPT square conjecture recently [50]. One can show that β is a PPT state of rank ten, and the range of β has only eight linearly independent product vectors. So β is entangled and violates Theorem 4. We have shown that such β and any state α satisfy Conjecture 3. Nevertheless, studying Conjecture 3 for the PPT entangled state β whose range is spanned by product vectors remains an open problem.

In the next section, we show a different method to detect the lower bound of the GE of quantum state, then we compare the range of detecting GE of state in the Example 7 and Example 8.

IV. THE LOWER BOUND OF DETECTING GE OF QUANTUM STATE

In this section we investigate the genuine entanglement of tripartite states in terms of the entanglement measure. As the computation of any proper entanglement measure is in general an NP-hard problem, it is crucial for the quantification of entanglement that reliable lower bounds can be derived. In this section, we present the lower bound of detecting GE of quantum state.

The GE concurrence is proved a well-defined measure [18]. For example, the GE concurrence is defined by $C_{GE}(|\psi\rangle) = \sqrt{\min\{1 - \text{Tr}\rho_1^2, 1 - \text{Tr}\rho_2^2, 1 - \text{Tr}\rho_3^2\}}$ for a pure state $|\psi\rangle \in \mathcal{B}(\mathcal{H}_1^d \otimes \mathcal{H}_2^d \otimes \mathcal{H}_3^d)$, where ρ_i is the reduced matrix for the i -th subsystem. And for mixed state $\rho \in \mathcal{B}(\mathcal{H}_1^d \otimes \mathcal{H}_2^d \otimes \mathcal{H}_3^d)$, the GE concurrence is $C_{GE}(\rho) = \min \sum_{p_\alpha, |\psi_\alpha\rangle} p_\alpha C_{GE}(|\psi_\alpha\rangle)$. The minimum is taken over all pure ensemble decompositions of ρ . We show that the GE concurrence satisfies the following fact.

Theorem 6 *For a tripartite qudit state ρ , the GE concurrence satisfies the following inequality,*

$$C_{GE}(\rho) \geq \max\left\{\frac{1}{2\sqrt{2}}\|T_\alpha^{(123)}\| - \frac{(d-1)}{d}\sqrt{\frac{d+1}{d}}, 0\right\}. \tag{23}$$

Proof. First, we consider pure states $\rho = |\psi\rangle\langle\psi|$. We have $\text{Tr}\rho^2 = 1$ and hence

$$\begin{aligned}
& \frac{1}{d^3} + \frac{1}{2d^2}(\sum (t_i^1)^2 + \sum (t_j^2)^2 + \sum (t_k^3)^2) \\
& + \frac{1}{4d}(\sum (t_{ij}^{12})^2 + \sum (t_{ik}^{13})^2 + \sum (t_{jk}^{23})^2) \\
& + \frac{1}{8}\sum (t_{ijk}^{123})^2 = 1.
\end{aligned} \tag{24}$$

We denote ρ_{jk} as the reduced density matrix for the subsystems $j \neq k = 1, 2, 3$. Using (4), we have

$$\text{Tr}\rho_1^2 = \frac{1}{d} + \frac{1}{2}\|T^{(1)}\|^2, \tag{25}$$

$$\text{Tr}\rho_{23}^2 = \frac{1}{d^2} + \frac{1}{2d}(\|T^{(2)}\|^2 + \|T^{(3)}\|^2) + \frac{1}{4}\|T^{(23)}\|^2, \tag{26}$$

$$\text{Tr}\rho_2^2 = \frac{1}{d} + \frac{1}{2}\|T^{(2)}\|^2, \tag{27}$$

$$\text{Tr}\rho_{13}^2 = \frac{1}{d^2} + \frac{1}{2d}(\|T^{(1)}\|^2 + \|T^{(3)}\|^2) + \frac{1}{4}\|T^{(13)}\|^2, \tag{28}$$

and

$$\text{Tr } \rho_3^2 = \frac{1}{d} + \frac{1}{2} \|T^{(3)}\|^2, \quad (29)$$

$$\text{Tr } \rho_{12}^2 = \frac{1}{d^2} + \frac{1}{2d} (\|T^{(1)}\|^2 + \|T^{(2)}\|^2) + \frac{1}{4} \|T^{(12)}\|^2. \quad (30)$$

Setting $C = \|T^{(123)}\|^2$ in (24), we obtain

$$\begin{aligned} \text{Tr } \rho^2 - \text{Tr } \rho_1^2 &= \left(\frac{3}{d^2} - \frac{2}{d^3} - \frac{1}{d}\right) + \frac{d-1-d^2}{2d^2} \|T^{(1)}\|^2 \\ &+ \frac{1}{2} \left(\frac{1}{d} - \frac{1}{d^2}\right) (\|T^{(2)}\|^2 + \|T^{(3)}\|^2) + \frac{1}{8} C. \end{aligned} \quad (31)$$

If we assume that $\|T^{(2)}\|^2 + \|T^{(3)}\|^2 = 0$, and we use $\|T^{(1)}\|^2 \leq \frac{2(d-1)}{d}$ of [41], then we obtain that $\text{Tr } \rho^2 - \text{Tr } \rho_1^2 \geq \frac{-(1+d)(d-1)^2}{d^3} + \frac{1}{8} C$, and the same lower bound of $\text{Tr } \rho^2 - \text{Tr } \rho_2^2$ and $\text{Tr } \rho^2 - \text{Tr } \rho_3^2$. Thus

$$\begin{aligned} C_{GE}^2(|\psi\rangle) &= \min\{1 - \text{Tr } \rho_1^2, 1 - \text{Tr } \rho_2^2, 1 - \text{Tr } \rho_3^2\} \\ &\geq \max\left\{\frac{-(1+d)(d-1)^2}{d^3} + \frac{1}{8} C, 0\right\}. \end{aligned} \quad (32)$$

Next we consider the mixed quantum state $\rho \in \mathcal{B}(\mathcal{H}_1^d \otimes \mathcal{H}_2^d \otimes \mathcal{H}_3^d)$. Let $\rho = \sum p_\alpha |\psi_\alpha\rangle\langle\psi_\alpha|$ be the optimal ensemble decomposition of ρ . We obtain that

$$\begin{aligned} C_{GE}(\rho) &= \sum p_\alpha C_{GE}(|\psi_\alpha\rangle) \\ &\geq \max\left\{\frac{1}{2\sqrt{2}} \|T_\alpha^{(123)}\| - \frac{(d-1)}{d} \sqrt{\frac{d+1}{d}}, 0\right\} \times \sum p_\alpha \\ &= \max\left\{\frac{1}{2\sqrt{2}} \|T^{(123)}\| - \frac{(d-1)}{d} \sqrt{\frac{d+1}{d}}, 0\right\}, \end{aligned} \quad (33)$$

where we have used the convexity of Frobenius norm and $\sqrt{x-y} \geq \sqrt{x} - \sqrt{y}$ for $x > y > 0$. \square

In the following we present two examples showing the power of Theorem 6.

Example 7 Consider the mixture of the GHZ state and W state in three-qubit quantum systems $\rho = \frac{1-x-y}{8} I + x|GHZ\rangle\langle GHZ| + y|W\rangle\langle W|$, where $|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$ and $|W\rangle = \frac{1}{\sqrt{3}}(|001\rangle + |010\rangle + |100\rangle)$.

To explain the example, we choose $\lambda_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$ and $\lambda_3 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ as generators λ -matrices of $SU(2)$. According to Theorem 6, then we have

$$\begin{aligned} C_{GE}(\rho) &= \sum p_\alpha C_{GE}(|\psi_\alpha\rangle) \\ &\geq \max\left\{\frac{1}{2\sqrt{2}} \|T^{(123)}(\rho)\| - \frac{1}{2} \sqrt{\frac{3}{2}}, 0\right\} \\ &= \max\left\{\frac{1}{2\sqrt{6}} (\sqrt{12x^2 + 11y^2} - 3), 0\right\}. \end{aligned} \quad (34)$$

From (34), we obtain that Theorem 6 can detect GE of state ρ when $\frac{1}{2\sqrt{6}} (\sqrt{12x^2 + 11y^2} - 3) \geq 0$ in Fig. 4. So we obtain that the lower bound of detecting GE of state ρ . (see Fig. 5).

Example 8 Consider a mixed state in three-qutrit quantum systems $\rho = \frac{1-x}{27} I + x|\psi\rangle\langle\psi|$, where $|\psi\rangle = \frac{1}{\sqrt{3}}(|012\rangle + |021\rangle + |111\rangle)$.




FIG. 4: Lower bound of GE concurrence in Theorem 6 of state ρ in Example 7.




FIG. 5: The lower bound for GE concurrence is equal to zero, then GE can be detected For $0.854794 < x \leq 1$ and $0.898272 < y \leq 1$ in Theorem 6.

To detect the genuine entanglement of ρ , then we respectively use the ways in Theorem 1, Theorem 2 and Theorem 6, the results are as follows,

(i) By using Theorem 1, we have

$$\begin{aligned}
 & M_k(\rho) - \frac{8}{9}\sqrt{\frac{2}{3}}(1 + 4\sqrt{2}) \\
 &= \frac{1}{3}(\|T_{123}\|_8 + \|T_{213}\|_8 + \|T_{312}\|_8) - \frac{8}{9}\sqrt{\frac{2}{3}}(1 + 4\sqrt{2}) \\
 &\approx 4.30179x - 3.628874 > 0.
 \end{aligned} \tag{35}$$

(ii) By using Theorem 2, we have

$$\|T_{i_1 i_2 i_3}\| - \frac{8}{3}\sqrt{\frac{2}{3}} \approx 2.17732(x - 1) > 0. \tag{36}$$

(iii) By using Theorem 6, we have

$$\begin{aligned}
 C_{GE}(\rho) &\geq \max\left\{\frac{1}{2\sqrt{2}}\|T^{(123)}\| - \frac{2}{3}\sqrt{\frac{4}{3}}, 0\right\} \\
 &\approx \max\{-0.7698 + 0.7698x, 0\}.
 \end{aligned} \tag{37}$$

In Fig. 6, the lower bound of GE concurrence in Theorem 1 can detect GE better than Theorem 2 and the Theorem 6.




FIG. 6: GE detection: dashed brown line by Theorem 6 and solid blue line in Theorem 2 cannot detect GE for the whole region of x . When $d = 3, k = 4$, Theorem 1 in this paper detect GE for $0.843573 < x \leq 1$.

To conclude, we point out that some results of this section have corrected some suspicious claims from [51]. First in Theorem 6, our article corrects the maximum lower bound for genuine multipartite entanglement concurrence in terms of the norms of the correlation tensors of $T_{\alpha}^{(123)}$. Next, in Example 7, we have corrected the lower bound for genuine multipartite entanglement concurrence. In Example 8, we have corrected the representation of $M(\rho)$ and the lower bound for GE concurrence. Then we obtain that the result of Theorem 6 cannot detect tripartite GE in Example 8.

V. CONCLUSIONS

In this paper, we have asked whether the product of two entangled states $\alpha_{AC_1}, \beta_{BC_2}$ is still tripartite genuine entangled, which has been formulated by Conjecture 3. Then we have discussed the realization of Conjecture 3 in experiment by using two Werner states. By restricting the interval of parameter p_1 and p_2 , we can detect the tripartite genuine entanglement. Besides, we also have corrected the lower bound for genuine multipartite entanglement concurrence of any quantum states, and have detected genuine entanglement in two examples by using it.

A direct open problem from this paper is to keep studying Cojecture 3 for region of parameter p of two bipartite NPT states and more general cases. However, it is also very interesting to find out a counterexample, because it shows the physical difference between bipartite and tripartite genuine entanglement.

Acknowledgments

Authors were supported by the NNSF of China (Grant No. 11871089), and the Fundamental Research Funds for the Central Universities (Grant Nos. KG12080401 and ZG216S1902).

Appendix A: GE detection of ρ by Other ways compare with Theorem 5

To compare the region of x for detecting GE of state ρ in Theorem 5, there are two different ways (i), (ii) as follows, (i) By using Theorem 2, we have

$$\|T_{i_1 i_2 i_3}\| - \frac{3}{2}\sqrt{\frac{5}{2}} = \sqrt{3} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x - \frac{3}{2}\sqrt{\frac{5}{2}} > 0 \quad (\text{A1})$$

where $p_1, p_2 \in [-1, -\frac{1}{2}]$. To maximum detect the GE, then we find the maximum slope of x in (A1). By analysing the maximum of the coefficient of x , we have the maximum of $\sqrt{3} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x$ is $\sqrt{6}$, when $p_1 = p_2 = -1$. then Theorem 2 can detect the GE of state ρ for $0.968246 < x \leq 1$.




FIG. 7: Under the condition that $p_1 = p_2 = -1$, then we have the region of detecting GE of state ρ is maximum. One finds in this figure that solid green line by Theorem 6 and solid blue line in Theorem 2 can detect GE of ρ in the same region $(0.968246, 1]$. The dashed brown line by Theorem 1 can detect GE of ρ in $(0.902646, 1]$.

On the other hand, in the region of x that GE of state ρ can be detected, we consider the boundary condition that can detect GE of state ρ . From (A1), we obtain that the GE of ρ can be detected when

$$\sqrt{3} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x \geq \frac{3}{2} \sqrt{\frac{5}{2}}. \quad (\text{A2})$$

Then we have the boundary condition that can detect GE of state ρ is

$$\sqrt{3} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x = \frac{3}{2} \sqrt{\frac{5}{2}}. \quad (\text{A3})$$

From (A3), then we obtain that negative correlation between p_1 and p_2 for $-1 \leq p_1, p_2 \leq -0.981475$. In the region of p_1, p_2 , the GE of state ρ can be detected. Specially, when $p_1 = p_2 = -1$, the region of GE detection of state ρ is maximum.

(ii) By using Theorem 6, we have

$$\|T^{(123)}\| = \sqrt{3} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x. \quad (\text{A4})$$

Then we obtain

$$\begin{aligned} C_{GE}(\rho) &\geq \max\left\{\frac{1}{2\sqrt{2}}\|T^{(123)}\| - \frac{3}{4}\sqrt{\frac{5}{4}}, 0\right\} \\ &= \max\left\{\frac{\sqrt{3}}{2\sqrt{2}} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x - \frac{3}{4}\sqrt{\frac{5}{4}}, 0\right\}. \end{aligned} \quad (\text{A5})$$

From the maximum of $\sqrt{3} \frac{\sqrt{p_2^2 + p_1 p_2^2 + p_1^2(1 + p_2 + 2p_2^2)}}{(2 + p_1)(2 + p_2)} x = \sqrt{6}$ when $p_1 = p_2 = -1$, then we have $\max\|T^{(123)}\| = \sqrt{6}x$. So we obtain that Theorem 6 can detect the GE of ρ for $0.968246 < x \leq 1$.

Thus, by comparing the region of x for detecting GE of state ρ in Fig. 7, we obtain that the Theorem 5 can detect GE of state ρ for the maximum region of x .

-
- [1] Ryszard Horodecki, Pawel Horodecki, Michal Horodecki, and Karol Horodecki. Quantum entanglement. 2007.
 - [2] Michael A. Nielsen and Isaac L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition*. 2011.
 - [3] David P. Divincenzo. Quantum computation. *Science*, 270(5234):255–261, 1995.
 - [4] Siddhartha Das, Stefan Buml, Marek Winczewski, and Karol Horodecki. Universal limitations on quantum key distribution over a network, 2019.

- [5] H. J. Briegel, D. E. Browne, W. Dr, R. Raussendorf, and M. Van Den Nest. Measurement-based quantum computation. *Nature Physics*, 111(21):65–118, 2009.
- [6] Robert Raussendorf and Hans J. Briegel. A one-way quantum computer. *Physical Review Letters*, 86(22):5188–5191.
- [7] Aditi Sen De and Ujjwal Sen. Quantum advantage in communication networks.
- [8] Saptarshi Roy, Tamoghna Das, and Aditi Sen De. Computable genuine multimode entanglement measure: Gaussian vs. non-gaussian, 2019.
- [9] Won Kyu Calvin Sun, Alexandre Cooper, and Paola Cappellaro. Improved entanglement detection with subspace witnesses, 2019.
- [10] Marcus Huber, Florian Mintert, Andreas Gabriel, and Beatrix C. Hiesmayr. Detection of high-dimensional genuine multipartite entanglement of mixed states. *Physical Review Letters*, 104(21):210501.
- [11] Marcus Huber and Ritabrata Sengupta. Witnessing genuine multipartite entanglement with positive maps. *Physical Review Letters*, 113(10):100501.
- [12] Julio I. De Vicente and Marcus Huber. Multipartite entanglement detection from correlation tensors. *Physical Review A*, 84(6):062306.
- [13] Marcus Huber, Mart Perarnau-Llobet, and Julio I. De Vicente. Entropy vector formalism and the structure of multidimensional entanglement in multipartite systems. *Physical Review A*, 88(4):109–112, 2013.
- [14] J. Sperl and W. Vogel. Multipartite entanglement witnesses. *Physical Review Letters*, 111(11):110503.
- [15] R. Augusiak and J Stasiska. Positive maps, majorization, entropic inequalities and detection of entanglement. *New Journal of Physics*, 11(5):053018, May 2009.
- [16] Bastian Jungnitsch, Tobias Moroder, and Otfried Ghne. Taming multiparticle entanglement. *Physical Review Letters*, 106(19):190502.
- [17] Valerie Coffman, Joydip Kundu, and William K. Wootters. Distributed entanglement. *Physical Review A*, 61(5):052306.
- [18] Zhi Hao Ma, Zhi-Hua Chen, Jing-Ling Chen, Christoph Spengler, Andreas Gabriel, and Marcus Huber. Measure of genuine multipartite entanglement with computable lower bounds. *Physical Review A*, 83(6):062325.
- [19] Zhi Hua Chen, Zhi-Hao Ma, Jing-Ling Chen, and Simone Severini. Improved lower bounds on genuine-multipartite-entanglement concurrence. *Physical Review A*, 85(6).
- [20] Yan Hong, Ting Gao, and Fengli Yan. Measure of multipartite entanglement with computable lower bounds. *Physical Review A*, 86(6):29940–29948, 2012.
- [21] Ting Gao, Fengli Yan, and S. J. Van Enk. On the permutationally invariant part of a density matrix and nonseparability of n-qubit states. *Physical Review Letters*, 112(18):180501–180501, 2014.
- [22] Jean Daniel Bancal, Nicolas Gisin, Yeong-Cherng Liang, and Stefano Pironio. Device-independent witnesses of genuine multipartite entanglement. *Physical Review Letters*, 106(25):250404.
- [23] Cecilia Lancien, Otfried Ghne, Ritabrata Sengupta, and Marcus Huber. Relaxations of separability in multipartite systems: Semidefinite programs, witnesses and volumes. *Journal of Physics A: Mathematical and Theoretical*, 48(50):505302, Nov 2015.
- [24] S M Giampaolo and B C Hiesmayr. Genuine multipartite entanglement in the cluster-ising model. *New Journal of Physics*, 16(9):093033, Sep 2014.
- [25] P. Van Loock and Samuel L. Braunstein. Multipartite entanglement for continuous variables: A quantum teleportation network. *Physical Review Letters*, 84(15):3482–3485.
- [26] L. A. Wu, M. S. Sarandy, and D. A. Lidar. Quantum phase transitions and bipartite entanglement. *Physical Review Letters*, 93(25):250404.
- [27] David P. Divincenzo, Peter W. Shor, John A. Smolin, Barbara M. Terhal, and Ashish V. Thapliyal. Evidence for bound entangled states with negative partial transpose. *Physical Review A*, 61(6):062312.
- [28] W. Dr, J. I. Cirac, M. Lewenstein, and D. Bruss. Distillability and partial transposition in bipartite systems. *Physical Review A*, 61(6):276–282, 1999.
- [29] B. Kraus, M. Lewenstein, and J. I. Cirac. Characterization of distillable and activatable states using entanglement witnesses. *Physical Review A*, 65(4):042327.
- [30] Somshubhro Bandyopadhyay and Vwani Roychowdhury. Class of n-copy undistillable quantum states with negative partial transposition. *Physical Review A*, 68(2):022319.
- [31] Reinaldo O. Vianna and Andrew C. Doherty. Distillability of werner states using entanglement witnesses and robust semidefinite programs. *Phys.rev.a*, 74(5):052306.
- [32] Lin Chen and Dragomirz. Dokovic. Distillability of non-positive-partial-transpose bipartite quantum states of rank four. *Phys.rev.a*, 94(5):052318.
- [33] Lieb and Elliott H. Proof of the strong subadditivity of quantum-mechanical entropy. *Journal of Mathematical Physics*, 14(12):1938.
- [34] Patrick Hayden, Richard Jozsa, Dnes Petz, and Andreas Winter. Structure of states which satisfy strong subadditivity of quantum entropy with equality. *Communications in Mathematical Physics*, 246(2):359–374.
- [35] Michael A Nielsen and Isaac L Chuang. Quantum computation and quantum information. *Mathematical Structures in Computer Science*, 17(6):1115–1115, 2007.
- [36] Maciej Demianowicz and Remigiusz Augusiak. An approach to constructing genuinely entangled subspaces of maximal dimension, 2019.
- [37] Otfried Ghne and Gza Tth. Entanglement detection. *Physics Reports*, 474(1):1–75, 2009.
- [38] Gen Kimura. The bloch vector for n -level systems. *Physics Letters A*, 314(5):339–349, 2003.
- [39] Sudhavathani Simon, S. P. Rajagopalan, and R. Simon. The structure of states and maps in quantum theory. *Pramana*

- Journal of Physics*, 73(3):471–483, 2009.
- [40] Werner and Reinhard F. Quantum states with einstein-podolsky-rosen correlations admitting a hidden-variable model. *Physical Review A*, 40(8):4277–4281.
 - [41] Julio I. De Vicente and Marcus Huber. Multipartite entanglement detection from correlation tensors. *Physical Review A*, 84(6):242–245, 2011.
 - [42] Yi Shen and Lin Chen. Construction of genuine multipartite entangled states.
 - [43] M. Barbieri, F. De Martini, G. Di Nepi, and P. Mataloni. Generation and characterization of werner states and maximally entangled mixed states by a universal source of entanglement. *Physical Review Letters*, 92(17):177901.
 - [44] H. Kumano, H. Nakajima, T. Kuroda, T. Mano, K. Sakoda, and I. Suemune. Nonlocal biphoton generation in a werner state from a single semiconductor quantum dot. *Physical Review B*, 91(20):205437.
 - [45] C. Cinelli, G. Di Nepi, F. De Martini, M. Barbieri, and P. Mataloni. Parametric source of two-photon states with a tunable degree of entanglement and mixing: Experimental preparation of werner states and maximally entangled mixed states. *Physical Review A*, 70(2):022321.
 - [46] Yong Sheng Zhang, Yun Feng Huang, Chuan Feng Li, and Guang Can Guo. Experimental preparation of the werner state via spontaneous parametric down-conversion. 66(6):317–322, 2002.
 - [47] David P. Divincenzo, Tal Mor, Peter W. Shor, John A. Smolin, and Barbara M. Terhal. Unextendible product bases, uncompletable product bases and bound entanglement. *Communications in Mathematical Physics*, 238(3):379–410.
 - [48] Lin Chen, Delin Chu, Lilong Qian, and Yi Shen. Separability of completely symmetric states in a multipartite system. *Physical Review A*, 99(3).
 - [49] Andrew C. Doherty, Pablo A. Parrilo, and Federico M. Spedalieri. Complete family of separability criteria. *Physical Review A*, 69(2):022308.
 - [50] Lin Chen, Yu Yang, and Wai-Shing Tang. Positive-partial-transpose square conjecture for. *Physical Review A*, 2019.
 - [51] Li Ming, Lingxia Jia, Wang Jing, Shuqian Shen, and Shao Ming Fei. Measure and detection of genuine multipartite entanglement for tripartite systems. *Phys.rev.a*, 96(5):052314, 2017.
 - [52] If a quantum state is still positive after partial transpose, then the state is PPT. Otherwise, it is NPT.