

## ON SARNAK'S DENSITY CONJECTURE AND ITS APPLICATIONS

KONSTANTIN GOLUBEV AND AMITAY KAMBER

ABSTRACT. Sarnak's Density Conjecture is an explicit bound on the multiplicities of non-tempered representations in a sequence of cocompact congruence arithmetic lattices in a semisimple Lie group, which is motivated by the work of Sarnak and Xue ([53]). The goal of this work is to discuss similar hypotheses, their interrelation and applications. We mainly focus on two properties — the spectral Spherical Density Hypothesis and the geometric Weak Injective Radius Property. Our results are strongest in the  $p$ -adic case, where we show that the two properties are equivalent, and both imply Sarnak's General Density Hypothesis. One possible application is that either the limit multiplicity property or the weak injective radius property imply Sarnak's Optimal Lifting Property ([52]). Conjecturally, all those properties should hold in great generality. We hope that this work will motivate their proofs in new cases.

## 1. INTRODUCTION

Let  $k$  be a local field (Archimedean or non-Archimedean), let  $G$  be the  $k$ -rational points of a semisimple algebraic group defined over  $k$ , let  $\Gamma_1 \subset G$  be a lattice and let  $\{\Gamma_N\}$  be a sequence of finite index subgroups of  $\Gamma_1$  with  $[\Gamma_1 : \Gamma_N] \rightarrow \infty$ . There are various results about multiplicities of irreducible representations appearing in the decomposition of  $L^2(\Gamma_N \backslash G)$  (e.g. [14, 54, 1]). An extremely strong property is the *very naive Ramanujan property*, stating that if  $\pi$  is non-tempered and non-trivial, then it does not appear in the decomposition. However, the very naive Ramanujan property is usually not true in high rank (see, e.g., [8]). Notice that we do not make a distinction between cusp forms and non-cusp forms — the *naive Ramanujan conjecture* states that cusp forms are tempered, and even this is not true ([32]). Moreover, even when expected to be true, the Generalized Ramanujan Conjecture seems to be out of reach by the existing methods.

Recently, Sarnak made a density conjecture which is an approximation to the very naive Ramanujan property and should serve as a replacement of it for applications. Some instances of this general idea were previously given for hyperbolic surfaces ([52, 27]) and for graphs ([6, 35]). Our goal here is to give a general framework for the proof of similar density conjectures and their use in applications.

---

Konstantin Golubev, ETH, k.golubev@gmail.com

Amitay Kamber, Einstein Institute of Mathematics, The Hebrew University of Jerusalem, amitay.kamber@gmail.com.

We give a geometric and somewhat elementary approach to the problem. An alternative approach based on deep results in the Langlands program may be found in an ongoing work of Shai Evra.

To state Sarnak's Density Conjecture, we first set some notations. Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  and let  $0 \neq v_0 \in V$  be a  $K$ -finite vector, where  $K$  is a maximal compact subgroup of  $G$ , which is good in the non-Archimedean case, in the sense of Bruhat and Tits ([7]). We let  $1 \leq p(\pi) \leq \infty$  be the infimum over  $p \geq 1$  such that the matrix coefficient  $\beta : G \rightarrow \mathbb{C}$ ,  $\beta(g) = \langle v_0, \pi(g)v_0 \rangle$  is in  $L^p(G)$ . It is a well-known fact that  $p(\pi)$  does not depend on the choice of  $v_0$ . Let  $\Pi(G)$  be the set of isomorphism classes of irreducible unitary representations of  $G$  endowed with the Fell topology. For a cocompact lattice  $\Gamma$  and  $(\pi, V) \in \Pi(G)$ , denote  $m(\pi, \Gamma) = \dim \text{Hom}_G(V, L^2(\Gamma \backslash G))$ , i.e., the number of times  $\pi$  occurs in the decomposition of  $L^2(\Gamma \backslash G)$ . For a subset  $A \subset \Pi(G)$ , denote  $M(A, \Gamma, p) = \sum_{\pi \in A, p(\pi) \geq p} m(\pi, \Gamma)$ .

**Conjecture 1.1** (Sarnak's Density Conjecture). *Let  $G$  be a real, semisimple, almost-simple and simply connected Lie group, let  $\Gamma_1$  be a cocompact arithmetic lattice of  $G$  and let  $\Gamma_N$  be a sequence of finite index congruence subgroups of  $\Gamma_1$ , with  $[\Gamma_1 : \Gamma_N] \rightarrow \infty$ . Then for every precompact subset  $A \subset \Pi(G)$  and  $\epsilon > 0$  there exists a constant  $C_{\epsilon, A}$  such that for every  $N$  and  $p > 2$ ,*

$$M(A, \Gamma, p) \leq C_{A, \epsilon} [\Gamma_1 : \Gamma_N]^{2/p + \epsilon}.$$

We refer to a sequence of lattices satisfying this multiplicity property as a sequence which satisfies the *General Density Hypothesis*. A similar conjecture came up in the work of Sarnak and Xue ([53]), but they only considered the case when  $A = \{\pi\}$  is a singleton. In such a case we say that the sequence of lattices satisfies the *Pointwise Multiplicity Hypothesis*.

We will prefer to work with a different spectral definition, the *Spherical Density Hypothesis*, which is easier to use for applications, and concerns only spherical representations. Let  $\Pi(G)_{\text{sph}} \subset \Pi(G)$  be the set of isomorphism classes of spherical representations, i.e., of irreducible unitary representations with a non-zero  $K$ -invariant vector. In the  $p$ -adic, or the rank 1 case,  $\{\pi \in \Pi(G)_{\text{sph}} : p(\pi) > 2\}$  is precompact, and the Spherical Density Hypothesis is simply the case when  $A = \{\pi \in \Pi(G)_{\text{sph}} : p(\pi) > 2\}$  in the General Density Hypothesis. When  $G$  is Archimedean of high rank,  $\{\pi \in \Pi(G)_{\text{sph}} : p(\pi) > 2\}$  is not necessarily precompact, so we associate to a spherical  $(\pi, V) \in \Pi(G)$  a number  $\lambda(\pi) \in \mathbb{R}_{\geq 0}$ , which is the eigenvalue of the Casimir Operator on the  $K$ -invariant vectors of  $\pi$ , and define:

**Definition 1.2.** The sequence  $\{\Gamma_N\}$  of cocompact lattices satisfies the *Spherical Density Hypothesis* if:

- In the  $p$ -adic or rank 1 case, for every  $\epsilon > 0$  there exists  $C_\epsilon$  such that for every  $N \geq 1$ ,  $p > 2$ ,

$$M\left(\pi \in \Pi(G)_{\text{sph}}, \Gamma_N, p\right) \leq C_\epsilon [\Gamma_1 : \Gamma_N]^{2/p + \epsilon}.$$

- In the general Archimedean case, there exists  $L > 0$  large enough, such that for every  $\epsilon > 0$  there exists  $C_\epsilon$  such that for every  $\lambda \geq 0$ ,  $N \geq 1$ ,  $p > 2$ ,

$$M\left(\left\{\pi \in \Pi(G)_{\text{sph}} : \lambda(\pi) \leq \lambda\right\}, \Gamma_N, p\right) \leq C_\epsilon (1 + \lambda)^L [\Gamma_1 : \Gamma_N]^{2/p+\epsilon}.$$

In the second case, the Spherical Density Hypothesis does not a priori follows from the General Density Hypothesis.

To state our main geometric definition, we need to set some more notations. We use the standard  $O, \Theta, o$  notations, where for example  $f(N, \epsilon) = O_\epsilon(g(N, \epsilon))$  says that for every  $\epsilon$  there exists  $C$  depending only on  $\epsilon$  and the group  $G$  such that  $f(N, \epsilon) \leq Cg(N, \epsilon)$  for  $N$  large enough. The notation  $f(N, \epsilon) \ll_\epsilon g(N, \epsilon)$  is the same as  $f(N, \epsilon) = O_\epsilon(g(N, \epsilon))$  and  $f(N, \epsilon) \asymp_\epsilon g(N, \epsilon)$  is the same as  $f(N, \epsilon) \ll_\epsilon g(N, \epsilon)$  and  $g(N, \epsilon) \ll_\epsilon f(N, \epsilon)$ .

We fix a Cartan decomposition  $G = KA_+K$  and an Iwasawa decomposition  $G = KP$ . Let  $\delta(\tilde{p})$  be the left modular character of  $P$  (see Section 4 for more details).

We define a length on  $G$  by first letting for  $a \in A_+$ ,  $l(a) = \log_q \delta(a)$ , where  $q$  is equal to  $e$  in the Archimedean case and to the size of the quotient field of  $k$  otherwise. Then we extend  $l : G \rightarrow \mathbb{R}_{\geq 0}$  using the Cartan decomposition, i.e.,  $l(k_1 a k_2) = l(a)$ . Finally, we define a metric on  $G/K$  by  $d(x, y) = l(x^{-1}y)$ .

The Weak Injective Radius Property is based on the lattice point counting approach of Sarnak and Xue ([53, Conjecture 2]). Given an element  $y = \Gamma_N y \in \Gamma_N \backslash \Gamma_1$ , we denote

$$\mathbf{N}(\Gamma_N, d_0, y) = \#\{\gamma \in \Gamma_N : l(y^{-1}\gamma y) \leq d_0\}.$$

**Definition 1.3.** The sequence  $\{\Gamma_N\}$  satisfies the *Weak Injective Radius Property* if for every  $0 \leq d_0 \leq 2 \log_q([\Gamma : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{y \in \Gamma_N \backslash \Gamma_1} \mathbf{N}(\Gamma_N, d_0, y) \ll_\epsilon [\Gamma_1 : \Gamma_N]^\epsilon q^{d_0(1/2+\epsilon)}.$$

This definition is somewhat different than [53, Conjecture 2]. For rank 1, it is slightly weaker (see Proposition 5.3), while for higher rank we use a different length. In this from, the Weak Injective Radius Property follows from the Spherical Density Hypothesis — see Theorem 1.6 below.

We can now state our intended application. First, we say that a sequence of lattices  $\{\Gamma_N\}$  has a Spectral Gap if there exists  $p_0 < \infty$  such that  $p(\pi) \leq p_0$  for every non-trivial spherical  $\pi \in \Pi(G)$  weakly contained in  $L^2(\Gamma_N \backslash G)$ . This definition captures the non-uniform case as well, and in the cocompact case we may replace "weakly contained" by  $m(\pi, \Gamma) > 0$ .

We look at the natural action  $\pi_N : \Gamma_1 \rightarrow \text{Aut}(\Gamma_N \backslash \Gamma_1)$ , defined by  $\pi_N(\gamma)(\Gamma_N \gamma') = \Gamma_N \gamma' \gamma^{-1}$ . Given  $x, y \in \Gamma_N \backslash \Gamma_1$ , we look for a "small" element  $\gamma \in \Gamma_1$  such that  $\pi_N(\gamma)x = y$ . A very general way of measuring how small is an element is by the Cartan Decomposition,  $G = KA_+K$ . For  $\gamma \in \Gamma \subset G$  we let  $a_\gamma$  be the element in  $A_+$  in the Cartan decomposition of  $\gamma$ . We also fix some norm  $\|\cdot\|_a$  on the underlying coroot space of  $A_+$ . By [17], the number of  $\gamma \in \Gamma_1$  with  $\|a_\gamma - a\|_a < \delta$  is  $\asymp_{\Gamma_1} q^{l(a)}$ . Therefore, the following definition is optimal.

**Definition 1.4.** The sequence  $\{\Gamma_N\}$  has the *Optimal Lifting Property* if for every  $\epsilon > 0$ , for every  $a \in A_+$  with  $l(a) \geq (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])$ ,

$$\#\left\{(x, y) \in (\Gamma_N \backslash \Gamma_1)^2 : \exists \gamma \in \Gamma_1 \text{ s.t. } \pi_N(\gamma)x = y, \|a_\gamma - a\|_{\mathfrak{a}} < \epsilon \|a\|_{\mathfrak{a}}\right\} = (1 - o_{\epsilon, \Gamma_1}(1)) [\Gamma_1 : \Gamma_N]^2.$$

Conjecturally, every sequence of congruence subgroups of an arithmetic lattice in an almost-simple and simply connected Lie group satisfies the Optimal Lifting Property. We refer to Conjecture 2.4 for a full statement.

The following two theorems show that our two main properties imply the Optimal Lifting Property.

**Theorem 1.5.** *Let  $\{\Gamma_N\}$  be a sequence of lattices having a Spectral Gap and satisfying the Weak Injective Radius Property. Then the sequence  $\{\Gamma_N\}$  has the Optimal Lifting Property.*

**Theorem 1.6.** *If the sequence  $\{\Gamma_N\}$  of cocompact lattices satisfies the Spherical Density Hypothesis, then  $\{\Gamma_N\}$  satisfies the Weak Injective Radius Property. Therefore, assuming also Spectral Gap, the Spherical Density Hypothesis implies the Optimal Lifting Property.*

The definition of Optimal Lifting is based on the main result in an influential letter of Sarnak ([52]), who proved the Optimal Lifting Property for principal congruence subgroups of  $SL_2(\mathbb{Z})$  by utilizing a version of the Spherical Density Hypothesis proved by Huxley ([33]).

In the cocompact case, one may relate the Optimal Lifting Property to almost-diameter of the quotient space as follows. If we give  $X_N = \Gamma_N \backslash G/K$  the quotient metric  $d$ , the Optimal Lifting Property implies that the distances between points of  $X_N$  are concentrated at the optimal location  $\log_q(\mu(X_N))$ , i.e. for every  $\epsilon > 0$ ,

$$\mu\left(\{(x, y) \in X_N \times X_N : d(x, y) < (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])\}\right) = (1 - o_\epsilon(1)) \mu^2(X_N).$$

This concentration of distances phenomena was proven for Ramanujan graphs by Sardari ([50]) and Lubetzky-Peres ([44]), who also related it to the cutoff phenomena. In higher dimensions, similar results for Ramanujan complexes appear in [34, 43]. Theorem 1.5 implies that one may get results which are almost as strong, as long as we assume only the far weaker Spherical Density Hypothesis or the Injective Radius Property.

The Weak Injective Radius Property is intended as the arithmetic, or geometric, input to our approach, and we discuss it further in Section 2. There are a few cases where it is known, most notably, following the work of Sarnak and Xue, for principal congruence subgroups of arithmetic lattices in  $SL_2(\mathbb{R})$  and  $SL_2(\mathbb{C})$  (see Subsection 2.4). In a companion paper by the second named author and Hagai Lavner, the Weak Injective Radius Property is proven for some non-principal congruence subgroups of  $SL_3(\mathbb{Z})$  (see Subsection 2.5), and it is closely related to the works [5, 4]. If we allow ourselves to relax the definition and add a parameter  $0 < \alpha \leq 1$  to it (see Subsection 2.2), it is quite straightforward to show that *principal* congruence subgroups of arithmetic groups satisfy the Injective Radius Property with some explicit parameter  $\alpha > 0$  (see Corollary 2.2). As a matter of fact, recent results in [1, 20] show that every sequence of

congruence subgroups satisfy the Weak Injective Radius property with some explicit parameter  $\alpha > 0$  (see Theorem 2.3). However, one must have  $\alpha = 1$  for the Optimal Lifting application.

**1.1. The Relations Between the Different Properties.** We already stated that the Spherical Density Hypothesis implies the Weak Injective Radius Property. For the deduction of spectral results from the Weak Injective Radius Property, we have partial results in the Archimedean case and full results in the  $p$ -adic case.

We believe that the following is true:

**Conjecture 1.7.** *The Weak Injective Radius Property implies both the General Density Hypothesis and the Spherical Density Hypothesis.*

In the  $p$ -adic case, one can choose sets covering  $\Pi(G)$  as follows. For a compact open subgroup  $K'$  of  $G$ , let  $\Pi(G)_{K'\text{-sph}}$  be the set of isomorphism classes of irreducible unitary representations with  $K'$ -invariant vectors. If we have a sequence  $\{K'_m\}$  of arbitrary small compact open subgroups (i.e., they generate the topology of  $G$  near the identity), then we have

$$\Pi(G) = \bigcup_m \Pi(G)_{K'_m\text{-sph}}.$$

We can now state:

**Theorem 1.8.** *Conjecture 1.7 is true when  $G$  is non-Archimedean. More Precisely, there exists a sequence  $\{K'_m\}$  consisting of arbitrarily small compact open subgroups of  $G$ , such that if the Weak Injective Radius Property holds for a sequence of cocompact lattices  $\{\Gamma_N\}$ , then for every  $N \geq 1$ ,  $m \geq 1$ ,  $p > 2$ ,  $\epsilon > 0$ ,*

$$M\left(\Pi(G)_{K'_m\text{-sph}}, \Gamma_N, p\right) \ll_{K'_m, \epsilon} [\Gamma_1 : \Gamma_N]^{2/p+\epsilon}.$$

In the Archimedean case, we do not know even whether the Weak Injective Radius Property implies the Spherical Density Hypothesis. However, for rank 1 it was essentially proven in [53] (see the remark after the statement of Theorem 3 in [53]):

**Theorem 1.9.** *If  $G$  is of rank one and  $\{\Gamma_N\}$  is a sequence of cocompact lattices, then the Weak Injective Radius Property implies the Spherical Density Hypothesis.*

For general representations in the Archimedean case, we have the following theorem. For rank 1 it was proven in [53, Theorem 3].

**Theorem 1.10.** *If the sequence  $\{\Gamma_N\}$  of cocompact lattices satisfies the Weak Injective Radius Property, then  $\{\Gamma_N\}$  satisfies the Pointwise Multiplicity Hypothesis.*

Figure 1.1 summarizes the different relations between our main properties for a sequence of cocompact lattices.

Let us end this introduction by stating some open problems this work leads to.




FIGURE 1.1. The relations between our main properties for a sequence of cocompact lattices.

The main open problem is to prove either the Spherical Density Hypothesis or the Injective Radius Property for new cases, which would lead to a proof of the Optimal Lifting Property. See Conjecture 2.4 for a general conjecture for the Archimedean case. Very few cases of this conjecture are known for groups of rank greater than 1. We remark that it seems that the problem is harder for principal congruence subgroups, and easier for group that are far from being normal, such as  $\Gamma_0(N)$  of  $SL_m(\mathbb{Z})$  (see Subsection 2.5). See, for example, the *Density Amplification* phenomena for graphs in [26].

In the more technical side, a main problem this work does not resolve is Conjecture 1.7, which would show, in particular, that the Spherical Density Hypothesis and the Injective Radius Property are indeed equivalent, also for the Archimedean high rank case. It relates in particular to the understanding of uniform lower bound on matrix coefficients, see, e.g., Conjecture 3.12.

Finally, we only discuss multiplicities for cocompact lattices. We strongly believe that the Weak Injective Radius Property has spectral implications for non-uniform lattices as well, e.g., for bounds of multiplicities of representations in the discrete spectrum. This problem is strongly related to the concentration of  $L^2$ -mass of non-tempered automorphic functions away from the cusp, in a uniform way. In hyperbolic spaces this problem is essentially solved, even for some discrete groups that are not lattices, thanks to the work of Gamburd on hyperbolic surfaces ([21]) and the work of Magee on general hyperbolic spaces ([48]). Therefore, Theorem 1.9 can be generalized to such cases.

**Structure of the Article.** In Section 2 we give some applications of our results, and state some open problems.

In Section 3 we present the main ideas behind the proofs.

In Section 4 we collect various results, mainly from representation theory. In particular, we discuss upper bounds for matrix coefficients, which are well understood. Using those upper bounds, we prove the essential Convolution Lemma 4.18.

In Section 5 we discuss the Weak Injective Radius Property Property and the spectral results it implies. We prove Proposition 3.9, which reduces Theorem 1.10, Theorem 1.9, and Theorem 1.8 to finding some explicit and strict lower bounds on matrix coefficients.

In Section 6 we prove Theorem 1.5.

In Section 7 we discuss the Spherical Density Hypothesis and the results it implies. We prove Theorem 1.6 and prove Theorem 7.4, which is a version of Theorem 1.5 which assumes the Spherical Density Hypothesis and has stronger implications.

In Section 8 we discuss Bernstein description of the Hecke algebra in the non-Archimedean case, and prove Theorem 3.11, which implies Theorem 1.8 together with Proposition 3.9

In Section 9 we discuss the theory of leading coefficients and prove Theorem 3.10, which implies Theorem 1.10 together with Proposition 3.9.

**Acknowledgments.** The authors are thankful to Amos Nevo and Peter Sarnak for their support of this project.

The first author is supported by the SNF grant 200020-169106 at ETH Zurich. This work is part of the Ph.D. thesis of the second author at the Hebrew University of Jerusalem, under the guidance of Prof. Alexander Lubotzky, and is supported by the ERC grant 692854.

## 2. APPLICATIONS AND OPEN PROBLEMS

**2.1. Ramanujan Graphs and Complexes.** We shortly note that the results of this paper, and in particular Theorem 1.5 (or the stronger Theorem 7.4) apply to Ramanujan Complexes, by which we mean here the situation when  $G$  is  $p$ -adic,  $\Gamma_1$  is cocompact, and no non-tempered and non-trivial spherical representation appears in the decomposition of  $L^2(\Gamma_N \backslash G)$ . In this case, the sequence  $\{\Gamma_N\}$  obviously satisfies the Spherical Density Hypothesis. The Ramanujan complexes themselves are the quotients of the Bruhat-Tits buildings of  $G$  by  $\Gamma_N$ . Such complexes were constructed (with the same definition) by Lubotzky, Samuels and Vishne (see [47, 46]). Similar results to Theorem 7.4 appear in [34, 43, 12].

Note that the definition we use is not the same as in the more modern approach to Ramanujan complexes, where one considers not only spherical representation but also representations with a non-trivial vector fixed by the Iwahori subgroup (See [45, Subsection 2.3] and the references within).

The standard way of proving that a complex is a Ramanujan complex is to use proved cases of the Langlands program (e.g. [41]), and eventually to apply Deligne's proof of the Weil conjectures ([15]). This approach has obvious limitations, and in particular it seems that one must assume that  $G$  is  $p$ -adic for it to succeed. We refer to [19] for some recent work on this subject, and to an ongoing and yet unpublished work of Shai Evra.

We will avoid diving deeper into this subject, as it is based on spectral input, unlike our approach which is geometric.

**2.2. Adding a Parameter to the Properties.** It is useful to add a parameter  $0 < \alpha \leq 1$  to the properties, with  $\alpha = 1$  being equivalent to the property without the parameter.

**Definition.** We say that the sequence  $\{\Gamma_N\}$  satisfies the *Weak Injective Radius Property with parameter  $\alpha$* , if for every  $0 \leq d_0 \leq 2\alpha \log_q([\Gamma : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{y \in \Gamma_N \setminus \Gamma_1} \mathbf{N}(\Gamma_N, d_0, y) \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon} q^{d_0(1/2+\epsilon)}.$$

**Definition.** We say that the sequence  $\{\Gamma_N\}$  satisfies the General Density Hypothesis with parameter  $\alpha$  if for every precompact subset  $A \subset \Pi(G)$  and every  $\epsilon > 0$ ,  $N$  and  $p \geq 2$ ,

$$M(A, \Gamma, p) \ll_{\epsilon, A} [\Gamma_1 : \Gamma_N]^{1-\alpha(1-2/p)+\epsilon}.$$

One may similarly define the Spherical Density Hypothesis with parameter  $\alpha$  and the Pointwise Density Hypothesis with parameter  $\alpha$ , by changing the exponent  $2/p$  to  $1 - \alpha(1 - 2/p)$ .

Figure 1.1 remains true if we replace the properties with their parameterized version, for the same parameter  $0 < \alpha \leq 1$ , with the exception of the derivation of the Optimal Lifting Property from the Spherical Density Hypothesis – we need  $\alpha = 1$  for this derivation to work.

In the work itself we work with the parameterized version of the properties.

**2.3. Congruence Subgroups of Arithmetic Groups.** Let  $\Gamma_1$  be an arithmetic lattice in a semisimple non-compact Lie group  $G$ . Following [53], by arithmetic we mean that  $G$  is defined over  $\mathbb{Q}$ ,  $\Gamma_1 \subset G(\mathbb{Q})$ , there is a  $\mathbb{Q}$ -embedding  $\rho : G \rightarrow GL_n$  and  $\Gamma_1$  is commensurable with  $\rho^{-1}(GL_n(\mathbb{Z}))$ .

In this case we may define a sequence of principle congruence subgroups  $\{\Gamma_N\}$  of  $\Gamma_1$  by letting

$$\Gamma_N = \Gamma_1(N) = \Gamma_1 \cap \rho^{-1}(\{A \in GL_n(\mathbb{Z}) : A \equiv I \pmod{N}\}).$$

It is a well known fact that such subgroups have (up to a finite index) injective radius which is logarithmic in the index (e.g., [53, Lemma 1] or [28, Proposition 16]).

Let us shortly give the argument — we assume by moving to a finite index in  $\Gamma_1$  that  $\Gamma_1 \subset \rho^{-1}(GL_n(\mathbb{Z}))$ . In such case  $\Gamma_N$  is normal in  $\Gamma_1$  and it is left to verify that for every  $d_0 \leq 2\alpha \ln([\Gamma_1 : \Gamma_N])$  and  $\epsilon > 0$ ,

$$\#\{\gamma \in \Gamma_N : l(\gamma) \leq d_0\} \ll_{\epsilon} e^{d_0(1/2+\epsilon)}.$$

We note that there exists a constant  $C$  such that for  $g \in G$  outside a compact set the length  $l(g)$  we defined satisfies

$$C^{-1} \ln(\|\rho(g)\|) \leq l(g)$$

where  $\|\cdot\| : GL_n(\mathbb{R}) \rightarrow \mathbb{R}_{\geq 0}$  is the maximal absolute value of an entry of the matrix. Since each element of  $\rho(\Gamma_N)$  which is not the identity has an entry of size  $N$ , We deduce that for every  $N$  large enough,

$$\{\gamma \in \Gamma_N : l(\gamma) < C \ln(N)\} = \{I\}.$$



On the other hand, obviously

$$[\Gamma_1 : \Gamma_N] \ll N^{n^2}$$

(and actually  $[\Gamma_1 : \Gamma_N] \ll_\epsilon N^{\dim G + \epsilon}$ ).

Combining the different estimates, for  $d_0 < 2\frac{C}{2n^2} \ln(\Gamma_1 : \Gamma_N) \leq 2\frac{C}{2n^2} \ln(N^{n^2}) = C \ln(N)$ , it holds that for  $N$  large enough,

$$\#\{\gamma \in \Gamma_N : l(\gamma) \leq d_0\} = 1.$$

Therefore, the Weak Injective Radius Property is satisfied with parameter  $\alpha = \frac{C}{2n^2}$ .

*Remark 2.1.* Pushing this argument further, we may take the parameter to be  $\alpha = \frac{C}{\dim G}$ . For example, for principal congruence subgroups of  $SL_2(\mathbb{Z})$  one gets this way the parameter  $\alpha = \frac{2}{3}$ , while  $\alpha = 1$  can be reached by a better analysis, see Subsection 2.4 below.

As a direct application of this fact we have:

**Corollary 2.2.** *Let  $G$  be Archimedean, let  $\Gamma_1$  be an arithmetic lattice and let  $\{\Gamma_N\}$  be the sequence of principle congruence subgroups of  $\Gamma_1$ . Then the sequence  $\{\Gamma_N\}$  satisfies the Weak Injective Radius Property with parameter  $\alpha = \alpha(\Gamma_1)$ .*

As a matter of fact this corollary can be easily be extended to principal congruence subgroups of  $S$ -arithmetic groups, once those are properly defined.

For arbitrary congruence subgroups, let us recall some of the results of [20] (the same result can be deduced from [1, Theorem 1.11 / Theorem 5.6]).

First, for  $\gamma \in \Gamma_1$ , we denote

$$c_{\Gamma_N}(\gamma) = |\{y \in \Gamma_N \setminus \Gamma_1 : y\gamma y^{-1} \in \Gamma_N\}|,$$

which is the number of fixed points of the right action of  $\gamma$  on  $\Gamma_N \setminus \Gamma_1$ . Then the Weak Injective Radius Property with parameter  $\alpha$  is equivalent to the fact that for every  $0 \leq d_0 \leq 2\alpha \log_q([\Gamma : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{l(\gamma) \leq d_0} c_{\Gamma_N}(\gamma) \ll_\epsilon [\Gamma_1 : \Gamma_N]^\epsilon q^{d_0(1/2+\epsilon)}.$$

A congruence subgroup is a subgroup of one of the groups  $\Gamma_1(N)$  as above.

**Theorem 2.3** (Following [20, Corollary 5.9]). *Let  $G$  be Archimedean, semisimple, almost-simple and simply connected, let  $\Gamma_1$  be an arithmetic lattice and let  $\{\Gamma_N\}$  be a sequence of congruence subgroups of  $\Gamma_1$ . Then there exists constants  $\mu > 0, c > 0$  depending only on  $\Gamma_1$  such that for every  $d_0 > 0, N \geq 1$  it holds that*

$$(2.1) \quad \frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{\gamma \in \Gamma_1, l(\gamma) \leq d_0} c_{\Gamma_N}(\gamma) \ll 1 + e^{cd_0} [\Gamma_1 : \Gamma_N]^{-\mu}.$$

*In particular, the injective radius property holds with parameter  $\alpha = \frac{\mu}{c}$ .*

*Proof.* By [20, Corollary 5.9], there exists  $c'$  such that for every  $\gamma \in \Gamma_1$  which does not belong to a proper normal subgroup of  $G$  it holds that

$$\frac{1}{[\Gamma_1 : \Gamma_N]} c_{\Gamma_N}(\gamma) \ll e^{c'l(\gamma)} [\Gamma_1 : \Gamma_N]^{-\mu}.$$

We remark that the dependence on  $\gamma$  in [20] is different, but it is obvious that  $e^{cl(\gamma)}$  for  $c$  large enough is an upper bound on it. By our assumptions on  $G$ , the number of  $\gamma \in \Gamma_1$  belonging to a proper normal subgroup is bounded.

By summing over all  $\gamma \in \Gamma_1$  with  $l(\gamma) \leq d_0$  we get

$$\begin{aligned} \frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{\gamma \in \Gamma_1, l(\gamma) \leq d_0} c_{\Gamma_N}(\gamma) &\ll \frac{[\Gamma_1 : \Gamma_N]}{[\Gamma_1 : \Gamma_N]} + e^{c'd_0} [\Gamma_1 : \Gamma_N]^{-\mu} \sum_{\gamma \in \Gamma_1, l(\gamma) \leq d_0} 1 \\ &\ll 1 + e^{(1+c')d_0} [\Gamma_1 : \Gamma_N]^{-\mu}, \end{aligned}$$

which implies Equation 2.1 with  $c = 1 + c'$ .

When  $d_0 \leq 2\frac{\mu}{c} \ln([\Gamma_1 : \Gamma_N])$ , or  $[\Gamma_1 : \Gamma_N]^\mu = e^{d_0 c/2}$  we get

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{\gamma \in \Gamma_1, l(\gamma) \leq d_0} c_{\Gamma_N}(\gamma) \ll e^{d_0/2},$$

which implies the Weak Injective Radius Property with parameter  $\alpha = \frac{\mu}{c}$ , as needed.  $\square$

In the cocompact case, Theorem 1.10 implies the Pointwise Multiplicity Hypothesis with parameter  $\alpha = \frac{\mu}{c}$ , namely

$$m(\pi, \Gamma_N) \ll_{\pi, \epsilon} [\Gamma_1 : \Gamma_N]^{1 - \frac{\mu}{c}(1 - 2/p(\pi)) + \epsilon}.$$

It should be compared with [1, Theorem 7.15], which states

$$m(\pi, \Gamma_N) \ll_{\pi} [\Gamma_1 : \Gamma_N]^{1 - \alpha(\pi)}.$$

Finally, we wish to make the following conjecture:

**Conjecture 2.4.** *Let  $G$  be Archimedean, semisimple, almost-simple and simply connected, let  $\Gamma_1$  be an arithmetic lattice and let  $\{\Gamma_N\}$  be a sequence of congruence subgroups of  $\Gamma_1$ . Then the sequence  $\{\Gamma_N\}$  satisfies the Weak Injective Radius Property with parameter  $\alpha = 1$ , and if  $\Gamma_1$  is cocompact then the sequence  $\{\Gamma_N\}$  satisfies the General Density Hypothesis with parameter  $\alpha = 1$ .*

*As a corollary, the sequence  $\{\Gamma_N\}$  has the Optimal Lifting Property.*

The conjecture generalizes Conjecture 1 and Conjecture 2 from the work of Sarnak and Xue ([53]).

Finally, a similar conjecture should also hold when  $G$  is  $p$ -adic, or in the  $S$ -arithmetic setting.

**2.4. The Work of Sarnak and Xue and its Implications.** Sarnak and Xue proved the Weak Injective Radius Property with parameter  $\alpha = 1$  for principal congruence subgroups of

cocompact arithmetic lattices in  $SL_2(\mathbb{R})$  and  $SL_2(\mathbb{C})$ , and  $\alpha = 5/6$  for principal congruence subgroups of cocompact arithmetic lattices for  $SU(2, 1)$ .

Let us explain their calculation, in a slightly more general case, which includes also lattices in  $p$ -adic groups.

We start with the injective radius property for the principal congruence subgroups of  $SL_2(\mathbb{Z})$ . In this case a simple argument says that we need to show that the number  $\mathbf{N}(T, \Gamma(N))$ , of solutions to  $ad - bc = 1$ , with  $a \equiv d \equiv 1 \pmod{N}$ ,  $b \equiv c \equiv 0 \pmod{N}$  and  $\max\{|a|, |b|, |c|, |d|\} \leq T$ , for  $T \leq N^3$ , is bounded by

$$\mathbf{N}(T, \Gamma(N)) \ll_{\epsilon} T^{1+\epsilon}.$$

This is done as follows (see also [21, Proposition 5.3]). From the congruence condition it follows that

$$a + d - 2 = -(a - 1)(d - 1) + bc \equiv 0 \pmod{N^2}.$$

One may therefore choose  $a+d$  in  $4(T/N^2 + 1)$  ways, and choose  $(a, d)$  in  $8(T/N^2 + 1)(T/N + 1)$  ways.

If  $ad \neq 1$ , from  $bc = 1 - ad$  and bounds on the divisor function, there are  $T^{\epsilon}$  ways of choosing  $bc$ . If  $ad = 1$  it is also simple to bound the number of possibilities by  $4(T/N + 1)$ . In total we get for  $T \leq N^3$

$$\mathbf{N}(T, \Gamma(N)) \ll_{\epsilon} T^{\epsilon} (T/N^2 + 1) (T/N + 1) \ll_{\epsilon} T^{1+\epsilon}.$$

The calculation above can be generalized to lattices coming from division algebras as follows. We follow [53, Section 3].

Let  $F$  be a number field and let  $p, q \in F$ , let  $A_{(F,p,q)}$  be the quaternion algebra with basis  $\{1, I, J, IJ\}$  over  $F$ , and the relations

$$I^2 = p, J^2 = q, IJ = -JI.$$

We have a norm and trace, defined by

$$N(a + bI + cJ + dIJ) = a^2 - pb^2 - qc^2 + pqd^2$$

$$\text{Tr}(a + bI + cJ + dIJ) = 2a$$

Let denote by  $v$  a place of  $F$ , which is a finite extension of  $\mathbb{Q}_p, \mathbb{R}$  or  $\mathbb{C}$ . The corresponding local field is denoted  $F_v$ .

We say that a place  $v$  of  $F$  is ramified if  $G(F_v) = \{\alpha \in A \otimes F_v : N(\alpha) = 1\} \subset M_{2 \times 2}(F_v)$  is a compact group. If a place  $v$  is unramified then

$$G(F_v) \cong SL_2(F_v).$$

We assume that the number of unramified real or complex places is 0 or 1. We let  $v_0$  be an unramified place, which is equal to the real or complex unramified place if there is one.

Let  $\mathcal{O}^{(v_0)} = \{a \in F : \forall v \neq v_0 \text{ finite, } |a|_v \leq 1\}$ . Notice that if  $v_0$  is real or complex then  $\mathcal{O}^{(v_0)} = \mathcal{O}$  is the ring of integers of  $F$ .

In such case, it is well known that

$$\Gamma = G\left(\mathcal{O}^{(v_0)}\right) \subset G(F_v) \cong SL_2(F_v)$$

is a lattice, which is cocompact if there are ramified places. Now, for an ideal  $\mathcal{I} \subset \mathcal{O}^{(v_0)}$ , let

$$\Gamma(\mathcal{I}) = \{\gamma \in \Gamma : \gamma \equiv 1 \pmod{\mathcal{I}}\}.$$

This is a normal subgroup in  $\Gamma$ , whose index is  $[\Gamma : \Gamma(\mathcal{I})] \asymp N^3(\mathcal{I})$ , where  $N(\mathcal{I})$  is the number of elements in  $\mathcal{O}^{(v_0)}/\mathcal{I}$ .

**Proposition 2.5.** *For  $T \leq N^3(\mathcal{I})$ , it holds that*

$$\left\{ \gamma = a + bI + cJ + dIJ \in \Gamma(\mathcal{I}) : \max\{|a|_{v_0}, |b|_{v_0}, |c|_{v_0}, |d|_{v_0}\} \leq T \right\} \ll_{\epsilon} T^{1+\epsilon}.$$

*Therefore, the subgroups  $\Gamma(\mathcal{I})$  satisfy the Weak Injective Radius Property with parameter  $\alpha = 1$ .*

We will not give a full proof of this estimate, and refer to the work of Sarnak and Xue. They prove by calculation that resembles the calculations for principal congruence subgroups of  $SL_2(\mathbb{Z})$  given above, that for every  $T$  the left hand side is bounded by

$$\ll_{\epsilon} \left( \frac{T^{2+\epsilon}}{N^3(\mathcal{I})} + \frac{T^{1+\epsilon}}{N(\mathcal{I})} + 1 \right),$$

which gives the required bound in our range.

*Remark 2.6.* Consider the quadratic form

$$H(x_1, x_2, x_3, x_4) = x_1^2 - px_2^2 - qx_3^2 + pqx_4^2.$$

Consider the number of solutions  $\mathbf{N}_H(T)$  to

$$H(x_1, x_2, x_3, x_4) = 1$$

with  $x_1, x_2, x_3, x_4 \in \mathcal{O}^{(v_0)}$ , subject to the conditions:

- For every  $v \neq v_0$  Archimedean it holds that  $|x_i|_v \leq C$  for some fixed  $C$  (this holds automatically by our ramification assumptions).
- $x_1 - 1 \equiv x_2 \equiv x_3 \equiv x_4 \pmod{\mathcal{I}}$ .
- $\max\{|x_1|_{v_0}, |x_2|_{v_0}, |x_3|_{v_0}, |x_4|_{v_0}\} \leq T$ .

The bound we seek is that for  $T \leq N^3(\mathcal{I})$ ,  $\mathbf{N}_H(T) \ll_{\epsilon} T^{1+\epsilon}$ .

The proof in [53, Section 3] actually works for every quadratic form with 4 variables, satisfying those 3 conditions.

One should compare this with the closely related lower bounds given by Sardari in [51], when  $K = \mathbb{Q}$ .

2.5. **On the Congruence Subgroups  $\Gamma_0(N)$ .** Consider the subgroup

$$\Gamma_0(N) = \left\{ \begin{pmatrix} * & * & * \\ * & * & * \\ a & b & * \end{pmatrix} \in SL_3(\mathbb{Z}) : a \equiv b \equiv 0 \pmod{N} \right\} \subset \Gamma_1 = SL_3(\mathbb{Z})$$

In a companion paper by the second named author and Hagai Lavner ([36]), the following is proven:

**Theorem 2.7.** *The sequence of lattices  $\Gamma_0(N)$ , for  $N$  prime, has the Weak Injective Radius Property (with parameter 1).*

*As a result, this sequence of lattices satisfies the Optimal Lifting Property.*

We refer to [36] for an interpretation of this result in terms of the action of  $SL_3(\mathbb{Z})$  on the projective plane over the field with  $N$  elements.

The work [36] is strongly influenced by a deep work of Blomer, Buttcane and Maga on the same subgroup ([5]). This result was recently generalized by Blomer to general  $SL_M(\mathbb{Z})$ , where the subgroup  $\Gamma_0(N)$  is similarly defined to be the set of matrices with the entries in the last row, except for the  $(M, M)$  entry, equal to 0 modulo  $N$ .

**Theorem 2.8** ([5, Theorem 4] for  $M = 3$ . [4] for general  $M$ ). *For  $\pi \in \Pi(SL_M(\mathbb{R}))_{sph}$ , let  $m_{cusp}(\pi, \Gamma_0(N))$  be the multiplicity of  $\pi$  in the cuspidal part of  $L^2(\Gamma_0(N) \backslash SL_M(\mathbb{R}))$ . Then for every compact  $A \subset \Pi(SL_M(\mathbb{R}))_{sph}$ , it holds that for every  $N$  prime,  $p > 2$ ,  $\epsilon > 0$ ,*

$$\sum_{\pi \in A, p(\pi) > 0} m_{cusp}(\pi, \Gamma_0(N)) \ll_{A, \epsilon} [\Gamma : \Gamma_0(N)]^{1-2(1-2/p)+\epsilon}.$$

Notice that it is very similar to a Density Hypothesis with parameter 2. However, the results of this theorem are slightly different than the Spherical Density Hypothesis in a number of ways: first,  $SL_3(\mathbb{Z})$  is not cocompact so our discussion does not apply to it. In particular, we have to deal with the continuous spectrum if we wish to deduce the Optimal Lifting Property. Secondly and less crucially, the dependence on the subset  $A$  is not explicit, as needed in the definition of Spherical Density and its applications for geometric phenomena.

2.6. **The Weak Injective Radius of Principal Congruence Subgroups of  $SL_n(\mathbb{Z})$ .** The weak injective radius conjecture for  $SL_n$  implies in particular that conjecturally

$$\#\{\gamma \in SL_n(\mathbb{Z}) : \gamma \equiv I \pmod{N}, \|\gamma\| \leq T\} \ll T^\epsilon N^\epsilon \left( \frac{T^{n^2-n}}{N^{n^2-1}} + T^{(n^2-n)/2} \right),$$

where  $\|\cdot\|$  is the maximal absolute value of an entry. One may also try to improve this estimate, in particular the “error” part  $T^{(n^2-n)/2}$ .

One of the results of [38], which is also cited in [53], states that

$$\#\{\gamma \in SL_n(\mathbb{Z}) : \gamma \equiv I \pmod{N}, \|\gamma\| \leq T\} \ll \frac{T^{n^2-n}}{N^{n^2-1}} \log T + 1.$$

As was pointed to us by Sarnak, the proof contains an error. As a matter of fact this naive estimate is actually *false*, for simple reasons. For example, in  $SL_2$ ,

$$\# \left\{ \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \in SL_2(\mathbb{Z}) : \gamma \equiv I \pmod{N}, \|\gamma\| \leq T \right\} \asymp T/N + 1.$$

and this is larger than  $\frac{T^2}{N^3} \log T$  in the range  $N(1+\epsilon) < T < (1-\epsilon)N^2$ . The same argument works for every  $n$ , but there are  $n(n-1)/2$  entries we can use. Therefore we have the lower bound:

$$\# \{ \gamma \in SL_n(\mathbb{Z}) : \gamma \equiv I \pmod{N}, \|\gamma\| \leq T \} \gg \left( \frac{T^{n^2-n}}{N^{n^2-1}} + \left( \frac{T}{N} \right)^{(n^2-n)/2} + 1 \right).$$

Up to  $T^\epsilon$ , this is also the upper bound for  $n=2$ . We conjecture that up to  $T^\epsilon$  this is also the upper bound for larger values of  $n$ .

### 3. MAIN IDEAS OF THE PROOFS

In this section we discuss some of the ideas of the proofs and the technical problems they lead to.

All the proofs are based on a reduction of the geometric properties into spectral ones. We first restate the Weak Injective Radius Property in terms of traces of operators.

**Definition 3.1.** For  $d_0 \in \mathbb{R}_{\geq 0}$ , let  $\chi_{d_0} \in C_c^\infty(K \backslash G / K)$  be a function equal to 1 for  $l(g) \leq d_0$ , equal to 0 for  $l(g) \geq d_0 + 1$  and for  $d_0 \leq l(g) \leq d_0 + 1$  it is defined between 0 and 1 so that  $\chi_{d_0} \in C_c^\infty(K \backslash G / K)$ .

Let  $\psi_{d_0}(g) \in C_c^\infty(K \backslash G / K)$  defined as  $\psi_{d_0}(g) = q^{(d_0-l(g))/2} \cdot \chi_{d_0}(g)$  for  $l(g) \geq 1$  and for  $0 \leq l(g) \leq 1$  it is defined between  $q^{(d_0-1)/2}$  and  $q^{d_0/2}$  so that  $\psi_{d_0}(g) \in C_c^\infty(K \backslash G / K)$ .

As  $\chi_{d_0}, \psi_{d_0} \in C_c^\infty(K \backslash G / K)$ , they act naturally on  $L^2(\Gamma_N \backslash G)$ , and moreover, have a trace ([23, Chapter 1], see also Subsection 4.6). The trivial eigenvalues of  $\chi_{d_0}, \psi_{d_0}$  satisfy  $q^{d_0} \ll \int_G \chi_{d_0}(g) dg \ll_\epsilon q^{d_0(1+\epsilon)}$  and  $q^{d_0} \ll \int_G \psi_{d_0}(g) dg \ll_\epsilon q^{d_0(1+\epsilon)}$ . Since  $\chi_{d_0}, \psi_{d_0} \in C_c(K \backslash G / K)$ , their action is actually on  $L^2(\Gamma_N \backslash G / K)$ .

The reason that we look at  $\psi_{d_0}$ , and one of the main reasons we use the length  $l(g)$ , is the following Convolution Lemma, which replaces the rank 1 case in [53, Lemma 3.1]. The lemma says that  $\psi_{d_0}$  provides an approximated upper bound for the convolution of  $\chi_{d_0/2}$  with itself:

**Lemma 3.2** (See Lemma 4.18). *It holds that  $c_{d_0} = \chi_{d_0} * \chi_{d_0} \in C_c^\infty(K \backslash G / K)$ , and for every  $g \in G$  and  $\epsilon > 0$ , satisfies the inequality*

$$c_{d_0}(g) \ll_\epsilon q^{d_0\epsilon} \psi_{2d_0}(g).$$

Using the pre-trace formula (see Subsection 4.7), we get:

**Proposition 3.3.** *The following are equivalent:*

- (1) *The sequence  $\{\Gamma_N\}$  satisfies the Weak Injective Radius Property with parameter  $\alpha$ .*

(2) For every  $0 \leq d_0 \leq 2\alpha \log_q([\Gamma_1 : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$(3.1) \quad \text{tr} \chi_{d_0} |_{L^2(\Gamma_N \backslash G)} \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{d_0(1/2+\epsilon)}.$$

(3) For every  $\epsilon > 0$ , for every  $h \in C_c(G)$  satisfying  $h(g) \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon} \psi_{2\alpha \log_q([\Gamma_1 : \Gamma_N])}(g)$ , it holds that

$$(3.2) \quad \text{tr} h |_{L^2(\Gamma_N \backslash G)} \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\alpha+\epsilon}.$$

The proposition allows one to prove the Weak Injective Radius property using the trace formula and Equation 3.1. It also shows that allows proving various multiplicity results using Equation 3.2.

By combining the Convolution Lemma 3.2, Spectral Gap, and a version of Equation 3.2, we prove Theorem 1.5. See Section 6.

The direction “spectral to geometrical” uses in a careful way Harish-Chandra’s upper bounds on spherical functions, and its generalization to arbitrary matrix coefficients of representations. Our analysis is closely related to the work of Ghosh, Gorodnik and Nevo on Diophantine exponents ([24, 25] and the reference therein). Very generally, the main difference between our analysis and theirs is that we assume Density, while they assume Spectral Gap. The new idea is to note that one applies the spectral estimates to characteristic functions of “small sets”, and if there are few “bad eigenvectors” they correlate poorly with such functions.

Let us state some explicit results. Let  $\Xi(g) = \int \delta(gk)^{-1/2} dk$  be Harish-Chandra’s function of  $G$ . This function satisfies  $|\Xi(g)| \ll_{\epsilon} q^{-l(g)(1/2+\epsilon)}$  (which is another motivation for the definition of  $l$ ). Now, combining the two theorems of [13] we get:

**Theorem 3.4** ([13]). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq 2$ , and let  $v_1, v_2 \in V$  be two  $K$ -finite vectors, such that  $\dim \text{span} K v_1 = d_1$ ,  $\dim \text{span} K v_2 = d_2$ . Then*

$$|\langle v_1, g v_2 \rangle| \leq \sqrt{d_1 d_2} \|v_1\| \|v_2\| \Xi(g).$$

This theorem is not enough however for our uses since we wish to use results for  $p(\pi)$  arbitrary. While [13] give results for arbitrary  $\pi$  and  $p$ , but they are not precise enough for our uses.

Let us state here a general theorem which generalizes the above. For  $2 \leq p \leq \infty$ , define  $\Xi_p(g) = \int \delta(gk)^{-1/p} dk$  as the  $p$ -th version Harish-Chandra’s function of  $G$ . This function satisfies  $\Xi_p(g) \ll \Xi(g) q^{-l(g)(1/p-1/2)} \ll_{\epsilon} q^{-l(g)(1/p+\epsilon)}$ . Then we have:

**Theorem 3.5** (See Theorem 4.10). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq p$ ,  $p \geq 2$ , and let  $v_1, v_2 \in V$  be two  $K$ -finite vectors, such that  $\dim \text{span} K v_1 = d_1$ ,  $\dim \text{span} K v_2 = d_2$ . Then*

$$|\langle v_1, g v_2 \rangle| \leq \sqrt{d_1 d_2} \|v_1\| \|v_2\| \Xi_p(g).$$

We further discuss such bounds in Subsection 4.2. Using Theorem 3.5 one may deduce various *upper bounds* on matrix coefficients, norms of operators and traces of operators. In particular, a useful bound is:

**Theorem 3.6** (See Corollary 4.14). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) = p$ ,  $p \geq 2$ . Then for every  $d_0 > 0$  and  $\epsilon > 0$*

$$\|\pi(\chi_{d_0})\| \ll_{\epsilon} q^{d_0(1-1/p+\epsilon)}.$$

Applying Theorem 3.6 in a careful way allows us to deduce Theorem 1.6.

To prove Theorem 1.10, Theorem 1.9 and Theorem 1.8, which deduce multiplicity bounds from the Weak Injective Radius Property, we use Equation 3.2 of Proposition 3.3. To deduce from it upper bounds on multiplicity, one needs *lower bounds* on traces on representations of functions  $h \in C_c^{\infty}(G)$ . The following definition and proposition capture the situation:

**Definition 3.7.** Let  $A \subset \Pi(G)$  be a precompact subset. We say that a family of functions  $\{f_{d_0}\} \subset C_c^{\infty}(G)$ ,  $d_0 \in \mathbb{R}$ ,  $d_0 \geq D$  is *good* for  $A$  if it holds that:

- (1) For every  $\pi \in A$ , and  $\epsilon > 0$ ,

$$q^{d_0(1-1/p(\pi)-\epsilon)} \ll_{A,\epsilon} \text{tr}(\pi(f_{d_0})).$$

- (2) It holds that for every  $g \in G$ ,  $\epsilon > 0$ ,  $f_{d_0}(g) \ll_{\epsilon} q^{d_0\epsilon} \psi_{d_0}(g)$ , where  $\psi_{d_0}(g)$  is from 3.1.

- (3) For every representation  $\pi' \in \Pi(G)$ , it holds that  $0 \leq \text{tr}(\pi'(f_{d_0}))$ .

*Remark 3.8.* If we assume the slightly stronger condition  $|f_{d_0}(g)| \ll_{\epsilon} q^{d_0\epsilon} \psi_{d_0}(g)$  and the fact that  $f_{d_0}$  is left and right  $K$ -finite, then one actually has by Theorem 3.5

$$\text{tr}(\pi(f_{d_0})) \ll_{\pi,\epsilon} q^{d_0(1-1/p(\pi)+\epsilon)},$$

so the lower bound is rather strict.

**Proposition 3.9.** *Let  $A \subset \Pi(G)$  be a precompact subset, and assume that it has a good family of functions. Under this condition, if the sequence  $\{\Gamma_N\}$  of cocompact lattices satisfies the Weak Injective Radius Property with parameter  $\alpha$  then for every  $N \geq 1$ ,  $p > 2$ ,  $\epsilon > 0$ ,*

$$M(A, \Gamma_N, p) \ll_{A,\epsilon} [\Gamma_1 : \Gamma_N]^{1-\alpha(1-2/p)+\epsilon}.$$

Finding general lower bounds on traces (uniformly for a family of representations) is not well studied. Two special cases, which appear (somewhat implicitly) in the work of Sarnak and Xue, correspond to Theorem 1.9 and Theorem 1.10:

- (1) In rank 1, one has a simple classification of spherical irreducible unitary representations. In the Archimedean case, for each  $2 < p \leq \infty$  there is at most a single spherical irreducible unitary representation  $(\pi, V)$  with  $p(\pi) = p$  (with a corresponding spherical function  $\Xi_p(g)$ ), and one can easily deduce lower bounds on the trace of



$f_{d_0} = \chi_{d_0/2} * \chi_{d_0/2}$ , and deduce Theorem 1.9. In the non-Archimedean case, Theorem 1.9 reduces to some statement on graphs. The details are given in Subsection 4.4. We remark that when the graphs are regular the proofs are also given in [26].

- (2) If one is interested in a single representation, one has the following, from which we deduce Theorem 1.10. It follows in the Archimedean case from the asymptotic behavior of leading exponents ([11, 39, Chapter VIII]). The non-Archimedean case is easier, and in any case follows from Theorem 3.11 below.

**Theorem 3.10** (See Section 9). *Let  $(\pi, V) \in \Pi(G)$ . Then the set  $A = \{\pi\}$  has a good family of functions.*

Finally, we provide the following theorem, which implies Theorem 1.8 together with Proposition 3.9.

**Theorem 3.11** (See Theorem 8.5). *Let  $G$  be non-Archimedean. Then there exists a set  $\{K'\}$  of arbitrarily small open-compact subgroups of  $G$ , such that for every  $K'$ ,  $\Pi(G)_{K'\text{-sph}}$  has a good family.*

The proof of Theorem 3.11 is based on two sources. The first is the connection between the Ihara graph Zeta function and expansion (see [31]), and the second is Bernstein description of the Hecke algebra  $C_c(K' \backslash G / K')$  (see [2]). A precise connection for  $(q+1)$ -regular graphs between  $p(\pi)$  for spherical function and poles of the Ihara zeta function may be found in [35]. In recent years, there were various generalizations of the graph zeta function to higher dimensional buildings (see e.g. [37] and the references within). In [34] the second named author generalized the connection between  $p(\pi)$  for representations  $\pi \in \Pi(G)$  with Iwahori-fixed vector and the poles of some generalized zeta function. By a slight variant of those ideas one may prove the special case of Theorem 3.11 when  $K'$  is the Iwahori subgroup. For more general  $K'$  we follow the same ideas, by using Bernstein's description of the Hecke algebra  $C_c(K' \backslash G / K')$ . See Section 8 for details.

If we consider only the spherical case, it would be useful if the functions  $f_{d_0}$  in the definition of a good family will be left and right  $K$ -invariant, i.e.,  $f_{d_0} \in C_c^\infty(K \backslash G / K)$ . Recently, Matz and Templier proved a similar theorem for  $G = PGL_n$  using the Satake isomorphism ([49]). However, their results are less precise- they find a spherical function  $f_{d_0} \in C_c^\infty(K \backslash G / K)$  which satisfies  $f_{d_0}(g) \ll_\epsilon q^{d_0 \epsilon} \psi_{d_0}(g)$ , with a lower bound  $q^{\beta d_0(1-1/p(\pi))} \ll \text{tr}(\pi(f_{d_0}))$  for some  $\beta < 1$ , instead of the optimal bound  $q^{d_0(1-1/p(\pi))} \ll \text{tr}(\pi(f_{d_0}))$ .

Let us finish this discussion with the following conjecture, which concerns only spherical function. for  $g \in G$ , let  $S(g)$  be the  $K$ -bi-invariant function such that  $\int_G f(g) dg = \int_K \int_K \int_{A_+} f(kak') S(a) da dk dk'$ . It holds that for  $g$  "far from the walls"  $S(g) \approx q^{l(g)}$  (see Subsection 4.1).

**Conjecture 3.12.** *There exists  $D > 0$ ,  $L > 0$  such that for every  $\epsilon > 0$  and every  $(\pi, V) \in \Pi(G)_{sph}$  (i.e. a unitary irreducible spherical representation) with  $p(\pi) > 2$ , if  $v \in V$ ,  $\|v\| = 1$  is  $K$ -fixed, then:*

(1) *In the non-Archimedean case, for  $d_0 > D$*

$$\int_{l(d) \leq d_0} S(g) |\langle v, \pi(g)v \rangle|^2 dg \gg_{\epsilon} q^{2d_0(1-1/p(\pi)-\epsilon)}.$$

(2) *In the Archimedean case, for  $d_0 > D$ ,*

$$\int_{l(d) \leq d_0} S(g) |\langle v, \pi(g)v \rangle|^2 dg \gg_{\epsilon} (\lambda(\pi) + 1)^{-L} q^{2d_0(1-1/p(\pi)-\epsilon)}.$$

The exponents in the conjecture are tight, as the corresponding upper bounds can be deduced from Theorem 3.5.

#### 4. PRELIMINARIES

**4.1. Distances and Length of Elements.** Besides our definition of length, the following is standard, see e.g., [24, Section 3]. We mainly follow [39] when  $G$  is Archimedean and [9] when  $G$  is non-Archimedean.

Let  $k$  be  $\mathbb{R}$  or a  $p$ -adic field, and  $|\cdot|_k : k \rightarrow \mathbb{R}_+$  its standard non-trivial valuation. Let  $G$  be a semisimple non-compact algebraic group over  $k$ , of  $k$ -rank  $r$ . Let  $T \cong G_m^r \subset G$  be a maximal  $k$ -split torus. The choice of  $T$  determines the set of weights  $X^*(T)$ , i.e., of rational characters of  $T$ . Let  $\Phi(T, G) \subset X^*(T)$  be the set of roots of  $G$  with respect to  $T$ .

In the real case, if  $T_0 \cong \{\pm 1\}^r$  is the maximal compact subgroup of  $T$ , the connected component of the identity of  $T/A \cong T/T_0$  is the Lie group of a Cartan subalgebra  $\mathfrak{a}$  of  $\mathfrak{g}$ , and we define  $\nu : T \rightarrow A \rightarrow \mathfrak{a} \cong \mathbb{R}^r$  by the logarithm map.

In the  $p$ -adic case, let  $T_0$  be a maximal compact subgroup of  $T$ . Then  $T/T_0 \cong \mathbb{Z}^r$ , this identification defines  $\nu : T \rightarrow \mathbb{Z}^r \subset \mathbb{R}^r$ , and we identify  $\mathbb{R}^r$  with  $\mathfrak{a}$ .

Let  $X^*(T)_{\mathbb{R}} \cong X^*(T) \otimes \mathbb{R}$  be the weight space. For an element  $\alpha \in X^*(T)$ , we let  $\chi_{\alpha} \in \mathfrak{a}^*$  be the linear functional defined such that  $|\alpha(t)|_k = q^{\chi_{\alpha}(\nu(t))}$ , where  $q = e$  in the Archimedean case and otherwise the size of the quotient field of  $k$ . For  $\alpha \in X^*(T)_{\mathbb{R}}$  we define  $\chi_{\alpha} \in \mathfrak{a}^*$  by extension of the action above. This isomorphism (as linear spaces) between  $X^*(T)_{\mathbb{R}}$  and  $\mathfrak{a}^*$  defines an isomorphism between the coweight space  $(X^*(T)_{\mathbb{R}})^*$  and  $\mathfrak{a}$ .

Choose an ordering on the root system which defines the positive roots  $\Phi_+ \subset \Phi(T, G)$  and let  $\Delta = \{\alpha_1, \dots, \alpha_r\} \subset X^*(T)_{\mathbb{R}} \cong \mathfrak{a}^*$  be the basis of  $\Phi$  with respect to this ordering. Let  $\{\omega_1, \dots, \omega_r\} \subset \mathfrak{a}$  be the set of fundamental dominant coweights, i.e. the dual basis to  $\Delta$ . The set  $\{\sum_{i=1}^r x_i \omega_i : x_i \geq 0\} \subset \mathfrak{a}$  is called the dominant sector, or the positive Weyl chamber. It is isomorphic to  $\mathfrak{a}/W$ , where  $W$  is the Weyl group of the root system.

Let  $\Phi_+^{\vee} \subset \mathfrak{a}$  be the corresponding co-root system, let  $\Delta^{\vee} = \{\alpha_1^{\vee}, \dots, \alpha_r^{\vee}\}$  be the co-basis of positive co-roots and let  $\{\omega_1^{\vee}, \dots, \omega_r^{\vee}\} \subset \mathfrak{a}^*$  be the set of fundamental dominant weights. We

define a partial ordering on  $\mathfrak{a}$  by  $\alpha \geq_{\mathfrak{a}} \alpha'$  if and only if  $\omega_i^{\vee}(\alpha) \geq \omega_i^{\vee}(\alpha')$  for every fundamental dominant weight, or alternatively  $\alpha - \alpha'$  is a non-negative sum of elements of  $\Delta^{\vee}$ .

Let  $P$  be the Borel subgroup with respect to the set of positive roots. It holds that  $P = MN$ , where  $M$  is the centralizer of  $T$  in  $G$  and  $N$  is the unipotent radical of  $P$  ([9, p. 134]). Let  $K$  be a maximal special compact open subgroup (i.e., in the  $p$ -adic case we choose it to be “good” in the sense of Bruhat and Tits [7]). The Iwasawa decomposition  $G = KP$  holds ([39, Proposition 1.2],[9, p. 140]). It holds that  $M = (M \cap K) \cdot T$ , and we extend  $\nu : M \rightarrow \mathbb{R}^r$  by  $\nu(k) = 1$  for  $k \in M \cap K$ .

Let us recall the Cartan decomposition  $G = KA_+K$ :

- In the real case, following [39, Theorem 5.20],  $A \cong T/T_0$  is the Lie group of the Cartan subalgebra  $\mathfrak{a}$  of  $\mathfrak{g}$ .  $A_+$  is the exponent of the closure of the dominant sector in  $\mathfrak{a}$ , i.e. the set of elements  $A_+ = \{t \in A : \forall \alpha \in \Phi_+, \alpha(t) \geq 1\}$ . It is isomorphic (as a set, using the exponential map) to the dominant sector in  $\mathfrak{a}$ .
- In the  $p$ -adic case, following [9, p. 140], let  $\Lambda \cong M/M^0$ , where  $M^0$  is the maximal compact subgroup of  $M$ . We identify elements of  $\Lambda$  with elements of  $M \subset P$ . Elements of the weight space are unramified characters of  $M$ , and therefore are characters of  $\Lambda$  as well. We let  $A_+ = \{\lambda \in \Lambda : \forall \alpha \in \Phi_+, \alpha(\lambda) \geq 1\}$ . The action  $\nu : T/T_0 \rightarrow \mathfrak{a}$  extends to  $\nu : M/M^0 \rightarrow \mathfrak{a}$ . Then  $A_+$  is isomorphic as a set with the intersection of  $\nu(\Lambda)$  with the dominant sector in  $\mathfrak{a}$ . It is also isomorphic with a subset of the special vertices in the dominant sector in an apartment in the Bruhat-Tits building of  $G$ .

In both cases, there exists a map  $\nu : A_+ \rightarrow \mathfrak{a}$ . It is also useful to extend it to a map  $H : G \rightarrow \mathfrak{a}$ , as follows – for  $g \in G$  let  $m \in M$  be its  $m$ -part in the Iwasawa decomposition and let  $H(g) = \nu(m)$ .

We will need the following fundamental technical lemma:

**Lemma 4.1.** *For  $a \in A_+$  and  $k \in K$*

$$H(ak) \leq_{\mathfrak{a}} H(a).$$

*Proof.* For the non-Archimedean case see [7, Proposition 4.4.4(i)]. For the Archimedean case see [22, Corollary 3.5.3].  $\square$

**Corollary 4.2.** *Let  $a, a', a'' \in A_+$ . If  $KaKa'K \cap Ka''K \neq \phi$  then  $\nu(a'') \leq_{\mathfrak{a}} \nu(a) + \nu(a')$ .*

*Proof.* We first notice that the following property of the  $H$ -function: for  $k \in K, g \in G, m \in M, n \in N$

$$H(kgmn) = H(g) + H(m)$$

Now, if  $KaKa'K \cap Ka''K \neq \phi$  then  $a'' = k_0 a k_1 a' k_2$ . Applying the Iwasawa decomposition to  $a' k_2$  we have  $a' k_2 = k'_2 m n$  with  $H(m) \leq_{\mathfrak{a}} H(a')$  by Lemma 4.1. Applying Lemma 4.1

again we have

$$\begin{aligned}\nu(a'') &= H(k_0 a k_1 k_2' m n) = H(k_0 a k_1 k_2') + H(m) \\ &\leq_{\mathfrak{a}} H(a) + H(a') = \nu(a) + \nu(a').\end{aligned}$$

□

Corollary 4.2 in the non-Archimedean case is [7, Proposition 4.4.4(iii)], and is deduced from Lemma 4.1 in the same way.

Let  $\delta(\tilde{p})$  be the left modular character of  $P$ , i.e. if  $d\tilde{p}$  is a left Haar measure on  $P$  then  $\delta(\tilde{p})d\tilde{p}$  is a right Haar measure. Normalize the measures so that for  $f \in C_c(G)$ ,  $\int_G f(g)dg = \int_K \int_P f(k\tilde{p})\delta(\tilde{p})d\tilde{p}dk = \int_K \int_P f(\tilde{p}k)d\tilde{p}dk$ ,  $\int_K dk = 1$ .

It can also be defined as follows:  $M$  acts by conjugation on the Lie algebra  $\mathfrak{n}$  of  $N$ . Then for  $m \in M$ ,  $\delta(m) = |\text{DetAd}_{\mathfrak{n}}(m)|_k$  ([9, p. 135], [39, Proposition 5.25]). Unwinding the definitions,  $\delta(m) = q^{2\rho(\nu(m))}$ , where  $\rho = \frac{1}{2} \sum_{\alpha \in \Phi_+} (\dim \mathfrak{g}_{\alpha}) \chi_{\alpha}$ . Here  $\mathfrak{g}_{\alpha}$  is the root space of  $\alpha$  in the Lie algebra  $\mathfrak{g}$  of  $G$ . Also recall that  $q = e$  for  $k = \mathbb{R}$  and otherwise is the size of the quotient field of  $k$ . As an example, for  $G = SL_n(\mathbb{R})$ , and the matrix  $a = \text{diag}(a_0, \dots, a_{n-1})$ ,  $\delta(a) = \prod a_i^{n-1-2i}$ .

We associate with each element  $a \in A_+ \subset G$  a length  $l : A_+ \rightarrow \mathbb{R}_{\geq 0}$  by  $l(a) = \log_q \delta(a) = 2\rho(a)$ . We extend  $l : G \rightarrow \mathbb{R}_{\geq 0}$  by  $l(kak') = l(a)$ . By definition,  $l$  is left and right  $K$ -invariant.

For  $a \in T$ , we may identify  $l(a)$  by the entropy (taken with logarithm in base  $q$ ) of the dynamical system of translation of  $\Gamma \backslash G$  by  $a$ , with respect to the Haar measure ( $\Gamma$  here is an arbitrary lattice, see [18, Theorem 7.9]). Using this fact, we have for  $a \in A$ ,  $l(a) = l(a^{-1})$  and therefore for every  $g \in G$ ,  $l(g) = l(g^{-1})$ . The same fact can be proven directly.

**Proposition 4.3.** *For  $g_1, g_2 \in G$ , it holds that  $l(g_1 g_2) \leq l(g_1) + l(g_2)$ .*

*Proof.* The proposition actually states that if  $KaKa'K \cap Ka''K \neq \emptyset$  for  $a, a', a'' \in A_+$ , then  $l(a'') \leq l(a) + l(a') = l(aa')$ . Since  $l(a) = 2\rho(a)$  it follows from Corollary 4.2. □

For  $a \in A_+$ , define  $S(a) = \int \delta(ak)dk$ . For  $f \in C_c(G)$ , we have  $\int f(g)dg = \int_K \int_K \int_{A_+} f(kgk')S(a)dkdk'da$ . We interpret  $S(a)$  as the measure of the ‘‘circle’’  $KaK$ . In the non-Archimedean case  $S(a) \approx \delta(a)$  ([9, p. 141]). In the Archimedean case, by [39, Proposition 5.28],

$$S(a) = \prod_{\alpha \in \Phi_+} (\sinh(\chi_{\alpha}(a)))^{\dim \mathfrak{g}_{\alpha}}.$$

Since  $\sinh(x) \approx_{\beta} e^x$  for  $x > \beta$ , for  $a \in A_+$  ‘‘far from the walls’’, i.e. with  $\chi_{\alpha}(a) > \beta$  for every  $\alpha \in \Delta$  (and therefore  $\chi_{\alpha}(a) > \beta$  for every  $\alpha \in \Phi_+$ ), we have  $S(a) \asymp \delta(a) = q^{l(a)}$ . Near the walls where  $\sinh x \approx x$  this approximation fails, but we still have  $S(a) \ll \delta(a)$ . In any case, if we choose some norm  $\|\cdot\|_{\mathfrak{a}}$  on  $A_+$  then for every  $\tau > 0$ ,  $a \in A_+$ ,  $\mu(\{g : \|a_g - a\|_{\mathfrak{a}} \leq \tau\}) \asymp_{\tau} q^{l(a)}$ , since the set  $\{a' \in A_+ : \|a' - a\|_{\mathfrak{a}} \leq \tau\}$  contains elements which are far enough (with respect to  $\tau$ ) from the walls.

We deduce that the size of balls for  $l \gg 1$ ,

$$(4.1) \quad q^l \ll \mu \left( \cup_{a': l(a') \leq l} K a' K \right) = \int_{a \in A_+ : l(a) \leq l} S(a) da \ll p(l) q^l \ll_{\epsilon} q^{l(1+\epsilon)},$$

for some polynomial  $p$ .

*Remark 4.4.* The literature has two popular choices of “distance” or “length” on  $G/K$  or  $G$ .

- (1) For  $G \subset SL_n$ , where  $K = G \cap K'$  for  $K'$  maximal compact in  $SL_n$ , one defines  $\tilde{l}(g) = \log \|g\|$ , where  $\|\cdot\|$  is some matrix norm on  $GL_n$ . Recall that we are mainly interested in distances as  $g \rightarrow \infty$ , so the specific choice of matrix norm does not matter. Such choice (without calling it a distance) is studied in [53, 17].
- (2) For  $G$  Archimedean, let  $\mathfrak{g}, \mathfrak{k}$  be the Lie algebras of  $G$  and  $K$ , and let  $B : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$  the Killing form of  $G$ . Let  $\mathfrak{p} = \{X \in \mathfrak{g} : B(X, Y) = 0 \forall Y \in \mathfrak{k}\}$ . Then  $B|_{\mathfrak{p} \times \mathfrak{p}}$  is positive definite.  $\mathfrak{p}$  can be identified with the tangent space of  $G/K$  at the identity and it defines a natural Riemannian structure on  $G/K$ , with length  $\hat{l}(g) = \hat{d}(g, 1)$ . See e.g. [14, Section 2], [53]. Similar distance is used in the  $p$ -adic case in [57, 2.3].

Since we mostly care about far distances in the group, and since  $K$  is compact, by the Cartan decomposition it suffices to compare  $l$  to other distances on  $A_+$ . In general,  $l$  and  $\tilde{l}$  looks like an  $L^1$ -norm on  $A_+$ , and  $\hat{l}$  looks like an  $L^2$ -norm.

Let us concentrate on  $G = SL_n(\mathbb{R})$  and  $\tilde{l}(g) = \ln(\|g\|_2)$ ,  $\|g\|_2^2 = \text{tr}(gg^t)$ . For  $G = SL_2(\mathbb{R})$ , its symmetric space is the hyperbolic plane with the standard metric of curvature  $-1$ , and  $l$  we defined above coincides with the hyperbolic metric. For example, consider the matrix  $g = \begin{pmatrix} e^{\frac{t}{2}} & 0 \\ 0 & e^{-\frac{t}{2}} \end{pmatrix}$ . Then  $l(g) = t$ , and  $\tilde{l}(g) = \frac{1}{2} \ln(e^t + e^{-t}) \approx \frac{t}{2}$ . In fact, for every  $g \in SL_2(\mathbb{R})$ ,  $l(g) - 2\tilde{l}(g) = O(1)$ , and  $l$  and  $\tilde{l}$  are equal up to additive constant.

For  $G = SL_3(\mathbb{R})$  it is no longer true. For  $g_1 = \begin{pmatrix} e^{t/3} & & \\ & e^{t/3} & \\ & & e^{-2t/3} \end{pmatrix}$ ,  $g_2 = \begin{pmatrix} e^{2t/3} & & \\ & e^{-t/3} & \\ & & e^{-t/3} \end{pmatrix}$ ,

$l(g_1) = l(g_2) = 2t$ . While  $\tilde{l}(g_1) \approx t/3$ ,  $l(g_2) \approx 2t/3$ . So the two distances  $l, \tilde{l}$  are not equivalent, but are only Lipschitz-equivalent, with  $\frac{3}{2}\tilde{l} \leq l + O(1) \leq 3\tilde{l}$ . However, if we chose  $\tilde{\tilde{l}}(g, I) = 2(\tilde{l}(g, I) + \tilde{l}(g^{-1}))$  then  $l(g) - 2\tilde{\tilde{l}}(g) = O(1)$ . This solution no longer works for  $SL_4(\mathbb{R})$ .

**4.2. Growth of Matrix Coefficients and Harish-Chandra's Bounds.** For  $2 \leq p \leq \infty$ , let  $\Xi_p(g) = \int \delta(gk)^{-1/p} dk$  be the  $p$ -th version of Harish-Chandra's function. Let  $\Xi(g) = \Xi_2(g)$  be standard Harish-Chandra's function. Note that since  $\Xi(g)$  is left and right  $K$ -invariant it only depends on  $a \in A_+$  from the Cartan decomposition of  $g$ .

An explicit upper bound on Harish-Chandra's function is given by the following Theorem:

**Theorem 4.5.** *We have for every  $g \in \mathfrak{g}$  and  $\epsilon > 0$ ,*

$$\Xi_p(g) \leq \Xi^{2/p}(g) \leq q^{(1/2-1/p)l(g)} \Xi(g) \leq C_2(G) ((l(g) + 1))^r q^{-l(g)/p} \ll_\epsilon q^{-l(g)(1/p-\epsilon)},$$

where  $C(G)$  depends only on  $G$ .

*Proof.* By the Cartan decomposition one may verify the Theorem for  $a \in A_+$ , where  $q^{l(a)} = \delta(a)$ . The first inequality follows from convexity, the second inequality follows from  $\Xi(g) \geq q^{-l(g)/2}$  (see Equation 4.4), and the fourth inequality is trivial. We are left with the third inequality.

For the Archimedean case see [30, Theorem 3] or [39, Proposition 7.15]. For the non-Archimedean case the standard reference is [56, 4.2.1], (where there is an assumption that  $\text{char } k = 0$ ), but the theorem is well known to experts. In any case, the general non-Archimedean result can be deduced from the results of [34] for arbitrary affine buildings.  $\square$

A representation  $(\pi, V)$  of  $G$  is called tempered if  $p(\pi) \leq 2$ . The following theorem is the standard reference for upper bounds on matrix coefficients:

**Theorem 4.6** ([13]). *Let  $(\pi, V)$  be a unitary irreducible tempered representation of  $G$  and let  $v_1, v_2 \in V$  be two  $K$ -finite vectors, such that  $\dim \text{Span} K v_1 = d_1$ ,  $\dim \text{Span} K v_2 = d_2$ . Then  $|\langle v_1, g v_2 \rangle| \leq \sqrt{d_1 d_2} \|v_1\| \|v_2\| \Xi(g)$ .*

The work [13] also provides a bound when  $p(\pi) > 2$ :

**Theorem 4.7** ([13]). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq 2k$ ,  $k \in \mathbb{N}$  and let  $v_1, v_2 \in V$  be two  $K$ -finite vectors, such that  $\dim \text{Span} K v_1 = d_1$ ,  $\dim \text{Span} K v_2 = d_2$ . Then  $|\langle v_1, g v_2 \rangle| \leq \sqrt{d_1 d_2} \|v_1\| \|v_2\| \Xi^{1/k}(g)$ .*

This theorem is not sufficient for this work since we need more precise bounds when  $p(\pi) \notin 2\mathbb{N}$ . The following theorem contains a general upper bounds that is good enough for all the applications of this paper.

**Theorem 4.8.** *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq p$ ,  $p \geq 2$ , and let  $v_1, v_2 \in V$  be two  $K$ -finite vectors, such that  $\dim \text{span} K v_1 = d_1$ ,  $\dim \text{span} K v_2 = d_2$ . Then*

$$|\langle v_1, g v_2 \rangle| \leq \sqrt{d_1 d_2} \|v_1\| \|v_2\| \Xi_p(g).$$

The main approximation we need can actually be deduced from earlier works, so Theorem 4.8 is not needed directly for us. Let us state two existing theorems that may serve as replacements for it.

We first consider bounds on matrix coefficients of a single representation  $\pi$ . In this case one has in the Archimedean case the theory of leading coefficients ([39, Chapter VIII]), which we describe in Subsection 9.1. From Theorem 9.1, and Theorem 9.2 we may deduce the following. In the  $p$ -adic case analogous results hold, although we couldn't find an explicit reference. It may be deduced from [10, Section 4].

**Theorem 4.9** (See Theorems 9.1,9.2). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq p$ ,  $p \geq 2$ , and let  $v_1, v_2 \in V$  be two  $K$ -finite vectors. Then for every  $\epsilon > 0$ ,*

$$|\langle v_1, gv_2 \rangle| \ll_{v_1, v_2, \pi, \epsilon} q^{-l(g)^{(1/p-\epsilon)}}.$$

Next we consider bounds on matrix coefficients when  $v_1, v_2$  are  $K$ -fixed. In such case (if  $v_1, v_2 \neq 0$ ) the representation is spherical, and well understood (see Subsection 4.4 below). From those bounds, we have:

**Theorem 4.10** (See [24, Section 3]). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq p$ , and let  $v_1, v_2 \in V$  be two  $K$ -fixed vectors. Then for every  $\epsilon > 0$ ,*

$$|\langle v_1, \pi(g)v_2 \rangle| \leq \|v_1\| \|v_2\| \Xi(g) \delta^{1/2-1/p} \ll_{\epsilon} \|v_1\| \|v_2\| q^{-l(g)^{(1/p-\epsilon)}}.$$

Let us provide a proof of Theorem 4.8 when  $v_1, v_2$  are  $K$ -fixed (in which case it closely resembles Theorem 4.10). Similar ideas can be used to give bounds for  $K$ -finite vectors as well, using the ideas in [13]. For simplicity and since we do not need the stronger theorem we restrict to the  $K$ -fixed case. Let us first prove the following lemma, which is based on the proof of [13, Theorem 2]. For  $f \in L^p(G)$  and  $g \in G$  we let  $gf \in L^p(G)$  be  $gf(g') = f(g^{-1}g')$ .

**Lemma 4.11.** *If  $f_1 \in L^{p/(p-1)}(G)$ ,  $f_2 \in L^p(G)$  are two  $K$ -fixed vectors, then  $|\langle f_1, gf_2 \rangle| \leq \|f_1\|_{p/(p-1)} \|f_2\|_p \Xi_p(g)$ .*

*Proof.* To avoid integrability questions we assume that  $f_1, f_2 \in C_c(G)$  and deduce the theorem by density. For  $f \in C_c(G)$  it holds that

$$\int_G f(x) dx = \int_K \int_P f(k\tilde{p}) \delta(p) d\tilde{p} dk,$$

so

$$\begin{aligned} |\langle f_1, gf_2 \rangle| &\leq \int_K \int_P |f_1(k\tilde{p})| |f_2(g^{-1}k\tilde{p})| \delta(\tilde{p}) d\tilde{p} dk = \\ &= \int_K \int_P |f_1(k\tilde{p})| \delta(\tilde{p})^{1/p'} |f_2(g^{-1}k\tilde{p})| \delta(\tilde{p})^{1/p} d\tilde{p} dk \\ &\leq \int_K \left( \int_P |f_1(k\tilde{p})|^{p'} \delta(\tilde{p}) d\tilde{p} \right)^{1/p'} \left( \int_P |f_2(g^{-1}k\tilde{p})|^p \delta(\tilde{p}) d\tilde{p} \right)^{1/p} dk. \end{aligned}$$

Since  $f_1$  is  $K$ -fixed and  $\int_K 1 dk = 1$  we have for every  $k \in K$ ,

$$\left( \int_P |f_1(k\tilde{p})|^{p'} \delta(\tilde{p}) d\tilde{p} \right)^{1/p'} = \|f_1\|_{p'}.$$

write  $g^{-1}k = k_0\tilde{p}_0$ . Then

$$\begin{aligned} \left( \int_P |f_2(g^{-1}k\tilde{p})|^p \delta(\tilde{p}) d\tilde{p} \right)^{1/p} &= \left( \int_P |f_2(k_0\tilde{p}_0\tilde{p})|^p \delta(\tilde{p}) d\tilde{p} \right)^{1/p} \\ &= \left( \int_P |f_2(\tilde{p}_0\tilde{p})|^p \delta(\tilde{p}_0\tilde{p}) d\tilde{p} \right)^{1/p} \delta(\tilde{p}_0)^{-1/p} \\ &= \left( \int_P |f_2(\tilde{p})|^p \delta(\tilde{p}) d\tilde{p} \right)^{1/p} \delta(\tilde{p}_0)^{-1/p} \\ &= \|f_2\|_p \delta(\tilde{p}_0)^{-1/p}. \end{aligned}$$

Since  $\delta$  is left  $K_0$ -fixed (by definition) and  $\tilde{p}_0 = k_0^{-1}g^{-1}k$

$$\begin{aligned} \int_{K_0} \left( \int_P |f_2(g^{-1}k\tilde{p})|^p \delta(\tilde{p}) d\tilde{p} \right)^{1/p} dk &= \|f_2\|_p \int \delta(g^{-1}k)^{-1/p} dk = \\ &= \|f_2\|_p \Xi_p(g). \end{aligned}$$

□

The lemma has a nice corollary: for  $a \in A_+$  let  $A_a$  be the operator  $A_a : C(K \backslash G) \rightarrow C(K \backslash G)$  defined by  $A_a f(g) = \int_K f(k^{-1}a^{-1}g) dk = \int_K \int_K f(k^{-1}a^{-1}k'g) dkdk'$ . Let us remark that we may define  $A_g$  for  $g \in G$ , but it only depends on the  $A_+$  component of  $g$  from the Cartan decomposition. Since  $A_a$  is a translation followed by an average, its  $L^p$ -norm is bounded by 1 on  $L^p(G) \cap C(K \backslash G)$ , and therefore it defines an operator  $A_a : L^p(K \backslash G) \rightarrow L^p(K \backslash G)$ .

**Corollary 4.12.** *The norm of  $A_a : L^p(K \backslash G) \rightarrow L^p(K \backslash G)$  is bounded by  $\Xi_p(a)$ .*

*Proof.* Let  $p' = p/(p-1)$ . Let  $f \in L^p(K \backslash G)$ . Let  $f_1 \in L^{p'}(K \backslash G)$  with  $\|f_1\|_{p'} = 1$  be such that  $\langle f_1, A_a f \rangle = \|A_a f\|_p$ . Then since  $f_1$  is left  $K$ -invariant  $\langle f_1, A_a f \rangle = \langle f_1, a f \rangle$ . Applying Lemma 4.11, we have

$$\|A_a f\|_p \leq \Xi_p(a) \|f\|_p,$$

as needed. □

Note that if  $(\pi, V)$  is a unitary representation of  $G$ , we may define  $\pi(A_a) : V^K \rightarrow V^K$  by the same arguments as above, as

$$(4.2) \quad \pi(A_a)v = \int_K \pi(ka)v dk = \int_K \int_K \pi(kak')v dkdk',$$



and  $\|A_a\|_V \leq 1$ . By standard arguments, this operation “commutes” with taking matrix coefficients – if we let for  $v_1, v_2 \in V^K$ ,  $\varphi_{v_1, v_2}(g) = \langle v_1, \pi(g)v_2 \rangle$ ,  $\varphi_{v_1, v_2}(g) \in L^\infty(G)$ , then  $A_a \varphi_{v_1, v_2}(g) = \varphi_{A_a v_1, v_2}(g)$ .

**Corollary 4.13.** *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq p$ , and let  $v_1, v_2 \in V$  be two  $K$ -fixed vectors. Then*

$$|\langle \pi(g)v_1, v_2 \rangle| \leq \|v_1\| \|v_2\| \Xi_p(g).$$

*Proof.* We may assume that  $v_1, v_2$  are of norm 1. Since both of them are left  $K$ -invariant  $\langle \pi(g)v_1, v_2 \rangle = \langle \pi(A_{a_g})v_1, v_2 \rangle$ , where  $a_g \in A_+$  is the  $A_+$  component of the Cartan decomposition of  $g$ . If  $\pi$  is irreducible, it is well known that the subspace  $V^K$  of  $K$ -fixed vectors is one dimensional. Therefore  $v_1$  is an eigenvector of  $\pi(A_{a_g})$  on  $V^K$ . Therefore  $c_{v_1, v_2}(g)$  is an eigenvector of  $A_{a_g}$  on  $L^{p+\epsilon}(g)$  for every  $\epsilon > 0$ , with eigenvalue  $\langle \pi(g)v_1, v_2 \rangle$ . As the norm of  $A_{a_g}$  on  $L^{p+\epsilon}(K \backslash G)$  is bounded by  $\Xi_{p+\epsilon}(g)$ , each of its eigenvalues is bounded by  $\Xi_{p+\epsilon}(g)$  as well. Therefore  $|\langle \pi(g)v_1, v_2 \rangle| \leq \Xi_{p+\epsilon}(g)$  for every  $\epsilon > 0$ . By taking  $\epsilon \rightarrow 0$  we deduce  $|\langle v_1, \pi(A_g)v_2 \rangle| \leq \Xi_p(g)$ , as required.  $\square$

**4.3. Upper Bounds on Operators.** Let  $(\pi, V)$  be a unitary representation of  $G$ . For  $a \in A_+$  there let  $\pi(A_a) : V^K \rightarrow V^K$  be from Equation 4.2. Similarly, for  $d_0 \in \mathbb{R}_{\geq 0}$ , let  $\chi_{d_0}, \psi_{d_0} \in C_c^\infty(G)$  be as in Definition 3.1 the introduction. Recall that  $\chi_{d_0}$  is a smooth approximations for the characteristic function of  $\{g : l(g) \leq d_0\}$  and  $\psi_{d_0}$  is a smooth approximation for  $q^{(d_0 - l(g))/2} \chi_{d_0}$ .

For  $h \in C_c(G)$  we define as usual  $\pi(h)v = \int_G h(g) \pi(g)v dg$ .

**Corollary 4.14.** *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$  with  $p(\pi) \leq p$ , and let  $v \in V$  be a  $K$ -fixed vector. Then*

$$\begin{aligned} \|\pi(A_a)v\|_V &\ll_\epsilon q^{-l(a)(1/p-\epsilon)} \|v\|_V \\ \|\pi(\chi_{d_0})v\|_V &\ll_\epsilon q^{d_0(1-1/p+\epsilon)} \|v\|_V \\ \|\pi(\psi_{d_0})v\|_V &\ll_\epsilon q^{d_0(1-1/p+\epsilon)} \|v\|_V. \end{aligned}$$

*Proof.* The bound on  $\pi(A_a)$  follows from Corollary 4.13, using the explicit bounds of Theorem 4.5.

We will only prove the estimate for  $\psi_{d_0}$ , since the proof for  $\chi_{d_0}$  is similar and a little easier.

$$\begin{aligned}
\|\pi(\psi_{d_0})v\| &\leq \int_{l(g) \leq d_0+1} q^{(d_0-l(g))/2} \|\pi(g)v\| dg \\
&\ll_\epsilon \int_{l(g) \leq d_0+1} q^{d_0/2-l(g)(1/2+1/p-\epsilon)} \|v\| dg \\
&= \int_K \int_K \int_{l(a) \leq d_0+1} S(a) q^{d_0/2-l(k_1 a k_2)(1/2+1/p-\epsilon)} \|v\| dk_1 da dk_1 \\
&\ll \int_{l(a) \leq d_0+1} q^{l(a)} q^{d_0/2-l(a)(1/2+1/p-\epsilon)} \|v\| da \\
&\ll_\epsilon \int_{l(a) \leq d_0+1} q^{d_0/2+l(a)(1/2-1/p+\epsilon)} \|v\| da \\
&\leq \int_{l(a) \leq d_0+1} q^{d_0/2+d_0(1/2-1/p+\epsilon)} \|v\| da \\
&\ll_\epsilon q^{d_0(1-1/p+\epsilon)} \|v\|_V.
\end{aligned}$$

In the last line we used the fact that  $\int_{l(a) \leq d_0+1} da \ll_\epsilon q^{d_0\epsilon}$ .  $\square$

**4.4. Spherical Functions and Lower Bounds.** Let us recall the definition of spherical functions. For  $\lambda \in a_{\mathbb{C}}^* = a^* \otimes \mathbb{C}$  dominant, we let  $\lambda_P : P \rightarrow \mathbb{C}^\times$  be  $\lambda_P(mn) = q^{\lambda(\nu(m))} \delta^{-1/2}(m) = q^{(\lambda-\rho)(\nu(m))}$ . Extend it to  $\tilde{\lambda} : G \rightarrow \mathbb{C}$  by  $\tilde{\lambda}(kp) = \lambda_P(p)$ . Finally, define the spherical function

$$\varphi_\lambda(g) = \int_K \tilde{\lambda}(gk) dk.$$

Note that  $\varphi_0 = \Xi$  and  $\varphi_{(1-2/p)\rho} = \Xi_p$  for  $2 \leq p \leq \infty$ .

The theory of spherical functions was developed by Harish-Chandra in the Archimedean case and by Satake in the non-Archimedean case (see [24, Subsection 3.2] and the reference therein). We will need some very basic properties of spherical functions.

Now let  $(\pi, V) \in \Pi(G)_{\text{sph}}$  be a unitary spherical representation, i.e. a unitary irreducible representation with a non-trivial  $K$ -fixed vector. If  $v_1, v_2$  are  $K$ -fixed, then there exists a dominant  $\lambda \in a_{\mathbb{C}}^* = a^* \otimes \mathbb{C}$ , such that

$$\langle v_1, \pi(g)v_2 \rangle = \langle v_1, v_2 \rangle \varphi_\lambda(g).$$

Let  $\omega_1, \dots, \omega_r \in \mathfrak{a}$  be the fundamental coweights. Then since  $\lambda$  is dominant  $\text{Re}\lambda(\omega_i) \geq 0$  for  $1 \leq i \leq r$ . By [39, Theorem 8.48] (the  $p$ -adic case being similar),  $p(\pi) \leq p$  if and only if

$$(4.3) \quad \text{Re}\lambda(\omega_i) \leq (1 - 2/p)\rho(\omega_i)$$

for every  $1 \leq i \leq r$ .

Upper bounds on Spherical functions are given as follows. By [24, Lemma 3.3], it holds that for  $a \in A_+$ ,  $k, k' \in K$  and  $\epsilon > 0$ ,

$$\varphi_\lambda(kak') = \varphi_\lambda(a) \ll q^{\operatorname{Re}\lambda(\nu(a))} \Xi(a) \ll_\epsilon q^{\operatorname{Re}\lambda(\nu(a))} q^{-l(a)(1/2-\epsilon)},$$

which together with Equation 4.3 gives a proof for Theorem 4.10.

Lower bounds on Spherical functions can be given in general if  $\operatorname{Re}\lambda = \lambda$ . By Lemma 4.1, if  $\operatorname{Re}\lambda = \lambda$  and  $\operatorname{Re}\lambda(\omega_i) \leq \rho(\omega_i)$  then for  $a \in A_+$ ,  $k, k' \in K$

$$(4.4) \quad \varphi_\lambda(kak') = \varphi_\lambda(a) = \int_K \tilde{\lambda}(ak) dk \geq \int_K \tilde{\lambda}(a) dk = \lambda_P(a).$$

In the case when  $G$  is Archimedean, it is well known that the dominant  $\lambda \in a_{\mathbb{C}}^*$  which occur this way for unitary spherical representations must satisfy  $-\bar{\lambda} = w\lambda$  for some  $w \in W$  ([40]).

If we moreover assume that  $G$  is of rank 1, then  $a_{\mathbb{C}}^* \cong \mathbb{C}$  and  $W = \{1, s\}$  acts by  $s\lambda = -\lambda$ . Therefore the only dominant  $\lambda \in a_{\mathbb{C}}^*$  which may occur satisfy either  $\operatorname{Re}\lambda = 0$  or  $\operatorname{Re}\lambda = \lambda$ . In the case  $\operatorname{Re}\lambda = 0$  the corresponding unitary representation satisfies  $p(\pi) = 2$ . If  $\operatorname{Re}\lambda = \lambda$  then  $\lambda = \alpha\rho$  with  $\alpha \geq 0$ . By Equation 4.3,  $\alpha \leq 1$ . Write  $\alpha = 1 - 1/p$ ,  $2 \leq p \leq \infty$ . Using 4.3 and the definition of  $\Sigma_p(g)$  we conclude that for  $p > 2$ , there is at most a single unitary irreducible representation  $(\pi, V)$  with a non-trivial  $K$ -fixed vector and  $p(\pi) = p$ . The corresponding spherical function is  $\Xi_p(g)$ . We conclude:

**Proposition 4.15.** *Let  $G$  be Archimedean of rank 1,  $p > 2$  and  $(\pi, V)$  a unitary irreducible representation of  $G$  with  $p(\pi) = p$ , having a non-trivial  $K$ -fixed vector  $v$ . Then  $\pi(A_a)v = \lambda_1 v$ ,  $\pi(\chi_{d_0})v = \lambda_2 v$ , with*

$$e^{-l(a)/p} \leq \lambda_1 \\ e^{d_0(1-1/p)} \ll \lambda_2 \text{ for } d_0 \geq 1.$$

Moreover, if  $h \in C_c(K \backslash G / K)$  satisfies  $h(g) \geq \chi_{d_0}(g)$  for every  $g \in G$ , then  $\pi(h)v = \lambda_h v$ , with  $\lambda_h \geq \lambda_2$ .

*Proof.* It is well known that in this case the set of all  $K$ -fixed vectors in  $V$  is one dimensional and equals  $\operatorname{span}\{v\}$ . Since  $\pi(A_a)v$ ,  $\pi(\chi_{d_0})v$ ,  $\pi(h)v$  are  $K$ -fixed we get  $\pi(A_a)v = \lambda_1 v$ ,  $\pi(\chi_{d_0})v = \lambda_2 v$  and  $\pi(h)v = \lambda_h v$ . Applying matrix coefficients we see that

$$\begin{aligned} \lambda_1 &= \Xi_p(a) \\ \lambda_2 &= \int_G \chi_{d_0}(g) \Xi_p(g) dg \\ \lambda_h &= \int_G h(g) \Xi_p(g) dg \geq \int_G \chi_{d_0}(g) \Xi_p(g) dg = \lambda_2. \end{aligned}$$

Using Inequality 4.4 we get

$$\Xi_p(g) \geq q^{-\frac{2}{p}\rho(\nu(a))} q^{-l(a)/p}.$$

And therefore  $\lambda_1 \geq q^{-l(a)/p}$  and for  $d_0 \geq 1$ ,

$$\begin{aligned} \lambda_2 &= \int_G \chi_{d_0}(g) \Xi_p(g) dg \geq \\ &\geq \int_0^{d_0} \sinh(t) e^{-t/p} dt \gg e^{(1-1/p)d_0}. \end{aligned}$$

□

Similar analysis can be done for the non-Archimedean rank 1 case. When the Bruhat-Tits Tree of  $G$  is not regular the some further calculations are needed. The analysis of this can be done by analyzing the slightly more general case of regular and bi-regular trees, which is a slight divergence fro the main topic. See [31] or [35] for an analysis of this case. We give here the details from a group theoretic point of view, since the exact results we need do not appear in those references.

Recall that we have a map  $\nu : M/M^0 \rightarrow \mathfrak{a}$  with a discrete image. Identify its image with  $\mathbb{Z}$  and use the usual bilinear form on  $\mathbb{R} \times \mathbb{R}$  to identify  $a_{\mathbb{C}}^* \cong \mathbb{C}$ . Then since  $\nu(m) \cong \mathbb{Z}$  the function  $\varphi_\lambda$  is equal to  $\varphi_{\lambda'}$  if  $\lambda = \lambda' + 2\pi i m \ln(q)$  for some  $m \in \mathbb{Z}$  (under the identification  $a_{\mathbb{C}}^* \cong \mathbb{C}$ ). Therefore there are 3 cases which may occur in unitary spherical representations:

- (1)  $\operatorname{Re}(\lambda) = 0$ . In such case  $p(\pi) = 2$  for a corresponding  $\pi$ .
- (2)  $\operatorname{Re}(\lambda) = \lambda$  and  $\lambda \neq 0$ . In such case if  $\lambda$  corresponds to a unitary representation  $(\pi, V)$  then necessarily  $\lambda = (1 - 2/p)\rho$  for  $2 < p \leq \infty$ , and  $p(\pi) = p$ . The spherical function is  $\Xi_p$ .
- (3)  $\operatorname{Re}(\lambda) + \pi i \ln(q) = \lambda$  and  $\lambda \neq \pi i \ln(q)$ . In such case if  $\lambda$  corresponds to a unitary representation  $(\pi, V)$  then necessarily  $\lambda = (1 - 2/p)\rho + \pi i \ln(q)$  for  $2 < p \leq \infty$ , and  $p(\pi) = p$ .

Further analysis requires separation to cases. If the Bruhat-Tits tree of  $G$  is regular then case 3 occurs when  $G$  acts transitively on the tree, and the spherical function equals  $X_p(g) c(g)$  where  $c : G \rightarrow \{\pm 1\}$  is the natural “color” representation corresponding to the possible coloring of the tree. If  $G$  does not act transitively on the Bruhat-Tits tree case 3 does not occur.

If the Bruhat-Tits building of  $G$  is not regular (i.e. is bi-regular) then case 3 give a new set of unitary spherical representations. Finding lower bounds for them requires direct computation. Let us do it in the contexts of bi-regular trees.

Let  $T$  be the Bruhat-Tits tree of  $G$ . Assume that  $T$  is  $(q_0 + 1, q_1 + 1)$ -regular, where  $q_0 > q_1$ . Let  $v_0$  be the vertex stabilized by  $K$ , which is  $q_0 + 1$  regular. The group  $G$  acts transitively on the vertices of even distance from  $v_0$ . It holds that  $q^{\rho(1)} = \sqrt{q_0 q_1}$  and for  $a \in A_+$ ,  $q^{l(a)} = (q_0 q_1)^{\nu(a)}$ . Since the tree is not regular  $l(a)$  does not give direct information about distances on the tree, but  $\nu(a)$  is half the natural distance on the tree between  $v_0$  and  $a \cdot v_0$ .

Using the identification  $a_{\mathbb{C}}^* \cong \mathbb{C}$ , we have  $\lambda : \mathbb{R} \rightarrow \mathbb{C}$  and let  $\theta = q^{(\rho-\lambda)(1)}$ . The relation between  $p(\pi)$  and  $\theta$  are  $|\theta| = (q_0 q_1)^{1-1/p}$ .




FIGURE 4.1. The values of  $\lambda(x)$  on the vertices of the Bruhat-Tits tree of  $G$  of even distance from  $v_0$ . Such vertices are marked by  $*$ .

The values of  $\theta$  which correspond to case 3 are for  $-q_0 < \theta < -q_1$  ([31, 35]).

Since  $\tilde{\lambda}$  and  $\varphi_\lambda$  are left  $K$  invariant, they define functions (also denoted  $\tilde{\lambda}, \varphi_\lambda$ ) on the vertices of even distance from  $v_0$ . The values of  $\tilde{\lambda}$  are shown in Figure 4.1.

Then if  $v_n$  is of distance  $2n$  from  $v_0$ , the value  $\varphi_\lambda(v_n)$  is the average of  $\tilde{\lambda}$  on the sphere of radius  $2n$  around  $v_0$ . Direct calculation gives

$$\begin{aligned}\varphi_\lambda(v_0) &= 1 \\ \varphi_\lambda(v_1) &= \frac{1}{(q_0 + 1)q_1} (\theta + (q_1 - 1) + q_0q_1\theta^{-1}) \\ \varphi_\lambda(v_2) &= \frac{1}{(q_0 + 1)q_1^2q_0} \\ &\quad \left( \theta^2 + (q_1 - 1)\theta + (q_0 - 1)q_1 + (q_1 - 1)q_0q_1\theta^{-1} + (q_0q_1\theta^{-1})^2 \right) \\ \varphi_\lambda(v_n) &= \frac{1}{(q_0 + 1)q_1(q_0q_1)^{n-1}} \\ &\quad \left( \theta^n + (q_1 - 1)\theta^{n-1} \sum_{k=1}^n (q_0q_1\theta^{-2})^{(k-1)} + (q_0 - 1)q_1\theta^{n-2} \sum_{k=1}^n (q_0q_1\theta^{-2})^{(k-1)} + (q_0q_1\theta^{-1})^n \right)\end{aligned}$$

For  $n$  even combine the sum of  $\varphi_\lambda(v_n)$  into pairs as follows:

$$\begin{aligned}\varphi_\lambda(v_n) &= \frac{1}{(q_0 + 1)q_1(q_0q_1)^{n-1}} \\ &\quad \left( \theta^n q_1^{-1} + (1 - q_1^{-1})(\theta^n + q_1\theta^{n-1}) + \sum_{k=2}^n \theta^{n-2} (q_0q_1\theta^{-2})^{(k-1)} ((q_0 - 1)q_1 + (q_1 - 1)q_0q_1\theta^{-1}) + (q_0q_1\theta^{-1})^n \right)\end{aligned}$$

Since  $-q_0 < \theta < -q_1$  and  $n$  is even we have  $\theta^n + q_1\theta^{n-1} > 0$  and  $(q_0 - 1)q_1 + (q_1 - 1)q_0q_1\theta^{-1} > 0$ . So in case 3 for  $n$  even,

$$\varphi_\lambda(v_n) \geq \frac{\theta^n q_1^{-1}}{(q_0 + 1)q_1(q_0q_1)^{n-1}}.$$

Returning to the group and applying the equalities  $|\theta| = (q_0q_1)^{1-1/p}$ ,  $q^{l(a)} = (q_0q_1)^{\nu(a)}$ , we get:

$$\varphi_\lambda(a) \gg q^{-l(a)/p}.$$

Combining all the different cases, we have:

**Proposition 4.16.** *Let  $G$  be non-Archimedean of rank 1,  $p > 2$  and  $(\pi, V)$  a unitary irreducible representation of  $G$  with  $p(\pi) = p$ , having a non-trivial  $K$ -fixed vector  $v$ . Then for  $a \in A_+$  with  $\nu(a) \in \mathbb{Z}$  even, we have  $\pi(A_a)v = \lambda_1 v$ , with*

$$q^{-l(a)/p} \ll \lambda_1.$$

We can finally conclude:

**Theorem 4.17.** *Let  $G$  be of rank 1. Then the set  $\Pi(G)_{sph,nt}$  of spherical non-tempered unitary representations has a good family of functions.*

*Proof.* For  $G$  Archimedean, we choose  $f_{d_0} = \chi_{d_0/2} * \chi_{d_0/2}$ . For  $G$  non-Archimedean, we choose  $a \in A_+$  with  $\nu(a) \in \mathbb{Z}$  even, and with  $l(a) = d_0/2 \pm O(1)$ , and choose  $f_{d_0} = A_a * A_a^*$ .

The first property of a good family follows from Proposition 4.15, Proposition 4.16 and the Convolution Lemma 4.18 below.

The second and third follows from simple properties of convolutions.  $\square$

**4.5. Convolution of Operators.** In this section we analyze the function  $\chi_{d_0} * \chi_{d_0}(g)$ . Similar analysis can be found for rank 1 in [53, Lemma 3.1]. Our analysis is less accurate, but is more abstract and works for every rank.

**Lemma 4.18** (Convolution Lemma). *It holds that  $c_{d_0} = \chi_{d_0} * \chi_{d_0} \in C_c(K \backslash G / K)$ , and satisfies the inequality*

$$c_{d_0}(g) \ll_{\epsilon} q^{d_0 \epsilon} \psi_{2d_0}(g).$$

*The same bound holds for the convolution of every  $f_1, f_2 \in C_c(K \backslash G / K)$  such that  $f_1(g), f_2(g) \ll_{\epsilon} q^{d_0 \epsilon} \chi_{d_0}(g)$ .*

*Proof.* The second statement is a consequence of the first, so we only need to prove the first statement.

The idea is to look at the action (by right convolution) of  $\chi_{d_0}$  on  $L^2(G)$ . By Lemma 4.11, and the same arguments as in Corollary 4.12, the norm of  $\chi_{d_0}$  on  $L^2(G)$  is bounded by  $\ll_{\epsilon} q^{d_0(1/2+\epsilon)}$ . Therefore the norm of  $c_{d_0}$  on  $L^2(G)$  is bounded by  $\ll_{\epsilon} q^{d_0(1+\epsilon)}$ . Now we need to use some continuity arguments to deduce pointwise bounds.

Notice that since  $\chi_{d_0} \in C_c(K \backslash G / K)$ , the same is true for  $c_{d_0}$ . In the non-Archimedean case the arguments are simpler- we look at the action of  $c_{d_0}$  on the characteristic function  $\mathbf{1}_K$  of  $K$ . Then

$$\|\mathbf{1}_K * c_{d_0}\|_{L^2(G)}^2 \ll_{\epsilon} q^{2d_0(1+\epsilon)} \|\mathbf{1}_K\|_{L^2(G)} = q^{2d_0(1+\epsilon)}.$$

But if  $c_{d_0}(g) = R$  then  $\mathbf{1}_K * c_{d_0}(g) = R$ , so

$$\|\mathbf{1}_K * c_{d_0}\|_{L^2(G)}^2 \geq \mu(KgK) R^2 \gg q^{l(g)} R^2.$$

Therefore  $c_{d_0}(g) = R \ll_{\epsilon} q^{d_0(1+\epsilon)-l(g)/2}$  as needed.

In the Archimedean case, assume that  $c_{d_0}(g) = R$ . Then  $c_{d_0+2}(g') \geq R$  for every  $g' \in G$  with  $|l(g) - l(g')| \leq 1$ . We consider  $\mathbf{1}_{B_1}$  where  $B_1$  is a the ball of radius 1 around the identity. It holds that

$$\|\mathbf{1}_{B_1} * c_{d_0+2}\|_{L^2(G)}^2 \ll_{\epsilon} q^{2d_0(1+\epsilon)} \|\mathbf{1}_{B_1}\|_{L^2(G)} \ll q^{2d_0(1+\epsilon)}.$$

It also holds that  $\mathbf{1}_{B_1} * c_{d_0+2}(g) \gg R$ , for  $g'$  satisfying  $|l(g) - l(g')| \leq 1$ , so

$$\|\mathbf{1}_K * c_{d_0+2}\|_{L^2(G)}^2 \gg \mu(KB_1gK) R^2 \gg q^{l(g)} R^2,$$

and  $c_{d_0}(g) = R \ll_{\epsilon} q^{d_0(1+\epsilon)-l(g)/2}$  as needed.  $\square$

**4.6. Traces of Operators on Irreducible Unitary Representations.** Our goal here in this section is to relate the lower and upper bounds of the previous sections to lower and upper bounds on traces.

Let us recall how to define traces of an operator  $h \in C_c(G)$  on a unitary irreducible representation  $(\pi, V)$ . We will only consider the case when  $h$  is right and left  $K$ -finite, which simplifies the theory. In such case there is an orthogonal projection  $e_h : C(K) \rightarrow C(K)$  such that  $h = e_h * h * e_h$  (see e.g. [13, proof of Theorem 2]). The orthogonal projection has a finite dimensional image, so it is supported on a finite number of  $K$ -types.

Since  $\pi$  is admissible, the image of  $\pi(h)$  is finite dimensional, and therefore have a trace ([39, Chapter X]), defined by

$$\mathrm{tr}\pi(h) = \sum_i \langle u_i, \pi(h) u_i \rangle,$$

where  $\{u_i\} \subset V$  is an orthonormal basis.

By uniform admissibility ([39, Theorem 10.2], [2]) and the fact the image of  $\pi(h)$  is supported on a finite number of  $K$ -types., the image of  $\pi(h)$  is of bounded dimension, depending only on the projection  $e_h$ .

Finally, recall that the norm of a finite dimensional operator is larger than the largest absolute value of an eigenvalue. We conclude:

**Proposition 4.19.** *Assume that  $h \in C_c(G)$  is right and left  $K$ -finite and  $(\pi, V) \in \Pi(G)$ . Then*

$$|\mathrm{tr}\pi(h)| \ll_{e_h} \|\pi(h)\|,$$

the bound depending only on the projection  $e_h$  such that  $e_h * h * e_h$ .

As an example, if  $h \in C_c(K \backslash G / K)$  is left and right  $K$ -invariant, the image of  $\pi(h)$  is of dimension 1 or 0. If the dimension is 0, obviously  $\mathrm{tr}\pi(h) = \pi(h) = 0$ . If the dimension is 1,  $V$  has a  $K$ -invariant vector  $v \in V$ ,  $\|v\| = 1$ , and

$$|\mathrm{tr}\pi(h)| = \|\pi(h)\| = |\langle v, \pi(h)v \rangle|.$$

One may also deduce lower bounds on traces of non-negative self-adjoint operators. It follows from the same considerations as in the finite dimensional case.

**Proposition 4.20.** *Assume that  $h \in C_c(G)$  is right and left  $K$ -finite and  $(\pi, V) \in \Pi(G)$ . Moreover assume that  $\pi(h)$  is self-adjoint and non-negative. Then*

$$\|\pi(h)\| = \sup_{v: \|v\|=1} \langle v, \pi(h)v \rangle \leq \mathrm{tr}\pi(h).$$

**4.7. The Pre-Trace formula.** Let us recall the simplest version of the trace formula, or the pre-trace formula ([23, Chapter 1]). Let  $h \in C_c(G)$  and let  $\Gamma \subset G$  be a cocompact lattice. Denote by  $\hat{h} \in C_c(G)$  the function  $\hat{h}(g) = h(g^{-1})$ . Then we have an operator



$h : L^2(\Gamma \backslash G) \rightarrow L^2(\Gamma \backslash G)$ , acting on  $f \in L^2(\Gamma \backslash G)$  by

$$\begin{aligned} (hf)(x) &= f * \hat{h}(x) = \int_G f(xg)h(g)dg = \int_G f(y)h(x^{-1}y)dy \\ &= \int_{\Gamma \backslash G} f(y) \left( \sum_{\gamma \in \Gamma} h(x^{-1}\gamma y) \right) dy = \int_{\Gamma \backslash G} K(x, y) f(y)dy, \end{aligned}$$

for  $K(x, y) = \sum_{\gamma \in \Gamma} h(x^{-1}\gamma y)$ . By [23, Chapter 1], if  $h$  is also self-adjoint then it has a trace on  $L^2(\Gamma \backslash G)$  and

$$\text{tr}h|_{L^2(\Gamma \backslash G)} = \int_{\Gamma \backslash G} \sum_{\gamma \in \Gamma} h(x^{-1}\gamma x) dx.$$

If moreover  $h \in C_c^\infty(G)$  and  $L^2(\Gamma \backslash G) \cong \bigoplus_{\pi \in \Pi(G)} m(\pi, \Gamma)$  is the decomposition into irreducible representations, then we have the pre-trace formula:

$$\begin{aligned} \text{tr}h|_{L^2(\Gamma \backslash G)} &= \int_{\Gamma \backslash G} \sum_{\gamma \in \Gamma} h(x^{-1}\gamma x) dx \\ &= \sum_{\pi \in \Pi(G)} m(\pi, \Gamma) \text{tr}\pi(h), \end{aligned}$$

where  $\text{tr}(\pi(h))$  is the usual trace on the representation space  $(\pi, V)$ .

The following lemma is immediate from the pre-trace formula but essential for our work.

**Lemma 4.21.** *If  $h_1, h_2 \in C_c(G)$  satisfy that  $h_1(g) \geq h_2(g)$  for every  $g \in G$  then  $\text{tr}h_1|_{L^2(\Gamma \backslash G)} \geq \text{tr}h_2|_{L^2(\Gamma \backslash G)}$ . In particular, if  $h_1(g) \geq 0$  then  $\text{tr}h_1|_{L^2(\Gamma \backslash G)} \geq 0$ .*

**4.8. Spectral Decomposition of a Characteristic Function of a Small Ball.** For  $x \in \Gamma_N \backslash \Gamma_1 \subset X_N = \Gamma_N \backslash G/K$ , let  $b_{x, \delta} \in L^2(X_N)$  be defined as follows:

- In the non-Archimedean case choose  $h_\delta \in C_c(K \backslash G/K)$  to be the characteristic function of  $K$ .
- For Archimedean case, choose  $\delta > 0$  such that  $l(\gamma) > 2\delta$  for every  $\gamma \in \Gamma_1$  such that  $l(\gamma) > 0$ . Choose a function  $h_\delta \in C_c^\infty(K \backslash G/K)$  such that:
  - $0 \leq h_\delta(g) \leq \frac{2}{\mu(B_\delta(e))}$  for all  $g \in G$ .
  - $h_\delta(g) = 0$  for  $l(g) > \delta$ .
  - $\int_G h_\delta(g)dg = 1$

Finally, let  $b_{x, \delta} \in L^2(X_N)$  be

$$b_{x, \delta}(y) = \sum_{\gamma \in \Gamma_N} h_\delta(x^{-1}\gamma y).$$

We notice that the sum is actually finite, and by the properties of  $\delta$  it is over at most  $|\Gamma_1 \cap K|$  elements. Therefore

$$\|b_{x, \delta}\|_2 \ll_{\delta, \Gamma_1} 1.$$

Let us remark that for our uses in the Archimedean rank 1 case one can simply choose

$$h_\delta(g) = \begin{cases} \frac{1}{\mu(B_\delta(e))} & l(g) \leq \delta \\ 0 & \text{else} \end{cases},$$

and for higher rank we make this choice so that one can apply the Paley-Wiener theorem for spherical functions due to Harish-Chandra, which is used below.

**Lemma 4.22.** *Let  $f \in L^2(X_N)$ . Then,*

$$\sum_{x \in \Gamma_N \setminus \Gamma_1} |\langle b_{x,\delta}, f \rangle|^2 \ll_{\Gamma_1, \delta} \|f\|_2^2.$$

*Moreover, in the Archimedean case, if  $f \in L^2(X_N)$  is the  $K$ -fixed function of some irreducible representation  $\pi \subset L^2(\Gamma_N \setminus G)$  with  $\lambda(\pi) = \lambda$ , then for every  $L' > 0$ ,*

$$\sum_{x \in \Gamma_N \setminus \Gamma_1} |\langle b_{x,\delta}, f \rangle|^2 \ll_{\Gamma_1, \delta, L'} (1 + \lambda)^{-L'} \|f\|_2^2.$$

*(The last result will only be used in the Archimedean rank  $\geq 2$  case).*

*Proof.* By our assumption on  $\delta$ , the balls  $B_\delta(x)$  for  $x \in \Gamma_N \setminus \Gamma_1$  are all either equal (with multiplicity at most  $|\Gamma_1 \cap K|$ ) or distinct, and each ball it holds by Cauchy-Schwartz that

$$\begin{aligned} \sum_{x \in \Gamma_N \setminus \Gamma_1} |\langle f, b_{x,\delta} \rangle|^2 &\leq \sum_{x \in \Gamma_N \setminus \Gamma_1} \|f|_{B_\delta(x)}\|_2^2 \|b_{x,\delta}\|_2^2 \leq |\Gamma_1 \cap K| \|f\|_2^2 \max_{x \in \Gamma_N \setminus \Gamma_1} \|b_{x,\delta}\|_2^2 \\ &\ll_{\delta, \Gamma_1} \|f\|_2^2 \end{aligned}$$

For the moreover part, note that

$$\langle f, b_{x,\delta} \rangle = f(x) \operatorname{tr} \pi(h_\delta).$$

It is well known (see [55] for an exact statement) that there is a constant  $M > 0$  such that

$$|f(x)| \ll_\delta (1 + \lambda)^M \|f|_{B_\delta(x)}\|_2^2.$$

By the Paley-Wiener theorem for spherical functions ([16, Subsection 3.4]),

$$\operatorname{tr} \pi(h_\delta) \ll_{L', \delta} (1 + \lambda)^{-L' - M}.$$

Combining both estimates we get the required inequality.  $\square$

## 5. THE WEAK INJECTIVE RADIUS PROPERTY

For most of this section, we will assume that  $\Gamma_1$  is cocompact, and will derive spectral results from the injective radius property. Later we will drop this assumption.

**5.1. Basic Results about the Weak Injective Radius Property.** Consider  $\chi_{d_0} \in C_c^\infty(G)$  from Section 3. It is self-adjoint since  $l(g) = l(g^{-1})$ . Since  $\chi_{d_0}$  is left and right  $K$ -invariant, it acts on  $L^2(\Gamma \setminus G/K)$ .

For  $x \in X = \Gamma \backslash G / K$  we denote

$$\mathbf{N}(\Gamma, d_0, x) = \# \{ \gamma \in \Gamma : l(x^{-1}\gamma x) \leq d_0 \},$$

which is an extension of the definition of  $\mathbf{N}(\Gamma, d_0, x)$  given in the introduction.

Note that by the definition of  $\mathbf{N}(\Gamma, d_0, x)$  it holds that for every  $x \in X$ ,

$$(5.1) \quad \mathbf{N}(\Gamma, d_0, x) \leq \sum_{\gamma \in \Gamma} \chi_{d_0}(x^{-1}\gamma x) \leq \mathbf{N}(\Gamma, d_0 + 1, x).$$

**Corollary 5.1** (Proof of Proposition 3.3). *The following are equivalent for a sequence  $\{\Gamma_N\}$  of cocompact lattices, and  $0 < \alpha \leq 1$ :*

(1) For every  $d_0 \leq 2\alpha \log_q([\Gamma : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$\text{tr} \chi_{d_0} |_{L^2(X_N)} = \int_{\Gamma_N \backslash G} \sum_{\gamma \in \Gamma} \chi_{d_0}(x^{-1}\gamma x) dx \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{d_0(1/2+\epsilon)}.$$

(2) The Weak Injective Radius Property with parameter  $\alpha$  — for every  $d_0 \leq 2\alpha \log_q([\Gamma_1 : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \int_{X_N} \mathbf{N}(\Gamma_N, d_0, x) dx \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon} q^{d_0(1/2+\epsilon)}.$$

(3) For every  $h \in C_c(G)$  self-adjoint and satisfying  $h(g) \ll_{\epsilon} [\Gamma : \Gamma_N]^{\epsilon} \psi_{\tilde{d}}(g)$  for  $\tilde{d} = 2\alpha \log_q([\Gamma : \Gamma_N])$ , it holds that

$$(5.2) \quad \text{tr} h |_{L^2(X_N)} \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{\tilde{d}(1/2+\epsilon)} \asymp [\Gamma_1 : \Gamma_N]^{1+\alpha+\epsilon}.$$

(4) For  $\tilde{d} = 2\alpha \log_q([\Gamma : \Gamma_N])$ , it holds that

$$\text{tr} \psi_{\tilde{d}} |_{L^2(X_N)} \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{\tilde{d}(1/2+\epsilon)} \asymp [\Gamma_1 : \Gamma_N]^{1+\alpha+\epsilon}.$$

*Proof.* The equivalence between (1) and (2) is a direct result of 5.1. The equivalence between (3) and (4) is a direct result of Lemma 4.21.

To show that (4) implies (1), note that for  $d_0 \leq \tilde{d} = 2\alpha \log_q([\Gamma : \Gamma_N])$ , it holds that  $\chi_{d_0} \ll q^{(d_0-\tilde{d})/2} \psi_{\tilde{d}}(g)$ . Then if (4) holds then

$$\begin{aligned} \text{tr} \chi_{d_0} |_{L^2(X_N)} &\ll q^{(d_0-\tilde{d})/2} \text{tr} \psi_{\tilde{d}} |_{L^2(X_N)} \\ &\ll q^{(d_0-\tilde{d})/2} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{\tilde{d}(1/2+\epsilon)} \\ &\ll [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{d_0(1/2+\epsilon)}. \end{aligned}$$

Finally, we prove that (1) implies (4). Note that for every  $g \in G$ ,

$$\psi_{\tilde{d}}(g) \ll \int_0^{\tilde{d}} q^{(\tilde{d}-d_0)/2} \chi_{d_0}(g) dd_0.$$

Then if (1) holds for every  $d_0 \leq \tilde{d}$  we have

$$\begin{aligned}
\mathrm{tr}\psi_{\tilde{d}} &\ll \int_0^{\tilde{d}} q^{(\tilde{d}-d_0)/2} \mathrm{tr}\chi_{d_0}|_{L^2(X_N)} dd_0 \\
&\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} \int_0^{\tilde{d}} q^{(\tilde{d}-d_0)/2} q^{d_0(1/2+\epsilon)} dd_0 \\
&= [\Gamma_1 : \Gamma_N]^{1+\epsilon} \tilde{d} q^{\tilde{d}(1/2+\epsilon)} \\
&\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{\tilde{d}(1/2+\epsilon)}.
\end{aligned}$$

□

## 5.2. Proof of Proposition 3.9.

*Proof.* Recall that we assume that the sequence  $\{\Gamma_N\}$  of cocompact lattices satisfies the Weak Injective Radius Property with parameter  $\alpha$ . Let  $d_0 = \alpha \log_q([\Gamma_1 : \Gamma_N])$  and let  $f_{d_0}$  from the definition of a good family. By Lemma 4.18 and Corollary 5.1,  $f_{d_0}$  has a trace on  $L^2(\Gamma_N \backslash G)$  and

$$\mathrm{tr}f_{d_0}|_{L^2(\Gamma_N \backslash G)} \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\alpha+\epsilon}.$$

Let us calculate the spectral side of the trace. From the second and first properties of a good family,

$$\begin{aligned}
\mathrm{tr}f_{d_0}|_{L^2(\Gamma_N \backslash G)} &\geq \sum_{\pi \in A} m(\pi, \Gamma_N) \mathrm{tr}\pi(f_{d_0}) \\
&\gg_{\epsilon, A} \sum_{\pi \in A} m(\pi, \Gamma_N) [\Gamma_1 : \Gamma_N]^{2\alpha(1-1/p(\pi)-\epsilon)} \\
&\gg_{\epsilon} M(A, \Gamma_N, p) [\Gamma_1 : \Gamma_N]^{2\alpha(1-1/p-\epsilon)}.
\end{aligned}$$

We deduce that for every  $N, p > 2, \epsilon > 0$ ,

$$M(A, \Gamma_N, p) \ll_{A, \epsilon} [\Gamma_1 : \Gamma_N]^{1-\alpha(1-2/p)+\epsilon},$$

as needed. □

**5.3. General Properties of the Weak Injective Radius.** In this subsection we prove some general results about the Weak Injective Radius Property.

**Proposition 5.2.** *Let  $\{\Gamma_N\}$  be a sequence of lattices. If  $\alpha$  is the Weak Injective Radius parameter of the sequence, then  $\alpha \leq 1$ .*

*Proof.* For simplicity we assume that the lattices  $\{\Gamma_N\}$  are cocompact so we can use Equation 5.2. The arguments can change to deal with the non-uniform case as well.

Note that Corollary 5.1 did not assume that  $\alpha \leq 1$ . It is therefore enough to prove that for every  $\delta > 0$  Equation 5.2 from Corollary 5.1 does not hold for  $\alpha = 1 + \delta$ .

Let  $d_0 = 2\alpha \log_q([\Gamma : \Gamma_N])$ ,  $d_1 = d_0/2$ , and  $c_{d_1} = \chi_{d_1} * \chi_{d_1}$ . By Lemma 4.18,  $c_{d_1} \ll_\epsilon [\Gamma_1 : \Gamma_N]^\epsilon \psi_{d_0}(g)$ , so the condition before Equation 5.2 holds, and

$$\mathrm{tr} c_{d_1}|_{L^2(X_N)} \ll_\epsilon [\Gamma_1 : \Gamma_N]^{2\alpha(1+\epsilon)}.$$

Since  $\chi_{d_1}$  is self-adjoint,  $\mathrm{tr} \pi(c_{d_1}) \geq 0$  for every  $\pi \in \Pi(G)$ , so

$$\mathrm{tr} c_{d_1}|_{L^2(X_N)} \leq \mathrm{tr}(\pi_{\mathrm{triv}}(c_{d_0})),$$

where  $\pi_{\mathrm{triv}}$  is the trivial representation. On the other hand, it holds that

$$\mathrm{tr}(\pi_{\mathrm{triv}}(c_{d_0})) = \int_G c_{d_1}(g) dg = \left( \int_G \chi_{d_1} dg \right)^2 dg \gg_\epsilon q^{2d_1(1-\epsilon)} = [\Gamma_1 : \Gamma_N]^{2\alpha(1-\epsilon)}.$$

As  $\alpha > 1$ , this contradicts Equation 5.2 for  $\epsilon > 0$  small enough and  $[\Gamma_1 : \Gamma_N]$  big enough.  $\square$

**Proposition 5.3.** *Let  $\{\Gamma_N\}$  be a sequence of lattices. Assume that the Weak Injective Radius Property holds with parameter  $\alpha = 1$ , i.e. for every  $0 \leq d_0 \leq 2 \log_q([\Gamma : \Gamma_N])$ ,  $\epsilon > 0$ ,*

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{y \in \Gamma_N \setminus \Gamma_1} \mathbf{N}(\Gamma_N, d_0, y) \ll_\epsilon [\Gamma_1 : \Gamma_N]^\epsilon q^{d_0(1/2+\epsilon)}.$$

*Then for every  $d_0 \geq 0, \epsilon > 0$  it holds that*

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{y \in \Gamma_N \setminus \Gamma_1} \mathbf{N}(\Gamma_N, d_0, y) \ll_\epsilon [\Gamma_1 : \Gamma_N]^\epsilon q^{d_0\epsilon} \left( \frac{q^{d_0}}{[\Gamma_1 : \Gamma_N]} + q^{d_0/2} \right).$$

*Proof.* It is enough to prove that for  $d_0 \geq 2 \log_q([\Gamma : \Gamma_N])$ ,

$$\sum_{y \in \Gamma_N \setminus \Gamma_1} \mathbf{N}(\Gamma_N, d_0, y) \ll_\epsilon q^{d_0(1+\epsilon)}.$$

First, we note that by the same arguments as above, for  $\delta > 0$  small enough,

$$\mathbf{N}(\Gamma, d_0 + 1, x) \ll_{\Gamma_1, \delta} \langle \chi_{d_0} b_{x, \delta}, b_{x, \delta} \rangle \ll_{\Gamma_1, \delta} \mathbf{N}(\Gamma, d_0 + 1, x).$$

Now, the fact that the Weak Injective Radius Property holds for  $\alpha = 1$  says that for  $\tilde{d} = \log_q([\Gamma_1 : \Gamma_N]) - 1$

$$\begin{aligned}
\sum_{x \in \Gamma_N \setminus \Gamma_1} \|\chi_{\tilde{d}} b_{x,\delta}\|_2^2 &= \sum_{x \in \Gamma_N \setminus \Gamma_1} \langle \chi_{\tilde{d}} b_{x,\delta}, \chi_{\tilde{d}} b_{x,\delta} \rangle \\
&= \sum_{x \in \Gamma_N \setminus \Gamma_1} \langle \chi_{\tilde{d}} * \chi_{\tilde{d}} b_{x,\delta}, b_{x,\delta} \rangle \\
&\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon} \sum_{x \in \Gamma_N \setminus \Gamma_1} \langle \psi_{2\tilde{d}} b_{x,\delta}, b_{x,\delta} \rangle \\
&\ll [\Gamma_1 : \Gamma_N]^{\epsilon} \sum_{x \in \Gamma_N \setminus \Gamma_1} \int_0^{2\tilde{d}} q^{(2\tilde{d}-d_0)/2} \langle \chi_{d_0} b_{x,\delta}, b_{x,\delta} \rangle dd_0 \\
&\ll [\Gamma_1 : \Gamma_N]^{\epsilon} \sum_{x \in \Gamma_N \setminus \Gamma_1} \int_0^{2\tilde{d}} q^{(2\tilde{d}-d_0)/2} \mathbf{N}(\Gamma, d_0 + 1, x) dd_0 \\
&\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} \int_0^{2\tilde{d}} q^{(2\tilde{d}-d_0)/2} q^{d_0/2} dd_0 \\
&\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} q^{\tilde{d}} \ll [\Gamma_1 : \Gamma_N]^{2+\epsilon}.
\end{aligned}$$

Now, for  $d_0 \geq 2\tilde{d} = 2(\log_q([\Gamma_1 : \Gamma_N]) - 1)$ , we use the bound

$$\chi_{d_0}(g) \ll (\chi_{\tilde{d}} * \chi_{d_0-2\tilde{d}+1} * \chi_{\tilde{d}})(g),$$

to deduce

$$\begin{aligned}
\sum_{x \in \Gamma_N \setminus \Gamma_1} \mathbf{N}(\Gamma, d_0, x) &\ll_{\delta} \sum_{x \in \Gamma_N \setminus \Gamma_1} \langle \chi_{d_0} b_{x,\delta}, b_{x,\delta} \rangle \\
&\ll \sum_{x \in \Gamma_N \setminus \Gamma_1} \langle \chi_{\tilde{d}} * \chi_{d_0-2\tilde{d}+1} * \chi_{\tilde{d}} b_{x,\delta}, b_{x,\delta} \rangle \\
&= \sum_{x \in \Gamma_N \setminus \Gamma_1} \langle \chi_{d_0-2\tilde{d}+1} * \chi_{\tilde{d}} b_{x,\delta}, \chi_{\tilde{d}} b_{x,\delta} \rangle \\
&\ll \sum_{x \in \Gamma_N \setminus \Gamma_1} \left\| \pi(\chi_{d_0-2\tilde{d}+1}) \right\| \|\chi_{\tilde{d}} b_{x,\delta}\|_2^2 \\
&\ll_{\epsilon} q^{(d_0-2\tilde{d}-1)(1+\epsilon)} \sum_{x \in \Gamma_N \setminus \Gamma_1} \|\chi_{\tilde{d}} b_{x,\delta}\|_2^2 \\
&\ll \frac{q^{d_0+\epsilon}}{[\Gamma_1 : \Gamma_N]^2} [\Gamma_1 : \Gamma_N]^{2+\epsilon} \ll q^{d_0+\epsilon}.
\end{aligned}$$

□

## 6. THE WEAK INJECTIVE RADIUS PROPERTY IMPLIES THE OPTIMAL LIFTING PROPERTY

In this Section we prove Theorem 1.5.

**6.1. Reduction to a Spectral Argument.** Recall that assuming the Weak Injective Radius Property and Spectral Gap, we should prove that for every  $\epsilon > 0$ , for every  $a \in A_+$  with  $l(a) \geq (1 + \epsilon) \log_q(\mu(X_N))$ ,

$$(6.1) \quad \#\left\{(x, y) \in (\Gamma_1/\Gamma_N)^2 : \exists \gamma \in \Gamma_1 \text{ s.t. } \pi_n(\gamma)x = y, \|a_\gamma - a\|_a < \epsilon \|a\|_a\right\} = (1 - o_\epsilon(1)) [\Gamma_1 : \Gamma_N]^2.$$

For  $(x, y) \in (\Gamma_1/\Gamma_N)^2$ ,  $a \in A_+$  and  $\epsilon > 0$ , we that  $\gamma \in \Gamma_1$  is *good for*  $(x, y, a, \epsilon)$  if  $\pi_n(\gamma)x = y$  and  $\|a_\gamma - a\|_a < \epsilon \|a\|_a$ .

**Lemma 6.1.** *Let  $(x, y) \in (\Gamma_1/\Gamma_N)^2$ , and assume that there is no good  $\gamma \in \Gamma_1$  for  $(x, y, a, \epsilon)$ . Identify  $x, y$  with elements  $x, y \in X_N = \Gamma_N \backslash G/K$ . Let  $f_a \in L^1(G)$  be a function supported on the set  $\{g \in G : \|a_g - a\|_a < \epsilon/2 \|a\|_a\}$ , and for  $\delta$  small enough with respect to  $\epsilon$  let  $b_{x,\delta}$  as in Subsection 4.8. Then*

$$f_a b_{x,\delta}(y) = 0.$$

Moreover, by choosing  $\delta$  small enough with respect to  $\epsilon$ , for every  $y' \in B_\delta(y)$  (i.e. in the ball of radius  $\delta$  around  $y$ ) it holds that

$$f_a b_{x,\delta}(y') = 0.$$

*Proof.* We think of  $f_a b_{x,\delta}$  as a  $\Gamma_N$ -invariant function on  $X = G$ , The support of  $f_a b_{x,\delta}$  is

$$\{y' \in G : \exists \gamma \in \Gamma_N, x' \in G, f_a(x'^{-1}\gamma y') > 0, d(x, x') < \delta\}.$$

Assume by contradiction that  $y' \in B_\delta(y)$  is in the support of  $f_a b_{x,\delta}$ , and let  $\gamma \in \Gamma_N$ ,  $x' \in G$  be such that  $f_a(x'^{-1}\gamma y') > 0$ ,  $d(x, x') < \delta$ . By the assumption on the support of  $f_a$ ,

$$\|a_\gamma - a\|_a < \epsilon/2 \|a\|_a.$$

Look at

$$\gamma' = x^{-1}\gamma y \in \Gamma_1.$$

Then  $a_{\gamma'} = a_{x^{-1}\gamma y}$ , and if  $\delta$  is small enough with respect to  $\epsilon$ , then  $\|a_{\gamma'} - a_\gamma\|_a < \epsilon/2 \|a\|_a$ . Therefore

$$\|a_{\gamma'} - a\|_a < \epsilon \|a\|_a,$$

and obviously  $\gamma'$  sends  $x \in \Gamma_N \backslash \Gamma_1$  to  $y \in \Gamma_N \backslash \Gamma_1$ .  $\square$

Let  $\pi \in L^2(X_N)$ , be the uniform probability distribution, i.e.,  $\pi(x) = \frac{1}{\mu(x_N)}$ .

**Lemma 6.2.** *The claim 6.1 holds if it holds that for every  $\epsilon > 0$ , for some  $\delta > 0$ , for every  $a \in A_+$  with  $l(a) > (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])$ , there exists a probability function  $f_a \in C_c^\infty(G)$*

supported on  $\{g \in G : \|a_g - a\|_a < \epsilon \|a\|_a\}$ , such that

$$(6.2) \quad \sum_{x \in \Gamma_1/\Gamma_N} \|f_a b_{x,\delta} - \pi\|_2^2 \stackrel{!}{=} o_{\epsilon,\delta}(1).$$

*Proof.* We assume 6.2 and prove that 6.1 holds.

Let  $\epsilon > 0$  be given and  $\delta > 0$  be small enough. For  $x, y \in (\Gamma_1/\Gamma_N)^2$ , by Lemma 6.1, if there is no good  $\gamma$  for  $(x, y, a, \epsilon/2)$ , then for  $y'$  in the  $\delta$ -neighborhood of  $y$  it holds that  $|(f_a b_{x,\delta} - \pi)(y')| = \pi(y') = \frac{1}{\mu(X_N)} \asymp_{X_1} [\Gamma_1 : \Gamma_N]^{-1}$ . Therefore, for a fixed  $x \in \Gamma_1/\Gamma_N$  each  $y$  without good  $\gamma$  contributes  $\gg_{\delta, X_1} [\Gamma_1 : \Gamma_N]^{-2}$  to  $\|f_a b_{x,\delta} - \pi\|_2^2$ .

Therefore,

$$\begin{aligned} & \# \left\{ (x, y) \in (\Gamma_1/\Gamma_N)^2 : \text{There is no good } \gamma \text{ for } (x, y, a, \epsilon/2) \right\} \\ & \ll_{\delta, X_1} \sum_{x \in \Gamma_1/\Gamma_N} \|f_a b_{x,\delta} - \pi\|_2^2 [\Gamma_1 : \Gamma_N]^2 \\ & = [\Gamma_1 : \Gamma_N]^2 \sum_{x \in \Gamma_1/\Gamma_N} \|f_a b_{x,\delta} - \pi\|_2^2 \\ & = o([\Gamma_1 : \Gamma_N]^2), \end{aligned}$$

where we used 6.2 in the last step. This implies 6.1 for  $\epsilon/2$ .  $\square$

The following lemma explains where Spectral Gap is used. Notice the difference between  $o_{\epsilon,\delta}(1)$  and  $O_{\epsilon,\delta,\epsilon_1}(q^{\epsilon_1 l(a)})$ .

**Lemma 6.3.** *Equation 6.1 follows from the following two conditions:*

- (1) *Spectral Gap holds for  $\{\Gamma_N\}$ .*
- (2) *For every  $\epsilon > 0$ , for some  $\delta > 0$ , for every  $\epsilon_1$ , for every  $a \in A_+$  with  $l(a) \geq (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])$ , there exists a probability function  $f_a \in C_c^\infty(G)$  supported on  $\{g \in G : \|a_g - a\|_a < \epsilon \|a\|_a\}$ , such that*

$$(6.3) \quad \sum_{x \in \Gamma_1/\Gamma_N} \|f_a b_{x,\delta}\|_2^2 \stackrel{!}{=} O_{\epsilon,\delta,\epsilon_1}(q^{\epsilon_1 l(a)}).$$

*Proof.* First we note that  $\|\pi\|_2^2 = \int_{X_N} \mu(X_N)^{-2} dx = \mu(X_N)^{-1} \asymp_{\Gamma_1} [\Gamma_1 : \Gamma_N]^{-1}$ . Therefore if Equation 6.3 holds then also

$$\sum_{x \in \Gamma_1/\Gamma_N} \|f_a b_{x,\delta} - \pi\|_2^2 \leq \sum_{x \in \Gamma_1/\Gamma_N} \|f_a b_{x,\delta}\|_2^2 + \sum_{x \in \Gamma_1/\Gamma_N} \|\pi\|_2^2 = O_{\epsilon,\delta,\epsilon_1}(q^{\epsilon_1 l(a)}),$$

which is similar to Equation 6.2, but  $o(1)$  is replaced with  $O_{\epsilon,\delta,\epsilon_1}(q^{\epsilon_1 l(a)})$ .

Let  $\epsilon' > 0$  and let  $f'_a = \chi_{\epsilon' l(a)} * f_a$ . Assuming  $\epsilon'$  is small enough,  $f'_a$  is supported on

$$\{g \in G : \|a_g - a\|_a < 2\epsilon \|a\|_a\}.$$

We will show that 6.2 holds for  $f'_a$ .



Notice that  $\chi_{\epsilon' \log_q([\Gamma_1 : \Gamma_N])} * \pi = \pi$  and  $f_a b_{x,\delta} - \pi \perp \pi$ . By the Spectral Gap assumption and Corollary 4.14, for some  $p' < \infty$ ,

$$\begin{aligned} \|f'_a b_{x,\delta} - \pi\|_2 &= \|\chi_{\epsilon' l(a)} (f_a b_{x,\delta} - \pi)\|_2 \\ &\ll q^{-p' \epsilon' l(a)} \|f_a b_{x,\delta} - \pi\|_2. \end{aligned}$$

Therefore,

$$\begin{aligned} \sum_{x \in \Gamma_1 / \Gamma_N} \|f'_a b_{x,\delta} - \pi\|_2^2 &\ll [\Gamma_1 : \Gamma_N]^{-2p' \epsilon'} \sum_{x \in \Gamma_1 / \Gamma_N} \|f_a b_{x,\delta} - \pi\|_2^2 \\ &\ll_{\epsilon, \delta, \epsilon_1} q^{-2p' \epsilon' l(a)} [\Gamma_1 : \Gamma_N]^{\epsilon_1}, \end{aligned}$$

and if we choose  $\epsilon_1$  small enough, then 6.2 holds. Applying Lemma 6.2, we get that 6.1 holds as well.  $\square$

**6.2. Completing the Proof of Theorem 1.5.** Recall that Theorem 1.5 states that Spectral Gap and the Weak Injective Radius Property implies the Optimal Lifting Property. In Lemma 6.3 we reduced it to some spectral statement, Equation 6.3. We now claim:

**Lemma 6.4.** *The Weak Injective Radius Property implies that for every  $\epsilon > 0$ , for some  $\delta > 0$ , for every  $a \in A_+$  with  $l(a) \geq \log_q([\Gamma_1 : \Gamma_N])$ , let  $f_a \in C_c^\infty(G)$  be a probability function on  $\{g \in G : \|a_g - a\|_a < \epsilon \|a\|_a\}$ , which is bounded by  $\ll_{\epsilon} q^{-l(a)} \asymp \mu(\{g \in G : \|a_g - a\|_a < \epsilon \|a\|_a\})^{-1}$ . Then*

$$\sum_{x \in \Gamma_1 / \Gamma_N} \|f_a b_{x,\delta}\|_2^2 \stackrel{!}{=} O_{\epsilon, \epsilon_1, \delta} \left( q^{-\epsilon_1 l(a)} \right).$$

The existence of such a probability function is a problems about smoothness and is obvious. Lemma 6.4 and Lemma 6.3 implies Theorem 1.5.

*Proof.* Let  $\epsilon > 0$ . We may choose  $\epsilon' > 0$  such that it holds that  $f_a(x) \ll q^{-(1+\epsilon')l(a)} \chi_{(1+\epsilon')l(a)}(x)$ , Therefore, by the Convolution Lemma 4.18,

$$\begin{aligned} \|f_a b_{x,\delta}\|_2^2 &\ll_{\epsilon} q^{-2(1+\epsilon')l(a)} \|\chi_{(1+\epsilon')l(a)} b_{x,\delta}\|_2^2 = q^{-2(1+\epsilon')l(a)} \langle \chi_{(1+\epsilon')l(a)} b_{x,\delta}, \chi_{(1+\epsilon')l(a)} b_{x,\delta} \rangle \\ &= q^{-2(1+\epsilon')l(a)} \langle \chi_{(1+\epsilon')l(a)} * \chi_{(1+\epsilon')l(a)} b_{x,\delta}, b_{x,\delta} \rangle \\ &\ll_{\epsilon_1} q^{-2(1+\epsilon'-\epsilon_1)l(a)} \langle \psi_{2(1+\epsilon')l(a)} b_{x,\delta}, b_{x,\delta} \rangle \\ &\ll q^{-2(1+\epsilon'-\epsilon_1)l(a)} \int_0^{2(1+\epsilon')l(a)} q^{(2(1+\epsilon')l(a)-d_0)/2} \langle \chi_{d_0} b_{x,\delta}, b_{x,\delta} \rangle d_0. \\ &\ll_{\delta} q^{-2(1+\epsilon'-\epsilon_1)l(a)} \int_0^{2(1+\epsilon')l(a)} q^{(2(1+\epsilon')l(a)-d_0)/2} N(x, \Gamma_N, d_0) d_0. \end{aligned}$$

Therefore, applying Proposition 5.3, we get

$$\begin{aligned} \sum_{x \in \Gamma_1 / \Gamma_N} \|f_a b_{x, \delta}\|_2^2 &\ll_{\delta, \epsilon, \epsilon_1} q^{-2(1+\epsilon'-\epsilon_1)l(a)} \int_0^{2(1+\epsilon')l(a)} q^{(2(1+\epsilon')l(a)-d_0)/2} \left( q^{d_0} + q^{d_0/2} [\Gamma_1 : \Gamma_N] \right) d_0 \\ &\ll_{\epsilon_1} q^{l(a)\epsilon_1} \left( 1 + q^{-l(a)/2} [\Gamma_1 : \Gamma_N] \right). \end{aligned}$$

Since  $l(a) \geq 2 \log_q([\Gamma_1 : \Gamma_N])$ , this is  $O(q^{l(a)\epsilon_1})$  as needed.  $\square$

## 7. THE SPECTRAL TO GEOMETRIC DIRECTION

**7.1. Some Technical Calculations.** For an ease of reference, we calculate here a couple of technical bounds.

**Lemma 7.1.** *Let  $G$  be non-Archimedean or rank 1. The following are equivalent for a sequence  $\{\Gamma_N\}$ :*

- (1) *The Spherical Density Hypothesis with parameter  $\alpha$ : for every  $\epsilon > 0$  and  $p > 2$*

$$M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p\right) \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1-\alpha(1-2/p)+\epsilon}.$$

- (2) *For every  $\epsilon > 0$*

$$\sum_{\pi \in \Pi(G)_{\text{sph}}, p(\pi) > 2} [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p(\pi_i))} m(\pi, \Gamma_N) \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon}.$$

*Proof.* The fact that (2) implies (1) is simple and left to the reader.

The fact that (1) implies (2) follows from a standard trick of integration by parts ([29, Theorem 421]):

$$\begin{aligned} &\sum_{\pi \in \Pi(G)_{\text{sph}}, p(\pi) > 2} [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p(\pi_i))} m(\pi, \Gamma_N) \\ &= \lim_{p_i \rightarrow 2, p_i > 2} M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p\right) [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p)} \\ &\quad + \int_2^{\infty} M(X_N, p) \frac{\partial}{\partial p} \left( [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p)} \right) dp \\ &= \lim_{p_i \rightarrow 2, p_i > 2} M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p\right) [\Gamma_1 : \Gamma_N]^{-1} \\ &\quad + \int_2^{\infty} M(X_N, p) 2p^{-2}\alpha \ln([\Gamma_1 : \Gamma_N]) [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p)} dp \\ &\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon} \left( 1 + \int_2^{\infty} p^{-2} dp \right) \asymp [\Gamma_1 : \Gamma_N]^{\epsilon}. \end{aligned}$$

□

For higher rank Archimedean we have:

**Lemma 7.2.** *Let  $G$  be Archimedean. The following are equivalent for a sequence  $\{\Gamma_N\}$ :*

- (1) *The Spherical Density Hypothesis with parameter  $\alpha$ : for some  $L > 0$  large enough and for every  $\lambda \geq 0$ ,  $N \geq 1$ ,  $p > 2$ ,  $\epsilon > 0$ ,*

$$M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p, \lambda\right) \ll_{\epsilon} (1 + \lambda)^L [\Gamma_1 : \Gamma_N]^{1 - \alpha(1 - 2/p) + \epsilon}$$

- (2) *For some  $L' > 0$  large enough and every  $\epsilon > 0$ ,*

$$\sum_{\pi \in \Pi(G)_{\text{sph}}, p(\pi) > 2} [\Gamma_1 : \Gamma_N]^{-1 + \alpha(1 - 2/p(\pi))} (1 + \lambda(\pi))^{-L'} m(\pi, \Gamma_N) \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon}.$$

*Proof.* The fact that (2) implies (1) is simple and is left to the reader.

The fact that (1) implies (2) is again done by integration by parts, with two variables. Let us state it formally. If  $((p_i, \lambda_i))_{i=1}^{\infty} \subset (2, \infty) \times (0, \infty)$  is a sequence of values without limit points,  $f(p, \lambda)$  is a non-negative smooth function, and  $M'(p, \lambda) = \#\{i : p_i \geq p, \lambda_i \leq \lambda\}$  then

$$\begin{aligned} \sum_i f(p_i, \lambda_i) &= \lim_{p \rightarrow 2} \lim_{\lambda \rightarrow \infty} M'(p, \lambda) f(p, \lambda) - \lim_{p \rightarrow 2} \int_0^{\infty} M'(p, \lambda) \frac{\partial}{\partial \lambda} f(p, \lambda) d\lambda \\ &\quad + \lim_{\lambda \rightarrow \infty} \int_2^{\infty} M'(p, \lambda) \frac{\partial}{\partial p} f(p, \lambda) dp \\ &\quad - \int_0^{\infty} \int_2^{\infty} M'(p, \lambda) \frac{\partial}{\partial \lambda} \frac{\partial}{\partial p} f(p, \lambda) dp d\lambda. \end{aligned}$$

Applying the integration by parts formula to the left hand side of (2), with  $M'(p, \lambda) = M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p, \lambda\right)$ ,  $f(p, \lambda) = [\Gamma_1 : \Gamma_N]^{-1 + \alpha(1 - 2/p)} (1 + \lambda)^{-L'}$ , we get

$$\begin{aligned} (7.1) \quad L.H.S &= \lim_{p \rightarrow 2} \lim_{\lambda \rightarrow \infty} M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p, \lambda\right) [\Gamma_1 : \Gamma_N]^{-1 + \alpha(1 - 2/p)} (1 + \lambda)^{-L'} \\ &\quad + \lim_{p \rightarrow 2} \int_0^{\infty} M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p, \lambda\right) [\Gamma_1 : \Gamma_N]^{-1 + \alpha(1 - 2/p)} L' (1 + \lambda)^{-L' - 1} d\lambda \\ &\quad + \lim_{\lambda \rightarrow \infty} \int_2^{\infty} M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p, \lambda\right) 2\alpha p^{-2} [\Gamma_1 : \Gamma_N]^{-1 + \alpha(1 - 2/p)} (1 + \lambda)^{-L'} dp \\ &\quad + \int_0^{\infty} \int_2^{\infty} M\left(\Pi(G)_{\text{sph}}, \Gamma_N, p, \lambda\right) 2\alpha p^{-2} [\Gamma_1 : \Gamma_N]^{-1 + \alpha(1 - 2/p)} L' (1 + \lambda)^{-L' - 1} dp d\lambda \end{aligned}$$

Note that by the spherical density, if we take  $p \rightarrow 2$ ,

$$M\left(\Pi(G)_{\text{sph}}, \Gamma_N, 2, \lambda\right) \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\epsilon} (1+\lambda)^{L''}.$$

Similar and more precise bound may be derived directly from Weyl's law ([16]).

Note also that we may assume that  $L' > L$ .

Therefore the first summand in Equation 7.1 is bounded by

$$\ll_{X_1, \epsilon} \lim_{\lambda \rightarrow 0} [\Gamma_1 : \Gamma_N]^{1+\epsilon} [\Gamma_1 : \Gamma_N] (1+\lambda)^{L''} (1+\lambda)^{-L'} = 0.$$

The second summand is bounded similarly by

$$\ll_{X_1, \epsilon} \int_0^{\infty} [\Gamma_1 : \Gamma_N]^{1+\epsilon} [\Gamma_1 : \Gamma_N]^{-1} (1+\lambda)^{L''} L' (1+\lambda)^{-L'-1} d\lambda \ll [\Gamma_1 : \Gamma_N]^{\epsilon},$$

The third summand is bounded by

$$\begin{aligned} &\ll_{\epsilon} \lim_{\lambda \rightarrow \infty} \int_2^{\infty} (1+\lambda)^L [\Gamma_1 : \Gamma_N]^{1-\alpha(1-2/p)+\epsilon} 2\alpha p^{-2} [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p)} (1+\lambda)^{-L'} dp \\ &= \lim_{\lambda \rightarrow \infty} (1+\lambda)^{L-L'} \int_2^{\infty} [\Gamma_1 : \Gamma_N]^{\epsilon} p^{-2} dp = 0, \end{aligned}$$

The final summand is bounded by

$$\begin{aligned} &\ll_{\epsilon} \int_0^{\infty} \int_2^{\infty} (1+\lambda)^L [\Gamma_1 : \Gamma_N]^{1-\alpha(1-2/p)+\epsilon} 2\alpha p^{-2} [\Gamma_1 : \Gamma_N]^{-1+\alpha(1-2/p)} L' (1+\lambda)^{-L'-1} dp d\lambda \\ &= [\Gamma_1 : \Gamma_N]^{\epsilon} \int_0^{\infty} L' (1+\lambda)^{L-L'-1} d\lambda \int_2^{\infty} p^{-2} dp \ll [\Gamma_1 : \Gamma_N]^{\epsilon}. \end{aligned}$$

Combining all the bounds we get (2).  $\square$

**7.2. Proof of Theorem 1.6.** We need to prove that the Spherical Density Hypothesis with parameter  $\alpha$  implies the Weak Injective Radius Property with parameter  $\alpha$ . The most natural proof of the claim is to analyze the spectral side of the trace formula for the function  $\chi_{d_0}$ . We will instead discretize and prove directly the Weak Injective Radius Property.

*Proof.* Recall that we should prove that for every  $d_0 \leq 2\alpha \log_q([\Gamma : \Gamma_N])$ ,  $\epsilon > 0$ ,

$$\frac{1}{[\Gamma_1 : \Gamma_N]} \sum_{x \in \Gamma_1/\Gamma_N} \mathbf{N}(\Gamma_N, d_0, x) dx \ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{\epsilon} q^{d_0(1/2+\epsilon)}.$$

For  $x \in \Gamma_N \setminus \Gamma_1$ , let  $b_{x, \delta} \in L^2(X_N)$  as in 4.8.

Notice that it holds for some  $C > 0$  that

$$\langle b_{x, \delta}, \chi_{d_0} b_{x, \delta} \rangle \ll_{\delta} \mathbf{N}(X_N, d_0, x) \ll_{\delta} \langle b_{x, \delta}, \chi_{d_0+C} b_{x, \delta} \rangle.$$

Let  $\{\pi_i\}_{i=1}^T$  be the set of irreducible subrepresentation of  $L^2(\Gamma_N \backslash G)$  with  $K$ -fixed vectors and  $p(\pi_i) > 2$  ( $T$  is finite in the  $p$ -adic or real rank 1 case, otherwise  $T$  may be  $\infty$ ). Recall that the set  $K$ -invariant vectors of each irreducible  $\pi_i$  is a one dimensional vector space. Let  $u_i \in L^2(X_N)$  be a  $K$ -invariant vector of  $\pi_i$  with  $\|u_i\| = 1$ . Let  $p_0 = 2$  and let  $V_0$  the orthogonal complement of  $\text{span}\{\pi\} \oplus (\oplus_i \text{span}\{u_i\})$  in  $L^2(X_N)$  ( $\pi$  here is the uniform probability function on  $X_N$ ). Note that the  $G$ -representation generated by  $V_0$  is (2-)tempered.

Decompose  $b_{x,\delta} = \pi + v_{0,x} + v_{1,x} + \dots$ , according to the decomposition  $L^2(X_N) = \text{span}\{\pi\} \oplus V_0 \oplus V_1 \oplus \dots$ , i.e for  $i = 1, 2, \dots$ ,  $v_{i,x} = \langle u_i, b_{x,\delta} \rangle u_i$ . Then,

$$(7.2) \quad \sum_{x \in \Gamma_1 / \Gamma_N} \langle b_{x,\delta}, \chi_{d_0+C} b_{x,\delta} \rangle = \sum_{x \in \Gamma_1 / \Gamma_N} \langle \pi, \chi_{d_0+C} \pi \rangle + \sum_{x \in \Gamma_1 / \Gamma_N} \langle v_{0,x}, \chi_{d_0+C} v_{0,x} \rangle + \sum_{x \in \Gamma_1 / \Gamma_N} \sum_{i=1}^T \langle v_{i,x}, \chi_{d_0+C} v_{i,x} \rangle.$$

The first summand in Equation 7.2 equals

$$\begin{aligned} \lambda_{\text{triv}}(\chi_{d_0+C}) [\Gamma_1 : \Gamma_N] \|\pi\|_2^2 &\ll_{\epsilon} q^{d_0(1+\epsilon)} [\Gamma_1 : \Gamma_N] \mu^{-1}(X_N) \ll_{X_1} q^{d_0(1+\epsilon)} \\ &\ll q^{d_0(1/2+\epsilon)} [\Gamma_1 : \Gamma_N], \end{aligned}$$

where  $\lambda_{\text{triv}}(\chi_{d_0+C})$  is the trivial eigenvalue of  $\chi_{d_0+C}$ , and we used the fact that  $d_0 \leq 2\alpha \log_q([\Gamma : \Gamma_N])$ .

Since  $V_0$  span a tempered representation, by Harish-Chandra's bounds the second summand in Equation 7.2 is bounded by

$$\ll_{\epsilon} q^{d_0(1/2+\epsilon)} \sum_{x \in \Gamma_1 / \Gamma_N} \|v_{0,x}\|_2^2 \ll q^{d_0(1/2+\epsilon)} [\Gamma_1 : \Gamma_N] \|b_{x,\delta}\|_2^2 \ll q^{d_0(1/2+\epsilon)} [\Gamma_1 : \Gamma_N].$$

To analyze the final summand in Equation 7.2, we first assume that  $G$  is  $p$ -adic or rank 1. By Lemma 4.22,  $\sum_{x \in \Gamma_1 / \Gamma_N} \|v_{i,x}\|_2^2 \ll_{\delta, \Gamma_1} 1$ . Write  $d_0 = 2\alpha' \log_q([\Gamma_1 : \Gamma_N])$ , for  $\alpha' \leq \alpha$ . Then

$$\begin{aligned} \sum_{x \in \Gamma_1 / \Gamma_N} \sum_{i=1}^T \langle v_{i,x}, \chi_{d_0+C} v_{i,x} \rangle &\ll_{\epsilon} \sum_{x \in \Gamma_1 / \Gamma_N} \sum_{i=1}^T q^{d_0(1-1/p(\pi_i)+\epsilon)} \|v_{i,x}\|_2^2 \\ &\ll_{\delta, \Gamma_1} \sum_{i=1}^T q^{d_0(1-1/p(\pi_i)+\epsilon)} \\ &\ll \sum_{\pi \in \Pi(G)_{\text{sph}}, p(\pi) > 2} m(\pi, \Gamma_N) q^{d_0(1-1/p(\pi)+\epsilon)} \\ &= \sum_{\pi \in \Pi(G)_{\text{sph}}, p(\pi) > 2} m(\pi, \Gamma_N) [\Gamma_1 : \Gamma_N]^{2\alpha'(1-1/p(\pi)+\epsilon)}. \end{aligned}$$

Applying Lemma 7.1 (for  $\alpha' \leq \alpha$ ) and arranging, we get

$$\ll_{\epsilon} [\Gamma_1 : \Gamma_N]^{1+\alpha'+\epsilon} = q^{d_0(1/2+\epsilon)} [\Gamma_1 : \Gamma_N].$$

For the Archimedean high-rank case, by Lemma 4.22, for  $L$  large enough

$$\sum_{x \in \Gamma_1/\Gamma_N} \|v_{i,x}\|_2^2 \ll_{\Gamma_1,\delta,L} (1 + \lambda(\pi_i))^{-L}.$$

The rest of the argument is as above, but using Lemma 7.2 instead of Lemma 7.1.  $\square$

*Remark 7.3.* The proof of Theorem 1.6 for hyperbolic spaces actually works for non-compact quotients as well. The reason is that the entire continuous spectrum is contained in  $V_0$  ([42]). For more details for hyperbolic surfaces, see [27].

### 7.3. A strong version of Theorem 1.5, assuming the Spherical Density Hypothesis.

The goal of this Subsection is to prove the following theorem:

**Theorem 7.4.** *Let  $\{\Gamma_N\}$  be a sequence satisfying the Spherical Density Hypothesis (with parameter  $\alpha = 1$ ) and Spectral Gap. Then for every  $\epsilon > 0$ , for every  $a \in A_+$  with  $l(a) \geq (1 + \epsilon) \log_q(\mu(X_N))$ ,*

$$\#\left\{(x, y) \in (\Gamma_1/\Gamma_N)^2 : \exists \gamma \in \Gamma_1 \text{ s.t. } \pi_n(\gamma)x = y, \|a_\gamma - a\|_a < \epsilon\right\} = (1 - o_\epsilon(1)) [\Gamma_1 : \Gamma_N]^2.$$

The result of Theorem 7.4 is stronger than in Theorem 1.5 – here we determine the  $A_+$ -component of the Cartan decomposition of  $\gamma$  in a far greater precision. In the non-Archimedean case it says that we may choose  $a \in A_+$  precisely.

As before, let  $\pi \in L^2(X_N)$ , be the uniform probability distribution, i.e.,  $\pi(x) = \frac{1}{\mu(x_N)}$ .

**Lemma 7.5.** *To prove Theorem 7.4, it suffices to prove under its assumptions that for  $a \in A_+$  with  $l(a) > (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])$*

$$(7.3) \quad \sum_{x \in \Gamma_1/\Gamma_N} \|A_a b_{x,\delta} - \pi\|_2^2 \stackrel{!}{=} o_{\epsilon,\delta}(1).$$

*Proof.* This is the same argument as in Lemma 6.2.

Let us first show that Equation 7.3 is immediate if  $M(X_N, p) = 0$  for  $p > 2$  (the Ramanujan case). If  $l(a) > (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])$  and then  $q^{-l(a)} \ll [\Gamma_1 : \Gamma_N]^{-(1+\epsilon)}$ . Then if we apply Corollary 4.14, for  $\epsilon'$  sufficiently small,

$$(7.4) \quad \begin{aligned} \|A_a b_{x_0,\delta} - \pi\|_2^2 &= \|A_a (b_{x_0,\delta} - \pi)\|_2^2 \ll_{\epsilon'} q^{-l(a)(1+\epsilon')} \|b_{x_0,\delta}\|_2^2 \\ &\ll [\Gamma_1 : \Gamma_N]^{-(1+\epsilon)(1-\epsilon')} = o([\Gamma_1 : \Gamma_N]^{-1}). \end{aligned}$$

Summing over  $x \in \rho_N^{-1}(x_0)$  we get the required bound. As a matter of fact, we proved that if  $M(X_N, p) = 0$  for  $p > 2$  then for every  $x \in \Gamma_1/\Gamma_N$

$$\#\{y \in \Gamma_1/\Gamma_N : \exists \gamma \in \Gamma_1 \text{ s.t. } \pi_n(\gamma)x = y, \|a_\gamma - a\|_a < \epsilon\} = (1 - o_\epsilon(1)) [\Gamma_1 : \Gamma_N].$$

The proof in the general case is basically the same as the proof of Theorem 1.6 in the previous subsection, and we use the same notations.

Let us quickly give the differences in the proofs.

The decomposition of  $b_{x,\delta}$  is the same, but instead of bounding

$$\sum_{x \in \Gamma_1/\Gamma_N} \langle b_{x,\delta}, \chi_{d_0+C} b_{x,\delta} \rangle,$$

we bound for  $l(a) > (1 + \epsilon) \log_q([\Gamma_1 : \Gamma_N])$

$$\sum_{x \in \Gamma_1/\Gamma_N} \|A_a(b_{x_0,\delta} - \pi)\|_2^2.$$

We get in the same way

$$(7.5) \quad \sum_{x \in \Gamma_1/\Gamma_N} \|A_a(b_{x_0,\delta} - \pi)\|_2^2 \ll \sum_{x \in \Gamma_1/\Gamma_N} \|A_a v_{0,x}\|_2^2 + \sum_{x \in \Gamma_1/\Gamma_N} \sum_{i=1}^T \|A_a v_{i,x}\|_2^2.$$

Instead of using the bound on  $\chi_{d_0}$ , we use the bound on  $A_a$ , which is very similar.

The first summand of 7.5 is bounded, for  $\epsilon'$  small enough, by

$$\begin{aligned} \sum_{x \in \Gamma_1/\Gamma_N} \|A_a v_{0,x}\|_2^2 &\ll_{\epsilon'} \sum_{x \in \Gamma_1/\Gamma_N} q^{-l(a)(1-\epsilon')} \|v_{0,x}\|_2^2 \\ &\ll_{\Gamma_1,\delta} [\Gamma_1 : \Gamma_N]^{-(1+\epsilon)(1-\epsilon')} [\Gamma_1 : \Gamma_N] \\ &= o(1). \end{aligned}$$

The second summand of 7.5 is bounded in the  $p$ -adic or rank 1 case by

$$\begin{aligned} \sum_{x \in \Gamma_1/\Gamma_N} \sum_{i=1}^T \|A_a v_{i,x}\|_2^2 &\ll_{\epsilon'} \sum_{x \in \Gamma_1/\Gamma_N} \sum_{i=1}^T q^{-l(a)(2/p(\pi_i)-\epsilon')} \|v_{i,x}\|_2^2 \\ &\ll_{\Gamma_1,\delta} \sum_{i=1}^T [\Gamma_1 : \Gamma_N]^{-(1+\epsilon)(2/p(\pi_i)-\epsilon')} [\Gamma_1 : \Gamma_N] \\ &\ll [\Gamma_1 : \Gamma_N]^{-\epsilon(2/p'-\epsilon')} \sum_{\pi \in \Pi(G)_{\text{sph}, p(\pi) > 2}} m(\pi, \Gamma_N) [\Gamma_1 : \Gamma_N]^{(1-2/p(\pi)+\epsilon')}, \end{aligned}$$

where  $p'$  satisfies  $p(\pi_i) \leq p'$ , by the spectral gap assumption. Using Lemma 7.1 we get for  $\epsilon'$  small enough

$$\sum_{x \in \Gamma_1/\Gamma_N} \sum_{i=1}^T \|A_a v_{i,x}\|_2^2 \ll_{\epsilon'} [\Gamma_1 : \Gamma_N]^{-\epsilon(2/p'-\epsilon')} [\Gamma_1 : \Gamma_N]^{\epsilon'} = o(1),$$

as needed.

The proof of the Archimedean high rank case is also similar.  $\square$

## 8. BERNSTEIN THEORY OF NON-BACKTRACKING OPERATORS

The results of this Section are the main technical contribution of this work. We base our work on the results of [2]. In this section  $G$  is assumed to be over a *non-Archimedean* field.

Let us first discuss some general results which follow from Harish-Chandra's bounds. Let  $K' \subset K$  be a compact open subgroup. For  $g \in G$  it holds that  $\mu(KgK)[K : K']^{-2} \leq \mu(K'gK') \leq \mu(KgK)$ , and therefore  $\mu(K'gK') \asymp_{K'} \mu(KgK) \asymp q^{l(g)}$ .

Consider the Hecke algebra  $H_{K'} = \mathbb{C}_c(K' \backslash G / K')$ . For  $g \in K' \backslash G / K'$ , denote  $h_{K',g} = \frac{1}{\mu(K')} K'gK'$ , and  $q_{K',g} = \mu(K'gK') \mu^{-1}(K')$ . As  $q_{K',g}$  is the number of right (or left)  $K'$  cosets in  $K'gK'$ , it is a natural number, and  $q_{K',g} \asymp_{K'} q^{l(g)}$ . As is well known by representation theory of  $p$ -adic groups ([9]), given a representation  $V$  of  $G$ , the Hecke algebra  $H_{K'}$  acts on the  $K'$ -fixed vectors  $V^{K'}$  of  $V$ . Bernstein's Decomposition of Elements in the Hecke Algebra

As a special case, we first discuss the Iwahori-Hecke algebra. Let  $I \subset K$  be the Iwahori-Hecke subgroup, i.e. the pointwise stabilizer of a chamber in the Bruhat-Tits building of  $G$ . Let  $W$  be the affine Weyl group of the root system of  $G$  (relative to the maximal  $k$ -torus  $T$ ) and  $\hat{W}$  the extended affine Weyl group. By the Iwahori decomposition we have  $G = I\tilde{W}I$ , where  $W \subset \tilde{W} \subset \hat{W}$ . For  $w \in \tilde{W}$ , denote  $q_w = \mu(IwI) / \mu(I)$  which is a natural number. Let  $H = C_c(I \backslash G / I)$  be the Iwahori-Hecke algebra of  $G$  and  $h_w \in H$  be the element  $\frac{1}{\mu(I)} IwI$ .

Let  $\beta_1, \dots, \beta_r \in \tilde{W}$  ( $r = \text{rank}G$ ) be some fixed multiples of the simple coweights of the root system of  $W$  (the simple coweight themselves belong to  $\hat{W}$ , so we cannot use them). Then  $h_{\beta_i}$  satisfies that  $h_{\beta_i}^m = h_{\beta_i^m}$ , i.e.  $h_{\beta_i}$  is a "non-backtracking operator" (when acting on the building  $\mathcal{B}$  of  $G$ , it is indeed a non-backtracking operator, or "collision free" in the notions of [43]). Then it holds that:

**Theorem 8.1** (See [34, Theorem 22.1]). *There exist two finite sets  $A, B \subset \tilde{W}$  such that each  $w \in \tilde{W}$  can be written uniquely as  $w = a\beta_1^{m_1} \cdot \dots \cdot \beta_r^{m_r}b$  with  $a \in A$ ,  $b \in B$ ,  $m_i \geq 0$ , and moreover  $l_{\tilde{W}}(w) = l_{\tilde{W}}(a) + \sum_{i=1}^r m_i l_{\tilde{W}}(\beta_i) + l_{\tilde{W}}(b)$ , where  $l_{\tilde{W}} : \tilde{W} \rightarrow \mathbb{N}$  is the length of the Coxeter group  $\tilde{W}$ .*

As a corollary, it holds that in the Iwahori-Hecke algebra,

$$h_w = h_a h_{\beta_1}^{m_1} \cdot \dots \cdot h_{\beta_r}^{m_r} h_b.$$

Let us now generalize. Let  $K' \subset G$  be a compact open subgroup. Consider the Hecke algebra  $H_{K'} = C_c(K' \backslash G / K')$ . For  $g \in K' \backslash G / K'$ , denote  $h_{K',g} = \frac{1}{\mu(K')} K'gK'$ ,  $q_{K',g} = \mu(K'gK') \mu^{-1}(K')$ .

Then it follows from the results of [2] (see also [3, Chapter II, Section 2]) that:

**Theorem 8.2** (Bernstein's Decomposition). *There exist arbitrarily small compact open subgroups  $K'$  such that for each  $K'$  there exist two finite sets  $A, B \subset K' \backslash G / K'$  and  $\beta_1, \dots, \beta_r \in K' \backslash G / K'$ ,  $r = \text{rank}_k G$ , such that:*

1. For each  $1 \leq i \leq r$  and  $m \geq 0$ ,  $h_{K',\beta_i}^m = h_{K',\beta_i^m}$ .
2. The operators  $h_{K',\beta_i}$ ,  $1 \leq i \leq r$ , commute.
3. For each  $g \in G$  there exist  $b \in B$ ,  $a \in A$  and  $m_i \geq 0$  such that

$$(8.1) \quad h_{K',g} = h_{K',a} h_{K',\beta_1}^{m_1} \cdot \dots \cdot h_{K',\beta_r}^{m_r} h_{K',b}.$$



*Remark 8.3.* Equation 8.1 is equivalent to the double coset decomposition

$$K'gK' = K'aK'\beta_1^{m_1}K' \cdots K'\beta_r^{m_r}K'bK' = K'a\beta_1^{m_1} \cdots \beta_r^{m_r}bK'.$$

Unlike in Theorem 8.2, there is no uniqueness in the claim.

*Proof.* We follow [3, Chapter II, Section 2]. Start from the Cartan decomposition  $KA_+K$ , where  $A_+ \subset P$  are the dominant elements in the lattice  $M/(M \cap K)$  ( $A_+$  is denoted  $\Lambda_+$  in [3]). Let  $\beta_1, \dots, \beta_r \in A_+$  be generators of a free semigroup  $\tilde{A}_+$  in  $A_+$  (they may be chosen so that their lift to  $M$  commute, not just as elements in  $M/(M \cap K)$ ). Let  $\mu_1, \dots, \mu_l \in A_+$  be elements such that  $A_+ = \cup_{i=1}^l \tilde{A}\mu_i$ , the union being disjoint.

By Bruhat's Theorem in [3], there exists arbitrary small compact open subgroups  $K' \subset K$  such that:

- $K'$  is normal in  $K$ .
- For every  $a, b \in A_+$  it holds that  $h_{K',a}h_{K',b} = h_{K',ab}$ , i.e.

$$(8.2) \quad K'aK'bK' = K'abK'.$$

Let  $x_1, \dots, x_m \in K$  be representatives of right cosets of  $K'$  in  $K$ . Since  $K'$  is normal in  $K$  they are also representatives of left cosets. Let  $A = \{x_1, \dots, x_r\}$  and let  $B = \{\mu_i x_j : i = 1, \dots, l, j = 1, \dots, r\}$ . By the Cartan decomposition for each  $g \in G$  there exists  $x \in A$ ,  $\mu x' \in B$  and  $m_1, \dots, m_r$  such that  $K'gK' = K'x\beta_1^{m_1} \cdots \beta_r^{m_r}\mu x'K'$ . It remains to show that

$$(8.3) \quad K'xK'\beta_1^{m_1}K' \cdots \beta_r^{m_r}K'\mu x'K' = K'x\beta_1^{m_1} \cdots \beta_r^{m_r}\mu x'K'.$$

Since  $\beta_1, \dots, \beta_r, \mu \in A_+$ ,  $K'\beta_1^{m_1}K' \cdots \beta_r^{m_r}K'\mu K' = K'\beta_1^{m_1} \cdots \beta_r^{m_r}\mu K'$  by Equation 8.2. Finally, since  $K'$  is normal in  $K$  and  $x, x' \in K$ ,  $K'x = xK'$  and  $K'x' = x'K'$ . Applying those equalities we get 8.3.  $\square$

**8.1. Non-Backtracking Operators and Temperedness.** We may now deduce:

**Theorem 8.4** (See also [34] for representations with a non-trivial Iwahori-invariant vector). *Let  $(\pi, V)$  be a unitary irreducible representation of  $G$ ,  $V^{K'}$  the  $K'$ -fixed vectors and  $V^{K'} \neq \{0\}$ . Consider the action of  $H_{K'} = C_c(K' \backslash G / K')$  on  $V^{K'}$ . Then  $\pi$  is  $p$ -tempered if and only if, for every  $1 \leq i \leq r$ , for every eigenvalue  $\lambda$  of  $\pi(h_{K',\beta_i})$  on  $V^{K'}$  it holds that  $|\lambda| \leq q_{K',\beta_i}^{1-1/p}$ .*

*Proof.* First, note that  $V$  is  $p$ -tempered if and only if for every  $0 \neq v_0 \in V^{K'}$  and  $p' > p$ ,

$$\begin{aligned} \int_G |\langle v_0, \pi(g) \cdot v_0 \rangle|^{p'} dg &= \sum_{g \in [K' \backslash G / K']} \mu(K'gK') \left( \frac{|\langle v_0, \pi(K'gK') v_0 \rangle|}{\mu(K'gK')} \right)^{p'} \\ &= \sum_{g \in [K' \backslash G / K']} (\mu(K'gK'))^{1-p'} (\mu(K') |\langle v_0, \pi(h_{K',g}) v_0 \rangle|)^{p'} \\ &\asymp \sum_{g \in [K' \backslash G / K']} (q_{K',g})^{1-p'} (|\langle v_0, \pi(h_{K',g}) v_0 \rangle|)^{p'} < \infty. \end{aligned}$$

The only if part is easier and does not require Bernstein's decomposition. For  $p = 2$  it is essentially the main result of [43].

Assume that some eigenvalue  $\lambda$  of  $h_{K',\beta_i}$  satisfies  $|\lambda| > q_{K',\beta_i}^{1-1/p}$ . Let  $v_0 \in V^K$  be an eigenvector of  $h_{K',\beta_i}$  with eigenvalue  $\lambda$ . Then, for  $p' > p$  such that  $|\lambda| \geq q_{K',\beta_i}^{1-1/p'}$ ,

$$\begin{aligned} \int_G |\langle v_0, \pi(g) \cdot v_0 \rangle|^{p'} dg &\gg \sum_{m=0}^{\infty} (q_{K',\beta_i^m})^{1-p} (|\langle v_0, \pi(h_{K',\beta_i^m}) v_0 \rangle|)^p \\ &= \sum_{m=0}^{\infty} (q_{K',\beta_i})^{m(1-p)} (|\langle v_0, \pi(h_{K',\beta_i})^m v_0 \rangle|)^p \\ &= \sum_{m=0}^{\infty} (q_{K',\beta_i})^{m(1-p)} (|\lambda|^m \|v_0\|^2)^p \\ &\geq \|v_0\|^{2p} \sum_{m=0}^{\infty} (q_{K',\beta_i})^{m((1-p)+p(1-1/p))} = \|v_0\|^{2p} \sum_{m=0}^{\infty} 1 = \infty, \end{aligned}$$

and  $V$  is not  $p$ -tempered.

We now prove the if part. One should prove that for  $p' > p$  matrix coefficients are in  $L^{p'}(G)$ . By Bernstein decomposition,

$$\begin{aligned} \int_G |\langle v_0, \pi(g) \cdot v_0 \rangle|^{p'} dg &\asymp \sum_{g \in [K' \backslash G / K']} (q_{K',g})^{1-p'} (|\langle v_0, \pi(h_{K',g}) v_0 \rangle|)^{p'} \\ &\leq \sum_{a \in A, b \in B} \sum_{m_1 \geq 0, \dots, m_r \geq 0} \left( q_{K',a} q_{K',\beta_1}^{m_1} \cdots q_{K',\beta_r}^{m_r} q_{K',b} \right)^{1-p'} \\ &\quad \cdot \left( \left| \langle \pi(h_{K',a})^* v_0, \pi(h_{K',\beta_1}^{m_1} \cdots h_{K',\beta_r}^{m_r}) \pi(h_{K',b}) v_0 \rangle \right| \right)^{p'} \\ &\ll_{K'} \sum_{a \in A, b \in B} \|\pi(h_{K',a})\| \|\pi(h_{K',b})\| \sum_{m_1 \geq 0, \dots, m_r \geq 0} \left( q_{K',\beta_1}^{m_1} \cdots q_{K',\beta_r}^{m_r} \right)^{1-p'} \\ &\quad \cdot \left\| \pi(h_{K',\beta_1}^{m_1} \cdots h_{K',\beta_r}^{m_r}) \right\|^{p'} \\ &\ll_{K'} \left( \sum_{m_1 \geq 0} q_{K',\beta_1}^{m_1(1-p')} \left\| \pi(h_{K',\beta_1}^{m_1}) \right\|^{p'} \right) \left( \sum_{m_2 \geq 0} q_{K',\beta_2}^{m_2(1-p')} \left\| \pi(h_{K',\beta_2}^{m_2}) \right\|^{p'} \right) \\ &\quad \cdots \left( \sum_{m_r \geq 0} q_{K',\beta_r}^{m_r(1-p')} \left\| \pi(h_{K',\beta_r}^{m_r}) \right\|^{p'} \right). \end{aligned}$$

Since all the eigenvalues of  $\pi(h_{K',\beta_i})$  are bounded by  $q_{K',\beta_i}^{1-1/p}$  it follows from the theory of matrix norms that the sum converges (note that  $\pi(h_{K',\beta_i})$  is usually not unitary or self-adjoint, so one should be a bit careful here).  $\square$

For  $d_0 \in \mathbb{R}_{\geq 0}$ , choose  $m_i = \frac{\log_q d_0}{2 \log_q \beta_i}$ , so  $q_{\beta_i}^{m_i} = d_0/2$  and denote  $f_{d_0} = \sum_{i=1}^r \left(h_{\beta_i}^{m_i}\right)^* h_{\beta_i}^{m_i}$ . The following theorem proves Theorem 3.11

**Theorem 8.5.** *The functions  $f_{d_0} \in H_{K'}$  are a good family for the set  $\Pi(G)_{K'\text{-sph}}$ :*

(1) *For every  $(\pi, V) \in \Pi(G)$  with a non-trivial  $K'$ -invariant vector, it holds that*

$$q^{d_0(1-1/p(\pi))} \ll_{K'} \text{tr}(\pi(f_{d_0})).$$

(2) *For every  $(\pi, V) \in \Pi(G)$  without a non-trivial  $K'$ -invariant vector, it holds that*

$$\text{tr}(\pi(f_{d_0})) = 0.$$

(3) *It holds that  $f_{d_0}(g) \ll_{\epsilon} q^{d_0 \epsilon} \psi_{d_0}(g)$ .*

*Proof.* The second part is obvious, since if there is non non-trivial  $K'$ -invariant vector then  $\pi(f_{d_0}) = 0$ .

The fact that  $f_{d_0}(g) \ll_{\epsilon} q^{d_0 \epsilon} \psi_{d_0}(g)$  follows from Lemma 4.18, as  $h_{\beta_i}^{m_i}(g) \ll \chi_{d_0/2}$ .

It remains to prove the first part. Since  $\pi(f_{d_0})$  is non-negative and self-adjoint it is diagonalizable on  $V^{K'}$  and all its eigenvalues are non-negative.

On the other hand, by uniform admissibility ([2]),  $\dim V^{K'} \ll_{K'} 1$ , so  $\text{tr} \pi(f_{d_0}) \asymp \|\pi(f_{d_0})\|$ . It holds that

$$\max_{i=1}^r \left\| h_{\beta_i}^{m_i} \right\|^2 \leq \|\pi(f_{d_0})\|.$$

Since  $V$  is not  $p'$ -tempered for  $p' < p$ , by Theorem 8.4, for some  $1 \leq i \leq r$  we have an eigenvalue  $\lambda$  of  $h_{\beta_i}$  with  $|\lambda| \geq q_{K', \beta_i}^{1-1/p}$ . Therefore,  $h_{\beta_i}^{m_i}$  has an eigenvalue  $\lambda^{m_i}$  with  $|\lambda^{m_i}| \geq q_{K', \beta_i}^{m_i(1-1/p)} \asymp_{K'} q^{l(\beta_i)m_i(1-1/p)} \asymp q^{d_0/2(1-1/p)}$ . Therefore  $\|\pi(f_{d_0})\|_{V^{K'}} \geq \left\| h_{\beta_i}^{m_i} \right\|_{V^{K'}}^2 \gg q^{d(1-1/p)}$ , as needed.  $\square$

## 9. LOWER BOUNDS ON MATRIX COEFFICIENTS FOR A SPECIFIC REPRESENTATION

In this section we prove Theorem 3.10. We assume that  $G$  is Archimedean and  $(\pi, V)$  is an irreducible unitary representation of  $G$ .

We will prove that  $\{\pi\}$  has a good family of functions. Let  $v \in V^{\tau}$ ,  $\|v\| = 1$ , belong to a fixed  $\tau$ -type of  $K$ , i.e.  $K$  acts on  $U_0 = \text{span}\{Kv\}$  as a  $K$ -irreducible representation  $\tau$ . In particular,  $v$  is  $K$ -finite. Let

$$g_{d_0/2}(g) = \chi_{d_0/2}(g) q^{l(g)p(\pi)-1} \overline{\langle v, \pi(g)v \rangle},$$

and let

$$f_{d_0} = g_{d_0/2}^* * g_{d_0/2}.$$

We first verify (2) and (3) of Definition 3.7. Since  $f_{d_0}$  is self-adjoint and non-negative, (2) follows.

By the theory of leading exponents sketched below,  $|\langle v, \pi(g)v \rangle| \ll_{\pi, v, \epsilon} q^{-l(g)(1/p(\pi)+\epsilon)}$ , so  $|g_{d_0/2}(g)| \ll q^{d_0\epsilon} \chi_{d_0/2}(g)$ . It follows that also  $|g_{d_0/2}^*(g)| \ll q^{d_0\epsilon} \chi_{d_0/2}(g)$ . By Lemma 4.18,  $|f_{d_0}(g)| \ll_{\epsilon} q^{d_0\epsilon} \psi_{d_0}(g)$ .

It remains to prove (1). We note that since  $f_{d_0}$  is self-adjoint and non-negative

$$\mathrm{tr}(\pi(f_{d_0})) \gg \|\pi(f_{d_0})\| = \|\pi(g_{d_0/2})\|^2$$

Now,

$$\begin{aligned} \langle v, \pi(g_{d_0/2})v \rangle &= \int_G g_{d_0/2}(g) \langle v, \pi(g)v \rangle \\ &= \int_G \chi_{d_0/2}(g) q^{l(g)p(\pi)^{-1}} \overline{\langle v, \pi(g)v \rangle} \langle v, \pi(g)v \rangle dg \\ &\geq \int_{l(g) \leq d_0/2} q^{l(g)p(\pi)^{-1}} |\langle v, \pi(g)v \rangle|^2 dg, \end{aligned}$$

and conclude that

$$\sqrt{\mathrm{tr}(\pi(f_{d_0}))} \gg \int_{l(g) \leq d_0/2} q^{l(g)p(\pi)^{-1}} |\langle v, \pi(g)v \rangle|^2 dg.$$

Therefore, to prove  $\mathrm{tr}(\pi(f_{d_0})) \gg q^{d_0(1-1/p(\pi)-\epsilon)}$ , one should prove (after changing  $d_0$  and  $d_0/2$ )

$$(9.1) \quad \int_{l(g) \leq d_0} q^{l(g)p(\pi)^{-1}} |\langle v, \pi(g)v \rangle|^2 dg \stackrel{!}{\gg}_{\pi, \epsilon} q^{d_0(1-p(\pi)^{-1}-\epsilon)}.$$

So far our proof is essentially the same as (part of) the proof of [53, Theorem 3], which only concerns rank 1.

Equation 9.1 follows from the theory of leading exponents. Before discussing this theory, we simplify the left hand side.

Applying the Cartan decomposition we get

$$\int_{l(g) \leq d_0} q^{l(g)p(\pi)^{-1}} |\langle v, \pi(g)v \rangle|^2 dg = \int_K \int_K \int_{a \in A_+, l(a) \leq d_0} q^{l(a)p(\pi)^{-1}} S(a) |\langle v, \pi(kak')v \rangle|^2 dkdk'da.$$

We identify  $a \in A_+ \subset \mathfrak{a}$  with coordinates  $(x_1, \dots, x_r)$ ,  $x_i \geq 0$  by  $(x_1, \dots, x_r) \rightarrow \sum_{i=1}^r x_i \omega_i$ , where  $\omega_1, \dots, \omega_r$  are fundamental coweights. Recall that for  $a \in A_+$ ,  $l(a) = 2\rho(a)$ . For  $\kappa > 0$ , denote by  $A_+^\kappa \subset A_+$  the set of  $a \in A_+$  with  $x_i > \kappa$ . Then for  $a \in A_+^\kappa$  it holds that

$S(a) \asymp_\kappa q^{l(a)} = q^{2l(a)}$ . Then:

$$\begin{aligned} & \int_K \int_K \int_{a \in A_+, l(a) \leq d_0} q^{l(a)p(\pi)^{-1}} S(a) |\langle v, \pi(kak')v \rangle|^2 dkdk' da \\ & \gg_\kappa \int_K \int_K \int_{a \in A_+^\kappa, l(a) \leq d_0} q^{2\rho(a)(1+p(\pi)^{-1})} |\langle v, \pi(kak')v \rangle|^2 dkdk' da, \end{aligned}$$

so we should prove that for some  $\kappa > 0$ ,

$$(9.2) \quad \int_K \int_K \int_{a \in A_+^\kappa, l(a) \leq d_0} q^{2\rho(a)(1+p(\pi)^{-1})} |\langle v, \pi(kak')v \rangle|^2 dkdk' da \stackrel{!}{\gg}_\epsilon q^{d_0(1-p(\pi)^{-1}-\epsilon)}.$$

**9.1. Leading Exponents.** We recall the Casselman-Harish Chandra-Milicic theory of *Leading Exponents*. We follow [39, Chapter VIII].

Inequality 9.2 is very similar to [39, Theorem 8.48,(b) implies (a)], which is one of the more technical parts of the theory. We will follow the same proof closely, while deriving an explicit expression. A side benefit is that the proofs below somewhat simplify the proof of [39, Theorem 8.48,(b) implies (a)].

By the theory of leading exponents, we may associate with an irreducible unitary representation (actually, to any irreducible admissible representation)  $(\pi, V)$  a finite subset called leading exponent  $\mathcal{F} \subset a^*$  of characters  $\nu : a \rightarrow \mathbb{C}$ , such that the following two useful theorems holds:

**Theorem 9.1** ([39, Theorem 8.47]). *The following are equivalent:*

- For  $\nu_0 : a \rightarrow \mathbb{R}$  a real character, every  $K$ -finite matrix coefficient  $\langle v, av \rangle$  for  $a \in A_+$  is bounded in absolute value by  $\ll_{\pi, v} e^{(\nu_0 - \rho)(a)} l(a)^N$  where  $N$  is some constant.
- For every  $\nu \in \mathcal{F}$ , and every fundamental weight  $\omega_i$ ,  $1 \leq i \leq r$ ,  $\text{Re} \nu(\omega_i) \leq \nu_0(\omega_i)$ .

**Theorem 9.2** ([39, Theorem 8.48]). *The following are equivalent:*

- Every  $K$ -finite matrix coefficient  $\phi(g) = \langle v, \pi(g)v \rangle$  is in  $L^p(G)$ .
- For every  $\nu \in \mathcal{F}$ , and every fundamental weight  $\omega_i$ ,  $1 \leq i \leq r$ ,  $\text{Re} \nu(\omega_i) < \left(1 - \frac{2}{p}\right) \rho(\omega_i)$ .

Note that the second theorem implies that  $p(\pi) = \min \left\{ p : \forall \nu \in \mathcal{F}, 1 \leq i \leq r, \text{Re} \nu(\omega_i) \leq \left(1 - \frac{2}{p}\right) \rho(\omega_i) \right\}$ , and that  $\phi(\pi) \notin L^{p(\pi)}(g)$ .

To state the main theorem about leading exponents, let us set some notations. Assume that  $0 \neq v \in V$  is  $K$ -finite, and let  $E_0 : V \rightarrow U_0$  be a projection onto a finite dimensional  $K$ -invariant subspace  $U_0 \subset V$  such that  $v \in U_0$ . We define  $F : A \rightarrow \text{End}_{\mathbb{C}}(U_0)$  by  $F(a) = E_0 \pi(a) E_0$ .

We denote by  $\mathcal{H}_{\text{End}U_0}$  the set of holomorphic functions  $f : D^r \rightarrow \text{End}(U_0)$ , where  $D = \{z \in \mathbb{C} : |z| < 1\}$  is the open unit ball. Each such function has convergent multiple power series, which is absolutely and uniformly convergent on compact subsets of  $D^r$ .

We identify  $a \in A_+ \subset \mathfrak{a}$  with coordinates  $(x_1, \dots, x_r)$ ,  $x_i \geq 0$  by  $(x_1, \dots, x_r) \rightarrow \sum_{i=1}^r x_i \omega_i$ , where  $\omega_1, \dots, \omega_r$  are the fundamental coweights.

We say that  $\nu, \nu' \in \mathfrak{a}^*$  are integrally equivalent if their  $\nu - \nu'$  difference is an integral combination of simple roots. If it is a non-negative integral combination of simple roots we write  $\nu' \leq \nu$ .

**Theorem 9.3** ([39, Theorem 8.32]). *There exists  $n_0 \in \mathbb{N}$ , a finite set  $\mathcal{F}'$  with  $\mathcal{F} \subset \mathcal{F}'$  and each  $\nu \in \mathcal{F}'$  satisfies  $\nu' \leq \nu$  for some  $\nu \in \mathcal{F}$ , such that for  $x_1 > 0, \dots, x_r > 0$  it holds that*

(9.3)

$$F(a) = F(x_1, \dots, x_r) = \sum_{\nu \in \mathcal{F}'} \sum_{1 \leq n_1 \leq n_2 \leq \dots \leq n_r \leq n} G_{\nu, n_1, \dots, n_r}(x_1, \dots, x_r) e^{(\nu - \rho)(x_1, \dots, x_r)} x_1^{n_1} \dots x_r^{n_r},$$

such that for  $\nu \in \mathcal{F}$ ,  $1 \leq n_i \leq n_0$ ,  $G_{\nu, n_1, \dots, n_r} : (0, \infty)^r \rightarrow \text{End}_{\mathbb{C}}(U_0)$  are functions given by  $G_{\nu, n_1, \dots, n_r}(x_1, \dots, x_r) = f_{\nu, n_1, \dots, n_r}(e^{-x_1}, \dots, e^{-x_r})$ , where  $f_{\nu, n_1, \dots, n_r} \in \mathcal{H}_{\text{End}U_0}$ .

Moreover, if  $f_{\nu, n_1, \dots, n_r} \neq 0$  then  $f_{\nu, n_1, \dots, n_r}(0, \dots, 0) \neq 0$  and for each  $\nu \in \mathcal{F}'$  there exists  $n_1, \dots, n_r$  with  $f_{\nu, n_1, \dots, n_r} \neq 0$ .

*Proof.* The theorem follows from [39, Theorem 8.32] and the discussion following it. Let us explain: by [39, Theorem 8.32],  $F$  has a decomposition like 9.3 with the set of leading exponents  $\mathcal{F}$  replaced by a different set  $\mathcal{F}'$ . By the power series expansion, we may assume the moreover part. By power series expansion, it gives a *unique* decomposition (see [39, Equation 8.52])

$$F(x_1, \dots, x_r) = \sum_{\nu \in \mathfrak{a}^*} F_{\nu - \rho}(x_1, \dots, x_r)$$

$$F_{\nu - \rho}(x_1, \dots, x_r) = \sum_{1 \leq n_1 \leq n_2 \leq \dots \leq n_r \leq n} c_{\nu, n_1, \dots, n_r} e^{(\nu - \rho)(x_1, \dots, x_r)} x_1^{n_1} \dots x_r^{n_r},$$

where each term  $\sum_{1 \leq n_1 \leq n_2 \leq \dots \leq n_r \leq n} G_{\nu, n_1, \dots, n_r}(x_1, \dots, x_r) e^{(\nu - \rho)(x_1, \dots, x_r)} x_1^{n_1} \dots x_r^{n_r}$  contributes into terms  $F_{\nu - \rho}$  with  $\nu' \leq \nu$ . The set of leading exponents are the maximal elements relative to  $\leq$  for  $\nu \in \mathcal{F}$  with  $F_{\nu - \rho} \neq 0$ . That immediately implies  $\mathcal{F} \subset \mathcal{F}'$ . Moreover each  $\nu' \in \mathfrak{a}^*$  with  $F_{\nu - \rho} \neq 0$  satisfies  $\nu' \leq \nu$  for  $\nu \in \mathcal{F}$ , which says that each  $\nu \in \mathcal{F}'$  satisfies  $\nu' \leq \nu$  for some  $\nu \in \mathcal{F}$ .  $\square$

We remark that Theorem 9.3 does not directly imply the upper bound given in Theorem 9.1, since it does not give bounds for  $x_i \rightarrow 0$ . Such bounds are available using asymptotic expansion near the walls ([39, Chapter VIII, Section 12]).

## 9.2. Some Technical Lemmas.

**Lemma 9.4** (Compare [39, Lemma B.24]). *Let  $f : \mathbb{R} \rightarrow \mathbb{C}$  be a function defined as  $f(x) = e^{\beta x} \sum_{i=1}^k c_i e^{-\alpha_i x} x^{n_i}$  with  $\alpha_i, c_i \in \mathbb{C}$ ,  $\text{Re}(\alpha_i) \geq 0$ ,  $\beta \in \mathbb{R}$ ,  $\beta \geq 0$ ,  $n_i \in \mathbb{N}$ . Assume that there is  $0 \leq i \leq k$  such that  $\text{Re}(\alpha_i) = 0$ ,  $c_i \neq 0$  and let  $n_0 = \max_{1 \leq i \leq k} \{n_i : \text{Re}(\alpha_i) = 0 \text{ and } c_i \neq 0\}$ .*

Then for  $T$  large enough,

$$\int_0^T |f(x)| dx \gg_f e^{\beta T} T^{n_0}.$$

Moreover, if we assume that  $n_0 = \max_{1 \leq i \leq k} \{n_i : \operatorname{Re}(\alpha_i) = 0\}$  then the underlying lower bound on  $T$  and the constants are continuous for small perturbations of the  $c_i$ .

*Proof.* During the proof,  $\gg$  may depend on  $f$ . Fix  $M$  large enough, depending on  $f$ , to be chosen later. Write

$$\int_0^T |f(x)| dx \geq \int_{T-M}^T |f(x)| dx.$$

Let  $\alpha_0 = \min_{1 \leq i \leq k} \{\operatorname{Re}(\alpha_i) : \operatorname{Re}(\alpha_i) \neq 0\}$ ,  $N = \max_{1 \leq i \leq k} \{n_i\}$ .

After re-arranging the summands, write  $f = f_0 + f_1$ , with  $f_0(x) = e^{\beta x} \sum_{i=1}^l c_i e^{-\alpha_i x} x^{n_0}$ ,  $f_1 = e^{\beta x} \sum_{i=l+1}^k c_i e^{-\alpha_i x} x^{n_i}$ , where for the summand  $1 \leq i \leq l$  contain all factors with  $\operatorname{Re}(\alpha_i) = 0$  and  $n_i = n_0$ . Then for  $\epsilon > 0$  small enough and  $T$  large enough,

$$\int_{T-M}^T |f_1(x)| dx \leq M \max_{T-M \leq x \leq T} \{f_1(x)\} \ll \left( e^{(\beta-\alpha)(T-M)} + e^{(\beta-\alpha)T} \right) T^N + e^{\beta T} T^{n_0-1} = o\left(e^{\beta T} T^{n_0}\right).$$

Now,

$$\begin{aligned} \int_{T-M}^T |f_0(x)| dx &= \int_{T-M}^T x^{n_0} e^{\beta x} \left| \sum_{i=1}^l c_i e^{-\alpha_i x} \right| dx \\ &\geq (T-M)^{n_0} e^{\beta(T-M)} \int_{T-M}^T \left| \sum_{i=1}^l c_i e^{-\alpha_i x} \right| dx \end{aligned}$$

Note that  $\left| \sum_{i=1}^l c_i e^{\alpha_i x} \right| \gg \left| \sum_{i=1}^l c_i e^{-\alpha_i x} \right|^2 = \sum_{i=1}^l |c_i|^2 + \sum_{1 \leq i \neq j \leq l} c_i \bar{c}_j e^{(\alpha_i - \alpha_j)x}$ , since this value is bounded.

$$\begin{aligned} \int_{T-M}^T \left| \sum_{i=1}^l c_i e^{\alpha_i x} \right| dx &\gg \int_{T-M}^T \left( \sum_{i=1}^l |c_i|^2 + \sum_{1 \leq i \neq j \leq l} c_i \bar{c}_j e^{(\alpha_i - \alpha_j)x} \right) dx \\ &\geq M \left( \sum_{i=1}^l |c_i|^2 \right) - \sum_{1 \leq i \neq j \leq l} \left| \frac{c_i \bar{c}_j}{\alpha_i - \alpha_j} \right| \\ &\gg M - O(1) \end{aligned}$$

For  $M$  large enough the last value is  $\gg 1$ , so

$$\int_{T-M}^T |f_0(x)| dx \gg e^{\beta T} T^{n_0}$$

and

$$\int_0^T |f(x)| dx \geq \int_{T-M}^T |f(x)| dx \geq \int_{T-M}^T |f_0(x)| dx - \int_{T-M}^T |f_1(x)| dx \gg e^{\beta T} T^{n_0}.$$

The moreover part of the argument follows from the proof.  $\square$

*Remark 9.5.* For  $\beta > 0$  our lower bound agree with a similar upper bound. For  $\beta = 0$  it is no longer true, but a similar proof will give the right lower bound  $T^{n_0+1}$ .

**Lemma 9.6.** *Let  $M$  be an open subset of a Lie group,  $F : M \times [R, \infty)$  a function defined by*

$$(9.4) \quad F(m, x) = \sum_{i=1}^k e^{s_i x} x^{n_i} F_i(m, x),$$

such that:  $s_i \in \mathbb{C}$ ,  $n_i \in \mathbb{N}$ , and  $F_i(m, x) = f_i(m, e^{-x})$  for some function  $f_i(m, z)$  real analytic on  $M \times \overline{D}_{e^{-R}}$ , where  $\overline{D}_r \subset \mathbb{C}$  is the closed ball of radius  $r$ , and holomorphic in the second variable. Further assume that for each  $1 \leq i \leq k$  there is  $m \in M^\circ$  with  $f_i(m, 0) \neq 0$ . Let  $s_0 = \max_{1 \leq i \leq k} \operatorname{Re} s_i$  and assume that  $s_0 \geq 0$ .

Then for  $T$  large enough

$$\int_M \int_R^T |F(s, r)| dr \gg e^{s_0 T}.$$

*Proof.* Decompose  $f_i(m, z) = c_i(m) + g_i(m, z)z$ , where  $g_i(m, z)$  is also holomorphic in  $|z| \leq e^{-R}$ . Then

$$F_i(m, r) = c_i(m) + e^{-r} G_i(m, r),$$

where  $G_i(m, r)$  is bounded for  $m, r \in M \times [R, \infty)$ . Therefore

$$(9.5) \quad F(m, r) = \sum_{i=1}^k e^{s_i r} r^{n_i} c_i(m) + \sum_{i=1}^k e^{(s_i-1)r} r^{n_i} G_i(m, r).$$

Without loss of generality,  $\operatorname{Re}(s_1) = s_0$  and  $n_1 = \max\{n_i : \operatorname{Re}(s_i) = s_0\}$ . Let  $m_0 \in M$  be a point with  $f_1(m_0, 0) \neq 0$ . Choose a small enough neighborhood  $M_0 \subset M$  of  $m_0$ . We have

$$\int_M \int_R^T |F(s, r)| dr \gg \int_{M_0} \int_R^T |F(s, r)| dr.$$

Since  $G_i(m, r)$  is bounded, for  $T$  large enough the second summand of Equation 9.5,

$$\int_{M_0} \int_R^T \left| \sum_{i=1}^k e^{(s_i-1)r} r^{n_i} G_i(m, r) \right| dr = o(e^{s_0 T}).$$



As for the first summand of Equation 9.5, by the moreover part of Lemma 9.4 and the fact that  $c_1(m_0) \neq 0$ , it holds that

$$\int_{M_0} \int_R^T \left| \sum_{i=1}^k e^{s_i r} r^{n_i} c_i(m) \right| dr \gg T^{n_1} e^{s_0 T}$$

and we are done.  $\square$

We can finally prove Equation 9.2.

*Proof.* Recall that we chose  $v \in V$ ,  $\|v\| = 1$  to span a representation  $\tau$  of  $K$ . We choose in Theorem 9.3  $U_0 = \text{span}Kv$ .

Using Theorem 9.2, choose  $\nu_0 \in \mathcal{F}'$  and  $1 \leq i \leq r$  such that  $\text{Re}\nu_0(\omega_i) = \left(1 - \frac{2}{p(\pi)}\right)\rho(\omega_i)$ . Without loss of generality  $i = 1$ . Moreover we assume that among all  $\nu \in \mathcal{F}'$  satisfying this condition,  $\nu_0$  has maximal  $0 \leq N_1 \leq n_0$  such that for some constants  $z_2, \dots, z_l \neq 0$ ,  $0 \leq n_2, \dots, n_r \leq n_0$ ,  $\lim_{z_1 \rightarrow 0} f_{\nu, N_1, \dots, n_k}(z_1, z_2, \dots, z_r) \neq 0$ , where  $f_{\nu, N_1, \dots, n_k}(z_1, z_2, \dots, z_r)$  is taken from Theorem 9.3 (there is a small technical fact here that  $\nu_0$  may belong to  $\mathcal{F}'$  and not  $\mathcal{F}$ ).

By Theorem 9.3, if we let  $M = K \times (0, \infty)^{r-1} \times K$ , we identify  $m \in M$  with  $m = (k_1, x_2, \dots, x_r, k_2)$ , and let  $G : M \times (0, \infty) \rightarrow \infty$  be

$$G(m, x_1) = \langle v, \pi(k_1 a(x_1, \dots, x_r) k_2) v \rangle,$$

then  $G(m, x_1)$  has the form 9.6, with  $s_1 = (\nu_0 - \rho)(\omega_1) = -\frac{2}{p(\pi)}\rho(\omega_1)$ ,  $n_1 = N_1$ . Note that  $G^2(m, x_1)$  also has this form, with  $s_1 = -\frac{4}{p(\pi)}\rho(\omega_1)$ . We Let

$$F(m, x_1) = e^{2\rho(\omega_1)(1+p(\pi)^{-1})x_1} G^2(m, x_1),$$

and  $F$  also has a similar form, with  $s_1 = 2\rho(\omega_1)\left(1 - p(\pi)^{-1}\right)$ . Let  $m_0 = (k_1, x_2, \dots, x_r, k_2)$  be a point where the condition of Lemma 9.6 holds. By Lemma 9.6, for a small neighborhood  $M_0$  of  $m_0$ , it holds for  $d_0$  large enough and some constant  $C$

$$\int_{M_0} \int_1^{(d_0-C)/2\rho(\omega_1)} |F(m, x_1)| dx_1 dm \gg e^{d_0(1-p(\pi)^{-1})}$$

Finally, for  $M_0, \kappa$  small enough, for each  $m = (k_1, x_2, \dots, x_r, k_2) \in M_0$  and  $1 \leq x_1 \leq (d_0 - C)/2\rho(\omega_1)$ , it holds that  $a = (x_1, \dots, x_r) \in A_+^\kappa$  and  $l(a) \leq d_0$ . Therefore

$$\begin{aligned} \int_K \int_K \int_{a \in A_+^\kappa, l(a) \leq d_0} q^{2\rho(a)(1-p(\pi)^{-1})} |\langle v, \pi(kak') v \rangle|^2 dk dk' da &\gg \int_{M_0} \int_0^{(d_0-1)/2\rho(\omega_1)} |F(m, x_1)| dx_1 dm \\ &\gg e^{d_0(1-p(\pi)^{-1})} \end{aligned}$$

as needed in Equation 9.2.  $\square$

## INDEX

The following notations appear throughout the paper.

- $k$  – a local field.
- $q$  – if  $k$  is Archimedean  $q = e$ . Otherwise  $q$  is the size of the quotient field of  $k$ .
- $G$  – the  $k$ -rational points of a semisimple algebraic group over  $k$ .
- $\Gamma$  – a lattice in  $G$ . If there is a sequence  $\{\Gamma_N\}$  of lattices then  $\Gamma_N$  is a finite index subgroup of  $\Gamma_1$ , with  $[\Gamma_1; \Gamma_N] \rightarrow \infty$ .
- $K$  – a good maximal compact subgroup of  $G$ .
- $X$  – the locally symmetric space  $\Gamma \backslash G/K$ .
- $\Pi(G)$  – the set of equivalence classes of irreducible unitary representations of  $G$ . A representation is usually denoted by  $(\pi, V)$ .
- $p(\pi)$  – the minimal  $p$  such that all  $K$ -finite matrix coefficients of  $(\pi, V)$  are in  $L^{p+\epsilon}(G)$ .
- $\lambda$  – an eigenvalue of the Casimir operator. Appears only in the Archimedean case.
- $l: G \rightarrow \mathbb{R}_{\geq 0}$  – a length function on  $G$ . A length is usually denoted  $d_0$ .
- $\chi_{d_0}$  – a smooth approximation for the characteristic set  $\{g \in G : l(g) \leq d_0\}$ .
- $\psi_{d_0}$  – a smooth approximation for  $q^{(d_0-l(g))/2} \chi_{d_0}$ .
- $b_{x_0, \delta}$  – for  $x \in \Gamma \backslash G/K$ ,  $\delta \in \mathbb{R}_{\geq 0}$ . In the non-Archimedean case it is the characteristic function of  $K$ . In the Archimedean case it is a smooth approximation for the characteristic function of a ball of radius  $\delta$  around  $X$ .

## REFERENCES

- [1] ABERT, M., BERGERON, N., BIRINGER, I., GELANDER, T., NIKOLOV, N., RAIMBAULT, J., AND SAMET, I. On the growth of  $L^2$ -invariants for sequences of lattices in Lie groups. *Annals of Mathematics* 185, 3 (2017), 711–790.
- [2] BERNSTEIN, J. All reductive p-adic groups are tame. *Functional Analysis and its applications* 8, 2 (1974), 91–93.
- [3] BERNSTEIN, J. Notes of lectures on Representations of p-adic Groups. [http://www.math.tau.ac.il/~bernstei/Unpublished\\_texts/unpublished\\_texts/Bernstein93new-harv.lect.from-chic.pdf](http://www.math.tau.ac.il/~bernstei/Unpublished_texts/unpublished_texts/Bernstein93new-harv.lect.from-chic.pdf) (1992).
- [4] BLOMER, V. Density theorems for  $GL(n)$ . *arXiv preprint arXiv:1906.07459* (2019).
- [5] BLOMER, V., BUTTCANE, J., AND MAGA, P. Applications of the Kuznetsov formula on  $GL(3)$  II: the level aspect. *Mathematische Annalen* 369, 1-2 (2017), 723–759.
- [6] BORDENAVE, C., AND LACONIN, H. Cutoff at the entropic time for random walks on covered expander graphs. *arXiv preprint arXiv:1812.06769* (2018).
- [7] BRUHAT, F., AND TITS, J. Groupes réductifs sur un corps local. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques* 41, 1 (1972), 5–251.
- [8] BURGER, M., LI, J.-S., AND SARNAK, P. Ramanujan duals and automorphic spectrum. *Bulletin of the American Mathematical Society* 26, 2 (1992), 253–257.

- [9] CARTIER, P. Representations of p-adic groups: a survey. In *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part (1979)*, vol. 1, pp. 111–155.
- [10] CASSELMAN, B. Introduction to admissible representations of p-adic groups. *unpublished notes* (1974).
- [11] CASSELMAN, W., AND MILICIC, D. Asymptotic behavior of matrix coefficients of admissible representations. *Duke Math. J* 49, 4 (1982), 869–930.
- [12] CHAPMAN, M., AND PARZANCHEVSKI, O. Cutoff on Ramanujan complexes and classical groups. *arXiv preprint arXiv:1901.09383* (2019).
- [13] COWLING, M., HAAGERUP, U., AND HOWE, R. Almost  $L^2$  matrix coefficients. *Journal für die Reine und Angewandte Mathematik* 387 (1988), 97–110.
- [14] DEGEORGE, D., AND WALLACH, N. Limit formulas for multiplicities in  $L^2(G/\gamma)$ . *Annals of Mathematics* (1978), 133–150.
- [15] DELIGNE, P. La conjecture de weil. i. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques* 43, 1 (1974), 273–307.
- [16] DUISTERMAAT, J., KOLK, J., AND VARADARAJAN, V. Spectra of compact locally symmetric manifolds of negative curvature. *Inventiones mathematicae* 52, 1 (1979), 27–93.
- [17] DUKE, W., RUDNICK, Z., SARNAK, P., ET AL. Density of integer points on affine homogeneous varieties. *Duke mathematical journal* 71, 1 (1993), 143–179.
- [18] EINSIEDLER, M., AND LINDENSTRAUSS, E. Diagonal actions on locally homogeneous spaces. *Homogeneous flows, moduli spaces and arithmetic* 155241 (2010).
- [19] EVRA, S., AND PARZANCHEVSKI, O. Ramanujan complexes and golden gates in  $PU(3)$ . *arXiv preprint arXiv:1810.04710* (2018).
- [20] FINIS, T., AND LAPID, E. An approximation principle for congruence subgroups. *Journal of the European Mathematical Society* 20, 5 (2018), 1075–1138.
- [21] GAMBURD, A. On the spectral gap for infinite index "congruence" subgroups of  $SL_2(\mathbb{Z})$ . *Israel Journal of Mathematics* 127, 1 (2002), 157–200.
- [22] GANGOLLI, R., AND VARADARAJAN, V. S. *Harmonic analysis of spherical functions on real reductive groups*, vol. 101 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1988.
- [23] GELFAND, I. M., GRAEV, M. I., AND PIATETSKI-SHAPIRO, I. *Representation theory and automorphic functions*. Saunders, 1968.
- [24] GHOSH, A., GORODNIK, A., AND NEVO, A. Diophantine approximation and automorphic spectrum. *International Mathematics Research Notices* 2013, 21 (2013), 5002–5058.
- [25] GHOSH, A., GORODNIK, A., AND NEVO, A. Best possible rates of distribution of dense lattice orbits in homogeneous spaces. *Journal für die reine und angewandte Mathematik (Crelles Journal)* (2014).
- [26] GOLUBEV, K., AND KAMBER, A. Cutoff on Graphs and The Sarnak-Xue Density of

- Eigenvalues. *arXiv preprint arXiv:1905.11165* (2019).
- [27] GOLUBEV, K., AND KAMBER, A. Cutoff on hyperbolic surfaces. *Geometriae Dedicata* (3 2019), 1–31.
  - [28] GUTH, L., AND LUBOTZKY, A. Quantum error correcting codes and 4-dimensional arithmetic hyperbolic manifolds. *Journal of Mathematical Physics* 55, 8 (2014), 082202.
  - [29] HARDY, G. H., AND WRIGHT, E. M. *An introduction to the theory of numbers*. Oxford university press, 1979.
  - [30] HARISH-CHANDRA. Spherical functions on a semisimple Lie group, I. *American Journal of Mathematics* (1958), 241–310.
  - [31] HASHIMOTO, K.-I. Zeta functions of finite graphs and representations of p-adic groups. *Automorphic forms and geometry of arithmetic varieties*. (1989), 211–280.
  - [32] HOWE, R. E., AND PIATETSKI-SHAPIRO, I. I. A counterexample to the "generalized Ramanujan conjecture for (quasi-) split groups. In *Proceedings of Symposia in Pure Mathematics, Volume 33, Part 1: Automorphic Forms, Representations and L-functions* (1979), vol. 1, American Mathematical Soc., pp. 315–322.
  - [33] HUXLEY, M. N. Exceptional eigenvalues and congruence subgroups. *The Selberg Trace Formula and Related Topics, Contemp. Math* 53 (1986), 341–349.
  - [34] KAMBER, A.  $L^p$ -expander complexes. *arXiv preprint arXiv:1701.00154* (2016).
  - [35] KAMBER, A.  $L^p$ -expander graphs. *Israel Journal of Mathematics* 234, 2 (2019), 863–905.
  - [36] KAMBER, A., AND LAVNER, H. Optimal Lifting for the Projective Action of  $SL_3(\mathbb{Z})$ . *arXiv preprint arXiv:1908.06682* (2019).
  - [37] KANG, M.-H., LI, W.-C. W., AND WANG, C.-J. The zeta functions of complexes from  $PGL(3)$ : a representation-theoretic approach. *Israel Journal of Mathematics* 177, 1 (2010), 335–348.
  - [38] KATZNELSON, Y. Singular matrices and a uniform bound for congruence groups of  $SL_n(\mathbb{Z})$ . *Duke Mathematical Journal* 69, 1 (1993), 121–136.
  - [39] KNAPP, A. W. *Representation Theory of Semisimple Groups: An Overview Based on Examples (PMS-36)*. Princeton university press, 2016.
  - [40] KOSTANT, B. On the existence and irreducibility of certain series of representations. *Bulletin of the American Mathematical Society* 75, 4 (1969), 627–642.
  - [41] LAFFORGUE, L. Chtoucas de drinfeld et correspondance de langlands. *Inventiones mathematicae* 147, 1 (2002), 1–241.
  - [42] LAX, P. D., AND PHILLIPS, R. S. The asymptotic distribution of lattice points in Euclidean and non-Euclidean spaces. *Journal of Functional Analysis* 46, 3 (1982), 280–350.
  - [43] LUBETZKY, E., LUBOTZKY, A., AND PARZANCHEVSKI, O. Random walks on Ramanujan complexes and digraphs. *arXiv preprint arXiv:1702.05452* (2017).
  - [44] LUBETZKY, E., AND PERES, Y. Cutoff on all Ramanujan graphs. *Geometric and Functional Analysis* 26, 4 (2016), 1190–1216.

- [45] LUBOTZKY, A. High dimensional expanders. *arXiv preprint arXiv:1712.02526* (2017).
- [46] LUBOTZKY, A., SAMUELS, B., AND VISHNE, U. Explicit constructions of Ramanujan complexes of type  $A_d$ . *European Journal of Combinatorics* 26, 6 (2005), 965–993.
- [47] LUBOTZKY, A., SAMUELS, B., AND VISHNE, U. Ramanujan complexes of type  $A_d$ . *Israel Journal of Mathematics* 149, 1 (2005), 267–299.
- [48] MAGEE, M. Quantitative spectral gap for thin groups of hyperbolic isometries. *Journal of the European Mathematical Society* 17, 1 (2015), 151–187.
- [49] MATZ, J., AND TEMPLIER, N. Sato-tate equidistribution for families of hecke-maass forms on  $SL(n, \mathbb{R})/SO(n)$ . *arXiv preprint arXiv:1505.07285* (2015).
- [50] SARDARI, N. T. Diameter of Ramanujan graphs and random Cayley graphs. *Combinatorica* 39, 2 (2019), 427–446.
- [51] SARDARI, N. T. Optimal strong approximation for quadratic forms. *Duke Mathematical Journal* (2019).
- [52] SARNAK, P. Letter to Stephen D. Miller and Naser Talebizadeh Sardari on optimal strong approximation by integral points on quadrics. 2015.
- [53] SARNAK, P., AND XUE, X. X. Bounds for multiplicities of automorphic representations. *Duke Math. J* 64, 1 (1991), 207–227.
- [54] SAUVAGEOT, F. Principe de densité pour les groupes réductifs. *Compositio Mathematica* 108, 2 (1997), 151–184.
- [55] SEEGER, A., AND SOGGE, C. D. Bounds for eigenfunctions of differential operators. *Indiana University Mathematics Journal* 38, 3 (1989), 669–682.
- [56] SILBERGER, A. G. *Introduction to harmonic analysis on reductive  $P$ -adic groups. (MN-23): Based on lectures by Harish-Chandra at The Institute for Advanced Study, 1971-73.* Princeton university press, 2015.
- [57] TITS, J. Reductive groups over local fields. In *Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part* (1979), vol. 1, pp. 29–69.