

MULTIPLICITY RESULTS FOR ELLIPTIC PROBLEMS INVOLVING NONLOCAL INTEGRODIFFERENTIAL OPERATORS WITHOUT AMBROSETTI-RABINOWITZ CONDITION

LAUREN MARIA MEZZOMO BONALDO, OLMPPIO HIROSHI MIYAGAKI AND ELARD JUREZ HURTADO

ABSTRACT. In this paper, we study the existence and multiplicity of weak solutions for a general class of elliptic equations (\mathcal{P}_λ) in a smooth bounded domain, driven by a nonlocal integrodifferential operator \mathcal{L}_{AK} with Dirichlet boundary conditions involving variable exponents without Ambrosetti and Rabinowitz type growth conditions. Using different versions of the Mountain Pass Theorem, as well as, the Fountain Theorem and Dual Fountain Theorem with Cerami condition, we obtain the existence of weak solutions for the problem (\mathcal{P}_λ) and we show that the problem treated has at least one nontrivial solution for any parameter $\lambda > 0$ small enough as well as that the solution blows up, in the fractional Sobolev norm, as $\lambda \rightarrow 0$. Moreover, for the case sublinear, by imposing some additional hypotheses on the nonlinearity $f(x, \cdot)$, by using a new version of the symmetric Mountain Pass Theorem due to Kajikiya [36], we obtain the existence of infinitely many weak solutions which tend to be zero, in the fractional Sobolev norm, for any parameter $\lambda > 0$. As far as we know, the results of this paper are new in the literature.

1. INTRODUCTION AND MAIN RESULTS

In recent years, the study of elliptic problems involving fractional operators with variable exponents has become the object of study of many researchers. Problems of this kind have a very interesting theoretic basis very interesting whose integrability and analytical structure require delicate proofing techniques as we can see in [7, 8, 29, 37]. As well as, they have concrete applications in the most diverse fields as optimization, finance, phase transitions, continuum mechanics, image process, game theory, crystal dislocation, quasi-geostrophic flows, peridynamic theory, see [5, 6, 11, 14, 15, 28, 30, 33, 47, 49, 54] and their references.

In this sense, in order to expand the results around this theory, in the present paper we are concerned with the existence and multiplicity of weak solutions for elliptic equations involving the nonlocal integrodifferential operators with variable exponents. Namely, we consider the following problem

$$\begin{cases} \mathcal{L}_{AK}u = \lambda f(x, u) & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases} \quad (\mathcal{P}_\lambda)$$

where $\lambda > 0$ is a real parameter, $\Omega \subset \mathbb{R}^N$, $N \geq 2$, is a smooth bounded domain, and to define the nonlocal integrodifferential operator \mathcal{L}_{AK} we will need the variable exponents $p(x) := p(x, x)$ for all $x \in \mathbb{R}^N$ with $p \in C(\mathbb{R}^N \times \mathbb{R}^N)$ satisfying:

p is symmetric, that is, $p(x, y) = p(y, x)$,

$$1 < p^- := \inf_{(x,y) \in \mathbb{R}^N \times \mathbb{R}^N} p(x, y) \leq \sup_{(x,y) \in \mathbb{R}^N \times \mathbb{R}^N} p(x, y) := p^+ < \frac{N}{s}, \quad s \in (0, 1), \quad (p_1)$$

and we consider the fractional critical variable exponent related to $p \in C(\mathbb{R}^N \times \mathbb{R}^N)$ defined by

$$p_s^*(x) = \frac{Np(x)}{N - sp(x)}.$$

2010 *Mathematics Subject Classification.* 35A15, 35B38, 35D3, 35J60, 35R11, 46E35, 47G20.

Key words and phrases. Nonlocal integrodifferential operator, fractional Sobolev space with variable exponents, Mountain Pass theorem, Fountain theorem, Dual Fountain theorem.

Department of Mathematics Universidade Federal de São Carlos. Carlos, São Carlos SP, Brazil. E-mail: laurenmbonaldo@hotmail.com. Supported by CAPES.

Department of Mathematics Universidade Federal de São Carlos. Carlos, São Carlos SP, Brazil. E-mail: ohmiyagaki@gmail.com Supported by CAPES.

Department of Mathematics, Universidade de Brasilia, Brasilia, DF, Brazil. E-mail: elardjh2@gmail.com Supported by CAPES.

The nonlocal integrodifferential operator \mathcal{L}_{AK} is defined on suitable fractional Sobolev spaces (see Section 2) by

$$\mathcal{L}_{AK}u(x) = P.V. \int_{\mathbb{R}^N} \mathcal{A}(u(x) - u(y))K(x, y) dy, \quad x \in \mathbb{R}^N,$$

where $P.V.$ is the principal value.

The map $\mathcal{A} : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function satisfying the next assumptions:

(a₁) \mathcal{A} is continuous, odd, and the map $\mathcal{A} : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$\mathcal{A}(t) := \int_0^{|t|} \mathcal{A}(\tau) d\tau$$

is strictly convex;

(a₂) There exist positive constants $c_{\mathcal{A}}$ and $C_{\mathcal{A}}$, such that for all $t \in \mathbb{R}$ and for all $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$

$$\mathcal{A}(t)t \geq c_{\mathcal{A}}|t|^{p(x,y)} \quad \text{and} \quad |\mathcal{A}(t)| \leq C_{\mathcal{A}}|t|^{p(x,y)-1};$$

(a₃) $\mathcal{A}(t)t \leq p^+ \mathcal{A}(t)$ for all $t \in \mathbb{R}$.

The kernel $K : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}^+$ be a measurable function satisfying the following property:

(K) There exist constants b_0 and b_1 , such that $0 < b_0 \leq b_1$,

$$b_0 \leq K(x, y)|x - y|^{N+sp(x,y)} \leq b_1 \text{ for all } (x, y) \in \mathbb{R}^N \times \mathbb{R}^N \text{ and } x \neq y.$$

We assume that $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function and satisfies:

(f₀) There exists a positive constant c_1 such that f satisfies the subcritical growth condition

$$|f(x, t)| \leq c_1(1 + |t|^{\vartheta(x)-1})$$

for all $(x, t) \in \Omega \times \mathbb{R}$, where $\vartheta \in C(\overline{\Omega})$, $1 < p^+ < \underline{\vartheta}^- \leq \vartheta(x) \leq \underline{\vartheta}^+ < p_s^*(x)$ for $x \in \overline{\Omega}$, and $\underline{\vartheta}^- := \inf_{x \in \overline{\Omega}} \vartheta(x)$, $\underline{\vartheta}^+ := \sup_{x \in \overline{\Omega}} \vartheta(x)$;

(f₁) $\lim_{|t| \rightarrow +\infty} \frac{F(x, t)}{|t|^{p^+}} = +\infty$ uniformly for almost everywhere a.e. $x \in \Omega$, that is, f is p^+ -superlinear at infinity, the function F is the primitive of f with respect to the second variable, that is,

$$F(x, t) := \int_{\Omega} f(x, \tau) d\tau;$$

(f₂) $f(x, t) = o(|t|^{p^+-1})$, as $t \rightarrow 0$, uniformly a.e. $x \in \Omega$;

(f₃) There exists a positive constant $c_{\star} > 0$ such that

$$\mathcal{G}(x, t) \leq \mathcal{G}(x, \tau) + c_{\star}$$

for all $x \in \Omega$, $0 < t < \tau$ or $\tau < t < 0$, where $\mathcal{G}(x, t) := tf(x, t) - p^+F(x, t)$.

With intending to find infinite solutions is natural to impose certain symmetry condition on the nonlinearity. In the sequel, we will assume the following assumption on f :

(f₄) f is odd in t , that is, $f(x, -t) = -f(x, t)$ for all $x \in \Omega$ and $t \in \mathbb{R}$.

In addition, to prove that the Euler Lagrange functional associated to the problem (\mathcal{P}_{λ}) verifies the Cerami condition $(C)_c$, we assume that the functions \mathcal{A} and K satisfy the following condition

(a₄) $\mathcal{H}(at) \leq \mathcal{H}(t)$ for all $t \in \mathbb{R}$ and $a \in [0, 1]$ where $\mathcal{H}(t) = p^+ \mathcal{A}(t) - \mathcal{A}(t)t$.

A typical mathematical generalization for \mathcal{A} and K verifying (a₁)-(a₄) and (K) respectively is $\mathcal{A}(t) = |t|^{p(x,y)-2}t$ and $K(x, y) = |x - y|^{-(N+sp(x,y))}$. When \mathcal{A} and K are these type, the operator \mathcal{L}_{AK} is the fractional $p(\cdot)$ -Laplacian operator $(-\Delta)_{p(\cdot)}^s$, which is defined by

$$(-\Delta)_{p(\cdot)}^s u(x) = P.V. \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p(x,y)-2}(u(x) - u(y))}{|x - y|^{N+sp(x,y)}} dy \text{ for all } x \in \mathbb{R}^N.$$

For further details on the fractional $p(\cdot)$ -Laplacian operator see for instance [7, 8, 29, 37]. Furthermore, it is worth to note that the assumptions (a₁)-(a₄) and (K) were similarly introduced in [3, 12, 13, 18, 22, 27, 31, 32, 39, 40, 50, 55].

In literature problem (\mathcal{P}_λ) was investigated primarily in the context of equations driven by the Laplacian (semilinear problems), i.e., was considered the following problem class

$$\begin{cases} -\Delta u &= \lambda f(x, u) & \text{in } \Omega, \\ u &= 0 & \text{in } \partial\Omega. \end{cases} \quad (1.1)$$

Particularly Ambrosetti and Rabinowitz in [2], were the first to use the Mountain Pass Theorem to prove that problems type above admits a solution when the following condition of the nonlinearity to $f(x, \cdot)$, well known in the literature as the Ambrosetti-Rabinowitz condition ((AR) condition, for short), is employed: there exist $\mu > 2$ and $M > 0$ such that

$$0 < \mu F(x, t) \leq f(x, t)t \text{ for all } x \in \Omega \text{ and for all } |t| \geq M \quad (1.2)$$

where $f : \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function and $F(x, t) = \int_0^t f(t, z) dz$. The (AR) condition is a tool essential to obtain weak solutions of superlinear problems. Moreover, their main role is to ensure the compactness, more specifically, the limitation of the Palais-Smale sequence required by minimax arguments. But, the (AR) condition is some restricted and eliminates some nonlinearities of $F(x, \cdot)$. Direct integration of (1.2) leads us to the following weak condition for the potential function $F(x, t)$,

$$F(x, t) \geq \alpha_1 |t|^\mu - \alpha_2 \text{ for all } x \in \overline{\Omega} \text{ and for all } t \in \mathbb{R} \text{ with } \alpha_1, \alpha_2 > 0. \quad (1.3)$$

However, even so, there are still many functions that are superlinear at infinity and that not satisfy the (AR) condition. Indeed, the condition (1.3) implies that the function $F(x, \cdot)$ displays at least one polynomial growth μ near to $\pm\infty$ and since that $\mu > p$

$$\lim_{|t| \rightarrow +\infty} \frac{F(x, t)}{|t|^p} = +\infty. \quad (1.4)$$

The function $f(x, t) = t \ln(1 + t)$ is a example of an function that satisfies the condition (1.4) but does not satisfy the condition (1.3) and therefore also not satisfies the condition (1.2).

In this sense, the study for problems involving not only Laplacian operator but also $p(\cdot)$ -Laplacian operator without the (AR) condition has become the focus for many researchers and the references in the literature have steadily increased. We mention some papers as [10, 31, 35, 43, 44, 45, 46] where the authors established at least nontrivial solution existence to the problems of type (1.1) without (AR) condition and in general, the main tool used to obter the results is Mountain Pass Theorem with Cerami condition.

Furthermore to this significant advance for local operators without the (AR) condition, recently, some researchers have started to study problems of type (1.1) nonlocal without the (AR) condition. More specifically, in the paper [56] considering $\mathcal{A}(t) = t$, $K(x, y) = |x - y|^{N+2s}$, and $\lambda = 1$, the authors proved that the problem (\mathcal{P}_λ) has infinite solutions using Fountain Theorem. In the paper [48] the author, for the case fractional p -Laplacian, $K(x, y) = |x - y|^{N+2s}$ and $\lambda = 1$ studies the existence of a weak solution to the problem (\mathcal{P}_λ) by Mountain Pass Theorem combined with the fractional MoserTrudinger inequality. Already in [58], for $\mathcal{A}(t) = t$ and some similar assumptions for the K kernel, the author proves that $\lambda_0 > 0$ exists such that the problem (\mathcal{P}_λ) supports two distinct weak solutions for every $\lambda \in (0, \lambda_0)$ using a new critical points result introduced by [41, Theorem 2.6].

Therefore, motivated by the above references, mainly by the papers [31, 46, 56, 58], the present paper is concerned about with the existence and multiplicity of solutions to the problem (\mathcal{P}_λ) without the (AR) condition. In particular, we extend, we complement and improve some of the references cited above once our problem (\mathcal{P}_λ) involves general nonlocal integrodifferential operators. Indeed, is important note that the operator \mathcal{L}_{AK} involving in problem (\mathcal{P}_λ) is, in particular, the fractional $p(\cdot)$ -Laplacian operator, so it is nonhomogeneous and problems involving this operator with variable exponents can not use Lagrange Multiplier Theorem, as can be done for operators involving constant exponents. It still is interesting to point that the operator \mathcal{L}_{AK} is no just a mere extension the fractional $p(\cdot)$ -Laplacian operator, because from conditions (a_1) - (a_4) and (\mathcal{K}) , map \mathcal{A} and kernel K are quite general and K includes singular kernels. Furthermore, we note that solution space for the problem (\mathcal{P}_λ) must be a fractional Sobolev space with variable exponents as we can see in [7, 8, 29, 37]. Thus, due to the characterization of our problem (\mathcal{P}_λ) it was required to define a new space in this paper. Then inspired by the spaces mentioned above and the

space $W_0^{s,p}(\Omega)$ (defined in [34]), we define the following space

$$\mathcal{W} = W_0^{s,p(\cdot,\cdot)} := \{u \in W^{s,p(\cdot,\cdot)}(\mathbb{R}^N) : u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega\},$$

which is a separable and reflexive Banach space and we prove an important result of compact and continuous embedding for \mathcal{W} , for more details see Subsection 2.2.

Moreover, as our problem has a structure variational and our objective is to prove existence, multiplicity and comporment of solution of the problem (\mathcal{P}_λ) , is essential verify that the operator Φ' (see Lemma 2.7) satisfies the property (S_+) , which is a property of compactness of the operator and usually is used for other property of compactness as for example the condition Palais-Smale or Cerami for the Euler Lagrange functional Ψ_λ defined in (2.3).

Lastly, we will obtain the existence of weak solutions to the problem (\mathcal{P}_λ) and, under some considerations, get a nontrivial solution for any parameter $\lambda > 0$ small enough such that the solution blows up, in the fractional Sobolev norm, as $\lambda \rightarrow 0^+$. For this, along of the paper, we will use different versions of the Mountain Pass Theorem, as well as, the Fountain and Dual Theorem with Cerami condition. Moreover, by imposing additional hypotheses on the nonlinearity $f(x, \cdot)$, we get the existence of infinitely many weak solutions which tend to be zero, in the fractional Sobolev norm, for any parameter $\lambda > 0$ by using the new version of the symmetric of the Pass Mountain Theorem introduced by Kajikya [36].

Our main results are the following Theorems.

Theorem 1.1. *Assume (a_1) - (a_4) , (\mathcal{K}) , and f satisfies (f_0) - (f_3) . Then problem (\mathcal{P}_λ) has at least one nontrivial weak solution in \mathcal{W} for all $\lambda > 0$.*

Theorem 1.2. *Assume (a_1) - (a_4) , (\mathcal{K}) , and f satisfies (f_0) - (f_4) . Then problem (\mathcal{P}_λ) has infinitely many solutions in \mathcal{W} for all $\lambda > 0$.*

Theorem 1.3. *Assume (a_1) - (a_4) , (\mathcal{K}) , and f satisfies (f_0) , (f_1) , (f_3) and (f_4) . Then, for each $\lambda \in \left(0, \frac{c_A b_0}{p^+}\right)$, the problem (\mathcal{P}_λ) has infinitely many weak solutions $u_k \in \mathcal{W}$, $k \in \mathbb{N}$ such that $\Psi_\lambda(u_k) \rightarrow +\infty$, as $k \rightarrow +\infty$. (Ψ_λ is defined (2.3))*

Theorem 1.4. *Assume (a_1) - (a_4) , (\mathcal{K}) , and f satisfies f satisfies (f_0) , (f_1) , (f_3) and (f_4) . Then, for each $\lambda \in \left(0, \frac{c_A b_0}{p^+}\right)$, the problem (\mathcal{P}_λ) has a sequence of weak solutions $v_k \in \mathcal{W}$, $k \in \mathbb{N}$ such that $\Psi_\lambda(v_k) < 0$, $\Psi_\lambda(v_k) \rightarrow 0$ as $k \rightarrow +\infty$.*

Theorem 1.5. *Assume (a_1) - (a_4) and (\mathcal{K}) . If f satisfies (f_0) , (f_1) , and (f_3) , moreover $f(x, 0) = 0$, $f(x, t) \geq 0$ a.e. $x \in \Omega$ and for all $t \geq 0$. Then there exists a positive constant $\bar{\lambda}$ such that problem (\mathcal{P}_λ) possesses at least one solution for all $\lambda \in (0, \bar{\lambda})$. Moreover*

$$\lim_{\lambda \rightarrow 0^+} \|u_\lambda\|_{\mathcal{W}} = +\infty.$$

($\|\cdot\|_{\mathcal{W}}$ is defined in Subsection(2.2))

Theorem 1.6. *Assume (a_1) - (a_3) , (\mathcal{K}) , and f satisfies (f_4) . In addition we will assume the following condition:*

(f_5) $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function and there exist positive constants C_0, C_1 such that

$$C_0|t|^{\mathbf{m}(x)-1} \leq f(x, t) \leq C_1|t|^{\mathbf{m}(x)-1} \text{ for all } x \in \bar{\Omega} \text{ and } t \geq 0,$$

where $\mathbf{m} \in C(\bar{\Omega})$ such that $1 < \mathbf{m}(x) < p^*(x)$ for all $x \in \bar{\Omega}$, with $\underline{\mathbf{m}}^+ < p^-$. Then problem (\mathcal{P}_λ) it has a sequence of on trivial solutions $u_k \in \mathcal{W}$, $k \in \mathbb{N}$ such that

$$\lim_{k \rightarrow +\infty} \|u_k\|_{\mathcal{W}} = 0$$

for all $\lambda > 0$.

We note that our results show that the main results of [31, 35, 42, 43, 46, 53, 59] remain valid for a wider class of nonlocal operators involving variable exponents. Moreover, as far as we are aware, there are no results on this approach even involving the fractional p -Laplacian problems as well as the fractional $p(\cdot)$ -Laplacian problems although we consider some the well-known technique.

This paper is organized as follows. In Section 2, we give some preliminary results about Lebesgue and fractional Sobolev spaces with variable exponents. Besides, we define the function space for the problem

(\mathcal{P}_λ), we give an important result of compact and continuous embedding, and we proof fundamentals properties of Euler Lagrange functional associated with the of problem (\mathcal{P}_λ). In Section 3, we prove the main theorems of this paper.

2. PRELIMINARIES

In this section, we give some preliminary results which will be used in the sequel to discuss the problem (\mathcal{P}_λ). Throughout the present paper c_i and C_i for $i = 1, 2, \dots$ will denote generic positive constants which may vary from line to line, but are independent of the terms which take part in any limit process.

2.1. Variable exponent Lebesgue space.

In this subsection, we briefly review some of the basic properties of the Lebesgue spaces with variable exponent, for more can be found in [1, 20, 21, 25, 24, 52] and the references therein.

Let $\Omega \subset \mathbb{R}^N$ ($N \geq 2$) an smooth bounded domain. We introduced

$$C^+(\overline{\Omega}) = \{h \in C(\overline{\Omega}) : h(x) > 1 \text{ for all } x \in \overline{\Omega}\}$$

and for $\sigma \in C^+(\overline{\Omega})$, denote

$$\underline{\sigma}^- := \inf_{x \in \overline{\Omega}} \sigma(x), \underline{\sigma}^+ := \sup_{x \in \overline{\Omega}} \sigma(x).$$

For $\sigma \in C^+(\overline{\Omega})$, we define the variable exponent Lebesgue space $L^{\sigma(\cdot)}(\Omega)$ as

$$L^{\sigma(\cdot)}(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable} : \exists \zeta > 0 : \int_{\Omega} \left| \frac{u(x)}{\zeta} \right|^{\sigma(x)} dx < +\infty \right\}. \quad (2.1)$$

The space $L^{\sigma(\cdot)}(\Omega)$ endowed with the Luxemburg norm,

$$\|u\|_{L^{\sigma(\cdot)}(\Omega)} := \inf \left\{ \zeta > 0 : \int_{\Omega} \left| \frac{u(x)}{\zeta} \right|^{\sigma(x)} dx \leq 1 \right\}.$$

is a separable and reflexive Banach space. If $\Omega = \mathbb{R}^N$, we simply denote the norm of any $u \in L^{\sigma(\cdot)}(\mathbb{R}^N)$ by $\|u\|_{L^{\sigma(\cdot)}(\mathbb{R}^N)}$. Note that, when σ is constant, the Luxemburg norm $\|\cdot\|_{L^{\sigma(\cdot)}(\Omega)}$ coincide with the standard norm $\|\cdot\|_{L^\sigma(\Omega)}$ of the Lebesgue space $L^\sigma(\Omega)$.

Denoting by $L^{\sigma'(\cdot)}(\Omega)$ the dual space of space $L^{\sigma(\cdot)}(\Omega)$, where $\frac{1}{\sigma(x)} + \frac{1}{\sigma'(x)} = 1$, for any $u \in L^{\sigma(\cdot)}(\Omega)$ and $v \in L^{\sigma'(\cdot)}(\Omega)$ the following Hölder type inequality holds

$$\int_{\Omega} |uv| dx \leq \left(\frac{1}{\underline{\sigma}^-} + \frac{1}{\underline{\sigma}'^-} \right) \|u\|_{L^{\sigma(\cdot)}(\Omega)} \|v\|_{L^{\sigma'(\cdot)}(\Omega)}.$$

If $\sigma_1, \sigma_2 \in C^+(\overline{\Omega})$, $\sigma_1(x) \leq \sigma_2(x)$ for all $x \in \overline{\Omega}$, then $L^{\sigma_2(\cdot)}(\Omega) \hookrightarrow L^{\sigma_1(\cdot)}(\Omega)$, and the embedding is continuous. An important role in manipulating the generalized Lebesgue spaces is played by the $\sigma(\cdot)$ -modular of the $L^{\sigma(\cdot)}(\Omega)$ space, which is the convex function $\rho_{\sigma(\cdot)} : L^{\sigma(\cdot)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\rho_{\sigma(\cdot)}(u) = \int_{\Omega} |u|^{\sigma(x)} dx,$$

along any function $u \in L^{\sigma(\cdot)}(\Omega)$.

Proposition 2.1. *Let $u \in L^{\sigma(\cdot)}(\Omega)$ and $(u_k)_{k \in \mathbb{N}} \subset L^{\sigma(\cdot)}(\Omega)$, then the following relations hold:*

- (a) *For $u \in L^{\sigma(\cdot)}(\Omega) \setminus \{0\}$, $\zeta = \|u\|_{L^{\sigma(\cdot)}(\Omega)}$ if and only if $\rho_{\sigma(\cdot)}\left(\frac{u}{\zeta}\right) = 1$;*
- (b) *$\|u\|_{L^{\sigma(\cdot)}(\Omega)} \geq 1 \Rightarrow \|u\|_{L^{\sigma(\cdot)}(\Omega)}^{\underline{\sigma}^-} \leq \rho_{\sigma(\cdot)}(u) \leq \|u\|_{L^{\sigma(\cdot)}(\Omega)}^{\underline{\sigma}^+}$;*
- (c) *$\|u\|_{L^{\sigma(\cdot)}(\Omega)} \leq 1 \Rightarrow \|u\|_{L^{\sigma(\cdot)}(\Omega)}^{\underline{\sigma}^+} \leq \rho_{\sigma(\cdot)}(u) \leq \|u\|_{L^{\sigma(\cdot)}(\Omega)}^{\underline{\sigma}^-}$;*

Moreover, $\|u_k - u\|_{L^{\sigma(\cdot)}(\Omega)} \rightarrow 0 \Leftrightarrow \rho_{\sigma(\cdot)}(u_k - u) \rightarrow 0 \Leftrightarrow u_k \rightarrow u$ in measure in Ω and $\rho_{\sigma(\cdot)}(u_k) \rightarrow \rho_{\sigma(\cdot)}(u)$. In particular, $\rho_{\sigma(\cdot)}$ is continuous in $L^{\sigma(\cdot)}(\Omega)$, and if furthermore $\sigma \in C^+(\overline{\Omega})$, then $\rho_{\sigma(\cdot)}$ is weakly lower semi-continuous.

2.2. The functional space \mathscr{W} and their properties.

Let $\Omega \subset \mathbb{R}^N$, $N \geq 2$, is a smooth bounded domain and $s \in (0, 1)$. We consider two variable exponents $q : \overline{\Omega} \rightarrow \mathbb{R}$ and $p : \overline{\Omega} \times \overline{\Omega} \rightarrow \mathbb{R}$, where both $q(\cdot)$ and $p(\cdot, \cdot)$ are continuous function. We assume that:

$$p \text{ is symmetric, this is, } p(x, y) = p(y, x),$$

$$1 < \underline{p}^- := \inf_{(x,y) \in \overline{\Omega} \times \overline{\Omega}} p(x, y) \leq p(x, y) \leq \sup_{(x,y) \in \overline{\Omega} \times \overline{\Omega}} p(x, y) := \underline{p}^+ < +\infty, \quad (q_1)$$

and

$$1 < \underline{q}^- := \inf_{x \in \overline{\Omega}} q(x) \leq q(x) \leq \sup_{x \in \overline{\Omega}} q(x) := \underline{q}^+ < +\infty. \quad (q_2)$$

The fractional Sobolev space with variable exponents is defined by

$$W^{s, q(\cdot), p(\cdot, \cdot)}(\Omega) := \left\{ u \in L^{q(\cdot)}(\Omega) : \int_{\Omega \times \Omega} \frac{|u(x) - u(y)|^{p(x, y)}}{\zeta^{p(x, y)} |x - y|^{N + sp(x, y)}} dx dy < +\infty, \text{ for some } \zeta > 0 \right\}$$

and we set

$$[u]_{\Omega}^{s, p(\cdot, \cdot)} = \inf \left\{ \zeta > 0; \int_{\Omega \times \Omega} \frac{|u(x) - u(y)|^{p(x, y)}}{\zeta^{p(x, y)} |x - y|^{N + sp(x, y)}} dx dy \leq 1 \right\}$$

the variable exponent Gagliardo seminorm. It is already known that $W^{s, q(\cdot), p(\cdot, \cdot)}(\Omega)$ is a separable and reflexive Banach space with the norm

$$\|u\|_{W^{s, q(\cdot), p(\cdot, \cdot)}(\Omega)} := \|u\|_{L^{q(\cdot)}(\Omega)} + [u]_{\Omega}^{s, p(\cdot, \cdot)},$$

see [4, 8, 37].

Remark 2.2. Throughout this paper, when $q(x) = p(x, x)$ we denote $p(x)$ instead of $p(x, x)$ and we will write $W^{s, p(\cdot, \cdot)}(\Omega)$ instead of $W^{s, p(\cdot), p(\cdot, \cdot)}(\Omega)$.

The next result is an consequence of [29, Theorem 3.2].

Corollary 2.3. Let $\Omega \subset \mathbb{R}^N$ a bounded Lipschitz domain, $s \in (0, 1)$, $p(x, y)$ and $p(x)$ be continuous variable exponents such that (q_1) - (q_2) be satisfied and that $sp^+ < N$. Then, for all $r : \overline{\Omega} \rightarrow (1, +\infty)$ a continuous function such that $p_s^*(x) > r(x)$ for all $x \in \overline{\Omega}$, the space $W^{s, p(\cdot, \cdot)}(\Omega)$ is continuously and compactly embedding in $L^{r(\cdot)}(\Omega)$.

Now, we consider the space

$$W^{s, p(\cdot, \cdot)}(\mathbb{R}^N) := \left\{ u \in L^{p(\cdot)}(\mathbb{R}^N) : \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^{p(x, y)}}{\zeta^{p(x, y)} |x - y|^{N + sp(x, y)}} dx dy < +\infty, \text{ for some } \zeta > 0 \right\}$$

where the space $L^{p(\cdot)}(\mathbb{R}^N)$ is defined analogous the space $L^{p(\cdot)}(\Omega)$. The corresponding norm for this space is

$$\|u\| := \|u\|_{L^{p(\cdot)}(\mathbb{R}^N)} + [u]_{\mathbb{R}^N}^{s, p(\cdot, \cdot)}.$$

The space $(W^{s, p(\cdot, \cdot)}(\mathbb{R}^N), \|\cdot\|)$ has the same properties that $(W^{s, p(\cdot, \cdot)}(\Omega), \|\cdot\|_{W^{s, p(\cdot, \cdot)}(\Omega)})$, this is, it is a reflexive and separable Banach space.

We define the space were will study the problem (\mathscr{P}_λ) . Let we will consider the variable exponents $p(x) := p(x, x)$ for all $x \in \mathbb{R}^N$ with $p \in C(\mathbb{R}^N \times \mathbb{R}^N)$ satisfying (p_1) and we denote by

$$\mathscr{W} = W_0^{s, p(\cdot, \cdot)} := \{u \in W^{s, p(\cdot, \cdot)}(\mathbb{R}^N) : u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega\}.$$

Note that \mathscr{W} is a closed subspace of $W^{s, p(\cdot, \cdot)}(\mathbb{R}^N)$, thus \mathscr{W} is a reflexive and separable Banach space with the norm

$$\|u\| := \|u\|_{L^{p(\cdot)}(\Omega)} + [u]_{\mathbb{R}^N}^{s, p(\cdot, \cdot)},$$

once the norms $\|\cdot\|_{L^{p(\cdot)}(\mathbb{R}^N)}$ and $\|\cdot\|_{L^{p(\cdot)}(\Omega)}$ coincide in \mathscr{W} .

The next Lemma shows that the space $(\mathscr{W}, \|\cdot\|)$ is equivalently defined with respect to the Gagliardo seminorm $[\cdot]_{\mathbb{R}^N}^{s, p(\cdot, \cdot)}$. See proof in Appendix 4.

Lemma 2.4. *Assume Ω be a smooth bounded domain in \mathbb{R}^N . Let $p(x) := p(x, x)$ for all $x \in \mathbb{R}^N$ with $p \in C(\mathbb{R}^N \times \mathbb{R}^N)$ satisfying (p_1) and $p_s^*(x) > p(x)$ for $x \in \mathbb{R}^N$. Then there exists $\zeta_1 > 0$ such that*

$$\|u\|_{L^{p(\cdot)}(\Omega)} \leq \frac{1}{\zeta_1} [u]_{\mathbb{R}^N}^{s, p(\cdot, \cdot)} \text{ for all } u \in \mathcal{W}.$$

Therefore, we will consider the space \mathcal{W} with norm $\|u\|_{\mathcal{W}} = [u]_{\mathbb{R}^N}^{s, p(\cdot, \cdot)}$ and we proof an important result of compact and continuous embedding of space \mathcal{W} as consequence of Corollary 2.3 and Lemma 2.4. See proof of Lemma 2.5 in Appendix 4.

Lemma 2.5. *Assume Ω be a smooth bounded domain in \mathbb{R}^N . Let $p(x) := p(x, x)$ for all $x \in \mathbb{R}^N$ with $p \in C(\mathbb{R}^N \times \mathbb{R}^N)$ satisfying (p_1) and $p_s^*(x) > p(x)$ for $x \in \mathbb{R}^N$. Assume that $r : \overline{\Omega} \rightarrow (1, +\infty)$ is a continuous function. Thus, the space $(\mathcal{W}, \|\cdot\|_{\mathcal{W}})$ is continuously and compactly embedding in $L^{r(\cdot)}(\Omega)$ for all $r(x) \in (1, p_s^*(x))$ for all $x \in \overline{\Omega}$.*

An important role in manipulating the fractional Sobolev spaces with variable exponent is played by the $(s, p(\cdot, \cdot))$ -convex modular function $\rho_{\mathcal{W}} : \mathcal{W} \rightarrow \mathbb{R}$ defined by

$$\rho_{\mathcal{W}}(u) = \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^{p(x, y)}}{|x - y|^{N + sp(x, y)}} dx dy.$$

The following proposition show the relationship between the norm $\|\cdot\|_{\mathcal{W}}$ and the $\rho_{\mathcal{W}}$ convex modular function.

Proposition 2.6. *For $u \in \mathcal{W}$ and $(u_k)_{k \in \mathbb{N}} \subset \mathcal{W}$, we have*

- (a) *For $u \in \mathcal{W} \setminus \{0\}$, $\zeta = \|u\|_{\mathcal{W}}$ if and only if $\rho_{\mathcal{W}}\left(\frac{u}{\zeta}\right) = 1$;*
- (b) *$\|u\|_{\mathcal{W}} \geq 1 \Rightarrow \|u\|_{\mathcal{W}}^{p^-} \leq \rho_{\mathcal{W}}(u) \leq \|u\|_{\mathcal{W}}^{p^+}$;*
- (c) *$\|u\|_{\mathcal{W}} \leq 1 \Rightarrow \|u\|_{\mathcal{W}}^{p^+} \leq \rho_{\mathcal{W}}(u) \leq \|u\|_{\mathcal{W}}^{p^-}$;*
- (d) *$\lim_{k \rightarrow +\infty} \|u_k - u\|_{\mathcal{W}} = 0 \Leftrightarrow \lim_{k \rightarrow +\infty} \rho_{\mathcal{W}}(u_k - u) = 0$;*
- (e) *$\lim_{k \rightarrow +\infty} \|u_k\|_{\mathcal{W}} = +\infty \Leftrightarrow \lim_{k \rightarrow +\infty} \rho_{\mathcal{W}}(u_k) = +\infty$.*

2.3. Auxiliary Result.

Lemma 2.7. *Assume that (a_1) - (a_3) , and (\mathcal{K}) is hold. We consider the functional $\Phi : \mathcal{W} \rightarrow \mathbb{R}$ defined by*

$$\Phi(u) = \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A}(u(x) - u(y))K(x, y) dx dy \text{ for all } u \in \mathcal{W},$$

has the following properties:

- (i) *The functional Φ is well defined on \mathcal{W} , is of class $C^1(\mathcal{W}, \mathbb{R})$, and its Gâteaux derivative is given by*

$$\langle \Phi'(u), v \rangle = \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A}(u(x) - u(y))(v(x) - v(y))K(x, y) dx dy \text{ for all } u, v \in \mathcal{W};$$

- (ii) *The functional Φ is weakly lower semicontinuous, that is, $u_k \rightharpoonup u$ in \mathcal{W} as $k \rightarrow +\infty$ implies that $\Phi(u) \leq \liminf_{k \rightarrow +\infty} \Phi(u_k)$;*
- (iii) *The functional $\Phi' : \mathcal{W} \rightarrow \mathcal{W}'$ is an operator of type (S_+) on \mathcal{W} , that is, if*

$$u_k \rightharpoonup u \text{ in } \mathcal{W} \text{ and } \limsup_{k \rightarrow +\infty} \langle \Phi'(u_k), u_k - u \rangle \leq 0, \tag{2.2}$$

then $u_k \rightarrow u$ in \mathcal{W} as $k \rightarrow +\infty$.

The proof of the result above will be referred to in Appendix 4.

Definition 2.8. *We say that $u \in \mathcal{W}$ is a weak solution to the problem (\mathcal{P}_λ) if, and only if,*

$$\int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A}(u(x) - u(y))(v(x) - v(y))K(x, y) dx dy = \lambda \int_{\Omega} f(x, u)v dx \text{ for all } v \in \mathcal{W}.$$

The weak solutions to the problem (\mathcal{P}_λ) coincide with the critical points of the Euler Lagrange functional $\Psi_\lambda : \mathcal{W} \rightarrow \mathbb{R}$ given by

$$\Psi_\lambda(u) = \Phi(u) - \lambda \int_{\Omega} F(x, u) dx \text{ for all } u \in \mathcal{W}. \quad (2.3)$$

Moreover, by Lemma 2.7 and standard arguments, the functional Ψ_λ is Frchet differentiable in $u \in \mathcal{W}$ and

$$\langle \Psi'_\lambda(u), v \rangle = \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A}(u(x) - u(y))(v(x) - v(y))K(x, y) dx dy - \lambda \int_{\Omega} f(x, u)v dx \text{ for all } v \in \mathcal{W}.$$

2.4. Standard results for the existence and multiplicity of weak solutions.

In this subsection, we enunciated some definitions and general theorems of the existence and multiplicity of weak solutions that we will use along of the paper to prove our main results, stated in Theorem 1.1, Theorem 1.2, Theorem 1.3, Theorem 1.4, Theorem 1.5, and Theorem 1.6.

Definition 2.9. *Let X be a real Banach space and $\Psi \in C^1(X, \mathbb{R})$. We say that Ψ satisfies the $(C)_c$ condition if for every sequence $(u_k)_{k \in \mathbb{N}} \subset X$ such that $\Psi(u_k) \rightarrow c$ and $\|\Psi'(u_k)\|_{X'}(1 + \|u_k\|_X) \rightarrow 0$, as $k \rightarrow +\infty$, has a convergent subsequence.*

The condition $(C)_c$, introduced by Cerami in [16, 17], is a little more weak version of the PalaisSmale $(PS)_c$ condition, a condition more common that we find in the literature. Thus, since the Deformation Theorem is still valid under the Cerami a condition it follows that the Mountain Pass Theorem, Fountain Theorem, and Dual Fountain Theorem under PalaisSmale $(PS)_c$ and $(PS)_c^*$ condition holds true also under this compactness.

The results below Mountain Pass Theorem, " \mathbb{Z}_2 - symmetric" version (for even functionals) Mountain Pass Theorem, Fountain Theorem and Dual Fountain Theorem can be seen respectively in [19, Theorem I], [51, Theorem 9.12], [45, Theorem 2.9], and [9, Theorem 2].

Theorem 2.10. *Let X be a real Banach space, let $\Psi : X \rightarrow \mathbb{R}$ be a functional of class $C^1(X, \mathbb{R})$ that satisfies the $(C)_c$ condition for any $c > 0$, $\Psi(0) = 0$, and the following conditions hold:*

- (i) *There exist positive constants ρ and \mathcal{R} such that $\Psi(u) \geq \mathcal{R}$ for any $u \in X$ with $\|u\|_X = \rho$;*
- (ii) *There exists a function $e \in X$ such that $\|e\|_X > \rho$ and $\Psi(e) < 0$.*

Then, the functional Ψ has a critical value $c \geq \mathcal{R}$, that is, there exists $u \in X$ such that $\Psi(u) = c$ and $\Psi'(u) = 0$ in X' .

Theorem 2.11. *Assume that X has infinite dimension and let $\Psi \in C^1(X, \mathbb{R})$ be a functional satisfying the $(C)_c$ condition as well as the following properties*

- i) $\Psi(0) = 0$, and there exist two constants $r, \rho > 0$ such that $\Psi|_{\partial B_r} \geq \rho$;
- ii) Ψ is even;
- iii) *For all finite dimensional subspace $\widehat{X} \subset X$ there exists $\mathcal{R} = \mathcal{R}(\widehat{X}) > 0$ such that*

$$\Psi(u) \leq 0 \text{ for all } u \in \widehat{X} \setminus B_{\mathcal{R}}(\widehat{X})$$

where $B_{\mathcal{R}}(\widehat{X}) = \{u \in \widehat{X} : \|u\|_X < \mathcal{R}\}$.

Then Ψ possesses an unbounded sequence of critical values.

Let X be a real, reflexive, and separable Banach space, it is known ([23, Chapter 4] or [57, Section 17] or [26]) that for a separable and reflexive Banach space there exist sequence $(e_l)_{l \in \mathbb{N}} \subset X$ and $(e_l^*)_{l \in \mathbb{N}} \subset X^*$ such that

$$X = \overline{\text{span}\{e_l : l = 1, 2, \dots\}}, \quad X^* = \overline{\text{span}\{e_l^* : l = 1, 2, \dots\}}^{\omega^*},$$

and

$$\langle e_i^*, e_l \rangle = \begin{cases} 1 & \text{se } i = l, \\ 0 & \text{se } i \neq l. \end{cases}$$

We denote

$$X_l = \text{span}\{e_l\}, \quad Y_j = \bigoplus_{l=1}^j X_l = \text{span}\{e_1, \dots, e_j\}, \quad \text{and } Z_j = \overline{\bigoplus_{l=j}^{\infty} X_l} = \overline{\text{span}\{e_j, e_{j+1}, \dots\}}.$$

Theorem 2.12. *Assume*

- (h₁) X is a Banach space, $\Psi \in C^1(X, \mathbb{R})$ is an even functional;
 If for every $j \in \mathbb{N}$ there exist $\rho_j > r_j > 0$ such that
 (h₂) $b_j := \inf\{\Psi(u) : u \in Z_j, \|u\|_X = r_j\} \rightarrow +\infty$ as $j \rightarrow +\infty$;
 (h₃) $a_j := \sup\{\Psi(u) : u \in Y_j, \|u\|_X = \rho_j\} \leq 0$;
 (h₄) Ψ satisfies the $(C)_c$ condition for every $c > 0$.

Then Ψ has a sequence of critical points $(u_j)_{j \in \mathbb{N}}$ such that $\Psi(u_j) \rightarrow +\infty$.

Definition 2.13. Let X be a separable and reflexive Banach space, $\Psi \in C^1(X, \mathbb{R})$, $c \in \mathbb{R}$. We say that Ψ satisfies the $(C)_c^*$ condition (with respect to Y_k), if any sequence $(u_k)_{k \in \mathbb{N}} \subset X$ for which $u_k \in Y_k$, for any $k \in \mathbb{N}$, $\Psi(u_k) \rightarrow c$ and $\|\Psi'_{|_{Y_k}}(u_k)\|_{X^*}(1 + \|u_k\|_X) \rightarrow 0$, as $k \rightarrow \infty$, contain a subsequence converging to a critical point of Ψ .

Theorem 2.14. Suppose (h₁). If for each $j \geq j_0$ there exist $\rho_j > r_j > 0$ such that

- (g₁) $a_j = \inf\{\Psi(u) : u \in Z_j, \|u\|_X = \rho_j\} \geq 0$;
 (g₂) $b_j = \sup\{\Psi(u) : u \in Y_j, \|u\|_X = r_j\} < 0$;
 (g₃) $d_j = \inf\{\Psi(u) : u \in Z_j, \|u\|_X \leq \rho_j\} \rightarrow 0$, as $j \rightarrow +\infty$;
 (g₄) Ψ satisfies the $(C)_c^*$ condition for every $c \in [d_{j_0}, 0]$.

Then Ψ has a sequence of negative critical values converging to 0.

We conclude this subsection defining the notion of Krasnoselskii genus and introducing a critical point theorem related to the a new version of the symmetric Mountain Pass Theorem studied by Kajikiya, see [36, Theorem 1].

Definition 2.15. Let X be a real Banach space and B a subset of X . B is said to be symmetric if $u \in B$ implies $-u \in B$. For a closed symmetric set B which does not contain the origin, we define a Krasnoselskii genus $\gamma(B)$ of B by the smallest integer j such that there exists an odd continuous mapping from B to $\mathbb{R}^j \setminus \{0\}$. If there does not exist such a j , we define $\gamma(B) = +\infty$. Moreover, we set $\gamma(\emptyset) = 0$.

Let us consider the following set,

$$\Gamma_j = \{B_j \subset X : B_j \text{ is closed, symmetric and } 0 \notin B_j \text{ such that the genus } \gamma(B_j) \geq j\}$$

Theorem 2.16. Let X be an infinite-dimensional space, $B \in \Gamma_j$, and $\Psi \in C^1(X, \mathbb{R})$ satisfying the following conditions

- (I1) $\Psi(u)$ is even, bounded from below, $\Psi(0) = 0$ and $\Psi(u)$ satisfies the Palais-Smale condition;
 (I2) For each $j \in \mathbb{N}$, there exists an $B_j \in \Gamma_j$ such that $\sup_{u \in B_j} \Psi(u) < 0$.

Then either (R1) or (R2) below holds

- (R1) There exists a sequence $(u_j)_{j \in \mathbb{N}}$ such that $\Psi'(u_j) = 0$, $\Psi(u_j) < 0$ and $(u_j)_{j \in \mathbb{N}}$ converges to zero.
 (R2) There exist two sequences $(u_j)_{j \in \mathbb{N}}$ and $(v_j)_{j \in \mathbb{N}}$ such that $\Psi'(u_j) = 0$, $\Psi(u_j) < 0$, $u_j \neq 0$, $\lim_{j \rightarrow +\infty} u_j = 0$,
 $\Psi'(v_j) = 0$, $\Psi(v_j) < 0$, $\lim_{j \rightarrow +\infty} v_j = 0$, and $(v_j)_{j \in \mathbb{N}}$ converges to a non-zero limit.

Remark 2.17. From Theorem 2.16 we have a sequence $(u_j)_{j \in \mathbb{N}}$ of critical points such that $\Psi(u_j) \leq 0$, $u_j \neq 0$, and $\lim_{j \rightarrow +\infty} u_j = 0$.

3. PROOF OF THEOREMS 1.1, 1.2, 1.3, 1.4, 1.5, AND 1.6

3.1. Proof of Theorem 1.1.

To prove Theorem 1.1 we need of Lemmas below and Theorem 2.10.

Lemma 3.1. Assume (a_1) - (a_3) , (\mathcal{K}) , and (f_0) - (f_2) are holds. Then we have the following assertions:

- (i) There exists $v \in \mathcal{W}$, $v > 0$, such that $\Psi_\lambda(tv) \rightarrow -\infty$ as $t \rightarrow +\infty$;
 (ii) There exist $r > 0$ and $\mathcal{R} > 0$ such that $\Psi_\lambda(u) \geq \mathcal{R}$ for any $u \in \mathcal{W}$ with $\|u\|_{\mathcal{W}} = r$.

Proof. (i) From (f_1) , it follows that for any $\mathcal{C} > 0$ there exists a constant $c_{\mathcal{C}} > 0$ such that

$$F(x, t) \geq \mathcal{C}|t|^{p^+} - c_{\mathcal{C}} \quad \text{for all } (x, t) \in \Omega \times \mathbb{R}. \quad (3.1)$$

Take $v \in \mathscr{W}$ with $v > 0$, for $t > 1$, by (3.1), (a₁), (a₂), and (K), we obtain

$$\Psi_\lambda(tv) \leq t^{p^+} \left[\frac{C_{\mathcal{A}b_1}}{p^-} \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|v(x) - v(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy - \lambda \mathcal{C} \int_{\Omega} v^{p^+} dx \right] + \lambda c_{\mathcal{C}} |\Omega| \quad (3.2)$$

where $|\Omega|$ denotes the Lebesgue measure of Ω . Hence, from (3.2) and taking \mathcal{C} large enough such that

$$\frac{C_{\mathcal{A}b_1}}{p^-} \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|v(x) - v(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy - \lambda \mathcal{C} \int_{\Omega} v^{p^+} dx < 0,$$

we have

$$\lim_{t \rightarrow +\infty} \Psi_\lambda(tv) = -\infty,$$

which completes the proof of (i).

(ii) First, since the embeddings $\mathscr{W} \hookrightarrow L^{p^+}(\Omega)$ and $\mathscr{W} \hookrightarrow L^{\vartheta(x)}(\Omega)$ are continuous (Lemma 2.5), there exist positive constants c_2, c_3 , such that

$$\|u\|_{L^{p^+}(\Omega)} \leq c_2 \|u\|_{\mathscr{W}}, \quad \|u\|_{L^{\vartheta(\cdot)}(\Omega)} \leq c_3 \|u\|_{\mathscr{W}} \quad \text{for all } u \in \mathscr{W}. \quad (3.3)$$

Now, let $0 < \varepsilon < \frac{c_{\mathcal{A}b_0}}{\lambda c_2^{p^+}}$. From (f₀) and (f₂), it follows that for all given $\varepsilon > 0$, there exists $c_\varepsilon > 0$, such that

$$F(x, t) \leq \frac{\varepsilon}{p^+} |t|^{p^+} + c_\varepsilon |t|^{\vartheta(x)} \quad \text{for all } (x, t) \in \Omega \times \mathbb{R}. \quad (3.4)$$

Now, Thus, for $u \in \mathscr{W}$ with $\|u\|_{\mathscr{W}} < 1$ sufficiently small, from (a₂), (a₃), (K), (3.3), and (3.4), we obtain

$$\Psi_\lambda(u) \geq \frac{\|u\|_{\mathscr{W}}^{p^+}}{p^+} (c_{\mathcal{A}b_0} - \lambda c_2^{p^+}) - \lambda c_\varepsilon c_3^{\vartheta^-} \|u\|_{\mathscr{W}}^{\vartheta^-}. \quad (3.5)$$

Therefore, since $\vartheta^- > p^+$ from (3.5) we can choose $\mathcal{R} > 0$ and $r > 0$ such that $\Psi_\lambda(u) \geq \mathcal{R} > 0$ for every $u \in \mathscr{W}$ and $\|u\|_{\mathscr{W}} = r$. This completes the proof of (ii). \square

Lemma 3.2. *Assume that the condition (a₁)-(a₄), (K), and f satisfies (f₀), (f₁) and (f₃). Then the functional Ψ_λ satisfies the $(C)_c$ condition at any level $c > 0$.*

Proof. Let $c \in \mathbb{R}$ and $(u_k)_{k \in \mathbb{N}} \subset \mathscr{W}$ be a $(C)_c$ sequence for Ψ_λ , that is,

$$\Psi_\lambda(u_k) \rightarrow c > 0 \quad \text{and} \quad \|\Psi'_\lambda(u_k)\|_{\mathscr{W}'} (1 + \|u_k\|_{\mathscr{W}}) \rightarrow 0 \quad \text{as } k \rightarrow +\infty. \quad (3.6)$$

Initially, we prove that the sequence $(u_k)_{k \in \mathbb{N}}$ is bounded in \mathscr{W} . Indeed, arguing by contradiction, up to a subsequence, still denoted by $(u_k)_{k \in \mathbb{N}}$, we suppose that $(u_k)_{k \in \mathbb{N}}$ is unbounded in \mathscr{W} . Define $\omega_k := \frac{u_k}{\|u_k\|_{\mathscr{W}}}$ for all $k \in \mathbb{N}$, then $(\omega_k)_{k \in \mathbb{N}} \subset \mathscr{W}$ and $\|\omega_k\|_{\mathscr{W}} = 1$. Thus, we can extract a subsequence, still denoted by $(\omega_k)_{k \in \mathbb{N}}$ and $\omega \in \mathscr{W}$ such that $\omega_k \rightarrow \omega$ in \mathscr{W} as $k \rightarrow +\infty$. From Lemma 2.5, it follows that

$$\omega_k(x) \rightarrow \omega(x) \text{ a.e. } x \in \Omega, \quad \omega_k \rightarrow \omega \text{ in } L^{p^+}(\Omega), \quad \text{and } \omega_k \rightarrow \omega \text{ in } L^{\vartheta(\cdot)}(\Omega) \text{ as } k \rightarrow +\infty. \quad (3.7)$$

We consider $\Omega_\star := \{x \in \Omega : \omega(x) \neq 0\}$. If $x \in \Omega_\star$, by (3.7), we have

$$|u_k(x)| = |\omega_k(x)| \|u_k\|_{\mathscr{W}} \rightarrow +\infty \text{ a.e. } x \in \Omega_\star \text{ as } k \rightarrow +\infty.$$

Therefore, by (f₁), we obtain for each $x \in \Omega_\star$

$$\lim_{k \rightarrow +\infty} \frac{F(x, u_k)}{|u_k|^{p^+}} \frac{|u_k|^{p^+}}{\|u_k\|_{\mathscr{W}}^{p^+}} = \lim_{k \rightarrow +\infty} \frac{F(x, u_k)}{|u_k|^{p^+}} |\omega_k|^{p^+} = +\infty. \quad (3.8)$$

Also, by (f₁) there exists $D > 0$ such that

$$\frac{F(x, t)}{|t|^{p^+}} > 1 \quad \text{for all } (x, t) \in \Omega \times \mathbb{R} \text{ with } |t| \geq D. \quad (3.9)$$

Since $F(x, t)$ is continuous on $\overline{\Omega} \times [-D, D]$, there exists a positive constant c_5 such that

$$|F(x, t)| \leq c_5 \quad \text{for all } (x, t) \in \overline{\Omega} \times [-D, D]. \quad (3.10)$$

Hence, by (3.9) and (3.10), we conclude that there is a constant c_6 such that

$$F(x, t) \geq c_6 \quad \text{for all } (x, t) \in \overline{\Omega} \times \mathbb{R},$$

which shows that

$$\frac{F(x, u_k) - c_6}{\|u_k\|_{\mathcal{W}}^{p^+}} \geq 0 \quad \text{for all } x \in \Omega \text{ and } k \in \mathbb{N},$$

that is,

$$\frac{F(x, u_k)}{|u_k|^{p^+}} |\omega_k|^{p^+} - \frac{c_6}{\|u_k\|_{\mathcal{W}}^{p^+}} \geq 0 \quad \text{for all } x \in \Omega \text{ and } k \in \mathbb{N}. \quad (3.11)$$

Now, by (3.6), (a₂), (a₃), and (K), we have

$$c \geq \frac{c_A b_0}{p^+} \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u_k(x) - u_k(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy - \lambda \int_{\Omega} F(x, u_k) dx + o_k(1).$$

Then,

$$\int_{\Omega} F(x, u_k) dx \geq \frac{c_A b_0}{\lambda p^+} \|u_k\|_{\mathcal{W}}^{p^-} - \frac{c}{\lambda} + \frac{o_k(1)}{\lambda} \rightarrow +\infty \text{ as } k \rightarrow +\infty. \quad (3.12)$$

On the other hand, from (a₁), (a₂), and (K), we also that

$$c \leq \frac{C_A b_1}{p^-} \|u_k\|_{\mathcal{W}}^{p^+} - \lambda \int_{\Omega} F(x, u_k) dx + o_k(1) \text{ for all } k \in \mathbb{N}.$$

Consequently, by (3.12), we achieve

$$\|u_k\|_{\mathcal{W}}^{p^+} \geq \frac{c p^-}{C_A b_1} + \frac{\lambda p^-}{C_A b_1} \int_{\Omega} F(x, u_k) dx - \frac{p^-}{C_A b_1} o_k(1) > 0, \quad (3.13)$$

for k large enough.

We claim that $|\Omega_{\star}| = 0$. Indeed, if $|\Omega_{\star}| \neq 0$, then by (3.8), (3.11), (3.13), and Fatou's Lemma, we have

$$\begin{aligned} +\infty &= \int_{\Omega_{\star}} \liminf_{k \rightarrow +\infty} \frac{F(x, u_k)}{|u_k|^{p^+}} |\omega_k(x)|^{p^+} dx - \int_{\Omega_{\star}} \limsup_{k \rightarrow +\infty} \frac{c_6}{\|u_k\|_{\mathcal{W}}^{p^+}} dx \\ &\leq \liminf_{k \rightarrow +\infty} \frac{\int_{\Omega} F(x, u_k) dx}{\frac{\lambda p^-}{C_A b_1} \int_{\Omega} F(x, u_k) dx - o_k(1)}. \end{aligned} \quad (3.14)$$

Therefore, from (3.13) and (3.14), we obtain that $+\infty \leq \frac{C_A b_1}{\lambda p^-}$, which is a contradiction. This proves that $|\Omega_{\star}| = 0$ and thus $\omega(x) = 0$ a.e. $x \in \Omega$.

Now as in [?], we define the continuous function $B_k : [0, 1] \rightarrow \mathbb{R}$ by $B_k(t) := \Psi_{\lambda}(tu_k)$. Since $B_k(t) := \Psi_{\lambda}(tu_k)$ is continuous in $[0, 1]$, we can say that for each $k \in \mathbb{N}$ there exists $t_k \in [0, 1]$ such that

$$\Psi_{\lambda}(t_k u_k) := \max_{t \in [0, 1]} B_k(t). \quad (3.15)$$

(If for $k \in \mathbb{N}$ is not unique we choose the smaller possible value). Note that $t_k > 0$ for all $k \in \mathbb{N}$. Indeed, passing to a subsequence if necessary, we have $\Psi_{\lambda}(u_k) \geq \frac{c}{2}$ for all $k \in \mathbb{N}$. So, if $t_k = 0$ for all $k \in \mathbb{N}$ it follows that

$$\Psi_{\lambda}(t_k u_k) = \Psi_{\lambda}(0) = 0, \quad (3.16)$$

however,

$$0 < \frac{c}{2} \leq \Psi_{\lambda}(u_k) \leq \max_{t \in [0, 1]} \Psi_{\lambda}(tu_k) = \Psi_{\lambda}(t_k u_k). \quad (3.17)$$

Thus, from (3.16) and (3.17), we obtain a contradiction.

If $t_k \in (0, 1)$, by (3.15), we infer that

$$\frac{d}{dt} \Big|_{t=t_k} \Psi_{\lambda}(tu_k) = 0 \quad \text{for all } k \in \mathbb{N}.$$

Moreover, if $t_k = 1$, by (3.6) we have $\langle \Psi'_{\lambda}(u_k), u_k \rangle = o_k(1)$. So we always have

$$\langle \Psi'_{\lambda}(t_k u_k), t_k u_k \rangle = t_k \frac{d}{dt} \Big|_{t=t_k} \Psi_{\lambda}(tu_k) = o_k(1).$$

Let $(r_j)_{j \in \mathbb{N}}$ be a positive sequence of real numbers such that $r_j > 1$ and $\lim_{j \rightarrow +\infty} r_j = +\infty$. Then $\|r_j \omega_k\|_{\mathscr{W}} = r_j > 1$ for all j and $k \in \mathbb{N}$. Fix $j \in \mathbb{N}$, since $\omega_k \rightarrow 0$ in $L^{\vartheta(\cdot)}(\Omega)$ and $\omega_k(x) \rightarrow 0$ a.e. $x \in \Omega$, as $k \rightarrow +\infty$, using the condition (f_0) , there exists a positive constant c_7 such that

$$\left| F(x, r_j \omega_k) \right| \leq c_7 \left(r_j |\omega_k(x)| + r_j^{\vartheta(x)} |\omega_k(x)|^{\vartheta(x)} \right) \quad (3.18)$$

and by continuity of the function F , we achieve

$$F(x, r_j \omega_k) \rightarrow F(x, r_j \omega) = 0 \text{ a.e. } x \in \Omega \text{ as } k \rightarrow +\infty, \quad (3.19)$$

for each $j \in \mathbb{N}$. Consequently, from (3.18), (3.19), and the Dominated Convergence Theorem, we obtain

$$\lim_{k \rightarrow +\infty} \int_{\Omega} F(x, r_j \omega_k) dx = 0 \quad (3.20)$$

for all $j \in \mathbb{N}$.

Since $\|u_k\|_{\mathscr{W}} \rightarrow +\infty$ as $k \rightarrow +\infty$, we also have either $\|u_k\|_{\mathscr{W}} > r_j$ or $\frac{r_j}{\|u_k\|_{\mathscr{W}}} \in (0, 1)$ for k large enough. Thus, using (3.15), (3.20), (a_2) , (a_3) , (\mathcal{K}) , and Proposition 2.6, we deduce that

$$\Psi_{\lambda}(t_k u_k) \geq \Psi_{\lambda} \left(\frac{r_j}{\|u_k\|_{\mathscr{W}}} u_k \right) = \Psi_{\lambda}(r_j \omega_k) \geq \frac{c_{\mathcal{A}} b_0 r_j^{p^-}}{p^+} - \lambda \int_{\Omega} F(x, r_j \omega_k) dx \quad (3.21)$$

for all k large enough.

Therefore, by (3.21) letting $k, j \rightarrow +\infty$, we conclude

$$\limsup_{k \rightarrow +\infty} \Psi_{\lambda}(t_k u_k) = +\infty. \quad (3.22)$$

Now, we affirm that $\limsup_{k \rightarrow +\infty} \Psi_{\lambda}(t_k u_k) \leq \delta$, for a suitable positive constant δ . Indeed, from (a_4) , (f_3) , (3.6), and for all k large enough, we have

$$\begin{aligned} \Psi_{\lambda}(t_k u_k) &= \Psi_{\lambda}(t_k u_k) - \frac{1}{p^+} \langle \Psi'_{\lambda}((t_k u_k)), t_k u_k \rangle + o_k(1) \\ &= \frac{1}{p^+} \mathcal{H}(t_k u_k) + \frac{\lambda}{p^+} \int_{\Omega} \mathcal{G}(x, t_k u_k) dx + o_k(1) \\ &\leq \frac{1}{p^+} \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{H}(u_k) K(x, y) dx dy + \frac{\lambda}{p^+} \int_{\Omega} (\mathcal{G}(x, u_k) + c_{\star}) dx + o_k(1) \\ &= \Psi(u_k) - \frac{1}{p^+} \langle \Psi'_{\lambda}(u_k), u_k \rangle + \frac{\lambda c_{\star} |\Omega|}{p^+} + o_k(1). \end{aligned}$$

Then,

$$\Psi_{\lambda}(t_k u_k) \longrightarrow c + \frac{\lambda c_{\star}}{p^+} |\Omega| \text{ as } k \longrightarrow +\infty. \quad (3.23)$$

Consequently, from (3.22) and (3.23) we obtain a contradiction. Therefore, the sequence $(u_k)_{k \in \mathbb{N}}$ is bounded in \mathscr{W} .

Now, with standard arguments, we prove that any $(C)_c$ sequence has a convergent subsequence. Since \mathscr{W} is a reflexive Banach space there exists $u \in \mathscr{W}$ such that, up to a subsequence still denoted by $(u_k)_{k \in \mathbb{N}}$ we obtain that $u_k \rightarrow u$ in \mathscr{W} and by Lemma 2.5, we achieve

$$u_k(x) \rightarrow u(x) \text{ a.e. } x \in \Omega, \quad u_k \rightarrow u \text{ in } L^{\vartheta(\cdot)}(\Omega), \text{ and } u_k \rightarrow u \text{ in } L^{m(\cdot)}(\Omega) \text{ as } k \rightarrow +\infty.$$

Hence, using the Hölder's inequality, we have

$$\left| \int_{\Omega} f(x, u_k)(u_k - u) dx \right| \leq C \|1 + |u_k|^{\vartheta(x)-1}\|_{L^{\vartheta'(\cdot)}(\Omega)} \|u_k - u\|_{L^{\vartheta(\cdot)}(\Omega)} \rightarrow 0 \text{ as } k \rightarrow +\infty. \quad (3.24)$$

and from (3.6), it follows that

$$\langle \Psi'_{\lambda}(u_k), u_k - u \rangle \rightarrow 0 \text{ as } k \rightarrow +\infty. \quad (3.25)$$

Thus, using (3.24) and (3.25), we get

$$\langle \Phi'(u_k), u_k - u \rangle = \lambda \int_{\Omega} f(x, u_k)(u_k - u) dx + \langle \Psi'_{\lambda}(u_k), u_k - u \rangle \rightarrow 0 \text{ as } k \rightarrow +\infty.$$

Thus, since the operator Φ' is of type (S_+) (see Lemma 2.7), we conclude that $u_k \rightarrow u$ in \mathscr{W} . Therefore, this proves that Ψ_λ satisfies the $(C)_c$ condition on \mathscr{W} and we finish the proof of Lemma. \square

Proof of Theorem 1.1. From Lemma 3.1 and Lemma 3.2, the Euler Lagrange functional Ψ_λ satisfies the geometric conditions the Mountain Pass Theorem. Moreover $\Psi_\lambda(0) = 0$. Therefore, by Theorem 2.10, the functional Ψ_λ has a critical value $c \geq \mathcal{R} > 0$, that is, exists $u \in \mathscr{W}$ such that problem (\mathcal{P}_λ) has at least one nontrivial weak solution in \mathscr{W} . \square

3.2. Proof of Theorem 1.2.

To prove Theorem 1.2 we use some Lemmas presented below and Theorem 2.11 what is " \mathbb{Z}_2 - symmetric" version (for even functionals) of the Mountain Pass.

Lemma 3.3. *Assume (a_1) - (a_3) , (\mathcal{K}) , and (f_0) - (f_2) are fulfilled. Then for each $\lambda > 0$, there exist $\mathcal{R}_1 > 0$ and $r_1 > 0$ such that $\Psi_\lambda(u) \geq \mathcal{R}_1$ for all $u \in \mathscr{W}$ with $\|u\|_{\mathscr{W}} = r_1$.*

Proof. The proof is as in the Lemma 3.1. \square

Lemma 3.4. *Assume (a_1) - (a_3) , (\mathcal{K}) , and (f_1) are fulfilled. For every finite dimensional subspace $\widehat{\mathscr{W}} \subset \mathscr{W}$ there exists $\mathcal{R}_2 = \mathcal{R}_2(\widehat{\mathscr{W}})$ such that*

$$\Psi_\lambda(u) \leq 0 \text{ for all } u \in \widehat{\mathscr{W}} \setminus B_{\mathcal{R}_2}(\mathscr{W}),$$

where $B_{\mathcal{R}_2}(\mathscr{W}) = \{u \in \widehat{\mathscr{W}} : \|u\|_{\mathscr{W}} < \mathcal{R}_2\}$.

Proof. Consider $\widehat{\mathscr{W}}$ be a finite dimensional subspace of \mathscr{W} and let $u \in \widehat{\mathscr{W}}$ with $\|u\|_{\mathscr{W}} = 1$ fixed. Thus, for all $t \geq 1$ using (a_1) , (a_2) , (\mathcal{K}) , (f_1) , (3.1), and Proposition 2.6, we get

$$\Psi_\lambda(tu) \leq \frac{C_{\mathcal{A}} b_1}{p^-} t^{p^+} - \lambda \mathcal{C} c_{p^+} t^{p^+} + \lambda c_{\mathcal{E}} |\Omega|.$$

Taking $\Pi(t) = \left[\frac{C_{\mathcal{A}} b_1}{p^-} - \lambda \mathcal{C} c_{p^+} \right] t^{p^+} + \lambda c_{\mathcal{E}} |\Omega|$, if \mathcal{C} is large enough, then $\Pi(t) \rightarrow -\infty$ as $t \rightarrow +\infty$.

Therefore, we have

$$\sup \{ \Psi_\lambda(u) : u \in \widehat{\mathscr{W}}, \|u\|_{\mathscr{W}} = \mathcal{R}_3 \} = \sup \{ \Psi_\lambda(\mathcal{R}_3 u) : u \in \widehat{\mathscr{W}}, \|u\|_{\mathscr{W}} = 1 \} \rightarrow -\infty$$

as $\mathcal{R}_3 \rightarrow +\infty$.

Hence, there exists $\mathcal{R}_2 > 0$ sufficiently large such that $\Psi_\lambda(u) \leq 0$ for all $u \in \widehat{\mathscr{W}}$ with $\|u\|_{\mathscr{W}} = \mathcal{R}_3$ and $\mathcal{R}_3 \geq \mathcal{R}_2$. \square

Proof of Theorem 1.2. From Lemma 3.1, the Euler Lagrange functional Ψ_λ satisfies the $(C)_c$ condition. Moreover, $\Psi_\lambda(0) = 0$ and Ψ_λ is even functional by conditions (a_1) and (f_4) . Therefore, using the Lemma 3.3, Lemma 3.4, and Theorem 2.11 we conclude the existence of an unbounded sequence of weak solutions to problem (\mathcal{P}_λ) and this completes the proof. \square

3.3. Proof of Theorem 1.3.

To prove the Theorem 1.3 we verify the hypotheses of Theorem 2.12 through the Lemmas presented below.

Since that \mathscr{W} is a separable and reflexive Banach space, using ([23, Chapter 4], or [57, Section 17]) or [26], there exist sequence $(e_l)_{l \in \mathbb{N}} \subset \mathscr{W}$ and $(e_l^*)_{l \in \mathbb{N}} \subset \mathscr{W}'$ such that

$$\mathscr{W} = \overline{\text{span}\{e_l : l = 1, 2, \dots\}}, \quad \mathscr{W}' = \overline{\text{span}\{e_l^* : l = 1, 2, \dots\}}^{\omega^*},$$

and

$$\langle e_i^*, e_l \rangle = \begin{cases} 1 & \text{se } i = l, \\ 0 & \text{se } i \neq l. \end{cases}$$

We denote

$$\mathscr{W}_l = \text{span}\{e_l\}, \quad Y_j = \bigoplus_{l=1}^j \mathscr{W}_l = \text{span}\{e_1, \dots, e_j\}, \quad \text{and } Z_j = \bigoplus_{l=j}^{\infty} \mathscr{W}_l = \overline{\text{span}\{e_j, e_{j+1}, \dots\}}.$$

Lemma 3.5. *If $\vartheta \in C^+(\overline{\Omega})$, $p^+ < \vartheta(x) < p_s^{*-}$ for all $x \in \overline{\Omega}$, denote*

$$\beta_j := \sup\{\|u\|_{L^{\vartheta(\cdot)}(\Omega)} : \|u\|_{\mathcal{W}} = 1, u \in Z_j\},$$

then $\lim_{j \rightarrow +\infty} \beta_j = 0$.

Proof. It is clear that, $0 \leq \beta_{j+1} \leq \beta_j$, thus $\beta_j \rightarrow \beta \geq 0$ as $j \rightarrow +\infty$. Let $u_j \in Z_j$ satisfy

$$\|u_j\|_{\mathcal{W}} = 1, \quad 0 \leq \beta_j - \|u_j\|_{L^{\vartheta(\cdot)}(\Omega)} < \frac{1}{j} \text{ for } j \in \mathbb{N}.$$

Since \mathcal{W} is reflexive there exist a subsequence of $(u_j)_{j \in \mathbb{N}}$ such that $u_j \rightharpoonup u$ as $j \rightarrow +\infty$. We claim $u = 0$. Indeed, for all $e_m^* \in \{e_l^* : l = 1, 2, \dots\}$, we obtain $\langle e_m^*, u_j \rangle = 0$, $j > m$. So $\langle e_m^*, u_j \rangle \rightarrow 0$ as $j \rightarrow \infty$, this concludes $\langle e_m^*, u \rangle = 0$ for all $e_m^* \in \{e_l^* : l = 1, 2, \dots\}$. Therefore, $u = 0$ and so $u_j \rightarrow 0$ as $j \rightarrow +\infty$. Since the embedding from $\mathcal{W} \hookrightarrow L^{\vartheta(x)}(\Omega)$ is compact, then $u_j \rightarrow 0$ in $L^{\vartheta(x)}(\Omega)$ as $j \rightarrow +\infty$. Hence we get $\beta_j \rightarrow 0$ as $j \rightarrow +\infty$. \square

Lemma 3.6. *Assume that $\Xi : \mathcal{W} \rightarrow \mathbb{R}$ is weakly-strongly continuous, namely, $u_k \rightharpoonup u$ implies $\Xi(u_k) \rightarrow \Xi(u)$, and $\Xi(0) = 0$. Then for each $\tau > 0$ and $j \in \mathbb{N}$ there exists*

$$\alpha_j := \sup\{|\Xi(u)| : u \in Z_j, \|u\|_{\mathcal{W}} < \tau\} < \infty.$$

Moreover, $\lim_{j \rightarrow +\infty} \alpha_j = 0$.

Proof. It is clear that, $0 \leq \alpha_{j+1} \leq \alpha_j$, thus $\alpha_j \rightarrow \alpha \geq 0$ as $j \rightarrow +\infty$. Let $u_j \in Z_j$, $\|u_j\|_{\mathcal{W}} \leq \tau$ such that

$$0 \leq \alpha_j - |\Xi(u_j)| < \frac{1}{j}.$$

Since \mathcal{W} is reflexive there exist a subsequence of $(u_j)_{j \in \mathbb{N}}$ such that $u_j \rightharpoonup u$. Proceeding as in the previous lemma, we obtain $u = 0$. The weak-strong continuity of Ξ , guarantees $\Xi(u_j) \rightarrow \Xi(0) = 0$. Therefore, $\alpha_j \rightarrow 0$ as $j \rightarrow +\infty$. \square

Lemma 3.7. *Assume (a_1) - (a_3) , (\mathcal{K}) , (f_0) , and (f_2) are satisfied. Moreover, let $\lambda \in \left(0, \frac{c_{\mathcal{A}} b_0}{p^+}\right)$. Then there exist $\rho_j > r_j > 0$ such that:*

- (h₂) $b_j := \inf\{\Psi_\lambda(u) : u \in Z_j, \|u\|_{\mathcal{W}} = r_j\} \rightarrow +\infty$ as $j \rightarrow +\infty$;
- (h₃) $a_j := \max\{\Psi_\lambda(u) : u \in Y_j, \|u\|_{\mathcal{W}} = \rho_j\} \leq 0$.

Proof. (h₂) First, notice that for all $u \in Z_j$ with $\|u\|_{\mathcal{W}} > 1$, using (a_2) , (a_3) , (\mathcal{K}) , and (f_0) , we infer

$$\begin{aligned} \Psi_\lambda(u) &\geq \frac{c_{\mathcal{A}} b_0}{p^+} \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy - \lambda c_1 \int_{\Omega} \left(|u| + \frac{|u|^{\vartheta(x)}}{\vartheta(x)} \right) dx \\ &\geq \frac{c_{\mathcal{A}} b_0}{p^+} \|u\|_{\mathcal{W}}^{p^-} - \lambda \frac{c_1 \beta_j^{\vartheta^+}}{\vartheta^-} \|u\|_{\mathcal{W}}^{\vartheta^+} - \lambda c_8 \|u\|_{\mathcal{W}}, \end{aligned} \quad (3.26)$$

for a constant $c_8 > 0$ and $\beta_j := \sup\{\|u\|_{L^{\vartheta(x)}(\Omega)} : \|u\|_{\mathcal{W}} = 1, u \in Z_j\}$.

Now, since $p^- \leq p^+ < \vartheta^+$, by the Lemma 3.5, it is easy see that $r_j := (c_1 \beta_j^{\vartheta^+})^{\frac{1}{p^- - \vartheta^+}} \rightarrow +\infty$ as $j \rightarrow +\infty$. Then, for j sufficiently large, $u \in Z_j$ with $\|u\|_{\mathcal{W}} = r_j > 1$, and by (3.26), we conclude

$$\Psi_\lambda(u) \geq \left(\frac{c_{\mathcal{A}} b_0}{p^+} - \lambda \right) r_j^{p^-} - \lambda c_8 r_j.$$

Therefore, since $p^- > 1$ and $\frac{c_{\mathcal{A}} b_0}{p^+} > \lambda$, we obtain that $b_j := \inf\{\Psi_\lambda(u) : u \in Z_j, \|u\|_{\mathcal{W}} = r_j\} \rightarrow +\infty$ as $j \rightarrow +\infty$.

(h₃) Note that for all $v \in Y_j$ with $\|v\|_{\mathcal{W}} = 1$, using (a_2) , (a_3) , (\mathcal{K}) , (f_0) and (3.1) for $t > 1$, we have

$$\Psi_\lambda(tv) \leq t^{p^+} \left(\frac{C_{\mathcal{A}} b_1}{p^-} \|v\|_{\mathcal{W}}^{p^+} - \lambda \mathcal{E} \int_{\Omega} |v|^{p^+} dx \right) + \lambda c_{\mathcal{E}} |\Omega|. \quad (3.27)$$

It is clear that we can choose $\mathcal{C} > 0$ large enough such that

$$\frac{C_{\mathcal{A}} b_1}{p^-} \|v\|_{\mathcal{W}}^{p^+} - \lambda \mathcal{C} \int_{\Omega} |v|^{p^+} dx < 0.$$

From (3.27), it follows that

$$\lim_{t \rightarrow +\infty} \Psi_{\lambda}(tv) = -\infty.$$

Therefore, there exists $t_0 > r_j > 1$ large enough such that $\Psi_{\lambda}(t_0 v) \leq 0$ and thus, if set $\rho_j = t_0$, we conclude that

$$a_j := \max\{\Psi_{\lambda}(u) : u \in Y_j; \|u\|_{\mathcal{W}} = \rho_j\} \leq 0.$$

□

Proof of Theorem 1.3. From conditions (a_1) and (f_4) the Euler Lagrange functional Ψ_{λ} is even functional and by Lemma 3.2, Ψ_{λ} satisfies the $(C)_c$ condition for every $c > 0$. Then, with that and by Lemma 3.7 all conditions of the Theorem 2.12 are satisfied. Therefore, we obtain a sequence of critical points $(u_k)_{k \in \mathbb{N}}$ in \mathcal{W} such that $\Psi_{\lambda}(u_k) \rightarrow +\infty$ as $k \rightarrow +\infty$. □

3.4. Proof of Theorem 1.4.

To prove Theorem 1.4 we use the Lemma presented below and we verify hypotheses Theorem 2.14.

Lemma 3.8. *Suppose that the hypotheses in Theorem 1.4 hold, then Ψ_{λ} satisfied the $(C)_c^*$ condition.*

Proof. Let $c \in \mathbb{R}$ and the sequence $(u_k)_{k \in \mathbb{N}} \subset \mathcal{W}$ be such that $u_k \in Y_k$ for all $k \in \mathbb{N}$, $\Psi_{\lambda}(u_k) \rightarrow c$ and $\|\Psi'_{\lambda|_{Y_{k_j}}}(u_k)\|_{\mathcal{W}'}(1 + \|u_k\|_{\mathcal{W}}) \rightarrow 0$, as $k \rightarrow +\infty$. Therefore, we have

$$c = \Psi_{\lambda}(u_k) + o_k(1) \text{ and } \langle \Psi'_{\lambda}(u_k), u_k \rangle = o_k(1),$$

where $o_k(1) \rightarrow 0$ as $k \rightarrow +\infty$. Analogously to the proof of Lemma 3.2, we can prove that the sequence $(u_k)_{k \in \mathbb{N}}$ is bounded in \mathcal{W} . Since \mathcal{W} is reflexive, we can extract a subsequence of $(u_k)_{k \in \mathbb{N}}$, denoted for $(u_{k_j})_{j \in \mathbb{N}}$, and $u \in \mathcal{W}$ such that $u_{k_j} \rightharpoonup u$ in \mathcal{W} as $j \rightarrow +\infty$.

On the other hand, as $\mathcal{W} = \overline{\cup_k Y_k} = \overline{\text{span}\{e_k : k \geq 1\}}$, we can choose $v_k \in Y_k$ such that $v_k \rightarrow u$ in \mathcal{W} as $k \rightarrow +\infty$. Hence, we conclude that

$$\langle \Psi'_{\lambda}(u_{k_j}), u_{k_j} - u \rangle = \langle \Psi'_{\lambda}(u_{k_j}), u_{k_j} - v_{k_j} \rangle + \langle \Psi'_{\lambda}(u_{k_j}), v_{k_j} - u \rangle.$$

Since $\Psi'_{\lambda|_{Y_{k_j}}}(u_{k_j}) \rightarrow 0$ and $u_{k_j} - v_{k_j} \rightarrow 0$ in Y_{k_j} , as $j \rightarrow +\infty$, we achieve

$$\lim_{j \rightarrow +\infty} \langle \Psi'_{\lambda}(u_{k_j}), u_{k_j} - u \rangle = 0.$$

Furthermore, using Hölder's inequality, we obtain that

$$\int_{\Omega} f(x, u_{k_j})(u_{k_j} - u) dx \rightarrow 0 \text{ as } j \rightarrow +\infty.$$

Therefore,

$$\langle \Phi'(u_{k_j}), u_{k_j} - u \rangle = \lambda \int_{\Omega} f(x, u_{k_j})(u_{k_j} - u) dx + \langle \Psi'_{\lambda}(u_{k_j}), u_{k_j} - u \rangle \rightarrow 0 \text{ as } j \rightarrow +\infty.$$

Consequently, since Φ' is of type (S_+) , it follows that $u_{k_j} \rightarrow u$ in \mathcal{W} as $j \rightarrow +\infty$. Then, we conclude that Ψ_{λ} satisfies the $(C)_c^*$ condition. Thus, we obtain that $\Psi'_{\lambda}(u_{k_j}) \rightarrow \Psi'_{\lambda}(u)$ as $j \rightarrow +\infty$.

Let us prove $\Psi'_{\lambda}(u) = 0$. Indeed, taking $\omega_l \in Y_l$, notice that when $k_j \geq l$, we achieve

$$\begin{aligned} \langle \Psi'_{\lambda}(u), \omega_l \rangle &= \langle \Psi'_{\lambda}(u) - \Psi'_{\lambda}(u_{k_j}), \omega_l \rangle + \langle \Psi'_{\lambda}(u_{k_j}), \omega_l \rangle \\ &= \langle \Psi'_{\lambda}(u) - \Psi'_{\lambda}(u_{k_j}), \omega_l \rangle + \langle \Psi'_{\lambda|_{Y_{k_j}}}(u_{k_j}), \omega_l \rangle, \end{aligned}$$

so, passing the limit on the right side of the equation above as $j \rightarrow +\infty$, we conclude

$$\langle \Psi'_{\lambda}(u), \omega_l \rangle = 0 \text{ for all } \omega_l \in Y_l.$$

Therefore, $\Psi'_{\lambda}(u) = 0$ in \mathcal{W}' and this show that Ψ_{λ} satisfies the $(C)_c^*$ condition for every $c \in \mathbb{R}$. □

Proof of Theorem 1.4. First we observe that from (a_1) , (f_3) , and Lemma 3.8 the Euler Lagrange functional Ψ_λ is even functional and satisfies the $(C)_c^*$ condition for all $c \in \mathbb{R}$.

Now we will show that the conditions (g_1) , (g_2) , and (g_3) of the Dual Fountain Theorem are satisfied:

(g_1) First we note that using (f_0) and Young's inequality, we have

$$|F(x, t)| \leq c_9(1 + |t|^{\vartheta(x)}) \quad (3.28)$$

for a positive constant c_9 . Moreover, as $\lambda < \frac{c_{\mathcal{A}}b_0}{p^+}$, we have

$$\lim_{j \rightarrow +\infty} \left(\frac{c_{\mathcal{A}}b_0}{p^+} - \lambda \right) (c_9\beta_j^{\vartheta^+})^{\frac{p^-}{p^- - \vartheta^+}} = +\infty.$$

Then, there exists $j_0 \in \mathbb{N}$ such that

$$\left(\frac{c_{\mathcal{A}}b_0}{p^+} - \lambda \right) (c_9\beta_j^{\vartheta^+})^{\frac{p^-}{p^- - \vartheta^+}} - \lambda c_9|\Omega| \geq 0 \quad \text{for all } j \geq j_0.$$

Taking $\rho_j = (c_9\beta_j^{\vartheta^+})^{\frac{1}{p^- - \vartheta^+}}$ for $j \geq j_0$. It is clear that $\rho_j > 1$ for all $j \in \mathbb{N}$, $j \geq j_0$, since $\lim_{j \rightarrow +\infty} \rho_j = +\infty$.

Using same the arguments of Theorem 1.3, for all $u \in Z_j$, with $\|u\|_{\mathcal{W}} = \rho_j$, and using (a_2) , (a_3) , (\mathcal{K}) , and (3.28), we conclude that

$$\Psi_\lambda(u) \geq \left(\frac{c_{\mathcal{A}}b_0}{p^+} - \lambda \right) (c_9\beta_j^{\vartheta^+})^{\frac{p^-}{p^- - \vartheta^+}} - \lambda c_9|\Omega| \geq 0.$$

So, we obtain

$$a_j := \inf\{\Psi_\lambda(u) : u \in Z_j, \|u\|_{\mathcal{W}} = \rho_j\} \geq 0.$$

(g_2) Since Y_j is finite dimensional all the norms are equivalent, there exists a constant $c_{10} > 0$ such that $\|u\|_{L^{p^+}(\Omega)} \geq c_{10}\|u\|_{\mathcal{W}}$ for all $u \in Y_j$. Then, from (a_1) , (a_2) , (\mathcal{K}) , (f_1) , (3.1), and Proposition 2.6, we infer

$$\Psi_\lambda(u) \leq \frac{C_{\mathcal{A}}b_1}{p^-} \|u\|_{\mathcal{W}}^{p^+} - \lambda c_{10} \mathcal{E}^{p^+} \|u\|_{\mathcal{W}}^{p^+} + \lambda c_{\mathcal{E}} |\Omega| \quad \text{for all } u \in Y_j \text{ with } \|u\|_{\mathcal{W}} \geq 1.$$

Let $\mathcal{B}(t) = \frac{C_{\mathcal{A}}b_1}{p^-} t^{p^+} - \lambda c_{10} \mathcal{E}^{p^+} t^{p^+} + \lambda c_{\mathcal{E}} |\Omega|$. However, we can choose $\mathcal{E} > 0$ large enough, such that

$$\lim_{t \rightarrow +\infty} \mathcal{B}(t) = -\infty.$$

Therefore, there exists $\bar{t} \in (1, +\infty)$ such that

$$\mathcal{B}(t) < 0 \quad \text{for all } t \in [\bar{t}, +\infty).$$

Consequently, $\Psi_\lambda(u) < 0$ for all $u \in Y_j$ with $\|u\|_{\mathcal{W}} = \bar{t}$. Then, choosing $r_j = \bar{t}$ for all $j \in \mathbb{N}$, we have

$$b_j := \max\{\Psi_\lambda(u) : u \in Y_j, \|u\|_{\mathcal{W}} = r_j\} < 0.$$

We observe that we can change j_0 on other more large, if necessary, so that $\rho_j > r_j > 0$ for all $j \geq j_0$.

(g_3) Since $Y_j \cap Z_j \neq \emptyset$ and $0 < r_j < \rho_j$, so, we have

$$d_j := \inf\{\Psi_\lambda(u) : u \in Z_j, \|u\|_{\mathcal{W}} \leq \rho_j\} \leq b_j := \max\{\Psi_\lambda(u) : u \in Y_j, \|u\|_{\mathcal{W}} = r_j\} < 0.$$

From (f_0) , we obtain $|F(x, u)| \leq c_{11}(|t| + |t|^{\vartheta(x)})$, for a constant positive c_{11} , and for all $(x, t) \in \Omega \times \mathbb{R}$. Consider, $\Sigma_1 : \mathcal{W} \rightarrow \mathbb{R}$ and $\Sigma_2 : \mathcal{W} \rightarrow \mathbb{R}$ defined by

$$\Sigma_1(u) = \int_{\Omega} \lambda c_{11} |u|^{\vartheta(x)} dx \quad \text{and} \quad \Sigma_2(u) = \int_{\Omega} \lambda c_{11} |u| dx. \quad (3.29)$$

We have $\Sigma_i(0) = 0$, $i = 1, 2$, and they are weakly-strongly continuous. Let us denote

$$\eta_j = \sup\{|\Sigma_1(u)| : u \in Z_j, \|u\|_{\mathcal{W}} \leq 1\}, \quad \xi_j = \sup\{|\Sigma_2(u)| : u \in Z_j, \|u\|_{\mathcal{W}} \leq 1\}.$$

Thus, since the embedding $\mathcal{W} \hookrightarrow L^{\vartheta(\cdot)}(\Omega)$ is compact it follows that by Lemma 3.6 that $\lim_{j \rightarrow +\infty} \eta_j = \lim_{j \rightarrow +\infty} \xi_j = 0$.

Now, consider $v \in Z_j$ with $\|v\|_{\mathscr{W}} = 1$ and $0 < t < \rho_j$. Then, from (a_1) , (a_2) , (a_3) , (\mathcal{K}) , and (3.29), we obtain

$$\Psi_\lambda(tv) \geq -\lambda \int_{\Omega} F(x, tv) dx \geq -\Sigma_1(tv) - \Sigma_2(tv),$$

and since

$$\Sigma_1(tv) \leq t^{\varrho^+} \Sigma_1(v) \text{ and } \Sigma_2(tv) = t \Sigma_2(v),$$

we achieve

$$\Psi_\lambda(tv) \geq -\rho_j^{\varrho^+} \Sigma_1(v) - \rho_j \Sigma_2(v) \geq -\rho_j^{\varrho^+} \eta_j - \rho_j \xi_j$$

for all $t \in (0, \rho_j)$ and $v \in Z_j$ with $\|v\|_{\mathscr{W}} = 1$. Thus, $d_j \geq -\rho_j^{\varrho^+} \eta_j - \rho_j \xi_j$ and as $d_j < 0$ for all $j \geq j_0$, we conclude that $\lim_{j \rightarrow +\infty} d_j = 0$.

Therefore, the conditions of Theorem 2.14 are satisfied. Consequently, there exists a sequence $(u_k)_{k \in \mathbb{N}} \subset \mathscr{W}$ of weak solutions of problem such that $\Psi_\lambda(u_k) < 0$ and $\Psi_\lambda(u_k) \rightarrow 0$ as $k \rightarrow +\infty$ for $\lambda \in \left(0, \frac{c_A b_0}{p^+}\right)$. \square

3.5. Proof of Theorem 1.5.

To prove Theorem 1.1 we need of Lemmas below and Theorem 2.10.

Lemma 3.9. *Suppose (a_1) - (a_3) and (f_1) holds. There is $v \in \mathscr{W} \setminus \{0\}$, such that $\lim_{t \rightarrow +\infty} \Psi_\lambda(tv) = -\infty$.*

Proof. The proof is as in the Lemma 3.1-(i). \square

Lemma 3.10. *Suppose (a_1) - (a_3) , (\mathcal{K}) , (f_0) , $f(x, 0) = 0$, and $f(x, t) \geq 0$ a.e. $x \in \Omega$ and for all $t \geq 0$ holds. Then, there exist $\bar{\lambda} > 0$, positive constants C_λ and ρ_λ for $\lambda \in (0, \bar{\lambda})$ such that $\lim_{\lambda \rightarrow 0^+} C_\lambda = +\infty$ and $\Psi_\lambda(u) > C_\lambda > 0$ whenever $\|u\|_{\mathscr{W}} = \rho_\lambda$.*

Proof. We consider $u \in \mathscr{W}$ with $\|u\|_{\mathscr{W}} > 1$. Then, using (3.28), Young's inequality, and Lemma 2.6, we conclude that

$$\Psi_\lambda(u) \geq \frac{c_A b_0}{p^+} \|u\|^{p^-} - \lambda c_{13} \|u\|^{\varrho^+} - \lambda c_9 |\Omega| \quad (3.30)$$

for constant positive c_{13} .

Let $\gamma \in \left(0, \frac{1}{\varrho^+ - p^-}\right)$ and $u \in \mathscr{W}$ such that $\|u\|_{\mathscr{W}} = \lambda^{-\gamma}$. We define $\rho_\lambda := \lambda^{-\gamma}$ and we observe that $\rho_\lambda > 1$ for λ small enough. Hence, from (3.30), we conclude

$$\Psi_\lambda(u) \geq \frac{c_A b_0}{p^+} \lambda^{-\gamma p^-} - c_{13} \lambda^{1-\gamma \varrho^+} - \lambda c_9 |\Omega|.$$

Since $\gamma < \frac{1}{\varrho^+ - p^-}$ it follows that $-\gamma p^- < 1 - \gamma \varrho^+$. Thus $C_\lambda := \frac{c_A b_0}{p^+} \lambda^{-\gamma p^-} - c_{13} \lambda^{1-\gamma \varrho^+} - \lambda c_9 |\Omega| \rightarrow +\infty$ as $\lambda \rightarrow 0^+$. Hence, there exists $\bar{\lambda} > 0$ small enough such that $C_\lambda > 0$ for all $\lambda \in (0, \bar{\lambda})$. Then we obtain that

$$\Psi_\lambda(u) \geq C_\lambda > 0 = \Psi_\lambda(0)$$

for all $u \in \mathscr{W}$ with $\|u\|_{\mathscr{W}} = \rho_\lambda = \lambda^{-\gamma}$ and $\lambda \in (0, \bar{\lambda})$. Therefore, C_λ verifies the assertion of the Lemma. \square

Proof of Theorem 1.5. From Lemma 3.2 the functional Ψ_λ satisfy the $(C)_c$ condition. Moreover, $\Psi_\lambda(0) = 0$. Then, by Lemma 3.9, Lemma 3.10, and Theorem 2.10 we obtain that there are constant $\bar{\lambda}$ and a nontrivial critical point u_λ for Ψ_λ with $\lambda \in (0, \bar{\lambda})$ such that

$$c = \Psi_\lambda(u_\lambda) \geq C_\lambda.$$

Then, from (a_1) , (a_2) , (\mathcal{K}) , (f_0) , (3.28), Proposition 2.6, and Lemma 2.5, we achieve

$$C_\lambda \leq \Psi_\lambda(u_\lambda) \leq \frac{C_A b_1}{p^-} \max\{\|u_\lambda\|_{\mathscr{W}}^{p^+}, \|u_\lambda\|_{\mathscr{W}}^{p^-}\} + \lambda c_{13} \max\{\|u_\lambda\|_{\mathscr{W}}^{\varrho^+}, \|u_\lambda\|_{\mathscr{W}}^{\varrho^-}\} + \lambda c_9 |\Omega|. \quad (3.31)$$

Hence, taking $\lambda \rightarrow 0^+$ in (3.31) as $C_\lambda \rightarrow +\infty$, we get

$$\lim_{\lambda \rightarrow 0^+} \|u_\lambda\|_{\mathscr{W}} = +\infty.$$

Therefore, we conclude the proof of Theorem 1.5. \square

3.6. Proof of Theorem 1.6.

To proof Theorem 1.6 it is enough to verify that Ψ_λ satisfies the hypotheses of Theorem 2.16.

Lemma 3.11. *Suppose (a_1) - (a_3) , (\mathcal{K}) , and (f_5) holds. Then the functional Ψ_λ is bounded from below and satisfies the (PS) condition.*

Proof. Let $u \in \mathscr{W}$ and suppose $\|u\|_{\mathscr{W}} > 1$. Then, from (a_2) , (a_3) , (\mathcal{K}) , (f_5) , Lemma 2.5 and Lemma 2.7 for each $\lambda > 0$, we infer that

$$\Psi_\lambda(u) \geq \frac{c_A b_0}{p^-} \|u\|_{\mathscr{W}}^{p^-} - \frac{\lambda c_{14}}{\underline{m}^-} \|u\|_{\mathscr{W}}^{\underline{m}^+}.$$

Since $\underline{m}^+ < p^-$, follows that Ψ_λ is coercive. Therefore, Ψ_λ is bounded from below. Now, let $(u_k)_{k \in \mathbb{N}}$ be a sequence in \mathscr{W} such that

$$\Psi_\lambda(u_k) \rightarrow c \text{ and } \Psi'_\lambda(u_k) \rightarrow 0 \text{ in } \mathscr{W}' \text{ as } k \rightarrow +\infty.$$

By coercivity of Ψ_λ , we have $(u_k)_{k \in \mathbb{N}}$ is bounded in \mathscr{W} . Hence up to a subsequence, still denoted by $(u_k)_{k \in \mathbb{N}}$, we have $u_0 \in \mathscr{W}$ such that $u_k \rightharpoonup u_0$ in \mathscr{W} and from Lemma 2.6 we infer that $u_k \rightarrow u_0$ in $L^{q(\cdot)}(\Omega)$ and $u_k(x) \rightarrow u_0(x)$ a.e. $x \in \Omega$, as $k \rightarrow +\infty$.

Since $\Psi'_\lambda(u_k) \rightarrow 0$ in \mathscr{W}' as $k \rightarrow +\infty$ and the sequence $(u_k)_{k \in \mathbb{N}}$ is bounded in \mathscr{W} , we have that

$$\langle \Psi'_\lambda(u_k), u_k - u_0 \rangle \rightarrow 0 \text{ as } k \rightarrow +\infty. \quad (3.32)$$

Now, using (f_5) and Hölder's inequality, we obtain that

$$\left| \int_{\Omega} f(x, u_k)(u_k - u_0) dx \right| \leq C_2 \left\| |u_k|^{m(x)-1} \right\|_{L^{\frac{m(\cdot)}{m(\cdot)-1}}(\Omega)} \|u_k - u_0\|_{L^{m(\cdot)}(\Omega)}.$$

Thus, taking into account that $u_k \rightarrow u_0$ in $L^{m(\cdot)}(\Omega)$ as $k \rightarrow +\infty$, we achieve

$$\int_{\Omega} f(x, u_k)(u_k - u_0) dx \rightarrow 0 \text{ as } k \rightarrow +\infty. \quad (3.33)$$

Therefore, from (3.32) and (3.33), we conclude that

$$\langle \Phi'(u_k), u_k - u_0 \rangle = \langle \Psi'_\lambda(u_k), u_k - u_0 \rangle - \lambda \int_{\Omega} f(x, u_k)(u_k - u_0) dx \rightarrow 0 \text{ as } k \rightarrow +\infty.$$

Since Φ' is of type (S_+) (see Lemma 2.7), we obtain $u_k \rightarrow u_0$ in \mathscr{W} as $k \rightarrow +\infty$, concluding the proof of the Palais-Smale condition. \square

Proof of Theorem 1.6. From (a_1) and (f_4) the Euler Lagrange functional Ψ_λ is even functional. Furthermore, by Lemma 3.11 Ψ_λ is bounded from below and satisfies the Palais-Smale condition. Hence the item (I1) of Theorem 2.16 is verified. Let us show that Ψ_λ satisfies (I2). Since \mathscr{W} is a reflexive and separable Banach space, for each $j \in \mathbb{N}$, consider a j -dimensional linear subspace $\mathscr{W}_j \subset C_0^\infty(\Omega)$ of \mathscr{W} . We define

$$\mathbb{S}_{\mathcal{R}_4}^j = \{u \in \mathscr{W}_j : \|u\|_{\mathscr{W}} = \mathcal{R}_4\}$$

where $\mathcal{R}_4 > 0$ will be determined later on. Since \mathscr{W}_j and \mathbb{R}^j are isomorphic and $\mathbb{S}_{\mathcal{R}_4}^j$ is homeomorphic to the $(j-1)$ -dimensional sphere \mathbb{S}^{j-1} by an odd mapping, it has genus j , i.e., $\gamma(\mathbb{S}_{\mathcal{R}_4}^j) = j$.

Now, from (a_1) , (a_2) , (\mathcal{K}) , and (f_5) , we obtain

$$\Psi_\lambda(u) \leq \frac{C_A b_1}{p^-} \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy - \lambda C_0 \int_{\Omega} |u|^{m(x)} dx. \quad (3.34)$$

By Proposition 2.6, if $\|u\|_{\mathscr{W}} < 1$, we have $\rho_{\mathscr{W}}(u) \leq \|u\|_{\mathscr{W}}^{p^-}$, and $\rho_{m(\cdot)}(u) \geq \|u\|_{L^{m(\cdot)}(\Omega)}^{\underline{m}^+}$ for every $u \in \mathscr{W}$. Moreover, since \mathscr{W}_j is a finite dimensional space, any norm in \mathscr{W}_j is equivalent to each other. Thus, there exist a constant $C(j) > 0$ such that $C(j)\|u\|_{\mathscr{W}}^{\underline{m}^+} \leq \int_{\Omega} |u|^{m(x)} dx$ for every $u \in \mathscr{W}_j$. Consequently, by (3.34), we get

$$\Psi_\lambda(u) \leq \|u\|_{\mathscr{W}}^{\underline{m}^+} \left(\frac{C_A b_1}{p^-} \|u\|_{\mathscr{W}}^{p^- - \underline{m}^+} - \lambda C(j) C_0 \right),$$

for every $u \in \mathscr{W}_j$ with $\|u\|_{\mathscr{W}} < 1$. Let $\mathcal{R}_5 \in (0, 1)$ such that

$$\frac{C_A b_1}{p^-} \mathcal{R}_5^{p^- - \mathfrak{m}^+} < \lambda C(j) C_0.$$

Thus, for all $0 < \mathcal{R}_4 < \mathcal{R}_5$ and $u \in \mathbb{S}_{\mathcal{R}_4}^j$, we have that

$$\Psi_\lambda(u) \leq \mathcal{R}_4^{\mathfrak{m}^+} \left(\frac{C_A b_1}{p^-} \mathcal{R}_4^{p^- - \mathfrak{m}^+} - \lambda C(j) C_0 \right) \leq \mathcal{R}_5^{\mathfrak{m}^+} \left(\frac{C_A b_1}{p^-} \mathcal{R}_5^{p^- - \mathfrak{m}^+} - C(j) C_0 \right) < 0 = \Psi_\lambda(0).$$

Therefore, we conclude that

$$\sup_{u \in \mathbb{S}_{\mathcal{R}_4}^j} \Psi_\lambda(u) < 0 = \Psi_\lambda(0).$$

Thus the hypotheses of Theorem 2.16 are satisfied and we conclude that there exists a sequence nontrivial weak solutions u_j in \mathscr{W} such that

$$\Psi_\lambda(u_j) \leq 0, \quad \Psi'_\lambda(u_j) = 0 \text{ and } \|u_j\|_{\mathscr{W}} \rightarrow 0 \text{ as } j \rightarrow +\infty.$$

□

4. APPENDIX

This section presents several results which are used in our paper.

Proof of Lemma 2.4. We consider the set $\mathbb{M} = \{u \in \mathscr{W}; \|u\|_{L^{p(\cdot)}(\Omega)} = 1\}$. Then to prove this Lemma it suffices to prove that

$$\inf_{u \in \mathbb{M}} [u]_{\mathbb{R}^N}^{s,p(\cdot,\cdot)} = \zeta_1 > 0.$$

Initially, we observe that $\zeta_1 \geq 0$ and we prove ζ_1 is attained in \mathbb{M} . Let $(u_k)_{k \in \mathbb{N}} \subset \mathbb{M}$ be a minimizing sequence, that is, $[u_k]_{\mathbb{R}^N}^{s,p(\cdot,\cdot)} \rightarrow \zeta_1$ as $k \rightarrow +\infty$. This implies that $(u_k)_{k \in \mathbb{N}}$ is bounded in \mathscr{W} and $L^{p(\cdot)}(\Omega)$, therefore in $W^{s,p(\cdot,\cdot)}(\Omega)$. Consequently up to a subsequence $u_k \rightarrow u_0$ in $W^{s,p(\cdot,\cdot)}(\Omega)$ as $k \rightarrow +\infty$. Thus, from Corollary 2.3, it follows that $u_k \rightarrow u_0$ in $L^{p(\cdot)}(\Omega)$ as $k \rightarrow +\infty$. We extend u_0 to \mathbb{R}^N by setting $u_0(x) = 0$ on $x \in \mathbb{R}^N \setminus \Omega$. This implies $u_k(x) \rightarrow u_0(x)$ a.e. $x \in \mathbb{R}^N$ as $k \rightarrow +\infty$. Hence by using Fatous Lemma, we have

$$\int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u_0(x) - u_0(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy \leq \liminf_{k \rightarrow +\infty} \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|u_k(x) - u_k(y)|^{p(x,y)}}{|x - y|^{N+sp(x,y)}} dx dy$$

which implies that

$$[u_0]_{\mathbb{R}^N}^{s,p(\cdot,\cdot)} \leq \liminf_{k \rightarrow +\infty} [u_k]_{\mathbb{R}^N}^{s,p(\cdot,\cdot)} = \zeta_1,$$

and thus $u_0 \in \mathscr{W}$. Moreover, $\|u_0\|_{L^{p(\cdot)}(\Omega)} = 1$ and then $u_0 \in \mathbb{M}$. In particular, $u_0 \neq 0$ and $[u_0]_{\mathbb{R}^N}^{s,p(\cdot,\cdot)} = \zeta_1 > 0$. □

Proof of Lemma 2.5. First we observe that by Lemma 2.4, for all $u \in \mathscr{W}$, we get

$$\|u\|_{W^{s,p(\cdot,\cdot)}(\Omega)} \leq \|u\|_{L^{p(\cdot)}(\Omega)} + \|u\|_{\mathscr{W}} \leq \left(\frac{1}{\zeta_1} + 1 \right) \|u\|_{\mathscr{W}}, \quad (4.1)$$

that is, \mathscr{W} is continuously embedded in $W^{s,p(\cdot,\cdot)}(\Omega)$, and by Corollary 2.3 we conclude that \mathscr{W} is continuously embedded in $L^{r(\cdot)}(\Omega)$. To prove that the embedding above is compact we consider $(u_k)_{k \in \mathbb{N}}$ a bounded sequence in \mathscr{W} . Using (4.1), follows that $(u_k)_{k \in \mathbb{N}}$ is bounded in $W^{s,p(\cdot,\cdot)}(\Omega)$. Hence by Corollary 2.3, we infer that there exists $u_0 \in L^{r(\cdot)}(\Omega)$ such that up to a subsequence $u_k \rightarrow u_0$ in $L^{r(\cdot)}(\Omega)$ as $k \rightarrow +\infty$. Since that $u_k = 0$ a.e. in $\mathbb{R}^N \setminus \Omega$ for all $k \in \mathbb{N}$, so we define $u_0 = 0$ a.e. in $\mathbb{R}^N \setminus \Omega$ and obtain that the convergence occurs in $L^{r(\cdot)}(\Omega)$. This completes the proof this Lemma. □

Proof of Lemma 2.7. (i) Using standard arguments prove this item.

(ii) From (i) the functional Φ is of class $C^1(\mathscr{W}, \mathbb{R})$, and by hypothesis (a_1) , the functional Φ' is monotone. Thus, by [38, Lemma 15.4] we conclude that $\langle \Phi'(u), u_k - u \rangle + \Phi(u) \leq \Phi(u_k)$ for all $k \in \mathbb{N}$.

Thus, since $u_k \rightarrow u$ in \mathscr{W} , as $k \rightarrow +\infty$ we obtain $\Phi(u) \leq \liminf_{k \rightarrow +\infty} \Phi(u_k)$. That is, the functional Φ is weakly lower semicontinuous.

(iii) Let $(u_k)_{k \in \mathbb{N}}$ be a sequence in \mathscr{W} as in the statement. We have that by (i), Φ' is a continuous functional. Therefore,

$$\lim_{k \rightarrow +\infty} \langle \Phi'(u), u_k - u \rangle = 0. \quad (4.2)$$

Now, we observe that

$$\langle \Phi'(u_k) - \Phi'(u), u_k - u \rangle = \langle \Phi'(u_k), u_k - u \rangle - \langle \Phi'(u), u_k - u \rangle \text{ for all } k \in \mathbb{N}. \quad (4.3)$$

Thus by (2.2), (4.2), and (4.3), we infer

$$\limsup_{k \rightarrow +\infty} \langle \Phi'(u_k) - \Phi'(u), u_k - u \rangle \leq 0. \quad (4.4)$$

Furthermore, since Φ is strictly convex by hypothesis (a_1) , Φ' is monotone (see [38, Lemma 15.4]), we have

$$\langle \Phi'(u_k) - \Phi'(u), u_k - u \rangle \geq 0 \text{ for all } k \in \mathbb{N}. \quad (4.5)$$

Therefore, by (4.4) and (4.5), we infer that

$$\lim_{k \rightarrow +\infty} \langle \Phi'(u_k) - \Phi'(u), u_k - u \rangle = 0. \quad (4.6)$$

Consequently, by (4.2), (4.3), and (4.6), we conclude

$$\lim_{k \rightarrow +\infty} \langle \Phi'(u_k), u_k - u \rangle = 0. \quad (4.7)$$

Since Φ is strictly convex, we get

$$\Phi(u) + \langle \Phi'(u_k), u_k - u \rangle \geq \Phi(u_k) \text{ for all } k \in \mathbb{N}. \quad (4.8)$$

Thus, by (4.7) and (4.8), we have

$$\Phi(u) \geq \lim_{k \rightarrow +\infty} \Phi(u_k). \quad (4.9)$$

Since Φ is weakly lower semicontinuous (see (ii)) and by (4.9), we conclude that

$$\Phi(u) = \lim_{k \rightarrow +\infty} \Phi(u_k). \quad (4.10)$$

On the other hand, by (4.6) the sequence $(\mathcal{U}_k(x, y))_{k \in \mathbb{N}}$ converge to 0 in $L^1(\mathbb{R}^N \times \mathbb{R}^N)$ as $k \rightarrow +\infty$, where

$$\mathcal{U}_k(x, y) := [\mathcal{A}(u_k(x) - u_k(y)) - \mathcal{A}(u(x) - u(y))] [(u_k(x) - u_k(y)) - (u(x) - u(y))] K(x, y) \geq 0.$$

Hence there exists a subsequence $(u_{k_j})_{j \in \mathbb{N}}$ of $(u_k)_{k \in \mathbb{N}}$ such that

$$\mathcal{U}_{k_j}(x, y) \rightarrow 0 \text{ a.e. } (x, y) \in \mathbb{R}^N \times \mathbb{R}^N \text{ as } j \rightarrow +\infty. \quad (4.11)$$

We denoted $\mu_j(x, y) = u_{k_j}(x) - u_{k_j}(y)$ and $\mu(x, y) = u(x) - u(y)$ for all $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$.

Claim a. If $\mathcal{U}_{k_j}(x, y) \rightarrow 0$ a.e. $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$, then $\mu_j(x, y) \rightarrow \mu(x, y)$ as $j \rightarrow +\infty$ for almost all $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$.

Indeed, fixed $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$ with $x \neq y$ we suppose by contradiction that the sequence $(\mu_j(x, y))_{j \in \mathbb{N}}$ is unbounded for $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$ fixed. Using (4.11) we get $\mathcal{U}_{k_j}(x, y) \rightarrow 0$ as $j \rightarrow +\infty$ in \mathbb{R} , consequently there is $M > 0$ such that for all $j \in \mathbb{N}$

$$\left| [\mathcal{A}(\mu_j(x, y)) - \mathcal{A}(\mu(x, y))] (\mu_j(x, y) - \mu(x, y)) K(x, y) \right| \leq M. \quad (4.12)$$

Thus, denoting

$$V_A := [\mathcal{A}(\mu_j(x, y)) \mu_j(x, y) + \mathcal{A}(\mu(x, y)) \mu(x, y)] K(x, y) \text{ for all } j \in \mathbb{N},$$

we get from (4.12) that

$$V_A \leq M + \mathcal{A}(\mu(x, y)) \mu_j(x, y) K(x, y) + \mathcal{A}(\mu_j(x, y)) \mu(x, y) K(x, y) \text{ for all } j \in \mathbb{N}.$$

So using (a_2) and (\mathcal{K}) in inequality above, we have for all $j \in \mathbb{R}$ that,

$$\begin{aligned} c_{\mathcal{A}}b_0 \frac{|\mu_j(x, y)|^{p(x, y)}}{|x - y|^{N+sp(x, y)}} + c_{\mathcal{A}}b_0 \frac{|\mu(x, y)|^{p(x, y)}}{|x - y|^{N+sp(x, y)}} \leq M + C_{\mathcal{A}}b_1 \frac{|\mu(x, y)|^{p(x, y)-1} |\mu_j(x, y)|}{|x - y|^{N+sp(x, y)}} \\ + C_{\mathcal{A}}b_1 \frac{|\mu_j(x, y)|^{p(x, y)-1} |\mu(x, y)|}{|x - y|^{N+sp(x, y)}}. \end{aligned} \quad (4.13)$$

Dividing (4.13) by $|\mu_j(x, y)|^{p(x, y)}$, we achieve

$$\begin{aligned} \frac{c_{\mathcal{A}}b_0}{|x - y|^{N+sp(x, y)}} + \frac{c_{\mathcal{A}}b_0 |\mu(x, y)|^{p(x, y)}}{|\mu_j(x, y)|^{p(x, y)} |x - y|^{N+sp(x, y)}} \leq \frac{M}{|\mu_j(x, y)|^{p(x, y)}} \\ + \frac{C_{\mathcal{A}}b_1 |\mu(x, y)|^{p(x, y)-1}}{|\mu_j(x, y)|^{p(x, y)-1} |x - y|^{N+sp(x, y)}} \\ + \frac{C_{\mathcal{A}}b_1 |\mu(x, y)|}{|\mu_j(x, y)| |x - y|^{N+sp(x, y)}} \end{aligned} \quad (4.14)$$

for all $j \in \mathbb{R}$. Since we are supposing that the sequence $(\mu_j(x, y))_{j \in \mathbb{N}}$ is unbounded, we can assume that $|\mu_j(x, y)| \rightarrow +\infty$ as $j \rightarrow +\infty$, then by (4.14) we obtain $c_{\mathcal{A}}b_0 \leq 0$ which is an contradiction.

Therefore, the sequence $(\mu_j(x, y))_{j \in \mathbb{N}}$ is bounded in \mathbb{R} and up to a subsequence $(\mu_j(x, y))_{j \in \mathbb{N}}$ converges to some $\nu \in \mathbb{R}$. Thus we obtain $\mu_j(x, y) \rightarrow \nu$ as $j \rightarrow +\infty$. Thence denoting

$$\mathcal{U}(x, y) := [\mathcal{A}(\nu) - \mathcal{A}(\mu(x, y))](\nu - (\mu(x, y)))K(x, y)$$

and using (a_1) we conclude that

$$\mathcal{U}_{k_j}(x, y) \rightarrow \mathcal{U}(x, y) \text{ as } j \rightarrow +\infty. \quad (4.15)$$

Consequently, by (4.11) and (4.15), we get

$$\mathcal{U}(x, y) = [\mathcal{A}(\nu) - \mathcal{A}(\mu(x, y))](\nu - (\mu(x, y)))K(x, y) = 0. \quad (4.16)$$

In this way, by strictly convexity of \mathcal{A} , (4.16) this occurs only if $\nu = \mu(x, y) = u(x) - u(y)$. Therefore, by uniqueness of limit

$$u_{k_j}(x) - u_{k_j}(y) = \mu_j(x, y) \rightarrow \mu(x, y) = u(x) - u(y) \text{ in } \mathbb{R} \text{ as } j \rightarrow +\infty \quad (4.17)$$

for almost all $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$.

Now we consider the sequence $(g_{k_j})_{j \in \mathbb{N}}$ in $L^1(\mathbb{R}^N \times \mathbb{R}^N)$ defined pointwise for all $j \in \mathbb{N}$ by

$$g_{k_j}(x, y) := \left[\frac{1}{2} \left(\mathcal{A}(\mu_j(x, y)) + \mathcal{A}(\mu(x, y)) \right) - \mathcal{A} \left(\frac{\mu_j(x, y) - \mu(x, y)}{2} \right) \right] K(x, y).$$

By convexity the map \mathcal{A} (see (a_1)), $g_{k_j}(x, y) \geq 0$ for almost all $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$. Furthermore, by continuity of map \mathcal{A} (see (a_1)) and (4.17), we have

$$g_{k_j}(x, y) \rightarrow \mathcal{A}(\mu(x, y))K(x, y) \text{ as } j \rightarrow +\infty \text{ for all } (x, y) \in \mathbb{R}^N \times \mathbb{R}^N.$$

Therefore, using this above information, (4.10), and Fatou's Lemma, we get

$$\Phi(u) \leq \liminf_{j \rightarrow +\infty} \int_{\mathbb{R}^N \times \mathbb{R}^N} g_{k_j}(x, y) dx dy = \Phi(u) - \limsup_{j \rightarrow +\infty} \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A} \left(\frac{\mu_j(x, y) - \mu(x, y)}{2} \right) K(x, y) dx dy.$$

Then

$$\limsup_{j \rightarrow +\infty} \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A} \left(\frac{\mu_j(x, y) - \mu(x, y)}{2} \right) K(x, y) dx dy \leq 0. \quad (4.18)$$

On the other hand, by (a_2) , (a_3) , (\mathcal{K}) , and Proposition 2.6, we infer that

$$\begin{aligned} \int_{\mathbb{R}^N \times \mathbb{R}^N} \mathcal{A} \left(\frac{\mu_j(x, y) - \mu(x, y)}{2} \right) K(x, y) dx dy \geq \frac{c_{\mathcal{A}}b_0}{2^{p^-} p^+} \inf \{ \|u_{k_j} - u\|_{\mathcal{W}^-}^{p^-}, \|u_{k_j} - u\|_{\mathcal{W}^+}^{p^+} \} \\ \geq 0 \text{ for all } j \in \mathbb{N}. \end{aligned} \quad (4.19)$$

Consequently, by (4.18) and (4.19), we achieve

$$\lim_{j \rightarrow +\infty} \|u_{k_j} - u\|_{\mathcal{W}} = 0.$$

Therefore, we can conclude that $u_{k_j} \rightarrow u$ in \mathcal{W} as $j \rightarrow +\infty$. Since $(u_{k_j})_{j \in \mathbb{N}}$ is an arbitrary subsequence of $(u_k)_{k \in \mathbb{N}}$, this shows that $u_k \rightarrow u$ as $k \rightarrow +\infty$ in \mathcal{W} , as required. \square

ACKNOWLEDGEMENTS

The first author was financed by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior(CAPES)-Finance Code 001. The second author was supported the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior(CAPES)-Finance Code 001. The third author named was partially supported by INCTMAT/CNPq/Brazil and CNPq/Brazil 307061/2018-3.

REFERENCES

- [1] C. O. Alves, M. C. Ferreira, *Nonlinear perturbations of a $p(x)$ -Laplacian equation with critical growth in \mathbb{R}^N* , Math. Nachr. 287 (2014), 849868.
- [2] A. Ambrosetti, P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis. 14 (1973), 349-381.
- [3] F. Andreu-Vailló, J. M. Mazn, J. D. Rossi, J. J. Toledo-Melero, *Nonlocal diffusion problems*, volume 165 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI; Real Sociedad Matemática Española, Madrid (2010).
- [4] E. Azroul, A. Benkirane, M. Shimi, *Eigenvalue problems involving the fractional $p(x)$ -Laplacian fractional operator*, Adv. Oper. Theory 4 (2019), 539-555.
- [5] G. Autuori, P. Pucci, *Existence of entire solutions for a class of quasilinear elliptic equations*, Nonlinear Differential Equations and Appl. 20 (2013), 9771009.
- [6] A. Bahrouni, *TrudingerMoser type inequality and existence of solution for perturbed nonlocal elliptic operators with exponential nonlinearity*, Comm. Pure Appl. Anal. 16 (2017), 243252.
- [7] A. Bahrouni, *Comparasion and sub-supersolution principles for the fractional $p(x)$ -Laplacian*, J. Math. Anal. Appl. 458 (2018), 1363-1372.
- [8] A. Bahrouni, V. D. Rădulescu, *On a new fractional Sobolev space and applications to nonlocal variational problems with variable exponent*, Discrete Contin. Dyn. Syst. 11 (2018), 379-389.
- [9] T. Bartsch, M. Willem, *On an elliptic equation with concave and convex nonlinearities*, Proc. Amer. Math. Soc. (11) 123 (1995), 35553561.
- [10] G. Bin, *On superlinear $p(x)$ -Laplacian like problem without Ambrsetti and Rabinowitz condition*, Bull. Korean Math. Soc. 51 (2014), 409-421.
- [11] G. M. Bisci, V. D. Rădulescu, *Ground state solutions of scalar field fractional Schrödinger equations*, Calc. Var. 54 (2015), 29853008.
- [12] G. M. Bisci, D. D. Repovš, *Multiple solutions for elliptic equations involving a general operator in divergence form*, Ann. Acad. Sci. Fenn. Math. 39 (2014), 259-273.
- [13] H. Brezis, H-M. Nguyen, *Non-local functionals related to the total variation and applications*, Image Processing. Ann. PDE 4 (2018), 77 pp.
- [14] C. Bucur, E. Valdinoci, *Nonlocal diffusion and applications*, Lect. Notes Unione Mat. Ital., vol.20, Springer/Unione Matematica Italiana, Cham/Bologna (2016).
- [15] L. Caffarelli, *Nonlocal equations, drifts and games*, Nonlinear Partial Differential Equations 7 (2012), 3752.
- [16] G. Cerami, *An existence criterion for the critical points on unbounded manifolds*, Istit. Lombardo Accad. Sci. Lett. Rend. A 112 (1978), 332336.
- [17] G. Cerami, *On the existence of eigenvalues for a nonlinear boundary value problem*, Ann. Mat. Pura Appl. (4) 124 (1980), 161179.
- [18] F. Colasuonno, P. Pucci, C. Varga, *Multiple solutions for an eigenvalue problem involving p Laplacian type operators*, Nonlinear Anal 75 (2012), 44964512.
- [19] D. G. Costa, C. A. Magalhães, *Existence results for perturbations of the p -Laplacian*, Nonlinear Anal. 24 (1995), 409-418.
- [20] L. Diening, P. Harjulehto, P. Hasto, M. Ruzicka, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Mathematics, Vol. 2017, Springer-Verlag, Heidelberg (2011).
- [21] D. E. Edmunds, J. Rákosník, *Sobolev embeddings with variable exponent*, Studia Math. 143 (2000), 267-293.
- [22] E. Emmrich, D. Puhst, *Measure-valued and weak solutions to the nonlinear peridynamic model in nonlocal elastodynamics*, Nonlinearity 28 (2014), 285307.
- [23] M. Fabian, P. Habala, P. Hájek, V. Montesinos, V. Zizler, *Banach Space Theory: The Basis for Linear and Nonlinear Analysis*, New York Springer (2011).
- [24] X. Fan, X. Han, *Existence and multiplicity of solutions for $p(x)$ -Laplacian equations in \mathbb{R}^N* , Nonlinear Anal. 59 (2004), 173-188.
- [25] X. Fan, Q. Zhang, *Existence of solutions for $p(x)$ -Laplacian Dirichlet problem*, Nonlinear Anal. 52 (2003), 1843-1852.
- [26] S. Fučik, O. John, J. Nečas, *On the existence of Schauder bases in Sobolev spaces*, Commentationes Mathematicae Universitatis Carolinae, 1, Vol. 13 (1972), 163-175.
- [27] C. Gal, M. Warma, *Bounded solutions for nonlocal boundary value problems on Lipschitz manifolds with boundary*, Adv. Nonlinear Stud. 16 (2016), 529550.

- [28] T. C. Halsey, *Electrorheological fluids*, Science 258 (1992), 761766.
- [29] K. Ho, Y. H. Kim, *A-priori bounds and multiplicity of solutions for nonlinear elliptic problems involving the fractional $p(\cdot)$ -Laplacian*, Nonlinear Analysis 188 (2019), 179201.
- [30] E. J. Hurtado, *Nonlocal diffusion equations involving the fractional $p(\cdot)$ -Laplacian*, Journal of Dynamics and Differential Equations (2019). doi.org/10.1007/s10884-019-09745-2.
- [31] E. J. Hurtado, O. H. Miyagaki, R. S. Rodrigues, *Existence and multiplicity of solutions for a class of elliptic equations without Ambrosetti-Rabinowitz type conditions*, Journal of Dynamics and Differential Equations 30 (2018), 405-432.
- [32] E. J. Hurtado, O. H. Miyagaki, R. S. Rodrigues, *Existence and asymptotic behaviour for a Kirchhoff type equation with variable critical growth exponent*, Milan J. Math. 85 (2017), 71102.
- [33] A. Iannizzotto, M. Squassina, *$\frac{1}{2}$ -Laplacian problems with exponential nonlinearity*, J. Math. Anal. Appl. 414 (2014), 372385.
- [34] A. Iannizzotto, M. Squassina, *Weyl-type laws for fractional p -eigenvalue problems*, Asymptot. Anal. 4 (2014), 233245.
- [35] C. Ji, *On the superlinear problem involving $p(x)$ -laplacian*, Electron. J. Qual. Theory Differ. Equ. 40 (2011), 1-9.
- [36] R. Kajikiya, *A critical point theorem related to the symmetric mountain-pass lemma and its applications to elliptic equations*, J. Funct. Anal. 225 (2005), 352370.
- [37] U. Kaufmann, J. D. Rossi, R. Vidal, *Fractional Sobolev spaces with variable exponents and fractional $p(x)$ -Laplacian*, Electron. J. Qual. Theory Differ. Equ. 76 (2017), 1-10.
- [38] O. Kavian, *Introduction à la Théorie des Points Critiques et Applications aux Problèmes Elliptiques*, Paris; New York Springer-Verlag (1993).
- [39] I. H. Kim, Y. H. Kim, *Mountain pass type solutions and positivity of the infimum eigenvalue for quasilinear elliptic equations with variable exponents*, Manuscripta Math. 147 (2015), 169191.
- [40] T. Kussi, G. Mingione, Y. Sire, *Nonlocal Equations with Measure Data*, Commun. Math. Phys. 337 (2015), 1317-1368.
- [41] J. Lee, Y. H. Kim, *Multiplicity results for nonlinear Neumann boundary value problems involving p -Laplace type operators*, Bound. Value Probl. 95 (2016), 125.
- [42] J. I. Lee, J. M. Kim, Y.H. Kim, J. Lee, *Multiplicity of weak solutions to non-local elliptic equations involving the fractional $p(x)$ -Laplacian*, Journal of Mathematical Physics, 61(1) (2020), 011505.
- [43] G. Li, C. Yang, *The existence of a nontrivial solution to a nonlinear elliptic boundary value problem of p -Laplacian type without the Ambrosetti-Rabinowitz condition*, Nonlinear Anal. 72 (2010), 4602-4613.
- [44] Z. Liu, Z. Q. Wang, *On the Ambrosetti-Rabinowitz superlinear condition*, Adv. Nonlinear Stud. 4 (2004), 563-574.
- [45] S. B. Liu, S. J. Li, *On superlinear problems without the Ambrosetti and Rabinowitz condition*, Nonlinear Anal. 73 (2010), 788795.
- [46] O. H. Miyagaki, M. A. S Souto, *Superlinear problems without Ambrosetti and Rabinowitz growth condition*, J. Differential Equations. 245 (2008), 3628-3638.
- [47] G. Palatucci, T. Kuusi, *Recent developments in the Nonlocal Theory, Book Series on Measure Theory*, De Gruyter Berlin (2016).
- [48] R. Pei, *Fractional p -laplacian equations with subcritical and critical exponential growth without the Ambrosetti-Rabinowitz condition*, Mediterranean Journal of Mathematics (2) 15 (2018).
- [49] P. Pucci, X. Mingqi, Z. Binlin, *Existence and multiplicity of entire solutions for fractional p -Kirchhoff equations*, Adv. Nonlinear Anal. 5 (2016), 2755.
- [50] P. Pucci, S. Saldi, *Multiple solutions for an eigenvalue problem involving nonlocal elliptic p -Laplacian operators*, Geometric Methods in PDEs - Springer INdAM Series - Vol. 13, G. Citti, M. Manfredini, D. Morbidelli, S. Polidoro, F. Uguzzoni Eds. (2015), pages 16.
- [51] P. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Regional Conf. Ser. inMath. 65. American Mathematical Society, Providence, RI, (1986).
- [52] V. D. Rădulescu, D. Repovš, *Partial differential equations with variable exponents: Variational Methods and Qualitative Analysis*, CRC Press, Taylor & Francis Group, Boca Raton, FL, (2015).
- [53] M. M. Rodrigues, *Multiplicity of solutions on a nonlinear eigenvalue problem for $p(x)$ -laplacian-like operators*, Mediterr. J. Math. 9 (2012), 211223.
- [54] R. Servadei, E. Valdinoci, *Mountain pass solutions for nonlocal elliptic operators*, J. Math. Anal. Appl. 389 (2012), 887898.
- [55] J. L. Vázquez, *The mathematical theories of diffusion: nonlinear and fractional diffusion*, In: Bonforte, M., Grillo, G. (eds.) Nonlocal and Nonlinear Diffusions and Interactions: New Methods and Directions. Lecture Notes in Mathematics 2186 (2017), 205278.
- [56] B. L. Zhang, G. Molica Bisci, R. Servadei, *Superlinear nonlocal fractional problems with infinitely many solutions*, Nonlinearity 28 No. 7 (2015), 2247-2264.
- [57] J. F. Zhao, *Structure Theory of Banach Spaces*, Wuhan University Press, Wuhan, (1991) (in Chinese).
- [58] Q. M. Zhou, *On a class of superlinear nonlocal fractional problems without Ambrosetti-Rabinowitz type conditions*, Electronic Journal of Qualitative Theory of Differential Equations 17 (2019), 1-12.
- [59] Q. M. Zhou, *On the superlinear problems involving $p(x)$ -Laplacian-like operators without AR-condition*, Nonlinear Analysis: Real World Applications. 21 (2015), 161-169.