

FUNCTORIALLY FINITE HEARTS, SIMPLE-MINDED SYSTEMS IN NEGATIVE CLUSTER CATEGORIES, AND NONCROSSING PARTITIONS

RAQUEL COELHO SIMÕES, DAVID PAUKSZTELLO, AND DAVID PLOOG

ABSTRACT. Let Q be an acyclic quiver and $w \geq 1$ be an integer. Let $\mathcal{C}_{-w}(\mathbf{k}Q)$ be the $(-w)$ -cluster category of $\mathbf{k}Q$. We show that there is a bijection between simple-minded collections in $\mathcal{D}^b(\mathbf{k}Q)$ lying in a fundamental domain of $\mathcal{C}_{-w}(\mathbf{k}Q)$ and w -simple-minded systems in $\mathcal{C}_{-w}(\mathbf{k}Q)$. This generalises the same result of Iyama-Jin in the case that Q is Dynkin. A key step in our proof is the observation that the heart \mathcal{H} of a bounded t-structure in a Hom-finite, Krull-Schmidt, \mathbf{k} -linear triangulated category \mathcal{D} is functorially finite in \mathcal{D} if and only if \mathcal{H} has enough injectives and enough projectives. We then establish a bijection between w -simple-minded systems in $\mathcal{C}_{-w}(\mathbf{k}Q)$ and positive w -noncrossing partitions of the corresponding Weyl group W_Q .

CONTENTS

Introduction	1
1. Preliminaries	3
2. Functorially finite hearts	6
3. Orthogonal collections	10
4. Simple-minded collections vs w -simple minded systems	11
5. Sincere orthogonal collections	17
6. Noncrossing partitions	22
Appendix A. Reduction of simple-minded collections revisited	25
References	27

INTRODUCTION

Cluster-tilting theory was introduced in [14] as an approach to categorifying the cluster algebras of Fomin and Zelevinsky in [23]. Since its inception, cluster-tilting theory has gone on to have widespread connections with many areas of mathematics.

Classical cluster-tilting theory takes place in an m -cluster category $\mathcal{C}_m(\mathbf{k}Q)$, for $m \geq 2$, which is an m -Calabi-Yau orbit category of the bounded derived category, $\mathcal{D}^b(\mathbf{k}Q)$, of the path algebra of a finite acyclic quiver Q , where \mathbf{k} is an algebraically closed field. One of the most fruitful connections has been between the representation theory of finite-dimensional algebras and Coxeter combinatorics. For example, let W be the Weyl group of type

2010 *Mathematics Subject Classification.* 18E30,16G10,05E10.

Key words and phrases. Simple-minded system, simple-minded collection, Calabi-Yau triangulated category, noncrossing partition, Riedtmann configuration.

Q . There are natural bijections between the sets of clusters and noncrossing partitions associated to W ; see [4, 41, 47]. There is in turn a bijection between the sets of clusters and cluster-tilting objects in the corresponding cluster category [15]. Beyond cluster-tilting theory, there are many further connections between the combinatorics of noncrossing partitions and representation theory, e.g. in the classification of wide subcategories and torsion theories [29], thick subcategories [38] and Cartan lattices [27], to name a few. For a broad treatment of the combinatorics of noncrossing partitions in finite type we refer the reader to [47].

Recently, there has been increasing interest in negative Calabi-Yau triangulated categories; see, for example, [12, 13, 16, 17, 18, 19, 20, 22, 26, 30, 32, 33, 36]. For an integer $w \geq 1$ and an acyclic quiver Q there is an orbit category $\mathcal{C}_{-w}(\mathbf{k}Q)$ of the bounded derived category $\mathbf{D}^b(\mathbf{k}Q)$ which is $(-w)$ -Calabi-Yau. This orbit category can be thought of as a ‘negative (Calabi-Yau) cluster category’. In this setting the analogue of m -cluster-tilting objects are so called w -simple-minded systems. Evidence supporting the viewpoint that w -simple-minded systems are a negative Calabi-Yau analogue of cluster-tilting objects is advanced by a growing body of work: [15, 16, 17, 18, 20, 22, 30, 32, 33].

It is therefore natural to ask about connections between negative cluster categories and Coxeter combinatorics. Previous work in this direction includes [15, 16, 30, 47]. Again let Q be an acyclic quiver with corresponding Weyl group W . In [16], the first author established a bijection between so-called positive noncrossing partitions of W and (1) -simple-minded systems in $\mathcal{C}_{-1}(\mathbf{k}Q)$ when Q is Dynkin. In [15], Buan, Reiten and Thomas obtained a bijection between simple-minded collections lying inside some ‘fundamental domain’ and w -noncrossing partitions, which was a forefather of the König-Yang correspondences [37]. Most recently, in [30], Iyama and Jin generalised [15] and [16] to obtain a bijection between w -simple-minded systems in $\mathcal{C}_{-w}(\mathbf{k}Q)$ and positive w -noncrossing partitions, again for Q Dynkin. This bijection proceeds via a bijection between w -simple-minded systems in $\mathcal{C}_{-w}(\mathbf{k}Q)$ and simple-minded collections lying in the fundamental domain \mathcal{F}_{-w} (see Section 1.2 for the precise definition) of $\mathbf{D}^b(\mathbf{k}Q)$; see [30, Theorem 1.2].

In this article we extend this bijection to the case that Q is an arbitrary quiver in the following main theorem.

Theorem A (Theorem 4.1). *Let Q be an acyclic quiver. The natural projection functor $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathcal{C}_{-w}(\mathbf{k}Q)$ induces a bijection*

$$\left\{ \begin{array}{l} \text{simple-minded collections of } \mathbf{D}^b(\mathbf{k}Q) \\ \text{contained in } \mathcal{F}_{-w} \end{array} \right\} \xleftrightarrow{1-1} \left\{ \begin{array}{l} w\text{-simple-minded systems} \\ \text{in } \mathcal{C}_{-w} \end{array} \right\}.$$

Using Theorem A, we are then able to obtain the following bijection involving noncrossing partitions for any acyclic quiver Q .

Theorem B (Theorem 6.4). *Let Q be an acyclic quiver. There is a bijection*

$$\left\{ \begin{array}{l} \text{positive } w\text{-noncrossing} \\ \text{partitions of } W_Q \end{array} \right\} \xleftrightarrow{1-1} \left\{ \begin{array}{l} w\text{-simple-minded systems} \\ \text{in } \mathcal{C}_{-w}(\mathbf{k}Q) \end{array} \right\}.$$

In order to obtain Theorem A, we require another observation that we believe holds independent interest and will be widely applicable. Let \mathbf{D} be a Hom-finite, Krull-Schmidt, \mathbf{k} -linear triangulated category and suppose that $\mathbf{H} \subseteq \mathbf{D}$ is the heart of a bounded t-structure. One can ask what is the relationship between the following properties of \mathbf{H} :

- \mathbf{H} is a length category with finitely many simple objects;
- \mathbf{H} has enough injective objects and enough projective objects; and,
- \mathbf{H} is a functorially finite subcategory of \mathbf{D} .

Of these three properties, the final property is the odd one out: it is the only property which takes the ambient triangulated category in which \mathbf{H} sits into account. As such a relationship between these properties is potentially very powerful. Unfortunately, Example 2.4 shows that there is no relationship, in general, between the first and the third properties. However, our third main theorem provides a relationship between the second and third properties of \mathbf{H} . It is this relationship which is a crucial ingredient in our proof of Theorem A.

Theorem C (Theorem 2.1). *Suppose $\mathbf{H} \subseteq \mathbf{D}$ is the heart of a bounded t -structure. Then \mathbf{H} is functorially finite in \mathbf{D} if and only if \mathbf{H} has enough injectives and enough projectives.*

We briefly sketch the structure of the paper. In Section 1, we recall the basic concepts we use in the paper. In Section 2 we prove Theorem C. Section 3 recalls the basic properties of orthogonal collections, simple-minded systems and simple-minded collections, and establishes a characterisation of simple-minded collections in terms of Riedtmann configurations that may be more widely applicable and illustrates the parallel between simple-minded collections and simple-minded systems. In Section 4, we prove our main theorem, Theorem A. Section 5 establishes a bijection between w -simple-minded systems and certain sincere orthogonal collections which is the crucial tool to pass from Theorem A to Theorem B, which is done in Section 6. Finally, in an appendix, we give an alternative proof of a theorem by Jin [33] on the reduction of simple-minded collections using the characterisation in terms of Riedtmann configurations, which avoids the passage to a Verdier localisation, and is more in the spirit of [20].

Notation convention. In abstract abelian and triangulated categories we will use lower case Roman letters to denote objects. When we specialise to module categories or derived categories of an algebra, we will use upper case to denote objects. The philosophy behind this is that in the latter case, these objects have elements.

Acknowledgments. The authors would like to thank Lidia Angeleri Hügel, Jorge Vitória and Alexandra Zvonareva for useful discussions. The authors are grateful to Peter Jørgensen for comments on a draft, and for pointing out the application of Theorem C given in Corollary 2.2. The first author is grateful to the European Union’s Horizon 2020 research and innovation programme for financial support through the Marie Skłodowska-Curie Individual Fellowship grant agreement number 838706.

1. PRELIMINARIES

To begin with \mathbf{D} will be a Hom-finite, Krull-Schmidt, \mathbf{k} -linear triangulated category over an algebraically closed field \mathbf{k} . The shift or suspension functor will be denoted by $\Sigma: \mathbf{D} \rightarrow \mathbf{D}$. Later we will specialise to the case that $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$ for a finite acyclic quiver Q . Abusing notation, if \mathbf{X} is a collection of objects of \mathbf{D} and d is an object of \mathbf{D} we will write $\mathrm{Hom}_{\mathbf{D}}(\mathbf{X}, d)$ to mean $\mathrm{Hom}_{\mathbf{D}}(x, d)$ where we take each $x \in \mathbf{X}$ in turn; likewise for $\mathrm{Hom}_{\mathbf{D}}(d, \mathbf{X})$. We shall assume all subcategories are full and strict.

For subcategories \mathbf{X} and \mathbf{Y} of \mathbf{D} , we write

$$\mathbf{X} * \mathbf{Y} = \{d \in \mathbf{D} \mid \text{there exists a triangle } x \rightarrow d \rightarrow y \rightarrow \Sigma x \text{ with } x \in \mathbf{X} \text{ and } y \in \mathbf{Y}\}.$$

We note that by the octahedral axiom, the $*$ product of subcategories is associative. A subcategory \mathbf{X} is *extension-closed* if $\mathbf{X} * \mathbf{X} = \mathbf{X}$. We denote by $\langle \mathbf{X} \rangle$, or sometimes by $\langle \mathbf{X} \rangle_{\mathbf{D}}$ when we need to emphasise the triangulated category in which we are working, the *extension closure* of \mathbf{X} , that is the smallest extension-closed subcategory of \mathbf{D} containing \mathbf{X} . The *right and left perpendicular categories* of \mathbf{X} are defined as follows:

$$\mathbf{X}^{\perp} = \{d \in \mathbf{D} \mid \mathrm{Hom}(\mathbf{X}, d) = 0\} \text{ and } {}^{\perp}\mathbf{X} = \{d \in \mathbf{D} \mid \mathrm{Hom}(d, \mathbf{X}) = 0\}.$$

In Section 5 we will also require a notion of perpendicular category that is more suited to abelian categories; see Definition 5.6.

An autoequivalence $\mathbb{S}: \mathbf{D} \rightarrow \mathbf{D}$ is called a Serre functor if for each $x, y \in \mathbf{D}$ there is an isomorphism,

$$\mathrm{Hom}(x, y) \simeq D \mathrm{Hom}(y, \mathbb{S}x),$$

which is natural in x and y , where $D = \mathrm{Hom}_{\mathbf{k}}(-, \mathbf{k})$. If \mathbf{D} has a Serre functor, it is unique up to isomorphism and we say \mathbf{D} satisfies *Serre duality*. For details we refer to [42].

Let $w \in \mathbb{Z}$. A triangulated category \mathbf{D} satisfying Serre duality is *w-Calabi-Yau* (or *w-CY*) if there is a natural isomorphism $\mathbb{S} \simeq \Sigma^w$, where \mathbb{S} is the Serre functor on \mathbf{D} .

1.1. Functorially finite subcategories and t-structures. Let \mathbf{X} be a subcategory of \mathbf{D} , and d an object in \mathbf{D} . A morphism $f: x \rightarrow d$, with $x \in \mathbf{X}$, is a *right X-approximation of d* if $\mathrm{Hom}(\mathbf{X}, f): \mathrm{Hom}(\mathbf{X}, x) \rightarrow \mathrm{Hom}(\mathbf{X}, d)$ is surjective. The morphism f is called *right minimal* if any $g: x \rightarrow x$ satisfying $fg = f$ is an automorphism. The morphism f is called a *minimal right X-approximation of d* if it is a right X-approximation of d and is right minimal. If $f: x \rightarrow d$ is a minimal right X-approximation and $h: x' \rightarrow d$ is a right X-approximation then x is isomorphic to a direct summand of x' [5].

If every object in \mathbf{D} admits a right X-approximation, then \mathbf{X} is said to be *contravariantly finite*. There are dual notions of (*minimal*) *left X-approximations* and *covariantly finite* subcategories. The subcategory \mathbf{X} of \mathbf{D} is called *functorially finite* if it is both contravariantly finite and covariantly finite.

A *t-structure* in \mathbf{D} consists of a pair of full subcategories (\mathbf{X}, \mathbf{Y}) such that $\mathrm{Hom}_{\mathbf{D}}(\mathbf{X}, \mathbf{Y}) = 0$, $\mathbf{D} = \mathbf{X} * \mathbf{Y}$ and $\Sigma \mathbf{X} \subseteq \mathbf{X}$ (equivalently, $\Sigma^{-1} \mathbf{Y} \subseteq \mathbf{Y}$). Its heart $\mathbf{H} = \mathbf{X} \cap \Sigma \mathbf{Y}$ is an abelian category [8]. A t-structure (\mathbf{X}, \mathbf{Y}) is called *bounded* if $\mathbf{D} = \bigcup_{i \in \mathbb{Z}} \Sigma^i \mathbf{X} = \bigcup_{i \in \mathbb{Z}} \Sigma^i \mathbf{Y}$. We recall the following standard characterisation of bounded t-structures; see, for example, [11, Lemma 3.2].

Lemma 1.1. *Let (\mathbf{X}, \mathbf{Y}) be a t-structure in \mathbf{D} with heart \mathbf{H} . The following conditions are equivalent:*

- (1) *the t-structure (\mathbf{X}, \mathbf{Y}) is bounded;*
- (2) *$\mathbf{X} = \bigcup_{n \geq 0} \Sigma^n \mathbf{H} * \Sigma^{n-1} \mathbf{H} * \dots * \mathbf{H}$ and $\mathbf{Y} = \bigcup_{n < 0} \Sigma^{-1} \mathbf{H} * \Sigma^{-2} \mathbf{H} * \dots * \Sigma^n \mathbf{H}$;*
- (3) *$\mathbf{D} = \bigcup_{n \geq m} \Sigma^n \mathbf{H} * \Sigma^{n-1} \mathbf{H} * \dots * \Sigma^{m+1} \mathbf{H} * \Sigma^m \mathbf{H}$.*

Note that for a t-structure (\mathbf{X}, \mathbf{Y}) , the inclusion $\mathbf{X} \rightarrow \mathbf{D}$ has a right adjoint and the inclusion $\mathbf{Y} \rightarrow \mathbf{D}$ has a left adjoint; these are given by the truncation functors. Therefore, the subcategory \mathbf{X} is always contravariantly finite and the subcategory \mathbf{Y} is always covariantly finite, and the approximations are even functorial.

Definition 1.2. A bounded t-structure (\mathbf{X}, \mathbf{Y}) in \mathbf{D} with heart \mathbf{H} is called *algebraic* if \mathbf{H} is a length category (every object has finite length, i.e. is Artinian as well as Noetherian) with only finitely many simple objects.

For example, the property of being an algebraic t-structure holds for module categories over finite-dimensional algebras.

1.2. Hereditary algebras and negative cluster categories. For this section, $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$ for some finite acyclic quiver Q . The main reference for the structure of derived categories of hereditary algebras (equivalently, path algebras of acyclic quivers) is [25].

Recall that an algebra A is *hereditary* if it is of global dimension 0 or 1, i.e. if the bifunctors $\mathrm{Ext}_A^n(-, -)$ are zero for $n \geq 2$. Typical examples are the path algebras $A = \mathbf{k}Q$. A well-known lemma says that each object of $\mathbf{D}^b(\mathbf{k}Q)$ decomposes as a direct sum of its cohomology. In particular, its Auslander–Reiten (AR) quiver has the following form:

\dots	$\Sigma^{-1}\mathrm{mod}(\mathbf{k}Q)$	$\mathrm{mod}(\mathbf{k}Q)$	$\Sigma\mathrm{mod}(\mathbf{k}Q)$	\dots
---------	--	-----------------------------	-----------------------------------	---------

Morphisms go from left to right, and since for two modules $M, N \in \mathrm{mod}(\mathbf{k}Q)$, one has $\mathrm{Hom}_{\mathbf{D}^b(\mathbf{k}Q)}(M, \Sigma^n N) = \mathrm{Ext}_{\mathbf{k}Q}^n(M, N)$, nonzero morphisms exist only from one degree to the next and not any higher. Moreover, there are no morphisms from one degree to a degree that is strictly lower.

In this article we will be interested in a certain orbit category of $\mathbf{D}^b(\mathbf{k}Q)$ which is constructed as follows; see [14]. Let $w \geq 1$ and $F = \Sigma^w \mathbb{S}$. We define the category

$$\mathbf{C}_{-w} = \mathbf{C}_{-w}(\mathbf{k}Q) := \mathbf{D}^b(\mathbf{k}Q)/F,$$

with the same objects as \mathbf{D} and whose morphisms are given by

$$\mathrm{Hom}_{\mathbf{C}_{-w}}(x, y) = \bigoplus_{n \in \mathbb{Z}} \mathrm{Hom}_{\mathbf{D}^b(\mathbf{k}Q)}(x, F^n y).$$

We write $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ for the natural projection functor. The orbit category \mathbf{C}_{-w} is a triangulated category (see [35]) which is $(-w)$ -Calabi-Yau; this uses that the algebra $\mathbf{k}Q$ is hereditary. Compare \mathbf{C}_{-w} with the construction of the *m-cluster category*, $\mathbf{D}^b(\mathbf{k}Q)/\Sigma^{-m}\mathbb{S}$, for $m \geq 2$, which is an m -Calabi-Yau triangulated category. As such we will refer to \mathbf{C}_{-w} as a *negative cluster category* or a $(-w)$ -*cluster category*.


It is often convenient to compute inside \mathbf{C}_{-w} using a so-called fundamental domain in $\mathbf{D}^b(\mathbf{k}Q)$. Let (\mathbf{X}, \mathbf{Y}) be the standard t-structure in $\mathbf{D}^b(\mathbf{k}Q)$, i.e. the t-structure with heart $\mathbf{H} = \mathrm{mod}(\mathbf{k}Q)$. We define the *fundamental domain* of \mathbf{C}_{-w} to be

$$\mathcal{F}_{-w} := \mathbf{X} \cap \Sigma^w \mathbf{Y}.$$

The natural projection functor $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ induces a bijection

$$\{\text{indecomposable objects in } \mathcal{F}_{-w}\} \xrightarrow{1-1} \{\text{indecomposable objects of } \mathbf{C}_{-w}\}.$$

Below we give a schematic of the fundamental domain \mathcal{F}_{-w} inside $\mathbf{D}^b(\mathbf{k}Q)$,



where the shaded grey region on the right-hand side comprises $\Sigma^w\text{inj}(\mathbf{k}Q)$, where $\text{inj}(\mathbf{k}Q)$ denotes the full subcategory of injective $\mathbf{k}Q$ -modules.

Finally, working inside \mathcal{F}_{-w} allows us to compute Hom spaces easily.

Lemma 1.3 ([30, Lemma 3.4]). *Suppose $x, y \in \mathcal{F}_{-w}$ and $0 \leq i \leq w$. Then*

$$\text{Hom}_{\mathcal{C}_{-w}}(x, \Sigma^{-i}y) = \text{Hom}_{\text{D}^b(\mathbf{k}Q)}(x, \Sigma^{-i}y) \oplus D \text{Hom}_{\text{D}^b(\mathbf{k}Q)}(y, \Sigma^{i-w}x).$$

2. FUNCTORIALLY FINITE HEARTS

In this section we establish an unexpected characterisation of algebraic t-structures: our third main theorem, which we state and prove first, relates homological properties of hearts to approximation properties. See [10, Proposition 3.2.2] for a similar characterisation in terms of co-t-structures. Here, \mathcal{D} will be a Hom-finite, Krull-Schmidt triangulated category.

Theorem 2.1. *Let \mathcal{H} be the heart of a bounded t-structure in \mathcal{D} . Then the following hold:*

- (1) *The heart \mathcal{H} is contravariantly finite in \mathcal{D} if and only if \mathcal{H} has enough injectives.*
- (2) *The heart \mathcal{H} is covariantly finite in \mathcal{D} if and only if \mathcal{H} has enough projectives.*
- (3) *The heart \mathcal{H} is functorially finite in \mathcal{D} if and only if \mathcal{H} has enough injectives and enough projectives.*

Proof. It suffices to prove (1) because (2) is the dual statement, and (3) is the combination of (1) and (2). Thus, let $(\mathcal{X}, \mathcal{Y})$ be a bounded t-structure in \mathcal{D} with heart \mathcal{H} .

Suppose \mathcal{H} is contravariantly finite in \mathcal{D} . We must show that every object of \mathcal{H} admits an injective envelope. Let $h \in \mathcal{H}$ and take a minimal right \mathcal{H} -approximation of Σh :

$$h' \rightarrow \Sigma h \rightarrow \Sigma e \rightarrow \Sigma h'.$$

By the triangulated Wakamatsu Lemma $\Sigma e \in \mathcal{H}^\perp$ (see e.g. [34, Lemma 2.1]). Rotating this triangle gives us the triangle $h \rightarrow e \rightarrow h' \rightarrow \Sigma h$, whence extension closure of \mathcal{H} means that $e \in \mathcal{H}$. Thus, $e \in \mathcal{H} \cap (\Sigma^{-1}\mathcal{H})^\perp$, i.e. $\text{Ext}_{\mathcal{H}}^1(-, e) = 0$ and e is injective. However, each short exact sequence $0 \rightarrow h' \rightarrow h \rightarrow h'' \rightarrow 0$ corresponds to a distinguished triangle $h' \rightarrow h \rightarrow h'' \rightarrow \Sigma h'$ and vice versa so that we have a short exact sequence $0 \rightarrow h \rightarrow e \rightarrow h' \rightarrow 0$, in which case $h \hookrightarrow e$ is an injective envelope.

Now for the converse suppose that \mathbf{H} has enough injectives. Since (\mathbf{X}, \mathbf{Y}) is a bounded t-structure, we have

$$\mathbf{D} = \bigcup_{i \geq j} \Sigma^i \mathbf{H} * \Sigma^{i-1} \mathbf{H} * \dots * \Sigma^{j+1} \mathbf{H} * \Sigma^j \mathbf{H}.$$

Since \mathbf{H} is the heart of a t-structure, we have $\mathrm{Hom}_{\mathbf{D}}(\mathbf{H}, \Sigma^{<0} \mathbf{H}) = 0$. Therefore, it is sufficient to show that \mathbf{H} is contravariantly finite in $\mathrm{susp}(\mathbf{H})$, where

$$\mathrm{susp}(\mathbf{H}) = \bigcup_{i \geq 0} \Sigma^i \mathbf{H} * \dots * \Sigma \mathbf{H} * \mathbf{H}.$$

We split the argument up into three steps.

Step 1. *For each $n \geq 0$, each object of $\Sigma^n \mathbf{H}$ admits a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$.*

When $n = 0$, this statement is trivial. To start the induction, suppose $n = 1$. For $h \in \mathbf{H}$ we show how to construct a right \mathbf{H} -approximation of Σh whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$. In \mathbf{H} , take an injective envelope of h and its cokernel, giving us a short exact sequence $0 \rightarrow h \rightarrow e \rightarrow c \rightarrow 0$ in \mathbf{H} . Rotating the corresponding triangle in \mathbf{D} gives us $c \rightarrow \Sigma h \rightarrow \Sigma e \rightarrow \Sigma c$, with $\Sigma e \in (\Sigma^{\leq 0} \mathbf{H})^\perp \cap \Sigma \mathbf{H}$ since e is injective. This tells us that the morphism $c \rightarrow \Sigma h$ is a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$.

Suppose $n > 1$ and each object of $\Sigma^{n-1} \mathbf{H}$ admits a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$. Let $h \in \mathbf{H}$ and take such a right \mathbf{H} -approximation of $\Sigma^{n-1} h$,

$$c_1 \rightarrow \Sigma^{n-1} h \rightarrow f_1 \rightarrow \Sigma c_1.$$

By induction, Σc_1 admits a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$:

$$c_2 \rightarrow \Sigma c_1 \rightarrow f_2 \rightarrow \Sigma c_2.$$

Applying the octahedral axiom to the composition $c_2 \rightarrow \Sigma c_1 \rightarrow \Sigma^n h$ gives the diagram

$$\begin{array}{ccccccc} & & f_1 & \xlongequal{\quad} & f_1 & & \\ & & \downarrow & & \downarrow & & \\ c_2 & \longrightarrow & \Sigma c_1 & \longrightarrow & f_2 & \longrightarrow & \Sigma c_2 \\ & & \downarrow & & \downarrow & & \parallel \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ c_2 & \longrightarrow & \Sigma^n h & \longrightarrow & f & \longrightarrow & \Sigma c_2 \\ & & \downarrow & & \downarrow & & \\ & & \Sigma f_1 & \xlongequal{\quad} & \Sigma f_1 & & \end{array}$$

Now since $\Sigma f_1 \in (\Sigma^{\leq 1} \mathbf{H})^\perp$ and $f_2 \in (\Sigma^{\leq 0} \mathbf{H})^\perp$, we get $f \in (\Sigma^{\leq 0} \mathbf{H})^\perp$, whence the triangle $c_2 \rightarrow \Sigma^n h \rightarrow f \rightarrow \Sigma c_2$ gives rise to a right \mathbf{H} -approximation of $\Sigma^n h$ whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$.

Step 2. *Each object of $\Sigma \mathbf{H} * \mathbf{H}$ admits a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$.*

Let $d \in \Sigma \mathbf{H} * \mathbf{H}$ and consider a decomposition $\Sigma h_1 \rightarrow d \rightarrow h_2 \rightarrow \Sigma^2 h_1$. By Step 1, we can take a right \mathbf{H} -approximation of Σh_1 ,

$$h \rightarrow \Sigma h_1 \rightarrow e \rightarrow \Sigma h,$$

where $e \in (\Sigma^{\leq 0} \mathbf{H})^\perp$. Consider the octahedral axiom applied to this triangle together with the decomposition triangle for d , which gives rise to the left-hand commutative diagram below, in which the indicated map is zero because $e \in (\Sigma^{\leq 0} \mathbf{H})^\perp$. Therefore, $f \simeq e \oplus h_2$.

Using the octahedral axiom for the composition $d \rightarrow f \simeq e \oplus h_2 \rightarrow e$, where the second morphism is the projection, gives the right-hand commutative diagram.

$$\begin{array}{ccccc}
& \Sigma^{-1}h_2 \simeq \Sigma^{-1}h_2 & & & \\
& \downarrow & & \downarrow^0 & \\
h \longrightarrow & \Sigma h_1 & \longrightarrow & e & \longrightarrow & \Sigma h \\
\parallel & \downarrow & & \downarrow & & \parallel \\
h \longrightarrow & d & \longrightarrow & f & \longrightarrow & \Sigma h \\
& \downarrow & & \downarrow & & \\
& h_2 \simeq h_2 & & & &
\end{array}
\quad
\begin{array}{ccccccc}
& & h_2 \simeq h_2 & & & & \\
& & \downarrow & & \downarrow & & \\
d \longrightarrow & e \oplus h_2 & \longrightarrow & \Sigma h & \longrightarrow & \Sigma d \\
\parallel & \downarrow & & \downarrow & & \parallel \\
d \longrightarrow & e & \longrightarrow & \Sigma h' & \longrightarrow & \Sigma d \\
& \downarrow & & \downarrow & & \\
& \Sigma h_2 \simeq \Sigma h_2 & & & &
\end{array}$$

From the right-hand diagram, we see that $h' \rightarrow d \rightarrow e \rightarrow \Sigma h'$ gives the required right \mathbf{H} -approximation.

Step 3. For each $n \geq 0$, each object of $\Sigma^n \mathbf{H} * \dots * \Sigma \mathbf{H} * \mathbf{H}$ admits a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$.

For $n = 0$, this is contained in Step 1. Therefore, assume $n > 0$ and assume, by induction, that any object of $\Sigma^{n-1} \mathbf{H} * \dots * \Sigma \mathbf{H} * \mathbf{H}$ admits a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$. Let $d \in \Sigma^n \mathbf{H} * \dots * \Sigma \mathbf{H} * \mathbf{H}$ and decompose it as $\Sigma d' \rightarrow d \rightarrow h \rightarrow \Sigma^2 d'$ with $d' \in \Sigma^{n-1} \mathbf{H} * \dots * \Sigma \mathbf{H} * \mathbf{H}$ and $h \in \mathbf{H}$. Take a right \mathbf{H} -approximation of d' ,

$$h' \rightarrow d' \rightarrow e' \rightarrow \Sigma h',$$

with $e' \in (\Sigma^{\leq 0} \mathbf{H})^\perp$. Therefore, $\Sigma e' \in (\Sigma^{\leq 1} \mathbf{H})^\perp$. Apply the octahedral axiom to the decomposition triangle for d together with the shifted approximation triangle for d' to obtain the left-hand commutative diagram below in which the first map of the second row is zero because $\Sigma e' \in (\Sigma^{\leq 1} \mathbf{H})^\perp$, making $f \cong \Sigma e' \oplus h$. Applying the octahedral axiom again, this time using the second column from the left-hand diagram and the other split triangle defining f gives the right-hand commutative diagram.

$$\begin{array}{ccccccc}
& & \Sigma h' \simeq \Sigma h' & & & & \\
& & \downarrow & & \downarrow & & \\
\Sigma^{-1}h \longrightarrow & \Sigma d' & \longrightarrow & d & \longrightarrow & h \\
\parallel & \downarrow & & \downarrow & & \parallel \\
\Sigma^{-1}h \xrightarrow{0} & \Sigma e' & \longrightarrow & f & \longrightarrow & h \\
& \downarrow & & \downarrow & & \\
& \Sigma^2 h' \simeq \Sigma^2 h' & & & &
\end{array}
\quad
\begin{array}{ccccccc}
& & h \simeq h & & & & \\
& & \downarrow & & \downarrow & & \\
d \longrightarrow & \Sigma e' \oplus h & \longrightarrow & \Sigma^2 h' & \longrightarrow & \Sigma d \\
\parallel & \downarrow & & \downarrow & & \parallel \\
d \longrightarrow & \Sigma e' & \longrightarrow & \Sigma d'' & \longrightarrow & \Sigma d \\
& \downarrow^0 & & \downarrow & & \\
& \Sigma h \simeq \Sigma h & & & &
\end{array}$$

Rotating the triangle comprising the second row of the right-hand diagram above, we get $d'' \rightarrow d \rightarrow \Sigma e' \rightarrow \Sigma d''$ with $d'' \in \Sigma \mathbf{H} * \mathbf{H}$ and $\Sigma e' \in (\Sigma^{\leq 1} \mathbf{H})^\perp$. By Step 2, we can take a right \mathbf{H} -approximation of d'' whose cone lies in $(\Sigma^{\leq 0} \mathbf{H})^\perp$:

$$h'' \rightarrow d'' \rightarrow e'' \rightarrow \Sigma h''.$$

Using the octahedral axiom once more, we get:

$$\begin{array}{ccccccc}
& & & e' & \xlongequal{\quad} & e' & \\
& & & \downarrow & & \downarrow & \\
h'' & \longrightarrow & d'' & \longrightarrow & e'' & \longrightarrow & \Sigma h'' \\
\parallel & & \downarrow & & \downarrow & & \parallel \\
h'' & \longrightarrow & d & \longrightarrow & e & \longrightarrow & \Sigma h'' \\
& & \downarrow & & \downarrow & & \\
& & \Sigma e' & \xlongequal{\quad} & \Sigma e' & &
\end{array}$$

Applying $\mathrm{Hom}(\Sigma^{\leq 0}\mathbf{H}, -)$ to the triangle in the second column shows that $e \in (\Sigma^{\leq 0}\mathbf{H})^\perp$. In particular, the map $h'' \rightarrow d$ is a right \mathbf{H} -approximation whose mapping cone lies in $(\Sigma^{\leq 0}\mathbf{H})^\perp$. \square

Using Theorem 2.1 we are able to establish the existence of Auslander–Reiten sequences.

Corollary 2.2. *Suppose \mathbf{D} is a Hom-finite, Krull-Schmidt triangulated category with Serre duality. Let \mathbf{H} be the heart of a bounded t -structure in \mathbf{D} .*

- (1) *If \mathbf{H} has enough injectives then any indecomposable object of \mathbf{H} which is not Ext-projective is the third object of an Auslander–Reiten sequence.*
- (2) *If \mathbf{H} has enough projectives then any indecomposable object of \mathbf{H} which is not Ext-injective is the first object of an Auslander–Reiten sequence.*

Proof. We only prove the first statement; the second is dual. Suppose $h \in \mathrm{ind}(\mathbf{H})$ is not Ext-projective, i.e. $\mathrm{Ext}_{\mathbf{H}}^1(h, -) = \mathrm{Hom}_{\mathbf{D}}(h, \Sigma -)|_{\mathbf{H}} \neq 0$. Then, since \mathbf{D} has Serre duality, by [42, Theorem I.2.4], \mathbf{D} has AR triangles. In particular, there is an AR triangle $x \rightarrow y \rightarrow h \rightarrow \Sigma x$ in \mathbf{D} . Since \mathbf{H} has enough injectives, by Theorem 2.1, \mathbf{H} is contravariantly finite in \mathbf{D} . Hence, x admits a minimal right \mathbf{H} -approximation. Thus we can now apply [34, Theorem 3.1] to conclude that there is an AR sequence $0 \rightarrow h' \rightarrow h'' \rightarrow h \rightarrow 0$ in \mathbf{H} . \square

Finally, we remark that Theorem 2.1 allows us to recognise module categories of finite-dimensional algebras via the approximation theory of the heart inside an ambient triangulated category. Here \mathbf{D} is again a Hom-finite \mathbf{k} -linear triangulated category.

Corollary 2.3. *Let (\mathbf{X}, \mathbf{Y}) be an algebraic t -structure in \mathbf{D} with heart \mathbf{H} . The heart \mathbf{H} is covariantly finite in \mathbf{D} if and only if $\mathbf{H} \simeq \mathrm{mod}(A)$ for a finite-dimensional \mathbf{k} -algebra A .*

Proof. By [6, Ch. II], \mathbf{H} has enough projectives if and only if $\mathbf{H} \simeq \mathrm{mod}(A)$ for some finite-dimensional \mathbf{k} -algebra A . The result now follows from Theorem 2.1. \square

The following example, however, shows that having enough projectives is not enough to detect algebraic hearts inside a Hom-finite Krull-Schmidt triangulated category.

Example 2.4. Let \mathbf{H} be a standard stable homogeneous tube. This is a length category in which every object is uniserial. Moreover, \mathbf{H} contains a single simple object, but no injective or projective objects except the zero object. It has infinitely many indecomposable objects. The category $\mathbf{D} = \mathbf{D}^b(\mathbf{H})$ is a 1-Calabi-Yau triangulated category in which the only torsion pairs are trivial or shifts of the standard t -structure, which is bounded, see [19, Theorem 9.1]. Since \mathbf{H} is length with one isoclass of simple objects, it is algebraic; but it doesn't have enough projectives or injectives and is therefore neither contravariantly finite nor covariantly finite in \mathbf{D} .

Example 2.5. Let $\mathbf{H} = \text{mod}(\mathbf{k}\tilde{A}_1)$ be the category of finite-dimensional representations of the Kronecker quiver. In the derived category $\mathbf{D} = \mathbf{D}^b(\mathbf{H})$, this heart is obviously algebraic. As is well known, \mathbf{D} is equivalent to the bounded derived category of coherent sheaves on the projective line \mathbb{P}^1 over \mathbf{k} , giving rise to another heart $\mathbf{H}' = \text{coh}(\mathbb{P}^1)$ in \mathbf{D} . The abelian category \mathbf{H}' has neither injective nor projective objects apart from 0. Hence by Theorem 2.1, the heart \mathbf{H}' is neither covariantly nor contravariantly finite in \mathbf{D} .

3. ORTHOGONAL COLLECTIONS

In this section we recall the various notions of orthogonal collections and then establish some useful characterisations of them. The main references for the definitions in this section are [16, 17, 18, 20, 22, 36].

Definitions 3.1. A collection of objects \mathbf{S} in \mathbf{D} is called a *1-orthogonal collection* (or simply *orthogonal collection*) if $\dim \text{Hom}_{\mathbf{D}}(x, y) = \delta_{xy}$ for every $x, y \in \mathbf{S}$. Let $w \geq 1$ be an integer. An orthogonal collection \mathbf{S} is said to be

- (i) (if $w > 1$) *w-orthogonal* if $\text{Hom}_{\mathbf{D}}(\Sigma^k x, y) = 0$ for $1 \leq k \leq w - 1$ and $x, y \in \mathbf{S}$;
- (ii) *∞ -orthogonal* if $\text{Hom}_{\mathbf{D}}(\Sigma^k x, y) = 0$ for $k \geq 1$ and $x, y \in \mathbf{S}$;
- (iii) a *w-simple-minded system* if it is *w-orthogonal* and $\mathbf{D} = \langle \mathbf{S} \rangle * \Sigma^{-1} \langle \mathbf{S} \rangle * \dots * \Sigma^{1-w} \langle \mathbf{S} \rangle$;
- (iv) a *simple-minded collection* if it is ∞ -orthogonal and $\mathbf{D} = \text{thick}_{\mathbf{D}}(\mathbf{S})$;
- (v) a *left w-Riedtmann configuration* if it is *w-orthogonal* and $\bigcap_{k=0}^{w-1} (\Sigma^k \mathbf{S})^\perp = 0$;
- (vi) a *right w-Riedtmann configuration* if it is *w-orthogonal* and $\bigcap_{k=0}^{w-1} (\Sigma^{-k} \mathbf{S})^\perp = 0$;
- (vii) a *w-Riedtmann configuration* if it is both a left *w-Riedtmann configuration* and a right *w-Riedtmann configuration*; and,
- (viii) an *∞ -Riedtmann configuration* if it is ∞ -orthogonal and ${}^\perp(\Sigma^{<0} \mathbf{S}) \cap (\Sigma^{\geq 0} \mathbf{S})^\perp = 0$.

In [40] an orthogonal collection is called a *system of orthogonal bricks*, in [22] a *set of (pairwise) orthogonal bricks* and in [3] a *semibrick*. In [30, 32, 33] *w-Riedtmann configurations* are called *(-w)-Calabi-Yau configurations* in light of [32, Theorem 6.2] which asserts that if \mathbf{S} is a *w-Riedtmann configuration* then $\mathbb{S}\Sigma^w \mathbf{S} = \mathbf{S}$.

Let $\mathbf{X} \subseteq \mathbf{D}$ be a collection of objects in \mathbf{D} . We set $(\mathbf{X})_1 = \mathbf{X}$ and $(\mathbf{X})_n = \mathbf{X} * (\mathbf{X})_{n-1}$. We now recall some basic properties of orthogonal collections.

Lemma 3.2. *Let \mathbf{S} be an orthogonal collection in \mathbf{D} . Then the following assertions hold:*

- (1) ([22, Lemma 2.7]) $(\mathbf{S})_n$ is closed under direct summands for each $n \geq 1$.
- (2) ([22, Lemma 2.3]) $\langle \mathbf{S} \rangle = \bigcup_{n \geq 1} (\mathbf{S})_n$.
- (3) ([20, Theorem 2.11] & [22, Theorem 3.3]) *If $\mathbf{S} \subseteq \mathbf{T}$ for an orthogonal collection \mathbf{T} in \mathbf{D} then $\langle \mathbf{S} \rangle$ is functorially finite in $\langle \mathbf{T} \rangle$.*

Lemma 3.2(2) means that the following definition makes sense.

Definition 3.3 ([22, Definition 2.5]). Let \mathbf{S} be an orthogonal collection in \mathbf{D} . The *\mathbf{S} -length* (or simply *length*) of $x \in \langle \mathbf{S} \rangle$ is the smallest natural number n such that $x \in (\mathbf{S})_n$.

Recall the following characterisation of *w-simple-minded systems* from [20].

Proposition 3.4 ([20, Proposition 2.13]). *Let \mathbf{S} be a collection of indecomposable objects in \mathbf{D} , and let $w \geq 1$ be an integer. The following conditions are equivalent:*

- (1) \mathbf{S} is a *w-simple-minded system*.
- (2) \mathbf{S} is a *right w-Riedtmann configuration* such that $\langle \mathbf{S} \rangle$ is covariantly finite in \mathbf{D} .

- (3) \mathbf{S} is a left w -Riedtmann configuration such that $\langle \mathbf{S} \rangle$ is contravariantly finite in \mathbf{D} .
(4) \mathbf{S} is a w -Riedtmann configuration such that $\langle \mathbf{S} \rangle$ is functorially finite in \mathbf{D} .

We next provide an analogue of Proposition 3.4 for simple-minded collections.

Proposition 3.5. *Let \mathbf{S} be a collection of indecomposable objects in \mathbf{D} . Then \mathbf{S} is a simple-minded collection if and only if \mathbf{S} is an ∞ -Riedtmann configuration such that $\text{susp } \mathbf{S}$ is contravariantly finite in \mathbf{D} and $\text{cosusp } \mathbf{S}$ is covariantly finite in \mathbf{D} .*

Proof. Suppose that \mathbf{S} is a simple-minded collection in \mathbf{D} . Since both simple-minded collections and ∞ -Riedtmann configurations are ∞ -orthogonal collections, we only need to check the other defining properties of ∞ -Riedtmann configurations. Then, by definition, $\text{thick}_{\mathbf{D}}(\mathbf{S}) = \mathbf{D}$ and by, e.g. [20, Lemma 2.7], we have

$$\mathbf{D} = \bigcup_{n>m} \Sigma^n \langle \mathbf{S} \rangle * \Sigma^{n-1} \langle \mathbf{S} \rangle * \dots * \Sigma^{m+1} \mathbf{S} * \Sigma^m \langle \mathbf{S} \rangle.$$

It therefore follows immediately that $(\text{susp } \mathbf{S}, \text{cosusp } \Sigma^{-1} \mathbf{S})$ is a bounded t-structure in \mathbf{D} , and, in particular, $\text{susp } \mathbf{S}$ is contravariantly finite and $\text{cosusp } \mathbf{S}$ is covariantly finite. Finally, $(\Sigma^{\geq 0} \mathbf{S})^{\perp} = (\text{susp } \mathbf{S})^{\perp} = \text{cosusp } \Sigma^{-1} \mathbf{S}$ and ${}^{\perp}(\Sigma^{< 0} \mathbf{S}) = {}^{\perp}(\text{cosusp } \Sigma^{-1} \mathbf{S}) = \text{susp } \mathbf{S}$, so that ${}^{\perp}(\Sigma^{< 0} \mathbf{S}) \cap (\Sigma^{\geq 0} \mathbf{S})^{\perp} = \text{susp } \mathbf{S} \cap \text{cosusp } \Sigma^{-1} \mathbf{S} = 0$.

Conversely, suppose that \mathbf{S} is an ∞ -Riedtmann configuration such that $\text{susp } \mathbf{S}$ is contravariantly finite in \mathbf{D} and $\text{cosusp } \mathbf{S}$ is covariantly finite in \mathbf{D} . To see that \mathbf{S} is a simple-minded collection we need only to check that $\text{thick}_{\mathbf{D}}(\mathbf{S}) = \mathbf{D}$. Let d be an object of \mathbf{D} . Since $\text{susp } \mathbf{S}$ is contravariantly finite in \mathbf{D} there is a decomposition triangle $x \rightarrow d \rightarrow y \rightarrow \Sigma x$ with respect to the t-structure $(\text{susp } \mathbf{S}, (\Sigma^{\geq 0} \mathbf{S})^{\perp})$. Now, since $\text{cosusp } \mathbf{S}$ is covariantly finite in \mathbf{D} there is a decomposition triangle of $y, \Sigma^{-1} v \rightarrow u \rightarrow y \rightarrow v$ with respect to the t-structure $({}^{\perp}(\Sigma^{< 0} \mathbf{S}), \text{cosusp } \Sigma^{-1} \mathbf{S})$. Note that $\text{cosusp } \Sigma^{-1} \mathbf{S} \subseteq (\Sigma^{\geq 0} \mathbf{S})^{\perp}$ so that $\Sigma^{-1} v \in (\Sigma^{\geq -1} \mathbf{S})^{\perp} \subseteq (\Sigma^{\geq 0} \mathbf{S})^{\perp}$. Since $(\Sigma^{\geq 0} \mathbf{S})^{\perp}$ is a perpendicular category, it is closed under extensions. It follows that $u \in {}^{\perp}(\Sigma^{< 0} \mathbf{S}) \cap (\Sigma^{\geq 0} \mathbf{S})^{\perp}$. Hence, since \mathbf{S} is ∞ -Riedtmann, we obtain that $u = 0$ and $y \simeq v$. In particular, we get

$$d \in \text{susp } \mathbf{S} * \text{cosusp } \Sigma^{-1} \mathbf{S} = \text{thick}_{\mathbf{D}}(\mathbf{S}).$$

Hence, $\mathbf{D} = \text{thick}_{\mathbf{D}}(\mathbf{S})$ and \mathbf{S} is a simple-minded collection. \square

Finally, we end this section with the following useful observation obtained by applying Theorem 2.1 in the context of simple-minded collections in bounded derived categories of finite-dimensional algebras.

Lemma 3.6. *Suppose $\mathbf{D} = \mathbf{D}^b(A)$ for some finite-dimensional \mathbf{k} -algebra A . If \mathbf{S} is a simple-minded collection in \mathbf{D} , then $\langle \mathbf{S} \rangle$ is functorially finite in \mathbf{D} .*

Proof. If \mathbf{S} is a simple-minded collection in \mathbf{D} then $\langle \mathbf{S} \rangle$ is the heart of a bounded t-structure which is equivalent to the category of finitely generated modules over a finite-dimensional algebra by [37, Proposition 5.4, Corollary 6.2]. Hence $\langle \mathbf{S} \rangle$ has enough projectives and enough injectives, so by Theorem 2.1, $\langle \mathbf{S} \rangle$ is functorially finite in \mathbf{D} . \square

4. SIMPLE-MINDED COLLECTIONS VS w -SIMPLE MINDED SYSTEMS

Let Q be a finite acyclic quiver and $w \geq 1$. Recall the construction of the negative cluster category $\mathbf{C}_{-w} = \mathbf{C}_{-w}(\mathbf{k}Q)$ as an orbit category of $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$ from Section 1.2. The aim of this section is to establish the following theorem, which is a generalisation of [30, Theorem 1.2], which was established when Q is simply-laced Dynkin.

Theorem 4.1. *Let Q be an acyclic quiver. The natural projection functor $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ induces a bijection*

$$\left\{ \begin{array}{c} \text{simple-minded collections of } \mathbf{D}^b(\mathbf{k}Q) \\ \text{contained in } \mathcal{F}_{-w} \end{array} \right\} \xleftrightarrow{1-1} \left\{ \begin{array}{c} w\text{-simple-minded systems} \\ \text{in } \mathbf{C}_{-w} \end{array} \right\}.$$

The proof of this theorem comes in two parts. We must first establish that the map induced by π is well defined and secondly that the map is surjective. Once we know that the map is well defined, injectivity follows immediately because the projection functor π induces a bijection between $\text{ind}(\mathbf{C}_{-w})$ and $\text{ind}(\mathcal{F}_{-w})$, where \mathcal{F}_{-w} is the fundamental domain; see Section 1.2 for notation.

4.1. The induced map is well defined. In this section, we establish the following.

Proposition 4.2. *Let Q be an acyclic quiver. The natural projection functor $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ induces a well-defined map*

$$\left\{ \begin{array}{c} \text{simple-minded collections of } \mathbf{D}^b(\mathbf{k}Q) \\ \text{contained in } \mathcal{F}_{-w} \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} w\text{-simple-minded systems} \\ \text{in } \mathbf{C}_{-w} \end{array} \right\}.$$

In [30, Proposition 3.6], Iyama and Jin show that there is a well-defined map

$$\left\{ \begin{array}{c} \text{simple-minded collections of } \mathbf{D}^b(\mathbf{k}Q) \\ \text{contained in } \mathcal{F}_{-w} \end{array} \right\} \longrightarrow \{w\text{-Riedtmann configurations in } \mathbf{C}_{-w}\}.$$

When Q is simply-laced Dynkin, by Proposition 3.4, this is enough to establish Proposition 4.2. However, more work is required for an arbitrary acyclic quiver Q . In fact, the following example illustrates that the map above is not surjective when Q is not simply-laced Dynkin.

Example 4.3. Let $Q = \tilde{A}_1$ be the Kronecker quiver and $w = 1$. Partition the set of homogeneous tubes into two nonempty, disjoint sets Λ and Ω and set

$$\mathbf{S} := \{S_\lambda \mid \lambda \in \Lambda\} \cup \{\Sigma S_\omega \mid \omega \in \Omega\},$$

where S_λ (resp. S_ω) denotes the quasi-simple modules lying on the mouth of the tubes indexed by Λ (resp. Ω). Then \mathbf{S} is a 1-Riedtmann configuration in $\mathbf{C}_{-1} := \mathbf{C}_{-1}(\mathbf{k}\tilde{A}_1)$; the required Hom-vanishing needs $\Lambda \neq \emptyset$ and $\Omega \neq \emptyset$. However, \mathbf{S} is not a simple-minded collection in $\mathbf{D} := \mathbf{D}^b(\mathbf{k}\tilde{A}_1)$ since $\text{thick}_{\mathbf{D}}(\mathbf{S}) \neq \mathbf{D}$. Note also that $\langle \mathbf{S} \rangle_{\mathbf{C}_{-1}}$ is not functorially finite in \mathbf{C}_{-1} and so \mathbf{S} is not a simple-minded system in \mathbf{C}_{-1} .

To prove Proposition 4.2, we need some lemmas. Recall $F = \Sigma^w \mathbf{S}: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{D}^b(\mathbf{k}Q)$.

Lemma 4.4. *Let \mathbf{S} be an ∞ -orthogonal collection of $\mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_{-w} .*

- (1) *The set $\{F^n \mathbf{S} \mid n \in \mathbb{Z}\}$ is an orthogonal collection in $\mathbf{D}^b(\mathbf{k}Q)$.*
- (2) *For $k \geq 1$, we have $\text{Ext}_{\mathbf{D}^b(\mathbf{k}Q)}^1(\mathbf{S}, F^k \mathbf{S}) = 0$.*

Proof. To prove (1), it is enough to show that $\text{Hom}_{\mathbf{D}}(S_1, F^n S_2) = 0$ for $S_1, S_2 \in \mathbf{S}$ and $n \in \mathbb{Z} \setminus \{0\}$ since \mathbf{S} is an orthogonal collection in $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$. For $n = 1$, we have $\text{Hom}_{\mathbf{D}}(S_1, F S_2) \simeq D \text{Hom}_{\mathbf{D}}(\Sigma^w S_2, S_1) = 0$, since $w \geq 1$ and \mathbf{S} is an ∞ -orthogonal collection.

Now consider $n = 2$. Recall that (\mathbf{X}, \mathbf{Y}) denotes the standard t-structure on \mathbf{D} . Since $S_1 \in \Sigma^w \mathbf{S}\mathbf{Y}$, there exists $Y \in \mathbf{Y}$ such that $S_1 = \Sigma^w \mathbf{S}Y$. Hence, $\text{Hom}_{\mathbf{D}}(S_1, F^2 S_2) \simeq D \text{Hom}_{\mathbf{D}}(\Sigma^w S_2, Y) = 0$, since S_2 , and so $\Sigma^w S_2$, lie in \mathbf{X} .

Now let $n > 2$. By the hereditary property, we have $F^n S_2 \in \mathbb{S}^n \Sigma^{nw} \mathbf{X} \subseteq \Sigma^{nw} \mathbf{X}$ and $S_1 \in \mathbb{S} \Sigma^w \mathbf{Y} \subseteq \Sigma^{w+1} \mathbf{Y}$. Therefore, $\mathrm{Hom}_{\mathbf{D}}(S_1, F^n S_2) \simeq \mathrm{Hom}_{\mathbf{D}}(Y, \Sigma^{(n-1)w-1} X)$, for some $Y \in \mathbf{Y}$ and $X \in \mathbf{X}$. Since $(n-1)w-1 \geq 1$, it follows that $\mathrm{Hom}_{\mathbf{D}}(S_1, F^n S_2) = 0$, again by the hereditary property.

Finally, let $n \leq -1$. Since $S_2 \in \Sigma^w \mathbb{S} \mathbf{Y}$, we get $F^n S_2 \in \Sigma^{(n+1)w} \mathbb{S}^{n+1} \mathbf{Y} \subseteq \Sigma^{(n+1)w} \mathbf{Y} \subseteq \mathbf{Y}$ by the hereditary property, the fact that \mathbf{Y} is the co-aisle of a t-structure, and $(n+1)w \leq 0$. It follows that $\mathrm{Hom}_{\mathbf{D}}(S_1, F^n S_2) = 0$ since $S_1 \in \mathbf{X}$.

Statement (2) follows from the ∞ -orthogonality property of \mathbf{S} when $k = 1$, and from the argument above for $n > 2$, when $k \geq 2$. \square

Corollary 4.5. *Let \mathbf{S} be an ∞ -orthogonal collection of $\mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_w . Then $\langle F^n \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \langle \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} \subseteq \langle \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \langle F^n \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}$, for all $n \geq 1$.*

Proof. Given $D \in \langle F^n \mathbf{S} \rangle_{\mathbf{D}} * \langle \mathbf{S} \rangle_{\mathbf{D}}$, we have a triangle $F^n S_1 \rightarrow D \rightarrow S_2 \xrightarrow{\alpha} \Sigma F^n S_1$, with $S_1, S_2 \in \langle \mathbf{S} \rangle$. It follows from Lemma 4.4(2) that $\alpha = 0$. Hence, the triangle splits, and $D \simeq S_2 \oplus F^n S_1 \in \langle \mathbf{S} \rangle_{\mathbf{D}} * \langle F^n \mathbf{S} \rangle_{\mathbf{D}}$. \square

Following Lemma 4.4 and Corollary 4.5, for an ∞ -orthogonal collection \mathbf{S} of $\mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_{-w} , we define the following extension-closed subcategory of $\mathbf{D}^b(\mathbf{k}Q)$:

$$\mathcal{E}_{\mathbf{S}} := \langle F^n \mathbf{S} \mid n \in \mathbb{Z} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}.$$

We now show that each object of $\mathcal{E}_{\mathbf{S}}$ admits a filtration by objects in $\{F^n \mathbf{S} \mid n \in \mathbb{Z}\}$.

Lemma 4.6. *Let \mathbf{S} be an ∞ -orthogonal collection of $\mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_{-w} . Then $\mathcal{E}_{\mathbf{S}}$ is closed under direct summands and*

$$\mathcal{E}_{\mathbf{S}} = \bigcup_{m < n} \langle F^m \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \langle F^{m+1} \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \cdots * \langle F^n \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}.$$

Proof. The first statement follows immediately from Lemma 4.4(1) and Lemma 3.2(1).

By Lemma 4.4(1) and Lemma 3.2(2), we have $\mathcal{E}_{\mathbf{S}} = \bigcup_{n \geq 1} (\mathbb{T})_n$, where $\mathbb{T} := \{F^n \mathbf{S} \mid n \in \mathbb{Z}\}$. Hence, for $D \in \mathcal{E}_{\mathbf{S}}$ there is a tower of the form

$$(1) \quad \begin{array}{ccccccccccc} 0 = D_0 & \longrightarrow & D_1 & \longrightarrow & D_2 & \longrightarrow & \cdots & \longrightarrow & D_{n-1} & \longrightarrow & D_n = D \\ & & \searrow & & \downarrow & & \searrow & & \downarrow & & \searrow \\ & & & & F^{i_1} S_1 & & & & F^{i_2} S_2 & & \cdots & & F^{i_{n-1}} S_{n-1} & & F^{i_n} S_n \end{array}$$

with $S_\ell \in \mathbf{S}$ for $1 \leq \ell \leq n$. By Corollary 4.5, we can re-order the indices in the tower above so that $i_1 \leq i_2 \leq \cdots \leq i_n$. Therefore, $\mathcal{E}_{\mathbf{S}} \subseteq \bigcup_{m < n} \langle F^m \mathbf{S} \rangle_{\mathbf{D}} * \langle F^{m+1} \mathbf{S} \rangle_{\mathbf{D}} * \cdots * \langle F^n \mathbf{S} \rangle_{\mathbf{D}}$, where $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$. The other inclusion is trivial. \square

Before we are able to prove Proposition 4.2 we need the following lemmas connecting the functorial finiteness of $\mathcal{E}_{\mathbf{S}}$ in $\mathbf{D}^b(\mathbf{k}Q)$ and the functorial finiteness of the extension closure of \mathbf{S} in \mathcal{C}_{-w} .

Lemma 4.7. *Let $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathcal{C}_{-w}$ be the natural projection functor. If \mathbf{S} is an ∞ -orthogonal collection of $\mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_{-w} , then $\pi(\mathcal{E}_{\mathbf{S}}) = \langle \mathbf{S} \rangle_{\mathcal{C}_{-w}}$.*

Proof. To see that $\pi(\mathcal{E}_{\mathbf{S}}) \subseteq \langle \mathbf{S} \rangle_{\mathcal{C}_{-w}}$, observe that, by Lemma 4.6, $D \in \mathcal{E}_{\mathbf{S}}$ admits a tower as in (1) with each $S_\ell \in \mathbf{S}$. Since $\pi(F^{i_\ell} S_\ell) = S_\ell \in \mathbf{S}$, applying π to the tower above gives a filtration of $\pi(D)$ in $\langle \mathbf{S} \rangle_{\mathcal{C}_{-w}}$.

Conversely, we need to show that if $D \in \pi^{-1}(\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}})$ then $D \in \mathcal{E}_{\mathbf{S}}$. It is enough to check for D indecomposable. Let $D \in \pi^{-1}(\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}})$ be indecomposable; note that $D \in F^k \mathcal{F}_{-w}$ for some $k \in \mathbb{Z}$. We proceed by induction on the \mathbf{S} -length of $\pi(D)$ in \mathbf{C}_{-w} . If $D \in \pi^{-1}(\mathbf{S})$, then $D = F^k \mathbf{S}$ for some $\mathbf{S} \in \mathbf{S}$. Hence, $D \in \{F^i \mathbf{S} \mid i \in \mathbb{Z}\} \subseteq \mathcal{E}_{\mathbf{S}}$. Suppose $n > 1$. Assume, by induction, that if $D \in \pi^{-1}(\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}})$ is such that $\pi(D)$ has \mathbf{S} -length $n - 1$ in \mathbf{C}_{-w} then $D \in \mathcal{E}_{\mathbf{S}}$ has $\{F^i \mathbf{S} \mid i \in \mathbb{Z}\}$ -length $n - 1$ in $\mathcal{E}_{\mathbf{S}}$. Now suppose $D \in \pi^{-1}(\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}})$ is such that $\pi(D)$ has \mathbf{S} -length n in \mathbf{C}_{-w} . Then, by Lemma 3.2(2), there is a triangle in \mathbf{C}_{-w} of the form

$$C \rightarrow \pi(D) \rightarrow S \rightarrow \Sigma C,$$

with $S \in \mathbf{S}$ and $C \in \langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ has \mathbf{S} -length $n - 1$ in \mathbf{C}_{-w} . This triangle is the image under π of a triangle in $\mathbf{D}^b(\mathbf{k}Q)$ of the form

$$F^i D' \rightarrow D \rightarrow F^j S \rightarrow \Sigma F^i D',$$

where $i \in \{k, k - 1\}$ and $j \in \{k, k + 1\}$, by Lemma 1.3. Since $\pi(F^i D') = C$, it follows by induction that $F^i D' \in \mathcal{E}_{\mathbf{S}}$ has $\{F^i \mathbf{S} \mid i \in \mathbb{Z}\}$ -length $n - 1$ in $\mathbf{D}^b(\mathbf{k}Q)$, whence $D \in \mathcal{E}_{\mathbf{S}}$ has $\{F^i \mathbf{S} \mid i \in \mathbb{Z}\}$ -length n . \square

Lemma 4.8. *Let $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ be the natural projection functor. Suppose \mathbf{S} is an ∞ -orthogonal collection of $\mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_{-w} . Then $\mathcal{E}_{\mathbf{S}}$ is contravariantly (resp. covariantly, functorially) finite in $\mathbf{D}^b(\mathbf{k}Q)$ if and only if $\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ is contravariantly (resp. covariantly, functorially) finite in \mathbf{C}_{-w} .*

Proof. We only establish the contravariantly finite statements; the covariantly finite statement is similar and the functorially finite statement follows from combining both.

Suppose $\mathcal{E}_{\mathbf{S}}$ is contravariantly finite in $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$. For $D \in \mathcal{F}_{-w}$, take a minimal right $\mathcal{E}_{\mathbf{S}}$ -approximation of D in \mathbf{D} and extend it to a triangle,

$$E_D \xrightarrow{\alpha} D \longrightarrow X_D \longrightarrow \Sigma E_D.$$

By the triangulated Wakamatsu lemma (see, e.g. [34, Lemma 2.1]), $X_D \in (\mathcal{E}_{\mathbf{S}})^\perp$. Applying π to this triangle, we get the following triangle in \mathbf{C}_{-w} :

$$\pi(E_D) \xrightarrow{\pi(\alpha)} \pi(D) \longrightarrow \pi(X_D) \longrightarrow \Sigma \pi(E_D).$$

By Lemma 4.7, we have $\pi(E_D) \in \langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$. By construction of \mathbf{C}_{-w} , see Section 1.2, we have $\mathrm{Hom}_{\mathbf{C}_{-w}}(\mathbf{S}, \pi(X_D)) = \bigoplus_{i \in \mathbb{Z}} \mathrm{Hom}_{\mathbf{D}}(F^i \mathbf{S}, X_D) = 0$ since $X_D \in (\mathcal{E}_{\mathbf{S}})^\perp$. Therefore, $\pi(\alpha)$ is a right $\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ -approximation of $D = \pi(D)$, and so $\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ is contravariantly finite in \mathbf{C}_{-w} .

Conversely, suppose $\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ is contravariantly finite in \mathbf{C}_{-w} . Let $D \in \mathbf{D}$ and consider the extension of a minimal right $\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ -approximation of $\pi(D)$ to a triangle in \mathbf{C}_{-w} ,

$$S_D \longrightarrow \pi(D) \longrightarrow Y_D \longrightarrow \Sigma S_D,$$

where $\mathrm{Hom}_{\mathbf{C}_{-w}}(\mathbf{S}, Y_D) = 0$ by the triangulated Wakamatsu lemma. Since the triangulated structure on \mathbf{C}_{-w} is induced by that of \mathbf{D} , this triangle is the image under π of a triangle,

$$E_D \xrightarrow{f} D \longrightarrow X_D \longrightarrow \Sigma E_D$$

in \mathbf{D} . It follows from Lemma 4.7 that $E_D \in \mathcal{E}_{\mathbf{S}}$. On the other hand, we have $0 = \mathrm{Hom}_{\mathbf{C}_{-w}}(\mathbf{S}, Y_D) = \bigoplus_{i \in \mathbb{Z}} \mathrm{Hom}_{\mathbf{D}}(F^i \mathbf{S}, X_D)$. Hence, $\mathrm{Hom}_{\mathbf{D}}(F^i \mathbf{S}, X_D) = 0$, for all $i \in \mathbb{Z}$, and so $X_D \in (\mathcal{E}_{\mathbf{S}})^\perp$. Therefore, the map f is a right $\mathcal{E}_{\mathbf{S}}$ -approximation of D , from which it follows that $\mathcal{E}_{\mathbf{S}}$ is contravariantly finite in \mathbf{D} . \square

We are almost ready to prove Proposition 4.2. First, we set up a final piece of notation. For integers $m \leq n$ and an ∞ -orthogonal collection \mathbf{S} , we set

$$\mathcal{E}_{\mathbf{S}}^{[m,n]} := \langle F^m \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \langle F^{m+1} \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \cdots * \langle F^n \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} \subseteq \mathcal{E}_{\mathbf{S}}.$$

Proof of Proposition 4.2. Let \mathbf{S} be a simple-minded collection in $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$ contained in \mathcal{F}_w . By the dual of [30, Proposition 3.6], $\pi(\mathbf{S})$ is a left w -Riedtmann configuration in \mathbf{C}_{-w} . Hence, by Proposition 3.4, it only remains to check that $\langle \mathbf{S} \rangle_{\mathbf{C}_{-w}}$ is contravariantly finite in \mathbf{C}_{-w} . Hence, by Lemma 4.8, it suffices to show that $\mathcal{E}_{\mathbf{S}}$ is contravariantly finite in \mathbf{D} .

Let $D \in \mathbf{D}$. There are only finitely many $i \in \mathbb{Z}$ with $\mathrm{Hom}_{\mathbf{D}}(F^i \mathbf{S}, D) \neq 0$ as $\mathbf{k}Q$ is hereditary. Let $m, n \in \mathbb{Z}$ such that $m \leq n$ and $\mathrm{Hom}_{\mathbf{D}}(F^{< m} \mathbf{S}, D) = 0 = \mathrm{Hom}_{\mathbf{D}}(F^{> n} \mathbf{S}, D)$. By Lemma 3.6, $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is contravariantly finite in $\mathbf{D}^b(\mathbf{k}Q)$. Hence, $\mathcal{E}_{\mathbf{S}}^{[m,n]}$ is contravariantly finite in $\mathbf{D}^b(\mathbf{k}Q)$ by [45, Lemma 5.3(1)]. Since any component of a nonzero morphism from an object of $\mathcal{E}_{\mathbf{S}}$ to D must originate from a summand $E \in \mathcal{E}_{\mathbf{S}}^{[m,n]}$, we have that any right $\mathcal{E}_{\mathbf{S}}^{[m,n]}$ -approximation of D is also a right $\mathcal{E}_{\mathbf{S}}$ -approximation. Hence, $\mathcal{E}_{\mathbf{S}}$ is contravariantly finite in \mathbf{D} . \square

4.2. The induced map is surjective. In order to complete the proof of Theorem 4.1, we have to show that the map induced by the natural projection functor $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ in Proposition 4.2 is surjective. To see this, we need a lemma, which first requires some notation. For an ∞ -orthogonal collection \mathbf{S} in $\mathbf{D}^b(\mathbf{k}Q)$ and an integer n , we define the following subcategories of $\mathcal{E}_{\mathbf{S}}$:

$$\begin{aligned} \mathcal{E}_{\mathbf{S}}^{\leq n} &:= \bigcup_{i \leq n} \langle F^i \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \langle F^{i+1} \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \cdots * \langle F^n \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}; \text{ and,} \\ \mathcal{E}_{\mathbf{S}}^{\geq n} &:= \bigcup_{i \geq n} \langle F^n \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \langle F^{n+1} \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)} * \cdots * \langle F^i \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}. \end{aligned}$$

Similarly, we also set $\mathcal{E}_{\mathbf{S}}^{< n} = \mathcal{E}_{\mathbf{S}}^{\leq n-1}$ and $\mathcal{E}_{\mathbf{S}}^{> n} = \mathcal{E}_{\mathbf{S}}^{\geq n+1}$.

Lemma 4.9. *Let \mathbf{S} be an ∞ -orthogonal collection in $\mathbf{D}^b(\mathbf{k}Q)$. Then*

- (1) $\langle \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}$ is contravariantly finite in $\mathcal{E}_{\mathbf{S}}^{\leq 0}$; and,
- (2) $\langle \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}$ is covariantly finite in $\mathcal{E}_{\mathbf{S}}^{\geq 0}$.

Proof. We only prove the first statement; the second statement is dual. Write $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$. Let $D \in \mathcal{E}_{\mathbf{S}}^{\leq 0}$. First note that if $\mathrm{Hom}_{\mathbf{D}}(\mathbf{S}, D) = 0$, then D admits a (minimal) right $\langle \mathbf{S} \rangle_{\mathbf{D}}$ -approximation, namely $S_D = 0 \rightarrow D$, whose cone $X_D \simeq D \in \mathbf{S}^{\perp}$. Therefore, we may assume that $\mathrm{Hom}_{\mathbf{D}}(\mathbf{S}, D) \neq 0$.

By Lemma 4.6, d admits a decomposition,

$$(2) \quad E^{< 0} \xrightarrow{\alpha} D \xrightarrow{\beta} E^0 \longrightarrow \Sigma E^{< 0},$$

with $E^{< 0} \in \mathcal{E}_{\mathbf{S}}^{< 0}$ and $E^0 \in \langle \mathbf{S} \rangle_{\mathbf{D}}$. We shall construct a right $\langle \mathbf{S} \rangle_{\mathbf{D}}$ -approximation of D by induction on the \mathbf{S} -length of E^0 in $\langle \mathbf{S} \rangle_{\mathbf{D}}$.

Suppose the \mathbf{S} -length of E^0 is one, i.e. $E^0 \in \mathbf{S}$, and suppose $\varphi: S \rightarrow D$ is a nonzero map with $S \in \mathbf{S}$. If $\beta\varphi = 0$, then $\varphi = 0$ because it factors through $E^{< 0}$ and, by Lemma 4.4, $\mathrm{Hom}_{\mathbf{D}}(\mathbf{S}, E^{< 0}) = 0$. Hence, $\beta\varphi \neq 0$, and since $S, E^0 \in \mathbf{S}$, $\beta\varphi$ must be an isomorphism because \mathbf{S} is an orthogonal collection in \mathbf{D} . Therefore, β is a split epimorphism and $D \simeq E^0 \oplus E^{< 0}$. It follows that $S \simeq E^0$ is the unique $S \in \mathbf{S}$, up to isomorphism, such

that $\text{Hom}_{\mathbb{D}}(S, D) \neq 0$. Hence, $E^0 \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} D$ is a right $\langle \mathbb{S} \rangle_{\mathbb{D}}$ -approximation of D , whose cone $X_D = E^{<0} \in \mathbb{S}^{\perp}$.

Now suppose the \mathbb{S} -length of E^0 is $n > 1$, i.e. there is a triangle $S' \rightarrow E^0 \rightarrow S \rightarrow \Sigma S'$ with $S \in \mathbb{S}$, and $S' \in \langle \mathbb{S} \rangle_{\mathbb{D}}$ of \mathbb{S} -length $n - 1$. Combining this triangle with (2), we get the octahedral diagram:

$$\begin{array}{ccccccc}
& & S' & \xlongequal{\quad} & S' & & \\
& & \downarrow & & \downarrow & & \\
D & \longrightarrow & E^0 & \longrightarrow & \Sigma E^{<0} & \longrightarrow & \Sigma D \\
\parallel & & \downarrow & & \downarrow & & \parallel \\
D & \longrightarrow & S & \longrightarrow & \Sigma A & \longrightarrow & \Sigma D \\
& & \downarrow & & \downarrow & & \\
& & \Sigma S' & \xlongequal{\quad} & \Sigma S' & &
\end{array}$$

The right-hand vertical triangle gives a decomposition of A in $\mathcal{E}_{\mathbb{S}}^{\leq 0}$, $E^{<0} \rightarrow A \rightarrow S' \rightarrow \Sigma E^{<0}$, in which the \mathbb{S} -length of S' is $n - 1$. Hence, by induction, A admits a right $\langle \mathbb{S} \rangle_{\mathbb{D}}$ -approximation $S_A \rightarrow A$, whose cone X_A lies in \mathbb{S}^{\perp} . Consider the following octahedral diagram:

$$(3) \quad \begin{array}{ccccccc}
& & \Sigma^{-1} S & \xlongequal{\quad} & \Sigma^{-1} S & & \\
& & \downarrow & & \downarrow & & \\
S_A & \longrightarrow & A & \longrightarrow & X_A & \longrightarrow & \Sigma S_A \\
\parallel & & \downarrow & & \downarrow & & \parallel \\
S_A & \xrightarrow{\alpha} & D & \longrightarrow & Y & \longrightarrow & \Sigma S_A \\
& & \downarrow & & \downarrow \beta & & \\
& & S & \xlongequal{\quad} & S & &
\end{array}$$

If $Y \in \mathbb{S}^{\perp}$ then α is a right $\langle \mathbb{S} \rangle_{\mathbb{D}}$ -approximation of D with cone lying in \mathbb{S}^{\perp} , and we are done. So suppose $Y \notin \mathbb{S}^{\perp}$. For $S_1 \in \mathbb{S}$, applying $\text{Hom}_{\mathbb{D}}(S_1, -)$ to the right-hand vertical triangle in (3), shows that the map $\text{Hom}_{\mathbb{D}}(S_1, \beta): \text{Hom}_{\mathbb{D}}(S_1, Y) \hookrightarrow \text{Hom}_{\mathbb{D}}(S_1, S)$ is injective because $X_A \in \mathbb{S}^{\perp}$. Hence, if $S_1 \neq S$, then $\text{Hom}_{\mathbb{D}}(S_1, Y) = 0$. Since $Y \notin \mathbb{S}^{\perp}$, it follows that we must have $\text{Hom}_{\mathbb{D}}(S, Y) \neq 0$ and $\text{Hom}_{\mathbb{D}}(S, \beta)$ is injective. Suppose $\gamma \in \text{Hom}_{\mathbb{D}}(S, Y)$ is nonzero. Then $\beta\gamma$ must be an isomorphism, which implies that β is a split epimorphism, whence $Y \simeq S \oplus X_A$.

Now, using the other split triangle, we get a new octahedral diagram:

$$\begin{array}{ccccccc}
& & S & \xlongequal{\quad} & S & & \\
& & \downarrow & & \downarrow & & \\
D & \longrightarrow & X_A \oplus S & \longrightarrow & \Sigma S_A & \longrightarrow & \Sigma D \\
\parallel & & \downarrow & & \downarrow & & \parallel \\
D & \longrightarrow & X_A & \longrightarrow & \Sigma S_D & \xrightarrow{-\Sigma g} & \Sigma D \\
& & \downarrow & & \downarrow & & \\
& & \Sigma S & \xlongequal{\quad} & \Sigma S & &
\end{array}$$

Clearly, $S_D \in \langle \mathbb{S} \rangle_{\mathbb{D}}$, and since $X_A \in \mathbb{S}^{\perp}$, g is a right $\langle \mathbb{S} \rangle_{\mathbb{D}}$ -approximation of D with cone in \mathbb{S}^{\perp} . This finishes the proof. \square

We are now ready to prove Theorem 4.1.

Proof of Theorem 4.1. By Proposition 4.2, this map is well defined and it is clearly injective since π gives a bijection between $\text{ind}(\mathcal{F}_{-w})$ and $\text{ind}(\mathcal{C}_{-w})$. It remains to show that π induces a surjection.

Let \mathbf{S} be a w -simple-minded system in \mathcal{C}_{-w} . The fact that the lift of \mathbf{S} to $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$, which will also be denoted by \mathbf{S} , is an ∞ -orthogonal collection in \mathbf{D} follows from the proof of [30, Theorem 1.2], as this part of the proof does not require Q to be Dynkin. Hence, it remains to show that $\text{thick}_{\mathbf{D}}(\mathbf{S}) = \mathbf{D}$.

First, we claim that $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is functorially finite in $\mathcal{E}_{\mathbf{S}}$. Indeed, on the one hand we have that $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is contravariantly finite in $\mathcal{E}_{\mathbf{S}}^{\leq 0}$ by Lemma 4.9. On the other hand, since $\mathcal{E}_{\mathbf{S}} = \mathcal{E}_{\mathbf{S}}^{\leq 0} * \mathcal{E}_{\mathbf{S}}^{> 0}$ by Lemma 4.6, and $\{F^n \mathbf{S} \mid n \in \mathbb{Z}\}$ is an orthogonal collection in \mathbf{D} , we have that $\mathcal{E}_{\mathbf{S}}^{\leq 0}$ is contravariantly finite in $\mathcal{E}_{\mathbf{S}}$. Hence, by transitivity, $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is contravariantly finite in $\mathcal{E}_{\mathbf{S}}$. Dually, using $\mathcal{E}_{\mathbf{S}}^{\geq 0}$, $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is also covariantly finite in $\mathcal{E}_{\mathbf{S}}$, and therefore $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is functorially finite in $\mathcal{E}_{\mathbf{S}}$. Since \mathbf{S} is a w -simple-minded system in \mathcal{C}_{-w} , we have that $\langle \mathbf{S} \rangle_{\mathcal{C}_{-w}}$ is functorially finite in \mathcal{C}_{-w} . It then follows by Lemma 4.8 and transitivity that $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is functorially finite in \mathbf{D} .

We will now show that $\text{thick}_{\mathbf{D}}(\mathbf{S}) = \mathbf{D}$. Since $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is functorially finite in \mathbf{D} , we have that $\langle \mathbf{S} \rangle_{\mathbf{D}}$ is contravariantly finite in $(\Sigma^{> 0} \mathbf{S})^{\perp}$ and covariantly finite in ${}^{\perp}(\Sigma^{< 0} \mathbf{S})$. And since $\mathbf{k}Q$ is hereditary, for any $D \in \mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$, we have $\text{Hom}_{\mathbf{D}}(\Sigma^i D, \mathbf{S}) = 0 = \text{Hom}_{\mathbf{D}}(\Sigma^i \mathbf{S}, D)$ for $i \gg 0$. Therefore, by [33, Proposition 3.2], we have that $({}^{\perp}(\Sigma^{< 0} \mathbf{S}), \text{cosusp } \mathbf{S})$ is a t-structure in \mathbf{D} .

Without loss of generality, we may assume that $0 \neq D \in \langle \Sigma^{\leq -w-2} \text{mod}(\mathbf{k}Q) \rangle_{\mathbf{D}}$. As a consequence, $\text{Hom}_{\mathbf{D}}(D, \Sigma^{\geq 0} \mathbf{S}) = 0$. Consider the decomposition triangle of D with respect to the t-structure $({}^{\perp}(\Sigma^{< 0} \mathbf{S}), \text{cosusp } \mathbf{S})$:

$$X_D \rightarrow D \rightarrow Y_D \rightarrow \Sigma X_D.$$

The morphism $X_D \rightarrow D$ is thus a minimal right ${}^{\perp}(\Sigma^{< 0} \mathbf{S})$ -approximation, which is therefore zero if and only if $X_D \simeq 0$. Suppose $X_D \neq 0$, then the morphism $X_D \rightarrow D$ must be nonzero. Since $\mathbf{k}Q$ is hereditary, this means that $X_D \in \langle \Sigma^{\leq -w-2} \text{mod}(\mathbf{k}Q) \rangle_{\mathbf{D}}$ also, in which case, $\text{Hom}_{\mathbf{D}}(X_D, \Sigma^{\geq 0} \mathbf{S}) = 0$. Hence $X_D \in {}^{\perp}(\Sigma^{\mathbb{Z}} \mathbf{S})$. Now, by the final part of the proof of [30, Theorem 1.2], we can conclude that $X_D \simeq 0$. Hence $D \simeq Y_D \in \text{cosusp } \mathbf{S} \subseteq \text{thick}_{\mathbf{S}}(\mathbf{D})$. It follows that $\text{thick}_{\mathbf{D}}(\mathbf{S}) = \mathbf{D}$, as required. \square

5. SINCERE ORTHOGONAL COLLECTIONS

Let Q be an acyclic quiver. In this section we establish a bijection between w -simple-minded systems in $\mathcal{C}_{-w}(\mathbf{k}Q)$ and sincere ∞ -orthogonal collections of $\mathbf{D}^b(\mathbf{k}Q)$ sitting in some truncation of the fundamental domain of $\mathcal{C}_{-w}(\mathbf{k}Q)$; see Theorem 5.8. This result generalises [16, Theorem 4.8], which established the same result in the case that $w = 1$ and Q is simply-laced Dynkin. In Section 6, we will use Theorem 5.8 to establish a bijection between w -simple-minded systems and positive w -noncrossing partitions in the Weyl group of the corresponding type. However, we believe that Theorem 5.8 holds independent representation-theoretic interest.

Before proceeding, we require some background on exceptional sequences.

5.1. Exceptional sequences. The notion of exceptional sequence goes back to the Moscow school in the 1980s; see e.g. [44]. Recall from [9] that an object e in a triangulated category \mathbf{D} is called *exceptional* if $\text{Hom}_{\mathbf{D}}(e, \Sigma^i e) = 0$ for all $i \neq 0$ and $\text{Hom}_{\mathbf{D}}(e, e) \simeq \mathbf{k}$. An

ordered collection of exceptional objects $\mathbf{E} = (e_1, \dots, e_r)$ of \mathbf{D} is called an *exceptional sequence* if $\mathrm{Hom}_{\mathbf{D}}(e_j, \Sigma^i e_k) = 0$ for all $i \in \mathbb{Z}$ and $j > k$. The exceptional sequence \mathbf{E} is called *complete* if $\mathrm{thick}_{\mathbf{D}}(\mathbf{E}) = \mathbf{D}$. If Q is an acyclic quiver with n vertices and $\mathbf{E} = (E_1, \dots, E_r)$ is an exceptional sequence in $\mathbf{D}^b(\mathbf{k}Q)$ then \mathbf{E} is complete if and only if $r = n$; see [15, 21, 43].

Exceptional sequences are often called ‘exceptional collections’. In this article we avoid this term to avoid ambiguity when considering orthogonal collections.

The following lemma gives a link between exceptional sequences and simple-minded collections in $\mathbf{D}^b(\mathbf{k}Q)$.

Lemma 5.1. *Let Q be an acyclic quiver with n vertices. If \mathbf{S} is a simple-minded collection in $\mathbf{D}^b(\mathbf{k}Q)$ then the objects of \mathbf{S} can be ordered into a (complete) exceptional sequence $\mathbf{E} = (E_1, \dots, E_n)$ in which the cohomological degrees of the E_i are weakly decreasing.*

Proof. By [15, Lemma 2.3], the objects of a $\mathrm{Hom}_{\leq 0}$ -configuration (for the definition, see [15, §2.2]) can be ordered into a complete exceptional sequence $\mathbf{E} = (E_1, \dots, E_n)$ in which the cohomological degrees of the E_i are weakly decreasing, and therefore strongly generate $\mathbf{D} = \mathbf{D}^b(\mathbf{k}Q)$. Hence the set of $\mathrm{Hom}_{\leq 0}$ -configurations in \mathbf{D} is a subset of the set of simple-minded collections. By [15, Theorem 2.4], the set of $\mathrm{Hom}_{\leq 0}$ -configurations in \mathbf{D} is in bijection with the set of silting objects in \mathbf{D} . However, the map from the set of $\mathrm{Hom}_{\leq 0}$ -configurations in \mathbf{D} to the set of silting objects in \mathbf{D} constructed in [15] coincides with the map from simple-minded collections in \mathbf{D} to silting objects in \mathbf{D} ; see [37, §5.6]. Hence the set of $\mathrm{Hom}_{\leq 0}$ -configurations in \mathbf{D} coincides with the set of simple-minded collections in \mathbf{D} , giving the lemma. \square

5.2. Sincere orthogonal collections. Let Q be an acyclic quiver and $w \geq 1$ be an integer. Write $\mathbf{H} := \mathrm{mod}(\mathbf{k}Q)$ and consider the functor

$$(4) \quad H: \mathbf{D}^b(\mathbf{k}Q) \longrightarrow \mathbf{H} \text{ given by } X \longmapsto \bigoplus_{i \in \mathbb{Z}} H^i(X),$$

where $H^i(-)$ denotes the i^{th} -cohomology of X with respect to the standard t-structure (\mathbf{X}, \mathbf{Y}) in $\mathbf{D}^b(\mathbf{k}Q)$.

Recall that a module $M \in \mathbf{H}$ is called *sincere* if $\mathrm{Hom}_{\mathbf{H}}(P, M) \neq 0$ for any projective module P . A set of modules $\mathbf{X} \subseteq \mathbf{H}$ will be called a *sincere set of modules* if for each projective module P there exists a module $X \in \mathbf{X}$ such that $\mathrm{Hom}_{\mathbf{H}}(P, X) \neq 0$. If \mathbf{X} is a finite set, then \mathbf{X} is a sincere set of modules if and only if $\bigoplus_{X \in \mathbf{X}} X$ is a sincere module. Note that there are equivalent formulations of sincerity using injective modules. Using sincerity, we make the following definition.

Definition 5.2. An ∞ -orthogonal collection $\mathbf{S} \subseteq \mathbf{D}^b(\mathbf{k}Q)$ will be called *w-sincere* if $\mathbf{S} \subseteq \mathbf{X} \cap \Sigma^w \mathbf{Y} \subseteq \mathcal{F}_{-w}$ and $\{H(S) \mid S \in \mathbf{S}\}$ is a sincere set of modules in \mathbf{H} . It will be called *exceptionally finite* if the objects of \mathbf{S} can be ordered into an exceptional sequence and $\langle \mathbf{S} \rangle_{\mathbf{D}^b(\mathbf{k}Q)}$ is functorially finite in $\mathbf{D}^b(\mathbf{k}Q)$.

Recall $\pi: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \mathbf{C}_{-w}$ is the canonical projection functor.

Proposition 5.3. *Let Q be an acyclic quiver and $w \geq 1$ an integer. Then there is a well-defined map*

$$\Theta: \left\{ \begin{array}{l} w\text{-Riedtmann configurations} \\ \text{in } \mathbf{C}_{-w} \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} w\text{-sincere } \infty\text{-orthogonal} \\ \text{collections in } \mathbf{D}^b(\mathbf{k}Q) \end{array} \right\},$$

$$\mathbf{S} \longmapsto \pi^{-1}(\mathbf{S}) \cap \mathbf{X} \cap \Sigma^w \mathbf{Y}$$

which restricts to a well-defined map

$$\Theta: \left\{ \begin{array}{l} w\text{-simple-minded} \\ \text{systems in } \mathcal{C}_{-w} \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} \text{exceptionally finite, } w\text{-sincere } \infty\text{-orthogonal} \\ \text{collections in } \mathcal{D}^b(\mathbf{k}Q) \end{array} \right\}.$$

Proof. Let \mathbf{S} be a w -Riedtmann configuration in \mathcal{C}_{-w} . Partition the lift of \mathbf{S} into \mathcal{F}_{-w} , $\pi^{-1}(\mathbf{S}) \cap \mathcal{F}_{-w}$, into $\mathbf{R} \cup \mathbf{T}$, where $\mathbf{R} \subseteq \mathbf{X} \cap \Sigma^w \mathbf{Y}$ and $\mathbf{T} \subseteq \Sigma^w \mathbf{X} \cap \mathbb{S} \Sigma^w \mathbf{Y}$. We need to show that \mathbf{R} is a w -sincere ∞ -orthogonal configuration. As in the proof of Theorem 4.1, we can invoke [30, Theorem 1.2] to see that \mathbf{R} is ∞ -orthogonal. By definition $\mathbf{R} \subseteq \mathbf{X} \cap \Sigma^w \mathbf{Y}$, so it only remains to show the sincerity part of the definition.

Suppose, for a contradiction, that $\{H(R) \mid R \in \mathbf{R}\}$ is not a sincere set. This means that there exists an indecomposable projective module P such that $\mathrm{Hom}_{\mathbf{D}}(P, \Sigma^{-i}R) = 0$ for each $R \in \mathbf{R}$ and $0 \leq i \leq w-1$, where $\mathbf{D} = \mathcal{D}^b(\mathbf{k}Q)$. Now, since $P \in \mathcal{F}_{-w}$ and $\mathbf{R} \subseteq \mathcal{F}_{-w}$, by Lemma 1.3 for $0 \leq i \leq w-1$, we have

$$\begin{aligned} \mathrm{Hom}_{\mathcal{C}_{-w}}(P, \Sigma^{-i}R) &\simeq \mathrm{Hom}_{\mathbf{D}}(P, \Sigma^{-i}R) \oplus D \mathrm{Hom}_{\mathbf{D}}(R, \Sigma^{i-w}P) \\ &\simeq D \mathrm{Hom}_{\mathbf{D}}(R, \Sigma^{i-w}P). \end{aligned}$$

Since $i-w < 0$ for each i , $\mathbf{R} \subseteq \mathcal{F}_{-w}$ and $P \in \mathbf{H}$, the hereditary property means that $\mathrm{Hom}_{\mathbf{D}}(R, \Sigma^{i-w}P) = 0$. Hence, $\mathrm{Hom}_{\mathcal{C}_{-w}}(P, \Sigma^{-i}R) = 0$ for each $R \in \mathbf{R}$ and each $0 \leq i \leq w-1$.

Now each $T \in \mathbf{T}$ can be written as $T = \Sigma^w T'$ for some $T' \in \mathbf{H}$. Again, using Lemma 1.3, for $0 \leq i \leq w-1$, we have

$$\begin{aligned} \mathrm{Hom}_{\mathcal{C}_{-w}}(P, \Sigma^{-i}T) &\simeq \mathrm{Hom}_{\mathbf{D}}(P, \Sigma^{-i}T) \oplus D \mathrm{Hom}_{\mathbf{D}}(T, \Sigma^{i-w}P) \\ &\simeq \mathrm{Hom}_{\mathbf{D}}(P, \Sigma^{w-i}T') \oplus D \mathrm{Hom}_{\mathbf{D}}(T', \Sigma^{i-2w}P). \end{aligned}$$

Since $w-i \geq 1$ and P is projective, we have $\mathrm{Hom}_{\mathbf{D}}(P, \Sigma^{w-i}T') = 0$. Since $i-2w < 0$, and \mathbf{H} is hereditary, we have $\mathrm{Hom}_{\mathbf{D}}(T', \Sigma^{i-2w}P) = 0$. It follows that $\mathrm{Hom}_{\mathcal{C}_{-w}}(P, \Sigma^{-i}T) = 0$ for each $0 \leq i \leq w-1$ and $T \in \mathbf{T}$. Since $0 \neq P \in \mathcal{C}_{-w}$, this contradicts the fact that \mathbf{S} is a w -Riedtmann configuration. Hence, $\{H(R) \mid R \in \mathbf{R}\}$ must be a sincere set, as required.

Finally, to see the restriction, suppose further that \mathbf{S} is a w -simple-minded system in \mathcal{C}_{-w} . Then $\langle \mathbf{R} \rangle_{\mathcal{C}_{-w}}$ is functorially finite in $\langle \mathbf{S} \rangle_{\mathcal{C}_{-w}}$ by Lemma 3.2(3). Hence, by Proposition 3.4 and transitivity of functorial finiteness, $\langle \mathbf{R} \rangle_{\mathcal{C}_{-w}}$ is functorially finite in \mathcal{C}_{-w} . By Lemmas 4.8 and 4.9, we see that $\langle \mathbf{R} \rangle_{\mathbf{D}}$ is functorially finite in \mathbf{D} . By Theorem 4.1 and Lemma 5.1, the objects of \mathbf{S} and hence \mathbf{R} can be ordered into an exceptional sequence, making \mathbf{R} exceptionally finite, as required. \square

Putting together Proposition 5.3 with Theorem 4.1 we obtain the following corollary.

Corollary 5.4. *Let Q be an acyclic quiver and $w \geq 1$ an integer. Then there is a well-defined map*

$$\Theta: \left\{ \begin{array}{l} \text{simple-minded collections of} \\ \mathcal{D}^b(\mathbf{k}Q) \text{ contained in } \mathcal{F}_{-w} \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} \text{exceptionally finite, } w\text{-sincere} \\ \infty\text{-orthogonal collections in } \mathcal{D}^b(\mathbf{k}Q) \end{array} \right\}.$$

$$\mathbf{S} \longmapsto \mathbf{S} \cap \mathbf{X} \cap \Sigma^w \mathbf{Y}$$

We now aim to show that the map Θ defined in Corollary 5.4 is a bijection. In order to establish this we need a special case of Jin's reduction of simple-minded collections [33, Theorem 3.1].

5.3. Reduction of simple-minded collections. In this section we present an explicit specialisation of [33, Theorem 3.1]. The following statement is related to [33, Theorem 3.1] in a way analogous to that Aihara-Iyama's silting reduction theorem [1, Theorem 2.37] is related to Iyama-Yang's silting reduction theorem [31, Theorem 3.7]. We include a proof for the convenience of the reader.

Proposition 5.5 (Specialisation of [33, Theorem 3.1]). *Let \mathcal{D} be a Hom-finite, Krull-Schmidt, \mathbf{k} -linear triangulated category. Suppose \mathcal{T} is an ∞ -orthogonal collection such that $\mathbf{thick}_{\mathcal{D}}(\mathcal{T})$ is functorially finite in \mathcal{D} . Then there is a bijection*

$$\Phi: \left\{ \begin{array}{l} \text{simple-minded collections} \\ \text{in } \mathcal{D} \text{ containing } \mathcal{T} \end{array} \right\} \xrightarrow{1-1} \left\{ \begin{array}{l} \text{simple-minded collections} \\ \text{in } (\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp} \end{array} \right\}.$$

In Appendix A we include another proof which does not rely on the functorial finiteness of the subcategory generated by \mathcal{T} or use Verdier localisation.

Proof. The following argument is analogous to [1, Theorem 2.37].

Let \mathcal{S} be a simple-minded collection in \mathcal{D} containing \mathcal{T} . For each $s \in \mathcal{S}$, consider the truncation triangle coming from the stable t-structure $(\mathbf{thick}_{\mathcal{D}}(\mathcal{T}), (\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp})$,

$$(5) \quad t_s \rightarrow s \rightarrow Ls \rightarrow \Sigma t_s,$$

where $L: \mathcal{D} \rightarrow (\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}$ is the left adjoint to the inclusion $(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp} \hookrightarrow \mathcal{D}$. Define $\Phi(\mathcal{S}) := \mathcal{R} = \{Ls \mid s \in \mathcal{S}\}$. Note that $Ls \neq 0$ if and only if $s \in \mathcal{S} \setminus \mathcal{T}$. By truncating t_s with respect to the bounded t-structure $(\mathbf{susp} \mathcal{T}, \mathbf{cosusp} \Sigma^{-1}\mathcal{T})$ in $\mathbf{thick}_{\mathcal{D}}(\mathcal{T})$, observe that the left $(\mathbf{cosusp} \Sigma^{-1}\mathcal{T})$ -approximation of t_s in $\mathbf{thick}_{\mathcal{D}}(\mathcal{T})$ also gives rise to a right $(\mathbf{thick}_{\mathcal{D}}(\mathcal{T}))$ -approximation of s , which shows that $t_s \in \mathbf{cosusp} \Sigma^{-1}\mathcal{T}$. Take $s_1, s_2 \in \mathcal{S}$ and consider the corresponding truncation triangle $t_i \rightarrow s_i \rightarrow Ls_i \rightarrow \Sigma t_i$ for $i = 1, 2$. Applying the functor $\mathbf{Hom}_{\mathcal{D}}(-, Ls_2)$ to the truncation triangle for s_1 and the functor $\mathbf{Hom}_{\mathcal{D}}(s_1, -)$ to the truncation triangle for s_2 shows that \mathcal{R} is an ∞ -orthogonal collection in $(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}$; here it is important to use the fact that $t_i \in \mathbf{cosusp} \Sigma^{-1}\mathcal{T}$.

Suppose $d \in (\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}$. Since, \mathcal{S} is a simple-minded collection in \mathcal{D} there are integers $n \geq m$ such that $d \in \Sigma^n \langle \mathcal{S} \rangle_{\mathcal{D}} * \cdots * \Sigma^m \langle \mathcal{S} \rangle_{\mathcal{D}}$. Applying the functor L to the corresponding tower for d shows that $d \in \Sigma^n \langle \mathcal{R} \rangle_{(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}} * \cdots * \Sigma^m \langle \mathcal{R} \rangle_{(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}}$. Hence $(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp} = \mathbf{thick}_{(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}}(\mathcal{R})$ and \mathcal{R} is a simple-minded collection in $(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}$. This shows that the map Φ is well defined.

We now construct a map

$$\Psi: \left\{ \begin{array}{l} \text{simple-minded collections} \\ \text{in } (\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp} \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} \text{simple-minded collections} \\ \text{in } \mathcal{D} \text{ containing } \mathcal{T} \end{array} \right\}.$$

First note that $\mathbf{cosusp} \mathcal{T}$ is covariantly finite in \mathcal{D} since $\mathbf{cosusp} \mathcal{T}$ is covariantly finite in $\mathbf{thick}_{\mathcal{D}}(\mathcal{T})$, which is in turn functorially finite in \mathcal{D} . Hence, there is an (unbounded) t-structure $({}^{\perp}(\Sigma^{\leq 0}\mathcal{T}), \mathbf{cosusp} \mathcal{T})$ in \mathcal{D} . Let \mathcal{R} be a simple-minded collection in $(\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp}$. For each $r \in \mathcal{R}$ take the truncation triangle with respect to the t-structure $({}^{\perp}(\Sigma^{\leq 0}\mathcal{T}), \mathbf{cosusp} \mathcal{T})$,

$$(6) \quad \Sigma^{-1}t_r \rightarrow s_r \rightarrow r \rightarrow t_r,$$

and set $\Psi(\mathcal{R}) = \mathcal{T} \cup \{s_r \mid r \in \mathcal{R}\}$. In a manner analogous to the argument above, one can show that $\Psi(\mathcal{R})$ is an ∞ -orthogonal collection in \mathcal{D} . For generation, take $d \in \mathcal{D}$ and truncate with respect to the stable t-structure $(\mathbf{thick}_{\mathcal{D}}(\mathcal{T}), (\Sigma^{\mathbb{Z}}\mathcal{T})^{\perp})$,

$$t_d \rightarrow d \rightarrow Ld \rightarrow \Sigma t_d,$$

and observe that $Ld \in \mathbf{thick}_{(\Sigma^{\mathbb{Z}}\mathbb{T})^\perp}(\mathbb{R}) = \mathbf{thick}_{\mathbb{D}}(\mathbb{R})$. It follows immediately that $d \in \mathbf{thick}_{\mathbb{D}}(\Psi(\mathbb{R}))$, whence $\Phi(\mathbb{R})$ is a simple-minded collection in \mathbb{D} , showing that Ψ is well defined.

Finally, to see that Φ and Ψ are mutually inverse, applying $\mathbf{Hom}_{\mathbb{D}}(-, t)$ for $t \in \mathbf{cosusp} \mathbb{T}$ to (5) reveals that the morphism $Ls \rightarrow \Sigma t$ is a left $(\mathbf{cosusp} \mathbb{T})$ -approximation of Ls , and applying $\mathbf{Hom}_{\mathbb{D}}(t, -)$ for $t \in \mathbf{thick}_{\mathbb{D}}(\mathbb{T})$ to (6) reveals that $\Sigma^{-1}t_r \rightarrow s_r$ is a right $(\mathbf{thick}_{\mathbb{D}}(\mathbb{T}))$ -approximation of s_r . \square

5.4. Bijectivity of Θ . To establish the bijectivity of the map occurring in Corollary 5.4 we require a notion of perpendicular category for abelian categories which is compatible with the one for derived categories. We recall the following from [24, 46].

Definition 5.6. Let \mathbb{H} be an abelian category and \mathbb{E} a collection of objects in \mathbb{H} . We define the *right perpendicular category* of \mathbb{E} by

$$\mathbb{E}^{\perp_{\mathbb{H}}} := \{X \in \mathbb{H} \mid \mathbf{Hom}_{\mathbb{H}}(\mathbb{E}, X) = 0 = \mathbf{Ext}_{\mathbb{H}}^1(\mathbb{E}, X)\}.$$

There is an analogous definition of *left perpendicular category*, ${}^{\perp_{\mathbb{H}}}\mathbb{E}$.

Lemma 5.7. Let Q be an acyclic quiver with n vertices, $\mathbb{H} = \mathbf{mod}(\mathbf{k}Q)$ and $\mathbb{E} = \{E_1, \dots, E_k\}$ be a collection of exceptional $\mathbf{k}Q$ -modules for $k \leq n$. Then $(\Sigma^{\mathbb{Z}}\mathbb{E})^\perp$ is equivalent to $\mathbf{D}^b(\mathbf{k}Q')$, where Q' is an acyclic quiver with $n - k$ vertices such that $\mathbb{E}^{\perp_{\mathbb{H}}} \simeq \mathbf{mod}(\mathbf{k}Q')$. Moreover, $\mathbb{H} \cap (\Sigma^{\mathbb{Z}}\mathbb{E})^\perp = \mathbb{E}^{\perp_{\mathbb{H}}}$.

Proof. Let $\mathbb{H} = \mathbf{mod}(\mathbf{k}Q)$ and $\mathbb{D} = \mathbf{D}^b(\mathbf{k}Q)$. For $X \in \mathbb{H}$, we have $\mathbf{Hom}_{\mathbb{D}}(\Sigma^i \mathbb{E}, X) = 0$ for all $i \in \mathbb{Z} \setminus \{-1, 0\}$ since $\mathbb{E} \subseteq \mathbb{H}$ and \mathbb{H} is hereditary. Therefore, for $X \in \mathbb{H}$ we have $X \in (\Sigma^{\mathbb{Z}}\mathbb{E})^\perp$ if and only if $X \in \mathbb{E}^{\perp_{\mathbb{H}}}$. In particular, it follows that $\Sigma^i \mathbb{H} \cap (\Sigma^{\mathbb{Z}}\mathbb{E})^\perp = \Sigma^i \mathbb{E}^{\perp_{\mathbb{H}}}$ for each $i \in \mathbb{Z}$. Now, by [46, Theorem 2.5], there is an equivalence of (abelian) categories $\mathbb{E}^{\perp_{\mathbb{H}}} \rightarrow \mathbf{mod}(\mathbf{k}Q')$. Since \mathbb{H} is hereditary, each object in \mathbb{D} decomposes into a direct sum of its cohomology, and thus this equivalence induces an equivalence of triangulated categories $(\Sigma^{\mathbb{Z}}\mathbb{E})^\perp \rightarrow \mathbf{D}^b(\mathbf{k}Q')$. \square

Theorem 5.8. Let Q be an acyclic quiver with n vertices and $w \geq 1$ an integer. Then there is a bijection

$$\Theta: \left\{ \begin{array}{l} \text{simple-minded collections of} \\ \mathbf{D}^b(\mathbf{k}Q) \text{ contained in } \mathcal{F}_{-w} \end{array} \right\} \xrightarrow{1-1} \left\{ \begin{array}{l} \text{exceptionally finite, } w\text{-sincere} \\ \infty\text{-orthogonal collections in } \mathbf{D}^b(\mathbf{k}Q) \end{array} \right\}.$$

$$\mathbb{S} \longmapsto \mathbb{S} \cap \mathbb{X} \cap \Sigma^w \mathbb{Y}$$

Proof. The strategy of the proof is based on [16, §4]. The map Θ is well defined by Corollary 5.4. We first show that Θ is surjective.

Let $\mathbb{T} = \{T_1, \dots, T_k\}$ be an exceptionally finite, w -sincere ∞ -orthogonal collection in $\mathbb{D} = \mathbf{D}^b(\mathbf{k}Q)$. Since the objects of \mathbb{T} can be ordered into an exceptional sequence we have $k \leq n$. To see that Θ is surjective, we need to find a simple-minded collection \mathbb{S} such that $\mathbb{T} \subseteq \mathbb{S}$ and $\mathbb{S} \setminus \mathbb{T} \subseteq \Sigma^{w+1}(\mathbb{H} \setminus \mathbf{inj}(\mathbb{H}))$. Let $\mathbb{E} = H(\mathbb{T})$, where H is the standard cohomology functor; see (4). By [9, Theorem 3.2], the subcategory $\mathbf{thick}_{\mathbb{D}}(\mathbb{T}) = \mathbf{thick}_{\mathbb{D}}(\mathbb{E})$ is functorially finite in \mathbb{D} . Hence, by Proposition 5.5, we are looking for simple-minded collections \mathbb{R} in $(\Sigma^{\mathbb{Z}}\mathbb{T})^\perp = (\Sigma^{\mathbb{Z}}\mathbb{E})^\perp$ such that $\Psi(\mathbb{R}) \setminus \mathbb{T} \subseteq \Sigma^{w+1}(\mathbb{H} \setminus \mathbf{inj}(\mathbb{H}))$, where Ψ is defined in the proof of Proposition 5.5.

Now, by Lemma 5.7, $(\Sigma^{\mathbb{Z}}\mathbb{T})^\perp \simeq \mathbf{D}^b(\mathbf{k}Q')$ for some acyclic quiver Q' with $n - k$ vertices. Let \mathbb{R}' be image of the set of simple $\mathbf{k}Q'$ -modules in $(\Sigma^{\mathbb{Z}}\mathbb{T})^\perp$ under the equivalence, and set $\mathbb{R} = \Sigma^{w+1}\mathbb{R}'$. Again, by Lemma 5.7, $\mathbb{R} \subseteq \Sigma^{w+1}\mathbb{H} \cap (\Sigma^{\mathbb{Z}}\mathbb{T})^\perp$. Since $\mathbb{T} \subseteq \mathbb{X} \cap \Sigma^w \mathbb{Y}$, by

the hereditary property we have that the object T_R in (6) is zero for each $R \in \mathbf{R}$, whence $S_R = E$ and $\Psi(\mathbf{R}) \setminus \mathbf{T} = \mathbf{R}$. Finally, since \mathbf{T} is w -sincere, for each injective $\mathbf{k}Q$ module I we have $\mathrm{Hom}_{\mathbf{D}}(\mathbf{T}, \Sigma^{w+1}I) \neq 0$, whence for each $R \in \mathbf{R}$ we have $R \notin \Sigma^{w+1}(\mathrm{inj}(\mathbf{k}Q))$. It follows that Θ is surjective.

To see that Θ is injective, suppose \mathbf{S} is a simple-minded collection of \mathbf{D} contained in \mathcal{F}_{-w} such that $\Theta(\mathbf{S}) = \mathbf{T}$. Let $\mathbf{P} = \mathbf{S} \setminus \mathbf{T}$ and note that $\mathbf{P} \subseteq \Sigma^{w+1}\mathbf{H} \cap (\Sigma^{\mathbb{Z}}\mathbf{T})^{\perp}$. It follows, by the hereditary property, that for each $P \in \mathbf{P}$ the object T_P in the truncation triangle (5) must be zero, whence $\Phi(\mathbf{P}) = \mathbf{P}$. In particular, by Proposition 5.5 and Lemma 5.7, \mathbf{P} is a simple-minded collection of $(\Sigma^{\mathbb{Z}}\mathbf{T})^{\perp}$ such that $\mathbf{P} \subseteq \Sigma^{w+1}\mathbf{E}^{\perp_{\mathbf{H}}}$, i.e. is concentrated in one degree with respect to the standard heart $\mathbf{E}^{\perp_{\mathbf{H}}} \simeq \mathrm{mod}(\mathbf{k}Q')$ in $(\Sigma^{\mathbb{Z}}\mathbf{T})^{\perp} \simeq \mathbf{D}^b(\mathbf{k}Q')$. Therefore, the objects of \mathbf{P} can be ordered into an exceptional sequence by Lemma 5.1. It now follows by [43, Theorem 3], which asserts that the unique orthogonal complete exceptional sequence in $\mathrm{mod}(\mathbf{k}Q')$ is that consisting of the simple $\mathbf{k}Q'$ -modules. Hence, $\mathbf{P} = \mathbf{R}$ and Θ is injective. \square

6. NONCROSSING PARTITIONS

Let Q be an acyclic quiver. In this section, we establish a bijection between w -simple-minded systems in $\mathbf{C}_{-w}(\mathbf{k}Q)$ and positive w -noncrossing partitions of the corresponding Weyl group W_Q . This bijection generalises the existing bijection in the case that Q is Dynkin, see [15, 16, 30]. The strategy follows that of [16]. We start with a brief review of the Weyl group of an acyclic quiver and noncrossing partitions. The main reference for the Weyl group is [43], while for noncrossing partitions we refer to [2].

6.1. Weyl group and noncrossing partitions. Consider the symmetric bilinear form on the Grothendieck group $K_0(\mathbf{k}Q)$ defined by

$([X], [Y]) := \dim \mathrm{Hom}_{\mathbf{k}Q}(X, Y) + \dim \mathrm{Hom}_{\mathbf{k}Q}(Y, X) - \dim \mathrm{Ext}_{\mathbf{k}Q}^1(X, Y) - \dim \mathrm{Ext}_{\mathbf{k}Q}^1(Y, X)$,
for $X, Y \in \mathrm{mod}(\mathbf{k}Q)$. Given $X \in K_0(\mathbf{k}Q)$ with $(X, X) \in \{-2, -1, 1, 2\}$, the *reflection* t_X along X is the isometry

$$t_X : K_0(\mathbf{k}Q) \longrightarrow K_0(\mathbf{k}Q), \quad t_X(Y) = Y - \frac{2(Y, X)}{(X, X)}X.$$

Let S_1, \dots, S_n be the simple $\mathbf{k}Q$ -modules and note that $([S_i], [S_i]) = 2$. The *Weyl group* W_Q associated to the quiver Q is the group of isometries generated by $R := \{t_{[S_1]}, \dots, t_{[S_n]}\}$. The set R is called the set of *simple reflections* of W_Q , and the set of all reflections in W_Q is denoted by T .

The *absolute length* of $w \in W_Q$, denoted by $\ell_T(w)$, is the minimum length of w written as a product of reflections. We call a minimum length expression for w written as a product of reflections a *T -reduced expression* of w ; we denote the set of all such expressions by $\mathrm{red}_T(w)$.

A *parabolic subgroup* of W_Q is a subgroup of W_Q generated by a subset of R . The following result, which holds for any Coxeter group W of finite rank, will be useful later.

Theorem 6.1 ([7, Theorem 1.4]). *Let W' be a parabolic subgroup of W . For each $u \in W'$, we have $\mathrm{red}_T(u) = \mathrm{red}_{T'}(u)$, where $T' = W' \cap T$ is the set of reflections in W' .*

Recall, e.g. from [29, §3.1], that a *Coxeter element* of W_Q is the product of all the simple reflections in some order; in [2], a Coxeter element is called a *standard Coxeter element*.

From now on, we will fix a Coxeter element $c \in W_Q$ such that the ordering of the product of simple reflections giving rise to c corresponds to an ordering of the simple $\mathbf{k}Q$ -modules into an exceptional sequence. Note that $\ell_T(c) = n$. We can now define (positive) w -noncrossing partitions for $w \geq 1$; see [2].

Definition 6.2. Let $w \geq 1$, $\mathbf{u} = (u_1, \dots, u_{w+1})$ be a $(w+1)$ -tuple of elements of W_Q and c be the Coxeter element fixed above. The tuple \mathbf{u} is said to be

- (i) a w -noncrossing partition if $c = u_1 u_2 \cdots u_{w+1}$ and $n = \ell_T(c) = \ell_T(u_1) + \cdots + \ell_T(u_{w+1})$; and,
- (ii) a positive w -noncrossing partition if it is a w -noncrossing partition such that the product $u_2 \cdots u_{w+1}$ does not lie in any proper parabolic subgroup.

The set of (positive) w -noncrossing partitions of W_Q with respect to c will be denoted by $\text{NC}_w^{(+)}(W_Q)$, with the fixed Coxeter element c implicitly understood.

We end this section with the following useful connection between exceptional sequences and expressions for the Coxeter element.

Theorem 6.3 ([28, Theorem 4.1]). *Let Q be an acyclic quiver. Then the following assertions hold.*

- (1) For any exceptional $\mathbf{k}Q$ -module E , we have $t_{[E]} \in T$.
- (2) If (X_1, \dots, X_n) is a complete exceptional sequence in $\text{mod}(\mathbf{k}Q)$, setting $t_i = t_{[X_i]}$, we have $c = t_1 \cdots t_n$.
- (3) If $c = t_1 \cdots t_n$, with $t_i \in T$, then $t_i = t_{[X_i]}$, for some exceptional $\mathbf{k}Q$ -module X_i , and (X_1, \dots, X_n) is a complete exceptional sequence.

It follows from Theorem 6.3 that the set of (positive) noncrossing partitions does not depend on the ordering of the set of simple $\mathbf{k}Q$ -modules into an exceptional sequence.

6.2. Simple-minded collections and noncrossing partitions. For an acyclic quiver Q and an integer $w \geq 1$, we recall the construction of a bijective map

$$(7) \quad \varphi: \text{NC}_w(W_Q) \xrightarrow{1-1} \left\{ \begin{array}{l} \text{simple-minded collections of } \text{D}^b(\mathbf{k}Q) \\ \text{contained in } \mathbf{X} \cap \Sigma^{w+1}\mathbf{Y} \end{array} \right\}$$

from [15, Theorem 7.3]; see [16] for a similarly constructed map in the case that Q is Dynkin. Let $\mathbf{u} = (u_1, \dots, u_{w+1}) \in \text{NC}_w(W_Q)$. We construct $\varphi(\mathbf{u})$ in two steps.

- We have $c = u_1 \cdots u_{w+1}$ and $n = \ell_T(u_1) + \ell_T(u_2) + \cdots + \ell_T(u_{w+1})$. For each $1 \leq i \leq w+1$, pick a T -reduced expression for u_i . The ordered product of these expressions gives rise to a T -reduced expression for c and by Theorem 6.3(3) we get a complete exceptional sequence $\mathbf{E} = (\mathbf{E}_1, \dots, \mathbf{E}_{w+1})$ in $\text{mod}(\mathbf{k}Q)$, where \mathbf{E}_i is the subsequence of \mathbf{E} corresponding to the T -reduced expression of u_i .
- Let \mathcal{C}_i be the smallest wide (=exact abelian extension-closed) subcategory of $\text{mod}(\mathbf{k}Q)$ containing \mathbf{E}_i . By [21, Lemma 5], \mathcal{C}_i is equivalent to $\text{mod}(\mathbf{k}Q_i)$ for some acyclic quiver Q_i with $\ell_T(u_i)$ vertices. Let \mathcal{S}_i be the set of simples in \mathcal{C}_i . Then we define

$$\varphi(\mathbf{u}) := \bigcup_{i=1}^{w+1} \Sigma^{w+1-i} \mathcal{S}_i.$$

The fact that $\varphi(\mathbf{u})$ is independent of the choice of T -reduced expression follows from [28, Theorem 4.3].

We now come to the main theorem of this section. The case when Q is Dynkin and $w = 1$ was given in [16, Theorem 5.7]. The case when Q is Dynkin and $w \geq 1$ is an integer was established in [15, Theorem 7.4] (see also [30, Theorem 1.1]), via bijections with m -clusters, and using a different description of positive w -noncrossing partitions. Before proceeding, we require a definition.

For a $\mathbf{k}Q$ -module X , the *support of X* is $\text{supp}(X) := \{a \in Q_0 \mid \text{Hom}(P_a, X) \neq 0\}$. Equivalently, $\text{supp}(X) = \{a \in Q_0 \mid S_a \text{ occurs in a composition series for } X\}$. For a set \mathbf{X} of $\mathbf{k}Q$ -modules the *support of \mathbf{X}* is $\text{supp}(\mathbf{X}) = \bigcup_{X \in \mathbf{X}} \text{supp}(X)$.

Theorem 6.4. *Let Q be an acyclic quiver. The map φ define in (7) restricts to a bijection*

$$\varphi: \text{NC}_w^+(W_Q) \xrightarrow{1-1} \left\{ \begin{array}{l} \text{simple-minded collections of } \mathbf{D}^b(\mathbf{k}Q) \\ \text{contained in } \mathcal{F}_{-w} \end{array} \right\}.$$

Proof. Let $\mathbf{u} = (u_1, u_2, \dots, u_{w+1}) \in \text{NC}_w^+(W_Q)$. Then, by the construction of the map φ in (7), $\varphi(\mathbf{u}) = \bigcup_{i=1}^{w+1} \Sigma^{w+1-i} \mathbf{S}_i$ is a simple-minded collection of $\mathbf{D}^b(\mathbf{k}Q)$ lying in $\mathbf{X} \cap \Sigma^{w+1} \mathbf{Y}$. To see that the restriction is well defined, by Theorem 5.8, it suffices to check that $\bigcup_{i=2}^{w+1} \Sigma^{w+1-i} \mathbf{S}_i \subseteq \mathbf{X} \cap \Sigma^w \mathbf{Y}$ is an exceptionally finite, w -sincere ∞ -orthogonal collection in $\mathbf{D}^b(\mathbf{k}Q)$. This set is clearly ∞ -orthogonal. Functorial finiteness and exceptionality follow immediately from Lemma 3.2(3) and Lemma 3.6, and Lemma 5.1, respectively. Only w -sincerity remains to be checked.

Suppose for a contradiction that $\bigcup_{i=2}^{w+1} \Sigma^{w+1-i} \mathbf{S}_i \subseteq \mathbf{X} \cap \Sigma^w \mathbf{Y}$ is not w -sincere. We claim that if E is an exceptional $\mathbf{k}Q$ -module, then $t_{[E]}$ lies in the parabolic subgroup generated by $\{t_{[S_a]} \in R \mid a \in \text{supp}(E)\}$. Indeed, consider the wide subcategory \mathcal{C}_E generated by the set of simple $\mathbf{k}Q$ -modules $\{S_a \mid a \in \text{supp}(E)\}$. The exceptional module E lies in \mathcal{C}_E and it can be extended to a complete exceptional sequence \mathbf{E} in \mathcal{C}_E (see [21, Lemma 1]). By the transitivity of the action of the braid group on the set of complete exceptional sequences (see [21] again), \mathbf{E} is obtained from the set $\{S_a \mid a \in \text{supp}(\mathbf{E})\}$ ordered into a complete exceptional sequence in \mathcal{C}_E , via a sequence of mutations corresponding to the braid group action. The claim then follows from [28, §2.2]. As a consequence, we have that $u_2 \cdots u_{w+1}$ lies in the parabolic subgroup generated by $\{t_{[S_a]} \in R \mid a \in \text{supp}(\bigcup_{i=2}^{w+1} \mathbf{E}_i)\}$. This subgroup is a proper parabolic subgroup since $\bigcup_{i=2}^{w+1} \mathbf{E}_i$ is not sincere, as $\text{supp}(\mathbf{S}_i) = \text{supp}(\mathcal{C}_i) = \text{supp}(\mathbf{E}_i)$, for each i . This contradicts the fact that \mathbf{u} is a positive w -noncrossing partition. Therefore, $\bigcup_{i=2}^{w+1} \Sigma^{w+1-i} \mathbf{S}_i \subseteq \mathbf{X} \cap \Sigma^w \mathbf{Y}$ is indeed w -sincere.

Since φ is the restriction of a bijection, it is clearly injective. It remains to check that φ is surjective. Let \mathbf{S} be a simple-minded collection contained in \mathcal{F}_{-w} . By Lemma 5.1, we can order the elements in \mathbf{S} into a (complete) exceptional sequence \mathbf{E} in which the cohomological degrees are weakly decreasing. For each $1 \leq i \leq w+1$, let \mathbf{E}_i the subsequence consisting of the elements of cohomological degree $i-1$, and \mathcal{C}_i be the smallest wide subcategory of $\text{mod}(\mathbf{k}Q)$ containing $H(\mathbf{E}_i)$, where $H: \mathbf{D}^b(\mathbf{k}Q) \rightarrow \text{mod}(\mathbf{k}Q)$ is the standard cohomology functor; see (4). Order the simple objects of \mathcal{C}_i into an exceptional sequence (cf. [43, Theorem 3]), and let u_i be the product of the corresponding reflections respecting the order of the exceptional sequence, which gives a T -reduced expression for u_i . Then $\mathbf{u} = (u_{w+1}, \dots, u_1)$ is a w -noncrossing partition and $\varphi(\mathbf{u}) = \mathbf{S}$.

Finally, we must check that \mathbf{u} is positive. Suppose \mathbf{u} is not positive. Then $u_w \cdots u_1$ lies in a proper parabolic subgroup W_J generated by $J \subsetneq R$. For each $1 \leq i \leq w$, each reflection appearing in the T -reduced expression of u_i above lies in W_J by Theorem 6.1. This means that for $1 \leq i \leq w$ the simple objects of \mathcal{C}_i do not have support at the

$$\begin{array}{ccccc}
& & s_f & \xrightarrow{\sigma} & s_x \\
& & \downarrow \alpha_f & & \downarrow \alpha_x \\
x & \xrightarrow{f} & y & \xrightarrow{g_1} & c_f & \xrightarrow{h_1} & \Sigma x \\
& & \downarrow \beta_f & & \downarrow \beta_x \\
& & z_f & \xrightarrow{h} & x\langle 1 \rangle \\
& & \downarrow \gamma_f & & \downarrow \gamma_x \\
& & \Sigma s_f & \xrightarrow{\Sigma \sigma} & \Sigma s_x
\end{array}$$

FIGURE 1. Triangles in \mathcal{Z} .

vertices of Q corresponding to the simple reflections at $R \setminus J$. Since the support of $H(\mathbf{E}_i)$ coincides with that of the simple objects in \mathcal{C}_i , it follows that $\mathbf{S} \cap \mathbf{X} \cap \Sigma^w \mathbf{Y}$ is not w -sincere, contradicting Theorem 5.8. Hence, \mathbf{u} must be positive and φ is surjective. \square

APPENDIX A. REDUCTION OF SIMPLE-MINDED COLLECTIONS REVISITED

In this appendix, we provide an alternative proof of [33, Theorem 3.1], which avoids using a Verdier localisation and is similar to the analogous result for simple-minded systems in [20, Section 6]. This proof was obtained by Alexandra Zvonareva and the first two authors before the preprint [33] appeared. We think it is still useful as an alternative approach.

Throughout, \mathbf{D} will be a Hom-finite, Krull-Schmidt, \mathbf{k} -linear triangulated category with shift functor $\Sigma: \mathbf{D} \rightarrow \mathbf{D}$. We will impose the following setup.

Setup A.1. Let \mathbf{S} be an ∞ -orthogonal collection of objects in \mathbf{D} and \mathcal{Z} a subcategory of \mathbf{D} satisfying the following conditions:

- (1) $\langle \mathbf{S} \rangle$ is covariantly finite in ${}^\perp(\Sigma^{<0}\mathbf{S})$ and contravariantly finite in $(\Sigma^{>0}\mathbf{S})^\perp$.
- (2) for $d \in \mathbf{D}$, we have $\mathrm{Hom}_{\mathbf{D}}(d, \Sigma^{\leq 0}\mathbf{S}) = 0 = \mathrm{Hom}_{\mathbf{D}}(\Sigma^{\geq 0}\mathbf{S}, d) = 0$; and,
- (3) $\mathcal{Z} := {}^\perp(\Sigma^{\leq 0}\mathbf{S}) \cap (\Sigma^{\geq 0}\mathbf{S})^\perp$.

In fact, if $\mathbf{D} = \mathbf{D}^b(A)$ for a finite-dimensional algebra A , the stronger condition that $\langle \mathbf{S} \rangle$ is functorially finite in $\mathbf{D}^b(A)$ holds by Lemmas 3.2(3) and 3.6.

We recall the following construction from [20, Section 4] (see also [33, 39]):

- For an object $z \in \mathcal{Z}$, a functor $\langle 1 \rangle: \mathcal{Z} \rightarrow \mathcal{Z}$ is defined on objects by taking the cone of a minimal right $\langle \mathbf{S} \rangle$ -approximation: $s_z \rightarrow \Sigma z \rightarrow z\langle 1 \rangle \rightarrow \Sigma s_z$. The functor $\langle 1 \rangle$ is defined on morphisms in the obvious way, and its quasi-inverse $\langle -1 \rangle$ by the dual construction; see [20, Lemma 3.6].
- For a morphism $f: x \rightarrow y$ in \mathcal{Z} , consider the triangle $x \xrightarrow{f} y \xrightarrow{g_1} c_f \xrightarrow{h_1} \Sigma x$ in \mathbf{D} together with the minimal right $\langle \mathbf{S} \rangle$ -approximation triangles of c_f and Σx in Figure 1. Note that applying $\mathrm{Hom}_{\mathbf{D}}(\Sigma^{>0}\mathbf{S}, -)$ to this triangle shows that both $c_f, \Sigma x \in (\Sigma^{>0}\mathbf{S})^\perp$ so that the required approximations exist.

Theorem A.2 ([33, Proposition 3.6]). *The category $(\mathcal{Z}, \langle 1 \rangle)$ admits a triangulated structure with standard triangles given by diagrams of the form $x \xrightarrow{f} y \rightarrow z_f \rightarrow x\langle 1 \rangle$.*

Proof. This can be proved in several ways. On one hand, observe that \mathbf{S} and \mathcal{Z} given in Setup A.1 satisfy [20, Lemma 6.3] so that [20, Theorem 4.1] can be applied. Otherwise,

one can proceed via Jin's construction in [33, Proposition 3.6], or Nakaoka's construction in [39, Theorem 4.15]. \square

Because there are two triangulated structures under consideration, we differentiate them as in [20, Section 6] with the following notation. Let \mathbf{X} and \mathbf{Y} be subcategories of \mathbf{Z} . We define

$$\mathbf{X} \star \mathbf{Y} := \{z \in \mathbf{Z} \mid \text{there exists a triangle } x \rightarrow z \rightarrow y \rightarrow x\langle 1 \rangle \text{ with } x \in \mathbf{X} \text{ and } y \in \mathbf{Y}\}.$$

We denote the extension closure of \mathbf{X} with respect to the triangulated structure in \mathbf{Z} by $\langle \mathbf{X} \rangle$. The usual notation $\mathbf{X} * \mathbf{Y}$ and $\langle \mathbf{X} \rangle$ keep their usual meanings in \mathbf{D} .

Before giving our alternative proof of [33, Theorem 3.1], we collect some preliminary statements.

Lemma A.3. *Let \mathbf{S} and \mathbf{Z} be as in Setup A.1. Suppose one of the two conditions holds:*

- (a) $\mathbf{S} \subseteq \mathbf{T}$ and \mathbf{T} is a simple-minded collection in \mathbf{D} ; in which case set $\mathbf{R} = \mathbf{T} \setminus \mathbf{S}$; or
- (b) $\mathbf{T} = \mathbf{S} \cup \mathbf{R}$, where \mathbf{R} is a simple-minded collection in \mathbf{Z} .

Then the following hold:

- (1) For $i \geq j$, $\{\mathbf{R}\langle i \rangle\} \star \{\mathbf{R}\langle i-1 \rangle\} \star \cdots \star \{\mathbf{R}\langle j \rangle\} \subseteq \langle \Sigma^i \mathbf{T} \rangle * \langle \Sigma^{i-1} \mathbf{T} \rangle * \cdots * \langle \Sigma^j \mathbf{T} \rangle$.
- (2) $\text{susp}_{\mathbf{D}} \mathbf{T} \cap \mathbf{Z} = \text{susp}_{\mathbf{Z}} \mathbf{R}$.
- (3) $\text{cosusp}_{\mathbf{D}} \mathbf{T} \cap \mathbf{Z} = \text{cosusp}_{\mathbf{Z}} \mathbf{R}$.
- (4) $\text{susp}_{\mathbf{D}} \mathbf{S}$ is contravariantly finite in \mathbf{D} .
- (5) $\text{cosusp}_{\mathbf{D}} \mathbf{S}$ is covariantly finite in \mathbf{D} .

Proof. Statement (1) can be argued as in Claims A and B in the proof of [20, Theorem 6.6]; see also [33, Lemma 3.4]. Statements (2) and (3) can be argued using [20, Lemma 6.5] as the start of an induction. Statements (4) and (5) follow immediately from conditions (1) and (2) in Setup A.1 using a standard argument; cf. the proof of Proposition 4.2. \square

Theorem A.4 ([33, Theorem 3.1]). *Let \mathbf{S} and \mathbf{Z} be as in Setup A.1. Then there is a bijection*

$$\begin{aligned} \{ \text{simple-minded collections in } \mathbf{D} \text{ containing } \mathbf{S} \} &\xrightarrow{1-1} \{ \text{simple-minded collections in } \mathbf{Z} \}. \\ \mathbf{T} &\longmapsto \mathbf{T} \setminus \mathbf{S} \end{aligned}$$

Proof. Let \mathbf{T} be a simple-minded collection in \mathbf{D} containing \mathbf{S} . We want to show that $\mathbf{R} := \mathbf{T} \setminus \mathbf{S}$ is a simple-minded collection in \mathbf{Z} . We will use the characterisation of simple-minded collections in Proposition 3.5.

Firstly, observe that \mathbf{R} is an ∞ -orthogonal collection in \mathbf{Z} , for instance, by using the dimension shifting argument in the proof of [20, Theorem 6.6].

The next step is to show that ${}^{\perp}(\mathbf{R}\langle < 0 \rangle) \cap (\mathbf{R}\langle \geq 0 \rangle)^{\perp} = 0$. Suppose $z \in \mathbf{Z}$ is such that $z \in {}^{\perp}(\mathbf{R}\langle < 0 \rangle) \cap (\mathbf{R}\langle \geq 0 \rangle)^{\perp}$. Since $\text{Hom}_{\mathbf{D}}(z\langle > 0 \rangle, \mathbf{R}) \simeq \text{Hom}_{\mathbf{D}}(\Sigma^{>0} z, \mathbf{R})$ and $\text{Hom}_{\mathbf{D}}(\mathbf{R}, z\langle \leq 0 \rangle) \simeq \text{Hom}_{\mathbf{D}}(\mathbf{R}, \Sigma^{\leq 0} z)$ by the dimension-shifting argument above, it follows that $z \in {}^{\perp}(\Sigma^{<0} \mathbf{R}) \cap (\Sigma^{\geq 0} \mathbf{R})^{\perp}$. Since $z \in \mathbf{Z}$, it follows that $z \in {}^{\perp}(\Sigma^{<0} \mathbf{T}) \cap (\Sigma^{\geq 0} \mathbf{T})^{\perp}$, and so $z = 0$, as \mathbf{T} is ∞ -Riedtmann in \mathbf{D} by Proposition 3.5.

Finally, we show that $\text{cosusp}_{\mathbf{Z}} \mathbf{R}$ is covariantly finite in \mathbf{Z} . By Lemma A.3(1), we have $\text{cosusp}_{\mathbf{Z}} \mathbf{R} \subseteq \text{cosusp}_{\mathbf{D}} \mathbf{T}$. By Lemma A.3(3), we have $(\text{cosusp}_{\mathbf{D}} \mathbf{T}) \cap \mathbf{Z} = \text{cosusp}_{\mathbf{Z}} \mathbf{R}$. Since \mathbf{T} is a simple-minded collection in \mathbf{D} , $\text{cosusp}_{\mathbf{D}} \mathbf{T}$ is covariantly finite in \mathbf{D} . Therefore, for $z \in \mathbf{Z} \subseteq \mathbf{D}$, we can take a decomposition triangle, $x \xrightarrow{f} z \longrightarrow t_z \longrightarrow \Sigma x$, in \mathbf{D}

with $x \in {}^\perp(\text{cosusp}_D \mathbf{T}) \subseteq {}^\perp(\Sigma^{\leq 0} \mathbf{S})$. An inspection of the resulting long exact sequence shows that $x \in (\Sigma^{\geq 0} \mathbf{S})^\perp$, and hence $x \in \mathbf{Z}$. Now taking the cone of a minimal right $\langle \mathbf{S} \rangle$ -approximation $s_f \rightarrow t_z$ of t_z gives a triangle in \mathbf{Z} :

$$x \xrightarrow{f} z \rightarrow z_f \rightarrow x\langle 1 \rangle.$$

We claim that the third term z_f lies in $(\text{cosusp}_D \mathbf{T}) \cap \mathbf{Z}$ and hence in $\text{cosusp}_Z \mathbf{R}$, which makes the triangle above into a decomposition triangle for z in \mathbf{Z} because $x \in {}^\perp(\text{cosusp}_D \mathbf{T}) \subseteq {}^\perp(\text{cosusp}_Z \mathbf{R})$. In particular, showing that $\text{cosusp}_Z \mathbf{R}$ is covariantly finite in \mathbf{Z} . The argument for the contravariant finiteness of $\text{susp}_Z \mathbf{R}$ in \mathbf{Z} is similar. Now applying Proposition 3.5, we conclude that \mathbf{R} is a simple-minded collection in \mathbf{Z} .

Therefore, we need to establish the claim above. Since $t_z \in \text{cosusp}_D \mathbf{T}$, there is a triangle $t' \rightarrow t_z \rightarrow t'' \rightarrow \Sigma t'$ with $t' \in \langle \mathbf{T} \rangle$ and $t'' \in \text{cosusp}_D \Sigma^{-1} \mathbf{T}$. Combining this triangle with the one coming from the minimal right $\langle \mathbf{S} \rangle$ -approximation gives the following commutative diagram by the octahedral axiom:

$$\begin{array}{ccccccc} & & s_f & \xlongequal{\quad} & s_f & & \\ & & g \downarrow & & \downarrow & & \\ \Sigma^{-1} t'' & \longrightarrow & t' & \longrightarrow & t_z & \longrightarrow & t'' \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ \Sigma^{-1} t'' & \longrightarrow & t_1 & \longrightarrow & z_f & \longrightarrow & t'' \\ & & \downarrow & & \downarrow & & \\ & & \Sigma s_f & \xlongequal{\quad} & \Sigma s_f & & \end{array}$$

Examining the long exact sequence obtained by applying $\text{Hom}_D(\mathbf{S}, -)$ to the lower horizontal triangle shows that $g: s_f \rightarrow t'$ is a right $\langle \mathbf{S} \rangle$ -approximation. If it were not minimal, then $s_f \rightarrow t_z$ would also fail to be minimal. Therefore $g: s_f \rightarrow t'$ is a minimal right $\langle \mathbf{S} \rangle$ -approximation and $t_1 \in \langle \mathbf{T} \rangle$ by [20, Theorem 2.11]. Hence, $z_f \in \text{cosusp}_D \mathbf{T}$, as required.

Conversely, suppose \mathbf{R} is a simple-minded collection in \mathbf{Z} . We will show that $\mathbf{T} := \mathbf{S} \cup \mathbf{R}$ is a simple-minded collection in \mathbf{D} . The proof that \mathbf{T} is an ∞ -orthogonal collection is similar to the corresponding argument above in the other direction.

Let d be an object in \mathbf{D} . By Lemma A.3(4) and (5), we have that $\text{susp}_D \mathbf{S}$ is contravariantly finite in \mathbf{D} and $\text{cosusp}_D \mathbf{S}$ is covariantly finite in \mathbf{D} . Decomposing first with respect to the t-structure $(\text{susp}_D \mathbf{S}, (\Sigma^{\geq 0} \mathbf{S})^\perp)$ and then the t-structure $({}^\perp(\Sigma^{\leq 0} \mathbf{S}), \text{cosusp}_D \mathbf{S})$ gives the following triangles,

$$x \rightarrow d \rightarrow y \rightarrow \Sigma x \quad \text{and} \quad z \rightarrow y \rightarrow v \rightarrow \Sigma z,$$

in which $z \in \mathbf{Z} = {}^\perp(\Sigma^{\leq 0} \mathbf{S}) \cap (\Sigma^{\geq 0} \mathbf{S})^\perp$, $v \in \text{cosusp}_D \mathbf{S}$ and $x \in \text{susp}_D \mathbf{S}$. This shows that $\mathbf{D} = (\text{susp}_D \mathbf{S}) * \mathbf{Z} * (\text{cosusp}_D \mathbf{S})$, cf. [33, Proposition 3.3]. Now, Lemma A.3(1), shows that $\mathbf{Z} = \text{thick}_Z(\mathbf{R}) \subseteq \text{thick}_D(\mathbf{T})$, from which it follows that $\mathbf{D} = \text{thick}_D(\mathbf{T})$, and we conclude that \mathbf{T} is a simple-minded collection. \square

REFERENCES

- [1] T. Aihara, O. Iyama, *Silting mutation in triangulated categories*, J. Lond. Math. Soc. (2) **85** (2012), no. 3, 633–668, also [arXiv:1009.3370](https://arxiv.org/abs/1009.3370)
- [2] D. Armstrong, *Generalized noncrossing partitions and combinatorics of Coxeter groups*, Mem. Am. Math. Soc. **202** (2009), no. 949, x+159 pp, also [arXiv:math/0611106](https://arxiv.org/abs/math/0611106).
- [3] S. Asai, *Semibricks*, Int. Math. Res. Not., DOI 10.1093/imrn/rny150 (2018), also [arXiv:1610.05860](https://arxiv.org/abs/1610.05860).

- [4] C. Athanasiadis, T. Brady, J. McCammond, J. Watt, *h-vectors of generalized associahedra and noncrossing partitions*, Int. Math. Res. Not. (2006), Art. ID 69705, 28pp, also [arXiv:math/0602293](#).
- [5] M. Auslander, S. O. Smalø, *Preprojective modules over Artin algebras*, J. Algebra **66** (1980), 61–122.
- [6] H. Bass, “Algebraic K -theory”, W. A. Benjamin, Inc., New York- Amsterdam, 1968.
- [7] B. Baumeister, M. Dyer, C. Stump, P. Wegener, *A note on the transitive Hurwitz action on decompositions of parabolic Coxeter elements*, Proc. Amer. Math. Soc. Ser. B **1** (2014), 149–154, also [arXiv:1402.2500](#).
- [8] A.A. Beilinson, J. Bernstein, P. Deligne, *Faisceaux pervers*, Astérisque **100** (1982).
- [9] A. I. Bondal, *Representations of associative algebras and coherent sheaves*, Math. USSR Izv. **34** (1990), no. 1, 23–42.
- [10] M. V. Bondarko, *On (well generated) weight structures, adjacent t -structures, and related (co)homological functors*, [arXiv:1611.00754](#).
- [11] T. Bridgeland, *Stability conditions on triangulated categories*, Ann. of Math. **166** (2007), 317–345, also [arXiv:math/0212237](#).
- [12] J. Brightbill, *The differential graded stable category of a self-injective algebra*, [arXiv:1811.08992](#).
- [13] J. Brightbill, *Perverse equivalences and dg stable combinatorics*, [arXiv:2001.00193](#).
- [14] A. B. Buan, R. Marsh, M. Reineke, I. Reiten, G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), no. 2, 572–618, also [arXiv:math/0402054](#).
- [15] A. B. Buan, I. Reiten, H. Thomas, *From m -clusters to m -noncrossing partitions via exceptional sequences*, Math. Z. **271** (2012), no. 3-4, 1117–1139, also [arXiv:1007.0928](#).
- [16] R. Coelho Simões, *Hom-configurations and noncrossing partitions*, J. Algebraic Combin. **35** (2012), no. 2, 313–343, also [arXiv:1012.1276](#).
- [17] R. Coelho Simões, *Hom-configurations in triangulated categories generated by spherical objects*, J. Pure and Appl. Algebra **219** (2015), no. 8, 3322–3336, also [arXiv:1312.4769](#).
- [18] R. Coelho Simões, *Mutations of simple-minded systems in Calabi-Yau categories generated by a spherical object*, Forum Math. **29**(5) (2017), 1065–1081, also [arXiv:1512.09321](#).
- [19] R. Coelho Simões, D. Pauksztello, *Torsion pairs in a triangulated category generated by a spherical object*, J. Algebra **448** (2016), 1–47, also [arXiv:1404.4623](#).
- [20] R. Coelho Simões, D. Pauksztello, *Simple-minded systems and reduction for negative Calabi-Yau triangulated categories*, Trans. Amer. Math. Soc. **373** (2020), no. 4, 2463–2498, also [arXiv:1808.02519](#).
- [21] W. Crawley-Boevey, *Exceptional sequences of representations of quivers*, In: V. Dlab, H. Lenzing, (eds.) Representations of Algebras, Proc. Ottawa 1992. Canadian Math. Soc. Conf. Proc, vol. 14, pp. 117–124. Amer. Math. Soc., Providence (1993).
- [22] A. Dugas, *Torsion pairs and simple-minded systems in triangulated categories*, Appl. Categ. Structures **23** (2015), no. 3, 507–526, also [arXiv:1207.7338](#).
- [23] S. Fomin, A. Zelevinsky, *Cluster algebras I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529, also [arXiv:math/0104151](#).
- [24] W. Geigle, H. Lenzing, *Perpendicular categories with applications to representations and sheaves*, J. Algebra **144** (1991), 273–343.
- [25] D. Happel, “Triangulated Categories in the Representation Theory of Finite Dimensional Algebras”, London Mathematical Society Lecture Note Series **119**, Cambridge University Press (1988).
- [26] T. Holm, P. Jørgensen, D. Yang, *Sparseness of t -structures and negative Calabi-Yau dimension in triangulated categories generated by a spherical object*, Bull. London Math. Soc. **45** (2013), no. 1, 120–130, also [arXiv:1108.2195](#).
- [27] A. Hubery, H. Krause, *A categorification of non-crossing partitions*, J. Eur. Math. Soc. **18** (2016), no. 10, 2273–2313, also [arXiv:1310.1907](#).
- [28] K. Igusa, R. Schiffler, *Exceptional sequences and clusters*, with an appendix written by the authors and H. Thomas, J. Algebra **323** (2010), no. 8, 2183–2202, also [arXiv:0901.2590](#).
- [29] C. Ingalls, H. Thomas, *Noncrossing partitions and representations of quivers*, Compositio Math. **145** (2009), 1533–1562, also [arXiv:math/0612219](#).
- [30] O. Iyama, H. Jin, *Positive Fuss-Catalan numbers and simple-minded systems in negative Calabi-Yau categories*, [arXiv:2002.09952](#).
- [31] O. Iyama, D. Yang, *Silting reduction and Calabi-Yau reduction of triangulated categories*, Trans. Amer. Math. Soc. **370** (2018), no. 11, 7861–7898, also [arXiv:1408.2678](#).
- [32] H. Jin, *Cohen-Macaulay differential graded modules and negative Calabi-Yau configurations*, [arXiv:1812.03737](#).

- [33] H. Jin, *Reductions of triangulated categories and simple-minded collections*, [arXiv:1907.05114](#).
- [34] P. Jørgensen, *Auslander-Reiten triangles in subcategories*, *J. K-Theory* **3** (2009), no. 3, 583–601, also [arXiv:math/0606647](#).
- [35] B. Keller, *On triangulated orbit categories*, *Doc. Math.* **10** (2005), 551–581, also [arXiv:math/0503240](#).
- [36] S. Koenig, Y. Liu, *Simple-minded systems in stable module categories*, *Q. J. Math.* **63**(3) (2012), 653–674, also [arXiv:1009.1427](#).
- [37] S. Koenig, D. Yang, *Silting objects, simple-minded collections, t-structures and co-t-structures for finite-dimensional algebras*, *Doc. Math.* **19** (2014), 403–438, also [arXiv:1203.5657](#).
- [38] C. Kohler, *Thick subcategories of finite algebraic triangulated categories*, [arXiv:1010.0146](#).
- [39] H. Nakaoka, *A simultaneous generalisation of mutation and recollement of cotorsion pairs on a triangulated category*, *Appl. Categ. Structures* **26** (2018), no. 3, 491–544, also [arXiv:1512.02173](#).
- [40] Z. Pogorzaly, *Algebras stably equivalent to self-injective special biserial algebras*, *Comm. Algebra* **22** (1994), no. 4, 1127–1160.
- [41] N. Reading, *Coxeter sortable elements and noncrossing partitions*, *Trans. Amer. Math. Soc.* **359** (2007), no. 12, 5931–5958, also [arXiv:math/0507186](#).
- [42] I. Reiten, M. van den Bergh, *Noetherian hereditary abelian categories satisfying Serre duality*, *J. Amer. Math. Soc.* **15** (2002), no. 2, 295–366, also [arXiv:math/9911242](#).
- [43] C. M. Ringel, *The braid group action on the set of exceptional sequences of a hereditary Artin algebra*, *Abelian Group Theory and Related Topics (Oberwolfach, 1993)*. *Contemp. Math.* **171**, 339–352. Amer. Math. Soc., Providence, RI, 1994.
- [44] A. N. Rudakov, *Exceptional collections, mutations and helices*, in “Helices and vector bundles”, 1–6, *London Math. Soc. Lecture Note Ser.* **148**, Cambridge University Press, Cambridge, 1990.
- [45] M. Saorín, A. Zvonareva, *Lifting of recollements and gluing of partial silting sets*, [arXiv:1809.03243](#).
- [46] A. Schofield, *Semi-invariants of quivers*, *J. London Math. Soc.* **43** (1991), 385–395.
- [47] C. Stump, H. Thomas, N. Williams, *Cataland: Why the Fuss?*, [arXiv:1503.00710](#).

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER, LA1 4YF, UNITED KINGDOM.

E-mail address: r.coelhosimoes@lancaster.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER, LA1 4YF, UNITED KINGDOM.

E-mail address: d.pauksztello@lancaster.ac.uk

INSTITUTT FOR MATEMATIKK OG FYSIKK, STAVANGER UNIVERSITET, STAVANGER, 4021 STAVANGER, NORWAY

E-mail address: david.ploog@uis.no