

Prevalence of ρ -irregularity and related properties

L. Galeati & M. Gubinelli
Institute of Applied Mathematics &
Hausdorff Center for Mathematics
University of Bonn
Germany

Email: {lucio.galeati,gubinelli}@iam.uni-bonn.de

21st April 2023

Abstract

We show that generic Hölder continuous functions are ρ -irregular. The property of ρ -irregularity has been first introduced by Catellier and Gubinelli (Stoc. Proc. Appl. 126, 2016) and plays a key role in the study of well-posedness for some classes of perturbed ODEs and PDEs. Genericity here is understood in the sense of *prevalence*. As a consequence we obtain several results on regularisation by noise “without probability”, i.e. without committing to specific assumptions on the statistical properties of the perturbations. We also establish useful criteria for stochastic processes to be ρ -irregular and study in detail the geometric and analytic properties of ρ -irregular functions.

Contents

1	Introduction	2
2	Preliminaries	6
2.1	Introduction to prevalence	6
2.2	Basic properties of ρ -irregularity and related concepts	8
2.3	Occupation measures and averaging operators	11
2.4	Fractional Brownian motion and local-nondeterminism	15
3	Main results	17
3.1	Statements	17
3.2	Applications to regularisation by noise for ODEs and PDEs	19
4	Criteria for ρ-irregularity of stochastic processes	24
4.1	General criteria	25
4.2	ρ -irregularity for β -(e)SLND Gaussian processes and examples	30
4.3	ρ -irregularity for α -stable processes	35
4.4	ρ -irregularity for β -LND Gaussian processes	36
5	Analytic properties of ρ-irregularity	39
5.1	Fourier dimension and Salem sets	39
5.2	ρ -irregular paths are rough	40
5.3	The general perturbation problem	43
5.4	The additive perturbation problem	47

A Auxiliary results	51
A.1 A primer on Young integration theory	51
A.2 Some useful tools	53

1 Introduction

In recent years there has been a lot of focus in understanding the role of noise in improving well-posedness of ordinary and partial differential equations (ODE/PDEs), or so called *regularisation by noise* phenomena, see [24] for a review. From the modelling point of view, the presence of external perturbations to otherwise autonomous evolutions is a very natural assumption. For the sake of abstraction from any specific origin of such perturbations, one is usually led to consider it *random*, i.e. model the noise as a stochastic process. This common approach introduces into the picture new considerations, some of which do to quite fit the original deterministic formulation:

1. It is not immediate to identify which types of noise are justified in any specific problem.
2. Measurability (or adaptedness) of solutions in the sense of stochastic processes (i.e. seen as random variables on a filtered probability space) must be required; probabilistic notions of uniqueness (like path-wise uniqueness) are obtained, which are not easy to compare to their deterministic counterparts.

Regarding 1., it is natural to require the noise to have statistical properties like Gaussianity or self-similarity, as they are associated to features of universality; as the perturbed problem has dynamical nature, it is also reasonable to impose Markovianity. These considerations make Brownian motion a natural candidate and a large set of theoretical tools is available to analyse the effect of Brownian perturbations to deterministic evolutions. For these reasons, this topic has a long and extensive history, see e.g. [67], [63], [44], [26], [17], [18], [5] and the references therein. More recently, other classes of random perturbations partially satisfying the above requirements have been analysed, like fractional Brownian motion, α -stable processes and more exotic variants, see e.g. [50], [45], [57], [21], [1].

In order to avoid the spurious technical problems of Point 2. one is led to modify the probabilistic setting. For example, in the Brownian setting, Davie and Flandoli introduced the stronger concept of *path-by-path uniqueness*, see [20], [24], [25]. Shaposhnikov obtained further results in this direction, see [60], [61] and the recent [62] with Wresch.

Here instead we follow the approach we initiated in [34] and consider the regularisation by noise problem from the point of view of *generic* perturbations, in particular without reference to any (specific) probabilistic setting.

We will say that a property holds for a generic path, or equivalently for *almost every* path, if it holds for the elements of a *prevalent* set. The notion of prevalence was first introduced by Christensen in [15] in the context of abelian Polish groups and later rediscovered independently by Hunt, Sauer and Yorke in [42] for complete metric vector spaces. It allows to define consistently “full Lebesgue measure sets” in infinite dimensional spaces, even if they don’t admit any σ -additive, translation invariant measure.

Prevalence has been used in different contexts in order to study the properties of generic functions of suitable regularity. For instance, it was proved in [41] that almost every continuous function is nowhere differentiable, while in [29], [30], the multi-fractal nature of generic Sobolev functions was shown. See [52] for a review. Recently, prevalence has also attracted attention in the study of dimension of graphs and images of almost every continuous function, see among others [28], [4].

One of the advantages of prevalence with respect to other notions of genericity is that it allows the use of probabilistic methods in the proofs. However the statements are fully non-probabilistic and the kind of problems one encounters in formulating prevalence results are quite distinct from those of a purely probabilistic setting, much more explored in the literature.

Our strategy and results are based on key ideas introduced by Catellier and one of the authors in [10], while studying the regularisation properties of fractional Brownian motion. In the following, we schematically recall some of the main results of that paper and how they fit in our framework.

Consider the perturbed ODE in \mathbb{R}^d in integral form given by

$$x_t = x_0 + \int_0^t b(s, x_s) ds + w_t \quad (1)$$

where $x_t = x(t)$, $w \in C^0(\mathbb{R}_+, \mathbb{R}^d)$ is our fixed perturbation and $b : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a time-dependent vector field (assume it to be continuous for the moment).

Due to the additive nature of the perturbation, there are no particular regularity requirements for w ; it is natural to ask for which classes of b equation (1) is well-posed and if, for suitable w , one can obtain well-posedness results in classes which are ill-posed when $w = 0$.


The work [10] singles out two analytic concepts linked to the well-posedness of (1):

1. The *averaged field* $T^w b$, given by


$$T^w b(t, x) = \int_0^t b(s, x + w_s) ds, \quad x \in \mathbb{R}^d, t \geq 0.$$

2. The notion of ρ -irregularity of a path w , defined precisely below.

Indeed, in order to study (1), i.e. to give it meaning and establish existence and uniqueness of solutions, together with a Lipschitz flow, it is enough to have good regularity estimates for the averaged function $T^w b$, even when the drift b is merely a distribution. Schematically:



In order to analyse the regularising effect of w on a collection of drifts b , the *averaging operator* $T^w : b \mapsto T^w b$ is then introduced; it is closely tied to the *occupation measure* μ^w , whose regularity is in turn quantified by the concept of ρ -irregularity:




Here E is a suitable Banach space, for instance in [10] Fourier–Lebesgue spaces are considered. The precise meaning of the above diagram is explained in detail in Sections 2.2 and 2.3.

Even if ρ -irregularity has been first introduced in the study of perturbed ODEs, it is a concept of independent interest and has been applied to establish regularisation by noise for several modulated PDEs, see [12], [14], [13]. More recently, a connection to the property of inviscid mixing for shear flows has been found in [35].

In the companion paper [34], we analyse in detail the regularity of the averaged field $T^w b$ for generic $w \in C^\delta$ when b is a *fixed* distribution. Here instead we focus on the regularity of the operator T^w for a generic $w \in C^\delta$, as quantified by the concept of ρ -irregularity. This has the advantage of showing that

a generic additive perturbation w regularises a huge class of ODEs, namely the ones of the form (1) associated to $b \in E$. On the other hand, it has the drawback that the space E is strictly smaller than the class of all possible drifts taken singularly. For instance in [34] time dependent distributions b , with only L^p -regularity in time, can be considered, while it is not possible to obtain regularity estimates for the operator T^w in that class, as explained in Section 3.1 therein.


In this paper, we expand the theory developed in [10] in several ways. For instance, we identify suitable locally-nondeterministic conditions (LND) which ensure ρ -irregular trajectories for a large class of stochastic Gaussian processes:



The concept of *local-nondeterminism* is explained in detail in Section 2.4. The above probabilistic result then allows to prove a much stronger (non probabilistic) prevalence statement, thanks to the fact that the LND condition above is invariant under additive deterministic perturbations.

Usually prevalence statements are established by means of *probes*, i.e. measures supported on finite dimensional subspaces. To the best of our knowledge, the only other work adopting a approach similar to ours, i.e. exploiting stochastic processes to establish prevalence results, is [4]. See also [54] and the references therein for the study of properties of fractional Brownian motion with deterministic drift (however without prevalence considerations).

We also study more in detail the geometric and analytic properties of ρ -irregular functions, like the Fourier dimension of their image. We establish that ρ -irregularity is indeed a concept of irregularity, as the terminology suggests, in a precisely quantifiable way:



Here δ^* is a critical scaling parameter which depends on ρ . The precise statements of our results are given in Section 3, but let us give here informally some of the main highlights:

- Almost every $w \in C^\delta([0, T]; \mathbb{R}^d)$ is ρ -irregular for any $\rho < (2\delta)^{-1}$. Moreover we introduce a stronger notion of irregularity, which allows us to prove that almost every $\varphi \in C^\delta([0, T]; \mathbb{R}^d)$ remains irregular under smooth additive perturbations. See Theorems 32-i) and 38.
- Almost every $w \in C^0([0, T]; \mathbb{R}^d)$ is infinitely irregular, in the sense that it is ρ -irregular for any $\rho < \infty$. The associated averaging operator is infinitely regularising, in the sense that T^w maps $S'(\mathbb{R}^d)$ into $C^\infty(\mathbb{R}^d)$. See Theorems 32-ii).
- Almost every $w \in C^0([0, T]; \mathbb{R}^d)$ has infinite regularisation effect on the ODE. Namely, it renders (1) well-posed and with a smooth flow for a large class of fields b , for instance for any $b \in H^\alpha(\mathbb{R}^d; \mathbb{R}^d)$ for any $\alpha \in \mathbb{R}$. See Theorem 40.

An overview of our results, combined with the path-wise theory developed in [10] [12], [14] discussed above, is given by the conceptual map of Figure 1.

While finalising the first version of this paper, we become aware of the preprint [39] which independently develop some result similar to ours. The authors also identify some LND conditions that ensure time-space regularity of μ^w and apply them in combination with the theory from [10] to establish that suitable stochastic processes are “infinitely regularising” the perturbed ODE. By introducing genericity into the picture our work carries such probabilistic considerations into a wider scope and quantifies




Figure 1: Scheme of the results

the regularisation effect precisely wrt. the Hölder regularity of the paths. We also answer some of the questions left open in [39], for instance on the link between regularity of μ^w and true roughness of w .

Let us finally mention the upcoming works [40, 58], further analysing notions of irregularity of paths and giving sufficient conditions to check them for stochastic processes; among other results, it seems the authors are able to give a negative answer to Conjecture 79 from Section 5.3.

Structure of the paper. Section 2 introduces all our main actors, namely prevalence, ρ -irregularity, occupation measures and averaging operators and describes the basic relations between them. Fractional Brownian motion (fBm) enters into the picture as a transverse measure for prevalence and local-nondeterminism (LND) as a key feature of fBm paths. The reader acquainted with these concepts can skip this preliminary section. With the above preparations, we can formulate in Section 3.1 the statements of our main results. Section 3.2 shows how they can be combined with already existing results to provide various “noiseless regularisation by noise” results both for ODEs and PDEs. Sections 4 and 5 constitute the main body of the paper and contain the proofs of the statements from Section 3.1. These sections split naturally into a probabilistic part, in which stochastic criteria for establishing ρ -irregularity are developed, and an analytic one, focused on geometric and analytical properties of ρ -irregular paths. We choose to put in the Appendix reminders of useful standard facts.

Notation. We will use the notation $a \lesssim b$ to mean that there exists a positive constant c such that

$a \leq c b$; we use the index $a \lesssim_x b$ to highlight the dependence $c = c(x)$. $a \sim b$ if and only if $a \lesssim b$ and $b \lesssim a$, similarly for $a \sim_x b$.

We will always work on a finite time interval $[0, T]$ unless stated otherwise. Whenever useful we adopt the convention that f_t stands for $f(t)$ for a function f indexed on $t \in [0, T]$, but depending on the context we will use both notations; similarly for the increments of $f_{s,t} = f_t - f_s$.

For $x \in \mathbb{R}^d$, $|x|$ denotes the Euclidean norm, $x \cdot y$ the scalar product, $\langle x \rangle = (1 + |x|^2)^{1/2}$. For any $R > 0$, B_R stands for $B(0, R) = \{x \in \mathbb{R}^d : |x| \leq R\}$, $\mathbb{S}^{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$.

We denote by $\mathcal{S}(\mathbb{R}^d; \mathbb{R}^m)$ and $\mathcal{S}'(\mathbb{R}^d; \mathbb{R}^m)$ respectively the spaces of vector-valued Schwarz functions and tempered distributions on \mathbb{R}^d , similarly $C_c^\infty(\mathbb{R}^d; \mathbb{R}^m)$ is the set of vector-valued smooth compactly supported functions; \hat{f} stands for the Fourier transform of $f \in \mathcal{S}'(\mathbb{R}^d)$. Standard Lebesgue spaces are denoted by $L^p(\mathbb{R}^d; \mathbb{R}^m)$. Whenever it doesn't create confusion, we will just write C_c^∞ and L^p for short.

Given a Banach space E , $\alpha \in (0, 1)$, $C^\alpha([0, T]; E) = C_t^\alpha E$ is the space of E -valued α -Hölder continuous functions, i.e. $f : [0, T] \rightarrow E$ such that

$$\|f\|_{C^\alpha E} := \|f\|_{C^0 E} + \llbracket f \rrbracket_{C^\alpha E} = \sup_{t \in [0, T]} \|f_t\|_E + \sup_{s \neq t \in [0, T]} \frac{\|f_{s,t}\|_E}{|t - s|^\alpha} < \infty.$$

A similar definition holds for $\text{Lip}([0, T]; E) = \text{Lip}_t E$. More generally, for a given modulus of continuity ω (possibly defined only in a neighbourhood of 0), we denote by $C^\omega([0, T]; E) = C^\omega E$ the set of all E -valued continuous functions with modulus of continuity ω . $C_t^0 E = C^0([0, T]; E)$ is the space of E -valued continuous functions, endowed with the supremum norm.

Whenever $E = \mathbb{R}^d$, we will refer to $w \in C_t^\alpha = C^\alpha([0, T]; \mathbb{R}^d)$ as a *path* and in this case we allow $\alpha \in [0, \infty)$ with the convention that $w \in C_t^\alpha$ it has continuous derivatives up to order $\lfloor \alpha \rfloor$ and $D^{\lfloor \alpha \rfloor} \varphi$ is $\{\alpha\}$ -Hölder continuous, where $\lfloor \alpha \rfloor$ and $\{\alpha\}$ denote integer and fractional part.

Whenever a stochastic process $X = \{X_t\}_{t \geq 0}$ is considered, even when it is not specified we imply the existence of an abstract underlying filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ such that \mathcal{F} and \mathcal{F}_t satisfy the usual assumptions and X_t is adapted to \mathcal{F}_t . If \mathcal{F}_t is said to be the natural filtration generated by X , then it is tacitly implied that it is actually its right continuous, normal augmentation. We denote by \mathbb{E} integration (equiv. expectation) w.r.t. the probability \mathbb{P} and by $\mathbb{E}[\cdot | \mathcal{G}]$ conditional expectation w.r.t. a σ -algebra \mathcal{G} .

Acknowledgements. We thank Nicolas Perkowski and Leonardo Tolomeo for useful discussions, leading respectively to Lemma 23 and Remark 24; we are also grateful to the anonymous referee for their very carefully reading and numerous insights, which highly improved the paper. The authors were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Hausdorff Center for Mathematics under Germany's Excellence Strategy – EXC-2047/1 – 390685813 and through CRC 1060 - projekt number 211504053.

2 Preliminaries

2.1 Introduction to prevalence

We follow the exposition and the terminology given in [42]. The reader might however keep in mind that in the following we will be only interested in the case of a Banach space E .

Definition 1. Let E be a complete metric vector space. A Borel set $A \subset E$ is said to be **shy** if there exists a measure μ such that:

- i. There exists a compact set $K \subset E$ such that $0 < \mu(K) < \infty$.
- ii. For every $v \in E$, $\mu(v + A) = 0$.

In this case, the measure μ is said to be transverse to A . More generally, a subset of E is *shy* if it is contained in a shy Borel set. The complement of a shy set is called a *prevalent set*.

Sometimes it is said more informally that the measure μ “witnesses” the prevalence of A^c in E . It follows immediately from part *i.* of the definition that if needed one can assume μ to be a compactly supported probability measure on E . If E is separable, then any probability measure on E is tight and therefore *i.* is automatically satisfied.

The following properties hold for prevalence:

1. If E is finite dimensional, then a set A is shy if and only if it has zero Lebesgue measure (cf. Fact 6 from [42]).
2. If A is shy, then so is $v + A$ for any $v \in E$ (Fact 1 from [42]).
3. Prevalent sets are dense (Fact 2' from [42]).
4. If $\dim(E) = +\infty$, then compact subsets of E are shy (Fact 8 from [42]).
5. Countable union of shy sets is shy; conversely, countable intersection of prevalent sets is prevalent (Fact 3" from [42]).

Remark 2. *Although it is not entirely trivial, it follows from the definition that a Borel set A cannot be at the same time both shy and prevalent. To see it, assume by contradiction this to be the case; then we can find two compactly supported probability measures μ, ν such that $\mu(v + A) = 0$ and $\nu(v + A^c) = 0$ for all $v \in E$, where A^c denotes the complement of A . It would then follow from Fact 3 from [42] that $(\mu * \nu)(v + A) = 0 = (\nu * \mu)(v + A^c)$ for all $v \in E$, yielding to $(\mu * \nu)(A)$ having at the same time both values 0 and 1.*

From now, whenever we say that a statement holds for a.e. $v \in E$, we mean that the set of elements of E for which the statement holds is a prevalent set; property 1. states that this is consistent with the finite dimensional case.

In the context of a function space E , it is natural to consider as probability measure the law induced by an E -valued stochastic process. Namely, given stochastic process W defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ taking values in a separable Banach space E , in order to show that a property \mathcal{P} holds for a.e. $f \in E$, it suffices to show that

$$\mathbb{P}(f + W \text{ satisfies property } \mathcal{P}) = 1 \quad \forall f \in E.$$

Clearly, we are assuming that the set $A = \{w \in E : w \text{ satisfies property } \mathcal{P}\}$ is Borel measurable; if E is not separable, then we need in addition to require that the law of W is tight, so that point *i.* of Definition 1 is satisfied.

Let us point out that, as a consequence of Properties 4. and 5. above, the set of all possible realisations of any probability measure μ on a separable infinite dimensional Banach E space (actually, any tight measure μ on an infinite dimensional Banach space E) is contained in a shy set, in the following sense: we can always find a Borel set $A \subset E$, which is a countable union of compacts, such that $\mu(A) = 1$, yet A is shy. If we are given another Borel set $\tilde{A} \subset E$, in order to verify its prevalence we really need to check the translation property that $\mu(\tilde{A} + f) = 1$ for any $f \in E$; if we only know that $\mu(\tilde{A}) = 1$, then \tilde{A} can be shy or prevalent (or neither).

This highlights the difference between a statement of the form “Property \mathcal{P} holds for a.e. f ” and for instance “Property \mathcal{P} holds for all Brownian trajectories”, where the last statement corresponds to $\mu(\text{Property } \mathcal{P} \text{ holds}) = 1$, μ being the Wiener measure on $C^0([0, 1])$. Indeed, the second statement doesn’t provide any information regarding whether the property might be prevalent or not; intuitively, the elements satisfying a prevalence statement are “many more” than just the realisations of the Wiener measure.

2.2 Basic properties of ρ -irregularity and related concepts

The concept of ρ -irregularity was introduced in [10] as an analytic property of continuous functions which allows to quantitatively measure both their irregularity and their smoothing effect on perturbations of ODEs.

Definition 3. Let $\gamma \in [0, 1]$, $\rho > 0$. A measurable path $w : [0, T] \rightarrow \mathbb{R}^d$ is (γ, ρ) -irregular if there exists a constant C such that

$$\left| \int_s^t e^{i\xi \cdot w_r} dr \right| \leq C |\xi|^{-\rho} |t - s|^\gamma \quad \forall \xi \in \mathbb{R}^d \setminus \{0\}, s, t \in [0, T]. \quad (2)$$

We denote by $\|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}$ the optimal constant C ; with the notation $\Phi_t^w(\xi) = \int_0^t e^{i\xi \cdot w_r} dr$, it holds

$$\|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} := \sup_{\xi \in \mathbb{R}^d, s \neq t} \frac{|\Phi_{s,t}^w(\xi)| |\xi|^\rho}{|t - s|^\gamma}.$$

We say that w is ρ -irregular if there exists $\gamma > 1/2$ such that w is (γ, ρ) -irregular.

We have the trivial bound $|\Phi_{s,t}^w(\xi)| \leq |t - s|$, so that (2) is always satisfied for $|\xi|$ small and the relevant information in the above definition is given by the uniform bound as $\xi \rightarrow \infty$. Therefore we can always without problems replace $|\xi|$ with another function with the same asymptotic behaviour (in the original definition from [10], $1 + |\xi|$ appeared, but we prefer here the choice $|\xi|$ for its better scaling properties).

Let us collect now some elementary facts on ρ -irregular functions.

Lemma 4. Let $w : [0, T] \rightarrow \mathbb{R}^d$ be a (γ, ρ) -irregular continuous path. Then the following hold:

- i. Symmetry invariance: $-w$ is (γ, ρ) -irregular with $\|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} = \|\Phi^{-w}\|_{\mathcal{W}^{\gamma, \rho}}$.
- ii. Translation invariance: for any $r \in [0, T]$, $w - w_r$ is (γ, ρ) -irregular with $\|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} = \|\Phi^{w - w_r}\|_{\mathcal{W}^{\gamma, \rho}}$.
- iii. Scaling invariance: for any $\lambda \in (0, 1)$, $w^\lambda(t) := \lambda^{-(1-\gamma)/\rho} w(\lambda t)$ is (γ, ρ) -irregular.
- iv. Rotation invariance: for any $O \in \text{SO}(d)$, Ow is (γ, ρ) -irregular with $\|\Phi^{Ow}\|_{\mathcal{W}^{\gamma, \rho}} = \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}$.
- v. More generally if $A \in \mathbb{R}^{d \times d}$ is invertible, then Aw is (γ, ρ) -irregular with

$$\|\Phi^{Aw}\|_{\mathcal{W}^{\gamma, \rho}} \leq \|(A^T)^{-1}\|^\rho \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}.$$

Proof. All the statements follow from elementary calculations; let us prove only iii. and v. . Fix $\lambda \in (0, 1)$, then

$$\Phi_{s,t}^{w^\lambda}(\xi) = \int_s^t e^{i\xi \cdot \lambda^{-(1-\gamma)/\rho} w_{\lambda r}} dr = \lambda^{-1} \int_{\lambda s}^{\lambda t} e^{i\lambda^{-(1-\gamma)/\rho} \xi \cdot w_r} dr = \lambda^{-1} \Phi_{\lambda s, \lambda t}^w(\lambda^{-(1-\gamma)/\rho} \xi),$$

so that

$$\frac{|\Phi_{s,t}^{w^\lambda}(\xi)| |\xi|^\rho}{|t - s|^\gamma} = \lambda^{-1} \frac{|\Phi_{\lambda s, \lambda t}^w(\lambda^{-(1-\gamma)/\rho} \xi)| |\xi|^\rho}{|t - s|^\gamma} = \frac{|\Phi_{\lambda s, \lambda t}^w(\lambda^{-(1-\gamma)/\rho} \xi)| |\lambda^{-(1-\gamma)/\rho} \xi|^\rho}{|\lambda t - \lambda s|^\gamma} \leq \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}.$$

Regarding v., similarly we have

$$\frac{|\Phi_{s,t}^{Aw}(\xi)| |\xi|^\rho}{|t - s|^\gamma} = \left| \int_s^t e^{i(A^T \xi) \cdot w_r} dr \right| \frac{|\xi|^\rho}{|t - s|^\gamma} \leq \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \left(\frac{|A^T \xi|}{|\xi|} \right)^{-\rho} \leq \|(A^T)^{-1}\|^\rho \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}.$$

□

Remark 5. There is a striking similarity between i.–iii. and properties of stochastic processes like self-similarity and stationarity; however the latter properties are of statistical nature, namely they preserve the law of the process, but not fixed trajectories, while ρ -irregularity is an analytical property which holds for deterministic trajectories. From property iii. we also deduce the existence of a critical scaling parameter associated to the pair (γ, ρ) given by

$$\delta_{\gamma, \rho}^* = \frac{1 - \gamma}{\rho}; \quad (3)$$

we will see in Theorem 36 how $\delta_{\gamma, \rho}^*$ relates to regularity of (γ, ρ) -irregular paths.

Remark 6. It's easy to see from the definition that (γ, ρ) -irregularity for w is equivalent to the following: there exists a constant C such that, for any $v \in \mathbb{S}^{d-1}$, $v \cdot w$ is (γ, ρ) -irregular with $\|\Phi^{v \cdot w}\|_{\mathcal{W}^{\gamma, \rho}} \leq C$. The latter condition is not equivalent to checking (γ, ρ) -irregularity of the coordinates $w^{(i)}$ (i.e. to $v = e_i$, $i = 1, \dots, d$): for instance if w is a 1-dimensional (γ, ρ) -irregular function and we define $\tilde{w}_t := (w_t, -w_t)$, then the single coordinates of \tilde{w} are (γ, ρ) -irregular but \tilde{w} is not, since $(1, 1) \cdot \tilde{w} \equiv 0$.

Lemma 7. Let w be (γ, ρ) -irregular, then for any $\theta \in [0, 1]$ it is also $(\gamma^\theta, \rho^\theta)$ -irregular for the choice $\gamma^\theta = 1 - \theta + \theta\gamma$, $\rho^\theta = \theta\rho$ and it holds $\|\Phi^w\|_{\mathcal{W}^{\gamma^\theta, \rho^\theta}} \leq \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}^\theta$.

Proof. The conclusion follows immediately by interpolating the two inequalities

$$|\Phi_{s,t}^w(\xi)| \leq |t - s|, \quad |\Phi_{s,t}^w(\xi)| \leq \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} |t - s|^\gamma |\xi|^{-\rho}.$$

□

The above lemma shows that we can always trade space regularity for time regularity, i.e. we can decrease the parameter ρ in order to increase γ . Let us also point out that

$$\delta_{\gamma, \rho}^* = \frac{1 - \gamma}{\rho} = \frac{1 - \gamma^\theta}{\rho^\theta} = \delta_{\gamma^\theta, \rho^\theta}^*, \quad (4)$$

i.e. the critical scaling parameter is left unchanged by this procedure.

In general dimension $d \geq 2$, it is in general difficult to establish if a given function is (γ, ρ) -irregular. This fact is one of the main motivations of our interest in establishing the prevalence of this property. The situation is different in the case $d = 1$, in which there are simple conditions to establish ρ -irregularity (at least for some values of ρ).

Proposition 8 (Proposition 1.4 from [12]). Let $w \in C^1([0, T]; \mathbb{R})$ be such that $\inf_t |w'_t| \geq \delta > 0$ and $w'' \in L^1$, then w is $(\gamma, 1 - \gamma)$ -irregular for any $\gamma \in (0, 1)$.

Let us also introduce the following concept of *exponential irregularity*, which is a novel contribution of the present work.

Definition 9. A measurable path $w : [0, T] \rightarrow \mathbb{R}^d$ is exponentially irregular if there exist positive constants c_1, c_2 and $\gamma \in (0, 1)$ such that

$$|\Phi_{s,t}^w(\xi)| \leq c_1 e^{-c_2 |\xi|} |t - s|^\gamma \quad \forall \xi \in \mathbb{R}^d \setminus \{0\}, s, t \in [0, T]. \quad (5)$$

Contrary to Definition 3, in Definition 9 we do not impose $\gamma > 1/2$; this is because, arguing by interpolation as in Lemma 7, it is easy to check that if (5) holds *some* $\gamma \in (0, 1)$, then the path w actually satisfies (5) for *all* $\gamma \in (0, 1)$, up to modifying the constant $c_i = c_i(\gamma)$, $i = 1, 2$. It is then also clear that exponential irregularity implies ρ -irregularity, for any value $\rho \in (0, +\infty)$.

We now prove that the ρ -irregularity and exponential irregularity properties define Borel sets, which is the first step in order to establish prevalence.

Lemma 10. For any $\rho > 0$, the set

$$A = \{w : [0, T] \rightarrow \mathbb{R}^d \mid w \text{ is } \rho\text{-irregular}\}$$

is Borel measurable w.r.t to the topology induced by any of the following norms: $\|\cdot\|_{L^p}$, $p \in [1, \infty]$, $\|\cdot\|_{C^0}$, $\|\cdot\|_{C^\alpha}$, $\alpha \in (0, 1)$.

Proof. We can write the set A as the following countable union:

$$A = \bigcup_{n,m \in \mathbb{N}} A_{n,m} = \bigcup_{n,m \in \mathbb{N}} \left\{ w : [0, T] \rightarrow \mathbb{R}^d \mid \sup_{\xi \in \mathbb{R}^d, s \neq t} \frac{|\Phi_{s,t}^w(\xi)| |\xi|^\rho}{|t-s|^{1/2+1/m}} \leq n \right\}.$$

It will be then sufficient to show that for every m, n the set $A_{m,n}$ is closed in the above topologies. We will actually show that it is closed under convergence in measure, which is weaker than any of the norms considered and therefore yields the conclusion.

Let w_k be a sequence of elements of $A_{m,n}$ such that $w_k \rightarrow w$ in measure, then by dominated convergence it's easy to see that for any fixed $s < t$, $\xi \in \mathbb{R}^d$ it holds $\Phi_{s,t}^{w_k}(\xi) \rightarrow \Phi_{s,t}^w(\xi)$. But then

$$\frac{|\Phi_{s,t}^w(\xi)| |\xi|^\rho}{|t-s|^{1/2+1/m}} = \lim_{k \rightarrow \infty} \frac{|\Phi_{s,t}^{w_k}(\xi)| |\xi|^\rho}{|t-s|^{1/2+1/m}} \leq n$$

and since the reasoning holds for any fixed $s < t$ and ξ we can conclude that $f \in A_{m,n}$ as well. \square

Remark 11. More generally, given a modulus of continuity φ and a function $F : \mathbb{R}^d \rightarrow \mathbb{R}^+$, the same proof shows that the set

$$B = \left\{ w : [0, T] \rightarrow \mathbb{R}^d \mid \sup_{\xi \in \mathbb{R}^d, s \neq t} \frac{|\Phi_{s,t}^w(\xi)| F(\xi)}{\varphi(|t-s|)} < \infty \right\}$$

is Borel measurable in any of the above topologies. The fact that exponential irregularity defines Borel sets is established similarly.

We conclude this section with a brief detour on Carathéodory functions and their connection with the exponential irregularity property. Here λ denotes the Lebesgue measure on \mathbb{R}^d , while \mathcal{L} denotes the Lebesgue measure on $[0, T]$.

Definition 12. A measurable function $f : [0, T] \rightarrow \mathbb{R}^d$ is a Carathéodory function if for any set $D \subset \mathbb{R}^d$ such that $\lambda(D) > 0$ it holds $\mathcal{L}(f^{-1}(D) \cap B) > 0$ for every (non empty) interval B .

Observe in particular that if f is Carathéodory, then it is unbounded on every interval and thus discontinuous.

Lemma 13. Let w be an exponentially irregular measurable path, then w is Carathéodory.

Proof. The statement follows immediately from the considerations given at the beginning of Section 6 from [6], see also Sections 11 and 28 from [37]. Let us briefly sketch the proof.

Denoting by $\mu_{s,t}^w$ the occupation measure associated to w , which is defined in Definition 14 below, it follows from the exponential irregularity property that we can find $c > 0$ such that for any $s < t$ it holds

$$\int_{\mathbb{R}^d} e^{c|\xi|} |\hat{\mu}_{s,t}^w(\xi)|^2 d\xi = \int_{\mathbb{R}^d} e^{c|\xi|} |\Phi_{s,t}^w(\xi)|^2 d\xi < \infty.$$

It then follows from the Paley–Wiener Theorem that $\mu_{s,t}^w$ is analytic and therefore it cannot vanish on any set $D \subset \mathbb{R}^d$ such that $\lambda(D) > 0$, namely for $B = [s, t]$ it holds

$$\mathcal{L}(w^{-1}(D) \cap [s, t]) = \int_D \mu_{s,t}^w(y) dy > 0.$$

\square

2.3 Occupation measures and averaging operators

So far we have discussed several properties of ρ -irregularity, but we haven't motivated the importance of such notion and its relation with regularisation by noise phenomena. It turns out that ρ -irregularity is closely tied to the *occupation measure* of the path w .

In the following $\mathcal{M}(\mathbb{R}^d)$ denotes the set of all finite Radon measures on \mathbb{R}^d , endowed with the total variation norm $\|\cdot\|_{\text{TV}}$; $\mathcal{M}_+(\mathbb{R}^d)$ is the closed subset of non-negative measures.

Definition 14. Given a measurable path $w : [0, T] \rightarrow \mathbb{R}^d$, we define its occupation measure as the family $(\mu_{s,t}^w)_{0 \leq s \leq t \leq T} \subset \mathcal{M}_+(\mathbb{R}^d)$ given by $\mu_{s,t}^w = w_*(\mathcal{L}_{[s,t]})$, namely

$$\int_{\mathbb{R}^d} f(y) \mu_{s,t}^w(dy) = \int_{[s,t]} f(w_r) dr \quad \forall f \in C_b^0(\mathbb{R}^d).$$

Observe that by definition $\mu_{s,t}^w = \mu_{0,t}^w - \mu_{0,s}^w$; for this reason we will identify the family $(\mu_{s,t}^w)_{s \leq t}$ with the map $\mu^w \in C^0([0, T]; \mathcal{M}_+(\mathbb{R}^d))$ given by $t \mapsto \mu_t^w = \mu_{0,t}^w$, so that $\mu_{s,t}^w$ represents an increment of μ^w .

Note that $\mu^w \in \text{Lip}([0, T]; \mathcal{M}_+(\mathbb{R}^d))$ with $\|\mu_{s,t}^w\|_{\text{TV}} = |t - s|$ and Gateaux derivative $\dot{\mu}_t^w = \delta_{w_t}$.

The Fourier transform of $\mu_{s,t}^w$ is given by

$$\widehat{\mu_{s,t}^w}(\xi) = \int_{\mathbb{R}^d} e^{-i\xi \cdot y} \mu_{s,t}^w(dy) = \int_s^t e^{-i\xi \cdot w_r} dr = \overline{\Phi_{s,t}^w(\xi)}$$

which shows that w is (γ, ρ) -irregular if and only if the map $t \mapsto \mu_t^w$ belongs to $C^\gamma([0, T]; \mathcal{FL}^{\rho, \infty}(\mathbb{R}^d))$, where the Fourier–Lebesgue spaces $\mathcal{FL}^{\rho, p}$ with $p \in [1, \infty]$ are defined by

$$\mathcal{FL}^{\rho, p} = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \langle \xi \rangle^\rho |\hat{f}(\xi)| \in L^p \right\},$$

see Appendix A.2 for their main properties. In particular we have

$$\|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \sim \|\mu^w\|_{C^\gamma \mathcal{FL}^{\rho, \infty}}. \quad (6)$$

Remark 15. We will mostly work with given measurable paths w , but both definitions of Φ^w and μ^w are not affected by changing w on an $\mathcal{L}_{[0, T]}$ -negligible subset of $[0, T]$; therefore they also makes sense when dealing with equivalence classes like $w \in L^p(0, T; \mathbb{R}^d)$ for $p \in [1, \infty]$. Similarly, it makes sense for w in an equivalence class to say that it is (γ, ρ) -irregular (resp. exponentially irregular).

Occupation measures are also closely related to averaging operators.

Definition 16. Let $w : [0, T] \rightarrow \mathbb{R}^d$ be a measurable bounded function, then we define the averaging operator associated to w as the family of operators $\{T_{s,t}^w, 0 \leq s \leq t \leq T\} \subset \mathcal{L}(\mathcal{S}'(\mathbb{R}^d); \mathcal{S}'(\mathbb{R}^d))$ given by

$$T_{s,t}^w b = \int_s^t b(\cdot + w_r) dr$$

or equivalently by duality as follows: for any $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and any $b \in \mathcal{S}'(\mathbb{R}^d)$ it holds

$$\langle T_{s,t}^w b, \varphi \rangle = \langle b, \int_s^t \varphi(\cdot - w_r) dr \rangle. \quad (7)$$

As before, $T_{s,t}^w = T_{0,t}^w - T_{0,s}^w$ and therefore we identify $(T_{s,t}^w)_{s \leq t}$ with the map $t \mapsto T_t^w = T_{0,t}^w$.

Remark 17. Differently from Definitions 3 and 14, in Definition 16 we have required that w is bounded; the reason for this is that otherwise it is unclear a priori if, for a given $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $\int_s^t \varphi(\cdot - w_r) dr \in$

$\mathcal{S}(\mathbb{R}^d)$ uniformly in $s < t$ and so if the above is a good definition. However by looking at the Fourier transform $\widehat{\varphi(\cdot - w_t)} = e^{-i\xi \cdot w_t} \hat{\varphi}$ one can check that $w \in L^\infty$ can be relaxed to requiring

$$\int_0^T |w_t|^N dt < \infty \quad \text{for all } N \in \mathbb{N},$$

namely $w \in L_t^p$ for all $p < \infty$. It is still true, analogously to Remark 15, that $T^w = T^{\tilde{w}}$ if $w = \tilde{w}$ up to $\mathcal{L}_{[0,T]}$ -negligible sets; for this reason from now on when dealing with T^w we will always implicitly assume $w \in L_t^\infty$.

Averaging operators can also be defined for time-dependent distributions, as done in [34]. In the time dependent case however we lose the following fundamental property which relates the averaging operator to the occupation measure.

Lemma 18. *Let $w \in L^\infty$, μ^w and T^w as above. Then for any $b \in \mathcal{S}'(\mathbb{R}^d)$, $T_{s,t}^w b = \tilde{\mu}_{s,t}^w * b$, where $\tilde{\mu}$ denotes the reflection of μ , i.e. $\tilde{\mu}_{s,t}^w(A) = \mu_{s,t}^w(-A)$.*

Proof. Observe that by definition of occupation measure, for any $s \leq t$ and any $R \geq \|w\|_{L^\infty}$, it holds $\text{supp } \mu_{s,t}^w \subset B_R$. In particular since $\mu_{s,t}^w$ is a measure with compact support, for any $b \in \mathcal{S}'(\mathbb{R}^d)$ the convolution $b * \mu_{s,t}^w$ is well defined. The same goes for $\tilde{\mu}_{s,t}^w$. Let $\varphi \in \mathcal{S}(\mathbb{R}^d)$, then for any $x \in \mathbb{R}^d$ it holds

$$\int_s^t \varphi(x - w_r) dr = \int \varphi(x - y) \mu_{s,t}^w(dy) = (\varphi * \mu_{s,t}^w)(x).$$

The conclusion for b follows by the duality formula (7) and $\langle b, \varphi * \mu_{s,t}^w \rangle = \langle \tilde{\mu}_{s,t}^w * b, \varphi \rangle$. \square

As a consequence, estimating the regularisation properties of T^w is equivalent to estimating the regularity of μ^w in suitable function spaces. This is exactly where the notion of ρ -irregularity comes into play.

Lemma 19. *Let $w \in L^\infty(0, T; \mathbb{R}^d)$ be (γ, ρ) -irregular. Then for any $\alpha \in \mathbb{R}$ and $p \in [1, \infty]$, the averaging operator T^w belongs to $C_t^\gamma \mathcal{L}(\mathcal{F}L^{\alpha,p}; \mathcal{F}L^{\alpha+\rho,p})$ and for any $b \in \mathcal{F}L^{\alpha,p}$ it holds*

$$\|T_{s,t}^w b\|_{\mathcal{F}L^{\alpha+\rho,p}} \lesssim |t-s|^\gamma \|b\|_{\mathcal{F}L^{\alpha,p}} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}. \quad (8)$$

Proof. The statement follows from the considerations given in the introduction of [10]. Alternatively, using Lemma 93 from Appendix A.2, it holds

$$\|T_{s,t}^w b\|_{\mathcal{F}L^{\alpha+\rho,p}} = \|\tilde{\mu}_{s,t}^w * b\|_{\mathcal{F}L^{\alpha+\rho,p}} \leq \|\tilde{\mu}_{s,t}^w\|_{\mathcal{F}L^{\rho,\infty}} \|b\|_{\mathcal{F}L^{\alpha,p}} \lesssim |t-s|^\gamma \|b\|_{\mathcal{F}L^{\alpha,p}} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}.$$

\square

Unfortunately, Fourier–Lebesgue spaces are not always very useful, with the exclusion of the scale $p = 2$, in which case $\mathcal{F}L^{\alpha,2} = H^\alpha$. We can however use Fourier–Lebesgue embeddings to deduce regularity for $\mu_{s,t}^w$ in other scales of spaces, which in turn imply different estimates for $T_{s,t}^w$. To this end, following [37], we introduce the concept of *occupation density*. In the probabilistic literature it is usually referred to as *local time* and we will indifferently use both terminologies.

Definition 20. *We say that a measurable $w : [0, T] \rightarrow \mathbb{R}^d$ admits an occupation density if for any $s < t$, $\mu_{s,t}^w$ is absolutely continuous w.r.t. $\mathcal{L}_{[s,t]}$, in which case we denote by $\ell_{s,t}^w$ its density, so that $\mu_{s,t}^w(dx) = \ell_{s,t}^w(x)dx$. As usual it holds $\ell_{s,t}^w = \ell_{0,t}^w - \ell_{0,s}^w$ and we set $\ell_{0,t}^w = \ell_t^w$. Sometimes we will also use the notation $\ell_t^w(x) = \ell^w(t, x)$.*

The spaces $B_{p,q}^s = B_{p,q}^s(\mathbb{R}^d)$ appearing in the next lemma are inhomogeneous Besov spaces, see [3]. Similar statements can be given for more classical Sobolev spaces $W^{k,p}$ or Bessel potential spaces $L^{s,p} = (1 - \Delta)^{-s/2} L^p$.

Lemma 21. *Let $w : [0, T] \rightarrow \mathbb{R}^d$ be a (γ, ρ) -irregular measurable path. Then:*

- i. If $\rho > d/2$, then w admits an occupation density $\ell^w \in C_t^\gamma L_x^2 \cap \text{Lip}_t L_x^1$.
- ii. If $\rho > d$, then ℓ^w is jointly continuous in (t, x) and $\ell^w \in C_t^\gamma C_x^0$.
- iii. If $\rho > d/2 + k$ for some $k \in \mathbb{N}$, then $\ell^w \in C_t^\gamma W_x^{k,2}$; in particular, if $w \in L^\infty$, then ℓ is compactly supported on $[0, T] \times \mathbb{R}^d$ and therefore $\ell^w \in C_t^\gamma W_x^{k,1}$.
- iv. As a consequence, if $\rho > d/2 + k$ for some $k \in \mathbb{N}$ and $w \in L^\infty$, then for any $\alpha \in \mathbb{R}$, $p, q \in [1, \infty]$ it holds

$$T^w \in C^\gamma([0, T]; \mathcal{L}(B_{p,q}^\alpha; B_{p,q}^{\alpha+k}))$$

where $B_{p,q}^\alpha$ denote Besov spaces. In particular, for any $b \in B_{p,q}^\alpha$ it holds

$$\|T_{s,t}^w b\|_{B_{p,q}^{\alpha+k}} \lesssim |t-s|^\gamma \|b\|_{B_{p,q}^\alpha} \|w\|_{L^\infty}^{d/2} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}} \quad \text{uniformly in } s < t.$$

Proof. By equation (6) and the Fourier–Lebesgue embedding $\mathcal{FL}^{\rho,\infty} \hookrightarrow \mathcal{FL}^{0,2} = L^2$ which holds for $\rho > d/2$, we deduce that $\mu^w \in C_t^\gamma L_x^2$, i.e. for any $s < t$ the measure $\mu_{s,t}^w$ can be identified with a function in L^2 , which is exactly $\ell_{s,t}^w$. Moreover $\mu_{s,t}^w$ is a positive measure with total variation $\|\mu_{s,t}^w\|_{\mathcal{M}} = t-s$, which implies that $\ell_{s,t}^w \in L_x^1$ with $\|\ell_{s,t}^w\|_{L^1} = t-s$. From this follows i.; ii. and the first part of iii. follow similarly by using the embeddings $\mathcal{FL}^{\rho,\infty} \hookrightarrow \mathcal{FL}^{0,1} \hookrightarrow C_b^0$, valid for $\rho > d$, and $\mathcal{FL}^{\rho,\infty} \hookrightarrow \mathcal{FL}^{\rho-d/2,2} \hookrightarrow W_x^{k,2}$, valid for $\rho - d/2 > k$. The second half of iii. follows from the fact that if $w \in L^\infty$, then $\mu_{s,t}^w$ is supported on $B_{\|w\|_{L^\infty}}$ and so we have the estimate $\|\ell_{s,t}^w\|_{W_x^{k,1}} \lesssim \|w\|_{L^\infty}^{d/2} \|\ell_{s,t}^w\|_{W_x^{k,2}}$. Finally statement iv. can be deduced from a combination of the previous estimate, the general Young-type inequality

$$\|f * g\|_{B_{p,q}^{\alpha+k}} \lesssim \|f\|_{B_{p,q}^\alpha} \|g\|_{W^{k,1}},$$

the identity $T_{s,t}^w b = \tilde{\mu}_{s,t}^w * b$ and the estimate $\|\mu_{s,t}^w\|_{W^{k,1}} \lesssim \|w\|_{L^\infty}^{d/2} \|\mu_{s,t}^w\|_{\mathcal{FL}^{\rho,\infty}} \lesssim |t-s|^\gamma \|\mu^w\|_{C^\gamma \mathcal{FL}^{\rho,\infty}}$. \square

Remark 22. We stated for simplicity Points iii. and iv. of Lemma 21 for $k \in \mathbb{N}$, so that the definition of $W^{k,1}$ is classical and unambiguous. However, up to coming up with a suitable concept for $W^{k,1}$ for $k \in (0, \infty) \setminus \mathbb{N}$ (e.g. replacing it with $B_{1,1}^k$), one can immediately extend the statements to the case of general $k \in \mathbb{R}_{\geq 0}$, since Fourier–Lebesgue embedding $\mathcal{FL}^\rho \hookrightarrow W^{k,1}$ and Young-type inequality for convolutions still hold.

The reader might wonder if the restriction $\rho > d/2$ appearing in Lemma 21 can be weakened, given that it does not appear when dealing with the scales of spaces $\mathcal{FL}^{\alpha,p}$ as in Lemma 19, nor in more probabilistic \mathbb{P} -a.s. statements where b is fixed and w is sampled as a stochastic process (cf. the results from [10, 34]). The next statement, which can be regarded as a useful rewriting of Remark 3.5 from [39], strongly suggests this not to be possible in general; indeed if the averaging operator T^w acts in a regularising way over the Besov–Hölder scales $B_{\infty,\infty}^\alpha$, then w must necessarily enjoy limited regularity.

Lemma 23. Let $w \in C^\delta([0, T]; \mathbb{R}^d)$ for some $\delta \in (0, 1]$; suppose that there exist $\alpha \in \mathbb{R}$, $\varepsilon > 0$ s.t. $T_{0,T}^w \in \mathcal{L}(B_{\infty,\infty}^\alpha, B_{\infty,\infty}^{\alpha+\varepsilon})$. Then it must hold $\delta \leq 1/d$.

Proof. Without loss of generality we can assume $\varepsilon \leq 1$. Observe that, since averaging and convolutions commute, i.e. $T^w(K * b) = K * (T^w b)$ (cf. Lemma 3.3 from [34]), if the hypothesis holds for *some* $\alpha \in \mathbb{R}$, then it holds for *all* $\alpha \in \mathbb{R}$ (one can take for instance $K = \Delta_j$ to be Littlewood–Paley blocks).

Next, choose $\alpha = \varepsilon/2$ and $p = 2d/\varepsilon$, so that we have the embedding $L^p \hookrightarrow B_{\infty,\infty}^{-\varepsilon/2}$, and observe that for any $f \in \mathcal{S}$ it holds

$$|\langle f, \tilde{\mu}_{0,T}^w \rangle| = |\langle f * \tilde{\mu}_{0,T}^w \rangle(0)| \lesssim \|f * \tilde{\mu}_{0,T}^w\|_{B_{\infty,\infty}^{\varepsilon/2}} = \|T_{0,T}^w f\|_{B_{\infty,\infty}^{\varepsilon/2}} \lesssim \|f\|_{B_{\infty,\infty}^{-\varepsilon/2}} \lesssim \|f\|_{L^p};$$

by density the inequality extends to all $f \in L^p$ and by duality we can conclude that there exists $g \in L^{p'}$, p' being the conjugate exponent to p , such that $\langle f, \tilde{\mu}_{0,T}^w \rangle = \langle f, \tilde{g} \rangle$ for all $f \in L^p$. This necessarily implies that $\mu_{0,T}^w(dx) = g(x)dx$; since w is continuous, $\mu_{0,T}^w$ is compactly supported, therefore it holds $g = \ell_{0,T}^w \in L^1 \cap L^{p'}$. Overall we have concluded that $\mu_{0,T}^w$ admits an occupation density $\ell_{0,T}^w \in L^1$.

Since $\mu_{0,T}^w(w([0,T])) = T$ and $\mu_{0,T}^w \ll \lambda$, where λ is the Lebesgue measure on \mathbb{R}^d , the set $w([0,T])$ must have positive Lebesgue measure; by standard facts from geometric measure theory, $w([0,T])$ then has Hausdorff dimension d . The conclusion then follows from the relation $d = \dim_H(w([0,T])) \leq \delta^{-1}$ (cf. Proposition 3.3 (a) from [23]). \square

Slightly anticipating upcoming results, it is interesting to observe that the restriction $\delta \leq 1/d$ matches almost exactly the sufficient condition $\rho > d/2$ given in Lemma 21-i. for the local time to exist, once we combine it with the fact (from Theorem 32 below) that a.e. $w \in C_t^\delta$ is ρ -irregular for any $\rho < 1/(2\delta)$.

Remark 24. It follows from Lemma 23 and the upcoming Theorem 32 that one can find many examples of paths $w : [0, T] \rightarrow \mathbb{R}^d$ whose averaging $T^w b$ has a regularizing effect when acting on the scales $\mathcal{FL}^{\alpha,p}$, but none on the scales $B_{\infty,\infty}^\alpha$. This issue boils down to the fact that in general convolutional operators with polynomially decaying Fourier transform are not necessarily nice Fourier multipliers, with strikingly similar examples coming from dispersive PDEs. Indeed, if one considers the linear wave PDE on \mathbb{R}^d

$$\partial_t^2 u = \Delta u, \quad u|_{t=0} = 0, \quad \partial_t u|_{t=0} = f,$$

then the solution at time $t > 0$ is given by the linear operator

$$T_t f = \frac{\sin(t|\nabla|)}{|\nabla|} f,$$

in the sense that T_t acts in Fourier space by multiplying \hat{f} by $\sin(t|\xi|)/|\xi|$. It is then clear that T_t maps H^s into H^{s+1} for any $s \in \mathbb{R}$; but if one tries to have a similar result in L^p -based results, this is not true in general, with explicit counterexamples given in [53]. What is more, when $d = 3$, T_t has the alternative harmonic mean representation $T_t f = \mu_t * f$, where μ_t is the unit measure on the sphere $S_t = \{x \in \mathbb{R}^d : |x| = t\}$, which is singular w.r.t. the Lebesgue measure; in particular, a similar argument to the one from Lemma 23 readily implies that T_t cannot map $B_{\infty,\infty}^\alpha$ into $B_{\infty,\infty}^{\alpha+\varepsilon}$ for any $\alpha \in \mathbb{R}$ and $\varepsilon > 0$. Roughly speaking T_t here behaves exactly like T_t^w would, in the case where the path w is 1-irregular but also δ -Hölder continuous with $\delta > 1/d$ (e.g. for $d = 3$, typical realization of fBm with $H \in (1/3, 1/2)$ would work by virtue of the upcoming Theorem 31).

In general, as pointed out in Remark 7 of [34], the operator T^w cannot regularise time-dependent fields $b = b(t, x)$, at least not uniformly in all $b \in C_t^0 E$ for suitable Banach spaces E . Intuitively, the reason is that the oscillations in time of b could compensate the oscillations of w and limit the regularisation effect. However if b is required to be sufficiently regular in t , it is still possible to obtain a regularisation effect, as we are going to show now.

Lemma 25. Let $w \in L^\infty$ be (γ, ρ) -irregular, $b \in C_t^\beta \mathcal{FL}^{\alpha,p}$ with $\beta > 1 - \gamma$. Then $T^w b \in C_t^\gamma \mathcal{FL}^{\alpha+\rho,p}$ and there exists a constant $C = C(\gamma + \beta, T) > 0$ such that

$$\|T_{s,t}^w b\|_{\mathcal{FL}^{\alpha+\rho,p}} \leq C \|b\|_{C_t^\beta \mathcal{FL}^{\alpha,p}} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}} |t - s|^\gamma \quad \text{for all } 0 \leq s \leq t \leq T. \quad (9)$$

Namely, the linear map $T^w : C_t^\beta \mathcal{FL}^{\alpha,p} \rightarrow C_t^\gamma \mathcal{FL}^{\alpha+\rho,p}$ is bounded with constant $C \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}$.

Proof. Let us first assume that b is a smooth function. In this case for any $[s, t] \subset [0, T]$ and any sequence Π of partitions of $[s, t]$ with infinitesimal mesh, it holds

$$\begin{aligned} T_{s,t}^w b(x) &= \int_s^t b(r, x + w_r) dr = \lim_{|\Pi| \rightarrow 0} \sum_i \int_{t_i}^{t_{i+1}} b(t_i, x + w_r) dr \\ &= \lim_{|\Pi| \rightarrow 0} \sum_i T_{t_i, t_{i+1}}^w [b(t_i, \cdot)](x) = \lim_{|\Pi| \rightarrow 0} \sum_i (b_{t_i} * \tilde{\mu}_{t_i, t_{i+1}}^w)(x) \end{aligned}$$

which implies the functional equality

$$T_{s,t}^w b = \lim_{|\Pi| \rightarrow 0} \sum_i b_{t_i} * \tilde{\mu}_{t_i, t_{i+1}}^w.$$

Since $b \in C_t^\beta \mathcal{FL}^{\alpha,p}$, $\mu^w \in C_t^\gamma \mathcal{FL}^{\rho,\infty}$, $\gamma + \beta > 1$ and the map $*$: $(f, g) \mapsto f * g$ is bilinear and bounded from $\mathcal{FL}^{\alpha,p} \times \mathcal{FL}^{\rho,\infty}$ into $\mathcal{FL}^{\alpha+\rho,p}$, it follows from Young integration in Banach spaces (see Appendix A.1) that

$$T_{s,t}^w b = \int_s^t b_r * \tilde{\mu}_{dr}^w,$$

as well as estimate (9), since $\|\tilde{\mu}^w\|_{C_t^\gamma \mathcal{FL}^{\rho,\infty}} = \|\mu^w\|_{C_t^\gamma \mathcal{FL}^{\rho,\infty}} \sim \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}$. The case of general b follows from approximation procedures: given $b \in C_t^\beta \mathcal{FL}^{\alpha,p}$, we can find a sequence b^n of smooth functions such that $\|b^n\|_{C_t^\beta \mathcal{FL}^{\alpha,p}} \leq \|b\|_{C_t^\beta \mathcal{FL}^{\alpha,p}}$ and that for any $\varepsilon > 0$ $b^n \rightarrow b$ locally in $C_t^{\beta-\varepsilon} \mathcal{FL}^{\alpha-\varepsilon,p}$; by properties of averaging, for any $s < t$ $T_{s,t}^w b^n$ converges to $T_{s,t}^w b$ weakly-* in $\mathcal{FL}^{\alpha,p}$. The conclusion then follows from taking the liminf as $n \rightarrow \infty$ on both sides of (9) applied to b^n and using the Fatou property of weak-* convergence. \square

Remark 26. *It is clear that the proof can be readapted in a more general setting: given E, F, G function spaces such that $*$: $(f, g) \mapsto f * g$ is a bilinear bounded map from $E \times F$ into G , if $\mu^w \in C_t^\gamma F$ and $\beta > 1 - \gamma$, then $T^w : C_t^\beta E \rightarrow C_t^\gamma G$ is a linear bounded map. This can be applied in combination with Lemma 21, obtaining regularising effects of T^w when E and G are taken in suitable Besov scales.*

2.4 Fractional Brownian motion and local-nondeterminism

The main tool in order to establish our prevalence results will be a family of measures associated to locally nondeterministic Gaussian processes. The main examples from this family are the laws μ^H of fractional Brownian motions (fBm) of Hurst parameter $H \in (0, 1)$. General references on fBm are [49] and [55].

A one dimensional fBm $(W_t^H)_{t \geq 0}$ of Hurst parameter $H \in (0, 1)$ is a centred Gaussian process with covariance

$$\mathbb{E}[W_t^H W_s^H] = \frac{1}{2}(|t|^{2H} + |s|^{2H} - |t - s|^{2H}).$$

Up to a multiplicative constant, it is the unique centred Gaussian process with stationary increments and H -self-similarity, i.e. with law invariant under the scaling $\tilde{W}_\lambda^H := \lambda^{-H} W_\lambda^H$ for any $\lambda > 0$.

When $H = 1/2$, it coincides with standard Brownian motion, while for $H \neq 1/2$ it is not a semimartingale nor a Markov process. However it shares many properties of Brownian motion, for instance fBm trajectories are \mathbb{P} -a.s. α -Hölder continuous for any $\alpha < H$ and nowhere α -Hölder continuous for any $\alpha \geq H$. A d -dimensional fBm W^H of Hurst parameter $H \in (0, 1)$ is an \mathbb{R}^d -valued Gaussian process with components given by independent one dimensional fBms.

Given a two-sided Brownian motion $(B_t)_{t \in \mathbb{R}}$, a fBm of parameter $H \neq 1/2$ can be constructed by the formula

$$W_t^H = c_H \int_{-\infty}^t [(t-r)_+^{H-1/2} - (-r)_+^{H-1/2}] dB_r \quad (10)$$

where $c_H = \Gamma(H + 1/2)^{-1}$ is a suitable renormalising constant and the integral is in the Itô sense. As a consequence, for any $0 \leq s < t$, the variable W_t^H decomposes into the sum $W_t^H = W_{s,t}^{1,H} + W_{s,t}^{2,H}$, where

$$W_{s,t}^{1,H} := c_H \int_s^t (t-r)^{H-1/2} dB_r, \quad W_{s,t}^{2,H} := \mathbb{E}[W_t^H | \mathcal{F}_s] = c_H \int_{-\infty}^s [(t-r)_+^{H-1/2} - (-r)_+^{H-1/2}] dB_r;$$

specifically, $W_{s,t}^{2,H}$ is \mathcal{F}_s -measurable, while $W_{s,t}^{1,H}$ is Gaussian, independent of \mathcal{F}_s and with variance

$$\text{Var}(W_{s,t}^{1,H}) = \tilde{c}_H |t - s|^{2H},$$

where $\tilde{c}_H = c_H^2/(2H)$. In particular this implies that

$$\text{Var}(W_t^H | \sigma(W_r^H, r \leq s)) \geq \text{Var}(W_t^H | \mathcal{F}_s) = \text{Var}(W_{s,t}^{1,H}) = \tilde{c}_H |t - s|^{2H}. \quad (11)$$

Equation (11) is known in the literature as a local-nondeterminism (LND) property. LND was first introduced by Berman in [7] in order to analyse the properties of the occupation measure (more precisely the local time) of Gaussian processes. Loosely speaking, it means that for any $s < t$, the increment $W_t^H - W_s^H$ contains a part which is independent of the the history of the path W^H up to time s and therefore makes the path W^H “intrinsically chaotic”.

There is now a huge literature on local-nondeterminism and several alternative definitions, which are not in general equivalent, see [64] for a survey; here we identify three types of LND which are closely tied with ρ -irregularity and exponential irregularity of sample paths of Gaussian processes. They will play a major role in the proofs respectively of Sections 4.2 and 4.4.

Definition 27. Let $(X_t)_{t \in [0, T]}$ be an \mathbb{R}^d -valued separable Gaussian process adapted to a given filtration \mathcal{F}_t . We say that X is strongly locally nondeterministic with parameter $\beta > 0$, X is β -SLND for short, if there exists $\delta > 0$ s.t.

$$\text{Var}(X_t | \mathcal{F}_s) \gtrsim |t - s|^{2\beta} I_d \quad \text{uniformly in } s, t \text{ such that } 0 < t - s < \delta. \quad (12)$$

Definition 28. Let $(X_t)_{t \in [0, T]}$ be an \mathbb{R}^d -valued separable Gaussian process adapted to a given filtration \mathcal{F}_t . We say that X is locally nondeterministic with parameter $\beta > 0$, X is β -LND for short, if for every integer $n \geq 2$ there exists positive constants c_n and δ_n such that

$$\text{Var} \left(\sum_{k=1}^n v_k \cdot (X_{t_{k+1}} - X_{t_k}) \right) \geq c_n \sum_{k=1}^n \text{Var}(v_k \cdot (X_{t_{k+1}} - X_{t_k})) \geq c_n \sum_{k=1}^n |v_k|^2 |t_{k+1} - t_k|^{2\beta} \quad (13)$$

for all ordered points $t_1 < t_2 < \dots < t_n$ with $t_n - t_1 < \delta_n$ and $v_k \in \mathbb{R}^m$.

Properly speaking, in the terminology of [64], Definition 27 is that of a one-sided strong local non-determinism, but we have preferred to adopt the terminology β -SLND for simplicity. It follows from Remark 2.3 from [64] that Definition 27 is strictly stronger than Definition 28, namely any β -SLND Gaussian process is also β -LND, while the converse is not true. Equation (11) implies that W^H is H -SLND.

Definition 29. Let $\{X_t\}_{t \in [0, T]}$ be an \mathbb{R}^d -valued separable Gaussian process adapted to a given filtration \mathcal{F}_t . We say that X is exponentially locally nondeterministic with parameter $\beta > 0$, X is β -eSLND for short, if there exists $\delta > 0$ s.t.

$$\text{Var}(X_t | \mathcal{F}_s) \gtrsim |\log(t - s)|^{-\beta} I_d \quad \text{uniformly in } s, t \text{ such that } 0 < t - s < \delta. \quad (14)$$

Remark 30. Definitions 27, 28 and 29 only involve the variance (resp. conditional variance) of the process X and are independent of its mean. This implies that they are all properties invariant under deterministic perturbations, namely if X is a β -(e)(S)LND process and f is a given measurable function, then $X + f$ is still β -(e)(S)LND. This can be interpreted as the chaoticity represented by local non-determinism being too strong to be disrupted by a deterministic additive perturbations; this fundamental feature will allow us to prove prevalence of ρ -irregularity and exponential irregularity.

We conclude this section by recalling the following result.

Theorem 31 (Theorem 1.4 from [10]). Let $H \in (0, 1)$ and denote by μ^H the law of a d -dimensional fBm of Hurst parameter H . Then for any $\rho < (2H)^{-1}$ there exists $\gamma > 1/2$ such that

$$\mu^H(w \in C^0([0, T]; \mathbb{R}^d) \mid w \text{ is } (\gamma, \rho)\text{-irregular}) = 1.$$

Let us point out that, combined with Lemma 7, this last theorem implies the existence of continuous (γ, ρ) -irregular functions for any choice of $\gamma \in (0, 1)$ and $\rho < \infty$.

3 Main results

3.1 Statements

We are now ready to present the main result of this paper. In the next statement, the space C_{slow}^∞ denotes the collection of all infinitely differentiable functions $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$ such that ψ and all its derivatives grow at most polynomially at infinity; instead we set $C_b^\infty = \cap_n C_b^n$ as the set of bounded smooth functions with bounded derivatives.

Theorem 32. *It holds that:*

- i. For any $\delta \in (0, \infty)$, almost every $\varphi \in C_t^\delta$ is ρ -irregular for any $\rho < (2\delta)^{-1}$. If $\delta \geq 1$, then in addition for any $k < \delta$, $D^{(k)}\varphi$ is ρ -irregular for any $\rho < (2(\delta - k))^{-1}$.*
- ii. Almost every $\varphi \in C_t^0$ is ρ -irregular for any $\rho < \infty$; in particular, its occupation measures $(\mu_{s,t}^\varphi) \subset C_c^\infty$ and its averaging operator T^w maps \mathcal{S}' into C_{slow}^∞ , as well as $B_{p,q}^s$ into C_b^∞ for any $s \in \mathbb{R}$, $p, q \in [1, \infty]$.*
- iii. For any $p \in [1, \infty)$, almost every $\varphi \in L_t^p$ is exponentially irregular; in particular it is Carathéodory and its occupation measures $(\mu_{s,t}^\varphi)$ are analytic.*

We avoid providing very similar statements, but like Point *ii.* above, Point *i.* in combinations with Lemmata 19, 21 and 25 provides several other prevalence statements in C_t^δ regarding the regularity of ℓ^w and the regularising effect of T^w acting on suitable function spaces.

The cornerstone in the proof of Theorem 32 is the following probabilistic result.

Theorem 33. *Let X be a continuous β -(S)LND Gaussian process with $\beta \in (0, \infty)$; then for any $\rho < (2\beta)^{-1}$ there exists $\gamma = \gamma(\rho, \beta) > 1/2$ such that X is (γ, ρ) -irregular with probability 1. Let X be a β -eSLND Gaussian with measurable, L^2 -integrable trajectories and $\beta \in (0, 1]$; then X is exponentially irregular with probability 1.*

The proof of Theorem 33 is given in Section 4. For simplicity, in the statement above no distinction between β -SLND and β -LND processes appears; in Sections 4.2 and 4.4 more refined results are given, so that the differences in the two cases become more evident.

Theorem 32 combined with geometric considerations implies also the following prevalence result.

Theorem 34. *Let $\delta \in [0, 1)$. The following hold:*

- i. Almost every $\varphi \in C_t^\delta$ has the property that*

$$\dim_F(\varphi([s, t])) = \dim_H(\varphi([s, t])) = \frac{1}{\delta} \wedge d \quad \forall [s, t] \subset [0, T].$$

- ii. If additionally $\delta < 1/(2d)$, then almost every $\varphi \in C_t^\delta$ has the property that, for all $[s, t] \subset \mathbb{R}^d$, $\varphi([s, t])$ contains an open set.*

In particular, for all $\delta \in [0, 1)$, the image of almost every function $\varphi \in C_t^\delta$ is a Salem set.

In the statement above, \dim_F and \dim_H denote respectively the Fourier and Hausdorff dimensions. The proof is given in Section 5.1, where also the definitions of \dim_F , \dim_H and Salem sets are recalled.

Remark 35. *Combining Theorem 33 with the reasoning from Theorem 34 gives an alternative proof of the fact that Fourier and Hausdorff dimension of images of intervals under a d -dimensional fBm are \mathbb{P} -a.s. $H^{-1} \wedge d$; see Theorem 4.29-b) from [48] (for $H = 1/2$) and Section 18 from [43] (for general $H \in (0, 1)$) for more classical proofs.*

The study of analytic properties of (γ, ρ) -irregular paths allows to show that, as the name suggests, they have a highly oscillatory behaviour; this can be related to other notions of roughness already existing in the literature.

Theorem 36. *Let w be (γ, ρ) -irregular, $\delta_{\gamma, \rho}^*$ defined as in (3); assume $\delta_{\gamma, \rho}^* \in (0, 1)$. Then for any $\delta > \delta_{\gamma, \rho}^*$, w is nowhere δ -Hölder continuous and it has infinite modulus of δ -Hölder roughness; for any $p < (\delta_{\gamma, \rho}^*)^{-1}$ and any interval $[s, t] \subset [0, T]$, w has infinite p -variation on $[s, t]$.*

The proof is given in Section 5.2, where also the concept of modulus of δ -Hölder roughness is recalled.

A relation between (γ, ρ) -irregularity and failure of δ -Hölder regularity was already observed for $d = 1$ in [12, Theorem 1.5]; our Theorem 36 is significantly sharper, as it avoids the additional constraint $\delta > 1 - \gamma$ as therein, holds in any dimension d and provides a nowhere δ -Hölder continuity condition (in particular, regularity will fail to hold at *all* points $t_0 \in [0, T]$, not just *some* point as in [12, Theorem 1.5]). Furthermore, Theorem 69, on which the proof is based, gives a more precise bound on the Lebesgue density of the points around any t_0 where an approximate δ -Hölder condition fails.

Quite nicely, Theorem 36 provides an alternative proof of Hunt's original results from [41].

Corollary 37. *Let $\delta \in [0, 1)$, then almost every $\varphi \in C_t^\delta$ is nowhere $(\delta + \varepsilon)$ -Hölder for any $\varepsilon > 0$.*

Proof. By Theorem 32, almost every $\varphi \in C_t^\delta$ has the following property: for any $\rho < 1/(2\delta)$, there exists $\gamma > 1/2$ such that φ is (γ, ρ) -irregular. It holds

$$\delta_{\gamma, \rho}^* = \frac{1 - \gamma}{\rho} < \frac{1}{2\rho}$$

which implies by Theorem 36 that any such function is nowhere $\tilde{\delta}$ -Hölder for any $\tilde{\delta} > 1/(2\rho)$. Taking $\rho = 1/(2\delta + 2\varepsilon)$ the conclusion follows. \square

In the current work we are also able to solve a problem left open in [10] and [12], which amounts to showing that if φ is ρ -irregular and $\psi \in C_t^\infty$, then $\varphi + \psi$ is ρ -irregular. We give a positive answer, up to strengthening the notion of ρ -irregularity.

Theorem 38. *Let φ be strongly ρ -irregular, then for any $\psi \in C_t^\infty$, $\varphi + \psi$ is $\tilde{\rho}$ -irregular for any $\tilde{\rho} < \rho$. For $\delta \in [0, 1)$, almost every $\varphi \in C_t^\delta$ is strongly ρ -irregular for any $\rho < (2\delta)^{-1}$.*

The proof is given in Section 5.4, where the concept of strong ρ -irregularity is introduced. It is a very technical concept, which is the reason why we have preferred not to introduce it in Section 2.2. We hope in the future simpler criteria will be developed for establishing the invariance of ρ -irregularity under additive perturbations.

Remark 39. *Let us discuss already here the optimality of Theorem 32, in light of Theorems 33-38 and the results presented in the rest of the paper.*

1. For $\delta \in (0, 1)$, optimality of the condition $\rho < (2\delta)^{-1}$ follows from Theorem 36: it must hold $\delta \leq \delta_{\gamma, \rho}^* < (2\rho)^{-1}$ in light of the condition $\gamma > 1/2$.
2. By applying Lemma 7, in the case $\delta = 0$ the result can be strengthened to the fact that almost every $\varphi \in C_t^0$ is (γ, ρ) -irregular for any $\gamma < 1$ and any $\rho < \infty$. Time regularity cannot be improved to ℓ^φ being differentiable in time, since we know that (in the weak sense)

$$\frac{d}{ds} \ell(s, \cdot)|_{s=t} = \delta_{\varphi_t} \quad \forall t \in [0, T].$$

Moreover $\{\ell_{s,t}^\varphi\} \subset C_c^\infty$ cannot be improved to $\ell_{s,t}^\varphi$ being analytic, since this would imply that $\varphi([s, t])$ is an unbounded set, thus violate continuity of φ .

3. One might wonder if, since by Lemma 7 we can always raise the value of γ by lowering the one of ρ , we can also do the opposite; in particular if, without imposing the restriction $\gamma > 1/2$, we can find functions $\varphi \in C^\delta$ which are (γ, ρ) -irregular for a pair (γ, ρ) satisfying $\delta \leq \delta_{\gamma, \rho}^*$ but also $\rho > (2\delta)^{-1}$. In the case $\delta > 1/d$, $\delta \in (0, 1]$ (thus implying $d \geq 2$) this possibility is ruled out by reasoning with Fourier dimensions (see Section 5.1 for more details), since it must hold

$$2\rho = \min(d, 2\rho) \leq \dim_F(\varphi([s, t])) \leq \dim_H(\varphi([s, t])) \leq \delta^{-1}$$

independently of the value of γ .

4. If $\delta \leq 1/d$, the problem posed above is currently open. The only information we are able to provide in this case is that for $d = 1$, by Proposition 8 there exist indeed C^1 functions (for instance $\varphi(t) = t$) which are $(\gamma, 1 - \gamma)$ -irregular for any $\gamma \in (0, 1)$.
5. Although Point 4. is open in terms of generic $\varphi \in C^\delta$, we are at least able to establish that fBm paths do not have this property, see Lemma 75 from Section 5.2.

We conclude this section with a proof of Theorem 32, based on the other results contained in the paper.

Proof of Theorem 32. We have already seen in Lemma 10 that both ρ -irregularity and exponential irregularity properties define Borel measurable sets in all of the above function spaces, so we only need to provide suitable measures “witnessing” their prevalence.

Let us start from the case $\delta \in (0, 1)$. Let μ^H to be the law on C_t^δ of a fBm W^H of parameter $H = \delta + \varepsilon$ for some $\varepsilon > 0$, which is tight on C_t^δ ; let $\varphi \in C_t^\delta$ be fixed. The process W^H is H -SLND and by Remark 30 so is $\varphi + W^H$; then by Theorem 33 it holds

$$\mu^H(\varphi + w \text{ is } \rho\text{-irregular for any } \rho < 1/(2H)) = \mathbb{P}(\varphi + W^H \text{ is } \rho\text{-irregular for any } \rho < 1/(2H)) = 1.$$

This implies that almost every $\varphi \in C_t^\delta$ is ρ -irregular for any $\rho < 1/(2\delta + 2\varepsilon)$; taking a sequence $\varepsilon_n \downarrow 0$ and using the fact that countable intersection of prevalent sets is still prevalent, we obtain the conclusion in this case.

Consider now the case $\delta \in [n, n + 1)$, $n \geq 1$; set $\delta = n + \theta$, $\theta \in [0, 1)$. Denote by μ^{n+H} the law of the process Y obtained by integrating n times an fBm, which is discussed at Point *iv.* of the list of examples from Section 4.2. Choose $H > \theta$, so that μ^{n+H} is tight in C_t^δ . Now fix $\varphi \in C_t^\delta$; by the discussion at Point *iv.*, the process Y is $(n + H)$ -SLND with $D^{(k)}Y$ being $(n + H - k)$ -SLND for any $k \in \{1, \dots, n\}$ therefore $Y + \varphi$ and $D^{(k)}(Y + \varphi) = D^{(k)}Y + D^{(k)}\varphi$ have the same properties by Remark 30. By applying Theorem 33 and arguing as in the previous point, taking a sequence $H_n \downarrow \delta$, the proof of claim *i.* is complete.

The first part of claim *ii.* is identical, relying this time on the fact that we can take any $H > 0$ and we obtain ρ -irregularity for any $\rho < 1/(2H)$, together with the property that countable intersection of prevalent sets is prevalent; the fact that $(\mu_{s,t}^\varphi) \subset C_c^\infty$ readily follows from $\text{supp } \mu_{s,t}^\varphi \subset B_{\|\varphi\|_{C^0}}$ and the Fourier–Lebesgue embeddings $\mathcal{FL}^{\rho,\infty} \hookrightarrow C^{\rho-d/2-\varepsilon}$. Concerning the second part of claim *ii.*, observing that $\tilde{\mu}_{s,t}^\varphi \in C_c^\infty \hookrightarrow \mathcal{S}$, we can combine Lemma 18 with Proposition 9.10 a) from [27] to deduce that T^φ maps \mathcal{S}' into C_{slow}^∞ ; the case of $B_{p,q}^s$ can be treated similarly, this time first embedding it into $B_{\infty,\infty}^{s-d/p} = C^{s-d/p}$ and then use standard Young-type estimates for convolutions.

The second part of claim *iii.* follows from Lemma 13, once we have shown the first part. For any $\beta \in (0, 1)$, denote by μ^β the law of (a measurable version of) the process X constructed in Proposition 55, which is β -eSLND. Let $\varphi \in L_t^p$; we can require φ to be an actual measurable function in its equivalence class, since the property of exponential irregularity does not depend on the chosen representative. By Remark 30, the process $\varphi + X$ is also β -eSLND and so by Theorem 33

$$\mu^\beta(\varphi + w \text{ is exponentially irregular}) = \mathbb{P}(\varphi + X \text{ is exponentially irregular}) = 1$$

which implies the conclusion. \square

3.2 Applications to regularisation by noise for ODEs and PDEs

In this section we show how our main result, Theorem 32, can be combined with already existing results in the literature, obtaining results on the regularising effect of almost every $w \in C_t^\delta$ on ODEs and PDEs.

As mentioned before, (γ, ρ) -irregularity is closely related to the regularising properties of the averaging operator T^w . Moreover, in order to develop a good solution theory for the perturbed ODE

$$\frac{d}{dt}x_t = b(x_t) + \frac{d}{dt}w_t \tag{15}$$

it is enough to have good regularity properties for $T^w b$. Results in this direction have been obtained in [10] and [34]. They can be combined with Lemma 19 and Theorem 32 as follows.

Theorem 40. *Suppose that either $T^w b \in C_t^\gamma C_x^2$ or $b \in L_{t,x}^\infty$ and $T^w b \in C_t^\gamma C_x^{3/2}$, for some $\gamma > 1/2$. Then for any $x_0 \in \mathbb{R}^d$ there exists a unique global solution to (15); moreover the ODE admits a locally Lipschitz flow. If, for some $n \geq 1$, $T^w b \in C_t^\gamma C_x^{n+1}$, or $b \in L_{t,x}^\infty$ and $T^w b \in C_t^\gamma C_x^{n+1/2}$, then the flow is locally C^n .*

Let $\delta \in [0, 1)$, then almost every $w \in C_t^\delta$ is such that for any $\alpha \in \mathbb{R}$, the following hold:

- i. if $\alpha > \max\{3/2 - (2\delta)^{-1}, 0\}$ or $\alpha > 2 - (2\delta)^{-1}$, then for any $b \in \mathcal{FL}^{\alpha,1}$ the perturbed ODE (15) is well posed and admits a locally Lipschitz flow;*
- ii. if $\alpha > \max\{n + 1/2 - (2\delta)^{-1}, 0\}$ or $\alpha > n + 1 - (2\delta)^{-1}$, for some $n \geq 1$, then for any $b \in \mathcal{FL}^{\alpha,1}$ the flow is locally C^n .*

In particular for almost every $\varphi \in C_t^0$, for any $\alpha \in \mathbb{R}$ and any $b \in \mathcal{FL}^{\alpha,1}$, the ODE (15) is well-posed and admits a C^∞ flow.

Proof. The first part of the statement comes from Remark 4.7 and Theorem 4.33 from [34].

Point *i.* then follows from a combination of Theorem 32-*i.*, Lemma 19 and the Fourier–Lebesgue embeddings $\mathcal{FL}^{\beta,1} \hookrightarrow C_b^\beta$ for $\beta \geq 0$. For instance in the case $\alpha > 2 - (2\delta)^{-1}$, a.e. $w \in C_t^\delta$ is (γ, ρ) -irregular for $\rho = 2 - \alpha < (2\delta)^{-1}$ and some $\gamma > 1/2$, implying that for any $b \in \mathcal{FL}^{\alpha,1}$ it holds $T^w b \in C_t^\gamma \mathcal{FL}^{\alpha+\rho,1} \hookrightarrow C_t^\gamma C_x^2$, so that we can apply the first part of the theorem. The case $\alpha > \max\{3/2 - (2\delta)^{-1}, 0\}$ is similar, only in this case by embeddings we have $b \in C_b^0$ to begin with, and our assumptions allow to verify that $T^w b \in C_t^\gamma C_x^{3/2}$. Point *ii.* can be established in a similar manner and the final claim comes from Theorem 32-*ii.* \square

The last part of the statement is an instance of the ∞ -regularising effect of generic continuous functions on (15). Applying Lemma 21, analogue statements can be given replacing Fourier–Lebesgue $\mathcal{FL}^{\alpha,1}$ with other scales like $B_{p,q}^s$ spaces. In addition, we can also consider time-dependent fields b , for instance such that $b \in C_t^{1/2} \mathcal{FL}^{\alpha,1}$, thanks to Lemma 25.

The theory developed for solving (15) can be also successfully applied to the study first order linear PDEs of the form

$$\partial_t u + b \cdot \nabla u + cu + \frac{dw}{dt} \cdot \nabla u = 0. \quad (16)$$

In general, if w is (γ, ρ) -irregular then it is nowhere differentiable, but assume for the moment $w \in C^1$. Setting $v(t, x) = u(t, x + w_t)$, $\tilde{b}(t, x) = b(t, x + w_t)$, $\tilde{c}(t, x) = c(t, x + w_t)$, u would be a classical solution of (16) if and only if v solved

$$\partial_t v + \tilde{b} \cdot \nabla v + \tilde{c}v = 0, \quad (17)$$

which is meaningful (under suitable assumptions on \tilde{b} , \tilde{c}) even when $w \in C^0$. For this reason we will say that u is a solution to (16) if and only if v solves (17), without properly defining the meaning of equation (16). Let us mention that in the context of rough path theory, if w admits a rough lift, then it is possible to give meaning to (16) and show the equivalence with (17), see [8]. If b and c are distributional but $T^w b$ and $T^w c$ have good regularity in $C_t^\gamma C_x^\beta$ -spaces, then it is possible to give meaning to (17) with Young integrals; in this case equation (17) (and consequently (16)) admit a natural interpretation in the sense of distributions, thus we will talk of weak solutions, cf. Definitions 5.5 and 5.13 from [34]; we refrain for giving here more details, for which we refer the interested reader to Section 5 from [34]. In the next statement, $\mathcal{M}_x = \mathcal{M}(\mathbb{R}^d)$ denotes the set of all finite Radon measures.

Theorem 41. *Let $\delta \in [0, 1)$ then almost every $w \in C_t^\delta$ is such that for any $\alpha \in \mathbb{R}$, the following hold:*

- i. if $\alpha > \max\{3/2 - (2\delta)^{-1}, 0\}$ or $\alpha > 2 - (2\delta)^{-1}$, then for any $b \in \mathcal{FL}^{\alpha,1}$ the transport PDE*

$$\partial_t u + b \cdot \nabla u + \frac{dw}{dt} \cdot \nabla u = 0$$

has a unique solution $u \in C_t^0 C_{\text{loc}}^1$ for any $u_0 \in C_x^2$;

ii. if for some $n \geq 1$, $\alpha > \max\{n + 1/2 - (2\delta)^{-1}, 0\}$ or $\alpha > n + 1 - (2\delta)^{-1}$, then for any $b \in \mathcal{FL}^{\alpha,1}$ and any $u_0 \in C_x^{n+1}$, the solution $u \in C_t^0 C_{\text{loc}}^n$;

iii. if $\alpha > \max\{3/2 - (2\delta)^{-1}, 0\}$ or $\alpha > 2 - (2\delta)^{-1}$, then for any $b \in \mathcal{FL}^{\alpha,1}$ the continuity equation

$$\partial_t u + \nabla \cdot (bu) + \frac{dw}{dt} \cdot \nabla u = 0$$

has a unique weak solution $u \in C_t^\gamma (C_x^1)^* \cap L_t^\infty \mathcal{M}_x$ for any $u_0 \in \mathcal{M}_x$.

Proof. It follows from the results of [34] that, if $b \in C_x^0$ and $T^w b \in C_t^\gamma C_x^{3/2}$ (cf. Proposition 5.1 therein) or $T^w b \in C_t^\gamma C_x^2$ (cf. Lemma 5.7 and Theorem 5.12) for some $\gamma > 1/2$, then existence and uniqueness of solutions to the transport PDE holds for any $u_0 \in C_x^2$ and we have the representation formula $u_t(x) = u_0(\phi_t^{-1}(x))$, where ϕ is the flow associated to the ODE (15). Arguing as in the proof of Theorem 40 it is easy to check that, under our assumptions on δ and α , a.e. $w \in C_t^\delta$ is ρ -irregular for the right range of parameters ρ needed to establish the required regularity of $T^w b$, proving Point i.

Point ii. follows from the representation formula for u and the higher regularity of the flow ϕ associated to the ODE, which comes from Theorem 40-ii.

Point iii. comes from similar reasonings, only this time applying the results from [34] for the continuity equation, which reduce again the problem to verify the regularity of $T^w b$; in particular one can invoke Proposition 5.3 therein in the case $b \in C_x^0$ and Lemma 5.15, Theorem 5.16 in the case of general b . \square

In the above examples, w enters the equation as a perturbation that can be reabsorbed by shifting the phase space, which is why the operator T^w appears. In the next examples instead w has the role of *modulating* a given group of transformations.

In the papers [12] and [14], the authors study the regularising properties of (γ, ρ) -irregular paths on non-linear dispersive PDEs of the general form

$$\frac{d}{dt} \varphi_t = A \varphi_t \frac{dw_t}{dt} + \mathcal{N}(\varphi_t), \quad (18)$$

where $w \in C_t^0$, $\varphi : D \rightarrow \mathbb{R}$ (or \mathbb{C}), A is a linear unbounded operator and \mathcal{N} is a nonlinearity (typically of polynomial type). Their cases of interest are:

1. (cubic NLS) Non-linear cubic Schrödinger, $D = \mathbb{T}^d$ or \mathbb{R}^d , $d = 1, 2$, $A = i\Delta$, $\mathcal{N}(\varphi) = i|\varphi|^2 \varphi$;
2. General NLS on $D = \mathbb{R}$ with $A = i\partial_x^2$, $\mathcal{N}(\varphi) = i|\varphi|^\mu \varphi$, $\mu \in (1, 4]$;
3. (dNLS) Non-linear derivative cubic Schrödinger on \mathbb{T} , $A = i\partial_x^2$, $\mathcal{N}(\varphi) = i\partial(|\varphi|^2 - \|\varphi\|_{L^2}^2) \varphi$;
4. (KdV) Korteweg–de Vries, $D = \mathbb{T}$ or \mathbb{R} , $A = \partial_x^3$, $\mathcal{N}(\varphi) = \partial_x(\varphi^2)$;
5. (mKdV) Modified Korteweg–de Vries, $D = \mathbb{T}$, $A = \partial_x^3$, $\mathcal{N}(\varphi) = \partial_x(\varphi^3 - 3\varphi\|\varphi\|_{L^2}^2)$.

From now on we will refer to these PDEs as *w-modulated equations*, e.g. *w-modulated cubic NLS*, *w-modulated KdV*, etc. In all cases 1.-5. above, although the original system would be of integrable nature, the presence of w doesn't allow to exploit any integrability features; moreover the group $\{e^{tA}\}_{t \in \mathbb{R}}$ associated to A acts isometrically on all H^α -spaces, thus doesn't provide a priori any regularisation.

In order to give meaning to (18), the authors adopt the mild formulation (which would be justified for $w \in C^1$ by the chain rule, but afterwards makes sense for any $w \in C^0$)

$$\varphi_t = U_t^w \varphi_0 + U_t^w \int_0^t (U_s^w)^{-1} \mathcal{N}(\varphi_s) ds$$

where $U_t^w = e^{w_t A}$; applying the change of variables $\psi_t = (U_t^w)^{-1} \varphi_t$, the equation becomes

$$\psi_t = \varphi_0 + \int_0^t (U_s^w)^{-1} \mathcal{N}(U_s^w \psi_s) ds. \quad (19)$$

It is finally possible to give meaning to (19), even in spaces H^α for which the original nonlinearity \mathcal{N} is not well defined, using the ρ -irregularity property to show that the operators $\{X_{s,t}\}_{s < t}$

$$X_{s,t}(\phi) = \int_s^t (U_r^w)^{-1} \mathcal{N}(U_r^w \phi) ds, \quad s < t$$

are continuous from H^α to itself (actually C^∞ since they are the monoid associated to an n -linear bounded operator). It is then imposed that φ is a solution to (18) if and only if the associated ψ solves (19). We refrain from giving further details on the topic and only point out that our Theorem 32 combined with their results give the following statements.

Theorem 42. *Let $\delta \in [0, 1)$. Then for almost every $w \in C^\delta([0, T])$, the w -modulated cubic NLS on \mathbb{T} and \mathbb{R} has a global solution in H^α for any $\alpha \geq 0$ and any $\varphi_0 \in H^\alpha$, which is unique in a suitable class; moreover the equation admits a locally Lipschitz continuous flow.*

Theorem 43. *Let $\delta \in [0, 2/3)$. Then:*

- i. For almost every $w \in C^\delta([0, T])$, the w -modulated KdV on \mathbb{T} has a unique local solution in H^α for any $\varphi_0 \in H^\alpha$ with $\alpha > -(2\delta)^{-1}$, which is global if $\alpha > -\min\{3/2, (4\delta)^{-1}\}$.*
- ii. For almost every $w \in C^\delta([0, T])$, the w -modulated KdV on \mathbb{R} has a unique local solution in H^α for any $\varphi_0 \in H^\alpha$ with $\alpha > -\min\{3/4, (2\delta)^{-1}\}$, which is global if $\alpha > -\min\{3/4, (4\delta)^{-1}\}$.*

Moreover for any $\delta \in [0, 1)$, for almost every $w \in C_t^\delta$, the w -modulated mKdV on \mathbb{T} has a unique local solution in H^α for any $\varphi_0 \in H^\alpha$ with $\alpha \geq 1/2$.

Proof of Theorems 42-43. Theorem 42 follows from a combination of Theorem 32-i) and Theorem 1.8 from [12]; indeed the condition $\delta < 1$ always ensures that a.e. $w \in C^\delta$ is ρ -irregular for some $\rho > 1/2$. Theorem 43 is instead implied by Theorem 32-i) and Theorems 1.6 from [14]. Indeed, Theorem 32-i) guarantees that a.e. $w \in C^\delta$ is (γ, ρ) -irregular, for γ arbitrarily close to $1/2$ and ρ arbitrarily close to $(2\delta)^{-1}$; the conclusion then follows by plugging the above consideration in the conditions for local and global wellposedness of solutions in H^α provided by Theorems 1.6 from [14], which are expressed in terms of (γ, ρ) . For instance, concerning Point i., local wellposedness holds for $\rho \sim (2\delta)^{-1} > 3/4$ (thus $\delta < 2/3$) and $\alpha > -\rho \sim -(2\delta)^{-1}$, while global existence holds for $\alpha > -\min\{3/2, \rho/(3-2\gamma)\} \sim -\min\{3/2, (4\delta)^{-1}\}$. Point ii. is similar. \square

Analogue statements can be obtained combining Theorem 32 with other results from the aforementioned papers, for instance Theorems 1.9 and 1.10 from [12] or Theorem 1.7 from [14].

In the setting of standard dispersive equations, a key role in establishing uniqueness of solutions is played by Strichartz estimates. In the paper [22], for a given path $w \in C^0([0, T])$, the authors study under which conditions the operator A given by

$$f \mapsto (Af)_t := \int_0^t |w_t - w_s|^{-\alpha} f_s ds$$

is bounded from $L^p(0, T)$ to $L^q(0, T)$ for suitable values of (p, q) ; the idea is to apply this kind of modulated Hardy–Littlewood–Sobolev inequality to obtain Strichartz estimates for the modulated semigroup

$$P_{s,t}\psi(x) = e^{i\Delta(w_t - w_s)}\psi(x) = \frac{1}{(4\pi(w_t - w_s))^{d/2}} \int_{\mathbb{R}^d} \exp\left(i \frac{|x - y|^2}{4(w_t - w_s)}\right) \psi(y) dy.$$

They only consider w sampled as a stochastic process, specifically a fBm of parameter $H \in (0, 1)$, but a closer inspection of the proof shows that their result (Theorem 1.1 from [22]) can be restated in an analytic fashion as follows.

Theorem 44. Suppose that $w \in C^0([0, T])$ admits an occupation density $\ell^w \in C_t^\beta C_x^0$ for some $\beta \in (0, 1)$. Then for any $p, q \in (1, \infty)$ and $\alpha \in (0, 1)$ satisfying

$$2 - \alpha = \frac{1}{p} + \frac{1}{q}$$

there exists a constant $C > 0$ such that for all $f \in L^p(0, T)$ and $g \in L^q(0, T)$ it holds

$$\left| \int_0^T \int_0^T f_t |w_t - w_s|^{-\alpha} g_s ds dt \right| \leq CT^{\alpha\beta} \|f\|_{L^p} \|g\|_{L^q}.$$

For any $\delta \in [0, 1/2)$, almost every $\varphi \in C^\delta([0, T])$ satisfies the above assumption for any $\beta < 1 - \delta$.

Proof. The proof of Theorem 1.1 in [22] is entirely analytical, as it follows closely the proof of the standard Hardy–Littlewood–Sobolev inequality from Lieb–Loss [46], but it requires a key property satisfied by fBm paths, given in Lemma 2.1 therein: if we define

$$W(r, T) := \sup_{t \in [0, T]} \int_0^T \mathbb{1}_{|w_t - w_s| < r} ds,$$

then there must exist a constant c such that

$$W(r, T) \leq 2rcT^\beta \quad \text{for all } r > 0. \quad (20)$$

It is not difficult to see that requirement (20) is equivalent to the request that $\ell_T^w(x) \leq cT^\beta$ for all $x \in \mathbb{R}$; indeed, assume first that $\ell_T^w(x) \leq cT^\beta$ holds, then

$$W(r, T) = \sup_{t \in [0, T]} \int_0^T \mathbb{1}_{w_s \in B(w_t, r)} ds = \sup_{t \in [0, T]} \int_{B(w_t, r)} \ell_T^w(x) dx \lesssim 2rcT^\beta.$$

On the other side, if w admits a continuous density ℓ^w and (20) holds, then

$$\ell_T^w(w_t) = \lim_{r \rightarrow 0} \frac{1}{2r} \int_{B(w_t, r)} \ell_T^w(x) dx \leq \lim_{r \rightarrow 0} \frac{1}{2r} W(r, T) \leq cT^\beta,$$

and since we know that ℓ_T^w is supported on $w([0, T])$, the above estimate extends to all $x \in \mathbb{R}^d$.

It is now clear that requirement (20) can be expressed in entirely analytical terms and so does the proof of Theorem 1.1 from [22], and the authors are only using the additional fact that almost every fBm trajectory satisfies (20). Another analytical condition ensuring that $\ell^w \in C_t^\beta C_x^0$ is the (γ, ρ) -irregularity property, since by Lemma 21, if $\rho > 1$, then $\ell^w \in C_t^\gamma C_x^0$. Moreover we can apply Lemma 7 to deduce that for any $\theta \in (0, 1)$ such that $\theta\rho > 1$, $\ell^w \in C_t^{1-\theta+\theta\gamma} C_x^0$, which implies that if w is ρ -irregular for any $\rho < \bar{\rho}$, $\bar{\rho} > 1$, then $\ell^w \in C_t^\beta C_x^0$ for any $\beta < 1 - (2\bar{\rho})^{-1}$. The conclusion follows applying the fact that for $\delta < 1/2$, almost every $\varphi \in C_t^\delta$ is ρ -irregular for any $\rho < (2\delta)^{-1}$. \square

Similarly, the proofs of Strichartz estimates and well-posedness for w -modulated NLS (Proposition 1.1 and Theorem 1.2 respectively) from [22] are entirely deterministic and only rely on the validity of the above modulated Hardy–Littlewood–Sobolev inequality; they can therefore be fully generalised to prevalence results, similarly to Theorems 42 and 43 above.

In [13], the authors provide regularity estimates for solutions to scalar conservation laws modulated by a path w of the form

$$\partial_t u + \sum_{i=1}^d \partial_{x_i} A^i(u) \circ \frac{dw_t^i}{dt} = 0 \quad \text{on } \mathbb{T}^d, \quad u(0) = u_0 \in L^\infty(\mathbb{T}^d). \quad (21)$$

They use the concept of (γ, ρ) -irregularity to show regularisation by noise phenomena whenever w is sampled as an fBm, but their results are of analytic (or path-by-path) type; before stating their result, let

us point out a simplification: given a (γ, ρ) -irregular $w \in C_t^\delta$, the authors impose on a suitable parameter λ , depending on another parameter $\nu \geq 1$, the condition

$$\lambda < \frac{\rho(\delta+1) - (1-\gamma)}{(\nu\rho \vee 1)(\delta+1) + (1-\gamma)} \wedge \frac{\rho + 2(\nu\rho \vee 1)}{(\nu\rho \vee 1)(2\delta+1) + (1-\gamma)} =: c_1 \wedge c_2.$$

Thanks to Theorem 36, we can actually simplify the above expression; we claim that $c_1 \leq c_2$. Since it must always hold $\delta \leq \delta_{\gamma, \rho}^*$, we have

$$\rho(\delta+1) - (1-\gamma) \leq \rho + \rho\delta_{\gamma, \rho}^* - (1-\gamma) \leq \rho;$$

to check that $c_1 \leq c_2$, it then suffices to verify that

$$\frac{\rho}{(\nu\rho \vee 1)(\delta+1) + 1 - \gamma} \leq \frac{\rho + 2(\nu\rho \vee 1)}{(\nu\rho \vee 1)(2\delta+1) + (1-\gamma)}$$

and after a few algebraic manipulations we see that this is equivalent to

$$\rho\delta \leq 2[(\nu\rho \vee 1)(\delta+1) + 1 - \gamma];$$

this is now clearly always true since $\nu \geq 1$, so that $\rho\delta \leq (\nu\rho \vee 1)\delta \leq 2(\nu\rho \vee 1)(\delta+1)$.

The main result of [13] can then be restated as follows:

Theorem 45 (Theorem 2.4 from [13]). *Let $w \in C_t^\delta$ be (γ, ρ) -irregular and let u be a quasi-solution to (21). Assume $A = (A^1, \dots, A^d) \in C^2(\mathbb{R}; \mathbb{R}^d)$ satisfies the following non-degeneracy condition: there exist $\nu \geq 1$ and $c > 0$ such that, for $A' = a = (a^1, \dots, a^d) \in C^1(\mathbb{R}; \mathbb{R}^d)$, it holds*

$$\inf_{v \in \mathbb{R}^d} \max_{i=1, \dots, d} |v_i(a^i(x) - a^i(y))| \geq c|x-y|^\nu \quad \text{for all } x, y \in \mathbb{R}.$$

Then there exists a constant $C = C(\|\Phi^w\|_{W^{\gamma, \rho}})$ such that for all $T > 0$ and all

$$\lambda < \frac{\rho(\delta+1) - (1-\gamma)}{(\nu\rho \vee 1)(\delta+1) + (1-\gamma)}$$

it holds

$$\int_0^T \|u_t\|_{W^{\lambda, 1}} dt \leq C(\|u_0\|_{L_x^1} + \|u\|_{L_{t,x}^1} + \|w\|_{C_t^\delta} \|a'(v)m\|_{TV}). \quad (22)$$

If u is an entropy solution then in addition

$$\|u_t\|_{W^{\lambda, 1}} < \infty \quad \text{for all } t > 0.$$

We avoid here providing all the details on the result above for which we refer the reader to [13]; let us only mention that the definition of quasi-solution to (21) requires the existence of a finite Radon measure m associated to u , which is the one appearing in estimate (22); u is an entropy solution if m is non-negative.

A few algebraic manipulations together with Theorem 32 imply the following result.

Corollary 46. *Let $\delta \in (0, 1)$. For almost every $\varphi \in C_t^\delta$, the statement of Theorem 45 holds for any*

$$\lambda < \frac{1}{(\nu \vee 2\delta)(\delta+1) + \delta}.$$

4 Criteria for ρ -irregularity of stochastic processes

This section is devoted to the study of probabilistic properties sufficient to ensure that a stochastic process has (γ, ρ) -irregular sample paths. It includes the proof of Theorem 33, which is the cornerstone for our main result prevalence Theorem 32, but develops several criteria which are of independent interest. In particular we establish ρ -irregularity for processes like fBm, α -stable process, Ornstein–Uhlenbeck as well as $X_t = \int_0^t B_s ds$; many of these process have appeared in regularisation by noise phenomena, see for instance [45], [2], [21], [11] among others.

4.1 General criteria

We provide here useful general criteria to establish ρ -irregularity for a given stochastic process, which will then be applied to several examples in the next section.

We adopt the following convention: although we always write statements to hold for any $\xi \in \mathbb{R}^d$, they must be interpreted as “for all ξ big enough”, i.e. $|\xi| \geq C$ for some universal deterministic constant $C > 0$, so that for instance expressions like $\log |\xi|$ are meaningful. We have seen that in the case of ρ -irregularity this is not an issue, since the only relevant information given by $\|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}$ is for big values of $|\xi|$. Similarly, for a modulus of continuity φ defined only on a neighbourhood of 0, t and s are tacitly assumed to be sufficiently close whenever $\varphi(|t-s|)$ appears.

The next statement is given in a general form, but keep in mind that our primary focus is the case $F(\xi) = |\xi|^\alpha$ for suitable values of α .

Theorem 47. *Let $(X_t)_{t \in [0,T]}$ be an \mathbb{R}^d -valued stochastic process with \mathbb{P} -a.s. measurable trajectories, $T < \infty$. Let $F : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ be a continuous function with the following properties:*

- i. *there exist constants $c_1, c_2 > 0$ such that $F(x) \leq c_1 F(y)$ whenever $|x - y| \leq c_2$;*
- ii. *F has at most polynomial growth, i.e. $F(\xi) \leq c_3 |\xi|^\alpha$ as $|\xi| \rightarrow \infty$ for some $\alpha < \infty$, $c_3 > 0$.*

Also assume that there exist positive constants μ, θ, δ such that the following hold:

- 1. *(Integrability condition)*

$$\mathbb{E} \left[\exp \left(\mu \int_0^T |X_t|^\theta dt \right) \right] < \infty; \quad (23)$$

- 2. *(Continuity condition)*

$$\sup_{\substack{0 < t-s < \delta \\ \xi \in \mathbb{R}^d}} \mathbb{E} \left[\exp \left(\mu \frac{|\Phi_{s,t}^X(\xi)|^2 |F(\xi)|^2}{|t-s|} \right) \right] < \infty. \quad (24)$$

Then, for the choice $\varphi(x) = \sqrt{x|\log x|}$, defining the random variable

$$Y := \sup_{s \neq t, \xi \in \mathbb{R}^d} \frac{|\Phi_{s,t}^X(\xi)| |F(\xi)|}{\varphi(|t-s|) \sqrt{\log |\xi|}},$$

there exists $\lambda > 0$ such that $\mathbb{E}[\exp(\lambda Y^2)] < \infty$.

Proof. First of all let us show that, starting from (24), we can find another constant $\tilde{\mu}$ such that the same bound holds taking the supremum over all $s < t$, without the restriction $|t-s| < \delta$. Let $[s, t]$ be such that $|t-s| > \delta$; we can split the interval $[s, t]$ in at most $n = \lfloor T/\delta \rfloor + 1$ intervals of the form $[t_i, t_{i+1}]$, of size at most δ ; we have the estimate

$$\begin{aligned} \mathbb{E} \left[\exp \left(\tilde{\mu} \frac{|\Phi_{s,t}^X(\xi)|^2 |F(\xi)|^2}{|t-s|} \right) \right] &\leq \mathbb{E} \left[\exp \left(\frac{n\tilde{\mu}}{\delta} \sum_i |\Phi_{t_i, t_{i+1}}^X(\xi)|^2 |F(\xi)|^2 \right) \right] \\ &= \mathbb{E} \left[\prod_i \exp \left(\frac{n\tilde{\mu}}{\delta} |\Phi_{t_i, t_{i+1}}^X(\xi)|^2 |F(\xi)|^2 \right) \right] \\ &\leq \prod_i \mathbb{E} \left[\exp \left(n^2 \tilde{\mu} \frac{|\Phi_{t_i, t_{i+1}}^X(\xi)|^2 |F(\xi)|^2}{|t_{i+1} - t_i|} \right) \right]^{1/n} \end{aligned}$$

and choosing $\tilde{\mu}$ such that $(\lfloor T/\delta \rfloor + 1)^2 \tilde{\mu} \leq \mu$ we obtain

$$\sup_{\substack{|t-s| > \delta \\ \xi \in \mathbb{R}^d}} \mathbb{E} \left[\exp \left(\tilde{\mu} \frac{|\Phi_{s,t}^X(\xi)|^2 |F(\xi)|^2}{|t-s|} \right) \right] < \infty.$$

From now on with a slight abuse we will denote by μ the new constant under which we have a bound of the form (24) over all possible $t \neq s$.

Let us define, for any $s \neq t$ and suitable $\lambda > 0$, the following quantity:

$$J_{s,t}(\lambda) := \sum_{n \in \mathbb{N}} 2^{-n} \sum_{\xi \in 2^{-n} \mathbb{Z}^d} 2^{-n(d+1)} (1 + |\xi|)^{-(d+1)} \exp \left(\lambda \frac{|\Phi_{s,t}^X(\xi)|^2 |F(\xi)|^2}{|t-s|} \right).$$

It follows from (24) that, for all $\lambda \leq \mu$, $\mathbb{E}[J_{s,t}(\lambda)] \leq K$ uniformly in s, t ; moreover by Jensen inequality it is clear that for any $\beta > 1$ it holds $J_{s,t}(\lambda)^\beta \lesssim J_{s,t}(\beta\lambda)$. Let us also define

$$Y_{s,t} := \frac{1}{|t-s|^{1/2}} \sup_{\xi \in \mathbb{R}^d} \frac{|\Phi_{s,t}^X(\xi)| |F(\xi)|}{\sqrt{\log |\xi|}}.$$

In order to conclude, it suffices to show that there exists $\lambda \in (0, \mu)$ such that $\mathbb{E}[\exp(\lambda Y_{s,t}^2)] \leq K$ uniformly in $s < t$; indeed, we can then apply Lemma 90 from Appendix A.2 for the choice $\alpha = 1/2$, $E = C^0(\mathbb{R}^d)$ and $Z : [0, T] \rightarrow E$ given by $Z_t(\xi) := \Phi_t^X(\xi) F(\xi) / \sqrt{\log |\xi|}$, to get the associated bound for Y .

Fix $\xi \in \mathbb{R}^d$. For any $n \in \mathbb{N}$, we can find $\tilde{\xi} \in 2^{-n} \mathbb{Z}^d$ such that $|\xi - \tilde{\xi}| \lesssim 2^{-n}$; for such $\tilde{\xi}$ it holds

$$\frac{|\Phi_{s,t}^X(\tilde{\xi})| |F(\tilde{\xi})|}{|t-s|^{1/2}} \lesssim \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda) + n + \log |\tilde{\xi}|}.$$

We can assume $\theta \in (0, 1]$; indeed, if (23) is satisfied for some $\theta > 1$, then by Hölder's inequality it also holds for $\theta = 1$. Combining the trivial estimates $|e^{i\xi \cdot x} - e^{i\tilde{\xi} \cdot x}| \leq 2$, $|e^{i\xi \cdot x} - e^{i\tilde{\xi} \cdot x}| \leq |\xi - \tilde{\xi}| |x|$, we then find

$$|\Phi_{s,t}^X(\tilde{\xi}) - \Phi_{s,t}^X(\xi)| \leq \int_s^t |e^{i\xi \cdot X_r} - e^{i\tilde{\xi} \cdot X_r}| dr \lesssim |\xi - \tilde{\xi}|^\theta \int_s^t |X_r|^\theta dr \leq |\xi - \tilde{\xi}|^\theta \|X\|_{L^\theta},$$

which interpolated together with $|\Phi_{s,t}^X(\xi)| \leq |t-s|$ gives

$$|\Phi_{s,t}^X(\tilde{\xi}) - \Phi_{s,t}^X(\xi)| \lesssim |t-s|^{1/2} |\xi - \tilde{\xi}|^{\theta/2} \|X\|_{L^\theta}^{\theta/2}.$$

Gathering everything together and using the fact that for n big enough it holds $F(\xi) \lesssim F(\tilde{\xi})$, we obtain

$$\begin{aligned} |\Phi_{s,t}^X(\xi)| |F(\xi)| &\lesssim |\Phi_{s,t}^X(\xi) - \Phi_{s,t}^X(\tilde{\xi})| |F(\xi)| + |\Phi_{s,t}^X(\tilde{\xi})| |F(\tilde{\xi})| \\ &\lesssim |t-s|^{1/2} |\xi - \tilde{\xi}|^{\theta/2} \|X\|_{L^\theta}^{\theta/2} F(\xi) \\ &\quad + |t-s|^{1/2} \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda) + n + \log |\tilde{\xi}|} \\ &\lesssim |t-s|^{1/2} \|X\|_{L^\theta}^{\theta/2} 2^{-n\theta/2} F(\xi) \\ &\quad + |t-s|^{1/2} \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda) + n + \log |\xi|} + c \end{aligned}$$

Therefore, choosing $n \sim \log |\xi|$, which is by assumption enough for $F(\xi) 2^{-n\theta/2} \lesssim 1$, we obtain

$$|\Phi_{s,t}^X(\xi)| |F(\xi)| \lesssim |t-s|^{1/2} \left[\|X\|_{L^\theta}^{\theta/2} + \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda) + \log |\xi|} \right].$$

Dividing by $\sqrt{\log |\xi|} |t-s|^{1/2}$ and taking the supremum we get

$$Y_{s,t} \lesssim \|X\|_{L^\theta}^{\theta/2} + \lambda^{-1/2} + \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda)},$$

and so there exists a constant C such that

$$\begin{aligned}\exp(\lambda Y_{s,t}^2) &\lesssim \exp(\lambda C \|X\|_{L^\theta}^\theta) J_{s,t}(\lambda)^C \\ &\lesssim \exp(2\lambda C \|X\|_{L^\theta}^\theta) + J_{s,t}(\lambda)^{2C} \\ &\lesssim \exp(2\lambda C \|X\|_{L^\theta}^\theta) + J_{s,t}(2\lambda C).\end{aligned}$$

Choosing λ such that $2\lambda C \leq \mu$ we therefore obtain a uniform bound for $\mathbb{E}[\exp(\lambda Y_{s,t}^2)]$ which by the above reasoning implies the conclusion. \square

Remark 48. *Going through the same proof, one can obtain a similar statement for F such that:*

- i. there exist constants $c_1, c_2, c_3 > 0$ such that $F(x) \leq c_1 F(c_2 y)$ whenever $|x - y| \leq c_3$;*
- ii. F has exponential-type growth, i.e. $\log F(\xi) \leq c_4 |\xi|^\alpha$ as $\xi \rightarrow \infty$ for some $\alpha < \infty$, $c_4 > 0$.*

Then under conditions (23) and (24) it is possible to find $c > 0$ such that defining

$$Y := \sup_{s \neq t, \xi \in \mathbb{R}^d} \frac{|\Phi_{s,t}^X(\xi)| F(c\xi)}{\varphi(|t-s|)|\xi|^\alpha} \quad (25)$$

the same conclusion as in Theorem 47 holds. The choice $F(\xi) = \exp(\lambda|\xi|^\alpha)$ satisfies the above requirements and in this case we can get rid of $|\xi|^\alpha$ in the denominator of (25) by changing c .

We immediately deduce the following result.

Corollary 49. *Let X be a process satisfying the assumptions of Theorem 47 for $F(\xi) = |\xi|^\alpha$. Then for any $\rho < \alpha$, there exists $\gamma = \gamma(\rho) > 1/2$ such that X is (γ, ρ) -irregular and moreover*

$$\mathbb{E}[\exp(\lambda \|\Phi^X\|_{\mathcal{W}^{\gamma, \rho}}^2)] < \infty \quad \forall \lambda \in \mathbb{R}. \quad (26)$$

Proof. Let Y be defined as in Theorem 47; combining the trivial estimate $|\Phi_{s,t}^X(\xi)| \leq |t-s|$ with

$$|\Phi_{s,t}^X(\xi)| \leq Y |\xi|^{-\alpha} \log^{1/2} |\xi| |t-s|^{1/2} \log |t-s|^{1/2}$$

by interpolation we obtain, for any fixed $\varepsilon > 0$,

$$\begin{aligned}|\Phi_{s,t}^X(\xi)| &\leq Y^{1-2\varepsilon} |\xi|^{-\alpha(1-2\varepsilon)} \log^{1/2-\varepsilon} |\xi| |t-s|^{1/2+\varepsilon} \log |t-s|^{1/2} \\ &\lesssim_\varepsilon Y^{1-2\varepsilon} |\xi|^{-\alpha(1-3\varepsilon)} |t-s|^{1/2+\varepsilon/2}\end{aligned}$$

so that setting $\rho = \alpha(1-3\varepsilon) < \alpha$, $\gamma = 1/2 + \varepsilon/2 > 1/2$, we obtain

$$\|\Phi^X\|_{\mathcal{W}_T^{\gamma, \rho}} \lesssim Y^{1-2\varepsilon}$$

which also implies that, for a suitable $C = C(\varepsilon)$, taking $\mu > 0$ small enough, it holds

$$\mathbb{E}[\exp(\mu \|\Phi^X\|_{\mathcal{W}^{\gamma, \rho}}^{2/(1-2\varepsilon)})] \leq \mathbb{E}[\exp(\mu C Y^2)] < \infty. \quad (27)$$

Since $\varepsilon > 0$, $2/(1-2\varepsilon) > 2$ and therefore from (27) we immediately deduce (26). The reasoning holds for any $\varepsilon > 0$, so we can invert the relations between ρ , ε and γ to deduce that for any given $\rho < \alpha$ we can take $\gamma(\rho) = 1/2 + (1 - \rho/\alpha)/6$. \square

Theorem 47 and Corollary 49 are well suited for establishing ρ -irregularity in several situations, as we will show in the next section. However, conditions (23) and (24) are in general difficult to check, due to their exponential nature. We present here a weaker version of Theorem 47 which relaxes condition (23); in Section 4.3 we will show how it is possible to relax condition (24) instead.

Corollary 50. Let $(X_t)_{t \in [0, T]}$ be an \mathbb{R}^d -valued stochastic process with \mathbb{P} -a.s. measurable trajectories, $T < \infty$; assume that it satisfies the continuity condition (24) from Theorem 47 for $F(\xi) = |\xi|^\alpha$ and that there exists $\theta > 0$ such that

$$\mathbb{P} \left(\int_0^T |X_t|^\theta dt < \infty \right) = 1.$$

Then for any $\rho < \alpha$ there exists $\gamma = \gamma(\rho) > 1/2$ such that \mathbb{P} -a.s. X is (γ, ρ) -irregular.

Proof. Let $f \in C_t^0$ be a deterministic continuous function which is α -irregular, whose existence is granted by Theorem 31.

For any $N \in \mathbb{N}$, set $A = \{\omega \in \Omega : \|X(\omega)\|_{L^\theta([0, T])} \leq N\}$ and define $X^N := \mathbb{1}_A X + \mathbb{1}_{A^c} f$. Then it is easy to check that by construction

$$\mathbb{E}[\exp(\lambda \|X^N\|_{L^\theta})] < \infty \quad \forall \lambda \in \mathbb{R}.$$

Moreover, letting μ, δ denote the constant under which X satisfies condition (24), for any $|s - t| < \delta$ we have

$$\begin{aligned} \mathbb{E} \left[\exp \left(\mu \frac{|\Phi_{s,t}^{X^N}(\xi)|^2 |\xi|^{2\alpha}}{|t-s|} \right) \right] &\leq \mathbb{E} \left[\exp \left(\mu \frac{|\Phi_{s,t}^X(\xi)|^2 |\xi|^{2\alpha}}{|t-s|} \right) \right] + \exp \left(\mu \frac{|\Phi_{s,t}^f(\xi)|^2 |\xi|^{2\alpha}}{|t-s|} \right) \\ &\leq \sup_{\substack{0 < s-t < \delta \\ \xi \in \mathbb{R}^d}} \mathbb{E} \left[\exp \left(\mu \frac{|\Phi_{s,t}^X(\xi)|^2 |\xi|^{2\alpha}}{|t-s|} \right) \right] + \exp(\mu \|\Phi^f\|_{\mathcal{W}^{1/2, \alpha}}^2) < \infty \end{aligned}$$

which implies that X^N also satisfies condition (24). Therefore, by Corollary 49, for fixed N and $\rho < \alpha$, X^N is \mathbb{P} -a.s. ρ -irregular; but then

$$\mathbb{P}(X \text{ is } \rho\text{-irregular}) \geq \mathbb{P}(X = X^N, X^N \text{ is } \rho\text{-irregular}) = 1 - \mathbb{P}(\|X\|_{L^\theta} > N) \rightarrow 1 \quad \text{as } N \rightarrow \infty.$$

□

The following lemma provides easy-to-check sufficient conditions for (24) to hold. It is a more general version of the technique developed in [10], Theorem 4.3.

Lemma 51. Let $(X_t)_{t \in [0, T]}$ be an \mathbb{R}^d -valued stochastic process with measurable trajectories and $f \in C^0([0, T] \times \mathbb{R}^d; \mathbb{C})$ such that $\|f\|_{C^0} \leq 1$. Assume that there exists a deterministic constant C_f such that for any $s < t$ it holds

$$\mathbb{P} \left(\left| \mathbb{E} \left[\int_s^t f(u, X_u) du \middle| \mathcal{F}_s \right] \right| \leq C_f \right) = 1. \quad (28)$$

Then there exists a constant $\mu > 0$ independent of f, s, t such that

$$\sup_{s \neq t} \mathbb{E} \left[\exp \left(\mu \frac{\left| \int_s^t f(u, X_u) du \right|^2}{C_f |t-s|} \right) \right] < \infty. \quad (29)$$

Proof. We can assume without loss of generality f to be real valued, since otherwise we can just apply separately estimates for $\Re f$ and $\Im f$, which still satisfy the above assumptions. Suppose first that $|t-s| \leq C_f$. Then we have the trivial estimate

$$\left| \int_s^t f(u, X_u) du \right|^2 \leq |t-s|^2 \leq |t-s| C_f,$$

which immediately implies the estimate (29) in this case.

Suppose now that $|t-s|/C_f \geq 1$. Let $n \in \mathbb{N}$ to be fixed later; define $t_k = s + k(t-s)/n$ for $k \in \{0, \dots, n\}$ and

$$Z_k = \mathbb{E} \left[\int_s^t f(u, X_u) du \middle| \mathcal{F}_{t_{k+1}} \right] - \mathbb{E} \left[\int_s^t f(u, X_u) du \middle| \mathcal{F}_{t_k} \right].$$

Setting $S_n = \sum_{k=0}^{n-1} Z_k$, S_n is a martingale and it holds

$$\int_s^t f(u, X_u) du = S_n + \mathbb{E} \left[\int_s^t f(u, X_u) du | \mathcal{F}_s \right].$$

In order to get a bound for such quantity, we want to apply Azuma-Hoeffding inequality to the martingale increments Z_k . We start by estimating the second term:

$$\left| \mathbb{E} \left[\int_s^t f(u, X_u) du | \mathcal{F}_s \right] \right| \leq C_f \leq C_f^{1/2} |t - s|^{1/2},$$

so that uniformly in $s < t$ we have

$$\exp \left(\mu \left| \mathbb{E} \left[\int_s^t f(u, X_u) du | \mathcal{F}_s \right] \right|^2 / (C_f |t - s|) \right) < \infty. \quad (30)$$

For the martingale increments instead we have

$$Z_k = \mathbb{E} \left[\int_{t_{k+1}}^t f(u, X_u) du | \mathcal{F}_{t_{k+1}} \right] - \mathbb{E} \left[\int_{t_k}^t f(u, X_u) du | \mathcal{F}_{t_k} \right] + \int_{t_k}^{t_{k+1}} f(u, X_u) du$$

and so by the hypothesis and $\|f\|_\infty \leq 1$ we have the \mathbb{P} -a.s. estimate

$$|Z_k| \leq 2C_f + \int_{t_k}^{t_{k+1}} |f(u, X_u)| du \leq 2C_f + |t - s|/n.$$

By Azuma-Hoeffding inequality, there exist universal constants $\mu, K > 0$ such that

$$\mathbb{E}[\exp(\mu |S_n|^2 / C_n)] < K$$

where the constant C_n is given by

$$C_n = \sum_{k=0}^{n-1} (2C_f + |t - s|/n)^2 \sim 4n C_f^2 + |t - s|^2/n.$$

Choosing n such that $n \sim |t - s|/C_f$ we obtain $C_n \sim |t - s|C_f$ and so for μ sufficiently small it holds

$$\mathbb{E}[\exp(\mu |S_n|^2 / C_f |t - s|)] < K.$$

where the estimate is uniform over s, t . Together with estimate (30), this implies the conclusion. \square

Remark 52. Lemma 51 is of independent interest, even outside the context of ρ -irregularity. For instance, let X be a Markov process with semigroup $P_t = e^{tA}$ and let $-\lambda$ be an eigenvalue of A with associated eigenvector f^λ ; more precisely, assume $\lambda > 0$ and let $f^\lambda \in L^\infty$ (normalised so that $\|f^\lambda\|_{L^\infty} = 1$) satisfy $Af^\lambda = -\lambda f^\lambda$. Then it holds

$$\left| \mathbb{E} \left[\int_s^t f^\lambda(X_u) du | \mathcal{F}_s \right] \right| = \left| \int_s^t (P_{u-s} f^\lambda)(X_s) du \right| = \left| \int_s^t e^{-\lambda(u-s)} f^\lambda(X_s) du \right| \leq \frac{1}{\lambda};$$

this implies for instance for $s = 0, t = 1$ that there exists a universal $\mu > 0$ such that

$$\sup_{\lambda > 0} \mathbb{E} \left[\exp \left(\mu \lambda \left| \int_0^1 f^\lambda(X_u) du \right|^2 \right) \right] < \infty.$$

4.2 ρ -irregularity for β -(e)SLND Gaussian processes and examples

We apply here the results of the previous section to prove part of Theorem 33. More quantitative results are given in the next two statements.

Theorem 53. *Let $\{X_t\}_{t \in [0, T]}$ be a \mathbb{R}^d -valued separable Gaussian process with measurable paths s.t.*

$$\int_0^T \mathbb{E}[|X_t|^2] dt < \infty; \quad (31)$$

suppose also that X is β -SLND for some $\beta \in (0, \infty)$. Then X satisfies the assumptions of Theorem 47 for $F(\xi) = |\xi|^{1/2\beta}$; therefore for any $\rho < (2\beta)^{-1}$ there exists $\gamma = \gamma(\rho) > 1/2$ such that X is (γ, ρ) -irregular

$$\mathbb{E}[\exp(\lambda \|\Phi^X\|_{\mathcal{W}^{\gamma, \rho}}^2)] < \infty \quad \forall \lambda \in \mathbb{R}.$$

Proof. It follows from (31) that X is an $L^2([0, T]; \mathbb{R}^d)$ -valued Gaussian process and therefore by Fernique Theorem we can conclude that there exists $\mu > 0$ such that

$$\mathbb{E}[\exp(\mu \|X\|_{L^2}^2)] < \infty,$$

which implies that condition (23) is satisfied. It remains to check condition (24), which we plan to do with the help of Lemma 51. Taking $f(t, x) = e^{i\xi \cdot x}$ for a fixed $\xi \in \mathbb{R}^d$, we need to estimate the associate constant $C_f = C_\xi$; it is enough to provide estimates whenever s and t both lie in an interval of size at most δ , δ being the parameter for which the β -SLND condition is satisfied. It holds

$$\begin{aligned} \left| \mathbb{E} \left[\int_s^t e^{i\xi \cdot X_u} du \middle| \mathcal{F}_s \right] \right| &\leq \int_s^t |\mathbb{E}[e^{i\xi \cdot X_u} | \mathcal{F}_s]| du \\ &= \int_s^t \exp \left(-\frac{1}{2} \xi \cdot (\text{Var}(X_u | \mathcal{F}_s) \xi) \right) du \\ &\leq \int_s^t \exp(-c|\xi|^2 |u - s|^{2\beta}) du \\ &\leq \int_0^\infty \exp(-c|\xi|^2 r^{2\beta}) dr \sim |\xi|^{-1/\beta}, \end{aligned}$$

where in the third passage we used the β -SLND property. Therefore $C_\xi \leq c|\xi|^{-1/\beta}$ and thus there exists $\mu > 0$ such that for any $s, t \in [0, T]$ satisfying $0 < s - t < \delta$ it holds

$$\mathbb{E} \left[\exp \left(\mu \frac{|\xi|^{1/\beta} |\Phi_{s,t}^X(\xi)|^2}{|t - s|} \right) \right] \leq \mathbb{E} \left[\exp \left(c\mu \frac{|\Phi_{s,t}^X(\xi)|^2}{C_\xi |t - s|} \right) \right] \leq K$$

uniformly in $\xi \in \mathbb{R}^d$. By Theorem 47 we obtain the conclusion for the choice $F(\xi) = |\xi|^{1/2\beta}$. The last statement follows from Corollary 49. \square

Proposition 54. *Let $\{X_t\}_{t \in [0, T]}$ be a \mathbb{R}^d -valued separable Gaussian process with measurable paths satisfying condition (31) which is β -eSLND for some $\beta \in (0, \infty)$. Then there exist constants $c, \lambda > 0$ such that defining*

$$Y := \sup_{s \neq t, \xi \in \mathbb{R}^d} \frac{|\Phi_{s,t}^X(\xi)| \exp(c|\xi|^{2/(1+\beta)})}{\varphi(|t - s|)}$$

with $\varphi(x) = \sqrt{x |\log x|}$, it holds

$$\mathbb{E}[\exp(\lambda Y^2)] < \infty.$$

In particular, if $\beta \leq 1$, then X is exponentially irregular.

Proof. As in the proof of Theorem 53, condition (31) implies condition (23) by Fernique Theorem. The rest of the proof is similar, relying on Lemma 51 applied to $f(t, x) = e^{i\xi \cdot x}$, only this time we want to apply Remark 48 for the choice $F(\xi) = \exp(|\xi|^{2/(1+\beta)})$, which requires a faster decay for $C_f = C_\xi$.

For any $s < t$ such that $|t - s| < \delta$, where δ is the parameter in the β -eSLND condition, it holds

$$\begin{aligned} \left| \mathbb{E} \left[\int_s^t e^{i\xi \cdot X_u} du \middle| \mathcal{F}_s \right] \right| &= \int_s^t \exp \left(-\frac{1}{2} \text{Var}(\xi \cdot X_u | \mathcal{F}_s) \right) du \\ &\leq \int_0^1 \exp(-c|\xi|^2 |\log r|^{-\beta}) dr \\ &\sim \int_0^{+\infty} \exp \left(-c|\xi|^2 x - x^{-\frac{1}{\beta}} \right) x^{-\frac{\beta+1}{\beta}} dx \\ &\lesssim \int_0^\infty \exp \left(-\tilde{c} \left(|\xi|^2 x + x^{-\frac{1}{\beta}} \right) \right) dx \end{aligned}$$

where in the third line we used the change of variables $x = |\log r|^{-\beta}$. By the general inequality $a + b \gtrsim a^\theta b^{1-\theta}$, valid for all $a, b > 0$ and $\theta \in (0, 1)$, it holds

$$|\xi|^2 x + x^{-\frac{1}{\beta}} \gtrsim |\xi|^{2(1-\theta)} x^{1-\theta \frac{\beta+1}{\beta}} = |\xi|^{2/(\beta+1)}$$

for the choice $\theta = \beta/(\beta+1)$; therefore there exists a constant c_1 such that

$$\left| \mathbb{E} \left[\int_s^t e^{i\xi \cdot X_u} du \middle| \mathcal{F}_s \right] \right| \lesssim \exp(-c_1 |\xi|^{2/(\beta+1)})$$

and by Lemma 51 we deduce that there exists $c_2 > 0$ such that

$$\sup_{s \neq t, \xi \in \mathbb{R}^d} \mathbb{E} \left[\exp \left(c_2 \frac{e^{c_1 |\xi|^{2/(\beta+1)}} |\Phi_{s,t}^X(\xi)|^2}{|t-s|} \right) \right] < \infty.$$

The conclusion then follows from Remark 48. In particular if $\beta \in (0, 1]$, then $2/(\beta+1) \geq 1$, which implies that X is exponentially irregular. \square

The rest of this section is devoted to providing examples of Gaussian processes with the above properties; in the following $(\mathcal{F}_t)_t$ denotes either the filtration generated by B or the filtration generated by a given process X which will be clear from the context. Here are some examples of β -SLND Gaussian processes:

- i. Let W^H be a d -dimensional fBm of Hurst parameter H . We have seen in Section 2.4 that it is H -SLND, therefore W^H is \mathbb{P} -a.s. ρ -irregular for any $\rho < (2H)^{-1}$, recovering the results from [10].
- ii. Let B be a standard Brownian motion in \mathbb{R}^d and let $A \in \mathbb{R}^{d \times d}$, $x_0 \in \mathbb{R}^d$, $\sigma \in \mathbb{R}_{>0}$ and a given measurable function $f : [0, T] \rightarrow \mathbb{R}^d$; consider a generalised d -dimensional Ornstein–Uhlenbeck process, solution to the SDE

$$dX_t = (-A X_t + f_t) dt + \sigma dB_t, \quad X_0 = x_0.$$

The explicit expression for X is given by

$$\begin{aligned} X_t &= e^{-tA} x_0 + \int_0^t e^{-(t-s)A} f_s ds + \sigma \int_0^t e^{-(t-s)A} dB_s \\ &= e^{-(t-s)A} X_s + \int_s^t e^{-(t-r)A} f_r dr + \sigma \int_s^t e^{-(t-r)A} dB_r \end{aligned}$$

which implies that we have a decomposition in \mathcal{F}_s -adapted and \mathcal{F}_s -independent parts given by $X_t = X_{s,t}^{(1)} + X_{s,t}^{(2)} := \mathbb{E}[X_t|\mathcal{F}_s] + (X_t - \mathbb{E}[X_t|\mathcal{F}_s])$ where

$$X_{s,t}^{(1)} = e^{-(t-s)A}X_s + \int_s^t e^{-(t-r)A}f_r dr, \quad X_{s,t}^{(2)} = \sigma \int_s^t e^{-(t-r)A}dB_r.$$

It follows that for any $s < t$ such that $|s - t| < \delta$ and any $v \in \mathbb{R}^d$ it holds

$$\text{Var}(X_t \cdot v | \mathcal{F}_s) = \text{Var}(X_{s,t}^{(2)} \cdot v) = \sigma^2 \int_s^t |e^{-(t-r)A^T} v|^2 dr = \sigma^2 \int_0^{t-s} |e^{-rA^T} v|^2 dr;$$

choosing δ small enough such that $\|I_d - e^{-rA^T}\| \leq 1/2$ for all $r < \delta$ we deduce that

$$\text{Var}(X_t \cdot v | \mathcal{F}_s) \gtrsim |v|^2 |t - s|,$$

namely $\text{Var}(X_t | \mathcal{F}_s) \gtrsim |t - s|I_d$. We conclude that X is ρ -irregular for any $\rho < 1$.

- iii. There is a general class of Gaussian processes for which a β -SLND condition holds, given by so called *moving averages of white noise*, as already observed by Berman (see Section 3 from [7]). Specifically, let $K : \mathbb{R}^+ \rightarrow \mathbb{R}^{d \times d}$ be a function such that $KK^T(t) \gtrsim |t|^{2\beta-1}I_d$, for some $\beta > 0$ and all t small enough and define the process

$$X_t := \int_0^t K(t-r)dB_r,$$

where B is a standard Bm in \mathbb{R}^d . Then for any $s < t$, we have a decomposition in \mathcal{F}_s -adapted and \mathcal{F}_s -independent parts given by

$$X_t = X_{s,t}^{(1)} + X_{s,t}^{(2)} = \int_0^s K(t-r)dB_r + \int_s^t K(t-r)dB_r$$

which implies that

$$\text{Var}(X_t | \mathcal{F}_s) = \int_s^t (KK^T)(t-r)dr = \int_0^{t-s} (KK^T)(u)du \gtrsim |t-s|^{2\beta}$$

for all s, t such that $|t-s|$ is small enough; we deduce that in this case we have ρ -irregularity for any $\rho < (2\beta)^{-1}$. The standard example for this type of processes is for the choice $K(t) = k(t)I_d$, where $k : \mathbb{R}_+ \rightarrow \mathbb{R}$ is such that $|k(t)| \gtrsim |t|^{\beta-1/2}$. For the choice $k(t) = |t|^{H-1/2}$ we obtain the d -dimensional Levy fBm (sometimes also referred to as type-II fBm), for which again we have $\rho < (2H)^{-1}$.

- iv. The class of moving averages is closed under integration. Given K and X as above, defining $Y := \int_0^\cdot X_s ds$, by stochastic Fubini it holds

$$Y_t = \int_0^t \int_0^s K(s-r)dB_r ds = \int_0^t \int_r^t K(s-r)ds dB_r = \int_0^t \tilde{K}(t-r)dB_r$$

for the choice $\tilde{K}(\cdot) = \int_0^\cdot K(s)ds$. For the choice $K(t) = |t|^{H-1/2}I_d$, i.e. X being a Levy fBm of parameter H , then $\tilde{K}(t) = c_H |t|^{(H+1)-1/2}$, which implies that Y is ρ -irregular for any $\rho < (2+2H)^{-1}$. Technically speaking standard fBm does not belong to the moving average class, but it is clear that we can apply the same reasoning to formula (10), obtaining

$$\begin{aligned} Y_t &= \int_0^t W_r^H dr = \left[\int_0^s W_r^H dr + \int_s^t \mathbb{E}[W_r^H | \mathcal{F}_s] dr \right] + \int_s^t (W_r^H - \mathbb{E}[W_r^H | \mathcal{F}_s]) dr \\ &= Y_{s,t}^{(1)} + c_H \int_s^t \int_s^r (r-u)^{H-1/2} dB_u dr \\ &= Y_{s,t}^{(1)} + \tilde{c}_H \int_s^t (t-u)^{(H+1)-1/2} dB_u = Y_{s,t}^{(1)} + Y_{s,t}^{(2)}. \end{aligned}$$

Arguing as above, we see that Y is $(1+H)$ -SLND and thus ρ -irregular for any $\rho < (2+2H)^{-1}$. Since W^H as $C^{H-\varepsilon}$ trajectories, Y has $C^{1+H-\varepsilon}$ trajectories, showing that it is still possible for differentiable paths to be ρ -irregular, for relatively small parameters ρ . The reasoning can be iterated: if Y is the process obtained by integrating n times W^H , so that $Y \in C^{n+H-}$ with full probability, then Y_t has a decomposition into \mathcal{F}_s -adapted and \mathcal{F}_s -independent parts given by

$$Y_t = Y_{s,t}^{(1)} + Y_{s,t}^{(2)}, \quad Y_{s,t}^{(2)} = Y_t - \mathbb{E}[Y_t | \mathcal{F}_s] = c_{n,H} \int_s^t (t-u)^{(H+n)-1/2} dB_u$$

which implies that it is $(n+H)$ -SLND; moreover for any $k \in \{0, \dots, n\}$, $D^k Y$ is $(H+n-k)$ -SLND.

- v. The following example is taken from [1] and it provides an explicit Gaussian process with continuous trajectories which are \mathbb{P} -a.s. ρ -irregular for all $\rho < \infty$. Functions with such properties can be also constructed by prevalence techniques, using the fact that countable intersection of prevalent sets is prevalent.

Let H_n be a sequence in $(0, 1)$ such that $H_n \downarrow 0$ and $\{B^{H_n}\}_n$ be a sequence of independent fBms in \mathbb{R}^d with parameters H_n , defined on an interval $[0, T]$; also consider a sequence λ_n of strictly positive numbers such that

$$\sum_n \lambda_n \mathbb{E}[\|B^{H_n}\|_{C^0}] < \infty$$

(for instance one can take $\lambda_n = (1 + \mathbb{E}[\|B^{H_n}\|_{C^0}])^{-1} n^{-2}$). Then it holds

$$\mathbb{E}\left[\sum_n \lambda_n \|B^{H_n}\|_{C^0}\right] < \infty$$

which implies that \mathbb{P} -a.s. the series $\sum_n \lambda_n B^{H_n}$ is absolutely convergent, thus uniformly convergent to an element of C_t^0 ; we denote such limit by Y , which is therefore a Gaussian variable on C_t^0 . By construction

$$\text{Var}(Y_t | \mathcal{F}_s) \geq \lambda_n^2 \text{Var}(B_t^{H_n} | \mathcal{F}_s) = \lambda_n^2 c_{H_n} |t-s|^{2H_n},$$

which implies that \mathbb{P} -a.s. Y is ρ -irregular for any $\rho < (2H_n)^{-1}$. As the reasoning holds for all n and $H_n \downarrow 0$ we conclude that Y is \mathbb{P} -a.s. ρ -irregular for all $\rho < \infty$.

We can also provide an example of a β -eSLND Gaussian process.

Proposition 55. *Let $\beta > 0$ and consider the d -dimensional Gaussian process X defined by*

$$X_t = \int_0^t (t-s)^{-1/2} |\log(t-s)|^{-\beta/2-1/2} dB_s \quad \forall t \in [0, 1/2] \quad (32)$$

where B is a standard Bm in \mathbb{R}^d . Then X admits a modification which is β -eSLND and satisfies the hypothesis of Proposition 54; moreover, X has trajectories in L_t^p for any $p < \infty$.

Proof. The process X is separable, as it is constructed from Bm, which is a separable process. Moreover, it is easy to check X is stochastically continuous and therefore by Proposition 3.2 from [19] it admits a measurable modification; from now on we will work with this modification. It holds

$$\text{Var}(X_t) = I_d \int_0^t (t-s)^{-1} |\log(t-s)|^{-\beta-1} ds = c_\beta |\log t|^{-\beta} \lesssim 1 \quad \forall t \in [0, 1/2]$$

which also implies by properties of Gaussian variables that we can find $\lambda > 0$ small enough s.t.

$$\mathbb{E}\left[\int_0^{1/2} \exp(\lambda |X_t|^2) dt\right] = \int_0^{1/2} \mathbb{E}[\exp(\lambda |X_t|^2)] dt < \infty$$

which implies that $X \in L_t^p$ for all $p \in [1, \infty)$. Finally, since X is the moving average of white noise associated to $K(r) = r^{-1/2} |\log r|^{-\beta/2-1/2}$, for any $s < t$ it holds

$$\text{Var}(X_t | \mathcal{F}_s) = I_d \int_0^{t-s} K^2(r) dr = I_d \int_0^{t-s} r^{-1} |\log r|^{-\beta-1} dr \sim |\log(t-s)|^{-\beta} I_d$$

which proves the β -eSLND property. \square

Remark 56. We have constructed the process X on the interval $[0, 1/2]$ for simplicity, but up to rescaling, a process with the same properties can be constructed on any finite interval $[0, T]$. Recall that for $\beta \leq 1$, X is exponentially irregular, thus Carathéodory and unbounded; therefore the L^p (actually, exponential) integrability obtained is optimal in this case. While we were working on the present paper, we became aware of the work-[39], in which the same process is independently introduced and called p -log Brownian motion, see Definition 4.1 from-[39]; the regime $p > 1/2$ therein corresponds to our $\beta > 0$. However, the authors of [39] establish infinite regularisation of the path X only in the case $p > 1$ (equivalently $\beta > 1$), see Corollary 4.4 from [39], and do not prove the exponential decay of $\Phi^X(\xi)$.

Further examples of ρ -irregular functions can be produced by combining a given β -SLND Gaussian process X with suitable deterministic functions.

Proposition 57. *The following hold:*

- a. Given a measurable $f : [0, T] \rightarrow \mathbb{R}^d$ and a β -SLND Gaussian process X , $Y := f + X$ is β -SLND; if moreover X satisfies condition (31) and $f \in L_t^\theta$ for some $\theta \in (0, \infty)$, then Y is also ρ -irregular for any $\rho < 1/(2\beta)$. A similar statement holds for f as above and X β -eSLND.
- b. Given a measurable $f : [0, T] \rightarrow \mathbb{R}$ satisfying $c^{-1} \leq |f_t| \leq c$ for a suitable constant $c > 0$ and a process X satisfying the assumptions of Theorem 53, $Y := fX$ also satisfies the assumptions and is therefore ρ -irregular for any $\rho < 1/(2\beta)$.
- c. Suppose X is a β -SLND Gaussian process with $\beta \in (0, 1]$ with trajectories in C_t^α and $A \in C^\gamma([0, T]; \mathbb{R}^{d \times d})$ is a deterministic function such that $\alpha + \gamma > 1$, satisfying

$$(A^T A)_t \geq c I_d \quad \forall t \in [0, T].$$

Then the process defined by the Young integral $Y = \int_0^\cdot A_s dX_s$ is also Gaussian, β -SLND, with trajectories in C_t^α ; it is ρ -irregular for any $\rho < 1/(2\beta)$.

Proof. Part a) follows from Remark 30 and the fact that if $f \in L_t^\theta$ and X satisfies the integrability assumptions, then so does $f + X$. Regarding b), it is clear that the process Y defined in this way is still Gaussian satisfying (31). The process Y is β -SLND since

$$\text{Var}(Y_t | \mathcal{F}_s) = |f_t|^2 \text{Var}(X_t | \mathcal{F}_s) \sim \text{Var}(X_t | \mathcal{F}_s) \gtrsim |t - s|^{2\beta} I_d.$$

It remains to prove c). By properties of Young integral (see Appendix A.1) $\varphi \mapsto \int_0^\cdot A d\varphi$ is a bounded linear map from C_t^α to itself, therefore Y is a Gaussian process on C_t^α since X is so; by Fernique theorem, it holds $\mathbb{E}[\exp(\lambda \|Y\|_{C^\alpha})] < \infty$ for all $\lambda \in \mathbb{R}$. By definition of Young integral, it holds

$$Y_t = Y_s + A_s(X_t - X_s) + R_{s,t}$$

where $|R_{s,t}| \lesssim |t - s|^{\alpha+\gamma} \|A\|_{C^\gamma} \|X\|_{C^\alpha}$. This implies that Y_t must have a decomposition in \mathcal{F}_s -adapted and \mathcal{F}_s -independent parts of the form $Y_t = Y_{s,t}^{(1)} + Y_{s,t}^{(2)} = \mathbb{E}[Y_t | \mathcal{F}_s] + (Y_t - \mathbb{E}[Y_t | \mathcal{F}_s])$, so that

$$Y_t^{(1)} = Y_s + A_s(\mathbb{E}[X_t | \mathcal{F}_s] - X_s) + \mathbb{E}[R_{s,t} | \mathcal{F}_s], \quad Y_{s,t}^{(2)} = A_s(X_t - \mathbb{E}[X_t | \mathcal{F}_s]) + R_{s,t} - \mathbb{E}[R_{s,t} | \mathcal{F}_s].$$

Since conditional expectation is a projection on $L^2(\Omega, \mathcal{F}, \mathbb{P})$, it holds

$$\mathbb{E}[|R_{s,t} - \mathbb{E}[R_{s,t} | \mathcal{F}_s]|^2] \leq \mathbb{E}[|R_{s,t}|^2] \lesssim |t - s|^{2(\alpha+\gamma)} \|A\|_{C^\gamma}^2 \mathbb{E}[\|X\|_{C^\alpha}^2] \lesssim |t - s|^{2(\alpha+\gamma)},$$

where $\alpha + \gamma > 1 \geq \beta$; the variance above is of order $o(|t - s|^{2\beta})$ as $|s - t| \sim 0$ and so the decay of $\text{Var}(Y_t|\mathcal{F}_s)$ is governed by

$$\text{Var}(A_s(X_t - \mathbb{E}[X_t|\mathcal{F}_s])) = A_s \text{Var}(X_t - \mathbb{E}[X_t|\mathcal{F}_s])A_s^T \gtrsim |t - s|^{2\beta} A_s A_s^T \gtrsim |t - s|^{2\beta}$$

whenever $|t - s|$ is small enough. This implies that Y is β -SLND and ρ -irregular for any $\rho < 1/(2\beta)$. \square

Remark 58. The above examples can be further combined together, for instance considering

$$Y_t := \int_0^t A_s dX_s + f_t X_t + g_t$$

for A, f, g satisfying the assumptions. One can moreover replace such deterministic objects by stochastic processes Z^i , independent of X , satisfying suitable regularity and integrability assumptions; this can be readily seen by conditioning on $(Z^i)_i$ first and applying the deterministic result. This allows to construct processes with ρ -irregular trajectories which are not Gaussian nor Markovian.

4.3 ρ -irregularity for α -stable processes

Section 4.2 deals exclusively with Gaussian processes, but the criteria developed in Section 4.1 apply in more general situations, including Markov processes. Here we treat the case of suitable α -stable processes.

Recall that a d -dimensional process X is a symmetric α -stable process, for some $\alpha \in (0, 2)$, with spherical measure μ (up to a renormalising constant) if it is a Lévy process such that, for any $s < t$,

$$\mathbb{E}[\exp(i\langle \xi, X_t - X_s \rangle)] = \exp\left(-(t - s) \int_{\mathbb{S}^{d-1}} |\langle \xi, z \rangle|^\alpha \mu(dz)\right).$$

See [59] for more details on the topic. From now on, we will say that X satisfies the *non-degeneracy condition* if there exists $c > 0$ such that

$$G(\xi) := \int_{\mathbb{S}^{d-1}} |\langle \xi, z \rangle|^\alpha \mu(dz) \geq c|\xi|^\alpha. \quad (33)$$

A similar condition has already appeared in the literature on regularisation by noise, see [21].

Proposition 59. *Let X be a symmetric α -stable process satisfying the non-degeneracy condition (33). Then X is \mathbb{P} -a.s. ρ -irregular for any $\rho < \alpha/2$.*

Proof. Let \mathcal{F}_t be the natural filtration associated to X ; for any $\xi \in \mathbb{R}^d$ and $s < t$, by the independence of increments and the non degeneracy condition, it holds

$$\left| \mathbb{E} \left[\int_s^t e^{i\langle \xi, X_r \rangle} dr \middle| \mathcal{F}_s \right] \right| = \left| e^{i\langle \xi, X_s \rangle} \int_s^t e^{-(r-s)G(\xi)} dr \right| = \int_0^{t-s} e^{-rG(\xi)} dr \leq \int_0^\infty e^{-cr|\xi|^\alpha} dr \sim |\xi|^{-\alpha}.$$

Applying Lemma 51 to the choice $f(t, x) = e^{i\xi \cdot x}$, since $C_f = C_\xi \lesssim |\xi|^{-\alpha}$, there exists $\lambda > 0$ s.t.

$$\sup_{s \neq t, \xi \in \mathbb{R}^d} \mathbb{E} \left[\exp \left(\lambda \frac{|\xi|^\alpha |\Phi_{s,t}^X(\xi)|^2}{|t - s|} \right) \right] < \infty.$$

We would like to conclude that the process X is ρ -irregular for any $\rho < \alpha/2$, but the integrability condition from Theorem 47 is not satisfied. However, the process X belongs \mathbb{P} -a.s. to L_t^θ for any $\theta < \alpha$ (see for instance Example 25.10 from [59]) and therefore we can apply Corollary 50 to obtain the conclusion. \square

Remark 60. The non-degeneracy condition (33) is for instance satisfied in the cases of μ_1 being the uniform measure on \mathbb{S}^{d-1} and $\mu_2 = \sum_{i=1}^d \delta_{e_i}$. The associated processes have respectively generators $\mathcal{L}_1 = (-\Delta)^{\alpha/2}$ and $\mathcal{L}_2 = \sum_i (-\partial_{x_i}^2)^{\alpha/2}$.

The previous examples generalises to anisotropic Markov processes, which show different irregularity behaviour in different directions. Property (34) below could be regarded as a notion of “ α -irregularity”.

Corollary 61. *Let $\alpha = (\alpha_1, \dots, \alpha_d) \in (0, 2)^d$ and let X be a d -dimensional process whose components $X^{(i)}$ are independent symmetric α_i -stable processes, so that*

$$\mathbb{E}[\exp(i\langle \xi, X_t - X_s \rangle)] = \exp\left(-|t - s| \sum_i |\xi_i|^{\alpha_i}\right).$$

Define $\|\xi\|_\alpha^2 = \sum_i |\xi_i|^{\alpha_i}$. Then setting $\varphi(x) = \sqrt{|x| \log x}$, it holds

$$\sup_{s \neq t, \xi \in \mathbb{R}^d} \frac{\|\xi\|_\alpha |\Phi_{s,t}^X(\xi)|}{\sqrt{\log |\xi|} \varphi(|t - s|)} < \infty \quad \mathbb{P}\text{-a.s.} \quad (34)$$

Proof. Going through the same calculations as in Proposition 59, we deduce that there exists $\lambda > 0$ such that

$$\sup_{s \neq t, \xi \in \mathbb{R}^d} \mathbb{E} \left[\exp \left(\lambda \frac{\|\xi\|_\alpha^2 |\Phi_{s,t}^X(\xi)|^2}{|t - s|} \right) \right] < \infty.$$

The conclusion then follows from Theorem 47 applied to the choice $F(\xi) = \|\xi\|_\alpha$ (together with a reasoning analogue to that of Corollary 50 in order to relax the integrability condition). \square

4.4 ρ -irregularity for β -LND Gaussian processes

We now turn to study the case of β -LND processes. Before proceeding further, let us explain the importance of β -LND in the literature and why they require a different treatment from the β -SLND case; for a broader overview, we refer to [37, 64] and the discussion in the introduction of [65].

The LND condition was historically the first one introduced by Berman [7], then extended to the case of random fields by Pitt [56]; it can be often checked using the *spectral measure* associated to the Gaussian field. Although the SLND property usually yields stronger results (e.g. for proving law of iterated logarithm for the local time of a Gaussian field), it is harder to check in practice, with the exception of the case where a *moving average representation* is available. For such reasons, also in view of possible (quite natural) future extensions of ρ -irregularity to the case of random fields, we find it useful to present a proof solely relying on the LND property here.

As already mentioned in Section 2.4, there are known examples of Gaussian processes X satisfy a LND condition but not a SLND one. Specifically, Cuzick in [16, Section 3] constructs a stationary centered process X , whose spectral measure F is described explicitly and thus characterizes it uniquely, such that for any $n \in \mathbb{N}$ there exists $c_n > 0$ such that, for any $t_1 < \dots < t_n$, it holds

$$\text{Var}(X_{t_n} | X_{t_1}, \dots, X_{t_{n-1}}) \geq c_n |t_n - t_{n-1}| \quad (35)$$

but also such that for any $s < t$ it holds

$$\text{Var}(X_t | X_r, r \leq s) = 0. \quad (36)$$

As argued in [16] and the references therein, condition (35) implies that X satisfies the 1-LND condition; on the other hand, not only (36) clearly implies that a β -SLND condition must fail for any $\beta \geq 0$, but it also informs us that the constants c_n appearing in (35) must satisfy $c_n \rightarrow 0$ as $n \rightarrow \infty$.

For this reason, in the case of β -LND processes, we will be unable to establish exponential estimates like those from Lemma 51, which are essential in order to apply Theorem 47: the behaviour of the constants c_n is unclear and we cannot make naive assumptions on them, cf. Remark 66 below. Therefore we need to proceed differently and find different ways to establish ρ -irregularity. The proof is very closed in spirit to the ones given by Kahane for fBm in Sections 17 and 18 of [43], which are based on combinatorial techniques combined with analytical properties of Fourier transform. Here the combinatorial argument is taken from [9], but it goes back to [51].

Theorem 62. Let X be a d -dimensional β -LND Gaussian process with continuous trajectories. Then for any $\gamma < 1/2$ and any $\rho < (2\beta)^{-1}$, X is (γ, ρ) -irregular with probability 1 and moreover

$$\mathbb{E}[\|\Phi^X\|_{\mathcal{W}_{\gamma, \rho}}^n] < \infty \quad \forall n \in \mathbb{N}.$$

Furthermore \mathbb{P} -a.s. X is ρ -irregular for all $\rho < (2\beta)^{-1}$.

The major drawback, compared to Theorem 53, is that in the β -LND case we lose exponential estimates, although we still have moments of any order. The proof is split in several parts; we start with two lemmas, of analytic and combinatorial nature respectively. We then provide the main estimate, given in Proposition 65, which allows to conclude the proof of Theorem 62.

The next lemma shows that, under suitable assumptions, we can obtain $\mathcal{FL}^{\rho, \infty}$ bounds starting from $\mathcal{FL}^{\rho, p}$ bounds for some $p < \infty$.

Lemma 63. Suppose $\varphi \in \mathcal{FL}^{\rho, p}$ has support contained in B_R , $\rho \geq 0$. Then $\varphi \in \mathcal{FL}^{\rho, \infty}$ and there exists a constant $C(\rho, p) > 0$ such that

$$\|\varphi\|_{\mathcal{FL}^{\rho, \infty}} \leq C R^{d/p} \|\varphi\|_{\mathcal{FL}^{\rho, p}}.$$

Proof. Up to a rescaling argument, it suffices to prove the statement in the case $R = 1$. Let $g \in C_c^\infty$ such that $g \equiv 1$ on $B(0, 1)$, then by assumption $\varphi = \varphi g$ and so $\hat{\varphi} = \hat{\varphi} * \hat{g}$. But then

$$\begin{aligned} (1 + |\xi|)^\rho |\hat{\varphi}(\xi)| &\leq (1 + |\xi|)^\rho \int |\hat{\varphi}(\xi - \eta)| |\hat{g}(\eta)| d\eta \\ &\lesssim_\rho \int (1 + |\xi - \eta|)^\rho |\hat{\varphi}(\xi - \eta)| (1 + |\eta|)^\rho |\hat{g}(\eta)| d\eta \\ &\lesssim \|\varphi\|_{\mathcal{FL}^{\rho, p}} \|g\|_{\mathcal{FL}^{\rho, p'}} \\ &\sim_p \|\varphi\|_{\mathcal{FL}^{\rho, p}} \end{aligned}$$

which gives the conclusion; in the above, p' denoted the conjugate exponent to p . \square

We now need to introduce some notations. Let $n \in \mathbb{N}$, S_{2n} denote the group of permutations of $\{1, \dots, 2n\}$; for $\sigma \in S_{2n}$, define $\Delta_\sigma = \Delta_{2n, \sigma, [s, t]} = \{(t_1, \dots, t_{2n}) \in [s, t]^{2n} : s \leq t_{\sigma(1)} \leq \dots \leq t_{\sigma(2n)} \leq t\}$. Moreover, for a given $(t_1, \dots, t_{2n}) \in \Delta_\sigma$, define $(a_k^\sigma)_{1 \leq k \leq 2n-1}$ by

$$\sum_{k=1}^n (X_{t_k} - X_{t_{k+n}}) = \sum_{k=1}^{2n-1} a_k^\sigma (X_{t_{\sigma(k+1)}} - X_{t_{\sigma(k)}})$$

and similarly $\varepsilon(k)$ by $\sum_{k=1}^n (X_{t_k} - X_{t_{k+n}}) = \sum_{k=1}^{2n} \varepsilon(k) X_{t_k}$ (equivalently, $\varepsilon(k) = 1$ for $k = 1, \dots, n$ and $\varepsilon(k) = -1$ for $k = n+1, \dots, 2n$). Then we have the following

Lemma 64. For $\sigma \in S_{2n}$ and $k = 1, \dots, 2n-1$ it holds $a_k^\sigma = -\sum_{l=1}^k \varepsilon(\sigma(l))$.

Proof. Let $\sigma \in S_{2n}$, then by definition we have $\sum_{k=1}^{2n} \varepsilon(k) X_{t_k} = \sum_{k=1}^{2n} \varepsilon(\sigma(k)) X_{t_{\sigma(k)}}$ and therefore, setting as a convention $a_0^\sigma = a_{2n}^\sigma = 0$, by definition of a_k^σ we have

$$\sum_{k=1}^{2n} \varepsilon(\sigma(k)) X_{t_{\sigma(k)}} = \sum_{k=1}^{2n-1} a_k^\sigma (X_{t_{\sigma(k+1)}} - X_{t_{\sigma(k)}}) = \sum_{k=1}^{2n-1} (a_{k-1}^\sigma - a_k^\sigma) X_{t_{\sigma(k)}}$$

which implies $a_{k-1}^\sigma - a_k^\sigma = \varepsilon(\sigma(k))$ for all k and thus the conclusion. \square

Proposition 65. Under the assumptions of Theorem 62, for any $n \in \mathbb{N}$ there exist $C_n > 0$ s.t.

$$\mathbb{E} \left[\left| \int_s^t e^{i\xi \cdot X_r} dr \right|^{2n} \right] \leq C_n |t - s|^n |\xi|^{-n/\beta} \quad \forall \xi \in \mathbb{R}^d, \quad s, t \in [0, T]. \quad (37)$$

Proof. Without loss of generality we can assume $|t - s| < \delta_{2n}$, where δ_{2n} is taken from the definition of β -LND; the general case then follows from reasoning as in the beginning of the proof of Theorem 47. We can take X of the form $X = m + Y$, where Y is a centred Gaussian β -LND process and $m = \mathbb{E}[X]$ is a deterministic continuous function. We have

$$\begin{aligned} \mathbb{E} \left[\left| \int_s^t e^{i\xi \cdot (m_r + Y_r)} dr \right|^{2n} \right] &= \left| \int_{[s,t]^{2n}} \mathbb{E} \left[\prod_{k=1}^n e^{i\xi \cdot (Y_{t_k} - Y_{t_{k+n}})} \right] \prod_{k=1}^n e^{i\xi \cdot (m_{t_k} - m_{t_{k+n}})} dt_1 \dots dt_{2n} \right| \\ &\leq \int_{[s,t]^{2n}} \left| \mathbb{E} \left[\prod_{k=1}^n e^{i\xi \cdot (Y_{t_k} - Y_{t_{k+n}})} \right] \right| dt_1 \dots dt_{2n} \\ &= \sum_{\sigma \in S_{2n}} \int_{\Delta_\sigma} \left| \mathbb{E} \left[\prod_{k=1}^n e^{i\xi \cdot (Y_{t_{\sigma(k)}} - Y_{t_{\sigma(k+n)}})} \right] \right| dt_{\sigma(1)} \dots dt_{\sigma(2n)} \\ &= \sum_{\sigma \in S_{2n}} \int_{\Delta_\sigma} \left| \mathbb{E} \left[\exp \left(i\xi \cdot \sum_{k=1}^{2n-1} a_k^\sigma (Y_{t_{\sigma(k+1)}} - Y_{t_{\sigma(k)}}) \right) \right] \right| dt_{\sigma(1)} \dots dt_{\sigma(2n)}; \end{aligned}$$

introducing the notation $s_k = t_{\sigma(k)}$, we therefore find

$$\begin{aligned} \mathbb{E} \left[\left| \int_s^t e^{i\xi \cdot (m_r + Y_r)} dr \right|^{2n} \right] &= \sum_{\sigma \in S_{2n}} \int_{\Delta_\sigma} \exp \left(-\frac{1}{2} \text{Var} \left(\xi \cdot \sum_{k=1}^{2n-1} a_k^\sigma (Y_{s_{k+1}} - Y_{s_k}) \right) \right) ds_1 \dots ds_{2n} \\ &\leq \sum_{\sigma \in S_{2n}} \int_{\Delta_\sigma} \exp \left(-c_{2n} |\xi|^2 \sum_{k=1}^{2n-1} |s_{k+1} - s_k|^{2\beta} (a_k^\sigma)^2 \right) ds_1 \dots ds_{2n}, \end{aligned}$$

where in the last passage we used the β -LND property. It is clear by definition that $\varepsilon(k) \in \{-1, 1\}$ for all k and therefore by Lemma 64 it follows that, for any odd k , $(a_k^\sigma)^2 \geq 1$. Estimating $(a_k^\sigma)^2$ trivially by 0 for even k , we obtain that, for any $\sigma \in S_{2n}$,

$$\begin{aligned} \int_{\Delta_\sigma} \exp \left(-c_{2n} |\xi|^2 \sum_{k=1}^{2n-1} |s_{k+1} - s_k|^{2\beta} (a_k^\sigma)^2 \right) ds_1 \dots ds_{2n} \\ \leq \int_{\Delta_\sigma} \exp \left(-c_{2n} |\xi|^2 \sum_{k=1}^n |s_{2k} - s_{2k-1}|^{2\beta} \right) ds_1 \dots ds_{2n}. \end{aligned}$$

Applying the change of variable $y_{2k} = s_{2k} - s_{2k-1}$ and estimating the integrals w.r.t. y_{2k} on suitable bounded intervals by the same integrals taken over $(0, \infty)$, we obtain

$$\begin{aligned} \int_{\Delta_\sigma} \exp \left(-c_{2n} |\xi|^2 \sum_{k=1}^n |s_{2k} - s_{2k-1}|^{2\beta} \right) ds_1 \dots ds_{2n} \\ \leq \int_{\Delta_\sigma} dt_1 \dots dt_n \left(\int_0^\infty e^{-c_{2n} |\xi|^2 |u|^{2\beta}} du \right)^n \\ \leq \frac{1}{n!} |t - s|^n \left(\int_0^\infty e^{-c_{2n} |\xi|^2 |u|^{2\beta}} du \right)^n. \end{aligned}$$

Summing over $\sigma \in S_{2n}$ we conclude that

$$\mathbb{E} \left[\left| \int_s^t e^{i\xi \cdot X_r} dr \right|^{2n} \right] \leq \frac{(2n)!}{n!} |t - s|^n \left(\int_0^\infty e^{-c_{2n} |\xi|^2 |u|^{2\beta}} du \right)^n \leq C_n |t - s|^n |\xi|^{-n/\beta}.$$

□

Remark 66. The constants c_{2n} , coming from the definition of β -LND, depend on n in an unspecified way; this is the reason why we are not able to obtain exponential integrability as in Theorem 53. We can overcome this difficulty by exploiting Lemma 63, up to the price of restricting ourselves to working with uniformly bounded paths and only obtaining moment estimates for the ρ -irregularity.

Proof of Theorem 62. By Proposition 65, for any $s, t \in [0, T]$ it holds

$$\begin{aligned}\mathbb{E}[\|\mu_{s,t}^X\|_{\mathcal{FL}^{\rho,2n}}^{2n}] &= \mathbb{E}\left[\int (1+|\xi|)^{2\rho n} |\Phi_{s,t}^X(\xi)|^{2n} d\xi\right] \\ &= \int (1+|\xi|)^{2\rho n} \mathbb{E}[|\Phi_{s,t}^X(\xi)|^{2n}] d\xi \\ &\lesssim_n |t-s|^n \int (1+|\xi|)^{2n\rho-n/\beta} d\xi \\ &\lesssim_{n,\rho} |t-s|^n\end{aligned}$$

whenever $2n\rho - n/\beta < -d$, namely $\rho < (1/\beta - d/n)/2$. Then by Kolmogorov continuity criterion (and the fact that $\mu_0^X \equiv 0$ by definition), for any $\rho < (1/\beta - d/n)/2$ and $\gamma < (1 - 1/n)/2$, \mathbb{P} -a.s. $\mu^X \in C_t^\gamma \mathcal{FL}_x^{\rho,2n}$ and

$$\mathbb{E}[\|\mu^X\|_{C^\gamma \mathcal{FL}^{\rho,2n}}^{2n}] < \infty.$$

By hypothesis, X is a Gaussian process with continuous trajectories, therefore by Fernique theorem $\|X\|_{C^0}$ admits moments of any order. Since $\text{supp}(\mu_{s,t}^X) \subset B_{\|X\|_{C^0}}$ uniformly in s, t , by Lemma 63 we deduce that

$$\|\mu^X\|_{C^\gamma \mathcal{FL}^{\rho,\infty}} \lesssim \|X\|_{C^0}^{d/2n} \|\mu^X\|_{C^\gamma \mathcal{FL}^{\rho,2n}}$$

and thus

$$\mathbb{E}[\|\Phi^X\|_{\mathcal{W}^{\gamma,\rho}}^n] = \mathbb{E}[\|\mu^X\|_{C^\gamma \mathcal{FL}^{\rho,\infty}}^n] \lesssim \mathbb{E}[\|X\|_{C^0}^d]^{1/2} \mathbb{E}[\|\mu^X\|_{C^\gamma \mathcal{FL}^{\rho,2n}}^{2n}]^{1/2} < \infty.$$

This implies that for any fixed n , with probability 1 X is (γ, ρ) -irregular for any $\gamma < (1 - 1/n)/2$ and $\rho < (1/\beta - d/n)/2$. We can then find a set of full probability such that X is (γ, ρ) -irregular for any $\rho < 1/(2\beta)$ and $\gamma < 1/2$; finally by the interpolation argument from Lemma 7 we can conclude that, with probability 1, X is ρ -irregular for any $\rho < 1/(2\beta)$. \square

We are finally ready to complete the

Proof of Theorem 33. The statement follows from a combination of Theorem 62 above, together with Theorem 53 and Proposition 54 from Section 4.2. \square

5 Analytic properties of ρ -irregularity

This section is devoted to the study of deterministic (γ, ρ) -irregular paths. It includes the proof of Theorems 36 and 34. In Section 5.3 we discuss also what we call the perturbation problem.

5.1 Fourier dimension and Salem sets

We highlight here the connection of ρ -irregularity and Fourier dimension and provide the proof of Theorem 34. This connection was already noticed in [14]. We start by recalling some facts of geometric measure theory, which can be found in [47].

Definition 67. Given $E \subset \mathbb{R}^d$ Borel, denote by $\mathcal{M}_+(E)$ the set of positive measures supported on E . The Fourier and Hausdorff dimension of E correspond respectively to

$$\dim_F(E) = \sup \left\{ \alpha \in [0, d] : \exists \mu \in \mathcal{M}_+(E), \mu \in \mathcal{FL}^{\alpha/2, \infty} \right\},$$

$$\dim_H(E) = \sup \left\{ \alpha \in [0, d] : \exists \mu \in \mathcal{M}_+(E), I^\alpha(\mu) < \infty \right\},$$

where

$$I^\alpha(\mu) := \int_{\mathbb{R}^{2d}} \frac{\mu(dx)\mu(dy)}{|x-y|^\alpha} = c_{\alpha,d} \int |\xi|^{\alpha-d} |\hat{\mu}(\xi)|^2 d\xi = c_{\alpha,d} \|\mu\|_{\mathcal{FL}^{\alpha/2-d/2, 2}}^2.$$

It is clear from the definition and the embedding $\mathcal{F}L^{s,\infty} \hookrightarrow \mathcal{F}L^{s-d/2-\varepsilon,2}$ that

$$0 \leq \dim_F(E) \leq \dim_H(E) \leq d; \quad (38)$$

moreover there are examples in which all inequalities are strict. This motivates the following definition.

Definition 68. A Borel set $E \subset \mathbb{R}^d$ is a Salem set if $\dim_F(E) = \dim_H(E)$.

If w is (γ, ρ) -irregular, it is clear that for any $[s, t] \subset [0, T]$ it holds

$$\mu_{s,t}^w \in \mathcal{F}L^{\rho,\infty}, \quad I^\alpha(\mu_{s,t}^w) \lesssim_\alpha |t-s|^{2\gamma} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}^2 \quad \forall \alpha < 2\rho.$$

In particular, since $\mu_{s,t}^w$ is a measure supported on $w([s, t])$, it holds

$$\min(d, 2\rho) \leq \dim_F(w([s, t])) \quad \forall [s, t] \subset [0, T].$$

On the other hand, recall that if $f \in C_t^\delta$ for some $\delta \in (0, 1]$, then for any $[s, t] \subset [0, T]$ it holds $\dim_H(f([s, t])) \leq \delta^{-1}$, see for instance Proposition 3.3 (a) from [23].

We are now ready to give the proof of Theorem 34.

Proof of Theorem 34. To prove Point *i.*, first assume $\delta \geq 1/d$. By Theorem 32, almost every $\varphi \in C_t^\delta$ is ρ -irregular for any $\rho < (2\delta)^{-1}$. It then follows from (38) and the considerations above that

$$2\rho = \min(d, 2\rho) \leq \dim_F(\varphi([s, t])) \leq \dim_H(\varphi([s, t])) \leq \delta^{-1};$$

since the inequality holds for all $\rho < (2\delta)^{-1}$, the conclusion follows. The case $\delta < 1/d$ is even more direct, since in this case we can find $\rho < (2\delta)^{-1}$ such that $2\rho > d$ and therefore we obtain

$$d = \min(d, 2\rho) \leq \dim_F(\varphi([s, t])) \leq \dim_H(\varphi([s, t])) \leq d.$$

Suppose now additionally that $\delta < (2d)^{-1}$, then almost every $\varphi \in C_t^\delta$ is ρ -irregular for some $\rho > d$; by Point *ii.* of Lemma 21 it follows that $\mu_{s,t}^\varphi$ admits a continuous occupation density $\ell_{s,t}^\varphi$. Therefore there exists $x \in \varphi([s, t])$ such that $\ell_{s,t}^\varphi(x) > \varepsilon > 0$ and by continuity the same must hold on an open ball $B(x, r)$ for some $r > 0$; this implies that $B(x, r) \subset \varphi([s, t])$. \square

It is possible to show that ρ -irregular paths cannot be δ -Hölder for δ too large reasoning by dimensionality, since otherwise it wouldn't be true that $\dim_F(w([s, t])) \geq \min(d, 2\rho)$; in the next section we are going to provide a much sharper result.

5.2 ρ -irregular paths are rough

The results of this section are inspired by the similar discussion carried out in Sections 9-11 of [37], in which it is shown that functions with sufficiently regular occupation densities must exhibit a quite erratic behaviour. Let us point out however that here we only assume the function w to be (γ, ρ) -irregular, which in general does not imply the existence of an occupation density. Theorem 36 follows from the results of this section and implies that the prevalence results from Theorem 32 are sharp (cf. Remark 39).

The next statement shows that (γ, ρ) -irregularity is indeed a notion of irregularity, in a sense that can be explicitly quantified. We recall to the reader that the critical parameter $\delta_{\gamma,\rho}^*$ is given by

$$\delta_{\gamma,\rho}^* = \frac{1-\gamma}{\rho}.$$

Throughout this section, we will always assume $\delta_{\gamma,\rho}^* \in (0, 1)$, which in view of Theorem 32 is the generic scenario for δ -Hölder continuous paths w with $\delta \in (0, 1)$, where $\delta_{\gamma,\rho}^*$ can get arbitrarily close to δ . In the case $\delta_{\gamma,\rho}^* > 1$ (resp. $\delta > 1$), many of the concepts introduced below don't have a direct analogue; here however, again by Theorem 32, we expect the path w in consideration to be such that the highest order derivative $D^{(n)}w$, for $n = \lfloor \delta \rfloor$, is again (γ', ρ') -irregular for some new parameters (γ', ρ') , with $\delta_{\gamma',\rho'}^* < 1$; the results presented here can then be applied successfully to $D^{(n)}w$.

Theorem 69. *Let w be a (γ, ρ) -irregular function. Then for any $\delta > \delta_{\gamma, \rho}^*$, w is nowhere δ -Hölder continuous. In particular, for any fixed $M > 0$ and any $s \in [0, T]$, the set of points t around s which satisfy an approximate δ -Hölder condition with constant M is a zero density set:*

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{\mathcal{L}(B(s, \varepsilon) \cap [0, T])} \mathcal{L}(t \in B(s, \varepsilon) \cap [0, T] : |w_t - w_s| \leq M|t - s|^\delta) = 0.$$

Proof. First consider the case of $\rho \leq d$. Let $\psi \in C_c^\infty(\mathbb{R}^d)$ be a radially symmetric, decreasing function such that $\psi \equiv 1$ on B_1 . Fix $M > 0$ and let us consider first $s \in (0, T)$, so that $B(s, \varepsilon) \cap [0, T] = (s - \varepsilon, s + \varepsilon)$ for ε small enough; up to translation we can assume $w_s = 0$. We have

$$\begin{aligned} \frac{1}{2\varepsilon} \mathcal{L}(t \in (s - \varepsilon, s + \varepsilon) : |w_t - w_s| \leq M|t - s|^\delta) &\leq \frac{1}{2\varepsilon} \mathcal{L}(t \in (s - \varepsilon, s + \varepsilon) : |w_t| \leq M\varepsilon^\delta) \\ &\leq \frac{1}{2\varepsilon} \mathcal{L}\left(t \in (s - \varepsilon, s + \varepsilon) : \psi\left(\frac{w_t}{M\varepsilon^\delta}\right) \geq 1\right) \\ &\leq \frac{1}{2\varepsilon} \int_{s-\varepsilon}^{s+\varepsilon} \psi\left(\frac{w_t}{M\varepsilon^\delta}\right) dt \\ &= \frac{1}{2\varepsilon} \int_{\mathbb{R}^d} \psi\left(\frac{y}{M\varepsilon^\delta}\right) \mu_{s-\varepsilon, s+\varepsilon}^w(dy) \\ &= \frac{1}{2} M^d \varepsilon^{d\delta-1} \int_{\mathbb{R}^d} \hat{\psi}(M\varepsilon^\delta \xi) \overline{\hat{\mu}_{s-\varepsilon, s+\varepsilon}^w}(\xi) d\xi \\ &\lesssim_M \varepsilon^{\gamma+\delta d-1} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \int_{\mathbb{R}^d} |\hat{\psi}(M\varepsilon^\delta \xi)| (1 + |\xi|)^{-\rho} d\xi. \end{aligned}$$

By Hölder inequality, for any $q > d/\rho \geq 1$, setting $1/p = 1 - 1/q$ it holds

$$\int_{\mathbb{R}^d} |\hat{\psi}(M\varepsilon^\delta \xi)| (1 + |\xi|)^{-\rho} d\xi \lesssim_{M, q} \varepsilon^{-\delta d/p} \|\hat{\psi}\|_{L^p}.$$

Therefore we obtain

$$\frac{1}{2\varepsilon} \mathcal{L}(t \in (s - \varepsilon, s + \varepsilon) : |w_t - w_s| \leq M|t - s|^\delta) \lesssim_{M, q} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \varepsilon^{\gamma+\delta d/q-1},$$

where the last quantity is infinitesimal as $\varepsilon \rightarrow 0$ for any q such that $d/q < \rho$ and $\gamma + \delta d/q > 1$. In particular if δ satisfies $\delta > \delta_{\gamma, \rho}^*$, then it's always possible to find such q , which gives the conclusion for $s \in [0, T]$. The reasoning at the endpoints $\{0, T\}$ is analogous: for instance in the case $s = 0$, similar calculations yield

$$\frac{1}{\varepsilon} \mathcal{L}(t \in (0, \varepsilon) : |w_t - w_0| \leq M|t|^\delta) \lesssim \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \varepsilon^{\gamma+\delta d/q-1}.$$

This concludes the proof in the case $\rho \leq d$. For $\rho > d$, by Lemma 7 and equation (4), we can find a new pair $(\tilde{\gamma}, \tilde{\rho})$ with $\tilde{\rho} \leq d$ such that w is $(\tilde{\gamma}, \tilde{\rho})$ -irregular and $\delta_{\gamma, \rho}^* = \delta_{\tilde{\gamma}, \tilde{\rho}}^*$; the conclusion then follows from the previous case. \square

Remark 70. *It is clear from the proof that the statement can be localised as follows. For a fixed $s \in [0, T]$, if the map $t \mapsto \mu_t^w$ satisfies an approximate γ -Holder condition in $\mathcal{FL}^{\rho, \infty}$ around s , namely there exist constants $C, r > 0$ such that*

$$\|\mu_{s, t}^w\|_{\mathcal{FL}^{\rho, \infty}} \leq C|t - s|^\gamma \quad \text{for all } t \in (s - r, s + r),$$

then w cannot satisfy an approximate δ -Holder condition around s for any $\delta > \delta_{\gamma, \rho}^$.*

Let us recall that for a given path $w \in C_t^0$, its p -variation on an interval $[s, t] \subset [0, T]$ is defined as

$$\|w\|_{p\text{-var}; [s, t]} = \left(\sup_{\Pi} \sum_i |w_{t_i, t_{i+1}}|^p \right)^{1/p}$$

where the supremum is taken over all possible finite partitions $\Pi = \{s = t_0 < \dots < t_n = t\}$ of $[s, t]$.

Corollary 71. *Let w be a (γ, ρ) -irregular function. Then for any $p < (\delta_{\gamma, \rho}^*)^{-1}$ and any $[s, t] \subset [0, T]$, $\|w\|_{p\text{-var}; [s, t]} = +\infty$.*

Proof. Since the ρ -irregularity property is scaling invariant and the p -variation is invariant under reparametrization, it suffices to show that if w is (γ, ρ) -irregular, then $\|w\|_{p\text{-var}, [0, 1]} = \infty$ for any p as above. Going through analogous computations to those of Theorem 69, it can be shown that for any $\delta > \delta_{\gamma, \rho}^*$ it holds

$$\lim_{\varepsilon \rightarrow 0^+} \sup_{s \in [0, 1-\varepsilon]} \frac{1}{\varepsilon} \mathcal{L}(t \in (s, s+\varepsilon) : |w_{s,t}| \leq \varepsilon^\delta) = 0.$$

In particular, for all $\varepsilon > 0$ small enough it holds

$$\sup_{s \in [0, 1-\varepsilon]} \mathcal{L}(t \in (s, s+\varepsilon) : |w_{s,t}| > \varepsilon^\delta) > \frac{\varepsilon}{2} > 0;$$

thus for any $s \in [0, 1-\varepsilon]$, there exists $t \in (s, s+\varepsilon)$ such that $|w_{s,t}| > \varepsilon^\delta$. Taking $n \sim 1/\varepsilon$, we can construct a partition $\{0 = t_0 < \dots < t_{2n} = 1\}$ such that $t_{2k} = k\varepsilon$ and $t_{2k+1} \in (t_{2k}, t_{2k+2})$ has the above property. We obtain

$$\|w\|_{p\text{-var}}^p \geq \sum_{k=0}^{2n-1} |w_{t_k, t_{k+1}}|^p \gtrsim \varepsilon^{p\delta-1},$$

since ε can be taken arbitrarily small, if $p < 1/\delta$ then $\|w\|_{p\text{-var}} = \infty$. As the reasoning holds for any $\delta > \delta_{\gamma, \rho}^*$, the conclusion follows. \square

Theorem 69 suggests that the behaviour of w is quite wild. This intuition can be captured by the following notions of irregularity introduced in [31] and nicely presented in [32].

Definition 72. *We say that a path $w \in C_t^\delta$ is rough at time s , $s \in (0, T)$, if*

$$\limsup_{t \rightarrow s} \frac{|v \cdot w_{s,t}|}{|t-s|^{2\delta}} = +\infty \quad \forall v \in \mathbb{S}^{d-1}.$$

A path w is truly rough if it is rough on some dense set of $[0, T]$.

Definition 73. *A path $w \in C_t^\delta$ is θ -Hölder rough for $\theta \in (0, 1)$ on scale ε_0 if there exists a constant $L := L_\theta(w) := L(\theta, \varepsilon_0, T; w) > 0$ such that for every $v \in \mathbb{S}^{d-1}$, $s \in [0, T]$ and $\varepsilon \in (0, \varepsilon_0]$ there exists $t \in [0, T]$ such that*

$$|t-s| < \varepsilon \quad \text{and} \quad |v \cdot (w_{s,t})| \geq L_\theta(w) \varepsilon^\theta.$$

The largest such value of L is called the modulus of θ -Hölder roughness of w .

Corollary 74. *Let w be a (γ, ρ) -irregular path; then for any $\theta > \delta_{\gamma, \rho}^*$, w is θ -Hölder rough with infinite modulus of θ -Hölder roughness.*

Proof. For simplicity we show all the properties for s away from the endpoints $\{0, T\}$, but it is easy to check how all the reasonings can be adapted in the other case. Recall that, if w is (γ, ρ) -irregular and $v \in \mathbb{S}^{d-1}$, then $v \cdot w$ is (γ, ρ) -irregular and $\|\Phi^{v \cdot w}\|_{\mathcal{W}^{\gamma, \rho}} \leq \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}$. The calculations in the proof of Theorem 69 show that, for any $\delta > \delta_{\gamma, \rho}^*$ and any $M > 0$,

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\varepsilon} \mathcal{L}\{t \in (s-\varepsilon, s+\varepsilon) : |v \cdot w_{s,t}| \geq M\varepsilon^\delta\} = 1,$$

where the rate of convergence only depends on M and $\|\Phi^{v \cdot w}\|_{\mathcal{W}^{\gamma, \rho}} \leq \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}$ and it is thus uniform in s and v . For fixed M we can then find $\varepsilon_0 = \varepsilon_0(M, \delta)$ sufficiently small such that, it holds

$$\frac{1}{2\varepsilon} \mathcal{L}\{t \in (s-\varepsilon, s+\varepsilon) : |v \cdot w_{s,t}| \geq M\varepsilon^\delta\} \geq \frac{1}{2},$$

for all $\varepsilon \in (0, \varepsilon_0]$, uniformly in s and v . Since the set has non-zero Lebesgue measure, it's always possible to find $t \in (s-\varepsilon, s+\varepsilon)$ such that $|v \cdot w_{s,t}| \geq M\varepsilon^\delta$, which shows that the definition of θ -Hölder roughness is satisfied with $\theta = \delta$ and $L \geq M$. By the arbitrariness of M we can conclude. \square

Proof of Theorem 36. Follows from a combination of Corollaries 71 and 74. \square

We conclude this subsection with the promised claim from Remark 39 that typical fBm paths cannot be (γ, ρ) -irregular for values $\rho > (2H)^{-1}$; the proof can also be readapted to consider other Gaussian processes.

Lemma 75. *Let W^H be a fBm of parameter H , then for any $s < t$ and any $\rho > (2H)^{-1}$ it holds*

$$\mathbb{E}[\|\mu_{s,t}\|_{\mathcal{FL}^{\rho,\infty}}^2] = \infty.$$

Proof. Up to rescaling, we can assume $s = 0$, $t = 1$. Since $\mathcal{FL}^{\rho,\infty} \hookrightarrow H^{\rho-d/2-}$, in order to conclude it suffices to show that

$$\mathbb{E}[\|\mu_1\|_{H^{1/(2H)-d/2}}^2] = \infty.$$

This quantity can now be computed explicitly:

$$\begin{aligned} \mathbb{E}[\|\mu_1\|_{H^{1/(2H)-d/2}}^2] &= \mathbb{E}\left[\int_{\mathbb{R}^d} |\xi|^{1/H-d} \left|\int_0^1 e^{i\xi \cdot W_s^H} ds\right|^2 d\xi\right] \\ &= \int_{\mathbb{R}^d} \int_{[0,1]^2} |\xi|^{1/H-d} \mathbb{E}[e^{i\xi \cdot (W_{s,t}^H)}] dt ds d\xi \\ &= \int_{\mathbb{R}^d} \int_{[0,1]^2} |\xi|^{1/H-d} \exp\left(-\frac{|\xi|^2 |t-s|^{2H}}{2}\right) dt ds d\xi \\ &= \int_{\mathbb{R}^d} |\tilde{\xi}|^{1/H-d} e^{-|\tilde{\xi}|^2/2} d\tilde{\xi} \cdot \int_{[0,1]^2} |t-s|^{-1} dt ds = \infty \end{aligned}$$

where in the last passage we use the change of variables $\tilde{\xi} = \xi|t-s|^H$. \square

5.3 The general perturbation problem

As in the previous section, here we will work exclusively with (γ, ρ) -irregular paths w such that $\delta_{\gamma,\rho}^* < 1$.

The perturbation problem was first introduced in Section 12 of [37], in the context of paths which admit an occupation density. In the case of ρ -irregularity, it can be reformulated as:

If w is ρ -irregular and φ is a sufficiently regular function, is $w + \varphi$ still ρ -irregular?

We address here the more general question:

Which classes of transformations preserve the property of (γ, ρ) -irregularity?

It follows from the results of the previous section that good candidates are transformations which preserve the very oscillatory behaviour of w , namely at least the property that

$$\limsup_{t \rightarrow s} \frac{|v \cdot w_{s,t}|}{|t-s|^\delta} = +\infty \quad \text{for all } s \in (0, T), v \in \mathbb{S}^{d-1}, \delta > \delta_{\gamma,\rho}^*. \quad (39)$$

Interestingly, it turns out that several transformations $F : C_t^0 \rightarrow C_t^0$ have the property that if w is (γ, ρ) -irregular, then $F(w)$ is $(\tilde{\gamma}, \tilde{\rho})$ -irregular with parameters such that $\delta_{\gamma,\rho}^* = \delta_{\tilde{\gamma},\tilde{\rho}}^*$, so that property (39) is preserved. In many cases we are however unable to show that $(\gamma, \rho) = (\tilde{\gamma}, \tilde{\rho})$, which remains a major open problem. A notable exception is given by the additive perturbations $F(w) = w + \varphi$ with $\varphi \in C_t^\infty$, whose treatment is postponed to the next subsection.

We start by showing that (γ, ρ) -irregularity is invariant under sufficiently regular time reparametrization.

Lemma 76. Let w be (γ, ρ) -irregular, $g \in C_t^\beta$ with $\beta + \gamma > 1$. Then

$$\left| \int_s^t e^{i\xi \cdot w_r} g_r dr \right| \lesssim \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \|g\|_{C^\beta} |t - s|^\gamma |\xi|^{-\rho} \quad \text{uniformly in } \xi \in \mathbb{R}^d.$$

In particular, for β as above, let $\tau : [0, T] \rightarrow [\tau(0), \tau(T)]$ be a $C_t^{1+\beta}$ -diffeomorphism, i.e. $\tau \in C_t^{1+\beta}$ is invertible on its image with inverse of class $C_t^{1+\beta}$. Then $\tilde{w}_r := w_{\tau^{-1}(r)}$ is also ρ -irregular and

$$\|\Phi^{\tilde{w}}\|_{\mathcal{W}^{\gamma, \rho}} \lesssim \|\tau^{-1}\|_{C^1}^\gamma \|\tau\|_{C^{1+\beta}} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}. \quad (40)$$

Proof. Let w, g be as above. Then by properties of Young integral it holds

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot w_r} g_r dr \right| &= \left| \int_s^t g_r d \left(\int_s^r e^{i\xi \cdot w_u} du \right) \right| \\ &\lesssim |g_s| \left| \int_s^t e^{i\xi \cdot w_r} dr \right| + |t - s|^{\beta+\gamma} \|g\|_{C^\beta} \left\| \int_s^\cdot e^{i\xi \cdot w_r} dr \right\|_{C^\gamma} \\ &\lesssim |t - s|^\gamma |\xi|^{-\rho} \|g\|_{C^\beta} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}, \end{aligned}$$

which gives the first claim. Applying the change of variables $\tilde{r} = \tau^{-1}(r)$, we then have

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot \tilde{w}_r} dr \right| &= \left| \int_{\tau^{-1}(s)}^{\tau^{-1}(t)} e^{i\xi \cdot w_r} \tau_r' dr \right| \lesssim |\tau^{-1}(t) - \tau^{-1}(s)|^\gamma |\xi|^{-\rho} \|\tau'\|_{C^\beta} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \\ &\lesssim |t - s|^\gamma |\xi|^{-\rho} \|\tau^{-1}\|_{C^1}^\gamma \|\tau\|_{C^{1+\beta}} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \end{aligned}$$

which implies (40). \square

Remark 77. We have only defined the notion of ρ -irregularity in terms of push-forward under w of the Lebesgue measure $\mathcal{L}_{[s,t]}$, but we could consider more generally the Fourier transform of the push-forward under w of a bounded Borel measure ν on $[0, T]$, namely

$$\widehat{w_* \nu_{s,t}}(\xi) = \int_{[s,t]} e^{i\xi \cdot w_r} d\nu(r).$$

The first part of previous lemma could then be interpreted as follows: if w is (γ, ρ) -irregular w.r.t. \mathcal{L} , then it is also (γ, ρ) -irregular w.r.t. any measure $\nu \ll \mathcal{L}$ with a sufficiently regular density g .

Remark 78. Lemma 76 can also be used to further enlarge the class of stochastic processes X which are ρ -irregular: given any such X and any random $C_t^{3/2}$ -diffeomorphism, $Y_t := X_{\tau^{-1}(t)}$ is still a ρ -irregular process.

Let us make some considerations based on the result above. Recall that if $f \in C_t^\delta$ for $\delta \in (0, 1)$ and τ is sufficiently regular (i.e. bi-Lipschitz), then $f \circ \tau^{-1}$ is still C_t^δ , but this is not true for a general homeomorphism τ . On the other hand, if $f \in C_t^0$ has finite $1/\delta$ -variation, then there exist a homeomorphism τ and $g \in C_t^\delta$ such that $f \circ \tau^{-1} = g$ (see for instance Proposition 5.15 from [33]). Moreover the $1/\delta$ -variation is a quantity invariant under time reparametrization. Lemma 76 suggests that the situation here could be similar: (γ, ρ) -irregularity is preserved only if the reparametrization is smooth enough, but there might exist another underlying property which is invariant under a larger class of homeomorphism τ . We formulate this as a conjecture.

Conjecture 79. For any pair (γ, ρ) , there exists a property \mathcal{P} such that:

1. For any $f \in C_t^0$ with property \mathcal{P} there exists a homeomorphism τ such that $g = f \circ \tau^{-1}$ is (γ, ρ) -irregular.
2. The property \mathcal{P} is invariant under time reparametrization.

In the rest of the section, we will address the perturbation problem only for transformations $z = F(w)$ with a very specific structure, which makes z locally look like w . The treatment is a bit abstract, but simple examples will be given in Remark 82.

Definition 80. Let w be (γ, ρ) -irregular. We say that $z : [0, T] \rightarrow \mathbb{R}^d$ is controlled by w with “derivative” z' if there exist $z' \in C^0([0, T]; \mathbb{R}^{d \times d})$ and $R \in C_2^\beta([0, T]; \mathbb{R}^d)$ with $\beta > \delta_{\gamma, \rho}^*$ such that

$$z_{s,t} = z'_s w_{s,t} + R_{s,t} \quad \text{for all } (s, t) \in \Delta_2.$$

Here $\Delta_2 = \{(s, t) : 0 \leq s < t \leq T\}$ and $R \in C_2^\beta([0, T]; \mathbb{R}^d)$ means that $R : \Delta_2 \rightarrow \mathbb{R}^d$ and it satisfies

$$\|R\|_\beta := \sup_{s < t} \frac{|R_{s,t}|}{|t - s|^\beta} < \infty.$$

The definition of controlled paths is usually given in the rough paths framework, see for instance [38] and [32]; however here we do not impose $w, z \in C_t^\alpha$ with $R \in C_2^{2\alpha}$ and we do not require w to admit a rough lift.

In order for Definition 80 to be meaningful, we need to verify that the pair (z', R) is unique; this follows from the condition $\beta > \delta_{\gamma, \rho}^*$ and property (39), which follows from Corollary 74. Indeed, let (\tilde{z}', \tilde{R}) be another such pair and set $A = z' - \tilde{z}'$, $B = R - \tilde{R}$. Choosing $\delta \in (0, 1)$ such that $\delta_{\gamma, \rho}^* < \delta < \beta$, for any $s \in (0, T)$ and any $v \in \mathbb{S}^{d-1}$ it holds

$$\limsup_{t \rightarrow s} \frac{|(A_s^T v) \cdot w_{s,t}|}{|t - s|^\delta} = \limsup_{t \rightarrow s} \frac{|B_{s,t}|}{|t - s|^\delta} \leq \|B\|_\beta \limsup_{t \rightarrow s} |t - s|^{\beta - \delta} = 0$$

which implies by (39) that $A_s^T v = 0$ for all $v \in \mathbb{S}^{d-1}$ and $s \in (0, T)$, thus $A \equiv 0$ and $B \equiv 0$ as well.

We will from now on assume in addition that there exists $c > 0$ such that

$$z'_s (z'_s)^T \geq c^2 I_d \quad \forall s \in [0, T]. \quad (41)$$

In particular, the above non-degeneracy condition implies that z satisfies property (39) as well.

Proposition 81. Let w be (γ, ρ) -irregular, z controlled by w with z' satisfying (41). Then there exists $\tilde{\gamma} > \gamma$ such that z is $(\tilde{\gamma}, \tilde{\rho})$ -irregular and $\tilde{\rho}$ is given by

$$\tilde{\rho} = \frac{\beta}{1 - \gamma + \beta} \rho - \frac{1 - \gamma}{1 - \gamma + \beta} > 0;$$

moreover $\delta_{\gamma, \rho}^* = \delta_{\tilde{\gamma}, \tilde{\rho}}^*$ and we have the estimate

$$\|\Phi^z\|_{\mathcal{W}^{\tilde{\gamma}, \tilde{\rho}}} \lesssim 1 + \|R\|_\beta + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}.$$

Proof. Let $\xi \in \mathbb{R}^d$, $|\xi| \geq 1$ be fixed. For any $s < t$, it holds

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot z_r} dr \right| &= \left| \int_s^t e^{i\xi \cdot z_{s,r}} dr \right| \\ &\leq \left| \int_s^t [e^{i\xi \cdot z_{s,r}} - e^{i\xi \cdot z'_s w_{s,r}}] dr \right| + \left| \int_s^t e^{i\xi \cdot z'_s w_{s,r}} dr \right| \\ &\lesssim \int_s^t |\xi| |R_{s,r}| dr + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} |\xi|^{-\rho} |t - s|^\gamma \\ &\lesssim \|R\|_\beta |\xi| |t - s|^{1+\beta} + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} |\xi|^{-\rho} |t - s|^\gamma. \end{aligned}$$

First assume that $|t - s|^{1-\gamma+\beta} |\xi|^{1+\rho} \leq 1$, so that $|\xi| |t - s|^{1+\beta} \leq |\xi|^{-\rho} |t - s|^\gamma$, then in this case we trivially get

$$\left| \int_s^t e^{i\xi \cdot z_r} dr \right| \lesssim (\|R\|_\beta + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}) |\xi|^{-\rho} |t - s|^\gamma. \quad (42)$$

Assume now that $|t-s|^{1-\gamma+\beta}|\xi|^{1+\rho} > 1$; choose $N \in \mathbb{N}$ such that $N^{1-\gamma+\beta} \sim |t-s|^{1-\gamma+\beta}|\xi|^{1+\rho}$ and split the interval $[s, t]$ in N subintervals of size $|t-s|/N$. Applying the previous estimate to each of them and summing over we obtain

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot z_r} dr \right| &\lesssim \|R\|_\beta N^{-\beta} |\xi| |t-s|^{1+\beta} + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}} N^{1-\gamma} |t-s|^\gamma |\xi|^{-\rho} \\ &\sim (\|R\|_\beta + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}) |t-s| |\xi|^{-\tilde{\rho}} \end{aligned} \quad (43)$$

where

$$\tilde{\rho} = \frac{\beta}{1-\gamma+\beta} \rho - \frac{1-\gamma}{1-\gamma+\beta} = \lambda \rho + \lambda - 1$$

for suitable choice of $\lambda \in (0, 1)$, which implies that $\tilde{\rho} < \rho$; moreover $\tilde{\rho} > 0$, since $\rho > (1-\lambda)/\lambda = (1-\gamma)/\beta$ thanks to the condition $\beta > \delta_{\gamma,\rho}^*$.

Next consider $\theta \in (0, 1)$ such that $\tilde{\rho} = \theta \rho$; when $|t-s|^{1-\gamma+\beta}|\xi|^{1+\rho} \leq 1$, interpolating between (42) and the trivial bound $|\int_s^t e^{i\xi \cdot z_r} dr| \leq |t-s|$, we obtain

$$\left| \int_s^t e^{i\xi \cdot z_r} dr \right| \lesssim (\|R\|_\beta + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}})^\theta |t-s|^{\tilde{\gamma}} |\xi|^{-\tilde{\rho}}, \quad (44)$$

where $\tilde{\gamma} = \theta \gamma + 1 - \theta$ is such that $\tilde{\gamma} > \gamma$ and $\delta_{\gamma,\rho}^* = \delta_{\tilde{\gamma},\tilde{\rho}}^*$ (cf. Lemma 7). Combining the bounds (43) and (44), using the simple bound $\max\{a^\theta, a\} \leq 1 + a$, we arrive at

$$\left| \int_s^t e^{i\xi \cdot z_r} dr \right| \lesssim_T (1 + \|R\|_\beta + c^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma,\rho}}) |\xi|^{-\tilde{\rho}} |t-s|^{\tilde{\gamma}} \quad \forall |\xi| \geq 1, s, t \in [0, T]$$

without conditions on $|t-s|^{1-\gamma+\beta}|\xi|^{1+\rho}$, which gives the conclusion. \square

Remark 82. If for instance w is ρ -irregular and $\beta \geq 1$, then we obtain that z is $\tilde{\rho}$ -irregular with

$$\tilde{\rho} \geq \frac{2}{3} \rho - \frac{1}{3}.$$

Examples of z satisfying the assumptions of Proposition 81 are the following:

- Take $z_t = \varphi_t w_t$ with $\varphi \in C^\beta([0, T]; \mathbb{R})$ satisfying $\varphi_t \geq c > 0$, then $z'_s = \varphi_s I_d$, $R_{s,t} = w_t \varphi_{s,t} \in C_2^\beta$ (here $\beta \leq 1$).
- Suppose $w \in C_t^\delta$ with $\delta \in (0, 1)$ and take $z_t = \int_0^t A_s dw_s$, where $A \in C^\alpha([0, T]; \mathbb{R}^{d \times d})$ satisfies (41), $\alpha + \delta > 1$ and the integral is defined in the Young sense. Then $z'_t = A_t$ and $\beta = \alpha + \delta > 1$.
- Finally, if $z = w + \varphi$ with $\varphi \in C_t^\beta$, then $z' \equiv I_d$ and $R_{s,t} = \varphi_{s,t} \in C_2^\beta$ (with $\beta \leq 1$); this case is however quite special and better estimates are available, see Section 5.4 below.

Let us highlight the difference between the purely analytical result of Proposition 81 compared to the probabilistic result of Proposition 57, in which instead we have examples of Gaussian processes which are ρ -irregular with parameter ρ invariant under any of the deterministic transformations from the list above.

There is another notable class of transformations which preserve some properties of the occupation measure μ^w . In this case however it is rather complicated to consider the (γ, ρ) -irregularity property and it is instead more natural to reason with occupation densities. Suppose that w admits an occupation density $\ell_{s,t}^w$ (which we know to be true by Lemma 21 if for instance w is ρ -irregular with $\rho > d/2$) and let $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a global diffeomorphism; define $z_t = F(w_t)$. Then z still admits an occupation density $\ell_{s,t}^z$, since

$$\int_s^t \varphi(z_r) dr = \int_s^t \varphi(F(w_r)) dr = \int_{\mathbb{R}^d} \varphi(F(x)) \ell_{s,t}^w(x) dx = \int_{\mathbb{R}^d} \varphi(x) |\det(DF^{-1}(x))| \ell_{s,t}^w(F^{-1}(x)) dx$$

which shows that

$$\mu_{s,t}^z(dx) = |\det(DF^{-1}(x))| \ell_{s,t}^w(F^{-1}(x)) dx = \ell_{s,t}^z(x) dx.$$

This also implies that $\ell_{s,t}^z$ inherits the regularity of $\ell_{s,t}^w$ and F ; for instance if $\ell^w \in C_t^\gamma L_x^2$, then

$$\begin{aligned} \|\ell_{s,t}^z\|_{L^2}^2 &= \int_{\mathbb{R}^d} |\det(DF^{-1}(x))|^2 |\ell_{s,t}^w(F^{-1}(x))|^2 dx \\ &= \int_{\mathbb{R}^d} |\det(DF(x))|^{-1} |\ell_{s,t}^w(x)|^2 dx \\ &\leq \|DF^{-1}\|_{L^\infty} \|\ell^w\|_{C^\gamma L^2}^2 |t-s|^{2\gamma}. \end{aligned}$$

Similar estimates hold if $\ell^w \in C_t^\gamma L_x^p$ or if $\ell^w \in C_t^\gamma C_x^0$.

5.4 The additive perturbation problem

In this section we treat for simplicity only the case $w \in C_t^\delta$ with $\delta \in (0, 1)$. In view of Theorem 32, we will always assume $\rho > 1/2$ (equivalently $(2\rho)^{-1} < 1$).

We first present a partial result, which is a slight improvement of Theorem 1.6 from [10].

Lemma 83. *Let w be (γ, ρ) -irregular and $\varphi \in C_t^\beta$ for some $\beta \in (0, 1]$ such that $\beta > \delta_{\gamma, \rho}^*$. Then for any choice of $\delta \leq \beta$ satisfying $1 - \gamma < \delta < \beta\rho$, $w + \varphi$ is $(\tilde{\gamma}, \tilde{\rho})$ -irregular for the choice*

$$\tilde{\gamma} = \gamma \left(1 - \frac{\delta}{\beta\rho}\right) + \frac{\delta}{\beta\rho}, \quad \tilde{\rho} = \rho - \frac{\delta}{\beta}, \quad (45)$$

and it holds

$$\|\Phi^{w+\varphi}\|_{\mathcal{W}^{\tilde{\gamma}, \tilde{\rho}}} \lesssim (1 + \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}})(1 + \|\varphi\|_{C^\beta}^{\delta/\beta}).$$

If w is ρ -irregular and $\beta > \max\{1/2, (2\rho)^{-1}\}$, then $w + \varphi$ is $\left(\frac{1}{2} + \frac{1}{4\beta\rho}, \rho - \frac{1}{2\beta}\right)$ -irregular with

$$\|\Phi^{w+\varphi}\|_{\mathcal{W}^{\tilde{\gamma}, \tilde{\rho}}} \lesssim (1 + \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}})(1 + \|\varphi\|_{C^\beta}^{1/2\beta}).$$

Proof. Since $\varphi \in C_t^\beta$, so does $e^{i\xi \cdot \varphi}$, for all $\xi \in \mathbb{R}^d$. For any $\delta > 1 - \gamma$ we can then apply the estimates from Young integration as follows:

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| &= \left| \int_s^t e^{i\xi \cdot \varphi_r} d \left(\int_s^r e^{i\xi \cdot w_u} du \right) \right| \\ &\lesssim_{\delta+\gamma} |e^{i\xi \cdot \varphi_s}| \left| \int_s^t e^{i\xi \cdot w_r} dr \right| + |t-s|^{\gamma+\delta} \|e^{i\xi \cdot \varphi}\|_{C^\delta} \left\| \int_s^\cdot e^{i\xi \cdot w_r} dr \right\|_{C^\gamma} \\ &\leq \left| \int_s^t e^{i\xi \cdot w_r} dr \right| + |t-s|^{\gamma+\delta} |\xi|^{-\rho} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \|e^{i\xi \cdot \varphi}\|_{C^\delta}. \end{aligned}$$

Now since $\varphi \in C_t^\beta$, by interpolation we have (we are using the hypothesis $\delta \leq \beta$)

$$|e^{i\xi \cdot \varphi_t} - e^{i\xi \cdot \varphi_s}| \leq 2, \quad |e^{i\xi \cdot \varphi_t} - e^{i\xi \cdot \varphi_s}| \leq |\xi| |t-s|^\beta \|\varphi\|_{C^\beta} \Rightarrow |e^{i\xi \cdot \varphi_t} - e^{i\xi \cdot \varphi_s}| \lesssim \|\varphi\|_{C^\beta}^{\delta/\beta} |\xi|^{\delta/\beta} |t-s|^\delta;$$

similarly for any $\theta \in (0, 1)$ it holds

$$\left| \int_s^t e^{i\xi \cdot w_r} dr \right| \lesssim \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}^\theta |t-s|^{\gamma\theta+1-\theta} |\xi|^{-\theta\rho}$$

and so putting everything together we obtain

$$\left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| \lesssim \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}}^\theta |t-s|^{\gamma\theta+1-\theta} |\xi|^{-\theta\rho} + |t-s|^{\gamma+\delta} |\xi|^{-\rho+\delta/\beta} \|\Phi^w\|_{\mathcal{W}^{\gamma, \rho}} \|\varphi\|_{C^\beta}^{\delta/\beta}.$$

Choosing $\theta \in (0, 1)$ such that $\theta\rho = \rho - \delta/\beta$, namely $\theta = 1 - \delta/(\beta\rho)$ we obtain the first statement. The second one simply follows from the assumption $\gamma > 1/2$, taking $\delta = 1/2$. \square

The partial result above implies that, even if we consider a perturbation $\varphi \in C_t^1$, we should expect a loss in spatial regularity of order $1/2$, which is only partially recovered by an improvement in time regularity of order $1/4\rho$. The new parameters $(\tilde{\gamma}, \tilde{\rho})$ given by Lemma 45 satisfy $\delta_{\tilde{\gamma}, \tilde{\rho}}^* = \delta_{\gamma, \rho}^*$, which implies that $w + \varphi$ still satisfies property (39), as can be checked directly using the fact that $\varphi \in C_t^\beta$ for some $\beta > \delta_{\gamma, \rho}^*$. This hints that the above result, while not being fully satisfactory, might be optimal, even if we cannot exclude the existence of other pairs (γ', ρ') with $\rho' > \tilde{\rho}$ such that $w + \varphi$ is (γ', ρ') -irregular.

The proof above cannot provide better results in the case $\varphi \in C_t^\beta$ with $\beta > 1$. Even if it were false in general that $w + \varphi$ is (γ, ρ) -irregular whenever w is so and $\varphi \in C_t^\beta$ with $\beta > \delta_{\gamma, \rho}^*$, we would at least expect the claim to be true whenever φ is C_t^∞ ; this is a conjecture left open in [10].

We can give it a positive answer, up to strengthening the notion of ρ -irregularity. Before giving the rigorous statement, let us give an intuition by considering the following case. Suppose that $\varphi \in C_t^{1+\beta}$ for some $\beta \in [0, 1]$ and suppose that w satisfies the following property: for any $a \in \mathbb{R}^d$, $t \mapsto w_t + at$ is ρ -irregular, uniformly in a , in the sense that $\sup_a \|\Phi^{w+at}\|_{\mathcal{W}^{\gamma, \rho}} < \infty$. Then we can improve the previous estimates as follows:

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| &= \left| e^{-i\xi \cdot (\varphi_s - s\varphi'_s)} \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| \\ &= \left| \int_s^t e^{i\xi \cdot (\varphi_{s,r} - \varphi'_s(r-s))} d \left(\int_s^r e^{i\xi \cdot (w_u + \varphi'_s u)} du \right) \right| \\ &\lesssim \|\Phi^{w+\varphi'_s t}\|_{\mathcal{W}^{\gamma, \rho}} |t-s|^\gamma |\xi|^{-\rho} + \|\Phi^{w+\varphi'_s t}\|_{\mathcal{W}^{\gamma, \rho}} |t-s| \|e^{i\xi \cdot (\varphi_{s,\cdot} - \varphi'_s(\cdot-s))}\|_{C^{1/2}} |\xi|^{-\rho} \\ &\lesssim_w |t-s|^\gamma |\xi|^{-\rho} + |t-s| |\xi|^{-\rho} \|e^{i\xi \cdot (\varphi_{s,\cdot} - \varphi'_s(\cdot-s))}\|_{C^{1/2}} \end{aligned}$$

where the last norm is taken over the interval $[s, t]$. As before, we can estimate it using simple interpolation arguments, only this time we have

$$\begin{aligned} |\varphi_{s,u} - \varphi'_s(u-s) - \varphi_{s,v} + \varphi'_s(v-s)| &= |\varphi_{u,v} - \varphi'_s(v-u)| \\ &= \left| \int_v^u \varphi'_{r,s} dr \right| \\ &\leq \int_v^u \|\varphi'\|_{C^\beta} |r-s|^\beta dr \\ &\lesssim_\varphi |u-v|^{1/2} |t-s|^{1/2+\beta} \end{aligned}$$

where we used the fact that $[v, u] \subset [s, t]$. Therefore we obtain

$$\left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| \lesssim |t-s|^\gamma |\xi|^{-\rho} + |t-s|^{3/2+\beta} |\xi|^{1-\rho}$$

and we can now reason as in the proof of Lemma 81, i.e. split the interval $[s, t]$ into N subintervals of size $|t-s|/N$, apply the estimate on such intervals, sum over N and choose $N \sim |\xi|^{1/(1+\beta)}$ to obtain

$$\left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| \lesssim |t-s|^\gamma |\xi|^{-\rho+1/2(1+\beta)}.$$

This shows that $w + \varphi$ is $(\rho - (2 + 2\beta)^{-1})$ -irregular. In particular, even if we are not able to recover ρ -irregularity, the loss of regularity for $\varphi \in C^{1+\beta}$ is now expected to be $(2 + 2\beta)^{-1}$, which suggests that more generally for $\varphi \in C^\beta$, $w + \varphi$ should be $(\rho - (2\beta)^{-1})$ -irregular, for any $\beta \in [1/2, +\infty)$.

This motivates the following definition; here $F(\xi) = |\xi|^\rho / \sqrt{\log |\xi|}$, $\psi(x) = \sqrt{|x| \log |x|}$.

Definition 84. We say that $w \in C_t^0$ is strongly ρ -irregular if the following holds: for any $n \in \mathbb{N}$, given $\eta \in \mathbb{R}^n$ and denoting by $g_r^\eta := \sum_{k=1}^n \eta_k r^k$, then

$$\sup_{\xi \in \mathbb{R}^d, \eta \in \mathbb{R}^n, s \neq t} \frac{\left| \int_s^t e^{i\xi \cdot w_r + i g_r^\eta} dr \right| F(\xi)}{\sqrt{\log(1 + |\eta|)} \psi(|t-s|)} < \infty. \quad (46)$$

The notion formalises the idea that the irregularity of w should be only mildly affected by polynomial perturbations of any degree; this allows to proceed as above by locally expanding a more general additive perturbation φ in its Taylor series, centred at s .

Theorem 85. *Let w be strongly ρ -irregular. Then for any $\varphi \in C_t^\beta$, $\beta \in [1/2, \infty)$ and for any $\tilde{\rho} < \rho$, $w + \varphi$ is $(\tilde{\rho} - 1/2\beta)$ -irregular. In particular, if $\varphi \in C_t^\infty$, then $w + \varphi$ is $\tilde{\rho}$ -irregular for any $\tilde{\rho} < \rho$.*

Proof. Let $\xi \in \mathbb{R}^d$, $0 \leq s < t \leq T$ be fixed. We write φ as its Taylor series of order $\lfloor \beta \rfloor$ centred in s plus a reminder term:

$$\varphi_r = p_r + R_r = \sum_{k=0}^{\lfloor \beta \rfloor} D^{(k)} \varphi_s \frac{(r-s)^k}{k!} + R_r \quad \forall r \in [s, t].$$

Observe that all the terms in p_r depending exclusively on s do not play any role when estimating $\left| \int_s^t e^{i\xi \cdot (w_r + p_r)} dr \right|$, since they just go outside the integral and provide a term with modulus 1; therefore we will systematically drop them in the calculations. Then

$$\xi \cdot p_r = \sum_{k=1}^{\lfloor \beta \rfloor} \xi \cdot D^{(k)} \varphi_s \frac{(r-s)^k}{k!} = \sum_{k=1}^{\lfloor \beta \rfloor} \xi \cdot D^{(k)} \varphi_s \sum_{j=1}^k \frac{r^j (-s)^{k-j}}{j!(k-j)!} = \sum_{j=1}^{\lfloor \beta \rfloor} \left(\sum_{k=j}^{\lfloor \beta \rfloor} \xi \cdot D^{(k)} \varphi_s \frac{(-s)^{k-j}}{j!(k-j)!} \right) r^j$$

which implies by the definition of strong ρ -irregularity for $n = \lfloor \beta \rfloor$ that

$$\left| \int_s^t e^{i\xi \cdot (w_r + p_r)} dr \right| \lesssim_{n,T} |\xi|^{-\rho} \sqrt{\log |\xi|} \sqrt{\log(1 + \|\xi\| \|\varphi\|_{C^\beta})} \psi(|t-s|).$$

By the usual interpolation arguments, we deduce that for any $\tilde{\rho} < \rho$ there exists $\gamma = \gamma(\tilde{\rho}) > 1/2$ such that $w_r + p_r$ is $\tilde{\rho}$ -irregular; from now on we work with such fixed pair $(\gamma, \tilde{\rho})$.

On the other hand, $\|R\|_{C([s,t];\mathbb{R}^d)} \lesssim \|\varphi\|_{C^\beta} |t-s|^\beta$ and that $\|R'\|_{C([s,t];\mathbb{R}^d)} \lesssim \|\varphi\|_{C^\beta} |t-s|^{\beta-1}$, where R' is the derivative of R ; by interpolation this gives $\|R\|_{C^{1/2}([s,t];\mathbb{R}^d)} \lesssim \|\varphi\|_{C^\beta} |t-s|^{\beta-1/2}$. Therefore we have

$$\begin{aligned} \left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| &= \left| \int_s^t e^{i\xi \cdot R_r} d \left(\int_s^r e^{i\xi \cdot (p_u + w_u)} du \right) \right| \\ &\lesssim \|\Phi^{p+w}\|_{\mathcal{W}_{\gamma,\tilde{\rho}}} |\xi|^{-\tilde{\rho}} (|t-s|^\gamma + \|e^{i\xi \cdot R}\|_{C^{1/2}} |t-s|) \\ &\lesssim_\varphi |t-s|^\gamma |\xi|^{-\tilde{\rho}} + |t-s|^{\beta+1/2} |\xi|^{1-\tilde{\rho}}. \end{aligned}$$

Splitting the integral $[s, t]$ in N subintervals of length $|t-s|/N$, applying the estimate on such subintervals and then summing over N we obtain

$$\left| \int_s^t e^{i\xi \cdot (w_r + \varphi_r)} dr \right| \lesssim N^{1/2} |t-s| |\xi|^{-\tilde{\rho}} + N^{-\beta+1/2} |t-s|^{\beta+1/2} |\xi|^{1-\tilde{\rho}}.$$

Choosing N such that $N \sim |\xi|^{1/\beta}$ allows to conclude. \square

Theorem 86. *For any $\delta \in (0, 1)$, almost every $\varphi \in C_t^\delta$ is strongly ρ -irregular for any $\rho < (2\delta)^{-1}$. Almost every $\varphi \in C_t^0$ is strongly ρ -irregular for any $\rho < \infty$.*

Proof. We only briefly sketch the proof as it mostly relies on the same techniques used to prove prevalence of ρ -irregularity and Theorem 47; it is easy to guess the our candidate transverse measures are the laws μ^H of fBm with parameter $H > \delta$. The proof that the strong ρ -irregularity property defines Borel sets in C_t^δ , $\delta \in [0, 1)$, is identical to the one of Lemma 10, so we will omit it. Now fix $\varphi \in C_t^\delta$; we need to show that

$$\mu^H(w \in C_t^\delta : \varphi + w \text{ is strongly } \rho\text{-irregular for any } \rho < (2H)^{-1}) = 1.$$

Once this is proven, the conclusion follows from the usual argument regarding countable intersection of prevalent sets. We switch to the probabilistic notation, W^H being an fBm with law μ^H .

Fix $n \in \mathbb{N}$. In the following, all estimates depend on n . Using the same technique as in Lemma 51 and Theorem 53, it can be shown that there exists $\bar{\lambda}$ such that

$$\sup_{\xi, s, t} \mathbb{E} \left[\exp \left(\frac{\bar{\lambda} \left| \int_s^t e^{i\xi \cdot (\varphi_r + W_r^H) + ig_r^\eta} dr \right|^2 |\xi|^{1/H}}{|t-s|} \right) \right] \leq K \quad \text{uniformly in } \eta \in \mathbb{R}^n.$$

Let us fix $s < t$ and define

$$J_{s,t}(\lambda) := \sum_{N \in \mathbb{N}} 2^{-N} \sum_{\xi \in 2^{-N} \mathbb{Z}^d, \eta \in 2^{-N} \mathbb{Z}^n} 2^{-N(d+n+2)} (1 + |\xi|)^{-(d+1)} (1 + |\eta|)^{-(n+1)} \times \\ \times \exp \left(\lambda \frac{\left| \int_s^t e^{i\xi \cdot (\varphi_r + W_r^H) + ig_r^\eta} dr \right|^2 |\xi|^{1/H}}{|t-s|} \right).$$

It is clear that $\mathbb{E}[J_{s,t}(\lambda)] \lesssim K$ for all $\lambda \leq \bar{\lambda}$ and that by Jensen inequality $J_{s,t}(\lambda)^\beta \lesssim J_{s,t}(\beta\lambda)$ for all $\beta \geq 1$. Let us also define

$$Y_{s,t} = \sup_{\xi \in \mathbb{R}^d, \eta \in \mathbb{R}^n} \frac{\left| \int_s^t e^{i\xi \cdot (\varphi_r + W_r^H) + ig_r^\eta} dr \right| |\xi|^{\frac{1}{2H}}}{\sqrt{\log(1 + |\eta|)} \sqrt{\log(1 + |\xi|)} |t-s|^{1/2}}.$$

In order to prove the theorem it suffices to show that there exists $\lambda > 0$ such that $\mathbb{E}[e^{\lambda Y_{s,t}^2}] \leq K$, since in that case we are in the conditions to apply Lemma 90. Let (ξ, η) be fixed, then for any N there exists $(\tilde{\xi}, \tilde{\eta}) \in (2^{-N} \mathbb{Z}^d, 2^{-N} \mathbb{Z}^n)$ such that $|(\xi, \eta) - (\tilde{\xi}, \tilde{\eta})| \lesssim 2^{-N}$ and we have

$$\left| \int_s^t (e^{i\xi \cdot (\varphi_r + W_r^H) + ig_r^\eta} - e^{i\tilde{\xi} \cdot (\varphi_r + W_r^H) + ig_r^{\tilde{\eta}}}) dr \right| \lesssim |t-s|^{1/2} (1 + \|\varphi + W^H\|_{L^1}^{1/2}) |(\xi, \eta) - (\tilde{\xi}, \tilde{\eta})|^{1/2}.$$

On the other hand, for $(\tilde{\xi}, \tilde{\eta})$ it holds

$$\left| \int_s^t e^{i\tilde{\xi} \cdot (\varphi_r + W_r^H) + ig_r^{\tilde{\eta}}} dr \right| \lesssim \lambda^{-1/2} |\tilde{\xi}|^{-1/2H} |t-s|^{1/2} \sqrt{\log J_{s,t}(\lambda) + N + \log(1 + |\tilde{\xi}|) + \log(1 + |\tilde{\eta}|)}$$

and therefore, putting everything together, after some computations we obtain

$$\left| \int_s^t e^{i\xi \cdot (\varphi_r + W_r^H) + ig_r^\eta} dr \right| |\xi|^{1/2H} \lesssim |t-s|^{1/2} 2^{-N/2} (1 + \|\varphi + W^H\|_{L^1}^{1/2}) |\xi|^{1/2H} \\ + |t-s|^{1/2} \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda) + N + \log(1 + |\eta|) + \log(1 + |\xi|) + c}.$$

Choosing N such that $2^{-N/2} \sim |\xi|^{1/2H}$, dividing by $\sqrt{\log(1 + |\eta|)} \sqrt{\log(1 + |\xi|)} |t-s|^{1/2}$ and taking the supremum we get

$$Y_{s,t} \lesssim 1 + \|\varphi + W^H\|_{L^1}^{1/2} + \lambda^{-1/2} + \lambda^{-1/2} \sqrt{\log J_{s,t}(\lambda)}$$

and so there exists a constant C such that

$$e^{\lambda Y_{s,t}^2} \lesssim e^{\lambda C(1 + \|\varphi + W^H\|_{L^1})} J_{s,t}(\lambda)^C \lesssim e^{2\lambda C(1 + \|\varphi + W^H\|_{L^1})} + J_{s,t}(\lambda)^{2C} \lesssim e^{2\lambda C(1 + \|\varphi + W^H\|_{L^1})} + J_{s,t}(2C\lambda).$$

Invoking Fernique theorem and choosing λ such that $2C\lambda < \bar{\lambda}$ we obtain the result for fixed n . Since n is arbitrary, we obtain the conclusion. \square

Remark 87. The same proof shows that any β -SLND Gaussian process with suitable integrability conditions is strongly $(2\beta)^{-1}$ -irregular with probability 1.

Proof of Theorem 38. Follows from a combination of Theorems 85 and 86. \square

A Auxiliary results

A.1 A primer on Young integration theory

Estimates involving Young integrals are frequently used throughout this work; for this reason we present here a brief account on the topic for the interested reader. Young integrals go back to [66], where a quantity of the form

$$\int_0^t f_s dg_s$$

is defined in terms of the limit of Riemann–Stieltjes sums, under the condition that f and g are respectively of finite p - and q -variation with $1/p + 1/q > 1$; for a modern account on the theory see for instance Section 6 of [33].

Here we restrict ourselves to f and g belonging to suitable Hölder spaces and we follow the modern approach of constructing Young integrals by means of the Sewing lemma.

Consider an interval $[0, T]$ and a Banach space E ; let Δ_n denote the n -simplex on $[0, T]$, so that $\Delta_n = \{(t_1, \dots, t_n) : 0 \leq t_1 \leq \dots \leq t_n \leq T\}$. Given a map $\Gamma : \Delta_2 \rightarrow E$, we define $\delta\Gamma : \Delta_3 \rightarrow E$ by

$$\delta\Gamma_{s,u,t} := \Gamma_{s,t} - \Gamma_{s,u} - \Gamma_{u,t}.$$

We say that $\Gamma \in C_2^{\alpha,\beta}([0, T]; E)$ if $\Gamma_{t,t} = 0$ for all $t \in [0, T]$ and $\|\Gamma\|_{\alpha,\beta} < \infty$, where

$$\|\Gamma\|_\alpha := \sup_{s < t} \frac{\|\Gamma_{s,t}\|_E}{|t - s|^\alpha}, \quad \|\delta\Gamma\|_\beta := \sup_{s < u < t} \frac{\|\delta\Gamma_{s,u,t}\|_E}{|t - s|^\beta}, \quad \|\Gamma\|_{\alpha,\beta} := \|\Gamma\|_\alpha + \|\delta\Gamma\|_\beta.$$

Let us remark that for a map $f : [0, T] \rightarrow E$, we still denote by $f_{s,t}$ the increment $f_t - f_s$.

Lemma 88 (Sewing lemma). *Let α, β be such that $0 < \alpha \leq 1 < \beta$. For any $\Gamma \in C_2^{\alpha,\beta}([0, T]; E)$ there exists a unique map $\mathcal{I}\Gamma \in C^\alpha([0, T]; E)$ such that $(\mathcal{I}\Gamma)_0 = 0$ and*

$$\left\| (\mathcal{I}\Gamma)_{s,t} - \Gamma_{s,t} \right\|_E \leq C \|\delta\Gamma\|_\beta |t - s|^\beta \quad (47)$$

where the constant C only depends on β . In particular, the map $\mathcal{I} : C_2^{\alpha,\beta} \rightarrow C^\alpha$ is linear and bounded and there exists a constant C' which only depends on β and T such that

$$\|\mathcal{I}\Gamma\|_{C^\alpha} \leq C' \|\Gamma\|_{\alpha,\beta}. \quad (48)$$

For given Γ , the map $\mathcal{I}\Gamma$ is characterised as the unique limit of Riemann–Stieltjes sums: for any $t > 0$

$$(\mathcal{I}\Gamma)_t = \lim_{|\Pi| \rightarrow 0} \sum_i \Gamma_{t_i, t_{i+1}}.$$

The notation above means that for any sequence of partitions $\Pi_n = \{0 = t_0 < t_1 < \dots < t_{k_n} = t\}$ with mesh $|\Pi_n| = \sup_{i=1, \dots, k_n} |t_i - t_{i-1}| \rightarrow 0$ as $n \rightarrow \infty$, it holds

$$(\mathcal{I}\Gamma)_t = \lim_{n \rightarrow \infty} \sum_{i=0}^{k_n-1} \Gamma_{t_i, t_{i+1}}.$$

For a proof, see Lemma 4.2 from [32]. With this tool at hand, it is possible to define Young integrals in a variety of situations; we start by the most general one.

Given two Banach spaces E and F , we denote by $\mathcal{L}(E; F)$ the set of all bounded linear operators from E to F , which is a Banach space with the norm

$$\|A\|_{\mathcal{L}(E; F)} = \sup_{\varphi \in E \setminus \{0\}} \frac{\|A\varphi\|_F}{\|\varphi\|_E}.$$

Corollary 89 (Young integral). *Let $A \in C^\alpha([0, T]; \mathcal{L}(E; F))$ and $\varphi \in C^\beta([0, T]; E)$ such that $\alpha, \beta \in [0, 1]$ and $\alpha + \beta > 1$. Then for any $t \in [0, T]$, the limit in F of the following Riemann–Stieltjes sums exists and is unique:*

$$\int_0^t A_{ds} \varphi_s := \lim_{|\Pi| \rightarrow 0} \sum_i A_{t_i, t_{i+1}} \varphi_{t_i}.$$

Moreover there exists a constant C which only depends on $\alpha + \beta$ such that

$$\left\| \int_s^t A_{dr} \varphi_r - A_{s,t} \varphi_s \right\|_F \leq C \|A\|_{C^\alpha \mathcal{L}} \|\varphi\|_{C^\beta E}. \quad (49)$$

If $A \in C^1([0, T]; \mathcal{L}(E, F))$, then

$$\int_s^t A_{dr} \varphi_r = \int_s^t A'_r \varphi_r dr \quad \text{for all } 0 \leq s < t \leq T.$$

Proof. Define the map $\Gamma : \Delta_2 \rightarrow F$ by setting $\Gamma_{s,t} := A_{s,t} \varphi_s$; by definition $\Gamma_{t,t} = 0$ and moreover

$$\begin{aligned} \|\Gamma_{s,t}\|_F &\leq \|A_{s,t}\|_{\mathcal{L}} \|\varphi_s\|_E \leq |t-s|^\alpha \|A\|_{C^\alpha \mathcal{L}} \|\varphi\|_{C^\beta E} \\ \|\delta \Gamma_{s,u,t}\|_F &= \|A_{u,t} \varphi_{s,u}\|_F \leq \|A_{u,t}\|_{\mathcal{L}} \|\varphi_{s,u}\|_E \leq \|A\|_{C^\alpha \mathcal{L}} \|\varphi\|_{C^\beta E} |t-s|^{\alpha+\beta} \end{aligned}$$

which implies that we can apply the Sewing lemma for such choice of Γ ; inequality (49) is then an immediate consequence of the above estimates combined with (47). The last statement follows from the fact that if $A \in C^1([0, T]; \mathcal{L}(E, F))$, then by standard theory for any $t > 0$ it holds

$$\int_0^t A'_r \varphi_r dr = \lim_{|\Pi| \rightarrow 0} \sum_i A_{t_i, t_{i+1}} \varphi_{t_i}.$$

□

Let us stress that in the above statements whether E and F are finite or infinite dimensional does not play any role and that the constants C and C' do not depend on them.

The most basic definition of Young integral is in the case $f \in C^\alpha([0, T]; \mathbb{R})$ and $g \in C^\beta([0, T]; \mathbb{R})$ with $\alpha + \beta > 1$, in which case for the choice $E = F = \mathbb{R}$ and the identification $\mathbb{R} = \mathcal{L}(\mathbb{R}; \mathbb{R})$ we can define both

$$\int_0^\cdot f_s dg_s \in C^\beta([0, T]; \mathbb{R}), \quad \int_0^\cdot g_s df_s \in C^\alpha([0, T]; \mathbb{R}).$$

In the case $f \in C^\alpha([0, T]; \mathbb{R}^{m \times n})$ and $g \in C^\beta([0, T]; \mathbb{R}^n)$, both above integrals can be defined, this time being \mathbb{R}^m -valued functions, either by reasoning component-by-component or using identifications between dual spaces ($v \in \mathbb{R}^n$ can be identified with the map $A \mapsto Av$ which is an element of $\mathcal{L}(\mathbb{R}^{m \times n}; \mathbb{R}^m)$).

Another important case is the following: let E, F and G be Banach spaces and let $A : E \times F \rightarrow G$ be a bilinear bounded map, i.e. such that

$$\|A\|_{\mathcal{L}^2(E \times F; G)} = \sup_{\substack{v \in E \setminus \{0\} \\ w \in F \setminus \{0\}}} \frac{\|A(v, w)\|_G}{\|v\|_E \|w\|_F} < \infty.$$

Then given $\varphi \in C^\alpha([0, T]; E)$ and $\psi \in C^\beta([0, T]; F)$ we can define respectively

$$\int_0^\cdot A(\varphi_s, d\psi_s) \in C^\beta([0, T]; G), \quad \int_0^\cdot A(d\varphi_s, \psi_s) \in C^\alpha([0, T]; G);$$

this follows from the fact that the map A allows to define an embedding $E \hookrightarrow \mathcal{L}(F; G)$ by $v \mapsto A(v, \cdot)$, similarly for $F \hookrightarrow \mathcal{L}(E; G)$, and that Hölder continuity of paths are preserved by these maps.

A.2 Some useful tools

We give here a chaining lemma which was frequently used in Section 4. It is a slight variation on Lemma 3.1 from [10].

Lemma 90. *Let E be a Banach space and let $X : [0, T] \rightarrow E$ be a continuous stochastic process such that, for some $\lambda > 0$,*

$$\mathbb{E} \left[\exp \left(\lambda \frac{\|X_t - X_s\|_E^2}{|t - s|^{2\alpha}} \right) \right] \leq C \quad \forall s \neq t \in [0, T].$$

Then \mathbb{P} -a.s. $X \in C^\varphi([0, T]; E)$ for the modulus of continuity $\varphi(x) = x^\alpha \sqrt{|\log x|}$ and there exists $\beta > 0$ such that

$$\mathbb{E}[\exp(\beta \|X\|_{C^\varphi E}^2)] < \infty.$$

In particular, if $X_0 \equiv 0$, then for any $\gamma < \alpha$ there exists $\beta > 0$ such that

$$\mathbb{E}[\exp(\beta \|X\|_{C^\gamma E}^2)] < \infty.$$

Proof. Without loss of generality we can assume $T = 1$. Also, we will only show that proof in the case $\alpha = 1/2$, the other cases being entirely analogue. Let us define the random variable

$$R(\lambda) = \sum_{n \in \mathbb{N}} \sum_{k=0}^{2^n-1} 2^{-2n} \exp \left(\mu \frac{\|X_{(k+1)2^{-n}} - X_{k2^{-n}}\|_E^2}{2^{-n}} \right).$$

Then it follows from the assumption that $\mathbb{E}[R(\lambda)] \leq C$. We can then apply Lemma 3.1 from [10] to deduce that there exist deterministic positive constants K, β such that

$$\exp \left(\beta \frac{\|X_t - X_s\|_E^2}{|t - s|} \right) \lesssim |t - s|^{-K} R(\lambda) \quad \forall s \neq t$$

which implies by taking the logarithm and dividing by $-\log |t - s|$ that

$$\exp \left(\beta \left(\sup_{s \neq t} \frac{\|X_t - X_s\|_E}{|t - s|^{1/2} \sqrt{-\log |t - s|}} \right)^2 \right) = \sup_{s \neq t} \exp \left(\beta \frac{\|X_t - X_s\|_E^2}{|t - s|(-\log |t - s|)} \right) \lesssim R(\lambda)$$

which yields the conclusion. Alternatively, it follows from the assumption that

$$\mathbb{E}[B] := \mathbb{E} \left[\int_{[0, T]^2} \exp \left(\lambda \frac{\|X_t - X_s\|_E^2}{|t - s|^{2\alpha}} \right) dt ds \right] < \infty$$

which implies that we can apply Garsia–Rodemich–Rumsey Theorem (see [36]) for the choice $\psi(x) = e^{\lambda x^2}$, $p(x) = x^\alpha$, which gives

$$\|X_t - X_s\|_E \lesssim \int_0^{|t-s|} \sqrt{\log B - \log u} u^{\alpha-1} du \lesssim \left(\sqrt{\log B} + \sqrt{-\log |t - s|} \right) |t - s|^\alpha$$

and from which we can again deduce that

$$\sup_{s \neq t} \frac{\|X_t - X_s\|_E}{|t - s|^\alpha \sqrt{-\log |t - s|}} \lesssim 1 + \sqrt{\log B}$$

and thus the bound $\exp(\beta \|X\|_{C^\varphi E}^2) \lesssim B^{C\beta}$, for a suitable constant $C > 0$ and any $\beta > 0$. Choosing $\beta = 1/C$, using the fact that $\mathbb{E}[B] < \infty$, we obtain the exponential integrability bound.

The final claim follows immediately. \square

We also present here some details on Fourier–Lebesgue spaces.

Definition 91. Let $\alpha \in \mathbb{R}$, $p \in [1, \infty]$; we define the Fourier–Lebesgue space $\mathcal{FL}^{\alpha,p}(\mathbb{R}^d)$ as

$$\mathcal{FL}^{\alpha,p}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) : \langle \xi \rangle^\alpha |\hat{f}(\xi)| \in L^p(\mathbb{R}^d) \right\}$$

where $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$. It is a Banach space endowed with the norm

$$\|f\|_{\mathcal{FL}^{\alpha,p}} = \|\langle \cdot \rangle^\alpha \hat{f}\|_{L^p}.$$

It follows immediately from the definition that we could replace $\langle \cdot \rangle$ with any other function having the same behaviour at infinity, for instance with $(1 + |\cdot|)$; $\langle \cdot \rangle$ is usually considered as it is the Fourier symbol associated to the operator $(I - \Delta)^{1/2}$. Here is a list of relations of Fourier–Lebesgue spaces with other known functional spaces:

- For any $\alpha \in \mathbb{R}$, $\mathcal{FL}^{\alpha,2}$ coincides the classical fractional Sobolev space $H^\alpha = (I - \Delta)^{\alpha/2} L^2$.
- By Hausdorff–Young inequality, for $p \in [1, 2]$ we have the embedding $L^p \hookrightarrow \mathcal{FL}^{0,p'}$, where p' denotes the conjugate exponent to p ; similarly for p as above, for the Bessel spaces $L^{\alpha,p} = (I - \Delta)^{\alpha/2} L^p$ we have $L^{\alpha,p} \hookrightarrow \mathcal{FL}^{\alpha,p'}$.
- In the case $f \in L^1$ the result is slightly stronger, namely \hat{f} is uniformly continuous, bounded and $\hat{f}(\xi) \rightarrow 0$ as $\xi \rightarrow \infty$ by Riemann–Lebesgue lemma; if f is a finite measure on \mathbb{R}^d , then \hat{f} is still uniformly continuous and bounded.
- We have the embedding $\mathcal{FL}^{0,1} \hookrightarrow C^0$ and more generally $\mathcal{FL}^{\alpha,1} \hookrightarrow C^\alpha$, where for $\alpha = n \in \mathbb{N}$ we mean the classical C^n space, while for α fractional or negative $C^\alpha = B_{\infty,\infty}^\alpha$, the latter being a Besov–Hölder space.
- Similarly by Hausdorff–Young for $p \in [1, 2]$ we have the embedding $\mathcal{FL}^{\alpha,p} \hookrightarrow L^{\alpha,p'}$.

There are also embeddings in different scales of Fourier–Lebesgue spaces.

Lemma 92. For any $q < p$ and any $\varepsilon > 0$ it holds

$$\mathcal{FL}^{\alpha,p} \hookrightarrow \mathcal{FL}^{\alpha-d(\frac{1}{q}-\frac{1}{p})-\varepsilon,q}.$$

Proof. For any $q < p$ and $s > 0$ we have

$$\|f\|_{\mathcal{FL}^{\alpha-s,q}} = \left(\int_{\mathbb{R}^d} (\langle \xi \rangle^\alpha |\hat{f}(\xi)|)^q \langle \xi \rangle^{-sq} d\xi \right)^{1/q} \leq \|f\|_{\mathcal{FL}^{\alpha,p}} \left(\int_{\mathbb{R}^d} \langle \xi \rangle^{-s\frac{pq}{p-q}} d\xi \right)^{\frac{1}{q}-\frac{1}{p}}$$

where the integral is convergent if and only if $-spq/(p-q) < -d$, namely

$$s > d \left(\frac{1}{q} - \frac{1}{p} \right).$$

□

The above statement can be combined with other embeddings like the ones mentioned above. For instance we have $\mathcal{FL}^{\alpha,\infty} \hookrightarrow \mathcal{FL}^{\alpha-d/2-\varepsilon,2} = H^{\alpha-d/2-\varepsilon,2}$ and $\mathcal{FL}^{\alpha,\infty} \hookrightarrow \mathcal{FL}^{\alpha-d-\varepsilon,1} \hookrightarrow C^{\alpha-d-\varepsilon}$.

One of the main motivations to introduce Fourier–Lebesgue spaces is that they behave nicely under convolution, due to the properties of Fourier transform.

Lemma 93. Let $f \in \mathcal{FL}^{\alpha,p}$, $g \in \mathcal{FL}^{\beta,q}$ with $\frac{1}{p} + \frac{1}{q} \leq 1$. Then $f * g \in \mathcal{FL}^{\alpha+\beta,r}$ where $\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$ and

$$\|f * g\|_{\mathcal{FL}^{\alpha+\beta,r}} \leq \|f\|_{\mathcal{FL}^{\alpha,p}} \|g\|_{\mathcal{FL}^{\beta,q}}.$$

Proof. By the properties of Fourier transform $\widehat{f * g} = \hat{f} \hat{g}$, therefore

$$\|f * g\|_{\mathcal{FL}^{\alpha+\beta,r}} = \left(\int_{\mathbb{R}^d} (\langle \xi \rangle^\alpha |\hat{f}(\xi)|)^r (\langle \xi \rangle^\beta |\hat{g}(\xi)|)^r d\xi \right)^{1/r} \leq \|f\|_{\mathcal{FL}^{\alpha,p}} \|g\|_{\mathcal{FL}^{\beta,q}}$$

where in the last passage we used the generalised Hölder inequality $\|\varphi\psi\|_{L^r} \leq \|\varphi\|_{L^p} \|\psi\|_{L^q}$ for r, p and q as above. □

It follows in particular from the above that any bounded Fourier symbol acts continuously on $\mathcal{FL}^{\alpha,p}$, for any choice of α and p . We also have $\mathcal{FL}^{\alpha,p} * \mathcal{FL}^{\beta,\infty} \hookrightarrow \mathcal{FL}^{\alpha+\beta,p}$.

References

- [1] Oussama Amine, David Baños, and Frank Proske. C^∞ -regularization by noise of singular ODE's. preprint *arXiv:1710.05760*, 2017.
- [2] Siva Athreya, Oleg Butkovsky, and Leonid Mytnik. Strong existence and uniqueness for stable stochastic differential equations with distributional drift. *The Annals of Probability*, 48(1):178–210, 2020.
- [3] Hajer Bahouri, Jean-Yves Chemin, and Raphaël Danchin. *Fourier analysis and nonlinear partial differential equations*, volume 343. Springer Science & Business Media, 2011.
- [4] Frédéric Bayart and Yanick Heurteaux. On the Hausdorff dimension of graphs of prevalent continuous functions on compact sets. In *Further Developments in Fractals and Related Fields*, pages 25–34. Springer, 2013.
- [5] Lisa Beck, Franco Flandoli, Massimiliano Gubinelli, and Mario Maurelli. Stochastic ODEs and stochastic linear PDEs with critical drift: regularity, duality and uniqueness. *Electronic Journal of Probability*, 24, 2019.
- [6] Simeon M. Berman. Harmonic analysis of local times and sample functions of Gaussian processes. *Transactions of the American Mathematical Society*, 143:269–281, 1969.
- [7] Simeon M Berman. Local nondeterminism and local times of Gaussian processes. *Indiana University Mathematics Journal*, 23(1):69–94, 1973.
- [8] R. Catellier. Rough linear transport equation with an irregular drift. *Stoch. Partial Differ. Equ. Anal. Comput.*, 4(3):477–534, 2016.
- [9] R. Catellier and M. Gubinelli. Averaging along irregular curves and regularisation of ODEs. Preprint *arXiv:1205.1735v1* (version 1), 2012.
- [10] Rémi Catellier and Massimiliano Gubinelli. Averaging along irregular curves and regularisation of ODEs. *Stochastic Processes and their Applications*, 126(8):2323–2366, 2016.
- [11] P. E. Chaudru de Raynal. Strong existence and uniqueness for degenerate SDE with Hölder drift. *Ann. Inst. Henri Poincaré Probab. Stat.*, 53(1):259–286, 2017.
- [12] K. Chouk and M. Gubinelli. Nonlinear PDEs with modulated dispersion I: Nonlinear Schrödinger equations. *Comm. Partial Differential Equations*, 40(11):2047–2081, 2015.
- [13] Khalil Chouk and Benjamin Gess. Path-by-path regularization by noise for scalar conservation laws. *Journal of Functional Analysis*, 277(5):1469–1498, 2019.
- [14] Khalil Chouk and Massimiliano Gubinelli. Nonlinear PDEs with modulated dispersion II: Korteweg–de Vries equation. *arXiv preprint arXiv:1406.7675*, 2014.
- [15] Jens Peter Reus Christensen. On sets of Haar measure zero in Abelian Polish groups. *Israel Journal of Mathematics*, 13(3-4):255–260, 1972.
- [16] Jack Cuzick. A lower bound for the prediction error of stationary Gaussian processes. *Indiana University Mathematics Journal*, 26(3):577–584, 1977.
- [17] Giuseppe Da Prato and Franco Flandoli. Pathwise uniqueness for a class of SDE in Hilbert spaces and applications. *Journal of Functional Analysis*, 259(1):243–267, 2010.
- [18] Giuseppe Da Prato, Franco Flandoli, Michael Röckner, and A. Yu Veretennikov. Strong uniqueness for SDEs in Hilbert spaces with nonregular drift. *The Annals of Probability*, 44(3):1985–2023, 2016.
- [19] Giuseppe Da Prato and Jerzy Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge University Press, 2014.

- [20] A. M. Davie. Uniqueness of solutions of stochastic differential equations. *Int. Math. Res. Not. IMRN*, (24), 2007.
- [21] Paul-Eric Chaudru de Raynal, Stephane Menozzi, and Enrico Priola. Weak well-posedness of multidimensional stable driven SDEs in the critical case. *Stochastics and Dynamics*, 20(6):2040004, 2020.
- [22] Romain Duboscq and Anthony Réveillac. On a stochastic Hardy–Littlewood–Sobolev inequality with application to Strichartz estimates for a noisy dispersion. *Annales Henri Lebesgue*, 5:263–274, 2022. doi:10.5802/ahl.122
- [23] Kenneth Falconer. *Fractal geometry*. John Wiley & Sons, Ltd., Chichester, third edition, 2014. Mathematical foundations and applications.
- [24] Franco Flandoli. *Random Perturbation of PDEs and Fluid Dynamic Models: École d’été de Probabilités de Saint-Flour XL–2010*, volume 2015. Springer Science & Business Media, 2011.
- [25] Franco Flandoli. Regularizing properties of Brownian paths and a result of Davie. *Stochastics and Dynamics*, 11(2-3):323–331, 2011.
- [26] Franco Flandoli, Massimiliano Gubinelli, and Enrico Priola. Well-posedness of the transport equation by stochastic perturbation. *Inventiones mathematicae*, 180(1):1–53, 2010.
- [27] Gerald B. Folland. *Real analysis: modern techniques and their applications*, volume 40. John Wiley & Sons, 1999.
- [28] Jonathan M. Fraser and James T. Hyde. The Hausdorff dimension of graphs of prevalent continuous functions. *Real Analysis Exchange*, 37(2):333–352, 2012.
- [29] Aurélia Fraysse. Regularity criteria for almost every function in Sobolev spaces. *Journal of Functional Analysis*, 258(6):1806–1821, 2010.
- [30] Aurélia Fraysse and Stéphane Jaffard. How smooth is almost every function in a Sobolev space? *Revista Matemática Iberoamericana*, 22(2):663–682, 2006.
- [31] Peter Friz and Atul Shekhar. Doob-Meyer for rough paths. *Bulletin of the Institute of Mathematics. Academia Sinica. New Series*, 8(1):73–84, 2013.
- [32] Peter K. Friz and Martin Hairer. *A Course on Rough Paths: With an Introduction to Regularity Structures*. Springer, 2014.
- [33] Peter K. Friz and Nicolas B. Victoir. *Multidimensional stochastic processes as rough paths: theory and applications*, volume 120. Cambridge University Press, 2010.
- [34] Lucio Galeati and Massimiliano Gubinelli. Noiseless regularisation by noise. *Rev. Mat. Iberoam.*, 38(2):433–502, 2022.
- [35] Lucio Galeati and Massimiliano Gubinelli. Mixing for generic rough shear flows. *arXiv:2107.12115*, 2021.
- [36] Adriano M. Garsia, Eugene Rodemich, H. Rumsey, and M. Rosenblatt. A real variable lemma and the continuity of paths of some Gaussian processes. *Indiana University Mathematics Journal*, 20(6):565–578, 1970.
- [37] Donald Geman and Joseph Horowitz. Occupation densities. *The Annals of Probability*, pages 1–67, 1980.
- [38] Massimiliano Gubinelli. Controlling rough paths. *Journal of Functional Analysis*, 216(1):86–140, 2004.

- [39] Fabian A. Harang and Nicolas Perkowski. C^∞ -regularization of ODEs perturbed by noise. *Stochastics and Dynamics*, 21(08):2140010, 2021.
- [40] Antoine Hocquet, Marco Romito, and Leonardo Tolomeo. About some notions of irregularity of paths. *To appear*, 2023+.
- [41] Brian R. Hunt. The prevalence of continuous nowhere differentiable functions. *Proceedings of the American mathematical Society*, 122(3):711–717, 1994.
- [42] Brian R. Hunt, Tim Sauer, and James A. Yorke. Prevalence: a translation-invariant “almost every” on infinite-dimensional spaces. *Bulletin of the American mathematical society*, 27(2):217–238, 1992.
- [43] Jean-Pierre Kahane. *Some random series of functions*. Volume 5. Cambridge University Press, 1993.
- [44] Nicolai V. Krylov and Michael Röckner. Strong solutions of stochastic equations with singular time dependent drift. *Probability theory and related fields*, 131(2):154–196, 2005.
- [45] Khoa Lê. A stochastic sewing lemma and applications. *Electronic Journal of Probability*, 25, 2020.
- [46] Elliott H. Lieb and Michael Loss. *Analysis*. Volume 14. American Mathematical Soc., 2001.
- [47] Pertti Mattila. *Fourier analysis and Hausdorff dimension*, volume 150. Cambridge University Press, 2015.
- [48] Peter Mörters and Yuval Peres. *Brownian motion*, volume 30 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2010. With an appendix by Oded Schramm and Wendelin Werner.
- [49] David Nualart. *The Malliavin calculus and related topics*. Volume 1995. Springer, 2006.
- [50] David Nualart and Youssef Ouknine. Regularization of differential equations by fractional noise. *Stochastic Processes and their Applications*, 102(1):103–116, 2002.
- [51] David Nualart, Carles Rovira, and Samy Tindel. Probabilistic models for vortex filaments based on fractional Brownian motion. *The Annals of Probability*, 31(4):1862–1899, 2003.
- [52] William Ott and James Yorke. Prevalence. *Bulletin of the American Mathematical Society*, 42(3), 2005.
- [53] Juan C. Peral. L^p estimates for the wave equation. *J. Functional Analysis*, 36(1):114–145, 1980.
- [54] Yuval Peres and Perla Sousi. Dimension of fractional Brownian motion with variable drift. *Probability Theory and Related Fields*, 165(3-4):771–794, 2016.
- [55] Jean Picard. Representation formulae for the fractional Brownian motion. In *Séminaire de probabilités XLIII*, pages 3–70. Springer, 2011.
- [56] Loren D. Pitt. Local times for Gaussian vector fields. *Indiana University Mathematics Journal*, 27(2):309–330, 1978.
- [57] Enrico Priola. Pathwise uniqueness for singular SDEs driven by stable processes. *Osaka Journal of Mathematics*, 49(2):421–447, 2012.
- [58] Marco Romito and Leonardo Tolomeo. Yet another notion of irregularity through small ball estimates. *arXiv:2207.02716*, 2022.
- [59] Ken-iti Sato. *Lévy processes and infinitely divisible distributions*. Cambridge University Press, 1999.
- [60] A. V. Shaposhnikov. Some remarks on Davie’s uniqueness theorem. *Proceedings of the Edinburgh Mathematical Society*, 59(4):1019–1035, 2016.

- [61] A. V. Shaposhnikov. Correction to the paper “Some remarks on Davie’s uniqueness theorem”. *arXiv preprint arXiv:1703.06598*, 2017.
- [62] Alexander Shaposhnikov and Lukas Wresch. Pathwise vs. path-by-path uniqueness. *Annales de l’Institut Henri Poincaré (B) Probabilités et statistiques*, 58(3):1640–1649, 2022.
- [63] Alexander Y. Veretennikov. On strong solutions and explicit formulas for solutions of stochastic integral equations. *Sbornik: Mathematics*, 39:387–403, 1981.
- [64] Yimin Xiao. Properties of local–nondeterminism of gaussian and stable random fields and their applications. In *Annales de la Faculté des sciences de Toulouse: Mathématiques*, volume 15, pages 157–193, 2006.
- [65] Yimin Xiao. Strong local nondeterminism and the sample path properties of Gaussian random fields. *Asymptotic theory in probability and statistics with applications*, pages 136–176, 2007.
- [66] Laurence C. Young. An inequality of the Hölder type, connected with Stieltjes integration. *Acta Mathematica*, 67:251–282, 1936.
- [67] Alexander K. Zvonkin. A transformation of the phase space of a diffusion process that removes the drift. *Mathematics of the USSR-Sbornik*, 22(1):129, 1974.