

CAUCHY THEORY AND MEAN-FIELD LIMIT FOR GENERAL VICSEK MODELS IN COLLECTIVE DYNAMICS

MARC BRIANT, SARA MERINO-ACEITUNO

ABSTRACT. In this paper we provide a local Cauchy theory both on the torus and in the whole space for general Vicsek dynamics at the particle and at the kinetic level. We consider rather general interaction kernels, nonlinear viscosity and nonlinear friction. We also establish the mean-field limit in the large particle limit. Particularly, we include normalised kernels which display a singularity when the flux of particles vanishes. Thus, in terms of the Cauchy theory for the kinetic equation, we extend to more general interactions and complete the program initiated in [30] (where the authors assume that the singularity does not take place) and in [27] (where the authors prove that the singularity does not happen in the space homogeneous case). Moreover, we derive an explicit lower time of existence as well as a global existence criterion that is applicable, among other cases, to obtain a long time theory for non-renormalised kernels and for the original Vicsek problem without any *a priori* assumptions.

Keywords: Vicsek model; Vicsek-Kolmogorov equation; collective dynamics; nonlinear Fokker-Planck equation on the sphere; normalized interaction kernels; mean-field limit, well-posedness.

Acknowledgements. S. Merino-Aceituno is supported by the Vienna Science and Technology Fund (WWTF) with a Vienna Research Groups for Young Investigators, grant VRG17-014.

The authors also thank Amic Frouvelle for useful discussions.

CONTENTS

1. Introduction	2
1.1. Motivation	2
1.2. Particular case: Vicsek model as in [20] and [3, 29]	2
1.3. General forms of the Vicsek model considered	5
1.4. Aims of the paper	6
1.5. Structure of the paper	6
2. Main results and strategy	6
2.1. Functional framework and notations	6
2.2. Kinetic point of view: strategy and results	8
2.3. Microscopic point of view and mean-field limit: strategy and results	11
3. The mesoscopic framework: kinetic differential equation	14
3.1. Linear equation: dependence on coefficients and averaging positivity	14
3.2. The local-in-time nonlinear Cauchy theory	19
3.3. Global existence and decay of the free energy	21
4. Mean-field limit: proofs of the results in Section 2.3	24

4.1. Preliminaries: functional framework and notations.	25
4.2. The non-singular dynamics.	25
4.3. Proof of Theorem 2.9 and Theorem 2.10 assuming Proposition 4.3	27
4.4. Proof of Proposition 4.3	29
Appendix A. Some known results on SDE	31
A.1. Results on existence of solutions and large-particle limits	31
A.2. Stratonovich to Ito's convention	33
References	34

1. INTRODUCTION

1.1. Motivation. The emergence of collective motions among a group of self-propelled agents is receiving a great deal of attention: flocks of birds [32], schools of fish [31], pedestrian dynamics [37], micro-swimmers [6]. Many models have been proposed to explain these collective behaviours. Among them, the Vicsek model [20, 44] is one of the most studied ones. In this model all agents move at a constant speed while trying to adopt the same velocity as their neighbours, up to some noise.

In this introductory section we present the discrete dynamics for the Vicsek model as presented in [20] as well as its corresponding kinetic equation for the time-evolution of the distribution of the particles. Up to now, there were no complete rigorous results on the existence of solutions and the derivation of the kinetic equation (mean-field limit) for the Vicsek dynamics presented in [20]. The reason for this is the presence of a singularity in the dynamics that is reached when the local average velocity of the agents vanishes.

In contrast to [20], one can define a Vicsek model without a singularity [3, 11, 13, 29], for different modelling choices. However, these different modelling choices for the Vicsek model have profound mathematical implications (like appearance of phase transitions). For this reason, in this article we investigate a general form of the Vicsek model that includes the forms presented in [20] (with a singularity), in [3, 29] (without singularity) as well as further extensions of the model like the ones in [4, 12, 13, 22, 28]. We give conditions to have existence of solutions for the equations and for the mean-field limit to hold for this general class of Vicsek models. All of this is detailed in the following sections.

1.2. Particular case: Vicsek model as in [20] and [3, 29]. In what follows we assume that the agents move in a domain Ω with Ω being either the d -dimensional torus \mathbb{T}^d or the whole space \mathbb{R}^d for $d \geq 1$. The orientations of the agents are given elements on the sphere \mathbb{S}^{d-1} . Throughout the article, for a given vector $\omega \in \mathbb{R}^d$ we denote $\mathbf{P}_{\omega^\perp}$ the orthogonal projection in \mathbb{R}^d onto $(\mathbb{R}\omega)^\perp$. The time-continuous Vicsek model [20] considers N agents characterised by their positions $X^{i,N}(t) \in \Omega$, $i = 1, \dots, N$ and orientations $\omega^{i,N}(t) \in \mathbb{S}^{d-1}$ over time $t \geq 0$. The evolution of the system is given by the following Stratonovich stochastic differential equation, where $c, \nu, \sigma > 0$ are positive constants:

$$(1.1) \quad dX^{i,N} = c\omega^{i,N} dt,$$

$$(1.2) \quad d\omega^{i,N} = \nu \nabla_{\omega^{i,N}}(\omega^{i,N} \cdot \bar{\omega}_i) dt + \mathbf{P}_{(\omega^{i,N})^\perp} \circ \sqrt{2\sigma} dB_t^i.$$

We explain next this system of equations. Eq. (1.1) describes the transport of the agents: agent i moves in the orientation $\omega^{i,N}$ at speed $c > 0$. Eq. (1.2) gives the evolution of the orientations over time. It includes two competing forces. On one hand the first term in the form of a gradient represents organised motion: agents try to adopt the same orientation. The gradient ∇_ω is the gradient on the sphere and the term $\bar{\omega}_i$ represents the average orientation of the neighbours around agent i . Therefore, the first term in the right-hand-side of Eq. (1.2) is a gradient flow which relaxes the value of the orientation $\omega^{i,N}$ towards the mean orientation of the neighbouring particles $\bar{\omega}_i$. The constant $\nu > 0$ gives the speed of this relaxation. We will comment later on the different choices to compute the average orientation $\bar{\omega}_i$. These different choices give rise to the different models in [20] and [3, 29].

On the other hand B_t^i , $i = 1, \dots, N$ are N independent Brownian motions in \mathbb{R}^d and they introduce noise in the dynamics, driving particles away from organised motion. The constant $\sigma > 0$ gives the intensity of the noise. The symbol 'o' is used to specify that Eq. (1.2) has to be understood in the Stratonovich sense. This ensures that $\omega_i(t) \in \mathbb{S}^{d-1}$ for all times where the solution is defined (this will be proven later).

Now, formally at least, one can compute the time-evolution for the distribution of agents $f = f(t, x, \omega)$ in space $x \in \Omega$ and orientations $\omega \in \mathbb{S}^{d-1}$ at time $t \geq 0$ as the number of particles $N \rightarrow \infty$ [20]. The dynamics for the distribution f is given by the following kinetic equation:

$$(1.3) \quad \partial_t f + c\omega \cdot \nabla_\omega f = \nabla_\omega \cdot [\nu \mathbf{P}_{\omega^\perp} \bar{\omega}_f f + \sigma \nabla_\omega f] =: L(f),$$

where $\nabla_\omega \cdot$ denotes the divergence on the sphere \mathbb{S}^{d-1} and where $\bar{\omega}_f(t, x)$ represents the average orientation of the particles around position x at time t . Notice that the projection term appears due to the fact that

$$\nabla_\omega(\omega \cdot \bar{\omega}) = \mathbf{P}_{\omega^\perp} \bar{\omega}, \quad \text{for } \bar{\omega} \in \mathbb{R}^d \text{ fixed.}$$

One can show that the operator L on the right-hand-side of (1.3) can be rewritten as a non-linear Fokker-Planck operator:

$$(1.4) \quad L(f) := \sigma \nabla_\omega \cdot \left[M_{\bar{\omega}_f} \nabla_\omega \left(\frac{f}{M_{\bar{\omega}_f}} \right) \right],$$

with

$$(1.5) \quad M_v(\omega) = \frac{\exp\left(\frac{\nu}{\sigma}(\omega \cdot v)\right)}{Z}, \quad Z = \int_{\mathbb{S}^{d-1}} \exp\left(\frac{\nu}{\sigma}(\omega \cdot v)\right) d\omega,$$

for any $v \in \mathbb{S}^{d-1}$. The function M_v is a probability density on the sphere named von-Mises distribution. Notice that its mean is in the orientation given by v . When $\sigma \rightarrow \infty$ (large-noise limit) the von-Mises distribution converges to the uniform distribution on the sphere.

We comment now on the different choices to define the average orientation around an agent i , denoted by $\bar{\omega}_i$ in Eq. (1.2). The choice considered has profound implications in the dynamics of the particles and in the derivation of the kinetic equation (1.3) (mean-field limit) and the derivation of macroscopic equations (equations for

the particle density and mean orientation). In the mathematics literature, two options were first considered given, in a compact way, by:

$$(1.6) \quad \bar{\omega}_i = \frac{J_i^N}{\alpha + (1 - \alpha)|J_i^N|}, \quad J_i^N(t) = \sum_{j \neq i}^N K(|X_i(t) - X_j(t)|) \omega_j(t).$$

The two options correspond to the cases $\alpha = 0$ (as presented in [20]) or $\alpha = 1$ (as presented in [3, 29]). The kernel $K \geq 0$ is an interaction kernel that represents the weights given to the neighbouring particles depending on their distance to particle i . A classical choice is to take K the indicator function of a ball of radius $R > 0$. The flux J_i^N is, indeed, an average of the orientations of the neighbouring particles. In the kinetic equation these choices define the operator $\bar{\omega}_f$ as:

$$(1.7) \quad \bar{\omega}_f(t, x) = \frac{\mathbf{J}_f(t, x)}{\alpha + (1 - \alpha)|\mathbf{J}_f(t, x)|}, \quad \mathbf{J}_f(t, x) = \int_{\Omega \times \mathbb{S}^{d-1}} K(|x - y|) \omega f(t, y, \omega) dy d\omega.$$

We compare next, the mathematical implications of these two choices:

Case $\alpha = 1$ (non-normalisation). If $\alpha = 1$, the average orientation $\bar{\omega}_i = J_i^N \notin \mathbb{S}^{d-1}$ is not a unit vector. This also holds in the kinetic equation (1.3) for $\bar{\omega}_f = \mathbf{J}_f \notin \mathbb{S}^{d-1}$ in (1.7). However, the case $\alpha = 1$ removes the singularity when $|\mathbf{J}_f| = 0$ or $|J_i^N| = 0$. In [29] the authors prove the well-posedness of the space-homogeneous kinetic equation in any Sobolev space and in [2] the authors prove the mean-field limit (here we will recover these results). As a counterpart, though, this choice for the average $\bar{\omega}_i$ makes the derivation of macroscopic equations for the density of the particles $\rho = \rho(t, x)$ and the mean orientation $\bar{\omega} = \bar{\omega}(t, x)$ of the agents more complex than in the case $\alpha = 0$. Specifically, two equilibria exist for the mean particle orientation $\bar{\omega}$ depending if $\mathbf{J}_f \neq 0$ or $\mathbf{J}_f = 0$ [12, 13, 29]. In loose terms, if $\mathbf{J}_f \neq 0$, then the von-Mises equilibria $M_{\mathbf{J}_f}$ is an equilibrium. The other equilibrium is given by the uniform distribution on the sphere, which corresponds to the case $\mathbf{J}_f = 0$ in $M_{\mathbf{J}_f}$. In different spatial regions (depending on the particle density ρ) one of these two equilibria is stable. Consequently, this gives rise to a bifurcation or phase transition: in some spatial regions the mean orientation of the agents is $\bar{\omega} = 0$ (corresponding to disordered dynamics) and in other spatial regions the mean orientation is given by a unit vector $|\bar{\omega}| = 1$ (corresponding to ordered dynamics or flocking). The interested reader can find all the details in [4, 12, 13, 29]. The presence of phase transitions enriches the dynamics, in the sense that allows for a wider variety of patterns to arise. Understanding phase transitions is also of great interest in the Physics literature [44].

Case $\alpha = 0$ (normalisation). The first mathematical works on the Vicsek model correspond to this case [20]. The choice $\alpha = 0$ comes with the difficulty of dealing with singularities when the flux $J_i^N = 0$ (another example of collective dynamics with singularities, different from the Vicsek model can be found in [41] for the Kuramoto model). However, assuming that the flux \mathbf{J}_f in the kinetic equation does not vanish along the dynamics, then the macroscopic equations can be obtained more easily than in the case $\alpha = 1$ because there is a unique equilibria: the von-Mises distribution in (1.5). Then the macroscopic equations for the particle density $\rho = \rho(t, x)$ and mean orientation $\bar{\omega} = \bar{\omega}(t, x) \in \mathbb{S}^{d-1}$ correspond to the Self-Organised Hydrodynamics (SOH) given in [20]. This was the first formal derivation of the SOH

dynamics. See also [35, 46] as well as [34] for later rigorous results. In this scenario a rigorous mean-field limit to derive the kinetic equation was missing as well as a Cauchy theory for the particle dynamics and the kinetic equation. We address these points in the present article:

Theorem 1.1 (Cauchy theory and mean-field for the Vicsek model with normalisation). *Suppose that $\alpha = 0$ (normalised case). Suppose that the kernel K is Lipschitz, bounded and that $f(0, x, \omega)$ satisfies assumptions (i) and (ii) in Theorem 2.3. Then, there exists a unique local-in-time solution for particle dynamics (1.1)-(1.2) and the kinetic equation (1.7). Moreover, for short times the kinetic equation can be obtained as the limit of the particle dynamics as the number of particles $N \rightarrow \infty$.*

The precise definition of ‘solution’ and of the limit can be found in Theorems 2.3, 2.9 and 2.10. As mentioned before, the difficulty in proving the previous theorem is to show that there exists at least some time interval $t \in [0, T]$ such that $\mathbf{J}_f \neq 0$ and that for N large enough $J_i^N \neq 0$ for any of the particles. To prove a lower bound in the flux $\mathbf{J}_f \neq 0$ is, therefore, key. Indeed, in [30] the authors manage to prove well-posedness for the kinetic equation (1.3) assuming precisely that *a priori* all solutions satisfy $|\mathbf{J}_f|(t, x, \omega) \geq a > 0$ for all times, positions and orientations. In [27] the authors prove well-posedness in the space homogeneous setting as long as initially $|\mathbf{J}_{f_0}| > 0$ (plus some other assumptions). This has been followed by further extensions in [36]. The space-homogeneous case corresponds to spatially local kernels, i.e., $K(x - y) = \delta_0(x - y)$ (delta distribution).

To conclude this part, the choice on how to define the average orientation $\bar{\omega}_i$ has profound mathematical and modelling implications (like dealing with singularities or with phase transitions). For this reason, in this article we will consider a general class of Vicsek models to encompass different modelling choices - existing or yet to be developed.

1.3. General forms of the Vicsek model considered. As we saw in the previous section, modelling choices in the Vicsek model may produce important differences in the mathematical properties of the equations. For this reason, in this article we consider a wide class of Vicsek models based not only on the choice of the averaged orientation (1.2) but also on the particular shape of the friction ν , the viscosity σ and the interaction kernel K . Specifically, we will allow ν and σ to be functions of f (the solution to the kinetic equation) at the kinetic level and functions of the empirical distribution (see Eq. (2.12)) at the particle level. The interaction kernel K can take the general form $K = K(t, x, x_*, \omega, \omega_*)$. The precise assumptions on ν , σ and K can be found in Section 2.1.

As a consequence, the results presented here apply to a wide breath of Vicsek-types models, like the ones in [12, 13, 22, 28]. In [12, 13] the authors consider functions $\nu = \nu(|\mathbf{J}_f|)$, $\sigma = \sigma(|\mathbf{J}_f|)$ that depend on the flux of the particles; in [28] the author considers a kernel K that is not isotropic; in [22] the authors consider a Vicsek-type model for alignment of discs. Beyond this, the approach presented here can be adapted to investigate other models like the ones in [4, 8, 9, 15, 14, 16, 17, 18, 19, 21, 38]. In [9] the authors couple the Vicsek model with a Kuramoto model. In [38] the author considers a Vicsek model with two species having different velocities. The models in [4, 15, 16, 17, 18] describe collective motion based on nematic alignment (where particles align by adopting the same direction of motion and not necessarily

the same orientation). BGK versions of Vicsek-type models are considered in [8, 14]. In [19] the authors couple the Vicsek model with Stokes equations to model microswimmers. Finally, in [21] a model for the persistent turning walker with curvature ‘alignment’ is presented.

1.4. Aims of the paper. In this paper, for a general form of the Vicsek model, we aim at:

- (i) establishing the well-posedness, at least locally in time, of both the agent-based model (Theorem 2.9) and the kinetic models (Theorem 2.3); in the latter, existence of solutions will be proven in Lebesgue spaces;
- (ii) rigorously proving the mean-field limit that links the agent-based dynamics with the kinetic equation in the limit $N \rightarrow +\infty$ (Theorem 2.10);
- (iii) deriving a criterion that characterises the global-in-time well-posedness of these systems (points (a) and (b) in Theorem 2.3) and apply it to several interesting cases (Section 3.3.1). Along the way we shall also open the discussion to long-time behaviour and construct a free energy but we shall not investigate further (Section 3.3.2).

We aim at giving constructive proofs rather than weak-compactness arguments and at working in Lebesgue spaces rather than more regular Sobolev spaces. Our strategy is thus to prove a fixed point contraction theory for the kinetic equation (2.3) and use this limiting equation to obtain the well-posedness of the stochastic system of equations (2.13) and the mean-field limit using the Sznitmann approach [3, 5, 43].

1.5. Structure of the paper. The paper is structured as follows. Section 2 describes our main results, first in the kinetic framework in Section 2.2 and second for the particle dynamics and its mean-field limit in Section 2.3. The proofs for the results in Section 2.2 are given in Section 3 and the proofs of the results in Section 2.3 are given in Section 4. In the Appendix A the reader can find known results on stochastic differential equations and mean-field limits that are used in Section 4 and that have been added here for the sake of completeness. The results in Appendix A are mainly from [5].

2. MAIN RESULTS AND STRATEGY

2.1. Functional framework and notations. First let us give some notations for functional spaces. We shall work on Lebesgue spaces indexed by the variable into consideration: for any p in $[1, +\infty)$ we denote

$$\|f\|_{L_\omega^p} = \left(\int_{\mathbb{S}^{d-1}} |f(\omega)|^p d\omega \right)^{\frac{1}{p}} \quad \text{and} \quad \|f\|_{L_x^p} = \left(\int_{\Omega} |f(x)|^p dx \right)^{\frac{1}{p}}$$

and for a time-dependent function for any $T > 0$

$$\|f\|_{L_t^p} = \left(\int_{\mathbb{R}^+} |f(t)|^p dt \right)^{\frac{1}{p}} \quad \text{and} \quad \|f\|_{L_{[0,T]}^p} = \left(\int_0^T |f(t)|^p dt \right)^{\frac{1}{p}}.$$

Finally we denote the several variable Lebesgue spaces for any q and r in $[1, +\infty)$

$$\|f\|_{L_{[0,T]}^r L_x^q L_\omega^p} = \left(\int_0^T \left(\int_{\Omega} \|f(t, x, \cdot)\|_{L_\omega^p}^q dx \right)^{\frac{r}{q}} dt \right)^{\frac{1}{r}}$$

with direct modifications for p, q or r being $+\infty$. Note that when two indexes are the same we shall use the short-hand notation $L_x^p L_\omega^p = L_{x,\omega}^p$.

For the mean-field limit results, we will consider the space $\mathcal{H} = (\Omega \times \mathbb{R}^d \times \mathcal{P}_2(\Omega \times \mathbb{R}^d))$, where $\mathcal{P}_2(\Omega \times \mathbb{R}^d)$ is the space of probability measures in $\Omega \times \mathbb{R}^d$ with finite second-order moment, i.e., $\mu \in \mathcal{P}_2(\Omega \times \mathbb{R}^d)$ if it fulfils

$$\int_{\Omega \times \mathbb{R}^d} |z|^2 \mu(dz) < \infty.$$

For $\mu, \mu' \in \mathcal{P}_2(\Omega \times \mathbb{R}^d)$, the 2-Wasserstein distance is given by

$$W_2(\mu, \mu') = \inf \left\{ \left[\int_{(\Omega \times \mathbb{R}^d)^2} |z - z'|^2 \pi(dz, dz') \right]^{1/2}; \right. \\ \left. \pi \in \mathcal{P}(\Omega \times \mathbb{R}^d)^2 \text{ with marginals } \mu \text{ and } \mu' \right\}.$$

The distance W_2 induces the topology of weak convergence of measures and the convergence of all the moments of order up to 2 [5] (see also [45, p. 83]). The space \mathcal{H} is a metric space with the distance $d_{\mathcal{H}}$ given by

$$d_{\mathcal{H}}\left((x, v, m), (y, w, p)\right) = |x - y| + |v - w| + W_2(m, p).$$

The hypotheses we shall make on the nonlinearities are the following.

(H1) The interactions are Lipschitz, regular and bounded in different variables :

$$\mathbf{K}(t, x, x_*, \omega, \omega_*) \in W_{t, x_*}^{1, \infty} W_{\omega_*}^{2, \infty} L_{x, \omega}^\infty;$$

(H2) The viscosity satisfies that for any $T > 0$, $\sigma : L_{[0, T], x, \omega}^2 \longrightarrow L_{[0, T], x, \omega}^\infty$ and is bounded from below and Lipschitz:

$$\forall (f, g) \in (L_{[0, T], x, \omega}^2)^2, \quad 0 < \sigma_0 \leq \sigma(f) \leq \sigma_\infty \\ \|\sigma(f) - \sigma(g)\|_{L_{[0, T], x, \omega}^\infty} \leq \sigma_{\text{lip}} \|f - g\|_{L_{[0, T], x, \omega}^2}.$$

(H3) The friction is local in time and Lipschitz that is $\nu(f)(t, x, v) = \nu(f(t, \cdot, \cdot))(x, v)$ where $\nu : L_{x, \omega}^2 \longrightarrow L_x^\infty W_\omega^{1, \infty}$:

$$\forall (f, g) \in (L_{x, \omega}^2)^2, \quad \|\nu(f)\|_{L_x^\infty W_\omega^{1, \infty}} \leq \nu_\infty \\ \|\nu(f) - \nu(g)\|_{L_{x, \omega}^\infty} \leq \nu_{\text{lip}} \|f - g\|_{L_{x, \omega}^2}.$$

(H4) For the mean-field limit results we will assume further that σ, ν and $\nabla_\omega \sigma$ are Lipschitz and bounded in the space \mathcal{H} defined above.

Remark 2.1. • Note that (H2) and (H3) are satisfied when σ and ν do not depend on f , which is the case in most models so far. Ultimately, one would want $\sigma(f)$ to be local in time and Lipschitz, like $\nu(f)$. We think that this could be achieved with standard parabolic regularity methods for more regular initial data and $\sigma(f)$. The main issue for closing a fixed point argument is the speed of convergence to 0 of the gain of regularity of the solutions f that we did not manage to quantify, see Proposition 3.3 and Remark 3.4.

- In applications, typically the functions $\nu = \nu(t, x, \omega, f)$ and $\sigma = \sigma(t, x, \omega, f)$ are of the form

$$(2.1) \quad \nu(t, x, \omega, f) = \int_{\Omega \times \mathbb{S}^{d-1}} \tilde{\nu}(t, x, \omega, x_*, \omega_*) f(t, x_*, \omega_*) dx_* d\omega_*,$$

for some function $\tilde{\nu}$, and analogously for σ . For example, in [20] the authors consider

$$\nu(t, x, \omega, f) = \omega \cdot \bar{\omega}_f(t, x),$$

with

$$(2.2) \quad \bar{\omega}_f(t, x) = \frac{\tilde{\mathbf{J}}_f}{|\tilde{\mathbf{J}}_f|}(t, x), \quad \tilde{\mathbf{J}}_f(t, x) = \int_{\Omega \times \mathbb{S}^{d-1}} K(x - x_*, \omega_*) f(t, x_*, \omega_*) dx_* d\omega_*.$$

And in [13] the authors consider ν and σ of the form

$$\nu = \nu(t, x, f) = \nu(|\tilde{\mathbf{J}}_f|(t, x)),$$

with $\tilde{\mathbf{J}}_f$ given by (2.2).

2.2. Kinetic point of view: strategy and results. We will consider the following kinetic equation for the distribution function $f = f(t, x, \omega)$ for $(t, x, \omega) \in \mathbb{R}_+ \times \Omega \times \mathbb{R}^d$ (with Ω either the d -dimensional torus \mathbb{T}^d or the full space \mathbb{R}^d):

$$(2.3) \quad \partial_t f + c\omega \cdot \nabla_x f = \nabla_\omega \cdot (\sigma(f) \nabla_\omega f) + \nabla_\omega \cdot (\nu(f) f \mathbf{P}_{\omega^\perp} [\Psi[f]]),$$

where $\nabla_\omega, \nabla_\omega \cdot$ are the gradient and divergence operators on the sphere; $c > 0$ is a positive constant and ν, σ, Ψ are given functions.

There exists an interaction kernel $\mathbf{K} : \mathbb{R}^+ \times \Omega^2 \times (\mathbb{S}^{d-1})^2 \rightarrow \mathbb{R}^d$ that defines the flux \mathbf{J} as

$$(2.4) \quad \mathbf{J}[f](t, x, \omega) = \int_{\Omega \times \mathbb{S}} \mathbf{K}(t, x, x_*, \omega, \omega_*) f(t, x_*, \omega_*) dx_* d\omega_*.$$

From the flux \mathbf{J} we define the functional Ψ as

$$(2.5) \quad \exists \alpha \in [0, 1], \quad \Psi[f](x, \omega) = \frac{\mathbf{J}[f](t, x, \omega)}{|\mathbf{J}[f](t, x, \omega)|_\alpha},$$

where

$$(2.6) \quad |\mathbf{J}[f](t, x, \omega)|_\alpha = \alpha + (1 - \alpha) |\mathbf{J}[f](t, x, \omega)|,$$

where $|\cdot|$ stands for the norm in \mathbb{R}^d . Note that when $\alpha = 0$ we talk about a normalised operator since $|\Psi[f]| = 1$ whereas for $\alpha = 1$ we have a complete kernel operator.

The orthogonal kernel non-linearity (2.3) we tackle is more general than the gradient-type interaction (2.7). Furthermore, it is also more physically relevant when one wants to derive the Kolmogorov-Vicsek type of kinetic equation from particle systems behaviour [2, 3]. The main issue for the kinetic part is the possible degeneracy of $|\Psi[f]|_\alpha$ as well as the nonlinearity of the dissipativity and viscosity that does not necessarily compensate the aforementioned degeneracy, unlike existing works in the literature. The key part of our strategy is to provide first a quantification of the possible time of degeneracy of the nonlinearity combined to a study of the dependencies over σ and ν at a linear level.

Remark 2.2. *In the literature, typically, the operator K is of the form $\mathbf{K}(x, x_*, \omega, \omega_*) = \mathbf{K}(x - x_*, \omega_*)$. In this case it holds that*

$$\mathbf{P}_{\omega^\perp}(\Psi[f]) = \nabla_\omega \left(\omega \cdot \int_{\Omega \times \mathbb{S}^{d-1}} \mathbf{K}(x - x_*, \omega_*) f(x_*, \omega_*) dx_* d\omega_* \right)$$

because for a vector \mathbf{X} in \mathbb{R}^d , $\nabla_\omega(\omega \cdot X) = \mathbf{P}_{\omega^\perp}(\mathbf{X})$. Therefore, in this case the Kolmogorov-Vicsek equation (2.3) can be rewritten as a gradient-type interaction

$$(2.7) \quad \partial_t f + c\omega \cdot \nabla_x f = \nabla_\omega \cdot (\sigma(f) \nabla_\omega f) + \nabla_\omega \cdot (\nu(f) f \nabla_\omega \psi[f]).$$

Theorem 2.3. *Let Ω being either \mathbb{T}^d or \mathbb{R}^d , α in $[0, 1]$ and σ , ν and \mathbf{K} satisfying the hypotheses (H1) – (H2) – (H3). Let p belong to $[2, +\infty]$ and f_0 be such that*

(i) f_0 is a non-negative function in $L^1_{x,\omega} \cap L^p_{x,\omega}$ with mass

$$\int_{\Omega \times \mathbb{S}^{d-1}} f_0(x, \omega) dx d\omega := M_0 > 0;$$

(ii) $\inf_{(x,\omega) \in \Omega \times \mathbb{S}^{d-1}} |\mathbf{J}[f_0](x, \omega)|_\alpha := J_0 > 0$, where $|\mathbf{J}[f_0]|_\alpha$ was defined in (2.6).

There exists a time $T_{\max} > 0$, independent of p , and a unique weak solution f in $L^2([0, T_{\max}), L^1_{x,\omega} \cap L^2_{x,\omega})$ to (2.3) with f_0 as initial datum. Moreover, f is non-negative on $[0, T_{\max})$, belongs to $L^\infty([0, T_{\max}), L^p_{x,\omega}) \cap L^2([0, T_{\max}), L^2_x H^1_\omega)$, preserves the mass M_0 and one of the following holds

(a) $T_{\max} = +\infty$

(b) $\lim_{t \rightarrow T_{\max}^-} \sup_{\Omega \times \mathbb{S}^{d-1}} \frac{\nu(f)}{|\mathbf{J}[f]|_\alpha} = +\infty$.

Remark 2.4. *Of important note are the following consequences.*

- Our Cauchy theory does not stand on any a priori assumption of the solutions and provide an explicit lower bound for T_{\max} (see (3.17) – (3.19)).
- The theorem above includes all the previous results made in L^2 or L^∞ , both in the non-homogeneous and the homogeneous case (it suffices to consider $\mathbf{K} = k(x_*)\mathbf{K}(\omega, \omega_*)$ with $\int_\Omega k = 1$).
- The global existence criterion offers direct global existence for non-normalized interactions $\alpha \neq 0$ but it also gives global existence for the original Vicsek equation with spatially-homogeneous kernel $\mathbf{K}(t, x, x_*, \omega, \omega_*) = \omega_*$ and fully homogeneous viscosity and negative friction (i.e. only f and time-dependent). Section 3.3.1 describes several general cases where global existence happens in the problematic and purely normalised case $\alpha = 0$.

Besides the issue of well-posedness, it is of great interest to understand the large time behaviour of the solutions. We recall that Section 3.3.1 proves global existence for different types of interactions and it also exhibits a free energy which decreases along the flow. One cannot expect a general theory since it heavily depends on the shape of the kernel \mathbf{K} but one can still wonder if there are equilibria and if they are attractive. For this purpose, kinetic equations with gradient non-linearities (2.7) are often used because one can extract an explicit free energy functional decreasing along time. Hence leading to the existence of equilibria and a hope for an asymptotic study

of the solutions. This has been tackled in [29, 10, 13] where the authors exhibited a decreasing energy functional

$$(2.8) \quad \mathcal{F}(t) = \int_{\mathbb{S}^{d-1}} f \ln(f) d\omega + \frac{1}{2} \int_0^{|\mathbf{J}_f|} \frac{\nu(s)}{\sigma(s)} ds$$

when σ and ν are solely functions of $|\mathbf{J}_f|$. Associated to the latter is an energy dissipation allowing to dig out explicitly the equilibria in the spatially homogeneous setting.

It appears that a free energy is also underlying in the general equation: (2.3) enjoys a free energy functional that decreases along the flow with an explicit energy dissipation. Namely,

$$(2.9) \quad \begin{aligned} \mathcal{F}[f](t) &= \int_{\Omega \times \mathbb{S}^{d-1}} f \ln(f) dx d\omega \\ &+ \int_0^t \int_{\Omega \times \mathbb{S}^{d-1}} f \left[\nabla_\omega \cdot (\nu(f) \mathbf{P}_{\omega^\perp}[\Psi[f]]) - \frac{\nu(f)^2}{\sigma(f)} |\mathbf{P}_{\omega^\perp}[\Psi[f]]|^2 \right] dx d\omega ds \end{aligned}$$

and

$$(2.10) \quad \mathcal{D}[f](t) = \int_{\Omega \times \mathbb{S}^{d-1}} \sigma(f) f \left| \nabla_\omega \ln(f) + \frac{\nu(f)}{\sigma(f)} \mathbf{P}_{\omega^\perp}[\Psi[f]] \right|^2 dx d\omega.$$

which will be proven, in Section 3.3, to satisfy along the flow

$$\frac{d}{dt} \mathcal{F}[f](t) = -\mathcal{D}[f](t).$$

Remark 2.5. *Let us emphasize that considering the spatially homogeneous case $\mathbf{K} = \omega_*$, (2.9) and (2.10) are the ones obtained in [10] with $|\mathbf{J}_f|^2$ instead of $|\mathbf{J}_f|$ (in case of constant viscosity and friction we also recover the original Doi-Onsager free energy [39, 23]). For general gradient kernel interactions $\psi[f]$ (2.7) that are symmetric in ω and ω_* the free energy (2.9) becomes*

$$\mathcal{F}(t) = \int_{\mathbb{S}^{d-1}} f \ln(f) d\omega + \frac{1}{2} \int_0^{\langle \psi[f], f \rangle_{L^2_{x,\omega}}} \frac{\nu(s)}{\sigma(s)} ds$$

for ν and σ being solely functions of $\langle \psi[f], f \rangle_{L^2_{x,\omega}}$ (which equals $|\mathbf{J}_f|^2$). Hence offering a new view on the gradient structure where the natural dependencies are, in fact, $\langle \psi[f], f \rangle_{L^2_{x,\omega}} = |\mathbf{J}_f|^2$.

One can immediately see that the energy dissipation vanishes on

$$(2.11) \quad \mathcal{E}_\infty = \left\{ f_\infty \geq 0 \in L_t^\infty L_x^p H_\omega^2, \quad \nabla_\omega (\ln f_\infty)(t, x, \omega) = -\frac{\nu(f_\infty)}{\sigma(f_\infty)} \mathbf{P}_{\omega^\perp}[\Psi[f_\infty]](t, x, \omega) \right\}$$

and when it vanishes so does the right-hand side of the kinetic equation (2.3). The latter with the decrease of $\mathcal{F}[f](t)$ is a good hope that one could get a La Salle's invariance principle - in the spirit of [29] - when solutions are globally defined: namely the solution draws closer to the set of local equilibria \mathcal{E}_∞ .

Remark 2.6. *If one establishes a La Salle's principle then necessarily, for a global equilibrium f_∞ the quantity $\frac{\nu(f_\infty)}{\sigma(f_\infty)} \mathbf{P}_{\omega^\perp}[\Psi[f_\infty]] = \nabla_\omega (\psi[f])$ must be a tangential gradient - and then $f_\infty(t, x, \omega) = A(t, x) e^{-\psi[f](t, x, \omega)}$ and we recover generalised von Mises*

equilibria of the previous [29, 10, 13]. Hence, as mentioned earlier, a gradient structure pops out naturally but does not solely concern $\mathbf{P}_{\omega^\perp}[\Psi[f]]$ as first considered in (2.7). It however is not the purpose of this article and we did not investigate further.

2.3. Microscopic point of view and mean-field limit: strategy and results.

We consider a system of N particles given by their positions $X_t^{i,N} \in \Omega$ and velocities $V_t^{i,N} \in \mathbb{R}^d$ over time $t \geq 0$. Notice that we shall consider that the equations are written for $v \in \mathbb{R}^d$ rather than in $\omega \in \mathbb{S}^{d-1}$, but we shall later prove that the dynamics are restricted to the sphere, thus recovering the expected spherical dynamics. We recall the empirical distribution of the process $(X_t^{i,N}, V_t^{i,N})_{1 \leq i \leq N}$:

$$(2.12) \quad \mu_t^N(x, v) := \frac{1}{N} \sum_{i=1}^N \delta_{(X_t^{i,N}, V_t^{i,N})}(x, v),$$

The evolution of the particles is given by the following Stratonovich stochastic differential equation:

$$(2.13a) \quad \begin{cases} dX_t^{i,N} = V_t^{i,N} dt, \\ dV_t^{i,N} = \nu(\mu_t^N) \mathbf{P}_{(V_t^{i,N})^\perp}(\Psi(X_t^{i,N}, V_t^{i,N}, \mu_t^N)) dt \\ \quad + \frac{1}{2} \mathbf{P}_{(V_t^{i,N})^\perp}[(\nabla_v \sigma(\mu_t^N))(X_t^{i,N}, V_t^{i,N})] dt \\ \quad + \sqrt{2\sigma(\mu_t^N)} \mathbf{P}_{(V_t^{i,N})^\perp} \circ dB_t^i, \\ X_t^{i,N}(t=0) = X_0^{i,N}, \quad V_t^{i,N}(t=0) = V_0^{i,N}, \end{cases}$$

where \mathbf{P}_{v^\perp} is the projection operator

$$\mathbf{P}_{v^\perp} = \text{Id} - \frac{v \otimes v}{|v|^2},$$

where Id is the identity matrix; $((B_t^i)_{t \geq 0})_{i=1, \dots, N}$ are independent Brownian motions in \mathbb{R}^d . The symbol ' \circ ' denotes that the stochastic differential equation (2.13) is in Stratonovich convention.

Remark 2.7. Notice that the term (2.13b) in system (2.13) appears so that we obtain the kinetic equation (2.3). This is just a technicality: system (2.13) in Itô's convention corresponds to:

$$(2.14a) \quad \begin{cases} dX^{i,N} = V^{i,N} dt, \\ dV^{i,N} = \nu(\mu^N) P_{(V^{i,N})^\perp}(\Psi(X^{i,N}, V^{i,N}, \mu^N)) dt \\ \quad + P_{(\bar{V}^{i,N})^\perp}[(\nabla_v \sigma(\mu^N))(\bar{X}^{i,N}, \bar{V}^{i,N})] dt \\ \quad + \sqrt{2\sigma(\mu^N)} P_{(V^{i,N})^\perp} dB_t^i \\ \quad - 2(d-1)\sigma(\mu^N) \frac{V^{i,N}}{|V^{i,N}|^2}, \\ X^{i,N}(t=0) = X_0^{i,N}, \quad V^{i,N}(t=0) = V_0^{i,N}. \end{cases}$$

See also Appendix A.2 for more details. Without the extra term (2.13b) we would obtain a kinetic equation where the operator in ω is of the form (see Eq. (A.3))

$$\Delta_\omega(\sigma(f)f) - \frac{1}{2}\nabla_\omega \cdot (\nabla_\omega \sigma(f)f).$$

But with the extra term the operator ω in the kinetic equation is of the form

$$\Delta_\omega(\sigma(f)f) - \nabla_\omega \cdot (\nabla_\omega \sigma(f)f) = \nabla_\omega \cdot (\sigma(f)\nabla_\omega f),$$

which is the operator that we are dealing with in (2.3). Notice that if we wanted to obtain just a factor $\Delta_\omega(\sigma(f)f)$ then the constant in front of the extra term (2.13b) should be $-1/2$ rather than $1/2$. Notice also that if $\sigma = \sigma(t, x, f)$ does not depend on ω , then the extra term (2.13b) does not appear. This is the case, for example, when $\sigma = \sigma(|\tilde{\mathbf{J}}_f|(t, x))$ for $\tilde{\mathbf{J}}_f$ defined in (2.2), see [13].

As announced before, the dynamics described by our agent-based model indeed force the velocities to have unit norm, as shown by the next lemma.

Lemma 2.8. *Suppose that $(X_t^{i,N}, V_t^{i,N})$ is a solution to the particle dynamics (2.13). Suppose that $|V_0^{i,N}| = 1$ for all $i = 1, \dots, N$, then it holds that*

$$|V_t^{i,N}| = 1, \quad \text{for all } i = 1, \dots, N,$$

for all times where the solution is defined.

Proof of Lemma 2.8. It is a direct check that

$$d|V_t^{i,N}|^2 = 0$$

using Stratonovich chain rule (see, for example, [26, p. 122]) and the fact that $V \cdot \mathbf{P}_{V^\perp} = 0$. \square

The potential degeneracy of Ψ breaks the Lipschitz regularity of the interactions and thus prevents standard agent-based well-posedness results to apply. There exist results for non-Lipschitz interactions [2] but their singularities differ from the one at stake in the present article.

As we can control the formation of the degeneracy at the mesoscopic level - see Theorem 2.3 - we shall in fact solve the microscopic system (2.13) when the number of agents N is sufficiently large. In the spirit of [3], we thus shall prove well-posedness of the system hand-in-hand with the mean-field limit with the aid of an auxiliary process which mixes microscopic dynamics with the mesoscopic distribution function f_t . The main differences with respect to [3] is that, first, we need to deal with the appearance of singularities in the functional Ψ defined in (2.5) when $\alpha = 0$ and $|\mathbf{J}|_0 = 0$ and, second, that the noise coefficient σ is not constant. Let us explicit the

auxiliary process $(\bar{X}_t^{i,N}, \bar{V}_t^{i,N})_{t \geq 0}$, $i = 1, \dots, N$, solution of

$$\begin{aligned}
 (2.15a) \quad & \left\{ \begin{aligned} d\bar{X}_t^{i,N} &= \bar{V}_t^{i,N} dt, \\ d\bar{V}_t^{i,N} &= \nu(f_t) \mathbf{P}_{(\bar{V}_t^{i,N})^\perp}(\Psi(\bar{X}_t^{i,N}, \bar{V}_t^{i,N}, f_t)) dt \\ &\quad + \frac{1}{2} \mathbf{P}_{(\bar{V}_t^{i,N})^\perp}[(\nabla_\omega \sigma(f))(\bar{X}_t^{i,N}, \bar{V}_t^{i,N})] dt \\ (2.15b) \quad &\quad + \sqrt{2\sigma(f_t)} \mathbf{P}_{(\bar{V}_t^{i,N})^\perp} \circ dB_t^i, \\ (2.15c) \quad & f_t = \text{law}(\bar{X}_t^{i,N}, \bar{V}_t^{i,N}), \\ (2.15d) \quad & \bar{X}_t^{i,N}(t=0) = X_0^{i,N}, \quad \bar{V}_t^{i,N}(t=0) = V_0^{i,N}, \end{aligned} \right.
 \end{aligned}$$

where the initial data and the Brownian processes $(B_t^i)_{t \geq 0}$ are the same ones as in (2.13). Notice that all the processes $(\bar{X}_t^{i,N}, \bar{V}_t^{i,N})$ are independent by construction and they all have the same law f_t .

We will prove the following two results for the particle approximation (2.13) to the kinetic equation (2.3).

Theorem 2.9 (Local-in-time existence and uniqueness of solutions). *Under the assumptions of Theorem 2.3 and assumption (H4), let f_0 be a probability measure on $\mathbb{R}^d \times \mathbb{S}^{d-1}$ with finite second moment in $x \in \mathbb{R}^d$ and let $(X_0^{i,N}, V_0^{i,N})$ for $i = 1, \dots, N$ be N independent random variables with law f_0 . Under the hypotheses of Theorem 2.3, the following holds:*

- (i) *There exists $T > 0$ such that for N large enough, there exists a pathwise unique solution for $t \in [0, T)$ to the SDE system (2.13) with initial data $(X_0^{i,N}, V_0^{i,N})$ for $i = 1, \dots, N$. Moreover, the solution is such that $|V_t^{i,N}| = 1$ for all times where the solution is defined.*
- (ii) *There exists $T > 0$ such that, there exists a pathwise unique solution to the auxiliary process (2.15) with initial data $(X_0^{i,N}, V_0^{i,N})$ for $i = 1, \dots, N$ and $|\bar{V}_t^{i,N}| = 1$ for all times where the solution is defined.*
- (iii) *There exists a local-in-time unique weak solution of the kinetic equation (2.3) with initial datum f_0 . The solution of the kinetic equation is the law of the process solution to the auxiliary System (2.15), wherever the solution is defined.*

Along with this well-posedness result for the agent-based model we rigorously derive the mean-field limit between the microscopic system (2.13) and (2.3).

Theorem 2.10 (Mean-field limit - propagation of chaos). *Under the assumptions of Theorem 2.3 and assumption (H4), for the respective solutions $(X_t^{i,N}, V_t^{i,N})_{t \geq 0}$ and $(\bar{X}_t^{i,N}, \bar{V}_t^{i,N})_{t \geq 0}$ of (2.13) and (2.15), there exists $T > 0$ such that, for N large enough, it holds*

$$(2.16) \quad \lim_{N \rightarrow \infty} \sup_{1 \leq i \leq N} \mathbb{E} \left[\sup_{0 \leq t \leq T} (|X_t^{i,N} - \bar{X}_t^{i,N}|^2 + |V_t^{i,N} - \bar{V}_t^{i,N}|^2) \right] = 0.$$

Remark 2.11. In [3] the Vicsek model with no normalisation ($\alpha = 1$) is investigated so there is no restriction on the time T or the value of N . Here Theorems 2.9 and 2.10 come with restrictions on the time interval $t \in [0, T]$ and are valid only for N large enough. These constraints are due to the fact that these theorems hold true as long as

$$|\mathbf{J}|(X_t^{i,N}, V_t^{i,N}, \mu_t^N) > 0,$$

and we can prove that this holds true only for short times and for N large enough.

3. THE MESOSCOPIC FRAMEWORK: KINETIC DIFFERENTIAL EQUATION

3.1. Linear equation: dependence on coefficients and averaging positivity.

The differential operator ∇_ω is the tangential gradient - also called Günter derivatives - on the sphere \mathbb{S}^{d-1} . It can be related to the standard gradient on \mathbb{R}^d by

$$\nabla_\omega = \nabla_v - \langle \nabla_v, \omega \rangle_{\mathbb{R}^d} \omega = \mathbf{P}_{\omega^\perp} [\nabla_v]$$

where ∇_v is the euclidean gradient for functions from \mathbb{R}^d to \mathbb{R}^d . First of all let us emphasize that the tangential gradient ∇_ω displays very different behaviour than the usual gradient. We give here three formulas that we shall use throughout the proofs. We refer the interested reader to [24, 1] for an introduction on tangential - Günter-derivatives and to [40, Appendix II] for the specific calculus of the following formulas (proven in dimension 3 but immediately extendable in dimension d). Integration by parts is allowed but generate an additional term in the direction of the orientation ω :

$$(3.1) \quad \int_{\mathbb{S}^{d-1}} \nabla_\omega (f(\omega)) g(\omega) d\omega = - \int_{\mathbb{S}^{d-1}} f(\omega) \nabla_\omega (g(\omega)) d\omega + (d-1) \int_{\mathbb{S}^{d-1}} \omega f g d\omega.$$

It also holds the following explicit tangential derivatives for each coordinate ω_i of $\omega = (\omega_1, \dots, \omega_d)$ on the sphere

$$(3.2) \quad \nabla_\omega (\omega_i) = (\delta_{ij} - \omega_i \omega_j)_{1 \leq j \leq d} \quad \text{and} \quad \Delta_\omega (\omega_i) = -(d-1) \omega_i$$

As mentioned in the introduction, (2.3) can be viewed as a nonlinear Fokker-Planck equation on the torus with local and nonlinear viscosity σ and drift ν . The purpose of this section is to study the associated linear equation when the nonlinearities are considered as given parameters. In other words we consider here the following differential problem

$$(3.3) \quad \partial_t f + c\omega \cdot \nabla_x f = \nabla_\omega \cdot (\bar{\sigma}(t, x, \omega) \nabla_\omega f + f \bar{\Psi}(t, x, \omega))$$

where $\bar{\sigma}$ and $\bar{\Psi}$ are given functions.

The issue of existence and uniqueness for (3.3) has been solved for instance in [7, Appendix A] for velocities in \mathbb{R} or \mathbb{R}^2 for constant $\bar{\sigma}$ with $\bar{\Psi}$ and $\nabla_\omega \cdot \bar{\Psi}$ being in $L_{x,\omega}^\infty$. However their methods are directly applicable for velocities in \mathbb{S}^{d-1} as shown in [30, Lemma 4.1] for constant $\bar{\sigma}$. The case of non-constant $\bar{\sigma}$ is a straightforward adaptation of their proofs if $\bar{\sigma}$ belongs to $L_{t,x,\omega}^\infty$ and is uniformly bounded from below by a constant $\bar{\sigma}_0 > 0$ (note that one could also adapt to tangential derivatives standard Galerkin type methods for linear parabolic equations [25, Chapter 6], as done in [29] in the spatially homogeneous case). We thus omit the proof and state:

Theorem 3.1. *Let $\bar{\sigma}$ be in $L_{t,x,\omega}^\infty$ uniformly bounded from below: $\bar{\sigma}(t, x, \omega) \geq \bar{\sigma}_0 > 0$ and let $\bar{\Psi}$ and $\nabla_\omega \cdot \bar{\Psi}$ be in $L_{t,x,\omega}^\infty$ with the orthogonal property $\mathbf{P}_{\omega^\perp}(\bar{\Psi}) = \bar{\Psi}$. Then for any f_0 in $L_{x,\omega}^2$ there exists a unique f in the space*

$$Y = \{u \in L_{t,x}^2 H_\omega^1, \quad \partial_t u + \omega \cdot \nabla_x u \in L_{t,x}^2 H_\omega^{-1}\}$$

solution to (3.3) with initial datum f_0 .

Moreover, if f_0 is non-negative then f is non-negative.

The next Section 3.1.1 is dedicated to L^p bounds and gain of regularity for solutions to (3.3) whilst Section 3.1.2 studies the dependences on the viscosity and friction. To conclude, Section 3.1.3 tackles the issue of an explicit lower bound on the vanishing time for velocity averaging densities.

3.1.1. *L^p bounds and gain of regularity.* The issue of $L_{x,\omega}^p$ estimates and gain of regularity had already been investigated in previous cited references for particular $\bar{\sigma}$ and $\bar{\Psi}$ and we provide here an adapted version that fits our general setting.

Proposition 3.2. *Let p belong to $[2, +\infty]$ and $f_0 \geq 0$ be in $L_{x,\omega}^1 \cap L_{x,\omega}^p$ with mass $\|f_0\|_{L_{x,\omega}^1} = M_0$. Under the assumptions of Theorem 3.1 on $\bar{\sigma}$ and $\bar{\Psi}$ the solution f to (3.3) is in $L_{x,\omega}^p$, preserves the mass M_0 and gains regularity in ω since it satisfies*

$$\forall t \geq 0, \quad \|f(t)\|_{L_{x,\omega}^p} \leq e^{C_p(\bar{\sigma}, \bar{\Psi})t} \|f_0\|_{L_{x,\omega}^p},$$

and

$$\int_0^t e^{C_p(\bar{\sigma}, \bar{\Psi})(t-s)} \left\| f^{\frac{p-2}{2}} \nabla_\omega f(s) \right\|_{L_{x,\omega}^2}^2 \|f(s)\|_{L_{x,\omega}^p}^{1-p} ds \leq \frac{2 \|f_0\|_{L_{x,\omega}^p}}{(p-1)\bar{\sigma}_0} e^{C_p(\bar{\sigma}, \bar{\Psi})t},$$

where

$$(3.4) \quad C_p(\bar{\sigma}, \bar{\Psi}) = \|\nabla_\omega \cdot \bar{\Psi}\|_{L_{t,x,\omega}^\infty} + \frac{\|\bar{\Psi}\|_{L_{t,x,\omega}^\infty}^2}{(p-1)\bar{\sigma}_0},$$

with $C_\infty = \lim_{p \rightarrow \infty} C_p$.

Proof of Proposition 3.2. Step 1: L^p bounds and gain of regularity. Note that since ∇_ω and $\bar{\Psi}$ are orthogonal to ω , we can perform on (3.3) integration by parts on \mathbb{S}^{d-1} as if we were working with classical derivatives. Indeed, the orthogonality to ω exactly cancels the additional term in the tangential derivatives integration by parts formula (3.1). It yields for any $p \geq 2$,

$$\begin{aligned} I &:= \frac{1}{p} \frac{d}{dt} \|f(t)\|_{L_{x,\omega}^p}^p = \int_{\Omega \times \mathbb{S}^{d-1}} \operatorname{sgn}(f(t, x, v)) |f(t, x, v)|^{p-1} \partial_t f(t, x, \omega) dx d\omega \\ &= - \int_{\Omega \times \mathbb{S}^{d-1}} \operatorname{sgn}(f) |f|^{p-1} \nabla_x (c\omega f) - \int_{\Omega \times \mathbb{S}^{d-1}} \bar{\sigma}(t, x, \omega) \nabla_\omega (\operatorname{sgn}(f) |f|^{p-1}) \cdot \nabla_\omega f \\ &\quad + \int_{\Omega \times \mathbb{S}^{d-1}} \operatorname{sgn}(f) |f|^{p-1} \nabla_\omega \cdot (f \bar{\Psi}) dx d\omega \end{aligned}$$

We use that the first integrand on the right-hand side can be written as a divergence in x : $\nabla_x \cdot (c\omega |f|^p)$ whereas the second one can be written as $(p - 1)\bar{\sigma}(t, x, \omega) |f|^{p-2} (\nabla_\omega f)^2$. This leads to the following upper bound

$$\begin{aligned} I &\leq - (p - 1)\bar{\sigma}_0 \left\| f^{\frac{p-2}{2}} \nabla_\omega f \right\|_{L^2_{x,\omega}}^2 + \int_{\Omega \times \mathbb{S}^{d-1}} (|f|^p \nabla_\omega \cdot \bar{\Psi} + \bar{\Psi} |f|^{p-1} |\nabla_\omega f|) dx d\omega \\ &\leq - (p - 1)\bar{\sigma}_0 \left\| f^{\frac{p-2}{2}} \nabla_\omega f \right\|_{L^2_{x,\omega}}^2 + \|\nabla_\omega \cdot \bar{\Psi}\|_{L^\infty_{t,x,\omega}} \|f\|_{L^p_{x,\omega}}^p + \|\bar{\Psi}\|_{L^\infty_{t,x,\omega}} \|f\|_{L^p_{x,\omega}}^{\frac{p}{2}} \left\| f^{\frac{p-2}{2}} \nabla_\omega f \right\|_{L^2_{x,\omega}} \\ &\leq - \left((p - 1)\bar{\sigma}_0 - \frac{\eta}{2} \|\bar{\Psi}\|_{L^\infty_{t,x,\omega}}^2 \right) \left\| f^{\frac{p-2}{2}} \nabla_\omega f \right\|_{L^2_{x,\omega}}^2 + \left(\|\nabla_\omega \cdot \bar{\Psi}\|_{L^\infty_{t,x,\omega}} + \eta^{-1} \right) \|f\|_{L^p_{x,\omega}}^p. \end{aligned}$$

Note that we used Cauchy-Schwarz and Young inequality for any $\eta > 0$. Choosing $\frac{p-1}{2}\bar{\sigma}_0 - \frac{\eta}{2} \|\bar{\Psi}\|_{L^\infty_{t,x,v}}^2 = 0$ end up with

$$\begin{aligned} \frac{d}{dt} \|f(t)\|_{L^p_{x,\omega}} &= \frac{1}{p} \frac{d}{dt} \|f(t)\|_{L^p_{x,\omega}}^p \|f(t)\|_{L^p_{x,\omega}}^{1-p} \\ &\leq -\frac{(p-1)}{2} \bar{\sigma}_0 \left\| f^{\frac{p-2}{2}} \nabla_\omega f \right\|_{L^2_{x,\omega}}^2 \|f\|_{L^p_{x,\omega}}^{1-p} + C_p \|f\|_{L^p_{x,\omega}}. \end{aligned}$$

From which we deduce thanks to Grönwall lemma that for all $t \geq 0$

$$(3.5) \quad \|f(t)\|_{L^p_{x,\omega}} + \frac{(p-1)}{2} \bar{\sigma}_0 \int_0^t e^{C_p(t-s)} \left\| f^{\frac{p-2}{2}}(s) \nabla_\omega f(s) \right\|_{L^2_{x,\omega}}^2 \|f(s)\|_{L^p_{x,\omega}}^{1-p} ds \leq e^{C_p t} \|f_0\|_{L^p_{x,\omega}}.$$

The latter is exactly the standard bounds in Proposition 3.2 for p in $[2, +\infty)$.

Step 2: Mass conservation and L^∞ bounds. Now let us suppose that f_0 is non-negative, which implies that f is non-negative at all time by Theorem 3.1. Hence, it holds that

$$\begin{aligned} \frac{d}{dt} \|f(t)\|_{L^1_{x,\omega}} &= \frac{d}{dt} \int_{\Omega \times \mathbb{S}^{d-1}} f(t, x, \omega) dx d\omega \\ &= \int_{\Omega \times \mathbb{S}^{d-1}} \nabla_x \cdot (c\omega f) + \nabla_\omega \cdot (\bar{\sigma} \nabla_\omega f + f \bar{\Psi}) dx d\omega = 0 \end{aligned}$$

because, again, the integration by parts (3.1) does not generate additional terms in the direction of ω . This concludes the preservation of the $L^1_{x,\omega}$ -norm.

The $L^\infty_{x,\omega}$ estimates follows straight from the limit p goes to $+\infty$, since $f(t)$ belongs to $L^1_{x,\omega}$. \square

3.1.2. Dependence on the coefficients. Our strategy to tackle the non-linear equation is *via* a fixed point argument. We thus need to understand how solutions to (3.3) differ from one another when they are associated to different coefficients $\bar{\sigma}$ and $\bar{\Psi}$. The main issue in establishing estimate on $\|f_1 - f_2\|_{L^p_{x,\omega}}$ of two different solutions relies on the fact that the gain of regularity proved in Proposition 3.2 is highly non-linear as soon as $p > 2$. Moreover, we did not manage to quantify the way the gain of regularity vanishes at initial time so we cannot close a direct L_t^∞ fixed point method. We shall thus only study the dependence on coefficient in $L^2_{t,x}$.

Proposition 3.3. *Let f_0 be a non-negative function in $L^1_{x,\omega} \cap L^2_{x,\omega}$. Suppose that $\bar{\sigma}_1, \bar{\sigma}_2$ and $\bar{\Psi}_1, \bar{\Psi}_2$ satisfy the assumptions of Theorem 3.1. Let f_1 and f_2 be the solutions of (3.3) associated respectively to the coefficient $(\bar{\sigma}_1, \bar{\Psi}_1)$ and $(\bar{\sigma}_2, \bar{\Psi}_2)$ with initial datum f_0 .*

Then the following holds for any $t \geq 0$,

$$\begin{aligned} \|(f_1 - f_2)(t)\|_{L^2_{x,\omega}}^2 &\leq \frac{4e^{3C_2 t} \|f_0\|_{L^2_{x,\omega}}^2}{\inf\{\bar{\sigma}_1 + \bar{\sigma}_2\}} \int_0^t \|(\bar{\Psi}_1 - \bar{\Psi}_2)(s)\|_{L^\infty_{x,\omega}}^2 ds \\ &\quad + \frac{2\|f_0\|_{L^2_{x,\omega}}}{\inf\{\bar{\sigma}_1 + \bar{\sigma}_2\}} \left[\int_0^t e^{C_2(t-s)} \left(\|\nabla_\omega f_1\|_{L^2_{x,\omega}}^2 + \|\nabla_\omega f_2\|_{L^2_{x,\omega}}^2 \right) ds \right] \|\bar{\sigma}_1 - \bar{\sigma}_2\|_{L^\infty_{t,x,\omega}}^2, \end{aligned}$$

where $C_2 = C_2(\bar{\sigma}_1, \bar{\Psi}_1) + C_2(\bar{\sigma}_2, \bar{\Psi}_2)$ defined in Proposition 3.2.

Remark 3.4. *The main difficulty with dealing with local-in-time $\sigma(f)$ appears in the second term on the right-hand side above. Indeed, from Proposition 3.2 we see that $\|\nabla_\omega f_i\|_{L^2_{x,\omega}}^2$ is integrable on $[0, t]$ and thus $\int_0^t \|\nabla_\omega f_i\|_{L^2_{x,\omega}}^2$ vanishes when t goes to 0, thus almost allowing to close a contraction fixed point argument. As we shall see later on, we however need an explicit independence over $\bar{\sigma}$ (other than $\inf \bar{\sigma}$ and $\sup \bar{\sigma}$) to avoid any nonlinear breakdown of contraction theorem in short times. As a result a $L^2([0, T], L^2_{x,\omega})$ setting is required to get an extra integration in time. This explains hypothesis (H2), but one could also ask for more regularity for σ and f_0 to explicitly estimate the convergence to 0 of $\int_0^t \|\nabla_\omega f_i\|_{L^2_{x,\omega}}^2$ with parabolic regularity.*

Proof of Proposition 3.3. To shorten notations we will use $\bar{\sigma}^+ = \bar{\sigma}_1 + \bar{\sigma}_2$, $\bar{\sigma}^- = \bar{\sigma}_1 - \bar{\sigma}_2$, $\bar{\Psi}^+ = \bar{\Psi}_1 + \bar{\Psi}_2$ and $\bar{\Psi}^- = \bar{\Psi}_1 - \bar{\Psi}_2$. Also we denote the constant constructed from (3.4) by $C_2 = \max\{C_2(\bar{\sigma}_1, \bar{\Psi}_1), C_2(\bar{\sigma}_2, \bar{\Psi}_2), C_2(\bar{\sigma}^+, \bar{\Psi}^+), C_2(\bar{\sigma}^-, \bar{\Psi}^-)\}$.

Using the algebraic identity $ab - cd = \frac{1}{2}[(a - c)(b + d) + (a + c)(b - d)]$ we find

$$\begin{aligned} \partial_t [f_1 - f_2] + c\omega \nabla_x \cdot [f_1 - f_2] &= \frac{1}{2} \nabla_\omega \cdot ((\bar{\sigma}_1 + \bar{\sigma}_2) \nabla_\omega [f_1 - f_2]) + \frac{1}{2} \nabla_\omega \cdot ([f_1 - f_2] (\bar{\Psi}_1 + \bar{\Psi}_2)) \\ &\quad + \frac{1}{2} \nabla_\omega \cdot ((\bar{\sigma}_1 - \bar{\sigma}_2) \nabla_\omega (f_1 + f_2)) + \frac{1}{2} \nabla_\omega \cdot [(f_1 + f_2) (\bar{\Psi}_1 - \bar{\Psi}_2)], \end{aligned}$$

We start by bounding the terms in $f_1 - f_2$ as in Proposition 3.2 and we obtain

$$(3.6) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|f_1 - f_2\|_{L^2_{x,\omega}}^2 &\leq -\frac{\bar{\sigma}_0^+}{4} \|\nabla_\omega (f_1 - f_2)\|_{L^2_{x,\omega}}^2 + C_2 \|f_1 - f_2\|_{L^2_{x,\omega}}^2 \\ &\quad + \frac{1}{2} \int_{\Omega \times \mathbb{S}^{d-1}} (f_1 - f_2) \nabla_\omega \cdot \left[\bar{\sigma}^- \nabla_\omega (f_1 + f_2) + \bar{\Psi}^- (f_1 + f_2) \right] dx d\omega. \end{aligned}$$

where $\bar{\sigma}_0^+ = \inf\{\bar{\sigma}_1 + \bar{\sigma}_2\}$. We bound the two terms inside the integral on the right-hand side. First thanks to an integration by parts (3.1).

$$\int_{\Omega \times \mathbb{S}^{d-1}} (f_1 - f_2) \nabla_\omega \cdot \left[\bar{\sigma}^- \nabla_\omega (f_1 + f_2) \right] dx d\omega = - \int_{\Omega \times \mathbb{S}^{d-1}} \bar{\sigma}^- \nabla_\omega (f_1 - f_2) \cdot \nabla_\omega (f_1 + f_2) dx d\omega.$$

Cauchy-Schwarz with Young's inequalities yields

$$(3.7) \quad \left| \int_{\Omega \times \mathbb{S}^{d-1}} (f_1 - f_2) \nabla_\omega \cdot [\bar{\sigma}^- \nabla_\omega (f_1 + f_2)] \right| \leq \frac{\bar{\sigma}_0^+}{8} \|\nabla_\omega (f_1 - f_2)\|_{L_{x,\omega}^2}^2 + \frac{4 \|\bar{\sigma}^-\|_{L_{x,\omega}^\infty}^2}{\bar{\sigma}_0^+} \|\nabla_\omega (f_1 + f_2)\|_{L_{x,\omega}^2}^2.$$

The second integrand in (3.6) is dealt with in the same way and we get

$$(3.8) \quad \left| \int_{\Omega \times \mathbb{S}^{d-1}} (f_1 - f_2) \nabla_\omega \cdot [\bar{\Psi}^- (f_1 + f_2)] \right| \leq \frac{\bar{\sigma}_0^+}{8} \|\nabla_\omega (f_1 - f_2)\|_{L_{x,\omega}^2}^2 + \frac{4 \|\bar{\Psi}^-\|_{L_{x,\omega}^\infty}^2}{\bar{\sigma}_0^+} \|(f_1 + f_2)\|_{L_{x,\omega}^2}^2.$$

We then plug (3.7) and (3.8) into (3.6) and get

$$\frac{1}{2} \frac{d}{dt} \|f_1 - f_2\|_{L_{x,\omega}^2}^2 \leq C_2 \|f_1 - f_2\|_{L_{x,\omega}^2}^2 + \frac{2 \|\bar{\sigma}^-\|_{L_{x,\omega}^\infty}^2}{\bar{\sigma}_0^+} \|\nabla_\omega (f_1 + f_2)\|_{L_{x,\omega}^2}^2 + \frac{2 \|\bar{\Psi}^-\|_{L_{x,\omega}^\infty}^2}{\bar{\sigma}_0^+} \|(f_1 + f_2)\|_{L_{x,\omega}^2}^2$$

on which we apply Grönwall lemma to obtain

$$(3.9) \quad \|f_1 - f_2\|_{L_{x,\omega}^2}^2 \leq \frac{2}{\bar{\sigma}_0^+} \|\bar{\sigma}^-\|_{L_{t,x,\omega}^\infty}^2 \int_0^t e^{C_2(t-s)} \|\nabla_\omega (f_1 + f_2)\|_{L_{x,\omega}^2}^2 ds + \frac{2}{\bar{\sigma}_0^+} \int_0^t e^{C_2(t-s)} \|(f_1 + f_2)\|_{L_{x,\omega}^2}^2 \|\bar{\Psi}^-(s)\|_{L_{x,\omega}^\infty}^2 ds.$$

To conclude we apply Proposition 3.2 to bound the second term on the right-hand side, which directly gives the expected result. \square

3.1.3. *Estimation of vanishing time for velocity averaging densities.* Now we turn to an explicit estimation of the vanishing time of velocity averaging densities.

Proposition 3.5. *Let p belong to $[2, +\infty]$ and $f_0 \geq 0$ be in $L_{x,\omega}^1 \cap L_{x,\omega}^p$. Let also $\bar{\sigma}$ and $\bar{\Psi}$ satisfy the assumptions of Theorem 3.1 supplemented with $\bar{\sigma}$ Lipschitz. Let f be the solution to (3.3) associated to f_0 . For any \mathbf{K} in $W_{t,x_*}^{1,\infty} W_{\omega_*}^{2,\infty} L_{x,\omega}^\infty$ denote*

$$\mathbf{J}(t, x, \omega) = (J_i(t, x, \omega))_{1 \leq i \leq d} = \int_{\Omega \times \mathbb{S}^{d-1}} f(t, x_*, \omega_*) \mathbf{K}(t, x, x_*, \omega, \omega_*) dx_* d\omega_*.$$

Then the following holds

$$\forall (t, x, \omega) \in \mathbb{R}^+ \times \Omega \times \mathbb{S}^{d-1}, \quad |\mathbf{J}(t, x, \omega)| \geq \sum_{i=1}^d |J_i(0, x, \omega)| - K_{i,\infty} \|f_0\|_{L_{x,\omega}^1} t$$

where we defined

$$(3.10) \quad K_{i,\infty} = \left(1 + |c| + \|\bar{\sigma}\|_{L_{t,x}^\infty W_\omega^{1,\infty}} + \|\bar{\Psi}\|_{L_{t,x,\omega}^\infty} \right) \|K_i\|_{W_{t,x_*}^{1,\infty} W_{\omega_*}^{2,\infty} L_{x,\omega}^\infty}.$$

Proof of Proposition 3.5. Since f_0 is non-negative and belongs to $L^1_{x,\omega} \cap L^p_{x,\omega}$ we deduce from Theorem 3.1 and Proposition 3.2 that $f(t)$ is non-negative and belongs to $L^1_{x,\omega} \cap L^p_{x,\omega}$ for all time. Hence, we can multiply (3.3) by $\mathbf{K}(x, x_*, \omega, \omega_*)$ and integrate by parts (3.1). Here again note that there are no additional terms along ω_* since $\overline{\Psi}$ and ∇_{ω_*} are both orthogonal to ω_* . This yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |J_i(t, x, \omega)|^2 &= J_i(t, x, \omega) \int_{\Omega \times \mathbb{S}^{d-1}} K_i(t, x, x_*, \omega, \omega_*) \frac{\partial f}{\partial t}(t, x_*, \omega_*) dx_* d\omega_* \\ &\quad + J_i(t, x, \omega) \int_{\Omega \times \mathbb{S}^{d-1}} f(t, x_*, \omega_*) \frac{\partial K_i}{\partial t}(t, x, x_*, \omega, \omega_*) dx_* d\omega_* \\ &= J_i(t, x, \omega) \int_{\Omega \times \mathbb{S}^{d-1}} f \left[\frac{\partial K_i}{\partial t} + c\omega \cdot \nabla_{x_*} K_i + \nabla_{\omega_*} \cdot (\overline{\sigma} \nabla_{\omega_*} K_i) - \overline{\Psi} \cdot \nabla_{\omega_*} K_i \right] dx_* d\omega_* \\ &\geq -|J_i(t, x, \omega)| \left(1 + |c| + \|\overline{\sigma}\|_{L^\infty_{t,x} W^{1,\infty}} + \|\overline{\Psi}\|_{L^\infty_{t,x,\omega}} \right) \|K_i\|_{W^{1,\infty}_{t,x_*} W^{2,\infty}_{\omega_*} L^\infty_{x,\omega}} \|f(t)\|_{L^1_{x,\omega}} \\ &\geq -K_{i,\infty} \|f_0\|_{L^1_{x,\omega}} |J_i(t, x, \omega)| \end{aligned}$$

where we used the conservation of mass along time. The latter implies that

$$|J_i(t, x, \omega)| \geq |J_i(0, x, \omega)| - K_{i,\infty} \|f_0\|_{L^1_{x,\omega}} t$$

which concludes the proof. \square

3.2. The local-in-time nonlinear Cauchy theory. The present section is devoted to the proof of Theorem 2.3 thanks to the linear study we presented in Sec. 3.1. We shall prove existence, uniqueness and global existence criterion to

$$(3.11) \quad \partial_t f + c\omega \cdot \nabla_x f = \nabla_\omega \cdot (\sigma(f) \nabla_\omega f) + \nabla_\omega \cdot (\nu(f) f \mathbf{P}_{\omega^\perp} [\Psi[f]]).$$

Proof of Theorem 2.3. We fix p in $[2, +\infty]$ and $f_0 \geq 0$ in $L^1_{x,\omega} \cap L^p_{x,\omega}$. We recall some assumptions of Theorem 2.3:

$$\begin{aligned} \int_{\Omega \times \mathbb{S}^{d-1}} f_0(x, \omega) dx d\omega &= M_0 > 0 \\ \inf_{(x,\omega) \in \Omega \times \mathbb{S}^{d-1}} |\mathbf{J}[f_0](x, \omega)|_\alpha &= J_0 > 0. \end{aligned}$$

The strategy is to apply a contraction fix point argument so we start by defining the Banach space on which we shall work. For any $M \geq M_0$ and $T > 0$ we define

$$(3.12) \quad \Gamma_T^M = \left\{ f \in L^2_{[0,T],x,\omega}, \quad \text{and} \quad \begin{array}{l} \text{ess sup}_{[0,T]} \|f(t)\|_{L^1_{x,\omega}} \leq M \\ \text{ess inf}_{[0,T] \times \Omega \times \mathbb{S}^{d-1}} |\mathbf{J}[f(t)](x, \omega)|_\alpha \geq \alpha + (1 - \alpha) \frac{J_0}{2} \end{array} \right\}.$$

For a function g in Γ_T^M we have that g belongs to $L^2_{[0,T],x,\omega}$ so by Fubini's theorem g belongs to $L^2([0, T], L^2_{x,\omega})$ and for almost every t in $[0, T]$ the function $g(t, \cdot, \cdot)$ belongs to $L^2_{x,\omega}$. Therefore $\sigma(g(t))$ and $\nu(g(t))$ are defined almost everywhere in $[0, T]$ and with hypotheses (H2) and (H3) one has

$$\sigma_0 \leq \sigma(g(t)) \leq \sigma_\infty \quad \text{and} \quad \|\nu(g(t))\|_{L^\infty_{x,\omega} W^{1,\infty}} \leq \nu_\infty \quad \text{almost everywhere in } [0, T].$$

Defining

$$\overline{\Psi}(t, x, \omega) = \nu(g(t))(x, \omega) \mathbf{P}_{\omega^\perp} [\Psi[g]]$$

hypotheses (H1) shows for $t \in [0, T]$

$$(3.13) \quad \begin{aligned} |\overline{\Psi}(t, x, \omega)| &\leq \nu_\infty \frac{\left| \int_{\Omega \times \mathbb{S}^{d-1}} g(t, x_*, \omega_*) \mathbf{K}(t, x, x_*, \omega, \omega_*) dx_* d\omega_* \right|}{|\mathbf{J}[g](t, x, \omega)|_\alpha} \\ &\leq \nu_\infty \begin{cases} \|\mathbf{K}\|_{L^\infty_{t,x,x_*,\omega,\omega_*}} M & \text{if } \alpha = 1 \\ \frac{1}{1-\alpha}, & \text{if } \alpha \neq 1 \end{cases} := C_M^1 \end{aligned}$$

and

$$(3.14) \quad \begin{aligned} |\nabla_\omega \cdot \overline{\Psi}(t, x, \omega)| &\leq |\nabla_\omega \nu(g) \cdot \mathbf{P}_{\omega^\perp} [\Psi[g]]| + |\nu(g) \nabla_\omega \cdot \mathbf{P}_{\omega^\perp} [\Psi[g]]| \\ &\leq C_M^1 + \nu_\infty C_\alpha M \|\mathbf{K}\|_{L^\infty_{t,x,x_*,\omega_*} W_\omega^{1,\infty}} := C_M^2 \end{aligned}$$

where $C_\alpha > 0$ only depends on α and equals 1 when $\alpha = 1$. Therefore thanks to Theorem 3.1 we are able to define

$$\begin{aligned} \Phi : \Gamma_T^M &\longrightarrow L^2_{[0,T],x,\omega} \quad , \\ g &\longmapsto \Phi(g) = f_g \end{aligned}$$

where f_g is the solution on $[0, T] \times \Omega \times \mathbb{S}^{d-1}$ to the linear equation

$$\partial_t f + c\omega \cdot \nabla_x f = \nabla_\omega \cdot (\sigma(g)(t, x, \omega) \nabla_\omega f + f \nu(g) \mathbf{P}_{\omega^\perp} [\Psi[g]])$$

associated to the initial datum f_0 .

We now show that for a specific T , Φ is in fact a contraction from Γ_T^M to Γ_T^M . Due to Proposition 3.2 and since f_0 belongs to $L^2_{x,\omega}$ we see that $\Phi(g)$ belongs to $L^\infty([0, T])L^2_{x,\omega} \subset L^2_{[0,T] \times \Omega \times \mathbb{S}^{d-1}}$ and also that for almost every $t \in [0, T]$

$$\|\Phi(g)(t)\|_{L^1_{x,\omega}} = M_0 < M.$$

Moreover, Proposition 3.5 gives that almost everywhere

$$(3.15) \quad \begin{aligned} |\mathbf{J}[\Phi(g)](t, x, \omega)| &\geq \sum_{i=1}^d |J_i[f](0, x, \omega)| - K_{i,\infty} M_0 t \\ &\geq |\mathbf{J}[f](0, x, \omega)| - K_\infty M_0 t \\ &\geq J_0 - K_\infty M_0 T. \end{aligned}$$

We defined $K_\infty = \sum_{i=1}^d K_{i,\infty}$ which by (3.10) explicitly gives

$$(3.16) \quad K_\infty(f_0) = (1 + |c| + \sigma_\infty + \sigma_{lip} + C_M^1) \sum_{n=1}^d \|K_i\|_{W_{t,x_*}^{1,\infty} W_{\omega_*}^{2,\infty} L_{x,\omega}^\infty}.$$

Therefore if we choose

$$(3.17) \quad T < T_0 := \frac{J_0}{2K_\infty M_0},$$

the lower bound (3.15) implies

$$\forall t \in [0, T], \quad |\mathbf{J}[\Phi(g)](t, x, \omega)|_\alpha \geq \alpha + (1 - \alpha) \frac{J_0}{2},$$

and we thus proved that for any $0 < T < T_0$, Φ maps Γ_T^M to itself.

It remains to prove that Φ is a contraction on Γ_T^M for T sufficiently small. By hypothesis (H2) we have

$$\|\sigma(g_1) - \sigma(g_2)\|_{L_{[0,T],x,\omega}^\infty} \leq \sigma_{\text{lip}} \|g_1 - g_2\|_{L_{[0,T],x,\omega}^2}.$$

Also, thanks to the Lipschitz property (H3) of the friction ν and the kernel form of $\Psi[g]$ one infers

$$\int_0^t \|\overline{\Psi}(g_1) - \overline{\Psi}(g_2)\|_{L_{x,\omega}^\infty}^2 \leq \Psi_{\text{lip}} \|g_1 - g_2\|_{L_{[0,t],x,\omega}^2}^2$$

where $\Psi_{\text{lip}} > 0$ is a constant only depending on $\|\mathbf{K}\|_{L_{t,x,x_*,\omega,\omega_*}^\infty}$, M , α , Ψ_0 and ν_{lip} . This inequality comes from the algebraic identity $ab - cd = \frac{(a-c)(b+d)}{2} + \frac{(a+c)(b-d)}{2}$.

We apply Proposition 3.3 to see that for any g_1 and g_2 in Γ_T^M

$$\|\Phi(g_1) - \Phi(g_2)\|_{L_{[0,T],x,\omega}^2}^2 \leq \Lambda(T) \|g_1 - g_2\|_{L_{[0,T],x,\omega}^2}^2$$

where

$$(3.18) \quad \Lambda(T) = \frac{2 \|f_0\|_{L_{x,\omega}^2}^2}{\sigma_0^2} T e^{3C_2 T} (\Psi_{\text{lip}} \sigma_0 + 2\sigma_{\text{lip}}).$$

Note that we used Proposition 3.2 to obtain explicit constants. To conclude it suffices to choose

$$(3.19) \quad T < \min \left\{ T_0, \sup_{t>0} \Lambda(t) \leq \frac{1}{4} \right\} := T_1$$

because then for any g_1 and g_2 in Γ_T^M we have proved

$$\|\Phi(g_1) - \Phi(g_2)\|_{L_{[0,T],x,\omega}^2} \leq \frac{1}{2} \|g_1 - g_2\|_{L_{[0,T],x,\omega}^2},$$

which implies that Φ is a contraction on Γ_T^M .

We thus proved the local existence and uniqueness of a solution f to the nonlinear kinetic equation (2.3) in $L_{[0,T_{\max}),x,\omega}^2$ with $T_{\max} \geq T_1$. The solution f belongs to $L^\infty([0, T_{\max}), L_{x,\omega}^p) \cap L^2([0, T_{\max}), L_x^2 H_\omega^1)$ due to Proposition 3.2 since $\sigma(f)$ and $\nu(f)$ are well-defined and satisfy the required assumptions, as we saw above. At last, thanks to the global in time Cauchy theory for the linear equation given by Theorem 3.1 we see that if $\limsup_{t \rightarrow T_{\max}^-} \sup_{\Omega \times \mathbb{S}^{d-1}} \frac{\nu(f)}{|\mathbf{J}[f]|_\alpha} < +\infty$, then we can apply our fixed point argument starting at T_{\max} and therefore T_{\max} must be $+\infty$. Which concludes the proof of Theorem 2.3. \square

3.3. Global existence and decay of the free energy. The present section focuses on some important cases where one is ensured to have globally defined solution to the fully non-linear kinetic equation (2.3). Then we get a look into the free energy and its energy dissipation.

3.3.1. *Important examples of global solutions.* Theorem 2.3 offers a direct and easy way to check if the solutions obtained are globally defined. Indeed, under the assumptions (H1) – (H2) – (H3), if one shows that $\inf_{\Omega \times \mathbb{S}^{d-1}} |\mathbf{J}[f]|_\alpha$ cannot vanish along the flow then indeed $\sup_{\Omega \times \mathbb{S}^{d-1}} \frac{|\nu(f)|}{|\mathbf{J}[f]|_\alpha}$ cannot explode in finite time.

The non-normalised case: $\alpha \neq 0$. The discussion above implies that when $\alpha \neq 0$ then $|\mathbf{J}[f]|_\alpha \geq \frac{1}{\alpha} > 0$ for any function f so that $\Psi[f]$ is well-defined. Therefore for any $\alpha \neq 0$ the solutions constructed in Theorem 2.3 are global in time. This is the case of the equations looked at in [39, 23, 29] among others.

The non-asymptotic normalised case: $\frac{\nu(f)}{|\mathbf{J}[f]|}$ prevents any explosion. Here we are interested in the normalised framework $\alpha = 0$ but when the coefficient $\nu(f)$ compensate a possible vanishing of $|\mathbf{J}[f]|$. Namely if we add the assumption that $f \mapsto \frac{\nu(f)}{|\mathbf{J}[f]|}$ is a bounded function when $|\mathbf{J}[f]|$ tends to zero. Again, in this particular case the solutions constructed in Theorem 2.3 are global in time. This is the case of equations looked at in [10, 13] where $\nu(f) = \nu(|\mathbf{J}_f|)$ with $\frac{\nu(y)}{y}$ bounded and is generalised here for instance to any $\nu(f) = \nu(|\mathbf{J}[f]|)$ with $\frac{\nu(y)}{y}$ bounded near zero.

The normalised case of kernel with one coordinate with a strict sign on a closed subset. In this paragraph we assume that the kernel $\mathbf{K}(t, x, x_*, \omega, \omega_*)$ is such that one of its coordinate K_i has a strict sign in the following sense: there exists a closed set $\Gamma \subset \Omega \times \mathbb{S}^{d-1}$ - note that $\Gamma = \Omega \times \mathbb{S}^{d-1}$ works as well - and a positive constant $k_i > 0$ such that the two following properties hold.

- (i) for all (t, x, ω) in $\mathbb{R}^+ \times \Omega \times \mathbb{S}^{d-1}$, $K_i(t, x, \cdot, \omega, \cdot)|_{\Gamma^c} = 0$ (this assumption being void when $\Gamma = \Omega \times \mathbb{S}^{d-1}$);
- (ii) either $\inf_{\mathbb{R}^+ \times \Omega \times \mathbb{S}^{d-1}} \inf_{\Gamma} K_i(t, x, \cdot, \omega, \cdot) \geq k_i$ or $\sup_{\mathbb{R}^+ \times \Omega \times \mathbb{S}^{d-1}} \sup_{\Gamma} K_i(t, x, \cdot, \omega, \cdot) \leq -k_i$.

Recalling the notations of Proposition 3.5 we shall show that such a property implies that $J_i(t, x, \omega) = \int_{\Omega \times \mathbb{S}^{d-1}} f(t, x_*, \omega_*) K_i(t, x, x_*, \omega, \omega_*) dx_* d\omega_*$ cannot vanish in finite time if it is non-zero at initial time. Note that this is trivial if Γ is the whole set since then, thanks to the positivity of f , either

$$\int_{\Omega \times \mathbb{S}^{d-1}} f K_i dx_* d\omega_* \geq k_i M_0 \quad \text{or} \quad \int_{\Omega \times \mathbb{S}^{d-1}} f K_i dx_* d\omega_* \leq -k_i M_0.$$

For a general closed set Γ we consider the positive case, the negative one being dealt in the same manner. Recalling the equality obtained in the proof of Proposition 3.5:

$$(3.20) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} |J_i(t, x, \omega)|^2 &= J_i \int_{\Omega \times \mathbb{S}^{d-1}} f \left[\frac{\partial K_i}{\partial t} + c\omega \cdot \nabla_{x_*} K_i + \nabla_{\omega_*} \cdot (\sigma \nabla_{\omega_*} K_i) - \nu \mathbf{P}_{\omega_*^\perp}[\Psi] \cdot \nabla_{\omega_*} K_i \right] \\ &= J_i \int_{\Gamma} f \left[\frac{\partial K_i}{\partial t} + c\omega \cdot \nabla_{x_*} K_i + \nabla_{\omega_*} \cdot (\sigma \nabla_{\omega_*} K_i) - \nu \mathbf{P}_{\omega_*^\perp}[\Psi] \cdot \nabla_{\omega_*} K_i \right] \end{aligned}$$

As we saw in Section 3.2 the integrand is uniformly bounded from above as long as $t < T_{\max}$ by $K_{i,\infty}$ defined by (3.10). As \widetilde{K}_i is always positive it follows that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |J_i(t, x, \omega)|^2 &\leq -K_{i,\infty} J_i(t, x, \omega) \int_{\Gamma} f(t, x_*, \omega_*) dx_* d\omega_* \\ &\leq -\frac{K_{i,\infty}}{k_i} J_i(t, x, \omega) \int_{\Gamma} f(t, x_*, \omega_*) K_i dx_* d\omega_* \\ &\leq -\frac{K_{i,\infty}}{k_i} |J_i(t, x, \omega)|^2. \end{aligned}$$

We infer thanks to Grönwall lemma that

$$\forall (t, x, \omega) \in [0, T_{\max}) \times \Omega, \times \mathbb{S}^{d-1}, \quad |J_i(t, x, \omega)|^2 \geq |J_i(0, x, \omega)|^2 e^{-\frac{K_{i,\infty}}{k_i} t} > 0.$$

If $J_i(0, x, \omega) \neq 0$ then $J_i(t, x, \omega)$ cannot vanish in finite time and again, in this particular case the solutions constructed in Theorem 2.3 are global in time.

The Vicsek kernel operator. We study the special case when $\mathbf{K}(t, x, x_*, \omega, \omega_*) = \omega_*$ and normalised $\alpha = 0$, as in [10, 13, 30, 27] for the spatially homogeneous equation. Note that here we also obtain the global in time for the non-spatially homogeneous setting. We also assume that ν and σ do not depend on x nor ω and that the friction ν is negative (standard in all previous studies).

Contrary to the setting above, the present kernel can degenerate in the sense that one non-zero coordinate of \mathbf{K} can possibly vanish later on but giving rise to another non-zero coordinate. As a consequence, the full $|\mathbf{K}|$ will not vanish in finite time. First we explicitly write

$$J_i(t, x, \omega) = J_i(t) = \int_{\Omega \times \mathbb{S}^{d-1}} \omega_{i*} f(t, x_*, \omega_*) dx_* d\omega_* \quad \text{and} \quad \Psi = \frac{\mathbf{J}}{|\mathbf{J}|}.$$

As before let us consider the computation below for $0 \leq t < T_{\max}$ and recall (3.20) which in the present setting yields

$$\frac{1}{2} \frac{d}{dt} |\mathbf{J}|^2 = \sum_{i=1}^d J_i(t) \int_{\Omega \times \mathbb{S}^{d-1}} f [\sigma(f)(t) \Delta_{\omega_*}(\omega_{*i}) - \nu(f)(t) \mathbf{P}_{\omega_*^\perp}[\Psi] \cdot \nabla_{\omega_*}(\omega_{*i})].$$

Thanks to the properties of the tangential derivatives we can compute directly with (3.2)

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |\mathbf{J}(t)|^2 &= -(d-1)\sigma(f)(t) |\mathbf{J}(t)|^2 - \nu(f)(t) \int_{\Omega \times \mathbb{S}^{d-1}} \frac{1}{|\mathbf{J}(t)|} \mathbf{P}_{\omega_*^\perp}[\mathbf{J}(t)] \cdot \nabla_{\omega_*}(\omega_* \cdot \mathbf{J}(t)) \\ &= -(d-1)\sigma(f)(t) |\mathbf{J}(t)|^2 - \frac{\nu(f)(t)}{|\mathbf{J}(t)|} \int_{\Omega \times \mathbb{S}^{d-1}} |\nabla_{\omega_*}(\omega_* \cdot \mathbf{J}(t))|^2 \\ &\geq -(d-1)\sigma_0 |\mathbf{J}(t)|^2. \end{aligned}$$

Note that we use the fact that the friction is negative. Using Grönwall inequality we conclude that $|\mathbf{J}(t)|^2 \geq |\mathbf{J}(0)|^2 e^{-2(d-1)\sigma_0 t}$.

These computations were already done in an homogeneous regularised setting in [27] and our local theory allows us to carry them out directly. Such a lower bound thus imply the non-vanishing of $|\mathbf{J}(t)|$ in finite time and thus leading again to global in time solution in this setting.

3.3.2. *Decay of the free energy.* We recall our definitions for the free energy and the energy dissipation (2.9) – (2.10)

$$\begin{aligned}\mathcal{F}[f](t) &= \int_{\Omega \times \mathbb{S}^{d-1}} f \ln(f) dx d\omega \\ &\quad + \int_0^t \int_{\Omega \times \mathbb{S}^{d-1}} f \left[\nabla_\omega \cdot (\nu(f) \mathbf{P}_{\omega^\perp}[\Psi[f]]) - \frac{\nu(f)^2}{\sigma(f)} |\mathbf{P}_{\omega^\perp}[\Psi[f]]|^2 \right] dx d\omega ds \\ \mathcal{D}[f](t) &= \int_{\Omega \times \mathbb{S}^{d-1}} \sigma(f) f \left| \nabla_\omega \ln(f) + \frac{\nu(f)}{\sigma(f)} \mathbf{P}_{\omega^\perp}[\Psi[f]] \right|^2 dx d\omega.\end{aligned}$$

Let f_0 , σ , ν and \mathbf{K} be as in Theorem 2.3. We now establish the decay of $\mathcal{F}[f](t)$:

$$(3.21) \quad \forall t \geq 0, \quad \frac{d}{dt} \mathcal{F}[f](t) = -\mathcal{D}[f](t).$$

It comes from direct computations. Indeed since $\nabla_{x/\omega} [1 + \ln f] = \frac{\nabla_{x/\omega} f}{f}$ it follows by the integrating by parts formula (3.1),

$$\begin{aligned}\frac{d}{dt} \int_{\Omega \times \mathbb{S}^{d-1}} f \ln f(t, x, \omega) dx d\omega &= \int_{\Omega \times \mathbb{R}^d} [1 + \ln f] \partial_t f(t, x, \omega) dx d\omega \\ &= \int_{\Omega \times \mathbb{S}^{d-1}} c\omega \cdot \nabla_x f dx d\omega \\ &\quad + \int_{\Omega \times \mathbb{S}^{d-1}} [\sigma(f) \nabla_\omega f + \nu(f) f \mathbf{P}_{\omega^\perp}[\Psi[f]]] \cdot \frac{\nabla_\omega f}{f} dx d\omega \\ &= - \int_{\Omega \times \mathbb{S}^{d-1}} \left[\sigma(f) \frac{|\nabla_\omega f|^2}{f} + \nu(f) \mathbf{P}_{\omega^\perp}[\Psi[f]] \cdot \nabla_\omega f \right] dx d\omega.\end{aligned}$$

Hence,

$$\begin{aligned}\frac{d}{dt} \mathcal{F}[f](t) &= - \int_{\Omega \times \mathbb{S}^{d-1}} \left[\sigma(f) \frac{|\nabla_\omega f|^2}{f} + 2\nu(f) \mathbf{P}_{\omega^\perp}[\Psi[f]] \cdot \nabla_\omega f + \frac{\nu(f)^2}{\sigma(f)} |\mathbf{P}_{\omega^\perp}[\Psi[f]]|^2 f \right] dx d\omega \\ &= -\mathcal{D}[f](t)\end{aligned}$$

as claimed above.

4. MEAN-FIELD LIMIT: PROOFS OF THE RESULTS IN SECTION 2.3

Let us first describe the strategy we are about to implement. At the core, we will use Sznitman approach [43] (which differs from classical BBGKY approaches [33]):

- First, we will prove theorems 2.9 and 2.10 for modified particle systems and kinetic equation where the functional ψ is substituted by a functional τ_0 which is Lipschitz and bounded and takes the same values of ψ whenever $|\mathbf{J}| \geq c_*/2 > 0$. To prove theorems 2.9 and 2.10 for these modified ‘non-singular’ equations, we will follow the same steps in [3] and the results in [5]. This first step corresponds to Proposition 4.3.
- Second, we will prove that there exists a time $T > 0$ such that, for N large enough, it holds that $|\mathbf{J}(\mu_t^N)| \geq c_*/2$ for $t \in [0, T)$. In this time scale, the solutions of the ‘non-singular’ equations are also solutions of the original equation, and therefore, we will have concluded the results.

4.1. Preliminaries: functional framework and notations. In this section we will work in the space \mathcal{H} defined in Section 2.1 and with the 2-Wasserstein distance also defined there. We will need some results on Wasserstein distances in the sequel, that we summarise next (these results and proofs can be found in [45]). In our setting, the Wasserstein distance of order 1 is given by

$$W_1(m, p) := \inf \left\{ \left[\int_{(\Omega \times \mathbb{R}^d)^2} |z - u| \pi(dz, du) \right]; \right. \\ \left. \pi \in \mathcal{P}(\Omega \times \mathbb{R}^d)^2 \text{ with marginals } m \text{ and } p \right\}.$$

This distance can be expressed in duality form using the Kantorovich-Rubinstein distance

$$(4.1) \quad W_1(m, p) = \sup_{\|\varphi\|_{Lip} \leq 1} \left\{ \int_{\Omega \times \mathbb{R}^d} \varphi(z) dm - \int_{\Omega \times \mathbb{R}^d} \varphi(z) dp \right\}.$$

Also, by Hölder's inequality, it holds

$$(4.2) \quad W_1 \leq W_2.$$

4.2. The non-singular dynamics. To prove Theorem 2.9 and 2.10, we will consider first modified versions of Systems (2.13) and (2.15) where the potential singularity at $|\mathbf{J}| = 0$ is avoided. However, the solution of these 'non-singular' systems will coincide with solutions of the original systems as long as the flux $|\mathbf{J}| \neq 0$. Being more precise, we construct $\gamma : \mathbb{R}^d \rightarrow \mathbb{R}^d$ Lipschitz and bounded with

$$(4.3) \quad \gamma(v) = v, \quad \text{when } |v| \leq 2$$

and we define a functional $\tau_0 : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that τ_0 is Lipschitz and bounded in \mathcal{H} whenever $|\mathbf{J}(x, \gamma(v), m)| \leq c_*/2$ and

$$(4.4) \quad \tau_0(x, v, m) = \frac{\mathbf{J}(x, \gamma(v), m)}{|\mathbf{J}(x, \gamma(v), m)|_\alpha}, \quad |\mathbf{J}(x, \gamma(v), m)| \geq c_*/2,$$

where $c_* > 0$ can be considered as small as needed. It turns out that τ_0 is Lipschitz and bounded in the whole \mathcal{H} :

Lemma 4.1. *The functions $r = r(x, v, m) := |\mathbf{J}|(x, \gamma(v), m) : \mathcal{H} \rightarrow \mathbb{R}_+$ and $\tau_0 = \tau_0(x, v, m) : \mathcal{H} \rightarrow \mathbb{R}_+$ are Lipschitz and bounded. Moreover, there exists a constant $c > 0$ such that*

$$(4.5) \quad \left| |\mathbf{J}(x', \gamma(v'), m')| - |\mathbf{J}(x, \gamma(v), m)| \right| \leq c(|x' - x| + |v' - v| + W_1(m', m)),$$

for all $(x, v, m), (x', v', m') \in \mathcal{H}$.

Proof of Lemma 4.1. We notice, first, that since γ is bounded, then $K(x, \gamma(v), m)$ is bounded in all the variables and, therefore, $r_f(z)$ is bounded in \mathcal{H} . Now, let $f, g \in \mathcal{P}_2(\Omega \times \mathbb{R}^d)$ and fix $z \in \Omega \times \mathbb{R}^d$, we show next that $r_f := r(z, f)$ is Lipschitz in f . Consider $z, z' \in \Omega \times \mathbb{R}^d$ and for $z = (x, v)$, denote $\tilde{z} = (x, \gamma(v))$. Now, we

consider (the integrals are in $\Omega \times \mathbb{R}^d$)

$$\begin{aligned}
|r_f(z) - r_g(z')| &= \left| \left| \int K(\tilde{z}, y) f(dy) \right| - \left| \int K(\tilde{z}', y) g(dy) \right| \right| \\
&\leq \left| \int [K(\tilde{z}, y) f(dy) - K(\tilde{z}', y) g(dy)] \right| \\
&\leq \|K\|_{Lip} |\tilde{z} - \tilde{z}'| + \left| \int K(\tilde{z}', y) [f(dy) - g(dy)] \right| \\
&\leq \|K\|_{Lip} (\|\gamma\|_{Lip} + 1) |z - z'| \\
&\quad + \|K\|_{Lip} \left| \sup_{\|\varphi\|_{Lip} \leq 1} \left(\int \varphi(y) f(dy) - \int \varphi(y) g(dy) \right) \right| \\
&= \|K\|_{Lip} (\|\gamma\|_{Lip} + 1) |z - z'| + \|K\|_{Lip} W_1(f, g) \\
&\leq \|K\|_{Lip} (\|\gamma\|_{Lip} + 1) |z - z'| + \|K\|_{Lip} W_2(f, g)
\end{aligned}$$

where in the second line we used the reverse triangle inequality; in the third line we have used that $\int_{\mathbb{R}^{2d}} f = 1$ and that K is Lipschitz; in the fourth line we used that γ is Lipschitz and choose an arbitrary $z_0 \in \mathbb{R}^{2d}$; the fifth line is given by (4.1); and the last inequality follows from (4.2). Therefore, we conclude that $r = r_f(z)$ is Lipschitz in \mathcal{H} .

Next, we also show that τ_0 is Lipschitz and bounded in \mathcal{H} . First, whenever $r \geq c_*/2$ we have that $(\alpha + (1-\alpha)r)^{-1}$ is also Lipschitz and bounded for all $\alpha \in [0, 1]$, therefore, τ_0 is product of Lipschitz and bounded functions, hence it is Lipschitz and bounded. \square

With the function τ_0 we define the *non-singular particle dynamics* as:

$$\begin{aligned}
(4.6a) \quad & \begin{cases} dX_t^{i,N} = V_t^{i,N} dt, \\ dV_t^{i,N} = \nu(\mu^N) \mathbf{P}_{(V^{i,N})^\perp}(\tau_0(X^{i,N}, V^{i,N}, \mu^N)) dt \\ \quad + \frac{1}{2} \mathbf{P}_{(\bar{V}^{i,N})^\perp}[(\nabla_v \sigma(\mu^N))(\bar{X}^{i,N}, \bar{V}^{i,N})] dt \\ (4.6b) \quad \quad \quad + \sqrt{2\sigma(\mu^N)} \mathbf{P}_{(V^{i,N})^\perp} \circ dB_t^i, \end{cases}
\end{aligned}$$

(this is exactly System (2.13) substituting the function Ψ by τ_0) and the *non-singular auxiliary process* is given by:

$$\begin{aligned}
(4.7a) \quad & \begin{cases} d\bar{X}_t^{i,N} = \bar{V}_t^{i,N} dt, \\ d\bar{V}_t^{i,N} = \nu(f) \mathbf{P}_{(\bar{V}^{i,N})^\perp}(\tau_0(\bar{X}^{i,N}, \bar{V}^{i,N}, f)) dt \\ \quad + \frac{1}{2} \mathbf{P}_{(\bar{V}^{i,N})^\perp}[(\nabla_v \sigma(f))(\bar{X}^{i,N}, \bar{V}^{i,N})] dt \\ (4.7b) \quad \quad \quad + \sqrt{2\sigma(f)} \mathbf{P}_{(\bar{V}^{i,N})^\perp} \circ dB_t^i, \\ (4.7c) \quad \quad \quad f_t = \text{law}(\bar{X}^{i,N}, \bar{V}^{i,N}), \end{cases}
\end{aligned}$$

(again this is exactly System (2.15) substituting the function Ψ by τ_0). Analogously, we also consider the *non-singular kinetic equation* in $\Omega \times \mathbb{S}^{d-1}$ given by

$$(4.8) \quad \partial_t f + \omega \cdot \nabla_x f = \nabla_\omega \cdot (\sigma(f) \nabla_\omega f) + \nabla_\omega \cdot (\nu(f) f \nabla_\omega \tau_0(f)).$$

Note that under the assumptions of Theorem 2.3, (4.8) has global existence and uniqueness of solutions in the spaces stated in Theorem 2.3.

Remark 4.2. Notice that we are assuming that ν , σ and $\nabla_v \sigma$ are bounded and Lipschitz for all $v \in \mathbb{R}^d$ rather than $v \in \mathbb{S}^{d-1}$ (see hypothesis (H4)). However, we just need that ν , σ and $\nabla_v \sigma$ to be Lipschitz and bounded in a neighbourhood of $|v| = 1$, as we can use regularising arguments as the one done to define the functional τ_0 .

Proposition 4.3. Theorems 2.9 and 2.10 hold replacing in the statements System (2.13) by System (4.6); System (2.15) by System (4.7); and the kinetic equation (2.3) by the kinetic equation (4.8). However the existence results are global in time and for any N . Theorem 2.10 holds for all $T > 0$.

We prove this Proposition in Section 4.4.

4.3. Proof of Theorem 2.9 and Theorem 2.10 assuming Proposition 4.3.

Assuming that Proposition 4.3 holds true, we can prove the main results. For this we will use the following:

Lemma 4.4. For each $t \in [0, T]$ let $(Z_N(t))_{N \geq 0}$ be a sequence of random variables such that

$$(4.9) \quad \sup_{t \in [0, T]} \mathbb{E}[|Z_N(t) - Z(t)|] \rightarrow 0, \quad \text{as } N \rightarrow \infty,$$

where $Z(t) \geq c_*$ for some $c_* > 0$ and for all $t \in [0, T]$. Then for all $\varepsilon > 0$ there exists $N_0(\varepsilon) > 0$ such that for all $t \in [0, T]$ and for all $N \geq N_0(\varepsilon)$ it holds that

$$Z_N \geq c - \varepsilon.$$

Proof of Lemma 4.4. The statement is proved by contradiction: suppose that there exists $\varepsilon > 0$ such that for all $k \geq 0$ there exist $t_k \in [0, T]$ and $N_k \geq k$ such that $Z_{N_k}(t_k) < c - \varepsilon$. Consider h a Lipschitz bounded function strictly increasing (e.g., the hyperbolic tangent), then, by assumption we have that

$$\mathbb{E}[h(Z(t_k))] - \mathbb{E}[h(Z_{N_k}(t_k))] \geq h(c) - h(c - \varepsilon) > 0,$$

which is strictly positive since h is strictly increasing by assumption. On the other hand we have that

$$\begin{aligned} 0 < h(c) - h(c - \varepsilon) &\leq \mathbb{E}[h(Z(t_k))] - \mathbb{E}[h(Z_{N_k}(t_k))] \\ &= \mathbb{E}[h(Z(t_k)) - h(Z_{N_k}(t_k))] \\ &\leq \|h\|_{Lip} \sup_{t \in [0, T]} \mathbb{E}[|Z(t) - Z_{N_k}(t)|] \rightarrow 0 \quad \text{as } k \rightarrow \infty, \end{aligned}$$

where we have used that h is Lipschitz and assumption (4.9). Therefore we reached a contradiction. \square

Lemma 4.5 (Convergence of the flux). *With the notations of Proposition 4.3 and assuming the latter holds then*

$$\sup_{t \in [0, T]} \mathbb{E}[|\mathbf{J}(X_t^{i,N}, V_t^{i,N}, \mu_t^N) - \mathbf{J}(\bar{X}_t, \bar{V}_t, f_t)|] \rightarrow 0 \quad \text{as } N \rightarrow \infty,$$

weakly (in distribution); where $(X^{i,N}, V^{i,N})$ is the solution to the non-singular System (4.6), μ^N the associated empirical distribution and f_t is the law of the process given by the auxiliary non-singular system (4.7).

Proof of Lemma 4.5. A computation shows (see proof of Lemma 4.1) that

$$(4.10) \quad \begin{aligned} \left| |\mathbf{J}(X_t^{i,N}, V_t^{i,N}, \mu_t^N)| - |\mathbf{J}(\bar{X}_t, \bar{V}_t, f_t)| \right| &\leq c \left(|X_t^{i,N} - \bar{X}_t^i| + |V_t^{i,N} - \bar{V}_t^i| \right. \\ &\quad \left. + \left| \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) [\mu_t^N(dy) - f_t(dy)] \right| \right), \end{aligned}$$

for some constant $c > 0$ and where we denoted $\bar{Z}_t^i = (\bar{X}_t^i, \bar{V}_t^i)$. Taking expectations and the supremum we have that

$$\begin{aligned} &\sup_{t \in [0, T]} \mathbb{E} \left[\left| |\mathbf{J}(X_t^{i,N}, V_t^{i,N}, \mu_t^N)| - |\mathbf{J}(\bar{X}_t, \bar{V}_t, f_t)| \right| \right] \\ &\leq c \left(\mathbb{E} \left[\sup_{t \in [0, T]} |X_t^{i,N} - \bar{X}_t|^2 + |V_t^{i,N} - \bar{V}_t|^2 \right] + \sup_{t \in [0, T]} \mathbb{E} \left[\left| \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) [\mu_t^N(dy) - f_t(dy)] \right|^2 \right] \right), \end{aligned}$$

where we have used Cauchy-Schwarz and Jensen's inequality. We have that

$$\begin{aligned} &\mathbb{E} \left[\left| \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) [\mu_t^N(dy) - f_t(dy)] \right|^2 \right] \\ &\leq \mathbb{E} \left[\left| \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) [\mu_t^N(dy) - \bar{\mu}_t^N(dy)] \right|^2 \right] + \mathbb{E} \left[\left| \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) [\bar{\mu}_t^N(dy) - f_t(dy)] \right|^2 \right] \\ &\leq \mathbb{E} \left[\frac{1}{N} \sum_{j=1}^N |K(\bar{Z}_t^i, Z_t^{j,N}) - K(\bar{Z}_t^i, \bar{Z}_t^j)|^2 \right] + \mathbb{E} \left[\left| \frac{1}{N} \sum_{j=1}^N K(\bar{Z}_t^i, \bar{Z}_t^j) - \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) f_t(dy) \right|^2 \right] \\ &\leq \|K\|_{Lip} \mathbb{E} \left[\frac{1}{N} \sum_{j=1}^N |Z_t^{j,N} - \bar{Z}_t^j|^2 \right] + \frac{C(T)}{N}, \end{aligned}$$

where $\bar{\mu}^N$ is the empirical distribution associated to the process $(\bar{Z}_t^1, \dots, \bar{Z}_t^N)$; the final bound is consequence of K being Lipschitz and a classical computation due to Sznitmann [43], where $C(T)$ is a finite constant that depends on T . Using this bound and the convergence of the particles (2.16) we have that

$$\sup_{t \in [0, T]} \mathbb{E} \left[\left| \int_{\mathbb{R}^{2d}} K(\bar{Z}_t^i, y) [\mu_t^N(dy) - f_t(dy)] \right|^2 \right] \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Applying this limit on expression (4.10) and again (2.16), we conclude the result. \square

Proof of theorems 2.9 and 2.10 (assuming that Proposition 4.3 holds). Let f_t be solution to our original kinetic equation (2.3), by Theorem 2.3 we know that this solution exists at least for short times and that there exists $c_* > 0$ and a time $T = T(c_*) > 0$ such that

$$|\mathbf{J}(t, x, \omega, f)| \geq c_* > 0.$$

Therefore, for $t \leq T(c_*)$ the non-singular-kinetic equation (4.8) coincides with the original kinetic equation (2.3). At the same time, by Lemmas 4.5 and 4.4, we have that for N large enough it holds that

$$|\mathbf{J}(t, X_t^{i,N}, V_t^{i,N}, \mu_t^N)| \geq c_*/2.$$

Therefore the original particle dynamics (2.13) coincide with the ones of the non-singular system (4.6). Consequently, for $t < T(c_*)$ and N large enough we have

that Theorems 2.9 and 2.10 hold for the particle dynamics (2.13) and the kinetic equation (2.3). \square

4.4. Proof of Proposition 4.3. The proof of Proposition 4.3 follows closely the methodology in [3] and [5] which are based on Sznitman approach [43]. The main difference is that the interaction rate ν and the noise coefficient σ are considered to be constant in [3]. In particular, this gives rise to an extra term in the equations (coming from (2.13b)).

Step 1. Regularised version of the non-singular dynamics

The non-singular particle dynamics (4.6) written in Itô's convention corresponds to (see Section A.2 for more details):

$$(4.11a) \quad \left\{ \begin{array}{l} dX^{i,N} = V^{i,N} dt, \\ dV^{i,N} = \nu(\mu^N) \mathbf{P}_{(V^{i,N})^\perp}(\tau_0(X^{i,N}, V^{i,N}, \mu^N)) dt \\ \quad + \sqrt{2\sigma(\mu^N)} \mathbf{P}_{(V^{i,N})^\perp} dB_t^i \\ \quad + (d-1)\sigma(\mu^N) \frac{V^{i,N}}{|V^{i,N}|^2} dt. \end{array} \right.$$

And the Itô formulation for the non-singular auxiliary process (4.7) is given by

$$(4.12a) \quad \left\{ \begin{array}{l} d\bar{X}^{i,N} = \bar{V}^{i,N} dt, \\ d\bar{V}^{i,N} = \nu(f) \mathbf{P}_{(\bar{V}^{i,N})^\perp}(\tau_0(\bar{X}^{i,N}, \bar{V}^{i,N}, f)) dt \\ \quad + \mathbf{P}_{(\bar{V}^{i,N})^\perp}[(\nabla_v \sigma(\mu^N))(\bar{X}^{i,N}, \bar{V}^{i,N})] dt \\ \quad + \sqrt{2\sigma(f)} \mathbf{P}_{(\bar{V}^{i,N})^\perp} dB_t^i \\ \quad + (d-1)\sigma(f) \frac{\bar{V}^{i,N}}{|\bar{V}^{i,N}|^2} dt, \\ (4.12b) \quad f_t = \text{law}(\bar{X}^{i,N}, \bar{V}^{i,N}). \end{array} \right.$$

Remark 4.6. Notice that the solutions of (4.11) and (4.12) fulfil $|V_t|^2 = |V_0|^2$ in the velocities for all times where the solution is defined (this is shown as in Lemma 2.8).

Step 2. Existence and uniqueness for the regularised particle system - Proof of part (i) of Theorem 2.9 for (4.6).

We consider now a regularised version of Systems (4.11) and (4.12) using two functions τ_1 and τ_2 both Lipschitz and bounded and satisfying:

$$\begin{aligned} \tau_1(v) &= \mathbf{P}_{v^\perp} = \text{Id} - \frac{v \otimes v}{|v|^2}, & \text{if } |v| \geq 1/2, \\ \tau_2(v) &= \frac{v}{|v|^2}, & \text{if } |v| \geq 1/2. \end{aligned}$$

With these functions we defined the *regularised particle dynamics* as

$$(4.13a) \quad \left\{ \begin{array}{l} dX^{i,N} = V^{i,N} dt, \\ dV^{i,N} = \nu(\mu^N) \tau_1(V^{i,N})(\tau_0)(X^{i,N}, V^{i,N}, \mu^N) dt \\ \quad + \tau_1(V^{i,N})[(\nabla_v \sigma(\mu^N))(\bar{X}^{i,N}, \bar{V}^{i,N})] dt \\ \quad + \sqrt{2\sigma(\mu^N)} \tau_1(V^{i,N}) dB_t^i \\ \quad + (d-1)\sigma(\mu^N)\tau_2(V^{i,N})dt. \end{array} \right.$$

(4.13b)

Remark 4.7. Notice that the functions τ_0 , τ_1 and τ_2 are introduced to regularise the original system, in the sense that we obtain a new system where all the coefficients are Lipschitz and bounded in \mathcal{H} . This regularity allows us to apply classical results of existence of solutions and convergence, as we will see next. At the same time, when $|V| = 1$ and $|\mathbf{J}| \geq c_*/2$ we recover the original equations.

Firstly, we have by Lemma 4.1 that τ_0 is a Lipschitz bounded function and, moreover, all the coefficients are also Lipschitz bounded (using that the product of Lipschitz bounded functions is Lipschitz bounded, see also Remark 4.2). So we have existence and uniqueness of pathwise solutions for (4.13) (see [5, Theorem 1.2], which, for completeness we have added in a simplified form in Theorem A.2 in the Appendix.).

Secondly, one can check as in Lemma 2.8 that $|V_t^{i,N}| = |V_0^{i,N}| = 1$. Therefore, the solution to the regularised system (4.13) is also a solution to the non-singular system (4.6).

Step 3. Existence and uniqueness for an auxiliary regularised process of (2.15) - Proof of part (ii) of Theorem 2.9 for (4.7).

Similarly as before, we consider a regularised version of the non-singular auxiliary process (4.7) given by:

$$(4.14a) \quad \left\{ \begin{array}{l} d\bar{X}^{i,N} = \bar{V}^{i,N} dt, \\ d\bar{V}^{i,N} = \nu(f) \tau_1(\bar{V}^{i,N})(\tau_0)(\bar{X}^{i,N}, \bar{V}^{i,N}, f) dt \\ \quad + \tau_1(\bar{V}^{i,N})[(\nabla_v \sigma(\mu^N))(\bar{X}^{i,N}, \bar{V}^{i,N})] dt \\ \quad + \sqrt{2\sigma(f)} \tau_1(\bar{V}^{i,N}) dB_t^i \\ \quad + (d-1)\sigma(f)\tau_2(\bar{V}^{i,N})dt \\ f_t = \text{law}(\bar{X}^{i,N}, \bar{V}^{i,N}). \end{array} \right.$$

(4.14b)

(4.14c)

(4.14d)

Since (4.14) has bounded and Lipschitz coefficients in \mathcal{H} (this can be seen as in the previous proof), it admits a pathwise unique global solution (this statement is shown in [5, Theorem 1.7], which is a generalisation of Sznitmann's strategy [43, Theorem 1.1] - see Theorem A.3 in the Appendix). We can prove as in Lemma 2.8 that $d|\bar{V}_t|^2 = 0$ so $|\bar{V}_t| = 1$ for all times. Therefore, the solution to (4.14) is also solution of the non-singular auxiliary system (4.7).

Step 4. Existence and uniqueness for the non-singular kinetic equation (4.8) - Proof of part (iii) of Theorem 2.9 for the non-singular case: (4.7) and (4.8)

Next we show that part (iii) of Theorem 2.9 holds for the non-singular auxiliary process (4.7) and the non-singular kinetic equation (4.8).

We can apply Theorem 2.3 to the non-singular kinetic equation (4.8). Moreover, since ν is bounded and τ_0 is also bounded we have global-in-time existence of solutions for (4.8).

Now, we also know that the kinetic equation (2.3) has existence and uniqueness of weak solutions in $L^\infty([0, T_{max}), L^1_{x,\omega} \cap L^p_{x,\omega})$ for p in $[2, \infty]$. Let f_t be the weak solution of the kinetic equation for $t \in [0, T_{max})$. We also know that for $c_* > 0$ small enough there is a time $T_{c_*} \leq T_{max}$ such that $|\mathbf{J}_{f_t}| > c_*/2$ for all $t \in [0, T_{c_*})$. Therefore, for $t \in [0, T_{c_*})$ the solution f_t of the kinetic equation is also solution of the non-singular kinetic equation (4.8).

The fact that the Fokker-Planck equation for the non-singular auxiliary process (\bar{X}_t, \bar{V}_t) (4.8) corresponds to the non-singular kinetic equation (4.8) is proven in [3] (this is an application of Itô's formula, see for example [43, Remark 1.2], [5, Remark 1.8] and Eq. (A.3)). The proof that (\bar{X}_t, \bar{V}_t) has law f_t (local-in-time) is carried out in the same way as in [3].

Step 5. Mean-field limit for the non-singular dynamics - Proof of Theorem 2.10 for non-singular dynamics (4.6) and (4.7).

The proof of Theorem 2.10 is an application of Sznitman approach, generalised in [5].

We can apply the result in [5, Theorem 1.10] (a condensed version can be found in Theorem A.5 in the Appendix).

Remark 4.8. *The control in the particle position is immediate from the one in the velocities:*

$$|X_T^{i,N} - \bar{X}_T^{i,N}|^2 = \left| \int_0^T V_s^{i,N} ds - \int_0^T \bar{V}_s^{i,N} \right|^2 \leq T \sup_{0 \leq t \leq T} |V_t^{i,N} - \bar{V}_t^{i,N}|^2.$$

APPENDIX A. SOME KNOWN RESULTS ON SDE

For the sake of completeness, we introduce in this section known results for stochastic differential equations that we used in the main part of the document.

A.1. Results on existence of solutions and large-particle limits. The results and proofs from this section can be found in a more general form in [5]. The original approach to proving the large-particle limit is due to Sznitmann and can be found in [43].

General stochastic differential equations. Consider a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ and an $(\mathcal{F}_t)_{t \geq 0}$ -Brownian motion B taking values in \mathbb{R}^m . Consider the stochastic differential equation giving the evolution for $Z_t \in \mathbb{R}^p$, for $t \geq 0$:

$$(A.1) \quad dZ_t = b(Z_t)dt + a(Z_t)dB_t,$$

with initial data $Z(t=0) = Z_0 \in \mathbb{R}^p$, where the coefficients $a : \mathbb{R}^p \rightarrow \mathbb{R}^{p \times m}$ and $b : \mathbb{R}^p \rightarrow \mathbb{R}^p$ are Lipschitz and bounded. (This is the form taken by the SDE (4.13) with $p = 2d$ and $Z_t = (X_t, V_t)$.)

Definition A.1 (See Definition 1.1 in [5]). *An $(\mathcal{F}_t)_{t \geq 0}$ -adapted continuous process $(Z_t)_{0 \leq t \leq T}$ is a solution to (A.1) if*

$$Z_t = Z_0 + \int_0^t b(Z_s) ds + \int_0^t a(Z_s) dB_s, \quad 0 \leq t \leq T.$$

Theorem A.2 (Adapted from Theorem 1.2 [5]). *Let us assume that $Z_0 \in L^2$ is independent of B and that the coefficients b and a are Lipschitz and bounded. Then, there exists a unique solution of (A.1). Moreover, $Z_t \in L^2$ for all $t < T$, with T finite.*

Results for non-linear SDE.

We consider a nonlinear SDE of the form

$$(A.2) \quad dZ_t = b(Z_t, \mathcal{L}(Z_t))dt + a(Z_t, \mathcal{L}(Z_t))dB_t.$$

where $\mathcal{L}(Z)$ denotes the distribution or law of the random element Z . Here we also assume that $b : \mathbb{R}^p \times \mathcal{P}_2(\mathbb{R}^p) \rightarrow \mathbb{R}^p$ and $a : \mathbb{R}^p \times \mathcal{P}_2(\mathbb{R}^p) \rightarrow \mathbb{R}^{p \times m}$ are bounded and Lipschitz. Specifically, in this case to be Lipschitz means that for all $z, z' \in \mathbb{R}^d$ and all $\mu, \mu' \in \mathcal{P}(\mathbb{R}^d)$ it holds that

$$|(b, a)(z, \mu) - (b, a)(z', \mu')| \leq c(|z - z'| + W^{(2)}(\mu, \mu')),$$

for some constant c (see Section 4.1 for more details). System (A.2) is the form taken by the system auxiliary system (4.14).

Theorem A.3 (From Theorem 1.7 in [5]). *Let us assume that $Z_0 \in L^2$ is independent of B , and that the coefficients b and a are bounded and Lipschitz. Then, there exists a unique solution to (A.2).*

For $\varphi \in C_b^2(\mathbb{R}^p)$, Itô's formula applied to the process Z_t gives (see Remark 1.8 in [5]):

$$(A.3) \quad \begin{aligned} \varphi(Z_t) = & \varphi(Z_0) + \int_0^t \left[\frac{1}{2} \text{trace}[a(Z_s, \mathcal{L}(Z_s))^T a(Z_s, \mathcal{L}(Z_s))] D^2 \varphi(Z_s) \right. \\ & \left. + b(Z_s, \mathcal{L}(Z_s)) D \varphi(Z_s) \right] ds + \int_0^t D \varphi(Z_s) a(Z_s, \mathcal{L}(Z_s)) dB_s, \end{aligned}$$

where the exponent T denotes the transpose and D^2 denotes the Hessian matrix.

Lemma A.4 (Law of large numbers: Lemma 1.9 in [5]). *Let $\mu \in \mathcal{P}_2(\mathbb{R}^p)$, and let $(Z_i)_{i \in \mathbb{N}}$ be a sequence of independent identically distributed random variables with common law μ . For each $N \geq 1$ denote by μ^N the empirical distribution associated to the first N elements of the sequence, i.e.*

$$\mu^N(z) = \frac{1}{N} \sum_{i=1}^N \delta_{Z_i}(z).$$

Then, it holds that

$$\lim_{N \rightarrow \infty} \mathbb{E}[W_2(\mu^N, \mu)]^2 = 0.$$

Let $z_i \in \mathbb{R}^p$ and $z'_i \in \mathbb{R}^p$ for $i = 1, \dots, N$, it holds ([5, (1.24)])

$$(A.4) \quad W^{(2)} \left(\frac{1}{N} \sum_{i=1}^N \delta_{z_i}, \frac{1}{N} \sum_{i=1}^N \delta_{z'_i} \right) \leq \left(\frac{1}{N} \sum_{i=1}^N |z_i - z'_i|^2 \right)^{1/2}.$$

Particle approximations (from [5, Section 1.3.4]).

Theorem A.5 (Extracted and adapted from Section 1.3.4 in [5]). *Consider*

$$(A.5) \quad Z_t^i = Z_0^i + \int_0^t b(Z_s^i, \mathcal{L}(Z_s^i)) ds + \int_0^t a(Z_s^i, \mathcal{L}(Z_s^i)) dB_s^i,$$

and

$$(A.6) \quad Z_t^{i,N} = Z_0^i + \int_0^t b(Z_s^{i,N}, \mu_s^N) ds + \int_0^t a(Z_s^{i,N}, \mu_s^N) dB_s^i,$$

where μ_s^N is the empirical distribution of the N particles. Assume that a, b are bounded and Lipschitz in $\mathbb{R}^p \times \mathcal{P}_2(\mathbb{R}^p)$. Then, it holds that

$$(A.7) \quad \lim_{N \rightarrow \infty} \sup_{1 \leq i \leq N} \mathbb{E} \left[\sup_{0 \leq t \leq T} |Z_t^{i,N} - Z_t^i|^2 \right] = 0.$$

A.2. Stratonovich to Ito's convention. We use the results presented in [26] as well as [42, V.30, Theorem (30.14)]. Particularly, if Z_t is solution to the Stratonovich SDE

$$(A.8a) \quad \begin{cases} dZ = b(Z, t)dt + a(Z, t) \circ dB_t, \\ (A.8b) \quad Z(t=0) = Z_0, \end{cases}$$

where B is an m - dimensional Brownian motion, and $b : \mathbb{R}^p \times [0, T] \rightarrow \mathbb{R}^p$ and $a : \mathbb{R}^p \times [0, T] \rightarrow M^{p \times m}$ (the space of real matrices of dimension $p \times m$) and are such that there is existence and uniqueness of solutions for the SDE (A.8). Then, Z_t is solution to the Itô SDE:

$$(A.9a) \quad \begin{cases} dZ = [b(Z, t) + \frac{1}{2}c(Z, t)] dt + a(Z, t)dB_t, \\ (A.9b) \quad Z(t=0) = Z_0, \end{cases}$$

where

$$c_i(z, t) = \sum_{k=1}^m \sum_{j=1}^n \frac{\partial a_{ik}}{\partial z_j}(z, t) a_{jk}(z, t).$$

In our case $a : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d \times d}$ with

$$a(x, v) = \begin{pmatrix} 0_{d \times d} \\ \eta(x, v) \end{pmatrix},$$

where $0_{d \times d}$ is a $d \times d$ zero-matrix and

$$\eta(x, v) = \alpha(x, v)P_{v^\perp}; \quad \alpha(x, v) = \sqrt{2\sigma(f)(x, v)}.$$

With this we have that $c : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$ with $c_i = 0$ for $i = 1, \dots, d$ since $\gamma_{jk} = 0$ for all $j = 1, \dots, d$ and any $k = 1, \dots, d$. Now, for $i = d+1, \dots, 2d$ we have that

$$c_i(x, v) = \sum_{k=1}^d \sum_{j=1}^d \frac{\partial \eta_{ik}}{\partial v_j}(x, v) \eta_{jk}(x, v).$$

Using that

$$\eta_{jk}(x, v) = \alpha(x, v) \left(\delta_{j=k} - \frac{v_j v_k}{|v|^2} \right),$$

we compute the previous expression and obtain that

$$c(x, v) = \left(0_{1 \times d}, 2(d-1)\sigma(f)(x, v) \frac{v}{|v|^2} + P_{v^\perp}[(\nabla_v \sigma(f))(x, v)] \right).$$

REFERENCES

- [1] BENDALI, A., AND TORDEUX, S. Extension of the Günter derivatives to Lipschitz domains and application to the boundary potentials of elastic waves. working paper or preprint, Nov. 2016.
- [2] BOLLEY, F., CAÑIZO, J. A., AND CARRILLO, J. A. Stochastic mean-field limit: Non-lipschitz forces and swarming. *Mathematical Models and Methods in Applied Sciences* 21, 11 (2011), 2179–2210.
- [3] BOLLEY, F., CAÑIZO, J. A., AND CARRILLO, J. A. Mean-field limit for the stochastic vicsek model. *Applied Mathematics Letters* 25, 3 (2012), 339 – 343.
- [4] BRIANT, M., MEUNIER, N., AND NAVORET, L. work in progress.
- [5] CARMONA, R. *Lectures on BSDEs, stochastic control, and stochastic differential games with financial applications*, vol. 1. SIAM, 2016.
- [6] DAVID, I., KOHNKE, P., LAGRIFFOUL, G., PRAUD, O., PLOUARBOUÉ, F., DEGOND, P., AND DRUART, X. Mass sperm motility is associated with fertility in sheep. *Animal reproduction science* 161 (2015), 75–81.
- [7] DEGOND, P. Global existence of smooth solutions for the vlasov-fokker-planck equation in 1 and 2 space dimensions. *Annales scientifiques de l'école Normale Supérieure* 19, 4 (1986), 519–542.
- [8] DEGOND, P., DIEZ, A., FROUVELLE, A., AND ACEITUNO, S. M. Phase transitions and macroscopic limits in a bgk model of body-attitude coordination. *arXiv preprint arXiv:1905.04885* (2019).
- [9] DEGOND, P., DIMARCO, G., AND MAC, T. B. N. Hydrodynamics of the kuramoto–vicsek model of rotating self-propelled particles. *Mathematical Models and Methods in Applied Sciences* 24, 02 (2014), 277–325.
- [10] DEGOND, P., FROUVELLE, A., AND LIU, J.-G. A note on phase transitions for the Smoluchowski equation with dipolar potential. In *Hyperbolic Problems: Theory, Numerics, Applications* (Padova, Italy, June 2012), vol. 8 of *Applied Mathematics*, AIMS, pp. 179–192.
- [11] DEGOND, P., FROUVELLE, A., AND LIU, J.-G. Macroscopic limits and phase transition in a system of self-propelled particles. *Journal of nonlinear science* 23, 3 (2013), 427–456.
- [12] DEGOND, P., FROUVELLE, A., AND LIU, J.-G. Macroscopic limits and phase transition in a system of self-propelled particles. *Journal of nonlinear science* 23, 3 (2013), 427–456.
- [13] DEGOND, P., FROUVELLE, A., AND LIU, J.-G. Phase transitions, hysteresis, and hyperbolicity for self-organized alignment dynamics. *Archive for Rational Mechanics and Analysis* 216, 1 (2015), 63–115.
- [14] DEGOND, P., FROUVELLE, A., MERINO-ACEITUNO, S., AND TRESCASES, A. Alignment of self-propelled rigid bodies: from particle systems to macroscopic equations. In *International workshop on Stochastic Dynamics out of Equilibrium* (2017), Springer, pp. 28–66.
- [15] DEGOND, P., FROUVELLE, A., MERINO-ACEITUNO, S., AND TRESCASES, A. Quaternions in collective dynamics. *Multiscale Modeling & Simulation* 16, 1 (2018), 28–77.
- [16] DEGOND, P., MANHART, A., AND YU, H. A continuum model for nematic alignment of self-propelled particles. *Discrete & Continuous Dynamical Systems-B* 22, 4 (2017), 1295–1327.
- [17] DEGOND, P., MANHART, A., AND YU, H. An age-structured continuum model for myxobacteria. *Mathematical Models and Methods in Applied Sciences* 28, 09 (2018), 1737–1770.
- [18] DEGOND, P., AND MERINO-ACEITUNO, S. Nematic alignment of self-propelled particles in the macroscopic regime. *arXiv preprint arXiv:1910.02865* (2019).
- [19] DEGOND, P., MERINO-ACEITUNO, S., VERGNET, F., AND YU, H. Coupled self-organized hydrodynamics and stokes models for suspensions of active particles. *Journal of Mathematical Fluid Mechanics* 21, 1 (2019), 6.
- [20] DEGOND, P., AND MOTSCH, S. Continuum limit of self-driven particles with orientation interaction. *Mathematical Models and Methods in Applied Sciences* 18, supp01 (2008), 1193–1215.
- [21] DEGOND, P., AND MOTSCH, S. A macroscopic model for a system of swarming agents using curvature control. *Journal of Statistical Physics* 143, 4 (2011), 685–714.

- [22] DEGOND, P., AND NAVORET, L. A multi-layer model for self-propelled disks interacting through alignment and volume exclusion. *Mathematical Models and Methods in Applied Sciences* 25, 13 (2015), 2439–2475.
- [23] DOI, M. Molecular dynamics and rheological properties of concentrated solutions of rodlike polymers in isotropic and liquid crystalline phases. *Journal of Polymer Science: Polymer Physics Edition* 19, 2 (1981), 229–243.
- [24] DUDUCHAVA, R. On poincaré, friedrichs and korns inequalities on domains and hypersurfaces. *arXiv preprint arXiv:1504.01677* (2015).
- [25] EVANS, L. C. *Partial differential equations*, second ed., vol. 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2010.
- [26] EVANS, L. C. *An introduction to stochastic differential equations*, vol. 82. American Mathematical Soc., 2012.
- [27] FIGALLI, A., KANG, M.-J., AND MORALES, J. Global well-posedness of the spatially homogeneous kolmogorov–vicsek model as a gradient flow. *Archive for Rational Mechanics and Analysis* (Sep 2017).
- [28] FROUVELLE, A. A continuum model for alignment of self-propelled particles with anisotropy and density-dependent parameters. *Mathematical Models and Methods in Applied Sciences* 22, 07 (2012), 1250011.
- [29] FROUVELLE, A., AND LIU, J.-G. Dynamics in a kinetic model of oriented particles with phase transition. *SIAM Journal on Mathematical Analysis* 44, 2 (2012), 791–826.
- [30] GAMBA, I. M., AND KANG, M.-J. Global weak solutions for kolmogorov–vicsek type equations with orientational interactions. *Archive for Rational Mechanics and Analysis* 222, 1 (Oct 2016), 317–342.
- [31] HEMELRIJK, C. K., AND HILDENBRANDT, H. Schools of fish and flocks of birds: their shape and internal structure by self-organization. *Interface focus* 2, 6 (2012), 726–737.
- [32] HILDENBRANDT, H., CARERE, C., AND HEMELRIJK, C. Self-organized aerial displays of thousands of starlings: a model. *Behavioral Ecology* 21, 6 (10 2010), 1349–1359.
- [33] JABIN, P.-E., AND WANG, Z. Mean field limit for stochastic particle systems. In *Active Particles, Volume 1*. Springer, 2017, pp. 379–402.
- [34] JIANG, N., LUO, Y.-L., AND ZHANG, T.-F. Coupled self-organized hydrodynamics and navier–stokes models: Local well-posedness and the limit from the self-organized kinetic-fluid models. *Archive for Rational Mechanics and Analysis* 236, 1 (2020), 329–387.
- [35] JIANG, N., XIONG, L., AND ZHANG, T.-F. Hydrodynamic limits of the kinetic self-organized models. *SIAM Journal on Mathematical Analysis* 48, 5 (2016), 3383–3411.
- [36] KANG, M.-J., AND MORALES, J. Dynamics of a spatially homogeneous vicsek model for oriented particles on the plane. *arXiv preprint arXiv:1608.00185* (2016).
- [37] MAURY, B., ROUDNEFF-CHUPIN, A., AND SANTAMBROGIO, F. A macroscopic crowd motion model of gradient flow type. *Mathematical Models and Methods in Applied Sciences* 20, 10 (2010), 1787–1821.
- [38] NAVORET, L. A two-species hydrodynamic model of particles interacting through self-alignment. *Mathematical Models and Methods in Applied Sciences* 23, 06 (2013), 1067–1098.
- [39] ONSAGER, L. The effects of shape on the interaction of colloidal particles. *Annals of the New York Academy of Sciences* 51, 4 (1949), 627–659.
- [40] OTTO, F., AND TZAVARAS, A. E. Continuity of velocity gradients in suspensions of rod-like molecules. *Comm. Math. Phys.* 277, 3 (2008), 729–758.
- [41] POYATO, D. Filippov flows and mean-field limits in the kinetic singular kuramoto model. *arXiv preprint arXiv:1903.01305* (2019).
- [42] ROGERS, L. C. G., AND WILLIAMS, D. *Diffusions, Markov processes and martingales: Volume 2, Itô calculus*, vol. 2. Cambridge university press, 2000.
- [43] SZNITMAN, A.-S. Topics in propagation of chaos. In *Ecole d’été de probabilités de Saint-Flour XIX1989*. Springer, 1991, pp. 165–251.
- [44] VICSEK, T., CZIRÓK, A., BEN-JACOB, E., COHEN, I., AND SHOCHET, O. Novel type of phase transition in a system of self-driven particles. *Physical review letters* 75, 6 (1995), 1226.
- [45] VILLANI, C. *Optimal transport: old and new*, vol. 338. Springer Science & Business Media, 2008.

- [46] ZHANG, T.-F., AND JIANG, N. A local existence of viscous self-organized hydrodynamic model. *Nonlinear Analysis: Real World Applications* 34 (2017), 495–506.

MARC BRIANT

UNIVERSITÉ DE PARIS,
MAP5, CNRS,
F-75006 PARIS, FRANCE

E-MAIL: briant.maths@gmail.com

SARA MERINO ACEITUNO

FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA,
OSKAR-MORGENSTERN-PLATZ 1, 1020 VIENNA, AUSTRIA
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SUSSEX
BRIGHTON, BN1 9RH, UNITED KINGDOM

E-MAIL: sara.merino@univie.ac.at