

# Model-independent constraints in inflationary magnetogenesis

Yuri Shtanov<sup>a,b</sup> and Mykhailo Pavliuk<sup>c</sup>

<sup>a</sup>Bogolyubov Institute for Theoretical Physics,  
Metrologichna St. 14-b, Kiev 03143, Ukraine

<sup>b</sup>Astronomical Observatory, Taras Shevchenko National University of Kiev,  
Observatorna St. 3, Kiev 04053, Ukraine

<sup>c</sup>Department of Physics, Taras Shevchenko National University of Kiev,  
Academician Glushkov Ave. 2, Kiev 03022, Ukraine

E-mail: [shtanov@bitp.kiev.ua](mailto:shtanov@bitp.kiev.ua), [pavliukconnection@gmail.com](mailto:pavliukconnection@gmail.com)

**Abstract.** We derive a simple model-independent upper bound on the strength of magnetic fields obtained in inflationary and post-inflationary magnetogenesis taking into account back-reaction and Schwinger effect. This bound turns out to be rather low for cosmologically interesting spatial scales. Somewhat higher upper bound is obtained if one assumes that some unknown mechanism suppresses the Schwinger effect in the early universe. Incidentally, we correct our previous estimates for this case.

**Keywords:** primordial magnetic fields, inflation

**ArXiv ePrint:** [2004.00947](https://arxiv.org/abs/2004.00947)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The setup</b>	<b>2</b>
<b>3</b>	<b>Amplification inside the Hubble radius</b>	<b>3</b>
<b>4</b>	<b>Back-reaction and Schwinger effect</b>	<b>5</b>
<b>5</b>	<b>Amplification outside the Hubble radius</b>	<b>7</b>
<b>6</b>	<b>The role of spectrum</b>	<b>10</b>
<b>7</b>	<b>Summary and discussion</b>	<b>12</b>

---

## 1 Introduction

Origin of magnetic fields, observed in different objects and on different spatial scales in the universe [1], remains to be unclear (for reviews, see [2–7]). One of the widely discussed possibilities is that they were generated at the inflationary stage; this naturally explains their large coherence length, which can be comparable to the size of the large-scale structure [8–12]. Numerous attempts at realising inflationary magnetogenesis were made by introducing non-trivial couplings of other fields to the electromagnetic field that break the conformal invariance of the latter. A general class of models that we consider here is described by a gauge-invariant action of the form, pioneered in [13, 14],

$$\mathcal{L}_{\text{em}} = -\frac{1}{4}I^2 \left( F_{\mu\nu} + \frac{1}{4}f\tilde{F}_{\mu\nu} \right) F^{\mu\nu}, \quad (1.1)$$

where  $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu}{}^{\alpha\beta}F_{\alpha\beta}$  is the Hodge dual of the electromagnetic strength tensor  $F_{\mu\nu}$ , and  $I$  and  $f$  are non-trivial functions of the cosmological time due to their possible dependence on the background fields such as the inflaton or the metric curvature.

Inflationary magnetogenesis theory is confronted by the issues of back-reaction and strong gauge coupling [15–18], and by the Schwinger effect [19–27] that all reduce its efficiency. Various models of evolution of  $I$  and  $f$  were tried in the literature to overcome these difficulties. In this paper, we would like to provide general model-independent upper limits imposed by the back-reaction and Schwinger effect on the possible strength of magnetic field on a particular comoving spatial scale. We will see that strong gauge coupling is automatically disfavoured by the Schwinger effect. Our investigation is similar in spirit to [15, 16] except that we will not impose a priori ansatz for the evolution of the couplings and also take into account the Schwinger effect.

Given a particular comoving spatial scale, characterised by wavenumber  $k$ , one can start amplifying electromagnetic field on that scale well before its Hubble-radius

crossing during inflation. One possible way of achieving this was demonstrated in our previous publications [28, 29] and will be reviewed below. It suffices to consider evolution of the chiral coupling  $f(\eta)$  in (1.1) which is linear in conformal time  $\eta$  in some time interval. Then, only the modes of one helicity in a narrow band with  $k \sim |f'(\eta)|$  will be exponentially amplified. It was mistakenly claimed in our previous papers [28, 29] that this amplification alone could produce large-scale magnetic fields of sufficiently large strengths today. What was overlooked in [28, 29] is that the peak of electromagnetic field in this process is attained by the moment of Hubble-radius crossing during inflation. The subsequent inflationary dilution of electromagnetic fields then makes them quite small by the end of inflation.

Hence, it is necessary to amplify the fields also outside the Hubble radius during and after inflation. At all stages, one should take into account the constraints imposed by back-reaction and Schwinger effect. This will be the subject of the present paper, in which a simple model-independent upper bound for the final magnetic field is derived. This upper bound appears to be quite low for the prospects of inflationary magnetogenesis. Somewhat higher upper bound is obtained if one assumes that some unknown mechanism suppresses the Schwinger effect in the early universe.

## 2 The setup

We work in a spatially flat cosmological model with the metric

$$ds^2 = a^2(\eta) (d\eta^2 - \delta_{ij} dx^i dx^j) , \quad (2.1)$$

where  $\eta$  is the conformal time, and  $a$  is the scale factor. Adopting the longitudinal gauge  $A_0 = 0$ ,  $\partial^i A_i = 0$  for the vector potential, from (1.1) one obtains the equation satisfied by the transverse field variable  $A_i$ :

$$(I^2 A'_i)' - I^2 \nabla^2 A_i + (I^2 f)' \epsilon_{ijk} \partial_j A_k = a^2 j_i , \quad (2.2)$$

where  $j_i$  is the electromagnetic current density,  $\epsilon_{ijk}$  is the spatial Kronecker alternating tensor with  $\epsilon_{123} = 1$ , and the prime denotes the derivative with respect to the conformal time  $\eta$ . Assuming the Ohm's law in a highly conducting medium with conductivity  $\sigma$ , we have

$$j_i = \sigma E_i = -\frac{\sigma}{a} A'_i . \quad (2.3)$$

In the spatial Fourier representation, and in the constant (in space and time) normalized helicity basis  $\{e_i^h(\mathbf{k})\}$ ,  $h = \pm 1$ , such that  $\mathbf{i}\mathbf{k} \times \mathbf{e}^h = hk\mathbf{e}^h$ , we have  $A_i = \sum_h \mathcal{A}_h e_i^h e^{i\mathbf{k}\cdot\mathbf{x}}$ . Then equations (2.2), (2.3) for the helicity components  $\mathcal{A}_h$ , imply

$$(I^2 \mathcal{A}'_h)' + a\sigma \mathcal{A}'_h + \left[ I^2 k^2 + hk (I^2 f)' \right] \mathcal{A}_h = 0 , \quad (2.4)$$

or, equivalently,

$$(I\mathcal{A}_h)'' + \frac{a\sigma}{I^2} (I\mathcal{A}_h)' + \left[ k^2 + hk \frac{(I^2 f)'}{I^2} - \frac{I''}{I} - \frac{a\sigma I'}{I^3} \right] I\mathcal{A}_h = 0 . \quad (2.5)$$

The spectral densities of quantum fluctuations of magnetic and electric field are characterized, respectively, by the standard relations<sup>1</sup>

$$\mathcal{P}_B = \frac{k^4}{8\pi^2 a^4} \sum_h |\mathcal{A}_h(\eta, k)|^2, \quad \mathcal{P}_E = \frac{k^4}{8\pi^2 a^4} \sum_h \left| \frac{\mathcal{A}'_h(\eta, k)}{k} \right|^2, \quad (2.6)$$

in which the amplitude of the vector potential is normalized so that  $I\mathcal{A}_h \sim e^{-ik\eta}$  as  $\eta \rightarrow -\infty$ . The factor in front of the sums in (2.6) is the spectral density of the normal vacuum fluctuations in each mode at the physical wavenumber  $k/a$ . The (nonrenormalized) spectral energy densities per logarithmic interval of wavenumbers are given by


$$\frac{d\rho_B}{d \ln k} = I^2 \mathcal{P}_B, \quad \frac{d\rho_E}{d \ln k} = I^2 \mathcal{P}_E. \quad (2.7)$$

### 3 Amplification inside the Hubble radius

In this paper, we are interested in general upper limits imposed by the back-reaction and Schwinger effect on the possible strength of magnetic field on a particular comoving spatial scale characterized by wavenumber  $k$ . We can start amplifying the field prior to Hubble-radius crossing during inflation. A model for such amplification was proposed in our works [28, 29]. Assuming constant (slowly evolving) kinetic coupling  $I$ , suppose that the chiral coupling evolves as (see Fig. 1)

$$f(\eta) = \frac{1}{2} \Delta f \tanh \left( \frac{2\eta - \eta_1 - \eta_2}{\Delta\eta} \right) + \text{const}, \quad (3.1)$$

in which  $\Delta\eta = \eta_2 - \eta_1 > 0$  is its temporal width. Then  $f'_0 \equiv f'((\eta_1 + \eta_2)/2) = \Delta f / \Delta\eta$ .



**Figure 1.** Evolution of the form (3.1).

<sup>1</sup>These are electromagnetic fields ‘felt’ by the charged matter, in the Lagrangian of which we assume the standard couplings of the form  $|(\partial - iqA)\psi|^2$ , where  $\psi$  is the matter field and  $q$  is its electric charge. We work in the system of units  $\hbar = c = 1$ .

Introducing the dimensionless time  $\tau = (2\eta - \eta_1 - \eta_2)/\Delta\eta$  and wavenumber  $p = k\Delta\eta/2$ , one can write the general solution of (2.4) (with zero electric conductivity) in terms of the Ferrers functions [30, Chapter 14]:

$$\mathcal{A}_h(\tau) = c_+ P_\nu^\mu(\tanh \tau) + c_- P_\nu^{-\mu}(\tanh \tau), \quad (3.2)$$

with

$$\mu = ip, \quad \nu = q - \frac{1}{2}, \quad q \equiv \frac{1}{2}\sqrt{1 + 2h\Delta f p}. \quad (3.3)$$

The asymptotics of the canonically normalized quantity  $I\mathcal{A}_h \sim e^{-ik\eta} = e^{-ip\tau}$  as  $\tau \rightarrow -\infty$  determines the constants  $c_+$  and  $c_-$  in (3.2). By considering the opposite asymptotics  $I\mathcal{A}_h \sim \alpha_k e^{-ip\tau} + \beta_k e^{ip\tau}$  as  $\tau \rightarrow \infty$  in (3.2), one reads off the Bogolyubov's coefficients  $\alpha_k$  and  $\beta_k$ . Skipping a simple calculation, we present here the result:

$$\alpha_k = \frac{\Gamma(1 - ip)\Gamma(-ip)}{\Gamma(\frac{1}{2} + q - ip)\Gamma(\frac{1}{2} - q - ip)}, \quad \beta_k = -i \frac{\cos(\pi q)}{\sinh(\pi p)}, \quad (3.4)$$

with the required property  $|\alpha_k|^2 - |\beta_k|^2 = 1$ . After the evolution of the coupling, the mean number of quanta in a given mode is

$$n_k = |\beta_k|^2 = \frac{\cos^2(\pi q)}{\sinh^2(\pi p)}. \quad (3.5)$$

For the helicity satisfying  $h\Delta f < 0$ , the quantity  $q$ , given by (3.3), is purely imaginary for  $p > 1/2|\Delta f|$ . In the approximation  $p \gg \max\{1/2|\Delta f|, 1/\pi\}$ , we then obtain

$$n_k \approx e^{2\pi(\sqrt{|\Delta f|p/2} - p)} = e^{\pi|\Delta f|(\sqrt{k/k_0} - k/k_0)}, \quad (3.6)$$

where  $k_0 = |f'_0|$  is the wavenumber at which spectrum (3.6) reaches unity on the slope of its exponential decline. The exponent of this expression reaches maximum at  $k_m = k_0/4$ , with the maximum mean occupation number  $n_m = e^{\pi|\Delta f|/4}$ , which is exponentially large for  $|\Delta f| \gg 1$ . Spectrum (3.5) is plotted in Fig. 2 on a logarithmic scale for a typical value  $|\Delta f| = 40$  (see Sec. 4). The mean occupation numbers for the opposite helicity can be neglected.


The spectral densities (2.6) are of comparable magnitudes, and, since one of the helicities is dominating in  $n_k$ , we have, using (3.6),

$$\mathcal{P}_B \approx \mathcal{P}_E \approx \frac{k^4}{4\pi^2 a^4 I^2} \left[ 1 + e^{\pi|\Delta f|(\sqrt{k/k_0} - k/k_0)} \right]. \quad (3.7)$$

We observe that the spectral densities are peaked at the central value  $k = k_m$  with width  $\Delta k \simeq k_m/\sqrt{|\Delta f|} \ll k_m$  for  $|\Delta f| \gg 1$ . Thus, electric and magnetic fields are generated in this scenario with similar spectra in the spectral region of amplification.

Using a Gaussian approximation to (3.7), one can estimate the total excess (over vacuum) of the electromagnetic energy density and the electromagnetic field strength after the end of the process of amplification:

$$\rho_{\text{em}} \simeq \frac{2k_m^4}{\pi^2 a^4} \frac{e^{\pi|\Delta f|/4}}{|\Delta f|^{1/2}}, \quad B \simeq E \simeq \frac{\sqrt{\rho_{\text{em}}}}{I}. \quad (3.8)$$



**Figure 2.** Spectrum (3.5) on a logarithmic scale for the helicity satisfying  $h\Delta f < 0$  and for  $|\Delta f| = 40$ .

The theory contains two free parameters,  $\Delta f$  and  $\Delta\eta$ , which can be adjusted to produce maximally helical electromagnetic fields of desirable strength with spectral density centred at a desirable wavenumber  $k_m = |\Delta f|/4\Delta\eta$ .

The mechanism of electromagnetic-field generation considered here is quite general and can take place at any stage of cosmological evolution where electric conductivity can be neglected. One can use it to amplify electromagnetic field at the inflationary stage prior to the Hubble-radius crossing for the relevant comoving spatial scale.

Solution (3.2) for the electromagnetic field at the centre of the spectral domain of amplification and in the interval  $\eta_1 < \eta < \eta_2$  grows exponentially:  $\mathcal{A}_h \propto e^{k_m\eta}$ . Assuming  $|k_m\eta_2| \lesssim 1$  and taking into account the prefactors  $\propto a^{-4}$  in (2.6), we conclude that the peak of the electromagnetic spectral density is reached at the time where  $|k_m\eta| = k_m/aH \approx 2$ , i.e., a little before the Hubble-radius crossing.<sup>2</sup>

## 4 Back-reaction and Schwinger effect

Regardless of a concrete model of electromagnetic-field amplification, its characteristics should meet the constraints imposed by the considerations of back-reaction and Schwinger effect.

The back-reaction constraint means that the electromagnetic energy density  $\rho_{\text{em}} = I^2(B^2 + E^2)/2$  at least should not exceed the energy density of the rest of matter  $\rho = M_{\text{P}}^2 H^2$ , where  $M_{\text{P}} = \sqrt{3/8\pi G} \approx 4.2 \times 10^{18}$  GeV is the reduced Planck mass expressed through the Newton's gravitational constant  $G$ . Hence, we should have

$$IE \simeq IB \lesssim M_{\text{P}} H. \quad (4.1)$$

<sup>2</sup>And not at the end of inflation, as was mistakenly assumed in [28, 29] with ensuing gross overestimation of the resulting magnetic field.

The electromagnetic energy density at the time of Hubble-radius crossing is estimated as  $\rho_{\text{em}} \simeq Q^2 H^4$  with the requirement  $Q \lesssim M_{\text{P}}/H$  for the amplification factor of electromagnetic field relative to the purely vacuum fluctuations on the Hubble scale. In the model of amplification considered in the previous section, from (3.8) taken at the Hubble-radius crossing, we have

$$Q \simeq \frac{e^{\pi|\Delta f|/8}}{|\Delta f|^{1/4}} \lesssim \frac{M_{\text{P}}}{H}. \quad (4.2)$$

For a typical value  $H \simeq 10^{-6} M_{\text{P}}$ , this requirement is translated as  $|\Delta f| \lesssim 40$ , which limiting value was chosen in the plot of Fig. 2.

Consider now the role of electric conductivity. For our model of amplification from the previous section, the second term in Eq. (2.4) can be neglected under the condition

$$\frac{a\sigma}{I^2} \ll \left| \frac{\mathcal{A}_h''}{\mathcal{A}_h'} \right| \lesssim k_{\text{m}} \quad \Rightarrow \quad \frac{\sigma}{I^2} \ll H, \quad (4.3)$$

since  $k_{\text{m}} = 2aH$  at the peak of electromagnetic field strength. In the opposite case,  $\sigma/I^2 \gg H$ , the term with conductivity in Eq. (2.4) will cause the derivative  $\mathcal{A}_h'$  to exponentially decrease on time scale much exceeding the Hubble time, making the mode  $\mathcal{A}_h$  approximately constant in time.

The main cause of electric conductivity during inflation is the Schwinger effect of production of charged particle-antiparticle pairs from the vacuum. For fermionic particles with mass  $m$  and electric charge  $q$ , the conductivity calculated in the approximation of homogeneous and slowly changing electromagnetic fields is given by [19, 27, 31, 32]

$$\sigma \simeq \frac{q^3}{6\pi^2} \frac{B}{H} \coth\left(\frac{\pi B}{E}\right) e^{-\pi m^2/qE}. \quad (4.4)$$

During inflation, the fermion mass  $m$  is determined by its Yukawa coupling and by the fluctuations of the Higgs field  $\phi \sim H$ , so that  $m = \gamma H$ , where  $\gamma \sim 10^{-6}$  for the electron. Hence, with the condition  $B \simeq E$ , inequality (4.3) leads to the constraint

$$\frac{q^3}{6\pi^2} \frac{E}{H^2} e^{-\pi\gamma^2 H^2/qE} \lesssim I^2. \quad (4.5)$$

For the uppermost admissible value of  $E/H^2 \simeq 6\pi^2 I^2/q^3$ , the exponent in this expression can be neglected, and we obtain the bound

$$E \lesssim \frac{6\pi^2}{q^3} I^2 H^2 \simeq 10^3 I^2 H^2. \quad (4.6)$$

where we have used  $q \approx \sqrt{4\pi/137} \approx 0.3$  for the unit charge.

In view of (4.1) and (4.6), we then have

$$B \simeq E \lesssim \min \left\{ \frac{M_{\text{P}} H}{I}, 10^3 I^2 H^2 \right\} \leq 10 M_{\text{P}}^2 \left( \frac{H}{M_{\text{P}}} \right)^{4/3}. \quad (4.7)$$

Constraint (4.7) at the Hubble-radius crossing would be insufficient to obtain reasonable magnetic fields on large spatial scales without further amplification. Indeed, if we assume adiabatic evolution  $B \propto a^{-2}$  until the present time, then, for the current magnetic field, we obtain an upper bound

$$\begin{aligned} B_0 &\lesssim 10M_{\text{P}}^2 \left(\frac{H_k}{M_{\text{P}}}\right)^{4/3} \left(\frac{a_k}{a_0}\right)^2 = 10 \left(\frac{M_{\text{P}}}{H_k}\right)^{2/3} \left(\frac{k}{a_0}\right)^2 \\ &\sim \frac{400}{\lambda^2} \left(\frac{M_{\text{P}}}{H_k}\right)^{2/3} \sim 10^{-54} \left(\frac{M_{\text{P}}}{H_k}\right)^{2/3} \left(\frac{\text{Mpc}}{\lambda}\right)^2 \text{ G}. \end{aligned} \quad (4.8)$$

Here,  $a_k = k/H_k$  and  $H_k$  are the scale factor and Hubble parameter, respectively, at the Hubble-radius crossing during inflation, and  $\lambda = 2\pi a_0/k$  is the present value of the comoving spatial scale corresponding to the wavenumber  $k$ . Such low values of the resulting magnetic field are insufficient for successful magnetogenesis in scenarios with reasonable energy scale of inflation. Hence, further amplification of magnetic field would be required on given comoving spatial scales after Hubble-radius crossing.

## 5 Amplification outside the Hubble radius

Further amplification of electromagnetic field outside the Hubble radius can be done by evolving the coupling  $I$  in (1.1), and we assume the usual conditions

$$\left\{ k^2, k \frac{|(I^2 f)'|}{I^2} \right\} \ll \frac{|I''|}{I} \quad (5.1)$$

to be valid well after Hubble-radius crossing. Under these conditions, the terms containing  $k$  in (2.5), hence, also in (2.4) can be neglected. Dropping also the terms with electric conductivity, we observe that equation (2.4) leads to the relation  $\mathcal{A}'_h \propto I^{-2}$ . Then, solution for the mode that is matched at the Hubble-radius crossing with  $\mathcal{A}_h \simeq \mathcal{A}'_h/k \simeq Q/I_k$  (ignoring the irrelevant phase), where  $Q$  is the factor of amplification of the mode and  $I_k$  is the value of  $I$  at the Hubble-radius crossing, is given by

$$\mathcal{A}_h \simeq \frac{Q}{I_k} \left[ 1 + k \int_{\eta_k}^{\eta} \left( \frac{I_k}{I(\tilde{\eta})} \right)^2 d\tilde{\eta} \right] = \frac{Q}{I_k} \left[ 1 + \int \left( \frac{I_k}{I} \right)^2 R_k dN \right]. \quad (5.2)$$

In the last expression, the integration variable was transformed as  $kd\eta = kda/a^2H = kdN/aH = R_k dN$ , where  $R_k = k/aH < 1$  and  $N = \ln a$  is the number of  $e$ -foldings.<sup>3</sup>

We would like to obtain the largest possible amplitude for  $\mathcal{A}_h$  in the end. The process of field amplification is limited by the considerations of back-reaction and Schwinger effect, similar to those discussed in the previous section. The back-reaction constraints have the form (4.1). As regards the Schwinger effect, we estimate the

<sup>3</sup>Note that, even if the coupling  $I$  stops evolving, mode (5.2) can still continue to grow as  $\mathcal{A}_h \propto R_k \propto 1/aH$ , with magnetic field evolving as  $B \propto |\mathcal{A}_h|/a^2 \propto 1/a^3H$ , which law was noted and used in [33–35].

related conductivity at all stages by expression (4.4) under the condition  $B \lesssim E$ . This can be justified, e.g., by integrating [27, Eq. (4.11)], although the real situation with the Schwinger current, especially after inflation, is not very clear and may be more complicated [25] (see also discussion in Sec. 7). Then, upper bound of the form (4.6) is applied for the electric field at all stages of cosmological evolution. We thus have the following constraints:

$$IE \lesssim M_{\text{P}}H \quad \Rightarrow \quad Q \frac{k^2 I_k}{a^2 I} \lesssim M_{\text{P}}H, \quad (5.3)$$

$$IB \lesssim M_{\text{P}}H \quad \Rightarrow \quad Q \frac{k^2 I}{a^2 I_k} \left[ 1 + \int \left( \frac{I_k}{I} \right)^2 R_k dN \right] \lesssim M_{\text{P}}H, \quad (5.4)$$

$$E \lesssim 10^3 I^2 H^2 \quad \Rightarrow \quad Q \frac{k^2 I_k}{a^2 I} \lesssim 10^3 I^3 H^2. \quad (5.5)$$

The integrand in the last expression of (5.2) reaches maximum somewhere on the interval before electric conductivity becomes significant, and this gives an estimate for the final amplitude up to a constant of order unity:

$$\mathcal{A}_h \simeq \frac{Q}{I_k} \int \left( \frac{I_k}{I} \right)^2 R_k dN \sim \frac{Q}{I_k} \left[ \left( \frac{I_k}{I} \right)^2 R_k \right]_{\text{max}}. \quad (5.6)$$

The unity in the brackets of (5.2) was neglected since we are interested in a scenario with significant amplification. It remains to determine the allowable maximum in (5.6).

By virtue of conditions (5.3) and (5.5), we have, respectively,

$$\left( \frac{I_k}{I} \right)^2 R_k \lesssim \frac{M_{\text{P}}^2}{Q^2 H^2} R_k^{-3}, \quad (5.7)$$

$$\left( \frac{I_k}{I} \right)^2 R_k \lesssim \frac{(10 I_k)^{3/2}}{Q^{1/2}} = \text{const}. \quad (5.8)$$

It is notable that the right-hand side of (5.7) monotonically increases with time. If the constant threshold on the right-hand side of (5.8) is reached before the conductivity from preheating becomes significant, then it is the Schwinger effect that stops the growth of the mode  $\mathcal{A}_h$  by creating plasma with sufficiently high conductivity.<sup>4</sup> In the opposite case, it is preheating that contributes predominantly to the plasma conductivity, and the final amplitude of  $\mathcal{A}_h$  is determined by the right-hand side of (5.7) somewhere prior to reheating.

Because of the monotonic growth of the right-hand side of (5.7), for given fixed values of  $Q$  and  $I_k$ , the maximal possible amplitude of  $\mathcal{A}_h$  is obtained if the threshold (5.8) is actually reached, and hence, the shut-down of amplification is caused by the Schwinger effect. The resulting magnitude of the mode is estimated as

$$\mathcal{A}_h \sim \frac{Q (10 I_k)^{3/2}}{I_k Q^{1/2}} = 10^{3/2} (Q I_k)^{1/2}. \quad (5.9)$$

---

<sup>4</sup>We note that the Schwinger particle production in this type of models of cosmological magnetogenesis has even been contemplated as an alternative to traditional preheating [25, 36].

The constant on the right-hand side of (5.8) in this case should not exceed the right-hand side of (5.7) at the moment of reheating, so that

$$(10QI_k)^{3/2} \leq \left( \frac{M_{\text{P}}^2}{H^2} R_k^{-3} \right)_r, \quad (5.10)$$

where the label ‘r’ denotes quantities evaluated at the moment of reheating.

The maximal possible amplitude of  $\mathcal{A}_h$  is then determined by (5.9) and by the maximal value of  $I_k$  satisfying (5.10). Finally, we obtain

$$\mathcal{A}_h \lesssim 10 \left[ \left( \frac{M_{\text{P}}}{H} \right)^{2/3} R_k^{-1} \right]_r. \quad (5.11)$$

Note that this result is independent of  $Q$ , the amplification factor prior to the Hubble-radius crossing at the inflationary stage. This is because the constants  $Q$  and  $I_k$  enter our relations (5.6)–(5.8) explicitly only in the combination  $QI_k$ , which is then bonded in (5.10).

Condition (5.4) at the moment of reheating implies  $R_k|_r = k/a_r H_r \lesssim 1$ , i.e., our above estimates are valid for the modes that are outside the Hubble radius by the moment of reheating. Also note that the value of  $I_r$  can in this case be found from (5.8) and (5.10) and turns out to be

$$I_r \simeq 10^{-1} \left( \frac{M_{\text{P}}}{H_r} \right)^{1/3} \gtrsim 1 \quad (5.12)$$

for  $H_r \lesssim 10^{-3} M_{\text{P}}$ . Hence  $I_k/I_r \gtrsim R_k^{-4}|_r \geq 1$ . Thus, we are in a safe weak-coupling regime for our gauge field, although we never imposed this constraint. It turns out that the strong-coupling case is automatically disfavoured by the Schwinger effect.

The current strength of the magnetic field on the scale of interest is then estimated from (2.6) and (5.11) as<sup>5</sup>

$$B_0 \lesssim \frac{k^2}{2\sqrt{2}\pi a_0^2} |\mathcal{A}_h| \lesssim \left( \frac{k}{a_0} \right) \left( \frac{a_r}{a_0} \right) M_{\text{P}}^{2/3} H_r^{1/3} \simeq \frac{kT_0}{a_0} g_r^{-1/6} \left( \frac{M_{\text{P}}}{T_r} \right)^{1/3}, \quad (5.13)$$

where  $g_r$  is the number of relativistic degrees of freedom in thermal equilibrium after reheating ( $g_r \approx 100$  in the Standard Model),  $T_r$  is the reheating temperature, and  $T_0 = 2.34 \times 10^{-4}$  eV is the current temperature of relic photons. Substituting here the physical numbers, we get

$$B_0 \lesssim 10^{-30} g_r^{-1/6} \left( \frac{M_{\text{P}}}{T_r} \right)^{1/3} \frac{\text{Mpc}}{\lambda} \text{G}. \quad (5.14)$$

---

<sup>5</sup>Assuming the law  $B \propto a^{-2}$  in the hot universe, we neglect the possible chiral and turbulence effects [7, 37–49] that may modify the evolution of the power spectrum.

It is instructive to see whether the Schwinger effect played a significant role in our upper bound. If we ignore it altogether, then the final amplitude for fixed  $I_k$  is estimated by

$$\mathcal{A}_h \lesssim \frac{Q}{I_k} \left[ \left( \frac{I_k}{I} \right)^2 R_k \right]_r = Q I_k \left( \frac{R_k}{I^2} \right)_r, \quad (5.15)$$

with the only constraint  $I_r \geq 1$  on the value of the coupling to avoid the strong-coupling regime. The maximal possible value of  $I_k$  is obtained from (5.7):  $I_k \lesssim M_{\text{P}} I / Q H R_k^2|_r$ , so that  $\mathcal{A}_h \lesssim M_{\text{P}} / (H R_k I)_r$ . Using the constraint  $I_r \geq 1$ , we get an upper bound

$$\mathcal{A}_h \lesssim \left( \frac{M_{\text{P}}}{H} R_k^{-1} \right)_r. \quad (5.16)$$

This result is larger than (5.11) by a factor  $10^{-1} (M_{\text{P}}/H_r)^{1/3}$ , which is significant for low temperatures of reheating. The estimate for the current magnetic field in this case is

$$B_0 \lesssim 10^{-31} g_r^{-1/3} \frac{M_{\text{P}}}{T_r} \frac{\text{Mpc}}{\lambda} \text{G}. \quad (5.17)$$

This estimate would be applicable if the Schwinger particle production were suppressed, e.g., by enhancement of the masses of charged particles (perhaps, by some non-trivial coupling to  $I$ ), as discussed in [34].

## 6 The role of spectrum

In the previous section, we derived the weakest possible upper bounds, considering electromagnetic field only around one particular scale  $k$ . If we take into account that magnetogenesis, in fact, takes place on all spatial scales after Hubble-radius crossing, then our estimates may be strengthened. Let us show that this happens only mildly, i.e., that estimate (5.14) can be saturated by the order of magnitude. We demonstrate this by constructing a scenario in which the spectrum for electric field remains to be peaked around one particular scale  $k_m$ .

The fastest possible growth of the integrand in (5.2) compatible with the back-reaction constraint (5.3) during inflation occurs under the law<sup>6</sup>

$$I \propto a^{-2} \propto \eta^2. \quad (6.1)$$

The solution for the mode is then

$$\mathcal{A}_h \propto \int \frac{d\eta}{I^2} = C_1 + C_2 a^3. \quad (6.2)$$

---

<sup>6</sup>It is remarkable that evolution of the form (6.1) is also obtained in numerical simulations in models of type (1.1) in which back-reaction of electric field on the inflaton dynamics is taken into account through the dependence of the coupling  $I$  on the inflaton field  $\phi$  [23]. In models of this kind, back-reaction becomes important as the energy density of electric field reaches the value of the order  $\rho_E \sim \epsilon \rho_{\text{inf}}$ , where  $\epsilon = M_{\text{P}}^2 [V'(\phi)/V(\phi)]^2$  is the small inflationary slow-roll parameter, and  $V(\phi)$  is the inflaton potential. Back-reaction effects of this kind, which are model-dependent and which we do not consider in this paper, would further reduce our upper bounds for the admissible electric (hence, also magnetic) field.

The fields in the growing mode evolve as  $E \propto a^2$  and  $B \propto E/a \propto a$ , growing with time.

Consider then scenario (6.1), (6.2) with preliminary amplification as described in Sec. 3. This evolution of  $\mathcal{A}_h$  takes place for all super-Hubble modes starting from the moment  $\eta_m$  of the beginning of evolution of  $I$ , which we take to be the time of Hubble-radius crossing of the mode with wavenumber  $k_m$ . By matching the initial conditions at  $a_m$ , we will have

$$\mathcal{A}_h \simeq \mathcal{A}_m \left[ 1 + \frac{k}{3k_m} \left( \frac{a}{a_m} \right)^3 \right] = \mathcal{A}_m \left[ 1 + \frac{k}{3k_m} \left( \frac{aH}{k_m} \right)^3 \right], \quad (6.3)$$

$$\frac{\mathcal{A}'_h}{k} \simeq \mathcal{A}_m \left( \frac{aH}{k_m} \right)^4, \quad k \leq k_m, \quad (6.4)$$

where  $\mathcal{A}_m = \mathcal{A}_m(\eta_m, k)$  is the mode at the moment  $\eta_m$ , and  $k_m = Ha_m$ .

As regards the modes with  $k > k_m$ , the amplitude  $|I\mathcal{A}_h|$  remains constant till the Hubble-radius crossing, where  $a_k = k/H$ , so that  $\mathcal{A}_h \propto 1/I \propto a^2$ , after which it evolves as (6.2). Hence,

$$\mathcal{A}_h \simeq \mathcal{A}_m \left( \frac{a_k}{a_m} \right)^2 \left( \frac{a}{a_k} \right)^3 = \mathcal{A}_m \frac{(aH)^3}{k_m^2 k}, \quad (6.5)$$

$$\frac{\mathcal{A}'_h}{k} \simeq 3\mathcal{A}_m \frac{(aH)^4}{k_m^2 k^2}, \quad k \geq k_m, \quad (6.6)$$

Equations (6.3)–(6.6) can be interpolated for all  $k$  as

$$\mathcal{A}_h \simeq \mathcal{A}_m \left[ 1 + \frac{k/k_m}{3 + (k/k_m)^2} \left( \frac{aH}{k_m} \right)^3 \right], \quad (6.7)$$

$$\frac{\mathcal{A}'_h}{k} \simeq \mathcal{A}_m \frac{1}{1 + k^2/3k_m^2} \left( \frac{aH}{k_m} \right)^4. \quad (6.8)$$

The spectral densities during inflation are then estimated as

$$\mathcal{P}_B = \frac{d\langle B^2 \rangle}{d\ln k} \simeq \frac{k^4}{8\pi^2 a^4} \left[ 1 + \frac{k/k_m}{3 + (k/k_m)^2} \left( \frac{aH}{k_m} \right)^3 \right]^2 \sum_h |\mathcal{A}_h(\eta_m, k)|^2, \quad (6.9)$$

$$\mathcal{P}_E = \frac{d\langle E^2 \rangle}{d\ln k} \simeq \frac{k^4}{8\pi^2 a^4} \frac{1}{(1 + k^2/3k_m^2)^2} \left( \frac{aH}{k_m} \right)^8 \sum_h |\mathcal{A}_h(\eta_m, k)|^2. \quad (6.10)$$

Due to the strong narrow peak of  $Q(k) \equiv |\mathcal{A}_h(\eta_m, k)|$  at  $k = k_m$ , the spectrum for electric field remains to be peaked at  $k = k_m$  and is flat in the region  $k \gg k_m$ . Magnetic field is suppressed on all relevant scales  $k \lesssim aH$  with respect to the electric one. This, then, does not strengthen much our estimates made in the previous section.

We also remark that, although our upper bounds (5.11) and (5.16) are independent of the value of  $Q$  at that particular wavenumber, the role of initial amplification inside the Hubble radius is significant because it shapes the initial power spectrum, allowing it to be peaked at a given wavenumber and thus making our upper bounds virtually the least upper bounds.

## 7 Summary and discussion

In this paper, we have derived an upper bound [see (5.14)]

$$B_0 \lesssim 10^{-30} g_r^{-1/6} \left( \frac{M_{\text{P}}}{T_r} \right)^{1/3} \frac{\text{Mpc}}{\lambda} \text{G} \quad (7.1)$$

for the outcome of inflationary and post-inflationary magnetogenesis based on Lagrangian (1.1). In doing this, we took into account the constraints from back-reaction and Schwinger effect. Although we did not require the condition  $I > 1$  of weak gauge coupling from the very beginning, it is automatically favoured in the end by the Schwinger effect. Our upper bound is conservative in that it takes into account only electromagnetic fields on a particular spatial scale of interest and can only be lowered by considering the whole power spectrum. Still, we have shown that there exists a scenario in which the power spectrum is peaked around the spatial scale of interest, which makes our bound, in some sense, virtually the least upper bound.

Our upper bound (7.1) allows one to appreciate the difficulty in producing sufficiently large magnetic fields on large spatial scales. Indeed, the lowest possible temperature of reheating lies in the MeV range [58], and even for such low reheating temperature, one gets  $B_0 \lesssim 10^{-23}$  G on the megaparsec scale. Estimate (7.1) is then valid for comoving spatial scales larger than the Hubble radius at this temperature, i.e., for  $\lambda \gtrsim 100$  pc.

In our treatment, we used the simple expression (4.4) for the electric conductivity induced by the Schwinger effect at the post-inflationary stage. Hoping that it gives correct estimates by order of magnitude, we should point out that recent elaborate calculations in the kinetic approach reveal an oscillatory and non-Markovian character of the electric current and field after inflation, failing to satisfy the simple Ohm's law and decaying slower than in the case of naïve application of (4.4) [25]. This may call for revisions of our estimates in more refined treatments of the Schwinger effect.

If the Schwinger effect for some reason is suppressed in the early universe (e.g., because the masses of charged particles are sufficiently enlarged by some non-trivial mechanism), then only back-reaction is important, and our estimate for the final magnetic field becomes [see (5.17)]

$$B_0 \lesssim 10^{-31} g_r^{-1/3} \frac{M_{\text{P}}}{T_r} \frac{\text{Mpc}}{\lambda} \text{G}. \quad (7.2)$$

This would allow one to obtain  $B_0 \sim 10^{15}$  G on the megaparsec scale by choosing  $T_r \sim 10^2$  GeV. In some specific scenarios, generation of magnetic fields of this order of magnitude requires even lower reheating temperatures, of the order of GeV [35, 59].

We did not take into account the contribution of electromagnetic field to the primordial power spectrum of energy-density fluctuations, but the corresponding constraints on the magnetic field [50] appear to be much weaker than those obtained in this paper.

Finally, we note that primordial helical hypermagnetic fields may also be responsible for generating baryon asymmetry of the universe [51–57]. This imposes a typical

constraint [55, 57]

$$B_0 \lesssim 10^{-21} \left( \frac{\text{Mpc}}{\lambda} \right)^{1/2} \text{ G} \quad (7.3)$$

on the admissible values of  $B_0$  on the spatial scale  $\lambda$  for the fields of *maximal* helicity that existed prior to the electroweak transition. This constraint is weaker than our upper bound (7.1) for not very small spatial scales.

## Acknowledgments

We are grateful to Eduard Gorbar for valuable comments and discussion. Y. S. acknowledges support from the National Academy of Sciences of Ukraine in frames of priority project “Fundamental properties of matter in the relativistic collisions of nuclei and in the early Universe” (No. 0120U100935) and from the scientific program “Astronomy and Space Physics” (project 19BF023-01) of the Taras Shevchenko National University of Kiev.

## References

- [1] U. Klein and A. Fletcher, *Galactic and Intergalactic Magnetic Fields*, Springer (2015).
- [2] D. Grasso and H. R. Rubinstein, *Magnetic fields in the early universe*, *Phys. Rept.* **348** (2001) 163 [astro-ph/0009061] [inSPIRE].
- [3] L. M. Widrow, *Origin of galactic and extragalactic magnetic fields*, *Rev. Mod. Phys.* **74** (2002) 775 [astro-ph/0207240] [inSPIRE].
- [4] M. Giovannini, *The magnetized universe*, *Int. J. Mod. Phys. D* **13** (2004) 391 [astro-ph/0312614] [inSPIRE].
- [5] A. Kandus, K. E. Kunze and C. G. Tsagas, *Primordial magnetogenesis*, *Phys. Rept.* **505** (2011) 1 [1007.3891] [inSPIRE].
- [6] R. Durrer and A. Neronov, *Cosmological magnetic fields: their generation, evolution and observation*, *Astron. Astrophys. Rev.* **21** (2013) 62 [1303.7121] [inSPIRE].
- [7] K. Subramanian, *The origin, evolution and signatures of primordial magnetic fields*, *Rept. Prog. Phys.* **79** (2016) 076901 [1504.02311] [inSPIRE].
- [8] F. Tavecchio, G. Ghisellini, L. Foschini, G. Bonnoli, G. Ghirlanda and P. Coppi, *The intergalactic magnetic field constrained by Fermi/Large Area Telescope observations of the TeV blazar 1ES 0229+200*, *Mon. Not. Roy. Astron. Soc.* **406** (2010) L70 [1004.1329] [inSPIRE].
- [9] S. 'i. Ando and A. Kusenko, *Evidence for gamma-ray halos around active galactic nuclei and the first measurement of intergalactic magnetic fields*, *Astrophys. J.* **722** (2010) L39 [1005.1924] [inSPIRE].
- [10] A. Neronov and I. Vovk, *Evidence for Strong Extragalactic Magnetic Fields from Fermi Observations of TeV Blazars*, *Science* **328** (2010) 73 [1006.3504] [inSPIRE].

- [11] K. Dolag, M. Kachelriess, S. Ostapchenko and R. Tomàs, *Lower limit on the strength and filling factor of extragalactic magnetic fields*, *Astrophys. J. Lett.* **727** (2011) L4 [1009.1782] [inSPIRE].
- [12] A. M. Taylor, I. Vovk and A. Neronov, *Extragalactic magnetic fields constraints from simultaneous GeV–TeV observations of blazars*, *Astron. Astrophys.* **529** (2011) A144 [1101.0932] [inSPIRE].
- [13] M. S. Turner and L. M. Widrow, *Inflation-produced, large-scale magnetic fields*, *Phys. Rev. D* **37** (1988) 2743 [inSPIRE].
- [14] B. Ratra, *Cosmological “seed” magnetic field from inflation*, *Astrophys. J.* **391** (1992) L1 [inSPIRE].
- [15] V. Demozzi, V. Mukhanov and H. Rubinstein, *Magnetic fields from inflation?*, *JCAP* **08** (2009) 025 [0907.1030] [inSPIRE].
- [16] F. R. Urban, *On inflating magnetic fields, and the backreactions thereof*, *JCAP* **12** (2011) 012 [1111.1006] [inSPIRE].
- [17] H. Bazrafshan Moghaddam, E. McDonough, R. Namba and R. H. Brandenberger, *Inflationary magneto-(non)genesis, increasing kinetic couplings, and the strong coupling problem*, *Class. Quant. Grav.* **35** (2018) 105015 [1707.05820] [inSPIRE].
- [18] R. Sharma, K. Subramanian and T. R. Seshadri, *Generation of helical magnetic field in a viable scenario of inflationary magnetogenesis*, *Phys. Rev. D* **97** (2018) 083503 [1802.04847] [inSPIRE].
- [19] T. Kobayashi and N. Afshordi, *Schwinger effect in 4D de Sitter space and constraints on magnetogenesis in the early universe*, *JHEP* **1410** (2014) 166 [1408.4141] [inSPIRE].
- [20] R. Sharma, S. Jagannathan, T. R. Seshadri and K. Subramanian, *Challenges in inflationary magnetogenesis: Constraints from strong coupling, backreaction, and the Schwinger effect*, *Phys. Rev. D* **96** (2017) 083511 [1708.08119] [inSPIRE].
- [21] C. Stahl, *Schwinger effect impacting primordial magnetogenesis*, *Nucl. Phys. B* **939** (2019) 95 [1806.06692] [inSPIRE].
- [22] H. Kitamoto, *Schwinger effect in inflaton-driven electric field*, *Phys. Rev. D* **98** (2018) 103512 [1807.03753] [inSPIRE].
- [23] O. O. Sobol, E. V. Gorbar, M. Kamarpour and S. I. Vilchinskii, *Influence of backreaction of electric fields and Schwinger effect on inflationary magnetogenesis*, *Phys. Rev. D* **98** (2018) 063534 [1807.09851] [inSPIRE].
- [24] O. O. Sobol, E. V. Gorbar and S. I. Vilchinskii, *Backreaction of electromagnetic fields and the Schwinger effect in pseudoscalar inflation magnetogenesis*, *Phys. Rev. D* **100** (2019) 063523 [1907.10443] [inSPIRE].
- [25] E. V. Gorbar, A. I. Momot, O. O. Sobol and S. I. Vilchinskii, *Kinetic approach to the Schwinger effect during inflation*, *Phys. Rev. D* **100** (2019) 123502 [1909.10332] [inSPIRE].
- [26] V. Domcke and K. Mukaida, *Gauge field and fermion production during axion inflation*, *JCAP* **11** (2018) 020 [1806.08769] [inSPIRE].

- [27] V. Domcke, Y. Ema and K. Mukaida, *Chiral anomaly, Schwinger effect, Euler–Heisenberg Lagrangian and application to axion inflation*, *JHEP* **2002** (2020) 055 [[1910.01205](#)] [[inSPIRE](#)].
- [28] Y. Shtanov, *Viable inflationary magnetogenesis with helical coupling*, *JCAP* **10** (2019) 008 [[1902.05894](#)] [[inSPIRE](#)].
- [29] Y. V. Shtanov and M. V. Pavliuk, *Inflationary magnetogenesis with helical coupling*, *Ukr. Phys. J.* **64** (2019) 1009 [[1911.10424](#)] [[inSPIRE](#)]. [[arXiv:1911.10424](#) [[astro-ph.CO](#)]].
- [30] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clark (eds), *NIST Handbook of Mathematical Functions*, NIST & Cambridge University Press (2010).
- [31] E. Bavarsad, C. Stahl and S. S. Xue, *Scalar current of created pairs by Schwinger mechanism in de Sitter spacetime*, *Phys. Rev. D* **94** (2016) 104011 [[1602.06556](#)] [[inSPIRE](#)].
- [32] T. Hayashinaka, T. Fujita and J. Yokoyama, *Fermionic Schwinger effect and induced current in de Sitter space*, *JCAP* **07** (2016) 010 [[1603.04165](#)] [[inSPIRE](#)].
- [33] T. Kobayashi, *Primordial magnetic fields from the post-inflationary universe*, *JCAP* **05** (2014) 040 [[1403.5168](#)] [[inSPIRE](#)].
- [34] T. Kobayashi and M. S. Sloth, *Early cosmological evolution of primordial electromagnetic fields*, *Phys. Rev. D* **100** (2019) 023524 [[1903.02561](#)] [[inSPIRE](#)].
- [35] T. Fujita and R. Durrer, *Scale-invariant helical magnetic fields from inflation*, *JCAP* **09** (2019) 008 [[1904.11428](#)] [[inSPIRE](#)].
- [36] W. Tangarife, K. Tobioka, L. Ubaldi and T. Volansky, *Dynamics of relaxed inflation*, *JHEP* **1802** (2018) 084 [[1706.03072](#)] [[inSPIRE](#)].
- [37] R. Banerjee and K. Jedamzik, *Evolution of cosmic magnetic fields: From the very early Universe, to recombination, to the present*, *Phys. Rev. D* **70** (2004) 123003 [[astro-ph/0410032](#)] [[inSPIRE](#)].
- [38] A. Boyarsky, J. Fröhlich and O. Ruchayskiy, *Self-consistent evolution of magnetic fields and chiral asymmetry in the early universe*, *Phys. Rev. Lett.* **108** (2012) 031301 [[1109.3350](#)] [[inSPIRE](#)].
- [39] T. Kahniashvili, A. G. Tevzadze, A. Brandenburg and A. Neronov, *Evolution of primordial magnetic fields from phase transitions*, *Phys. Rev. D* **87** (2013) 083007 [[1212.0596](#)] [[inSPIRE](#)].
- [40] A. Saveliev, K. Jedamzik and G. Sigl, *Evolution of helical cosmic magnetic fields as predicted by magnetohydrodynamic closure theory*, *Phys. Rev. D* **87** (2013) 123001 [[1304.3621](#)] [[inSPIRE](#)].
- [41] A. Boyarsky, J. Fröhlich and O. Ruchayskiy, *Magnetohydrodynamics of chiral relativistic fluids*, *Phys. Rev. D* **92** (2015) 043004 [[1504.04854](#)] [[inSPIRE](#)].
- [42] Y. Hirono, D. Kharzeev and Y. Yin, *Self-similar inverse cascade of magnetic helicity driven by the chiral anomaly*, *Phys. Rev. D* **92** (2015) 125031 [[1509.07790](#)] [[inSPIRE](#)].
- [43] E. V. Gorbar, I. A. Shovkovy, S. Vilchinskii, I. Rudenok, A. Boyarsky and

- O. Ruchayskiy, *Anomalous Maxwell equations for inhomogeneous chiral plasma*, *Phys. Rev. D* **93** (2016) 105028 [1603.03442] [inSPIRE].
- [44] M. Sydorenko, O. Tomalak and Y. Shtanov, *Magnetic fields and chiral asymmetry in the early hot universe*, *JCAP* **10** (2016) 018 [1607.04845] [inSPIRE].
- [45] E. V. Gorbar, I. Rudenok, I. A. Shovkovy and S. Vilchinskii, *Anomaly-driven inverse cascade and inhomogeneities in a magnetized chiral plasma in the early Universe*, *Phys. Rev. D* **94** (2016) 103528 [1610.01214] [inSPIRE].
- [46] P. Pavlović, N. Leite and G. Sigl, *Chiral magnetohydrodynamic turbulence*, *Phys. Rev. D* **96** (2017) 023504 [1612.07382] [inSPIRE].
- [47] A. Brandenburg, J. Schober, I. Rogachevskii, T. Kahniashvili, A. Boyarsky, J. Fröhlich, O. Ruchayskiy and N. Kleeorin, *The turbulent chiral magnetic cascade in the early universe*, *Astrophys. J. Lett.* **845** (2017) L21 [1707.03385] [inSPIRE].
- [48] I. Rogachevskii, O. Ruchayskiy, A. Boyarsky, J. Fröhlich, N. Kleeorin, A. Brandenburg and J. Schober, *Laminar and turbulent dynamos in chiral magnetohydrodynamics. I. Theory*, *Astrophys. J.* **846** (2017) 153 [1705.00378] [inSPIRE]. [arXiv:1705.00378 [physics.plasm-ph]].
- [49] J. Schober, I. Rogachevskii, A. Brandenburg, A. Boyarsky, J. Fröhlich, O. Ruchayskiy and N. Kleeorin, *Laminar and turbulent dynamos in chiral magnetohydrodynamics. II. Simulations*, *Astrophys. J.* **858** (2018) 124 [1711.09733] [inSPIRE].
- [50] T. Markkanen, S. Nurmi, S. Räsänen and V. Vennin, *Narrowing the window of inflationary magnetogenesis*, *JCAP* **06** (2017) 035 [1704.01343] [inSPIRE].
- [51] K. Bamba, *Baryon asymmetry from hypermagnetic helicity in dilaton hypercharge electromagnetism*, *Phys. Rev. D* **74** (2006) 123504 [hep-ph/0611152] [inSPIRE].
- [52] M. M. Anber and E. Sabancilar, *Hypermagnetic fields and baryon asymmetry from pseudoscalar inflation*, *Phys. Rev. D* **92** (2015) 101501(R) [1507.00744] [inSPIRE].
- [53] T. Fujita and K. Kamada, *Large-scale magnetic fields can explain the baryon asymmetry of the Universe*, *Phys. Rev. D* **93** (2016) 083520 [1602.02109] [inSPIRE].
- [54] K. Kamada and A. J. Long, *Baryogenesis from decaying magnetic helicity*, *Phys. Rev. D* **94** (2016) 063501 [1606.08891] [inSPIRE].
- [55] K. Kamada and A. J. Long, *Evolution of the baryon asymmetry through the electroweak crossover in the presence of a helical magnetic field*, *Phys. Rev. D* **94** (2016) 123509 [1610.03074] [inSPIRE].
- [56] D. Jiménez, K. Kamada, K. Schmitz and X.-J. Xu, *Baryon asymmetry and gravitational waves from pseudoscalar inflation*, *JCAP* **12** (2017) 011 [1707.07943] [inSPIRE].
- [57] V. Domcke, B. von Harling, E. Morgante and K. Mukaida, *Baryogenesis from axion inflation*, *JCAP* **10** (2019) 032 [1905.13318] [inSPIRE].
- [58] S. Hannestad, *What is the lowest possible reheating temperature?*, *Phys. Rev. D* **70** (2004) 043506 [astro-ph/0403291] [inSPIRE].
- [59] T. Fujita and R. Namba, *Pre-reheating magnetogenesis in the kinetic coupling model*, *Phys. Rev. D* **94** (2016) 043523 [1602.05673] [inSPIRE].