

# The G-dynamics of the QCPB theory

Gen WANG \*

*School of Mathematical Sciences, Xiamen University,  
Xiamen, 361005, P.R.China*

## Abstract

In this paper, we further study the G-dynamics newly emerged in the covariant dynamics defined by the QCPB theory. Then we want to seek the precise formulations of the G-dynamics to calculate the practical quantum problems. We explicitly verify how the role of G-dynamics plays in quantum mechanics, and then we propose three new operators based on the G-dynamics: D-operator, T-operator and G-operator, we find that the non-Hermitian operators are inevitable to appear in the framework of the QCPB. Then the eigenvalue equation of them are discussed accordingly by using the Schrödinger equation and geometrinetic energy operator (GEO) which leads to the Ri-operator, respectively, we obtain some useful results between two quantum situations. Meanwhile, we present generalized covariant wave equation based on the G-operator. The GEO for quantum harmonic oscillator as an application is further calculated.

---

\*720195765863, email: Wg1991dream@163.com  
sterces ofo neila aiv siunew gang-pwc-zgy.

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	The Schrödinger equation . . . . .	2
<b>2</b>	<b>Generalized geometric commutator and GAC</b>	<b>3</b>
2.1	QCPB and quantum geobacket (QGB) . . . . .	4
2.2	Quantum covariant Hamiltonian system . . . . .	5
2.3	Covariant dynamics, generalized Heisenberg equation, G-dynamics . . . . .	6
2.4	Imaginary geomenergy . . . . .	6
2.5	The QCPB for quantum harmonic oscillator . . . . .	7
2.6	Geomentum operator . . . . .	8
2.7	The geoperator . . . . .	8
<b>3</b>	<b>G-dynamics</b>	<b>9</b>
3.1	D-operator, T-operator and G-operator . . . . .	9
<b>4</b>	<b>Geometrinetic energy operator (GEO) and Ri-operator</b>	<b>10</b>
4.1	The G-dynamics with Schrödinger equation . . . . .	11
<b>5</b>	<b>The cases of G-dynamics</b>	<b>14</b>
5.1	For classical Hamiltonian operator . . . . .	14
5.2	For Ri-operator . . . . .	15
5.3	G-operator for Ri-operator . . . . .	17
5.4	Generalized covariant wave equation . . . . .	18
<b>6</b>	<b>GEO for generalized quantum harmonic oscillator</b>	<b>19</b>
<b>7</b>	<b>Conclusions</b>	<b>21</b>

## 1 Introduction

### 1.1 The Schrödinger equation

In quantum mechanics, the form of the Schrödinger equation depends on the physical situation. The most general form is the time-dependent Schrödinger equation, which gives a description of a system evolving with time [1,2]

$$\sqrt{-1}\hbar\frac{\partial\psi(r,t)}{\partial t} = \hat{H}^{(cl)}\psi = \left(-\frac{\hbar^2}{2m}\nabla^2 + V\right)\psi(r,t) \quad (1)$$

where  $\partial/\partial t$  symbolizes a partial derivative with respect to time  $t$ ,  $\psi$  is the wave function of the quantum system,

$$\hat{H}^{(cl)} = -\frac{\hbar^2}{2m}\nabla^2 + V \quad (2)$$

is the Hamiltonian operator in terms of the coordinates. The time-dependent Schrödinger equation described above predicts that wave functions can form standing waves, called stationary states. Stationary states can also be described by a simpler form of the Schrödinger equation, the time-independent Schrödinger equation

$$\widehat{H}^{(cl)}\psi = \left( -\frac{\hbar^2}{2m}\nabla^2 + V \right) \psi = E\psi$$

The energy and momentum operators, which are respectively shown as

$$\widehat{H}^{(cl)} = \sqrt{-1}\hbar\frac{\partial}{\partial t}, \quad \widehat{p}^{(cl)} = -\sqrt{-1}\hbar\nabla \quad (3)$$

The Schrödinger equation in its general form is shown as (1).

With the non-universal validity of the QPB that has been replaced by the complete QCPB theory. In this paper, on the foundation of the Schrödinger equation, we mainly explore the properties of the G-dynamics in the covariant dynamics defined by the QCHS in QCPB, we obtain the more specific expressions of the G-dynamics in terms of the coordinates to make the G-dynamics practical in the quantum mechanics as it develops.

## 2 Generalized geometric commutator and GAC

Let  $a$  and  $b$  be any two operators, their commutator is formally defined by  $[a, b]_{cr} = ab - ba$ . Note that the operator for commutator can be any mathematical form to be appeared for the calculation, such as a function, vector, differential operator, partial differential operator, even a number in a number field, and so on, it can be arbitrarily chosen according to our needs.

Let  $a$  and  $b$  be any elements of any algebra, or any operators, their generalized geometric commutator is formally defined by the following.

**Definition 1** (GGC). [3] *A generalized geometric commutator (GGC) of arity two  $a, b$  is formally given by*

$$[a, b] = [a, b]_{cr} + G(s; a, b)$$

*The geomutator is*

$$G(s; a, b) = a[s, b]_{cr} - b[s, a]_{cr}$$

*satisfying  $G(s; a, b) = -G(s; b, a)$ , where  $s$  is a geometric structure function given by domain.*

Some properties of the geomutator are given by

$$G(s, a + b, c + d) = G(s, a, c) + G(s, b, d) + G(s, b, c) + G(s, a, d)$$

$$G(s, a + b, c) = G(s, a, c) + G(s, b, c)$$

$$G(s, a, c + d) = G(s, a, c) + G(s, a, d)$$

for operators  $a, b, c, d$ .

Obviously, in some sense, the anticommutator also needs to be generalized as the commutator does, Analogously, denote by  $\{a, b\}_{ir} = ab + ba$  the anticommutator.

**Definition 2.** *The geometric anticommutator (GAC) of any two elements  $a$  and  $b$  is defined by*

$$\{a, b\} = \{a, b\}_{ir} + Z(s, a, b)$$

where  $Z(s, a, b) = Z(s, b, a)$  is called anti-geomutator,  $s$  is geometric function.

As the definition above stated,

**Definition 3.** *The anti-geomutator can be taken as*

$$Z(s, a, b) = (a : s : b) + \{a, b\}_{ir,s}$$

where  $(a : s : b) = asb + bsa$ , and  $s$  is the geometric function created by the environment.

As a result of the symmetry of anti-geomutator, geometric anticommutator then follows the symmetry  $\{a, b\} = \{b, a\}$ . Actually, the anti-geomutator can be expressed in the form

$$\begin{aligned} Z(s, a, b) &= a\{s, b\}_{ir} + b\{s, a\}_{ir} \\ &= (a : s : b) + \{a, b\}_{ir,s} \end{aligned}$$

As seen, this form is very similar to the quantum geometric bracket, this is why we need to generalize the anticommutator. There has a clear property of the anti-geomutator given by

$$\begin{aligned} Z(s, a + b, c + d) &= (a + b)\{s, c + d\}_{ir} + (c + d)\{s, a + b\}_{ir} \\ &= Z(s, a, c) + Z(s, a, d) + Z(s, b, c) + Z(s, b, d) \end{aligned}$$

for operators  $a, b, c, d$ .

## 2.1 QCPB and quantum geobacket (QGB)

As [3] stated, the quantum covariant Poisson bracket (QCPB) is defined by generalized geometric commutator (GGC) while quantum geometric bracket (QGB) is given based on the geomutator. More precisely,

**Definition 4** (QCPB). [3] *The QCPB is generally defined as*

$$[\hat{f}, \hat{g}] = [\hat{f}, \hat{g}]_{QPB} + G(s, \hat{f}, \hat{g})$$

in terms of quantum operator  $\hat{f}$ ,  $\hat{g}$ , where  $G(s, \hat{f}, \hat{g}) = -G(s, \hat{g}, \hat{f})$  is called quantum geometric bracket.

It is zero if and only if  $\hat{f}$  and  $\hat{g}$  covariant commute, i.e.  $[\hat{f}, \hat{g}] = 0$ . It is remarkable to see that the QCPB representation admits a dynamical geometric bracket formula on the manifold.

**Definition 5** (Quantum geometric bracket (QGB)). [3] *The quantum geometric bracket is*

$$G(s, \hat{f}, \hat{g}) = \hat{f}[s, \hat{g}]_{QPB} - \hat{g}[s, \hat{f}]_{QPB}$$

where  $s$  represents the globally condition of space.

Let's assert the role of the structure function  $s$  that is a geometric structure function given by domain based on the generalized geometric commutator in definition 4, it means that structure function  $s$  is only determined by the environment, or spacetime, or manifolds, the domain, ect, from this viewpoint, the environment joins the physical process, the influence of the environment now based on new theory can't be ignored, it's naturally necessary to be considered in a physical process.

**Definition 6.** [3] *The covariant equilibrium equation is given by  $[\hat{f}, \hat{g}] = 0$ , i.e.,*

$$[\hat{f}, \hat{g}]_{QPB} + G(s, \hat{f}, \hat{g}) = 0$$

for operators  $\hat{f}$ ,  $\hat{g}$ .

## 2.2 Quantum covariant Hamiltonian system

This section will give the covariant dynamics which contains two different sub-dynamics: the generalized Heisenberg equation and G-dynamics [3]. It tells that the generalized Heisenberg equation has improved the classical Heisenberg equation by considering the quantum geometric bracket. Firstly, let's give the definition of quantum covariant Hamiltonian system (QCHS) based on the QCPB as definition 4 defined.

In the beginning, let us start with the QCPB by taking  $\hat{g} = \hat{H}$  into consideration, then it formally yields below definition.

**Definition 7 (QCHS).** [3] *The QCHS defined by QCPB in terms of quantum operator  $\hat{f}$ ,  $\hat{H}$  is generally given by*

$$[\hat{f}, \hat{H}] = [\hat{f}, \hat{H}]_{QPB} + G(s, \hat{f}, \hat{H})$$

where  $G(s, \hat{f}, \hat{H})$  is quantum geobacket .

**Definition 8.** [3] *The quantum geobacket for a Hamiltonian operator  $\hat{H}$  and operator  $\hat{f}$  is given by*

$$G(s, \hat{f}, \hat{H}) = \hat{f}[s, \hat{H}]_{QPB} - \hat{H}[s, \hat{f}]_{QPB}$$

and  $s$  is a real structural function only associated with the structure space.

Manifestly, this operator  $\hat{f}$  covariantly commutes with  $H$ . By our construction, the QCHS defined by the QCPB is a transition for the further development of a complete and covariant theory that can naturally generalize the Heisenberg equation. Since  $\hat{f}$  is an observable,  $s$  is assumed to be real function as well. In this way, we have successfully given a complete description of the Heisenberg equation.

**Definition 9.** [3] *The covariant equilibrium equation is given by  $[\hat{f}, \hat{H}] = 0$ , i.e.,*

$$[\hat{f}, \hat{H}]_{QPB} + G(s, \hat{f}, \hat{H}) = 0$$

for operators  $\hat{f}$ ,  $\hat{H}$ , then  $\hat{f}$  is called quantum covariant conserved quantity.

Furthermore, there is a certain case given by  $[\hat{H}, \hat{H}] = 0$ , when  $\hat{f} = \hat{H}$  based on definition 9. It is clear to see that  $\hat{H}$  is a quantum covariant conserved quantity.

### 2.3 Covariant dynamics, generalized Heisenberg equation, G-dynamics

In this section, we will briefly review the entire theoretical framework of quantum covariant Hamiltonian system defined by the quantum covariant Poisson bracket totally based on the paper [3]. More precisely, the time covariant evolution of any observable  $\hat{f}$  in the covariant dynamics is given by both the generalized Heisenberg equation of motion and G-dynamics.

**Theorem 1.** [3] *The covariant dynamics, the generalized Heisenberg equation, G-dynamics can be formally formulated as*

**The covariant dynamics:**  $\frac{\mathcal{D}\hat{f}}{dt} = \frac{1}{\sqrt{-1}\hbar} [\hat{f}, \hat{H}]$

**The generalized Heisenberg equation:**  $\frac{d\hat{f}}{dt} = \frac{1}{\sqrt{-1}\hbar} [\hat{f}, \hat{H}]_{QPB} - \frac{1}{\sqrt{-1}\hbar} \hat{H} [s, \hat{f}]_{QPB}$

**G-dynamics:**  $\hat{w} = \frac{1}{\sqrt{-1}\hbar} [s, \hat{H}]_{QPB}$ .

respectively, where  $\frac{\mathcal{D}}{dt} = \frac{d}{dt} + \hat{w}$  is covariant time operator, and  $\hat{H}$  is the Hamiltonian and  $[\cdot, \cdot]$  denotes the GGC of two operators.

### 2.4 Imaginary geomenergy

In this section, the imaginary geomenergy is defined based on the G-dynamics, by using this new concept, we can better give a presentation for the covariant dynamics, ect.

**Definition 10.** [3] *The imaginary geomenergy is defined by  $E^{(\text{Im})}(\hat{w}) = \sqrt{-1}\hbar\hat{w}$ , where  $\hat{w}$  means G-dynamics.*

It is clear that the covariant dynamics leads us to a complete quantum system in which the state of the system is represented by two separated dynamic quantum system, the covariant dynamics can be totally pictured in the form below

$$\begin{aligned} \sqrt{-1}\hbar \frac{\mathcal{D}}{dt} \hat{f} &= [\hat{f}, \hat{H}] = [\hat{f}, \hat{H}]_{QPB} + G(s, \hat{f}, \hat{H}) \\ &= \sqrt{-1}\hbar \frac{d}{dt} \hat{f} + \sqrt{-1}\hbar \hat{f} \hat{w} \end{aligned}$$

With the imaginary geomenergy defined above, then covariant dynamics is rewritten in the form

$$\sqrt{-1}\hbar \frac{\mathcal{D}}{dt} \hat{f} = [\hat{f}, \hat{H}] = \sqrt{-1}\hbar \frac{d}{dt} \hat{f} + \hat{f} E^{(\text{Im})}(\hat{w})$$

As a consequence of the imaginary geomenergy, we can say that imaginary geomenergy is a new kind of Hamiltonian operator.

## 2.5 The QCPB for quantum harmonic oscillator

In this section, we simply review some basic results given by the paper [3] in QCPB. As quantum mechanics illustrated, the commutator relations may look different than in the Schrödinger picture, because of the time dependence of operators. The time evolution of those operators depends on the Hamiltonian of the system. Considering the one-dimensional quantum harmonic oscillator,

$$\hat{H}^{(cl)} = \frac{\hat{p}^{(cl)2}}{2m} + \frac{m\omega^2 x^2}{2} \quad (4)$$

the evolution of the position and momentum operators is given by:

$$\begin{aligned} \frac{d}{dt}x(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(cl)}, x(t)]_{QPB} = \frac{\hat{p}^{(cl)}}{m} \\ \frac{d}{dt}\hat{p}^{(cl)}(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(cl)}, \hat{p}^{(cl)}(t)]_{QPB} = -m\omega^2 x \end{aligned} \quad (5)$$

As for the application of the QPB or the commutation, there are some many fields including the physics and mathematics, and so on.

As a certainly example, we will now start by briefly reviewing the quantum mechanics of a one-dimensional quantum harmonic oscillator (4), and see how the QCPB can be incorporated using GGC in the covariant quantization procedure. With the Hamiltonian given by (4), let's use the QCPB to recalculate the (5), we can see how difference emerges. More specifically, the covariant dynamics in terms of the position reads

$$\frac{\mathcal{D}}{dt}x(t) = \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(cl)}, x(t)] = \frac{\hat{p}^{(cl)}(t)}{m} + \frac{\sqrt{-1}}{\hbar} x(t) \frac{\hat{p}^{(cl)2}s + 2\hat{p}^{(cl)}s\hat{p}^{(cl)}}{2m}$$

By direct computation, the G-dynamics is given by

$$\hat{w}^{(cl)} = \frac{\sqrt{-1}}{\hbar} \frac{\hat{p}^{(cl)2}s + 2\hat{p}^{(cl)}s\hat{p}^{(cl)}}{2m} \quad (6)$$

And the generalized Heisenberg equation with respect to  $x$  follows

$$\begin{aligned} \frac{d}{dt}x(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(cl)}, x(t)]_{QPB} + \frac{\sqrt{-1}}{\hbar} \hat{H}^{(cl)}[s, x(t)]_{QPB} \\ &= \frac{\hat{p}^{(cl)}(t)}{m} \end{aligned}$$

In the same way, the covariant dynamics for the classical momentum operator is

$$\begin{aligned} \frac{\mathcal{D}}{dt}\hat{p}^{(cl)}(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(cl)}, \hat{p}^{(cl)}(t)] \\ &= -m\omega^2 x - \hat{H}^{(cl)} \frac{ds}{dx} + \hat{p}^{(cl)}(t) \hat{w}^{(cl)} \end{aligned}$$

Accordingly, the generalized Heisenberg equation in terms of the  $\hat{p}^{(cl)}$  appears

$$\begin{aligned} \frac{d}{dt}\hat{p}^{(cl)}(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(cl)}, \hat{p}^{(cl)}(t)]_{QPB} + \frac{\sqrt{-1}}{\hbar} \hat{H}^{(cl)}[s, \hat{p}^{(cl)}(t)]_{QPB} \\ &= -m\omega^2 x - \hat{H}^{(cl)} \frac{ds}{dx} \end{aligned}$$

## 2.6 Geomentum operator

**Definition 11.** [3] Let  $M$  be a smooth manifold represented by structural function  $s$ , then geomentum operator is defined as

$$\hat{p} = -\sqrt{-1}\hbar D$$

where  $D = \nabla + \nabla s$ . The component is  $\hat{p}_j = -\sqrt{-1}\hbar D_j$  in which  $D_j = \partial_j + \partial_j s$  holds,  $\partial_j = \frac{\partial}{\partial x_j}$ .

Note that the geomentum operator is a revision of the classical momentum operator given by (3).

**Theorem 2** (Geometric canonical quantization rules). [3] Geometric equal-time canonical commutation relation is

$$[\hat{x}_i, \hat{p}_j] = \sqrt{-1}\hbar D_j x_i$$

where  $[\cdot, \cdot] = [\cdot, \cdot]_{QPB} + G(s, \cdot, \cdot)$  is QCPB.

Geometric canonical commutation relation can be expressed in a specific form

$$[\hat{x}_i, \hat{p}_j] = \sqrt{-1}\hbar \left( \delta_{ij} + x_i \frac{\partial}{\partial x_j} s \right)$$

In other words, it also can be rewritten as  $[x_i, \hat{p}_j] = \sqrt{-1}\hbar \theta_{ij}$ , where  $\theta_{ij} = \delta_{ij} + x_i \partial_j s$ , and  $\partial_j = \frac{\partial}{\partial x_j}$ .

## 2.7 The geoperator

**Definition 12** (Geoperator). [4] Let  $\hat{X}$  be a Hermitian operator, then geoperator can be defined as

$$\hat{X}^{(s)} = \hat{X} + u$$

where  $u = \hat{X}s = I(\hat{X}, s)$  is coupling interaction between the observable  $\hat{X}$  and the environment  $s$ .

Obviously, the geometric operator is a extension of the Hermitian operator. Note that the interaction term  $u = \hat{X}s = I(\hat{X}, s) \neq 0$  actually is a precise function with respect to the spacetime, as we state, the structure function  $s$  generated by the space or manifolds as an environment variable only associate with the space or the manifolds, and it's independent to the wave function.

Note that the definition of the geoperator is reasonable, in particular, the evidence comes from the coupling interaction between the observable  $\hat{X}$  and the environment variable  $s$  that is described by  $u = \hat{X}s = I(\hat{X}, s)$  in which the environment variable  $s$  satisfies the G-dynamics in the theorem 1, it also can be seen from (6) in the one-dimensional quantum harmonic oscillator. In fact, geomentum operator is one of the geoperator if the Hermitian operator  $\hat{X} = \hat{p}^{(cl)} = -\sqrt{-1}\hbar \nabla$ .



### 3 G-dynamics

In this section, G-dynamics as a new dynamical form appears in the quantum mechanics, we mainly focus on the G-dynamics expressed by theorem 1, that is,  $\hat{w} = \frac{1}{\sqrt{-1}\hbar} \left[ s, \hat{H} \right]_{QPB}$ . As [3] already stated, the G-dynamics is only induced by the structure function  $s$ , and meanwhile, it's independent to the other observables. There is a clear fact of the G-dynamics taken as a different form by choosing different Hamiltonian operators, in one word, various Hamiltonian operators correspond to the multiple G-dynamics. Therefore, for a given wave function  $\psi$ , the eigenvalue equation of operator is accordingly given by  $\hat{w}\psi = w\psi$ , where  $w$  is the eigenvalue. By using definition 10, it gets corresponding imaginary geomenergy given by definition 10, that is,  $E^{(Im)} = \sqrt{-1}\hbar\hat{w}$ , thusly, it then has energy spectrum given by  $E^{(Im)}\psi = \sqrt{-1}E^{(g)}\psi$ , where  $E^{(g)} = \hbar w$ . Obviously, we can see that the G-dynamics is completely determined by the structure function  $s$ , it implies that the G-dynamics represents the properties of the spatial manifolds, in other words, the spatial manifolds has abundant activities. As a result of this point, we need to seek more clues to unlock this quantum characters. The classical Hamiltonian operator (3) or (2) is the first to bear the brunt to be considered in above formula.

#### 3.1 D-operator, T-operator and G-operator

In this section, the discussions on the most general form will be done, we use the G-dynamics  $\hat{w}$  to define the T-operator, and then the imaginary geomenergy is used with the classical Hamiltonian operator (3) to construct the G-operator as a non-Hermitian operator to deeply study some properties of the G-dynamics.

To start with the definition of the D-operator and the T-operator,

**Definition 13.** *The D-operator is given by  $\widehat{D}_t = \partial_t \pm \hat{w}$ , and then T-operator is defined as  $\widehat{N}_t = \sqrt{-1}\widehat{D}_t$ , where  $\hat{w}$  is the G-dynamics.*

Note that D-operator  $\widehat{D}_t = \partial_t \pm \hat{w}$  means that there are two kinds of D-operators. More precisely, we take one of them into consideration as reality needed. Its eigenvalue equation of D-operator is given by  $\widehat{D}_t\psi = D_t\psi$ .

As a result of the definition 13, that naturally leads to bond the imaginary geomenergy and the classical Hamiltonian operator (3) together to construct a new energy operator, accordingly, we can derive following G-operator.

**Definition 14.** *The G-operator can be defined as  $\widehat{H}^{(gr)} = \hbar\widehat{N}_t$ .*

Thusly, then the G-operator in details is given by

$$\begin{aligned} \widehat{H}^{(gr)} &= \widehat{H}^{(cl)} + E^{(Im)} = \widehat{H}^{(cl)} + \sqrt{-1}\hbar\hat{w} \\ &= \sqrt{-1}\hbar(\partial_t + \hat{w}) \\ &= \sqrt{-1}\hbar\widehat{D}_t \end{aligned}$$

That T-operator is then expressed as  $\widehat{H}^{(gr)}/\hbar = \widehat{N}_t$ . Thusly, the eigenvalue equation  $\widehat{N}_t\psi = N_t\psi$  follows. As we stated, G-dynamics  $\hat{w}$  is a Hermitian operator in the most of time, it

implies that the the G-operator is a non-Hermitian operator. As a result of this point, we say that the eigenvalue of the G-operator must be in a complex form.

**Theorem 3.** *The eigenvalue of G-operator is  $E^{(gr)} = \hbar N_t$ , where the eigenvalue of T-operator is*

$$N_t = E^{(gr)}/\hbar = E^{(cl)}/\hbar + \sqrt{-1}E^{(g)}/\hbar$$

where  $E^{(g)} = \hbar w$ .

*Proof.* According to the definition 14 of G-operator, the eigenvalue equation can be more specifically expressed below

$$\begin{aligned} \widehat{H}^{(gr)}\psi &= \widehat{H}^{(cl)}\psi + E^{(Im)}\psi = \widehat{H}^{(cl)}\psi + \sqrt{-1}\hbar\widehat{w}\psi \\ &= \sqrt{-1}\hbar(\partial_t + \widehat{w})\psi = \sqrt{-1}\hbar\widehat{D}_t\psi \\ &= \sqrt{-1}\hbar D_t\psi \\ &= \sqrt{-1}\hbar(\partial_t\psi + w\psi) \\ &= (E^{(cl)} + \sqrt{-1}\hbar w)\psi \\ &= E^{(gr)}\psi \end{aligned} \tag{7}$$

Hence, we obtain the eigenvalue of the G-operator that is given by  $E^{(gr)} = \sqrt{-1}\hbar D_t$ , it's rewritten as  $E^{(gr)} = E^{(cl)} + \sqrt{-1}E^{(g)}$  by denoting  $E^{(g)} = \hbar w$ , and the eigenvalue of the D-operator  $\widehat{D}_t$  follows

$$D_t = E^{(gr)}/\sqrt{-1}\hbar = E^{(cl)}/\sqrt{-1}\hbar + w$$

Similarly, the eigenvalue of the T-operator  $\widehat{N}_t$  also emerges as follows  $N_t = E^{(gr)}/\hbar$ , where  $E^{(g)}/\hbar = w$ , therefore, the eigenvalue of the operator  $\widehat{N}_t$  can be rewritten as

$$N_t = E^{(gr)}/\hbar = E^{(cl)}/\hbar + \sqrt{-1}E^{(g)}/\hbar$$

Note that the eigenvalue of the G-operator can be rewritten in a form  $E^{(gr)} = \hbar N_t$ . Therefore, we complete the proof as desired.  $\square$

Notice that the eigenvalue of G-operator and T-operator are complex form, these are the features of the non-Hermitian operator. As G-dynamics stated, it's generated by the structure function, and it forms the G-operator, it reveals that there exists a complete Schrödinger equation such that the structure function is naturally involved in the equation.

## 4 Geometrinetic energy operator (GEO) and Ri-operator

As a result of the G-operator, we use geomentum operator 11 to reconstruct the new Hamiltonian operator. In order to smoothly build such complete new Hamiltonian operator, we firstly need to define a corresponding kinetic energy operator associated with the structure function  $s$ , therefore, let's rewrite geomentum operator as  $\dot{P}^{(ri)} = -\sqrt{-1}\hbar D$  from definition 11, then we give the definition below based on the geomentum operator.

**Definition 15** (Geometrinetic energy operator (GEO)). *Geometrinetic energy operator (GEO) is defined as*

$$\hat{T}^{(ri)} = \frac{\hat{P}^{(ri)2}}{2m} = -\frac{\hbar^2}{2m} (\Delta + 2\nabla s \cdot \nabla) + T^{(c)}$$

where  $T^{(c)} = -\frac{\hbar^2}{2m} (\Delta s + \nabla s \cdot \nabla s)$  is called structural c-energy.

More precisely, the derivation of the geometrinetic energy operator can be given by logically computation. Since the square of generalized gradient operator that is calculated as

$$D^2\psi = \Delta\psi + 2\nabla s \cdot \nabla\psi + \psi (\Delta s + \nabla s \cdot \nabla s)$$

Accordingly, it yields an second order operator

$$D^2 = \Delta + 2\nabla s \cdot \nabla + (\Delta s + \nabla s \cdot \nabla s) \quad (8)$$

Then the eigenvalue equation of the geometrinetic energy operator  $\hat{T}^{(ri)}$  with respect to the wave function  $\psi$  follows

$$\hat{T}^{(ri)}\psi = \frac{\hat{P}^{(ri)2}}{2m}\psi = -\frac{\hbar^2}{2m} (\Delta\psi + 2\nabla s \cdot \nabla\psi) + \psi T^{(c)}$$

**Definition 16.** *Ri-operator is defined as*

$$\hat{H}^{(ri)} = \hat{T}^{(ri)} + V$$

where  $\hat{T}^{(ri)}$  is the geometrinetic energy operator.

As a result, Ri-operator can be rewritten in the form

$$\hat{H}^{(ri)} = \hat{H}^{(cl)} - \frac{\hbar^2}{m} \nabla s \cdot \nabla + T^{(c)} \quad (9)$$

by using the classical Hamiltonian operator (2).

## 4.1 The G-dynamics with Schrödinger equation

This section, we will mainly discuss the eigenvalue equation of G-dynamics based on the Schrödinger equation (1), it nicely proves that G-operator and T-operator are natural results on G-dynamics. Rewriting the G-dynamics that is originally proposed as

$$E^{(Im)}/\sqrt{-1}\hbar = \hat{w} = \frac{1}{\sqrt{-1}\hbar} [s, \hat{H}]_{QPB}$$

where the Hamiltonian operator  $\hat{H}$  in it is completely determined by the physical truth.

**Theorem 4.** *The eigenvalue of the G-dynamics  $\hat{w}^{(cl)}$  based on Schrödinger equation (1) in time parameter or Hamiltonian operator (3) is given by  $w^{(a)} = -\partial_t s$ .*

*Proof.* Let's take  $H = \widehat{H}^{(cl)}$ , then, more precisely,

$$\widehat{w}^{(cl)}\psi = \frac{1}{\sqrt{-1}\hbar} \left[ s, \widehat{H}^{(cl)} \right]_{QPB} \psi = \frac{1}{\sqrt{-1}\hbar} \left( s\widehat{H}^{(cl)}\psi - \widehat{H}^{(cl)}(s\psi) \right)$$

Plugging the Schrödinger equation (1) into above eigenvalue equation, we obtain

$$\sqrt{-1}\hbar\partial_t\psi = \widehat{H}^{(cl)}\psi, \quad \sqrt{-1}\hbar\partial_t(s\psi) = \widehat{H}^{(cl)}(s\psi)$$

and the eigenvalue equation of G-dynamics  $\widehat{w}^{(cl)}$  as an operator is

$$\begin{aligned} \widehat{w}^{(cl)}\psi &= \frac{1}{\sqrt{-1}\hbar} \left[ s, \widehat{H}^{(cl)} \right]_{QPB} \psi \\ &= \frac{1}{\sqrt{-1}\hbar} \left( s\sqrt{-1}\hbar\partial_t\psi - \sqrt{-1}\hbar\partial_t(s\psi) \right) \\ &= (s\partial_t\psi - \partial_t(s\psi)) \\ &= -\psi\partial_t s \end{aligned} \tag{10}$$

We denote  $\widehat{w}^{(cl)}\psi = w^{(q)}\psi$  here. Thusly, we certainly get the specific eigenvalue for  $\widehat{w}^{(cl)}$  given by  $w^{(q)} = -\partial_t s$ .  $\square$

Note that theorem 4 has indicated a fact that for any function form given for the G-dynamics  $\widehat{w}$  in such condition, the eigenvalue is always dependent on the structure function.

**Theorem 5.** *The heat equation for the structure function  $s$  based on the Schrödinger equation (1) is given by*

$$\partial_t s = \frac{\sqrt{-1}\hbar}{m} (\Delta s/2 + \nabla s \cdot \nabla \ln \psi)$$

*Proof.* In the one hand, theorem 4 has evidently given. In another hand, we take the (2) into the G-dynamics  $\widehat{w}$ , and it deduces a result

$$\begin{aligned} \widehat{w}^{(cl)}\psi &= \frac{1}{\sqrt{-1}\hbar} \left[ s, \widehat{H}^{(cl)} \right]_{QPB} \psi \\ &= \frac{1}{\sqrt{-1}\hbar} \left[ s, -\frac{\hbar^2}{2m}\Delta + V \right]_{QPB} \psi \\ &= -\frac{\hbar}{2\sqrt{-1}m} [s, \Delta]_{QPB} \psi \\ &= \frac{\hbar}{2\sqrt{-1}m} (2\nabla s \cdot \nabla + \Delta s) \psi \end{aligned}$$

According to (10), it yields

$$\widehat{w}^{(cl)}\psi = -\psi\partial_t s = \frac{\hbar}{2\sqrt{-1}m} (\Delta s + 2\nabla s \cdot \nabla) \psi$$

It's definite to form a heat equation associated with the structure function

$$\partial_t s = \frac{\sqrt{-1}\hbar}{2m\psi} (\psi\Delta s + 2\nabla s \cdot \nabla\psi) = \frac{\sqrt{-1}\hbar}{m} (\Delta s/2 + \nabla s \cdot \nabla \ln \psi)$$

Then, the proof is completed.  $\square$

**Corollary 1.** *The G-dynamics in terms of the Hamiltonian operator (2) in coordinates can be expressed as*

$$\hat{w}^{(cl)} = \frac{\hbar}{2\sqrt{-1}m} (\Delta s + 2\nabla s \cdot \nabla) \quad (11)$$

As we can see, the G-dynamics on such expression given by the (11) is totally determined by the structure function. Geometrinetic energy operator by using the G-dynamics  $\hat{w}^{(cl)}$  can be rewritten as

$$\hat{T}^{(ri)} = \hat{T}^{(cl)} - E^{(Im)} (\hat{w}^{(cl)}) - E^{(s)}/2$$

where the imaginary geomenergy is  $E^{(Im)} (\hat{w}^{(cl)}) = \sqrt{-1}\hbar\hat{w}^{(cl)}$ , and  $\hat{T}^{(cl)} = -\frac{\hbar^2}{2m}\Delta$ , and  $E^{(s)} = \frac{\hbar^2}{m}\nabla s \cdot \nabla s$ . As a consequence, based on the theorem 5, we can get a corollary shown as follows.

**Corollary 2.** *If wave function  $\psi$  satisfies both (1) and heat equation 5, then*

$$\begin{cases} \partial_t s = \frac{\sqrt{-1}\hbar}{m} (\Delta s/2 + \nabla s \cdot \nabla \ln \psi) \\ \sqrt{-1}\hbar\partial_t \psi = -\frac{\hbar^2}{2m}\Delta\psi + V\psi \end{cases}$$

Take a deep analysis to above two equations, obviously, it yields

$$\sqrt{-1}\hbar\psi\partial_t s = -\frac{\hbar^2}{2m} (\psi\Delta s + 2\nabla s \cdot \nabla\psi)$$

It follows the bonding equation

$$\sqrt{-1}\hbar (\partial_t \psi + \psi\partial_t s) = -\frac{\hbar^2}{2m} (\Delta\psi + 2\nabla s \cdot \nabla\psi + \psi\Delta s) + V\psi$$

Definitely, it can be rewritten as

$$\begin{aligned} & \sqrt{-1}\hbar (\partial_t + \partial_t s) \psi \\ &= -\frac{\hbar^2}{2m} (\Delta + 2\nabla s \cdot \nabla + \Delta s + \nabla s \cdot \nabla s) \psi \\ & \quad + \left( V + \frac{\hbar^2}{2m} \nabla s \cdot \nabla s \right) \psi \\ &= \hat{H}^{(ri)}\psi + \frac{\hbar^2}{2m}\psi\nabla s \cdot \nabla s \end{aligned}$$

Or in the form

$$\sqrt{-1}\hbar (\partial_t - \hat{w}^{(cl)}) \psi = \hat{H}^{(ri)}\psi + E^{(s)}\psi/2 \quad (12)$$

where  $E^{(s)} = \frac{\hbar^2}{m}\nabla s \cdot \nabla s$ , it evidently implies that  $E^{(s)}$  plays the role of potential energy. In fact, the (12) supports a statement to the definition 14. In other words, such definition 14 is a natural extension for the classical results, in the later discussions, we can surely say about this points.

**Definition 17.** *D-operator based on the Schrödinger equation (1) is  $\widehat{D}_t^{(cl)} = \partial_t - \hat{w}^{(cl)}$ .*

Actually, the D-operator also can be taken in another form

$$\widehat{D}_t^{(cl)} = \partial_t + \widehat{w}^{(cl)} \quad (13)$$

that is,  $-\widehat{w}^{(cl)} \rightarrow \widehat{w}^{(cl)}$  on above definition. According to the G-operator given by definition 14, thusly, it yields

**Definition 18.** *G-operator is then defined as*

$$\widehat{H}^{(gr)} = \hbar \widehat{N}_t^{(cl)} = \widehat{H}^{(cl)} - E^{(Im)}(\widehat{w}^{(cl)})$$

based on definition 17, where the imaginary geomenergy is  $E^{(Im)}(\widehat{w}^{(cl)}) = \sqrt{-1}\hbar\widehat{w}^{(cl)}$ .

As a consequence of the definition 14, we figure out that eigenvalue theorem below.

**Theorem 6.** *The eigenvalue of the G-operator in definition 18 on Schrödinger equation (1) is*

$$E^{(gr)} = E^{(cl)} - \sqrt{-1}\hbar w^{(q)}$$

*Proof.* For a given wave function  $\psi$ , the eigenvalue equation of the G-operator on Schrödinger equation (1) is expressed as

$$\begin{aligned} \widehat{H}^{(gr)}\psi &= \hbar \widehat{N}_t^{(cl)}\psi = \widehat{H}^{(cl)}\psi - E^{(Im)}(\widehat{w}^{(cl)})\psi \\ &= \widehat{H}^{(cl)}\psi - \sqrt{-1}\hbar\widehat{w}^{(cl)}\psi \\ &= E^{(cl)}\psi - \sqrt{-1}\hbar\psi w^{(q)} \\ &= (E^{(cl)} - \sqrt{-1}\hbar w^{(q)})\psi \\ &= E^{(gr)}\psi \end{aligned}$$

where  $\widehat{w}^{(cl)}\psi = w^{(q)}\psi$  has been used for the proof. Then, we complete the proof.  $\square$

In fact, if D-operator is taken according to (13), then eigenvalue of the G-operator on Schrödinger equation (1) is

$$E^{(gr)} = E^{(cl)} + \sqrt{-1}\hbar w^{(q)}$$

This is exactly the characters of conjugate complex numbers.

**Corollary 3.** *Based on theorem 6, there exists a pure imaginary geomenergy form  $E^{(Im)}(\pm w^{(q)}) = \pm\sqrt{-1}\hbar w^{(q)}$  induced by G-dynamics.*

## 5 The cases of G-dynamics

### 5.1 For classical Hamiltonian operator

To consider quantum harmonic oscillator (4), as the G-dynamics of it shown by (6),

$$\widehat{w}^{(cl)} = \frac{\sqrt{-1}}{\hbar} \frac{\widehat{p}^{(cl)2} s}{2m} + \frac{\sqrt{-1}}{m\hbar} \widehat{p}^{(cl)} s \widehat{p}^{(cl)}$$

The pure imaginary geomenergy form is

$$E^{(\text{Im})}(\hat{w}^{(cl)}) = \sqrt{-1}\hbar\hat{w}^{(cl)} = -\frac{\hat{p}^{(cl)2}s}{2m} - \frac{1}{m}\hat{p}^{(cl)}s\hat{p}^{(cl)}$$

Let's see how its characteristic equation shows, it's given by

$$\begin{aligned}\hat{w}^{(cl)}\psi &= \frac{\sqrt{-1}}{\hbar}\psi\frac{\hat{p}^{(cl)2}s}{2m} + \frac{\sqrt{-1}}{m\hbar}\hat{p}^{(cl)}s\hat{p}^{(cl)}\psi \\ &= -\frac{\sqrt{-1}\hbar}{2m}(\psi\Delta s + 2\nabla s \cdot \nabla\psi) \\ &= \frac{\hbar}{2\sqrt{-1}m}(\Delta s + 2\nabla s \cdot \nabla)\psi\end{aligned}$$

where  $\Delta = \nabla^2$ . Subsequently, it derives the G-dynamics (11), or in the form  $\frac{2\sqrt{-1}m}{\hbar}\hat{w}^{(cl)} = \Delta s + 2\nabla s \cdot \nabla$ , by using (8), we get its another expression in terms of the G-dynamics shown as

$$D^2 = \frac{2\sqrt{-1}m}{\hbar}\hat{w}^{(cl)} + \Delta + \nabla s \cdot \nabla s$$

## 5.2 For Ri-operator

**Theorem 7.** *The G-dynamics in  $\hat{H}^{(ri)}$  in coordinates is*

$$\hat{w}^{(ri)} = \hat{w}^{(cl)} + w^{(s)}$$

where  $w^{(s)} = \frac{\hbar}{\sqrt{-1}m}\nabla s \cdot \nabla s$  is denoted.

*Proof.* For this case, let's consider  $\hat{H}^{(ri)} = \hat{T}^{(ri)} + V$  given by Ri-operator 16, then the G-dynamics can be calculated as

$$\hat{w}^{(ri)} = \frac{1}{\sqrt{-1}\hbar}[s, \hat{H}^{(ri)}]_{QP B} = \frac{1}{\sqrt{-1}\hbar}[s, \hat{T}^{(ri)} + V]_{QP B}$$

where  $V$  is a real potential function. Thusly, we obtain

$$\begin{aligned}[s, \hat{T}^{(ri)}]_{QP B}\psi &= -\frac{\hbar^2}{2m}[s, \Delta + 2\nabla s \cdot \nabla]_{QP B}\psi \\ &= -\frac{\hbar^2}{2m}[s, \Delta]_{QP B}\psi - \frac{\hbar^2}{m}[s, \nabla s \cdot \nabla]_{QP B}\psi\end{aligned}$$

where

$$\begin{aligned}[s, \Delta]_{QP B}\psi &= s\Delta\psi - \Delta(s\psi) \\ &= -2\nabla s \cdot \nabla\psi - \psi\Delta s \\ &= -(2\nabla s \cdot \nabla + \Delta s)\psi\end{aligned}$$

and

$$[s, \nabla s \cdot \nabla]_{QPB} \psi = -\psi \nabla s \cdot \nabla s$$

Therefore, we get

$$\left[ s, \widehat{T}^{(ri)} \right]_{QPB} = \frac{\hbar^2}{2m} (2\nabla s \cdot \nabla + \Delta s + 2\nabla s \cdot \nabla s)$$

Thusly,

$$\begin{aligned} \widehat{w}^{(ri)} &= \frac{1}{\sqrt{-1}\hbar} \left[ s, \widehat{T}^{(ri)} \right]_{QPB} = \frac{\hbar}{2\sqrt{-1}m} (2\nabla s \cdot \nabla + \Delta s + 2\nabla s \cdot \nabla s) \\ &= \frac{\hbar}{2\sqrt{-1}m} (2\nabla s \cdot \nabla + \Delta s) + \frac{\hbar}{\sqrt{-1}m} \nabla s \cdot \nabla s \\ &= \widehat{w}^{(cl)} + w^{(s)} \end{aligned} \quad (14)$$

where  $w^{(s)} = \frac{\hbar}{\sqrt{-1}m} \nabla s \cdot \nabla s$  is denoted. □

Obviously,  $\widehat{w}^{(ri)} - \widehat{w}^{(cl)} = w^{(s)}$  holds. The theorem 7 can be rewritten in a form

$$E^{(\text{Im})}(\widehat{w}^{(ri)}) = \sqrt{-1}\hbar\widehat{w}^{(ri)} = \sqrt{-1}\hbar\widehat{w}^{(cl)} + E^{(s)}$$

where  $E^{(s)} = \frac{\hbar^2}{m} \nabla s \cdot \nabla s = \sqrt{-1}\hbar w^{(s)}$ , or expressed by the imaginary geomenergy

$$E^{(\text{Im})}(\widehat{w}^{(ri)}) = E^{(\text{Im})}(\widehat{w}^{(cl)}) + E^{(s)}$$

where  $E^{(\text{Im})}(\widehat{w}^{(cl)}) = \sqrt{-1}\hbar\widehat{w}^{(cl)}$  is the imaginary geomenergy in terms of  $\widehat{w}^{(cl)}$ . Correspondingly, the structural  $c$ -energy can be given in a form

$$\begin{aligned} T^{(c)} &= -\frac{\hbar^2}{2m} (\Delta s + |\nabla s|^2) = -\frac{\hbar^2}{2m} \Delta s - \frac{\hbar^2}{2m} |\nabla s|^2 \\ &= -\frac{\hbar^2}{2m} \Delta s - E^{(s)}/2 \end{aligned}$$

where it results in  $T^{(c)} + E^{(s)}/2 = -\frac{\hbar^2}{2m} \Delta s$ . There is more we need to say clearly,

$$\widehat{w}^{(ri)} = \widehat{w}^{(cl)} + w^{(s)} = \widehat{w}^{(cl)} - \sqrt{-1}a \nabla s \cdot \nabla s$$

where  $a = \hbar/m$ . Then the conjugate transpose of the  $\widehat{w}^{(ri)}$  is given by

$$\widehat{w}^{(ri)\dagger} = \widehat{w}^{(cl)\dagger} + \sqrt{-1}a \nabla s \cdot \nabla s$$

Accordingly, it yields  $\widehat{w}^{(ri)} + \widehat{w}^{(ri)\dagger} = \widehat{w}^{(cl)} + \widehat{w}^{(cl)\dagger} = 2\widehat{w}^{(cl)}$ . Meanwhile, let  $\widehat{w}^{(g)}$  be denoted as NG operator to fit

$$\widehat{w}^{(g)} = \widehat{w}^{(ri)} - \widehat{w}^{(ri)\dagger} = -2\sqrt{-1}a \nabla s \cdot \nabla s \neq 0$$

Obviously, this NG operator breaks the Hermiticity as usually understood



### 5.3 G-operator for Ri-operator

To analyze G-dynamics for both situations together, we can obtain some useful results.

$$\begin{aligned}\widehat{w}^{(ri)}\psi &= \widehat{w}^{(cl)}\psi + w^{(s)}\psi \\ &= \frac{\hbar}{2\sqrt{-1}m}(\Delta s + 2\nabla s \cdot \nabla)\psi + \frac{\hbar}{\sqrt{-1}m}\psi\nabla s \cdot \nabla s\end{aligned}$$

Accordingly, it leads to the imaginary geomenergy

$$\sqrt{-1}\hbar\widehat{w}^{(ri)}\psi = \frac{\hbar^2}{2m}(2\nabla s \cdot \nabla + \Delta s + 2\nabla s \cdot \nabla s)\psi$$

and with the Schrödinger equation (1), it generates

$$\begin{aligned}\sqrt{-1}\hbar\partial_t\psi - \sqrt{-1}\hbar\widehat{w}^{(ri)}\psi &= \sqrt{-1}\hbar(\partial_t - \widehat{w}^{(ri)})\psi \\ &= -\frac{\hbar^2}{2m}\Delta\psi + V\psi - \frac{\hbar^2}{2m}(2\nabla s \cdot \nabla + \Delta s + 2\nabla s \cdot \nabla s)\psi \\ &= -\frac{\hbar^2}{2m}(\Delta + 2\nabla s \cdot \nabla + \Delta s + \nabla s \cdot \nabla s)\psi + (V - E^{(s)}/2)\psi \\ &= \widehat{H}^{(ri)}\psi - E^{(s)}\psi/2\end{aligned}$$

As a result of equation (12) together, we get a similar equations given by

$$\begin{aligned}\sqrt{-1}\hbar(\partial_t - \widehat{w}^{(cl)})\psi &= \widehat{H}^{(ri)}\psi + E^{(s)}\psi/2 \\ \sqrt{-1}\hbar(\partial_t - \widehat{w}^{(ri)})\psi &= \widehat{H}^{(ri)}\psi - E^{(s)}\psi/2\end{aligned}\tag{15}$$

And then it rightly reflects the theorem 7.

**Corollary 4.** *Ri-operator expressed by the G-operator is given by*

$$\widehat{H}^{(ri)} = \widehat{H}^{(gr)} + E^{(s)}/2$$

*Proof.* Since the (9) and theorem 7 hold, then

$$\begin{aligned}\widehat{H}^{(ri)} &= \widehat{H}^{(cl)} - \frac{\hbar^2}{m}\nabla s \cdot \nabla + T^{(c)} \\ &= \widehat{H}^{(cl)} - \sqrt{-1}\hbar\widehat{w}^{(ri)} + E^{(s)}/2\end{aligned}$$

With the definition 14 of G-operator, it has

$$\widehat{H}^{(gr)} = \widehat{H}^{(cl)} - \sqrt{-1}\hbar\widehat{w}^{(ri)} = \sqrt{-1}\hbar\widehat{D}_t^{(ri)}$$

where D-operator here is  $\widehat{D}_t^{(ri)} = \partial_t - \widehat{w}^{(ri)}$ , thus, then we complete the proof.  $\square$

Conversely, note that the G-operator in corollary 4 is also expressed as

$$\widehat{H}^{(gr)} = \widehat{H}^{(ri)} - E^{(s)}/2\tag{16}$$

By using Ri-operator in corollary 4, (15) can be simplified as

$$\begin{aligned}\sqrt{-1}\hbar(\partial_t - \hat{w}^{(cl)})\psi &= \hat{H}^{(gr)}\psi + E^{(s)}\psi \\ \sqrt{-1}\hbar(\partial_t - \hat{w}^{(ri)})\psi &= \hat{H}^{(gr)}\psi\end{aligned}\tag{17}$$

where G-operator in terms of the Ri-operator is definitely chosen as

$$\hat{H}^{(gr)} = \sqrt{-1}\hbar(\partial_t - \hat{w}^{(ri)}) = \sqrt{-1}\hbar\hat{D}_t^{(ri)}$$

and  $\hat{D}_t^{(ri)} = \partial_t - \hat{w}^{(ri)}$ . As a matter of fact, these two equations are equivalent to each other based on the theorem 7.

## 5.4 Generalized covariant wave equation

Correspondingly, the definition of generalized covariant wave equation as a revision of classic Schrödinger equation follows based on geometrinetic energy operator.

**Theorem 8.** *The generalized covariant wave equation is defined as*

$$\hbar\hat{N}_t^{(ri)}\psi = \hat{H}^{(gr)}\psi$$

by using (16), where the D-operator is taken as  $\hat{D}_t^{(ri)} = \partial_t - \hat{w}^{(ri)}$ .

Apparently, the generalized covariant wave equation can be given by employing (17),

$$\sqrt{-1}\hbar(\partial_t - \hat{w}^{(cl)} - E^{(s)}/\sqrt{-1}\hbar)\psi = \hat{H}^{(gr)}\psi$$

where  $\hat{w}^{(ri)} = \hat{w}^{(cl)} + E^{(s)}/\sqrt{-1}\hbar$  is distinct to be seen from the theorem 7. More precisely about the generalized covariant wave equation are listed

$$\begin{aligned}\hat{H}^{(gr)} &= -\frac{\hbar^2}{2m}(\Delta + 2\nabla s \cdot \nabla + \Delta s + 2\nabla s \cdot \nabla s) = \hat{H}^{(ri)} - E^{(s)}/2 \\ \hat{w}^{(ri)} &= \frac{\hbar}{2\sqrt{-1}m}(2\nabla s \cdot \nabla + \Delta s + 2\nabla s \cdot \nabla s) = \hat{w}^{(cl)} + E^{(s)}/\sqrt{-1}\hbar\end{aligned}$$

Obviously, the generalized covariant wave equation is a natural complete extension of the classic Schrödinger equation.

Supported by the definition 12, we consider geoperator in terms of the Hamiltonian  $\hat{H}$ , let's see what we get in this way. Thusly, we have  $\hat{H}^{(s)} = \hat{H} + u_H$  that can be called the geomenergy operator, where  $u_H = \hat{H}s$  is coupling interaction between the observable  $\hat{H}$  and the environment  $s$ . In order to better understand this case, we precisely consider the classical Hamiltonian operator (3)  $\hat{H}^{(cl)}$  for the geomenergy operator that is expressed as  $\hat{H}^{(s)} = \hat{H}^{(cl)} + u_{H^{(cl)}}$ , where  $u_{H^{(cl)}} = \hat{H}^{(cl)}s$ , more specifically, the coupling interaction for the classical Hamiltonian operator (3)  $\hat{H}^{(cl)}$  and environment is given by  $u_{H^{(cl)}} = \hat{H}^{(cl)}s = -E^{(Im)}(w^{(q)})$ , then,  $E^{(Im)}(w^{(q)}) = -u_{H^{(cl)}}$ , as a result, the geomenergy operator can be rewritten as  $\hat{H}^{(s)} = \hat{H}^{(cl)} - E^{(Im)}(w^{(q)})$ , when we refer to the (12), we can obtain self-consistent result  $\hat{H}^{(s)}\psi = \sqrt{-1}\hbar(\partial_t - \hat{w}^{(cl)})\psi$  for a given wave function  $\psi$ , that is to say,  $\hat{H}^{(s)}\psi = \hat{H}^{(ri)}\psi + E^{(s)}\psi/2$  holds, accordingly, it yields  $\hat{H}^{(s)}\psi - E^{(s)}\psi/2 = \hat{H}^{(ri)}\psi$  which is exactly the generalized covariant wave equation as theorem 8 given.

## 6 GEO for generalized quantum harmonic oscillator

The covariant time evolution of operators  $x(t), \hat{p}^{(ri)}(t)$  depends on the Hamiltonian of the system. Considering the one-dimensional generalized quantum harmonic oscillator based upon geomentum operator 11, it's given by

$$\hat{H}^{(ri)} = \frac{\hat{p}^{(ri)2}}{2m} + \frac{m\omega^2 x^2}{2} \quad (18)$$

where  $V(x) = \frac{1}{2}m\omega^2 x^2$ , the geomentum operator in one-dimensional is  $\hat{p}^{(ri)} = -\sqrt{-1}\hbar \frac{D}{dx}$ , and  $\frac{D}{dx} = d/dx + ds/dx$ . Accordingly, the covariant evolution of the position and geomentum operator based on the QCHS given by definition 7 are respectively given by:

$$\begin{aligned} \frac{D}{dt}x(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(ri)}, x(t)] \\ \frac{D}{dt}\hat{p}^{(ri)}(t) &= \frac{\sqrt{-1}}{\hbar} [\hat{H}^{(ri)}, \hat{p}^{(ri)}(t)] \end{aligned} \quad (19)$$

where  $[\cdot, \cdot]$  denotes the GGC of two operators. More precisely, the concrete computational process of the first covariant evolution terms of the position is shown as

$$\left[ \hat{H}^{(ri)}, x \right] = \left[ \hat{H}^{(ri)}, x \right]_{QPB} + G\left(s, \hat{H}^{(ri)}, x\right)$$

Hence, the direct computation leads to a series of results

$$\begin{aligned} \left[ \hat{H}^{(ri)}, x \right]_{QPB} &= \left[ \frac{\hat{p}^{(ri)2}}{2m}, x \right]_{QPB} = \frac{1}{2m} [\hat{p}^{(ri)2}, x]_{QPB} \\ &= \frac{-\hbar^2}{2m} \left[ \frac{d^2}{dx^2} + 2\frac{ds}{dx} \frac{d}{dx}, x \right]_{QPB} \\ &= \frac{-\hbar^2}{m} \frac{D}{dx} \end{aligned}$$

where

$$\hat{p}^{(ri)2} = -\hbar^2 \left( \frac{d^2}{dx^2} + 2\frac{ds}{dx} \frac{d}{dx} + \frac{d^2 s}{dx^2} + \left( \frac{ds}{dx} \right)^2 \right)$$

and the QGB in terms of  $\hat{H}^{(ri)}, x$  is

$$\begin{aligned} G\left(s, \hat{H}^{(ri)}, x\right) &= -x \left[ s, \hat{H}^{(ri)} \right]_{QPB} = x \left[ \hat{H}^{(ri)}, s \right]_{QPB} \\ &= -\frac{\hbar^2}{m} x \frac{ds}{dx} \frac{d}{dx} - \frac{\hbar^2}{2m} x \left( \frac{d^2}{dx^2} s + 2 \left( \frac{ds}{dx} \right)^2 \right) \end{aligned}$$

Then the QCPB about  $\hat{H}^{(ri)}, x$  specifically shows

$$\left[ \hat{H}^{(ri)}, x \right] = \left[ \hat{H}^{(ri)}, x \right]_{QPB} + G\left(s, \hat{H}^{(ri)}, x\right)$$

$$\begin{aligned}
&= -\frac{\hbar^2}{m} \frac{D}{dx} - \frac{\hbar^2}{m} x \frac{ds}{dx} \frac{d}{dx} - \frac{\hbar^2}{2m} x \left( \frac{d^2}{dx^2} s + 2 \left( \frac{ds}{dx} \right)^2 \right) \\
&= -\frac{\hbar^2}{m} \left( \frac{D}{dx} + x \frac{ds}{dx} \frac{d}{dx} + \frac{1}{2} x \left( \frac{d^2}{dx^2} s + 2 \left( \frac{ds}{dx} \right)^2 \right) \right)
\end{aligned}$$

As a result, the covariant evolution of the position is given by

$$\begin{aligned}
\frac{\mathcal{D}}{dt} x(t) &= \frac{\sqrt{-1}}{\hbar} [\widehat{H}^{(ri)}, x] \\
&= -\frac{\sqrt{-1}\hbar}{m} \left( \frac{D}{dx} + x \frac{ds}{dx} \frac{d}{dx} + \frac{1}{2} x \left( \frac{d^2}{dx^2} s + 2 \left( \frac{ds}{dx} \right)^2 \right) \right)
\end{aligned}$$

In other words, the covariant evolution of the position based on the QCHS is

$$\frac{\mathcal{D}}{dt} x(t) = \frac{\sqrt{-1}}{\hbar} [\widehat{H}^{(ri)}, x] = \frac{\widehat{p}^{(ri)}}{m} + x\widehat{w}^{(ri)}$$

where G-dynamics is equal to

$$\widehat{w}^{(ri)} = -\frac{\sqrt{-1}\hbar}{2m} \left( 2 \frac{ds}{dx} \frac{d}{dx} + \frac{d^2}{dx^2} s + 2 \left( \frac{ds}{dx} \right)^2 \right) \quad (20)$$

Definitely, G-dynamics (20) in terms of  $\widehat{w}^{(ri)}$  is the equivalent expression of the (14) in one dimensional case. Accordingly, the imaginary geomenergy follows

$$E^{(\text{Im})}(\widehat{w}^{(ri)}) = \sqrt{-1}\hbar\widehat{w}^{(ri)} = \frac{\hbar^2}{2m} \left( 2 \frac{ds}{dx} \frac{d}{dx} + \frac{d^2}{dx^2} s + 2 \left( \frac{ds}{dx} \right)^2 \right)$$

Similarly, the covariant evolution of the geomomentum operator is

$$\frac{\mathcal{D}}{dt} \widehat{p}^{(ri)}(t) = \frac{\sqrt{-1}}{\hbar} [\widehat{H}^{(ri)}, \widehat{p}^{(ri)}(t)]$$

The calculation process is as follows

$$\begin{aligned}
[\widehat{H}^{(ri)}, \widehat{p}^{(ri)}]_{QPB} &= [V, \widehat{p}^{(ri)}]_{QPB} = \frac{1}{2} m \omega^2 [x^2, \widehat{p}^{(ri)}]_{QPB} \\
&= \frac{1}{2} m \omega^2 [x^2, \widehat{p}^{(ri)}]_{QPB} = \sqrt{-1}\hbar m \omega^2 x
\end{aligned}$$

The QGB in terms of  $\widehat{H}^{(ri)}, \widehat{p}^{(ri)}$  is

$$G(s, \widehat{H}^{(ri)}, \widehat{p}^{(ri)}) = \widehat{H}^{(ri)} [s, \widehat{p}^{(ri)}]_{QPB} - \widehat{p}^{(ri)} [s, \widehat{H}^{(ri)}]_{QPB}$$

where  $[s, \widehat{p}^{(ri)}]_{QPB} = \sqrt{-1}\hbar \frac{ds}{dx}$  has been used, and the QGB in details becomes

$$G(s, \widehat{H}^{(ri)}, \widehat{p}^{(ri)}) = \sqrt{-1}\hbar \widehat{H}^{(ri)} \frac{ds}{dx} - \widehat{p}^{(ri)} [s, \widehat{H}^{(ri)}]_{QPB}$$

As a consequence, the QCPB here is given by

$$\begin{aligned} [\widehat{H}^{(ri)}, \widehat{p}^{(ri)}] &= [\widehat{H}^{(ri)}, \widehat{p}^{(ri)}]_{QPB} + G\left(s, \widehat{H}^{(ri)}, \widehat{p}^{(ri)}\right) \\ &= \sqrt{-1}\hbar m\omega^2 x + \sqrt{-1}\hbar \widehat{H}^{(ri)} \frac{ds}{dx} - \widehat{p}^{(ri)} \left[ s, \widehat{H}^{(ri)} \right]_{QPB} \end{aligned}$$

It turns out that the covariant evolution of the geomentum operator is derived as follows

$$\begin{aligned} \frac{\mathcal{D}}{dt} \widehat{p}^{(ri)} &= \frac{\sqrt{-1}}{\hbar} [\widehat{H}^{(ri)}, \widehat{p}^{(ri)}] \\ &= -m\omega^2 x - \widehat{H}^{(ri)} \frac{ds}{dx} - \frac{\sqrt{-1}}{\hbar} \widehat{p}^{(ri)} \left[ s, \widehat{H}^{(ri)} \right]_{QPB} \end{aligned}$$

where the G-dynamics reads

$$\widehat{w}^{(ri)} = -\frac{\sqrt{-1}}{\hbar} \left[ s, \widehat{H}^{(ri)} \right]_{QPB}$$

which is equivalent to the (20) as a precise expression. The G-dynamics seen from the (20) can evidently say how it forms and works associated with the structure function.

## 7 Conclusions

In this paper, we have proposed a series of new concepts based on the analysis of the G-dynamics included in the covariant dynamics defined by the QCHS in QCPB. On the solid foundation of the QCPB, we have gained some valuable consequences centred on the G-dynamics. By using the G-dynamics, we define new operators such as the D-operator, T-operator, G-operator, Ri-operator etc, we find their complex connections, in particular, we mainly focus on the their eigenvalues equation. With the help of the Schrödinger equation, we analyze the G-dynamics and its related properties. We concretely discuss the different cases of the G-dynamics, in this way, we present the extension of the Schrödinger equation, naturally. As an implication, we derive the generalized quantum harmonic oscillator and the covariant dynamics on it based on the QCHS theory.

## References

- [1] Greiner W. Relativistic quantum mechanics. wave equations [M]. Beijing: World Book Inc, 2003.
- [2] Nikolenko N V. On the complete integrability of the nonlinear Schrödinger equation [J]. Functional Analysis and Its Applications, 1976, 10(3): 209-220.
- [3] Wang G. Generalized geometric commutator theory and quantum geometric bracket and its uses [J]. arXiv:2001.08566.
- [4] Wang G. A revision for Heisenberg uncertainty relation based on environment variable in the QCPB theory [J]. arXiv:2003.07203.