# Masses and Decay widths of Charmonium states in presence of strong magnetic fields

Amruta Mishra<sup>\*</sup>

Department of Physics, Indian Institute of Technology, Delhi, Hauz Khas, New Delhi – 110 016, India

S.P. Misra<sup>†</sup>

Institute of Physics, Bhubaneswar - 751005, India

# Abstract

The masses and decay widths of charmonium states are studied in the presence of strong magnetic fields. The mixing between the pseudoscalar and vector charmonium states at rest is observed to lead to appreciable negative (positive) shifts in the masses of the pseudoscalar (longitudinal component of the vector) charmonium states in vacuum/hadronic medium in the presence of high magnetic fields. The pseudoscalar and vector charmonium masses in the hadronic medium, calculated in an effective chiral model from the medium changes of a scalar dilaton field, have additional significant modifications due to the mixing effects. The masses of the D and D mesons in the magnetized hadronic matter are calculated within the chiral effective model. The partial decay widths of the vector charmonium state to DD are computed using a field theoretical model for composite hadrons with quark/antiquark constituents, and are compared to the decay widths calculated using an effective hadronic Lagrangian. The effects of the mixing are observed to lead to significant contributions to the masses of the pseusoscalar and vector charmonium states, and an appreciable increase in the decay width  $\psi(3770) \rightarrow D\bar{D}$  at large values of the magnetic fields. These studies of the charmonium states in strong magnetic fields should have observable consequences on the dilepton spectra, as well as on the production of the open charm mesons and the charmonium states in ultra relativistic heavy ion collision experiments.

<sup>\*</sup>Electronic address: amruta@physics.iitd.ac.in

<sup>&</sup>lt;sup>†</sup>Electronic address: misrasibaprasad@gmail.com

#### I. INTRODUCTION

The study of properties of hadrons under extreme conditions, e.g., high temperatures and/or densities is a topic of intense research in strong interaction physics. This is due to its relevance in the ultra relativistic heavy ion collision experiments at various high energy particle accelerators, as well as in the study of the bulk matter of astrophysical objects, e.g., neutron stars. In the recent past, there have been a lot of studies on the in-medium properties of the heavy flavour mesons [1], as these can have observable consequences in high energy heavy ion collision experiments. The heavy quarkonium (charmonium and bottomonium) states have been studied using the potential models [2-6], the QCD sum rule approach [7-10], the Quark Meson cooupling (QMC) model [11], the coupled channel approach [12], and a chiral effective model [13, 14]. The heavy quarkonium (charmonium and bottomonium) state in the presence of a gluon field has been studied in Refs. [15-17]. Assuming the heavy quark (Q) and antiquark  $(\bar{Q})$  in the quarkonium state to be bound by color Coulomb potential, and the  $Q\bar{Q}$  separation to be small compared to the scale of the gluonic fluctuations, the mass shift in the quarkonium state in the leading order is observed to be proportional to to the medium modifications of the scalar gluon condensate. Using the leading order formula and the linear density approximation for the gluon condensate in the nuclear medium, the mass modifications of the charmonium states have been studied in Ref. [18]. The in-medium masses of the charmonium states have been computed within a chiral effective model [13, 14], from the medium change of a dilaton scalar field, which simulates the gluon condensate within the hadronic model.

The magnetic fields created in non-central ultra relativistic heavy ion collision experiments, have been estimated to be huge, e.g.,  $eB \sim 2m_{\pi}^2$  at Relativistic Heavy Ion Collider (RHIC) at BNL and  $eB \sim 15m_{\pi}^2$  at Large Hadron Collider (LHC) at CERN [19]. This has led to a lot of work on the study of effects of strong magnetic fields on the properties of the hadrons. The strong magnetic fields produced in the high energy heavy ion collisons rapidly drop after the collision. This leads to induced currents, which slow down the decrease in the magnetic field. The time evolution of the magnetic field [19] is still an open question, and needs the solutions of the magnetohydrodynamic equations, with a proper estimate of the electrical conductivity of the medium. In the (near) central collisions, the impact parameter is small, the magnetic field produced is weak and the created medium is dense. On the other hand, strong magnetic fields

are created in the peripheral ultra relativistic heavy ion collisions and the formed medium has low density. The effects of the magnetic fields on the heavy quarkonium (charmonium and bottomonium) states can have observable consequences as these are formed in heavy ion collisions at the early stage, when the magnetic fields can still be large. The open charm mesons [20-23] as well as the charmonium states [24-26] have been studied in the presence of magnetic fields. There is mixing of the pseudoscalar and vector mesons in the presence of magnetic fields, which modifies the properties of the charmonium states [25–28]. The masses of the charmonium states in the presence of magnetic fields have been studied in a consistent manner using QCD sum rule approach, incorporating the mixing of the pseudoscalar and vector charmonium states in the hadron spectral function on the phenomenological side, as well as, including the effects of magnetic field on the OPE (operator product expansion) side [25, 26]. The QCD sum rule approach [25, 26], as well as, a study of the charm-anticharm bound state described by an effective potential and solving the Schrödinger equation in the presence of an external magnetic field [27] show that the charmonium masses have dominant contributions from the mixing effects. A study of the mixing effects on the formation time of the charmonia are observed to lead to delayed (faster) formation time of the vector (pseudoscalar) charmonium states [28]. The  $J/\psi - \eta_c$  as well as  $\psi' - \eta'_c$  mixings and the faster formation time of the pseudoscalar mesons, might show as peaks in the dilepton spectra, due to 'anomalous' decay modes, e.g.,  $\eta_c, \eta'_c \to l^+ l^-$  and can act as a probe of the existence of strong magnetic field at the early stage [28].

The masses of vector charmonium states in a (magnetized) hadronic medium have been studied using a chiral effective model [13, 14, 24]. These are calculated from the medium modification of a scalar dilaton field which mimics the gluon condensates of QCD in the effective hadronic model. Within the chiral effective model, the modifications of the masses of the open charm  $(D, \bar{D})$  mesons in the (magnetized) hadronic medium arise from their interactions with the baryons and scalar mesons, and have been studied in Refs. [13, 14, 23]. The in-medium decay widths of the vector charmonium states to  $D\bar{D}$  have been computed from the mass modifications of the  $D, \bar{D}$  and charmonium states in Refs. [13, 14, 29, 30] using a light quark pair creation model, namely the  ${}^{3}P_{0}$  model [31–34], as well as using a field theoretical model with composite hadrons with quark/antiquark constituents [35]. In the  ${}^{3}P_{0}$  model, the decay

of the charmonium state to  $D\bar{D}$  proceeds with creation of a light quark antiquark pair in  ${}^{3}P_{0}$  state and the light quark (antiquark) combines with the charm antiquark (quark) of the decaying charmonium state to form the D and  $\overline{D}$  mesons [29, 32, 34]. The light quark pair creation model [32] describes the observed branching ratios of the decay of  $\psi(4040)$  to DD,  $(D\bar{D}^* + \bar{D}D^*)$ , and  $D^*\bar{D}^*$ , which can not be explained using a naive quark spin counting. The  ${}^{3}P_{0}$  model, where the internal structure of the charmonium state as well as the open charm mesons are taken into account, is observed to lead to strong suppression of the decay modes of  $\psi(4040)$  to  $D\bar{D}$  as well as  $(D\bar{D}^* + \bar{D}D^*)$ , as compared to the final state  $D^*\bar{D}^*$ , as has been observed experimentally. The matrix element for the decay of charmonium state to the open charm mesons, depends on the magnitude of the momentum,  $|\mathbf{p}|$  of the outgoing meson, which is given in terms of the masses of the decaying and the produced particles. The decay amplitude for  $\psi(4040)$  in the  ${}^{3}P_{0}$  model is observed to be extremely small for the value of  $|\mathbf{p}|$  corresponding to the decay modes  $D\bar{D}$  and  $(D\bar{D}^* + \bar{D}D^*)$ , as compared to the final state  $D^*\bar{D}^*$  [32] and leads to good agreement of the experimentally observed branching ratios of these modes. The medium dependence of the decay width of the charmonium state to  $D\bar{D}$  was studied using the  ${}^{3}P_{0}$  model in Ref. [29] from the mass modifications of the D and  $\overline{D}$  mesons, which in turn modify the magnitude of the momentum of the D(D) meson. The decay widths were computed assuming harmonic oscillator wave functions for the charmonium as well as Dand D mesons [34]. For certain values of the momentum of the D(D) meson, the decay widths were observed to vanish, and the dependence on  $|\mathbf{p}|$  was observed to be drastically different from the results obtained when the internal structure of the mesons are not taken into account [29]. The open flavour strong decays in vacuum have been studied extensively in the light sector [31, 33] as well as heavy flavour mesons [32, 34] accounting for the quark (antiquark) substructure of the hadrons, using the light quark aniquark pair creation in the  ${}^{3}P_{0}$  model. The in-medium decay widths of the charmonium states to  $D\bar{D}$  in hadronic matter have been studied from the mass modifications of the D and  $\overline{D}$  mesons as well as of the charmonium states calculated using a chiral effective model [13, 14, 35]. In the present work, the effects of the magnetic field on the masses of the charmonium states are investigated accounting for the mixing of the pseudoscalar and vector mesons, and, their in-medium decay widths are studied using a field theoretic model of composite hadrons [35] and compared with the results obtained using an effective hadronic model.

In the present work, the masses of the pseudoscalar ( $\eta_c \equiv \eta_c(1S)$  and  $\eta'_c \equiv \eta_c(2S)$ ) as well as the vector charmonium states  $(J/\psi, \psi(2S) \equiv \psi(3686)$  and  $\psi(1D) \equiv \psi(3770)$ ) in vacuum/hadronic matter are computed accounting for the mixing of the pseudoscalar and the vector charmonium states in the presence of strong magnetic fields. In the hadronic medium, the charmonium masses are calculated within the chiral effective model in the presence of a magnetic field [24], with additional contributions from the mixing effects. The mixing is observed to lead to significant modifications to these charmonium masses. The decay widths of the charmonium states to the open charm mesons  $(D\bar{D})$  are computed using a field theoretical model of composite hadrons with quark/antiquark constituents [35]. These decay widths are compared with the results obtained from an effective hadronic model.

The outline of the paper is as follows. In section II, we briefly describe the chiral effective model used to compute the masses of the (pseudoscalar and vector) charmonium states as well as the open charm meson masses in the magnetized matter. In the presence of a magnetic field, the mixings between the pseudoscalar and the vector mesons are taken into account through an effective hadronic interaction. In section III, we describe the decay of the charmonium states to  $D\bar{D}$  using a model for composite hadrons as well as an effective hadronic model. In subsection A, we describe the field theoretical model with composite hadrons with quark/antiquark constituents. This model is used in the present work to compute the partial decay widths of the charmonium states to  $D\bar{D}$  in vacuum/hadronic matter in the presence of strong magnetic fields. The charmonium decay widths to  $D\bar{D}$  as calculated using the field theoretic model for composite hadrons are compared to the results obtained from an effective hadronic Lagrangian as described in subsection B. In section IV, we discuss the results obtained in the present investigation of the charmonium masses as well as charmonium decay widths, accounting for the mixing of the pseudoscalar and vector charmonium states in the presence of strong magnetic fields. In section V, we summarize the findings of the present study.

## **II. CHARMONIUM MASSES IN STRONG MAGNETIC FIELDS**

In this section, we investigate the mass modifications of the vector and pseudoscalar charmonium masses in the presence of strong magnetic fields. The masses of these heavy charm quarkantiquark bound states are calculated in the magnetized nuclear matter from the in-medium gluon condensates, which are simulated by a scalar dilaton field within a chiral effective model. The mixing of the pseudoscalar and vector charmonium states in the presence of an external magnetic field is studied using an effective Lagrangian interaction term [26] and is observed to be the dominant contribution to the mass shifts for these charmonium states.

The Lagrangian density of the chiral effective model, in the presence of a magnetic field, is given as [36]

$$\mathcal{L} = \mathcal{L}_{kin} + \sum_{W} \mathcal{L}_{BW} + \mathcal{L}_{vec} + \mathcal{L}_{0} + \mathcal{L}_{scalebreak} + \mathcal{L}_{SB} + \mathcal{L}_{mag}^{B\gamma},$$
(1)

where,  $\mathcal{L}_{kin}$  corresponds to the kinetic energy terms of the baryons and the mesons,  $\mathcal{L}_{BW}$ contains the interactions of the baryons with the meson, W (scalar, pseudoscalar, vector, axialvector meson),  $\mathcal{L}_{vec}$  describes the dynamical mass generation of the vector mesons via couplings to the scalar fields and contains additionally quartic self-interactions of the vector fields,  $\mathcal{L}_0$  contains the meson-meson interaction terms,  $\mathcal{L}_{scalebreak}$  is a scale invariance breaking logarithmic potential given in terms of a scalar dilaton field,  $\chi$  and  $\mathcal{L}_{SB}$  describes the explicit chiral symmetry breaking. The term  $\mathcal{L}_{mag}^{B\gamma}$ , describes the interaction of the baryons with the electromagnetic field, which includes a tensorial interaction ~  $\bar{\psi}_i \sigma^{\mu\nu} F_{\mu\nu} \psi_i$ , whose coefficients account for the anomalous magnetic moments of the baryons [23].

The trace of the energy momentum tensor of QCD is equated to that of the chiral effective model to obtain a relation between the scalar gluon condensate  $\left\langle \frac{\alpha_s}{\pi} G^a_{\mu\nu} G^{\mu\nu a} \right\rangle$  and the dilaton field  $\chi$  of the the scale breaking term  $\mathcal{L}_{scalebreak}$ . The in-medium masses of the charmonium states are hence obtained from the medium changes of the dilaton field. The dilaton field,  $\chi$ is solved along with the scalar fields (non-strange scalar-isoscalar, field  $\sigma$ , non-strange scalar isovector,  $\delta$ , the strange field  $\zeta$ ) in the magnetized hadronic matter, from the coupled equations of motion of these fields. The values of the scalar fields are used to obtain the in-medium masses of the D and  $\overline{D}$  mesons as well as the charmonium states [23, 24, 30].

The mass shifts of the (pseudoscalar and vector) charmonium states, computed from the medium change of the dilaton field calculated within the chiral effective model are given as [13, 14]

$$\Delta m_{P,V} = \frac{4}{81} (1-d) \int d|\mathbf{k}|^2 \langle |\frac{\partial \psi(\mathbf{k})}{\partial \mathbf{k}}|^2 \rangle \frac{|\mathbf{k}|}{|\mathbf{k}|^2/m_c + \epsilon} \left(\chi^4 - \chi_0^4\right),\tag{2}$$

where

$$\langle |\frac{\partial \psi(\mathbf{k})}{\partial \mathbf{k}}|^2 \rangle = \frac{1}{4\pi} \int |\frac{\partial \psi(\mathbf{k})}{\partial \mathbf{k}}|^2 d\Omega.$$
(3)

In equation (2), the value of d, introduced in the logarithmic scale breaking term  $\mathcal{L}_{scalebreak}$ , is equal to  $(2N_f)/(11N_c)$  for the QCD  $\beta$  function calculated at the one loop level (with  $N_c$  and  $N_f$  as the number of colors and flavors), but taken as a parameter in the effective chiral model. The wave functions of the charmonium states,  $\psi(\mathbf{k})$  are assumed to be harmonic oscillator wave functions [13, 14, 18],  $m_c$  is the mass of charm quark,  $m_{P,V}^{vac}$  is the vacuum mass of the pseudoscalar (vector) charmonium state, and  $\epsilon = 2m_c - m_{P,V}^{vac}$  is the binding energy of the charm-anticharm bound state.

The mixings of the pseudoscalar  $(P \equiv \eta_c(1S), \eta_c(2S))$  and vector  $(V \equiv J/\psi, \psi(2S), \psi(1D))$ charmonium states are taken into account through the interaction [26]

$$\mathcal{L}_{PV\gamma} = \frac{g_{PV}}{m_{av}} e \tilde{F}_{\mu\nu} (\partial^{\mu} P) V^{\nu}, \qquad (4)$$

where  $m_{av} = (m_V + m_P)/2$ ,  $m_P$  and  $m_V$  are the masses for the pseudoscalar and vector charmonium states,  $\tilde{F}_{\mu\nu}$  is the dual electromagnetic field. In the hadronic medium, the charmonium masses are calculated from the medium modification of the dilaton field, using equation (2) within the chiral effective model. In equation (4), the coupling parameter  $g_{PV}$  is fitted from the observed value of the radiative decay width,  $\Gamma(V \to P + \gamma)$  given as

$$\Gamma(V \to P\gamma) = \frac{e^2}{12} \frac{g_{PV}^2 p_{cm}^3}{\pi m_{av}^2},\tag{5}$$

where,  $p_{cm} = (m_V^2 - m_P^2)/(2m_V)$  is the magnitude of the center of mass momentum in the final state. The masses of the pseudoscalar and the longitudinal component of the vector mesons including the mixing effects are given by

$$m_{P,V^{||}}^{(PV)} = \frac{1}{2} \left( M_{+}^{2} + \frac{c_{PV}^{2}}{m_{av}^{2}} \mp \sqrt{M_{-}^{4} + \frac{2c_{PV}^{2}M_{+}^{2}}{m_{av}^{2}}} + \frac{c_{PV}^{4}}{m_{av}^{4}} \right), \tag{6}$$

where  $M_{+}^{2} = m_{P}^{2} + m_{V}^{2}$ ,  $M_{-}^{2} = m_{V}^{2} - m_{P}^{2}$  and  $c_{PV} = g_{PV}eB$ . The effective Lagrangian term given by equation (4) has been observed to lead to the mass modifications of the longitudinal  $J/\psi$  and  $\eta_{c}$  due to the presence of the magnetic field, which agree extermely well with a study of these charmonium states using a QCD sum rule approach incorporating the mixing effects [25, 26]. The decay widths of the vector charmonium states to  $D\bar{D}$  in the presence of magnetic fields are calculated from the mass modifications of these charmonium states and the open charm mesons in the magnetized nuclear matter. The D and  $\bar{D}$  meson masses are calculated within the chiral effective model from their interactions with the nucleons and scalar mesons in the magnetized nuclear matter. The lowest Landau level contributions are retained for the charged D and  $\bar{D}$  mesons in the presence of the external magnetic field. The charmonium decay widths are calculated using a field theoretical model of composite hadrons with quark/antiquark constituents as well as an effective hadronic Lagrangian, as described in the following section.

#### III. DECAY WIDTHS OF CHARMONIUM STATES TO $D\bar{D}$

The in-medium decay widths of the vector charmonium states to  $D\bar{D}$  in the presence of strong magnetic fields are computed using field theoretic model of composite hadrons with quark/antiquark constituents, and compared to the results using an effective hadronic Lagrangian, when the internal structure of these mesons are ignored. The models used for the calculation of the charmonium decay widths are described in the following.

#### A. Model of composite hadrons with quark/antiquark constituents:

We investigate the charmonium decay widths to the open charm (D and D) mesons in nuclear matter in the presence of strong magnetic fields using a field theoretical model with composite hadrons [37–39]. The model describes the hadrons comprising of quark and antiquark constituents. The constituent quark field operators of the hadron in motion are constructed from the constituent quark field operators of the hadron at rest, by a Lorentz boosting. Similar to the MIT bag model [40], where the quarks (antiquarks) occupy specific energy levels inside the hadron, it is assumed in the present model for the composite hadrons that the quark/antiquark constituents carry fractions of the mass (energy) of the hadron at rest (in motion) [37, 38].

With explicit constructions of the charmonium state and the open charm mesons, the decay width is calculated using the light quark antiquark pair creation term of the free Dirac Hamiltonian for constituent quark field [35]. The relevant part of the quark pair creation term is through the  $dd(u\bar{u})$  creation for decay of the charmonium state,  $\Psi$ , to the final state  $D^+D^-(D^0\bar{D^0})$ . For  $\Psi \to D(\mathbf{p}) + \bar{D}(\mathbf{p}')$ , this pair creation term is given as

$$\mathcal{H}_{q^{\dagger}\tilde{q}}(\mathbf{x}, t=0) = Q_q^{(p)}(\mathbf{x})^{\dagger} (-i\boldsymbol{\alpha} \cdot \nabla + \beta M_q) \tilde{Q}_q^{(p')}(\mathbf{x})$$
(7)

where,  $M_q$  is the constituent mass of the light quark, q = (u, d). The subscript q of the field operators in equation (7) refers to the fact that the light antiquark,  $\bar{q}$  and light quark, q are the constituents of the D and  $\bar{D}$  mesons with momenta  $\mathbf{p}$  and  $\mathbf{p}'$  respectively in the final state of the decay of the charmonium state,  $\Psi$ .

The charmonium state,  $\Psi$  with spin projection m, at rest is written as

$$|\Psi_m(\mathbf{0})\rangle = \int d\mathbf{k} c_r^{\ i}(\mathbf{k})^{\dagger} u_r a_m(\Psi, \mathbf{k}) \tilde{c_s}^{\ i}(-\mathbf{k}) v_s |vac\rangle, \tag{8}$$

where, *i* is the color index of the charm quark/antiquark operators,  $u_r$  and  $v_s$  are the two component spinors for the quark and antiquark. The expressions for  $a_m(\Psi, \mathbf{k})$  are given in terms of the wave functions (assumed to be harmonic oscillator type) for the charmonium states [35, 41].

The  $D(D^+, D^0)$  and  $\overline{D}(D^-, \overline{D}^0)$  states, with finite momenta are contructed in terms of the constituent quark field operators, obtained from the quark field operators of these mesons at rest through a Lorentz boosting [39]. These states, assuming harmonic oscillator wave functions, are explicitly given as

$$D(\mathbf{p})\rangle = \frac{1}{\sqrt{6}} \left(\frac{R_D^2}{\pi}\right)^{3/4} \int d\mathbf{k} \exp\left(-\frac{R_D^2 \mathbf{k}^2}{2}\right) c_r^{\ i} (\mathbf{k} + \lambda_2 \mathbf{p})^{\dagger} u_r^{\dagger} \tilde{q}_s^{\ i} (-\mathbf{k} + \lambda_1 \mathbf{p}) v_s d\mathbf{k},\tag{9}$$

$$|\bar{D}(\mathbf{p}')\rangle = \frac{1}{\sqrt{6}} \left(\frac{R_D^2}{\pi}\right)^{3/4} \int d\mathbf{k} \exp\left(-\frac{R_D^2 \mathbf{k}^2}{2}\right) q_r^{\ i} (\mathbf{k} + \lambda_1 \mathbf{p}')^{\dagger} u_r^{\dagger} \tilde{c_s}^{\ i} (-\mathbf{k} + \lambda_2 \mathbf{p}') v_s d\mathbf{k}, \quad (10)$$

where, q = (d, u) for  $(D^+, D^-)$  and  $(D^0, \overline{D^0})$  respectively.

۵

In equations (9) and (10),  $\lambda_1$  and  $\lambda_2$  are the fractions of the mass (energy) of the  $D(\bar{D})$ meson at rest (in motion), carried by the constituent light (d,u) antiquark (quark) and the constituent heavy charm quark (antiquark), with  $\lambda_1 + \lambda_2 = 1$ . The values of  $\lambda_1$  and  $\lambda_2$  are calculated by assuming the binding energy of the hadron as shared by the quark (antquark) to be inversely proportional to the quark (antiquark) mass [38]. The energies of the light antiquark (quark) and heavy charm quark (antiquark),  $\omega_i = \lambda_i m_D (i = 1, 2)$ , are assumed to be [35, 38]

$$\omega_1 = M_q + \frac{\mu}{M_q} \times BE, \quad \omega_2 = M_c + \frac{\mu}{M_c} \times BE, \tag{11}$$

where  $BE = (m_D - M_c - M_q)$  is the binding energy of  $D(\bar{D})$  meson, with  $M_c$  and  $M_q$  as the masses of the constitutent charm and light quark (antiquark), and,  $\mu$  is the reduced mass of the heavy-light quark-antiquark system (the  $D(\bar{D})$  meson), defined by  $1/\mu = 1/M_q + 1/M_c$ . The reason for making this assumption comes from the example of hydrogen atom, which is the bound state of the proton and the electron. As the mass of proton is much larger as compared to the mass of the electron, the binding energy contribution from the electron is  $\frac{\mu}{m_e} \times BE \simeq BE$  of hydrogen atom, and the contribution from the proton is  $\frac{\mu}{m_p} \times BE$ , which is negligible as compared to the total binding energy of hydrogen atom, since  $m_p >> m_e$ . With this assumption, the binding energies of the heavy-light mesons, e.g.,  $D(\bar{D})$  mesons [35] and,  $B(\bar{B})$  mesons [42], mostly arise from the contribution from the light quark (antiquark).

To compute the decay width of the charmonium state,  $\Psi$  to DD, we evaluate the matrix element of the light quark-antiquark pair creation part of the Hamiltonian, between the initial charmonium state and the final state for the reaction  $\Psi \to D(\mathbf{p}) + \overline{D}(\mathbf{p}')$  as given by

$$\langle D(\mathbf{p})|\langle \bar{D}(\mathbf{p}')| \int \mathcal{H}_{d^{\dagger}\bar{d}}(\mathbf{x},t=0) d\mathbf{x} |\Psi_m(\vec{0})\rangle = \delta(\mathbf{p}+\mathbf{p}') A^{\Psi}(|\mathbf{p}|) p_m, \tag{12}$$

where,

$$A^{\Psi}(|\mathbf{p}|) = 6c_{\Psi} \exp[(a_{\Psi}b_{\Psi}^{2} - R_{D}^{2}\lambda_{2}^{2})\mathbf{p}^{2}] \cdot \left(\frac{\pi}{a_{\Psi}}\right)^{3/2} \left[F_{0}^{\Psi} + F_{1}^{\Psi}\frac{3}{2a_{\Psi}} + F_{2}^{\Psi}\frac{15}{4a_{\Psi}^{2}}\right].$$
 (13)

In the above, the parameters  $a_{\Psi}$ ,  $b_{\Psi}$  and  $c_{\Psi}$  are given in terms of  $R_D$  and  $R_{\Psi}$ , which are the strengths of the harmonic oscillator wave functions for the  $D(\bar{D})$  and the charmonium states, and  $F_i^{\Psi}(i = 0, 1, 2)$  are polynomials in  $|\mathbf{p}|$ , the magnitude of the momentum of the outgoing  $D(\bar{D})$  meson [35]. With  $\langle f|S|i \rangle = \delta_4(P_f - P_i)M_{fi}$ , we have

$$M_{fi} = 2\pi (-iA^{\Psi}(|\mathbf{p}|)p_m.$$
(14)

The expression of the decay width of the charmonium state,  $\Psi$  to DD, as calculated in the present model for composite hadrons, without accounting for the mixing effects, is given by

$$\Gamma(\Psi \to D(\mathbf{p})\bar{D}(-\mathbf{p})) = \gamma_{\Psi}^{2} \frac{1}{2\pi} \int \delta(m_{\Psi} - p_{D}^{0} - p_{\bar{D}}^{0}) |M_{fi}|_{\mathrm{av}}^{2} \cdot 4\pi |\mathbf{p}_{D}|^{2} d|\mathbf{p}_{D}| \\
= \gamma_{\Psi}^{2} \frac{8\pi^{2}}{3} |\mathbf{p}|^{3} \frac{p_{D}^{0}(|\mathbf{p}|)p_{\bar{D}}^{0}(|\mathbf{p}|)}{m_{\Psi}} A^{\Psi}(|\mathbf{p}|)^{2}$$
(15)

In the above,  $p_{D(\bar{D})}^{0}(|\mathbf{p}|) = (m_{D(\bar{D})}^{2} + |\mathbf{p}|^{2})^{1/2}$ , and the expression for  $A^{\Psi}(|\mathbf{p}|)$  is given by equation (13). The parameter,  $\gamma_{\Psi}$ , in the expression for the charmonium decay width, is a measure of the coupling strength for the creation of the light quark antiquark pair, to produce the  $D\bar{D}$  final state. This parameter is adjusted to reproduce the vacuum decay widths of  $\psi(3770)$  to  $D^+D^$ and  $D^0\bar{D^0}$  [35]. The decay width of the charmonium state is observed to have the dependence on the magnitude of the 3-momentum of the produced  $D(\bar{D})$  meson,  $|\mathbf{p}|$ , as a polynomial part multiplied by an exponential term. The medium modification of the charmonium decay width is studied due to the mass modifications of the charmonium state, the D and  $\bar{D}$  mesons through  $|\mathbf{p}|$ , which is given as,

$$|\mathbf{p}| = \left(\frac{m_{\psi}^2}{4} - \frac{m_D^2 + m_{\bar{D}}^2}{2} + \frac{(m_D^2 - m_{\bar{D}}^2)^2}{4m_{\Psi}^2}\right)^{1/2}.$$
(16)

The expression for the charmonium decay width given by equation (15) is for the case when the mixing of the pseudoscalar and the longitudinal component of the vector mesons is not taken into account, and in equation (16), the masses of the charmonium and open charm mesons are the effective masses in the hadronic matter in the presence of a magnetic field.

When we include the mixing effect, the expression for the decay width is given as

$$\Gamma^{PV}(\Psi \to D(\mathbf{p})\bar{D}(-\mathbf{p})) = \gamma_{\Psi}^{2} \frac{8\pi^{2}}{3} \left[ \left( \frac{2}{3} |\mathbf{p}|^{3} \frac{p_{D}^{0}(|\mathbf{p}|) p_{\bar{D}}^{0}(|\mathbf{p}|)}{m_{\Psi}} A^{\Psi}(|\mathbf{p}|)^{2} \right) + \left( \frac{1}{3} |\mathbf{p}|^{3} \frac{p_{D}^{0}(|\mathbf{p}|) p_{\bar{D}}^{0}(|\mathbf{p}|)}{m_{\Psi}^{PV}} A^{\Psi}(|\mathbf{p}|)^{2} \right) \left( |\mathbf{p}| \to |\mathbf{p}| (m_{\Psi} = m_{\Psi}^{PV}) \right) \right].$$
(17)

In the above, the first term corresponds to the transverse polarizations for the charmonium state,  $\Psi$ , whose masses remain unaffected by the mixing of the pseudoscalar and vector charmonium states. The second term in (17) corresponds to the longitudinal component of the charmonium state whose mass is modified due to mixing with the pseudoscalar meson in the presence of the magnetic field, as given by equation (6).

#### B. Effective hadronic model

To calculate the decay width of the vector charmonium state to  $D\bar{D}$ , we use the effective interaction Lagrangian

$$\mathcal{L}_{had} = ig_{\Psi D\bar{D}}\Psi^{\mu}((\partial_{\mu}\bar{D})D - \bar{D}(\partial_{\mu}D)), \tag{18}$$

where,  $D = (D^0, D^+)^T$  and  $\overline{D} = (\overline{D^0}, D^-)$ . The above Lagrangian is motivated by the hidden gauge approach [43]. It might be noted here that another approach of introducing the massive vector mesons as gauge bosons using the minimal substitution leads to an additional term which is a contact interaction  $\sim \Psi^{\mu}\Psi_{\mu}\overline{D}D$  [18]. These methods are shown to be consistent when both the vector and axial vector mesons are included [44]. Using the interaction Lagrangian as given by equation (18), the decay width of the vector charmonium state at rest decaying to  $D\overline{D}(D^+D^-, D^0\overline{D^0})$  is obtained as

$$\Gamma_{\rm had}(\Psi \to D\bar{D}) = \frac{g_{\Psi D\bar{D}}^2}{6\pi m_{\Psi}^2} |\mathbf{p}|^3, \tag{19}$$

where  $|\mathbf{p}|$  is the magnitude of the momentum of the outgoing  $D(\bar{D})$  meson, given by equation (16). The coupling constant  $g_{\Psi D\bar{D}}$  for  $\Psi \to D\bar{D}$  is fitted from the measured decay width in vacuum. When the masses of the D and  $\bar{D}$  mesons are taken to be the same, the formula for the decay width reduces to the expression [29]

$$\left(\Gamma_{\rm had}(\Psi \to D\bar{D})\right)_{\rm approx} = \frac{g_{\Psi D\bar{D}}^2}{48\pi m_{\Psi}^2} (m_{\Psi}^2 - 4m_D^2)^{3/2},\tag{20}$$

for each of the channels  $\psi \to D^+ D^-$  and  $\Psi \to D^0 \overline{D^0}$ , giving the expression for the total decay width to be twice of the expression given by equation (20) [29].

In the presence of the magnetic field, there is mixing of the pseudoscalar and vector mesons leading to the masses of the longitudinal component of the vector charmonium state,  $\Psi$  to be modified. This leads to the decay width of  $\Psi \to D\bar{D}$  to be modified to

$$\Gamma_{had}^{PV}(\Psi \to D(\mathbf{p})\bar{D}(-\mathbf{p})) = \frac{g_{\Psi D\bar{D}}^2}{6\pi m_{\Psi}^2} \left[ \frac{2}{3} \left( |\mathbf{p}|(m_{\Psi}) \right)^3 + \frac{1}{3} \left( |\mathbf{p}|(m_{\Psi} \to m_{\Psi}^{PV}) \right)^3 \right],$$
(21)

where the first term corresponds to the contributions from the transverse components and the second term from the longitudinal component of the charmonium state  $\Psi$ .

## IV. RESULTS AND DISCUSSIONS

In the present work, the masses and the partial decay widths of the charmonium states to  $D\bar{D}$  are investigated in the presence of strong magnetic fields. These are studied taking the effects of the mixing of the pseudoscalar and vector mesons into consideration, in the presence

of magnetic fields. The created magnetic fields in the peripheral ultra-relavistic heavy ion collsion experiments, e.g., at RHIC and LHC, are huge, and the matter resulting from the high energy collision is of (extremely) low density. In the present work, the effects of magnetic field on the charmonium states are studied at zero density as well as at the nuclear matter saturation density in the magnetized (asymmetric) nuclear matter. In the nuclear medium, the masses of the charmonium states and the open charm mesons in the presence of a magnetic field are calculated using a chiral effective model. The in-medium charmonium masses are studied in the magnetized nuclear matter as arising from the medium changes in a scalar dilaton field,  $\chi$ , which mimics the gluon condensates of QCD in the effective hadronic model [24]. The masses of the open charm mesons in the magnetized nuclear matter are modified due to their interactions with the nucleons as well as the scalar mesons,  $\sigma$ ,  $\zeta$ , and  $\delta$  [23]. These scalar fields and the dilaton field  $\chi$  are solved from their coupled equations of motion. The charmonium and open charm meson masses are obtained from the changes in the scalar fields in the isospin asymmetric nuclear matter (with isospin asymmetry parameter,  $\eta$  defined as  $\eta = (\rho_n - \rho_p)/(2\rho_B)$ ), in the presence of magnetic fields using the chiral effective model [30]. The effects of the mixing of the pseudoscalar and longitudinal vector mesons are taken into account, which are observed to lead to significant contributions to the charmonium masses. These, as we shall see later, have appreciable effects on the partial decay width of the  $\psi(3770)$  to DD at high magnetic fields.

The mass modifications of the pseudoscalar ( $\eta_c \equiv \eta_c(1S)$  and  $\eta'_c \equiv \eta_c(2S)$ ) and the vector  $(J/\psi, \psi(2S) \equiv \psi(3686), \psi(1D) \equiv \psi(3770))$  charmonium states are investigated in the presence of magnetic fields. These masses are studied accounting for the mixing of the pseudoscalar and vector mesons in the presence of magnetic fields, described by the effective interaction Lagrangian term given by equation (4). For the charmonium states at rest, the mixing effect leads to a drop (increase) in the mass of the pseudoscalar meson (the longitudinal component of the vector charmonium state) as given by equation (6), and these are observed to be the dominant modifications to the charmonium masses.

The in-medium masses of the vector charmonium states  $(J/\psi, \psi(2S), \psi(1D))$  have been studied from the medium change of the gluon condensate, using the leading order mass formula in Ref. [18] and using the chiral effective model in Refs. [13, 14, 24] in (magnetized) hadronic matter. The mass modifications of the charmonium states within the chiral effective model




FIG. 1: (Color online) The masses of the pseudoscalar mesons ( $\eta_c \equiv \eta_c(1S)$  and  $\eta'_c \equiv \eta_c(2S)$ ) and the longitudinal components of the vector charmonium states ( $J/\psi, \psi(2S) \equiv \psi(3686), \psi(1D) \equiv \psi(3770)$ ) are plotted as functions of  $eB/m_{\pi}^2$  for  $\rho_B = 0$  as well as at  $\rho_B = \rho_0$  in symmetric nuclear matter ( $\eta=0$ ). The effects of the mixing between the pseudoscalar and the vector charmonium states ( $J/\psi$  with  $\eta_c$ ,  $\psi(2S)$  with  $\eta_c(2S)$  and  $\psi(1D)$  with  $\eta_c(2S)$ ) on the charmonium masses are shown and compared to the case of not including the mixing effects (shown as dotted lines).




FIG. 2: (Color online) Same as fig. 1, for asymmetric nuclear matter with  $\eta=0.5$ .

arise from the modifications of a scalar dilaton field which simulates the gluon condensates of QCD. These in-medium masses were studied assuming the harmonic oscillator wave functions for the charm-anticharm bound states. The values of the strength parameter,  $\beta \equiv 1/R$ , of the charmonium wave functions were fitted to the rms radii of  $J/\psi$ ,  $\psi(2S)$  and  $\psi(1D)$  to be (0.47 fm)<sup>2</sup>, (0.96 fm)<sup>2</sup> and 1 fm<sup>2</sup> respectively [18], yielding their values as 513 MeV, 384 MeV and 368 MeV [13, 14]. In the present work, the mass modifications for the pseudoscalar states,




FIG. 3: (Color online) The masses of D mesons ( $D^+$  and  $D^0$ ) are plotted as functions of  $eB/m_{\pi}^2$ . These masses are shown for the case of  $\rho_B = 0$ , as well as for  $\rho_B = \rho_0$  in symmetric ( $\eta=0$ ) and asymmetric (with  $\eta=0.5$ ) nuclear matter.

 $\eta_c \equiv \eta_c(1S)$  and  $\eta'_c \equiv \eta_c(2S)$  in the nuclear matter are investigated using the chiral effective model. The values of the harmonic oscillator strengths for  $\eta_c$  and  $\eta'_c$  are obtained as 535 MeV and 394.6 MeV respectively, assuming these states to be lying in a straight line with the states  $J/\psi$  and  $\psi(2S)$  in the mass versus  $\beta$  graph. In the absence of a magnetic feild, the masses




FIG. 4: (Color online) Same as fig. 3, for  $D^-$  and  $\overline{D^0}$ .

for the pseudoscalar mesons  $\eta_c$  and  $\eta'_c$  are modified to 2977.26 (2977.47) and 3548.55 (3551.46) MeV at  $\rho_B = \rho_0$  for symmetric (asymmetric with  $\eta=0.5$ ) nuclear matter from their vacuum masses of 2983.9 MeV and 3637.5 MeV respectively. The mass shift of  $\eta_c$  in symmetric nuclear matter at the nuclear matter saturation density of around 6.6 MeV may be compared to the values of 5 MeV [9] and 5.69 MeV [10] using a QCD sum rule approach, and of 3 MeV as calculated from  $\eta_c$ -nucleon scattering length [45]. The mass drop in  $J/\psi$  of around 8.6 MeV




FIG. 5: (Color online) Decay widths of  $\psi(3770)$  to (I)  $D^+D^-$ , (II)  $D^0\bar{D^0}$ , and the total of these two channels (I+II), are plotted as functions of  $eB/m_{\pi}^2$ . These are shown for the case of  $\rho_B = 0$ , as well as, for  $\rho_B = \rho_0$ , in symmetric ( $\eta=0$ ) and asymmetric (with  $\eta=0.5$ ) nuclear matter. The panels (a), (c) and (e) show the results when the mixing with  $\eta'_c$  is not taken into account, and (b), (d) and (f) correspond to the results including the mixing effect. These results are compared with the decay widths obtained using a hadronic Lagrangian given by equation (18), shown as dotted lines.

at  $\rho_B = \rho_0$  in symmetric nuclear matter obtained using the chiral effective model [13] may be compared with the mass drop of around 8 MeV calculated using second order Stark effect in QCD with the gluon condensate calculated in the linear density approximation [18] as well as the mass shifts of 4 MeV and 7 MeV using QCD sum rule calculations with OPE upto dimension six and four in Refs. [8] and [9] respectively. The mass shifts of the excited states  $\psi(3686)$  and  $\psi(3770)$  obtained in the chiral effective model [13, 14] are observed to be similar to the values of 100 and 140 MeV using the QCD second order Stark effect [18], which are much larger than the mass shift of  $J/\psi$ .

In figure 1, the effects of the magnetic field on the masses of the charmonium states are shown for zero baryon density as well as for  $\rho_B = \rho_0$  in symmetric ( $\eta=0$ ) nuclear matter. In panel (a), the masses of the pseudoscalar charmonium state,  $\eta_c$  and the longitudinal component of  $J/\psi$ as modified due to the effect of mixing of these states, are shown as functions of  $eB/m_{\pi}^2$ . These masses are calculated by using the equation (6). The parameter  $g_{PV} \equiv g_{\eta_c J/\psi}$  is evaluated to be 2.094 from the observed radiative decay width  $\Gamma(J/\psi \rightarrow \eta_c \gamma)$  in vacuum of 92.9 keV [46], using equation (5). The dotted lines correspond to the case when these mixing effects are not taken into account. For  $\rho_B=0$ , without the mixing effects, the masses of the charmonium states are unaffected by the magnetic field, and remain at the vacuum values of the masses of  $J/\psi$  and  $\eta_c$  of 3097 and 2983.9 MeV respectively. The mixing effect is observed to lead to appreciable modifications to their masses at high magnetic fields, with an increase in the mass (in MeV) of  $J/\psi^{\parallel}$ , the longitudinal component of  $J/\psi$ , by about 9 (31.5) and a drop in the mass of  $\eta_c$  of around 8.9 (30) at eB=5 (10)  $m_{\pi}^2$  respectively. The mass difference (in MeV) of  $J/\psi$  and  $\eta_c$  of 113 in vacuum is thus modified to 131 (174.5) at value of the magnetic field eBas 5 (10)  $m_{\pi}^2$  for the  $J/\psi^{||}$  and  $\eta_c$  mesons. Such a level repulsion of the  $J/\psi^{||}$  and  $\eta_c$  states has also been observed by incorporating the effects of a magnetic field consistently in the Operator product expansion (OPE) as well as in the phenomenological side within a QCD sum rule calculation [25, 26]. An appropriate form of the spectral ansatz is used to describe the mixing of the pseudoscalar and longitudinal vector mesons, retaining the effects of external magnetic field up to second order in eB. In Refs. [25, 26], the masses of the longitunal  $J/\psi$  and  $\eta_c$  in the presence a magnetic field calculated in the QCD sum rule approach are compared with the masses from the effective hadronic interaction

$$m_{P,V||_{approx}}^{2} = m_{P,V}^{2} \mp \frac{(g_{PV}eB)^{2}}{m_{V}^{2} - m_{P}^{2}},$$
(22)

obtained from the expressions given by (6) by retaining terms up to the second order in eB and the leading order in  $\frac{(m_V - m_P)}{(m_V + m_P)}$  [26]. It is observed that the masses of the longitudinal  $J/\psi$  and  $\eta_c$ agree remarkably well with the second order results of the masses obtained from the effective hadronic interaction up to  $eB \sim 5m_{\pi}^2 (\sim 0.1 GeV^2)$  with slight deviation at higher values of eB. Also, the second order results for the masses of the  $J/\psi^{\parallel}$  and  $\eta_c$  given by equation (22) are observed to agree extremely well with the general expressions given by equation (6) with marginal variations at higher values of magnetic field [26]. In Refs. [25, 26], for comparing the results of the QCD sum rules with the second order results of the effective hadronic model, the vacuum masses (in MeV) of  $J/\psi$  and  $\eta_c$  of 3092 and 3025 respectively in the QCD sum rule approach, are also taken as the vacuum masses for these mesons to compute the masses of  $J/\psi^{||}$ and  $\eta_c$  due to their mixing. One can see from the second order expression for the masses given by equation (22) that the mixing effect leads to an increase (drop) in the mass of the longitudinal  $J/\psi$  ( $\eta_c$ ) meson. The level repulsion of these states was also studied using a potential model approach by solving the Schrodinger equation in the presence of an external magnetic field [27]. The masses of heavy quarkonium states have been studied including the effects of mixing of the longitudinal vector and pseudoscalar quarkonium states. The Schrödinger equation for the heavy quark antiquark system is solved in the presence of a magnetic field, assuming a Cornell potential along with spin-spin interaction. In the presence of an external magnetic field, the center of mass momentum of the quarkonium state is no longer a conserved quantity. The masses of the lowest charmonium states  $J/\psi$  and  $\eta_c$  are obtained by solving Schrödinger equation for given values of eB and the center of mass momentum,  $\langle P_{kin} \rangle$  [27]. For  $\langle P_{kin} \rangle = 0$ , there is observed to be an increase (drop) of  $J/\psi^{||}(\eta_c)$  mass of the order of 80 (50) MeV for  $eB = 10m_{\pi}^2 \sim 0.2 \text{GeV}^2$  as is expected due to the spin-mixing [27]. These values may be compared to the values of around 31.5 (30) for the positive (negative) mass shifts of  $J/\psi^{||}(\eta_c)$ in the present investigation. The mixing of the  $\eta_c$  with the longitudinal  $J/\psi$  can also have observational consequences on the dilepton spectra. The probability of finding the longitudinal  $J/\psi$  in the state  $\eta_c$  due to their mixing has been calculated to be around 12 (26)% for  $eB \sim$ 5 (10)  $m_{\pi}^2$  in Ref. [27]. This should show as a suppression of dileptons arising from the longitudinal  $J/\psi$  (by about 12 (26)%), which should instead show as decay from  $\eta_c$  in the dilepton spectra. The increase in the mass of  $J/\psi^{||}$  in the presence of strong magnetic fields should also show as a suppression in the  $J/\psi$  production [27].

The masses of  $J/\psi^{\parallel}$  and  $\eta_c$  are plotted for  $\rho_B = \rho_0$  in symmetric nuclear matter in panel (b) in figure 1. In the absence of mixing effects, the masses are observed to vary marginally with magnetic field. The mass difference (in MeV) of  $J/\psi$  and  $\eta_c$  of around 111 at  $\rho_B = \rho_0$  without mixing effect, is observed to be modified to 132 (175.7) for the value of eB as 5 (10)  $m_{\pi}^2$  in the presence of mixing for the longitudinal  $J/\psi$  and  $\eta_c$  mesons. These values are observed to be similar to the values of the mass differences of these states of 131 (174.5) MeV for eB=5 (10)  $m_{\pi}^2$  for  $\rho_B=0$ .

The masses of  $\psi^{\parallel}(2S)$  and  $\eta'_c \equiv \eta_c(2S)$  in the presence of a magnetic field are shown for  $\rho_B=0$  in panel (c) in figure 1. The value for the mixing coupling parameter of these states,  $g_{PV} \equiv g_{\eta'_c \psi(2S)}$  is obtained to be 3.184, fitted from the observed vacuum value of the decay width  $\Gamma(\psi(2S) \to \eta'_c \gamma)$  of 0.2058 keV [46]. For  $\rho_B = \rho_0$  as shown in panel (d), the mixing effect is observed to be much more prominent as compared to the zero density case. This is due to the reason that the mass modification due to mixing is larger for a smaller difference in the masses of the pseudoscalar and vector mesons. As can be seen from equation (22), the shifts in the masses of the pseudoscalar and vector mesons are inversely proportional to  $(m_V^2 - m_P^2)$ , and hence inversely proportional to the mass difference of these mesons. The mass splittings (in MeV) of the  $\psi^{\parallel}(2S)$  and  $\eta'_c$  are observed to be around 90.5 (174.9) for eB=5(10)  $m_{\pi}^2$  at  $\rho_B = \rho_0$  and 97 (174.88) for  $\rho_B = 0$  which are much larger as compared to the mass splitting of 48.5 in vacuum for zero magnetic field. As might be observed from the panels (c) and (d), the mass splitting for zero magnetic field is smaller for the  $\rho_B = \rho_0$  case as compared to  $\rho_B = 0$ . However, the mass splitting is observed to be similar for zero as well as nuclear matter saturation density for the considered magnetic fields, due to the larger mass shifts for the longitudinal  $\psi(2S)$  and  $\eta'_c$  for  $\rho_B = \rho_0$  compared to zero density.

The masses of  $\eta'_c$  and the longitudinal component of  $\psi(1D) \equiv \psi(3770)$  in the presence of a magnetic field are plotted for zero density and for  $\rho_B = \rho_0$  in symmetric nuclear matter in panels (e) and (f) respectively in figure 1, including the contributions from the mixing effects. The value for  $g_{PV} \equiv g_{\eta'_c \psi(1D)}$  is obtained as 7.657 from the observed vacuum value of the decay width  $\Gamma(\psi(2S) \to \eta'_c \gamma)$  of 24.48 keV [46]. Due to the smaller difference in the masses of the  $\eta'_c$  and  $\psi(3770)$  in the magnetized nuclear matter, the mixing effects are observed to lead to larger shifts in the masses of these mesons at  $\rho_B = \rho_0$  as compared to zero density. This is similar to the larger mass shifts of  $\eta'_c$  and  $\psi(3686)$  due to the mixing effects, at  $\rho_B = \rho_0$  as compared to zero density. The mass splittings of the  $\psi^{\parallel}(1D)$  and  $\eta'_c$  are observed to be around 188.2 (332.7) for  $eB = 5(10) m_{\pi}^2$  at  $\rho_B = \rho_0$  and 213.7 (346.2) for  $\rho_B = 0$ , which is much larger than the mass difference of 135.5 in vacuum for zero magnetic field. As can be observed from the figure, the mass of the longitudinal  $\psi(3770)$  has dominant positive contributions at high magnetic fields due to the mixing effects. This leads to significant modification of the partial decay width of  $\psi(3770)$  to  $D\bar{D}$  at high magnetic fields, as we shall see later.

The mixing of the longitudinal component of the vector charmonium state with the pseudoscalar mesons can lead to 'anomalous'  $\eta_c$  and  $\eta'_c$  peaks, in addition to the  $J/\psi$ ,  $\psi(3686)$  and  $\psi(3770)$ -like peaks in the dilepton spectra, and can probe the existence of early magnetic field. A study of the formation times of the charmonium states due to the mixing effects of  $J/\psi - \eta_c$  as well as  $\psi(3686) - \eta'_c$  is observed to lead to a faster (slower) formation of the pseudoscalar (vector) charmonium state. The early formation of the  $\eta_c$  and  $\eta'_c$  can probe the magnetic field at the early stage when the magnetic field can still be large.

The charmonium masses are plotted for asymmetric nuclear matter with  $\eta=0.5$  in figure 2. The effects of the isospin asymmetry on the charmonium masses are observed to be small at the nuclear matter saturation density, as calculated within the chiral effective model. This, in turn, leads to small modifications to the charmonium masses in the isospin asymmetric matter (with  $\eta=0.5$ ) as compared to the symmetric nuclear matter, in the presence of mixing effects. For example, the masses (in MeV) of  $\psi(3686)$  and  $\psi(3770)$  at the nuclear matter saturation density, and for  $eB = 10m_{\pi}^2$ , the highest magnetic field considered in the present work, are observed to be modified from 3650.57 and 3808.45 for symmetric nuclear matter to the values 3662.2 and 3821.4 for asymmetric nuclear matter with  $\eta=0.5$ . It might be noted here that the contributions to the masses of the excited charmonium states  $\psi(3686)$  and  $\psi(3770)$ , at zero density as well as at  $\rho_B = \rho_0$  in nuclear matter, due to the mixing with  $\eta_c$  are negligible for the magnetic fields considered in the present work. This is expected from the larger mass differences of these vector mesons from the mass of  $\eta_c$ , which give negligible shifts to these charmonium masses due to mixing with  $\eta_c$ , as is evident from equation (22). The excited states  $\psi(3686)$  and  $\psi(3770)$  thus have dominant contributions to their masses arising from the mixing with  $\eta'_c$  in the presence of high magnetic fields, as has been discussed above.

The mass splittings of the longitudinal component of the vector  $(V^{||})$  and the pseudoscalar (P) charmonium states are observed to much larger for high magnetic fields than the values in vacuum at zero magnetif field, due to the effects of mixing. For  $J/\psi^{||} - \eta_c$ ,  $\psi^{||}(3686) - \eta'_c$  and  $\psi^{||}(3770) - \eta'_c$  mixings, the values of mass difference in these states (in MeV) of 113, 48.5 and 135.5 for vacuum at zero magnetic field are modified to around 130 (175), 90 (175) and 188 (333) for  $eB = 5(10)m_{\pi}^2$ . These large mass splittings should show as peaks corresponding to these states in the dilepton spectra in ultrarelativistic heavy ion collision experiments and can probe the existence of magnetic field in the early stage.

In figures 3 and 4, the masses of the  $D(D^+, D^0)$  mesons and  $\overline{D}(D^-, \overline{D}^0)$  mesons, are plotted as functions of  $eB/m_{\pi}^2$ , for  $\rho_B = 0$  as well as for  $\rho_B = \rho_0$  with  $\eta = 0$  and  $\eta = 0.5$ . In the nuclear matter in the presence of a magnetic field, these masses are calculated within the chiral effective model, from their interactions with the nucleons and scalar mesons. The charged open charm mesons  $(D^+ \text{ and } D^-)$  have additional positive shifts in their masses due to contributions from the Landau levels in the presence of a magnetic field. On the other hand, the neutral  $D^0$  and  $D^0$  mesons are observed to have very small changes in their masses due to the magnetic field. The D and D masses in nuclear matter have been calculated in presence of magnetic fields using the chiral effective model in Ref. [23], without accounting for the medium modification of the scalar dilaton field, which simulates the gluon condensates of QCD. Using the chiral effective model, the mass modifications of the charmonium states arise due to the medium changes of the dilaton field. The in-medium decay widths of the charmonium states to DDhave been studied using the mass modifications of the charmonium states [24] as well as the open charm mesons in the magnetized nuclear matter [30]. In the present work, the vector charmonium masses have been studied including the mixing of the longitudinal component of these states with the pseudoscalar charmonium states. The mass of the longitudinal component of the vector charmonium state has an appreciable positive shift due to the pseudoscalar-vector meson mixing in the presence of the magnetic field. The lowest charmonium state decaying to  $D\bar{D}$  is  $\psi(3770)$  at  $\rho_B = \rho_0$  as well as at zero baryon density. As has already been mentioned, the charmonium decay width,  $\Gamma(\psi(3770) \rightarrow D\bar{D})$  is calculated using the light quark pair creation term of the free Dirac Hamiltonian expressed in terms of the constituent quark operators, using explicit constructions for the charmonium and the open charm (D and  $\bar{D}$ ) mesons. The matrix element for the calculation of the decay width is multiplied with a factor  $\gamma_{\psi}$ , which gives the strength of the light quark pair creation leading to the decay of the charmonium state to  $D\bar{D}$  in the magnetized hadronic medium [35]. The value of  $\gamma_{\psi}$  to be 1.35, is chosen so as to reproduce the decay widths of  $\psi(3770) \rightarrow D^+D^-$  and  $\psi(3770) \rightarrow D^0\bar{D}^0$  in vacuum, to be around 12 MeV and 16 MeV respectively [35]. The constituent quark masses for the light quarks (u and d) are taken to be 330 MeV and for the charm quark, the value is taken to be  $M_c = 1600$  MeV [35].

The partial decay widths of  $\psi(3770)$  to  $D\bar{D}(D^+D^-, D^0\bar{D}^0)$ , are calculated using a field theoretic model for composite hadrons as described in section III A. These are plotted in figure 5 as functions of  $eB/m_{\pi}^2$  for  $\rho_B = \rho_0$  as well as at zero baryon density. To examine the effects of the compositeness of the hadrons on the charmonium decay widths in the presence of strong magnetic fields, the decay widths of  $\psi(3770) \rightarrow D\bar{D}$  as calculated using a field theoretic model of hadrons with quark/antiquark constituents are compared with the results obtained using the effective hadronic Lagrangian [43] given by equation (18) (shown as the dotted lines). The value of the coupling constant  $g_{\psi D\bar{D}}$  in the interaction hadronic Lagrangian is taken as 14.5, which gives the vacuum values of the decay widths of  $\psi(3770) \rightarrow D\bar{D}$  to be around 28 MeV, with the values of 12 and 16 MeV in the decay channels to  $D^+D^-$  and  $D^0\bar{D^0}$  respectively. In panel (a) in figure 5, the partial decay widths of  $\psi(3770)$  to (I)  $D^+D^-$ , (II)  $D^0\bar{D^0}$ , and the total of these two channels (I+II), are plotted for  $\rho_B = 0$ , without taking into account the pseudoscalar-vector meson  $(\eta'_c - \psi(3770))$  mixing. The decay width for  $\psi(3770)$  decaying to the charged  $D\bar{D}$  is observed to drop with increase in the magnetic field up to eB equal to  $3m_{\pi}^2$ , when it becomes zero and remains zero for larger values of the magnetic field. This is due to the reason that the charged D and D mesons have higher values for their masses in the presence of a magnetic field, due to positive Landau level contributions to their masses. On the other hand, the decay wdith to the neutral DD pair is observed to be unaffected by the magnetic field in the absence of the mixing effects as seen in panel (a). Panel (b) shows the partial decay widths when the mass modification of the longitudinal component of  $\psi(3770)$  is taken into account due to mixing with  $\eta'_c$  in the presence of a magnetic field. This leads to an appreciable increase in the decay to neutral  $D^0 \overline{D^0}$  with rise in the magnetic field, whereas the decay to the  $D^+D^-$  is observed to have an initial drop up to around  $eB \sim 3m_\pi^2$  beyond which it is observed to be almost constant (with a value of around 7 MeV) when calculated using the field theoretic model of composite hadrons. There are observed to be appreciable deviations in the decay widths computed using the model for composite hadrons from the calculations within the effective hadronic model at high values of the magnetic fields in the presence of mixing effects. As can be seen from the equations (15) and (19), the dependence of the charmonium decay widths to  $D\bar{D}$  on the magnitude of the momentum of the outgoing are given as a polynomial multiplied by an exponential function in the model of composite hadrons, whereas, there is a  $|\mathbf{p}|^3$  dependence in the effective hadronic model. These are modified to the expressions given by equations (17) and (21) in the presence of magnetic fields, as the longitudinal components of the vector charmonium states are modified due to mixing with the pseudoscalar charmonium states. The in-medium decay widths of charmonium states to  $D\bar{D}$ have been studied due to mass modifications of the open charm mesons in hadronic matter for zero magnetic field, using the compositeness of the hadrons using the  ${}^{3}P_{0}$  model [14, 29] as well as using the field theoretic model for composite hadrons [35] as considered in the present work, are observed to vanish at certain densities (so called nodes). This is due to the dependence of these decay widths on the magnitude of the momentum of the outgoing  $D(\bar{D})$  meson, ||, as a polynomial multiplied by an exponential function, which vanishes for the values of the  $|\mathbf{p}|$ corresponding to these densities. On the other hand, the effective hadronic model as given by (18), due to the  $|\mathbf{p}|^3$  as given by equation (19) is observed to lead to a monotonic increase with increase in  $|\mathbf{p}|$  (corresponding to smaller values of the  $D(\bar{D})$  mass in the medium). The results for the charmonium decay widths obtained from the effective hadronic model are significantly different from the behaviour as obtained in the  ${}^{3}P_{0}$  model [29]. In the present work, the increase in the value of  $|\mathbf{p}|$  in the presence of magnetic fields is due to the appreciable increase in the mass of the longitudinal component of the charmonium state,  $\psi(3770)$  due to mixing with the pseudoscalar meson,  $\eta_c'$ , which is observed to lead to much larger increase in the decay width of  $\psi(3770) \rightarrow D^0 \bar{D^0}$  obtained from the effective hadronic model, as compared to the values calculated in the field theoretic model. The presence of a magnetic field leads to increase in the masses of the  $D^{\pm}$  mesons due to the Landau level contributions, which is observed as a drop in the decay width up to the value of  $eB \sim 3m_{\pi}^2$ , beyond which it is observed to remain almost constant (increases) with further increase in the magnetic field, in the model for composite hadrons (effective hadronic model) as can be seen in panel (b) for zero baryon density and including the effect of mixing of the longitudinal  $\psi(3770)$  with  $\eta'_c$ .

The decay widths of  $\psi(3770) \rightarrow D\bar{D}$  for baryon density  $\rho_B = \rho_0$  in magnetized nuclear matter are shown without and with the mixing effects, in panels (c) and (d) for isospin asymmetry parameter,  $\eta=0$  and in panels (e) and (f) for  $\eta=0.5$  in figure 5. In the absence of mixing effects, these decay widths are obtained from the modifications of the masses of the charmonium state  $\psi(3770)$  as well as the D and D mesons in magnetized nuclear matter using the chiral effective model. In the absence of mixing effects, the decay width to charged  $D^+D^-$  is observed to be around 0.5 MeV at zero magnetic field and vanishes for eB higher than  $m_{\pi}^2$ , when the drop in the charmonium mass in the medium as well as the increase in the masses of the charged open charm mesons due to the Landau contributions in the presence of the magnetic field kinematically forbids the decay of  $\psi(3770)$  to  $D^+D^-$ . The decay to the neutral  $D\bar{D}$  is observed to increase with magnetic field and the rise is much higher in the effective hadronic model due to the  $|\mathbf{p}|^3$  dependence of the decay width. The panels (e) and (f) show the results for the decay widths for asymmetric nuclear matter with  $\eta = 0.5$  for  $\rho_B = \rho_0$ . In the absence of mixing effects, there is observed to be contribution to the decay width of  $\psi(3770)$  only from the channel of decay to the charged DD pair. The increase in the masses of the final state charged open charm mesons with magnetic field due to the Landau contributions is observed as a drop and vanishing of the decay wdith of  $\psi(3770) \rightarrow D^+D^-$  for eB higher than around 2.3  $m_{\pi}^2$ . In the presence of the mixing effects, due to increase in the mass of the longitudinal  $\psi(3770)$ , there is observed to be contribution from the partial decay width to the neutral open charm meson pair for eB higher than 2  $m_{\pi}^2$ . For larger values of the magnetic field, the decay width is observed to be dominated by the contribution from the neutral DD channel. However, the decay widths obtained within the effective hadronic model, shown as dotted lines, are observed to have a much sharper rise as compared to the results obtained using the model of composite hadrons.

The decay of the charmonium state  $\psi(3770) \rightarrow D\bar{D}$  is observed to be dominated by the decay to the neutral  $D\bar{D}$  pair as compared to  $D^+D^-$ , for large values of the magnetic fields, both for the zero baryon density, as well as for  $\rho_B = \rho_0$  in (asymmetric) nuclear matter. This

should lead to the neutral open charm mesons  $(D^0 \text{ and } \overline{D^0})$  to be more abundant as compared to the charged  $D^{\pm}$  mesons. The decay width is observed to increase with the magnetic field. This should lead to suppression in the yield of  $\psi(3770)$  at large values of the magnetic fields, hence of the  $J/\psi$ , as the excited charmonium states are a major source of  $J/\psi$ .

### V. SUMMARY

In the present work, the charmonium masses as well as the decay widths of the vector charmonium states to  $D\bar{D}$  are investigated in the presence of strong magnetic fields. The study can be of relevance to the observables in ultra-relativistic heavy ion collision experiments. The estimated magnetic fields created in the peripheral heavy ion collisions, e.g., at RHIC and at LHC, are huge, where the formed matter is (extremely) low in density. In the present work, we study the effects of the magnetic fields on the masses and decay widths of the charmonium states at zero as well as at nuclear matter saturation density.

The mixing of the pseudoscalar mesons and the vector mesons in the presence of strong magnetic fields are studied using an effective Lagrangian interaction. The effects from the mixing are observed to have dominant contributions to the masses of the charmonium states, with an increase (drop) for the longitudinal component of the vector (pseudoscalar) charmonium state in vacuum/nuclear matter. In the nuclear medium, the masses of the vector charmonium states  $(J/\psi, \psi(3686), \psi(3770))$  and the pseudoscalar states  $\eta_c$  and  $\eta'_c$  are studied as arising from the medium changes of the scalar dilaton field (which simulates the gluon condensates of QCD) using a chiral effective model. These masses charmonium states are additionally modified in the nuclear medium due to the mixing effects. The mixing of the pseudoscalar mesons,  $\eta_c$  and  $\eta_c'$  with the longitudinal vector (V) charmonium states, lead to large positive (negative) shifts in the masses of  $V^{\parallel}$  (pseudoscalar) mesons at high magnetic fields. The finite probability of the longitudinal component of the vector meson to be in the pseudoscalar state should lead to suppression of the dileptons from the decay of  $V^{||}$ , which should instead arise from the decay of the pseudoscalar meson. The mixing effect can thus show as peaks in the dilepton spectra due to 'anomalous' decay modes, e.g.,  $\eta_c, \eta_c' \to l^+ l^-$  in addition to the  $V \to l^+ l^-$  and can probe the magnetic field at the early stage.

The in-medium decay widths of charmonium state to  $D\bar{D}$  in the presence of strong mag-

netic fields are computed using a field theoretical model for composite hadrons with quark (antiquark) constituents. The matrix element for computation of the charmonium decay width is calculated from the free Dirac Hamiltonian of the constituent quarks, using the explicit constructions of the charmonium state, the D and the  $\overline{D}$  mesons. The decay widths are computed from the modifications of the masses of the charmonium state and the open charm mesons in vacuum/nuclear medium in the presence of strong magnetic fields. In the nuclear medium, the charmonium masses as well as the D and D masses are calculated within a chiral effective model. As has already been mentioned, the charmonium masses are calculated from the medium modification of a scalar dilaton field, and, the masses of the D and  $\overline{D}$  mesons are calculated within the model due to their interactions with the nucleons and the scalar mesons in the magnetized nuclear matter, with additonal Landau contributions to the masses of the charged  $D^{\pm}$  mesons in the presence of the magnetic field. At  $\rho_B=0$  as well as at nuclear matter saturation density, the lowest charmonium state which can decay to  $D\bar{D}$  is  $\psi(3770)$ . The mass of longitudinal  $\psi(3770)$  is observed to have appreciable positive contribution due to mixing with  $\eta_c'$  in the presence of strong magnetic fields, which is seen to have significant modification to the partial decay width,  $\Gamma(\psi(3770) \rightarrow D\bar{D})$ . The decay of  $\psi(3770)$  to  $D^+D^-$  are observed to be quite suppressed as compared to decay to  $D^0 \overline{D^0}$ , in the presence of strong magnetic fields. This is due to the higher masses of the charged mesons  $(D^+, D^-)$  due to contributions from the Landau levels. This should lead to the production of the neutral  $D^0$  and  $\overline{D^0}$  mesons to be more abundant as compared to the  $D^{\pm}$  mesons from  $\psi(3770)$ . The increase of the decay width of  $\psi(3770)$  at high magnetic fields should show as a suppression in the yield of  $\psi(3770)$  and hence of  $J/\psi$ , as  $\psi(3770)$  is a major source for production of  $J/\psi$ .

To summarize, the present study of the masses of the charmonium states in strong magnetic fields, which has dominant contributions from the mixing of the longitudinal vector meson – pseudoscalar mesons can be a probe for the early magnetic field as the mixing can show as pseudoscalar like peaks in addition to the vector meson peaks in the dilepton spectra, and the study of charmonium decay width to  $D\bar{D}$ , which is observed to be enhanced in strong magnetic fields, predominantly in the decay mode to neutral  $D\bar{D}$ , should have observable consequences on the suppression in the yield of  $\psi(3770)$  (and hence of  $J/\psi$ ) and higher yield for the neutral open charm ( $D^0$ ,  $\bar{D}^0$ ) as compared to the charged  $D^{\pm}$  mesons due to the presence of strong magnetic fields created in non-central ultra-relativistic heavy ion collision experiments.

One of the authors (AM) is grateful to ITP, University of Frankfurt, for warm hospitality and acknowledges financial support from Alexander von Humboldt Stiftung when this work was initiated.

- [1] A. Hosaka, T. Hyodo, K. Sudoh, Y. Yamaguchi, S. Yasui, Prog. Part. Nucl. Phys. 96, 88 (2017).
- [2] E.Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.M. Yan, Phys. Rev. D 17, 3090 (1978).
- [3] E.Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.M. Yan, Phys. Rev. D 21, 203 (1980).
- [4] L. Kluberg and H. Satz, in Relativistic Heavy Ion Physics, edited by R. Stock, Landolt-Börnstein
   Group I Elementary Particles, Nuclei and Atoms Vol. 23 (Springer, Berlin, Heidelberg, 2010).
- [5] A. Mocsy, P. Petreczky and M. Strickland, Int. Jour. Mod. Phys. A 28, 1340012 (2013).
- [6] S.F. Radford and W.W. Repko, Phys.Rev D 75, 074031 (2007).
- [7] Arata Hayashigaki, Phys. Lett. B 487, 96 (2000); T. Hilger, R. Thomas and B. Kämpfer, Phys. Rev. C bf 79, 025202 (2009); T. Hilger, B. Kämpfer and S. Leupold, Phys. Rev. C 84, 045202 (2011); S. Zschocke, T. Hilger and B. Kämpfer, Eur. Phys. J. A 47 151 (2011)
- [8] Sugsik Kim, Su Houng Lee, Nucl. Phys. A 679, 517 (2001).
- [9] F. Klingl, S. Kim, S. H. Lee, P. Morath and W. Weise, Phys. Rev. Lett. 82, 3396 (1999).
- [10] Arvind Kumar and Amruta Mishra, Phys. Rev. C 95, 065206 (2010)).
- [11] G. Krein, A.W. Thomas, K. Tsushima, Prog. Part. Nucl. Phys. 100, 161 (2018).
- [12] R. Molina, D. Gamermann, E. Oset, and L. Tolos, Eur. Phys. J A 42, 31 (2009); L. Tolos, R. Molina, D. Gamermann, and E. Oset, Nucl. Phys. A 827 249c (2009).
- [13] Arvind Kumar and Amruta Mishra, Phys. Rev. C 81, 065204 (2010).
- [14] Arvind Kumar and Amruta Mishra, Eur. Phys. A 47, 164 (2011).
- [15] M.E. Peskin, Nucl. Phys. **B156**, 365 (1979).
- [16] G. Bhanot and M.E. Peskin, Nucl. Phys. **B156**, 391 (1979).
- [17] M.B.Voloshin, Nucl. Phys. B 154, 365 (1979).
- [18] Su Houng Lee and Che Ming Ko, Phys. Rev. C 67, 038202 (2003).
- [19] K. Tuchin, Adv. High Energy Physics 2013, 490495 (2013).

- [20] P. Gubler, K. Hattori, S. H. Lee, M. Oka, S. Ozaki and K. Suzuki, Phys. Rev. D 93, 054026 (2016).
- [21] C. S. Machado, F. S. Navarra. E. G. de Oliveira and J. Noronha, Phys. Rev. D 88, 034009 (2013).
- [22] C. S. Machado, R.D. Matheus, S.I. Finazzo and J. Noronha, Phys. Rev. D 89, 074027 (2014).
- [23] Sushruth Reddy P, Amal Jahan CS, Nikhil Dhale, Amruta Mishra, J. Schaffner-Bielich, Phys. Rev. C 97, 065208 (2018).
- [24] Amal Jahan CS, Nikhil Dhale, Sushruth Reddy P, Shivam Kesarwani, Amruta Mishra, Phys. Rev. C 98, 065202 (2018).
- [25] S. Cho, K. Hattori, S. H. Lee, K. Morita and S. Ozaki, Phys. Rev. Lett. **113**, 122301 (2014).
- [26] S. Cho, K. Hattori, S. H. Lee, K. Morita and S. Ozaki, Phys. Rev. D 91, 045025 (2015).
- [27] J. Alford and M. Strickland, Phys. Rev. D 88, 105017 (2013).
- [28] K. Suzuki and S. H. Lee, Phys. Rev. C 96, 035203 (2017).
- [29] B. Friman, S. H. Lee and T. Song, Phys. Lett, B 548, 153 (2002).
- [30] Amruta Mishra, Amal Jahan C.S., Shivam Kesarwani, Haresh Raval, Shashank Kumar, Jitendra Meena, Eur. Phys. J. A 55, 99 (2019).
- [31] A. Le Yaouanc, L. Oliver, O. Pene and J.-C. Raynal, Phys. Rev. D 8 2223, (1973); ibid, Phys. Rev. D 9, 1415 (1974); ibid, Phys. Rev. D 11, 1272 (1975).
- [32] A. Le Yaouanc, L. Oliver, O. Pene and J.-C. Raynal, Phys. Lett. B 71 397 (1977).
- [33] E.S.Ackleh, T. Barnes and E. S. Swanson, Phys. Rev. D 54, 6811, 1996.
- [34] T. Barnes, F. E. Close, P. R. Page and E. S. Swanson, Phys. Rev. D 55, 4157 (1997).
- [35] Amruta Mishra, S. P. Misra and W. Greiner, Int. J. Mod. Phys. E 24, 155053 (2015).
- [36] P. Papazoglou, D. Zschiesche, S. Schramm, J. Schaffner-Bielich, H. Stöcker, and W. Greiner, Phys. Rev. C 59, 411 (1999).
- [37] S. P. Misra, Phys. Rev. D 18, 1661 (1978).
- [38] S. P. Misra, Phys. Rev. D 18, 1673 (1978).
- [39] S. P. Misra and L. Maharana, Phys. Rev. D 18, 4103 (1978).
- [40] A. Chodos, R. L. Jaffe, K. Johnson and C. B. Thorn, Phys. Rev. D 10, 2599 (1974).
- [41] S.P.Misra, K. Biswal and B. K. Parida, Phys. Rev. D 21, 2029 (1980).
- [42] Amruta Mishra and S. P. Misra, Phy. Rev. C 95, 065206 (2017).

- [43] Z.w Lin, C.M. Ko, B. Zhang, Phys. Rev. C 61, 024904 (2000); S.H. Lee, C. Song, H. Yabu, Phys.
   Lett. B 341, 407 (1995).
- [44] U. G. Meissner, Phys. Rep. 161, 213 (1988).
- [45] A. B. Kaidalov, P.E. Volkovitsky, Phys. Rev. Lett. 69, 3155 (1992).
- [46] M. Tanabashi et al (Particle Data Group), Phys. Rev. D 98, 030001 (2018) and 2019 update.