

On the shifted Littlewood-Richardson coefficients and the Littlewood-Richardson coefficients

Duc Khanh Nguyen

Abstract

We give a new interpretation of the shifted Littlewood-Richardson coefficients $f_{\lambda\mu}^\nu$ (λ, μ, ν are strict partitions). The coefficients $g_{\lambda\mu}$ which appear in the decomposition of Schur Q -function Q_λ into the sum of Schur functions $Q_\lambda = 2^{l(\lambda)} \sum_{\mu} g_{\lambda\mu} s_\mu$ can be considered as a special case of $f_{\lambda\mu}^\nu$ (here λ is a strict partition of length $l(\lambda)$). We also give another description for $g_{\lambda\mu}$ as the cardinal of a subset of a set that counts Littlewood-Richardson coefficients $c_{\mu^t\mu}^{\tilde{\lambda}}$. This new point of view allows us to establish connections between $g_{\lambda\mu}$ and $c_{\mu^t\mu}^{\tilde{\lambda}}$. More precisely, we prove that $g_{\lambda\mu} = g_{\lambda\mu^t}$, and $g_{\lambda\mu} \leq c_{\mu^t\mu}^{\tilde{\lambda}}$. We conjecture that $g_{\lambda\mu}^2 \leq c_{\mu^t\mu}^{\tilde{\lambda}}$ and formulate some conjectures on our combinatorial models which would imply this inequality if it is valid.

2020 Mathematics Subject Classification. 05E10, 05E05 (primary), 14M15 (secondary).

Key words and phrases. Young tableaux, Schur functions, Littlewood-Richardson coefficients, tableau switching, Grassmannians, Schubert varieties, shifted tableaux, Schur Q -functions, shifted Littlewood-Richardson coefficients, Lagrangian Grassmannians.

1 Introduction

Let λ, μ, ν be partitions. Let $l(\lambda)$ be the length of λ , and s_λ be the Schur function associated to the partition λ . The Littlewood-Richardson coefficients $c_{\lambda\mu}^\nu$ appear in the expansion (see [Ful97])

$$s_\lambda s_\mu = \sum_{\nu} c_{\lambda\mu}^\nu s_\nu. \quad (1)$$

If now λ, μ, ν are strict partitions, let Q_λ be the shifted Schur Q -function associated to the strict partition λ . The shifted Littlewood-Richardson coefficients appear in the expansion (see [Ste89])

$$Q_\lambda Q_\mu = \sum_{\nu} 2^{l(\lambda)+l(\mu)-l(\nu)} f_{\lambda\mu}^\nu Q_\nu. \quad (2)$$

For any strict partition λ , and a partition μ of the same integer, the coefficients $g_{\lambda\mu}$ appear in the decomposition (see [Ste89])

$$Q_\lambda = 2^{l(\lambda)} \sum_{\mu} g_{\lambda\mu} s_\mu. \quad (3)$$

The coefficients $g_{\lambda\mu}$ can be considered as shifted Littlewood-Richardson coefficients by the identity (see [Ste89])

$$g_{\lambda\mu} = f_{\lambda\delta}^{\mu+\delta}, \quad (4)$$

where $\delta = (l, l-1, \dots, 1)$ with $l = l(\mu)$.

There were several developments beyond the Littlewood-Richardson rule. For example,

- Zelevinsky [Zel81] expressed the coefficients $c'_{\lambda\mu}$ as the number of pictures between μ and ν/λ .
- Remmel and Whitney [RW84] described $c'_{\lambda\mu}$ as the number of standard tableaux of shape λ constructed by some rules from the reverse filling of the skew shape ν/μ . There are also a similar version by Chen, Garsia, Remmel [CGR84] but they replaced λ with ν and ν/μ with $\lambda * \mu$.
- White [Whi81] showed that the set of tableaux in the construction of Remmel and Whitney [RW84] can be understood from a different point of view. It arises from Robinson-Schensted insertion of reading words of column-strict tableaux of a fixed skew shape.

There are new approaches that come from geometry: the algorithm by Liu [Liu10] and the rule of Ravi Vakil [Vak06] etc.

The theory and methods for shifted Littlewood-Richardson coefficients are also developed parallelly with the theory of Littlewood-Richardson coefficients. Based on the work of Worley [Wor84], Sagan [Sag87], Stembridge [Ste89], there are several versions of the shifted Littlewood-Richardson rule for $f'_{\lambda\mu}$, for example, the works of Serrano [Ser10] and Shimozono [Shi99] and so on. The shifted Littlewood-Richardson rule given by Stembridge [Ste89] is also re-obtained by using the theory of crystal bases for the quantum queer superalgebra (see [GJK⁺14] and [CK18]). In [CNO14], the authors established the bijections between three models for shifted Littlewood-Richardson coefficients in [Ste89], [Ser10] and [GJK⁺14].

In this article, instead of re-interpreting the shifted Littlewood-Richardson rule given by Stembridge [Ste89], we use this rule to obtain a new combinatorial models for the coefficients $f'_{\lambda\mu}$ and $g_{\lambda\mu}$. The advantage of the new results allows us to compute the coefficients easier and to realize the connections with Littlewood-Richardson coefficients. The motivation of our work comes from the work of P. Belkale, S. Kumar and N. Ressayre [BKR12]. The main results in the article [BKR12] raised up some first clues about relations between shifted Littlewood-Richardson coefficients with Littlewood-Richardson coefficients. N. Ressayre conjectures an inequality between them in [Res19]. We do not use the approach from geometry as in [BKR12], but we try to develop the combinatorial model of Stembridge [Ste89] to discover the bridge between coefficients. To be more precise, we describe the results as follows.

Our first result, Theorem 4.2 is a new combinatorial model for the shifted Littlewood-Richardson coefficients. This is analogous to Remmel and Whitney's work [RW84]. The combinatorial model proposed by Shimozono in [Shi99] is analogous to White's model [Whi81], arising from Sagan's shifted insertion [Sag87]. Despite the case of Littlewood-Richardson coefficients where Remmel and Whitney's construction is identified with White's construction, our construction and Shimozono's construction do not produce the same model. Since $g_{\lambda\mu}$ can be considered as a shifted Littlewood-Richardson coefficient, we obtain a new model for $g_{\lambda\mu}$ in Theorem 4.3.

Our second result, Theorem 5.3 is also a new combinatorial interpretation of the coefficients $g_{\lambda\mu}$. More precisely, let $\tilde{\lambda}$ be the partition such that its Young diagram is the union of shifted diagram corresponding to λ and its reflection through the main diagonal. Let μ^t be the conjugate partition of μ . We prove that $g_{\lambda\mu}$ is the cardinal of a subset of a set that

counts the coefficients $c_{\mu^t\mu}^{\tilde{\lambda}}$. This implies Theorem 5.5 that

$$g_{\lambda\mu} \leq c_{\mu^t\mu}^{\tilde{\lambda}}. \tag{5}$$

We conjecture a stronger inequality (see Conjecture 5.6)

$$g_{\lambda\mu}^2 \leq c_{\mu^t\mu}^{\tilde{\lambda}}. \tag{6}$$

Using a computer program, we checked this conjecture on a lot of examples. Based on our combinatorial model for the coefficients $g_{\lambda\mu}$, we formulate Conjecture 5.7 whose validity implies Conjecture 5.6. An evidence for Conjecture 5.7 is that it implies easily the equality

$$g_{\lambda\mu} = g_{\lambda\mu^t}. \tag{7}$$

The equality (7) might be well known among experts, nevertheless we include a geometric proof in Proposition 3.6.

The article contains four sections. In the first section, we collect some basic background about the theory of Young tableaux, and related models for Littlewood-Richardson coefficients. In the second section, we present the theory of shifted tableaux, and related models, some interpretations for shifted Littlewood-Richardson coefficients. The last two sections present our two main results on the coefficients $f_{\lambda\mu}^{\nu}$ and $g_{\lambda\mu}$.

Contents

1	Introduction	1
2	The Littlewood-Richardson coefficients	4
2.1	Young tableaux	4
2.2	Row-insertion and product tableau	4
2.3	Sliding and jeu de taquin	5
2.4	The Robinson-Schensted-Knuth correspondence	6
2.5	Littlewood-Richardson rule	7
2.6	Tableau switching	10
2.7	The symmetry of Littlewood-Richardson coefficients	12
3	The shifted Littlewood-Richardson coefficients	12
3.1	Shifted tableaux	12
3.2	Shifted jeu de taquin	13
3.3	Shifted Littlewood-Richardson rule	14
3.4	Geometric interpretation of the coefficients $f_{\lambda\mu}^{\nu}$ and $g_{\lambda\mu}$	15
3.5	Application to the identity $g_{\lambda\mu} = g_{\lambda\mu^t}$	16
4	A new combinatorial models for the coefficients $f_{\lambda\mu}^{\nu}$	17
5	On the coefficients $g_{\lambda\mu}$	21
5.1	A new interpretation of the coefficients $g_{\lambda\mu}$	21
5.2	Inequality $g_{\lambda\mu} \leq c_{\mu\mu^t}^{\tilde{\lambda}}$	23
5.3	Inequality $g_{\lambda\mu}^2 \leq c_{\mu^t\mu}^{\tilde{\lambda}}$	24
	Acknowledgements	29

2 The Littlewood-Richardson coefficients

In this section, we present Young tableaux, and related models for Littlewood-Richardson coefficients.

2.1 Young tableaux

For a **partition** λ , we write $\lambda = (\lambda_1, \lambda_2, \dots)$ with $\lambda_1 \geq \lambda_2 \geq \dots$. If $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ with $\lambda_l > 0$ and $\sum_{i=1}^l \lambda_i = n$, we write $l(\lambda) = l$, $|\lambda| = n$.

Each partition λ is presented by a **Young diagram** $Y(\lambda)$.

Example 2.1.

$$Y((3, 2)) = \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array}$$

The reflection $\sigma(Y)$ through the main diagonal of a Young diagram Y is also a Young diagram. The **conjugate partition** λ^t of λ is defined by $\sigma(Y(\lambda)) = Y(\lambda^t)$.

A **semistandard Young tableau** of shape λ is a filling of the Young diagram $Y(\lambda)$ by the ordered alphabet $\{1 < 2 < \dots\}$ such that:

(Y1) The entries in each column are strictly increasing.

(Y2) The entries in each row are weakly increasing.

Let $\nu = (\nu_1, \nu_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$ be two partitions. We say that ν is bigger than μ if and only if $\nu_i \geq \mu_i$ for all i , and we write $\nu \geq \mu$. In this case, we define the **skew Young diagram** $Y(\nu/\mu)$ as the result of removing boxes in the Young diagram $Y(\mu)$ from the Young diagram $Y(\nu)$. We write $|\nu/\mu| = |\nu| - |\mu|$. A **skew Young tableau** T of skew shape ν/μ is a result of filling the skew Young diagram $Y(\nu/\mu)$ by the ordered alphabet $\{1 < 2 < \dots\}$ satisfying the rules (Y1) and (Y2).

The **word** $w(T)$ of a Young tableau T is defined to be the sequence obtained by reading the rows of T from left to right, starting from bottom to top. A Young tableau of skew shape ν/μ is said to be a **standard skew Young tableau** if its word is a permutation of the word $12\dots|\nu/\mu|$. The **transpose** of a standard skew Young tableau T is also a standard skew Young tableau and it is denoted by T^t .

2.2 Row-insertion and product tableau

For a Young tableau T and a positive integer x , we recall **row-insertion** x to T from [Ful97]. The result of row-insertion x to T is a Young tableau, is denoted by $T \leftarrow x$.

Example 2.2.

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 2 \\ \hline 3 & 4 & \\ \hline \end{array} \leftarrow 1 = \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 4 & \\ \hline 3 & & \\ \hline \end{array}$$

Let T and U be Young tableaux, the **product tableau** $T.U$ is defined by

$$T.U := (\dots((T \leftarrow x_1) \leftarrow x_2) \leftarrow \dots \leftarrow x_{n-1}) \leftarrow x_n, \quad (8)$$

where $w(U) = x_1 x_2 \dots x_n$.

Example 2.3. *Let*

$$T = \begin{array}{|c|c|c|} \hline 1 & 2 & 2 \\ \hline 3 & 4 & \\ \hline \end{array} \quad \text{and} \quad U = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 2 & 4 & \\ \hline \end{array}$$

then $w(U) = 24123$ and

$$T.U = \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 2 & 2 & 2 & 3 \\ \hline 2 & 4 & 4 & & & \\ \hline 3 & & & & & \\ \hline \end{array}$$

2.3 Sliding and jeu de taquin

For the skew Young diagram $Y(\nu/\mu)$, an **inner corner** of $Y(\nu/\mu)$ is a box in the Young diagram $Y(\mu)$ such that the boxes below and to the right are not in $Y(\mu)$. An **outside corner** is a box in the Young diagram $Y(\nu)$ such that the boxes below and to the right are not in $Y(\nu)$.

Let T be a skew Young tableau of skew shape ν/μ . Let b be an inner corner of ν/μ . We recall **sliding** b out of T from [Ful97]. The result of applying sliding b out of T gives us a new skew Young tableau T' of skew shape ν'/μ' such that $|\nu'| = |\nu| - 1$, $|\mu'| = |\mu| - 1$. Choose a random inner corner b' of T' and do sliding b' out of T' as before, we get a new skew Young tableau T'' of skew shape ν''/μ'' such that $|\nu''| = |\nu| - 2$, $|\mu''| = |\mu| - 2$. So repeat the process as many times as possible, we finally get a Young tableau and the process will terminate. There is a fact that the Young tableau we get does not depend on the choice of random inner corners in each step. The final tableau we have obtained is called the **rectification** of T and it is denoted by $Rect(T)$. The whole process we apply on T to get $Rect(T)$ is called the **jeu de taquin**.

Lemma 2.4. *Let T and U be skew Young tableaux. If $w(T) = w(U)$ then $Rect(T) = Rect(U)$.*

Example 2.5. *Let*

$$T = \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 1 & 2 & 2 & \\ \hline 3 & 4 & & \\ \hline \end{array} \quad \text{and} \quad U = \begin{array}{|c|c|c|c|c|} \hline & & & & 1 \\ \hline & 1 & 2 & 2 & \\ \hline & 4 & & & \\ \hline 3 & & & & \\ \hline \end{array}$$

The process of applying the jeu de taquin on T can be visualized as follows:


$$\begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 1 & 2 & 2 & \\ \hline 3 & 4 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 1 & 2 & 2 & \\ \hline 3 & 4 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline & 1 & 2 \\ \hline 1 & 2 & \\ \hline 3 & 4 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 4 & \\ \hline 3 & & \\ \hline \end{array}$$

where the boxes in red are chosen to be slid. Hence,

$$\text{Rect}(T) = \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 4 & \\ \hline 3 & & \\ \hline \end{array}$$

One can easily check that $\text{Rect}(U) = \text{Rect}(T)$.

Let T and U be Young tableaux. We denote $T * U$ the new skew Young tableau which is defined as follows:



We have another point of view about the product tableau $T.U$.

Lemma 2.6. *Let T and U be Young tableaux. We have $T.U = \text{Rect}(T * U)$.*

2.4 The Robinson-Schensted-Knuth correspondence

A two-rowed array is defined by

$$w = \begin{pmatrix} u_1 & \dots & u_n \\ v_1 & \dots & v_n \end{pmatrix}, \quad (9)$$

with u_i 's and v_i 's are in two independent alphabets. We say that w is in **lexicographic order** if

1. $u_1 \leq u_2 \leq \dots \leq u_n$.
2. If $u_{k-1} = u_k$ for some k , then $v_{k-1} \leq v_k$.

The **Robinson-Schensted-Knuth correspondence** is mentioned in [Ful97], which set

up an one-to-one correspondence between a two-rowed array in lexicographic order $\begin{pmatrix} u_1 & \dots & u_n \\ v_1 & \dots & v_n \end{pmatrix}$

and a pair of tableaux of the same shape $\begin{pmatrix} Q \\ P \end{pmatrix}$. We write

$$\begin{pmatrix} u_1 & \dots & u_n \\ v_1 & \dots & v_n \end{pmatrix} \xleftrightarrow{RSK} \begin{pmatrix} Q \\ P \end{pmatrix}. \quad (10)$$

Example 2.7. *Let w be the two-rowed array*

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 2 & 3 & 6 & 7 & 4 & 5 & 1 \end{pmatrix}. \quad (11)$$

The tableaux P and Q are

$$P = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 4 & 5 \\ \hline 2 & 7 & & \\ \hline 6 & & & \\ \hline \end{array} \text{ and } Q = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & & \\ \hline 7 & & & \\ \hline \end{array}$$

2.5 Littlewood-Richardson rule

A Young tableau T is said to have **content** $\gamma = (\gamma_1, \gamma_2, \dots)$ if γ_i is the number of entries i in the tableau T . We write

$$x^T = x^\gamma = x_1^{\gamma_1} x_2^{\gamma_2} \dots \quad (12)$$

For each partition λ , the **Schur function** s_λ in variables x_1, x_2, \dots is defined as the sum of x^T , where T runs over the semistandard Young tableaux of shape λ . Let $\Lambda = \bigoplus_{n \geq 0} \Lambda^n$ be the **graded ring of symmetric functions** in the variables x_1, x_2, \dots with coefficients in \mathbb{Z} . The following set is a \mathbb{Z} -basis of Λ^n

$$\{s_\lambda \mid \lambda \text{ is a partition of } n\}.$$

The integers $c'_{\lambda\mu}$ for each partitions λ, μ, ν defined by

$$s_\lambda s_\mu = \sum_{\nu} c'_{\lambda\mu} s_\nu, \quad (13)$$

are called **Littlewood-Richardson coefficients**.

For any Young tableaux V_0 of shape ν , let $\mathcal{T}(\lambda, \mu, V_0)$ be the set

$$\{(\Lambda, U) \mid \Lambda, U \text{ are Young tableaux of shapes } \lambda, \mu, \text{ respectively and } \Lambda.U = V_0\}. \quad (14)$$

For any tableau U_0 of shape μ , let $\mathcal{S}(\nu/\lambda, U_0)$ be the set

$$\{\text{Skew tableaux } S \text{ of skew shape } \nu/\lambda \text{ such that } \text{Rect}(S) = U_0\}. \quad (15)$$

We can describe a canonical one-to-one correspondence between $\mathcal{T}(\lambda, \mu, V_0)$ and $\mathcal{S}(\nu/\lambda, U_0)$ as follows:

1. Let (Λ, U) be an element of the set $\mathcal{T}(\lambda, \mu, V_0)$. Suppose that

$$\begin{pmatrix} U \\ U_0 \end{pmatrix} \xleftrightarrow{RSK} \begin{pmatrix} u_1 & \dots & u_m \\ v_1 & \dots & v_m \end{pmatrix}, \quad (16)$$

where $m = |\mu|$. Let S be the new skew tableau obtained by placing u_1, \dots, u_m into the new boxes while doing row-insertion v_1, \dots, v_m into Λ . Then S is an element of $\mathcal{S}(\nu/\lambda, U_0)$.

2. Let S be an element of $\mathcal{S}(\nu/\lambda, U_0)$. Let Λ' be an arbitrary Young tableau of shape λ . Put an order on the letters in Λ' and S in such a way that all letters in Λ' are smaller than those in S . Now, suppose that

$$\begin{pmatrix} V_0 \\ \Lambda' \cup S \end{pmatrix} \xleftrightarrow{RSK} \begin{pmatrix} t_1 & \dots & t_n & u_1 & \dots & u_m \\ x_1 & \dots & x_n & v_1 & \dots & v_m \end{pmatrix}, \quad (17)$$

where $n = |\lambda|$. Then we can construct a tableau Λ such that $(\Lambda, U) \in \mathcal{T}(\lambda, \mu, V_0)$ by

$$\begin{pmatrix} t_1 & \dots & t_m \\ x_1 & \dots & x_m \end{pmatrix} \xleftrightarrow{RSK} \begin{pmatrix} \Lambda \\ \Lambda' \end{pmatrix}. \quad (18)$$


3. Denote by $\mathcal{F}_{\nu/\lambda, U_0}^{\lambda, \mu, V_0}$ the map that sends (Λ, U) in $\mathcal{T}(\lambda, \mu, V_0)$ to S in $\mathcal{S}(\nu/\lambda, U_0)$. This map is a bijection between two sets.

A method to compute the set $\mathcal{S}(\nu/\lambda, U_0)$ is explained in subsection 2.7. In particular case when U_0 is the Young tableau \mathcal{U}_μ of shape μ whose all entries in the k^{th} row are equal to k , one can compute explicitly the set $\mathcal{S}(\nu/\lambda, \mathcal{U}_\mu)$ by the model of Remmel and Whitney [RW84] as follows:

1. With the skew shape ν/λ , we number the boxes from top to bottom and right to left in each row by $1, 2, \dots, |\nu/\lambda|$, respectively. The result is called the **reverse filling of the skew shape ν/λ** . We denote it by $T_{\nu/\lambda}$.
2. Define $\mathcal{O}(\nu/\lambda)$ to be the set of Young tableaux T of size $|\nu/\lambda|$, constructed from $T_{\nu/\lambda}$ satisfying the following conditions:

(R1) If k and $k + 1$ appear in the same row of $T_{\nu/\lambda}$, then $k + 1$ appears weakly above and strictly right of k in T .

(R2) If h appears in the box directly below k in $T_{\nu/\lambda}$, then h appears strictly below and weakly left of k in T .



3. Let $\mathcal{O}_\mu(\nu/\lambda)$ be the set of all tableaux T in $\mathcal{O}(\nu/\lambda)$ of shape μ . For each T in $\mathcal{O}_\mu(\nu/\lambda)$, we construct a word $x_{|\mu|} \dots x_1$, where x_k is the row where k belongs to in the tableau T . There exists a unique skew Young tableau T^* of skew shape ν/λ such that $w(T^*)$ is the word we have created.
4. It is proved that the set $\mathcal{S}(\nu/\lambda, \mathcal{U}_\mu)$ is the set of all skew tableaux T^* where T runs over the set of all tableaux in $\mathcal{O}_\mu(\nu/\lambda)$.

Theorem 2.8. *Let λ, μ, ν be partitions. Let T_0 be a Young tableau of shape ν and U_0 be a Young tableau of shape μ . We have $c'_{\lambda\mu} = \#\mathcal{O}_\mu(\nu/\lambda) = \#\mathcal{S}(\nu/\lambda, U_0) = \#\mathcal{T}(\lambda, \mu, V_0)$.*

The example below shows how to compute the Littlewood-Richardson coefficients and the three models presented in Theorem 2.8.

Example 2.9. Set $\lambda = (3, 2, 1, 1)$, $\mu = (4, 2, 1)$ and $\nu = (6, 4, 2, 1, 1)$. Then

$$T_{\nu/\lambda} = \begin{array}{|c|c|c|c|c|c|} \hline & & & 3 & 2 & 1 \\ \hline & & 5 & 4 & & \\ \hline & 6 & & & & \\ \hline & & & & & \\ \hline 7 & & & & & \\ \hline \end{array}$$

All tableaux of the set $\mathcal{O}_\mu(\nu/\lambda)$ are

$$\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 7 \\ \hline 4 & 5 & & \\ \hline 6 & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 6 \\ \hline 4 & 5 & & \\ \hline 7 & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 5 \\ \hline 4 & 6 & & \\ \hline 7 & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 5 \\ \hline 4 & 7 & & \\ \hline 6 & & & \\ \hline \end{array}$$

Hence, $c'_{\lambda\mu} = 4$. We have

$$\mathcal{U}_\mu = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 1 \\ \hline 2 & 2 & & \\ \hline 3 & & & \\ \hline \end{array}$$

All tableaux of the set $\mathcal{S}(\nu/\lambda, \mathcal{U}_\mu)$ are

$$\begin{array}{|c|c|c|c|c|c|} \hline & & & 1 & 1 & 1 \\ \hline & & 2 & 2 & & \\ \hline & 3 & & & & \\ \hline & & & & & \\ \hline 1 & & & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|c|c|} \hline & & & 1 & 1 & 1 \\ \hline & & 2 & 2 & & \\ \hline & 1 & & & & \\ \hline & & & & & \\ \hline 3 & & & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|c|c|} \hline & & & 1 & 1 & 1 \\ \hline & & 1 & 2 & & \\ \hline & 2 & & & & \\ \hline & & & & & \\ \hline 3 & & & & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|c|c|} \hline & & & 1 & 1 & 1 \\ \hline & & 1 & 2 & & \\ \hline & 3 & & & & \\ \hline & & & & & \\ \hline 2 & & & & & \\ \hline \end{array}$$

Set

$$V_0 = \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 2 & 3 & 4 & 5 \\ \hline 2 & 6 & 6 & 7 & & \\ \hline 3 & 7 & & & & \\ \hline 4 & & & & & \\ \hline 5 & & & & & \\ \hline \end{array} \quad \text{and} \quad \Lambda' = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline 4 & & \\ \hline \end{array}$$

with order that $1 < 2 < 3 < 4 < 1 < 2 < 3$. Set

$$S = \begin{array}{|c|c|c|c|c|c|} \hline & & & 1 & 1 & 1 \\ \hline & & 2 & 2 & & \\ \hline & 3 & & & & \\ \hline & & & & & \\ \hline 1 & & & & & \\ \hline \end{array} \in \mathcal{S}(\nu/\lambda, \mathcal{U}_\mu)$$

The two-rowed array corresponding to the pair $\begin{pmatrix} V_0 \\ \Lambda' \cup S \end{pmatrix}$ is

$$\begin{pmatrix} 1112234 & 1111223 \\ 1562432 & 1377465 \end{pmatrix}. \tag{19}$$

The tableaux Λ and U such that $(\Lambda, U) \in \mathcal{T}(\lambda, \mu, V_0)$ corresponding to S are

$$\Lambda = \begin{array}{|c|c|c|} \hline 1 & 2 & 2 \\ \hline 3 & 6 & \\ \hline 4 & & \\ \hline 5 & & \\ \hline \end{array} \quad \text{and} \quad U = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 4 & 5 \\ \hline 6 & 7 & & \\ \hline 7 & & & \\ \hline \end{array}$$

2.6 Tableau switching

In this subsection, we recall the definition and basic properties of the switching procedure. The main reference is the article [BSS96] by G. Benkart, F. Sottile, J. Stroomer.

For each skew shape γ , we define a **perforated tableau** T of shape γ to be a result of filling some boxes in $Y(\gamma)$ with integers such that:

(PT1) The entries in each column are strictly increasing.

(PT2) The entries in the northwest of each entry t of T are less than or equal to t .

Let S, T be perforated tableaux of shape γ . We say that S, T **completely fill** γ if all boxes in $Y(\gamma)$ are filled by entries of S or T , and no box is filled twice. We then call $S \cup T$ a **perforated pair** of shape γ .

Let $S \cup T$ be a perforated pair of shape γ . Let s in S and t in T be adjacent integers, t is below or in the right of s . We define **switching** $s \leftrightarrow t$ by interchanging s and t such that after the action, both perforated tableau of shape γ filled by entries t , and perforated tableau of shape γ filled by entries s satisfy the conditions (PT1) and (PT2).

Choose a random pair (s, t) in $S \cup T$ such that we can do the switching $s \leftrightarrow t$. Repeat this process until there are no more pair (s, t) in $S \cup T$ that can be switched $s \leftrightarrow t$. The result is a new perforated pair $T' \cup S'$ of shape γ , where S' is the perforated tableau filled by entries s and T' is the perforated tableau filled by entries t . The point is that the resulting pair $T' \cup S'$ does not depend on the choices, it is denoted by ${}^S T \cup S_T$ (see [BSS96]). The process we have done to produce ${}^S T \cup S_T$ from $S \cup T$ is called the **switching procedure**. The map that sends $S \cup T$ to ${}^S T \cup S_T$ is called the **switching map**.

The example below visualizes switching procedure.

Example 2.10. Let $\gamma = (4, 3, 3, 2)/(2, 1)$. The tableau S with red entries and the tableau T with blue entries below are perforated tableaux of shape γ .

$$S = \begin{array}{|c|c|c|c|} \hline & & 1 & \\ \hline & & & \\ \hline 1 & & 2 & \\ \hline & 3 & & \\ \hline \end{array} \quad T = \begin{array}{|c|c|c|c|} \hline & & & -1 \\ \hline & -2 & -2 & \\ \hline & 1 & & \\ \hline 2 & & & \\ \hline \end{array} \quad S \cup T = \begin{array}{|c|c|c|c|} \hline & & 1 & -1 \\ \hline & -2 & -2 & \\ \hline 1 & 1 & 2 & \\ \hline 2 & 3 & & \\ \hline \end{array}$$

Look at the entries inside the circles below


$$\begin{array}{|c|c|c|c|} \hline & & 1 & -1 \\ \hline & -2 & -2 & \\ \hline (1) & (1) & 2 & \\ \hline (2) & 3 & & \\ \hline \end{array}$$

We see that we can just switch $1 \leftrightarrow 1$, but we cannot switch $1 \leftrightarrow 2$. Indeed, after switching $1 \leftrightarrow 1$, we get

		1	-1
	-2	-2	
1	1	2	
2	3		

The new tableau formed by the red entries and the new tableau formed by blue entries satisfy the conditions (PT1), (PT2). But after switching $1 \leftrightarrow 2$, the new tableau formed by the blue entries does not satisfy the condition (PT2).

Here is the visualization of switching procedure with starting point $S \cup T$ (we choose pairs in circles to switch).



Hence,

$$S_T = \begin{array}{|c|c|c|c|} \hline & & -2 & -1 \\ \hline & -2 & & \\ \hline 1 & & & \\ \hline 2 & & & \\ \hline \end{array} \quad \text{and} \quad S_T = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & 1 & \\ \hline & 1 & 2 & \\ \hline & 3 & & \\ \hline \end{array}$$

Let S, T be skew tableaux. We say that T extends S if T has skew shape ν/λ and S has shape λ/μ for some partitions $\nu \geq \lambda \geq \mu$. The following theorem is a collection of some important properties in Theorem 2.2 and Theorem 3.1 in the article [BSS96].

Theorem 2.11. *Let S, T be skew Young tableaux such that T extends S . Then*

1. S_T and ${}^S T$ are skew Young tableaux, S_T extends ${}^S T$.
2. ${}^S T \cup S_T$ has the same shape as $S \cup T$.
3. $\text{Rect}(S) = \text{Rect}(S_T)$.
4. $\text{Rect}(T) = \text{Rect}({}^S T)$.
5. The switching map $S \cup T \mapsto {}^S T \cup S_T$ is an involution.

Example 2.12. *Let*

$$S = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline 4 & & \\ \hline \end{array} \quad \text{and} \quad T = \begin{array}{|c|c|c|c|c|c|} \hline & & & 1 & 1 & 1 \\ \hline & & 2 & 2 & & \\ \hline & 3 & & & & \\ \hline & & & & & \\ \hline 1 & & & & & \\ \hline \end{array}$$

Then T extends S and

$$s_T = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 1 \\ \hline 2 & 2 & & \\ \hline 3 & & & \\ \hline \end{array} \quad \text{and} \quad S_T = \begin{array}{|c|c|c|c|c|c|} \hline & & & & 1 & 1 \\ \hline & & & 1 & 2 & \\ \hline & & 2 & & & \\ \hline 3 & & & & & \\ \hline 4 & & & & & \\ \hline \end{array}$$

2.7 The symmetry of Littlewood-Richardson coefficients

The tableau switching provides a bijective proof of the symmetry of Littlewood-Richardson coefficients

$$c_{\lambda\mu}^\nu = c_{\mu\lambda}^\nu. \quad (20)$$

Indeed, let Λ_0 be a Young tableau of shape λ and U_0 be a Young tableau of shape μ . We can describe a one-to-one correspondence between $\mathcal{S}(\nu/\mu, \Lambda_0)$ and $\mathcal{S}(\nu/\lambda, U_0)$ by tableau switching as follows:

1. Let S be an element of $\mathcal{S}(\nu/\lambda, U_0)$. The switching map sends $\Lambda_0 \cup S$ to ${}^{\Lambda_0}S \cup (\Lambda_0)_S$. By Theorem 2.11, we have ${}^{\Lambda_0}S = U_0$ and $\text{Rect}((\Lambda_0)_S) = \Lambda_0$. Hence, $(\Lambda_0)_S \in \mathcal{S}(\nu/\mu, \Lambda_0)$.
2. By Theorem 2.11, the switching map is an involution. Hence, the map that sends S to $(\Lambda_0)_S$ is a bijection between $\mathcal{S}(\nu/\lambda, U_0)$ and $\mathcal{S}(\nu/\mu, \Lambda_0)$. We denote this map by $\mathcal{B}_{\nu/\mu, \Lambda_0}^{\nu/\lambda, U_0}$.

Let V_0 and W_0 be Young tableaux of shape ν . The composition of the bijections below

$$\mathcal{T}(\lambda, \mu, V_0) \xrightarrow{\mathcal{F}_{\nu/\lambda, U_0}^{\lambda, \mu, V_0}} \mathcal{S}(\nu/\lambda, U_0) \xrightarrow{\mathcal{B}_{\nu/\mu, \Lambda_0}^{\nu/\lambda, U_0}} \mathcal{S}(\nu/\mu, \Lambda_0) \xrightarrow{\left(\mathcal{F}_{\nu/\mu, \Lambda_0}^{\mu, \lambda, W_0}\right)^{-1}} \mathcal{T}(\mu, \lambda, W_0) \quad (21)$$

gives us a bijection between the set $\mathcal{T}(\lambda, \mu, V_0)$ and the set $\mathcal{T}(\mu, \lambda, W_0)$. We denote this map by $\mathcal{S}_{V_0, U_0, \Lambda_0, W_0}^{\lambda, \mu, \nu}$.

Remark 2.13. Subsection 2.5 provides an algorithm to determine the set $\mathcal{S}(\nu/\lambda, \mathcal{U}_\mu)$. Applying then $\mathcal{B}_{\nu/\mu, \Lambda_0}^{\nu/\lambda, \mathcal{U}_\mu}$, we get an algorithm to compute $\mathcal{S}(\nu/\mu, \Lambda_0)$ for any Λ_0 .

3 The shifted Littlewood-Richardson coefficients

In this section, we present the definition and Stembridge's models, geometric points of view for shifted Littlewood-Richardson coefficients.

3.1 Shifted tableaux

A partition $\lambda = (\lambda_1, \lambda_2, \dots)$ is said to be **strict** if $\lambda_1 > \lambda_2 > \dots$.

Each strict partition λ is presented by a **shifted diagram** $sY(\lambda)$ that is a collection of boxes such that:

(SD1) The leftmost boxes of each row are in the main diagonal.

(SD2) The number of boxes from top row to bottom row are $\lambda_1, \lambda_2, \dots$, respectively.

A **shifted tableau** T of shifted shape λ is a result of filling the shifted diagram $sY(\lambda)$ by the ordered alphabet $\{1' < 1 < 2' < 2 < \dots\}$ such that

(T1) The entries in each column and in each row are weakly increasing.

(T2) The entries k' in each row are strictly increasing.


(T3) The entries k in each column are strictly increasing.

The shifted tableau T is said to have **content** $\gamma = (\gamma_1, \gamma_2, \dots)$ if γ_i is the number of i or i' in T . We write


$$x^T = x^\gamma = x_1^{\gamma_1} x_2^{\gamma_2} \dots \quad (22)$$

Let $\nu = (\nu_1, \nu_2, \dots)$ and $\mu = (\mu_1, \mu_2, \dots)$ be two strict partitions with $\nu \geq \mu$. We define the **skew shifted diagram** $sY(\nu/\mu)$ as the result of removing boxes in shifted diagram $sY(\mu)$ from shifted diagram $sY(\nu)$. A **skew shifted tableau** T of skew shifted shape ν/μ is a result of filling the shifted diagram $sY(\nu/\mu)$ by the ordered alphabet $\{1' < 1 < 2' < 2 < \dots\}$ satisfying the rules (T1), (T2) and (T3). The **content** of a skew shifted tableau T is defined by the same way as for a shifted tableau.

Example 3.1. Let $\lambda = (4, 2, 1)$. Then the shifted diagram $sY(\lambda)$ is



And




is a shifted tableau of shifted shape $(4, 2, 1)$. The content of T is $(1, 4, 1, 1)$.

3.2 Shifted jeu de taquin


For the skew shifted diagram $sY(\nu/\mu)$, we also define **inner corners** and **outside corners** by the same way as for the case of skew Young diagrams. Let T be a skew shifted tableau of skew shifted shape ν/μ without entries k' . Let b be an inner corner of skew shifted diagram $sY(\nu/\mu)$, we define **shifted sliding** b out of T , and **shifted jeu de taquin** on T , **shifted rectification** of T which we denote by $sRect(T)$, by the same way as for the case of skew Young tableaux.

Here is an example of shifted jeu de taquin.

Example 3.2. Set



The process of applying the shifted jeu de taquin on T can be visualized as follows:



where the boxes in red are chosen to be slid. Hence,

$$sRect(T) = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 5 \\ \hline & 4 & & \\ \hline \end{array}$$

3.3 Shifted Littlewood-Richardson rule

The **Schur Q -function** $Q_\lambda = Q_\lambda(x)$ in variables x_1, x_2, \dots is defined as the sum of x^T where T runs over the shifted tableaux of shape λ . Since every coefficient in Q_λ is divisible by $2^{l(\lambda)}$, we can define a formal power series with integer coefficients

$$P_\lambda(x) = 2^{-l(\lambda)} Q_\lambda(x). \quad (23)$$

We define the **power-sum symmetric function** p_r with $r \geq 1$ by

$$p_r = x_1^r + x_2^r + \dots \quad (24)$$

For each partition $\lambda = (\lambda_1, \lambda_2, \dots)$, we define

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \dots \quad (25)$$

The following set is a \mathbb{Z} -basis of Λ^n

$$\{p_\lambda \mid \lambda \text{ is a partition of } n\}.$$

Let $\Omega_{\mathbb{Q}} = \bigoplus_{n \geq 0} \Omega_{\mathbb{Q}}^n$ be the graded subalgebra of $\Lambda_{\mathbb{Q}} = \mathbb{Q} \otimes_{\mathbb{Z}} \Lambda$ generated by $1, p_1, p_3, p_5, \dots$. Let $\Omega = \Omega_{\mathbb{Q}} \cap \Lambda$ be the \mathbb{Z} -hyper subring of $\Omega_{\mathbb{Q}}$. We write $\Omega = \bigoplus_{n \geq 0} (\Omega_{\mathbb{Q}}^n \cap \Lambda)$ as a graded ring. Since

$$\{P_\lambda \mid \lambda \text{ is a strict partition of } n\}$$

is a \mathbb{Z} -basis of $\Omega_{\mathbb{Q}}^n \cap \Lambda$, we can define integers $f_{\lambda\mu}^\nu$ for each strict partitions λ, μ, ν by

$$P_\lambda P_\mu = \sum_{\nu} f_{\lambda\mu}^\nu P_\nu. \quad (26)$$

The integers $f_{\lambda\mu}^\nu$ are called the **shifted Littlewood-Richardson coefficients**.

For any (skew) shifted tableau T , we define the **word** $w(T)$ to be the sequence obtained by reading the rows of T from left to right, starting from bottom to top.

Given a word $w = w_1 w_2 \dots w_n$ over the alphabet $\{1' < 1 < 2' < 2 < \dots\}$, we define a sequence of statistics $m_i(j)$ ($0 \leq j \leq 2n, i \geq 1$) as follows:

$$\begin{aligned} m_i(j) &= \text{multiplicity of } i \text{ among } w_n \dots w_{n-j+1} & (0 \leq j \leq n), \\ m_i(j) &= \text{multiplicity of } i' \text{ among } w_1 \dots w_{j-n} \\ &+ \text{multiplicity of } i \text{ among } w_n \dots w_1 & (n < j \leq 2n). \end{aligned}$$

We say that the word w is a **shifted lattice word** if, whenever $m_i(j) = m_{i-1}(j)$, the next letter to be read after j^{th} step (it is w_{n-j} if $0 \leq j < n$ and w_{j+1-n} if $n \leq j < 2n$) is not be i, i' if $0 \leq j < n$ and not be $i-1, i'$ if $n \leq j < 2n$.

Stembridge in [Ste89] obtained a shifted analogue of the Littlewood-Richardson rule as follows.

Theorem 3.3. *Let λ, μ, ν be strict partitions. Then the coefficient $f_{\lambda\mu}^\nu$ is the number of skew shifted tableaux T of skew shifted shape ν/μ and content λ satisfying*

(F1) *The leftmost letter of $\{i, i'$ in $w(T)\}$ is unmarked ($1 \leq i \leq l(\lambda)$).*

(F2) *The word $w(T)$ is a shifted lattice word.*

For each strict partition λ and partition μ of the same integer n , let $g_{\lambda\mu}$ be the integer defined by

$$P_\lambda = \sum_{|\mu|=n} g_{\lambda\mu} s_\mu. \quad (27)$$

In the proof of Theorem 9.3 in [Ste89], Stembridge used the fact that

$$g_{\lambda\mu} = f_{\lambda\delta}^{\mu+\delta}, \quad (28)$$

where

$$\mu = (\mu_1, \mu_2, \dots, \mu_l) \text{ with } l = l(\mu), \quad (29)$$

$$\delta = (l, l-1, \dots, 1), \quad (30)$$

$$\mu + \delta = (\mu_1 + l, \mu_2 + l - 1, \dots, \mu_l + 1). \quad (31)$$

With the identity (28), he obtained an explicit interpretation of $g_{\lambda\mu}$ as in the following theorem.

Theorem 3.4. *Let λ be a strict partition and μ be a partition. Then the coefficient $g_{\lambda\mu}$ is the number of skew shifted tableaux T of shape μ and content λ satisfying*

(G1) *The leftmost letter of $\{i, i'$ in $w(T)\}$ is unmarked ($1 \leq i \leq l(\lambda)$).*

(G2) *The word $w(T)$ is a shifted lattice word.*

A skew shifted tableau of skew shifted shape ν/μ is said to be **standard** if its word is a permutation of the word $12 \dots |\nu/\mu|$. The following result can be translated equivalently from Lemma 8.4 in the article [Ste89] of J. Stembridge.

Theorem 3.5. *Let λ, μ, ν be strict partitions. Choose a standard shifted tableau \mathcal{T}_λ of shifted shape λ . Then the coefficient $f_{\lambda\mu}^\nu$ is the number of standard skew shifted tableaux S of skew shifted shape ν/μ such that $sRect(S) = \mathcal{T}_\lambda$.*

3.4 Geometric interpretation of the coefficients $f_{\lambda\mu}^\nu$ and $g_{\lambda\mu}$

Let V be a complex vector space of dimension $m+n$. The set $Gr(m, V)$ of linear subspaces of dimension m in V is called a **complex Grassmannian**. Fix a complete flag of V

$$\mathcal{F} : 0 = V_0 \subset \dots \subset V_i \subset \dots \subset V_{m+n} = V, \quad (32)$$

where each V_i is a vector subspace of V of dimension i . To each partition $\lambda = (\lambda_1, \dots, \lambda_m)$ with $\lambda_m \geq 0$, contained in the $m \times n$ rectangle, we associate the Schubert variety

$$X_\lambda(\mathcal{F}) = \{W \in Gr(m, V) \mid \dim(W \cap V_{n+i-\lambda_i}) \geq i \ (1 \leq i \leq m)\}. \quad (33)$$

The Poincare dual class of $X_\lambda(\mathcal{F})$ is denoted by σ_λ and called a Schubert class. Then σ_λ is an element of $H^{2|\lambda|}(Gr(m, V))$. We have (see [Ful97])

$$H^*(Gr(m, V)) = \bigoplus_{\lambda \text{ is a partition contained in the } m \times n \text{ rectangle}} \mathbb{Z}\sigma_\lambda. \quad (34)$$

Now, let V be a complex vector space V of dimension $2n$, endowed with a nondegenerate skew-symmetric bilinear form ω . A subspace W of V is isotropic if the form ω vanishes on it, i.e., $\omega(v, w) = 0$ for all $v, w \in W$. A maximal isotropic subspace of V is called **Lagrangian**. The set $LG(n, V)$ of Lagrangian subspaces in V is called the **Lagrangian Grassmannian**. Fix a complete isotropic flag of V

$$\mathcal{L} : 0 = V_0 \subset \cdots \subset V_i \subset \cdots \subset V_n \subset V, \quad (35)$$

where each V_i is a vector subspace of V , of dimension i and V_n is Lagrangian. To each strict partition $\lambda = (\lambda_1, \dots, \lambda_l)$ with $\lambda_i > 0$, contained in $(n, n-1, \dots, 1)$, we associate the Schubert variety

$$Y_\lambda(\mathcal{L}) = \{W \in LG(n, V) \mid \dim(W \cap V_{n+1-\lambda_i}) \geq i \ (1 \leq i \leq l)\}. \quad (36)$$

The Poincare dual class of $Y_\lambda(\mathcal{L})$ is denoted by θ_λ and called a Schubert class. Then θ_λ is an element of $H^{2|\lambda|}(LG(n, V))$. We have (see [Pra91])

$$H^*(LG(n, V)) = \bigoplus_{\lambda \text{ is a strict partition contained in } (n, n-1, \dots, 1)} \mathbb{Z}\theta_\lambda, \quad (37)$$

and

$$\theta_\lambda \theta_\mu = \sum_{\nu} 2^{l(\lambda)+l(\mu)-l(\nu)} f_{\lambda\mu}^\nu \theta_\nu. \quad (38)$$

There is a canonical embedding $\iota : LG(n, V) \rightarrow Gr(n, V)$. The map ι induces the ring homomorphism $\iota^* : H^*(Gr(n, V)) \rightarrow H^*(LG(n, V))$. For each partition μ contained in the $n \times n$ rectangle, we have (see [Pra00])

$$\iota^*(\sigma_\mu) = \sum_{\lambda \text{ is a strict partition contained in } (n, n-1, \dots, 1)} g_{\lambda\mu} \theta_\lambda. \quad (39)$$

3.5 Application to the identity $g_{\lambda\mu} = g_{\lambda\mu^t}$

Proposition 3.6. *Let λ be a strict partition and μ be a partition. Then $g_{\lambda\mu} = g_{\lambda\mu^t}$.*

Proof. Let V be a complex vector space of dimension $2n$, endowed with a nondegenerate skew-symmetric bilinear form ω . For each subspace W of V , set

$$W^{\perp\omega} = \{v' \in V \text{ such that } \omega(v', v) = 0 \text{ for all } v \in W\}, \quad (40)$$

$$W^\perp = \{f \in V^* \text{ such that } f(v) = 0 \text{ for all } v \in W\}. \quad (41)$$

Fix a complete isotropic flag of V

$$\mathcal{L} : 0 = V_0 \subset \cdots \subset V_i \subset \cdots \subset V_n \subset V. \quad (42)$$

Then we can extend \mathcal{L} to a complete flag \mathcal{F} of V as follow

$$\mathcal{F} : 0 = V_0 \subset \cdots \subset V_i \subset \cdots \subset V_{2n} = V, \quad (43)$$

where $V_{n+i} = (V_{n-i})^{\perp\omega}$ for each $i = 1, \dots, n$. Moreover, the flag \mathcal{F}^\perp defined below is a complete flag of V^*

$$\mathcal{F}^\perp : 0 = (V_{2n})^\perp \subset \cdots \subset (V_{2n-i})^\perp \subset \cdots \subset (V_0)^\perp = V^*. \quad (44)$$

Then flag \mathcal{L}^\perp defined below is a complete isotropic flag of V^*

$$\mathcal{L}^\perp : 0 = (V_{2n})^\perp \subset \cdots \subset (V_{2n-i})^\perp \subset \cdots \subset (V_n)^\perp \subset V^*. \quad (45)$$

We define an isomorphism $\eta : Gr(n, V) \rightarrow Gr(n, V^*)$ by $W \mapsto W^\perp$. By [GH78], we know that

$$\eta(X_\mu(\mathcal{F})) = X_{\mu^t}(\mathcal{F}^\perp). \quad (46)$$

Hence, the map η induces the ring isomorphism $\eta^* : H^*(Gr(n, V^*)) \rightarrow H^*(Gr(n, V))$ with

$$\eta^*(\sigma_\mu) = \sigma_{\mu^t}. \quad (47)$$

The restriction of η on $LG(n, V)$ is also an isomorphism and we still denote it by η . We have

$$\eta(Y_\lambda(\mathcal{L})) = Y_\lambda(\mathcal{L}^\perp). \quad (48)$$

Indeed,

1. For each $W \in Y_\lambda(\mathcal{L})$, we have $W^{\perp\omega} = W$, and

$$\dim(W \cap V_{n+1-\lambda_i})^{\perp\omega} = 2n - 1 + \lambda_i - \dim(W \cap V_{n-1+\lambda_i}). \quad (49)$$

Then we can rewrite (36) as

$$Y_\lambda(\mathcal{L}) = \{W \in LG(n, V) \mid \dim(W \cap V_{n-1+\lambda_i}) \geq i + \lambda_i - 1 \ (1 \leq i \leq l)\}. \quad (50)$$

2. Now, for any $W \in Y_\lambda(\mathcal{L})$ given by (50), we have

$$\dim(W \cap V_{n-1+\lambda_i})^\perp = 2n + 1 - \lambda_i - \dim(W^\perp \cap V_{n-1+\lambda_i}^\perp). \quad (51)$$

Then $W^\perp \in Y_\lambda(\mathcal{L}^\perp)$ given by (36).

Hence, the map η induces the ring isomorphism $\eta^* : H^*(LG(n, V^*)) \rightarrow H^*(LG(n, V))$ with

$$\eta^*(\theta_\lambda) = \theta_\lambda. \quad (52)$$

We have

$$\eta^* \iota^* = \iota^* \eta^*. \quad (53)$$

Apply η^* on both sides of the equality (39), with the help of (47), (52), (53), we get

$$\eta^*(\iota^*(\sigma_\mu)) = \sum_{\lambda \text{ is a strict partition contained in } (n, n-1, \dots, 1)} g_{\lambda\mu} \theta_\lambda \quad (54)$$

$$= \sum_{\lambda \text{ is a strict partition contained in } (n, n-1, \dots, 1)} g_{\lambda\mu^t} \theta_\lambda. \quad (55)$$

It implies $g_{\lambda\mu} = g_{\lambda\mu^t}$. □

4 A new combinatorial models for the coefficients $f_{\lambda\mu}^\nu$


Given a skew shifted shape ν/μ , we number the boxes from top to bottom and right to left in each row by $1, 2, \dots, |\nu/\mu|$, respectively. The result is called the **shifted reverse filling of the skew shifted shape** ν/μ . We denote it by $\tilde{T}_{\nu/\mu}$.

For each $k = 1, 2, \dots, |\nu/\mu|$, let k^* to be meant k or k' .

We now let $\tilde{\mathcal{O}}(\nu/\mu)$ be the set of all tableaux T of size $|\nu/\mu|$ of unshifted shape constructed from $\tilde{T}_{\nu/\mu}$, satisfying the following conditions:

(C1) If k and $k + 1$ appear in the same row of $\tilde{T}_{\nu/\mu}$, then $(k + 1)^*$ appears weakly above k or $(k + 1)^*$ appears strictly above k' in T .

(C2) If h appears in the box directly below k in $\tilde{T}_{\nu/\mu}$, then h^* appears weakly below k' or h^* appears strictly below k in T .



(C3) T is filled by the alphabet $\{1' < 1 < 2' < 2 < \dots < |\nu/\mu|' < |\nu/\mu|\}$ such that only one of k or k' appears in T for each $k = 1, 2, \dots, |\nu/\mu|$. The rightmost letter in each row of T is unmarked.

(C4) For each $j = 1, 2, \dots, n - 1$, let $T^{j\downarrow}$ be the result of T by removing the boxes with entries k' or $k > j$ if exists. Suppose that the shape of $T^{j\downarrow}$ is (τ_1, τ_2, \dots) . Then $\tau_1 \geq \tau_2 \geq \dots$ and if $\tau_{i-1} = \tau_i$ for some i , the entry $(j + 1)^*$ does not belong to the i^{th} row of T .

(C5) For each $j = n, n - 1, \dots, 2$, let $T^{j\uparrow}$ be the result of T by changing k' to k with $k \geq j$, removing the boxes with entries $k' < j$ if exists. Suppose that the shape of $T^{j\uparrow}$ is (τ_1, τ_2, \dots) . Then $\tau_1 \geq \tau_2 \geq \dots$ and if $\tau_{i-1} = \tau_i$ for some i , the entry $j - 1$ does not belong to the $(i - 1)^{\text{th}}$ row of T and the entry $(j - 1)'$ does not belong to the i^{th} row of T .

Example 4.1. We illustrate how the tableaux $T^{j\downarrow}$ and $T^{j\uparrow}$ come from a tableau T and the conditions (C4) and (C5) work. Let T be the following tableau

$$T = \begin{array}{|c|c|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 5' & 8' & 10 \\ \hline 6 & 7' & 9 & & & & \\ \hline 11 & 12 & & & & & \\ \hline \end{array}$$

We can easily compute the tableaux $T^{j\downarrow}$ for $j = 1, 2, \dots, 11$ and check that all of them satisfy the condition (C4). For example, here is the tableau $T^{10\downarrow}$ with shape $\tau = (5, 2)$ (the boxes in pink mean to be removed)

$$\begin{array}{|c|c|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 5' & 8' & 10 \\ \hline 6 & 7' & 9 & & & & \\ \hline 11 & 12 & & & & & \\ \hline \end{array} = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 10 \\ \hline 6 & 9 & & & \\ \hline & & & & \\ \hline \end{array}$$

For $j = 12, 11, \dots, 2$, we can also compute $T^{j\uparrow}$ but there is some tableaux $T^{j\uparrow}$ that does not satisfy the condition (C5), one of which is the tableau $T^{10\uparrow}$ given below (the boxes in pink mean to be removed and there is no letter k' such that $k \geq 10$)

$$\begin{array}{|c|c|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 5' & 8' & 10 \\ \hline 6 & 7' & 9 & & & & \\ \hline 11 & 12 & & & & & \\ \hline \end{array} = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 10 \\ \hline 6 & 9 & & & \\ \hline 11 & 12 & & & \\ \hline \end{array}$$

The shape of $T^{10\uparrow}$ is $\tau = (5, 2, 2)$. We see that $\tau_2 = \tau_3 = 2$ and the letter 9 belongs to the second row of T .

Theorem 4.2. Let λ, μ, ν be strict partitions. Then the coefficient $f_{\lambda\mu}^\nu$ is the number of the tableaux T in $\tilde{\mathcal{O}}(\nu/\mu)$ of shape λ .

Proof. Let $\tilde{\mathcal{S}}_\lambda(\nu/\mu)$ be the set of tableaux in Theorem 3.3. Let $\tilde{\mathcal{O}}_\lambda(\nu/\mu)$ be the set of tableaux in the set $\tilde{\mathcal{O}}(\nu/\mu)$ of shape λ .

Let $T \in \tilde{\mathcal{S}}_\lambda(\nu/\mu)$ with $w(T) = w_1 w_2 \dots w_{|\nu/\mu|}$. We associate T with an unique tableau T' of unshifted shape by the rules: For each $i = |\nu/\mu|, \dots, 2, 1$, we have

- If $w_i = k$, then $|\nu/\mu| + 1 - i$ appears in the k^{th} row of T' .
- If $w_i = k'$, then $(|\nu/\mu| + 1 - i)'$ appears in the k^{th} row of T' .

We can easily check that $T' \in \tilde{\mathcal{O}}_\lambda(\nu/\mu)$. Indeed,

- T has content λ if and only if T' has shape λ .
- The conditions (T1), (T2) and (T3) of T are equivalent to the conditions (C1) and (C2) of T' .
- The condition (F1) of T is equivalent to the condition (C3) of T' .
- The condition (F2) of T is equivalent to the conditions (C4) and (C5) of T' .

Hence, we can define an injection $\phi : \tilde{\mathcal{S}}_\lambda(\nu/\mu) \rightarrow \tilde{\mathcal{O}}_\lambda(\nu/\mu)$, $T \mapsto T'$.

Moreover, for each $T' \in \tilde{\mathcal{O}}_\lambda(\nu/\mu)$, we associate T' with an unique tableau T of skew shifted shape ν/μ and word $w(T) = w_1 w_2 \dots w_{|\nu/\mu|}$ by the rule: for each $j = |\nu/\mu|, \dots, 2, 1$, we have

- If j appears in the k^{th} row of T' , then $w_{|\nu/\mu|+1-j} = k$.
- If j' appears in the k^{th} row of T' , then $w_{|\nu/\mu|+1-j} = k'$.

The equivalence of the conditions we have already shown implies that $T \in \tilde{\mathcal{S}}_\lambda(\nu/\mu)$. So we can define an injection $\psi : \tilde{\mathcal{O}}_\lambda(\nu/\mu) \rightarrow \tilde{\mathcal{S}}_\lambda(\nu/\mu)$, $T' \mapsto T$. Moreover, $\phi\psi = Id$. Hence, ϕ is a bijection between two sets and $f_{\lambda\mu}^\nu = \#\tilde{\mathcal{S}}_\lambda(\nu/\mu) = \#\tilde{\mathcal{O}}_\lambda(\nu/\mu)$. \square

Theorem 4.3. Let λ be a strict partition and μ be a partition. Then the coefficient $g_{\lambda\mu}$ is the number of the tableaux T in $\tilde{\mathcal{O}}(\mu + \delta/\delta)$ of shape λ .

Proof. This follows from Theorem 4.2 and identity (28). \square

We illustrate the method to compute the coefficients $f_{\lambda\mu}^\nu$ through an example.

Example 4.4. Set $\lambda = (3, 2), \mu = (3, 2), \nu = (5, 3, 2)$.

(1) The shifted reverse filling of the skew shifted shape ν/μ is

$$\tilde{T}_{\nu/\mu} = \begin{array}{|c|c|c|c|c|} \hline & & & 2 & 1 \\ \hline & & & 3 & \\ \hline & & 5 & 4 & \\ \hline \end{array}$$

(2) To construct the tableaux T' in $\tilde{\mathcal{O}}_\lambda(\nu/\mu)$, we first use three conditions (C1), (C2) and (C3). Then check the results if they satisfy the conditions (C4) and (C5) or not.

1. We start with 1^* , there are two possibilities, they are $1'$ and 1 . But if $1'$ appears in the tableau T' then the next position of 2^* will be in the row above the first row by the condition (C1). It is impossible. Hence, just only one case that 1 appears in T' . Then the next two possibilities by the condition (C1) are

$$\boxed{1 \ 2} \quad \boxed{1 \ 2'}$$

2. For the second case, by the condition (C2), there are four possibilities below

$$\begin{array}{|c|c|} \hline 1 & 2' \\ \hline 3' & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2' \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2' & 3 \\ \hline & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2' & 3' \\ \hline & & \\ \hline \end{array}$$

- The last one cannot happen since the tableau T' has shape $\lambda = (3, 2)$. Then we consider $3'$ as the rightmost letter in the first row of T' and it should be 3 to satisfy the condition (C3).
- The second one also cannot happen because the next position of 4^* will be in the row below the row of 3 by the condition (C2). It cannot produce a tableau of shape $\lambda = (3, 2)$ later.
- For the third one, the next position of 4^* is based on the condition (C2). To produce the shape $\lambda = (3, 2)$ later, it will be as follows:

$$\begin{array}{|c|c|c|} \hline 1 & 2' & 3 \\ \hline 4 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2' & 3 \\ \hline 4' & & \\ \hline \end{array}$$

- For the first one, the next position of 4^* is based on the conditions (C2) and (C3). To produce the shape $\lambda = (3, 2)$ later, will be as follows:

$$\begin{array}{|c|c|} \hline 1 & 2' \\ \hline 3' & 4 \\ \hline \end{array}$$


Continue until the end on the remaining cases by similar arguments, we finally can find the tableaux of shape $\lambda = (3, 2)$ satisfying all conditions (C1), (C2) and (C3) as follows:

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3' & 4 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2' & 5 \\ \hline 3' & 4 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2' & 3 \\ \hline 4 & 5 & \\ \hline \end{array}$$


We can check that only first two tableaux above satisfy the conditions (C4) and (C5). Hence,

$$f_{\lambda\mu}^\nu = 2.$$


We automatically find out the set of skew shifted tableaux described in Theorem 3.3 by using the bijection we mentioned in the proof of Theorem 4.2. Here they are



Remark 4.5. The model of Remmel and Whitney for Littlewood-Richardson coefficients [RW84], i.e., the set $\mathcal{O}_\mu(\nu/\lambda)$ in Theorem 2.8, has another interpretation by White [Whi81]. Namely, each tableau in the set $\mathcal{O}_\mu(\nu/\lambda)$ can be considered as the recording tableau of the word rewritten in inverse order of a tableau in the set $\mathcal{S}(\nu/\lambda, \mathcal{U}_\mu)$. Our model, i.e., the set $\tilde{\mathcal{O}}_\lambda(\nu/\mu)$, is analogous to Remmel and Whitney's model but for shifted Littlewood-Richardson coefficients. In [Shi99], M. Shimozono gave an analogous model to the White's model in [Whi81] but for shifted Littlewood-Richardson coefficients. Our model and Shimozono's model are totally different. For example, with $\lambda = (3, 2)$, $\mu = (3, 2)$ and $\nu = (5, 3, 2)$, our model $\tilde{\mathcal{O}}_\lambda(\nu/\mu)$ consists of the elements below



However, Shimozono's model consists of the elements below



In conclusion, the Shimozono's model is not the same as our model.


5 On the coefficients $g_{\lambda\mu}$

In this section, we present our second result. Namely, we present a new interpretation of the coefficients $g_{\lambda\mu}$ as a subset of a set that counts Littlewood-Richardson coefficients. As corollaries, we can compute the coefficients $g_{\lambda\mu}$ by models for Littlewood-Richardson coefficients. We will prove and conjecture inequalities between the coefficients and also state some conjectures that explain the hidden structure behind them.

5.1 A new interpretation of the coefficients $g_{\lambda\mu}$


For any (skew) shifted tableau T without entries k' , let $s(T)$ be the new (skew) tableau which is defined as follow:

1. Create an image of T by the symmetry through its main diagonal.
2. Combine the image we have created with T by gluing them along the main diagonal as the image below.



Let ν/μ be the skew shifted shape of T , then we denote the shape of $s(T)$ by $\widetilde{\nu}/\mu$.

Example 5.1.



The following result is a restatement of Proposition 5.4 in the article [Hai89] of M. D. Haiman.

Proposition 5.2. *Let T be a skew shifted tableau without entries k' . Then we have $s(sRect(T)) = Rect(s(T))$.*


For any strict partition λ of n , let \mathcal{T}_λ be the shifted tableau of shifted shape λ , obtained by putting numbers $1, 2, \dots, n$ in the boxes of shifted diagram $sY(\lambda)$ from left to right, starting from top to bottom. Let $\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$ be the subset of $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$ of all pairs (T, U) such that $T = U^t$.

Theorem 5.3. *Let λ be a strict partition and μ be a partition. Then $g_{\lambda\mu} = \#\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$.*

Proof. We have $g_{\lambda\mu} = f_{\lambda\delta}^{\mu+\delta}$. By Theorem 3.5, it is the number of standard skew shifted tableaux S of skew shifted shape $(\mu + \delta)/\delta$ such that $sRect(S) = \mathcal{T}_\lambda$. The condition $sRect(S) = \mathcal{T}_\lambda$, by Proposition 5.2 is equivalent to the condition

$$s(\mathcal{T}_\lambda) = Rect(s(S)). \quad (56)$$

The tableau $s(S)$ has form



where U is a standard Young tableau of shape μ . Since $s(S)$ and $U^t * U$ have the same word, then by Lemma 2.4 and Lemma 2.6, we have


$$s(\mathcal{T}_\lambda) = Rect(U^t * U) = U^t \cdot U. \quad (57)$$

It is clear that U is uniquely determined by S . Hence, $g_{\lambda\mu}$ is the number of pairs (T, U) in the set $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$ such that $T = U^t$. \square

Theorem 5.3 gives a way to compute the coefficients $g_{\lambda\mu}$.

Example 5.4. *Let $\lambda = (5, 2)$ and $\mu = (4, 2, 1)$. Since $\mu^t = (3, 2, 1, 1)$, we can re-use the computation in Example 2.9. The elements in the set $\mathcal{S}(\tilde{\lambda}/\mu^t, \mathcal{U}_\mu)$ with the corresponding*

elements in the set $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$ are (the elements in the subsets $\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$ are marked by coloring all boxes in green).



Hence, $g_{\lambda\mu} = 2$.

5.2 Inequality $g_{\lambda\mu} \leq c_{\mu\mu^t}^{\tilde{\lambda}}$

In this subsection, we prove a conjecture of N. Ressayre [Res19]: $g_{\lambda\mu} \leq c_{\mu\mu^t}^{\tilde{\lambda}}$. The conjecture was based on the facts below

$$g_{\lambda\mu} = 1 \text{ implies } c_{\mu\mu^t}^{\tilde{\lambda}} = 1, \quad (58)$$

$$g_{\lambda\mu} \neq 0 \text{ implies } c_{\mu\mu^t}^{\tilde{\lambda}} \neq 0. \quad (59)$$

The conclusions (58), (59) are versions of Theorem 1.4 and Proposition 1.6 in the article [BKR12] for the case $G = Sp(2n, \mathbb{C})$. Indeed, with the notations in the article [BKR12], by [Res12], in the case $G = Sp(2n, \mathbb{C})$, G/P is the Lagrangian Grassmannian $LG(n, \mathbb{C}^{2n})$, the

corresponding Levi group is $GL(n)$, and W^P is parametrized by strict partitions. Suppose that $w \in W^P$ corresponds to strict partition λ , then χ_w corresponds to the partition $\tilde{\lambda}$. Let λ^\vee be the strict partition corresponding to the completion of $sY(\lambda)$ in $sY((n, n-1, \dots, 1))$. Let $\lambda_1, \lambda_2, \lambda_3$ be the strict partitions corresponding to $w_1, w_2, w_3 \in W^P$, respectively in Theorem 1.4 in [BKR12]. Since the structure constants for the singular cohomology and the deformed cohomology \odot_0 in this case are the same, the theorem says that for any positive integer k , we have

$$f_{\lambda_1 \lambda_2}^{\lambda_3^\vee} = 1 \text{ implies } c_{\widetilde{k\lambda_1 k\lambda_2}}^{\widetilde{k\lambda_3^\vee}} = 1. \quad (60)$$

In particular, when $\lambda_1 = \lambda$, $\lambda_2 = \delta$, $\lambda_3^\vee = \mu + \delta$, the left-hand side of (60) becomes

$$1 = f_{\lambda \delta}^{\mu + \delta} = g_{\lambda \mu}.$$

With $k = 1$, the right-hand side of (60) becomes

$$\begin{aligned} 1 &= c_{\widetilde{\lambda \delta}}^{\widetilde{\mu + \delta}} = \#\mathcal{S}(\widetilde{\mu + \delta / \delta}, s(\mathcal{T}_\lambda)) \\ &= \#\mathcal{S}(\mu^t * \mu, s(\mathcal{T}_\lambda)) \\ &= \#\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda)) \\ &= c_{\mu^t \mu}^{\tilde{\lambda}}. \end{aligned}$$

Hence, we get the conclusion (58). Similarly, the conclusion (59) follows Proposition 1.6 in [BKR12].

Theorem 5.5. *Let λ be a strict partition and μ be a partition. Then $g_{\lambda \mu} \leq c_{\mu^t \mu}^{\tilde{\lambda}}$.*

Proof. By Theorem 5.3 and Theorem 2.8, we have

$$g_{\lambda \mu} = \overline{\#\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))} \leq \#\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda)) = c_{\mu^t \mu}^{\tilde{\lambda}}. \quad (61)$$

□

5.3 Inequality $g_{\lambda \mu}^2 \leq c_{\mu^t \mu}^{\tilde{\lambda}}$

In this subsection, we propose a stronger conjectural inequality than Theorem 5.5. We provide some examples to support this conjecture. Indeed, we formulate a conjecture on combinatorial models whose validity implies the first conjecture.

Conjecture 5.6. *Let λ be a strict partition and μ be a partition. Then $g_{\lambda \mu}^2 \leq c_{\mu^t \mu}^{\tilde{\lambda}}$.*

To compute the decomposition of P_λ into Schur functions s_μ by computer, we use the code below. For example, with $\lambda = (4, 2)$, we have

$$P_{(4,2)} = s_{(2,2,1,1)} + s_{(2,2,2)} + s_{(3,1,1)} + 2s_{(3,2,1)} + s_{(3,3)} + s_{(4,1,1)} + s_{(4,2)}. \quad (62)$$

The code in SageMath (online version: <https://cocalc.com/>) is

```
Sym = SymmetricFunctions(FractionField(QQ['t']))
SP = Sym.hall_littlewood(t=-1).P();
s = Sym.schur();
s(SP([4,2]))
```

The result appears in computer is

$$s[2, 2, 1, 1] + s[2, 2, 2] + s[3, 1, 1, 1] + 2*s[3, 2, 1] + s[3, 3] + s[4, 1, 1] + s[4, 2].$$

To compute the Littlewood-Richardson coefficients by computer, we use the code below. For example, for $\lambda = (4, 2)$, $\mu = \mu^t = (3, 2, 1)$, we have

$$c_{\mu^t \mu}^{\tilde{\lambda}} = 4. \tag{63}$$

The code is

```
import sage.libs.lrcalc.lrcalc as lrcalc
A = [5,4,2,1]
B = [3,2,1]
C = [3,2,1]
lrcalc.lrcoef(A,B,C)
```

The result appears in computer is

4.

We check the conjecture for all strict partitions λ such that $|\lambda| \leq 11$. By Theorem 5.5, we just need to check the cases $g_{\lambda\mu} > 1$. Here is the data of computations on computer.

$ \lambda $	strict partitions λ such that there exists $g_{\lambda\mu} > 1$	μ such that $g_{\lambda\mu} > 1$	$g_{\lambda\mu}$	$c_{\mu^t \mu}^{\tilde{\lambda}}$
11	(9,2)	(3, 2, 1, 1, 1, 1, 1)	2	4
11	(9,2)	(4, 2, 1, 1, 1, 1)	2	4
11	(9,2)	(5, 2, 1, 1, 1, 1)	2	4
11	(9,2)	(6, 2, 1, 1, 1)	2	4
11	(9,2)	(7, 2, 1, 1)	2	4
11	(9,2)	(8, 2, 1)	2	4
11	(8,3)	(3, 2, 2, 1, 1, 1, 1)	2	4
11	(8,3)	(4, 2, 1, 1, 1, 1, 1)	2	4
11	(8,3)	(4, 2, 2, 1, 1, 1)	2	4
11	(8,3)	(4, 3, 1, 1, 1, 1)	2	4
11	(8,3)	(5, 2, 1, 1, 1, 1)	2	4
11	(8,3)	(5, 2, 2, 1, 1)	2	4
11	(8,3)	(5, 3, 1, 1, 1)	2	4
11	(8,3)	(6, 2, 1, 1, 1)	2	4
11	(8,3)	(6, 2, 2, 1)	2	4
11	(8,3)	(6, 3, 1, 1)	2	4
11	(8,3)	(7, 2, 1, 1)	2	4
11	(8,3)	(7, 3, 1)	2	4
11	(7,4)	(3, 2, 2, 2, 1, 1)	2	4
11	(7,4)	(4, 2, 2, 1, 1, 1)	2	4
11	(7,4)	(4, 2, 2, 2, 1)	2	4
11	(7,4)	(4, 3, 2, 1, 1)	2	4
11	(7,4)	(5, 2, 1, 1, 1, 1)	2	4
11	(7,4)	(5, 2, 2, 1, 1)	2	4
11	(7,4)	(5, 3, 1, 1, 1)	2	4
11	(7,4)	(5, 3, 2, 1)	2	4
11	(7,4)	(5, 4, 1, 1)	2	4
11	(7,4)	(6, 2, 1, 1, 1)	2	4
11	(7,4)	(6, 3, 1, 1)	2	4
11	(7,4)	(6, 4, 1)	2	4

11	(7,3,1)	(3, 3, 2, 1, 1, 1)	2	6
11	(7,3,1)	(4, 2, 2, 1, 1, 1)	2	5
11	(7,3,1)	(4, 3, 1, 1, 1, 1)	2	5
11	(7,3,1)	(4, 3, 2, 1, 1)	3	13
11	(7,3,1)	(5, 2, 2, 1, 1)	2	5
11	(7,3,1)	(5, 3, 1, 1, 1)	2	5
11	(7,3,1)	(5, 3, 2, 1)	3	13
11	(7,3,1)	(6, 2, 2, 1)	2	5
11	(7,3,1)	(6, 3, 1, 1)	2	5
11	(7,3,1)	(6, 3, 2)	2	6
11	(6,4,1)	(3, 3, 2, 2, 1)	2	6
11	(6,4,1)	(4, 2, 2, 2, 1)	2	5
11	(6,4,1)	(4, 3, 2, 1, 1)	3	14
11	(6,4,1)	(4, 3, 2, 2)	2	7
11	(6,4,1)	(4, 3, 3, 1)	2	4
11	(6,4,1)	(4, 4, 2, 1)	2	7
11	(6,4,1)	(5, 2, 2, 1, 1)	2	5
11	(6,4,1)	(5, 3, 1, 1, 1)	2	5
11	(6,4,1)	(5, 3, 2, 1)	3	14
11	(6,4,1)	(5, 4, 1, 1)	2	5
11	(6,4,1)	(5, 4, 2)	2	6
11	(6,3,2)	(4, 3, 2, 1, 1)	3	10
11	(6,3,2)	(4, 3, 2, 2)	2	4
11	(6,3,2)	(4, 3, 3, 1)	2	5
11	(6,3,2)	(4, 4, 2, 1)	2	4
11	(6,3,2)	(5, 3, 2, 1)	3	10
11	(5,4,2)	(4, 3, 2, 1, 1)	2	4
11	(5,4,2)	(4, 3, 2, 2)	2	5
11	(5,4,2)	(4, 3, 3, 1)	2	5
11	(5,4,2)	(4, 4, 2, 1)	2	5
11	(5,4,2)	(5, 3, 2, 1)	2	4
10	(8,2)	(3, 2, 1, 1, 1, 1, 1)	2	4
10	(8,2)	(4, 2, 1, 1, 1, 1)	2	4
10	(8,2)	(5, 2, 1, 1, 1)	2	4
10	(8,2)	(6, 2, 1, 1)	2	4
10	(8,2)	(7, 2, 1)	2	4
10	(7,3)	(3, 2, 2, 1, 1, 1)	2	4
10	(7,3)	(4, 2, 1, 1, 1, 1)	2	4
10	(7,3)	(4, 2, 2, 1, 1)	2	4
10	(7,3)	(4, 3, 1, 1, 1)	2	4
10	(7,3)	(5, 2, 1, 1, 1)	2	4
10	(7,3)	(5, 2, 2, 1)	2	4
10	(7,3)	(5, 3, 1, 1)	2	4
10	(7,3)	(6, 2, 1, 1)	2	4
10	(7,3)	(6, 3, 1)	2	4
10	(6,4)	(3, 2, 2, 2, 1)	2	4
10	(6,4)	(4, 2, 2, 1, 1)	2	4
10	(6,4)	(4, 3, 2, 1)	2	4
10	(6,4)	(5, 2, 1, 1, 1)	2	4

10	(6,4)	(5, 3, 1, 1)	2	4
10	(6,4)	(5, 4, 1)	2	4
10	(6,3,1)	(3, 3, 2, 1, 1)	2	6
10	(6,3,1)	(4, 2, 2, 1, 1)	2	5
10	(6,3,1)	(4, 3, 1, 1, 1)	2	5
10	(6,3,1)	(4, 3, 2, 1)	3	13
10	(6,3,1)	(5, 2, 2, 1)	2	5
10	(6,3,1)	(5, 3, 1, 1)	2	5
10	(6,3,1)	(5, 3, 2)	2	6
10	(5,4,1)	(4, 3, 2, 1)	2	7
10	(5,3,2)	(4, 3, 2, 1)	3	9
9	(7,2)	(3, 2, 1, 1, 1, 1)	2	4
9	(7,2)	(4, 2, 1, 1, 1)	2	4
9	(7,2)	(5, 2, 1, 1)	2	4
9	(7,2)	(6, 2, 1)	2	4
9	(6,3)	(3, 2, 2, 1, 1)	2	4
9	(6,3)	(4, 2, 1, 1, 1)	2	4
9	(6,3)	(4, 2, 2, 1)	2	4
9	(6,3)	(4, 3, 1, 1)	2	4
9	(6,3)	(5, 2, 1, 1)	2	4
9	(6,3)	(5, 3, 1)	2	4
9	(5,3,1)	(3, 3, 2, 1)	2	6
9	(5,3,1)	(4, 2, 2, 1)	2	5
9	(5,3,1)	(4, 3, 1, 1)	2	5
9	(5,3,1)	(4, 3, 2)	2	6
8	(6,2)	(3, 2, 1, 1, 1)	2	4
8	(6,2)	(4, 2, 1, 1)	2	4
8	(6,2)	(5, 2, 1)	2	4
8	(5,3)	(3, 2, 2, 1)	2	4
8	(5,3)	(4, 2, 1, 1)	2	4
8	(5,3)	(4, 3, 1)	2	4
7	(5,2)	(3, 2, 1, 1)	2	4
7	(5,2)	(4, 2, 1)	2	4
6	(4,2)	(3,2,1)	2	4
< 6	\emptyset	\emptyset		

Conjecture 5.7. *We have*

1. *The restriction of the map $\mathcal{S}_{s(\mathcal{T}_\lambda), \mathcal{U}_\mu, \mathcal{U}_{\mu^t}, s(\mathcal{T}_\lambda)}^{\mu^t, \mu, \bar{\lambda}}$ on the set $\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$ is a bijection onto the set $\overline{\mathcal{T}(\mu, \mu^t, s(\mathcal{T}_\lambda))}$.*
2. *The elements of the set $\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$ have the form (U_α^t, U_α) , with index α . Let (V_α, V_α^t) be the image of (U_α^t, U_α) through the bijection $\mathcal{S}_{s(\mathcal{T}_\lambda), \mathcal{U}_\mu, \mathcal{U}_{\mu^t}, s(\mathcal{T}_\lambda)}^{\mu^t, \mu, \bar{\lambda}}$. Let (U_α^t, U_α) and (U_β^t, U_β) be elements of the set $\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$. If (U_α^t, U_β) is not in the set $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$, then (V_α, V_β^t) is in the set $\mathcal{T}(\mu, \mu^t, s(\mathcal{T}_\lambda))$.*

Remark 5.8. *Thanks to Theorem 5.3, the validity of Conjecture 5.7 1. implies the equality $g_{\lambda\mu} = g_{\lambda\mu^t}$, which was proved in Proposition 3.6.*

Proposition 5.9. *Suppose that Conjecture 5.7 holds. Then we have $g_{\lambda\mu}^2 \leq c_{\mu^t\mu}^{\bar{\lambda}}$.*


Proof. We suppose that (U_α^t, U_α) and (U_β^t, U_β) are elements in $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$. We construct an element $(\widetilde{U_\alpha^t}, \widetilde{U_\beta})$ is still in the set as follows:

1. If (U_α^t, U_β) belongs to $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$, then we set $(\widetilde{U_\alpha^t}, \widetilde{U_\beta}) = (U_\alpha^t, U_\beta)$.
2. If (U_α^t, U_β) does not belong to $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$, then by Conjecture 5.7, (V_α, V_β^t) belongs to $\mathcal{T}(\mu, \mu^t, s(\mathcal{T}_\lambda))$. Set $(\widetilde{U_\alpha^t}, \widetilde{U_\beta})$ is the image of (V_α, V_β^t) through the bijection $\left(\mathcal{S}_{s(\mathcal{T}_\lambda), \mathcal{U}_\mu, \mathcal{U}_{\mu^t}, s(\mathcal{T}_\lambda)}^{\mu^t, \mu, \bar{\lambda}}\right)^{-1}$.

The set of all pairs $(\widetilde{U_\alpha^t}, \widetilde{U_\beta})$ we have constructed is a subset of $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$. Since its cardinal is $g_{\lambda\mu}^2$, we have $g_{\lambda\mu}^2 \leq c_{\mu^t\mu}^{\bar{\lambda}}$. \square

We can see the conjecture through following example.

Example 5.10. Let $\lambda = (5, 2)$ and $\mu = (4, 2, 1)$. The correspondence between elements in $\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))$ and elements in $\mathcal{T}(\mu, \mu^t, s(\mathcal{T}_\lambda))$ is showed below (the elements in the subsets $\overline{\mathcal{T}(\mu^t, \mu, s(\mathcal{T}_\lambda))}$ and $\overline{\mathcal{T}(\mu, \mu^t, s(\mathcal{T}_\lambda))}$ are marked by coloring all boxes in green).



Acknowledgments

The author would like to express his sincere gratitude to his supervisors Prof. Nicolas Ressayre and Prof. Kenji Iohara for suggesting the subject and for many useful discussion, inspiring ideas during the work. He also grateful to their correction and their teaching how to write better, understandable and clearly.

References

- [BKR12] Prakash Belkale, Shrawan Kumar, and Nicolas Ressayre. A generalization of Fulton’s conjecture for arbitrary groups. *Math. Ann.*, 354(2):401–425, 2012.
- [BSS96] Georgia Benkart, Frank Sottile, and Jeffrey Stroomer. Tableau switching: algorithms and applications. *J. Combin. Theory Ser. A*, 76(1):11–43, 1996.
- [CGR84] Y. M. Chen, A. M. Garsia, and J. Remmel. Algorithms for plethysm. In *Combinatorics and algebra (Boulder, Colo., 1983)*, volume 34 of *Contemp. Math.*, pages 109–153. Amer. Math. Soc., Providence, RI, 1984.
- [CK18] Seung-II Choi and Jae-Hoon Kwon. Crystals and Schur P -positive expansions. *Electron. J. Combin.*, 25(3):Paper No. 3.7, 27, 2018.
- [CNO14] Seung-II Choi, Sun-Young Nam, and Young-Tak Oh. Bijections among combinatorial models for shifted Littlewood-Richardson coefficients. *J. Combin. Theory Ser. A*, 128:56–83, 2014.
- [Ful97] William Fulton. *Young tableaux*, volume 35 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
- [GH78] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics.
- [GJK⁺14] Dimitar Grantcharov, Ji Hye Jung, Seok-Jin Kang, Masaki Kashiwara, and Myungho Kim. Crystal bases for the quantum queer superalgebra and semi-standard decomposition tableaux. *Trans. Amer. Math. Soc.*, 366(1):457–489, 2014.
- [Hai89] Mark D. Haiman. On mixed insertion, symmetry, and shifted Young tableaux. *J. Combin. Theory Ser. A*, 50(2):196–225, 1989.
- [Liu10] Ricky Ini Liu. An algorithmic Littlewood-Richardson rule. *J. Algebraic Combin.*, 31(2):253–266, 2010.
- [Pra91] Piotr Pragacz. Algebro-geometric applications of Schur S - and Q -polynomials. In *Topics in invariant theory (Paris, 1989/1990)*, volume 1478 of *Lecture Notes in Math.*, pages 130–191. Springer, Berlin, 1991.
- [Pra00] Piotr Pragacz. Addendum: “A generalization of the Macdonald-You formula” [J. Algebra 204 (1998), no. 2, 573–587; MR1624487 (99g:05181)]. *J. Algebra*, 226(1):639–648, 2000.
- [Res12] Nicolas Ressayre. A cohomology-free description of eigencones in types A, B, and C. *Int. Math. Res. Not. IMRN*, (21):4966–5005, 2012.

- [Res19] N. Ressayre. Private communication, 2019.
- [RW84] J. B. Remmel and R. Whitney. Multiplying Schur functions. *J. Algorithms*, 5(4):471–487, 1984.
- [Sag87] Bruce E. Sagan. Shifted tableaux, Schur Q -functions, and a conjecture of R. Stanley. *J. Combin. Theory Ser. A*, 45(1):62–103, 1987.
- [Ser10] Luis Serrano. The shifted plactic monoid. *Math. Z.*, 266(2):363–392, 2010.
- [Shi99] Mark Shimozono. Multiplying Schur Q -functions. *J. Combin. Theory Ser. A*, 87(1):198–232, 1999.
- [Ste89] John R. Stembridge. Shifted tableaux and the projective representations of symmetric groups. *Adv. Math.*, 74(1):87–134, 1989.
- [Vak06] Ravi Vakil. A geometric Littlewood-Richardson rule. *Ann. of Math. (2)*, 164(2):371–421, 2006. Appendix A written with A. Knutson.
- [Whi81] Dennis E. White. Some connections between the Littlewood-Richardson rule and the construction of Schensted. *J. Combin. Theory Ser. A*, 30(3):237–247, 1981.
- [Wor84] Dale Raymond Worley. *A theory of shifted Young tableaux*. ProQuest LLC, Ann Arbor, MI, 1984. Thesis (Ph.D.)—Massachusetts Institute of Technology.
- [Zel81] A. V. Zelevinsky. A generalization of the Littlewood-Richardson rule and the Robinson-Schensted-Knuth correspondence. *J. Algebra*, 69(1):82–94, 1981.

Universite Lyon, University Claude Bernard Lyon 1, CNRS UMR 5208, Institut Camille Jordan, 43 Boulevard du 11 Novembre 1918, F-69622 Villeurbanne cedex, France
 E-mail: khanh.mathematic@gmail.com