

# SYMPLECTIC 4-MANIFOLDS ADMIT WEINSTEIN TRISECTIONS

PETER LAMBERT-COLE, JEFFREY MEIER, AND LAURA STARKSTON

ABSTRACT. We prove that every symplectic 4-manifold admits a trisection that is compatible with the symplectic structure in the sense that the symplectic form induces a Weinstein structure on each of the three sectors of the trisection. Along the way, we show that a (potentially singular) symplectic braided surface in  $\mathbb{C}\mathbb{P}^2$  can be symplectically isotoped into bridge position.

## 1. INTRODUCTION

A trisection of a 4-manifold is a decomposition into three basic pieces with nice overlaps, which allows one to study the smooth 4-manifold using curves on a surface up to isotopy and certain combinatorial equivalence moves. A trisection can be viewed as an alternative to a handle decomposition, the advantage being that curves on a surface are simpler than the framed links which serve as attaching spheres of handles. Although trisections are designed to study smooth topology, we can ask in which ways a trisection can keep track of additional geometric structures on a 4-manifold. In this article, the geometric structure of interest is a symplectic structure.

In the case that  $(X, \omega)$  is a symplectic 4-manifold, it is natural to look for trisections on  $X$  that are compatible in some way with  $\omega$ . The standard trisection of  $\mathbb{C}\mathbb{P}^2$  (see Example 2.3) can be constructed via toric geometry and therefore interacts nicely with the symplectic geometry. The contact type property of the boundaries of the three sectors in  $\mathbb{C}\mathbb{P}^2$  played an important role in the trisection-theoretic proof of the Thom conjecture [Lam18]. Further progress relating trisections and symplectic structures was made by Gay: Every closed symplectic manifold admits a Lefschetz pencil, and Gay described how to construct a trisection from a pencil [Gay16]; see also related work on Lefschetz fibrations by Castro-Ozbagci [CO19].

Weinstein provided a method of building symplectic manifolds through handle attachments [Wei91], but the construction has a significant limitation. The handles for a symplectic manifold of dimension  $2n$  can have index at most  $n$ , and for symplectic 4-manifolds, even the 2-handles can only be attached with certain framings (less than the maximal Thurston-Bennequin framing of a Legendrian realization of the attaching circle; see [GS99, Chapter 11] and [Gom98] for details). There is no general method to extend a symplectic structure over a 4-dimensional 3-handle or 4-handle. In particular, any closed symplectic 4-manifold cannot be built solely from Weinstein's handle construction.

The main result of this article demonstrates a new advantage that trisections enjoy over handle decompositions, namely compatibility with a symplectic structure on a closed manifold. In [LM18], the first two authors proposed the notion of a *Weinstein trisection* of  $(X, \omega)$  as the appropriate generalization of the standard trisection of  $\mathbb{C}\mathbb{P}^2$ , and conjectured that every symplectic 4-manifold admits a Weinstein trisection. In this paper, we prove that conjecture.

---

2010 *Mathematics Subject Classification.* 53D05; 57M12; 57R15.

*Key words and phrases.* 4-manifold, trisection, bridge trisection, branched cover, symplectic structure, Weinstein structure, quasiholomorphic curves.

**Theorem 1.1.** *Every closed, symplectic 4–manifold  $(X, \omega)$  admits a Weinstein trisection.*

The proof of Theorem 1.1 relies on important results of Auroux [Aur00] and Auroux-Katzarkov [AK00] that every closed, symplectic 4–manifold admits a quasiholomorphic branched covering map  $f: X \rightarrow \mathbb{C}\mathbb{P}^2$ . Trisections are naturally suited for studying branched coverings of 4–manifolds (e.g. [BCKM19, RT16, LM18, CK17]), and the basic strategy is to pull back the Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$  via the singular branched covering map  $f$ .

To carry this strategy out, we must first symplectically isotope the branch locus, which is possibly singular, into *bridge position* with respect to a genus one Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$ . Auroux and Katzarkov proved that the branch locus can be assumed to be braided with respect to the standard linear pencil [AK00]. We show in Section 5 that all such curves can be symplectically isotoped into bridge position.

Putting this result (Theorem 5.9) together with the fact that any smooth symplectic submanifold (or with singularities modeled on a complex algebraic curve) in  $\mathbb{C}\mathbb{P}^2$  can be assumed to be braided (see Remark 5.4), we obtain the following theorem, which may shed some light on the study of the symplectic isotopy problem using bridge trisections.

**Theorem 1.2.** *Every smooth (or algebraically singular) symplectic surface in  $\mathbb{C}\mathbb{P}^2$  is symplectically isotopic to one in bridge position with respect to a genus one Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$ .*

Once we know that we can symplectically isotope our branch locus into bridge position, to prove that every symplectic manifold has a Weinstein trisection, it remains to show that (a) pulling back the (topological) trisection of  $\mathbb{C}\mathbb{P}^2$  by  $f$  gives a topological trisection of  $X$ , and (b) pulling back the Weinstein structure on each sector by  $f$  gives a Weinstein structure compatible with the global symplectic form. These tasks are carried out in Sections 3 and 6, respectively. We conclude with a discussion of examples in Section 7.

**Acknowledgements.** This project arose thanks to the support of an AIM SQuaRE grant. The authors would like to express their gratitude to the American Institute of Mathematics for providing an ideal collaborative environment, as well as acknowledge Paul Melvin, Juanita Pinzón-Caicedo, and Alex Zupan for their input at the beginning of the project. JM was supported by NSF grant DMS-1933019. LS was supported by NSF grant DMS-1904074. The authors also thank the Max Planck Institute for Mathematics for its hospitality.

## 2. DEFINITIONS, NOTIONS OF EQUIVALENCE, AND OPEN QUESTIONS

In this section, we give formal definitions for the concepts from the introduction, discuss ways in which two Weinstein trisections could be considered equivalent, and present a handful of open questions that we hope will direct and encourage future development.

A *trisection*  $\mathbb{T}$  of a 4–manifold  $X$  is a decomposition  $X = Z_1 \cup Z_2 \cup Z_3$  into three pieces such that, for each  $\lambda \in \mathbb{Z}_3$ ,

- (1) Each  $Z_\lambda$  is diffeomorphic to  $\natural^{k_\lambda}(S^1 \times B^3)$ , for some  $k_\lambda$ ,
- (2) Each double intersection  $H_\lambda = Z_{\lambda-1} \cap Z_\lambda$  is a 3–dimensional, genus  $g$  handlebody, and
- (3) The triple intersection  $\Sigma = Z_1 \cap Z_2 \cap Z_3$  is a closed, genus  $g$  surface, with  $\Sigma = \partial H_\lambda$ .

The surface  $\Sigma$  is called the *core* of the trisection, and the genus of the core is the *genus* of the trisection.

Gay and Kirby showed that every closed, oriented, smooth 4–manifold  $X$  admits a trisection  $\mathbb{T}$  and that any two trisections  $\mathbb{T}$  and  $\mathbb{T}'$  of  $X$  have a common stabilization [GK16]. Trisections of low genus

have been classified [MZ17b, MSZ16], and trisections for many familiar 4-manifolds, including complex hyper-surfaces in  $\mathbb{C}\mathbb{P}^3$ , have been described [LM18].

Suppose that  $X$  admits a symplectic structure  $\omega$  and a trisection  $\mathbb{T}$ . We want to understand whether the symplectic form is compatible with the 1-handlebody structure on each of the three sectors  $Z_\lambda$ . Weinstein's handle construction gives a notion of compatibility between a symplectic structure and a 1-handlebody structure [Wei91].

We now give a more analytic way to keep track of Weinstein's handle structure. A *Weinstein structure* on a 4-manifold  $W$  is a quadruple  $(W, \omega, V, \phi)$  where  $W$  is a smooth manifold with boundary,  $\omega$  is the symplectic structure,  $V$  is a Liouville vector field for  $\omega$  which is outwardly transverse to the boundary, and  $\phi: W \rightarrow \mathbb{R}$  is a Morse function such that  $V$  is gradient-like for  $\phi$ . To say a vector field  $V$  is Liouville for  $\omega$  means that the 1-form  $\eta := \iota_V \omega$  obtained by taking the interior product of  $\omega$  by  $V$  (i.e.  $\eta(\cdot) = \omega(V, \cdot)$ ) satisfies  $d\eta = \omega$ . To say that  $V$  is gradient-like for  $\phi$  implies that the zeros of  $V$  are the critical points of  $\phi$  and  $d\phi(V) > 0$  whenever  $V \neq 0$ . The data of the function  $\phi$  is less important than the Liouville vector field  $V$  (because  $V$  is what must be compatible with the symplectic form and different choices of  $\phi$  for a fixed  $V$  are equivalent), but the *existence* of some function  $\phi$  for which  $V$  is gradient-like is important. The existence of  $\phi$  is equivalent to asking that the zeros of  $V$  are locally modeled on the gradient of a Morse function, and  $V$  should have no limit cycles (no periodic orbits and no oriented cyclic loops of trajectories). Therefore, we can focus solely on the Liouville vector field  $V$ , as long as we control its trajectories and zeros. A key feature of a Weinstein structure is that it induces a contact structure on the boundary of the handlebody. The contact form is given by the restriction of  $\eta$  to the boundary.

Each piece  $Z_\lambda$  of the trisection is a 4-dimensional 1-handlebody and therefore admits a (sub-critical) Weinstein structure  $(Z_\lambda, \omega_\lambda, V_\lambda, \phi_\lambda)$ . Thus, one can ask whether such Weinstein structures on the  $Z_\lambda$  can be chosen compatibly with the global symplectic form  $\omega$  on  $X$ .

**Definition 2.1.** A *Weinstein trisection* of a symplectic 4-manifold  $(X, \omega)$  consists of a trisection  $\mathbb{T}$ , with induced decomposition  $X = Z_1 \cup Z_2 \cup Z_3$ , and a Weinstein structure  $(Z_\lambda, \omega|_{Z_\lambda}, V_\lambda, \phi_\lambda)$  on each sector using the restriction of the symplectic form  $\omega$ . In this case, we say  $\mathbb{T}$  and  $\omega$  are *compatible*.

**Remark 2.2.** Note that the sectors  $Z_\lambda$  are manifolds with boundary *and corners*, where the corner is along the trisection surface  $\Sigma$ . Typically, we define a Weinstein domain structure on a manifold with boundary, requiring the Liouville vector field to be outwardly transverse to the boundary. In this situation with corners, we take a sequence of approximating hypersurfaces approaching the boundary which smooth the corner (in the canonical manner), and require that the Liouville vector field is transverse to sufficiently nearby smoothings of the boundary.

**Example 2.3.** Consider  $\mathbb{C}\mathbb{P}^2$  with homogeneous coordinates  $[z_1 : z_2 : z_3]$ , and consider the subsets

$$\begin{aligned} Z_\lambda &= \{[z_1 : z_2 : z_3] \mid |z_\lambda|, |z_{\lambda+1}| \leq |z_{\lambda+2}|\} \\ H_\lambda &= \{[z_1 : z_2 : z_3] \mid |z_\lambda| \leq |z_{\lambda+1}| = |z_{\lambda+2}|\}. \end{aligned}$$

This decomposition yields a Weinstein trisection  $\mathbb{T}_0$  for  $(\mathbb{C}\mathbb{P}^2, \omega_{FS})$  first given in [LM18], where  $\omega_{FS}$  denotes the Fubini-Study symplectic form. This trisection is compatible with the toric structure on  $\mathbb{C}\mathbb{P}^2$ , and Figure 1 shows the decomposition in the image of the moment map.

The corners of  $\partial Z_\lambda$  can be smoothed by approximating  $\partial Z_\lambda$  by  $f_{\lambda, N}^{-1}(1)$  where

$$f_{\lambda, N}(z_\lambda, z_{\lambda+1}) := \frac{1}{N}(|z_\lambda|^2 + |z_{\lambda+1}|^2) + |z_\lambda|^{2N} + |z_{\lambda+1}|^{2N}$$

for  $N \gg 0$  (as in [Lam18]).




FIGURE 1. The standard genus one Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$  coming from toric geometry.

**Definition 2.4.** Two Weinstein trisections  $(X, \mathbb{T}, \omega_0)$  and  $(X, \mathbb{T}, \omega_1)$  are *isotopic* if there exists a family of Weinstein trisections  $(X, \mathbb{T}, \omega_t)$ , with  $t \in [0, 1]$ , such that  $\omega_t$  is a family of symplectic forms interpolating between  $\omega_0$  and  $\omega_1$  with  $[\omega_t] \in H^2(X; \mathbb{R})$  independent of  $t$ .

Given such a family of symplectic forms, Moser's trick implies that there exists an ambient isotopy  $\Psi_t: X \rightarrow X$  such that  $\Psi_0 = id$  and  $\Psi_t^*(\omega_t) = \omega_0$ . Therefore, we could equivalently consider the family  $(X, \mathbb{T}_t, \omega_0)$  where the trisection  $\mathbb{T}_t = \Psi_t(\mathbb{T})$  varies through an isotopy and the symplectic form is fixed.

Even if two symplectic structures are isotopic, and both are compatible with the same trisection, it is not clear whether the isotopy between them can remain compatible with the fixed trisection.

**Question 2.5.** *Suppose that  $(X, \mathbb{T}, \omega_0)$  and  $(X, \mathbb{T}, \omega_1)$  are Weinstein trisections and that  $\omega_0$  and  $\omega_1$  are isotopic via a family  $\omega_t$  such that  $[\omega_t] \in H^2(X; \mathbb{R})$  is independent of  $t$ . Are  $(X, \mathbb{T}, \omega_0)$  and  $(X, \mathbb{T}, \omega_1)$  necessarily isotopic as Weinstein trisections?*

Recall that the *trisection genus* of a 4-manifold  $X$  is the minimum value  $g$  such that  $X$  admits a trisection with core surface of genus  $g$ .

**Definition 2.6.** The *Weinstein trisection genus* of a symplectic 4-manifold  $(X, \omega)$  is the minimum value  $g$  such that  $(X, \omega)$  admits a Weinstein trisection with core surface of genus  $g$ .

**Question 2.7.** *Is the Weinstein trisection genus of a symplectic 4-manifold  $(X, \omega)$  always the same as the trisection genus of  $X$ ?*

Note that there are examples of symplectic 4-manifolds whose Weinstein trisection genus (for some symplectic form) equals their trisection genus. These include complex hypersurfaces in  $\mathbb{C}\mathbb{P}^3$  and the elliptic surfaces  $E(n)$ ; compare [LM18, Theorem 1.1] with Theorem 6.1 and Example 7.1 below.

Given a genus  $g$  trisection  $\mathbb{T}$  for a 4-manifold  $X$ , there is a natural process, called *stabilization*, that can be applied to produce a genus  $g + 1$  trisection  $\mathbb{T}'$  of  $X$ . Gay and Kirby proved that any two trisections of  $X$  have a common stabilization. (See [MSZ16] and [GK16] for details.) If we start with a genus  $g$  Weinstein trisection, we can perform this stabilization in a way that yields a genus  $g + 1$  Weinstein trisection.

**Proposition 2.8.** *Let  $(X, \mathbb{T}, \omega)$  be a genus  $g$  Weinstein trisection, then any trisection stabilization can be performed such that the resulting  $g + 1$  trisection is compatible with  $\omega$ .*

*Proof.* The stabilization operation of a trisection can be understood in the following manner. Choose one sector  $Z_\lambda$ . Then  $H_{\lambda+1}$  is the handlebody which is not contained in  $\partial Z_\lambda$ . Let  $a$  be a properly-embedded, boundary-parallel arc in  $H_{\lambda+1}$ . Any stabilization of the trisection is obtained from some

such choice of arc, by adding a regular neighborhood of  $a$  to  $Z_\lambda$  and deleting this regular neighborhood from  $Z_{\lambda\pm 1}$ . This has the effect of adding a 1-handle to  $Z_\lambda$ . It does not change the topology of  $Z_{\lambda\pm 1}$  because the arc  $a$  lay in the boundary of each of these sectors, so deleting its neighborhood simply carves out a “bite” from their boundaries. The new pairwise intersections are still 3-dimensional handlebodies when  $a$  is boundary parallel.

Now we consider how this construction interacts with the symplectic form. Any choice of arc  $a$  is isotropic, so it has a standard symplectic neighborhood in  $(X, \omega)$ . This neighborhood can be identified with Weinstein’s model for a symplectic 1-handle attachment (identifying  $a$  with the core of the 1-handle). In particular, the Weinstein structure on  $Z_\lambda$  extends over any sufficiently small neighborhood of  $a$  compatibly with the fixed symplectic form  $\omega$ .




FIGURE 2. Dimensionally reduced neighborhood  $N$  of the arc  $a$  giving a Weinstein stabilization.

To ensure that the Weinstein structures on  $Z_{\lambda\pm 1}$  remain Weinstein after deleting the neighborhood of  $a$ , we choose the neighborhood  $N$  of  $a$  so that the “bite” is sufficiently *shallow*. We will simply restrict the original Weinstein structure on  $Z_{\lambda\pm 1}$  to  $Z_{\lambda\pm 1} \setminus N$ . As long as the boundary  $\partial(Z_{\lambda\pm 1} \setminus N)$  is sufficiently  $C^1$ -close to the boundary  $\partial Z_{\lambda\pm 1}$ , the Liouville vector field  $V_{\lambda\pm 1}$  will still be outwardly transverse to the boundary. To choose  $N$  to satisfy this, choose symplectic coordinates  $(x_1, y_1, x_2, y_2)$  on a neighborhood of  $a$  in  $X$  such that  $H_{\lambda+1} = \{y_2 = 0\}$  and  $a = \{y_1 = x_2 = y_2 = 0\}$ . Let

$$N = \{(x_1, y_1, x_2, y_2) \mid y_1^2 + x_2^2 \leq h(|y_2|)^2, y_2 \in [-\varepsilon, \varepsilon]\}$$

where  $h: [0, \varepsilon] \rightarrow [0, \delta]$  is a monotonically decreasing function. See Figure 2 for a dimensionally reduced (there is only one dimension representing both the  $y_1$  and  $x_2$  directions) picture for the shape of  $N$ . Choosing  $\varepsilon/\delta$  sufficiently small ensures that  $N$  is sufficiently shallow so the new boundary is  $C^1$ -close to the old boundary. Note that because of the absolute value in the argument for  $h$  in the definition of  $N$ ,  $N$  has boundary and corners, but this is expected in a trisection, and we smooth corners as discussed above.  $\square$

Even if the answer to Question 2.5 is negative, it could be possible that after stabilization, the answer becomes positive.

**Question 2.9.** *Suppose that  $(X, \mathbb{T}, \omega_0)$  and  $(X, \mathbb{T}, \omega_1)$  are Weinstein trisections and that  $\omega_0$  and  $\omega_1$  are isotopic via a family  $\omega_t$  such that  $[\omega_t] \in H^2(X; \mathbb{R})$  is independent of  $t$ . Does there exist a common stabilization  $\mathbb{T}'$  such that  $(X, \mathbb{T}', \omega_0)$  and  $(X, \mathbb{T}', \omega_1)$  are isotopic through Weinstein trisections?*

More generally, we can ask whether Gay-Kirby’s common stabilization theorem holds in the Weinstein setting.

**Question 2.10.** *Suppose  $(X, \mathbb{T}, \omega)$  and  $(X, \mathbb{T}', \omega')$  are Weinstein trisections such that  $\omega = \omega'$  (or, alternatively, such that  $\omega$  and  $\omega'$  are isotopic). Is there a Weinstein trisection  $(X, \mathbb{T}'', \omega)$  such that  $\mathbb{T}''$  is a stabilization of both  $\mathbb{T}$  and  $\mathbb{T}'$ ?*

A long-standing point of interest in symplectic geometry is the question of how to compare different symplectic structures on a fixed 4-manifold. The ability to stabilize Weinstein trisection provides a possible avenue to address this question in the following way. Given two symplectic structures  $\omega_1$  and  $\omega_2$  on  $X$ , we can find Weinstein trisections  $\mathbb{T}_1$  and  $\mathbb{T}_2$  for  $X$  that are compatible with  $\omega_1$  and  $\omega_2$ , respectively. By stabilizing, we obtain Weinstein trisections  $\mathbb{T}'_1$  and  $\mathbb{T}'_2$  that are isotopic as trisections, but may or may not be isotopic via a family of Weinstein trisections. Thus, this allows us to compare the symplectic structures  $\omega_1$  and  $\omega_2$  on  $X$  relative to a fixed trisection  $\mathbb{T}'$  of  $X$ . To this end, Questions 2.5, 2.9, and 2.10 are relevant.

### 3. SINGULAR BRIDGE TRISECTIONS AND BRANCHED COVERINGS

In this section, we discuss singular surfaces in 4-manifolds and their branched coverings, along with their compatibility with symplectic structures and trisections.

#### 3.1. Singular bridge trisections.

Let  $X$  be a closed, orientable, smooth 4-manifold. We call a closed surface  $\mathcal{S} \subset X$  a *singular surface-link* provided that

- (1)  $\mathcal{S}$  is smoothly immersed away from finitely many singular points; and
- (2) in a small neighborhood of each singular point,  $\mathcal{S}$  is homeomorphic to the cone on a knot in  $S^3$ .

The transverse self-intersections of  $\mathcal{S}$  are called *nodes*, and a singular point is called a (*simple*) *cusplike* if it is given as the cone on a trefoil knot.

For a singular surface-link to be *symplectic*, we want its tangent spaces to be symplectic subspaces where they are defined, and at singular points, we want specific local models.

**Definition 3.1.** A *symplectic singular surface* in a symplectic 4-manifold  $(X, \omega)$  is the image  $Q = g(S)$  of a map  $g: S \rightarrow X$ , with a finite set of points  $P = \{p_1, \dots, p_n\} \subset S$ , satisfying the following properties.

- (1)  $g|_{S \setminus P}$  is a smooth immersion.
- (2)  $\omega$  is positive on  $dg_p(T_p S)$  for  $p \in S \setminus P$ .
- (3) For each  $i \in \{1, \dots, n\}$ , there exists a neighborhood  $B_i$  of  $g(p_i)$  in  $X$ , and a diffeomorphism  $\Psi: B_i \rightarrow U_i \subset (\mathbb{C}^2, \omega_{std})$  such that if  $N_i$  is the connected component of  $g^{-1}(B_i)$  in  $S$  then  $\Psi \circ g(N_i) \subset U_i \subset \mathbb{C}^2$  is cut out by a complex polynomial. (Each branch of the singular surface is smoothly modeled on a complex algebraic curve.)
- (4) For each  $p \in P$ , let  $T_p$  be the limiting tangent space  $T_p = \lim_{p_k \rightarrow p} dg_{p_k}(T_{p_k} S)$ ; then,  $\omega$  is positive on  $T_p$ .

A 1-parameter family  $Q_t = g_t(S)$  of symplectic singular surfaces in  $(X, \omega)$  such that the links of the singular points remain smoothly isotopic knots/links throughout the family, will be called a *topologically equisingular symplectic isotopy*.

Note that the analytic type of the singularities may change. In particular, if  $q_0 \in Q_0$  and  $q_1 \in Q_1$  are corresponding singularities in such a family, there need *not* exist neighborhoods  $q_i \in U_i \subset X$  and a symplectomorphism  $\Phi: U_0 \rightarrow U_1$  such that  $\Phi(Q_0 \cap U_0) = Q_1 \cap U_1$ . Therefore, there need not be an ambient symplectic isotopy inducing this isotopy of surfaces. In other words, there may not be a family of symplectomorphisms  $\Psi_t: (X, \omega) \rightarrow (X, \omega)$  such that  $\Psi_t(Q_0) = Q_t$ . However, the next lemma shows that the local singularities are the only obstruction to finding such an ambient symplectic isotopy.

**Lemma 3.2.** *Let  $Q_t = g_t(S)$  be a topologically equisingular symplectic isotopy, for  $t \in [0, 1]$ . Let  $\{U_t^i\}_{i=1}^N$  be small neighborhoods of the singular points of  $Q_t$ . Then there exists a family of symplectomorphisms  $\Psi_t: (X, \omega) \rightarrow (X, \omega)$  such that*

$$\Psi_t(Q_0) \setminus \left( \bigcup_{i=1}^N U_t^i \right) = Q_t \setminus \left( \bigcup_{i=1}^N U_t^i \right).$$

*Proof.* The case when  $Q_t$  is a smooth symplectic submanifold is proven in [Aur97, Proposition 4] using a Moser argument (in this case there are no neighborhoods  $U_t^i$  of singular points). We review this argument inserting the necessary modifications for the singular case.

The first step is to find neighborhoods  $N_t$  of  $Q_t$  and symplectomorphisms  $\psi_t: N_0 \rightarrow N_t$  such that  $\psi_t(Q_0)$  agrees with  $Q_t$  outside of the neighborhoods  $U_t^i$ . Outside of any small neighborhood of the singular points, where  $Q_t$  is smooth and symplectic, it has a standard symplectic neighborhood (because the symplectic normal bundle is uniquely determined by the smooth normal bundle). This defines  $N_t$  and  $\psi_t$  away from the singular points. Each sufficiently small neighborhood of a singular point has a Darboux chart (though the precise model for the intersection of this Darboux chart with  $Q_t$  may vary). These standard pieces can be glued together compatibly along a contactomorphism of  $(S^3, \xi_{std})$  along the boundary of the Darboux chart. If the topological singularity type is fixed for all  $t$ , the transverse link of the singularity is the same up to transverse isotopy for all  $t$ . Since any transverse isotopy can be realized by a contact isotopy, we can assume that the gluing aligns the smooth piece of  $Q_t$  with the singular part of  $Q_t$ . Note that every contactomorphism of  $(S^3, \xi_{std})$  extends to a symplectomorphism of  $(B^4, \omega_{std})$  so while we may not have control over the precise image of  $Q_t$  near its singular points, we do know that we can extend the standard neighborhood on the smooth part to a symplectomorphism  $\psi_t: N_0 \rightarrow N_t$  such that  $\psi_t(Q_0) \setminus (\cup_i U_t^i) = Q_t \setminus (\cup_i U_t^i)$ .

Next, we extend  $\psi_t$  to a diffeomorphism  $F_t: X \rightarrow X$  such that  $F_0$  is the identity. Because this extension was arbitrary, we do not expect  $\omega_t := F_t^*(\omega)$  to equal  $\omega$  outside of the neighborhood  $N_0$  for  $t \neq 0$ , but we will use Moser's trick to construct another family of diffeomorphisms  $\tau_t: X \rightarrow X$  such that  $\tau_t^*(F_t^*\omega) = \omega$ , such that  $\tau_t(F_t(Q_0)) = F_t(Q_0)$ . Note that  $F_t(Q_t)$  agrees by construction with  $Q_t$  outside of the neighborhoods  $U_t^i$ .

In Moser's trick,  $\tau_t$  is defined as the integral flow of a vector field  $V_t$  such that  $d(\iota_{V_t}\omega_t) = -\frac{d\omega_t}{dt}$ . Moser's method verifies that this condition on  $V_t$  is sufficient to ensure that  $\tau_t^*\omega_t = \omega$ , but we need to know that there exists such a  $V_t$  whose flow preserves  $Q_0$ . By non-degeneracy, finding  $V_t$  is equivalent to finding  $\alpha_t = \iota_{V_t}\omega_t$  such that  $d\alpha_t = -\frac{d\omega_t}{dt}$ . We want  $V_t$  to vanish at the singularities of  $Q_0$  and to be tangent to  $Q_0$  at smooth points. Since  $\omega_t = \omega$  in a neighborhood of  $Q_0$  (because  $F_t$  agrees with the symplectomorphism  $\psi_t$  here), these conditions are equivalent to asking that  $\alpha_t$  vanishes at the singularities of  $Q_0$  and  $T_p Q_0^{\perp \omega} \subset \ker(\alpha_t)_p$  at each smooth point  $p \in Q_0$ . We know that  $-\frac{d\omega_t}{dt}$  is zero in cohomology so it has some family of primitives  $\beta_t$ . We will modify  $\beta_t$  to a 1-form  $\alpha_t$ , differing from  $\beta_t$  by an exact form and satisfying the desired tangency and vanishing criteria along  $Q_0$ .

First, choose a 1-form  $\delta_t$  on  $Q_0$  such that  $\delta_t = 0$  in an  $2\varepsilon$ -neighborhood of the singularities such that  $i^*[\beta_t] = [\delta_t] \in H^1(D_0; \mathbb{R})$  where  $i: Q_0 \rightarrow N_0$  is the inclusion. Let  $\pi: N_0 \setminus (\cup_i B^i) \rightarrow Q_0 \setminus (\cup_i U_0^i)$  be the symplectic orthogonal projection of the neighborhood of  $Q_0$  to  $Q_0$  outside a  $\varepsilon$ -neighborhood of the singularities (we can extend  $\pi$  arbitrarily near the singularities). In the  $\varepsilon$ -neighborhoods of the singularities, let  $\gamma_t = 0$  and elsewhere let  $\gamma_t = \pi^*\delta_t$ . Then  $\gamma_t$  are smooth 1-forms,  $i^*[\gamma_t] = [\delta_t] = i^*[\beta_t]$  so  $[\gamma_t] = [\beta_t]$  since  $i$  is a homotopy equivalence. Additionally,  $\gamma_t$  satisfies  $(T_x C_0)^{\perp \omega} \subseteq \ker \gamma_t$  at all points and  $\gamma_t = 0$  at the singularities of  $Q_0$ . Now write  $\gamma_t - \beta_t = df_t$  for functions  $f_t: U_0 \rightarrow \mathbb{R}$ , and choose arbitrary extensions  $\tilde{f}_t: M \rightarrow \mathbb{R}$ . Then  $\alpha_t = \beta_t + d\tilde{f}_t$  agrees with  $\gamma_t$  in  $U_0$  (so  $\alpha_t$  has the required vanishing and kernel conditions along  $C_0$ ) and it satisfies  $d\alpha_t = d\beta_t = -\frac{d\omega_t}{dt}$  as required.  $\square$

Next we explain how singular surfaces can interact well with a trisection.

Let  $L = L_1 \sqcup \cdots \sqcup L_n$  be a link in  $S^3 = \partial B^4$  given as the split union of non-split links  $L_i$ . Let  $\mathcal{D} \subset B^4$  be a collection of disks with  $\partial \mathcal{D} = L$ . We call  $\mathcal{D}$  a *singular disk-tangle* for  $L$  if

- (1) each component of  $\mathcal{D}$  with unknotted boundary is a smoothly embedded, boundary parallel disk,
- (2) each component of  $\mathcal{D}$  with knotted boundary is a cone on that knot, and
- (3) the components of  $\mathcal{D}$  are disjoint when their boundaries are split and intersect transversely – away from all cone points – otherwise.

For example, if  $L$  were the split union of a the torus link  $T(9, 6)$ , whose components are three trefoils, and the Hopf link  $T(2, 1)$ , then  $\mathcal{D}$  would consist of three cones on trefoils, which intersect pairwise transversely in six points, split union two smoothly embedded disks, which intersect transversely in a single point. Note that this is a generalization of the notion of a trivial disk-tangle appearing elsewhere in the literature [CK17, LM18, MZ17a, MZ18].

We extend the notion of a singular disk-tangle to immersed collections of disks  $\mathcal{D}$  in  $\mathbb{t}^k(S^1 \times B^3)$  by requiring that  $\mathcal{D}$  be contained in a properly embedded 4–ball.

**Definition 3.3.** Let  $X$  be a 4–manifold equipped with a trisection  $\mathbb{T}$ , and let  $\mathcal{S} \subset X$  be a singular surface-link. We say that  $\mathcal{S}$  is in *bridge trisected position* with respect to  $\mathbb{T}$  if, for each  $\lambda \in \mathbb{Z}_3$ ,

- (1)  $\mathcal{T}_\lambda = H_\lambda \cap \mathcal{S}$  is a trivial tangle, and
- (2)  $\mathcal{D}_\lambda = Z_\lambda \cap \mathcal{S}$  is a singular disk-tangle for the link  $K_\lambda = \mathcal{T}_\lambda \cup \mathcal{T}_{\lambda+1}$ .

We refer to the  $\mathcal{D}_\lambda$  as *patches* and to the  $\mathcal{T}_\lambda$  as *seams*. The decomposition

$$(X, \mathcal{S}) = (Z_1, \mathcal{D}_1) \cup (Z_2, \mathcal{D}_2) \cup (Z_3, \mathcal{D}_3)$$

is called a *singular bridge trisection*.

A straight-forward adaptation of the techniques of [MZ18] to the setting of singular surface-links reveals that any singular surface-link can be put in bridge trisected position with respect to any trisection. (The main idea: Treat the singular points as minima and follow the Morse-theoretic construction of the bridge trisected position.) In Section 5, we prove a stronger result for symplectic braided surfaces, showing that they can be *symplectically* isotoped to lie in bridge trisected position in  $\mathbb{C}\mathbb{P}^2$ .

### 3.2. Singular branched coverings.

We refer the reader to [Zud08] a nice exposition of branched coverings. Most branched coverings considered here will be *simple* (meaning they are modeled by an involution near non-singular branch points), since these are the coverings produced by Auroux and Katzarkov [Aur00, AK00]. However, we will also consider some cyclic branched coverings of higher order when studying particularly nice examples (cf. Definition 3.4 and Section 7). Moreover, some results, such as Theorem 4.1, hold for more general branched coverings. We begin with the following general definition.

**Definition 3.4.** Let  $X$  be a 4–manifold and let  $\mathcal{S}$  in  $X$  be a singular surface-link. A proper map  $f: \tilde{X} \rightarrow X$  is called a *singular branched covering* of  $X$  along  $\mathcal{S}$  if the following two conditions hold:

- (1) Away from the singular points in  $X$ ,  $f$  is a branched covering map.
- (2) In a 4–ball neighborhood  $B \subset X$  of a singular point,  $f^{-1}(B) = \tilde{B}_1 \sqcup \cdots \sqcup \tilde{B}_n$  is the disjoint union of 4–balls such that  $f|_{\tilde{B}_i}: \tilde{B}_i \rightarrow B$  is either the identity or the cone on a branched covering map from  $S^3$  to itself.



Note that the existence of a singular branched covering of  $X$  branched along a singular surface-link  $\mathcal{S}$  restricts the singularities of  $\mathcal{S}$  significantly. In particular, the link of any singularity must be a knot or link that admits a branched covering to the disjoint union of copies of  $S^3$ . On the other hand, there are many examples of such links, including 2-bridge links. We will now discuss explicitly the most important cases for the present work.

**Examples 3.5.** We describe some important examples of singular branched coverings in coordinates. Since we will mainly consider *simple* singular branched coverings in this paper, the following three examples exhaust the relevant local models describing our singular branched covering maps.

- (1) First, consider the standard branching model, described by the map  $f: \mathbb{C}^2 \rightarrow \mathbb{C}^2$  given in complex coordinates by  $f(x, y) = (x^2, y)$ . The 3-sphere  $S_\varepsilon = \{|z|^2 + |w|^2 = \varepsilon\}$  in the codomain has preimage  $f^{-1}(S_\varepsilon) = \{|x|^4 + |y|^2 = \varepsilon\}$  homeomorphic to a 3-sphere, which can be seen by changing coordinates by the scaling map

$$x \mapsto \frac{x}{\sqrt{|x|}}.$$

The singular points of  $f$  occur when  $x = 0$ . The intersection of  $\{x = 0\}$  with  $f^{-1}(S_\varepsilon)$  is an unknot. Thus,  $f|_{S_\varepsilon}$  represents the 2-fold cyclic covering of  $S^3$  over itself, branched along the unknot, and  $f$  can be thought of as the cone of the map  $f_{S_\varepsilon}$ , though in this case, the cone can be treated as a smooth, trivial disk bounded by the unknot.

- (2) Next, we describe the model for the cusp, which is given in complex coordinates by the function

$$f(x, y) = (x^3 - xy, y).$$

The critical points of  $f$  occur where the derivative

$$df_{(x,y)} = \begin{bmatrix} 3x^2 - y & -x \\ 0 & 1 \end{bmatrix}$$

fails to have full rank.

This occurs when  $3x^2 - y = 0$ . Therefore, the critical set of this map is  $\{y = 3x^2\}$ . The critical values of  $f$  are thus the image of this set, namely the points  $(-2x^3, 3x^2)$  for  $x \in \mathbb{C}$ . With coordinates  $(z, w)$  on the codomain, the critical values are the zero locus of the polynomial  $27z^2 - 4w^3$ . Thus the branch locus is parameterized as  $\{(-2c^3, 3c^2) \mid c \in \mathbb{C}\}$  and has a singular point at  $(0, 0)$ . Observe that for each value  $c$ ,  $f^{-1}(-2c^3, 3c^2) = \{(c, 3c^2), (-2c, 3c^2)\}$ . The points  $(c, 3c^2)$  are critical points and have multiplicity two. On the other hand, the points  $(-2c, 3c^2)$  (the locus  $y = \frac{3}{4}x^2$ ) are not critical points when  $c \neq 0$ , but they do project to critical values in the branch locus. This occurs because the branched covering is irregular.

To view this as a cone over a branched covering of  $S^3$ , consider again the 3-sphere  $S_\varepsilon$ , and let  $\tilde{S}_\varepsilon = f^{-1}(S_\varepsilon) = \{|x^3 - xy|^2 + |y|^2 = \varepsilon\}$ . The singular curve  $\{27z^2 - 4w^3 = 0\}$  intersects  $S_\varepsilon$  along the right-handed trefoil  $K$ , and  $f$  is the cone of the restriction  $f|_{\tilde{S}_\varepsilon}: \tilde{S}_\varepsilon \rightarrow S_\varepsilon$ .

In a neighborhood of any point on  $K \subset S_\varepsilon$ ,  $f|_{\tilde{S}_\varepsilon}$  is given by the standard branching model described above. It follows that  $f|_{\tilde{S}_\varepsilon}$  is a 3-fold, simple covering of  $S_\varepsilon$ , branched along  $K$ . Such a covering is determined by a map  $\rho: \pi_1(S_\varepsilon \setminus \nu(K)) \rightarrow \mathfrak{S}_3$  sending meridians of  $K$  to transpositions. (See [Zud08] for definitions and details.) Since such surjections are unique up to conjugation for the trefoil, it follows that  $f|_{\tilde{S}_\varepsilon}$  is the (only) irregular 3-fold branched cover over the trefoil. Thus  $f^{-1}(S_\varepsilon) \cong S^3$ , as desired; the cusp map is the cone on this branched covering.

- (3) Finally, consider the quotient map  $f: \mathbb{C}_1^2 \sqcup \mathbb{C}_2^2 \rightarrow \mathbb{C}^2$  defined by  $f(x_1, y_1) = (x_1^2, y_1)$  for  $(x_1, y_1) \in \mathbb{C}_1^2$  and  $f(x_2, y_2) = (x_2, y_2^2)$  for  $(x_2, y_2) \in \mathbb{C}_2^2$ . This can also be thought of as the quotient map by

the equivalence relation where  $(x_1, y_1) \sim (-x_1, y_1)$ ,  $(x_2, y_2) \sim (x_2, -y_2)$ , and  $(x_1, y_1) \sim (x_2, y_2)$  if  $x_1^2 = x_2$  and  $y_1 = y_2^2$  for any  $(x_1, y_1) \in \mathbb{C}_1^2$  and  $(x_2, y_2) \in \mathbb{C}_2^2$ . Over points  $(z, w) \in \mathbb{C}^2$ , this map is four-to-one when  $z \neq 0$  and  $w \neq 0$ , is three-to-one when either  $z = 0$  or  $w = 0$  (but not both), and is two-to-one when  $(z, w) = (0, 0)$ .

We can view this map as the disjoint union of cones of branched coverings from  $S^3$  to  $S^3$  as follows. For  $i = 1, 2$ , the restriction  $f_i$  of  $f$  to  $\mathbb{C}_i$  is simply the map from Example (1) above (up to swapping the coordinates in  $\mathbb{C}_2^2$ ). Each of these is a cone, so  $f$  is the disjoint union of cones on these maps

Finally, note that in this case the link  $S_\varepsilon \cap \{zw = 0\}$  is simply the Hopf link, the lift  $f^{-1}(S_\varepsilon)$  is the disjoint union of two 3-spheres, and the restriction  $f|_{f^{-1}(S_\varepsilon)}$  is a 4-fold covering of  $S^3$ , branched along the Hopf link. The preimage of the Hopf link is a pair of Hopf links in the two covering 3-spheres, and each component of the link downstairs is double covered by one component upstairs and single covered by one component upstairs. This covering is determined by the map  $\rho: \pi_1(S_\varepsilon) \rightarrow \mathcal{S}_4$  that sends the meridians of the Hopf link to commuting transpositions, say (12) and (34). Again, this map is unique up to conjugation, but this time it is not surjective. The non-surjectivity of  $\rho$  corresponds to the fact that the branched covering is disconnected.

Note: Reversing the orientation on  $\mathbb{C}^2$  to disagree with the complex orientation changes this to the model for a negative node.

Our use of singular branched coverings here comes from their connection with symplectic manifolds. The following definition captures the key result that we need from Auroux's work.

**Definition 3.6.** A map  $f: X \rightarrow \mathbb{C}\mathbb{P}^2$  is a *symplectic singular branched covering* if at any point it is modeled on one of the three following maps.

- (1) local diffeomorphism:  $(z_1, z_2) \mapsto (z_1, z_2)$
- (2) cyclic branched covering:  $(z_1, z_2) \mapsto (z_1^d, z_2)$
- (3) cusp covering:  $(z_1, z_2) \mapsto (z_1^3 - z_1 z_2, z_2)$

Moreover, the branch locus  $R \subset X$  (where  $f$  cannot be modeled by the local diffeomorphism) and its image  $f(R)$  must be (singular) symplectic submanifolds, and the 2-form  $\tilde{\omega}_t := tf^*\omega_{FS} + (1-t)k\omega$  is a symplectic form for all  $t \in [0, 1)$ , where  $k$  is the degree of  $f$ .

The branched covering and cusp covering models were examined in detail in Examples 3.5 above. Note that these models are local in the domain of the map, and the images of these local charts in the co-domain  $\mathbb{C}\mathbb{P}^2$  may overlap. In particular, the image of the branching locus may self-intersect (either positively or negatively), as in the node in Example 3.5 (3).

**Theorem 3.7** ([Aur00]). *Every closed symplectic 4-manifold admits a symplectic singular branched covering.*

**Remark 3.8.** The main theorem statement in [Aur00] is that every closed symplectic 4-manifold admits an  $\varepsilon$ -holomorphic singular covering. The  $\varepsilon$ -holomorphic condition is related to an almost complex structure  $J$  which is compatible with the symplectic form. This condition means that the coordinate charts used to identify  $f$  locally with the models take  $J$  very close to the standard integrable complex structure on  $\mathbb{C}^2$ . For the purposes of this paper, we want to deal directly with the symplectic form instead of a compatible almost complex structure. If  $\varepsilon < 1$ , the branch locus  $R$  and its image  $f(R)$  are (singular) symplectic submanifolds. Auroux also proved that a  $\varepsilon$ -holomorphic singular covering also satisfies the symplectic condition that  $\tilde{\omega}_t := tf^*\omega_{FS} + (1-t)k\omega$  is symplectic for  $t \in [0, 1)$  as [Aur00, Proposition 11], so Theorem 3.7 follows from combining Theorem 1 and Proposition 11 in [Aur00]. By

Moser's theorem  $(X, \tilde{\omega}_t)$  is symplectomorphic to  $(X, k\omega)$  for  $t < 1$ . The pull-back form  $f^*\omega_{FS}$  is not itself symplectic because it is degenerate along the branch locus. However, the condition that  $\tilde{\omega}_t$  is symplectic for  $t < 1$  shows that  $f^*\omega_{FS}$  is arbitrarily close to a symplectic form that is symplectomorphic to  $(X, k\omega)$ . In fact, there is an explicit small perturbation of  $f^*\omega_{FS}$  which is localized near the branch locus that yields a symplectic form directly from the smooth map  $f$ . We will discuss and use this perturbation in detail in Section 6.

**Remark 3.9.** Auroux actually shows that such symplectic singular branched coverings exist without needing the branched covering model for  $d > 2$ . The  $d = 2$  (simple) branched covering model and the cusp covering model are the generic singularities that occur in holomorphic maps from one complex 2-dimensional manifold to another. We allow the non-generic  $d > 2$  branched covering model because we can sometimes find a map  $f$  with significantly simpler branch locus if we allow the branched covering model with higher values of  $d$ . Examples of this are included in Section 7.

#### 4. SINGULAR BRANCHED COVERS PULL BACK TRISECTIONS

In this section we prove that the pull back of a trisection under a singular branched cover is a trisection. This part is purely about the smooth topology and holds for general singular branched coverings. We will address the symplectic aspects in Section 6.

**Theorem 4.1.** *Let  $X$  be a 4-manifold, and let  $\mathbb{T}$  be a trisection on  $X$  given by the decomposition*

$$X = Z_1 \cup Z_2 \cup Z_3.$$

*Let  $f: \tilde{X} \rightarrow X$  be a singular branched covering with branch locus the singular surface-link  $\mathcal{S}$  in  $X$ . Assume that  $\mathcal{S}$  is in bridge trisected position with respect to  $\mathbb{T}$ . Then, letting  $\tilde{Z}_\lambda = f^{-1}(Z_\lambda)$ , the decomposition*

$$\tilde{X} = \tilde{Z}_1 \cup \tilde{Z}_2 \cup \tilde{Z}_3$$

*is a trisection of  $\tilde{X}$ , provided  $\tilde{X}$  is connected.*

*Proof.* Let  $H_\lambda = Z_\lambda \cap Z_{\lambda-1}$ , and let  $\Sigma = \partial H_\lambda$ . Let  $\tilde{H}_\lambda = f^{-1}(H_\lambda)$ , and let  $\tilde{\Sigma} = f^{-1}(\Sigma)$ . Note that  $\tilde{H}_\lambda = \tilde{Z}_\lambda \cap \tilde{Z}_{\lambda-1}$  and  $\tilde{\Sigma} = \tilde{Z}_1 \cap \tilde{Z}_2 \cap \tilde{Z}_3$ .

Consider the restriction  $f|_{\tilde{\Sigma}}: \tilde{\Sigma} \rightarrow \Sigma$ . This map is a covering of the closed, orientable surface  $\Sigma$ , branched along the bridge points  $\mathbf{x} = \Sigma \cap \mathcal{S}$ . It follows that  $\tilde{\Sigma}$  is a closed, orientable surface. A priori,  $\tilde{\Sigma}$  may be disconnected, however.

Next, consider the restriction  $f|_{\tilde{H}_\lambda}$ . This map is a covering of the handlebody  $H_\lambda$ , branched along the seams  $\mathcal{T}_\lambda = H_\lambda \cap \mathcal{S}$ , which form a trivial tangle  $(H_\lambda, \mathcal{T}_\lambda)$ . Such a trivial tangle is homeomorphic to  $(S, \mathbf{y}) \times I$ , where  $S$  is a compact, orientable surface with boundary and  $\mathbf{y} \in \text{Int}(S)$  is a collection of points. (This is the definition of a trivial tangle.) It follows that  $f|_{\tilde{H}_\lambda}$  has the structure of a covering of  $S$ , branched along the points  $\mathbf{y}$ , crossed with  $I$ . Since such a cover of  $S$  is, again, a compact, orientable surface with boundary, we see that  $\tilde{H}_\lambda$  is a handlebody (or at least a disjoint union of handlebodies). Note that  $\partial \tilde{H}_\lambda = \tilde{\Sigma}$ , so  $\tilde{\Sigma}$  is connected if and only if the  $\tilde{H}_\lambda$  are.

Finally, consider the restriction  $f|_{\tilde{Z}_\lambda}$ . It remains to see that  $\tilde{Z}_\lambda$  is a 4-dimensional 1-handlebody. Break  $Z_\lambda$  up into two pieces based on the components of the disk-tangle  $\mathcal{D}_\lambda$ . Recall that each connected component of  $\mathcal{D}_\lambda$  (as a subset of  $Z_\lambda$ ) has boundary a knot or link. For each connected component  $D$ , consider a properly embedded 4-ball  $B_D$  that splits that component from the rest of  $\mathcal{D}_\lambda$ . Let  $Z^{\text{branch}}$  denote the disjoint union of these 4-balls. Let  $Z^{\text{rest}}$  denote  $Z_\lambda \setminus \text{Int}(Z^{\text{branch}})$ . Note that  $Z^{\text{rest}}$  is diffeomorphic to  $Z_\lambda$ ; since each component of  $Z^{\text{branch}}$  is a 4-ball with half of its boundary on  $\partial Z_\lambda$ ,

the removal of  $Z^{\text{branch}}$  from  $Z_\lambda$  doesn't change its diffeomorphism type. Let  $\tilde{Z}^{\text{branch}} = f^{-1}(Z^{\text{branch}})$ , and let  $\tilde{Z}^{\text{rest}} = f^{-1}(Z^{\text{rest}})$ .

By definition, the restriction  $f|_{f^{-1}(B_D)}: f^{-1}(B_D) \rightarrow B_D$  is a map of a disjoint union of 4-balls  $\tilde{B}_1, \dots, \tilde{B}_l$  to the 4-ball  $B_D$ , such that each map  $f_j: \tilde{B}_j \rightarrow B_D$  is given as the cone on a (possibly trivial) covering of  $S^3$  by itself, branched along a (possibly empty) subset of the components of the link  $\partial D$ . Note that the 3-sphere boundary of  $B_D$  is divided into two 3-balls,  $\partial B_D = K \cup L$  where  $K = B_D \cap Z^{\text{rest}}$  and  $L = B_D \cap \partial Z_\lambda$ . Note that  $K$  is disjoint from the branch locus and  $L$  contains the link  $\partial D$ . Then for each connected component  $\tilde{B}_j$  of  $f^{-1}(B_D)$ , the boundary 3-sphere of  $\tilde{B}_j$  is decomposed into  $\tilde{K}_j = \tilde{B}_j \cap f^{-1}(K)$  and  $\tilde{L}_j = \tilde{B}_j \cap f^{-1}(L)$ . Because  $K$  is a 3-ball disjoint from the branch locus,  $\tilde{K}_j$  is a disjoint union of 3-balls.  $\tilde{L}_j$  is a 3-sphere with a collection of 3-balls removed. Now, each 4-ball  $\tilde{B}_j$  is attached to  $\tilde{Z}^{\text{rest}}$  along  $\tilde{K}_j$ , a disjoint union of 3-balls. If  $\tilde{K}_j$  consists of a single 3-ball, then the attachment of  $\tilde{B}_j$  adds only a trivial collar to  $\tilde{Z}^{\text{rest}}$ . If  $\tilde{K}_j$  consists of exactly two 3-balls, attaching  $\tilde{B}_j$  to  $\tilde{Z}^{\text{rest}}$  amounts to attaching a 4-dimensional 1-handle. In general, when  $\tilde{K}_j$  consists of  $m$  3-balls, the attachment of  $\tilde{B}_j$  to  $\tilde{Z}^{\text{rest}}$  is equivalent to attaching  $m - 1$  4-dimensional 1-handles.

It follows that  $\tilde{Z}_\lambda = \tilde{Z}^{\text{rest}} \cup \tilde{Z}^{\text{branch}}$  is obtained by attaching 4-dimensional 1-handles to  $\tilde{Z}^{\text{rest}}$ . We next show that  $\tilde{Z}^{\text{rest}}$  is a 4-dimensional 1-handlebody (or a disjoint union thereof), which will imply that  $\tilde{Z}_\lambda$  is a 4-dimensional 1-handlebody (or disjoint union thereof), as desired.

The restriction  $f|_{\tilde{Z}^{\text{rest}}}: \tilde{Z}^{\text{rest}} \rightarrow Z^{\text{rest}}$  is an unbranched covering of  $Z^{\text{rest}}$ . Since  $Z^{\text{rest}}$  is a 4-dimensional neighborhood of a graph,  $\tilde{Z}^{\text{rest}}$  is also a 4-dimensional neighborhood of a graph. It follows that  $\tilde{Z}^{\text{rest}}$ , hence  $\tilde{Z}_\lambda$  is a 4-dimensional 1-handlebody, or a disjoint union thereof.

To complete the proof, we must argue that  $\tilde{\Sigma}$ ,  $\tilde{H}_\lambda$ , and  $\tilde{Z}_\lambda$  are each connected. However, we have shown that  $\tilde{\Sigma}$  is a disjoint union of closed surfaces; that  $\tilde{H}_\lambda$  is a 3-dimensional handlebody with  $\partial\tilde{H}_\lambda = \tilde{\Sigma}$ ; and that  $\tilde{Z}_\lambda$  is a 4-dimensional 1-handlebody with  $\partial\tilde{Z}_\lambda = \tilde{H}_\lambda \cup_{\tilde{\Sigma}} \tilde{H}_{\lambda+1}$ . (That the boundaries are as claimed follows from the fact that the restrictions  $f|_{\tilde{\Sigma}}$ ,  $f|_{\tilde{H}_\lambda}$ , and  $f|_{\tilde{Z}_\lambda}$  are all proper maps.) Therefore,  $\tilde{\Sigma}$ ,  $\tilde{H}_\lambda$ ,  $\tilde{Z}_\lambda$ , and  $\tilde{X}$  must have the same number of connected components, as desired.  $\square$

In light of this proposition, we call the trisection of  $\tilde{X}$  obtained by pulling back  $\mathbb{T}$  via  $f$  the *pullback trisection* and we denote it by  $f^*(\mathbb{T})$ .

**Remark 4.2.** We note that the trisection parameters  $\tilde{g}$  and  $\tilde{\mathbf{k}} = (\tilde{k}_1, \tilde{k}_2, \tilde{k}_3)$  corresponding to  $f^*(\mathbb{T})$  can be calculated in practice. The core  $\tilde{\Sigma}$  is a branched covering of the core  $\Sigma$ , so  $g(\tilde{\Sigma})$  can be related to  $g(\Sigma)$  via the Riemann-Hurwitz Formula by considering the local degrees of the covering  $f$ . Since  $\partial\tilde{Z}_\lambda$  is a cover of  $\partial Z_\lambda$ , branched along the link  $\partial\mathcal{D}_\lambda$ , the parameter  $\tilde{k}_\lambda$  is related to the parameter  $k_\lambda$  by understanding each of the coverings of a disjoint union of copies of  $S^3$  over  $S^3$ , branched along the split components of this link. In the case that  $f$  is a symplectic singular branched covering, these components are Hopf links and trefoils, so this computation can be done by knowing the number of each and the local degrees.

## 5. PUTTING SYMPLECTIC BRAIDED SURFACES IN BRIDGE POSITION

A refinement of Theorem 3.7 was proven by Auroux and Katzarkov [AK00], which ensures that we find a symplectic singular branched covering  $f: X \rightarrow \mathbb{C}\mathbb{P}^2$  with the additional property that the image of the branch locus sits in a nice position with respect to the standard linear pencil on  $\mathbb{C}\mathbb{P}^2$ . In this

section we will show that we can symplectically isotope such a surface so that it is in bridge position with respect to a genus one Weinstein trisection on  $\mathbb{C}\mathbb{P}^2$ .

The standard linear pencil on  $\mathbb{C}\mathbb{P}^2$  we will work with is the map  $\pi: \mathbb{C}\mathbb{P}^2 \setminus \{[0:0:1]\} \rightarrow \mathbb{C}\mathbb{P}^1$  given by  $\pi([z_0:z_1:z_2]) = [z_0:z_1]$ . We will often work in the affine chart where  $z_0 \neq 0$  with coordinates  $(z_1, z_2)$ . The restriction of  $\pi$  to this chart is the map  $\pi: \mathbb{C}^2 \rightarrow \mathbb{C}$  where  $\pi(z_1, z_2) = z_1$ .

**Definition 5.1** ([AK00]). A *braided surface* in  $\mathbb{C}\mathbb{P}^2$  is a singular surface  $Q = g(S)$  for  $g: S \rightarrow \mathbb{C}\mathbb{P}^2$  such that

- (1)  $[0:0:1] \notin Q$ ;
- (2)  $Q$  is positively transverse to the fibers of the pencil  $\pi$ , except at finitely many points where it becomes non-degenerately positively tangent to the fibers (locally modeled on  $z_2^2 = z_1$  where  $\pi$  is the projection to the  $z_1$  coordinate);
- (3) the only singularities of  $Q$  are (simple) cusps and transverse double points which may be either positive or negative self-intersections;
- (4) the cusps are positively transverse to the fibers of  $\pi$ ; and
- (5) the “special points” (cusps, double points, and tangency points) all lie in distinct fibers of  $\pi$ .

A *symplectic braided surface* is a braided surface which is a symplectic singular surface. We will say that two (symplectic) braided surfaces are *isotopic through (symplectic) braided surfaces* if there is a one parameter family of (symplectic) braided surfaces connecting them.

**Remark 5.2.** Auroux and Katzarkov call approximately holomorphic braided surfaces “quasiholomorphic curves.”

**Theorem 5.3** ([AK00]). *Every closed symplectic 4-manifold admits a symplectic singular branched covering, such that the image of the branch locus in  $\mathbb{C}\mathbb{P}^2$  is a symplectic braided surface.*

**Remark 5.4.** This theorem goes beyond Auroux’s original Theorem 3.7 because when the branch locus in  $\mathbb{C}\mathbb{P}^2$  has negative nodes, it is not automatic that the surface can be assumed to be braided with respect to a linear pencil. However it is worth noting that any smooth symplectic surface, or symplectic surface whose singularities are symplectically modeled on complex curve singularities (which we will call an *algebraically singular symplectic surface*), can be assumed to be braided with respect to a linear pencil on  $\mathbb{C}\mathbb{P}^2$ . More specifically, any smooth (or algebraically singular) symplectic surface  $Q$  can be realized as a  $J$ -holomorphic curve for some almost complex structure  $J$  which is compatible with the standard symplectic form  $\omega_{FS}$  on  $\mathbb{C}\mathbb{P}^2$ . The space  $\mathcal{J}^{\omega_{FS}}(Q)$  of such  $J$  which make  $Q$   $J$ -holomorphic is contractible and in particular non-empty (see [GS19, Lemma 3.4] for the singular case). Moreover for any such  $J \in \mathcal{J}^{\omega_{FS}}(Q)$ , we obtain a pencil of  $J$ -holomorphic lines through the point  $[0:0:1]$ , and (after a generic perturbation of  $Q$ )  $Q$  will be braided with respect to this pencil. Moreover, there is a symplectomorphism taking this  $J$ -holomorphic pencil to the standard complex linear pencil. Therefore, up to a symplectomorphism of  $\mathbb{C}\mathbb{P}^2$  (which is always symplectically isotopic to the identity), every symplectic surface which is smooth or algebraically singular is braided.

Now we will show that we can isotope a symplectic braided surface into bridge position through symplectic braided surfaces. To keep the braided condition, we will need to keep track of the interaction of  $Q$  with the fibration  $\pi: \mathbb{C}\mathbb{P}^2 \setminus \{[0:0:1]\} \rightarrow \mathbb{C}\mathbb{P}^1$ .

If  $\phi: \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{R}_{>0}$  is any smooth function, we obtain a fiber preserving diffeomorphism of  $\mathbb{C}\mathbb{P}^2$  by the map  $\Phi: \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{C}\mathbb{P}^2$ ,  $\Phi([z_0:z_1:z_2]) = [z_0:z_1:\phi([z_0:z_1])z_2]$ . If  $Q$  is braided, then  $\Phi(Q)$  is still braided. However, if  $Q$  is symplectic, in general it is possible that  $\Phi(Q)$  may no longer be symplectic. Note that when  $\phi \equiv 1$ ,  $\Phi$  is the identity. The case when  $\phi$  is constant is particularly easy to understand.

**Lemma 5.5.** *Let  $\Phi_c([z_0 : z_1 : z_2]) = [z_0 : z_1 : cz_2]$ .*

- (1) *If  $Q = g(S)$  with  $g: S \rightarrow \mathbb{CP}^2 \setminus \{[0 : 0 : 1]\}$  is a symplectic braided surface, and  $c \leq 1$ , then  $\Phi_c(Q)$  is also a symplectic braided with respect to  $\omega_{FS}$ .*
- (2) *If  $g: S' \rightarrow \mathbb{CP}^2 \setminus \{[0 : 0 : 1]\}$  is positively transverse to the fibers of  $\pi$  and  $S'$  is compact then there exists a  $c$  sufficiently small such that  $\Phi_c \circ g(S')$  is symplectic (with respect to  $\omega_{FS}$ ).*

*Proof.* The Fubini-Study symplectic form in affine coordinates  $(z_1, z_2)$  is given by  $\omega_{FS} = -\frac{1}{4}dd^c(\log(1 + |z_1|^2 + |z_2|^2))$ . Letting  $z_j = x_j + iy_j$ , and expanding out this 2-form, we can calculate that

$$\Phi_c^*(\omega_{FS}) = (1 - c^2)dx_1 \wedge dy_1 + c^2\omega_{FS}.$$

Therefore

$$(\Phi_c \circ g)^*(\omega_{FS}) = g^*(\Phi_c^*\omega_{FS}) = g^*((1 - c^2)dx_1 \wedge dy_1 + c^2\omega_{FS}) = (1 - c^2)g^*(dx_1 \wedge dy_1) + c^2g^*\omega_{FS}.$$

If  $g$  is braided,  $g^*(dx_1 \wedge dy_1) \geq 0$ . (This is a rephrasing of the condition that at each point  $g$  is either positively tangent to the fibers of  $\pi$  or  $d(\pi \circ g) = 0$ .)

Part (1) then follows at points where  $g$  is an immersion because  $g^*(dx_1 \wedge dy_1) \geq 0$  and  $g^*\omega_{FS} > 0$  implies  $(\Phi_c \circ g)^*(\omega_{FS}) > 0$ . At cusp singularities where  $g^*\omega_{FS} = 0$ , this argument does not apply. However, if  $p \in S$  is a cusp point of  $Q$ , the limiting tangent space to  $\Phi_c(Q)$  at  $\Phi_c(g(p))$  is  $d\Phi_c(T)$  where  $T$  is the limiting tangent space to  $Q$  at  $g(p)$ . Then

$$\omega_{FS}|_{d\Phi_c(T)} = \Phi_c^*(\omega_{FS})|_T = ((1 - c^2)dx_1 \wedge dy_1 + c^2\omega_{FS})|_T$$

This is positive because  $(dx_1 \wedge dy_1)|_T > 0$  (since the cusp is positively transverse to the fibers of  $\pi$ ) and  $\omega_{FS}|_T > 0$  (since  $Q$  was symplectically braided).

The hypothesis in part (2) that  $g$  is positively transverse to the fibers of  $\pi$  is equivalent to saying that  $g^*(dx_1 \wedge dy_1) > 0$ . Let  $A$  denote a fixed area form on  $S'$ . By compactness and the fact that  $dx_1 \wedge dy_1 > 0$  on the tangent spaces to  $g(S')$ , we can choose  $\varepsilon > 0$  and  $K > 0$  such that  $g^*(dx_1 \wedge dy_1) \geq \varepsilon A$  and  $g^*(\omega_{FS}) \geq -KA$ . Choosing  $c$  small enough so that  $c^2 < \frac{\varepsilon}{\varepsilon + K}$  implies that

$$(\Phi_c \circ g)^*(\omega_{FS}) \geq ((1 - c^2)\varepsilon - Kc^2)A > 0$$

so  $\Phi_c \circ g(S')$  is symplectic.  $\square$

In addition to the isotopies above which fix each fiber of  $\pi$ , we will need an isotopy which sends fibers to fibers but moves the critical values of the projection of the surface into a controlled region with respect to the trisection.

**Lemma 5.6.** *If  $Q$  is a symplectic braided surface, then it is isotopic through symplectic braided surfaces to a symplectic braided surface such that every critical value of  $\pi \circ F$  lies on the equator of  $\mathbb{CP}^1$ ,  $|z_1| = |z_0|$ .*

*Proof.* Let  $c_1, \dots, c_n \in \mathbb{CP}^1$  be the critical values of the projection  $\pi|_Q: Q \rightarrow \mathbb{CP}^1$ . Choose small disjoint neighborhoods  $U_j$  of  $c_j$ . Let  $\psi_t: \mathbb{CP}^1 \rightarrow \mathbb{CP}^1$  be an isotopy such that  $\psi_t|_{U_j}$  is a projective unitary transformation and  $\psi_1(c_j)$  lies on the equator of  $\mathbb{CP}^1$ . Let

$$\Psi_t: \mathbb{CP}^2 \setminus \{[0 : 0 : 1]\} \rightarrow \mathbb{CP}^2 \setminus \{[0 : 0 : 1]\}$$

be defined by  $\Psi_t([z_0 : z_1 : z_2]) = [\psi_t([z_0 : z_1]) : z_2]$ . Then  $\pi \circ \Psi_t = \psi_t \circ \pi$ . Therefore the critical values of the restriction of  $\pi$  to  $\Psi_1(Q)$  all lie on the equator of  $\mathbb{CP}^1$ . Also,  $\Psi_t(Q)$  is a braided surface for all  $t$ . Finally,  $\Psi_t$  is a symplectomorphism on the neighborhoods  $\pi^{-1}(U_j)$  of the critical fibers because it is a unitary transformation there.

It is possible that  $\Psi_t(Q)$  will fail to be symplectic for some  $t > 0$ , but we do know that  $\Psi_t(\pi^{-1}(U_j) \cap Q)$  will be symplectic for each  $j$ . Let  $Q_j = Q \cap \pi^{-1}(U_j)$  and let  $Q' = Q \setminus (\cup_j Q_j)$ . Since  $\Psi_t$  is fiber preserving,  $\Psi_t(Q')$  is a compact surface which is everywhere transverse to the fibers of  $\pi$ . Then by Lemma 5.5(2), there exists a sufficiently small value of  $c_t > 0$  such that  $\Phi_{c_t}(\Psi_t(Q'))$  is symplectic. Since  $[0, 1]$  is compact, we can choose a uniform  $0 < c \leq 1$  such that  $\Phi_c(\Psi_t(Q'))$  is symplectic for all  $t \in [0, 1]$ . By Lemma 5.5(1),  $\Phi_c(\Psi_t(Q_j))$  is also symplectic because  $\Psi_t(Q_j)$  is symplectic. Therefore  $\Phi_c \circ \Psi_t$  is a symplectic isotopy from  $\Phi_c(Q)$  to  $\Phi_c \circ \Psi_1(Q)$ . Additionally,  $\Phi_{1+t(c-1)}(Q)$  is a family of symplectic braided surfaces for  $t \in [0, 1]$ , interpolating between  $Q$  and  $\Phi_c(Q)$ . Concatenating these yields the required isotopy from  $Q$  to  $\Phi_c(\Psi_1(Q))$ .  $\square$

Recall the genus one Weinstein trisection  $\mathbb{T}_0$  for  $\mathbb{C}\mathbb{P}^2$  described in Section 1 and shown schematically as the moment map image of the standard toric action in Figure 1. In affine coordinates  $(z_1, z_2)$ , the preimage of equator of  $\mathbb{C}\mathbb{P}^1$  is  $\{|z_1| = 1\}$ . Thus, the handlebody  $H_2 = \{|z_1| = 1, |z_2| \leq 1\}$  is contained in the preimage of the equator.

**Lemma 5.7.** *Let  $Q = g(S)$  for  $g: S \rightarrow \mathbb{C}\mathbb{P}^2$  be a symplectic braided surface, such that the critical values of  $\pi \circ g$  lie on the equator of  $\mathbb{C}\mathbb{P}^1$ . Then there exists a map  $\phi: \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{R}_{>0}$ , which is constant on neighborhoods  $\{V_j\}$  of the critical values of  $\pi \circ g$ , such that the induced fiber preserving diffeomorphism  $\Phi$  has  $\Phi(Q)$  in singular bridge position with respect to the standard trisection on  $\mathbb{C}\mathbb{P}^2$ .*

*Proof.* By first applying  $\Phi_c$  induced by a small constant  $\phi = c$ , we can rescale  $Q$  so its moment image projects towards the bottom of the triangle of Figure 1 and thus we may assume that  $Q \cap Z_3 = \emptyset$ . If all the critical points of  $Q$  lie above the equator of  $\mathbb{C}\mathbb{P}^1$ , they will all be contained in  $H_2$ . Away from the critical points,  $Q$  intersects  $H_2$  transversally (since  $H_2$  is contained in the union of a 1-parameter family of fibers of  $\pi$ ). Therefore, away from the singular fibers  $Q \cap H_2$  is a braid, transverse to the meridional disks.

More generally, we can look at the braid in the nearby open solid tori  $V_r := \{|z_1| = r\}$  for  $r \in (1 - \varepsilon, 1 + \varepsilon)$ . Note that in the adjacent open solid tori,  $V_r$  for  $r \neq 1$ , the intersection  $Q \cap V_r$  is a smooth braid, by transversality. For each type of singularity, there is a different local movie of braids (where  $r$  is the time direction of the movie) that becomes singular in  $H_2 \subset V_1$ .

At a tangency, a positive half-twist (single crossing) is added to the braid after passing through the tangency point – Figure 3. At a positive node, a negative crossing switches to a positive crossing (so overall one full positive twist is added between the two strands) – Figure 4. A negative node switches a positive crossing to a negative one (adding one full negative twist between two strands). At a cusp, three positive half-twists are added to the braid after passing through the cusp – Figure 5. These all follow from calculating the braid monodromy directly from the complex local models for these three singularity types.




FIGURE 3. The movie showing a braided surface near a tangency.




FIGURE 4. The movie showing a braided surface near a node.




FIGURE 5. The movie showing a braided surface near a cusp.

Note that for  $r < 1$ , the intersection of  $V_r$  with  $H_1$  is a torus with fixed  $z_2$  radial coordinate  $|z_2| = 1$ . Similarly, for  $r > 1$ , the intersection of  $V_r$  with  $H_3$  is the torus where  $|z_2| = |z_1| = r$ . Initially because  $Q \cap Z_3 = \emptyset$ ,  $Q$  does not intersect  $H_1$  or  $H_3$ . This surface is certainly not in bridge position, because the intersection of  $Q$  with  $H_2$  is generally singular, non-trivially braided, and  $Q$  does not intersect the core surface  $\Sigma$ .

We will perform a fiber preserving isotopy  $\Phi$  to push the neighborhoods of the singular points into the interior of  $Z_3$ . We will also push some smoothly braided parts of  $Q \cap H_2$  into the interior of  $Z_3$  to address the non-trivial braiding of  $Q \cap H_2$ , so that in the end  $Q \cap H_2$  will become a trivial tangle.

The fiber preserving isotopy allows us to modify the  $z_2$  radial coordinate. In Figure 6, this direction is depicted as vertical, while the time coordinate  $r$  varies as you move from one box to another horizontally. The isotopy  $\Phi$  of  $Q$  will be chosen with the goal of making the resulting  $Q \cap H_2$  a trivial tangle, but we will also need to keep track of how these isotopies modify  $Q \cap H_1$  and  $Q \cap H_3$  to verify that these will also be trivial tangles.

The isotoped version of  $Q$  in the case of the tangency is depicted as a movie in Figure 6. Replacing the braid movie in Figure 6 with the braid movies for the node and cusp models would give the corresponding movie for the local isotopy near each of these singularities. Additionally, for every crossing in the braid, we will push a neighborhood of that crossing into  $Z_3$ , plugging in the trivial movie on that crossing into the movie of Figure 6. To put this all together, we concatenate these models according to the singular braid we started with in  $H_2$ . Note that the models are inserted into disjoint regions of the torus.

In order to see the intersection of  $Q$  with  $H_1$  and  $H_3$ , observe that, in the movie,  $H_1$  is sliced into concentric tori and  $Q$  intersects the torus in each slice in a finite collection of points. Dropping the radial  $z_2$  coordinate, we draw  $H_1$  and  $H_3$ , tracking the movie of points as a tangle for each model in Figure 7. In the movie of points in the tori  $H_1 \cap V_r$  for  $r \leq 1$  or  $H_3 \cap V_r$  for  $r \geq 1$ , pairs of points join together and disappear to form arcs of the tangle. The track of the points in this movie is the projection of these arcs to the torus. Because arcs of the same color can be projected up to isotopy as






FIGURE 6. A fiber preserving isotopy of a neighborhood of a tangency point of  $Q$  with the fibers of the pencil  $\pi$ . The top row shows the surface at the beginning of the isotopy, and the bottom row shows the surface at the end of the isotopy.




FIGURE 7. (Top) The pieces of the tangles in  $\overline{H}_1$  and  $H_3$  after isotoping  $Q$  near a tangency. (Note that  $\partial Z_3 = \overline{H}_1 \cup_{\Sigma} H_3$ .) Gluing these pieces together gives an unknot. (Bottom, left to right) The projections of the corresponding tangles in  $\Sigma$  for crossings, tangencies, nodes, and cusps, respectively. Note that in this figure  $\Sigma$  is the oriented boundary of  $H_3$  and the oppositely oriented boundary of  $\overline{H}_1$ .

embeddings, the tangles  $Q \cap H_1$  and  $Q \cap H_3$  are trivial. Note that for the total surface there will be many disjoint copies of these models inserted in disjoint regions of the torus.

**Warning:** When drawing pictures and projections, one needs to be very careful to keep track of which orientation one is using on each handlebody. *Our convention is that  $H_\lambda$  is positively oriented as part of the boundary of  $Z_\lambda$  and that  $\Sigma$  is the positively oriented boundary of  $H_\lambda$ .* When we look at a braid in  $V_r$ , we think of  $V_r$  as a parallel copy of  $\overline{H}_2$ , with boundary  $\overline{\Sigma}$ . As we pass singularities, the braid obtains positive twists (when viewed in  $\overline{H}_2$  which is the orientation it inherits as the boundary of  $Z_1$ ). In particular, the natural projection of the positive braid is to  $\overline{\Sigma}$ . Pushing parts of this braid into  $Z_3$  gradually, we trace out the intersection of  $Q$  with  $H_3$ . We can then project this trace onto  $\overline{\Sigma}$  and reflect to obtain the projection to  $\Sigma$ . See Figure 8 for the case of the cusp.

Note that when we have the projections of the arcs from  $H_1$ ,  $H_2$ , and  $H_3$  all on  $\Sigma$ , we can think of one set of arcs (say the blue arcs of  $H_2$ ) as being isotoped to lie exactly on  $\Sigma$ . Pushing the red arcs slightly up and the green arcs slightly down is consistent with viewing all three sets of arcs inside a




FIGURE 8. (Top row) Pushing the neighborhood of the cusp into  $Z_3$ . (Middle rows) Tracing out the intersection with  $H_3$ . (Bottom row) Projecting the tangle to  $\overline{\Sigma}$  and its reflection  $\Sigma$ .

neighborhood of  $\Sigma$  in  $H_3 \cup_{\Sigma} \overline{H}_1$ . In particular, the knot formed by the union of the red and green arcs in the bottom-right frame of Figure 8 gives a positive trefoil with this convention. Since  $H_3 \cup_{\Sigma} \overline{H}_1$  is the oriented boundary of  $Z_3$ , we see that this positive trefoil is the link of the cusp singularity in  $\partial Z_3$  as expected.  $\square$

**Lemma 5.8.** *If  $\mathbb{T}_0$  is the standard trisection of  $\mathbb{C}\mathbb{P}^2$ , then its image under the rescaling,  $\Phi_c(\mathbb{T}_0)$  is a genus one Weinstein trisection.*

*Proof.* The three sectors of the standard trisection are given in homogeneous coordinates by  $[z_0 : z_1 : z_2]$  by

$$Z_{\lambda} = \{|z_{\lambda}|, |z_{\lambda+1}| \leq |z_{\lambda+2}|\},$$

for  $i = 0, 1, 2$  (where the indices are considered mod 3). The effect of the rescaling map  $\Phi_c$ , in terms of homogeneous coordinates on the target  $[w_0 : w_1 : w_2]$  is  $w_0 = z_0$ ,  $w_1 = z_1$ ,  $w_2 = cz_2$ . Therefore the image of the trisection under the rescaling is given by the three sectors

$$Z'_1 = \{|w_1| \leq |w_0|, |w_2| \leq c|w_0|\}, \quad Z'_2 = \{|w_0| \leq |w_1|, |w_2| \leq c|w_1|\}, \quad \text{and} \quad Z'_3 = \{|w_0|, |w_1| \leq \frac{1}{c}|w_2|\}$$

Note that the Fubini-Study form on any affine chart is explicitly exact:  $\omega_{FS} = d(-d^{\mathbb{C}}(\log(1+|w|^2)))$ . The Liouville vector field  $V$  dual to the primitive  $\eta = -d^{\mathbb{C}}(\log(1+|w|^2))$  has the property that it is obtained from the radial vector field  $V_0$  by rescaling by a positive function,  $V = hV_0$ . We can see this as follows. Let  $f_0(w) = |w|^2$  and let  $\eta_0 = -d^{\mathbb{C}}f_0$ . Then  $\eta_0$  is the standard Liouville form for the Darboux symplectic form  $\omega_0 = d\eta_0$  and its vector field dual is the radial vector field  $V_0 = x_1\partial_{x_1} + y_1\partial_{y_1} + x_2\partial_{x_2} + y_2\partial_{y_2}$  (with  $\iota_{V_0}d\eta_0 = \eta_0$ ). Now, the chain rule shows that  $\eta = g\eta_0$  where  $g(w) = (1+|w|^2)^{-1}$  and  $\omega_{FS} = d\eta = dg \wedge \eta_0 + g d\eta_0$ . Then if  $V = hV_0$  is a multiple of the radial vector field by a function we have

$$\iota_V \omega_{FS} = h \iota_{V_0} (dg \wedge \eta_0 + g d\eta_0) = h(dg(V_0)\eta_0 + g\eta_0).$$

Therefore by choosing  $h$  appropriately so that  $h(dg(V_0) + g) = g$ , we get that  $V = hV_0$  is the vector field dual to the Liouville form  $\eta$ .

Each of our three rescaled sectors lies in one of the three affine charts, so it suffices to show that the corresponding Liouville vector field  $V$  (or equivalently the radial vector field  $V_0$ ) is outwardly transverse to the boundary of the sector (after smoothing the corner).

In the affine chart where  $w_0 = 1$ , we have the sector  $\{|w_1| \leq 1, |w_2| \leq c\}$  with boundary

$$\{|w_1|^2 = 1, |w_2| \leq c\} \cup \{|w_1| \leq 1, |w_2|^2 = c^2\}.$$

The radial vector field  $V_0$  is transverse to the first piece because  $d(|w_1|^2)(V_0) = |w_1|^2 = 1 > 0$  and on the second piece because  $d(|w_2|^2)(V_0) = |w_2|^2 = c > 0$ . We can find an approximation smoothing the corner by looking at the hypersurface

$$Y = \left\{ \frac{1}{N} \left( |w_1|^2 + \frac{1}{c^2} |w_2|^2 \right) + |w_1|^{2N} + \frac{1}{c^{2N}} |w_2|^{2N} = 1 \right\}$$

The vector field  $V_0$  is positively transverse to  $Y$  because

$$\begin{aligned} & d \left( \frac{1}{N} \left( |w_1|^2 + \frac{1}{c^2} |w_2|^2 \right) + |w_1|^{2N} + \frac{1}{c^{2N}} |w_2|^{2N} \right) (V_0) \\ &= \frac{1}{N} d(|w_1|^2)(V_0) + \frac{1}{Nc^2} d(|w_2|^2)(V_0) + N|w_1|^{2N-2} d(|w_1|^2)(V_0) + \frac{N}{c^{2N}} |w_2|^{2N-2} d(|w_2|^2)(V_0) \end{aligned}$$

which is a positive combination of positive quantities (when  $w_1 = 0$  or  $w_2 = 0$  some of the quantities are 0 and others are positive, otherwise all terms in the sum are positive).

In the other affine charts, the computations are similar to show that the radial vector field  $V_0$ , and thus the Liouville vector field  $V$ , is positively transverse to the (smoothed) boundary of the sectors.  $\square$

Now we have all the tools to deduce the main result of this section.

**Theorem 5.9.** *If  $g: S \rightarrow \mathbb{C}\mathbb{P}^2$  is a symplectic braided surface in  $\mathbb{C}\mathbb{P}^2$ , then there is an ambient symplectic isotopy  $H_t: \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{C}\mathbb{P}^2$  such that  $H_1(g(S))$  is in bridge position with respect to a genus one Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$ .*

*Proof.* By Lemma 5.6, we get a symplectic isotopy  $\Psi_t: \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{C}\mathbb{P}^2$  such that  $\Psi_0 = id$  and the critical values of  $\pi \circ \Psi_1 \circ g$  are points  $r_1, \dots, r_n$  on the equator of  $\mathbb{C}\mathbb{P}^1$ .

Then by Lemma 5.7, there exists a function  $\phi: \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{R}_{>0}$  such that

- $\phi$  is constant on open neighborhoods  $V_j$  of  $r_j$  for  $j = 1, \dots, n$ , and
- for the map  $\Phi$  induced by  $\phi$ ,  $\Phi(Q)$  is in bridge position with respect to the standard trisection on  $\mathbb{C}\mathbb{P}^2$ .

Let

$$C = \sup_{[z_0:z_1] \in \mathbb{C}\mathbb{P}^1} (\phi([z_0:z_1])).$$

Then  $\Phi_{1/C} \circ \Phi(Q)$  is in bridge position with respect to  $\Phi_{1/C}(\mathbb{T}_0)$ . Let  $\phi_t = \frac{t}{C}\phi + 1 - t$ . Then for  $t \in [0, 1]$ ,  $\phi_t: \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{R}_{>0}$  induces a fiber preserving map  $\Phi_t: \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{C}\mathbb{P}^2$ . Note that  $\phi_t$  is constant and less than or equal to 1 in  $V_j$ , for  $j = 1, \dots, n$ . Since  $\Psi_1(g(S))$  is symplectic, Lemma 5.5(1) implies that  $\Phi_t(\Psi_1(g(S))) \cap \pi^{-1}(V_j)$  is symplectic for all  $t \in [0, 1]$  and  $j = 1, \dots, n$ .

Outside of  $\pi^{-1}(V_j)$  it is possible that  $\Phi_t(\Psi_1(g(S)))$  fails to be symplectic at some time  $t$ . Let  $S' = S \setminus (\pi \circ \Psi_1 \circ g)^{-1}(V_j)$ . Note that  $\pi = \pi \circ \Phi_t$ . Because the critical values of  $\pi \circ \Phi_t \circ \Psi_1 \circ g$  are contained in the neighborhoods  $V_j$ , the map  $\Phi_t \circ \Psi_1 \circ g: S' \rightarrow \mathbb{C}\mathbb{P}^2$  is positively transverse to the fibers of  $\pi$  and  $S'$  is compact. Therefore by Lemma 5.5(2), there exists a sufficiently small  $0 < c_t \leq 1$  such that the induced constant rescaling map  $\Phi_{c_t}$  has  $\Phi_{c_t}(\Phi_t(\Psi_1(g(S'))))$  is symplectic. Since  $[0, 1]$  is compact, we can choose  $c = \inf_{t \in [0, 1]} \{c_t\} > 0$  such that  $\Phi_c(\Phi_t(\Psi_1(g(S'))))$  is symplectic for all  $t \in [0, 1]$ . Then  $\Phi_c(\Phi_t(\Psi_1(g(S))))$  is symplectic for all  $t \in [0, 1]$  because it is symplectic outside  $\pi^{-1}(V_j)$  by the choice of  $c$  and it is symplectic inside  $\pi^{-1}(V_j)$  because  $\Phi_t(\Psi_1(g(S)))$  was symplectic and  $\Phi_c$  preserves symplecticness here by Lemma 5.5(1).

Using all this information, we can construct the isotopy through symplectic braided surfaces from  $g: S \rightarrow \mathbb{C}\mathbb{P}^2$  to a symplectic braided surface in bridge position with respect to a genus one Weinstein trisection on  $\mathbb{C}\mathbb{P}^2$  as the concatenation of the following isotopies. First, we isotope  $g(S)$  to  $\Psi_1(g(S))$  using  $\Psi_t$  from Lemma 5.6. Then we isotope  $\Psi_1(g(S))$  to  $\Phi_c(\Psi_1(g(S)))$  through the isotopy of constant rescaling maps  $\Phi_{1-t(1-c)}$ ; this is an isotopy through symplectic braided surfaces by Lemma 5.5(1). Finally, we concatenate with the isotopy  $\Phi_c \circ \Phi_t \circ \Psi_1 \circ g: S \rightarrow \mathbb{C}\mathbb{P}^2$  described above. Because  $\Phi_1$  has the property that  $\Phi_1 \circ \Psi_1 \circ g(S)$  is in bridge position with respect to the trisection  $\Phi_{1/C}(\mathbb{T}_0)$  on  $\mathbb{C}\mathbb{P}^2$ , we have that  $\Phi_c \circ \Phi_1 \circ \Psi_1 \circ g(S)$  is in bridge position with respect to the image  $\Phi_{c/C}(\mathbb{T}_0)$  of the standard trisection under  $\Phi_{c/C}$ . This is a genus one Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$  by Lemma 5.8.

By Lemma 3.2, there is an ambient symplectic isotopy  $H_t: \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{C}\mathbb{P}^2$  such that  $H_t(g(S))$  agrees with this family of symplectic braided surfaces, outside of an arbitrarily small neighborhood of the singular points. In the neighborhood of the singular points,  $H_t(g(S))$  will have the same topological type of the singularity (node/cusp) but the analytic type may slightly differ. By choosing the neighborhoods of the singularities sufficiently small, we can assume that these neighborhoods are contained on the interior of  $Z'_3 := \Phi_c(Z_3)$  and therefore  $H_1(g(S))$  is also in bridge position.

□

## 6. LIFTING THE WEINSTEIN STRUCTURE

Given a singular symplectic branched covering  $f: (X, \omega) \rightarrow (\mathbb{C}\mathbb{P}^2, \omega_{FS})$ , Auroux constructs a symplectic structure  $\omega_f$  on  $X$  which is an arbitrarily small perturbation of  $f^*\omega_{FS}$  [Aur00, Proposition 10]. Note that because  $f$  has critical points,  $f^*\omega_{FS}$  cannot be non-degenerate along the critical locus. To remedy this,  $\omega_f$  is built by adding a small multiple of an exact form which is supported near the critical locus of  $f$ . Although  $\omega_f$  is not identical to our original symplectic form  $\omega$ , we will see in Lemma 6.2 that  $(X, k\omega)$  and  $(X, \omega_f)$  are symplectomorphic and this suffices to show that a Weinstein trisection of  $(X, \omega_f)$  induces a Weinstein trisection of  $(X, \omega)$ .

**Theorem 6.1.** *Suppose that  $Q \subset \mathbb{C}\mathbb{P}^2$  is a symplectic singular surface in bridge position with respect to a Weinstein trisection  $\mathbb{T}_0$  of  $\mathbb{C}\mathbb{P}^2$ . Let  $f: X \rightarrow \mathbb{C}\mathbb{P}^2$  be a singular branched covering with the same local models as a symplectic singular branched covering (local diffeomorphism, cyclic branching, and the cusp). Then  $f^*\mathbb{T}_0$  is a Weinstein trisection for  $(X, \omega_f)$ .*

*Proof.* We know from Theorem 4.1 that  $f^*\mathbb{T}_0$  is smoothly a trisection. We will verify that each sector with its restricted symplectic form is a Liouville domain filling  $\#^k(S^1 \times S^2)$ . In general a Weinstein structure is stronger than a Liouville structure: a Weinstein structure requires that the Liouville vector field is gradient-like for some Morse-function. However when the boundary is  $\#^k(S^1 \times S^2)$ , the contact structure is planar, so for any Liouville filling, after possibly adding a trivial collar to the boundary, the symplectic structure is compatible with a Weinstein structure by [Wen10] (relevant arguments on holomorphic disk filling of  $\#^k(S^1 \times S^2)$  can also be found in [Eli90, CE12]). Because the skeleton will lie on the interior of the domain (since the Liouville vector field is transverse to the boundary), the addition of the trivial collar does not make a difference for our purposes. Since there is a unique Weinstein structure on this filling by [CE12], we can deform the Liouville vector field through Liouville vector fields for the fixed symplectic structure which are all positively transverse to the boundary until it agrees with the standard Weinstein structure.

Branched coverings have been studied in both symplectic and contact topology. The elemental question of how to lift the symplectic or contact structures from the base to the cover is answered in [Aur00, Proposition 10] and [Gei08, Theorem 7.5.4] in the symplectic and contact cases respectively. In our case, we want to lift the symplectic filling structure of  $(S^3, \xi_{std})$  on  $(Z_\lambda, \omega_{FS}|_{Z_\lambda})$  to a symplectic filling structure on  $(\tilde{Z}_\lambda, \omega_f)$  of  $\#^{k_\lambda}(S^1 \times S^2)$  with its unique tight contact structure. The key point we will see is that the constructions of symplectic and contact structures on the branch cover are compatible with each other.

Because the branch locus in  $\mathbb{C}\mathbb{P}^2$  is in bridge trisected position, its intersection with  $\partial Z_\lambda$  is a smooth link. The singular points of the branch locus are away from  $\partial Z_\lambda$ . Let  $i: \partial Z_\lambda \rightarrow Z_\lambda$  be the inclusion. Let  $\omega_\lambda := \omega_{FS}|_{Z_\lambda}$ , and let  $\eta_\lambda$  be the Liouville form ( $\eta_\lambda = \omega_\lambda(V_\lambda, \cdot)$ ) which satisfies  $d\eta_\lambda = \omega_\lambda$ . The transversality of the Liouville vector field implies that  $\alpha_\lambda := i^*\eta_\lambda$  is a contact form on  $\partial Z_\lambda$  (see the end of this proof for how to understand this equivalence if you are not already familiar). Because the branch locus is symplectic, this link  $Q \cap \partial Z_\lambda$  is a transverse link with respect to the contact planes  $\xi_\lambda = \ker \alpha_\lambda$ .

The symplectic form  $\omega_f$  on the cover  $X$  is constructed as follows. Initially we start with the form  $f^*\omega_{FS}$ . This is certainly closed since  $d(f^*\omega_{FS}) = f^*d\omega_{FS} = 0$ , but it fails to be non-degenerate near the singular points of  $f$ . At points where  $f$  is a local diffeomorphism,  $f^*\omega_{FS}$  will be non-degenerate, but at singular points where  $f$  is modeled on a cyclic branched covering  $f(z_1, z_2) = (z_1^d, z_2)$  or a cusp  $f(z_1, z_2) = (z_1^3 - z_1 z_2, z_2)$ ,  $f^*\omega_{FS}$  has kernel in the  $\{dz_2 = 0\}$  direction. Therefore, we need to add

something to  $f^*\omega_{FS}$  to achieve non-degeneracy. Choose coordinate charts where the local models for  $f$  hold, such that the coordinate charts containing a cusp model are disjoint from  $\partial\tilde{Z}_\lambda$  for all  $\lambda \in \mathbb{Z}_3$ . The ramification locus  $R$  in each coordinate chart is the set  $\{z_1 = 0\}$  which is locally parameterized by the coordinate  $z_2$ .

Following Auroux [Aur00, Proposition 10], choose a radius  $\rho > 0$  such that the polydisk  $B_{2\rho}(0) \times B_{2\rho^2}(0)$  is contained in the local coordinate patches with cusp models, and the polydisk  $B_{2\rho}(0) \times B_{2\rho}(0)$  is contained in the local coordinate patches in cyclic branch covering models. Let  $\{U_j\}$  be an cover of the ramification locus  $R$  by such polydisks. Let  $\chi_1^j$  be a cut-off function on  $\mathbb{C}$  that is 1 on  $B_\rho(0) \subset \mathbb{C}$  and 0 outside  $B_{2\rho}(0)$ . Let  $\chi_2^j$  be a cut-off function on  $\mathbb{C}$ , is 1 on  $B_{\rho^2}(0) \subset \mathbb{C}$  and 0 outside  $B_{2\rho^2}(0)$ . Choose these cut-offs such that, using the coordinate models in the  $U_j$ , we have that  $\{\chi_{s_j}^j(z_2)\}$  (where  $s_j = 1$  for a coordinate patch with a cyclic branching chart and  $s_j = 2$  for a coordinate patch with a cusp model) is a partition of unity on  $R$  subordinate to  $\{U_j \cap R\}$ .

On each  $U_j$  with coordinates  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$ , define

$$\phi_j = \frac{1}{2}\chi_1^j(z_1)\chi_{s_j}^j(z_2)(x_1dy_1 - y_1dx_1)$$

where  $s_j = 1$  for a coordinate patch with a cyclic branching chart and  $s_j = 2$  for a coordinate patch with a cusp model. (Note that Auroux uses  $x_1dy_1$  instead of  $\frac{1}{2}(x_1dy_1 - y_1dx_1)$  as a primitive for the  $dx_1 \wedge dy_1$ , but this choice does not affect the computation.) Where the cut-off functions are identically 1,  $d\phi_j = dx_1 \wedge dy_1$ . This is positive on the kernel of  $df$ , which makes up for the non-degeneracy of  $f^*\omega_{FS}$ . Finally, observe that because of the cut-off functions, each  $\phi_j$  extends by zero to the entire manifold. Let  $\phi = \sum_j \phi_j$ .

Finally define  $\omega_f := f^*\omega_{FS} + \varepsilon d\phi$ . Then for  $\varepsilon$  sufficiently small, Auroux verifies that  $\omega_f$  is non-degenerate [Aur00, Proposition 10]. Note that Auroux only considers cyclic branched covering models  $(z_1, z_2) \mapsto (z_1^d, z_2)$  when  $d = 2$ , but the computation is unaffected by using higher values of  $d$  because the kernel of  $df$  is the same and  $\phi$  is constructed using the coordinates upstairs.

Now restrict  $\omega_f$  to  $\tilde{Z}_\lambda$ . There,

$$\omega_f = f^*\omega_\lambda + \varepsilon d\phi|_{\tilde{Z}_\lambda} = f^*d\eta_\lambda + \varepsilon d\phi = d(f^*\eta_\lambda + \varepsilon\phi)$$

Let  $i: \partial\tilde{Z}_\lambda \rightarrow \tilde{Z}_\lambda$  be the inclusion. Restricting this primitive to the boundary gives an induced 1-form on  $\partial\tilde{Z}_\lambda$ ,  $i^*(f^*\eta_\lambda + \varepsilon\phi) = f^*\alpha_\lambda + i^*\varepsilon\phi$ , which we can verify is contact. Since the ramification locus intersects  $\partial\tilde{Z}_\lambda$  transversally along a link,  $i^*\varepsilon\phi$  is supported in a neighborhood of this ramification link. The coordinate  $z_1$  in each  $U_j$  provides a coordinate on the normal bundle to this ramification link. The natural contact form on the branched cover constructed in [Gei08, Theorem 7.5.4] is (up to a constant factor)  $f^*\alpha_\lambda + g(r)r^2d\theta$  where  $(r, \theta)$  are polar coordinates on the normal bundle ( $z_1 = re^{i\theta}$ ) and  $g(r)$  is a cut-off function. Noting that  $(x_1dy_1 - y_1dx_1) = r^2d\theta$ , we find that the same computation verifies that  $i^*(f^*\eta_\lambda + \varepsilon\phi)$  is a contact form on  $\partial\tilde{Z}_\lambda$ .

Note that because the primitive  $\eta_f := f^*\eta_\lambda + \varepsilon\phi$  for  $\omega_f|_{\tilde{Z}_\lambda}$  restricts to a positive contact form on the boundary, it gives a Liouville vector field  $V_\lambda$  (defined by  $\omega_f(V_\lambda, \cdot) = \eta_f$ ) which is outwardly transverse to the boundary defined by  $\iota_{V_\lambda}(\omega_f|_{\tilde{Z}_\lambda}) = \eta_f$ . To understand why, let  $i: \partial\tilde{Z}_\lambda \rightarrow \tilde{Z}_\lambda$  be the inclusion and  $\alpha_f = i^*\eta_f$ . The claim is that the property that  $V_\lambda$  be outwardly transverse to the boundary is equivalent to the condition that  $\alpha_f \wedge d\alpha_f$  is a positive volume form. This is because  $\alpha_f \wedge d\alpha_f = i^*\eta_f \wedge i^*\omega_f = i^*(\iota_{V_\lambda}\omega_f \wedge \omega_f)$ . Since  $\omega_f \wedge \omega_f$  is a positive volume form,  $V_\lambda$  is outwardly transverse to  $\partial\tilde{Z}_\lambda$  if and only if there is a positively oriented frame  $(\zeta_1, \zeta_2, \zeta_3)$  for  $T\partial\tilde{Z}_\lambda$  such that

$\omega_f \wedge \omega_f(V_\lambda, \zeta_1, \zeta_2, \zeta_3) > 0$ . But

$$\omega_f \wedge \omega_f(V_\lambda, \zeta_1, \zeta_2, \zeta_3) = \iota_{V_\lambda} \omega_f \wedge \omega_f(\zeta_1, \zeta_2, \zeta_3) = \alpha_f \wedge d\alpha_f(\zeta_1, \zeta_2, \zeta_3)$$

so this condition is equivalent to asking that  $\alpha_f \wedge d\alpha_f > 0$ .

Therefore  $(\tilde{Z}_\lambda, \omega_f|_{\tilde{Z}_\lambda}, \eta_\lambda)$  is a Liouville filling of its boundary, and as discussed above, since the boundary is  $\#^{k_\lambda}(S^1 \times S^2)$ , we can conclude that  $(\tilde{Z}_\lambda, \omega_f|_{\tilde{Z}_\lambda})$  is a Weinstein domain.  $\square$

**Lemma 6.2.** *Let  $f : (X, \omega) \rightarrow (\mathbb{C}\mathbb{P}^2, \omega_{FS})$  be a symplectic singular branched cover, where  $X$  is a closed 4-manifold. For  $\omega_f = f^*\omega_{FS} + \varepsilon d\phi$  as above with  $\varepsilon$  sufficiently small,  $(X, \omega_f)$  and  $(X, k\omega)$  are symplectomorphic, where  $k$  is the degree of the branch cover.*

*Proof.* The cohomology classes of  $\omega_f$  and  $k\omega$  are the same because  $[\omega_f] = [f^*\omega_{FS} + \varepsilon d\phi] = [f^*\omega_{FS}] = k[\omega]$  since  $f$  has degree  $k$ . By definition of a symplectic singular branched cover, the 2-forms

$$t f^* \omega_0 + (1-t) k \omega$$

are symplectic for all  $t \in [0, 1)$ . Therefore for  $\varepsilon$  sufficiently small, the 2-forms

$$\omega_t := t(f^*\omega_0 + \varepsilon d\phi) + (1-t)k\omega$$

are still symplectic for all  $t \in [0, 1)$ . For  $t = 1$ ,  $\omega_1 = \omega_f$ . Therefore we have a 1-parameter family of symplectic forms  $\omega_t$  in the same cohomology class interpolating between  $\omega_f$  and  $k\omega$ , so by Moser's theorem[Mos65], there is an ambient isotopy  $\Psi_t : X \rightarrow X$  such that  $\Psi_t^*(\omega_t) = \omega_0 = k\omega$ . Therefore  $\Psi_1$  gives the required symplectomorphism.  $\square$

We conclude this section with the proof of our main result.

**Theorem 1.1.** *Every closed, symplectic 4-manifold  $(X, \omega)$  admits a Weinstein trisection.*

*Proof of Theorem 1.1.* Let  $(X, \omega)$  be a closed, symplectic 4-manifold. By Theorem 5.3, there is a singular symplectic branched covering  $f : X \rightarrow \mathbb{C}\mathbb{P}^2$  such that the branch locus  $Q$  is a symplectic braided surface.

By Theorem 5.9, we may assume after a symplectomorphism of  $\mathbb{C}\mathbb{P}^2$  that  $Q$  is in bridge position with respect to a genus one trisection  $\mathbb{T}_0$  of  $\mathbb{C}\mathbb{P}^2$ .

By Theorem 4.1, the preimages  $W_\lambda = f^{-1}(Z_\lambda)$  of the sectors  $Z_\lambda$  of the trisection  $\mathbb{T}_0$  give a trisection  $\tilde{\mathbb{T}} = f^*(\mathbb{T}_0)$  of  $X$ .

By Theorem 6.1,  $X$  carries a symplectic structure  $\omega_f$  such that the trisection  $\mathbb{T} = f^*\mathbb{T}_0$  is Weinstein with respect to  $\omega_f$ . By Lemma 6.2, there is a symplectomorphism  $\Psi : (X, \omega_f) \rightarrow (X, k\omega)$ , where  $k$  is the degree of the covering. Therefore  $\Psi_*(\mathbb{T})$  gives a Weinstein trisection on  $(X, k\omega)$ .

Let  $W_\lambda$  be the three sectors of  $\Psi_*(\mathbb{T})$  on  $X$  for  $\lambda = 1, 2, 3$  and let  $\omega_\lambda = \omega|_{W_\lambda}$ . Then there are Weinstein structures  $(W_\lambda, k\omega_\lambda, V_\lambda, \phi_\lambda)$ . The Weinstein conditions on  $V_\lambda$  and  $\phi_\lambda$  are that  $d\iota_{V_\lambda}(k\omega_\lambda) = k\omega_\lambda$  and  $V_\lambda$  is gradient-like for  $\phi_\lambda$ . Because  $k$  is a constant, this implies  $d\iota_{V_\lambda}(\omega_\lambda) = \omega_\lambda$ . In other words  $V_\lambda$  is also a Liouville vector field for  $\omega_\lambda$ . Therefore globally rescaling the symplectic form by a constant, we still have Weinstein structures on the three sectors  $(W_\lambda, \omega_\lambda, V_\lambda, \phi_\lambda)$ . Therefore  $\Psi_*(\mathbb{T})$  is a Weinstein trisection on  $(X, \omega)$ .  $\square$

## 7. EXAMPLES

In this section we describe a number of examples of Weinstein trisections.

**7.1. Weinstein trisections for complex hypersurfaces in  $\mathbb{C}\mathbb{P}^3$ .** Let  $S_d$  denote the degree  $d$  complex hypersurface in  $\mathbb{C}\mathbb{P}^3$ . Let  $\mathbb{T}_{S_d}$  denote the efficient trisection of  $S_d$  constructed in [LM18, Section 4] as the  $d$ -fold cyclic covering of the genus one trisection of  $\mathbb{C}\mathbb{P}^2$ , branched along an efficient bridge trisection of a curve  $\mathcal{C}_d$  of degree  $d$  in  $\mathbb{C}\mathbb{P}^2$ . A priori, the efficient bridge trisection given in [LM18] corresponds to a surface  $\mathcal{S}$  that is smoothly isotopic to  $\mathcal{C}_d$ . However, by [Lam, Proposition 3.3],  $\mathcal{S}$  is isotopic through (transverse) bridge trisected surfaces to a symplectic surface. Now, by Theorem 6.1, we have that the trisections  $\mathbb{T}_{S_d}$  are Weinstein.

**7.2. A Weinstein trisection for  $\mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$ .** Consider the the surface  $\mathcal{C}$  described by the shadow diagram in Figure 9(a). (See [LM18] for details regarding shadow diagrams.) This diagram corresponds to a singular bridge trisection, with each patch given as the cone on a right-handed trefoil. We encourage the reader to verify that each of the links

$$(S_\lambda^3, K_\lambda) = (H_\lambda, \mathcal{T}_\lambda) \bigcup_{(\Sigma, \mathbf{x})} \overline{(H_{\lambda+1}, \mathcal{T}_{\lambda+1})}$$

is a right-handed trefoil: For each  $\lambda \in \mathbb{Z}_3$ , find an orientation-preserving homeomorphism from the shadow diagram onto the genus one Heegaard surface in  $S^3$  such that  $H_\lambda$  is inside the torus and  $\overline{H_{\lambda+1}}$  is outside, and consider the knot obtained by perturbing the interiors of the shadow arcs into their respective solid tori.

To see that this surface has degree four, we briefly recall how the homology of  $\mathbb{C}\mathbb{P}^2$  can be understood from its trisection diagram (cf. [FKSZ18]). Following [LM18, Subsection 6.2], we note that

$$H_2(X; \mathbb{Z}) = L_\gamma \cap (L_\alpha + L_\beta),$$

where  $L_\lambda \subset H_1(\Sigma; \mathbb{Z})$  is the subspace spanned by the curve  $\lambda$ . Note that  $[\gamma] = -[\alpha] - [\beta]$  in  $H_1(\Sigma; \mathbb{Z})$ . It follows that  $[\mathbb{C}\mathbb{P}^1] = [\gamma] = -[\alpha] - [\beta]$ . Let  $\phi$  denote the triangular region of  $\Sigma \setminus (\alpha \cup \beta \cup \gamma)$  containing the positive bridge point, together with a compressing disk bounded by each of the  $\lambda$  in  $H_\lambda$ . Then,  $\phi$  is a 2-chain in  $\mathbb{C}\mathbb{P}^2$  representing  $[\mathbb{C}\mathbb{P}^1]$  in  $H_2(\mathbb{C}\mathbb{P}^2; \mathbb{Z})$ .

To apply Proposition 6.1 of [LM18], we need a shadow diagram for  $\mathcal{C}$  where the shadow arcs are disjoint from the cut curves for their respective solid tori. This could be accomplished by isotoping the cut curves (cf. Figure 5(c)), but then we would need to recalculate  $\phi$ . Instead, we perturb the bridge trisection to be 4-bridge, as shown in Figure 5(b). Now, it follows that  $[\mathbb{C}\mathbb{P}^1] \cdot [\mathcal{C}] = 4$ , since there are now four positive bridge points in the triangular region of  $\phi$ .

Using the techniques of Section 3 of [Lam], combined with a slight generalization of [Lam, Propositions 3.2 and 3.3] to a setting that includes cusp singularities, it can be shown that  $\mathcal{C}$  is isotopic through singular (transverse) bridge trisections to a symplectic surface. Thus,  $\mathcal{C}$  is a symplectic tri-cuspidal quartic. (Note that by [GS19] there is a unique symplectic isotopy class of tri-cuspidal quartics.) In what follows, we will describe a singular, irregular, 3-fold branched covering

$$f: \mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2} \rightarrow \mathbb{C}\mathbb{P}^2$$

that is branched along the tri-cuspidal quartic  $\mathcal{S}$ . Applying Theorem 4.1, we will lift a genus one Weinstein trisection of  $\mathbb{C}\mathbb{P}^2$  to a genus two trisection of  $\mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$ , and this lifted trisection will be Weinstein by Theorem 6.1.

Figure 5(c) is a doubly pointed trisection diagram for  $(\mathbb{C}\mathbb{P}^2, \mathcal{C})$  that is obtained from Figure 5(a) by isotoping the  $\lambda$ -curves to be disjoint from the shadow arcs and forgetting the shadow arcs. (The arcs can be recovered uniquely by connecting the two bridge points in the complement of the respective  $\lambda$ -curve; see [GM18] for more discussion of doubly pointed trisection diagrams and the connection to knotted spheres in 4-manifolds and note the relevant connection here that  $\mathcal{C}$  is rational.) Figure 5(d)



is an isotopic doubly pointed trisection diagram that has the feature that the  $\alpha$ , is now parallel to an edge of the square representing the torus.

Figure 5(e), which is just this new doubly pointed trisection diagram with disk neighborhoods of the bridge points removed, can be used to calculate  $\pi_1(\mathbb{CP}^2 \setminus \nu(\mathcal{C}))$  as follows. Choose as base-point the corner(s) of the square, let the classes of the horizontal and vertical edges be  $a$  and  $b$ , respectively. Connect the punctures to the corners with whiskers as shown, and denote the loops following these arcs and encircling the punctures  $x$  and  $y$ , respectively. Then, the compressing disk bounded by  $\alpha$  gives the relation  $a = 1$ , the compressing disk bounded by  $\beta$  gives the relation  $y^{-1}a^{-1}b^{-1}x^{-1}bayb = 1$ , and the compressing disk bounded by  $\gamma$  gives the relation  $y^{-1}b^{-1}y^{-1}a^{-1}b^{-1}x^{-1}b = 1$ . We also get the relation  $ayba^{-1}b^{-1}x^{-1} = 1$  from the core surface. From all this, we get the presentation

$$\pi_1(\mathbb{CP}^2 \setminus \nu(\mathcal{C})) = \langle b, y \mid yby = byb, y^2 = b^{-2} \rangle,$$

noting that  $x = y$ . The fact that this choice of base-point and whiskers yields  $x = y$  and  $a = 1$  will make drawing the branched cover easier, since this presentation gives rise to a particularly nice representation to  $\mathcal{S}_3$ .

**Remark 7.1.** We remark in passing that this group is closely related to several well-known groups via the following sequence of quotients:

$$\mathcal{B}_3 \cong \pi(T_{3,2}) \twoheadrightarrow \pi(\mathcal{C}) \twoheadrightarrow \pi(\mathcal{P}_{T_{3,2}}) \cong \mathrm{PSL}_2(\mathbb{Z}).$$

Here,  $\mathcal{B}_3$  is the braid group on three strands,  $\pi(T_{3,2})$  is the group of the right-handed trefoil,  $\pi(\mathcal{C})$  is the group of the tri-cuspidal quartic,  $\pi(\mathcal{P}_{T_{3,2}})$  is the group of the knotted projective plane in  $S^4$  obtained by forming the connected sum of the spun trefoil with the standard cross-cap, and  $\mathrm{PSL}_2(\mathbb{Z})$  is the modular group.

Consider the representation  $\rho: \pi_1(\mathbb{CP}^2 \setminus \nu(\mathcal{C})) \rightarrow \mathcal{S}_3$ , where  $y \mapsto (12)$  and  $b \mapsto (23)$ . Note that we must have  $a \mapsto 1$  and  $x \mapsto (12)$ . We remark that this representation is the unique (up to conjugation) representation onto  $\mathcal{S}_3$ , since any such representation is determined by choosing distinct transpositions for  $\rho(y)$  and  $\rho(b)$ . Let  $f^\circ: X^\circ \rightarrow \mathbb{CP}^2 \setminus \nu(\mathcal{C})$  be the irregular 3-fold (unbranched) covering corresponding to  $\rho$ . This covering can be extended to a (non-singular) branched covering over  $\mathbb{CP}^2 \setminus \nu$  (the cusps of  $\mathcal{C}$ ). On the boundary of the neighborhood of the cusps, this branched covering induces the unique irregular 3-fold covering of  $S^3$  over its self with branch set the right-handed trefoil, as in Examples 3.5(2). Thus, we can cone off over the cusps to obtain a singular branched covering  $f: X \rightarrow \mathbb{CP}^2$  with only cusp singularities. In particular,  $f$  satisfies the hypotheses of Theorem 6.1. Note that  $f$  is the unique such map, by our analysis of  $\rho$ .

To describe a trisection of the cover  $X$ , we first describe  $\tilde{\Sigma} = f^{-1}(\Sigma)$ . Let  $\Sigma^\circ$  denote the core surface  $\Sigma$ , cut along an arc connecting the punctures and along a curve parallel to  $\alpha$ . Then  $\Sigma^\circ$  is a punctured annulus whose core curve is given by  $a$ . Let  $\tilde{\Sigma}^\circ$  denote three copies of  $\Sigma^\circ$ , and note that the restriction  $f|_{\mathrm{Int}(\tilde{\Sigma}^\circ)}$  can be thought of as the three-to-one map  $\mathrm{Int}(\tilde{\Sigma}^\circ) \rightarrow \mathrm{Int}(\Sigma^\circ)$ , since  $\rho(a) = 1$ .

The three sheets,  $\tilde{\Sigma}_1^\circ$ ,  $\tilde{\Sigma}_2^\circ$ , and  $\tilde{\Sigma}_3^\circ$ , of  $\tilde{\Sigma}^\circ$  are shown in Figure 9(f). To recover  $\tilde{\Sigma}$ , we must understand the identifications of these sheets described in this figure. In the left sheet, the sides of  $\tilde{\Sigma}_1^\circ$  corresponding to  $a$  are re-glued as they are in  $\Sigma$ ; since  $\rho(b) = (23)$ , the curve  $b$  lifts to a curve in this first sheet. Now, the first sheet is a punctured torus with boundary coming from the cut along the arc connecting the punctures. This puncture is identified, as depicted, with the corresponding puncture in the second sheet. This is because this puncture came from cutting open  $\Sigma^\circ$  along an arc that connected the punctures  $x$  and  $y$ , each of whose image under  $\rho$  is  $(12)$ . For the same reason, in the third sheet, the cut along this arc is re-glued; the curves  $x$  and  $y$  lift to curves in the third sheet. Finally, the two  $a$ -curves of the third sheet are identified with the two  $a$ -curves of the second sheet, as shown. The




FIGURE 9. (a)–(d) Diagrams for a tri-cuspidal quartic. (e) The diagram from (c) with neighborhoods of the bridge points removed. (f) The three sheets corresponding the irregular 3-fold cover of the punctured surface. (g) A trisection diagram (after identification) for the irregular 3-fold branched cover of the tri-cuspidal quartic.

relevant fact here is that  $\rho(b) = (23)$ , so  $b$  is double covered by a curve contained in the union of the second and third sheets, which is a punctured torus. Finally, the meridional punctures  $x$  and  $y$  are filled in in the third sheet; this gives a branched covering of surfaces.

By Theorem 4.1, the genus one trisection of  $\mathbb{C}\mathbb{P}^2$  will lift to a trisection of  $\mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$ . However, we will now check explicitly that the cover of the core surface described above extends over the other pieces of the trisection. It remains to check that the map  $f: \tilde{\Sigma} \rightarrow \Sigma$  described above can be extended to three branched coverings of handlebodies. Since  $f^{-1}(\lambda)$  is a collection of curves for each  $\lambda$ , it suffices to check that each such collection is a cut system for  $\tilde{\Sigma}$ . We saw above that  $\tilde{\Sigma}$  is the union of two punctured tori, hence a genus two surface. So, we need that  $f^{-1}(\lambda)$  contains two non-parallel, non-separating curves for each  $\lambda$ . This is verifiable in Figure 5(f); each  $f^{-1}(\lambda)$  consists of three curves, two of which are parallel. In Figure 5(g), a redundant  $\lambda$ -curve has been deleted, and isotopy of the remaining two curves has been performed.

The result is that Figure 5(f) is a genus two trisection diagram for  $\mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$ . To verify this last claim, one could appeal to [MZ17b] to deduce that the diagram is either reducible or corresponds to  $S^2 \times S^2$ . Then, one could calculate homology following [FKSZ18] or [LM18, Subsection 6.2]. Alternatively, one could perform handleslides to find a reducing curve (a separating curve disjoint from all six cut curves) and verify explicitly that the two resulting summands correspond to  $\mathbb{C}\mathbb{P}^2$  and  $\overline{\mathbb{C}\mathbb{P}^2}$ .

## REFERENCES

- [AK00] Denis Auroux and Ludmil Katzarkov, *Branched coverings of  $\mathbb{C}\mathbb{P}^2$  and invariants of symplectic 4-manifolds*, Invent. Math. **142** (2000), no. 3, 631–673. MR 1804164
- [Aur97] D. Auroux, *Asymptotically holomorphic families of symplectic submanifolds*, Geom. Funct. Anal. **7** (1997), no. 6, 971–995. MR 1487750
- [Aur00] Denis Auroux, *Symplectic 4-manifolds as branched coverings of  $\mathbb{C}\mathbb{P}^2$* , Invent. Math. **139** (2000), no. 3, 551–602. MR 1738061
- [BCKM19] Ryan Blair, Patricia Cahn, Alexandra Kjachukova, and Jeffrey Meier, *A note on three-fold branched covers of  $S^4$* , arXiv e-prints (2019), arXiv:1909.11788.
- [CE12] Kai Cieliebak and Yakov Eliashberg, *From Stein to Weinstein and back*, American Mathematical Society Colloquium Publications, vol. 59, American Mathematical Society, Providence, RI, 2012, Symplectic geometry of affine complex manifolds. MR 3012475
- [CK17] Patricia Cahn and Alexandra Kjachukova, *Singular branched covers of four-manifolds*, arXiv e-prints (2017), arXiv:1710.11562.
- [CO19] Nickolas A. Castro and Burak Ozbagci, *Trisections of 4-manifolds via Lefschetz fibrations*, Math. Res. Lett. **26** (2019), no. 2, 383–420. MR 3999550
- [Eli90] Yakov Eliashberg, *Filling by holomorphic discs and its applications*, Geometry of low-dimensional manifolds, 2 (Durham, 1989), London Math. Soc. Lecture Note Ser., vol. 151, Cambridge Univ. Press, Cambridge, 1990, pp. 45–67. MR 1171908
- [FKSZ18] Peter Feller, Michael Klug, Trenton Schirmer, and Drew Zemke, *Calculating the homology and intersection form of a 4-manifold from a trisection diagram*, Proc. Natl. Acad. Sci. USA **115** (2018), no. 43, 10869–10874. MR 3871789
- [Gay16] David T. Gay, *Trisections of Lefschetz pencils*, Algebr. Geom. Topol. **16** (2016), no. 6, 3523–3531. MR 3584265
- [Gei08] Hansjörg Geiges, *An introduction to contact topology*, Cambridge Studies in Advanced Mathematics, vol. 109, Cambridge University Press, Cambridge, 2008. MR 2397738
- [GK16] David Gay and Robion Kirby, *Trisecting 4-manifolds*, Geom. Topol. **20** (2016), no. 6, 3097–3132. MR 3590351
- [GM18] David Gay and Jeffrey Meier, *Doubly pointed trisection diagrams and surgery on 2-knots*, arXiv e-prints (2018), arXiv:1806.05351.
- [Gom98] Robert E. Gompf, *Handlebody construction of Stein surfaces*, Ann. of Math. (2) **148** (1998), no. 2, 619–693. MR 1668563
- [GS99] Robert E. Gompf and András I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics, vol. 20, American Mathematical Society, Providence, RI, 1999. MR 1707327 (2000h:57038)

- [GS19] Marco Golla and Laura Starkston, *The symplectic isotopy problem for rational cuspidal curves*, arXiv e-prints (2019), arXiv:1907.06787.
- [Lam] Peter Lambert-Cole, *Symplectic surfaces and bridge position*, arXiv e-prints, arXiv:1904.05137, arXiv:1904.05137.
- [Lam18] ———, *Bridge trisections in  $CP^2$  and the Thom Conjecture*, arXiv e-prints (2018), arXiv:1807.10131.
- [LM18] Peter Lambert-Cole and Jeffrey Meier, *Bridge trisections in rational surfaces*, arXiv e-prints (2018), arXiv:1810.10450.
- [Mos65] Jürgen Moser, *On the volume elements on a manifold*, Trans. Amer. Math. Soc. **120** (1965), 286–294. MR 182927
- [MSZ16] Jeffrey Meier, Trent Schirmer, and Alexander Zupan, *Classification of trisections and the Generalized Property R Conjecture*, Proc. Amer. Math. Soc. **144** (2016), no. 11, 4983–4997. MR 3544545
- [MZ17a] Jeffrey Meier and Alexander Zupan, *Bridge trisections of knotted surfaces in  $S^4$* , Trans. Amer. Math. Soc. **369** (2017), no. 10, 7343–7386. MR 3683111
- [MZ17b] ———, *Genus-two trisections are standard*, Geom. Topol. **21** (2017), no. 3, 1583–1630. MR 3650079
- [MZ18] ———, *Bridge trisections of knotted surfaces in 4-manifolds*, Proc. Natl. Acad. Sci. USA **115** (2018), no. 43, 10880–10886. MR 3871791
- [RT16] J. Hyam Rubinstein and Stephan Tillmann, *Multisections of piecewise linear manifolds*, arXiv e-prints (2016), arXiv:1602.03279.
- [Wei91] Alan Weinstein, *Contact surgery and symplectic handlebodies*, Hokkaido Math. J. **20** (1991), no. 2, 241–251. MR 1114405
- [Wen10] Chris Wendl, *Strongly fillable contact manifolds and J-holomorphic foliations*, Duke Math. J. **151** (2010), no. 3, 337–384. MR 2605865
- [Zud08] Daniele Zuddas, *Branched coverings and 4-manifolds*, Ph.D. thesis, Scuola Normale Superiore Pisa, 2008.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY  
 MAX PLANCK INSTITUTE FOR MATHEMATICS

*E-mail address:* `plc@mpim-bonn.mpg.de`

*URL:* <https://www.math.gatech.edu/~plambertcole3>

DEPARTMENT OF MATHEMATICS, WESTERN WASHINGTON UNIVERSITY

*E-mail address:* `jeffrey.meier@wwu.edu`

*URL:* <https://jeffreymeier.org>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS

*E-mail address:* `lstarkston@math.ucdavis.edu`

*URL:* <https://www.math.ucdavis.edu/~lstarkston/>