

Remarks on computing Green functions

Frank Lübeck*

April 3, 2020

Abstract

We describe the computation of generalized Green functions and 2-parameter Green functions for finite reductive groups.

1 Introduction

Let \mathbf{G} be a connected reductive algebraic group over an algebraic closure $\overline{\mathbb{F}}_q$ of a finite field in characteristic p , defined over \mathbb{F}_q , and let $F : \mathbf{G} \rightarrow \mathbf{G}$ be the corresponding Frobenius morphism.

We are interested in (complex) class functions on unipotent conjugacy classes of the finite group $G(q) := \mathbf{G}^F$ of F -fixed points of \mathbf{G} . The aim is to compute explicitly the values of all ordinary irreducible characters of $G(q)$ on unipotent classes (which is an important step to find the full character table of $G(q)$). Here we consider one part of this problem which is the computation of Green functions.

Lusztig defined certain class functions on unipotent classes, called *generalized Green functions*, in terms of an induction functor for character sheaves. These functions occur in a character formula for characteristic functions of character sheaves, see [Lus85, 8.3, 8.5].

Deligne and Lusztig [DL76] and Lusztig [Lus76, 1.,2.] also introduced for F -stable Levi subgroups \mathbf{M} of \mathbf{G} a twisted induction R_M^G from class functions on $M(q)$ to those of $G(q)$, using l -adic cohomology of certain varieties. A character formula for this map involves certain 2-parameter Green functions which are parameterized by the unipotent classes of $M(q)$ and of $G(q)$.

In [Lus90] Lusztig showed (under some restrictions on p and q) that on the level of class functions on unipotent classes both types of induction are the same. From this one can conclude that the 2-parameter Green functions can be computed once the generalized Green functions are known for $M(q)$ and $G(q)$.

The generalized Green functions can be computed in many cases. So, in these cases we can compute 2-parameter Green functions and this yields an essential ingredient for evaluating the character formula for R_M^G .

*The authors acknowledges support from the DFG within the TRR 195.

In section 2 we describe in more detail the relation between generalized Green functions and 2-parameter Green functions.

In section 3 we collect what is known about the computation of generalized Green functions. There are two main steps. The first uses the known generalized Springer correspondence to compute the generalized Green functions as linear combinations of certain functions which have non-zero values only on elements lying in a fixed unipotent class of the algebraic group \mathbf{G} . The second is to determine these functions.

In section 4 we consider the groups $G(q) = \text{Spin}_8(q)$ as an example. In that case the known general results do not fully determine the generalized Green functions. But we show how to find the 2-parameter Green functions for some maximal Levi subgroups and the missing parameters for the generalized Green functions at the same time.

2 Computing 2-parameter Green functions from generalized Green functions

2.1 2-parameter Green functions

Let \mathbf{G} , F , $G(q)$ be as in the introduction. Let \mathcal{C} be the set of unipotent conjugacy classes of $G(q)$. Furthermore, let \mathbf{M} be an F -stable Levi subgroup of \mathbf{G} , $M(q) := \mathbf{M}^F$ and \mathcal{C}' be the unipotent classes of $M(q)$.

Lusztig [Lus76, 1.,2.] defined a twisted induction from $M(q)$ to $G(q)$ which defines a linear map R_M^G from the class functions on $M(q)$ to the class functions on $G(q)$. If f is a class function on $M(q)$ then there is a character formula which on unipotent classes $c \in \mathcal{C}$ of $G(q)$ is (we sometimes write class functions as functions on conjugacy classes):

$$(R_M^G f)(c) = \sum_{c' \in \mathcal{C}'} g_M^G(c, c') f(c'), \quad (1)$$

where $g_M^G(c, c')$ is the alternating sum of the traces of $(u, u') \in c \times c'$ on certain modules defined via l -adic cohomology divided by $|C_{M(q)}(u')|$ for a $u' \in c'$.

We call g_M^G the 2-parameter Green functions. For $M(q) = T(q)$ a maximal torus and so $c' = \{1\}$ these include the ordinary Green functions $Q_T^G(u) = g_T^G(u, 1)$ introduced in [DL76]. (Note that we use a slightly different normalization compared to the functions $Q_M^G(u, u') = \frac{1}{|c'|} g_M^G(c, c')$ in [DM91, 12.1], and to the functions $\gamma_M^G(u, u') = \frac{|M(q)|}{|c'|} g_M^G(c, c')$ in [Lus90, 1.7].)

In general, these functions are difficult to compute directly from their definition.

But in the special case that $\mathbf{M} = F(\mathbf{M})$ is the Levi complement of an F -stable parabolic subgroup $\mathbf{P} = \mathbf{M}\mathbf{U}$ with unipotent radical \mathbf{U} the module from l -adic cohomology has a simple structure, namely it is the permutation module on $G(q)/U(q)$ on which $G(q)$ is acting by left multiplication and $M(q)$ by right

multiplication, see [DM91, 11.1]. So, in this case the trace of $(g, m) \in G(q) \times M(q)$ is just the number of fixed points under above action. It is easy to see that with each fixed point $xU(q)$ also $xyU(q)$ is a fixed point for all $y \in C_{M(q)}(m)$. This shows that

$$\text{if } \mathbf{M} \text{ is contained in an } F\text{-stable parabolic, then all } g_M^G(c, c') \in \mathbb{Z}_{\geq 0}. \quad (2)$$

In particular, for $\mathbf{M} = \mathbf{G}$, and so $\mathbf{U} = \{1\}$ we get that $g_G^G(c, c') = \delta_{c, c'}$ for all $c, c' \in \mathcal{C}$.

Conjecture: the values of $g_M^G(c, c')$ are always algebraic integers.

Furthermore, if we know a set of linearly independent unipotent class functions for $M(q)$ of cardinality $|\mathcal{C}'|$ as well as their images under R_M^G then we can use equation (1) to compute all $g_M^G(c, c')$. Such a set of functions is provided by the generalized Green functions which we recall now.

2.2 Generalized Green functions

Let \mathbf{G}, F be as above. Lusztig has defined in [Lus85, 8.3] a set of unipotent class functions on $G(q)$, called generalized Green functions. These are partitioned into subsets which belong to triples $(\mathbf{L}, C, \mathcal{E})$ of a Levi subgroup \mathbf{L} of \mathbf{G} , a unipotent class C of \mathbf{L} and a cuspidal local system \mathcal{E} on C , up to \mathbf{G} -conjugacy. Assume that \mathbf{L} is an F -stable Levi complement of an F -stable parabolic subgroup, $F(C) = C$ and that $F^*\mathcal{E} \cong \mathcal{E}$.

Fix an isomorphism $\tau : F^*\mathcal{E} \rightarrow \mathcal{E}$ (unique up to scalar). The Frobenius morphism F induces an action on the group $W_{\mathbf{L}} = N_{\mathbf{G}}(\mathbf{L})/\mathbf{L}$. The $G(q)$ -conjugacy classes of F -stable \mathbf{G} -conjugates of \mathbf{L} are parameterized by the F -conjugacy classes of $W_{\mathbf{L}}$. For $w \in W_{\mathbf{L}}$ let \mathbf{L}_w be a corresponding twisted Levi subgroup, C_w the conjugate of C in \mathbf{L}_w and \mathcal{E}_w the conjugate local system. The isomorphism τ can also be conjugate to an isomorphism $\tau_w : F^*\mathcal{E}_w \rightarrow \mathcal{E}_w$ (see [Lus85, 10.6]). Lusztig defines for each such $(\mathbf{L}_w, C_w, \mathcal{E}_w, \tau_w)$ (up to $G(q)$ -conjugacy) a unipotent class function $Q_{\mathbf{L}_w, C_w, \mathcal{E}_w, \tau_w}^G$ on $G(q)$.

These functions form a basis of the space of class functions on unipotent classes of $G(q)$, see [Lus86b, 25.4]. More precisely, they fulfill orthogonality relations with respect to the scalar product $(f_1, f_2) := 1/|G(q)| \sum_{u \in G(q)_{uni}} f_1(u) \overline{f_2(u)}$ for class functions f_1, f_2 on $G(q)$: the scalar product of different Green functions is 0 and (see [Lus85, 9.11]):

$$(Q_{\mathbf{L}_w, C_w, \mathcal{E}_w, \tau_w}^G, Q_{\mathbf{L}_w, C_w, \mathcal{E}_w, \tau_w}^G) = \frac{|W_{\mathbf{L}}^{wF}| \cdot q^d}{|Z^0(\mathbf{L})^F|}, \quad (3)$$

where $Z^0(\mathbf{L})$ is the connected center of \mathbf{L} and $d = \dim C - \dim \mathbf{L} + \dim Z^0(\mathbf{L})$.

Under some restriction on the characteristic p of \mathbf{G} Lusztig showed in [Lus90, 1.14] that for sufficiently large q (depending on the Dynkin type of \mathbf{G}) we have

$$Q_{\mathbf{L}_w, C_w, \mathcal{E}_w, \tau_w}^G = R_L^G(Q_{\mathbf{L}_w, C_w, \mathcal{E}_w, \tau_w}^{L_w}). \quad (4)$$

The restriction on p is no longer needed because of results by Shoji [Sho95, I 1.11, 1.12] and Lusztig [Lus12]. The restriction on q can at least be dropped for the case $\mathbf{L} = \mathbf{T}$ a maximal torus, see [Sho95, II 5.5], or more generally for all $(\mathbf{L}, C, \mathcal{E})$ which still occur for the group $\mathbf{G}/Z(\mathbf{G})$ with connected center, see [Sho96, 4.].

Now let \mathbf{M} be any F -stable Levi subgroup of \mathbf{G} . Let $(L_v, C_v, \mathcal{E}_v, \tau_v)$ be a tuple as above, but now \mathbf{L} is an F -stable Levi subgroup of an F -stable parabolic subgroup of \mathbf{M} , and $v \in N_{\mathbf{M}}(\mathbf{L})/\mathbf{L}$.

Using equation (4) and the transitivity of twisted induction (see [DM91, 11.5]) we get

$$R_M^G(Q_{L_v, C_v, \mathcal{E}_v, \tau_v}^M) = R_M^G R_L^M(Q_{L_v, C_v, \mathcal{E}_v, \tau_v}^{L_v}) = Q_{L_v, C_v, \mathcal{E}_v, \tau_v}^G. \quad (5)$$

So, if we know the generalized Green functions of $M(q)$ and of $G(q)$ and if we can identify each Green function $Q_{L_v, C_v, \mathcal{E}_v, \tau_v}^G$ with one of $G(q)$ (that is, we can identify the $w \in W_{\mathbf{L}}$ such that \mathbf{L}_v and \mathbf{L}_w are $G(q)$ -conjugate, and we can compare τ_v and τ_w), then we can compute the 2-parameter Green functions via (1). The relations (3) make it easy to invert the matrix of values of generalized Green functions of $M(q)$.

3 Computing generalized Green functions

It is now possible to compute generalized Green functions in many cases, and otherwise to compute at least a good "approximation". Here is a sketch of the method.

As in the last section let $(\mathbf{L}, C, \mathcal{E})$ be a triple of a Levi subgroup of \mathbf{G} , a unipotent class C of \mathbf{L} and \mathcal{E} be a cuspidal local system on C . We discuss those generalized Green functions corresponding to this triple (an important case is $(\mathbf{T}, \{1\}, \mathbb{Q}_l)$ which yields the ordinary Green functions of $G(q)$ which are needed to compute the Deligne-Lusztig characters).

3.1 The generalized Springer correspondence

In [Lus84] Lusztig associates a geometric object to \mathbf{G} and (C, \mathcal{E}) whose decomposition into simple objects is described in terms of the irreducible characters of the relative Weyl group $W_{\mathbf{L}} = N_{\mathbf{G}}(\mathbf{L})/\mathbf{L}$. The simple object associated to a character $\chi \in \text{Irr}(W_{\mathbf{L}})$ is an object defined by a pair (C', \mathcal{E}') of a unipotent class C' and a \mathbf{G} -invariant irreducible local system \mathcal{E}' on C' .

This construction leads to a bijective map, called the generalized Springer correspondence,

$$\dot{\bigcup}_{(\mathbf{L}, C, \mathcal{E})} \text{Irr}(W_{\mathbf{L}}) \rightarrow \mathcal{N}_{\mathbf{G}}, \quad (6)$$

where $\mathcal{N}_{\mathbf{G}}$ is the set of pairs (C', \mathcal{E}') of a unipotent class C' in \mathbf{G} and a \mathbf{G} -equivariant irreducible local system \mathcal{E}' on C' modulo \mathbf{G} -conjugacy; the left hand side is a disjoint union over all triples $(\mathbf{L}, C, \mathcal{E})$ as above modulo \mathbf{G} -conjugacy.

The generalized Springer correspondence has been explicitly determined in [Lus84], [LS85], [Spa85] and [Lus19] except for two small ambiguities in composition factors of type E_8 in characteristic 3 (and in [Sho97, 8.6] is a correction of [Lus84, 12.3]).

To understand how the Springer correspondence can be described, we give some more details about the sets on both sides of the map.

The set $\mathcal{N}_{\mathbf{G}}$: This set is a disjoint union of the subsets which collect the pairs (C', \mathcal{E}') for one fixed unipotent class C' . So, fix C' and $u \in C'$. Let $A_{\mathbf{G}}(u) = C_{\mathbf{G}}(u)/C_{\mathbf{G}}^0(u)$ be the component group of the centralizer of u in \mathbf{G} . Then the \mathbf{G} -equivariant irreducible local systems \mathcal{E}' on C' can be parameterized by the irreducible representations of $A_{\mathbf{G}}(u)$. For simple \mathbf{G} of simply-connected type parameterizations of the unipotent classes and descriptions of the corresponding component groups are known; for classical groups in terms of partitions for Jordan block sizes and sometimes additional invariants, and for exceptional types in a more ad hoc manner.

Let \mathbf{L} be a Levi subgroup of \mathbf{G} . Up to conjugacy we can assume that it is a standard Levi subgroup; such that the Weyl group of \mathbf{L} is a Coxeter group with set I of Coxeter generators which is a subset of Coxeter generators S of the Weyl group W of \mathbf{G} . In the case that \mathbf{L} has a cuspidal pair (C, \mathcal{E}) as above the relative Weyl group $W_{\mathbf{L}} = N_{\mathbf{G}}(\mathbf{L})/\mathbf{L}$ is also a Coxeter group, see [Lus84, Thm. 9.2]. It has Coxeter generators labeled by the complement $S \setminus I$; for $j \in S \setminus I$ the relative Weyl group of \mathbf{L} in the Levi subgroup corresponding to $I \cup \{j\}$ is of order 2 and its generator is one of the Coxeter generators of $W_{\mathbf{L}}$. So, using the usual labelings of irreducible characters of finite Coxeter groups there is a good parameterization for the left hand side of the Springer correspondence (6).

The articles mentioned above contain explicit descriptions of the Springer correspondence for all simple simply-connected groups. The general case can be reduced to these results. First, the natural quotient map $\mathbf{G} \rightarrow \mathbf{G}/Z^0(\mathbf{G})$ is a bijection on unipotent elements and since $Z^0(\mathbf{G})$ is contained in the connected centralizer of any element, the component groups do not change. The description of the generalized Springer correspondence is the same for \mathbf{G} and $\mathbf{G}/Z^0(\mathbf{G})$. So we can assume that \mathbf{G} is semisimple. Then there is a covering map $\mathbf{G}_1 \times \mathbf{G}_2 \times \cdots \times \mathbf{G}_k \rightarrow \mathbf{G}$ where the \mathbf{G}_i ($1 \leq i \leq k$) are simple simply-connected groups and the kernel of the map is a finite subgroup of the center of the left hand side. Let $\tilde{u} = (\tilde{u}_1, \dots, \tilde{u}_k) \in \mathbf{G}_1 \times \cdots \times \mathbf{G}_k$ be a unipotent element with image u in \mathbf{G} , and let A_i be the component group of \tilde{u}_i in \mathbf{G}_i . Then \tilde{u} has component group $A_1 \times \cdots \times A_k$. Let Z_i be the (finite) centre of \mathbf{G}_i , there is a natural map $Z_i \rightarrow A_i$, let \bar{Z}_i be its image. The covering map is a bijection on unipotent elements and maps centralizers of unipotent elements to centralizers, but the component groups can become smaller. If $K \leq Z_1 \times \cdots \times Z_k$ is the kernel of the covering and \bar{K} is its image in $\bar{Z}_1 \times \cdots \times \bar{Z}_k$ then the component group of $u \in \mathbf{G}$ is $A_{\mathbf{G}}(u) = A_1 \times \cdots \times A_k/\bar{K}$. So, $\text{Irr}(A_{\mathbf{G}}(u))$ is in bijection with the characters in $\text{Irr}(A_1 \times \cdots \times A_k)$ which have \bar{K} in their kernel. The generalized Springer correspondence for \mathbf{G} is given by the part of the generalized Springer correspondence of $G_1 \times \cdots \times G_k$ where

the image is a pair (C', \mathcal{E}') for which \mathcal{E}' corresponds to one of the characters of the component group as just described. The generalized Springer correspondence of $G_1 \times \cdots \times G_k$ is easily described componentwise. So, in addition to the description of the generalized Springer correspondence for all simple simply-connected groups \mathbf{G} the maps $Z(\mathbf{G}) \rightarrow A_{\mathbf{G}}(u)$ for all unipotent elements of \mathbf{G} must be known for this reduction. This information is implicitly contained in the description of the Springer correspondence (in terms of characters of the centers).

3.2 Generalized Springer correspondence and Frobenius action

Now we take also the Frobenius morphism of \mathbf{G} into account. We are only interested in the part of the generalized Springer correspondence which belongs to triples $(\mathbf{L}, C, \mathcal{E})$ whose \mathbf{G} -conjugacy class is F -stable. In that case we can assume that for our representative in this class we have $F(\mathbf{L}) = \mathbf{L}$ is contained in an F -stable parabolic subgroup, $F(C) = C$ and for $u \in C^F$ the induced F -action on $A_{\mathbf{L}}(u)$ and so $\text{Irr}(A_{\mathbf{L}}(u))$ fixes the character corresponding to \mathcal{E} . There is an induced action of F on $W_{\mathbf{L}}$, we only consider the generalized Springer correspondence for the characters of $W_{\mathbf{L}}$ which are F -stable. Then, the image of all of these F -stable characters for all F -stable $(\mathbf{L}, C, \mathcal{E})$ consists exactly of the F -stable pairs $(C', \mathcal{E}') \in \mathcal{N}_{\mathbf{G}}$.

Let $\iota = (C', \mathcal{E}') \in \mathcal{N}_{\mathbf{G}}$ be an F -stable pair. This means $F(C') = C'$ so that there is a $u \in (C')^F = C' \cap G(q)$, and that there is an isomorphism $\tau : F^*\mathcal{E}' \rightarrow \mathcal{E}'$ (unique up to a nonzero scalar). Then F induces an automorphism on the component group $A_{\mathbf{G}}(u)$. The Lang-Steinberg theorem yields that the $G(q)$ -conjugacy classes contained in $(C')^F$ are parameterized by the F -conjugacy classes of $A_{\mathbf{G}}(u)$. If $a \in A_{\mathbf{G}}(u)$ we write C'_a for the corresponding $G(q)$ -conjugacy class (this labelling depends on the choice of u). Let χ be the character of $A_{\mathbf{G}}(u)$ that parameterizes \mathcal{E}' , then χ is F -stable and can be extended to the semidirect product $A_{\mathbf{G}}(u) \rtimes \langle F \rangle$ (here F means the automorphism induced on $A_{\mathbf{G}}(u)$), this class function is nonzero and unique on the coset $A_{\mathbf{G}}(u)F$ up to multiplication with an $|F|$ -th root of unity. To τ there is an associated class function Y_{ι} on $G(q)$ which is zero outside $(C')^F$ and has value

$$Y_{\iota}(C'_a) = c \cdot \chi(a^F) \text{ for } a \in A_{\mathbf{G}}(u) \quad (7)$$

for some non-zero constant c which depends on τ and the chosen extension of χ . See [Lus86b, 24.2] or [Sho06, 1.3] for more details.

Let $(\mathbf{L}, C, \mathcal{E})$ be an F -stable triple as before. Fix an isomorphism $\tau : F^*\mathcal{E} \rightarrow \mathcal{E}$ and fix for each F -stable irreducible character χ of $W_{\mathbf{L}}$ an extension to the coset $W_{\mathbf{L}}F$ (in [Lus86a, 17.2] Lusztig chooses such extensions which he calls preferred extensions). Let $\iota \in \mathcal{N}_{\mathbf{G}}$ be the image of χ under the Springer correspondence. Lusztig's construction of the generalized Green functions 2.2 shows how these data determine a unique function Y_{ι} as in the last paragraph (that is, determine the constant c), see again [Lus86b, 24.2] and [Sho06, 1.3]. The Lusztig-Shoji algorithm allows to compute the generalized Green functions corresponding to $(\mathbf{L}, C, \mathcal{E})$ as

linear combinations of these class functions Y_i , it is described in [Lus86b, 24.4]. The input data for this algorithm are: the generalized Springer correspondence, the character values of the extensions of the F -stable χ to $W_{\mathbf{L}}F$, the orders of the $Z^0(\mathbf{L}_w)^F$ of the twisted groups \mathbf{L}_w , and the dimensions of the classes C' in the image of the Springer correspondence (the dimensions of classes are given in [Spa82, I.]). The algorithm does not only yield the Green functions but also the precise orders of the classes $C'_a \subset G(q)$.

What is missing to find the generalized Green functions as explicit class functions on $G(q)$ is to somehow fix the isomorphisms τ and to determine the functions Y_i corresponding to the choices made. This is complicated, but was worked out in many cases, at least for certain triples $(\mathbf{L}, C, \mathcal{E})$, by various authors, in particular by Shoji, see [Sho06], [Sho07] and the references given there. Recently, Geck [Ge20] has determined the ordinary Green functions for larger rank exceptional types in small characteristic, only type E_8 was missing and this will be handled in [L20].

So far, the Y_i were not yet determined for: Spin-groups in odd characteristic (the part that does not appear in SO), SU for small primes (depending on the rank, for non-ordinary Green functions [Sho06, Thm. 4.4]), exceptional groups (the part that does not appear in the adjoint type).


For computations we use our own implementation of the Springer correspondence and of the reduction to simply connected groups. For an overview of basic computations with root data see [BL13, 2.].

4 Examples for $G(q) = \text{Spin}_8(q)$, q odd

In this section we consider the untwisted simple simply connected groups \mathbf{G} of type D_4 in odd characteristic. The fixed point groups under the Frobenius are isomorphic to the spin groups $G(q) = \mathbf{G}^F = \text{Spin}_8(q)$. We will also consider some Levi subgroups \mathbf{M} of \mathbf{G} of semisimple rank 3.

We are interested in the generalized Green functions of $G(q)$ and $M(q)$ and the 2-parameter Green functions associated to $M(q) \leq G(q)$.

We label the Dynkin diagram of \mathbf{G} as follows:



Corresponding to the graph automorphism which permutes the nodes $(1, 2, 4)$ cyclically, there is also an automorphism of \mathbf{G} of order 3.

We choose this example to contribute to a project to compute the full character table for the groups $\text{Spin}_8(q)$. Also, we wanted to compare the method described here to another approach described in [RM20]. That alternative method is more elementary but needs some tricky computations with explicit unipotent elements.

4.1 Generalized Springer correspondence

We will need the generalized Springer correspondence for simple groups of type A and D , this was determined in [Lus84] and [LS85].

For type D_4 there are four triples $(\mathbf{L}, C, \mathcal{E})$ to consider, \mathbf{L} is a maximal torus or a Levi subgroup corresponding to one of the subsets of nodes $\{1, 4\}$, $\{2, 4\}$ and $\{1, 2\}$. In the first case $C = \{1\}$ and \mathcal{E} is constant. In the other cases C is the regular unipotent class of \mathbf{L} and for $u \in C$ we have $A_{\mathbf{L}}(u)$ is cyclic of order 2. Here \mathcal{E} corresponds to the non-trivial character of $A_{\mathbf{L}}(u)$.

\mathbf{G} has 12 unipotent conjugacy classes C' , they are all F -stable and contain representatives $v \in C' \cap G(q)$ such that F acts trivially on $A_{\mathbf{G}}(v)$. For three classes $A_{\mathbf{G}}(v) \cong Z_2 \times Z_2$ is non-cyclic of order 4, for seven classes $A_{\mathbf{G}}(v) \cong Z_2$ is of order 2, and for the remaining two classes $A_{\mathbf{G}}(v)$ is trivial. So, altogether there are 28 unipotent conjugacy classes in $G(q)$.

The relative Weyl groups $W_{\mathbf{L}}$ for \mathbf{L} not a torus are all of type B_2 with Dynkin diagram $\bullet \leftarrow \bullet$; here the left node 1 corresponds in the Dynkin diagram of D_4 to node 2, 1, 3, when \mathbf{L} is defined by the nodes $\{1, 4\}$, $\{2, 4\}$, $\{1, 2\}$, respectively (note that this is not symmetric under the triality automorphism, the third case corresponds to the SO_8 quotient of \mathbf{G} and the other two to the two half spin quotients of \mathbf{G} which in rank 4 are isomorphic to SO_8).

The Frobenius induces the identity map on all $W_{\mathbf{L}}$. So we get 13 (ordinary) Green functions and three times 5 generalized Green functions for the cases where \mathbf{L} is of type $A_1 + A_1$.

4.2 The Y_{ι} -functions

The spin group \mathbf{G} has a non-cyclic center of order 4. For any subgroup $Z = \langle z \rangle$ of order 2 of this center the factor group is $\mathbf{G}/Z \cong SO_8(\overline{\mathbb{F}}_q)$.

For groups of type $\mathbf{H} = SO_N$ Shoji [Sho07, Thm. 4.2] showed the following: First he chooses for each unipotent class C' a specific $H(q)$ -class $C'_1 \subseteq (C')^F$ (its elements are called split elements). For a cuspidal pair in $(\mathbf{L}, C, \mathcal{E})$ one can now normalize the isomorphism $\tau : F^*\mathcal{E} \rightarrow \mathcal{E}$ by the condition that the characteristic function of τ has a specific positive real value on split elements. Shoji shows that in this case the values of all Y_{ι} (where $\iota = (C', \mathcal{E}')$ is in the image of the $(\mathbf{L}, C, \mathcal{E})$ -part of the Springer correspondence) is $\chi(1)$ on split elements of C' where $\chi \in \text{Irr}(A_{\mathbf{H}}(u))$ corresponds to \mathcal{E}' . In other words, when choosing $u \in (C')^F$ as split element then the scalar in equation (7) is always $c = 1$.

Unfortunately, a similar description is not available for spin groups. But we can deduce much from the SO_8 -case. If for a unipotent class C' in \mathbf{G} with $u \in C'$ the image of z in $A_{\mathbf{G}}(u)$ is trivial, then the image of u in \mathbf{G}/Z has the same component group. In this case we can find a split element in $(C')^F$ as a preimage of a split element in \mathbf{G}/Z . If the component group of u has order 2, then only one of the three possible elements z has trivial image in the component group, so a split element in C' is well defined. But there are also three classes with component

group of order 4 and in these cases the image of the center is equal to $A_{\mathbf{G}}(u)$. If we choose two different elements z as above then there is a unique $G(q)$ class in $(C')^F$ which is mapped to the split class of $\mathbf{G}/\langle z \rangle$ in both cases. It is not clear if this class is also mapped to split elements in $\mathbf{G}/\langle z \rangle$ for the third possibility of z . To conclude: We have a similar statement about the Y_ℓ for \mathbf{G} as for SO_8 , except that for the three unipotent classes with component group of order 4 one of the non-trivial Y_ℓ is only determined up to sign.

4.3 Levi subgroups \mathbf{M} of type $A_1 + A_1 + A_1$

We consider the Levi subgroups \mathbf{M} which correspond to the subset $\{1, 2, 4\}$ in the Dynkin diagram of \mathbf{G} . Here $W_{\mathbf{M}} = N_{\mathbf{G}}(\mathbf{M})/\mathbf{M}$ is of order 2, so that we have two $G(q)$ -conjugacy classes of F -stable conjugates of \mathbf{M} . The tables of generalized Green functions are the same for both types, we get them from the generalized Green functions for $\mathrm{SL}_2(q)$ via the simply connected covering of $\mathbf{M}/Z^0(\mathbf{M})$ by a direct product of three copies of SL_2 . The triples $(\mathbf{L}, C, \mathcal{E})$ with cuspidal pairs are the same as for \mathbf{G} , so \mathbf{L} is a maximal torus or one of the subgroups corresponding to the subgraphs $\{1, 4\}$, $\{2, 4\}$, $\{1, 2\}$ of the Dynkin diagram. We remark that the split element chosen to normalize the isomorphism $\tau : F^*\mathcal{E} \rightarrow \mathcal{E}$ is the same in the cases of SL and Spin (the class of $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ in each SL_2 -component). The Weyl group of \mathbf{M} is elementary abelian of order 8, so we get 8 ordinary Green functions. The relative Weyl group $N_{\mathbf{M}}(\mathbf{L})/\mathbf{L}$ is of order 2, so for each \mathbf{L} of type $A_1 + A_1$ we get 2 generalized Green functions. Together we have 14 unipotent classes and generalized Green functions for each $M(q)$.

The matrix of values of generalized Green functions is easy to invert using (3). To find the 2-parameter Green functions with (5) we need to determine the fusion of the F -conjugacy classes in $N_{\mathbf{M}}(\mathbf{L})/\mathbf{L}$ into the F -conjugacy classes of $W_{\mathbf{L}}$ in each case.

We computed the 2-parameter Green functions first for the split version of \mathbf{M} and introduced three indeterminates a_{10} , a_{22} and a_{27} for the unknown signs in three Y_ℓ . Then we could use the positivity and integrality property (2) to determine the values of these indeterminates. For example, the value

$$g_M^G((2.2.2, 4), (3221, 4)) = \frac{1}{4}(a_{10} - 1)(q^4 + q^3)$$

shows that $a_{10} = 1$ (otherwise, the expression above would be negative).

For c, c' both regular there is a value

$$\frac{1}{4}(-a_{27} + 1)$$

which shows that $a_{27} = 1$ (otherwise the value would not be integral).

And for a c of type (53) and c' regular we find a value

$$\frac{1}{4}(q + a_{22} - 2)$$

This shows that $a_{22} = 1$ if $q \equiv 1 \pmod{4}$ and $a_{22} = -1$ if $q \equiv 3 \pmod{4}$. So, the values of the generalized Green functions for $\text{Spin}_8(q)$ depend on the congruence class of q modulo 4 (while everything we did so far was only depending on the congruence modulo 2).

To conclude: we have found the generalized Green functions for $\text{Spin}_8(q)$ for odd q , and can now compute the 2-parameter Green functions for the split and non-split version of \mathbf{M} .

4.4 Tables

Here are tables of 2-parameter Green functions. The rows are labelled by the unipotent classes of $M(q)$. They are described in terms of the Jordan block sizes in the three SL_2 -components, this determines the class in the algebraic group \mathbf{M} , and a number which counts the $M(q)$ -classes of such elements; the first corresponds to the class of split elements.

The columns are labelled by the unipotent classes of $G(q)$. Here we use the Jordan block sizes of images in SO_8 . In some cases there are two classes with the same Jordan block sizes, we add a + character if the class has representatives in the Levi subgroup of the subdiagram $\{1, 3, 4\}$ and a - otherwise. Again, there is also a counter for the $G(q)$ -classes in a \mathbf{G} -class, and the first indicates the class of split elements.

Many values of the 2-parameter Green functions factorize with cyclotomic polynomials as factors. In the tables we print P_i for the i -th cyclotomic polynomial evaluated at q (so $P_1 = (q - 1)$, $P_2 = (q + 1)$, and so on).

The first table is the case of split \mathbf{M} of type $A_1 + A_1 + A_1$ with $|Z^0(\mathbf{M})^F| = q - 1$, and the second is the twisted version of \mathbf{M} with $|Z^0(\mathbf{M})^F| = q + 1$.

Open question

For the generalized Green functions corresponding to Levi subgroups of type $A_1 + A_1$ the identification in (4) was only shown for large enough q .

$$M(q) = A_1(q) + A_1(q) + A_1(q) + T(q-1) \leq \text{Spin}_8(q), \quad a_{22} \in \{-1, 1\} \text{ and } a_{22} = q \pmod{4}$$

11

	11111111,1	221111,1	2222+,1	2222+,2	2222-,1	2222-,2	311111,1	311111,2	3221,1	3221,2	3221,3	3221,4	3311,1	3311,2
11.11.11,1	P2P3P4^2P6	q^4+3*q^3+3*q^2+q+1	P2P4	P2P4	P2P4	P2P4	P2P4	P2P4	P2	P2	P2	P2	1	1
11.11.2,1	.	q^4P2	q^2P4	q^2P4	2*q	.
11.2.11,1	.	q^4P2	q^2P4	q^2P4	2*q	.
11.2.2,1	q^2P2P4	.	.	.	q^2	q^2	.	.	qP1	.
11.2.2,2	q^2P2P4	q^2	q^2	qP1	.
2.11.11,1	.	q^4P2	.	.	q^2P4	q^2P4	2*q	.
2.11.2,1	.	.	q^2P2P4	q^2	.	.	q^2	qP1	.
2.11.2,2	.	.	.	q^2P2P4	q^2	q^2	q^2	.	qP1	.
2.2.11,1	q^2P2P4	.	q^2	.	q^2	.	qP1	.
2.2.11,2	q^2P2P4	.	q^2	.	q^2	qP1	.
2.2.2,1	q^3P2	.	.	.	1/4*qP1^2	1/4*qP2^2
2.2.2,2	q^3P2	.	1/4*qP1^2	1/4*qP2^2
2.2.2,3	q^3P2	1/4*qP1^2	1/4*qP2^2
2.2.2,4	q^3P2	.	.	.	1/4*qP1^2	1/4*qP2^2
	44+,1	44+,2	44-,1	44-,2	5111,1	5111,2	53,1	53,2	53,3	53,4	71,1	71,2	71,3	71,4
11.11.11,1
11.11.2,1	1	1
11.2.11,1	1	1
11.2.2,1	.	.	P2	.	.	.	1	1
11.2.2,2	.	.	.	P2	1
2.11.11,1	.	.	1	1
2.11.2,1	P2	1	.	.	.	1	.	.	.
2.11.2,2	.	P2	1	1
2.2.11,1	P2	.	1/2*(a22+1)	1/2*(-a22+1)	1/2*(a22+1)	1/2*(-a22+1)	1/2*(a22+1)	.	.	.
2.2.11,2	P2	1/2*(-a22+1)	1/2*(a22+1)	1/2*(-a22+1)	1/2*(a22+1)	.	.	.
2.2.2,1	1/2*P1P2	.	1/2*P1P2	.	1/2*P1P2	.	1/4*(q-a22-4)	1/4*(q+a22-2)	1/4*(q-a22)	1/4*(q+a22-2)	1/4*(q-a22)	1/4*(q+a22-2)	1	.
2.2.2,2	.	1/2*P1P2	.	1/2*P1P2	1/2*P1P2	.	1/4*(q-a22)	1/4*(q+a22-2)	1/4*(q-a22-4)	1/4*(q+a22-2)	1/4*(q-a22-4)	1/4*(q+a22-2)	.	1
2.2.2,3	1/2*P1P2	.	.	1/2*P1P2	.	1/2*P1P2	1/4*(q+a22-2)	1/4*(q-a22)	1/4*(q+a22-2)	1/4*(q-a22-4)	1/4*(q-a22-4)	.	.	1
2.2.2,4	.	1/2*P1P2	1/2*P1P2	.	.	1/2*P1P2	1/4*(q+a22-2)	1/4*(q-a22-4)	1/4*(q+a22-2)	1/4*(q-a22)	1/4*(q-a22)	.	1	.

$$M(q) = A_1(q) + A_1(q) + A_1(q) + T(q+1) \leq \text{Spin}_8(q), \quad a_{22} \in \{-1, 1\} \text{ and } a_{22} = q \pmod{4}$$

12

	11111111,1	221111,1	2222+,1	2222+,2	2222-,1	2222-,2	311111,1	311111,2	3221,1	3221,2	3221,3	3221,4	3311,1	3311,2
11.11.11,1	-P1P3P4^2P6	q^4-3*q^3+3*q^2-q+1	-P1P4	-P1P4	-P1P4	-P1P4	-P1P4	-P1P4	-P1	-P1	-P1	-P1	1	1
11.11.2,1	.	-q^4P1	q^2P4	q^2P4	-2*q
11.2.11,1	.	-q^4P1	q^2P4	q^2P4	-2*q
11.2.2,1	-q^2P1P4	.	.	.	q^2	q^2	.	.	.	qP2
11.2.2,2	-q^2P1P4	q^2	q^2	.	qP2
2.11.11,1	.	-q^4P1	.	.	q^2P4	q^2P4	-2*q
2.11.2,1	.	.	-q^2P1P4	q^2	.	.	q^2	.	qP2
2.11.2,2	.	.	.	-q^2P1P4	q^2	q^2	.	.	qP2
2.2.11,1	-q^2P1P4	.	q^2	.	q^2	.	.	qP2
2.2.11,2	-q^2P1P4	.	q^2	.	q^2	.	qP2
2.2.2,1	q^3P1	.	.	.	-1/4*qP1^2	-1/4*qP2^2
2.2.2,2	q^3P1	.	-1/4*qP1^2	-1/4*qP2^2
2.2.2,3	q^3P1	-1/4*qP1^2	-1/4*qP2^2
2.2.2,4	q^3P1	.	.	.	-1/4*qP1^2	-1/4*qP2^2
	44+,1	44+,2	44-,1	44-,2	5111,1	5111,2	53,1	53,2	53,3	53,4	71,1	71,2	71,3	71,4
11.11.11,1
11.11.2,1	1	1
11.2.11,1	1	1
11.2.2,1	.	.	.	-P1	1	.	1	.	.	.
11.2.2,2	.	.	-P1	.	.	.	1	1
2.11.11,1	.	.	1	1
2.11.2,1	.	-P1	1	1
2.11.2,2	-P1	1	.	.	.	1	.	.	.
2.2.11,1	-P1	1/2*(-a22+1)	1/2*(a22+1)	1/2*(-a22+1)	1/2*(a22+1)	1/2*(a22+1)	.	.	.
2.2.11,2	-P1	.	1/2*(a22+1)	1/2*(-a22+1)	1/2*(a22+1)	1/2*(-a22+1)	1/2*(a22+1)	.	.	.
2.2.2,1	.	1/2*P1P2	.	1/2*P1P2	.	1/2*P1P2	1/4*(-q+a22)	1/4*(-q-a22-2)	1/4*(-q+a22-4)	1/4*(-q-a22-2)	1	.	.	.
2.2.2,2	1/2*P1P2	.	1/2*P1P2	.	.	1/2*P1P2	1/4*(-q+a22-4)	1/4*(-q-a22-2)	1/4*(-q+a22)	1/4*(-q-a22-2)	.	.	1	.
2.2.2,3	.	1/2*P1P2	1/2*P1P2	.	1/2*P1P2	.	1/4*(-q-a22-2)	1/4*(-q+a22-4)	1/4*(-q-a22-2)	1/4*(-q+a22)	.	.	.	1
2.2.2,4	1/2*P1P2	.	.	1/2*P1P2	1/2*P1P2	.	1/4*(-q-a22-2)	1/4*(-q+a22)	1/4*(-q-a22-2)	1/4*(-q+a22-4)	.	1	.	.

References

- [BL13] O. Brunat and F. Lübeck. “On defining characteristic representations of finite reductive groups”. *J. Algebra* 395 (2013), pp. 121–141.
- [DL76] P. Deligne and G. Lusztig. “Representations of reductive groups over finite fields”. *Ann. of Math. (2)* 103.1 (1976), pp. 103–161.
- [DM91] F. Digne and J. Michel. *Representations of finite groups of Lie type*. Vol. 21. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1991, pp. iv+159.
- [Ge20] M. Geck. “Computing Green functions in small characteristic”. *J. Algebra*, to appear (2020).
- [L20] F. Lübeck. “Green functions of $E_8(q)$ in small characteristic” (in preparation).
- [LS85] G. Lusztig and N. Spaltenstein. “On the generalized Springer correspondence for classical groups”. *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*. Vol. 6. Adv. Stud. Pure Math. North-Holland, Amsterdam, 1985, pp. 289–316.
- [Lus12] G. Lusztig. “On the cleanness of cuspidal character sheaves”. *Mosc. Math. J.* 12.3 (2012), pp. 621–631, 669.
- [Lus19] G. Lusztig. “On the generalized Springer correspondence”. *Representations of reductive groups*. Vol. 101. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2019, pp. 219–253.
- [Lus76] G. Lusztig. “On the finiteness of the number of unipotent classes”. *Invent. Math.* 34.3 (1976), pp. 201–213.
- [Lus84] G. Lusztig. “Intersection cohomology complexes on a reductive group”. *Invent. Math.* 75.2 (1984), pp. 205–272.
- [Lus85] G. Lusztig. “Character sheaves. II, III”. *Adv. in Math.* 57.3 (1985), pp. 226–265, 266–315.
- [Lus86a] G. Lusztig. “Character sheaves. IV”. *Adv. in Math.* 59.1 (1986), pp. 1–63.
- [Lus86b] G. Lusztig. “Character sheaves. V”. *Adv. in Math.* 61.2 (1986), pp. 103–155.
- [Lus90] G. Lusztig. “Green functions and character sheaves”. *Ann. of Math. (2)* 131.2 (1990), pp. 355–408.
- [RM20] G. Malle and E. Rotilio. *The 2-parameter Green functions for 8-dimensional spin groups*. 2020. arXiv: [2003.14231 \[math.RT\]](https://arxiv.org/abs/2003.14231).
- [Sho06] T. Shoji. “Generalized Green functions and unipotent classes for finite reductive groups. I”. *Nagoya Math. J.* 184 (2006), pp. 155–198.

- [Sho07] T. Shoji. “Generalized Green functions and unipotent classes for finite reductive groups. II”. *Nagoya Math. J.* 188 (2007), pp. 133–170.
- [Sho95] T. Shoji. “Character sheaves and almost characters of reductive groups. I, II”. *Adv. Math.* 111.2 (1995), pp. 244–313, 314–354.
- [Sho96] T. Shoji. “On the computation of unipotent characters of finite classical groups”. Vol. 7. 3. *Computational methods in Lie theory* (Essen, 1994). 1996, pp. 165–174.
- [Sho97] T. Shoji. “Unipotent characters of finite classical groups”. *Finite reductive groups (Luminy, 1994)*. Vol. 141. *Progr. Math.* Birkhäuser Boston, Boston, MA, 1997, pp. 373–413.
- [Spa82] N. Spaltenstein. *Classes unipotentes et sous-groupes de Borel*. Vol. 946. *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1982, pp. ix+259.
- [Spa85] N. Spaltenstein. “On the generalized Springer correspondence for exceptional groups”. *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*. Vol. 6. *Adv. Stud. Pure Math.* North-Holland, Amsterdam, 1985, pp. 317–338.