

# The Differential Geometry of the Orbit space of Extended Affine Jacobi Group $A_n$

Guilherme F. Almeida

SISSA, via Bonomea 265, Trieste, Italy

November 1, 2021

Dedicated to the memory of Professor Boris Dubrovin.

## Abstract

We define certain extensions of Jacobi groups of  $A_n$ , prove an analogue of Chevalley Theorem for their invariants, and construct a Dubrovin Frobenius structure on it orbit space.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Topological quantum field theory . . . . .	2
1.2	Orbit space of reflection groups and its extensions . . . . .	3
1.3	Hurwitz space/ Orbit space correspondence . . . . .	3
1.4	Results . . . . .	4
<b>2</b>	<b>Review of Dubrovin-Frobenius manifolds</b>	<b>9</b>
2.1	Basic definitions . . . . .	9
2.2	Intersection form . . . . .	11
2.3	Reconstruction . . . . .	12
2.4	Monodromy of Dubrovin-Frobenius manifold . . . . .	12
2.5	Dubrovin Connection . . . . .	12
2.6	Differential geometry preliminaries . . . . .	13
2.7	Semisimple Dubrovin Frobenius manifolds . . . . .	15
<b>3</b>	<b>Review of Dubrovin-Frobenius manifold on Hurwitz spaces</b>	<b>17</b>
3.1	Hurwitz spaces . . . . .	17
3.2	Bidifferential $W$ . . . . .	18
3.3	Reconstruction of Dubrovin Frobenius manifold . . . . .	19

---

<b>4</b>	<b>Differential geometry of the orbit space of extended Jacobi group <math>A_n</math></b>	<b>20</b>
4.1	The Group $\mathcal{J}(\tilde{A}_n)$ . . . . .	20
4.2	Jacobi forms of $\mathcal{J}(\tilde{A}_n)$ . . . . .	22
4.3	Proof of the Chevalley theorem . . . . .	29
4.4	Intersection form . . . . .	33
4.5	The second metric of the pencil . . . . .	41
4.6	Flat coordinates of $\eta$ . . . . .	49
4.7	The extended ring of Jacobi forms . . . . .	56
4.8	Christoffel symbols of the intersection form . . . . .	58
4.9	Unit and Euler vector field of the orbit space of $\mathcal{J}(\tilde{A}_n)$ . . . . .	64
4.10	Discriminant locus and the monodromy of the orbit space of $\mathcal{J}(\tilde{A}_n)$ . . . . .	68
4.11	Construction of WDVV solution . . . . .	70

# 1 Introduction

Dubrovin Frobenius manifold is a geometric interpretation of a remarkable system of differential equations, called WDVV equations [10]. Since the beginning of the nineties, there has been a continuous exchange of ideas from fields that are not trivially related to each other, such as: Topological quantum field theory, non-linear waves, singularity theory, random matrices theory, integrable systems, and Painleve equations. Dubrovin Frobenius manifolds theory is a bridge between them.

## 1.1 Topological quantum field theory

The connections made by Dubrovin Frobenius manifolds theory works because all the mentioned theories are related with some WDVV equation. In [10], Dubrovin showed that many constructions of Topological field theories (TFT) can be deduced from the geometry of Dubrovin-Frobenius manifolds. For instance, one of the main objects to be computed in TFT are correlation functions, which are mean values of physical quantities. Since in TFT it is possible to have infinite many correlation functions, an efficient way to compute all of them is encoding all the correlators in a single function, called partition function. In [19], Konsevitch proved that a partition function of a specific TFT can be obtained from the solution of KdV hierarchy, which is an example of integrable hierarchy, i.e. an infinite list of integrable partial differential equations. This discovery opened a new field of research in mathematical physics, because for this case, it was found an effective way to compute exactly all the correlation functions due to its integrable system nature. In [14], Dubrovin and Zhang constructed a method to derive an integrable hierarchy from the data of Dubrovin Frobenius manifold, furthermore, in many important examples, they were able to relate these integrable hierarchy with partition functions of some TFT.

## 1.2 Orbit space of reflection groups and its extensions

In [10], Dubrovin point out that, WDVV solutions with certain good analytic properties are related with partition functions of TFT. Afterwards, Dubrovin conjectured that WDVV solutions with certain good analytic properties are in one to one correspondence with discrete groups. This conjecture is supported in ideas which come from singularity theory, because in this setting there exist a integrable systems/ discrete group correspondence. Furthermore, in minimal models as Gepner chiral rings there exist a correspondence between physical models and discrete groups. In [18], Hertling proved that a particular class of Dubrovin-Frobenius manifold, called polynomial Dubrovin-Frobenius manifold is isomorphic to orbit space of a finite Coxeter group, which are spaces such that its geometric structure is invariant under the finite Coxeter group. In [6], [7],[9], [10], [12], [13], [33], there are many examples of WDVV solutions that are associated with orbit spaces of natural extensions of finite Coxeter groups such as extended affine Weyl groups, and Jacobi groups. Therefore, the construction of Dubrovin Frobenius manifolds on orbit space of reflection groups and its extensions is a prospective project of the classification of WDVV solutions. In addition, WDDV solutions arising from orbit spaces may also have some applications in TFT or some combinatorial problem, because previously these relation was demonstrated in some examples such as the orbit space of the finite Coxeter group  $A_1$ , and the extended affine Weyl group  $A_1$  [11], [14].

## 1.3 Hurwitz space/ Orbit space correspondence

There are several others non-trivial connections that Dubrovin Frobenius manifolds theory can make. For example, Hurwitz spaces is the one of the main source of examples of Dubrovin Frobenius manifolds. Hurwitz spaces  $H_{g,n_0,n_1,\dots,n_m}$  are moduli space of covering over  $\mathbb{CP}^1$  with a fixed ramification profile. More specifically,  $H_{g,n_0,n_1,\dots,n_m}$  : is moduli space of pairs

$$\{C_g, \lambda : C_g \mapsto \mathbb{CP}^1\}$$

where  $C_g$  is a compact Riemann surface of genus  $g$ ,  $\lambda$  is meromorphic function with poles in

$$\lambda^{-1}(\infty) = \{\infty_0, \infty_1, \dots, \infty_m\}.$$

Moreover,  $\lambda$  has degree  $n_i + 1$  near  $\infty_i$ . Hurwitz space with a choice of specific Abelian differential, called quasi-momentum or primary differential, give rise to a Dubrovin Frobenius manifold, see section 3 for details. In some examples, the Dubrovin Frobenius structure of Hurwitz spaces are isomorphic to Dubrovin Frobenius manifolds associated with orbit spaces of suitable groups. For instance, the orbit space of the finite Coxeter group  $A_n$  is isomorphic to the Hurwitz space  $H_{0,n}$ , furthermore, orbit space of the extended affine Weyl group  $\tilde{A}_n$  and of the Jacobi group  $\mathcal{J}(A_n)$  are isomorphic to the Hurwitz spaces  $H_{0,n-1,0}$  and  $H_{1,n}$  respectively. Motivated by these examples, we construct the following diagram

$$\begin{array}{ccc} H_{0,n} \cong \text{Orbit space of } A_n & \xrightarrow{1} & H_{0,n-1,0} \cong \text{Orbit space of } \tilde{A}_n \\ \downarrow 2 & & \downarrow 4 \\ H_{1,n} \cong \text{Orbit space of } \mathcal{J}(A_n) & \xrightarrow{3} & H_{1,n-1,0} \cong ? \end{array}$$

---

From the Hurwitz space side, the vertical lines 2 and 4 mean that we increase the genus by 1, and the horizontal line means that we split one pole of order  $n + 1$  in a simple pole and a pole of order  $n$ . From the orbit space side, the vertical line 2 means that we are doing an extension from the finite Coxeter group  $A_n$  to the Jacobi group  $\mathcal{J}(A_n)$ , the line horizontal line 1 means that we are extending the Orbit space of  $A_n$  to the extended affine Weyl group  $\tilde{A}_n$ . Therefore, one might ask if the line 3 and 4 would imply an orbit space interpretation of the Hurwitz space  $H_{1,n-1,0}$ . The main goal of this thesis is to define a new class of groups such that its orbit space carries Dubrovin-Frobenius structure of  $H_{1,n-1,0}$ . The new group is called extended affine Jacobi group  $A_n$ , and denoted by  $\mathcal{J}(\tilde{A}_n)$ . This group is an extension of the Jacobi group  $\mathcal{J}(A_n)$ , and of the extended affine Weyl group  $\tilde{A}_n$ .

## 1.4 Results

The main goal of this thesis is to derive the Dubrovin Frobenius structure of the Hurwitz space  $H_{1,n-1,0}$  from the data of the group  $\mathcal{J}(\tilde{A}_n)$ . First of all, we define the group  $\mathcal{J}(\tilde{A}_n)$ . Recall that the group  $A_n$  acts on  $\mathbb{C}^n$  by permutations, then the group  $\mathcal{J}(\tilde{A}_n)$  is an extension of the group  $A_n$  in the following sense:

**Proposition 1.1.** The group  $\mathcal{J}(\tilde{A}_n) \ni (w, t, \gamma)$  acts on  $\Omega := \mathbb{C} \oplus \mathbb{C}^{n+2} \oplus \mathbb{H} \ni (u, v, \tau)$  as follows

$$\begin{aligned} w(u, v, \tau) &= (u, wv, \tau) \\ t(u, v, \tau) &= (u - \langle \lambda, v \rangle_{\tilde{A}_n} - \frac{1}{2} \langle \lambda, \lambda \rangle_{\tilde{A}_n} \tau, v + \lambda \tau + \mu, \tau) \\ \gamma(u, v, \tau) &= (u + \frac{c \langle v, v \rangle_{\tilde{A}_n}}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{a\tau + b}{c\tau + d}) \end{aligned} \tag{1.1}$$

where  $w \in A_n$  acts by permutations in the first  $n + 1$  variables of  $\mathbb{C}^{n+2} \ni v = (v_0, v_1, \dots, v_n, v_{n+1})$ ,

$$\begin{aligned} t &= (\lambda, \mu) \in \mathbb{Z}^{n+2}, \\ \begin{pmatrix} a & b \\ c & d \end{pmatrix} &\in SL_2(\mathbb{Z}), \\ \langle v, v \rangle_{\tilde{A}_n} &= \sum_{i=0}^n v_i^2 \Big|_{\sum v_i=0} - n(n+1)v_{n+1}^2. \end{aligned}$$

See section 4.1 for details.

In order to define any geometric structure in a orbit space, first it is necessary to define a notion of invariant  $\mathcal{J}(\tilde{A}_n)$  sections. For this purpose, we generalise the ring of invariant functions used in [6], [7] for the group  $\mathcal{J}(A_n)$ , which are called Jacobi forms. This notion was first defined in [16] by Eichler and Zagier for the group  $\mathcal{J}(A_1)$ , and further it was generalised for the group  $\mathcal{J}(A_n)$  in [32] by Wirthmuller. Furthermore, an explicit base of generators were derived in [6], [7] by Bertola. The Jacobi forms used in this thesis are defined by

**Definition 1.1.** The weak  $\tilde{A}_n$ -invariant Jacobi forms of weight  $k$ , order  $l$ , and index  $m$  are functions

on  $\Omega = \mathbb{C} \oplus \mathbb{C}^{n+2} \oplus \mathbb{H} \ni (u, v_0, v_1, \dots, v_{n+1}, \tau) = (u, v, \tau)$  which satisfy

$$\begin{aligned} \varphi(w(u, v, \tau)) &= \varphi(u, v, \tau), \quad A_n \text{ invariant condition} \\ \varphi(t(u, v, \tau)) &= \varphi(u, v, \tau) \\ \varphi(\gamma(u, v, \tau)) &= (c\tau + d)^{-k} \varphi(u, v, \tau) \\ E\varphi(u, v, \tau) &:= -\frac{1}{2\pi i} \frac{\partial}{\partial u} \varphi(u, v, \tau) = m\varphi(u, v, \tau), \quad \text{Euler vector field} \end{aligned} \tag{1.2}$$

Moreover, the weak  $\tilde{A}_n$ -invariant Jacobi forms are meromorphic in the variable  $v_{n+1}$  on a fixed divisor, in contrast with the Jacobi forms of the group  $\mathcal{J}(A_n)$ , which are holomorphic in each variable, see details on the definition 4.2. The ring of weak  $\tilde{A}_n$ -invariant Jacobi forms gives the notion of Euler vector field, indeed, the vector field defined in the last equation of (1.2) measures the degree of Jacobi form, which coincides with the index. The differential geometry of the orbit space of the group  $\mathcal{J}(\tilde{A}_n)$  should be understood as the space such that its sections are written in terms of Jacobi forms. Then, in order for this statement to make sense, we prove a Chevalley type theorem, which is

**Theorem 1.2.** The trigraded algebra of Jacobi forms  $J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)} = \bigoplus_{k, l, m} J_{k, l, m}^{\tilde{A}_n}$  is freely generated by  $n+1$  fundamental Jacobi forms  $(\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n)$  over the graded ring  $E_{\bullet, \bullet}$ .

$$J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)} = E_{\bullet, \bullet}[\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n], \tag{1.3}$$

where

$$E_{\bullet, \bullet} = J_{\bullet, \bullet, 0} \text{ is the ring of coefficients.}$$

More specifically, the ring of function  $E_{\bullet, \bullet}$  is the space of functions  $f(v_{n+1}, \tau)$  such that for fixed  $\tau$ , the functions  $\tau \mapsto f(v_{n+1}, \tau)$  is an elliptic function.

Moreover,  $(\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n)$  are given by

**Corollary 1.2.1.** The functions  $(\varphi_0, \varphi_1, \dots, \varphi_n)$  obtained by the formula

$$\begin{aligned} \lambda^{\tilde{A}_n} &= e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(z - v_i + v_{n+1}, \tau)}{\theta_1^n(z, \tau) \theta_1(z + (n+1)v_{n+1}, \tau)} \\ &= \varphi_n \wp^{n-2}(z, \tau) + \varphi_{n-1} \wp^{n-3}(z, \tau) + \dots + \varphi_2 \wp(z, \tau) \\ &\quad + \varphi_1 [\zeta(z, \tau) - \zeta(z + (n+1)v_{n+1}, \tau) + \zeta((n+1)v_{n+1})] + \varphi_0 \end{aligned} \tag{1.4}$$

are Jacobi forms of weight  $0, -1, -2, \dots, -n$  respectively, index 1, and order 0. Here  $\theta_1$  is the Jacobi theta 1 function and  $\zeta$  is the Weierstrass zeta function.

This lemma realises the functions  $(\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n, v_{n+1}, \tau)$  as coordinates of the orbit space of  $\mathcal{J}(\tilde{A}_n)$ . The unit vector field is chosen to be

$$e = \frac{\partial}{\partial \varphi_0}, \tag{1.5}$$

because  $\varphi_0$  is the basic generator with maximum weight degree, see the sections 4.2, 4.3 for details. The last ingredient we need to construct is the flat pencil metric associated with the orbit space of  $\mathcal{J}(\tilde{A}_n)$ ,

which is two compatible flat metric  $g^*$  and  $\eta^*$ , which are sections of the cotangent bundle of the orbit space of  $\mathcal{J}(\tilde{A}_n)$  such that

$$g^* + \lambda \eta^*$$

is also flat, and the linear combination of its Christoffel symbols

$$\Gamma_{g^*}^{ij} + \lambda \Gamma_{\eta^*}^{ij}$$

is the Christoffel symbol of the flat pencil of metrics (see section 2.6 for the details) [9], [10]. The natural candidate to be one of the metrics of the pencil is the invariant metric of the group  $\mathcal{J}(\tilde{A}_n)$ . This metric is given by

$$g^* = \sum_{i,j} A_{ij}^{-1} \frac{\partial}{\partial v_i} \otimes \frac{\partial}{\partial v_j} - n(n+1) \frac{\partial}{\partial v_{n+1}} \otimes \frac{\partial}{\partial v_{n+1}} + \frac{\partial}{\partial \tau} \otimes \frac{\partial}{\partial u} + \frac{\partial}{\partial u} \otimes \frac{\partial}{\partial \tau}, \quad (1.6)$$

where

$$A_{ij} = \begin{pmatrix} 2 & 1 & 1 & \dots & 1 \\ 1 & 2 & 1 & \dots & 1 \\ 1 & 1 & 2 & \dots & 1 \\ 1 & 1 & 1 & \dots & 1 \\ 1 & 1 & 1 & \dots & 2 \end{pmatrix}$$

This metric is called intersection form. The second metric is given by

$$\eta^* := Lie_e g^*,$$

and it is denoted by Saito metric due to K.Saito, who was the first to define this metric for the case of finite Coxeter group [26]. One of the main technical problems of the thesis is to prove that the Saito metric  $\eta^*$  is flat. For this purpose, we construct a generating function for the coefficients of the metric  $\eta^*$  in the coordinates  $(\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n, v_{n+1}, \tau)$ . We prove the following.

**Corollary 1.2.2.** Let  $\eta^*(d\varphi_i, d\varphi_j)$  be given by

$$\eta^*(d\varphi_i, d\varphi_j) := \frac{\partial g^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0}. \quad (1.7)$$

The coefficient  $\eta^*(d\varphi_i, d\varphi_j)$  is recovered by the generating formula

$$\begin{aligned} & \sum_{k,j=0}^{n+1} \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \tilde{\eta}^*(d\varphi_i, d\varphi_j) \wp(v)^{(k-2)} \wp(v')^{(j-2)} = \\ & = 2\pi i (D_\tau \lambda(v) + D_\tau \lambda(v')) + \frac{1}{2} \frac{\wp'(v) + \wp'(v')}{\wp(v) - \wp(v')} \left[ \frac{d\lambda(v')}{dv'} - \frac{d\lambda(v)}{dv} \right] \\ & - \frac{1}{n(n+1)} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p)}{\partial v_{n+1}} \right) \frac{\partial \lambda(p')}{\partial v_{n+1}} - \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p')}{\partial v_{n+1}} \right) \\ & + \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_j}{\partial v_{n+1}} \right) \frac{\partial \varphi_k}{\partial v_{n+1}} \\ & + \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_k}{\partial v_{n+1}} \right), \end{aligned} \quad (1.8)$$

where

$$\begin{aligned}\tilde{\eta}^*(d\varphi_i, d\varphi_j) &= \eta^*(d\varphi_i, d\varphi_j), \quad i, j \neq 0, \\ \tilde{\eta}^*(d\varphi_0, d\varphi_j) &= \eta^*(d\varphi_i, d\varphi_j) + 4\pi i k_j \varphi_j.\end{aligned}\tag{1.9}$$

See section 4.4 for the definition of  $D_\tau$  and for further details. Thereafter, we extract the coefficients  $\eta^*(d\varphi_i, d\varphi_j)$  from the generating function (1.8)

**Theorem 1.3.** The coefficients  $\eta^*(d\varphi_i, d\varphi_j)$  can be obtained by the formula,

$$\begin{aligned}\tilde{\eta}^*(d\varphi_i, d\varphi_j) &= (i+j-2)\varphi_{i+j-2}, \quad i, j \neq 0 \\ \tilde{\eta}^*(d\varphi_i, d\varphi_0) &= 0, \quad i \neq 0, \quad i \neq 1, \\ \tilde{\eta}^*(d\varphi_1, d\varphi_j) &= 0, \quad j \neq 0, \\ \tilde{\eta}^*(d\varphi_1, d\varphi_0) &= \wp((n+1)v_{n+1})\varphi_1.\end{aligned}\tag{1.10}$$

Inspired by the construction of the flat coordinates of the Saito metric done in [26], we construct explicitly the coordinates  $t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau$  by the following formulae

$$\begin{aligned}t^\alpha &= \frac{n}{n+1-\alpha} (\varphi_n)^{\frac{n+1-\alpha}{n}} (1 + \Phi_{n-\alpha})^{\frac{n+1-\alpha}{n}}, \\ t^0 &= \varphi_0 - \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_1 + 4\pi i g_1(\tau) \varphi_2,\end{aligned}\tag{1.11}$$

where

$$\begin{aligned}(1 + \Phi_i)^{\frac{n+1-\alpha}{n}} &= \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \Phi_i^d, \\ \Phi_i^d &= \sum_{i_1+i_2+\dots+i_d=i} \frac{\varphi_{(n-i_1)}}{\varphi_n} \dots \frac{\varphi_{(n-i_d)}}{\varphi_n}.\end{aligned}\tag{1.12}$$

The Theorem 1.3 together with the formulae (1.11), and some extra auxiliary technical lemmas, implies the flatness of the Saito metric  $\eta$ .

**Theorem 1.4.** Let  $(t^0, t^1, t^2, \dots, t^n)$  be defined in (1.11), and  $\eta^*$  the Saito metric. Then,

$$\begin{aligned}\eta^*(dt^\alpha, dt^{n+3-\beta}) &= -(n+1)\delta_{\alpha\beta}, \quad 2 \leq \alpha, \beta \leq n \\ \eta^*(dt^1, dt^\alpha) &= 0, \\ \eta^*(dt^0, dt^\alpha) &= 0, \\ \eta^*(dt^i, d\tau) &= -2\pi i \delta_{i0}, \\ \eta^*(dt^i, dv_{n+1}) &= -\frac{\delta_{i1}}{n+1}.\end{aligned}\tag{1.13}$$

Moreover, the coordinates  $t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau$  are the flat coordinates of  $\eta^*$ . See details in sections 4.5 and 4.6.

A remarkable fact to point out is that, even though the intersection form  $g^*$  and their Levi Civita connection are  $\mathcal{J}(\tilde{A}_n)$  invariant sections, their coefficients  $g^{ij}, \Gamma_k^{ij}$  are not Jacobi forms, but they live in an extension of the Jacobi forms ring. Hence, we have the following lemma

---

**Lemma 1.5.** The coefficients of the intersection form  $g^{\alpha\beta}$  and its Christoffel symbol  $\Gamma_{\gamma}^{\alpha\beta}$  on the coordinates  $t^0, t^1, \dots, t^n, v_{n+1}, \tau$  belong to the ring  $\tilde{E}_{\bullet, \bullet}[t^0, t^1, \dots, t^n, \frac{1}{t^n}]$ , where  $\tilde{E}_{\bullet, \bullet}$  is a suitable extension of the ring  $E_{\bullet, \bullet}$ .

This lemma 1.5 is important because it gives a tri-graded ring structure for the coefficients  $g^{\alpha\beta}$  and  $\Gamma_{\gamma}^{\alpha\beta}$ . In particular, the lemma 1.5 implies that  $g^{\alpha\beta}$  and  $\Gamma_{\gamma}^{\alpha\beta}$  are eigenfunctions of the Euler vector field given by the last equation (1.2). Using this fact, one can prove that  $g^{\alpha\beta}$  and  $\Gamma_{\gamma}^{\alpha\beta}$  are at most linear on the variable  $t^0$ , and this fact together with Theorem 1.4 proves that  $g^*, \eta^*$  form a flat pencil metric, and consequently we can prove

**Lemma 1.6.** Let the intersection form be (1.6), the unit vector field be (1.5), and the Euler vector field be given by the last equation of (1.2). Then, there exist a function

$$F(t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau) = -\frac{(t^0)^2 \tau}{4\pi i} + \frac{t^0}{2} \sum_{\alpha, \beta \neq 0, \tau} \eta_{\alpha\beta} t^{\alpha} t^{\beta} + G(t^1, t^2, \dots, t^n, v_{n+1}, \tau), \quad (1.14)$$

such that

$$\begin{aligned} \text{Lie}_E F &= 2F + \text{quadratic terms}, \\ \text{Lie}_E (F^{\alpha\beta}) &= g^{\alpha\beta}, \\ \frac{\partial^2 G(t^1, t^2, \dots, t^n, v_{n+1}, \tau)}{\partial t^{\alpha} \partial t^{\beta}} &\in \tilde{E}_{\bullet, \bullet}[t^1, t^2, \dots, t^n, \frac{1}{t^n}], \end{aligned} \quad (1.15)$$

where

$$F^{\alpha\beta} = \eta^{\alpha\alpha'} \eta^{\beta\beta'} \frac{\partial F^2}{\partial t^{\alpha'} \partial t^{\beta'}}. \quad (1.16)$$

Using the lemma 1.6 with some more technical results, we obtain our final result

**Theorem 1.7.** A suitable covering of the orbit space  $(\mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H}) / \mathcal{J}(\tilde{A}_n)$  with the intersection form (1.6), unit vector field (1.5), and Euler vector field given by the last equation of (1.2) has a Dubrovin Frobenius manifold structure. Moreover, a suitable covering of  $\mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} / \mathcal{J}(\tilde{A}_n)$  is isomorphic as Dubrovin Frobenius manifold to a suitable covering of the Hurwitz space  $H_{1, n-1, 0}$ .

See section 4.11 for details.

The results of this thesis would be important because of the following

1. The Hurwitz spaces  $H_{1, n-1, 0}$  are classified by the group  $\mathcal{J}(\tilde{A}_n)$ , hence we increase the knowledge of the WDVV/ discrete group correspondence. In particular the WDVV solutions associated with this orbit spaces contains a kind of elliptic function in an exceptional variable, which is exotic in theory of WDVV solutions, since most of the known examples are polynomial or polynomial with exponential function. Recently, the case  $\mathcal{J}(\tilde{A}_1)$  attracted the attention of experts due to its application in integrable systems [15], [17], [24].
2. It is well known that Hurwitz spaces are related to matrix models, then, if one derive the associated matrix model of the Hurwitz spaces  $H_{1, n-1, 0}$ , we would immediately classify it by the group  $\mathcal{J}(\tilde{A}_n)$ .



- 
3. Even though the orbit space of  $\mathcal{J}(\tilde{A}_n)$  is locally isomorphic as Dubrovin Frobenius manifold to the Hurwitz space  $H_{1,n-1,0}$ , these two spaces are not necessarily the same. Indeed, the Dubrovin Frobenius manifold associated to Hurwitz spaces is a local construction, because it is defined in a domain of a solution of a Darboux-Egoroff system. On another hand, orbit spaces are somehow global objects, because their ring of invariant functions are polynomial over a suitable ring. In addition, the notion of invariant function gives information about the non-cubic part of the WDDV solution associated with the orbit space, see the last equation of (1.15) for instance.
  4. The orbit space construction of the group  $\mathcal{J}(\tilde{A}_n)$  can be generalised to the other classical finite Coxeter groups as  $B_n, D_n$ . Hence, these orbit spaces could give rise to a new class of Dubrovin Frobenius manifolds. Furthermore, the associated integrable hierarchies of this new class of Dubrovin Frobenius manifolds could have applications in Gromov-Witten theory and combinatorics.

The thesis is organised in the following way. In section 2, we recall the basic definitions of Dubrovin-Frobenius manifolds. In section 3, we recall the Dubrovin Frobenius manifold construction on Hurwitz spaces. In section 4, we defined extended affine Jacobi group  $\mathcal{J}(\tilde{A}_n)$  and, we construct Dubrovin-Frobenius structure on the orbit spaces of  $\mathcal{J}(\tilde{A}_n)$ . Furthermore, we show that the orbit space of the group  $\mathcal{J}(\tilde{A}_n)$  is isomorphic, as Dubrovin-Frobenius manifold, to the Hurwitz-Frobenius manifold  $\tilde{H}_{1,n-1,0}$  [10], [27]. See theorem 4.58 for details. The section 4 generalises the results of section [1] for the group  $\mathcal{J}(\tilde{A}_n)$ .

## Acknowledgements

I am grateful to Professor Boris Dubrovin for proposing this problem, for his remarkable advice and guidance. I would like also to thank Prof. Davide Guzzetti, and Prof. Marco Bertola for helpful discussions, and guidance of this paper. Moreover, the author acknowledges the support of the H2020-MSCA-RISE-2017 PROJECT No. 778010 IPADEGAN.

## 2 Review of Dubrovin-Frobenius manifolds

### 2.1 Basic definitions

We recall the basic definitions of Dubrovin Frobenius manifold, for more details [10].

**Definition 2.1.** A Frobenius Algebra  $\mathcal{A}$  is an unital, commutative, associative algebra endowed with an invariant non degenerate bilinear pairing

$$\eta(, ) : \mathcal{A} \otimes \mathcal{A} \mapsto \mathbb{C},$$

invariant in the following sense:

$$\eta(A \bullet B, C) = \eta(A, B \bullet C),$$

$$\forall A, B, C \in \mathcal{A}.$$

---

**Definition 2.2.**  $M$  is smooth or complex Dubrovin-Frobenius manifold of dimension  $n$  if a structure of Frobenius algebra is specified on any tangent plane  $T_t M$  at any point  $t \in M$ , smoothly depending on the point such that:

1. The invariant inner product  $\eta(,)$  is a flat metric on  $M$ . The flat coordinates of  $\eta(,)$  will be denoted by  $(t^1, t^2, \dots, t^n)$ .
2. The unity vector field  $e$  is covariantly constant w.r.t. the Levi-Civita connection  $\nabla$  for the metric  $\eta(,)$

$$\nabla e = 0 \quad (2.1)$$

3. Let

$$c(u, v, w) := \eta(u \bullet v, w) \quad (2.2)$$

(a symmetric 3-tensor). We require the 4-tensor

$$(\nabla_z c)(u, v, w) \quad (2.3)$$

to be symmetric in the four vector fields  $u, v, w, z$ .

4. A vector field  $E$  must be determined on  $M$  such that:

$$\nabla \nabla E = 0 \quad (2.4)$$

and that the corresponding one-parameter group of diffeomorphisms acts by conformal transformations of the metric  $\eta$ , and by rescalings on the Frobenius algebras  $T_t M$ . Equivalently:

$$[E, e] = -e, \quad (2.5)$$

$$\begin{aligned} \mathcal{L}_E \eta(X, Y) &:= E\eta(X, Y) - \eta([E, X], Y) - \eta(X, [E, Y]) \\ &= (2 - d)\eta(X, Y), \end{aligned} \quad (2.6)$$

$$\begin{aligned} \mathcal{L}_E c(X, Y, Z) &:= Ec(X, Y, Z) - c([E, X], Y, Z) - c(X, [E, Y], Z) \\ &\quad - c(X, Y, [E, Z]) = (3 - d)c(X, Y, Z). \end{aligned} \quad (2.7)$$

The Euler vector  $E$  can be represented as follows:

**Lemma 2.1.** If the grading operator  $Q := \nabla E$  is diagonalizable, then  $E$  can be represented as:

$$E = \sum_{i=1}^n ((1 - q_i)t_i + r_i)\partial_i \quad (2.8)$$

We now define scaling exponent as follows:

**Definition 2.3.** A function  $\varphi : M \mapsto \mathbb{C}$  is said to be quasi-homogeneous of scaling exponent  $d_\varphi$ , if it is a eigenfunction of Euler vector field:

$$E(\varphi) = d_\varphi \varphi \quad (2.9)$$

---

**Definition 2.4.** The function  $F(t), t = (t^1, t^2, \dots, t^n)$  is a solution of WDVV equation if its third derivatives

$$c_{\alpha\beta\gamma} = \frac{\partial^3 F}{\partial t^\alpha \partial t^\beta \partial t^\gamma} \quad (2.10)$$

satisfy the following conditions:

1.

$$\eta_{\alpha\beta} = c_{1\alpha\beta}$$

is constant nondegenerate matrix.

2. The function

$$c_{\alpha\beta}^\gamma = \eta^{\gamma\delta} c_{\alpha\beta\delta}$$

is structure constant of associative algebra.

3.  $F(t)$  must be quasi-homogeneous function

$$F(c^{d_1} t^1, \dots, c^{d_n} t^n) = c^{d_F} F(t^1, \dots, t^n)$$

for any nonzero  $c$  and for some numbers  $d_1, \dots, d_n, d_F$ .

**Lemma 2.2.** Any solution of WDVV equations with  $d_1 \neq 0$  defined in a domain  $t \in M$  determines in this domain the structure of a Dubrovin-Frobenius manifold. Conversely, locally any Dubrovin-Frobenius manifold is related to some solution of WDVV equations.

## 2.2 Intersection form

**Definition 2.5.** Let  $x = \eta(X, \cdot), y = \eta(Y, \cdot) \in \Gamma(T^*M)$  where  $X, Y \in \Gamma(TM)$ . An induced Frobenius algebra is defined on  $\Gamma(T^*M)$  by:

$$x \bullet y = \eta(X \bullet Y, \cdot).$$

**Definition 2.6.** The intersection form is a bilinear pairing in  $T^*M$  defined by:

$$(\omega_1, \omega_2)^* := \iota_E(\omega_1 \bullet \omega_2)$$

where  $\omega_1, \omega_2 \in T^*M$  and  $\bullet$  is the induced Frobenius algebra product in the cotangent space. the intersection form will be denoted by  $g^*$ .

**Remark 1:** Let  $x = \eta(X, \cdot), y = \eta(Y, \cdot) \in \Gamma(T^*M)$ . Then:

$$g^*(x, y) = \eta(X \bullet Y, E) = c(X, Y, E).$$

**Remark 2:** It is possible to prove that the tensor  $g^*$  defines a bilinear form on the tangent bundle that is almost everywhere non degenerate [10].

**Proposition 2.3.** The metric  $g^*$  is flat, and  $\forall \lambda \in \mathbb{C}$ , the contravariant metric  $\eta^*(\cdot, \cdot) + \lambda g^*(\cdot, \cdot)$  is flat, and the contravariant connection is  $\nabla^\eta + \lambda \nabla^g$ , where  $\nabla^\eta, \nabla^g$  are the contravariant connections of  $\eta^*$  and  $g^*$  respectively. The family of metrics  $\eta^*(\cdot, \cdot) + \lambda g^*(\cdot, \cdot)$  is called Flat pencil of metrics.

---

**Lemma 2.4.** The induced metric  $\eta^*$  on the cotangent bundle  $T^*M$  can be written as Lie derivative with respect the unit vector field  $e$  of the intersection form  $g^*$ . i.e

$$\eta^* = \mathcal{L}_e g^*. \quad (2.11)$$

**Lemma 2.5.** The correspondent WDVV solution  $F(t^1, \dots, t^n)$  of the Dubrovin-Frobenius manifold works as potential function for  $g^*$ . More precisely:

$$g^*(dt^i, dt^j) = (1 + d - q_i - q_j) \nabla_{(dt^i)^\sharp} \nabla_{(dt^j)^\sharp} F. \quad (2.12)$$

where the form  $(dt^j)^\sharp$  is the image of  $dt^j$  by the isomorphism induced by the metric  $\eta$ .

### 2.3 Reconstruction

Suppose that given a Dubrovin-Frobenius manifold  $M$ , only the following data are known: intersection form  $g^*$ , unit vector field  $e$ , Euler vector field  $E$ . From the previous lemmas we can reconstruct the Dubrovin-Frobenius manifold by setting:

$$\eta^* = \mathcal{L}_e g^*. \quad (2.13)$$

Then, we find the flat coordinates of  $\eta$  as homogeneous functions, and the structure constants by imposing:

$$g^*(dt^i, dt^j) = (1 + d - q_i - q_j) \nabla_{(dt^i)^\sharp} \nabla_{(dt^j)^\sharp} F. \quad (2.14)$$

Therefore, it is possible to compute the Free-energy by integration. Of course, we may have obstructions when,  $1 + d = q_i + q_j$ .

### 2.4 Monodromy of Dubrovin-Frobenius manifold

The intersection form  $g$  of a Dubrovin-Frobenius manifold is a flat almost everywhere nondegenerate metric. Let

$$\Sigma = \{t \in M : \det(g) = 0\}$$

Hence, the linear system of differential equations determining  $g^*$ -flat coordinates

$$g^{\alpha\epsilon}(t) \frac{\partial^2 x}{\partial t^\beta \partial t^\epsilon} + \Gamma_{\beta}^{\alpha\epsilon}(t) \frac{\partial x}{\partial t^\epsilon} = 0$$

has poles, and consequently its solutions  $x_a(t^1, \dots, t^n)$  are multivalued, where  $(t^1, \dots, t^n)$  are flat coordinates of  $\eta$ . The analytical continuation of the solutions  $x_a(t^1, \dots, t^n)$  has monodromy corresponding to loops around  $\Sigma$ . This gives rise to a monodromy representation of  $\pi_1(M \setminus \Sigma)$ , which is called Monodromy of the Dubrovin-Frobenius manifold.

### 2.5 Dubrovin Connection

In the theory of Dubrovin Frobenius manifold, there is another way to associate a monodromy group on it. Consider the following deformation of the Levi-Civita connection defined on a Dubrovin Frobenius manifold  $M$

$$\tilde{\nabla}_u v := \nabla_u v + zu \bullet v, \quad u, v \in \Gamma(TM), \quad (2.15)$$

where  $\nabla$  is the Levi-Civita connection of the metric  $\eta$ ,  $\bullet$  is the Frobenius product, and  $z \in \mathbb{CP}^1$ . Then, the following connection defined in  $M \times \mathbb{CP}^1$

$$\begin{aligned}\tilde{\nabla}_u v &:= \nabla_u v + zu \bullet v, \\ \tilde{\nabla}_{\frac{d}{dz}} \frac{d}{dz} &= 0, \quad \tilde{\nabla}_v \frac{d}{dz} = 0, \\ \tilde{\nabla}_{\frac{d}{dz}} v &= \partial_z v + E \bullet v - \frac{1}{z} \mu(v).\end{aligned}\tag{2.16}$$

where  $\mu$  is the diagonal matrix be given by

$$\mu_{\alpha\beta} = (q_\alpha - \frac{d}{2})\delta_{\alpha\beta}.\tag{2.17}$$

The monodromy representation arise by considering the solutions of the flat coordinate systems

$$\tilde{\nabla} d\tilde{t} = 0.\tag{2.18}$$

After doing some Gauge transformations in the system (2.18), and writing it in matricidal form. The system 2.18 takes the form

$$\begin{aligned}\frac{dY}{dt^\alpha} &= zC_\alpha Y, \\ \frac{dY}{dz} &= \left(U + \frac{\mu}{z}\right)Y,\end{aligned}\tag{2.19}$$

where

$$C_\alpha^\gamma = c_{\alpha\beta}^\gamma, \quad U_\beta^\gamma = E^\epsilon c_{\beta\epsilon}^\gamma.\tag{2.20}$$

## 2.6 Differential geometry preliminaries

In order to derive the Dubrovin Frobenius manifolds, we recall some results related with Riemannian geometry of the contravariant "metric"  $g^{ij}$ . By metric, I mean symmetric, bilinear, non-degenerate. In coordinates, let the metric

$$g_{ij}dx^i dx^j$$

and its induced contravariant metric

$$g^{ij} \frac{\partial}{\partial x_i} \otimes \frac{\partial}{\partial x_j}$$

The Levi Civita connection is uniquely specified by

$$\nabla_k g_{ij} = \partial_k g_{ij} - \Gamma_{ki}^s g_{sj} - \Gamma_{kj}^s g_{is} = 0,\tag{2.21}$$

or

$$\nabla_k g^{ij} = \partial_k g^{ij} - \Gamma_{ks}^i g^{sj} - \Gamma_{ks}^j g^{is} = 0,\tag{2.22}$$

and

$$\Gamma_{ij}^k = \Gamma_{ji}^k.\tag{2.23}$$

The Christoffel symbol can be written as

$$\Gamma_{ij}^k = g^{ks} (\partial_i g_{sj} + \partial_j g_{is} - \partial_s g_{ij}).$$

---

But, for our purpose it will be more convenient to use

$$\Gamma_k^{ij} := -g^{is}\Gamma_{sk}^j. \quad (2.24)$$

Then, the equations (2.21), (2.22), and (2.23) are equivalent to

$$\begin{aligned} \partial_k g^{ij} &= \Gamma_k^{ij} + \Gamma_k^{ji}, \\ g^{is}\Gamma_s^{jk} &= g^{js}\Gamma_s^{ik}. \end{aligned} \quad (2.25)$$

Introducing the operators

$$\begin{aligned} \nabla^i &= g^{is}\nabla_s, \\ \nabla^i \xi_k &= g^{is}\partial_s \xi_k + \Gamma_k^{is} \xi_s. \end{aligned} \quad (2.26)$$

The curvature tensor  $R_{ijk}^l$  of the metric measures noncommutativity of the operators  $\nabla_i$  or, equivalently  $\nabla^i$

$$(\nabla_i \nabla_j - \nabla_j \nabla_i) \xi_k = R_{ijk}^l \xi_l \quad (2.27)$$

where

$$R_{ijk}^l = \partial_i \Gamma_{jk}^l - \partial_j \Gamma_{ik}^l + \Gamma_{is}^l \Gamma_{jk}^s - \Gamma_{js}^l \Gamma_{ik}^s \quad (2.28)$$

We say that the metric is flat if the curvature of it vanishes. For a flat metric local flat coordinates  $p_1, \dots, p_n$  exist such that in these coordinates the metric is constant and the components of the Levi-Civita connection vanish. Conversely, if a system of flat coordinates for a metric exists then the metric is flat. The flat coordinates are determined uniquely up to an affine transformation with constant coefficients. They can be found from the following system

$$\nabla^i \partial_k p = g^{is} \partial_s \partial_k p + \Gamma_k^{is} \partial_s p = 0. \quad (2.29)$$

The correspondent Riemman tensor for the contravariant metric  $g^{ij}$  can be written as

$$R_l^{ijk} := g^{is} g^{jt} R_{stl}^k = g^{is} \left( \partial_s \Gamma_l^{jk} - \partial_l \Gamma_s^{jk} \right) + \Gamma_s^{ij} \Gamma_l^{sk} - \Gamma_s^{ik} \Gamma_l^{sj}. \quad (2.30)$$

The aim of this section is to construct a Dubrovin Frobenius structure on the orbit space of  $A_n$ . The strategy to achieve this goal is based on the derivation of a WDVV solution from the geometric data of the orbit space  $A_n$ . More specifically, the WDVV solution will be derived from the flat pencil structure which the orbit space of  $A_n$  naturally has.

**Definition 2.7.** [9] Two metrics  $(g^*, \eta^*)$  form a flat pencil if:

1. The metric

$$g_\lambda^{ij} := g^{ij} + \lambda \eta^{ij}, \quad (2.31)$$

is flat for arbitrary  $\lambda$ .

---

2. The Levi-Civita connection of the metric (2.31) has the form

$$\Gamma_{k,\lambda}^{ij} := \Gamma_{k,g}^{ij} + \lambda \Gamma_{k,\eta}^{ij}$$

where  $\Gamma_{k,g}^{ij}, \Gamma_{k,\eta}^{ij}$  are the Levi-Civita connection of  $g^*$ , and  $\eta^*$  respectively.

The main source of flat pencil metric is the following lemma

**Lemma 2.6.** [9] If for a flat metric  $g^*$  in some coordinate  $a_2, a_3, \dots, a_{n+1}$  both the coefficients of the metric  $g^{ij}$  and Levi Civita connection  $\Gamma_k^{ij}$  are linear in the coordinate  $a_{n+1}$ , and if  $\det(g^*) \neq 0$ , then, the metric

$$g^{ij} + \lambda \frac{\partial g^{ij}}{\partial a_{n+1}} \quad (2.32)$$

form a flat pencil. The corresponding Levi-Civita connection have the form

$$\Gamma_{k,g}^{ij} := \Gamma_k^{ij}, \quad \Gamma_{k,\eta}^{ij} := \frac{\partial \Gamma_k^{ij}}{\partial a_{n+1}} \quad (2.33)$$

Hence, our goal is first to construct a flat globally well defined metric in the orbit space of  $A_n$  such that there exist coordinates  $a_2, a_3, \dots, a_{n+1}$  in which both the metric and its Christoffel symbols are at most linear on  $a_{n+1}$ .

The following lemma shows that flat pencil structure is almost the same as Dubrovin Frobenius structure.

**Lemma 2.7.** [9] For a flat pencil metric  $g^{\alpha\beta}, \eta^{\alpha\beta}$  there exist a vector field  $f = f^\gamma \partial_\gamma$  such that the tensor

$$\Delta^{\alpha\beta\gamma} = \eta^{\alpha\delta} \Gamma_{\delta,g}^{\beta\gamma} - g^{\alpha\delta} \Gamma_{\delta,\eta}^{\beta\gamma} \quad (2.34)$$

and the metric  $g^{\alpha\beta}$  have the following form

$$\begin{aligned} \Delta^{\alpha\beta\gamma} &= \eta^{\alpha\mu} \eta^{\beta\nu} \partial_\mu \partial_\nu f^\gamma \\ g^{\alpha\beta} &= \eta^{\alpha\mu} \partial_\mu f^\beta + \eta^{\beta\nu} \partial_\nu f^\alpha + c \eta^{\alpha\beta} \end{aligned} \quad (2.35)$$

for some constant  $c$ . The vector field  $f$  should satisfy

$$\Delta_\epsilon^{\alpha\beta} \Delta_\delta^{\epsilon\gamma} = \Delta_\epsilon^{\alpha\gamma} \Delta_\delta^{\epsilon\beta} \quad (2.36)$$

where

$$\begin{aligned} \Delta_\gamma^{\alpha\beta} &= \eta_{\gamma\epsilon} \Delta^{\alpha\beta\epsilon} = \eta^{\alpha\mu} \partial_\mu \partial_\gamma f^\beta \\ (\eta^{\alpha\epsilon} g^{\beta\delta} - g^{\alpha\epsilon} \eta^{\beta\delta}) \partial_\epsilon \partial_\delta f^\gamma &= 0 \end{aligned} \quad (2.37)$$

Conversely, for any metric  $\eta^{\alpha\beta}$  and for  $f$  solution of the system (2.36) and (2.37) the metrics  $\eta^{\alpha\beta}$  and  $g^{\alpha\beta}$  form a flat pencil metric.

## 2.7 Semisimple Dubrovin Frobenius manifolds

**Definition 2.8.** A Frobenius algebra is called semisimple if it does not have nilpotent, i.e. if  $a \neq 0$  implies

$$a^m \neq 0, \quad \text{for any } m \in \mathbb{Z}. \quad (2.38)$$

---

**Lemma 2.8.** [10] Let  $A$  be a semisimple Frobenius algebra, then there exist a base  $e_1, e_2, \dots, e_n$  of  $A$ , such that the Frobenius product  $\bullet$  in this base is described by

$$e_i \bullet e_j = \delta_{ij} e_i. \quad (2.39)$$

**Definition 2.9.** A point in a Dubrovin Frobenius manifold is called semisimple, if the Frobenius algebra in its tangent space is semisimple.

**Remark 2.1.** Note that semisimplicity is an open condition.

**Lemma 2.9.** [10] In a neighbourhood of a semi semisimple point, there exist local coordinates  $(u_1, u_2, \dots, u_n)$  such that

$$\frac{\partial}{\partial u_i} \bullet \frac{\partial}{\partial u_j} = \delta_{ij} \frac{\partial}{\partial u_i}. \quad (2.40)$$

The coordinates  $(u_1, u_2, \dots, u_n)$  are called canonical coordinates.

**Lemma 2.10.** [10] Let  $M$  a semisimple Dubrovin Frobenius manifold, on the canonical coordinates  $(u_1, u_2, \dots, u_n)$  the intersection form, Euler vector field, and unit vector field can be written as

$$\begin{aligned} g^{ii} &= u_i \eta^{ii} \delta_{ij}, \\ e &= \sum_{i=1}^n \frac{\partial}{\partial u_i}, \\ E &= \sum_{i=1}^n u_i \frac{\partial}{\partial u_i}. \end{aligned} \quad (2.41)$$

**Proposition 2.11.** [10] In a neighborhood of a semisimple point all the roots  $(u_1, u_2, \dots, u_n)$  of the characteristic equation

$$\det(g^{\alpha\beta} - u\eta^{\alpha\beta}) = 0 \quad (2.42)$$

are simple. They are canonical coordinates in this neighbourhood. Conversely, if the roots of the characteristic equation are simple in a point  $p \in M$ , then  $p \in M$  is a semisimple point on the Frobenius manifold and  $(u_1, u_2, \dots, u_n)$  are canonical coordinates in the neighbourhood of this point.

**Definition 2.10.** [10] A diagonal metric on a  $n$ -dimensional manifold

$$\eta = \sum_{i=1}^n \eta_{ii} du_i^2 \quad (2.43)$$

is called potential, if there exist a function  $U(u_1, u_2, \dots, u_n)$  such that

$$\eta_{ii} = \frac{\partial U}{\partial u_i}. \quad (2.44)$$

**Definition 2.11.** [10] A potential diagonal flat metric  $\eta$  on a  $n$ -dimensional manifold is called Darboux-Egoroff metric.

**Lemma 2.12.** [10] Let be  $\eta$  a diagonal potential metric on a  $n$ -dimensional manifold

$$\eta = \sum_{i=1}^n \eta_{ii} du_i^2. \quad (2.45)$$



---

Then, the metric (2.45) is Darboux-Egoroff iff its rotational coefficients  $\beta_{ij}$

$$\beta_{ij} = \frac{\partial_j \sqrt{\eta_{ii}}}{\sqrt{\eta_{jj}}} \quad (2.46)$$

satisfy the system of equations

$$\begin{aligned} \partial_k \beta_{ij} &= \beta_{ik} \beta_{kj}, \\ \sum_{k=1}^n \partial_k \beta_{ij} &= 0. \end{aligned} \quad (2.47)$$

### 3 Review of Dubrovin-Frobenius manifold on Hurwitz spaces

#### 3.1 Hurwitz spaces

The main reference of this section are [10] and [27].

**Definition 3.1.** The Hurwitz space  $H_{g,n_0,\dots,n_m}$  is the moduli space of curves  $C_g$  of genus  $g$  endowed with a  $N$  branched covering,  $\lambda : C_g \mapsto \mathbb{CP}^1$  of  $\mathbb{CP}^1$  with  $m+1$  branching points over  $\infty \in \mathbb{CP}^1$  of branching degree  $n_i + 1, i = 0, \dots, m$ .

**Definition 3.2.** Two pairs  $(C_g, \lambda)$  and  $(\tilde{C}_g, \tilde{\lambda})$  are said Hurwitz-equivalent if there exist an analytic isomorphic  $F : C_g \mapsto \tilde{C}_g$  such that

$$\lambda \circ F = \tilde{\lambda}. \quad (3.1)$$

Roughly speaking, Hurwitz spaces  $H_{g,n_0,\dots,n_m}$  are moduli spaces of meromorphic functions which realise a Riemann surface of genus  $g$   $C_g$  as covering over  $\mathbb{CP}^1$  with a fixed ramification profile.

**Example 1:**

A generic point of the Hurwitz space  $H_{0,n}$  is

$$H_{0,n} = \{\lambda(p, x_0, x_1, x_2, \dots, x_n) = \prod_{i=0}^n (p - x_i) : \sum_{i=0}^n x_i = 0\} \quad (3.2)$$

**Example 2:**

A generic point of the Hurwitz space  $H_{0,n-1,0}$  is

$$H_{0,n} = \{\lambda(p, a_2, a_3, \dots, a_{n+1}, a_{n+2}) = p^n + a_2 p^{n-2} + \dots + a_n p + a_{n+1} + \frac{a_{n+2}}{p}\}. \quad (3.3)$$

**Example 3:**

A generic point of the Hurwitz space  $H_{1,n}$  is

$$H_{1,n} = \{\lambda(p, u, v_0, v_1, \dots, v_n, \tau) = e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(p - v_i, \tau)}{\theta_1^{n+1}(v, \tau)} : \sum_{i=0}^n v_i = 0\} \quad (3.4)$$

here  $\theta_1$  is the Jacobi theta function, see (??).

**Example 4:**

A generic point of the Hurwitz space is  $H_{1,n-1,0}$  is

$$H_{1,n-1,0} = \{\lambda(p, u, v_0, v_1, \dots, v_n, v_{n+1}, \tau) = e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(p - v_i, \tau)}{\theta_1^n(v, \tau) \theta_1(v + (n+1)v_{n+1}, \tau)} : \sum_{i=0}^n v_i = -(n+1)v_{n+1}\} \quad (3.5)$$

---

The covering  $\tilde{H} = \tilde{H}_{g,n_0,\dots,n_m}$  consist of the set of points

$$(C_g; \lambda; k_0, \dots, k_m; a_1, \dots, a_g, b_1, \dots, b_g) \in \tilde{H}_{g,n_0,\dots,n_m}$$

where  $C_g, \lambda$  are the same as above,  $a_1, \dots, a_g, b_1, \dots, b_g \in H_1(C_g, \mathbb{Z})$  are the canonical symplectic basis, and  $k_0, \dots, k_m$  are roots of  $\lambda$  near  $\infty_0, \infty_1, \dots, \infty_m$  of the orders  $n_0 + 1, n_1 + 1, \dots, n_m + 1$ . resp.,

$$k_i^{n_i+1}(P) = \lambda(P), \quad P \text{ near } \infty_i.$$

### 3.2 Bidifferential $W$

**Definition 3.3.** [27] Let  $P, Q \in C_g$ . The meromorphic Bidifferential  $W$  is given by

$$W(P, Q) = d_P d_Q \log E(P, Q), \quad (3.6)$$

where  $E(P, Q)$  is the prime form on the Riemann surface  $C_g$ . Alternatively, it can be characterised by the following properties

1. symmetric meromorphic differential in  $C_g \times C_g$ , with second order pole on  $P = Q$  with biresidue 1
- 2.

$$\begin{aligned} \int_{a_k} W(P, Q) &= 0; \\ \int_{b_k} W(P, Q) &= 2\pi i \omega_k(P) \end{aligned} \quad (3.7)$$

where  $\{\omega_k(P)\}$  are the normalized base of holomorphic differentials, i.e.  $\int_{a_j} \omega_k(P) = \delta_{ij}$

The dependence of the bidifferential  $W$  on branch points of the Riemann surface is given by the Rauch variational formulas

$$\frac{\partial W(P, Q)}{\partial u_i} = \frac{1}{2} W(P, P_i) W(P_i, Q), \quad (3.8)$$

where  $W(P, P_i)$  is the evaluation of  $W(P, Q)$  at  $Q = P_i$  with respect to the standard local parameter  $x_j(Q) = \sqrt{\lambda - \lambda(Q)}$

$$W(P, P_i) = \left. \frac{W(P, Q)}{dx_j(Q)} \right|_{Q=P_i} \quad (3.9)$$

A remarkable consequence of the Rauch variational formula is that it induce a flat metric in the Hurwitz space. Indeed,

**Proposition 3.1.** [27] Let be the metric

$$ds_W^2 = \sum_{i=1}^n \left( \oint_l h(Q) W(Q, P_i) \right)^2 (du_i)^2, \quad (3.10)$$

where  $l$  is a smooth contour in the Riemann surface such that  $P_i \notin l$ , and  $h(Q)$  is a smooth function independent of  $\{u_i\}$ . Then, the rotational coefficients of (3.10) satisfies a Darboux-Egoroff system in lemma 2.12.

For particular choices of the function  $h(Q)$  the metrics (3.10) coincides with the metrics induced by the primary differentials see section 3.3 and [10] for details. This fact is remarkable, because, it shows that from only the data of the Hurwitz space, one can construct a flat metric for the desired Dubrovin Frobenius manifold.

### 3.3 Reconstruction of Dubrovin Frobenius manifold

Over the space  $\tilde{H}_{g,n_0,\dots,n_m}$ , it is possible to introduce a Dubrovin-Frobenius structure by taking as canonical coordinates the ramification points  $(u_1, u_2, \dots, u_n)$  of  $H_{g,n_0,\dots,n_m}$ . The Dubrovin-Frobenius structure is specified by the following objects:

$$\text{multiplication } \partial_i \bullet \partial_j = \delta_{ij} \partial_i, \text{ where } \partial_i = \frac{\partial}{\partial u_i}, \quad (3.11)$$

$$\text{Euler vector field } E = \sum_i u_i \partial_i, \quad (3.12)$$

$$\text{unit vector field } e = \sum_i \partial_i, \quad (3.13)$$

and the metric  $\eta$  defined by the formula

$$ds_\phi^2 = \sum \text{res}_{P_i} \frac{\phi^2}{d\lambda} (du_i)^2, \quad (3.14)$$

where  $\phi$  is some primary differential of the underlying Riemann surface  $C_g$ . Note that the Dubrovin-Frobenius manifold structure depends on the meromorphic function  $\lambda$ , and on the primary differential  $\phi$ . The list of possible primary differential  $\phi$  is in [10].

Consider a multivalued function  $p$  on  $C$  by taking the integral of  $\phi$

$$p(P) = v.p \int_{\infty_0}^P \phi$$

The principal value is defined by omitting the divergent part, when necessary, because  $\phi$  may be divergent at  $\infty_0$ , as function of the local parameter  $k_0$ . Indeed the primary differentials defined on [10] may diverge as functions of  $k_i$ .

$$\phi = dp.$$

Let  $\tilde{H}_\phi$  be the open domain in  $\tilde{H}$  specifying by the condition

$$\phi(P_i) \neq 0.$$

**Theorem 3.2.** [10] For any primary differential  $\phi$  of the list in [10] the multiplication (3.11), the unity (3.13), the Euler vector field (3.12), and the metric (3.14) determine a structure of Dubrovin Frobenius manifold on  $\tilde{H}_\phi$ . The corresponding flat coordinates  $t_A$ ,  $A = 1, \dots, N$  consist of the five parts

$$t_A = (t^{i,\alpha}, i = 0, \dots, m, \alpha = 1, \dots, n_i; p^i, q^i, i = 1, \dots, m; r^i, s^i, i = 1, \dots, g) \quad (3.15)$$

where

$$t^{i,\alpha} = \text{res}_{\infty_i} k_i^{-\alpha} p d\lambda \quad i = 0, \dots, m, \alpha = 1, \dots, n_i \quad (3.16)$$

$$p^i = v.p \int_{\infty_0}^{\infty_i} dp \quad i = 1, \dots, m. \quad (3.17)$$

$$q^i = -\text{res}_{\infty_i} \lambda dp \quad i = 1, \dots, m. \quad (3.18)$$

$$r^i = \int_{a_i} dp \quad i = 1, \dots, g. \quad (3.19)$$

$$s^i = -\frac{1}{2\pi i} \int_{b_i} \lambda dp \quad i = 1, \dots, g. \quad (3.20)$$

---

Moreover, function  $\lambda = \lambda(p)$  is the superpotential of this Dubrovin Frobenius manifold, i.e. we have the following formulas to compute the metric  $\eta = \langle, \rangle$ , the intersection form  $g^* = (, )$  and the structure constants  $c$ .

$$\langle \partial', \partial'' \rangle = - \sum \text{res}_{d\lambda=0} \frac{\partial'(\lambda) \partial''(\lambda)}{d\lambda} dp \quad (3.21)$$

$$(\partial', \partial'') = - \sum \text{res}_{d \text{Log} \lambda=0} \frac{\partial'(\text{Log} \lambda) \partial''(\text{Log} \lambda)}{d \text{Log} \lambda} dp \quad (3.22)$$

$$c(\partial', \partial'', \partial''') = - \sum \text{res}_{d\lambda=0} \frac{\partial'(\lambda) \partial''(\lambda) \partial'''(\lambda)}{d\lambda} dp \quad (3.23)$$

**Remark 3.1.** The Dubrovin Frobenius structure on Hurwitz spaces depend on a choice of suitable primary differentials. Dropping this suitable choice implies that we typically lose the quasi homogeneous condition of the WDVV equation or the fact the unit is covariant constant.

## 4 Differential geometry of the orbit space of extended Jacobi group $A_n$

This section is dedicated to generalise the group  $\mathcal{J}(\tilde{A}_1)$  defined in [1] for arbitrary  $n$ , this new class of groups will be denoted by  $\mathcal{J}(\tilde{A}_n)$ . From the data of the group  $\mathcal{J}(\tilde{A}_n)$ , we will construct the Dubrovin Frobenius manifold in the orbit space of  $\mathcal{J}(\tilde{A}_n)$ . Furthermore, this Dubrovin Frobenius manifold will be locally isomorphic to the Hurwitz space  $H_{1,n-1,0}$ . In the section 1.4, there is a scheme of the technical steps that we should take to built the desired Dubrovin Frobenius manifold.

### 4.1 The Group $\mathcal{J}(\tilde{A}_n)$

In this section, we define the group  $\mathcal{J}(\tilde{A}_n)$ . In order to understand the motivation of this group see 1.4.

Consider the  $A_n$  in the following extended space

$$L^{\tilde{A}_n} = \{(z_0, z_1, \dots, z_n, z_{n+1}) \in \mathbb{Z}^{n+2} : \sum_{i=0}^n z_i = 0\}.$$

The action of  $A_n$  on  $L^{\tilde{A}_n}$  is given by

$$w(z_0, z_1, z_2, \dots, z_{n-1}, z_n, z_{n+1}) = (z_{i_0}, z_{i_1}, z_{i_2}, \dots, z_{i_{n-1}}, z_{i_n}, z_{n+1})$$

permutations in the first  $n+1$  variables. Moreover,  $A_n$  also acts on the complexification of  $L^{\tilde{A}_n} \otimes \mathbb{C}$ . Let

the quadratic form  $\langle, \rangle_{\tilde{A}_n}$  be given by

$$\begin{aligned}
\langle v, v \rangle_{\tilde{A}_n} &= v^T M_{\tilde{A}_n} v \\
&= v^T \begin{pmatrix} 2 & 1 & 1 & \dots & 1 & 0 \\ 1 & 2 & 1 & \dots & 1 & 0 \\ 1 & 1 & 2 & \dots & 1 & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot & 0 \\ 1 & 1 & 1 & \dots & 2 & 1 \\ 1 & 1 & 1 & \dots & 2 & 0 \\ 0 & 0 & 0 & \dots & 0 & -n(n+1) \end{pmatrix} v \\
&= \sum_{i=0}^{n-1} A_{ij} v_i v_j - n(n+1) v_{n+1}^2.
\end{aligned} \tag{4.1}$$

Consider the following group  $L^{\tilde{A}_n} \times L^{\tilde{A}_n} \times \mathbb{Z}$  with the following group operation

$$\begin{aligned}
&\forall (\lambda, \mu, k), (\tilde{\lambda}, \tilde{\mu}, \tilde{k}) \in L^{\tilde{A}_n} \times L^{\tilde{A}_n} \times \mathbb{Z} \\
&(\lambda, \mu, k) \bullet (\tilde{\lambda}, \tilde{\mu}, \tilde{k}) = (\lambda + \tilde{\lambda}, \mu + \tilde{\mu}, k + \tilde{k} + \langle \lambda, \tilde{\lambda} \rangle_{\tilde{A}_n})
\end{aligned}$$

Note that  $\langle, \rangle_{\tilde{A}_n}$  is invariant under  $A_n$  group, then  $A_n$  acts on  $L^{\tilde{A}_n} \times L^{\tilde{A}_n} \times \mathbb{Z}$ . Hence, we can take the semidirect product  $A_n \ltimes (L^{\tilde{A}_n} \times L^{\tilde{A}_n} \times \mathbb{Z})$  given by the following product.

$$\begin{aligned}
&\forall (w, \lambda, \mu, k), (\tilde{w}, \tilde{\lambda}, \tilde{\mu}, \tilde{k}) \in A_n \times L^{\tilde{A}_n} \times L^{\tilde{A}_n} \times \mathbb{Z}, \\
&(w, \lambda, \mu, k) \bullet (\tilde{w}, \tilde{\lambda}, \tilde{\mu}, \tilde{k}) = (w\tilde{w}, w\lambda + \tilde{\lambda}, w\mu + \tilde{\mu}, k + \tilde{k} + \langle \lambda, \tilde{\lambda} \rangle_{\tilde{A}_n}).
\end{aligned}$$

Denoting  $W(\tilde{A}_n) := A_n \ltimes (L^{\tilde{A}_n} \times L^{\tilde{A}_n} \times \mathbb{Z})$ , we can define

**Definition 4.1.** The Jacobi group  $\mathcal{J}(\tilde{A}_n)$  is defined as a semidirect product  $W(\tilde{A}_n) \rtimes SL_2(\mathbb{Z})$ . The group action of  $SL_2(\mathbb{Z})$  on  $W(\tilde{A}_n)$  is defined as

$$\begin{aligned}
Ad_\gamma(w) &= w, \\
Ad_\gamma(\lambda, \mu, k) &= (a\mu - b\lambda, -c\mu + d\lambda, k + \frac{ac}{2} \langle \mu, \mu \rangle_{\tilde{A}_n} - bc \langle \mu, \lambda \rangle_{\tilde{A}_n} + \frac{bd}{2} \langle \lambda, \lambda \rangle_{\tilde{A}_n}),
\end{aligned}$$

for  $(w, t = (\lambda, \mu, k)) \in W(\tilde{A}_n), \gamma \in SL_2(\mathbb{Z})$ . Then the multiplication rule is given as follows

$$(w, t, \gamma) \bullet (\tilde{w}, \tilde{t}, \tilde{\gamma}) = (w\tilde{w}, t \bullet Ad_\gamma(w\tilde{t}), \gamma\tilde{\gamma}).$$

Let us use the following identification  $\mathbb{Z}^{n+1} \cong L^{\tilde{A}_n}, \mathbb{C}^{n+1} \cong L^{\tilde{A}_n} \otimes \mathbb{C}$  that is possible due to maps

$$\begin{aligned}
(v_0, \dots, v_{n-1}, v_{n+1}) &\mapsto (v_0, \dots, v_{n-1}, -\sum_{i=0}^n v_i, v_{n+1}), \\
(v_0, \dots, v_{n-1}, v_n, v_{n+1}) &\mapsto (v_0, \dots, v_{n-1}, v_{n+1}).
\end{aligned}$$

Then the action of Jacobi group  $\mathcal{J}(\tilde{A}_n)$  on  $\Omega := \mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H}$  is given as follows

---

**Proposition 4.1.** The group  $\mathcal{J}(\tilde{A}_n) \ni (w, t, \gamma)$  acts on  $\Omega := \mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} \ni (u, v, \tau)$  as follows

$$\begin{aligned} w(u, v, \tau) &= (u, wv, \tau) \\ t(u, v, \tau) &= (u - \langle \lambda, v \rangle_{\tilde{A}_n} - \frac{1}{2} \langle \lambda, \lambda \rangle_{\tilde{A}_n} \tau, v + \lambda \tau + \mu, \tau) \\ \gamma(u, v, \tau) &= (\phi + \frac{c \langle v, v \rangle_{\tilde{A}_n}}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{a\tau + b}{c\tau + d}) \end{aligned} \quad (4.2)$$

The proof of the proposition 4.1 follows from the proposition 1.1 of [6] with small adaptation with the extra trivial action in the exceptional variable  $v_{n+1}$  with respect the  $A_n$  action.

## 4.2 Jacobi forms of $\mathcal{J}(\tilde{A}_n)$

In order to understand the differential geometry of orbit space, first we need to study the algebra of the invariant functions. Informally, every time that there is a group  $W$  acting on a vector space  $V$ , one could think the orbit spaces  $V/W$  as  $V$ , but you should remember yourself that it is only allowed to use the  $W$ -invariant sections of  $V$ . Hence, motivated by the definition of Jacobi forms of group  $A_n$  defined in [32], and used in the context of Dubrovin-Frobenius manifold in [6],[7], and generalising also the meromorphic Jacobi forms introduced in [1] we give the following:

**Definition 4.2.** The weak  $\tilde{A}_n$ -invariant Jacobi forms of weight  $k$ , order  $l$ , and index  $m$  are functions on  $\Omega = \mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} \ni (u, v', v_{n+1}, \tau) = (u, v, \tau)$  which satisfy

$$\begin{aligned} \varphi(w(u, v, \tau)) &= \varphi(u, v, \tau), \quad A_n \text{ invariant condition} \\ \varphi(t(u, v, \tau)) &= \varphi(u, v, \tau) \\ \varphi(\gamma(u, v, \tau)) &= (c\tau + d)^{-k} \varphi(u, v, \tau) \\ E\varphi(u, v, \tau) &:= -\frac{1}{2\pi i} \frac{\partial}{\partial u} \varphi(u, v, \tau) = m\varphi(u, v, \tau) \end{aligned} \quad (4.3)$$

Moreover,

1.  $\varphi$  is locally bounded functions on  $v'$  as  $\Im(\tau) \mapsto +\infty$  (weak condition).
2. For fixed  $u, v', \tau$  the function  $v_{n+1} \mapsto \varphi(u, v', v_{n+1}, \tau)$  is meromorphic with poles of order at most  $l + 2m$  on  $v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$ .
3. For fixed  $u, v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}, \tau$  the function  $(i \neq n+1) v_i \mapsto \varphi(u, v', v_{n+1}, \tau)$  is holomorphic.
4. For fixed  $u, v', v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$ . the function  $\tau \mapsto \varphi(u, v', v_{n+1}, \tau)$  is holomorphic.

The space of  $\tilde{A}_n$ -invariant Jacobi forms of weight  $k$ , order  $l$ , and index  $m$  is denoted by  $J_{k,l,m}^{\tilde{A}_n}$ , and  $J_{\bullet,\bullet,\bullet}^{\mathcal{J}(\tilde{A}_n)} = \bigoplus_{k,l,m} J_{k,l,m}^{\tilde{A}_n}$  is the space of Jacobi forms  $\tilde{A}_n$  invariant.

**Remark 4.1.** The condition  $E\varphi(u, v', v_{n+1}, \tau) = m\varphi(u, v', v_{n+1}, \tau)$  implies that  $\varphi(u, v', v_{n+1}, \tau)$  has the following form

$$\varphi(u, v', v_{n+1}, \tau) = f(v', v_{n+1}, \tau) e^{2\pi i m u}$$

See also remarks ??.

---

The main result of section is the following.

**The ring of  $\tilde{A}_n$  invariant Jacobi forms is polynomial over a suitable ring  $E_{\bullet,\bullet} := J_{\bullet,\bullet,0}^{\mathcal{J}(\tilde{A}_n)}$  on suitable generators  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$ .**

Before state precisely the theorem, I will define the objects  $E_{\bullet,\bullet}, \varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$ .

The ring  $E_{\bullet,l} := J_{\bullet,l,0}^{\mathcal{J}(\tilde{A}_n)}$  is the space of meromorphic Jacobi forms of index 0 with poles of order at most  $l$  on  $v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$ , by definition. The sub-ring  $J_{\bullet,0,0}^{\mathcal{J}(\tilde{A}_n)} \subset E_{\bullet,\bullet}$  has a nice structure, indeed:

**Lemma 4.2.** The sub-ring  $J_{\bullet,0,0}^{\mathcal{J}(\tilde{A}_n)}$  is equal to  $M_{\bullet} := \bigoplus M_k$ , where  $M_k$  is the space of modular forms of weight  $k$  for the full group  $SL_2(\mathbb{Z})$ .

*Proof.* Using the Remark 4.1, we know that functions  $\varphi(u, v', v_{n+1}, \tau) \in J_{\bullet,0,0}^{\mathcal{J}(\tilde{A}_n)}$  can not depend on  $u$ , then  $\varphi(u, v', v_{n+1}, \tau) = \varphi(v', v_{n+1}, \tau)$ . Moreover, for fixed  $v_{n+1}, \tau$  the functions  $v_i \mapsto \varphi(v', v_{n+1}, \tau)$  are holomorphic elliptic function for any  $i \neq n+1$ . Therefore, by Liouville theorem, these function are constant in  $v'$ . Similar argument shows that these function do not depend on  $v_{n+1}$ , because  $l+2m=0$ , i.e there is no pole. Then,  $\varphi = \varphi(\tau)$  are standard holomorphic modular forms.  $\square$

**Lemma 4.3.** If  $\varphi \in E_{\bullet,\bullet} = J_{\bullet,\bullet,0}^{\mathcal{J}(\tilde{A}_n)}$ , then  $\varphi$  depends only on the variables  $v_{n+1}, \tau$ . Moreover, if  $\varphi \in J_{0,l,0}^{\mathcal{J}(\tilde{A}_n)}$  for fixed  $\tau$  the function  $\tau \mapsto \varphi(v_{n+1}, \tau)$  is a elliptic function with poles of order at most  $l$  on  $v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$ .

*Proof.* The proof follows essentially in the same way of the lemma 4.2, the only difference is that now we have poles on  $v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$ . Then, we have depedence in  $v_{n+1}$ .  $\square$

As a consequence of lemma 4.3, the function  $\varphi \in E_{k,l} = J_{k,l,0}^{\mathcal{J}(\tilde{A}_n)}$  has the following form

$$\varphi(v_{n+1}, \tau) = f(\tau)g(v_{n+1}, \tau)$$

where  $f(\tau)$  is holomorphic modular form of weight  $k$ , and for fixed  $\tau$ , the function  $v_{n+1} \mapsto g(v_{n+1}, \tau)$  is an elliptic function of order at most  $l$  on the poles  $v_{n+1} = \frac{j}{n} + \frac{l\tau}{n}, 0 \leq l, j \leq n-1 \bmod \mathbb{Z} \oplus \tau\mathbb{Z}$ .

Note that a natural way to produce meromorphic Jacobi form is by using rational functions of holomorphic Jacobi forms. Starting from now, we will denote the Jacobi forms related with the Jacobi group  $\mathcal{J}(A_{n+1})$  with the upper index  $\mathcal{J}(A_{n+1})$ , for instance

$$\varphi^{\mathcal{J}(A_{n+1})},$$

and the Jacobi forms related with the Jacobi group  $\mathcal{J}(\tilde{A}_n)$  with the with the upper index  $\mathcal{J}(\tilde{A}_n)$

$$\varphi^{\mathcal{J}(\tilde{A}_n)}.$$

In [6], Bertola found basis of the algebra of Jacobi form by producing a holomorphic Jacobi form of type  $A_n$  as product of theta functions.

$$\varphi_{n+2}^{\mathcal{J}(A_{n+1})} = e^{-2\pi i u} \prod_{i=1}^{n+2} \frac{\theta_1(z_i, \tau)}{\theta_1'(0, \tau)}. \quad (4.4)$$

Afterwards, Bertola defined a recursive operator to produce the remaining basic generators. In order to recall the details see section ???. Our strategy will follow the same logic of Bertola method, we use theta functions to produce a basic generator and thereafter, we produce a recursive operator to produce the remaining part.

**Lemma 4.4.** Let be  $\varphi_{n+2}^{\mathcal{J}(A_{n+1})}(u_1, z_1, z_2, \dots, z_n, \tau)$  the holomorphic  $A_{n+1}$ -invariant Jacobi forms which correspond to the algebra generator of maximal weight degree, in this case degree  $n+2$ . More explicitly,

$$\varphi_{n+2}^{\mathcal{J}(A_2)} = e^{-2\pi i u_1} \prod_{i=1}^{n+2} \frac{\theta_1(z_i, \tau)}{\theta_1'(0, \tau)}. \quad (4.5)$$

Let be  $\varphi_2^{\mathcal{J}(A_1)}(u_2, z_{n+1}, \tau)$  the holomorphic  $A_1$ -invariant Jacobi form which correspond to the algebra generator of maximal weight degree, in this case degree 2.

$$\varphi_2^{\mathcal{J}(A_1)} = e^{-2\pi i u_2} \frac{\theta_1(z_{n+1}, \tau)^2}{\theta_1'(0, \tau)^2}. \quad (4.6)$$

Then, the function

$$\varphi_n^{\mathcal{J}(\tilde{A}_n)} = \frac{\varphi_{n+2}^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}} \quad (4.7)$$

is meromorphic Jacobi form of index 1, weight  $-n$ , order 0.

*Proof.* For our convenience, we change the labels  $u_2 - u_1, z_1, z_2, \dots, z_{n+2}$  to

$$\begin{aligned} u &= u_2 - u_1, \\ z_1 &= v_0 - v_{n+1}, \\ z_2 &= v_1 - v_{n+1}, \\ &\vdots \\ &\vdots \\ z_{n+1} &= -\sum_{i=0}^n v_i - v_{n+1}, \\ z_{n+2} &= (n+1)v_{n+1}. \end{aligned} \quad (4.8)$$

Then (4.7) has the following form

$$\varphi_n^{\mathcal{J}(\tilde{A}_n)}(u, v_0, v_1, \dots, v_{n+1}, \tau) = e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1((n+1)v_{n+1}, \tau)} \quad (4.9)$$

Let us prove each item separated.

#### 1. $A_n$ invariant

The  $A_n$  group acts on (4.9) by permuting its roots, thus (4.9) remains invariant under this operation.

#### 2. Translation invariant

Recall that under the translation  $v \mapsto v + m + n\tau$ , the Jacobi theta function transform as [6], [31]:

$$\theta_1(v_i + \mu_i + \lambda_i \tau, \tau) = (-1)^{\lambda_i + \mu_i} e^{-2\pi i(\lambda_i v_i + \frac{\lambda_i^2}{2} \tau)} \theta_1(v_i, \tau). \quad (4.10)$$

Then substituting the transformation (4.10) into (4.9), we conclude that (4.9) remains invariant.



### 3. $SL_2(\mathbb{Z})$ invariant

Under  $SL_2(\mathbb{Z})$  action the following function transform as

$$\frac{\theta_1(\frac{v_i}{c\tau+d}, \frac{a\tau+d}{c\tau+d})}{\theta_1'(0, \frac{a\tau+d}{c\tau+d})} = (c\tau+d)^{-1} \exp(\frac{\pi i c v_i^2}{c\tau+d}) \frac{\theta_1(v_i, \tau)}{\theta_1'(0, \tau)}. \quad (4.11)$$

Then, substituting (4.11) in (4.9), we get

$$\varphi_n^{\mathcal{J}(\tilde{A}_n)} \mapsto \frac{\varphi_n^{\mathcal{J}(\tilde{A}_n)}}{(c\tau+d)^n}$$

### 4. Index 1

$$\frac{1}{2\pi i} \frac{\partial}{\partial u} \varphi_n^{\mathcal{J}(\tilde{A}_n)} = \varphi_n^{\mathcal{J}(\tilde{A}_n)}. \quad (4.12)$$

### 5. Analytic behavior

Note that  $\varphi_n^{\mathcal{J}(\tilde{A}_n)} \theta_1^2((n+1)v_{n+1}, \tau)$  is holomorphic function in all the variables  $v_i$ . Therefore  $\varphi_n^{\mathcal{J}(\tilde{A}_n)}$  are holomorphic functions on the variables  $\{v_i\}$ , and meromorphic function in the variable  $v_{n+1}$  with poles on  $\frac{j}{n} + \frac{l\tau}{n}, j, l = 0, 1, \dots, n-1$  of order 2, i.e  $l = 0$ , since  $m = 1$ .

□

In order to define the desired recursive operator, it is necessary to enlarge the domain of the Jacobi forms from  $\mathbb{C} \oplus \mathbb{C}^n \oplus \mathbb{H} \ni (u, v_0, v_1, \dots, v_{n+1}, \tau)$  to  $\mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} \ni (u, v_0, v_1, \dots, v_{n+1}, p, \tau)$ . In addition, we define lift a of Jacobi forms defined in  $\mathbb{C} \oplus \mathbb{C}^2 \oplus \mathbb{H}$  to  $\mathbb{C} \oplus \mathbb{C}^3 \oplus \mathbb{H}$  as

$$\varphi(u, v_0 - v_{n+1}, v_1 - v_{n+1}, \dots, (n+1)v_{n+1}, \tau) \mapsto \hat{\varphi}(p) := \varphi(u, v_0 - v_{n+1} + p, v_1 - v_{n+1} + p, \dots, (n+1)v_{n+1} + p, \tau)$$

A convenient way to do computation in these extended Jacobi forms is by using the following coordinates

$$\begin{aligned} s &= u + ng_1(\tau)p^2, \\ z_1 &= v_0 - v_{n+1} + p, \\ z_2 &= v_1 - v_{n+1} + p, \\ &\vdots \\ &\vdots \\ z_{n+1} &= -\sum_{i=0}^n v_i - v_{n+1} + p, \\ z_{n+2} &= (n+1)v_{n+1} + p, \\ \tau &= \tau. \end{aligned} \quad (4.13)$$

The bilinear form  $\langle v, v \rangle_{\tilde{A}_1}$  is extended to

$$\langle (z_1, z_2, \dots, z_{n+2}), (z_1, z_2, \dots, z_{n+2}) \rangle_E = \sum_{i=1}^{n+1} z_i^2 - z_{n+2}^2, \quad (4.14)$$

The action of the Jacobi group  $\tilde{A}_n$  in this extended space is

$$\begin{aligned}\hat{w}_E(u, v, p, \tau) &= (u, w(v), p, \tau) \\ t_E(u, v, p, \tau) &= \left( u - \langle \lambda, v \rangle_E - \frac{1}{2} \langle \lambda, \lambda \rangle_E \tau + k, v + p + \lambda \tau + \mu, \tau \right) \\ \gamma_E(u, v, p, \tau) &= \left( u + \frac{c \langle v, v \rangle_E}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{p}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right)\end{aligned}\tag{4.15}$$

**Proposition 4.5.** Let be  $\varphi \in J_{k,m,\bullet}^{\mathcal{J}(\tilde{A}_n)}$ , and  $\hat{\varphi}$  the correspondent extended Jacobi form. Then,

$$\left. \frac{\partial}{\partial p} (\hat{\varphi}) \right|_{p=0} \in J_{k-1,m,\bullet}^{\mathcal{J}(\tilde{A}_n)}.\tag{4.16}$$

*Proof.* 1.  **$A_n$ -invariant**

The vector field  $\frac{\partial}{\partial p}$  in coordinates  $s, z_1, z_2, \dots, z_{n+2}, \tau$  reads

$$\frac{\partial}{\partial p} = \sum_{i=1}^{n+2} \frac{\partial}{\partial z_i} + 2ng_1(\tau)p \frac{\partial}{\partial u}$$

Moreover, in the coordinates  $s, z_1, z_2, \dots, z_{n+1}, z_{n+2}, \tau$  the  $A_n$  group acts by permuting  $z_1, z_2, \dots, z_{n+1}$ . Then

$$\begin{aligned}\left. \frac{\partial}{\partial p} (\varphi(s, z_2, z_1, z_3, \tau)) \right|_{p=0} &= \left( \sum_{i=1}^{n+2} \frac{\partial}{\partial z_i} \right) (\varphi(s, z_{i_1}, z_{i_2}, \dots, z_{i_{n+1}}, z_{n+2}, \tau)) \Big|_{p=0} \\ &= \left( \sum_{i=1}^{n+2} \frac{\partial}{\partial z_i} \right) (\varphi(s, z_1, z_2, \dots, z_{n+1}, z_{n+2}, \tau)) \Big|_{p=0}.\end{aligned}$$

2. **Translation invariant**

$$\begin{aligned}& \left. \frac{\partial}{\partial p} (\varphi(u - \langle \lambda, v \rangle_E - \langle \lambda, \lambda \rangle_E \tau, v + p + \lambda \tau + \mu, \tau)) \right|_{p=0} \\ &= \left. \frac{\partial}{\partial p} \langle \lambda, v \rangle_E \right|_{p=0} \varphi(u, v, \tau) + \frac{\partial \varphi}{\partial p} \left( u - \langle \lambda, v \rangle_{\tilde{A}_n} - \frac{1}{2} \langle \lambda, \lambda \rangle_{\tilde{A}_n} \tau + k, v + \lambda \tau + \mu, \tau \right) \\ &= \frac{\partial \varphi}{\partial p} \left( u - \langle \lambda, v \rangle_{\tilde{A}_n} - \frac{1}{2} \langle \lambda, \lambda \rangle_{\tilde{A}_n} \tau + k, v + \lambda \tau + \mu, \tau \right) \\ &= \left. \frac{\partial \varphi}{\partial p} (u, v, \tau) \right|_{p=0}.\end{aligned}$$

3.  **$SL_2(\mathbb{Z})$  equivariant**

$$\begin{aligned}& \left. \frac{\partial}{\partial p} \left( \varphi \left( u + \frac{c \langle v, v \rangle_E}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{p}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \right) \right|_{p=0} \\ &= \frac{c}{2(c\tau + d)} \left. \frac{\partial}{\partial p} \langle v, v \rangle_E \right|_{p=0} \varphi(u, v, \tau) + \frac{1}{c\tau + d} \frac{\partial \varphi}{\partial p} \left( u + \frac{c \langle v, v \rangle_{\tilde{A}_n}}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{p}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\ &= \frac{1}{c\tau + d} \frac{\partial \varphi}{\partial p} \left( u + \frac{c \langle v, v \rangle_{\tilde{A}_n}}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{p}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \\ &= \frac{1}{(c\tau + d)^k} \left. \frac{\partial \varphi}{\partial p} (u, v, \tau) \right|_{p=0}.\end{aligned}$$

Then,

$$\left. \frac{\partial \varphi}{\partial p} \left( u + \frac{c \langle v, v \rangle_E}{2(c\tau + d)}, \frac{v}{c\tau + d}, \frac{p}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right) \right|_{p=0} = \frac{1}{(c\tau + d)^{k-1}} \left. \frac{\partial \varphi}{\partial p} (u, v, \tau) \right|_{p=0}.$$

#### 4. Index 1

$$\frac{1}{2\pi i} \frac{\partial}{\partial u} \frac{\partial}{\partial p} \hat{\varphi} = \frac{1}{2\pi i} \frac{\partial}{\partial p} \frac{\partial}{\partial u} \hat{\varphi} = \frac{\partial}{\partial p} \hat{\varphi}.$$

□

**Corollary 4.5.1.** The function

$$\begin{aligned} & \left[ e^{z \frac{\partial}{\partial p}} \left( e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(p + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(p + (n+1)v_{n+1}, \tau)} \right) \right] \Big|_{p=0} \\ &= \varphi_n^{\mathcal{J}(\tilde{A}_n)} + \varphi_{n-1}^{\mathcal{J}(\tilde{A}_n)} z + \varphi_{n-2}^{\mathcal{J}(\tilde{A}_n)} z^2 + \dots + \varphi_0^{\mathcal{J}(\tilde{A}_n)} z^n + O(z^{n+1}), \end{aligned} \quad (4.17)$$

is a generating function for the Jacobi forms  $\varphi_n^{\mathcal{J}(\tilde{A}_n)}, \varphi_{n-1}^{\mathcal{J}(\tilde{A}_n)}, \varphi_{n-2}^{\mathcal{J}(\tilde{A}_n)}, \dots, \varphi_0^{\mathcal{J}(\tilde{A}_n)}$ , where

$$\varphi_k^{\mathcal{J}(\tilde{A}_n)} := \frac{\partial^{n-k}}{\partial p^{n-k}} \left( \hat{\varphi}_n^{\mathcal{J}(\tilde{A}_n)} \right) \Big|_{p=0}. \quad (4.18)$$

*Proof.* Acting  $\frac{\partial}{\partial p}$   $k$  times in  $\varphi_n^{\mathcal{J}(\tilde{A}_n)}$ , we have

$$\left[ \frac{\partial^k}{\partial p^k} \left( e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(p + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(p + (n+1)v_{n+1}, \tau)} \right) \right] \Big|_{p=0} \in J_{-n+k, 1, \bullet}^{\mathcal{J}(\tilde{A}_n)}.$$

□

**Corollary 4.5.2.** The generating function can be written as

$$\left[ e^{z \frac{\partial}{\partial p}} \left( e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(p + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(p + (n+1)v_{n+1}, \tau)} \right) \right] \Big|_{p=0} = e^{-2\pi i(u + ng_1(\tau)z^2)} \frac{\prod_{i=0}^n \theta_1(z + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(z + (n+1)v_{n+1}, \tau)}. \quad (4.19)$$

*Proof.*

$$\begin{aligned} & \left[ e^{z \frac{\partial}{\partial p}} \left( e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(p + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(p + (n+1)v_{n+1}, \tau)} \right) \right] \Big|_{p=0} = \\ &= \left[ e^{z \frac{\partial}{\partial p}} \left( e^{2\pi i(s + ng_1(\tau)p^2)} \frac{\prod_{i=0}^n \theta_1(p + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(p + (n+1)v_{n+1}, \tau)} \right) \right] \Big|_{p=0} \\ &= e^{-2\pi i(u + ng_1(\tau)z^2)} \frac{\prod_{i=0}^n \theta_1(z + v_i - v_{n+1}, \tau)}{\theta_1'(0, \tau)^n \theta_1(z + (n+1)v_{n+1}, \tau)}. \end{aligned} \quad (4.20)$$

□

The next lemma is one of the main points of this section, because this lemma identify the orbit space of the group  $\mathcal{J}(\tilde{A}_n)$  with the Hurwitz space  $H_{1,n-1,0}$ . This relationship is possible due to the construction of the generating function of the Jacobi forms of type  $\tilde{A}_n$ , which can be completed to be the Landau-Ginzburg superpotential of  $H_{1,n-1,0}$  as follows

**Lemma 4.6.** There is local biholomorphism between the orbit space  $\mathcal{J}(\tilde{A}_n)$  and  $H_{1,n-1,0}$ , i.e the space of elliptic functions with 1 pole of order  $n$ , and one simple pole.

*Proof.* The correspondence is realized by the map:

$$[(u, v_0, v_1, \dots, v_{n-1}, v_{n+1}, \tau)] \longleftrightarrow \lambda(v) = e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(z - v_i, \tau)}{\theta_1^n(v, \tau) \theta_1(v + (n+1)v_{n+1}, \tau)} \quad (4.21)$$

Note that this map is well defined and one to one. Indeed:

### 1. Well defined

Note that proof that the map does not depend on the choice of the representant of  $[(\phi, v_0, v_1, \dots, v_{n-1}, v_{n+1}, \tau)]$  is equivalent to prove that the function (4.21) is invariant under the action of  $\mathcal{J}(\tilde{A}_n)$ . Indeed

### 2. $A_n$ invariant

The  $A_n$  group acts on (4.21) by permuting its roots, thus (4.21) remains invariant under this operation.

### 3. Translation invariant

Recall that under the translation  $v \mapsto v + m + n\tau$ , the Jacobi theta function transform as [6], [31]:

$$\theta_1(v_i + \mu_i + \lambda_i \tau, \tau) = (-1)^{\lambda_i + \mu_i} e^{-2\pi i(\lambda_i v_i + \frac{\lambda_i^2}{2} \tau)} \theta_1(v_i, \tau) \quad (4.22)$$

Then substituting the transformation (4.22) into (4.21), we conclude that (4.21) remains invariant.

### 4. $SL_2(\mathbb{Z})$ invariant

Under  $SL_2(\mathbb{Z})$  action the following function transform as

$$\frac{\theta_1(\frac{v_i}{c\tau+d}, \frac{a\tau+d}{c\tau+d})}{\theta_1'(0, \frac{a\tau+d}{c\tau+d})} = \exp(\frac{\pi i c v_i^2}{c\tau+d}) \frac{\theta_1(v_i, \tau)}{\theta_1'(0, \tau)} \quad (4.23)$$

Then substituting the transformation (4.23) into (4.21), we conclude that (4.21) remains invariant.

### 5. Injectivity

Two elliptic functions are equal if they have the same zeros and poles with multiplicity.

### 6. Surjectivity

Any elliptic function can be written as rational functions of Weierstrass sigma function up to a multiplication factor [31]. By using the formula to relate Weierstrass sigma function and Jacobi theta function

$$\sigma(v_i, \tau) = \frac{\theta_1(v_i, \tau)}{\theta_1'(0, \tau)} \exp(-2\pi i g_1(\tau) v_i^2) \quad (4.24)$$

□

**Corollary 4.6.1.** The functions  $(\varphi_0^{\tilde{A}_n}, \varphi_1^{\tilde{A}_n}, \dots, \varphi_n^{\tilde{A}_n})$  obtained by the formula

$$\begin{aligned} \lambda^{\tilde{A}_n} &= e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(z - v_i + v_{n+1}, \tau)}{\theta_1^n(z, \tau) \theta_1(z + (n+1)v_{n+1}, \tau)} \\ &= \varphi_n^{\tilde{A}_n} \wp^{n-2}(z, \tau) + \varphi_{n-1}^{\tilde{A}_n} \wp^{n-3}(z, \tau) + \dots + \varphi_2^{\tilde{A}_n} \wp(z, \tau) \\ &\quad + \varphi_1^{\tilde{A}_n} [\zeta(z, \tau) - \zeta(z + (n+1)v_{n+1}, \tau) + \varphi_0^{\tilde{A}_n}] \end{aligned} \quad (4.25)$$

are Jacobi forms of weight  $0, -1, -2, \dots, -n$  respectively, index 1, and order 0.

*Proof.* Let us prove each item separated.

#### 1. $A_n$ invariant, translation invariant

The l.h.s of (4.25) are  $A_n$  invariant, and translation invariant by the lemma (4.6). Then, by the uniqueness of Laurent expansion of  $\lambda^{\tilde{A}_n}$ , we have that  $\varphi_i^{\tilde{A}_n}$  are  $A_n$  invariant, and translation invariant.

## 2. $SL_2(\mathbb{Z})$ equivariant

The l.h.s of (4.25) are  $SL_2(\mathbb{Z})$  invariant, but the Weierstrass functions of the r.h.s have the following transformation law

$$\wp^{(k-2)}\left(\frac{z}{c\tau+d}, \frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^k \wp^{(k-2)}(z, \tau). \quad (4.26)$$

Then,  $\varphi_k^{\tilde{A}_n}$  must have the following transformation law

$$\varphi_k^{\tilde{A}_n}\left(u + \frac{c\langle v, v \rangle_{\tilde{A}_n}}{2(c\tau+d)}, \frac{v}{c\tau+d}, \frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^{-k} \varphi_k^{\tilde{A}_n}(u, v, \tau). \quad (4.27)$$

## 3. Index 1

$$\frac{1}{2\pi i} \frac{\partial}{\partial u} \lambda^{\tilde{A}_n} = \lambda^{\tilde{A}_n}. \quad (4.28)$$

Then

$$\frac{1}{2\pi i} \frac{\partial}{\partial u} \varphi_i^{\tilde{A}_n} = \varphi_i^{\tilde{A}_n}. \quad (4.29)$$

## 4. Analytic behavior

Note that  $\lambda^{\tilde{A}_n} \theta_1^2((n+1)v_{n+1}, \tau)$  is holomorphic function in all the variables  $v_i$ . Therefore  $\varphi_i^{\tilde{A}_n}$  are holomorphic functions on the variables  $v_0, v_1, \dots, v_{n-1}$ , and meromorphic function in the variable  $(n+1)v_{n+1}$  with poles on  $\frac{j}{n} + \frac{l\tau}{n}, j, l = 0, \dots, n-1$  of order 2, i.e  $l = 0$ , since  $m = 1$  for all  $\varphi_i^{\tilde{A}_n}$ .

□

## 4.3 Proof of the Chevalley theorem

At this stage, the principal theorem can be state in precise way as follows.

**Theorem 4.7.** The trigraded algebra of Jacobi forms  $J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)} = \bigoplus_{k,l,m} J_{k,l,m}^{\tilde{A}_n}$  is freely generated by  $n+1$  fundamental Jacobi forms  $(\varphi_0^{\tilde{A}_n}, \varphi_1^{\tilde{A}_n}, \dots, \varphi_n^{\tilde{A}_n})$  over the graded ring  $E_{\bullet, \bullet}$

$$J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)} = E_{\bullet, \bullet} \left[ \varphi_0^{\tilde{A}_n}, \varphi_1^{\tilde{A}_n}, \varphi_2^{\tilde{A}_n}, \dots, \varphi_n^{\tilde{A}_n} \right]. \quad (4.30)$$

Before proving this Theorem, some auxiliary lemmas will be necessary.

**Proposition 4.8.** [6] The functions  $(\varphi_0, \varphi_2, \dots, \varphi_{n+1})$  obtained by the formula

$$\begin{aligned} \lambda^{\mathcal{J}(A_n)}(z) &= e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(z - v_i, \tau)}{\theta_1^{n+1}(z, \tau)} \\ &= \varphi_{n+1}^{\mathcal{J}(A_n)} \wp^{n-1}(z, \tau) + \varphi_n^{\mathcal{J}(A_n)} \wp^{n-2}(z, \tau) + \dots + \varphi_2^{\mathcal{J}(A_n)} \wp(z, \tau) + \varphi_0^{\mathcal{J}(A_n)}, \end{aligned} \quad (4.31)$$

are Jacobi forms of the Jacobi group  $\mathcal{J}(A_n)$  of weight  $0, -2, \dots, -n-1$  respectively, index 1.

---

**Proposition 4.9.** [6] Let the Jacobi forms  $\varphi_0, \dots, \varphi_{n+1}$  be defined by (4.31), then the lowest term of the Taylor expansion in the variables  $\{v_i\}$  are given by

$$\begin{aligned}
\varphi_{n+1}^{\mathcal{J}(A_n)} &= a_{n+1} + O(\|v\|^n), \\
\varphi_n^{\mathcal{J}(A_n)} &= a_n + O(\|v\|^{n-1}) \\
\varphi_{n-1}^{\mathcal{J}(A_n)} &= a_{n-1} + O(\|v\|^{n-2}) \\
&\vdots \\
&\vdots \\
\varphi_2^{\mathcal{J}(A_n)} &= a_2 + O(\|v\|^4) \\
\varphi_1^{\mathcal{J}(A_n)} &= a_1 = 0 \\
\varphi_0^{\mathcal{J}(A_n)} &= a_0 + O(\|v\|^2)
\end{aligned} \tag{4.32}$$

where  $a_2, a_3, \dots, a_{n+1}$  are the elementary symmetric polynomials defined by the formula

$$\begin{aligned}
\lambda^{A_n}(p) &= \prod_{i=0}^n (p - x_i) \\
&= p^{n+1} + a_2 p^{n-1} + a_3 p^{n-2} + \dots + a_n p + a_{n+1}.
\end{aligned} \tag{4.33}$$

**Lemma 4.10.** Let  $\{\varphi_i^{\tilde{A}_n}\}$  be set of functions given by the formula (4.25), and  $\{\varphi_j^{A_{n+1}}\}$  given by (4.31), then

$$\begin{aligned}
\varphi_{n+2}^{\mathcal{J}(A_{n+1})} &= \varphi_n^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)}, \\
\varphi_{n+1}^{\mathcal{J}(A_{n+1})} &= \varphi_{n-1}^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)} + a_{n-1}^n \varphi_n^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)}, \\
\varphi_{n+2}^{\mathcal{J}(A_{n+1})} &= \varphi_{n-2}^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)} + a_{n-2}^{n-1} \varphi_{n-1}^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)} + a_{n-2}^n \varphi_n^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)}, \\
&\vdots \\
&\vdots \\
\varphi_2^{\mathcal{J}(A_{n+1})} &= \varphi_0^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)} + \sum_{j=1}^n a_0^j \varphi_j^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)}, \\
\varphi_0^{\mathcal{J}(A_{n+1})} &= \sum_{j=0}^n a_{-1}^j \varphi_j^{\tilde{A}_n} \varphi_2^{\mathcal{J}(A_1)}.
\end{aligned} \tag{4.34}$$

where  $\varphi_2^{\mathcal{J}(A_1)}$  is defined on (4.6) for  $z_{n+2} = (n+1)v_{n+1}$ , and  $a_i^j = a_i^j(v_{n+1}, \tau)$  are elliptic functions on  $v_{n+1}$ .

*Proof.* Note the following relation

$$\begin{aligned}
\frac{\lambda^{\mathcal{J}(A_{n+1})}}{\lambda^{\mathcal{J}(\tilde{A}_n)}} &= \frac{\theta_1(z - (n+1)v_{n+1}, \tau) \theta_1(z + (n+1)v_{n+1}, \tau)}{\theta_1^2(z, \tau)} e^{-2\pi i u_2} \\
&= \varphi_2^{\mathcal{J}(A_1)} \wp(z, \tau) - \varphi_2^{\mathcal{J}(A_1)} \wp((n+1)v_{n+1}, \tau)
\end{aligned}$$

Hence,

$$\begin{aligned}
& \varphi_{n+2}^{\mathcal{J}(A_{n+1})} \wp^{n-2}(z, \tau) + \varphi_{n+1}^{\mathcal{J}(A_{n+1})} \wp^{n-3}(z, \tau) + \dots + \varphi_2^{\mathcal{J}(A_{n+1})} \wp(z, \tau) + \varphi_0^{\mathcal{J}(A_{n+1})} \\
&= (\varphi_n^{\mathcal{J}(\tilde{A}_n)} \wp^{n-2}(z, \tau) + \varphi_{n-1}^{\mathcal{J}(\tilde{A}_n)} \wp^{n-3}(z, \tau) + \dots + \varphi_2^{\mathcal{J}(\tilde{A}_n)} \wp(z, \tau) \\
&+ \varphi_1^{\mathcal{J}(\tilde{A}_n)} [\zeta(z, \tau) - \zeta(z + (n+1)v_{n+1}, \tau) + \varphi_0^{\mathcal{J}(\tilde{A}_n)}] (\varphi_2^{\mathcal{J}(A_1)} \wp(z, \tau) - \varphi_2^{\mathcal{J}(A_1)} \wp((n+1)v_{n+1}, \tau)).
\end{aligned} \tag{4.35}$$

Then, the desired result is obtained by doing a Laurent expansion in the variable  $z$  in both side of the equality.  $\square$

As a consequence of the previous lemma, we have

**Corollary 4.10.1.** The Jacobi forms  $\{\varphi_i^{\mathcal{J}(\tilde{A}_n)}\}$  are algebraically independent.

*Proof.* Suppose that there exist polynomial  $h(x_0, x_1, \dots, x_n)$  not identically 0, such that

$$h(\varphi_0^{\mathcal{J}(\tilde{A}_n)}, \varphi_1^{\mathcal{J}(\tilde{A}_n)}, \varphi_2^{\mathcal{J}(\tilde{A}_n)}, \dots, \varphi_n^{\mathcal{J}(\tilde{A}_n)}) = 0$$

then, because  $J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)}$  is graded ring  $h(x_0, x_1, \dots, x_n)$  should be 0 in each homogeneous component  $h_m(x_0, x_1, \dots, x_n)$  of index  $m$ . Let  $\tilde{h}_m := (\varphi_2^{\mathcal{J}(A_1)})^m h_m(\varphi_0^{\mathcal{J}(\tilde{A}_n)}, \varphi_1^{\mathcal{J}(\tilde{A}_n)}, \varphi_2^{\mathcal{J}(\tilde{A}_n)}, \dots, \varphi_n^{\mathcal{J}(\tilde{A}_n)})$ . Let us expand the functions  $\varphi_i^{\mathcal{J}(\tilde{A}_n)}$  in the variables  $v_i$ , then  $\tilde{h}_m$  vanishes iff its vanishes in each order of this expansion.

From equations (4.32), we know that the lowest term of the taylor expansion of  $\varphi_{n+2}^{\mathcal{J}(A_{n+1})}$  are the elementary symmetric polynomials. Using lemma 4.10, we conclude that the lowest term of  $\varphi_j^{\mathcal{J}(\tilde{A}_n)}$  is the same as the lowest term of  $\varphi_{j+2}^{\mathcal{J}(A_{n+1})}$ , but those terms are exactly the elementary symmetric polynomials. But the elementary symmetric polynomials are algebraically independent, then they can not solve any polynomial equation. Lemma proved.  $\square$

**Corollary 4.10.2.**

$$E_{\bullet, \bullet} \left[ \varphi_0^{\mathcal{J}(\tilde{A}_n)}, \varphi_1^{\mathcal{J}(\tilde{A}_n)}, \varphi_2^{\mathcal{J}(\tilde{A}_n)}, \dots, \varphi_n^{\mathcal{J}(\tilde{A}_n)} \right] = E_{\bullet, \bullet} \left[ \frac{\varphi_0^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}}, \frac{\varphi_2^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}}, \dots, \frac{\varphi_n^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}} \right]$$

Moreover, we have the following lemma

**Lemma 4.11.** Let  $\varphi \in J_{\bullet, \bullet, m}^{\mathcal{J}(\tilde{A}_n)}$ , then  $\varphi \in E_{\bullet, \bullet} \left[ \frac{\varphi_0^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}}, \frac{\varphi_2^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}}, \dots, \frac{\varphi_n^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}} \right]$ .

*Proof.* Let  $\varphi \in J_{\bullet, \bullet, m}^{\mathcal{J}(\tilde{A}_n)}$ , then the function  $\frac{\varphi}{\varphi_n^{\mathcal{J}(\tilde{A}_n)}}$  is an elliptic function on the variables  $(v_0, v_1, \dots, v_{n-1}, v_{n+1})$  with poles on  $v_i - v_{n+1}, (n+1)v_{n+1}$ . Expanding the function  $\frac{\varphi}{\varphi_n^{\mathcal{J}(\tilde{A}_n)}}$  in the variables  $v_0, v_1, \dots, v_{n-1}$  we get

$$\begin{aligned}
\frac{\varphi}{\varphi_n^{\mathcal{J}(A_n)}} &= \sum_{i=0}^{n-1} a_m^i \wp^{(m-2)}(v_i - v_{n+1}) + \sum_{i=0}^{n-1} a_{m-1}^i \wp^{(m-3)}(v_i - v_{n+1}) + \dots \\
&+ \sum_{i=0}^{n-1} a_1^i \zeta^{(m-2)}(v_i - v_{n+1}) + b(v_{n+1}, \tau)
\end{aligned}$$

But the function  $\frac{\varphi}{\varphi_n^{(A_n)}}$  is invariant under the permutations of the variables  $v_i$ , then

$$\begin{aligned} \frac{\varphi}{\varphi_n^{(A_n)}} &= a_m \sum_{i=0}^{n-1} \wp^{(m-2)}(v_i - v_{n+1}) + a_{m-1} \sum_{i=0}^{n-1} \wp^{(m-3)}(v_i - v_{n+1}) + .. \\ &\quad + a_1 \sum_{i=0}^{n-1} \zeta^{(m-2)}(v_i - v_{n+1}) + b(v_{n+1}, \tau) \end{aligned} \quad (4.36)$$

Now we complete this function to  $A_{n+1}$  invariant function by summing and subtracting the following function in e.q (4.36)

$$\begin{aligned} f(v_{n+1}, \tau) &= a_m \sum_{i=0}^{n-1} \wp^{(m-2)}((n+1)v_{n+1}) + a_{m-1} \sum_{i=0}^{n-1} \wp^{(m-3)}((n+1)v_{n+1}) + .. \\ &\quad + a_1 \sum_{i=0}^{n-1} \zeta^{(m-2)}((n+1)v_{n+1}) \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\varphi}{\varphi_n^{(A_n)}} &= a_m \left( \sum_{i=0}^{n-1} \wp^{(m-2)}(v_i - v_{n+1}) + \wp^{(m-2)}((n+1)v_{n+1}) \right) \\ &\quad + a_{m-1} \sum_{i=0}^{n-1} (\wp^{(m-3)}(v_i - v_{n+1}) + \wp^{(m-3)}((n+1)v_{n+1})) + .. \\ &\quad + a_1 \sum_{i=0}^{n-1} (\zeta^{(m-2)}(v_i - v_{n+1}) + \zeta^{(m-2)}((n+1)v_{n+1})) + \tilde{b}(v_{n+1}, \tau) \end{aligned} \quad (4.37)$$

To finish the proof note the following

1. The functions  $\varphi_{n+2}^{(A_{n+1})} [\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  are  $A_{n+1}$  by construction,
2. The functions  $\varphi_{n+2}^{(A_{n+1})} [\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  are invariant under the action of  $(\mathbb{Z} \oplus \tau\mathbb{Z})^{2n+2}$ , because  $\varphi_{n+2}^{(A_{n+1})}$  is invariant, and  $\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  are elliptic functions.
3. The functions  $\varphi_{n+2}^{(A_{n+1})} [\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  are equivariant under the action of  $SL_2(\mathbb{Z})$ , because  $\varphi_{n+2}^{(A_{n+1})}$  is equivariant, and  $\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  are elliptic functions.
4. The function  $\varphi_{n+2}^{(A_{n+1})}$  has zeros on  $v_i - v_{n+1}, (n+1)v_{n+1}$  of order  $m$ , and  $\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  has poles on  $v_i - v_{n+1}, (n+1)v_{n+1}$  of order  $j+2 \leq m$ . Then, the functions  $\varphi_{n+2}^{(A_{n+1})} [\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})]$  are holomorphic.
5. We conclude that  $g_j := \varphi_{n+2}^{(A_{n+1})} [\wp^{(j)}(v_i - v_{n+1}) + \wp^{(j)}((n+1)v_{n+1})] \in J_{\bullet, \bullet}^{(A_{n+1})}$ . Hence,

$$\varphi = \sum_{i=1}^m a_i \frac{g_i}{(\varphi_2^{(A_1)})^m} + \tilde{b}(v_{n+1}, \tau) \left( \frac{\varphi_{n+2}^{(A_{n+1})}}{\varphi_2^{(A_1)}} \right)^m \in E_{\bullet, \bullet} \left[ \frac{\varphi_0^{(A_{n+1})}}{\varphi_2^{(A_1)}}, \frac{\varphi_2^{(A_{n+1})}}{\varphi_2^{(A_1)}}, \dots, \frac{\varphi_n^{(A_{n+1})}}{\varphi_2^{(A_1)}} \right]. \quad (4.38)$$

□



*Proof.*

$$J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)} \subset E_{\bullet, \bullet} \left[ \frac{\varphi_0^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}}, \frac{\varphi_2^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}}, \dots, \frac{\varphi_n^{\mathcal{J}(A_{n+1})}}{\varphi_2^{\mathcal{J}(A_1)}} \right] = E_{\bullet, \bullet} \left[ \varphi_0^{\mathcal{J}(\tilde{A}_n)}, \varphi_1^{\mathcal{J}(\tilde{A}_n)}, \varphi_2^{\mathcal{J}(\tilde{A}_n)}, \dots, \varphi_n^{\mathcal{J}(\tilde{A}_n)} \right] \subset J_{\bullet, \bullet, \bullet}^{\mathcal{J}(\tilde{A}_n)}. \quad (4.39)$$

□

## 4.4 Intersection form

In the next sections, we introduce the minimal geometric data to reconstruct a Dubrovin Frobenius manifold structure as it was already announced in the subsection 1.4. The first step of this process is to build the invariant  $\mathcal{J}(A_n)$  invariant sections : intersection form, Unit vector field and Euler vector field, and from this data, we reconstruct the Dubrovin Frobenius structure of the Hurwitz space  $H_{1,n-1,0}$ . In this section, we start by introducing the intersection form of the orbit space of the group  $\mathcal{J}(\tilde{A}_n)$ , which is defined to be a natural extensions of the intersection forms of the orbit space of the extended affine Weyl group  $\tilde{A}_n$  [13], and of the Jacobi group  $\mathcal{J}(A_n)$  [6], [7]. In this section, we prove that the intersection form can be understood as an invariant section of the orbit space of  $\mathcal{J}(\tilde{A}_n)$ . Moreover, we define a new metric on the orbit space of  $\mathcal{J}(\tilde{A}_n)$ , which are closed related with the intersection form and it is denoted by  $\tilde{M}$ , further, we construct a generating function of the coefficients for this new metric  $\tilde{M}$  in the coordinates  $(\varphi_0, \varphi_1, \dots, \varphi_n, v_{n+1}, \tau)$ .

**Remark 4.2.** From this point, we will use  $(\varphi_0, \varphi_1, \dots, \varphi_n)$  to denote the Jacobi forms of the group  $\mathcal{J}(\tilde{A}_n)$  again, since there will not be anymore Jacobi form from others Jacobi groups.

**Definition 4.3.** Let

$$\begin{aligned} g &= \sum_{i=0}^n dv_i^2 \Big|_{\sum_{i=0}^n v_i=0} - n(n+1)dv_{n+1}^2 + 2dud\tau, \\ &= \sum_{i,j=0}^{n-1} A_{ij} dv_i dv_j - n(n+1)dv_{n+1}^2 + 2dud\tau, \\ &= \sum_{i,j=0}^{n+1} g_{ij} dv_i dv_j + 2dud\tau. \end{aligned} \quad (4.40)$$

where  $A_{ij}$  is defined in (4.1). The intersection form is given by

$$g^* = \sum_{i,j=0}^{n-1} A_{ij}^{-1} \frac{\partial}{\partial v_i} \otimes \frac{\partial}{\partial v_j} - \frac{1}{n(n+1)} \frac{\partial}{\partial v_{n+1}} \otimes \frac{\partial}{\partial v_{n+1}} + \frac{\partial}{\partial u} \otimes \frac{\partial}{\partial \tau} + \frac{\partial}{\partial \tau} \otimes \frac{\partial}{\partial u}. \quad (4.41)$$

**Proposition 4.12.** The intersection form (4.40) is invariant under the first two actions of (4.2), and behaves a modular form of weight 2 under the last action of (4.2).

*Proof.* The first action of (4.2) only acts on the variables  $v_i$ , the intersection form is invariant under this action because  $A_{ij} dv_i dv_j$  is invariant under permutation.

Under the second action of (4.2), the differentials transform as:

$$\begin{aligned} dv_i &\mapsto dv_i + \lambda_i d\tau, \\ du &\mapsto du - \langle \lambda, \lambda \rangle d\tau - 2 \sum_{i=0}^{n+1} g_{ij} \lambda_j dv_i, \\ d\tau &\mapsto d\tau. \end{aligned} \tag{4.42}$$

Hence:

$$\begin{aligned} g_{ij} dv_i dv_j &\mapsto g_{ij} dv_i dv_j + 2g_{ij} \lambda_i dv_j d\tau + g_{ij} \lambda_i \lambda_j d\tau^2, \\ 2dud\tau &\mapsto 2dud\tau - \langle \lambda, \lambda \rangle d^2\tau - 2 \sum_{i=0}^{n-1} g^{ij} \lambda_j dv_i d\tau. \end{aligned} \tag{4.43}$$

Then:

$$\sum_{i,j=0}^{n+1} g_{ij} dv_i dv_j + 2dud\tau \mapsto \sum_{i,j=0}^{n+1} g_{ij} dv_i dv_j + 2dud\tau. \tag{4.44}$$

Let us show that the metric has conformal transformation under the third action of transformations (4.2):

$$\begin{aligned} dv_i &\mapsto \frac{dv_i}{c\tau + d} - \frac{cv_i d\tau}{(c\tau + d)^2}, \\ d\tau &\mapsto \frac{d\tau}{(c\tau + d)^2}, \\ du &\mapsto du + \frac{cg_{ij} v_i dv_j}{c\tau + d} - \frac{c\langle v, v \rangle d\tau}{2(c\tau + d)^2}. \end{aligned} \tag{4.45}$$

Then:

$$\begin{aligned} g_{ij} dv_i dv_j &\mapsto \frac{g_{ij} dv_i dv_j}{(c\tau + d)^2} - \frac{2cg_{ij} v_i dv_j d\tau}{(c\tau + d)^3} + \frac{g_{ij} v_i v_j d\tau^2}{(c\tau + d)^4}, \\ 2dud\tau &\mapsto \frac{2dud\tau}{(c\tau + d)^2} + \frac{2cg_{ij} v_i dv_j d\tau}{(c\tau + d)^3} - \frac{c\langle v, v \rangle d\tau^2}{(c\tau + d)^4}. \end{aligned} \tag{4.46}$$

Then,

$$\sum_{i,j=0}^{n+1} g_{ij} dv_i dv_j + 2dud\tau \mapsto \frac{\sum_{i,j=0}^{n+1} g_{ij} dv_i dv_j + 2dud\tau}{(c\tau + d)^2}. \tag{4.47}$$

□

An efficient way to compute all  $g^*(d\varphi_i, d\varphi_j)$  is by collecting all of them in a generating function. Note that  $(d\varphi_i, d\varphi_j)$  is not a Jacobi form, and this fact makes the computation more difficult. Hence, in order to circle around this problem, we define the following coefficients in (4.49).

**Lemma 4.13.** Let  $\varphi_i \in J_{k_i, m_i}^{A_n}$ , then the metric given by

$$\frac{1}{\eta^{2i+2j}} g^*(d\eta^{2i} \varphi_i, d\eta^{2j} \varphi_j) \frac{\partial}{\partial \varphi_i} (\eta^{2i} \cdot) \otimes \frac{\partial}{\partial \varphi_j} (\eta^{2j} \cdot) \tag{4.48}$$

is invariant under the first two actions of (4.2), and behaves a modular form of weight 2 under the last action of (4.2).

Moreover, the coefficients of the metric (4.48)

$$\begin{aligned} M(d\varphi_i, d\varphi_j) &:= \frac{1}{\eta^{2i+2j}} g^*(d\eta^{2i}\varphi_i, d\eta^{2j}\varphi_j) \\ &= g^*(d\varphi_i, d\varphi_j) - 4\pi i g_1(\tau) (k_i m_j + k_j m_i) \varphi_i \varphi_j, \end{aligned} \quad (4.49)$$

belong to  $J_{k_i+k_j-2, m_i+m_j}^{A_n}$ .

*Proof.* The metric (4.48) is invariant under the first two actions of (4.2), because proposition 4.12, and because  $\eta$  do not change under this action. The equivariance with respect to the  $SL_2(\mathbb{Z})$  follows again from proposition 4.12, and from the fact that the transformation laws of  $\eta$  get canceled.

The equation (4.49) follows from the chain rule, from the identity

$$\frac{\eta'}{\eta}(\tau) = g_1(\tau) \quad (4.50)$$

□

**Proposition 4.14.** [6] Let  $E^k$  the space of elliptic function of weight  $k$ . The elliptic connection  $D_\tau : E^K \mapsto E^k$  is linear map defined by

$$D_\tau F(v, \tau) = \partial_\tau F(v, \tau) - 2k g_1(\tau) F(v, \tau) - \frac{1}{2\pi i} \frac{\theta'_1(v, \tau)}{\theta_1(v, \tau)} F'(v, \tau), \quad (4.51)$$

where  $F(v, \tau) \in E^k$ .

In order to compute the coefficient of  $M^*(d\varphi_i, d\varphi_j)$  it will be necessary to define an extended intersection form  $g$ .

**Definition 4.4.** The extended metric  $g$  is defined by

$$\tilde{g} = \sum_{i=0}^n dv_i^2 \Big|_{\sum_{i=0}^n v_i=0} - n(n+1) dv_{n+1}^2 + 2dud\tau + ndp^2 + 4ng_1(\tau) pdpd\tau + 2ng'_1(\tau) p^2 d\tau^2, \quad (4.52)$$

or alternatively,

$$\tilde{g} = \sum_{i=1}^{n+1} dz_i^2 - dz_{n+2}^2 + 2dsd\tau, \quad (4.53)$$

where  $(s, z_1, \dots, z_{n+2}, \tau)$  is given by (4.13). The extended intersection form read as

$$\tilde{g}^* = \sum_{i,j} A_{ij}^{-1} \frac{\partial}{\partial v_i} \otimes \frac{\partial}{\partial v_j} - \frac{1}{n(n+1)} \frac{\partial}{\partial v_{n+1}} \otimes \frac{\partial}{\partial v_{n+1}} + \frac{1}{n} \frac{\partial}{\partial p} \otimes \frac{\partial}{\partial p} + \frac{\partial}{\partial s} \otimes \frac{\partial}{\partial \tau} + \frac{\partial}{\partial \tau} \otimes \frac{\partial}{\partial s} \quad (4.54)$$

The following technical result proved by Bertola in [6] will be useful to prove the subsequent results.

**Proposition 4.15.** [6] The following formula holds

$$\left( \frac{\alpha''(x)}{\alpha(x)} + \frac{\alpha''(y)}{\alpha(y)} - \frac{\alpha'(x)\alpha'(y)}{\alpha(x)\alpha(y)} \right) = -4\pi i \frac{\partial_\tau \alpha(x-y)}{\alpha(x-y)} + 2 \frac{\alpha'(x-y)}{\alpha(x-y)} \left[ \frac{\alpha'(x)}{\alpha(x)} - \frac{\alpha'(y)}{\alpha(y)} \right]. \quad (4.55)$$

where  $\alpha(p)$  is given by the second equation (4.59).

The desired generating function is a consequence of the following lemmas.

---

**Lemma 4.16.** Let  $\Phi(p)$  be given by

$$\Phi(p) = e^{-2\pi i u - 2\pi i p^2 n g_1(\tau)} \prod_{i=1}^{n+1} \frac{\theta_1(z_i, \tau)}{\theta_1'(0, \tau)} \frac{\theta_1'(0, \tau)}{\theta_1(z_{n+2}, \tau)}, \quad (4.56)$$

and  $\widetilde{M}$  the extended modified intersection form

$$\widetilde{M}(d\Phi(p), d\Phi(p')) = \frac{1}{\eta^{4n+4}} \widetilde{g}^* \left( d \left( \eta^{2n+2} \Phi(p) \right), d \left( \eta^{2n+2} \Phi(p') \right) \right), \quad (4.57)$$

then,

$$\begin{aligned} e^{2\pi i n(p^2 + p'^2)} \widetilde{M}(d\Phi(p), d\Phi(p')) &= \\ &= 2\pi i n \frac{\nabla_\tau \alpha(p - p')}{\alpha(p - p')} + \frac{\alpha'(p - p')}{\alpha(p - p')} \left[ P(p) \frac{dP(p')}{dp'} - P(p') \frac{dP(p)}{dp} \right], \end{aligned} \quad (4.58)$$

where

$$\begin{aligned} \nabla_\tau F(v, \tau) &= \frac{1}{\eta^{2k}} \frac{\partial (\eta^{2k} F)}{\partial \tau}, \quad F \in E^k, \\ \alpha(p) &= \frac{\theta_1(p, \tau)}{\theta_1'(0, \tau)}, \\ P(p) &= e^{-2\pi i u} \prod_{i=1}^{n+1} \frac{\theta_1(z_i, \tau)}{\theta_1'(0, \tau)} \frac{\theta_1'(0, \tau)}{\theta_1(z_{n+2}, \tau)}. \end{aligned} \quad (4.59)$$

*Proof.*

$$\begin{aligned} e^{2\pi i n g_1(\tau)(p^2 + p'^2)} \widetilde{M}(d\Phi(p), d\Phi(p')) &= \\ &= e^{2\pi i g_1(\tau)n(p^2 + p'^2)} \widetilde{M} \left( d \left( e^{-2\pi i n g_1(\tau)p^2} P(p) \right), d \left( e^{-2\pi i n g_1(\tau)p'^2} P(p') \right) \right) \end{aligned} \quad (4.60)$$

Note that

$$\begin{aligned} \frac{\partial}{\partial p} &= \sum_{i=1}^{n+2} \frac{\partial}{\partial z_i} + 2n p g_1(\tau) \frac{\partial}{\partial u}, \\ \frac{\partial}{\partial v_{n+1}} &= \sum_{i=1}^{n+1} \frac{\partial}{\partial z_i} + (n+1) \frac{\partial}{\partial z_{n+2}}, \\ \frac{\partial}{\partial \tau} &= \frac{\partial}{\partial \tilde{\tau}} + n p^2 g_1'(\tau) \frac{\partial}{\partial u}. \end{aligned}$$

Hence,

$$\begin{aligned} e^{2\pi i n g_1(\tau)p^2} \frac{\partial}{\partial p} \left( e^{-2\pi i n g_1(\tau)p^2} P(p) \right) &= \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} - \frac{\partial P(p)}{\partial z_{n+2}}, \\ e^{2\pi i n g_1(\tau)p^2} \frac{\partial}{\partial \tau} \left( e^{-2\pi i n g_1(\tau)p^2} P(p) \right) &= \frac{\partial P(p)}{\partial \tilde{\tau}}. \end{aligned} \quad (4.61)$$

Substituting (4.61) in (4.60) we get

$$\begin{aligned}
& e^{2\pi i g_1(\tau)(p^2+p'^2)} \widetilde{M}(d\Phi(p), d\Phi(p')) = \\
& = \sum_{i,j=0}^{n-1} A_{ij}^{-1} \frac{\partial P(p)}{\partial v_i} \frac{\partial P(p')}{\partial v_j} + \frac{1}{n} \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \sum_{i=1}^{n+1} \frac{\partial P(p')}{\partial z_i} - \frac{1}{n(n+1)} \frac{\partial P(p)}{\partial v_{n+1}} \frac{\partial P(p')}{\partial v_{n+1}} \\
& - \frac{1}{n} \frac{\partial P(p)}{\partial z_{n+2}} \sum_{i=1}^{n+1} \frac{\partial P(p')}{\partial z_i} - \frac{1}{n} \frac{\partial P(p')}{\partial z_{n+2}} \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} + \frac{1}{n} \frac{\partial P(p)}{\partial z_{n+2}} \frac{\partial P(p')}{\partial z_{n+2}} \\
& - 2\pi i P(p) \nabla_\tau P(p') - 2\pi i P(p') \nabla_\tau P(p) \\
& = \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \frac{\partial P(p')}{\partial z_i} - 2\pi i P(p) \nabla_\tau P(p') - 2\pi i P(p') \nabla_\tau P(p).
\end{aligned} \tag{4.62}$$

Using the following identity in (4.62)

$$\begin{aligned}
& \sum_{i,j=0}^{n-1} A_{ij}^{-1} \frac{\partial P(p)}{\partial v_i} \frac{\partial P(p')}{\partial v_j} + \frac{1}{n} \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \sum_{i=1}^{n+1} \frac{\partial P(p')}{\partial z_i} \\
& = \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \frac{\partial P(p')}{\partial z_i} + \frac{1}{n(n+1)} \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \sum_{i=1}^{n+1} \frac{\partial P(p')}{\partial z_i},
\end{aligned} \tag{4.63}$$

we get

$$\begin{aligned}
& e^{2\pi i g_1(\tau)(p^2+p'^2)} \widetilde{M}(d\Phi(p), d\Phi(p')) = \\
& = \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \frac{\partial P(p')}{\partial z_i} - \frac{\partial P(p)}{\partial z_{n+2}} \frac{\partial P(p')}{\partial z_{n+2}} - 2\pi i P(p) \nabla_\tau P(p') - 2\pi i P(p') \nabla_\tau P(p),
\end{aligned} \tag{4.64}$$

we now compute

$$\begin{aligned}
& \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \frac{\partial P(p')}{\partial z_i} - 2\pi i P(p) \nabla_\tau P(p') - 2\pi i P(p') \nabla_\tau P(p) = \\
& = \sum_{i=1}^{n+1} \frac{\partial P(p)}{\partial z_i} \frac{\partial P(p')}{\partial z_i} - \frac{2\pi i}{\eta^{2n}} \left[ P(p) \frac{\partial (\eta^{2n} P(p'))}{\partial \tau} + P(p') \frac{\partial (\eta^{2n} P(p))}{\partial \tau} \right] \\
& = -\frac{1}{2} \left( \sum_{i=1}^{n+1} \frac{\alpha''(z_i)}{\alpha(z_i)} + \frac{\alpha''(w_i)}{\alpha(w_i)} - 2 \frac{\alpha'(z_i) \alpha'(w_i)}{\alpha(z_i) \alpha(w_i)} \right) P(p) P(p') + 4\pi i g_1 n P(p) P(p') \\
& + \frac{1}{2} \left( \frac{\alpha''(z_{n+2})}{\alpha(z_{n+2})} + \frac{\alpha''(w_{n+2})}{\alpha(w_{n+2})} - 2 \frac{\alpha'(z_{n+2}) \alpha'(w_{n+2})}{\alpha(z_{n+2}) \alpha(w_{n+2})} \right) P(p) P(p'),
\end{aligned} \tag{4.65}$$

where  $z_i := z_i(v_i, p)$ ,  $w_i := z_i(v_i, p')$ . Substituting (4.55) in (4.65)

$$\begin{aligned}
& = -\frac{1}{2} \sum_{i=1}^{n+1} \left( \frac{\alpha''(z_i)}{\alpha(z_i)} + \frac{\alpha''(w_i)}{\alpha(w_i)} - 2 \frac{\alpha'(z_i) \alpha'(w_i)}{\alpha(z_i) \alpha(w_i)} \right) P(p) P(p') + 4\pi i g_1 n P(p) P(p') \\
& + \frac{1}{2} \left( \frac{\alpha''(z_{n+2})}{\alpha(z_{n+2})} + \frac{\alpha''(w_{n+2})}{\alpha(w_{n+2})} - 2 \frac{\alpha'(z_{n+2}) \alpha'(w_{n+2})}{\alpha(z_{n+2}) \alpha(w_{n+2})} \right) P(p) P(p') \\
& = 2\pi i n \frac{\nabla_\tau \alpha(p-p')}{\alpha(p-p')} + \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ P(p) \frac{dP(p')}{dp'} - P(p') \frac{dP(p)}{dp} \right].
\end{aligned} \tag{4.66}$$

□

---

**Lemma 4.17.** For the coefficients  $\tilde{M}$  of intersection form, we have the following formula

$$\begin{aligned}
& \sum_{k,j} C_k(p) C_j(p') \tilde{M}(d\varphi_k, d\varphi_j) = \\
& = 2\pi i n \frac{\nabla_\tau \alpha(p-p')}{\alpha(p-p')} \lambda(p) \lambda(p') + \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ P(p) \frac{dP(p')}{dp'} - P(p') \frac{dP(p)}{dp} \right] \\
& - \sum_{k,j} \tilde{M}(dC_k(p), dC_j(p')) \varphi_{n+1-k} \varphi_{n+1-j}
\end{aligned} \tag{4.67}$$

where  $C_k(p)$  is given by

$$\begin{aligned}
C_k(p) &= \frac{(-1)^k}{(k-1)!} \alpha^n(p) \wp^{(k-2)}(p), \\
\wp^{-1}(p) &= \zeta(p) - \zeta(p + (n+1)v_{n+1}) + (n+1)\zeta(v_{n+1}), \\
\wp^{-2}(p) &= 1,
\end{aligned} \tag{4.68}$$

and

$$\begin{aligned}
\tilde{M}(dC_k(p), dC_j(p')) &:= -2\pi i (C_k(p) \nabla_\tau C_j(p') + C_j(p') \nabla_\tau C_k(p)) + \frac{1}{n} \frac{\partial C_k(p)}{\partial p} \frac{\partial C_j(p')}{\partial p'} \\
&- \frac{1}{n(n+1)} \frac{\partial C_k(p)}{\partial v_{n+1}} \frac{\partial C_j(p')}{\partial v_{n+1}} + \sum_{k,j=0}^n C_k(p) C_j(p') \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial \varphi_k}{\partial v_{n+1}}.
\end{aligned} \tag{4.69}$$

*Proof.* Note that  $P(p) = \sum_{k=0}^{n+1} C_k(p) \varphi_k$ , then

$$\begin{aligned}
& e^{2\pi i n(p^2+p'^2)} \tilde{M}(d\Phi(p), d\Phi(p')) = \\
& = \sum_{j,k} \tilde{M}(d(C_k(p)\varphi_k), d(C_j(p')\varphi_j)) \\
& = \sum_{j,k} C_k(p) C_j(p') \tilde{M}(d\varphi_k, d\varphi_j) + \sum_{j,k} C_j(p') \varphi_k \tilde{M}(dC_k(p), d\varphi_j) \\
& + \sum_{j,k} C_k(p) \varphi_j \tilde{M}(d\varphi_k, dC_j(p')) + \sum_{j,k} \varphi_k \varphi_j \tilde{M}(dC_k(p), dC_j(p')) \\
& = \sum_{j,k} C_k(p) C_j(p') \tilde{M}(d\varphi_k, d\varphi_j) - 2\pi i \sum_{j,k} C_j(p') \varphi_k \varphi_j \nabla_\tau C_k(p) \\
& - 2\pi i \sum_{j,k} C_k(p) \varphi_j \varphi_k \nabla_\tau C_j(p') + \frac{1}{n} \sum_{j,k} \varphi_k \varphi_j \frac{\partial C_k(p)}{\partial p} \frac{\partial C_j(p')}{\partial p'} \\
& - \frac{1}{n(n+1)} \sum_{j,k} \varphi_k \varphi_j \frac{\partial C_k(p)}{\partial v_{n+1}} \frac{\partial C_j(p')}{\partial v_{n+1}} - \frac{1}{n(n+1)} \sum_{j,k} C_k(p) \varphi_j \frac{\partial \varphi_k}{\partial v_{n+1}} \frac{\partial C_j(p')}{\partial v_{n+1}} \\
& - \frac{1}{n(n+1)} \sum_{j,k} C_j(p') \varphi_k \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial C_k(p)}{\partial v_{n+1}}.
\end{aligned} \tag{4.70}$$

Then, isolating  $\sum_{k,j} C_k(p) C_j(p') \tilde{M}(d\varphi_k, d\varphi_j)$

$$\begin{aligned}
& \sum_{k,j} C_k(p) C_j(p') \tilde{M}(d\varphi_k, d\varphi_j) = \\
& = 2\pi i n \frac{\nabla_\tau \alpha(p-p')}{\alpha(p-p')} \lambda(p) \lambda(p') + \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ P(p) \frac{dP(p')}{dp'} - P(p') \frac{dP(p)}{dp} \right] \\
& - \sum_{k,j} \tilde{M}(dC_k(p), dC_j(p')) \varphi_{n+1-k} \varphi_{n+1-j}.
\end{aligned} \tag{4.71}$$

□

**Theorem 4.18.** The coefficient of  $M^*(d\varphi_i, d\varphi_j)$  is recovered by the generating formula

$$\begin{aligned}
& \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} M^*(d\varphi_i, d\varphi_j) \wp(v)^{(k-2)} \wp(v')^{(j-2)} = \\
& = 2\pi i (\lambda(v') D_\tau \lambda(v) + \lambda(v) D_\tau \lambda(v')) - \frac{1}{n+1} \frac{d\lambda(v)}{dv} \frac{d\lambda(v')}{dv'} \\
& + \frac{1}{2} \frac{\wp'(v) + \wp'(v')}{\wp(v) - \wp(v')} \left[ \lambda(v) \frac{d\lambda(v')}{dv'} - \frac{d\lambda(v)}{dv} \lambda(v') \right] - \frac{1}{n} \lambda'(p) \lambda'(p') \\
& - \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial \lambda(p')}{\partial v_{n+1}} + \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial \varphi_k}{\partial v_{n+1}}.
\end{aligned} \tag{4.72}$$

*Proof.* We start by dividing the expression (4.71) by  $\alpha^n(p)\alpha^n(p')$

$$\begin{aligned}
& \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} M^*(d\varphi_i, d\varphi_j) \wp(v)^{(k-2)} \wp(v')^{(j-2)} = \\
& = 2\pi i n \frac{\nabla_\tau \alpha(p-p')}{\alpha(p-p')} \lambda(p) \lambda(p') + \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ \lambda(p) \frac{dP(p')}{dp'} \frac{1}{\alpha^n(p')} - \lambda(p') \frac{dP(p)}{dp} \frac{1}{\alpha^n(p)} \right] \\
& - \sum_{k,j} \frac{\widetilde{M}(dC_k(p), dC_j(p'))}{\alpha^n(p)\alpha^n(p')} \varphi_{n+1-k} \varphi_{n+1-j} \\
& = (1) - (2).
\end{aligned} \tag{4.73}$$

Computing separately

$$\begin{aligned}
(1) & := 2\pi i n \frac{\nabla_\tau \alpha(p-p')}{\alpha(p-p')} \lambda(p) \lambda(p') + \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ \lambda(p) \frac{dP(p')}{dp'} \frac{1}{\alpha^n(p')} - \lambda(p') \frac{dP(p)}{dp} \frac{1}{\alpha^n(p)} \right] = \\
& = n \left( 2\pi i \frac{\nabla_\tau \alpha(p-p')}{\alpha(p-p')} - \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ \frac{\alpha'(p)}{\alpha(p)} - \frac{\alpha'(p')}{\alpha(p')} \right] \right) \lambda(p) \lambda(p') \\
& + \frac{\alpha'(p-p')}{\alpha(p-p')} \left[ \lambda(p) \frac{d\lambda(p')}{dp'} - \lambda(p') \frac{d\lambda(p)}{dp} \right].
\end{aligned} \tag{4.74}$$

and

$$\begin{aligned}
(2) &:= \sum_{k,j} \frac{\widetilde{M}(dC_k(p), dC_j(p'))}{\alpha^n(p)\alpha^n(p')} \varphi_{n+1-k} \varphi_{n+1-j} = \\
&= -2\pi i (\lambda(p) \nabla_\tau \lambda(p') + \lambda(p') \nabla_\tau \lambda(p)) - 2\pi i n \left[ 4g_1 + \frac{\partial_\tau \alpha(p)}{\alpha(p)} + \frac{\partial_\tau \alpha(p')}{\alpha(p')} \right] \lambda(p) \lambda(p') \\
&+ \frac{1}{n} \left[ \frac{\partial \lambda(p)}{\partial p} + n \frac{\alpha'(p)}{\alpha(p)} \lambda(p) \right] \left[ \frac{\partial \lambda(p')}{\partial p'} + n \frac{\alpha'(p')}{\alpha(p')} \lambda(p') \right] \\
&- \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial \lambda(p')}{\partial v_{n+1}} + \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial \varphi_k}{\partial v_{n+1}} \\
&= -2\pi i (\lambda(p) \nabla_\tau \lambda(p') + \lambda(p') \nabla_\tau \lambda(p)) \\
&- n \left[ 8\pi i g_1 + 2\pi i \left( \frac{\partial_\tau \alpha(p)}{\alpha(p)} + \frac{\partial_\tau \alpha(p')}{\alpha(p')} \right) - \frac{\alpha'(p)\alpha'(p')}{\alpha(p)\alpha(p')} \right] \lambda(p) \lambda(p') \\
&+ \frac{1}{n} \frac{\partial \lambda(p)}{\partial p} \frac{\partial \lambda(p')}{\partial p'} + \frac{\alpha'(p)}{\alpha(p)} \lambda(p) \frac{\partial \lambda(p')}{\partial p'} + \frac{\alpha'(p')}{\alpha(p')} \lambda(p') \frac{\partial \lambda(p)}{\partial p} \\
&- \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial \lambda(p')}{\partial v_{n+1}} + \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial \varphi_k}{\partial v_{n+1}} \\
&= -2\pi i (\lambda(p) D_\tau \lambda(p') + \lambda(p') D_\tau \lambda(p)) \\
&- n \left[ 8\pi i g_1 + 2\pi i \left( \frac{\partial_\tau \alpha(p)}{\alpha(p)} + \frac{\partial_\tau \alpha(p')}{\alpha(p')} \right) - \frac{\alpha'(p)\alpha'(p')}{\alpha(p)\alpha(p')} \right] \lambda(p) \lambda(p') \\
&+ \frac{1}{n} \frac{\partial \lambda(p)}{\partial p} \frac{\partial \lambda(p')}{\partial p'} + \left( \frac{\alpha'(p)}{\alpha(p)} - \frac{\alpha'(p')}{\alpha(p')} \right) \left[ \lambda(p) \frac{d\lambda(p')}{dp'} - \lambda(p') \frac{d\lambda(p)}{dp} \right] \\
&- \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial \lambda(p')}{\partial v_{n+1}} + \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial \varphi_k}{\partial v_{n+1}}.
\end{aligned} \tag{4.75}$$

Computing (1)-(2), and using equation (4.55), we obtain

$$\begin{aligned}
&\sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} M^*(d\varphi_i, d\varphi_j) \wp(v)^{(k-2)} \wp(v')^{(j-2)} = \\
&= 2\pi i (\lambda(v') D_\tau \lambda(v) + \lambda(v) D_\tau \lambda(v')) - \frac{1}{n+1} \frac{d\lambda(v)}{dv} \frac{d\lambda(v')}{dv'} \\
&+ \frac{1}{2} \frac{\wp'(v) + \wp'(v')}{\wp(v) - \wp(v')} \left[ \lambda(v) \frac{d\lambda(v')}{dv'} - \frac{d\lambda(v)}{dv} \lambda(v') \right] - \frac{1}{n} \lambda'(p) \lambda'(p') \\
&\frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial \lambda(p')}{\partial v_{n+1}} - \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial \varphi_k}{\partial v_{n+1}}.
\end{aligned} \tag{4.76}$$

□

**Corollary 4.18.1.** Let  $\tilde{\eta}^*(d\varphi_i, d\varphi_j)$  and  $\eta^*(d\varphi_i, d\varphi_j)$  be given by

$$\begin{aligned}
\tilde{\eta}^*(d\varphi_i, d\varphi_j) &:= \frac{\partial M^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0}, \\
\eta^*(d\varphi_i, d\varphi_j) &:= \frac{\partial g^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0}.
\end{aligned} \tag{4.77}$$



The coefficient of  $\tilde{\eta}^*(d\varphi_i, d\varphi_j)$  is recovered by the generating formula

$$\begin{aligned}
& \sum_{k,j=0}^{n+1} \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \tilde{\eta}^*(d\varphi_i, d\varphi_j) \wp(v)^{(k-2)} \wp(v')^{(j-2)} = \\
& = 2\pi i (D_\tau \lambda(v) + D_\tau \lambda(v')) + \frac{1}{2} \frac{\wp'(v) + \wp'(v')}{\wp(v) - \wp(v')} \left[ \frac{d\lambda(v')}{dv'} - \frac{d\lambda(v)}{dv} \right] \\
& - \frac{1}{n(n+1)} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p)}{\partial v_{n+1}} \right) \frac{\partial \lambda(p')}{\partial v_{n+1}} - \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p')}{\partial v_{n+1}} \right) \\
& + \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_j}{\partial v_{n+1}} \right) \frac{\partial \varphi_k}{\partial v_{n+1}} \\
& + \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_k}{\partial v_{n+1}} \right)
\end{aligned} \tag{4.78}$$

Moreover,

$$\begin{aligned}
\tilde{\eta}^*(d\varphi_i, d\varphi_j) &= \eta^*(d\varphi_i, d\varphi_j), \quad i, j \neq 0, \\
\tilde{\eta}^*(d\varphi_0, d\varphi_j) &= \eta^*(d\varphi_i, d\varphi_j) + 4\pi i k_j \varphi_j.
\end{aligned} \tag{4.79}$$

*Proof.* Just differentiate equation (4.72) with respect  $\varphi_0$ , and use the equation (4.25).  $\square$

**Corollary 4.18.2.** The metric  $\tilde{\eta}^*$  and  $\eta^*$  defined in (4.77) is invariant under the second action of (4.2), furthermore, behave as modular form of weight 2 under the last action of (4.2).

*Proof.* The metric  $\tilde{\eta}^*$  and  $\eta^*$  are given by

$$\begin{aligned}
\tilde{\eta}^* &= Lie_{\frac{\partial}{\partial \varphi_0}} M \\
\eta^* &= Lie_{\frac{\partial}{\partial \varphi_0}} g^*
\end{aligned} \tag{4.80}$$

The fact that  $\frac{\partial}{\partial \varphi_0}, g^*, M^*$  is invariant under the second action of (4.2), furthermore, behave as modular form of weight 2 under the last action of (4.2) give the desired result.  $\square$

## 4.5 The second metric of the pencil

In the (4.77), it was defined the second metric  $\eta$ , furthermore, it was derived a generating function for it. The main goal of this section is to extract the coefficients  $\eta(d\varphi_i, d\varphi_j)$  from its generating function. In order to do this extraction, some auxiliaries lemmas are needed.

**Lemma 4.19.** Let  $\varphi_1, \varphi_n$  be defined on (4.25), then

$$\begin{aligned}
\varphi_1 &= e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(v_i + n v_{n+1})}{\theta_1^n((n+1)v_{n+1}) \theta_1'(0)}, \\
\varphi_n &= e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(-v_i + v_{n+1})}{\theta_1(-(n+1)v_{n+1}) \theta_1'(0)^n}.
\end{aligned} \tag{4.81}$$

*Proof.*

$$\begin{aligned}
\varphi_1 &= \lim_{p \rightarrow -(n+1)v_{n+1}} (p + (n+1)v_{n+1}) \lambda(p) = -e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1^n(-v_i - n v_{n+1})}{\theta_1^n(-(n+1)v_{n+1}) \theta_1'(0)} = e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(v_i + n v_{n+1})}{\theta_1^n((n+1)v_{n+1}) \theta_1'(0)}, \\
\varphi_n &= \lim_{p \rightarrow 0} p^n \lambda(p) = e^{2\pi i u} \frac{\prod_{i=0}^n \theta_1(-v_i + v_{n+1})}{\theta_1(-(n+1)v_{n+1}) \theta_1'(0)^n}.
\end{aligned}$$

□

**Lemma 4.20.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$  be defined on (4.25), then

$$\begin{aligned}\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_i}{\partial v_{n+1}} \right) &= 0, \quad i > 1, \\ \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_1}{\partial v_{n+1}} \right) &= n, \\ \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_0}{\partial v_{n+1}} \right) &= -n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})}.\end{aligned}\tag{4.82}$$

*Proof.* Computing  $\frac{\partial \varphi_n}{\partial v_{n+1}}$  by using the first equation of (4.81)

$$\frac{\partial \varphi_n}{\partial v_{n+1}} = \left( \sum_{i=0}^n \frac{\theta'_1(-v_i + v_{n+1})}{\theta_1(-v_i + v_{n+1})} - (n+1) \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right) \varphi_n.\tag{4.83}$$

Recall the recursive relation between  $\{\varphi_i\}$  in (4.18)

$$\varphi_i = \frac{\partial^{(n-i)} \varphi_n(p)}{\partial p^{n-i}} \Big|_{p=0}.\tag{4.84}$$

In particular,

$$\varphi_{n-1} = \frac{\partial \varphi_n(p)}{\partial p} \Big|_{p=0} = \left( \sum_{i=0}^n \frac{\theta'_1(-v_i + v_{n+1})}{\theta_1(-v_i + v_{n+1})} - \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right) \varphi_n,\tag{4.85}$$

Then,

$$\frac{\partial \varphi_n}{\partial v_{n+1}} = \varphi_{n-1} - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_n,\tag{4.86}$$

consequently

$$\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_n}{\partial v_{n+1}} \right) = 0.$$

Suppose that for  $i > 1$ , we have

$$\varphi_i = f(-v_0 + v_{n+1}, -v_1 + v_{n+1}, \dots, (n+1)v_{n+1}, \tau) \varphi_n,\tag{4.87}$$

where  $f(-v_0 + v_{n+1}, -v_1 + v_{n+1}, \dots, (n+1)v_{n+1}, \tau)$  is an elliptic function on the variables  $v_0, v_1, \dots, v_{n+1}$  with zeros on  $-v_i + v_{n+1}$  and poles on  $(n+1)v_{n+1}$ . Consider the extended  $\varphi_i(p)$  as

$$\varphi_i(p) = f(p - v_0 + v_{n+1}, p - v_1 + v_{n+1}, \dots, p + (n+1)v_{n+1}, \tau) \varphi_n.$$

The action of the vector fields  $\frac{\partial}{\partial p}$  and  $\frac{\partial}{\partial v_{n+1}}$  in  $\varphi_i(p)$  and  $\varphi_i$  are given by

$$\begin{aligned}\frac{\partial \varphi_i(p)}{\partial p} \Big|_{p=0} &= \frac{\partial f}{\partial p} \Big|_{p=0} \varphi_n + f \frac{\partial \varphi_n(p)}{\partial p} \Big|_{p=0}, \\ \frac{\partial \varphi_i}{\partial v_{n+1}} &= \frac{\partial f}{\partial v_{n+1}} \varphi_n + f \frac{\partial \varphi_n}{\partial v_{n+1}}.\end{aligned}$$

Note that

$$\frac{\partial f}{\partial p} \Big|_{p=0} - \frac{\partial f}{\partial v_{n+1}} = b(v_{n+1}, \tau),$$

where  $b((n+1)v_{n+1}, \tau)$  is an elliptic function on  $(n+1)v_{n+1}$ , because  $\frac{\partial f}{\partial p}\big|_{p=0}$  and  $\frac{\partial f}{\partial v_{n+1}}$  are elliptic functions with the same Laurent tail in the variables  $-v_i + v_{n+1}$ . Hence, using equation (4.84)

$$\begin{aligned}\frac{\partial \varphi_i}{\partial v_{n+1}} &= \frac{\partial \varphi_i(p)}{\partial p}\bigg|_{p=0} + h(v_{n+1}, \tau)\varphi_n, \\ &= \varphi_{i-1} + h(v_{n+1}, \tau)\varphi_n.\end{aligned}$$

Then,

$$\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_i}{\partial v_{n+1}} \right) = 0, \quad i > 1,$$

Computing  $\varphi_0$

$$\begin{aligned}\varphi_0 &= \frac{\partial \varphi_1}{\partial p}\bigg|_{p=0} \\ &= \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right] \varphi_1.\end{aligned}$$

Computing  $\frac{\partial \varphi_1}{\partial v_{n+1}}$  in terms of  $\varphi_0$

$$\begin{aligned}\frac{\partial \varphi_1}{\partial v_{n+1}} &= \left[ \sum_{i=0}^n n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - n(n+1) \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right] \varphi_1 \\ &= n\varphi_0 - n \frac{\theta_1((n+1)v_{n+1})}{\theta'_1((n+1)v_{n+1})} \varphi_1.\end{aligned}\tag{4.88}$$

Then,

$$\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_1}{\partial v_{n+1}} \right) = n.\tag{4.89}$$

Computing  $\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_0}{\partial v_{n+1}} \right)$

$$\begin{aligned}\frac{\partial \varphi_0}{\partial v_{n+1}} &= \left[ n \sum_{i=0}^n \frac{\partial^2 \log \theta_1(v_i + nv_{n+1})}{\partial v_{n+1}^2} - n(n+1) \frac{\partial^2 \log \theta_1((n+1)v_{n+1})}{\partial v_{n+1}^2} \right] \varphi_1 \\ &\quad + \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right] \frac{\partial \varphi_1}{\partial v_{n+1}} \\ &= \left[ n \sum_{i=0}^n \frac{\partial^2 \log \theta_1(v_i + nv_{n+1})}{\partial v_{n+1}^2} - n(n+1) \frac{\partial^2 \log \theta_1((n+1)v_{n+1})}{\partial v_{n+1}^2} \right] \varphi_1 \\ &\quad + \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right] \left[ -n\varphi_0 - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_1 \right] \\ &= \left[ n \sum_{i=0}^n \frac{\partial^2 \log \theta_1(v_i + nv_{n+1})}{\partial v_{n+1}^2} - n(n+1) \frac{\partial^2 \log \theta_1((n+1)v_{n+1})}{\partial v_{n+1}^2} \right] \varphi_1 \\ &\quad - n \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right]^2 \varphi_1 - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_0 \\ &= - \frac{\partial^2 \log \theta_1((n+1)v_{n+1})}{\partial v_{n+1}^2} \varphi_1 - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_0 + na\varphi_1.\end{aligned}\tag{4.90}$$

where  $a$  is defined by

$$a = \sum_{i=0}^n \frac{\partial^2 \log \theta_1(v_i + nv_{n+1})}{\partial v_{n+1}^2} - \frac{\partial^2 \log \theta_1((n+1)v_{n+1})}{\partial v_{n+1}^2} - \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right]^2. \quad (4.91)$$

Note that  $na\varphi_1$  can not be proportional to  $b\varphi_0$ , for any  $b = b(v_{n+1}, \tau)$  elliptic function in the variable  $v_{n+1}$ . Indeed, if

$$na\varphi_1 = b \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right] \varphi_1,$$

we obtain,

$$na = b \left[ \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right]. \quad (4.92)$$

Analysing the Laurent tail in  $v_i + nv_{n+1}$  of (4.91)

$$a = -2 \sum_{i \leq j} \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} \frac{\theta'_1(v_j + nv_{n+1})}{\theta_1(v_j + nv_{n+1})} + 2 \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \sum_{i=0}^n \frac{\theta'_1(v_i + nv_{n+1})}{\theta_1(v_i + nv_{n+1})} + \text{regular terms}, \quad (4.93)$$

then, the first term of the equation (4.93) implies that the left-hand side and right hand side of the equation (4.92) have a different Laurent tail which is an absurd. Hence,

$$\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_0}{\partial v_{n+1}} \right) = -n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})}.$$

□

**Corollary 4.20.1.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$  be defined on (4.25) and the metric  $\eta^*$  defined in (4.77), then

$$\begin{aligned} \eta^*(d\varphi_i, dv_{n+1}) &= 0, \quad i > 1, \\ \eta^*(d\varphi_1, dv_{n+1}) &= -\frac{1}{n+1}, \\ \eta^*(d\varphi_0, dv_{n+1}) &= -\frac{1}{n+1} \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})}. \end{aligned} \quad (4.94)$$

*Proof.*

$$\begin{aligned} \eta^*(d\varphi_i, dv_{n+1}) &= \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_i}{\partial v_{n+1}} \right) = 0, \quad i > 1, \\ \eta^*(d\varphi_1, dv_{n+1}) &= -\frac{1}{n(n+1)} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_1}{\partial v_{n+1}} \right) = -\frac{1}{n+1}, \\ \eta^*(d\varphi_0, dv_{n+1}) &= -\frac{1}{n(n+1)} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_0}{\partial v_{n+1}} \right) = -\frac{1}{n+1} \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})}. \end{aligned}$$

□

**Lemma 4.21.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$  be defined on (4.25) and the metric  $\eta^*$  defined in (4.77), then

$$\begin{aligned} \eta^*(d\varphi_i, d\tau) &= 0, \quad i \neq 0, \\ \eta^*(d\varphi_0, d\tau) &= -2\pi i. \end{aligned} \quad (4.95)$$

*Proof.*

$$\eta^*(d\varphi_i, d\tau) = -2\pi i \frac{\partial \varphi_i}{\partial \varphi_0} = \delta_{i0}.$$

□

**Theorem 4.22.** Let  $\eta^*(d\varphi_i, d\varphi_j)$  be defined in (4.77), then its coefficients can be obtained by the formula

$$\begin{aligned} \tilde{\eta}^*(d\varphi_i, d\varphi_j) &= (i+j-2)\varphi_{i+j-2}, \quad i, j \neq 0 \\ \tilde{\eta}^*(d\varphi_i, d\varphi_0) &= 0, \quad i \neq 0, \quad i \neq 1, \\ \tilde{\eta}^*(d\varphi_1, d\varphi_j) &= 0, \quad j \neq 0, \\ \tilde{\eta}^*(d\varphi_1, d\varphi_0) &= \wp((n+1)v_{n+1})\varphi_1. \end{aligned} \tag{4.96}$$

*Proof.* We start by dividing the right hand side of the expression (4.78) in two parts:

$$\begin{aligned} (a) &:= 2\pi i (D_\tau \lambda(v) + D_\tau \lambda(v')) + \frac{1}{2} \frac{\wp'(v) + \wp'(v')}{\wp(v) - \wp(v')} \left[ \frac{d\lambda(v')}{dv'} - \frac{d\lambda(v)}{dv} \right], \\ (b) &:= \frac{1}{n(n+1)} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p)}{\partial v_{n+1}} \right) \frac{\partial \lambda(p')}{\partial v_{n+1}} + \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p')}{\partial v_{n+1}} \right) \\ &\quad - \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_j}{\partial v_{n+1}} \right) \frac{\partial \varphi_k}{\partial v_{n+1}} \\ &\quad - \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_k}{\partial v_{n+1}} \right). \end{aligned} \tag{4.97}$$

Consider the equation (4.25) written in a concise way as follows

$$\lambda(v) = \sum_{k=0}^n \frac{(-1)^k}{(k-1)!} \varphi_k \wp^{k-2}(v). \tag{4.98}$$

Substituting (4.98) in the first equation of (4.97)

$$\begin{aligned} (a) &= \sum_{k=0}^n \frac{(-1)^k}{(k-1)!} \varphi_k \left[ 2\pi i (D_\tau \wp^{n-1-k}(v) + D_\tau \wp^{n-1-k}(v')) \right] \\ &\quad + \sum_{k=0}^n \frac{(-1)^k}{(k-1)!} \varphi_k \left[ (\zeta(v-v') + \zeta(v') - \zeta(v)) (\wp^{n-k}(v') - \wp^{n-k}(v)) \right]. \end{aligned} \tag{4.99}$$

Dividing the expression once more

$$\begin{aligned} (a)_1 &= \sum_{k=2}^n \frac{(-1)^k}{(k-1)!} \varphi_k \left[ 2\pi i (D_\tau \wp^{n-1-k}(v) + D_\tau \wp^{n-1-k}(v')) \right] \\ &\quad + \sum_{k=2}^n \frac{(-1)^k}{(k-1)!} \varphi_k \left[ (\zeta(v-v') + \zeta(v') - \zeta(v)) (\wp^{n-k}(v') - \wp^{n-k}(v)) \right], \\ (a)_2 &:= \varphi_1 \left[ 2\pi i (D_\tau \wp^{-1}(v) + D_\tau \wp^{-1}(v')) \right] \\ &\quad + \varphi_1 \left[ (\zeta(v-v') + \zeta(v') - \zeta(v)) (\wp^{-1}(v') - \wp^{-1}(v)) \right]. \end{aligned} \tag{4.100}$$

---

Expanding the left-hand side of (4.78), we get

$$\begin{aligned} & \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial M^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \\ &= \sum_{k,j=0}^n \frac{\partial M^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0} \frac{1}{v^k (v')^j} + \text{Other terms}, \end{aligned} \quad (4.101)$$

where "Other terms" in the equation (4.101) means positive powers of either  $v$  or  $v'$ . For convenience, define

$$\begin{aligned} (1) : &= 2\pi i \left( D_\tau \wp^{(k-2)}(v) + D_\tau \wp^{(j-2)}(v') \right) \\ &+ (\zeta(v - v') + \zeta(v') - \zeta(v)) \left( \wp^{(j-1)}(v') - \wp^{(k-1)}(v) \right) \end{aligned} \quad (4.102)$$

In order to better to compute (4.102), consider the analytical behaviour of the term

$$D_\tau \wp^{k-2}(v) = \partial_\tau \wp^{k-2}(v) - 2(k-2)g_1(\tau) \wp^{k-2}(v) - \frac{1}{2\pi i} \frac{\theta'_1(v, \tau)}{\theta_1(v, \tau)} \wp^{k-1}(v). \quad (4.103)$$

The term  $\partial_\tau \wp^{k-2}(v)$  in (4.103) is holomorphic, therefore, it does not contribute for the Laurent tail. The term

$$2(k-2)g_1(\tau) \wp^{k-2}(v) \quad (4.104)$$

also do not contribute, because the full expression (4.103) behaves as modular form under the  $SL_2(\mathbb{Z})$ , but (4.104) is clear a quasi-modular form, since it contains  $g_1(\tau)$ . Hence, (4.104) is canceled with the Laurent tail of

$$\frac{1}{2\pi i} \frac{\theta'_1(v, \tau)}{\theta_1(v, \tau)} \wp^{k-1}(v). \quad (4.105)$$

To sum up, the analytical behaviour avior of (4.103) is essentially given by (4.105), under this consideration, and by using the equation

$$\begin{aligned} \zeta(v, \tau) &= \frac{\theta'_1(v, \tau)}{\theta_1(v, \tau)} - 4\pi i g_1(\tau) v \\ &= \frac{1}{v} + O(v^3), \end{aligned} \quad (4.106)$$

the equation (4.102) became

---


$$\begin{aligned}
(1) &= -\zeta(v)\wp^{k-1}(v) - \zeta(v')\wp^{k-1}(v') \\
&+ (\zeta(v-v') + \zeta(v') - \zeta(v)) (\wp^{k-1}(v') - \wp^{k-1}(v)) + \text{Other terms} \\
&= \zeta(v-v') (\wp^{k-1}(v') - \wp^{k-1}(v)) - \zeta(v)\wp^{k-1}(v') - \zeta(v')\wp^{k-1}(v) + \text{Other terms} \\
&= \frac{1}{v-v'} \left( \frac{(-1)^k k!}{v'^{k+1}} - \frac{(-1)^k k!}{v^{k+1}} \right) - \frac{1}{v} \frac{(-1)^k k!}{v'^{k+1}} - \frac{1}{v'} \frac{(-1)^k k!}{v^{k+1}} + \text{Other terms} \\
&= (-1)^k k! \left( \frac{1}{v-v'} \frac{v^{k+1} - v'^{k+1}}{(v'v)^{k+1}} \right) - \frac{1}{v} \frac{(-1)^k k!}{v'^{k+1}} - \frac{1}{v'} \frac{(-1)^k k!}{v^{k+1}} + \text{Other terms} \\
&= (-1)^k k! \left( \sum_{j=0}^k \frac{v^{k-j} v'^j}{(v'v)^{k+1}} \right) - \frac{1}{v} \frac{(-1)^k k!}{v'^{k+1}} - \frac{1}{v'} \frac{(-1)^k k!}{v^{k+1}} + \text{Other terms} \tag{4.107} \\
&= (-1)^k k! \left( \sum_{j=0}^k \frac{1}{v^{1+j} v'^{k+1-j}} \right) - \frac{1}{v} \frac{(-1)^k k!}{v'^{k+1}} - \frac{1}{v'} \frac{(-1)^k k!}{v^{k+1}} + \text{Other terms} \\
&= (-1)^k k! \left( \sum_{j=1}^k \frac{1}{v^{1+j} v'^{k+1-j}} \right) + \text{Other terms} \\
&= (-1)^k k! \left( \sum_{j=2}^{k+1} \frac{1}{v^j v'^{k+2-j}} \right) + \text{Other terms.}
\end{aligned}$$

Substituting (4.107) in right-hand side of (4.99)

$$\begin{aligned}
&\sum_{k=0}^n \frac{(-1)^k}{(k-1)!} \varphi_k [2\pi i (D_\tau \wp^{k-2}(v) + D_\tau \wp^{k-1}(v'))] \\
&+ \sum_{k=0}^n \frac{(-1)^k}{(k-1)!} \varphi_k [(\zeta(v-v') + \zeta(v') - \zeta(v)) (\wp^{k-1}(v') - \wp^{k-1}(v))] \\
&= \sum_{k=0}^n \sum_{j=2}^{k+1} \frac{(k) \varphi_k}{v^j v'^{k+2-j}} + \text{Other terms} \tag{4.108} \\
&= \sum_{k=0}^n \sum_{j=2}^{n+1} \frac{(k+j) \varphi_{k+j}}{v^j v'^{k+2}} + \text{Other terms} \\
&= \sum_{k=2}^{n+2} \sum_{j=2}^{n+1} \frac{(k+j-2) \varphi_{k+j-2}}{v^j v'^k} + \text{Other terms.}
\end{aligned}$$

Computing the second expression of (4.100)

$$\begin{aligned}
(a)_2 &:= \varphi_1 [2\pi i (D_\tau \wp^{-1}(v) + D_\tau \wp^{-1}(v'))] \\
&+ \varphi_1 \left[ (\zeta(v - v') + \zeta(v') - \zeta(v)) (\wp^{-1'}(v') - \wp^{-1'}(v)) \right] \\
&= \varphi_1 [\zeta(v) (-\wp(v) + \wp(v + (n+1)v_{n+1})) + \zeta(v') (-\wp(v') + \wp(v' + (n+1)v_{n+1}))] \\
&+ \varphi_1 (\zeta(v - v') + \zeta(v') - \zeta(v)) (\wp(v) - \wp(v + (n+1)v_{n+1})) \\
&+ \varphi_1 (\zeta(v - v') + \zeta(v') - \zeta(v)) (-\wp(v') + \wp(v' + (n+1)v_{n+1})) + \text{Other terms} \\
&= \varphi_1 \zeta(v - v') (\wp(v) - \wp(v + (n+1)v_{n+1}) - \wp(v') + \wp(v' + (n+1)v_{n+1})) \\
&+ \varphi_1 [\zeta(v) (-\wp(v') + \wp(v' + (n+1)v_{n+1})) + \zeta(v') (-\wp(v) + \wp(v + (n+1)v_{n+1}))] + \text{Other terms} \\
&= \varphi_1 \zeta(v - v') (-\wp(v + (n+1)v_{n+1}) + \wp(v' + (n+1)v_{n+1})) \\
&+ \varphi_1 [\zeta(v) \wp(v' + (n+1)v_{n+1}) + \zeta(v') \wp(v + (n+1)v_{n+1})] + \widetilde{(a)}_2 + \text{Other terms},
\end{aligned} \tag{4.109}$$

where  $\widetilde{(a)}_2$  are terms that were already computed in (4.108). To compute the second expression in (4.97), consider

$$\frac{\partial \lambda(v)}{\partial v_{n+1}} = \sum_{k=0}^n \frac{(-1)^k}{(k-1)!} \frac{\partial \varphi_k}{\partial v_{n+1}} \wp^{(k-2)}(v) + (n+1) \varphi_1 [\wp(v + (n+1)v_{n+1}) - \wp((n+1)v_{n+1})], \tag{4.110}$$

and

$$\begin{aligned}
\frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(v)}{\partial v_{n+1}} \right) &= \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_1}{\partial v_{n+1}} \right) [\zeta(v) - \zeta(v + (n+1)v_{n+1}) + \zeta((n+1)v_{n+1})] + \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_0}{\partial v_{n+1}} \right) \\
&= n [\zeta(v) - \zeta(v + (n+1)v_{n+1}) + \zeta((n+1)v_{n+1})] - n \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \\
&= n [\zeta(v) - \zeta(v + (n+1)v_{n+1})].
\end{aligned} \tag{4.111}$$

Substituting (4.110) and (4.111) in the the second expression in (4.97)

$$\begin{aligned}
(b) &:= \frac{1}{n(n+1)} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p)}{\partial v_{n+1}} \right) \frac{\partial \lambda(p')}{\partial v_{n+1}} + \frac{1}{n(n+1)} \frac{\partial \lambda(p)}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \lambda(p')}{\partial v_{n+1}} \right) \\
&- \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_j}{\partial v_{n+1}} \right) \frac{\partial \varphi_k}{\partial v_{n+1}} \\
&- \frac{1}{n(n+1)} \sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial \varphi_j}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_0} \left( \frac{\partial \varphi_k}{\partial v_{n+1}} \right) \\
&= \varphi_1 [\zeta(v') - \zeta(v' + (n+1)v_{n+1})] [\wp(v + (n+1)v_{n+1}) - \wp((n+1)v_{n+1})] \\
&+ \varphi_1 [\zeta(v) - \zeta(v + (n+1)v_{n+1})] [\wp(v' + (n+1)v_{n+1}) - \wp((n+1)v_{n+1})].
\end{aligned} \tag{4.112}$$



Subtracting (4.109) and (4.112)

$$\begin{aligned}
(a)_2 - (b) &= -\varphi_1 [\zeta(v') - \zeta(v' + (n+1)v_{n+1})] [\wp(v + (n+1)v_{n+1}) - \wp((n+1)v_{n+1})] \\
&\quad - \varphi_1 [\zeta(v) - \zeta(v + (n+1)v_{n+1})] [\wp(v' + (n+1)v_{n+1}) - \wp((n+1)v_{n+1})] \\
&\quad + \varphi_1 \zeta(v - v') (-\wp(v + (n+1)v_{n+1}) + \wp(v' + (n+1)v_{n+1})) \\
&\quad + \varphi_1 [\zeta(v)\wp(v' + (n+1)v_{n+1}) + \zeta(v')\wp(v + (n+1)v_{n+1})] + \widetilde{(a)}_2 + \text{Other terms} \\
&= \varphi_1 \wp((n+1)v_{n+1}) [\zeta(v') - \zeta(v' + (n+1)v_{n+1})] + \varphi_1 \wp((n+1)v_{n+1}) [\zeta(v) - \zeta(v + (n+1)v_{n+1})] \\
&\quad + \varphi_1 [\zeta(v - v')] (\wp(v + (n+1)v_{n+1}) + \wp(v' + (n+1)v_{n+1})) \\
&\quad + \varphi_1 [\zeta(v' + (n+1)v_{n+1})] (\wp(v + (n+1)v_{n+1})) \\
&\quad + \varphi_1 [\zeta(v + (n+1)v_{n+1})] (\wp(v' + (n+1)v_{n+1})) \\
&\quad + \widetilde{(a)}_2 + \text{Other terms} \\
&= \varphi_1 \wp((n+1)v_{n+1}) [\zeta(v') - \zeta(v' + (n+1)v_{n+1})] + \varphi_1 \wp((n+1)v_{n+1}) [\zeta(v) - \zeta(v + (n+1)v_{n+1})] \\
&\quad + \widetilde{(a)}_2 + \text{Other terms.}
\end{aligned} \tag{4.113}$$

Summing (4.113) and (4.108), we have

$$\begin{aligned}
&\sum_{k,j=0}^n \frac{(-1)^{k+j}}{(k-1)!(j-1)!} \frac{\partial M^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0} \wp(v)^{(k-2)} \wp(v')^{(j-2)} \\
&= \sum_{k,j=0}^n \frac{\partial M^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0} \frac{1}{v^k (v')^j} + \text{Other terms,} \\
&= \sum_{k=2}^{n+2} \sum_{j=2}^{n+1} \frac{(k+j-2) \varphi_{k+j-2}}{v^j v'^k} \\
&\quad + \varphi_1 \wp((n+1)v_{n+1}) [\zeta(v') - \zeta(v' + (n+1)v_{n+1})] \\
&\quad + \varphi_1 \wp((n+1)v_{n+1}) [\zeta(v) - \zeta(v + (n+1)v_{n+1})] \\
&\quad + \text{Other terms.}
\end{aligned}$$

Hence, we get the desired result.  $\square$

**Corollary 4.22.1.** Let  $\eta^*(d\varphi_i, d\varphi_j)$  be defined in (4.77), then its coefficients can be obtained by the formula

$$\begin{aligned}
\eta^*(d\varphi_i, d\varphi_j) &= (i+j-2)\varphi_{i+j-2}, \quad i, j \neq 0 \\
\eta^*(d\varphi_i, d\varphi_0) &= 0, \quad i \neq 0, \quad i \neq 1, \\
\eta^*(d\varphi_1, d\varphi_j) &= 0, \quad j \neq 0, \\
\eta^*(d\varphi_1, d\varphi_0) &= \frac{\partial^2 \log(\theta_1(n+1)v_{n+1})}{\partial v_{n+1}^2} \varphi_1.
\end{aligned} \tag{4.114}$$

## 4.6 Flat coordinates of $\eta$

This section is dedicated to construct the flat coordinates of  $\eta$  and its relationship with the invariant coordinates  $\varphi_0, \varphi_1, \dots, \varphi_n, v_{n+1}, \tau$ . Our strategy will be an adaptation of the work done in [26]. The

---

flatness of the Saito metric  $\eta$  is proved in the Theorem 4.34.

Let  $t^1, t^2, \dots, t^n$  be given by the following generating function

$$v(z) = \frac{-1}{n} (t^n z + t^{n-1} z^2 + \dots + t^2 z^{n-1} + t^1 z^n + O(z^{n+1})), \quad (4.115)$$

be defined by the following condition

$$\lambda(v) = \frac{1}{z^n}.$$

Moreover,

$$t^0 = \varphi_0 - \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_1 + 4\pi i g_1(\tau) \varphi_2. \quad (4.116)$$

**Lemma 4.23.** The following identity holds

$$\frac{n}{n+1-\alpha} \operatorname{res}_{p=\infty} \left( \lambda^{\frac{n+1-\alpha}{n}}(p) dp \right) = - \operatorname{res}_{\lambda=\infty} \left( v(z) \lambda^{\frac{1-\alpha}{n}} d\lambda \right) \quad (4.117)$$

*Proof.* Consider the integration by parts

$$\frac{n}{n+1-\alpha} \int \left( \lambda^{\frac{n+1-\alpha}{n}}(p) dp \right) = p \lambda^{\frac{n+1-\alpha}{n}} - \int p \lambda^{\frac{1-\alpha}{n}} d\lambda \quad (4.118)$$

Lemma proved. □

**Lemma 4.24.** The functions  $t^1, t^2, \dots, t^n$  defined in (4.115) can be obtained by the formula

$$t^\alpha = - \operatorname{res}_{\lambda=\infty} \left( v(z) \lambda^{\frac{1-\alpha}{n}} d\lambda \right). \quad (4.119)$$

*Proof.* Let

$$z = \left( \frac{1}{\lambda} \right)^{\frac{1}{n}},$$

then,

$$\begin{aligned} v(z) \lambda^{\frac{1-\alpha}{n}} d\lambda &= \left( \frac{1}{n} \right) (t^n z + t^{n-1} z^2 + \dots + t^2 z^{n-1} + t^1 z^n + O(z^{n+1})) z^{\alpha-1} n z^{-n-1} dz \\ &= \left( \sum_{\beta=1}^n t^\beta z^{n+1-\beta} + O(z^{n+1}) \right) z^{\alpha-n-2} dz \\ &= \left( \sum_{\beta=1}^n t^\beta z^{\alpha-\beta-1} + O(z^{\alpha-1}) \right) dz. \end{aligned}$$

Hence, the residue is different from 0, when  $\alpha = \beta$ , resulting in this way the desired result. □

**Corollary 4.24.1.** The coordinate  $t^n$  can be written in terms of the coordinates  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_{n+1}$  as

$$t^n = n (\varphi_n)^{\frac{1}{n}}. \quad (4.120)$$

*Proof.*

$$\begin{aligned}
t^n &= n \operatorname{res}_{v=0} \lambda^{\frac{1}{n}}(v) dv \\
&= n \operatorname{res}_{v=0} \left( \frac{\varphi_n}{v^n} + \frac{\varphi_{n-1}}{v^{n-1}} + \dots + \frac{\varphi_2}{v^2} + \frac{\varphi_1}{v} + O(1) \right)^{\frac{1}{n}} dv \\
&= n \operatorname{res}_{v=0} \frac{(\varphi_n)^{\frac{1}{n}}}{v} (1 + O(v))^{\frac{1}{n}} dv = n (\varphi_n)^{\frac{1}{n}}.
\end{aligned}$$

□

**Lemma 4.25.** Let  $t^1, t^2, \dots, t^n$  be defined in (4.115), then

$$t^\alpha = \frac{n}{n+1-\alpha} (\varphi_n)^{\frac{n+1-\alpha}{n}} (1 + \Phi_{n-\alpha})^{\frac{n+1-\alpha}{n}}, \quad (4.121)$$

where

$$\begin{aligned}
(1 + \Phi_i)^{\frac{n+1-\alpha}{n}} &= \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \Phi_i^d, \\
\Phi_i^d &= \sum_{i_1+i_2+\dots+i_d=i} \frac{\varphi_{(n-i_1)}}{\varphi_n} \dots \frac{\varphi_{(n-i_d)}}{\varphi_n}.
\end{aligned} \quad (4.122)$$

*Proof.*

$$\begin{aligned}
t^\alpha &= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} \left( \lambda^{\frac{n+1-\alpha}{n}}(v) dv \right) \\
&= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} \left( \frac{\varphi_n}{v^n} + \frac{\varphi_{n-1}}{v^{n-1}} + \dots + \frac{\varphi_2}{v^2} + \frac{\varphi_1}{v} + O(1) \right)^{\frac{n+1-\alpha}{n}} dv \\
&= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} \left( \frac{\varphi_n}{v^n} \right)^{\frac{n+1-\alpha}{n}} \left( 1 + \frac{\varphi_{n-1}}{\varphi_n} v + \frac{\varphi_{n-2}}{\varphi_n} v^2 + \dots + \frac{\varphi_2}{\varphi_n} v^{n-2} + \frac{\varphi_1}{\varphi_n} v^{n-1} + O(v^{n+1}) \right)^{\frac{n+1-\alpha}{n}} dv \\
&= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} \left( \frac{\varphi_n}{v^n} \right)^{\frac{n+1-\alpha}{n}} \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \left( 1 + \frac{\varphi_{n-1}}{\varphi_n} v + \dots + \frac{\varphi_1}{\varphi_n} v^{n-1} + O(v^{n+1}) \right)^d dv \\
&= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} \left( \frac{\varphi_n}{v^n} \right)^{\frac{n+1-\alpha}{n}} \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \sum_{j_1+\dots+j_n=d} \frac{d!}{j_1! j_2! \dots j_n!} \prod_{i=1}^{n-1} \left( \frac{\varphi_{n-i} v^i}{\varphi_n} \right)^{j_i} (O(v^n))^{j_n} dv \\
&= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} (\varphi_n)^{\frac{n+1-\alpha}{n}} \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \sum_{j_1+\dots+j_n=d} \frac{d!}{j_1! j_2! \dots j_n!} \prod_{i=1}^{n-1} \left( \frac{\varphi_{n-i}}{\varphi_n} \right)^{j_i} v^{(\sum_{i=1}^{n-1} i j_i - n - 1 + \alpha)} dv \\
&\quad + O(1) \\
&= \frac{n}{n+1-\alpha} (\varphi_n)^{\frac{n+1-\alpha}{n}} \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \sum_{\substack{j_1+\dots+j_n=d \\ j_1+2j_2+3j_3+\dots+(n-1)j_{n-1}=n-\alpha}} \frac{d!}{j_1! j_2! \dots j_n!} \prod_{i=1}^{n-1} \left( \frac{\varphi_{n-i}}{\varphi_n} \right)^{j_i} \\
&= \frac{n}{n+1-\alpha} (\varphi_n)^{\frac{n+1-\alpha}{n+1}} \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \sum_{i_1+\dots+i_d=n-\alpha} \frac{\varphi_{(n-i_1)}}{\varphi_{n+1}} \dots \frac{\varphi_{(n-i_d)}}{\varphi_n} \\
&= \frac{n}{n+1-\alpha} (\varphi_n)^{\frac{n+1-\alpha}{n}} \sum_{d=0}^{\infty} \binom{\frac{n+1-\alpha}{n}}{d} \Phi_{n-\alpha}^d \\
&= \frac{n}{n+1-\alpha} (\varphi_n)^{\frac{n+1-\alpha}{n}} (1 + \Phi_{n-\alpha})^{\frac{n+1-\alpha}{n}}.
\end{aligned}$$

□

---

**Corollary 4.25.1.** The coordinate  $t^1$  can be written in terms of the coordinates  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$  as

$$t^1 = \varphi_1 \quad (4.123)$$

*Proof.* Using the relation (4.121) for  $\alpha = 1$

$$t^1 = \varphi_n (1 + \Phi_{n-1}) = \varphi_n \sum_{i_1=n-1} \frac{\varphi_{n-i_1}}{\varphi_n} = \varphi_1. \quad (4.124)$$

□

**Lemma 4.26.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$  and  $\lambda(v)$  be defined in (4.25), then

$$k\varphi_k = \operatorname{res}_{v=0} k\lambda v^{k-1} dv. \quad (4.125)$$

*Proof.*

$$\begin{aligned} \operatorname{res}_{v=0} k\lambda v^{k-1} dv &= \operatorname{res}_{v=0} k \left( \frac{\varphi_n}{v^n} + \frac{\varphi_{n-1}}{v^{n-1}} + \dots + \frac{\varphi_k}{v^k} + \dots + \frac{\varphi_2}{v^2} + \frac{\varphi_1}{v} + O(1) \right) v^{k-1} dv \\ &= k\varphi_k. \end{aligned}$$

□

**Lemma 4.27.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$  and  $\lambda(v)$  be defined in (4.25), then

$$k\varphi_k = - \operatorname{res}_{\lambda=\infty} v^k d\lambda. \quad (4.126)$$

*Proof.* Using formula (4.126) and integration by parts

$$k\varphi_k = \operatorname{res}_{v=0} k\lambda v^{k-1} dv = - \operatorname{res}_{\lambda=\infty} v^k d\lambda.$$

□

**Lemma 4.28.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n$ ,  $\lambda(v)$  be defined in (4.25) and  $(t^1, \dots, t^n)$  be defined in (4.115), then

$$k\varphi_k = \frac{(-1)^k}{n^{k-1}} T_n^k, \quad (4.127)$$

where

$$T_n^k = \sum_{i_1 + \dots + i_k = n} t^{(n+1-i_1)} \dots t^{(n+1-i_k)}. \quad (4.128)$$

*Proof.* Let  $z := \left(\frac{1}{\lambda}\right)^{\frac{1}{n}}$ , then by using equation (4.126):

$$\begin{aligned} - \operatorname{res}_{\lambda=\infty} v^k d\lambda &= \operatorname{res}_{z=0} \frac{nv^k(z)dz}{z^{n+1}} \\ &= \operatorname{res}_{z=0} \frac{(-1)^k}{n^{k-1}} (t^n z + t^{n-1} z^2 + \dots + t^2 z^{n-1} + O(z^{n+1}))^k \frac{dz}{z^{n+1}} \\ &= \operatorname{res}_{z=0} \frac{(-1)^k}{n^{k-1}} \sum_{j_1+j_2+\dots+j_n+k=n} (t^n z)^{j_1} (t^{n-1} z^2)^{j_2} \dots (t^2 z^{n-1})^{j_n} (O(z^{n+1}))^{j_{n+2}} \frac{dz}{z^{n+1}} \\ &= \frac{(-1)^k}{n^{k-1}} \sum_{\substack{j_1+j_2+\dots+j_n=k \\ j_1+2j_2+3j_3+\dots+(n)j_n=n}} \frac{k!}{j_1!j_2!\dots j_n!} (t^n)^{j_1} (t^{n-1})^{j_2} \dots (t^2)^{j_n} \\ &= \frac{(-1)^k}{n^{k-1}} \sum_{i_1+\dots+i_k=n} t^{(n+1-i_1)} \dots t^{(n+1-i_k)} \\ &= \frac{(-1)^k}{n^{k-1}} T_n^k. \end{aligned}$$

□

**Lemma 4.29.** Let  $T_n^k$  be defined in (4.128), then

$$\frac{\partial T_n^k}{\partial t^\alpha} = k T_{\alpha-1}^{k-1}. \quad (4.129)$$

*Proof.*

$$\begin{aligned} \frac{\partial T_n^k}{\partial t^\alpha} &= \frac{\partial}{\partial t^\alpha} \left( \sum_{i_1 + \dots + i_k = n} t^{(n+1-i_1)} \dots t^{(n+1-i_k)} \right) \\ &= \sum_{i_1 + \dots + i_k = n} k \delta_{n+1-i_k, \alpha} t^{(n+1-i_1)} \dots t^{(n+1-i_{k-1})} \\ &= k \sum_{i_1 + \dots + i_{k-1} = \alpha-1} t^{(n+1-i_1)} \dots t^{(n+1-i_{k-1})} \\ &= k T_{\alpha-1}^{k-1}. \end{aligned}$$

□

At this stage, we are able to compute the coefficients of  $\eta(dt^\alpha, dt^\beta)$ .

**Theorem 4.30.** Let  $(t^1, \dots, t^n)$  defined in (4.115), and  $\eta^*$  defined in (4.77). Then,

$$\eta^*(dt^\alpha, dt^{n+3-\beta}) = n \delta_{\alpha\beta}. \quad (4.130)$$

*Proof.* If  $i, j \neq 0$

$$\eta^*(d\varphi_i, d\varphi_j) = \sum_{\alpha=1}^n \sum_{\beta=1}^n \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_j}{\partial t^\beta} \eta^*(dt^\alpha, dt^\beta) \quad (4.131)$$

Using (4.129) and (4.127), we get

$$\frac{\partial \varphi_k}{\partial t^\alpha} = \frac{(-1)^k}{n^{k-1}} k T_{\alpha-1}^{k-1} \quad (4.132)$$

$$\sum_{\alpha=1}^n \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_{n+3-j}}{\partial t^{n+3-\alpha}} = \sum_{\alpha=1}^n \frac{(-1)^{i-j+n+1}}{n^{i-j+n+1}} T_{\alpha-1}^{i-1} T_{n+2-\alpha}^{n+2-j} \quad (4.133)$$

Using the second of the equation (??) in (4.133)

$$\begin{aligned} \sum_{\alpha=1}^n \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_{n+3-j}}{\partial t^{n+3-\alpha}} &= \frac{T_{n+1}^{n+1+i-j}}{n} \\ &= \frac{(n+1+i-j)}{n} \varphi_{(n+1+i-j)} \end{aligned} \quad (4.134)$$

Note that the following identity holds

$$\sum_{\alpha=2}^{n+1} \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_{n+3-j}}{\partial t^{n+3-\alpha}} = \sum_{\alpha=2}^{n+1} \sum_{\alpha=2}^{n+1} \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_{n+3-j}}{\partial t^{n+3-\beta}} \delta_{\alpha\beta}. \quad (4.135)$$

---

On another hand, using the first equation of (4.96), we have

$$\begin{aligned}
\eta^*(d\varphi_i, d\varphi_{n+3-j}) &= (n+1+i-j)\varphi_{n+1+i-j} \\
&= n \sum_{\alpha=2}^{n+1} \sum_{\beta=2}^{n+1} \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_{n+3-j}}{\partial t^{n+3-\beta}} \delta_{\alpha\beta} \\
&= \sum_{\alpha=2}^{n+1} \sum_{\beta=2}^{n+1} \frac{\partial \varphi_i}{\partial t^\alpha} \frac{\partial \varphi_{n+3-j}}{\partial t^{n+3-\beta}} \eta^*(dt^\alpha, dt^{n+3-\beta})
\end{aligned}$$

Then, we obtain

$$\eta^*(dt^\alpha, dt^{n+3-\beta}) = n\delta_{\alpha\beta}.$$

□

**Lemma 4.31.** Let  $(t^1, \dots, t^n)$  be defined in (4.115), and  $\eta^*$  be defined in (4.77). Then,

$$\begin{aligned}
\eta^*(dt^i, d\tau) &= -2\pi i \delta_{i0}, \\
\eta^*(dt^i, dv_{n+1}) &= -\frac{\delta_{i1}}{n+1}.
\end{aligned} \tag{4.136}$$

*Proof.* Using corollary 4.20.1 and lemma 4.21, we have

$$\eta^*(dt^i, d\tau) = \frac{\partial t^i}{\partial \varphi_j} \eta^*(d\varphi_j, d\tau) = -2\pi i \delta_{i0}.$$

In addition, if  $i \neq 0$

$$\begin{aligned}
\eta^*(dt^i, dv_{n+1}) &= \frac{\partial t^i}{\partial \varphi_j} \eta^*(d\varphi_j, dv_{n+1}) = 0, \quad i \neq 1, \\
\eta^*(dt^1, dv_{n+1}) &= \eta^*(d\varphi_1, dv_{n+1}) = -\frac{1}{n+1}.
\end{aligned}$$

Computing  $dt^0$  with respect the variables  $\varphi_i$  by using (4.116),

$$\begin{aligned}
dt^0 &= d\varphi_0 - (n+1) \frac{\partial^2 \log(\theta_1((n+1)v_{n+1}))}{\partial v_{n+1}^2} \varphi_1 dv_{n+1} - \frac{\partial \log(\theta_1((n+1)v_{n+1}))}{\partial v_{n+1}} d\varphi_1 \\
&\quad + 4\pi i g'_1(\tau) \varphi_2 d\tau + 4\pi i g_1(\tau) d\varphi_2.
\end{aligned} \tag{4.137}$$

Hence,

$$\begin{aligned}
\eta^*(dt^0, dv_{n+1}) &= \eta^*(d\varphi_0, dv_{n+1}) - \frac{\partial \log(\theta_1((n+1)v_{n+1}))}{\partial v_{n+1}} \eta^*(d\varphi_1, dv_{n+1}) \\
&= 0.
\end{aligned}$$

□

**Lemma 4.32.** Let  $t^0$  be defined in (4.116), and  $\eta^*$  be defined in (4.77). Then,

$$\eta^*(dt^0, dt^\alpha) = 0, \quad \alpha \neq 0. \tag{4.138}$$

---

*Proof.* Using the definition of  $\eta^*$  in equation (4.77), formula (4.96), and (4.79)

If  $i > 1$ ,

$$\begin{aligned}
\eta^*(dt^0, d\varphi_i) &= \eta^*(d\varphi_0, d\varphi_i) + 4\pi i g_1(\tau) \eta^*(d\varphi_2, d\varphi_i) + 4\pi i g_1'(\tau) \eta^*(d\tau, d\varphi_i) \\
&= \eta^*(d\varphi_0, d\varphi_i) + 4\pi i g_1(\tau) \eta^*(d\varphi_2, d\varphi_i) \\
&= \tilde{\eta}^*(d\varphi_0, d\varphi_i) - 4\pi i g_1(\tau) k_i \varphi_i + 4\pi i g_1(\tau) k_i \varphi_i = 0 \\
&= -4\pi i g_1(\tau) k_i \varphi_i + 4\pi i g_1(\tau) k_i \varphi_i = 0
\end{aligned}$$

Then, if  $\alpha > 1$

$$\eta^*(dt^0, dt^\alpha) = \sum_{\alpha=2}^n \frac{\partial t^\alpha}{\partial \varphi_i} \eta^*(dt^0, d\varphi_i) = 0.$$

Computing  $\eta^*(dt^0, dt^1)$  by using (4.137)

$$\begin{aligned}
\eta^*(dt^0, dt^1) &= \eta^*(d\varphi_0, d\varphi_1) - (n+1) \frac{\partial^2 \log(\theta_1((n+1)v_{n+1}))}{\partial v_{n+1}^2} \varphi_1 \eta^*(dv_{n+1}, d\varphi_1) \\
&= 0.
\end{aligned}$$

□

Instead of considering the coefficients of the metric  $\eta^*$ , let us investigate the flatness of its inverse  $\eta$  in order to make the computations shorter.

**Lemma 4.33.** The metric

$$\sum_{\alpha=2}^n \eta(dt^\alpha, dt^{n+3-\alpha}) dt^\alpha dt^{n+3-\alpha} - 2(n+1) dt^1 dv_{n+1} - \frac{1}{\pi i} dt^0 d\tau \quad (4.139)$$

is invariant under the second action of (4.2).

*Proof.* Under the second action of (4.2), we have that

$$\sum_{\alpha=2}^n \eta(dt^\alpha, dt^{n+3-\alpha}) dt^\alpha dt^{n+3-\alpha}$$

are invariant under the second action of (4.2), because the relationship between  $t^i$  and  $\varphi_i$  be given by (4.121), and the fact that the Jacobi forms  $\{\varphi_i\}$  are invariant under the second action of (4.2).  $t^0$  and  $v_{n+1}$  have the following transformation law

$$\begin{aligned}
t^0 &\mapsto t^0 - 2\pi i(n+1)\lambda_{n+1}t^1 \\
v_{n+1} &\mapsto v_{n+1} + \lambda_{n+1}\tau + \mu_{n+1}
\end{aligned} \quad (4.140)$$

Hence, its differentials are

$$\begin{aligned}
dt^0 &\mapsto dt^0 - 2\pi i(n+1)\lambda_{n+1}dt^1 \\
dv_{n+1} &\mapsto dv_{n+1} + \lambda_{n+1}d\tau
\end{aligned} \quad (4.141)$$

Substituting (4.141) in (4.139) we get the desired result.

□

---

**Theorem 4.34.** Let  $(t^0, t^1, t^2, \dots, t^n)$  defined in (4.115), and  $\eta^*$  defined in (4.77). Then,

$$\begin{aligned}
\eta^*(dt^\alpha, dt^{n+3-\beta}) &= -(n+1)\delta_{\alpha\beta}, \quad 2 \leq \alpha, \beta \leq n \\
\eta^*(dt^1, dt^\alpha) &= 0, \\
\eta^*(dt^0, dt^\alpha) &= 0, \\
\eta^*(dt^i, d\tau) &= -2\pi i \delta_{i0}, \\
\eta^*(dt^i, dv_{n+1}) &= -\frac{\delta_{i1}}{n+1}.
\end{aligned} \tag{4.142}$$

Moreover, the coordinates  $t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau$  are the flat coordinates of  $\eta^*$ .

*Proof.* The theorem is already proved for  $\alpha, \beta \in \{2, \dots, n\}$  in theorem 4.30, and for the rest in the lemma 4.32 and 4.31. The only missing part is to prove

$$\eta^*(dt^0, dt^0) = 0. \tag{4.143}$$

Recall that from corollary 4.18.2, the metric  $\eta^*$  is invariant under the second action of (4.2). Moreover, the same statement is valid for (4.139), because of lemma 4.33. However, the tensor  $dt^0 \otimes dt^0$  have a non-trivial transformation law under this action. Hence, if the coefficient of the component  $dt^0 \otimes dt^0$  is different from 0, we have a contradiction with corollary 4.18.2.

□

**Corollary 4.34.1.** The metric  $\eta^*(d\varphi_i, d\varphi_j) := \frac{\partial g^*(d\varphi_i, d\varphi_j)}{\partial \varphi_0}$  is triangular, and non degenerate.

**Definition 4.5.** Let  $\eta^* = \eta^{\alpha\beta} \frac{\partial}{\partial t^\alpha} \otimes \frac{\partial}{\partial t^\beta}$  defined in (4.77). The metric defined by

$$\eta = \eta_{\alpha\beta} dt^\alpha \otimes dt^\beta \tag{4.144}$$

is denoted by  $\eta$ .

## 4.7 The extended ring of Jacobi forms

The flat coordinates of the Saito metric  $\eta$  of the orbit space of  $\mathcal{J}(\tilde{A}_n)$  does not live in the orbit space of  $\mathcal{J}(\tilde{A}_n)$ , but live in a suitable covering of this orbit space. The main goal of this section is in describing this covering as the space such that the ring of functions of this covering is a suitable extension of the ring of Jacobi forms.

**Lemma 4.35.** The coordinates  $(t^0, t^1, t^2, \dots, v_{n+1}, \tau)$  defined on (4.115) have the following transformation laws under the action of the group  $\mathcal{J}(\tilde{A}_n)$ : They transform as follows under the second action of (4.2):

$$\begin{aligned}
t^0 &\mapsto t^0 - 2\pi i(n+1)\lambda_{n+1}t^1 \\
t^\alpha &\mapsto t^\alpha, \quad \alpha \neq 0 \\
v_{n+1} &\mapsto v_{n+1} + \mu_{n+1} + \lambda_{n+1}\tau \\
\tau &\mapsto \tau
\end{aligned} \tag{4.145}$$



Moreover, they transform as follows under the third action (4.2)

$$\begin{aligned}
t^0 &\mapsto t^0 + \frac{2c \sum_{\alpha, \beta \neq 0, \tau} \eta_{\alpha\beta} t^\alpha t^\beta}{c\tau + d} \\
t^\alpha &\mapsto \frac{t^\alpha}{c\tau + d}, \quad \alpha \neq 0 \\
v_{n+1} &\mapsto \frac{v_{n+1}}{c\tau + d} \\
\tau &\mapsto \frac{a\tau + b}{c\tau + d}
\end{aligned} \tag{4.146}$$

*Proof.* Note that the term  $\Phi_i^d$  equation (4.122) has weight  $+i$ , then using that  $\varphi_n$  has weight  $-n$ , we have that the weight of  $t^\alpha$  for  $\alpha \neq 1$  must have weight  $-1$  due to (4.121). The transformation law of  $t^1$  follows from the transformation law of  $\frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})}$  and  $g_1(\tau)$

$$\begin{aligned}
\frac{\theta'_1((n+1)v_{n+1} + (n+1)\lambda_{n+1}\tau + \mu_{n+1}, \tau)}{\theta_1((n+1)v_{n+1} + (n+1)\lambda_{n+1}\tau + \mu_{n+1}, \tau)} &= \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} - 2\pi i(n+1)\lambda_{n+1}, \\
\frac{\theta'_1(\frac{(n+1)v_{n+1}}{c\tau+d}, \frac{a\tau+b}{c\tau+d})}{\theta_1(\frac{(n+1)v_{n+1}}{c\tau+d}, \frac{a\tau+b}{c\tau+d})} &= \frac{\theta'_1((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} + 2\pi i c(n+1)v_{n+1}, \\
g_1(\frac{a\tau+b}{c\tau+d}) &= (c\tau+d)^2 g_1(\tau) + 2c(c\tau+d),
\end{aligned} \tag{4.147}$$

and by using equation (4.127) for  $k = 2$ .

□

In addition, from the formula (4.122) it is clear that the multivaluedness of  $(t^1, \dots, t^n)$  comes from  $(\varphi_n)^{\frac{1}{n}}$ . Therefore, the coordinates lives in a suitable covering over the orbit space of the group  $\mathcal{J}(\tilde{A}_n)$ . This covering is obtained by forgetting to act the Coxeter group  $A_n$  and the  $SL_2(\mathbb{Z})$  action, and the translation action  $v_{n+1} \mapsto v_{n+1} + \lambda_{n+1}\tau + \mu_{n+1}$  of  $\mathcal{J}(\tilde{A}_n)$  on  $\mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H}$ . The only remaining part of the  $\mathcal{J}(\tilde{A}_n)$  action are the translations

$$v_i \mapsto v_i + \lambda_i \tau + \mu_i, \quad i \neq n+1.$$

Hence, the coordinates  $(t^1, \dots, t^n)$  live in  $n$ -dimensional tori with fixed symplectic base of the torus homology, a fixed chamber in the tori parametrised by  $(v_{n+1}, \tau)$ , and with a branching divisor  $Y := \{\varphi_n = 0\}$ . Another way to describe this covering is using the flat coordinates of the intersection form  $(u, v_0, v_1, \dots, v_{n+1}, \tau)$ , and to fix a lattice  $\tau$ , a representative of the action

$$v_{n+1} \mapsto v_{n+1} + \lambda_{n+1}\tau + \mu_{n+1}, \tag{4.148}$$

and a representative of the  $A_n$  action. Then, the desired covering of the orbit space of the group  $\mathcal{J}(\tilde{A}_n)$  is defined by

$$\mathbb{C} \oplus \widetilde{\mathbb{C}^{n+1}} \oplus \mathbb{H} / \mathcal{J}(\tilde{A}_n) := \mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} / (\mathbb{Z}^n \oplus \tau \mathbb{Z}^n), \tag{4.149}$$

where  $\mathbb{Z}^n \oplus \tau \mathbb{Z}^n$  acts on  $\mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H}$  by

$$\begin{aligned}
v_i &\mapsto v_i + \lambda_i \tau + \mu_i, \quad i \neq n+1, \\
u &\mapsto u - 2A_{ij}\lambda_i v_j - A_{ij}\lambda_i \lambda_j \tau, \\
v_{n+1} &\mapsto v_{n+1}, \\
\tau &\mapsto \tau.
\end{aligned} \tag{4.150}$$

where  $A_{ij}$  is given by (??).

**Remark 4.3.** Note that due to the lemma 4.6 the covering space 4.149 is isomorphic to a suitable covering over the Hurwitz space  $H_{1,n-1,0}$ . The covering over  $H_{1,n-1,0}$  is given by a fixation of base of the homology in the tori generated by the lattice  $(1, \tau)$ , fixation of root  $\lambda$  (4.25) near  $\infty$ , and a fixation of a logarithm root.

In order to manipulate the geometric objects of this covering, it is more convenient to use their ring of functions. Hence, we define:

**Definition 4.6.** The extended ring of Jacobi forms with respect the ring of coefficients is the following ring

$$\tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n], \quad (4.151)$$

where

$$\tilde{E}_{\bullet, \bullet} = E_{\bullet, \bullet} \oplus \{g_1(\tau)\} \oplus \left\{ \frac{\theta_1'((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \right\}. \quad (4.152)$$

**Lemma 4.36.** The coefficients of the intersection form  $g^{ij}$  on the coordinates  $\varphi_0, \varphi_1, \dots, \varphi_n, v_{n+1}, \tau$  belong to the ring  $\tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ .

*Proof.* It is a consequence of the formula (4.49). □

**Lemma 4.37.** The coefficients of the intersection form  $g^{\alpha\beta}$  on the coordinates  $t^0, t^1, \dots, t^n, v_{n+1}, \tau$  belong to the ring  $\tilde{E}_{\bullet, \bullet}[t^0, t^1, \dots, t^n, \frac{1}{t^n}]$ .

*Proof.* Using the transformation law of  $g^{\alpha\beta}$

$$g^{\alpha\beta} = \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g(d\varphi_i, d\varphi_j), \quad (4.153)$$

we realise the term  $\frac{\partial t^\alpha}{\partial \varphi_i}$  as polynomial in  $t^0, t^1, \dots, t^n, \frac{1}{t^n}$  due to the relations (4.121) and (4.127). □

## 4.8 Christoffel symbols of the intersection form

In this section, we consider the Christoffel symbols of the intersection form (4.41). In particular, we prove that this Christoffel symbols, in coordinates  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n, v_{n+1}, \tau$ , live in  $\tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ . In addition, we will show that the Christoffel symbols depend at most linear in  $\varphi_0$  in this coordinates.

Recall that the Christoffel symbols  $\Gamma_k^{ij}(\varphi)$  associated with the intersection form  $g^*$  is given in terms of the conditions (2.25).

**Lemma 4.38.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n, v_{n+1}, \tau$  be defined in (4.25), then  $\Gamma_j^{ii}$  depend at most linear on  $\varphi_0$ .

*Proof.* Using the first condition of (2.25)

$$\partial_k g^{ii} = 2\Gamma_k^{ii}$$

Recall that due to the corollary 4.22.1, the metric  $g^{ij}$  depend at most linear on  $\varphi_0$ . Then,

$$2 \frac{\partial^2 \Gamma_k^{ii}}{\partial \varphi_0^2} = \partial_0^2 \partial_k g^{ii} = \partial_k \partial_0^2 g^{ii} = 0.$$

□

**Lemma 4.39.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n, v_{n+1}, \tau$  be defined in (4.25), then

$$\begin{aligned} \Gamma_j^{i\tau} &= 0, \\ \Gamma_k^{\tau k} &= -2\pi i \frac{\delta_{jk}}{k}. \end{aligned} \quad (4.154)$$

*Proof.* Let  $\Gamma_k^{ij}(x)$ , in the coordinates  $x_1, \dots, x_n$ , and  $\Gamma_l^{pq}(y)$  in the coordinates  $y_1, \dots, y_n$ , then the transformation law of the Christoffel symbol defined in the cotangent bundle is the following

$$\Gamma_k^{ij}(x) = \frac{\partial x^i}{\partial y^p} \frac{\partial x^j}{\partial y^q} \frac{\partial y^l}{\partial x^k} \Gamma_l^{pq}(y) + \frac{\partial x^i}{\partial y^p} \frac{\partial}{\partial x^k} \left( \frac{\partial x^j}{\partial y^q} \right) g^{pq}(y). \quad (4.155)$$

In particular, the  $\Gamma_k^{ij}(\varphi)$  in the coordinates  $(\varphi_0, \varphi_1, \dots, \varphi_n, v_{n+1}, \tau)$  could be derived from the Christoffel symbol in the coordinates  $v_0, v_1, \dots, v_{n+1}, \tau$  which is 0. Then,

$$\Gamma_k^{ij}(\varphi) = \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \varphi_k} \left( \frac{\partial \varphi_j}{\partial v_q} \right) g^{pq}(v). \quad (4.156)$$

Computing  $\Gamma_j^{iv_{n+1}}$ ,

$$\begin{aligned} \Gamma_k^{i\tau}(\varphi) &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \varphi_k} \left( \frac{\partial \tau}{\partial v_q} \right) g^{pq}(v) \\ &= -2\pi \varphi_i \frac{\partial}{\partial \varphi_k} (1) = 0. \end{aligned} \quad (4.157)$$

Computing  $\Gamma_j^{v_{n+1}i}$  by using the first condition of (2.25),

$$\begin{aligned} \Gamma_j^{\tau k}(\varphi) &= \partial_j g^{k\tau} - \Gamma_j^{k\tau} \\ &= \partial_j g^{k\tau} = -2\pi i \frac{\delta_{jk}}{k}. \end{aligned} \quad (4.158)$$

□

**Lemma 4.40.** Let  $\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n, v_{n+1}, \tau$  be defined in (4.25), then

$$\begin{aligned} \Gamma_j^{iv_{n+1}} &= 0, \\ \Gamma_j^{v_{n+1}i} &= \frac{\partial g^{iv_{n+1}}}{\partial \varphi_j} \in \tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n], \\ \Gamma_{v_{n+1}}^{ij} &\in \tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]. \end{aligned} \quad (4.159)$$

Moreover, these Christoffel symbols are at most linear on  $\varphi_0$ .

*Proof.* Let  $\Gamma_k^{ij}(x)$ , in the coordinates  $x_1, \dots, x_n$ , and  $\Gamma_l^{pq}(y)$  in the coordinates  $y_1, \dots, y_n$ , then the transformation law of the Christoffel symbol defined in the cotangent bundle is the following

$$\Gamma_k^{ij}(x) = \frac{\partial x^i}{\partial y^p} \frac{\partial x^j}{\partial y^q} \frac{\partial y^l}{\partial x^k} \Gamma_l^{pq}(y) + \frac{\partial x^i}{\partial y^p} \frac{\partial}{\partial x^k} \left( \frac{\partial x^j}{\partial y^q} \right) g^{pq}(y). \quad (4.160)$$

In particular, the  $\Gamma_k^{ij}(\varphi)$  in the coordinates  $(\varphi_0, \varphi_1, \dots, \varphi_n, v_{n+1}, \tau)$  could be derived from the Christoffel symbol in the coordinates  $v_0, v_1, \dots, v_{n+1}, \tau$  which is 0. Then,

$$\Gamma_k^{ij}(\varphi) = \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \varphi_k} \left( \frac{\partial \varphi_j}{\partial v_q} \right) g^{pq}(v). \quad (4.161)$$

Computing  $\Gamma_k^{iv_{n+1}}$ ,

$$\begin{aligned} \Gamma_k^{iv_{n+1}}(\varphi) &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \varphi_k} \left( \frac{\partial v_{n+1}}{\partial v_q} \right) g^{pq}(v) \\ &= \frac{\partial \varphi_i}{\partial v_{n+1}} \frac{\partial}{\partial \varphi_k} (1) g^{v_{n+1}v_{n+1}}(v) = 0. \end{aligned} \quad (4.162)$$

Computing  $\Gamma_k^{v_{n+1}i}$  by using the first condition of (2.25),

$$\begin{aligned} \Gamma_k^{v_{n+1}i}(\varphi) &= \partial_k g^{iv_{n+1}} - \Gamma_j^{iv_{n+1}} \\ &= \partial_k g^{iv_{n+1}}. \end{aligned} \quad (4.163)$$

Since,  $g^{iv_{n+1}} \in \tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ , we have that  $\Gamma_k^{v_{n+1}i}(\varphi) \in \tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ . In addition, since the metric  $\eta^{iv_{n+1}}$  is independent of  $\varphi_0$  due to the corollary 4.22.1, we have that  $\partial_k g^{iv_{n+1}}$  is at most linear on  $\varphi_0$

Computing  $\Gamma_{v_{n+1}}^{ij}$ ,

$$\begin{aligned} \Gamma_{v_{n+1}}^{ij}(\varphi) &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_{n+1}} \left( \frac{\partial \varphi_j}{\partial v_q} \right) g^{pq}(v) \\ &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial \varphi_j}{\partial v_{n+1}} \right) g^{pq}(v) \end{aligned} \quad (4.164)$$

In the subsequent computation, whenever appears a function depend only on  $v_{n+1}, \tau$ , we will call it by  $h(v_{n+1}, \tau)$ , because for our purpose, it is enough to prove that the subsequent function belong to the ring  $\tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ .

If  $j > 1$ , using equation (4.86),

$$\begin{aligned} \Gamma_{v_{n+1}}^{ij}(\varphi) &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial \varphi_j}{\partial v_{n+1}} \right) g^{pq}(v) \\ &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (\varphi_{j-1} + h(v_{n+1}, \tau) \varphi_n) g^{pq}(v) \\ &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (\varphi_{j-1} + h(v_{n+1}, \tau) \varphi_n) g^{pq}(v) \\ &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (\varphi_{j-1}) g^{pq}(v) + h(v_{n+1}, \tau) \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (\varphi_n) g^{pq}(v) \\ &\quad + \varphi_n \frac{\partial \varphi_i}{\partial v_{n+1}} \frac{\partial}{\partial v_{n+1}} (h(v_{n+1}, \tau)) g^{v_{n+1}v_{n+1}}(v) \\ &= g^{i(j-1)}(\varphi) + h(v_{n+1}, \tau) g^{in}(\varphi_n) + \frac{h'(v_{n+1}, \tau)}{n(n+1)} (\varphi_i + g(v_{n+1}, \tau) \varphi_n) \varphi_n. \end{aligned} \quad (4.165)$$

If  $j = 1$ , using equation (4.88)

$$\begin{aligned}
\Gamma_{v_{n+1}}^{i1}(\varphi) &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial \varphi_1}{\partial v_{n+1}} \right) g^{pq}(v) \\
&= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (n\varphi_0 + h(v_{n+1}, \tau)\varphi_1) g^{pq}(v) \\
&= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (n\varphi_0 + h(v_{n+1}, \tau)\varphi_1) g^{pq}(v) \\
&= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (n\varphi_0) g^{pq}(v) + h(v_{n+1}, \tau) \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (\varphi_1) g^{pq}(v) \\
&\quad + \varphi_1 \frac{\partial \varphi_i}{\partial v_{n+1}} \frac{\partial}{\partial v_{n+1}} (h(v_{n+1}, \tau)) g^{v_{n+1}v_{n+1}}(v) \\
&= ng^{i0}(\varphi) + h(v_{n+1}, \tau)g^{i1}(\varphi_1) + \frac{h'(v_{n+1}, \tau)}{n(n+1)} (\varphi_i + h_3(v_{n+1}, \tau)\varphi_n) \varphi_1.
\end{aligned} \tag{4.166}$$

If  $j = 0$ , using equation (4.90)

$$\begin{aligned}
\Gamma_{v_{n+1}}^{i0}(\varphi) &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial \varphi_0}{\partial v_{n+1}} \right) g^{pq}(v) \\
&= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (-nh_1(v_{n+1}, \tau)\varphi_0 + h_2(v_{n+1}, \tau)\varphi_1) g^{pq}(v) \\
&= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (-nh_1(v_{n+1}, \tau)\varphi_0 + h_2(v_{n+1}, \tau)\varphi_1) g^{pq}(v) \\
&= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (-nh_1(v_{n+1}, \tau)\varphi_0) g^{pq}(v) + h_2(v_{n+1}, \tau) \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} (\varphi_1) g^{pq}(v) \\
&\quad + \varphi_1 \frac{\partial \varphi_i}{\partial v_{n+1}} \frac{\partial}{\partial v_{n+1}} (h_2(v_{n+1}, \tau)) g^{v_{n+1}v_{n+1}}(v) \\
&= -nh_1(v_{n+1}, \tau)g^{i0}(\varphi) + h_2(v_{n+1}, \tau)g^{i1}(\varphi_n) + \frac{h'_2(v_{n+1}, \tau)}{n(n+1)} (\varphi_i + h_3(v_{n+1}, \tau)\varphi_n) \varphi_1 \\
&\quad - n \frac{h'_1(v_{n+1}, \tau)}{n(n+1)} (\varphi_i + h_3(v_{n+1}, \tau)\varphi_n) \varphi_0.
\end{aligned} \tag{4.167}$$

Hence  $\Gamma_{v_{n+1}}^{ij}(\varphi) \in \tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ , furthermore, it is at most linear on  $\varphi_0$ .

□

**Proposition 4.41.** The Christoffel symbols  $\Gamma_k^{ij}(\varphi)$  belong to the ring  $\tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ .

*Proof.* Note that the invariance of the Jacobi form  $\varphi_i$  with respect the first two actions of (4.2), and equivariance by the third one implies that the differential  $d\varphi_i$  is invariant under the first two actions of (4.2), and behaves as follows under the  $SL_2(\mathbb{Z})$

$$d\varphi_i \mapsto \frac{d\varphi_i}{(c\tau + d)^{k_i}} - \frac{c\varphi_i}{(c\tau + d)^{k_i+1}} \tag{4.168}$$

Therefore the Christoffel symbol  $\Gamma_k^{ij}$

$$\nabla_{(d\varphi_i)^\#} d\varphi_j = \Gamma_k^{ij} d\varphi_k \tag{4.169}$$

is a Jacobi form if  $\varphi_i$  has weight 0. Hence, doing the change of coordinates

$$\varphi_i \mapsto \hat{\varphi}_i := \eta^{2i}(\tau)\varphi_i, \tag{4.170}$$

we have that the Christoffel symbol  $\hat{\Gamma}_k^{ij}$

$$\frac{1}{\eta^{2i+2j}} \nabla_{(d\hat{\varphi}_i)^\#} d\hat{\varphi}_j = \hat{\Gamma}_k^{ij} d\hat{\varphi}_k \quad (4.171)$$

is a Jacobi form.

Comparing  $\hat{\Gamma}_k^{ij}$  with  $\Gamma_k^{ij}$

$$\begin{aligned} \nabla_{(d\hat{\varphi}_j)^\#} d\hat{\varphi}_i &= \nabla_{(2jg_1\eta^{2j}\varphi_j d\tau + \eta^{2j}d\varphi_j)^\#} (2ig_1\eta^{2i}\varphi_i d\tau + \eta^{2i}d\varphi_i) \\ &= \nabla_{(2jg_1\eta^{2j}\varphi_j d\tau)^\#} (2ig_1\eta^{2i}\varphi_i d\tau) + \nabla_{(2jg_1\eta^{2j}\varphi_j d\tau)^\#} (\eta^{2i}d\varphi_i) \\ &\quad + \nabla_{(\eta^{2j}d\varphi_j)^\#} (2ig_1\eta^{2i}\varphi_i d\tau) + \nabla_{(\eta^{2j}d\varphi_j)^\#} (\eta^{2i}d\varphi_i) \\ &= 2jg_1\eta^{2j}\varphi_j g^{l\tau} \nabla_{\frac{\partial}{\partial\varphi_l}} (2ig_1\eta^{2i}\varphi_i d\tau) + 2jg_1\eta^{2j}\varphi_j g^{l\tau} \nabla_{\frac{\partial}{\partial\varphi_l}} (\eta^{2i}d\varphi_i) \\ &\quad + \eta^{2j}g^{lj} \nabla_{\frac{\partial}{\partial\varphi_l}} (2ig_1\eta^{2i}\varphi_i d\tau) + \eta^{2j}g^{lj} \nabla_{\frac{\partial}{\partial\varphi_l}} (\eta^{2i}d\varphi_i) \\ &= 4ijg_1'g_1\varphi_i\eta^{2i+2j}\varphi_j g^{\tau\tau} d\tau + 4i^2jg_1^3\eta^{2i+2j}\varphi_j\varphi_i g^{\tau\tau} d\tau + 4ijg_1\eta^{2i+2j}\varphi_j g^{i\tau} d\tau \\ &\quad + 4ijg_1^2\eta^{2i+2j}\varphi_j g^{\tau\tau} d\varphi_i + 2jg_1\eta^{2i+2j}\varphi_j \Gamma_k^{i\tau} d\varphi_k + 4i^2g_1^2\eta^{2i+2j}\varphi_i g^{\tau j} d\tau \\ &\quad + 2ig_1'\eta^{2i+2j}\varphi_i g^{lj} d\tau + 2ig_1\eta^{2i+2j}g^{ij} d\tau + 2ig_1\varphi_i\eta^{2i+2j}\Gamma_k^{i\tau} d\varphi_k \\ &\quad + \eta^{2i+2j}g_1 g^{l\tau} d\varphi_i + \eta^{2i+2j}\Gamma_k^{ji} d\varphi_k. \end{aligned} \quad (4.172)$$

Dividing the equation (4.172) by  $\eta^{2i+2j}$  and isolating  $\Gamma_k^{ji} d\varphi_k$ , we have

$$\begin{aligned} \Gamma_k^{ji} d\varphi_k &= -4ijg_1'g_1\varphi_i\varphi_j g^{\tau\tau} d\tau - 4i^2jg_1^3\varphi_j\varphi_i g^{\tau\tau} d\tau + 4ijg_1\eta^{2i+2j}\varphi_j g^{i\tau} d\tau \\ &\quad - 4ijg_1^2\varphi_j g^{\tau\tau} d\varphi_i - 2jg_1\varphi_j \Gamma_k^{i\tau} d\varphi_k - 4i^2g_1^2\varphi_i g^{\tau j} d\tau \\ &\quad - 2ig_1'\varphi_i g^{lj} d\tau - 2ig_1\eta^{2i+2j}g^{ij} d\tau - 2ig_1\varphi_i \Gamma_k^{i\tau} d\varphi_k \\ &\quad - g_1 g^{l\tau} d\varphi_i + \hat{\Gamma}_k^{ji} d\varphi_k. \end{aligned} \quad (4.173)$$

Since the differential forms has a vector space structure and the right hand side of (4.173) depends only on  $g^{ij}$ ,  $g_1(\tau)$ ,  $\varphi_i$ , and  $\Gamma_k^{i\tau}$  which belongs to the ring  $\tilde{E}_{\bullet,\bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$ , the desired result is proved.  $\square$

**Lemma 4.42.** The Christoffel symbols  $\Gamma_k^{ij}(\varphi)$  depend linearly on  $\varphi_0$ .

*Proof.* The result was already proved, when  $k = v_{n+1}$  in the lemma 4.40. The proposition 4.41 gives to the space of Christoffel symbols the structure of graded algebra, in particular we can compute the degree  $m$  regarding to the algebra of Jacobi forms. Let  $\phi \in \tilde{E}_{\bullet,\bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$  with index  $m_\phi$  and weight  $k_\phi$ , then we write

$$\begin{aligned} \deg_m \phi &= m_\phi, \\ \deg_k \phi &= k_\phi. \end{aligned} \quad (4.174)$$

If  $k \neq \tau, v_{n+1}$ ,

$$\deg_m \Gamma_k^{ij} = \deg_m \left( \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \varphi_k} \left( \frac{\partial \varphi_j}{\partial v_q} \right) g^{pq}(v) \right) = 1. \quad (4.175)$$

Therefore,  $\Gamma_k^{ij}$  is at most linear on  $\varphi_0$ . If  $k = \tau$ ,

$$\deg_k \Gamma_\tau^{ij} = \deg_k \left( \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \tau} \left( \frac{\partial \varphi_j}{\partial v_q} \right) g^{pq}(v) \right) = -i - j + 4.$$

Suppose that  $\Gamma_\tau^{ij}$  contains a the term  $a(v_{n+1}, \tau) \varphi_0^2$ , where  $a(v_{n+1}, \tau)$  is an elliptic function in  $v_{n+1}$ , then

$$\deg_k a(v_{n+1}, \tau) = -i - j + 4 > 0.$$

The possible Christoffel symbols that could depend on  $\varphi_0^2$  are

$$\Gamma_\tau^{04}, \Gamma_\tau^{40}, \Gamma_\tau^{13}, \Gamma_\tau^{31}, \Gamma_\tau^{22}, \Gamma_\tau^{21}, \Gamma_\tau^{12}, \Gamma_\tau^{20}, \Gamma_\tau^{02}, \Gamma_\tau^{11}, \Gamma_\tau^{10}, \Gamma_\tau^{01}, \Gamma_\tau^{00}. \quad (4.176)$$

But  $\Gamma_\tau^{22}, \Gamma_\tau^{11}, \Gamma_\tau^{00}$  is linear on  $\varphi_0$  due to lemma 4.38.

Computing  $\Gamma_\tau^{ij}$

$$\Gamma_\tau^{ij} = \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial \tau} \left( \frac{\partial \varphi_j}{\partial v_q} \right) g^{pq}(v) = \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial \varphi_j}{\partial \tau} \right) g^{pq}(v) \quad (4.177)$$

In order to compute it, recall there exist a relationship between the holomorphic Jacobi forms of  $A_{n+1}$  type and the meromorphic Jacobi forms of  $\tilde{A}_n$  type given by (4.34). Moreover, in (4.32) it was demonstrated that the lowest degree term in the Taylor expansion of  $\varphi_i^{\mathcal{J}(A_{n+1})}$  with respect the variables  $v_0, v_1, \dots, v_{n+1}$  are the elementary symmetric polynomials  $a_i(v_0, v_1, \dots, v_{n+1})$  of degree  $i$ . Hence, using the equations (4.32) and (4.34), we can estimate the degree of the lowest degree term of the meromorphic Jacobi forms of  $\tilde{A}_n$ , more specifically,

$$\begin{aligned} \varphi_n^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} &= a_{n+2}(v) + b_{n+2}(v_{n+1}, \tau) a_{n+3}(v) + O(\|v\|^{n+4}), \\ \varphi_{n-1}^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} + b_{n-1}^n \varphi_n^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{A_1} &= a_{n+1}(v) + b_{n+1}(v_{n+1}, \tau) a_{n+2}(v) + O(\|v\|^{n+3}), \\ \varphi_{n-2}^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} + b_{n-2}^{n-1} \varphi_{n-1}^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} + a_{n-2}^n \varphi_n^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} &= a_n(v) + b_{n+2}(v_{n+1}, \tau) a_{n+1}(v) + O(\|v\|^{n+2}), \\ &\vdots \\ \varphi_0^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} + \sum_{j=1}^n b_0^j \varphi_j^{\mathcal{J}(\tilde{A}_n)} \varphi_2^{\mathcal{J}(A_1)} &= a_2(v) + b_3(v_{n+1}, \tau) a_3(v) + O(\|v\|^4). \end{aligned} \quad (4.178)$$

Note that the Christoffel symbol depend on  $\varphi_0$  iff it contains the term  $a_2^2(v)$  in its expansion. Our strategy is to show that the Christoffel symbols (4.176) contains only higher order polynomials in its expansions. Computing the lowest degree term in the expansion of (4.177)

$$\begin{aligned} \Gamma_\tau^{01} &= \frac{\partial \varphi_i}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial \varphi_j}{\partial \tau} \right) g^{pq}(v) \\ &= \frac{\partial a_{i+2}}{\partial v_p} \frac{\partial}{\partial v_q} \left( \frac{\partial b_{j+1}(v_{n+1}, \tau)}{\partial \tau} a_{j+3} \right) g^{pq}(v) + \dots \\ &= \frac{\partial a_{i+2}}{\partial v_p} \frac{\partial b_{j+1}(v_{n+1}, \tau)}{\partial \tau} \frac{\partial a_{j+3}}{\partial v_q} g^{pq}(v) + \dots \\ &= \frac{\partial b_{j+1}(v_{n+1}, \tau)}{\partial \tau} a_{i+j+3} + \dots \end{aligned} \quad (4.179)$$

Therefore, for  $i + j > 1$ , we have that the associated Christoffel symbol do not depend on  $\varphi_0^2$ . It remains to check only  $\Gamma_\tau^{01}$  and  $\Gamma_\tau^{10}$ . Computing  $\Gamma_\tau^{10}$  by using the second equation of (2.25) for  $i = 1, j = 0, k = 0$

$$\begin{aligned} \sum_{l=0}^{n+2} g^{1l} \Gamma_l^{00} &= \sum_{l=0}^{n+2} g^{0l} \Gamma_l^{10} \\ &= \sum_{l=0}^{n+1} g^{0l} \Gamma_l^{10} + g^{0\tau} \Gamma_\tau^{10}. \end{aligned}$$

Isolating  $\Gamma_\tau^{10}$ ,

$$\Gamma_\tau^{10} = \frac{1}{2\pi i \varphi_0} \left[ \sum_{l=0}^{n+2} g^{1l} \Gamma_l^{00} - \sum_{l=0}^{n+1} g^{0l} \Gamma_l^{10} \right], \quad (4.180)$$

we have that the right hand side of (4.180) depend at most linear on  $\varphi_0$ . Moreover, using the first equation (2.25), we have

$$\begin{aligned} \partial_0^2 \Gamma_\tau^{10} &= \partial_\tau \partial_0^2 g^{10} - \partial_0^2 \Gamma_\tau^{10} \\ &= \partial_\tau \partial_0^2 g^{10} = 0. \end{aligned}$$

Lemma proved. □

## 4.9 Unit and Euler vector field of the orbit space of $\mathcal{J}(\tilde{A}_n)$

The aim of this section is to define the Unit and Euler vector field and its respective actions on the geometric data of the orbit space of  $\mathcal{J}(\tilde{A}_n)$ . Further, these objects will be fundamental to define the unit of the Frobenius algebra of the desired Dubrovin Frobenius structure, and to give a quasi homogeneous property to the desired WDVV solution.

**Definition 4.7.** The Euler vector field with respect the orbit space  $\mathcal{J}(\tilde{A}_n)$  is defined by the last equation of (4.3), i.e

$$E := -\frac{1}{2\pi i} \frac{\partial}{\partial u} \quad (4.181)$$

**Definition 4.8.** A  $f$  is quasi homogeneous of degree  $d$  if it is an eigenfunction of the Euler vector field (4.181) with eigenvalue  $d$ , i.e.

$$E(f) = df.$$

**Lemma 4.43.** Let  $\lambda, \varphi_0, \dots, \varphi_n, \varphi_{n+1} = v_{n+1}, \varphi_{n+2} = \tau$  be defined in (4.25) and  $(t^0, \dots, t^n, v_{n+1}, \tau)$  the flat coordinates of eta defined in (4.115). Then,

$$\begin{aligned} E(\lambda) &= \lambda, \\ E(\varphi_i) &= d_i \varphi_i, \\ E(t^\alpha) &= d_\alpha t^\alpha, \end{aligned} \quad (4.182)$$



where

$$\begin{aligned}
d_i &= 1, \quad i < n+1, \\
d_i &= 0, \quad i \geq n+1, \\
d_\alpha &= \frac{n+1-\alpha}{n}, \quad \alpha \neq 0, \\
d_0 &= 1.
\end{aligned} \tag{4.183}$$

*Proof.* Recall that the function  $\lambda$  is given by

$$\begin{aligned}
\lambda &= e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(z - v_i + v_{n+1}, \tau)}{\theta_1^n(z, \tau) \theta_1(z + (n+1)v_{n+1})} \\
&= \varphi_n \wp^{n-2}(z, \tau) + \varphi_{n-1} \wp^{n-3}(z, \tau) + \dots + \varphi_2 \wp(z, \tau) \\
&\quad + \varphi_1 [\zeta(z, \tau) - \zeta(z + (n+1)v_{n+1}, \tau)] + \varphi_0.
\end{aligned}$$

Hence,

$$\begin{aligned}
\lambda &= \frac{1}{2\pi i} \frac{\partial}{\partial u} (\lambda) \\
&= E(\varphi_n) \wp^{n-2}(z, \tau) + E(\varphi_{n-1}) \wp^{n-3}(z, \tau) + \dots + E(\varphi_2) \wp(z, \tau) \\
&\quad + E(\varphi_1) [\zeta(z, \tau) - \zeta(z + (n+1)v_{n+1}, \tau)] + E(\varphi_0),
\end{aligned}$$

therefore,  $E(\varphi_i) = \varphi_i$ , and  $E(v_{n+1}) = E(\tau) = 0$ .

Recall that  $t^\alpha$  can be written in terms of equation (4.119) or in more convenient way

$$\begin{aligned}
t^\alpha &= \frac{n}{n+1-\alpha} \operatorname{res}_{v=0} \lambda^{\frac{n+1-\alpha}{n}}(v) dv, \quad \alpha \neq 0, \\
t^0 &= \varphi_0 - \frac{\theta_1'((n+1)v_{n+1})}{\theta_1((n+1)v_{n+1})} \varphi_1 + 4\pi i g_1(\tau) \varphi_2.
\end{aligned} \tag{4.184}$$

Applying the Euler vector in (4.184) we get the desired result.  $\square$

**Corollary 4.43.1.** The Euler vector field (4.181) in the flat coordinates of  $\eta^*$  has the following form

$$E := \sum_{\alpha=0}^n d_\alpha t^\alpha \frac{\partial}{\partial t^\alpha}, \tag{4.185}$$

where

$$\begin{aligned}
d_\alpha &= \frac{n+1-\alpha}{n}, \quad \alpha \neq 0, \\
d_0 &= 1.
\end{aligned} \tag{4.186}$$

*Proof.* Recall that

$$E = \frac{1}{2\pi i} \frac{\partial}{\partial u} = E(t^\alpha) \frac{\partial}{\partial t^\alpha} = \sum_{\alpha=0}^n d_\alpha t^\alpha \frac{\partial}{\partial t^\alpha}.$$

$\square$

---

**Lemma 4.44.** The Euler vector field (4.181) acts on the vector fields  $\frac{\partial}{\partial t^\alpha}$ ,  $\frac{\partial}{\partial \varphi_i}$  and differential forms  $dt^\alpha$ ,  $d\varphi_i$  as follows:

$$\begin{aligned} Lie_E d\varphi_i &= d_i d\varphi_i, \\ Lie_E dt^\alpha &= d_\alpha dt^\alpha, \\ Lie_E \frac{\partial}{\partial \varphi_i} &= -d_i \frac{\partial}{\partial \varphi_i}, \\ Lie_E \frac{\partial}{\partial t^\alpha} &= -d_\alpha \frac{\partial}{\partial t^\alpha}. \end{aligned} \tag{4.187}$$

*Proof.* Recall that the Lie derivative acts in vector fields by using the Lie bracket and in differential forms by the use of Cartan's magic formula

$$\begin{aligned} Lie_E \frac{\partial}{\partial t^\alpha} &= \left[ E, \frac{\partial}{\partial t^\alpha} \right], \\ Lie_E dt^\alpha &= dE(dt^\alpha) + E(d^2 t^\alpha) = dE(dt^\alpha). \end{aligned} \tag{4.188}$$

Using (4.188) and (4.182), we obtain the desired result.  $\square$

**Lemma 4.45.** The intersection form  $g^{ij}$  defined in (4.40) and its Christoffel symbol  $\Gamma_k^{ij}$  in the coordinates  $\varphi_0, \dots, \varphi_n, \varphi_{n+1} = v_{n+1}, \varphi_{n+2} = \tau$  defined in (4.25) are weighted polynomials in the variables  $\varphi_0, \dots, \varphi_n$ , with degrees

$$\deg(g^{ij}) = d_i + d_j, \quad \deg(\Gamma_k^{\alpha\beta}) = d_i + d_j - d_k. \tag{4.189}$$

*Proof.* The function  $g^{ij}$  and  $\Gamma_k^{ij}$  belong to the ring  $\tilde{E}_{\bullet, \bullet}[\varphi_0, \varphi_1, \dots, \varphi_n]$  due to 4.13 and 4.41. The degrees are computed by using the following formulae

$$\begin{aligned} E(g^{ij}(\varphi)) &= E\left(\frac{\partial \varphi_i}{\partial v^l} \frac{\partial \varphi_j}{\partial v^m} g^{lm}(v)\right) \\ &= E\left(\frac{\partial \varphi_i}{\partial v^l}\right) \frac{\partial \varphi_j}{\partial v^m} g^{lm}(v) + \frac{\partial \varphi_i}{\partial v^l} E\left(\frac{\partial \varphi_j}{\partial v^m} g^{lm}(v)\right) \\ &= \frac{\partial E(\varphi_i)}{\partial v^l} \frac{\partial \varphi_j}{\partial v^m} g^{lm}(v) + \frac{\partial \varphi_i}{\partial v^l} \frac{\partial E(\varphi_j)}{\partial v^m} g^{lm}(v) \\ &= (d_i + d_j) \frac{\partial \varphi_i}{\partial v^l} \frac{\partial \varphi_j}{\partial v^m} g^{lm}(v). \end{aligned}$$

and

$$\begin{aligned} E(\Gamma_k^{ij}(\varphi)) &= E\left(\frac{\partial \varphi_i}{\partial v^l} \frac{\partial}{\partial \varphi_k} \left(\frac{\partial \varphi_j}{\partial v^m}\right) g^{lm}(v)\right) \\ &= E\left(\frac{\partial \varphi_i}{\partial v^l}\right) \frac{\partial}{\partial \varphi_k} \left(\frac{\partial \varphi_j}{\partial v^m}\right) g^{lm}(v) + \frac{\partial \varphi_i}{\partial v^l} E\left(\frac{\partial}{\partial \varphi_k} \left(\frac{\partial \varphi_j}{\partial v^m}\right) g^{lm}(v)\right) \\ &= \frac{\partial E(\varphi_i)}{\partial v^l} \frac{\partial}{\partial \varphi_k} \left(\frac{\partial \varphi_j}{\partial v^m}\right) g^{lm}(v) + \frac{\partial \varphi_i}{\partial v^l} \frac{\partial}{\partial \varphi_k} \left(\frac{\partial E(\varphi_j)}{\partial v^m}\right) g^{lm}(v) \\ &\quad - d_k \frac{\partial \varphi_i}{\partial v^l} \frac{\partial}{\partial \varphi_k} \left(\frac{\partial \varphi_j}{\partial v^m}\right) g^{lm}(v) \\ &= (d_i + d_j - d_k) \frac{\partial \varphi_i}{\partial v^l} \frac{\partial}{\partial \varphi_k} \left(\frac{\partial \varphi_j}{\partial v^m}\right) g^{lm}(v). \end{aligned}$$

$\square$

**Lemma 4.46.** The intersection form  $g^{\alpha\beta}$  defined in (4.40) in the coordinates  $(t^0, \dots, t^n, v_{n+1}, \tau)$  defined in (4.115) and its Christoffel symbol  $\Gamma_\gamma^{\alpha\beta}$  are weighted polynomials in the variables  $t^0, t^1, \dots, t^n, \frac{1}{t^n}$  with degrees

$$\begin{aligned} \deg(g^{\alpha\beta}) &= d_\alpha + d_\beta, \\ \deg(\Gamma_\gamma^{\alpha\beta}) &= d_\alpha + d_\beta - d_\gamma. \end{aligned} \quad (4.190)$$

*Proof.* Lemma 4.37 guarantee that  $g^{\alpha\beta} \in \tilde{E}_{\bullet, \bullet}[t^0, t^1, \dots, t^n, \frac{1}{t^n}]$ . Using the formula

$$\begin{aligned} E(g^{\alpha\beta}) &= E\left(\frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi)\right) \\ &= E\left(\frac{\partial t^\alpha}{\partial \varphi_i}\right) \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi) + \frac{\partial t^\alpha}{\partial \varphi_i} E\left(\frac{\partial t^\beta}{\partial \varphi_j}\right) g^{ij}(\varphi) + \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} E(g^{ij}(\varphi)) \\ &= \frac{\partial E(t^\alpha)}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi) - d_i \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi) + \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial E(t^\beta)}{\partial \varphi_j} g^{ij}(\varphi) - d_j \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi) \\ &\quad + (d_i + d_j) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi) \\ &= (d_\alpha + d_\beta) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} g^{ij}(\varphi). \end{aligned}$$

The Christoffel symbol  $\Gamma_\gamma^{\alpha\beta}$  is given by the following

$$\Gamma_\gamma^{\alpha\beta} = \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} \frac{\partial \varphi_k}{\partial t^\gamma} \Gamma_k^{ij} + \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial}{\partial t^\gamma} \left( \frac{\partial t^\beta}{\partial \varphi_j} \right) g^{ij}.$$

$\Gamma_j^{ij}, \frac{\partial \varphi_k}{\partial t^\gamma} \in \tilde{E}_{\bullet, \bullet}[t^0, t^1, \dots, t^n]$  due to Lemma 4.41 and equations (4.132), (4.129). But using (4.121), we realise that  $\frac{\partial t^\alpha}{\partial \varphi_i}$  is polynomial in  $t^1, t^2, \dots, t^n, \frac{1}{t^n}$  due to equations (4.127) and (4.122). Therefore,  $\Gamma_\gamma^{\alpha\beta}$  are weighted polynomials in the variables  $t^0, t^1, \dots, t^n, \frac{1}{t^n}$ . Computing the degree of  $\Gamma_\gamma^{\alpha\beta}$

$$\begin{aligned} E(\Gamma_\gamma^{\alpha\beta}) &= E\left(\frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} \frac{\partial \varphi_k}{\partial t^\gamma} \Gamma_k^{ij} + \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial}{\partial t^\gamma} \left( \frac{\partial t^\beta}{\partial \varphi_j} \right) g^{ij}\right) \\ &= (d_\alpha - d_i) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} \frac{\partial \varphi_k}{\partial t^\gamma} \Gamma_k^{ij} + (d_\beta - d_j) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} \frac{\partial \varphi_k}{\partial t^\gamma} \Gamma_k^{ij} \\ &\quad + (d_k - d_\gamma) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial t^\beta}{\partial \varphi_j} \frac{\partial \varphi_k}{\partial t^\gamma} \Gamma_k^{ij} + (d_\alpha - d_i) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial}{\partial t^\gamma} \left( \frac{\partial t^\beta}{\partial \varphi_j} \right) g^{ij} \\ &\quad + (d_\beta - d_j) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial}{\partial t^\gamma} \left( \frac{\partial t^\beta}{\partial \varphi_j} \right) g^{ij} + (d_\alpha - d_i) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial}{\partial t^\gamma} \left( \frac{\partial t^\beta}{\partial \varphi_j} \right) g^{ij} \\ &\quad + (d_i + d_j) \frac{\partial t^\alpha}{\partial \varphi_i} \frac{\partial}{\partial t^\gamma} \left( \frac{\partial t^\beta}{\partial \varphi_j} \right) g^{ij} \\ &= (d_\alpha + d_\beta - d_\gamma) \Gamma_\gamma^{\alpha\beta}. \end{aligned}$$

□

**Definition 4.9.** The Unit vector field with respect the orbit space  $\mathcal{J}(\tilde{A}_n)$  is the vector associated to the invariant coordinate  $\varphi_0$  defined in (4.25), i.e

$$e := \frac{\partial}{\partial \varphi_0}. \quad (4.191)$$

---

**Lemma 4.47.** The Unit vector field (4.181) in the flat coordinates of  $\eta^*$  has the following form

$$e = \frac{\partial}{\partial t^0}. \quad (4.192)$$

*Proof.*

$$\frac{\partial}{\partial \varphi_0} = \frac{\partial t^\alpha}{\partial \varphi_0} \frac{\partial}{\partial t^\alpha} = \frac{\partial}{\partial t^0}.$$

□

**Lemma 4.48.** Let the metric  $\eta^*$  be defined on (4.77) and the Euler vector field (4.181). Then,

$$Lie_E \eta^{\alpha\beta} = (d_\alpha + d_\beta - d_1) \eta^{\alpha\beta}. \quad (4.193)$$

#### 4.10 Discriminant locus and the monodromy of the orbit space of $\mathcal{J}(\tilde{A}_n)$

This section relates the critical points of the (4.25) with the zeros of the determinant of the intersection form  $g^*$  (4.41). Further, in section 4.11, it will be built a Frobenius algebra in the sections of the orbit space of  $\mathcal{J}(\tilde{A}_n)$ , furthermore, the intersection form  $g^*$  (4.41) can be realised as the multiplication by the Euler vector field (4.181). The results of this section will imply that the intersection form  $g^*$  (4.41) is diagonalisable with eigenvalues generically different from 0, and this is equivalent to the Frobenius algebra be semisimple. Moreover, we can realise the isomorphism of orbit space of  $\mathcal{J}(\tilde{A}_n)$  with the Hurwitz space  $H_{1,n-1,0}$  as a Dubrovin Frobenius manifolds, see Theorem 4.59 for details.

**Definition 4.10.** Let  $g^*$  the metric defined on (4.41). The discriminant locus of the orbit space of  $\mathcal{J}(\tilde{A}_n) \subset \mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} / \mathcal{J}(\tilde{A}_n)$  is defined by

$$\Sigma = \{x \in \mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} / \mathcal{J}(\tilde{A}_n) : \det(g^*) = 0\}. \quad (4.194)$$

**Lemma 4.49.** The fixed points of the action  $\mathcal{J}(\tilde{A}_n)$  belong to the discriminant locus (4.194).

*Proof.* Note that the fixed points of the action  $\mathcal{J}(\tilde{A}_n)$  on  $\mathbb{C} \oplus \mathbb{C}^{n+1} \oplus \mathbb{H} \ni (u, v_0, v_1, \dots, v_{n-1}, v_{n+1}, \tau)$  are the fixed points of the action  $A_n$  on  $\mathbb{C}^n \ni (v_0, v_1, \dots, v_{n-1})$ . Therefore, the fixed points are the hyperplanes

$$v_i = v_j \quad i, j \in \{0, 1, \dots, n-1\}. \quad (4.195)$$

The intersection form (4.41) is given by

$$\begin{aligned} g &= \sum_{i=0}^n dv_i^2 \Big|_{\sum_{i=0}^n v_i=0} - n(n+1)dv_{n+1}^2 + 2dud\tau \\ &= \sum_{i,j=0}^{n-1} A_{ij} dv_i dv_j - n(n+1)dv_{n+1}^2 + 2dud\tau \end{aligned} \quad (4.196)$$

The intersection form is given by

$$g^* = \sum_{i,j=0}^{n-1} A_{ij}^{-1} \frac{\partial}{\partial v_i} \otimes \frac{\partial}{\partial v_j} - \frac{1}{n(n+1)} \frac{\partial}{\partial v_{n+1}} \otimes \frac{\partial}{\partial v_{n+1}} + \frac{\partial}{\partial u} \otimes \frac{\partial}{\partial \tau} + \frac{\partial}{\partial \tau} \otimes \frac{\partial}{\partial u} \quad (4.197)$$

became degenerate on the hyperplanes (4.195), because two columns of the matrix  $A_{ij}^{-1}$  became proportional.

□

---

**Lemma 4.50.** The function  $\lambda(p, u, v_0, v_1, \dots, v_{n+1}, \tau)$  defined on (4.25) has simple critical points if and only if  $(u, v_0, v_1, \dots, v_{n+1}, \tau)$  is a fixed point of the action  $\mathcal{J}(\tilde{A}_n)$ .

*Proof.* Using the local isomorphism given by 4.21

$$[(u, v_0, v_1, \dots, v_{n-1}, v_{n+1}, \tau)] \longleftrightarrow \lambda(v) = e^{-2\pi i u} \frac{\prod_{i=0}^n \theta_1(z - v_i, \tau)}{\theta_1^n(v, \tau) \theta_1(v + (n+1)v_{n+1}, \tau)}, \quad (4.198)$$

we can realise the discriminant locus as the space of parameters of  $\lambda(p, u, v_0, v_1, \dots, v_{n+1}, \tau)$  such that  $\lambda(p, u, v_0, v_1, \dots, v_{n+1}, \tau)$  has repeated roots. In these cases  $\lambda(p, u, v_0, v_1, \dots, v_{n+1}, \tau)$  has non simple critical points. □

**Definition 4.11.** The canonical coordinates  $(u_1, u_2, \dots, u_{n+2})$  of the orbit space  $\mathcal{J}(\tilde{A}_n)$  is given by the following relation

$$\begin{aligned} \lambda(q_i) &= u_i, \\ \lambda'(q_i) &= 0. \end{aligned} \quad (4.199)$$

**Lemma 4.51.** The determinant of the intersection form  $g^*$  defined on (4.40) is proportional to  $\prod_{i=1}^{n+2} u_i$ .

*Proof.* If  $u_i = \lambda(q_i, u, v_0, \dots, v_{n+1}, \tau) = 0$ , we have that  $\det g^* = 0$  due to lemma (4.50), then  $u_i$  are zeros of the equation  $\det g^* = 0$ . □

**Proposition 4.52.** In the canonical coordinates  $(u_1, u_2, \dots, u_{n+2})$  the unit vector field (4.191), the Euler vector field (4.181), and the intersection form (4.40) have the following form

$$\begin{aligned} g^{ii} &= u^i \eta^{ii} \delta_{ij}, \\ e &= \sum_{i=1}^{n+2} \frac{\partial}{\partial u_i}, \\ E &= \sum_{i=1}^{n+2} u_i \frac{\partial}{\partial u_i}. \end{aligned} \quad (4.200)$$

where  $\eta^{ii}$  are the coefficients of second metric  $\eta^*$  in canonical coordinates.

*Proof.* Note that  $g^*$  is diagonalisable with distinct eigenvalues if the following equation

$$\det(\eta_{\alpha\mu} g^{\mu\beta} - u \delta_\alpha^\beta) = 0, \quad (4.201)$$

has only simple roots. Since  $\det(\eta_{\alpha\mu}) \neq 0$ , the equation (4.201) is equivalent to

$$\det(g^{\alpha\beta} - u \eta^{\alpha\beta}) = 0. \quad (4.202)$$

Using that  $\eta^{\alpha\beta} = \partial_0 g^{\alpha\beta}$ , we have that

$$\det(g^{\alpha\beta} - u \eta^{\alpha\beta}) = \det(g^{\alpha\beta}(t^0 - u, t^1, t^2, t^3, \dots, t^n, v_{n+1}, \tau)) = 0. \quad (4.203)$$

Due to the lemma 4.51 the equation (4.202) has  $n + 2$  distinct roots

$$u^i = t^1 - y^i(t^1, t^2, t^3, \dots, t^n, v_{n+1}, \tau). \quad (4.204)$$

In the coordinates  $(u^1, u^2, \dots, u^{n+2})$  the matrix  $g_j^i$  is diagonal, then

$$g^{ij} = u^i \eta^{ij} \delta_{ij} \quad (4.205)$$

and the unit vector field have the following form

$$\frac{\partial}{\partial t^1} = \sum_{i=1}^{n+2} \frac{\partial u_i}{\partial t^1} \frac{\partial}{\partial u_i} = \sum_{i=1}^{n+2} \frac{\partial}{\partial u_i} \quad (4.206)$$

Moreover, since

$$[E, e] = \left[ \sum_{\alpha=0}^n d_\alpha t^\alpha \frac{\partial}{\partial t^\alpha}, \frac{\partial}{\partial t^1} \right] = -e, \quad (4.207)$$

the Euler vector field in the coordinates  $(u^1, u^2, \dots, u^{n+2})$  takes the following form

$$E = \sum_{i=1}^{n+2} u^i \frac{\partial}{\partial u_i}. \quad (4.208)$$

Lemma proved. □

#### 4.11 Construction of WDVV solution

The main aim of this section is to extract a WDVV equation from the data of the group  $\mathcal{J}(\tilde{A}_n)$ .

**Lemma 4.53.** The orbit space of  $\mathcal{J}(\tilde{A}_n)$  carries a flat pencil of metrics

$$g^{\alpha\beta}, \quad \eta^{\alpha\beta} := \frac{\partial g^{\alpha\beta}}{\partial t^0} \quad (4.209)$$

with the correspondent Christoffel symbols.

$$\Gamma_\gamma^{\alpha\beta}, \quad \eta^{\alpha\beta} := \frac{\partial \Gamma_\gamma^{\alpha\beta}}{\partial t^0} \quad (4.210)$$

*Proof.* The metric (4.209) satisfies the hypothesis of Lemma 2.7 which proves the desired result. □

**Lemma 4.54.** Let the intersection form be (4.40), unit vector field be (4.191), and Euler vector field be (4.181). Then, there exist a function

$$F(t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau) = -\frac{(t^0)^2 \tau}{4\pi i} + \frac{t^0}{2} \sum_{\alpha, \beta \neq 0, \tau} \eta_{\alpha\beta} t^\alpha t^\beta + G(t^1, t^2, \dots, t^n, v_{n+1}, \tau), \quad (4.211)$$

such that

$$\begin{aligned} Lie_E F &= 2F + \text{quadratic terms}, \\ Lie_E (F^{\alpha\beta}) &= g^{\alpha\beta}, \\ \frac{\partial^2 G(t^1, t^2, \dots, t^n, v_{n+1}, \tau)}{\partial t^\alpha \partial t^\beta} &\in \tilde{E}_{\bullet, \bullet} [t^1, t^2, \dots, t^n, \frac{1}{t^n}], \end{aligned} \quad (4.212)$$

where

$$F^{\alpha\beta} = \eta^{\alpha\alpha'} \eta^{\beta\beta'} \frac{\partial F^2}{\partial t^{\alpha'} \partial t^{\beta'}}. \quad (4.213)$$

---

*Proof.* Let  $\Gamma_{\gamma}^{\alpha\beta}(t)$  the Christoffel symbol of the intersection form (4.40) in the coordinates the flat coordinates of  $\eta^*$ , i.e  $t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau$ . According to the lemma 2.7, we can represent  $\Gamma_{\gamma}^{\alpha\beta}(t)$  as

$$\Gamma_{\gamma}^{\alpha\beta}(t) = \eta^{\alpha\epsilon} \partial_{\epsilon} \partial_{\gamma} f^{\beta}(t). \quad (4.214)$$

Using the relations (4.190), (4.187) and lemma 4.48

$$\begin{aligned} Lie_E(\Gamma_{\gamma}^{\alpha\beta}(t)) &= Lie_E(\eta^{\alpha\epsilon}) \partial_{\epsilon} \partial_{\gamma} f^{\beta}(t) + \eta^{\alpha\epsilon} Lie_E(\partial_{\epsilon} \partial_{\gamma} f^{\beta}(t)) \\ &= (d_{\alpha} + d_{\epsilon} - d_1) \eta^{\alpha\epsilon} \partial_{\epsilon} \partial_{\gamma} f^{\beta}(t) + (-d_{\epsilon} - d_{\gamma}) \eta^{\alpha\epsilon} \partial_{\epsilon} \partial_{\gamma} Lie_E(f^{\beta}(t)) \\ &= (d_{\alpha} + d_{\beta} - d_{\gamma}) \eta^{\alpha\epsilon} \partial_{\epsilon} \partial_{\gamma} f^{\beta}(t). \end{aligned}$$

Then, by isolation  $Lie_E(f^{\beta}(t))$  we get

$$Lie_E(f^{\beta}(t)) = (d_{\beta} + d_1) f^{\beta} + A_{\sigma}^{\beta} t^{\sigma} + B^{\beta}, \quad A_{\sigma}^{\beta}, B^{\beta} \in \mathbb{C}. \quad (4.215)$$

Considering the second relation of (2.25) for  $\alpha = \tau$

$$g^{\tau\sigma} \Gamma_{\sigma}^{\beta\gamma} = g^{\beta\sigma} \Gamma_{\sigma}^{\tau\gamma}, \quad (4.216)$$

and using lemma 4.21 and 4.39, we have.

$$-2\pi i d_{\sigma} t^{\sigma} \eta^{\beta\epsilon} \partial_{\sigma} \partial_{\epsilon} f^{\gamma} = -2\pi i d_{\sigma} \delta_{\sigma}^{\gamma} g^{\beta\sigma}, \quad (4.217)$$

which is equivalent to

$$Lie_E(\eta^{\beta\epsilon} \partial_{\epsilon} f^{\gamma}) = d_{\gamma} g^{\beta\gamma}. \quad (4.218)$$

Using (4.215) in the equation (4.218), we have

$$(d_{\beta} + d_{\gamma}) \eta^{\beta\epsilon} \partial_{\epsilon} f^{\gamma} = d_{\gamma} g^{\beta\gamma}. \quad (4.219)$$

If  $\gamma \neq v_{n+1}, \tau$ , we define

$$F^{\gamma} = \frac{f^{\gamma}}{d_{\gamma}}, \quad (4.220)$$

and note that  $g^{\beta\gamma}$  is symmetric with respect the indices  $\beta, \gamma$ . Hence,

$$(d_{\beta} + d_{\gamma}) \eta^{\beta\epsilon} \partial_{\epsilon} F^{\gamma} = (d_{\beta} + d_{\gamma}) \eta^{\gamma\epsilon} \partial_{\epsilon} F^{\beta}, \quad (4.221)$$

which is the integrability condition for

$$F^{\gamma} = \eta^{\gamma\mu} \partial_{\mu} F. \quad (4.222)$$

In order to extract information from  $\gamma = \tau$ , take  $\beta = \tau$  in equation (4.219)

$$\begin{aligned} d_{\gamma} \eta^{\tau 0} \partial_0 f^{\gamma} &= d_{\gamma} g^{\tau\gamma} \\ -2\pi i d_{\gamma} \partial_0 f^{\gamma} &= -2\pi i d_{\gamma} t^{\gamma} \end{aligned} \quad (4.223)$$

which is equivalent to

$$\eta^{\gamma\epsilon} \partial_{\epsilon} \partial_0 F = t^{\gamma},$$

inverting  $\eta^{\gamma\epsilon}$

$$\partial_\alpha \partial_0 F = \eta_{\alpha\gamma} t^\gamma, \quad (4.224)$$

integrating equation (4.224), we obtain

$$F(t^0, t^1, t^2, \dots, t^n, v_{n+1}, \tau) = -\frac{(t^0)^2 \tau}{4\pi i} + \frac{t^0}{2} \sum_{\alpha, \beta \neq 0, \tau} \eta_{\alpha\beta} t^\alpha t^\beta + G(t^1, t^2, \dots, t^n, v_{n+1}, \tau). \quad (4.225)$$

Substituting the equation (4.225) in the (4.219) for  $\gamma \neq v_{n+1}, \tau$ , we get

$$\begin{aligned} g^{\beta\gamma} &= (d_\beta + d_\gamma) \eta^{\beta\epsilon} \eta^{\gamma\mu} \partial_\epsilon \partial_\mu F, \\ &= Lie_E(F^{\beta\gamma}) \end{aligned} \quad (4.226)$$

Since  $g^{\beta\gamma}$  is a symmetric matrix the equation (4.226) is equivalent to the second equation of (4.212) for either  $\beta$  and  $\gamma$  different from  $v_{n+1}, \tau$ . Therefore, the missing part of the second equation of (4.212) is only for the cases  $\beta = \gamma = v_{n+1}$  and  $\beta = \gamma = \tau$ . Moreover, the intersection form  $g^{\beta\gamma}$  is proportional to the Hessian of the equation (4.225) for either  $\beta$  and  $\gamma$  different from  $v_{n+1}, \tau$ . Recall that from the data of a Hessian, we can reconstruct uniquely a function up to quadratic terms, therefore, by defining

$$\begin{aligned} Lie_E \left( \frac{\partial^2 F}{\partial t^{12}} \right) &= g^{v_{n+1}v_{n+1}}, \\ Lie_E \left( \frac{\partial^2 F}{\partial t^{12}} \right) &= g^{\tau\tau}. \end{aligned} \quad (4.227)$$

Just the second equation of (4.227) need to be proved, since the first equation defines the coefficients of the Hessian  $\frac{\partial^2 F}{\partial t^{12}}$ , in another words, it defines  $\frac{\partial^2 G}{\partial t^{12}}$ . The second equation must be compatible with the equation (4.225), then substituting (4.225) in the second equation of (4.227).

$$\begin{aligned} Lie_E \left( \frac{\partial^2 F}{\partial t^{12}} \right) &= Lie_E \left( \frac{\tau}{2\pi i} \right) \\ &= 0 = g^{\tau\tau}. \end{aligned}$$

Hence, we proved the second equation (4.212). Substituting the equation (4.225) in the second equation (4.212) for  $\alpha, \beta \neq \tau$

$$\begin{aligned} Lie_E(F^{\alpha\beta}) &= Lie_E \left( \eta^{\alpha\alpha'} \eta^{\beta\beta'} \frac{\partial F^2}{\partial t^{\alpha'} \partial t^{\beta'}} \right) \\ &= Lie_E \left( \eta^{\alpha\alpha'} \eta^{\beta\beta'} \frac{\partial G^2}{\partial t^{\alpha'} \partial t^{\beta'}} \right) \\ &= g^{\alpha\beta} \in \tilde{E}_{\bullet, \bullet}[t^1, t^2, \dots, t^n, \frac{1}{t^n}] \end{aligned}$$

Hence, the second equation (4.212) prove the third equation of (4.212).

Substituting (4.225) in (4.215), (4.222)

$$\begin{aligned} Lie_E(f^\beta) &= Lie_E(\eta^{\beta\epsilon} \partial_\epsilon F) \\ &= Lie_E(\eta^{\beta\epsilon} \partial_\epsilon F) \partial_\epsilon F + \eta^{\beta\epsilon} Lie_E(\partial_\epsilon F) \\ &= (d_\beta + d_\epsilon - d_1) \eta^{\beta\epsilon} \partial_\epsilon F \partial_\epsilon F + \eta^{\beta\epsilon} \partial_\epsilon Lie_E(F) - d_\epsilon \eta^{\beta\epsilon} \partial_\epsilon F \\ &= (d_\beta - d_1) \eta^{\beta\epsilon} \partial_\epsilon F \partial_\epsilon F + \eta^{\beta\epsilon} \partial_\epsilon Lie_E(F) \\ &= (d_\beta + d_1) \eta^{\beta\epsilon} \partial_\epsilon F + A_\sigma^\beta t^\sigma + B^\beta \end{aligned}$$



---

Hence, isolating  $Lie_E(F)$

$$\eta^{\beta\epsilon}\partial_\epsilon Lie_E(F) = 2\eta^{\beta\epsilon}\partial_\epsilon F + A_\sigma^\beta t^\sigma + B^\beta,$$

inverting  $\eta^{\beta\epsilon}$

$$\partial_\alpha Lie_E(F) = 2\partial_\alpha F + \eta_{\alpha\beta} A_\sigma^\beta t^\sigma + \eta_{\alpha\beta} B^\beta,$$

integrating

$$Lie_E(F) = 2F + \eta_{\alpha\beta} A_\sigma^\beta t^\sigma + \eta_{\alpha\beta} B^\beta t^\alpha,$$

Lemma proved. □

**Lemma 4.55.** Let be

$$c_{\alpha\beta\gamma} = \frac{\partial F^3}{\partial t^\alpha \partial t^\beta \partial t^\gamma}, \quad (4.228)$$

then,

$$c_{\alpha\beta}^\gamma = \eta^{\gamma\epsilon} c_{\alpha\beta\epsilon} \quad (4.229)$$

is a structure constant of a commutative algebra given by the following rule in the flat coordinate of  $\eta$

$$\partial_\alpha \bullet \partial_\beta = c_{\alpha\beta}^\gamma \partial_\gamma \quad (4.230)$$

such that

$$\eta(\partial_\alpha \bullet \partial_\beta, \partial_\gamma) = \eta(\partial_\alpha, \partial_\beta \bullet \partial_\gamma), \quad \textbf{Frobenius condition.} \quad (4.231)$$

*Proof.*

### 1. Commutative

The product defined in (4.230) is commutative, because its structure constant (4.229) is symmetric with respect its indices  $\alpha, \beta, \gamma$  due to the commutative behaviour of the partial derivatives  $\frac{\partial}{\partial t^\alpha}, \frac{\partial}{\partial t^\beta}, \frac{\partial}{\partial t^\gamma}$ .

### 2. Frobenius condition

$$\begin{aligned} \eta(\partial_\alpha \bullet \partial_\beta, \partial_\gamma) &= c_{\alpha\beta}^\epsilon \eta(\partial_\epsilon, \partial_\gamma) \\ &= c_{\alpha\beta}^\epsilon \eta_{\epsilon\gamma} \\ &= c_{\alpha\beta\gamma} \\ &= c_{\beta\gamma}^\epsilon \eta_{\alpha\epsilon} = \eta(\partial_\alpha, \partial_\beta \bullet \partial_\gamma). \end{aligned}$$

Lemma proved. □

**Lemma 4.56.** The unit vector field be defined in (4.191) is the unit of the algebra defined in lemma 4.55.

---

*Proof.* Substituting (4.222) and (4.220) in (4.214), we obtain

$$\Gamma_{\gamma}^{\alpha\beta} = d_{\beta} c_{\gamma}^{\alpha\beta}, \quad (4.232)$$

where

$$c_{\gamma}^{\alpha\beta} = \eta^{\alpha\mu} \eta^{\beta\epsilon} c_{\epsilon\mu\gamma}. \quad (4.233)$$

Substituting  $\alpha = \tau$  in (4.232) and using lemma 4.39

$$\begin{aligned} \Gamma_{\gamma}^{\tau\beta} &= -2\pi i d_{\beta} \delta_{\gamma}^{\beta}, \\ &= d_{\beta} c_{\gamma}^{\tau\beta}. \end{aligned}$$

Then,

$$c_{0\gamma}^{\beta} = \delta_{\gamma}^{\beta}.$$

Computing

$$\partial_0 \bullet \partial_{\gamma} = c_{0\gamma}^{\beta} \partial_{\beta} = \partial_{\gamma}.$$

Lemma proved. □

**Lemma 4.57.** The algebra defined in lemma 4.55 is associative and semisimple.

*Proof.* Recall that the Christoffel symbol  $\Gamma_{\gamma}^{\alpha\beta}$  is proportional to the structure constant of the algebra defined in lemma 4.55 for  $\beta \neq v_{n+1}, \tau$

$$\Gamma_{\gamma}^{\alpha\beta} = d_{\beta} c_{\gamma}^{\alpha\beta}.$$

Then, using (2.36), we obtain

$$\Gamma_{\sigma}^{\alpha\beta} \Gamma_{\epsilon}^{\sigma\gamma} = \Gamma_{\sigma}^{\alpha\gamma} \Gamma_{\epsilon}^{\sigma\beta} \quad (4.234)$$

Substituting (4.232) in (4.234), we have

$$c_{\sigma}^{\alpha\beta} c_{\epsilon}^{\sigma\gamma} = c_{\sigma}^{\alpha\gamma} c_{\epsilon}^{\sigma\beta}, \quad \text{for } \beta, \gamma \neq v_{n+1}, \tau.$$

If  $\beta = \tau$ ,

$$\begin{aligned} c_{\sigma}^{\alpha\tau} c_{\epsilon}^{\sigma\gamma} &= -2\pi i \delta_{\sigma}^{\alpha} c_{\epsilon}^{\sigma\gamma} \\ &= -2\pi i c_{\epsilon}^{\alpha\gamma} \\ &= -2\pi i \delta_{\epsilon}^{\sigma} c_{\sigma}^{\alpha\gamma} \\ &= c_{\sigma}^{\alpha\gamma} c_{\epsilon}^{\sigma\tau}. \end{aligned}$$

In order to prove the associativity for  $\beta = v_{n+1}$ , note that the multiplication by the Euler vector field is almost the same of the intersection form  $g^*$ . Indeed,

$$\begin{aligned} E \bullet \partial_{\alpha} &= t^{\sigma} c_{\sigma\alpha}^{\beta} \partial_{\beta} = t^{\sigma} \partial_{\sigma} (\eta^{\beta\mu} \partial_{\alpha} \partial_{\mu} F) \partial_{\beta} = \\ &= (d_{\alpha} - d_{\beta}) \eta^{\beta\mu} \partial_{\alpha} \partial_{\mu} F \partial_{\beta} = \eta_{\alpha\mu} g^{\mu\beta} \partial_{\beta} \end{aligned} \quad (4.235)$$

Then,

$$E^\sigma c_\sigma^{\alpha\beta} = g^{\alpha\beta}. \quad (4.236)$$

Using the relation (4.236) in the coordinates  $(u^1, u^2, \dots, u^{n+2})$ , we have

$$u^i \eta^{ij} \delta_{ij} = u^l \eta^{lm} \eta^{jn} c_{lmn}, \quad (4.237)$$

differentiating both side of the equation (4.237) with respect  $t^1$

$$c_{ij}^k = \delta_{ij}, \quad (4.238)$$

which proves that the algebra is associative and semisimple.  $\square$

Recall of the covering space of the orbit space of  $\mathcal{J}(\tilde{A}_n)$  defined in (4.149), see section 4.7 for details.

**Theorem 4.58.** The covering  $\mathbb{C} \oplus \widetilde{\mathbb{C}^n} \oplus \mathbb{H} / \mathcal{J}(\tilde{A}_n)$  with the intersection form (4.40), unit vector field (4.191), and Euler vector field (4.181) has a Dubrovin Frobenius manifold structure.

*Proof.* The function (4.211) satisfy a WDVV equation due to the lemmas 4.54, 4.55, 4.56, 4.57.  $\square$

Taking the same covering taking in the orbit space of  $\mathcal{J}(\tilde{A}_n)$  in the Hurwitz space  $H_{1,n-1,0}$ , fixing a symplectic base of cycle, a chamber in the tori where the variable  $v_{n+1}$  lives, and branching root of  $\varphi_n$ , denoting this covering by

$$\tilde{H}_{1,n-1,0}, \quad (4.239)$$

we obtain

**Theorem 4.59.** The Dubrovin Frobenius structure of the covering space  $\mathbb{C} \oplus \widetilde{\mathbb{C}^n} \oplus \mathbb{H} / \mathcal{J}(\tilde{A}_n)$  is isomorphic as Dubrovin Frobenius manifold to the covering  $\tilde{H}_{1,n-1,0}$ .

*Proof.* Both the orbit space  $\mathcal{J}(\tilde{A}_n)$  and the Hurwitz space  $H_{1,n-1,0}$  has the same intersection form, Euler vector, unit vector field due to proposition 4.52, lemma 4.50 and 4.51 From this data, one can reconstruct the WDVV solution by using the relation

$$F^{\alpha\beta} = \eta^{\alpha\alpha'} \eta^{\beta\beta'} \frac{\partial^2 F}{\partial t^{\alpha'} \partial t^{\beta'}} = \frac{g^{\alpha\beta}}{\deg g^{\alpha\beta}}. \quad (4.240)$$

Theorem proved.  $\square$

## References

- [1] G. F. Almeida, *Differential Geometry of Orbit space of Extended Affine Jacobi Group  $A_1$*  preprint arxiv:1907.01436 (2020).
- [2] A. Belavin, V. Belavin, *Flat structures on the deformations of Gepner chiral rings*, J. High Energy Phys., 10, 128 (2016)

- 
- [3] A. Belavin, V. Belavin, *Frobenius manifolds, integrable hierarchies and minimal Liouville gravity*, JHEP 09 (2014) 151.
  - [4] A. Belavin, D. Gepner and Y. Kononov, *Flat coordinates for Saito Frobenius manifolds and String theory*, arXiv:1510.06970
  - [5] V. Belavin, *Unitary minimal Liouville gravity and Frobenius manifolds*, JHEP 07 (2014) 129.
  - [6] M. Bertola, *Frobenius manifold structure on orbit space of Jacobi groups; Parts I*, Diff. Geom. Appl. 13, (2000), 19-41.
  - [7] M. Bertola, *Frobenius manifold structure on orbit space of Jacobi groups; Parts II*, Diff. Geom. Appl. 13, (2000), 213-23.
  - [8] Bourbaki N., *Groupes et Algebres de Lie, Chapitres 4, 5 et 6*, Masson, Paris-New York-Barcelone-Milan-Mexico-Rio de Janeiro, 1981.
  - [9] B. Dubrovin, *Differential geometry of the space of orbits of a Coxeter group*, Preprint SISSA-29/93/FM (February 1993), hep-th/9303152.
  - [10] B. Dubrovin, *Geometry of 2D topological field theories*, in: M. Francaviglia and S. Greco, eds., Integrable Systems and Quantum Groups, Montecatini Terme, 1993, Lecture Notes in Math. 1620 (Springer, Berlin, 1996) 120–384.
  - [11] B. Dubrovin, *Hamiltonian perturbations of hyperbolic PDEs: from classification results to the properties of solutions*, New Trends in Mathematical Physics (Selected Contributions of the XVth Int. Congress on Mathematical Physics) ed V Sidoravicius (Netherlands: Springer) pp 231–76.
  - [12] B. Dubrovin, I. A. B. Strachan, Y. Zhang, D. Zuo, *Extended affine Weyl groups of BCD type, Frobenius manifolds and their Landau-Ginzburg superpotentials*, Advances in Mathematics, Volume 351, 2019, Pages 897-946.
  - [13] B. Dubrovin and Y. Zhang, *Extended affine Weyl groups and Frobenius manifolds*, Comp. Math. 111 (1998) 167–219.
  - [14] B. Dubrovin and Y. Zhang, *Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants*, preprint, available at <http://arxiv.org/math.DG/0108160>.
  - [15] M. Cutimanco, and V. Shramchenko, *Explicit examples of Hurwitz Frobenius manifolds in genus one*, J. Math. Phys. 61, 013501 (2020).
  - [16] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, (Birkhauser, Boston, 1985).
  - [17] E. V. Ferapontov, and M. V. Pavlov, L. Xue, *Second-order integrable Lagrangians and WDVV equations*, arXiv:2007.03768, 2020.
  - [18] C. Hertling, *Multiplication on the tangent bundle*, math.AG/9910116.

- 
- [19] M. Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, Comm. Math. Phys. 147 (1992), 1-23.
  - [20] Krichever I.M, *The  $\tau$ -function of the universal Whitham hierarchy, matrix models and topological field theories*, Comm. Pure Appl. Math. 47 (1994) 437 - 475.
  - [21] E. K. Morrison, and I. A. B. Strachan, *Modular Frobenius manifolds and their invariant flows*, International Mathematics Research Notices, Volume 2011, Issue 17, 2011, Pages 3957–3982.
  - [22] E. K. Morrison, and I. A. B. Strachan, *Polynomial modular Frobenius manifolds*, Phys. D 241 (2012), 2145–2155, arXiv:1110.4021.
  - [23] M. V. Pavlov, S. P. Tsarev., *Tri-Hamiltonian structures of the Egorov systems*, Funct. Anal. and Appl. 37:1 (2003), 32-45.
  - [24] S. Romano, *4-dimensional Frobenius manifolds and Painleve' VI*, preprint arxiv 1209.3959 (2012).
  - [25] S. Romano, *Frobenius structures on double Hurwitz spaces*, Int. Math. Res. Notices, doi:10.1093/imrn/rnt215 (2013), available at 1210.2312.
  - [26] Saito K., Yano T., and Sekeguchi J., *On a certain generator system of the ring of invariants of a finite reflection group*, Comm. in Algebra 8(4) (1980) 373 - 408.
  - [27] V. Shramchenko, *Deformations of Frobenius structures on Hurwitz spaces*, International Mathematics Research Notices, Volume 2005, Issue 6, 2005, Pages 339–387.
  - [28] V. Shramchenko, *Riemann-Hilbert problem associated to Frobenius manifold structures on Hurwitz spaces: irregular singularity*, Duke Math. J. 144, no. 1, 1-52 (2008)
  - [29] I. A. B. Strachan, *Frobenius submanifolds*, J. Geom. Phys. 38:3-4 (2001), 285-307.
  - [30] I. A. B. Strachan, *Frobenius manifolds: natural submanifolds and induced bi-hamiltonian structures*, Diff. Geom. Appl. 20:1 (2004), 67-99.
  - [31] E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis*, IV ed. (Cambridge University Press, Cambridge, 1980).
  - [32] K. Wirthmüller, *Root systems and Jacobi forms*, Comp. Math. 82 (1992) 293–354.
  - [33] D. Zuo, *Frobenius manifolds and a new class of extended affine Weyl groups of A-type*, Lett. Math. Phys. 110 (2020), no. 7.

E-mail address: galmeida@sissa.it.