# **REJECTION LEMMA AND ALMOST SPLIT SEQUENCES**

YURIY A. DROZD

To the memory of Volodya Kirichenko

ABSTRACT. We study the behaviour of almost split sequences and Auslander–Reiten quivers of an order under rejection of bijective modules as defined in [7]. In particular, we establish relations of stable categories and almost split sequences for an order A and the order A' obtained from A by such rejection. These results are specified for Gorenstein and Frobenius cases.

## Contents

1.	Introduction	1
2.	Orders, lattices and duality	2
3.	Bijective lattices and Gorenstein orders	7
4.	Bass orders	11
5.	Stable categories	14
6.	Almost split sequences	17
7.	Gorenstein and Frobenius cases	21
References		23

## 1. INTRODUCTION

Bijective modules and "rejection lemma" [7] play an important role in the theory of orders and lattices, as well as Gorenstein (that is, self-bijective) orders (see, for instance, [7, 8, 12, 13]). On the other hand, now the importance of almost split sequences and Auslander–Reiten quivers is doubtful. In this paper we consider the behaviour of almost split sequences and Auslander–Reiten quivers under rejection of bijective modules. Namely, in Section 2 we recall general facts on orders, lattices and duality. Our considerations are a bit more general, since the basic commutative ring is not necessarily discrete valuation ring, though in fact all main results from the "classical" theory, as in [5], remain valid. In Section 3 we introduce bijective lattices, rejection lemma and Gorenstein orders and establish some basic results about them.

<sup>1991</sup> Mathematics Subject Classification. 16G30, 16G70, 16H20.

Key words and phrases. Orders, lattices, almost split sequences, Auslander–Reiten transpose, Gorenstein orders, bijective lattices, rejection lemma.

#### YURIY A. DROZD

In particular, we find out which lattices become projective and injective after rejection (Theorem 3.8). Section 4 is devoted to *Bass orders*, i.e. such that all their overrings are Gorenstein. The main result here is Theorem 4.3, which is a substantial generalization of the criterion for an order to be Bass from [7]. In Section 5 we consider stable categories and relate the stable category of an order A to that of the order A' obtained by the rejection of a bijective module (Theorem 5.4). In Section 6 we study almost split sequences and find out how almost split sequences over A can be described in terms of A'-lattices (Proposition 6.3 and Theorem 6.4). Finally, in Section 7 we specify the preceding results for Gorenstein and Frobenius cases.

This paper is devoted to the memory of my friend, colleague and co-author Vladimir Kirichenko.

## 2. Orders, lattices and duality

We denote by mM the direct sum of m copies of a module M. The formulae  $M \supset N$  and  $N \subset M$  mean that N is a *proper subset* of M.

In what follows R is a complete local reduced noetherian ring of Krull dimension 1 with the maximal ideal  $\mathfrak{m}$ , the residue field  $\Bbbk = R/\mathfrak{m}$  and the total ring of fractions K. It follows from [4] that the ring A is Cohen-Macaulay. We denote by *R*-mod the category of finitely generated *R*-modules and by R-lat its full subcategory of R-lattices, that is of torsion free R-modules or such modules M that the natural map  $M \to K \otimes_R M$  is an embedding. We write KM instead of  $K \otimes_R M$  and identify M with  $1 \otimes M \subseteq KM$ . In this case *R*-lattices coincide with maximal Cohen–Macaulay *R*-modules. As R is complete, it has a *canonical module* [4, Corollary 3.3.8], that is an *R*-lattice  $\omega_R$  such that inj.dim<sub>R</sub>  $\omega_R = 1$  and  $\operatorname{Ext}^1_{\mathbf{R}}(\mathbb{k}, \omega_R) = \mathbb{k}$ . The functor  $D: M \mapsto \operatorname{Hom}_{R}(M, \omega_{R})$  is an *exact duality* on the category R-lat [4, Theorem 3.3.10]. It means that if  $0 \to N \xrightarrow{\alpha} M \xrightarrow{\beta} L \to 0$  is an exact sequence of lattices, the sequence  $0 \to DL \xrightarrow{D\beta} DM \xrightarrow{D\alpha} DN \to 0$  is also exact, and the natural map  $M \to DDM$  is an isomorphism. As  $\operatorname{End}_R(\omega_R) \simeq \operatorname{End}_R R \simeq R$ and  $\operatorname{End}_K KM \simeq K \operatorname{End}_R M$  for every Cohen-Macaulay module M, we have that  $K\omega_R \simeq K$  and we identify  $\omega_R$  with its image in K. Note also that K is a direct product of fields  $K = \prod_{i=1}^{s} K_i$ , where  $K_i$  is the field of fractions of the ring  $R/\mathfrak{p}_i$  and  $\mathfrak{p}_i$  runs through minimal prime ideals of R.

An *R*-order is, by definition, a semiprime *R*-algebra *A* which is an *R*-lattice. Recall that semiprime means that *A* has no nilpotent ideals. Then *KA* is a semisimple *K*-algebra and they say that *A* is an *R*-order in *KA*. We denote by Z(A) the center of *A* and call *A* central if the natural map  $R \to Z(A)$  is an isomorphism. If *A* is connected, i.e. does not decomposes as a ring, its center is local and vice versa. We denote by *A*-mod the category of finitely generated *R*-modules and by *A*-lat its full subcategory of *A*-lattices, i.e. (left) *A*-modules which are *R*-lattices. The restriction of the duality functor *D* onto *A*-lat gives an exact duality of *A*-lat onto  $A^{\text{op}}$ -lat, which

we consider as the category of right A-lattices. Set  $\omega_A = \operatorname{Hom}_R(A, \omega_R)$ . It is an A-bimodule and, for any A-lattice M (left or right), its dual DM is identified with  $\operatorname{Hom}_A(M, \omega_A)$ . We say finite module instead of module of finite length and denote by  $\ell_A(M)$  the length of such module. We call the width of a lattice M and denote it by  $\operatorname{wd}_A(M)$  the length  $\ell_{KA}(KM)$ . One easily sees that  $\operatorname{wd}_A(M)$  is the maximal integer m such that M contains a direct sum of m nonzero submodules, or, equivalently, contains a chain of submodules  $M = M_0 \supset M_1 \supset \ldots \supset M_m$  such that  $M_i/M_{i+1}$  is a lattice for  $0 \leq i < m$ . A lattice M of width 1 is called L-irreducible.<sup>1</sup> If the ring R is fixed, we often say order instead of R-order.

As the ring R is complete, any finite R-algebra (i.e. finitely generated as R-module) is semiperfect [15]. Therefore, the category of finitely generated modules over a finite R-algebra A is Krull–Schmidt. In particular, any finitely generated projective A-module is isomorphic to a direct summand of A and there is a one-to-one correspondence between indecomposable finitely generated projective A-modules (called *principal* A-modules) and simple A-modules, which maps a principle A-module P to  $P/\mathfrak{r}P$ , where  $\mathfrak{r} = \operatorname{rad} A$ . For every finitely generated A-module M there is an epimorphism  $\pi : P \to M$  with projective P and Ker  $\pi \subseteq \mathfrak{r}P$ . The module P is unique up to isomorphism. It is called the *projective cover* of M and denoted by  $P_A(M)$ . Sometimes the epimorphism  $\pi$  is also called a projective cover of M, though it is only defined up to an automorphism of P. Obviously,  $\pi$  induces an isomorphism  $P/\mathfrak{r}P \xrightarrow{\sim} M/\mathfrak{r}M$ .

An overring of an R-order A is an R-order A' such that  $A \subset A' \subset KA$ . Then A'/A is a finite module and A-lat is a full subcategory of A'-lat. An order is said to be maximal if it has no overrings. An overring of A which is a maximal order is called a maximal overring of A. An overmodule of an A-lattice M is an A-lattice M' such that  $M \subset M' \subset KM$ . If A' is an overring of A and M is a A-lattice considered as a submodule in KM, then the A'-module A'M is defined and is an overmodule of M.

The following fact seems to be well-known. If R is a discrete valuation ring, it is proved in [5]. The general case easily reduces to this one, though we have not found any source in the literature.

**Proposition 2.1.** Every *R*-order *A* has a maximal overring. The center of a maximal order is a product of discrete valuation rings. A connected maximal order has a unique indecomposable lattice (up to isomorphism). Conversely, if an order *A* has a unique indecomposable lattice, it is maximal.

*Proof.* We may suppose A connected. Then its center Z(A) is also local and complete. Every overring of A is a Z(A)-order, so we may suppose that Z(A) = R. Then Z(KA) = K. Let S be the integral closure of R in K. As R is local and complete, it is an *excellent ring* [17]. In particular, S is a finitely generated R-module. As it is integrally closed, it is a direct product

<sup>&</sup>lt;sup>1</sup>One often call such a lattice *irreducible*, but we will use this word in another situation.

of discrete valuation rings. The ring SA is an S-order and an overring of A. It splits into a direct product of orders whose centers are discrete valuation rings. Then [5, Theorem 26.5] inplies that SA, hence also A, has a maximal overring A' and Z(A') = S. Now the remaining assertions also follow from [5].

As the algebra KA is semisimple, every finitely generated KA-module embeds into a finitely generated free module. It easily implies that any Alattice M embeds into a free A-module. Thus A-lattices are just submodules of free modules (*torsionless modules* in the sense of Bass [3]).

**Proposition 2.2.** Let  $I \in A$ -lat. The following conditions are equivalent:

- (1)  $\operatorname{inj.dim}_A I = 1$ .
- (2)  $\operatorname{Ext}^{1}_{A}(M, I) = 0$  for any  $M \in A$ -lat.
- (3)  $\operatorname{Ext}_{A}^{i}(M, I) = 0$  for any  $M \in A$ -lat and any  $i \geq 1$ .
- (4) Any exact sequence  $0 \to I \to N \to M \to 0$ , where  $M \in A$ -lat splits.
- (5)  $I \simeq DP$ , where P is a finitely generated projective  $A^{\text{op}}$ -module.
- (6) I is a direct summand of  $m\omega_A$  for some m.

We call a lattice I satisfying these conditions L-injective. If an L-injective lattice is indecomposable, we call it *coprincipal*.

*Proof.*  $(3) \Rightarrow (2)$  and  $(2) \Leftrightarrow (4)$  are obvious.

 $(2) \Rightarrow (3)$  since in a projective resolution

$$\cdots \to P_n \xrightarrow{d_n} P_{n-1} \xrightarrow{d_{n-1}} \cdots \to P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \to M \to 0$$

of M all modules  $M_i = \operatorname{Im} d_i$  are lattices and  $\operatorname{Ext}^i_A(M, I) \simeq \operatorname{Ext}^1_A(M_{i-1}, I)$ for i > 1.

 $(4) \Rightarrow (5)$ . By duality, (4) means that every exact sequence  $0 \rightarrow M \rightarrow N \rightarrow DI \rightarrow 0$  splits. As there is such a sequence with projective N, it implies that P = DI is projective and  $I \simeq DP$ .

 $(5) \Rightarrow (6)$ . Since a projective module P is a direct summand of a free module mA, the module I = DP is a direct summand of  $D(mA) = m\omega_A$ .

 $(6) \Rightarrow (2)$ . Let M be an A-lattice. Consider an exact sequence  $0 \rightarrow N \rightarrow P \rightarrow M \rightarrow 0$  with projective P. As all these modules are lattices, the induced sequence

$$0 \to \operatorname{Hom}_A(M, \omega_A) \to \operatorname{Hom}_A(P, \omega_A) \to \operatorname{Hom}_A(N, \omega_A) \to 0$$

is also exact, whence  $\operatorname{Ext}_{A}^{1}(M, \omega_{A}) = 0$ . Therefore, the same holds for  $m\omega_{A}$  and for its direct summand I.

 $(3) \Leftrightarrow (1)$ . It is known that

inj.dim 
$$I = \sup \{i \mid \operatorname{Ext}_{A}^{i}(A/L, I) \neq 0 \text{ for some left ideal } L\} =$$
  
=  $\sup \{i \mid \operatorname{Ext}_{A}^{i-1}(L, I) \neq 0 \text{ for some left ideal } L\}.$ 

As any ideal is a lattice,  $(3) \Rightarrow (1)$ . On the contrary, if (1) holds and M is a lattice, embed it into a projective module P. Then  $\operatorname{Ext}_{A}^{i}(M, I) = \operatorname{Ext}_{A}^{i+1}(P/M, I) = 0$  if  $i \ge 1$ , so (3) holds.  $\Box$ 

The category A-lat becomes an exact category [14] if we consider as exact pairs (or conflations) usual short exact sequences, i.e. all triples  $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ , where  $\alpha = \text{Ker }\beta$  and  $\beta = \text{Cok }\alpha$ . Therefore, deflations are epimorphisms and inflations are monomorphisms with torsion free cokernels (we will often use this term). This exact category contains enough projectives and injectives. Namely, projectives are just usual finitely generated projective A-modules, while injectives are L-injective lattices. To obtain a conflation  $M \to I \to N$  with L-injective M, one just has to dualize an exact sequence  $0 \to L \to P \to DM \to 0$  with projective P.

For lattices M, N we write  $M \searrow N$  (respectively,  $N \nearrow M$ ) if there is an epimorphism  $rM \to N$  (respectively, an inflation  $N \to rM$ ) for some r. For instance,  $A \searrow M$  and, dually,  $M \nearrow \omega_A$  for any lattice M. We write  $N \in M$  if N is a direct summand of rM for some r and  $M \bowtie N$  if both  $M \in N$  and  $N \in M$ . Recall that CM(A) is a Krull–Schmidt category. So, if N is indecomposable,  $N \in M$  means that N is a direct summand of Mand  $M \bowtie N$  means that M and N have the same set of indecomposable direct summands. Note that the relations  $\searrow$ ,  $\nearrow$  and  $\in$  are transitive.

**Definition 2.3.** Let M be an A-lattices,  $E = \operatorname{End}_A M$  and  $O(M) = \operatorname{End}_E M$ . If the natural map  $A \to O(M)$  is an isomorphism, we say that M is a *strict* A-lattice. Obviously, then M is faithful.

Certainly, O(M) is an overring of  $A / \operatorname{Ann}_A M$ . Due to the Burnside density theorem [9, Theorem 2.6.7], then O(M) can be identified with the subset  $\{a \in KA / \operatorname{Ann} KM \mid aM \subseteq M\}$ . In particular, a faithful A-lattice M is strict if and only if  $\{a \in KA \mid aM \subseteq M\} = A$ . If M and N are faithful and  $M \searrow N$  or  $N \nearrow M$ , then  $O(N) \supseteq O(M)$ .

**Proposition 2.4.** For every A-lattice M there is an exact sequence

$$(2.1) 0 \to O(M) \to nM \to mM$$

for some m, n. In particular, M is strict if and only if there is an exact sequence

(2.2) 
$$0 \to A \xrightarrow{\alpha} nM \xrightarrow{\beta} mM,$$

that is  $A \nearrow M$ .

Proof. If  $E = \operatorname{End}_A M$ , there is an exact sequence of E-modules  $mE \to nE \to M \to 0$ . Applying the functor  $\operatorname{Hom}_E(-, M)$ , we obtain the exact sequence (2.1). If M is strict, it coincides with (2.2). On the contrary, let an exact sequence (2.2) exists. Then M is faithful. We identify A with  $N = \{u \in nM \mid \beta(u) = 0\}$  If  $a \in O(M)$ , it is clear that  $aN \subseteq N$ , whence  $a \in A$ .

**Corollary 2.5.** An A-lattice M is strict if and only if there is an exact sequence

$$(2.3) mM \to nM \to \omega_A \to 0,$$

that is  $M \searrow \omega_A$ .

We will also use another kind of duality analogous to the Matlis duality [16].

**Theorem 2.6.** Let  $T_R = K\omega_R/\omega_R$ . Denote  $\hat{M} = \text{Hom}_R(M, T_R)$ . The functor  $M \mapsto \hat{M}$  induces an exact duality between the categories of noetherian and artinian *R*-modules.

Proof. Step 1. We denote by  $\gamma_M$  the natural map  $M \to \hat{M}$ . Any KRmodule V is an injective R-module and  $\operatorname{Hom}_R(V, M) = 0 = \operatorname{Hom}_R(L, V)$ for any noetherian M and any torsion R-module L. As inj.dim<sub>R</sub>  $\omega_R = 1$ ,  $T_R$  is also an injective R-module. So the functor  $M \mapsto \hat{M}$  is exact. If an R-module L is torsion, apply the functor  $\operatorname{Hom}_R(L, -)$  to the exact sequence  $0 \to \omega_R \to K\omega_R \to T_R \to 0$ . We obtain that  $\hat{L} \simeq \operatorname{Ext}^1_R(L, \omega_R)$ . In particular,  $\hat{R} = \hat{T}_R \simeq \operatorname{Ext}^1_R(T_R, \omega_R)$ . Apply to the same exact sequence the functor  $\operatorname{Hom}_R(-, \omega_R)$ . It gives  $R = \operatorname{Hom}_R(\omega_R, \omega_R) \simeq \operatorname{Ext}^1_R(T_R, \omega_R) = \hat{T}_R$ . Thus  $\gamma_R$  and  $\gamma_{T_R}$  are isomorphisms. Now the usual observation using a free presentation  $mR \to nR \to M \to 0$  shows that  $\gamma_M$  is an isomorphism for any noetherian R-module M.

Step 2. Show that the module  $N = \hat{M}$  is artinian if M is noetherian. Indeed, if  $N_1 \subset N$ , this embedding induces a surjection  $M = \hat{N} \xrightarrow{\alpha} \hat{N}_1$  with  $\operatorname{Ker} \alpha \simeq \widehat{N/N_1}$ . Moreover, if  $N_2 \subset N_1$ , we have surjections  $M \xrightarrow{\alpha} \hat{N}_1 \xrightarrow{\beta} \hat{N}_2$  such that  $\operatorname{Ker} \beta \alpha \supset \operatorname{Ker} \alpha$ . Thus any descending chain of submodules in  $\hat{M}$  gives an ascending chain of submodules in M. Therefore, there are no infinite descending chains in  $\hat{M}$ . In particular, the module  $T_R = \hat{R}$  is artinian.

Step 3. Let now the module N be artinian. It contains a simple submodule U. As  $\operatorname{Hom}_R(U, T_R) \neq 0$  and  $T_R$  is injective, there is a nonzero homomorphism  $\alpha_0 : N \to T_R$ . As  $\operatorname{Ker} \alpha_0$  is also artinian, there is a non-zero homomorphism  $\operatorname{Ker} \alpha_0 \to T_R$ , which extends to a homomorphism  $\alpha' : N \to T_R$ . Let  $\alpha_1 = \begin{pmatrix} \alpha_0 \\ \alpha' \end{pmatrix} : N \to 2T_R$ . Then  $\operatorname{Ker} \alpha_1 \subset \operatorname{Ker} \alpha_0$ . Iterating this procedure, we obtain homomorphisms  $\alpha_k : N \to kT_R$  such that  $\operatorname{Ker} \alpha_{k+1} \subset \operatorname{Ker} \alpha_k$  if  $\operatorname{Ker} \alpha_k \neq 0$ . As N is artinian, there is an embedding  $\beta : N \to mT_R$  for some m. As  $\operatorname{Cok} \beta$  is also artinian, we have an exact sequence  $0 \to N \to mT_R \to nT_R$ . Since the map  $\gamma_{T_R}$  is an isomorphism, it implies that  $\gamma_N$  is also an isomorphism. The observation analogous to Step 2 shows that  $\hat{N}$  is noetherian, which accomplishes the proof.  $\Box$ 

Obviously, if we restrict this duality to A-modules, we obtain a duality between the categories of left (right) noetherian and right (left) artinian Amodules. One easily sees that the category of lattices is then mapped to the category of artinian modules without finite quotients.

The duality  $M \mapsto \hat{M}$  is closely related to the duality D.

**Proposition 2.7.** Let  $0 \to M \xrightarrow{\alpha} N \to L \to 0$  be an exact sequence of A-modules, where M, N are lattices and L is finite. Then there is an exact

 $\mathbf{6}$ 

sequence  $0 \to DN \xrightarrow{D\alpha} DM \to \hat{L} \to 0$ . In particular, if M is a maximal submodule in N, DN is a minimal overmodule of DM and vice versa.

*Proof.* We have already seen in Step 1 of the previous proof that  $\hat{L} \simeq \operatorname{Ext}_{A}^{1}(L,\omega_{A})$ . Note also that  $\operatorname{Hom}_{A}(L,\omega_{A}) = 0$ . So we obtain the result if we apply to the given exact sequence the functor  $\operatorname{Hom}_{A}(_{-},\omega_{A})$ .

Let M be an A-lattice,  $\mathbf{r} = \operatorname{rad} A$ . As  $(DM)\mathbf{r}$  is the intersection of maximal submodules of DM, its dual  $M^{\mathfrak{r}} = D((DM)\mathbf{r})$  is the sum of all minimal overmodules of M. If  $\pi : P \xrightarrow{\pi} DM$  is a projective cover of DM, its dual  $D\pi : M \to DP$  is an inflation  $\iota : M \to I$  such that I is L-injective and  $\iota$  induces an isomorphism  $I^{\mathfrak{r}}/I \to M^{\mathfrak{r}}/M$ . We call it (and sometimes the map  $\iota$ ) the *L-injective envelop* of M. We also define iterated overmodules  $M^{\mathfrak{r}*k}$  setting  $M^{\mathfrak{r}*1} = M^{\mathfrak{r}}$  and  $M^{\mathfrak{r}*(k+1)} = (M^{\mathfrak{r}*k})^{\mathfrak{r}}$ . Obviously,  $M^{\mathfrak{r}*k} =$  $D((DM)\mathfrak{r}^k)$ . As a principal A-module P has one maximal submodule  $\mathfrak{r}P$ , a coprincipal A-lattice I has one minimal overmodule  $I^{\mathfrak{r}}$ .

## 3. BIJECTIVE LATTICES AND GORENSTEIN ORDERS

Let A be an R-order,  $\mathfrak{r} = \operatorname{rad} A$ . In this section we always suppose that A is connected.

**Definition 3.1.** An A-lattice B is called *bijective* [7] if it both projective and L-injective.

The main property of bijective lattices is the so called *rejection lemma* (cf. [7, Lemma 2.9]).

**Lemma 3.2.** Let B be a bijective A-lattice. Either there is a unique overring A' such that every A-lattice M is isomorphic to  $B' \oplus M'$ , where M' is an A'-lattice and  $B' \oplus B$ , or A is hereditary and  $A \oplus B$  (then  $M \oplus B$  for any A-lattice M).

We say that A' is obtained from A by rejection of B and denote it by  $A^{-}(B)$ . Obviously, if B is indecomposable,  $A^{-}(B)$  is a minimal proper overring of A.

Remark 3.3. By duality, DB is a bijective right A-lattice and every right A-lattice N is isomorphic to  $B' \oplus N'$ , where  $B' \oplus B$  and N' is a right A'-lattice.

Proof. If  $M \in B$ , then M is projective. Thus if  $M \in B$  for every A-lattice M, A is hereditary. So we can suppose that there are lattices M such that  $M \notin B$ . Certainly, then there are faithful lattices with this property. If M is a strict A-lattice, then  $A \nearrow M$ . As B is projective, also  $B \nearrow M$ , whence  $B \in M$ , since B is L-injective. Let  $A' = \bigcap_M O(M)$ , where M runs through all faithful A-lattices that does not have a direct summand  $B' \in B$ . There is a finite set of lattices  $M_1, M_2, \ldots, M_n$  such that A' = O(N), where  $N = \bigoplus_{i=1}^n M_i$ . If N is strict,  $B \in N$ , which is impossible. Hence  $A' \supset A$  and every faithful A-lattice M without direct summands  $B' \in B$  is an

A'-lattice. Let M be any A-lattice that has no direct summands  $B' \in B$ and  $U_1, U_2, \ldots, U_s$  be all non-isomorphic simple KA-modules. If M is not faithful, at least one of them, say  $U_1$ , does not occur as a direct summand of KM. We claim that there is a A-lattice  $L \subset U_1$  such that  $L \notin B$ . Then we replace M by  $M \oplus L$  and, continuing this procedure, obtain a faithful A-lattice M' without direct summands  $B' \in B$  such that M is its direct summand. Therefore, M', hence also M is an A'-lattice.

Suppose that  $L \in B$  for every A-lattice  $L \subset U_1$ . Let C be a simple component of KA such that  $U_1$  is a C-module,  $A_1$  be the projection of A onto C. If M is any  $A_1$ -lattice, it has a chain of submodules whose factors are submodules of  $U_1$ . Hence it is projective, so  $A_1$  is hereditary and is a direct factor of A. It implies that  $A_1 = A$  and KA = C is a simple K-algebra, so  $M \in B$  for every A-lattice.

To describe the structure of  $A^{-}(B)$ , we need some simple lemmas.

- **Lemma 3.4.** (1) Let P be a principal A-module. If all modules  $\mathbf{r}^i P$  are indecomposable and projective, A is hereditary and every indecomposable A-lattice is isomorphic to some  $\mathbf{r}^i P$ .
  - (2) Let I be a coprincipal A-lattice. If all modules  $I^{\mathfrak{r}*i}$  are indecomposable and L-injective, A is hereditary and every indecomposable A-lattice is isomorphic to some  $I^{\mathfrak{r}*i}$ .
  - (3) Let P be a principal A-module. If  $\mathfrak{r}P \simeq P$ , the order A is maximal and P is a unique indecomposable A-lattice.
  - (4) Let I be a coprincipal A-lattice. If  $I^{\mathfrak{r}} \simeq I$ , the order A is maximal. and I is a unique indecomposable A-lattice

*Proof.* (1) The conditions imply that  $\mathfrak{r}^{i+1}P$  is a unique maximal submodule of  $\mathfrak{r}^i P$ . Therefore, any submodule of P coincides with some  $\mathfrak{r}^i P$ , hence is projective and indecomposable. Then KP is a simple KA-module, so there is a simple component C of the algebra KA such that KP is a KAmodule. If V is any C-module, it is a multiple of KP. Hence if  $M \subset V$  is a lattice, it has a chain of submodules whose factors are submodules of KP. It implies that M is projective. In particular, the projection  $A_1$  of A onto Cis projective, so is a direct summand of A as A-module. It obviously implies that  $A_1$  is a direct factor of A, so  $A = A_1$ .

(3) If  $\mathfrak{r}P \simeq P$ , then  $\mathfrak{r}^k P \simeq P$  for all k, so all of them are principal. Just as in (1), it implies that  $A = A_1$  and P is a unique indecomposable A-lattice. Therefore, A is maximal.

(4) is dual to (3).

**Lemma 3.5.** We suppose that A is non-hereditary. Let B be an indecomposable bijective A-lattice,  $A' = A^{-}(B)$ . Then  $B^{\mathfrak{r}} \not\simeq B$  as well as  $\mathfrak{r}B \not\simeq B$ ,  $B^{\mathfrak{r}}$  is projective and  $\mathfrak{r}B$  is L-injective as A'-lattice.

*Proof.*  $B^{\mathfrak{r}} \not\simeq B$  as well as  $\mathfrak{r}B \not\simeq B$  by Lemma 3.6. Therefore, they are A'-lattices and  $A'B = B^{\mathfrak{r}}$ . The principal A-module B is a direct summand of

<sup>(2)</sup> is dual to (1).

A, so  $A \simeq B \oplus M$  for some M. Then  $A' = A'A \simeq A'B \oplus A'M = B^{\mathfrak{r}} \oplus A'M$ , hence  $B^{\mathfrak{r}}$  is projective over A'. By duality,  $\mathfrak{r}B$  is L-injective over A'. 

- Lemma 3.6. (1) Let P be a principal A-module, M be its minimal overmodule. Then M is either indecomposable or splits as  $M_1 \oplus M_2$ , where  $M_1, M_2$  are indecomposable. In the latter case,  $\mathfrak{r}P = \mathfrak{r}M_1 \oplus$  $\mathfrak{r}M_2$  and neither  $M_1$  nor  $M_2$  is projective.
  - (2) Let I be a coprincipal A-lattice, M be its maximal submodule. Then M is either indecomposable or splits as  $M_1 \oplus M_2$ , where  $M_1, M_2$  are indecomposable. In the latter case,  $I^{\mathfrak{r}} = M_1^{\mathfrak{r}} \oplus M_2^{\mathfrak{r}}$  and neither  $M_1$ nor  $M_2$  is L-injective.
  - (3) Let B be an indecomposable bijective A-lattice. Then its maximal submodule and minimal overmodule decompose simultaneously. Moreover, if  $\mathfrak{r}B$  is L-injective,  $B^{\mathfrak{r}}$  is projective and vice versa.

*Proof.* (1) As  $P \supseteq \mathfrak{r}M \supseteq \mathfrak{r}P$ ,  $\ell_A(M/\mathfrak{r}M) \leq 2$ , hence M is either indecomposable or splits as  $M_1 \oplus M_2$ , where  $M_1, M_2$  are indecomposable. In the latter case,  $\ell_A(M_1/\mathfrak{r}M_1) = 1$ , so  $N = \mathfrak{r}M_1 \oplus M_2 \neq P$  is a maximal submodule in  $M, N \cap P = \mathfrak{r}P$  and  $M_1/\mathfrak{r}M_1 \simeq M/N \simeq P/\mathfrak{r}P$ . Since  $M_1 \not\simeq P$ , it cannot be projective. The same applies to  $M_2$ . Moreover, in this case  $\ell_A(M/\mathfrak{r}M) = 2$ , whence  $\mathfrak{r}P = \mathfrak{r}M = \mathfrak{r}M_1 \oplus \mathfrak{r}M_2$ .

(2) follows by duality.

(3) By (1) and (2), if  $B^{\mathfrak{r}}$  is indecomposable, so is  $\mathfrak{r}B$  and vice versa. Suppose that  $\mathfrak{r}B$  is L-injective. Then it is indecomposable, hence  $B = (\mathfrak{r}B)^{\mathfrak{r}}$ is a unique minimal overmodule of  $\mathfrak{r}B$ . Then B is also a unique maximal submodule of  $B^{\mathfrak{r}}$ . Therefore, there is an epimorphism  $\pi: P \to B^{\mathfrak{r}}$ , where P is principal. If  $P \simeq B$ ,  $\pi$  is an isomorphism. If  $P \not\simeq B$ , it is an A'module. By Lemma 3.5,  $B^{\mathfrak{r}}$  is projective as A'-module, hence  $\pi$  splits, so is an isomorphism. In both cases  $B^{\mathfrak{r}}$  is projective over A. The converse follows by duality. 

**Definition 3.7.** Let *B* be a bijective *B*-lattice.

- (1) A *B*-link is a set of indecomposable lattices  $\{B_1, B_2, \ldots, B_l\}$  such that
  - $B_i \in B$  for all  $i = 1, \ldots, l$ ,
  - $B_i = \mathfrak{r} B_{i-1}$  for  $i = 2, \ldots, l$  (equivalently,  $B_{i-1} = B_i^{\mathfrak{r}}$ ),
  - $\mathfrak{r}B_l \notin B$  and  $B_1^{\mathfrak{r}} \notin B$ .
- (2) For an indecomposable A-lattice M define  $M^{\pm,B}$  as follows:
  - If  $M \notin B$ , then  $M^{\pm,B} = P$ ;
  - If  $M \in \{B_1, B_2, \dots, B_l\}$ , where  $\{B_1, B_2, \dots, B_l\}$  is a *B*-link, then  $M^{+,B} = B_1^{\mathfrak{r}}$  and  $M^{-,B} = \mathfrak{r}B_l$ . We denote by  $\iota_M^B$  the embedding  $M^{-,B} \to M^{+,B}$ .

**Theorem 3.8.** Suppose that the order A is non-hereditary. Let B be a bijective A-lattice,  $A' = A^{-}(B)$ . If  $A = \bigoplus_{i=1}^{n} P_i$ , where  $P_i$  are indecomposable,

then  $A' = \bigoplus_{i=1}^{n} P_i^{+,B}$ . In particular, all modules  $P_i^{+,B}$  are projective as A'-modules and every principal A'-module is isomorphic to a direct summand of some  $P_i^{+,B}$ .

Remark 3.9. Dually, if  $\omega_A = \bigoplus_{i=1}^n I_i$ , where  $I_i$  are indecomposable, then  $\omega_{A'} = \bigoplus_{i=1}^n I_i^{-,B}$ . In particular, all lattices  $I_i^{-,B}$  are L-injective as A'-lattices.

*Proof.* We write  $P'_i$  instead of  $P^{+,B}_i$ . Obviously, we may suppose that  $B = \bigoplus_{j=1}^{m} B_i$ , where all  $B_i$  are indecomposable and non-isomorphic. We use induction by m. Let m = 1, i.e. B is indecomposable. By Lemma 3.5,  $B^{\mathsf{r}} \not\simeq B$ , so  $B' = B^{\mathsf{r}}$  is an A'-lattice and A'B = B'. If P is principal and  $P \not\simeq B$ , then P' = P and it is an A'-lattice, so A'P = P. Therefore,  $A' = A'A = \bigoplus_{i=1}^{n} P'_i$ .

Suppose that the theorem holds for m-1 summands. If  $B_i^{\mathfrak{r}} \in B$  for all i, then  $B_1^{\mathfrak{r}*k} \in B$  for all k, hence A is hereditary by Lemma 3.4, which is impossible. Thus we can suppose that  $B_1^{\mathfrak{r}} \notin B$ . Set  $A_1 = A^-(B_1)$ ,  $\mathfrak{r}_1 = \operatorname{rad} A_1$ . Then  $A^-(B) = A_1^-(B(1))$ , where  $B(1) = \bigoplus_{i=2}^m B_i$ . If  $\mathfrak{r} B_1 = B_2 \in B$ , then  $B_1$  is its unique minimal overmodule. As  $B_1^{\mathfrak{r}}$  is a unique minimal overmodule of  $B_1$  and  $B_1$  is not an  $A_1$ -module, it implies that  $B_2^{\mathfrak{r}_1} = B_1^{\mathfrak{r}}$ , when we consider  $A_1$ -modules. Therefore,  $M^{+,B} = M^{+,B(1)}$  for every  $A_1$ -lattice M. If  $P_i \simeq B_1$  for  $i \leq r$  and  $P_i \neq B_1$  for i > r, then  $A_1 = A^-(B_1) = (\bigoplus_{i=1}^r P_i') \oplus (\bigoplus_{i=r+1}^n P_i)$ . Moreover,  $P_i'^{+,B(1)} = P_i'$  for  $i \leq r$  and  $P_i^{+,B(1)} = P_i'$  for i > r. By the induction hypothesis,  $A^-(B) = \bigoplus_{i=1}^n P_i'$ .

Now we introduce the class of orders which is the main in this paper. We follow the paper [10]. The following result is an obvious corollary of Propositions 2.2 and 2.4 and Corollary 2.5.

**Proposition 3.10.** Let A be an R-order. The following conditions are equivalent:

- (1) A is L-injective as left A-lattice.
- (2) A is L-injective as right A-lattice.
- (3)  $A \in M$  for every strict A-lattice M.
- (4)  $\omega_A \in M$  for every strict A-lattice M.
- (5) If M is a strict A-lattice, then  $M \searrow N$  for any A-lattice N.
- (6) If M is a strict A-lattice, then  $N \nearrow M$  for any A-lattice N.
- (7) Every projective (finitely generated) A-module is L-injective.
- (8) Every L-injective A-lattice is projective.

If these conditions hold, A is called a *Gorenstein order* [7].

Obviously, every hereditary order A is Gorenstein. If A is not hereditary, we denote by  $A^-$  the order  $A^-(A)$ . It is obtained by rejection of all projective modules. In the Gorenstein case Theorem 3.8 can be essentially simplified using the following result.

**Lemma 3.11.** Let A be a non-hereditary Gorenstein order, B be an indecomposable bijective A-lattice. Then neither  $B^{\mathfrak{r}}$  nor  $\mathfrak{r}B$  is projective (or, the same, L-injective).

Proof. Suppose that  $P = B^{\mathfrak{r}}$  is projective, hence bijective. By Lemma 3.6, it is indecomposable, hence  $\mathfrak{r}P = B$ . Let  $N = P^{\mathfrak{r}}$ . Then  $\mathfrak{r}N \supseteq \mathfrak{r}P = B$ . If  $\mathfrak{r}N = B$ , then  $B^{\mathfrak{r}} \supseteq N$ , which is impossible. Therefore,  $\mathfrak{r}N = P$ , so  $N/\mathfrak{r}N$  is a simple module. Then there is a surjection  $P' \to N$ , where P' is a principal module, hence a surjection  $\mathfrak{r}P' \to P$ . Thus P is a direct summand of  $\mathfrak{r}P'$ . By Lemma 3.6,  $\mathfrak{r}P' \simeq P$ , whence  $P' \simeq N$ , so  $N = B^{\mathfrak{r}*2}$  is also bijective. Going on, we see that all lattices  $B^{\mathfrak{r}*k}$  are bijective. By Lemma 3.4, A is hereditary, which is impossible, so  $B^{\mathfrak{r}}$  cannot be projective. The assertion about  $\mathfrak{r}B$  is just dual.

**Corollary 3.12.** Let A be a non-hereditary Gorenstein order,  $A = \bigoplus_{i=1}^{n} P_i$ , where  $P_i$  are indecomposable,  $P'_i = P^{\mathfrak{r}}_i$  and B be a bijective A-lattice. Suppose that  $P_i \in B$  for  $i \leq k$  and  $P_i \notin B$  for i > k. Then  $A^-(B) = (\bigoplus_{i=1}^{k} P'_i) \oplus (\bigoplus_{i=k+1}^{n} P_i)$ . Moreover,  $\mathfrak{r}P_i$  and  $P^{\mathfrak{r}}_i$  are  $A^-(B)$ -lattices for all i. In particular,  $A^- = \bigoplus_{i=1}^{k} P'_i$ ,  $\mathfrak{r}$  and  $A^{\mathfrak{r}}$  are  $A^-$ -lattices (both left and right).

*Proof.* Immediately follows from Theorem 3.8 and Lemma 3.11.  $\Box$ 

For Gorenstein orders the rejection lemma 3.2 can be inverted.

**Proposition 3.13.** If A is Gorenstein, every minimal overring of A is of the form  $A^{-}(B)$ , where B is an indecomposable bijective A-lattice.

*Proof.* If each indecomposable projective (or, the same, bijective) A-lattice is actually an A'-lattice, then A' = A. Therefore, there is an indecomposable bijective A-lattice B which is not an A'-lattice. Then  $A' \supseteq A^-(B)$ . As A' is minimal,  $A' = A^-(B)$ .

## 4. Bass orders

Recall that an order A is called a *Bass order* [10] if all its overrings (including A itself) are Gorenstein. The results of the preceding section imply the following criterion (cf. [7, Theorem 3.1]).

**Proposition 4.1.** The following conditions are equivalent:

- (1) A is a Bass order.
- (2)  $M \searrow O(M)$  for every A-lattice M.
- (3) For any two A-lattices M, N, if  $M \searrow N$ , then  $N \nearrow M$ .

(4) For any two A-lattices M, N, if  $N \nearrow M$ , then  $M \searrow N$ .

Therefore, any order that is Morita equivalent to a Bass order is also Bass.

**Example 4.2.** (1) Every hereditary order is a Bass order.

(2) If every ideal of A has 2 generators, A is a Bass order. It follows from [18] in the case when R is a discrete valuation ring, but the proof in the general case is the same.

#### YURIY A. DROZD

(3) Let  $\Delta$  be a maximal order in a skewfield,  $\mathfrak{d} = \operatorname{rad} \Delta$ ,  $B(m, \Delta)$  be the subring of Mat $(2, \Delta)$  consisting of such matrices  $(a_{ij})$  that  $a_{12} \in \mathfrak{d}^m$ . It is also a Bass order (hereditary for m = 1). We write symbolically  $B(k, \Delta) = \begin{pmatrix} \Delta & \mathfrak{d}^k \\ \Delta & \Delta \end{pmatrix}$ .

Actually, it is proved in [10] that every Bass order is either hereditary, or Morita equivalent to a local order such that every its ideal has 2 generators, or Morita equivalent to some  $B(\Delta, m)$ . We will obtain this description as a corollary of the following result, which generalizes [7, Theorem 3.3].

**Theorem 4.3.** Let A be a connected non-maximal order, P be an indecomposable bijective A-lattice and  $A_1 = A^-(P)$ . If  $P^{\mathfrak{r}} \simeq \mathfrak{r}P$ , the following holds.

- (1) There are chains of overmodules  $P = P_0 \subset P_1 \subset P_2 \subset \ldots \subset P_m$ and overrings  $A = A_0 \subset A_1 \subset A_2 \subset \ldots \subset A_m$  such that for every  $0 \leq i < m$ 
  - (a)  $P_{i+1} = P_i^{\mathfrak{r}_i} \simeq \mathfrak{r}_i P_i$ , where  $\mathfrak{r}_i = \operatorname{rad} P_i$ .
  - (b)  $P_i$  is an indecomposable bijective  $A_i$ -lattice which is not projective over  $A_{i-1}$  (hence over A) if  $i \neq 0$ .
  - (c)  $A_i$  is not maximal and  $A_{i+1} = A_i^-(P_i)$ .
- (2) If such a chain is of maximal length, then  $A_m$  is a hereditary order, has at most 2 non-isomorphic indecomposable lattices and every indecomposable A-lattice is isomorphic either to  $P_i$  for some  $0 \le i < m$ or to a direct summand of  $P_m$ .
- (3) A is Morita equivalent either to a local Bass order  $E = (\operatorname{End}_A P)^{\operatorname{op}}$ or to the Bass order  $B(k, \Delta)$  for some k and  $\Delta$ .

The condition  $P^{\mathfrak{r}} \simeq \mathfrak{r}P$  holds if  $P^{\mathfrak{r}}$  has no L-injective summands as Alattice, but is L-injective as A<sub>1</sub>-lattice, or, by duality, if  $\mathfrak{r}P$  has no projective summands as A-module but is projective as A<sub>1</sub>-module.

Recall also that, by Lemma 3.11,  $P^{t}$  cannot have L-injective summands if A is Gorenstein.

*Proof.* First, we prove the last claim. Theorem 3.8 implies that L-injective lattices over  $A_1$  are either L-injective over A or direct summands of  $\mathfrak{r}P$ . If  $P^{\mathfrak{r}}$  had no summands L-injective over A but is L-injective over  $A_1$ , every direct summand of  $P_1$  is isomorphic to a direct summand of  $\mathfrak{r}P$ . By Lemma 3.6, either  $P^{\mathfrak{r}}$  and  $\mathfrak{r}P$  are indecomposable or  $P^{\mathfrak{r}} = L_1 \oplus L_2$  and  $\mathfrak{r}P = \mathfrak{r}L_1 \oplus \mathfrak{r}L_2$ , where  $L_1, L_2, \mathfrak{r}L_1, \mathfrak{r}L_2$  are indecomposable. It implies that  $P^{\mathfrak{r}} \simeq \mathfrak{r}P$ .

Let  $P_1 = P^{\mathfrak{r}} \simeq \mathfrak{r}P$ . As A is not maximal,  $P_1 \not\simeq P$  by Lemma 3.4. So there are chains of overrings and overmodules satisfying (a),(b),(c): for instance,  $P = P_0 \subset P_1 = P^{\mathfrak{r}}$  and  $A = A_0 \subset A_1 = A^-(P)$ . As there are no infinite chains of overrings, consider a chain of maximal length m with these properties. Lemma 3.5 and Theorem 3.8 imply that

- $P_m$  is a bijective  $A_m$ -module, but is not projective over  $A_{m-1}$  (hence over A) if  $i \neq 0$ .
- If i < m, every indecomposable A-module either is isomorphic to one of the modules  $P_0, P_1, \ldots, P_i$  or is an  $A_{i+1}$ -module.
- Every principal  $A_i$ -module is either projective over A or isomorphic to a direct summand of  $P_i$  (hence to  $P_i$  if i < m).

If i < m,  $P_{i-1} \neq \mathfrak{r}_i P_i$ , since  $P_{i-1}$  is not an  $A_i$ -lattice, but  $\mathfrak{r}_i P_i \supseteq \mathfrak{r}_{i-1} P_{i-1}$ . If  $\mathfrak{r}_i P_i = \mathfrak{r}_{i-1} P_{i-1} \simeq P_i$ ,  $A_i$  is maximal, which is impossible. Therefore,  $\mathfrak{r}_i P_i \cap P_{i-1} = \mathfrak{r}_{i-1} P_{i-1}$  and  $\mathfrak{r}_i P_i + P_{i-1} = P_i$ , whence

(4.1) 
$$P_i/\mathfrak{r}_i P_i \simeq P_{i-1}/\mathfrak{r}_{i-1} P_{i-1} \simeq P_{i-2}/\mathfrak{r}_{i-2} P_{i-2} \simeq \cdots \simeq P/\mathfrak{r} P.$$

As  $\mathfrak{r}_i P_i \simeq P_{i+1}$  and  $\mathfrak{r}_{i-1} P_{i-1} \simeq P_i$ , we also have that

$$(4.2) P_{i+1}/P_i \simeq P_i/P_{i-1} \simeq P_{i-1}/P_{i-2} \simeq \cdots \simeq P_1/P$$

Suppose first that  $A_m$  decomposes:  $P_m = L_1 \oplus L_2$ , where  $L_1$  and  $L_2$  are indecomposable and non-projective over  $A_{m-1}$  (hence over A) by Lemma 3.6. As  $\mathfrak{r}_{i-1}P_m = \mathfrak{r}_{i-1}L_1 \oplus \mathfrak{r}_{i-1}L_2 \simeq L_1 \oplus L_2$  and  $\mathfrak{r}_{i-1}L_1, \mathfrak{r}_{i-1}L_2$  are indecomposable, either  $\mathfrak{r}_{i-1}L_1 \simeq L_1$  and  $\mathfrak{r}_{i-1}L_2 \simeq L_2$  or  $\mathfrak{r}_{i-1}L_1 \simeq L_2$  and  $\mathfrak{r}_{i-1}L_2 \simeq L_1$ . In both cases all submodules of  $L_1$  and  $L_2$  are projective, isomorphic either to  $L_1$  or to  $L_2$ . Therefore, all indecomposable  $A_m$ -lattices are isomorphic to  $L_1$ or to  $L_2$ ,  $A_m$  is hereditary and  $P_0, P_1, \ldots, P_{m-1}, L_1, L_2$  are all indecomposable A-lattices. Hence  $A_0, A_1, \ldots, A_{m-1}$  are all non-hereditary overrings of A, so A is Bass. P is a unique principal A-module, so A is Morita equivalent to the local Bass ring E.

Let now  $P_m$  be indecomposable. Note that  $P_{m-1} \supseteq \mathfrak{r}_{i-1}P_m \supseteq \mathfrak{r}_{i-1}P_{m-1}$ . Suppose that  $P_m$  is projective as  $A_{m-1}$ -module. Then  $\mathfrak{r}_{i-1}P_m = P_{m-1}$ . Conversely, if  $\mathfrak{r}_{i-1}P_m = P_{m-1}$ , i.e.  $\ell_{A_{m-1}}(P_m/\mathfrak{r}_{i-1}P_m) = 1$ , there is a surjection  $\varphi: P' \to P_m$ , where P' is a principal  $A_{m-1}$ -module. If  $P' = P_{m-1}$ , then  $\varphi$  is an isomorphism, since  $wd(P_{m-1}) = wd(P_m)$ . Otherwise P' is an  $A_m$ -module, thus  $P' \simeq P_m$ , since  $P_m$  is also projective over  $A_m$ . Thus  $P_m$  is projective over  $A_{m-1}$ , hence also over A. As  $\mathfrak{r}_{m-1}P_m \simeq P_{m-1}$  and  $\mathfrak{r}_{m-1}P_{m-1} \simeq P_m$ , Lemma 3.4 implies that  $A_{m-1}$  is hereditary and  $P_{m-1}, P_m$ are all its indecomposable modules. Let  $\Delta = \operatorname{End}_A P_m$ ,  $\mathfrak{d} = \operatorname{rad} \Delta$ . It is a maximal order and also  $\operatorname{End}_A P_{m-1} \simeq \Delta$  [5]. Since  $P_m \not\simeq P$ , the quotients  $P_m/P_{m-1}$  and  $P/\mathfrak{r}P$  are not isomorphic. From (4.1) and (4.2) it follows that, for every i < m,  $P_{i-1}$  is a unique maximal submodule of  $P_i$  such that  $P_i/P_{i-1} \simeq P_m/P_{m-1}$ . Therefore,  $\varphi(P_{i-1}) \subseteq P_{i-1}$  for every endomorphism  $\varphi \in \operatorname{End}_A P_i$ , hence  $\operatorname{End}_A P_i \simeq \Delta$  for all *i*, in particular,  $\operatorname{End}_A P \simeq \Delta$ . As P and  $P_m$  are all principal A-modules, A is Morita equivalent to  $\tilde{A} = (\operatorname{End}_A(P \oplus P_m))^{\operatorname{op}}$ . Since any  $\Delta$ -ideal (left or right) coincides with  $\mathfrak{d}^k$  for some k,

$$\tilde{A} \simeq \begin{pmatrix} \Delta & \mathfrak{d}^k \\ \mathfrak{d}^l & \Delta \end{pmatrix} \simeq \begin{pmatrix} \Delta & \mathfrak{d}^{k+l} \\ \Delta & \Delta \end{pmatrix} = B(k+l,\Delta)$$

for some k, l.

Let now  $P_m$  be indecomposable and not projective over  $A_{m-1}$ . Then  $\mathbf{r}_{m-1}P_m = \mathbf{r}_{m-1}P_{m-1}$  and  $P_m \supset \mathbf{r}_m P_m \supseteq \mathbf{r}_{m-1}P_{m-1}$ . If  $\mathbf{r}_m P_m = \mathbf{r}_{m-1}P_{m-1} \simeq P_m$ , then  $A_m$  is maximal and  $P_m$  is a unique indecomposable  $A_m$ -lattice. Therefore,  $P_0, P_1, \ldots, P_m$  are all indecomposable A-lattices,  $A_0, A_1, \ldots, A_m$ are all overrings of A and A is Bass. Moreover, P is a unique principal A-module and A is Morita equivalent to E.

If  $P_m$  is indecomposable, not projective over  $A_{m-1}$  and  $P_{m-1} \neq \mathfrak{r}_m P_m \neq \mathfrak{r}_{i-1}P_{i-1}$ , then  $\mathfrak{r}_m P_m$  is a minimal overmodule of  $\mathfrak{r}_{m-1}P_{m-1} \simeq P_m$ . Therefore,  $\mathfrak{r}_m P_m \simeq P_m^{\mathfrak{r}_m}$ , so, if we set  $P_{m+1} = P_m^{\mathfrak{r}_m}$ ,  $A_{m+1} = A_m^-(P_m)$ , we obtain a longer chain of overrings and overmodules satisfying the conditions (a),(b),(c), which is impossible. It accomplishes the proof.  $\Box$ 

**Corollary 4.4** ([7, Theorem 3.3]). Let A be a connected Gorenstein order. If one of its minimal overrings is also Gorenstein, then A is Bass and is either hereditary, or Morita equivalent to a local Bass order, or Morita equivalent to an order  $B(k, \Delta)$ .

*Proof.* It follows from Theorem 4.3 together with Lemma 3.11 and Proposition 3.13.  $\Box$ 

**Corollary 4.5** ([7, Proposition 3.7]). Let A be a local Gorenstein order,  $A' = A^{-}(A)$  be its minimal overring. If A' is not local, A' is hereditary and A is Bass.

Proof. By Proposition 3.13,  $A' = A^{-}(A)$ . If A' is not local, then  $A' = P_1 \oplus P_2$ , where both  $P_i$  are indecomposable projective A'-modules and both  $\mathfrak{r}_P_i$  are indecomposable L-injective A'-lattices. In particular, rad  $A' = \mathfrak{r}$ . Let  $P'_1$  be a minimal overmodule of  $P_1$  and M be a maximal submodule of  $P'_1$ . Then  $M = P_1$ : otherwise,  $M \cap P_1 = \mathfrak{r}_P_1$ , hence M is a minimal overmodule of  $\mathfrak{r}_P_1$ , which is impossible, since  $P_1$  is a unique minimal overmodule of  $\mathfrak{r}_P_1$ . Thus  $P_1$  is a unique maximal submodule of  $P'_1$ , so there is an epimorphism  $\varphi : P \to P'_1$  for some indecomposable projective A-module P. If  $P = P_1, \varphi$  is an isomorphism. If  $P = P_2, \varphi$  induces an epimorphism  $\varphi' : \mathfrak{r}_P_2 \to \mathfrak{r}_P'_1 = P_1$ . As  $\mathfrak{r}_P_2$  is indecomposable,  $\varphi'$  is an isomorphism, hence so is  $\varphi$ . Therefore, either  $P'_1 \simeq P_1$  or  $P'_1 \simeq P_2$ . Just in the same way, if  $P'_2$  is a minimal overmodule of  $P_2$ , then either  $P'_2 \simeq P_1$  or  $P'_2 \simeq P_2$ . Now Lemma 3.4 implies that A' is hereditary, thus A is Bass.

## 5. Stable categories

- **Definition 5.1.** (1) Let  $\mathcal{C}$  be an additive category,  $\mathfrak{S}$  be a set of morphisms from  $\mathcal{C}$ . Denote by  $\langle \mathfrak{S} \rangle$  the ideal of  $\mathcal{C}$  generated by  $\mathfrak{S}$ , i.e. consisting of morphisms of the form  $\sum_{i=1}^{k} \alpha_i \sigma_i \beta_i$ , where  $\sigma_i \in \mathfrak{S}$ . The quotient  $\mathcal{C}/\langle \mathfrak{S} \rangle$  is denoted by  $\mathcal{C}^{\mathfrak{S}}$ . Its objects are those from  $\mathcal{C}$  and the sets of morphisms from M to N are  $\operatorname{Hom}_{\mathcal{C}}^{\mathfrak{S}}(M,N) = \operatorname{Hom}_{\mathcal{C}}(M,N)/\mathfrak{S}(M,N)$ , where  $\mathfrak{S}(M,N) = \langle \mathfrak{S} \rangle \cap \operatorname{Hom}_{\mathcal{C}}(M,N)$ .
  - (2) The category A-mod<sup> $\langle 1_A \rangle$ </sup> is denoted by <u>A-mod</u> and its sets of morphisms are denoted by <u>Hom<sub>A</sub>(M, N)</u>. Obviously, it coincides with

A-mod<sup> $\mathfrak{P}$ </sup> for  $\mathfrak{P} = \{1_{P_1}, 1_{P_2}, \ldots, 1_{P_n}\}$ , where  $P_1, P_2, \ldots, P_n$  is a complete list of non-isomorphic principal A-modules. If A is an order, the full subcategory of A-mod<sup> $\langle 1_A \rangle$ </sup> consisting of A-lattices coincides with A-lat<sup> $\langle 1_A \rangle$ </sup> and is denoted by <u>A-lat</u>. We call it the *stable category* of the order A.

(3) Dually, the category  $A \operatorname{-lat}^{\langle 1_{\omega_A} \rangle}$  is denoted by  $\overline{A} \operatorname{-lat}$  and its sets of morphisms are denoted by  $\overline{\operatorname{Hom}}_A(M, N)$ . Obviously, it coincides with  $A \operatorname{-lat}^{\mathfrak{I}}$  for  $\mathfrak{I} = \{1_{I_1}, 1_{I_2}, \ldots, 1_{I_n}\}$ , where  $I_1, I_2, \ldots, I_n$  is a complete list of non-isomorphic coprincipal A-lattices. We call it the costable category of the order A.

The duality D induces a duality between the categories <u>A-lat</u> and  $\overline{A^{\text{op}}}$ -lat. If A is Gorenstein, the stable and costable categories coincide.

Note that all R-modules  $\underline{\text{Hom}}_A(M, N)$  and  $\text{Hom}_A(M, N)$  are of finite length. Moreover, we can estimate there annihilators.

**Lemma 5.2.** Let  $A_0$  be a hereditary (for instance, maximal) overring of A,  $\mathfrak{c} = \operatorname{Ann}_R(A_0/A)$ . Then  $\mathfrak{c}^2 \operatorname{\underline{Hom}}_A(M, N) = \mathfrak{c}^2 \operatorname{\overline{Hom}}_A(M, N) = 0$  for any  $M, N \in \operatorname{CM}(A)$ .

Proof. Let M and N be A-lattices,  $\lambda, \mu \in \mathfrak{c}$ . Consider  $A_0 M \subset KM$ . Then  $\lambda A_0 M \subseteq M$ . As  $A_0$  is hereditary,  $A_0 M$  is a projective  $A_0$ -module. Hence  $A_0 M$  is a direct summand of a free  $A_0$ -module F', which can be identified with  $A_0 F$ , where F is a free A-module. Any homomorphism  $f : M \to N$  extends to a homomorphism  $A_0 M \to A_0 N$ , hence to a homomorphism  $g : F' \to A_0 N$ . Moreover,  $F \supseteq \lambda F' \supseteq \lambda M$ , and  $\operatorname{Im}(\mu g) \subseteq \mu A_0 N \subseteq N$ . Therefore, the map  $\lambda \mu f$  can be considered as the composition

$$M \xrightarrow{\lambda} \lambda M \hookrightarrow F \xrightarrow{\mu g|_F} N.$$

So  $\lambda \mu f$  factors through a projective module and its image in  $\underline{\operatorname{Hom}}_A(M, N)$  is zero. By duality, the same is true for  $\overline{\operatorname{Hom}}_A(M, N)$ .

There are two important functors on stable categories. Let  $\pi : P \to M$ be a projective cover of a finitely generated A-module M,  $\Omega M = \text{Ker }\pi$ . Note that  $\Omega M$  is always an A-lattice, non-zero if M is not projective. If M is a non-projective lattice,  $\Omega M$  is not L-injective (otherwise  $\pi$  splits). If  $\pi': P' \to M'$  is a projective cover of M', any homomorphism  $\alpha : M \to M'$ can be lifted to a homomorphism  $P \to P'$ , hence induces a homomorphism  $\gamma : \Omega M \to \Omega M'$ . If  $\gamma'$  comes from another lifting of  $\alpha$ , one easily checks that  $\gamma - \gamma'$  factors through P. Hence, the class of  $\gamma$  in <u>A-mod</u> or in <u>A-lat</u> is well defined and  $\Omega$  can be considered as endofunctor on the stable category. Using L-injective envelops, we can define the analogous functor  $\Omega'$  on <del>A</del>-lat. If A is Gorenstein, a projective cover of M is also an L-injective envelop of  $\Omega M$ , hence  $\Omega'$  is a quasi-inverse of the functor  $\Omega$ .

Let now  $P_1 \xrightarrow{\psi} P_0 \xrightarrow{\varphi} M \to 0$  be a minimal projective presentation of a finitely generated A-module M, i.e. an exact sequence, where  $P_0, P_1$  are projective, Ker  $\varphi \subseteq \mathfrak{r}P_0$  and Ker  $\psi \subseteq \mathfrak{r}P_1$ . Apply to this sequence the functor  $\vee = \operatorname{Hom}_A(-, A)$ . We obtain the exact sequence of right modules

(5.1) 
$$0 \to M^{\vee} \xrightarrow{\varphi^{\vee}} P_0^{\vee} \xrightarrow{\psi^{\vee}} P_1^{\vee} \to \operatorname{Tr} M \to 0,$$

where  $\operatorname{Tr} M = \operatorname{Cok} \psi^{\vee}$ . Again one easily checks that in this way we obtain a functor  $\operatorname{Tr} : \underline{A \operatorname{-mod}} \to \underline{A^{\operatorname{op}}\operatorname{-mod}}$ . As the natural map  $P \to P^{\vee \vee}$  is an isomorphism for every finitely generated projective P, we have an isomorphism of functors  $\mathbf{1}_{\underline{A}\operatorname{-mod}} \xrightarrow{\sim} \operatorname{Tr}^2$ . Note that if M is a lattice, it can happen that  $\operatorname{Tr} M$  is not.

There is a natural map  $M^{\vee} \otimes_A N \to \operatorname{Hom}_A(M, N)$ , which maps  $u \otimes v$  to the homomorphism  $x \mapsto u(x)v$ . One easily sees [2] that its image coincides with  $\mathfrak{P}(M, N)$ . From the exact sequence (5.1) it follows that  $\operatorname{Tor}_1^A(\operatorname{Tr} M, N) \simeq \operatorname{Hom}_A(M, N)$ .

We will study the behaviour of <u>A-lat</u> and <del>A-lat</del> under rejection of bijective lattices.

**Lemma 5.3.** Suppose that the order A is not maximal. Let B be an indecomposable bijective A-lattice,  $A' = A^{-}(B)$ , M, N be A'-lattices.

- (1) The natural restriction maps  $\gamma_+$ : Hom<sub>A</sub>( $B^{\mathfrak{r}}, M$ )  $\rightarrow$  Hom<sub>A</sub>(B, M) and  $\gamma_-$ : Hom<sub>A</sub>( $M, \mathfrak{r}B$ )  $\rightarrow$  Hom<sub>A</sub>(M, B) are bijective.
- (2) A homomorphism  $\alpha : M \to N$  factors through B if and only if it factors through the embedding  $\mathfrak{r}B \to B^{\mathfrak{r}}$ .

*Proof.* (1) Since  $B/\mathfrak{r}B$  is a finite module, the map  $\gamma_{-}$  is injective. Since M does not have B as a direct summand,  $\operatorname{Im} \alpha \subseteq \mathfrak{r}B$  for any  $\alpha : M \to B$ . Hence  $\gamma_{-}$  is bijective. The assertion about  $\gamma_{+}$  is just dual.

(2) is an obvious consequence of (1).

**Theorem 5.4.** Let A be a non-hereditary order, B be a bijective A-lattice,  $P_1, P_2, \ldots, P_n$  be a complete list of non-isomorphic principal A-modules,  $I_1, I_2, \ldots, I_n$  be a complete list of non-isomorphic coprincipal A-lattices and  $A' = A^-(B)$ . Set  $\mathfrak{P}^B = \{\iota_{P_i}^B \mid 1 \leq i \leq n\}$  and  $\mathfrak{I}^B = \{\iota_{I_i}^B \mid 1 \leq i \leq n\}$ . Then  $\underline{A \cdot \text{lat}} \simeq A' \cdot \text{lat}^{\mathfrak{P}_B}$  and  $\overline{A \cdot \text{lat}} \simeq A' \cdot \text{lat}^{\mathfrak{I}_B}$ .

Actually, it means that, defining <u>A-lat</u> (respectively, <del>A-lat</del>) we may replace A by A' and, for each B-link  $B_1, B_2, \ldots, B_l$ , replace in  $\mathfrak{P}$  (respectively, in  $\mathfrak{I}$ ) all maps  $\mathbb{1}_{B_i}$   $(1 \leq i \leq l)$  by the embeddings  $\mathfrak{r}B_l \to B_1^{\mathfrak{r}}$ .

*Proof.* If B is indecomposable, the assertion follows from Lemma 5.3. Then the general case is obtained by induction on the number of non-isomorphic indecomposable direct summands of B using Theorem 3.8.  $\Box$ 

**Corollary 5.5.** Let A be a non-hereditary Gorenstein order,  $P_1, P_2, \ldots, P_n$ be a complete list of non-isomorphic principal A-modules,  $\iota_i$  be the embedding  $\mathfrak{r}P_i \to P_i^{\mathfrak{r}}$ ,  $A' = A^-(A)$ . Then <u>A-lat</u>  $\simeq A'$ -lat<sup> $\mathfrak{P}'$ </sup>, where  $\mathfrak{P}' = {\iota_1, \iota_2, \ldots, \iota_n}$ .

*Proof.* It follows from Theorem 5.4 and Lemma 3.12.

### 6. Almost split sequences

Recall some definitions and results (cf. [2]). Let A be an order,  $\alpha : N \to M$ and  $\beta : M \to N$  are homomorphisms of lattices, where M is indecomposable.

# **Definition 6.1.** (1) $\alpha$ is called *right almost split* if the following conditions hold:

- (a)  $\alpha$  is a non-split epimorphism;
- (b) any homomorphism  $\xi : X \to M$  which is not a split epimorphism factors through  $\alpha$ ;

(c) if  $\varphi : N \to N$  is such that  $\alpha \varphi = \alpha$ , then  $\varphi$  is an isomorphism. Note that if (a) and (b) holds, then either (c) holds or  $N = N_0 \oplus N_1$ , where  $N_0 \subset \text{Ker } \alpha$  and  $\alpha|_{N_1}$  is right almost split.

- (2)  $\beta$  is called *left almost split* if the following conditions hold:
  - (a)  $\beta$  is a non-split inflation;
  - (b) any homomorphism  $\xi : X \to M$  which is not a split monomorphism factors through  $\beta$ ;

(c) if  $\varphi : N \to N$  is such that  $\varphi \beta = \beta$ , then  $\varphi$  is an isomorphism. Note that If (a) and (b) holds, then either (c) holds or  $N = N_0 \oplus N_1$ , where Im  $\beta \subset N_1$  and  $\beta$  is left almost split considered as a map  $M \to N_1$ .

- (3) An exact sequence of A-lattices  $\varepsilon : 0 \to L \xrightarrow{\beta} N \xrightarrow{\alpha} M \to 0$ , where M and L are indecomposable, is called an *almost split sequence* if the following equivalent conditions hold:
  - (a)  $\alpha$  is right almost split;
  - (b)  $\beta$  is left almost split;
  - (c) for every homomorphism  $\xi : X \to M$ , which is not a split epimorphism, the exact sequence  $\varepsilon \xi$  splits;
  - (d) for every homomorphism  $\eta: L \to X$ , which is not a split inflation, the exact sequence  $\eta \varepsilon$  splits.

Here  $\varepsilon \xi$  (respectively,  $\eta \varepsilon$ ) is the pull-back of the exact sequence  $\varepsilon$  along  $\xi$  (respectively, the push-down of  $\varepsilon$  along  $\eta$ ).

Obviously, a right (or left) almost split morphism, if exists, is unique up to isomorphism of N. In the same way, an almost split sequence with a fixed term M (or L), if exists, is unique, up to isomorphism of L (respectively, of M). Actually, in the category A-lat it exists for every non-projective indecomposable M, as well as for every non-L-injective indecomposable L. It can be proved following literally to [1]. We recall the main steps.

The functor  $\tau_A = D\Omega \operatorname{Tr} : \underline{A \operatorname{-lat}} \to A \operatorname{-lat}$  is called the Auslander-Reiten transpose. Just as in [1, Proposition 1.1], one proves that

$$\operatorname{Ext}_{A}^{1}(N, \tau_{A}M) \simeq \operatorname{Hom}_{A}(\widetilde{M}, N).$$

Let M be an indecomposable non-projective A-lattice. Then the ring  $\Lambda = \underline{\operatorname{Hom}}_A(M, M)$  is local. Dually,  $\underline{\operatorname{Hom}}_A(M, M)$  has a unique minimal  $\Lambda$ -submodule U. If u is a non-zero element of U, then  $u(\lambda) = 0$  for every non-invertible  $\lambda \in \Lambda$ . If  $\xi : X \to M$  is not a split epimorphism, then  $\xi\varphi$  is not invertible for every  $\varphi : M \to X$ , whence  $u\xi(\varphi) = u(\xi\varphi) = 0$ , i.e.  $u\xi = 0$ . Therefore, the same holds for the corresponding extension  $\varepsilon \in \operatorname{Ext}^1_A(M, \tau_A M)$ , thus the extension

(6.1) 
$$\varepsilon: 0 \to \tau_A M \xrightarrow{\beta} E \xrightarrow{\alpha} M \to 0,$$

is an almost split sequence. Note that, if  $0 \to L \to N \to M \to 0$  is an almost split sequence, so is its dual  $0 \to DM \to DN \to DL \to 0$ . Therefore, if  $L = \tau_A M$ , then  $DM \simeq \tau_A DL$  and  $M \simeq D\tau_A DL \simeq \Omega \operatorname{Tr} DM$ . So the functor  $\tau_A$  has a quasi-inverse  $\tau_A^{-1} = \Omega \operatorname{Tr} D : \overline{A}\operatorname{-lat} \to \underline{A}\operatorname{-lat}$ .

Let  $M = \bigoplus_{i} M_{j}$  and  $N = \bigoplus_{i} N_{i}$ , where  $M_{j}$  and  $N_{i}$  are indecomposable A-lattices. Denote by  $\operatorname{Rad}_A(M, N)$  the set of homomorphisms  $\varphi: M \to N$ such that all components  $\varphi_{ij}: M_j \to N_i$  are non-isomorphisms. Obviously, we obtain an ideal of the category A-lat called its radical. So we can define its degrees  $\operatorname{Rad}_A^n$   $(n \in \mathbb{N})$  and  $\operatorname{Rad}_A^\infty = \bigcap_{n=1}^\infty \operatorname{Rad}_A^n$ . The homomorphisms from  $\operatorname{Rad}_A(M, N) \setminus \operatorname{Rad}_A^2(M, N)$  are called *irreducible*. The quotient  $_NV_M = \operatorname{Rad}_A(M, N) / \operatorname{Rad}_A^2(M, N)$  is a finite dimensional vector space over the residue field k. In particular, if the lattice M is indecomposable,  $F_M = {}_M V_M$  is a skewfield, and for any N both  ${}_N V_M$  and  ${}_M V_N$ are finite dimensional vector spaces over  $F_M$  (respectively, right and left). Let A-ind be the set of isomorphisms classes of indecomposable A-lattices. The set  $\{F_M, NV_M \mid M, N \in A \text{-ind}\}$  is called the Auslander-Reiten species of the order A and denoted by  $AR_A$ . It is indeed a k-species in the sense of [6], since all  $F_M$  are skewfields and  ${}_NV_M$  is an  $F_N$ - $F_M$ -bimodule. If the residue field k is algebraically closed, so  $F_M = k$  for any indecomposable M, this species is usually written as a quiver whose vertices are  $M \in A$ -ind and there are  $d_{NM}$  arrows from M to N, where  $d_{NM} = \dim_{\mathbb{K}}({}_{N}V_{M})$ . It is called the Auslander-Reiten quiver of A. Obviously, the species of  $A^{\text{op}}$ -lat is  $(F_M^{\text{op}}, {}_MV_N^{\text{op}})$ , where  ${}_MV_N^{\text{op}} = {}_NV_M$ . So in the Auslander-Reiten quiver one only has to revert all arrows.

If the lattice M is indecomposable and non-projective, the definition of an almost split sequence shows that every homomorphism from  $\operatorname{Rad}_A(N, M)$ , as well as every homomorphism from  $\operatorname{Rad}_A(\tau_A M, N)$  factors through the term E of the sequence (6.1). Hence, if  $E = \bigoplus_{i=1}^r E_i$  with indecomposable  $E_i$ , then  $_MV_N = 0 = _NV_{\tau_A M}$  if  $N \not\simeq E_i$  for all  $1 \leq i \leq r$ , while both  $_MV_{E_i}$  and  $_{E_i}V_{\tau_A M}$  are non-zero. In the case of the Auslander–Reiten quiver, there are only arrows from each of  $E_i$  to M and from  $\tau_A M$  to each of  $E_i$ . Note also that if  $\alpha_i$  are the components of  $\alpha$  and  $\beta_i$  are the components of  $\beta$  in the sequence 6.1, then  $\sum_{i=1}^r \alpha_i \beta_i = 0$ .

If P is principal, the image of any homomorphism  $N \to P$ , which is not a split epimorphism, belongs to  $\mathfrak{r}P$ . Therefore, if  $\mathfrak{r}P = \bigoplus_{i=1}^{r} E_i$  with indecomposable  $E_i$ , the only non-zero spaces  ${}_PV_N$  are  ${}_PV_{E_i}$ . Dually, if I is coprincipal and  $I^{\mathfrak{r}} = \bigoplus_{i=1}^r E_i$  with indecomposable  $E_i$ , the only non-zero spaces  ${}_NV_I$  are  ${}_{E_i}V_I$ .

If the lattices M and N are not projective, every homomorphism from  $\mathfrak{P}(M, N)$  is in  $\operatorname{Rad}_A^2(M, N)$ . So we can consider the stable Auslander-Reiten species (or the stable Auslander-Reiten quiver) <u>AR</u><sub>A</sub> whose objects are nonprojective indecomposable lattices and the bimodules  ${}_MV_N$  are the same as in AR<sub>A</sub>. Dually, the costable Auslander-Reiten species (or costable Auslander-Reiten quiver) <u>AR</u><sub>A</sub> is defined, consisting of non-L-injective indecomposable lattices. The functor  $\tau_A$  induces the Auslander-Reiten translation  $\underline{AR}_A \xrightarrow{\sim} \overline{AR}_A$ . Again, in Gorenstein case stable and costable species (or quivers) coincide.

We will use the following fact about irreducible morphisms between indecomposable lattices. Perhaps, it is known, though we have not found it in the literature.

**Proposition 6.2.** Let M, N be indecomposable lattices,  $\alpha : N \to M$  is an irreducible morphism. There are two possibilities:

- (1)  $\alpha$  is an isomorphism of N onto a direct summand of a maximal submodule of M.
- (2)  $\alpha$  is an epimorphism and there is a submodule  $L \subset N$  such that N/L is a lattice,  $L + \text{Ker } \alpha = N$  and  $L \cap \text{Ker } \alpha$  is an L-irreducible lattice.

Proof. Let  $M' = \operatorname{Im} \alpha$ ,  $\iota$  be the embedding  $M' \to M$  and  $\pi$  be the projection  $N \to M'$ . If  $M = \bigoplus_{i=1}^{m} M_i$ , where  $M_i$  are indecomposable,  $\iota_i$  and  $\pi_i$  are the components of  $\iota$  and  $\pi$  with respect to this decomposition. Then  $\alpha = \sum_{i=1}^{m} \iota_i \pi_i$ . As  $\alpha$  is irreducible, at least one of  $\iota_i$  or  $\pi_i$  must be invertible. Suppose that one of  $\iota_i$  is invertible. Then m = 1 and  $\alpha$  is an epimorphism. If Ker  $\alpha$  is L-irreducible, we can set  $L = \operatorname{Ker} \alpha$ . If Ker  $\alpha$  is not L-irreducible, it contains an L-irreducible sublattice S such that N/S is a lattice (take the intersection of Ker  $\alpha$  with a simple KA-submodule in  $K \operatorname{Ker} \alpha$ ). Then  $\alpha$  factors through the map  $\overline{\pi} : N/S \to N/\operatorname{Ker} \alpha \simeq M$ . Therefore,  $\overline{\pi}$  must be a split epimorphism, so  $N/S \simeq N/\operatorname{Ker} \alpha \oplus N/L$  for some  $L \supset S$  (in particular, N/L is a lattice). It actually means that  $L + \operatorname{Ker} \alpha = N$  and  $L \cap \operatorname{Ker} \alpha = S$ , which gives the possibility (2).

If one of  $\pi_i$  is invertible, then all other  $\pi_j = 0$  and  $\alpha$  is a monomorphism. If M' is a maximal submodule of M containing Im  $\alpha$ , then  $\alpha$  factors through the embedding Im  $\alpha \to M'$ , hence the latter must split. It gives the possibility (1).

We study the behaviour of these constructions under rejection of bijective lattices. First, a simple observation.

**Proposition 6.3.** Let B be a bijective A-lattice,  $A' = A^{-}(B)$ , M be an A'-lattice.

(1) If  $\alpha : N \to M$  is right almost split in A'-lat, it is so in A-lat.

(2) If  $\beta: M \to N$  is left almost split in A'-lat, it is so in A-lat.

(3) If  $0 \to L \to M \to N \to 0$  is an almost split sequence in A' -lat, it is so in A -lat.

*Proof.* (1) Let  $\xi \in \text{Hom}_A(X, M)$  be not a split epimorphism. If  $X \notin B$ , it is an A'-lattice, so  $\xi$  factors through  $\alpha$ . If  $X \in B$ , it is projective, so  $\xi$  also factors through  $\alpha$ .

- (2) by duality.
- (3) follows from (1) or (2).  $\Box$

The following theorem describes the "Auslander–Reiten behaviour" of new projective modules over the order  $A^{-}(B)$ .

**Theorem 6.4.** Let B be an indecomposable bijective A-lattice,  $A' = A^-(B)$ . Suppose that  $B^{\mathfrak{r}}$  is not projective over A (equivalently,  $\mathfrak{r}B$  is not L-injective over A).

(1) If  $B^{\mathfrak{r}}$  decomposes:  $B^{\mathfrak{r}} = M_1 \oplus M_2$ , there are almost split sequences

$$0 \to \mathfrak{r} M_1 \to B \to M_2 \to 0, 0 \to \mathfrak{r} M_2 \to B \to M_1 \to 0.$$

In particular,  $\tau_A M_1 = \mathfrak{r} M_2$  and  $\tau_A M_2 = \mathfrak{r} M_1$ .

(2) If  $B^{\mathfrak{r}}$  is indecomposable, then  $\tau_A B^{\mathfrak{r}} = \mathfrak{r}B$ ,  $B^{\mathfrak{r}}$  has a maximal submodule  $X \neq B$  and there is an almost split sequence

$$(6.2) 0 \to \mathfrak{r}B \to B \oplus X \xrightarrow{\alpha} B^{\mathfrak{r}} \to 0.$$

In particular,  $\tau_A B^{\mathfrak{r}} = \mathfrak{r} B$ .

Proof.  $B^{\mathfrak{r}}$  is projective and  $\mathfrak{r}B$  is L-injective over A' by Lemma 3.5. Let M be a direct summand of  $B^{\mathfrak{r}}$ ,  $N = \tau_A M$  and  $0 \to N \to E \to M \to 0$  be an almost split sequence in A-lat. If N were not L-injective as A'-lattice, there were an almost split sequence  $0 \to N \to E' \to M' \to 0$  in A'-lat. By Proposition 6.3, it were also an almost split sequence in A-lat, whence  $M' \simeq M$ , which is impossible, since M is projective over A'. Thus  $\tau_A M$  is L-injective as A'-lattice, but not as A-lattice. Therefore, it is a direct summand of  $\mathfrak{r}B$ . In particular, if  $B^{\mathfrak{r}}$  is indecomposable,  $\tau_A B^{\mathfrak{r}} = \mathfrak{r}B$ .

There is an irreducible morphism  $B \to M$ , hence B must be a direct summand of E, so  $E = B \oplus X$ . If  $B^{\mathfrak{r}} = M_1 \oplus M_2$ , there is an exact sequence  $0 \to \mathfrak{r}M_1 \to B \to M_2 \to 0$ . As  $KB \simeq KM_1 \oplus KM_2$ , X = 0. If B is indecomposable,  $KX \simeq KB$ . Hence Proposition 6.2 implies that in the almost split sequence (6.2) the restriction of  $\alpha$  on X is an isomorphism onto a maximal submodule of  $B^{\mathfrak{r}}$  which cannot coincide with B.

Remark 6.5. (1) It can happen that in case (1)  $M_1 \simeq M_2$  and in case (2)  $X \simeq B$ . If  $X \neq B$ , then it is an A'-lattice and  $X = \mathfrak{r}'B^{\mathfrak{r}}$ , where  $\mathfrak{r}' = \operatorname{rad} A'$ . If  $X \simeq B$ , then  $\mathfrak{r}'B^{\mathfrak{r}} = \mathfrak{r}B^{\mathfrak{r}}$ .

(2) By Lemma 3.11, the condition " $B^{\mathfrak{r}}$  is not projective" always holds if A is connected, Gorenstein and non-hereditary.

## 7. Gorenstein and Frobenius Cases

If the order A is Gorenstein, the functor  $\vee : M \mapsto M^{\vee} = \operatorname{Hom}_A(M, A)$ is an exact duality  $A\operatorname{-lat} \to A^{\operatorname{op}}\operatorname{-lat}$ . Combining it with the duality  $D : A^{\operatorname{op}}\operatorname{-lat} \to A\operatorname{-lat}$ , we obtain the Nakayama equivalence  $\mathcal{N} = D \vee : A\operatorname{-lat} \to A\operatorname{-lat}$ . It maps projective modules to projective, thus can also be considered as the functor on stable categories  $\underline{A\operatorname{-lat}} \to \underline{A\operatorname{-lat}}$ . The following result is an analogue of [2, Proposition IV.3.6].

**Proposition 7.1.** If the order A is Gorenstein, the functors  $\tau_A$ ,  $\Omega N$  and  $N\Omega$  are isomorphic.

*Proof.* Let M be a non-projective A-lattice. Consider an exact sequence

$$0 \to N \xrightarrow{\alpha} P_1 \xrightarrow{\beta} P_0 \xrightarrow{\gamma} M \to 0,$$

where  $P_1 \xrightarrow{\beta} P_0 \xrightarrow{\gamma} M \to 0$  is a minimal projective presentation of M. It gives the exact sequence

$$0 \to M^{\vee} \xrightarrow{\gamma^{\vee}} P_0^{\vee} \xrightarrow{\beta^{\vee}} P_1^{\vee} \xrightarrow{\alpha^{\vee}} N^{\vee} \to 0.$$

Thus  $N^{\vee} \simeq \operatorname{Tr} M$  and  $\Omega \operatorname{Tr} M \simeq \operatorname{Im} \beta^{\vee}$ . Now the exact sequence

 $0 \to D(\operatorname{Im} \beta^{\vee}) \to P_0^{\vee \vee} \to DM^{\vee} \to 0$ 

shows that  $\tau_A M \simeq D(\operatorname{Im} \beta^{\vee}) \simeq \Omega \mathcal{N} M$ . One easily sees that this construction is functorial in M, so it gives an isomorphism  $\tau_A \simeq \Omega \mathcal{N}$ . Since  $\mathcal{N}$  is exact and maps projective modules to projective, it commutes with  $\Omega$ , i.e.  $\Omega \mathcal{N} \simeq \mathcal{N} \Omega$ .

Let  $A \simeq \bigoplus_{i=1}^{s} m_i P_i$ , where  $P_1, P_2, \ldots, P_s$  are pairwise non-isomorphic principal left A-modules. Then also  $A \simeq \bigoplus_{i=1}^{s} m_i P_i^{\vee}$  as right A-module,  $A \simeq \bigoplus_{i=1}^{s} m_i D P_i^{\vee}$  as left A-module, and  $D P_1^{\vee}, D P_2^{\vee}, \ldots, D P_s^{\vee}$  are all pairwise non-isomorphic coprincipal left A-modules. Therefore, A is Gorenstein if and only if there is a permutation  $\nu$  such that  $P_i \simeq D P_{\nu i}^{\vee}$  for all  $i = 1, 2, \ldots, s$ . The permutation  $\nu$  is called the Nakayama permutation.

**Definition 7.2.** An order A is called *Frobenius* if  $A \simeq DA$  as left A-module. It is called *symmetric* if  $A \simeq DA$  as A-bimodule.

Obviously, A is Frobenius if and only if it is Gorenstein and  $m_i = m_{\nu i}$  for all  $i = 1, 2, \ldots, s$ , where  $\nu$  is the Nakayama permutation. One easily sees that in this case also  $A \simeq DA$  as right A-module, so the definition of Frobenius orders is left-right symmetric.

**Definition 7.3.** Let M be a left A-module,  $\sigma$  be an automorphism of A. We denote by  ${}^{\sigma}M$  the left A-module such that it coincides with M as a group, but, for every  $a \in A$  and  $x \in M$ , the product ax in  ${}^{\sigma}M$  coincides with the product  $\sigma(a)x$  in M. Analogously  $N^{\sigma}$  is defined for a right A-module N and  ${}^{\rho}M^{\sigma}$  is defined for an A-bimodule M, where  $\rho$  is also an automorphism of A. If  $\rho$  or  $\sigma$  are identity, it is omitted and we write, respectively,  $M^{\sigma}$  or  ${}^{\rho}M$ .

#### YURIY A. DROZD

One easily sees that the maps  $x \mapsto \rho^{-1}(x)$  and  $x \mapsto \sigma^{-1}(x)$  give isomorphisms of A-bimodules, respectively,  $\rho A^{\sigma} \simeq A^{\rho^{-1}\sigma}$  and  $\rho A^{\sigma} \simeq \sigma^{-1}\rho A$ .

**Proposition 7.4.** A is Frobenius if and only if there is an automorphism  $\sigma \in \operatorname{Aut} A$  such that  $DA \simeq A^{\sigma}$  as A-bimodule. Moreover, there is an invertible element  $s \in KA$  such that  $\sigma(a) = s^{-1}as$  for all A.

Proof. Obviously, if such an automorphism exists, A is Frobenius. Suppose that A is Frobenius and let  $\varphi : A \xrightarrow{\sim} \Delta$  be an isomorphism of left Amodules, where  $\Delta = DA$ . It induces an isomorphism of left KA-modules  $K\varphi : KA \xrightarrow{\sim} K\Delta$ . Since KA is semisimple, it is symmetric as K-algebra, [5, 9.8] i.e. there is an isomorphisms of KA-bimodules  $\theta : KA \xrightarrow{\sim} K\Delta$ . The composition  $\theta^{-1} \cdot K\varphi$  is an automorphism of KA as of left KA-module, hence there is an invertible element  $s \in KA$  such that  $\theta^{-1} \cdot K\varphi(x) = xs$  for every  $x \in KA$ . In particular,  $\varphi(x) = \theta(xs)$  for every  $x \in A$ , so  $\Delta = \theta(As)$ . It implies that  $As = \theta^{-1}(\Delta)$  is a two-sided A-module, so  $sA \subseteq As$  or  $sAs^{-1} \subseteq A$ . Therefore,  $sAs^{-1} = A$  and  $s^{-1}As = A$ . Moreover,

$$\varphi(xa) = \theta(xas) = \theta(xs \cdots^{-1} as) = \theta(xs)s^{-1}as = \varphi(x)s^{-1}as.$$

Hence  $\varphi$  is an isomorphism of A-bimodules  $A^{\sigma} \xrightarrow{\sim} \Delta$ , where  $\sigma(a) = s^{-1}as$ .

One can check that the element s above is defined up to a multiplier of the form  $q\lambda$ , where q and  $\lambda$  are invertible element, respectively, from A and from the center of KA.

**Corollary 7.5.** Let A be a Frobenius order,  $\sigma \in \text{Aut } A$  be as in Proposition 7.4,  $\mathcal{N}$  be the Nakayama equivalence. There are functorial isomorphisms:

- $DM \simeq (M^{\vee})^{\sigma}$  for any left A-lattice M and  $DN \simeq {}^{\sigma^{-1}}(N^{\vee})$  for every right A-lattice N;
- $\mathcal{N}M \simeq {}^{\sigma^{-1}}M$  and  $\tau_A M \simeq \Omega({}^{\sigma^{-1}}M) \simeq {}^{\sigma^{-1}}(\Omega M)$  for every left Alattice M.

In particular, if A is symmetric,  $\mathcal{N} \simeq \text{Id}$  and  $\tau_A \simeq \Omega$ .

*Proof* is obvious.

**Corollary 7.6.** Let A be a Gorenstein order,  $\mathfrak{r} = \operatorname{rad} A$ ,  $P_1, P_2, \ldots, P_s$  be a complete set of non-isomorphic principal A-modules,  $\omega_i = DP_i^{\vee}$  (then  $\omega_1, \omega_2, \ldots, \omega_s$  is a complete set of non-isomorphic coprincipal A-modules). Set  $A' = A^-(A)$ ,  $P'_i = P^{\mathfrak{r}}_i$  and  $\omega'_i = \mathfrak{r}\omega_i$ . Then  $\tau_A P'_i \simeq \omega'_{\nu i}$ , where  $\nu$  is the Nakayama permutation.

*Proof.* It follows from Theorem 6.4.

**Corollary 7.7.** Let G be a finite group, A be a block of the group ring  $\mathbb{Z}_pG$ . It is a symmetric  $\mathbb{Z}_p$ -order. Set  $A' = A^-(A)$ . Then, for every non-projective A-lattice M (or, the same, for every A'-lattice M),

$$\hat{H}^{n}(G,M) \simeq \hat{H}^{n+1}(G,\tau_{A}M) \simeq \hat{H}^{n-1}(G,\tau_{A}^{-1}M).$$

*Proof.* It follows from Corollary 7.5 and Proposition 6.3.

Note that  $\tau_A M = \tau_{A'} M$  if M is not projective over A'. Otherwise  $\tau_A M$  is given by Corollary 7.6. In some cases the structure of the Auslander–Reiten species  $AR_{A'}$  can be calculated explicitly. Then it gives the values of the cohomologies. An example, when G is the Kleinian 4-group, can be found in [11].

#### References

- Auslander, M., Reiten, I., Almost split sequences for Cohen-Macaulay modules. Math. Ann., 277, 345–349 (1987).
- [2] Auslander, M., Reiten, I., Smalø, S. Representation Theory of Artin Algebras. Cambridge University Press,
- Bass, H., Finitistic dimension and a homological generalization of semiprimary rings. Trans. Am. Math. Soc. 95, 466–488 (1960).
- [4] Bruns, W., Herzog, J., Cohen-Macaulay Rings. Cambridge University Press, 1993.
- [5] Curtis, Ch. W., Reiner, I., Methods of Representation Theory. Vol. 1. John Wiley & Sons, 1981.
- [6] Dlab, V., Ringel, C. M., Indecomposable representations of graphs and algebras. Mem. Amer. Math. Soc., 173 (1976).
- [7] Drozd, Yu. A., Kirichenko, V. V., On quasi-Bass orders. Izv. Akad. Nauk SSSR. Ser. Mat., 36, 328–370 (1972).
- [8] Drozd, Yu. A., Kirichenko, V. V., Primary orders with a finite number of indecomposable representations. Izv. Akad. Nauk SSSR. Ser. Mat., 37, 715–736 (1973).
- [9] Drozd, Yu. A., Kirichenko, V. V., Finite Dimensional Algebras. Springer, 1994.
- [10] Drozd, Yu. A., Kirichenko, V. V., Roiter, A. V., On hereditary and Bass orders. Izv. Akad. Nauk SSSR. Ser. Mat., 31, 1415–1436 (1967).
- [11] Drozd, Yu., Plakosh, A., Cohomologies of the Kleinian 4-group. Arch. Math. (2020) DOI:https://doi.org/10.001/s00013-020-01451-6 (arXiv:1910.03868 [math.Gr]).
- [12] Hijikata, H., Nishida, K., Bass orders in non semisimple algebras. J. Math. Kyoto Univ., 34, 797–837 (1994).
- [13] Hijikata, H., Nishida, K., Primary orders of finite representation type. J. Algebra 192, 592–640 (1997).
- [14] Keller, B., Derived categories and their use. Handbook of Algebra, vol. 1, 671–701. Elsevier, 1996.
- [15] Lam, T.Y., A First Course in Noncommutative Rings. Springer, 1991.
- [16] Matlis, E., Injective modules over Noetherian rings. Pacific J. Math., 8, 511–528 (1958).
- [17] Matsumura, H., Commutative Algebra. The Benjamin/Cummings Publishing Company, 1980.
- [18] Roiter, A. V., Analogue of a theorem of Bass for representation modules of noncommutative orders. Dokl. Akad. Nauk SSSR, 168, 1261-1264 (1966).

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, TERESCHENKIVSKA 3, KYIV 01001, UKRAINE

*E-mail address*: y.a.drozd@gmail.com *URL*: http://www.imath.kiev.ua/~drozd