

# THE $\ell^p$ -METRIZATION OF FUNCTORS WITH FINITE SUPPORTS

T. BANAKH, V. BRYDUN, L. KARCHEVSKA, M. ZARICHNYI

ABSTRACT. Let  $p \in [1, \infty]$  and  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  be a functor with finite supports in the category  $\mathbf{Set}$  of sets. Given a non-empty metric space  $(X, d_X)$ , we introduce the distance  $d_{FX}^p$  on the functor-space  $FX$  as the largest distance such that for every  $n \in \mathbb{N}$  and  $a \in Fn$  the map  $X^n \rightarrow FX$ ,  $f \mapsto Ff(a)$ , is non-expanding with respect to the  $\ell^p$ -metric  $d_{X^n}^p$  on  $X^n$ . We prove that the distance  $d_{FX}^p$  is a pseudometric if and only if the functor  $F$  preserves singletons;  $d_{FX}^p$  is a metric if  $F$  preserves singletons and one of the following conditions holds: (1) the metric space  $(X, d_X)$  is Lipschitz disconnected, (2)  $p = 1$ , (3) the functor  $F$  has finite degree, (4)  $F$  preserves supports. We prove that for any Lipschitz map  $f : (X, d_X) \rightarrow (Y, d_Y)$  between metric spaces the map  $Ff : (FX, d_{FX}^p) \rightarrow (FY, d_{FY}^p)$  is Lipschitz with Lipschitz constant  $\text{Lip}(Ff) \leq \text{Lip}(f)$ . If the functor  $F$  is finitary, has finite degree (and preserves supports), then  $F$  preserves uniformly continuous function, coarse functions, coarse equivalences, asymptotically Lipschitz functions, quasi-isometries (and continuous functions). For many dimension functions we prove the formula  $\dim F^p X \leq \text{deg}(F) \cdot \dim X$ . Using injective envelopes, we introduce a modification  $\check{d}_{FX}^p$  of the distance  $d_{FX}^p$  and prove that the functor  $\check{F}^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$ ,  $\check{F}^p : (X, d_X) \mapsto (FX, \check{d}_{FX}^p)$ , in the category  $\mathbf{Dist}$  of distance spaces preserves Lipschitz maps and isometries between metric spaces.

---

1991 *Mathematics Subject Classification.* 54B30; 54E35.

*Key words and phrases.* functor, distance, monoid, Hausdorff distance, finite support, dimension.

## CONTENTS

1. Introduction	3
2. Some notations and conventions	4
3. Distance spaces	4
4. Functions between distance spaces	5
5. The injective envelope of a distance space	7
6. Preliminaries on functors	13
7. Introducing the distance $d_{FX}^p$	14
8. The distance $d_{FX}^p$ between elements with small supports	14
9. Short linking chains	15
10. Equivalent definitions of the distance $d_{FX}^p$	17
11. The relation between the distances $d_{FX}^p$ and $d_{FX}^q$	19
12. Preserving Lipschitz functions by the functor $F^p$	20
13. Preserving isometries by the functor $F^p$	20
14. The functor $F^p f$ and the continuity modulus of functions	24
15. Upper bounds for $d_{FX}^p$	25
16. Lower bounds on the distance $d_{FX}^p$	27
17. Preservation of continuous functions by the functor $F^p$	31
18. Metric properties of the maps $\xi_X^a : X \rightarrow FX$ .	32
19. Preserving the compactness by the functor $F^p$	35
20. Preserving the completeness by the functor $F^p$	35
21. The functor $F^p$ and the metric entropy of distance spaces	40
22. The functor $F^p$ and the entropy dimensions of distance spaces	43
23. The functor $F^p$ and the Hausdorff dimension of distance spaces	45
24. The functor $F^p$ and the controlled dimensions of distance spaces	46
25. Concluding theorems on the properties of the distance $d_{FX}^p$	54
26. The $\check{\ell}^p$ -metrization $\check{F}^p$ of the functor $F$	56
27. The $\ell^p$ -metrization and natural transformations of functors	60
28. The metrizations of the functor of $n$ -th power	62
29. Metrizations of the functor of $M$ -valued finitary functions	62
30. The metrizations of the hyperspace functor	69
References	74

## 1. INTRODUCTION

Let **Set** be the category whose objects are sets and morphisms are functions between sets, and **Metr** be the category whose objects are metric spaces and morphisms are arbitrary maps between metric spaces. The category **Metr** is a subcategory of the category **Dist** whose objects are distance spaces and morphisms are arbitrary maps between distance spaces.

By a *distance space* we understand a pair  $(X, d_X)$  consisting of a set  $X$  and a *distance*  $d_X$  on  $X$ , which is a function  $d_X : X \times X \rightarrow [0, \infty]$  satisfying three well-known axioms:

- $d_X(x, x) = 0$ ,
- $d_X(x, y) = d_X(y, x)$ ,
- $d_X(x, z) \leq d_X(x, y) + d_X(y, z)$ ,

holding for any points  $x, y, z \in X$ . Here we assume that  $\infty + x = \infty = x + \infty$  for any  $x \in [0, \infty]$ .

An alternative term for a distance is an  $\infty$ -pseudometric. A distance  $d_X$  on a set  $X$  is called an  $\infty$ -*metric* if  $d_X(x, y) > 0$  for any distinct points  $x, y \in X$ . A distance  $d_X$  is a *pseudometric* if  $d_X(x, y) < \infty$  for any points  $x, y \in X$ .

In this paper, for every functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  with finite supports, every distance space  $(X, d_X)$  and every  $p \in [1, \infty]$ , we introduce a distance  $d_{FX}^p$  on the functor-space  $FX$  and study its properties. The distance  $d_{FX}^p$  is defined as the largest distance on  $FX$  such that for every  $n \in \mathbb{N}$  and  $a \in Fn$  the map  $\xi_X^a : X^n \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding with respect to the  $\ell^p$ -distance on the finite power  $X^n$  of  $X$ . The correspondence  $F^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$ ,  $F^p : (X, d_X) \mapsto (FX, d_{FX}^p)$ , determines a functor such that  $U \circ F^p = F \circ U$  where  $U : \mathbf{Dist} \rightarrow \mathbf{Set}$  is the (forgetful) functor assigning to each distance space  $(X, d)$  its underlying set  $X$ . So,  $F^p$  is a lifting of the functor  $F$  to the category **Dist**, called the  $\ell^p$ -*metrization* of the functor  $F$ . In the paper we establish some properties of the functor  $F^p$ ; in particular, we prove that this functor preserves Lipschitz functions between distance spaces. In Theorem 25.1 we prove that for any metric space  $(X, d_X)$  the distance  $d_{FX}^p$  is a metric if the functor  $F$  preserves singletons and one of the following conditions holds: (1) the distance space  $(X, d_X)$  is Lipschitz disconnected, (2)  $p = 1$ , (3)  $F$  has finite degree, (4)  $F$  preserves supports. If the functor  $F$  is finitary, has finite degree (and preserves supports), then  $F$  preserves uniformly continuous function, coarse functions, coarse equivalences, asymptotically Lipschitz functions, quas-isometries (and continuous functions). For many dimension functions we prove the formula  $\dim F^p X \leq \deg(F) \cdot \dim X$ . Using the injective envelopes, we also define a modification  $\check{d}_{FX}^p$  of the distance, which has some additional properties, for example, preserves isometries between distance spaces. The properties of the functors  $F^p$  and  $\check{F}^p$  are summed up in Theorems 25.1, 25.2, 26.1.

In the last three sections we will study the  $\ell^p$ -metrizations of some concrete functors with finite supports: of  $n$ th power, of free abelian (Boolean) group, and of free semilattice.

The results obtained in this paper can be considered as a development of the results of Shukel [29] and Radul [27] who studied the  $\ell^p$ -metrizations of normal functors for  $p = \infty$ . An axiomatic approach to metrization of functors was developed by Fedorchuk [19], see [32, §3.9].

## 2. SOME NOTATIONS AND CONVENTIONS

By  $\omega = \{0, 1, 2, \dots\}$  we denote the set of all finite ordinals. Each finite ordinal  $n \in \omega$  is identified with the set  $n := \{0, \dots, n-1\}$  of smaller ordinals. Cardinals are identified with the smallest ordinals of given cardinality.

For a set  $X$  and a cardinal  $\kappa$  we use the following notations:

- $\kappa^+$  is the smallest cardinal, which is larger than  $\kappa$ ;
- $|X|$  is the cardinality of  $X$ ;
- $[X]^\kappa := \{A \subseteq X : |A| = \kappa\}$ ;
- $[X]^{<\kappa} := \{A \subseteq X : |A| < \kappa\}$ ;
- $X^\kappa$  is the family of functions from  $\kappa$  to  $X$ ;
- $X^{<\omega} = \bigcup_{n \in \omega} X^n$  the family of all finite sequences of elements of  $X$ .

For a function  $f : X \rightarrow Y$  between sets and subsets  $A \subseteq X$ ,  $B \subseteq Y$  by  $f \upharpoonright A$  we denote the restriction of  $f$  to  $A$ , by  $f[A] := \{f(a) : a \in A\} \subseteq Y$  the image of the set  $A$  under the function  $f$ , and by  $f^{-1}[B] = \{x \in X : f(x) \in B\}$  the preimage of  $B$ . Such notations are convenient for handling functions on natural numbers  $n := \{0, \dots, n-1\}$ . In this case for a function  $f : n \rightarrow Y$  and element  $i \in n$  we have that  $f(i)$  is an element of  $Y$  but  $f[i]$  is the subset  $\{f(x) : x \in i\}$  of  $Y$ . For two sets  $A, B$  we denote by  $A \Delta B$  their symmetric difference  $(A \cup B) \setminus (A \cap B)$ .

We extend the basic arithmetic operations to the extended half line  $[0, \infty]$  letting  $x + \infty = \infty + x = \infty$  and  $\infty \cdot 0 = 0 = x - x$  for all  $x \in [0, \infty]$ , and

$$x \cdot \infty = \infty \cdot x = \infty^x = \infty - x = |x - \infty| = |\infty - x| = \infty$$

for any  $x \in (0, \infty]$ .

## 3. DISTANCE SPACES

In this section we present some basic definitions related to distance spaces.

We recall that a *distance space* is a pair  $(X, d_X)$  consisting of a set  $X$  and a *distance*  $d_X$  on  $X$ , which is a function  $d_X : X \times X \rightarrow [0, \infty]$  such that

- $d_X(x, x) = 0$ ,
- $d_X(x, y) = d_X(y, x)$ ,
- $d_X(x, z) \leq d_X(x, y) + d_X(y, z)$ ,

for any points  $x, y, z \in X$ .

A distance  $d_X$  on a set  $X$  is called

- an  $\infty$ -*metric* if  $d_X(x, y) > 0$  for any distinct points  $x, y \in X$ ;
- a *pseudometric* if  $d_X(x, y) < \infty$  for any points  $x, y \in X$ .

Each set  $X$  carries the  $\{0, \infty\}$ -valued  $\infty$ -metric  $d : X \times X \rightarrow \{0, \infty\}$  defined by

$$d(x, y) = \begin{cases} 0, & x = y; \\ \infty, & x \neq y. \end{cases}$$

An important example of a distance space is the extended real line  $\overline{\mathbb{R}} = [-\infty, +\infty]$ , endowed with the  $\infty$ -metric

$$d_{\overline{\mathbb{R}}}(x, y) = \begin{cases} |x - y|, & \text{if } x, y < \infty; \\ 0, & \text{if } x = y; \\ \infty, & \text{otherwise.} \end{cases}$$

For every set  $X$  the distance  $d_{\mathbb{R}}$  induces the sup-distance  $d_{\mathbb{R}^X}$  of the space  $\mathbb{R}^X$  of all functions from  $X$  to  $\mathbb{R}$ . The distance  $d_{\mathbb{R}^X}$  is defined by the formula

$$d_{\mathbb{R}^X}(f, g) = \sup_{x \in X} d_{\mathbb{R}}(f(x), g(x)).$$

Every distance  $d_X$  on a set  $X$  induces the *Hausdorff distance*

$$d_{HX}(A, B) = \begin{cases} \max\{\sup_{a \in A} \inf_{b \in B} d_X(a, b), \sup_{b \in B} \inf_{a \in A} d_X(a, b)\}, & \text{if } A \neq \emptyset \neq B, \\ 0, & \text{if } A = B, \\ \infty, & \text{otherwise,} \end{cases}$$

on the set  $HX$  of all subsets of  $X$ . It is a standard exercise to check that  $d_{HX}$  satisfies the three axioms of distance.

For a point  $x \in X$  of a distance space  $(X, d_X)$  and a number  $r \in [0, \infty]$  let

$$O(x; r) := \{y \in X : d_X(x, y) < r\} \text{ and } O[x; r] := \{y \in X : d_X(x, y) \leq r\}$$

be the open and closed balls of radius  $r$  around  $x$ , respectively.

For a subset  $A \subseteq X$  let

$$O[A; r] = \bigcup_{a \in A} O(a, r)$$

be the  $\varepsilon$ -neighborhood of  $A$  in  $(X, d_X)$ .

The family  $\{O(x; \infty) : x \in X\}$  is the partition of the distance space  $(X, d_X)$  into *pseudometric components* (observe that the restriction of the distance  $d_X$  to any pseudometric component  $O(x; \infty)$  is a pseudometric).

For a subset  $A$  of a distance space  $(X, d_X)$ , the numbers

$$\begin{aligned} \text{diam}(A) &:= \sup(\{0\} \cup \{d_X(x, y) : x, y \in A\}), \\ \bar{d}_X(A) &:= \sup(\{0\} \cup \{d_X(x, y) : x, y \in A, d_X(x, y) < \infty\}) \text{ and} \\ \underline{d}_X(A) &:= \inf(\{\infty\} \cup \{d_X(x, y) : x, y \in A, x \neq y\}) \end{aligned}$$

are called the *diameter*, the *real diameter* and the *separatedness number* of  $A$ .

A subset  $A$  of a distance space  $X$  is *bounded* if  $\text{diam}(A) < \infty$ .

A distance space  $(X, d_X)$  is called *separated* if  $\underline{d}_X(X) > 0$ .

Also for two subsets  $A, B \subseteq X$  we put

$$\underline{d}_X(A, B) = \inf(\{\infty\} \cup \{d_X(a, b) : a \in A, b \in B\}).$$

It is easy to see that  $\underline{d}_X(A, B) \leq d_{HX}(A, B)$  for any nonempty sets  $A, B \subseteq X$ .

For a family  $\mathcal{U}$  of subsets of a distance space  $X$  we put

$$\begin{aligned} \text{mesh}(\mathcal{U}) &:= \sup(\{0\} \cup \{\text{diam}(U) : U \in \mathcal{U}\}) \text{ and} \\ \underline{d}_X(\mathcal{U}) &= \inf(\{\infty\} \cup \{\underline{d}_X(U, V) : U, V \in \mathcal{U}, U \neq V\}). \end{aligned}$$

#### 4. FUNCTIONS BETWEEN DISTANCE SPACES


In this section we introduce some classes of functions between distance spaces.

A function  $f : X \rightarrow Y$  between distance spaces  $(X, d_X), (Y, d_Y)$  is called

- *an isometry* if  $d_Y(f(x), f(y)) = d_X(x, y)$  for every  $x, y \in X$ ;
- *non-expanding* if  $d_Y(f(x), f(y)) \leq d_X(x, y)$  for every  $x, y \in X$ ;

- *Lipschitz* if there exists a real number  $L$  such that  $d_Y(f(x), f(y)) \leq L \cdot d_X(x, y)$  for all  $x, y \in X$ ; the smallest such constant  $L$  is denoted by  $\text{Lip}(f)$  and called the *Lipschitz constant* of  $f$ ;
- *asymptotically Lipschitz* if there exist positive real numbers  $L, C$  such that  $d_Y(f(x), f(y)) \leq L \cdot d_X(x, y) + C$  for any  $x, y \in X$ ;
- *continuous* if for any  $x \in X$  and  $\varepsilon \in (0, \infty]$  there exists  $\delta \in (0, \infty]$  such that for any  $y \in X$  with  $d_X(x, y) \leq \delta$  we have  $d_X(f(x), f(y)) \leq \varepsilon$ ;
- *uniformly continuous* (or else *microform*) if for any  $\varepsilon \in (0, \infty]$  there exists  $\delta \in (0, \infty]$  such that for any  $x, y \in X$  with  $d_X(x, y) \leq \delta$  we have  $d_X(f(x), f(y)) \leq \varepsilon$ ;
- *coarse* (or else *macroform*) if for every  $\delta \in [0, \infty)$  there exists  $\varepsilon \in [0, \infty)$  such that for any  $x, y \in X$  with  $d_X(x, y) \leq \delta$  we have  $d_X(f(x), f(y)) \leq \varepsilon$ ;
- *duoform* if  $f$  is both microform and macroform;
- *bornologous* if for any bounded set  $B$  in  $X$  the image  $f(B)$  is bounded in  $Y$ ;
- a *coarse equivalence* if  $f$  is coarse and there exists a coarse map  $g : Y \rightarrow X$  such that  $\sup_{x \in X} d_X(x, f \circ g(x)) < \infty$  and  $\sup_{y \in Y} d_Y(y, f \circ g(y)) < \infty$ ;
- a *quasi-isometry* if  $f$  is an asymptotically Lipschitz map and there exists an asymptotically Lipschitz map  $g : Y \rightarrow X$  such that  $\sup_{x \in X} d_X(x, f \circ g(x)) < \infty$  and  $\sup_{y \in Y} d_Y(y, f \circ g(y)) < \infty$ .

The implications between these properties are described in the following diagram.



**Remark 4.1.** An isometry between distance spaces needs not be injective. On the other hand, an isometry between  $\infty$ -metric spaces is necessarily injective.

For a function  $f : X \rightarrow Y$  between distance spaces, its *continuity modulus* is the function  $\omega_f : (0, \infty] \rightarrow [0, \infty]$  defined by

$$\omega_f(\varepsilon) := \sup\{d_Y(f(x), f(y)) : x, y \in X \wedge d_X(x, y) < \varepsilon\} \text{ for } \varepsilon \in (0, \infty].$$

The definition implies that the continuity modulus  $\omega_f$  is a non-decreasing function from  $(0, \infty]$  to  $[0, \infty]$ .

Many properties of functions can be expressed via the continuity modulus.

In particular, a function  $f : X \rightarrow Y$  between distance spaces is

- non-expanding iff  $\omega_f(\varepsilon) \leq \varepsilon$  for all  $\varepsilon \in (0, \infty)$ ;
- Lipschitz iff there exists a real number  $L$  such that  $\omega_f(\varepsilon) \leq L \cdot \varepsilon$  for all  $\varepsilon \in (0, \infty)$ ;

- asymptotically Lipschitz iff there exist real numbers  $L, C$  such that  $\omega_f(\omega) \leq L \cdot \varepsilon + C$  for all  $\varepsilon \in (0, \infty)$ ;
- microform iff for any  $\varepsilon \in (0, \infty)$  there exists  $\delta \in (0, \infty)$  such that  $\omega_f(\delta) \leq \varepsilon$ ;
- macroform iff for any  $\delta \in (0, \infty)$  there exists  $\varepsilon \in (0, \infty)$  such that  $\omega_f(\delta) \leq \varepsilon$ .

The following two propositions are well-known for metric spaces, see [15, Lemma 1.1].

**Proposition 4.2.** *For any distance space  $(X, d_X)$ , the map*

$$i : X \rightarrow \bar{\mathbb{R}}^X, \quad i : x \mapsto d_X(x, \cdot),$$

*is an isometry.*

*Proof.* Given any point  $x \in X$  let  $d_x : X \rightarrow \bar{\mathbb{R}}$  be the function defined by  $d_x(y) = d_X(x, y)$  for  $y \in X$ . For any  $x, y, z \in X$ , the triangle inequality for the distance  $d_X$  implies  $d_{\bar{\mathbb{R}}}(d_x(z), d_y(z)) \leq d_X(x, y)$  and then  $d_{\bar{\mathbb{R}}^X}(d_x, d_y) \leq d_X(x, y)$ . On the other hand,  $d_{\bar{\mathbb{R}}^X}(d_x, d_y) \geq d_{\bar{\mathbb{R}}}(d_x(x), d_x(y)) = d_X(x, y)$ . Therefore,  $d_{\bar{\mathbb{R}}^X}(d_x, d_y) = d_X(x, y)$ , and the map  $i : X \rightarrow \bar{\mathbb{R}}^X, i : x \mapsto d_X(x, \cdot)$ , is an isometry.  $\square$

**Proposition 4.3.** *For any isometry  $i : X \rightarrow Y$  between distance spaces and any Lipschitz function  $f : X \rightarrow \bar{\mathbb{R}}$ , there exists a Lipschitz function  $\bar{f} : Y \rightarrow \bar{\mathbb{R}}$  such that  $\bar{f} \circ i = f$  and  $\text{Lip}(\bar{f}) = \text{Lip}(f)$ .*

*Proof.* For every  $y \in Y$ , consider the set  $O(y, \infty) = \{z \in Y : d_Y(z, y) < \infty\}$  and observe that  $\mathcal{O} = \{O(y, \infty) : y \in Y\}$  is a partition of  $Y$  into pseudometric subspaces (called the *pseudometric components* of  $Y$ ). For every  $B \in \mathcal{O}$  we will define a Lipschitz map  $\bar{f}_B : B \rightarrow \bar{\mathbb{R}}$  with Lipschitz constant  $\text{Lip}(\bar{f}_B) \leq \text{Lip}(f)$  such that  $\bar{f}_B \circ i|_B = f|_B$ .

The Lipschitz property of  $f$  implies that  $f[i[X] \cap B]$  is contained in one of the sets  $\mathbb{R}, -\infty$  or  $+\infty$ . If  $f[i[X] \cap B] \subseteq \{\pm\infty\}$ , then let  $\bar{f}_B : B \rightarrow \{\pm\infty\}$  be a constant map. It remains to consider the case  $f[i[X] \cap B] \subseteq \mathbb{R}$ . If  $i[X] \cap B$  is empty, then let  $\bar{f}_B : B \rightarrow \mathbb{R}$  be any (constant) map. If  $i[X] \cap B$  is not empty, then define a map  $\bar{f}_B : B \rightarrow \mathbb{R}$  by the explicit formula

$$\bar{f}_B(y) = \inf_{x \in i^{-1}[B]} (f(x) + \text{Lip}(f) \cdot d_X(i(x), y)) \quad \text{for any } y \in B.$$

It is easy to see that  $\bar{f}_B(i(x)) = f(x)$  for any  $x \in i^{-1}[B]$ . Repeating the argument of the proof of Lemma 1.1 in [15], we can show that the function  $\bar{f}_B : B \rightarrow \bar{\mathbb{R}}$  is well-defined, Lipschitz, and has Lipschitz constant  $\text{Lip}(\bar{f}_B) \leq \text{Lip}(f)$ . Finally, define the function  $\bar{f} : Y \rightarrow \bar{\mathbb{R}}$  letting  $\bar{f}|_B = \bar{f}_B$  for every  $B \in \mathcal{O}$ , and observe that  $\bar{f}$  has the desired property.  $\square$

## 5. THE INJECTIVE ENVELOPE OF A DISTANCE SPACE

In this section we extend the classical Isbell construction [24] (see also [16], [22], [26, §3]) of the injective envelope of a metric space to the category of distance spaces. Injective envelopes will be applied in Section 26 for defining the  $\ell^p$ -metrization of functors.

For a distance space  $X$  with distance  $d_X$ , let  $\Delta X$  be the set of all functions  $f : X \rightarrow \bar{\mathbb{R}}$  such that

$$f(x) + f(y) \geq d_X(x, y) \quad \text{for all } x, y \in X.$$

For  $x = y$  we have  $0 = d_X(x, x) \leq 2 \cdot f(x)$ , which means that functions  $f \in \Delta X$  take their values in the extended half-line  $\bar{\mathbb{R}}_+ = [0, \infty]$ .

The space  $\Delta X$  is endowed with the  $\infty$ -metric  $d_{\Delta X}$  inherited from  $\bar{\mathbb{R}}^X$ :

$$d_{\Delta X}(f, g) = \sup_{x \in X} d_{\bar{\mathbb{R}}}(f(x), g(x)) \quad \text{for } f, g \in \Delta X.$$

The set  $\Delta X$  also carries a natural partial order  $\leq$  defined by  $f \leq g$  iff  $f(x) \leq g(x)$  for all  $x \in X$ . For two function  $f, g \in \Delta X$  we write  $f < g$  if  $f \leq g$  and  $f \neq g$ .

A function  $f \in \Delta X$  is called *extremal* if  $f$  is a minimal element of the partially ordered set  $(\Delta X, \leq)$ . Let  $EX$  be the distance subspace of  $\Delta X$  consisting of extremal functions. By  $d_{EX}$  we denote the sup-distance of the space  $EX$ . The distance  $d_{EX}$  is an  $\infty$ -metric, being a restriction of the  $\infty$ -metric  $d_{\mathbb{R}X}$ .

The distance space  $EX$  is called the *injective envelope* of the distance space  $X$ . Now we establish some important properties of injective envelopes, generalizing well-known properties of injective envelopes of metric spaces.

**Lemma 5.1.** *For every  $z \in X$ , the function*

$$d_z : X \rightarrow [0, \infty], \quad d_z : x \mapsto d_X(x, z),$$

*is extremal.*

*Proof.* The triangle inequality for the distance  $d_X$  guarantees that the function  $d_z$  belongs to  $\Delta X$ . Assuming that  $d_z$  is not extremal, we can find a function  $f \leq d_z$  in  $\Delta X$  such that  $f(x) < d_z(x)$  for some  $x \in X$ . Then the number  $f(x)$  is finite,  $f(z) \leq d_z(z) = 0$  and hence

$$d_z(x) = d_X(x, z) \leq f(x) + f(z) = f(x) + 0 = f(x) < d_z(x),$$

which is a desired contradiction.  $\square$

**Lemma 5.2.** *The function  $e_X : X \rightarrow EX$ ,  $e_X : x \mapsto d_x$ , is a well-defined isometry between the distance spaces  $X$  and  $EX$ .*

*Proof.* By Lemma 5.1, the function  $e_X$  is well-defined. For any points  $x, y, z \in X$ , by the triangle inequality, we have

$$d_x(z) = d_X(x, z) \leq d_X(x, y) + d_X(y, z) = d_x(y) + d_y(z)$$

and hence  $d_{\mathbb{R}}(d_x(z), d_y(z)) \leq d_X(x, y)$ . Taking the supremum by  $z \in X$ , we obtain

$$d_{EX}(d_x, d_y) = \sup_{z \in X} d_{\mathbb{R}}(d_x(z), d_y(z)) \leq d_X(x, y),$$

which means that the map  $e_X : X \rightarrow EX$  is non-expanding.

On the other hand,

$$d_X(x, y) = d_{\mathbb{R}}(d_x(x), d_y(x)) \leq \sup_{z \in X} d_{\mathbb{R}}(d_x(z), d_y(z)) = d_{EX}(d_x, d_y)$$

witnessing that the map  $e_X : X \rightarrow EX$ , is an isometry.  $\square$

**Lemma 5.3.** *There exists a non-expanding map  $p : \Delta X \rightarrow EX$  such that  $p(f) \leq f$  for every  $f \in \Delta X$ .*

*Proof.* A subset  $B \subseteq X$  is called  $\infty$ -bounded if  $d_X(a, b) < \infty$  for any points  $a, b \in B$ . Denote by  $\mathcal{B}_\infty(X)$  the family of all  $\infty$ -bounded sets in  $X$ .

For any  $\infty$ -bounded set  $A \in \mathcal{B}_\infty(X)$ , consider the subset

$$\Delta_A X = \{f \in \Delta X : \infty \notin f[A], \quad f[X \setminus A] \subseteq \{\infty\}\}.$$

**Claim 5.4.**  $\Delta X = \bigcup_{A \in \mathcal{B}_\infty(X)} \Delta_A X$ .



*Proof.* Given any function  $f \in \Delta X$ , consider the set  $A = f^{-1}[\mathbb{R}]$ . We claim that  $A \in \mathcal{B}_\infty(X)$ . In the opposite case, we would find two points  $a, b \in A$  with  $d_X(a, b) = \infty$  and obtain the contradiction:

$$\infty > f(a) + f(b) \geq d_X(a, b) = \infty.$$

The definition of the set  $A$  ensures that  $f \in \Delta_A X$ .  $\square$

It is easy to see that for any distinct sets  $A, B \in \mathcal{B}_\infty(X)$  and any functions  $f \in \Delta_A X$ ,  $g \in \Delta_B X$ , we have  $d_{\Delta X}(f, g) = \infty$ . Therefore, it suffices for any set  $A \in \mathcal{B}_\infty(X)$  to construct a non-expanding function  $p_A : \Delta_A X \rightarrow EX$  such that  $p_A(f) \leq f$  for every  $f \in \Delta_A X$ .

If the set  $A$  is empty, then  $\Delta_A X$  is a singleton and any map  $p_A : \Delta_A X \rightarrow EX$  is non-expanding and has  $p_A(f) \leq f$  for the unique element  $f \in \{\infty\}^X = \Delta_A X$ .

So, we assume that  $A$  is not empty. In this case we can fix a point  $a \in A$  and conclude that  $A \subseteq B$  where  $B = \{x \in X : d_X(x, a) < \infty\}$ .

**Claim 5.5.** *There exists a non-expanding map  $q_A : \Delta_A X \rightarrow \Delta_B X$  such that  $q_A(f) \leq f$  for any  $f \in \Delta_A X$ .*

*Proof.* Consider the map  $q_A : \Delta_A X \rightarrow \Delta_B X$  assigning to each function  $f \in \Delta_A X$  the function  $g : X \rightarrow \mathbb{R}$  defined by

$$g(x) = \begin{cases} f(a) + d_X(a, x) & \text{if } x \in B \setminus A; \\ f(x) & \text{otherwise.} \end{cases}$$

We claim that  $g \in \Delta X$ . Given any points  $x, y \in X$ , we should check that  $g(x) + g(y) \geq d_X(x, y)$ . If  $x$  or  $y$  do not belong to  $B$ , then  $g(x) + g(y) = \infty \geq d_X(x, y)$  and we are done. So, assume that  $x, y \in B$ . If  $x, y \notin B \setminus A$ , then  $g(x) + g(y) = f(x) + f(y) \geq d_X(x, y)$ . If  $x, y \in A$ , then  $g(x) + g(y) = 2f(a) + d_X(a, x) + d_X(a, y) \geq d_X(x, y)$ . If  $x \in B \setminus A$  and  $y \in A$ , then  $g(x) + g(y) = d_X(a, x) + f(a) + f(y) \geq d_X(a, x) + d_X(a, y) \geq d_X(x, y)$ . By analogy we can treat the case  $x \in A$  and  $y \in B \setminus A$ . Therefore,  $g \in \Delta X$ . The definition of  $g$  guarantees that  $g \in \Delta_B X$  and  $g \leq f$ .

It remains to show that the map  $q_A$  is non-expanding. Choose arbitrary functions  $f, f' \in \Delta_A X$ , and consider the functions  $g = q_A(f)$  and  $g' = q_A(f')$ . If  $x \notin B \setminus A$ , then  $d_{\mathbb{R}}(g(x), g'(x)) = d_{\mathbb{R}}(f(x), f'(x)) \leq d_{\Delta X}(f, f')$ . If  $x \in B \setminus A$ , then

$$d_{\mathbb{R}}(g(x), g'(x)) = d_{\mathbb{R}}(f(a) + d_X(a, x), f'(a) + d_X(a, x)) \leq d_{\mathbb{R}}(f(a), f'(a)) \leq d_{\Delta X}(f, f').$$

$\square$

Let  $E_B X := EX \cap \Delta_B X$ .

**Claim 5.6.** *A function  $f : X \rightarrow \mathbb{R}$  belongs to  $E_B X$  if and only if  $f(X \setminus B) \subseteq \{\infty\}$  and  $f(x) = \sup_{y \in B} (d_X(x, y) - f(y)) < \infty$  for every  $x \in B$ .*

*Proof.* To prove the “only if” part, assume that  $f \in E_B X$ . Then  $f \in \Delta_B X$  and hence  $f(X \setminus B) \subseteq \{\infty\}$  and  $\infty > f(x) \geq \sup_{y \in B} (d_X(x, y) - f(y))$  for all  $x \in B$ . Assuming that  $f(x) \neq \sup_{y \in B} (d_X(x, y) - f(y))$  for some  $x \in B$ , we conclude that  $f(x) > \sup_{y \in B} (d_X(x, y) - f(y))$ . If  $s := \sup_{y \in B} (d_X(x, y) - f(y)) < 0$ , then  $s + f(y) \geq d_X(x, y)$  for every  $y \in X$  and the function  $g = f + \frac{1}{2}s < f$  belongs to  $\Delta_B X$  witnessing that  $f$  is not extremal. This contradiction shows that  $s \geq 0$ .

Consider the function  $g : X \rightarrow \mathbb{R}$  such that  $g \upharpoonright X \setminus \{x\} = f \upharpoonright X \setminus \{x\}$  and  $g(x) = s \geq 0$ . Then  $g < f$ . We claim that  $g \in \Delta_B X$ . Given two distinct points  $y, z \in X$  we should check that  $g(y) + g(z) \geq d_X(y, z)$ . If  $y, z \in X \setminus \{x\}$ , then this follows from the equality

$g|_{X \setminus \{x\}} = f|_{X \setminus \{x\}}$ . If  $y = z$ , then  $g(y) + g(z) = 2s \geq 0 = d_X(y, z)$ . If  $y = x$  and  $z \neq x$ , then  $g(y) + g(z) \geq d_X(y, z) - f(z) + f(z) = d_X(y, z)$ . By analogy we can treat the case  $z = x$  and  $y \neq x$ . The function  $g < f$  witnesses that  $f \notin E_B(X)$ , which is a contradiction showing that  $f(x) = \sup_{y \in B} (d_X(x, y) - f(y))$  for all  $x \in B$ .

Next, we prove the ‘‘if’’ part. Choose any function  $f : X \rightarrow \bar{\mathbb{R}}$  such that  $f(X \setminus B) \subseteq \{\infty\}$  and  $f(x) = \sup_{y \in B} (d_X(x, y) - f(y)) < \infty$  for all  $x \in B$ . Then  $f(x) + f(y) \geq d_X(x, y)$  for all  $x \in X$  and hence  $f \in \Delta_B X$ . Assuming that  $f \notin E_B X$ , we can find a function  $g \leq f$  in  $\Delta X$  such that  $g(x) < f(x)$  for some  $x \in X$ . Since  $\infty > f(a) + g(x) \geq g(a) + g(x) \geq d_X(a, x)$ , the point  $x$  belongs to the pseudometric component  $B$  of  $X$ . The inclusion  $g \in \Delta X$  and the inequality  $g \leq f$  imply

$$\sup_{y \in B} (d_X(x, y) - f(y)) \leq \sup_{y \in B} (d_X(x, y) - g(y)) \leq g(x) < f(x),$$

which contradicts our assumption. This contradiction shows that  $f \in E_B X$ .  $\square$

**Claim 5.7.** *There exists a non-expanding map  $p_B : \Delta_B X \rightarrow E_B X$  such that  $p_B(f) \leq f$  for any  $f \in \Delta_B X$ .*

*Proof.* We modify the argument of Dress [16, §1.9], following the lines of the proof of Proposition 2.1 in [26]. For every  $f \in \Delta_B X$ , define the function  $f^* : X \rightarrow \bar{\mathbb{R}}$  letting  $f^*(x) = \sup_{z \in B} (d_X(x, z) - f(z))$  for  $x \in B$  and  $f^*(x) = \infty$  for  $x \in X \setminus B$ . The inclusion  $f \in \Delta_B X$  implies  $f^* \leq f$ .

For every pair of points  $x, y \in X$ , the definition of  $f^*$  implies  $f(x) + f^*(y) \geq d_X(x, y)$  and  $f^*(x) + f(y) \geq d_X(x, y)$ . Then the function  $q(f) = \frac{1}{2}(f + f^*)$  belongs to  $\Delta_B X$  and  $q(f) \leq f$ .

For any functions  $f, g \in \Delta_B X$  and  $x \in B$ , we have

$$g^*(x) = \sup_{z \in B} (d_X(x, z) - f(z) + f(z) - g(z)) \leq f^*(x) + d_{\Delta X}(f, g)$$

and hence  $d_{\bar{\mathbb{R}} X}(f^*, g^*) \leq d_{\Delta X}(f, g)$  and finally

$$d_{\Delta X}(q(f), q(g)) \leq \frac{1}{2}d_{\Delta X}(f, g) + \frac{1}{2}d_{\Delta X}(f^*, g^*) \leq d_{\Delta X}(f, g).$$

Iterating the map  $q$ , we obtain for every  $f \in \Delta_B X$  a decreasing function sequence  $g(f) \geq q(f^2) \geq \dots$  in  $\Delta_B X$ , then define  $p_B(f)$  as the pointwise limit. Clearly,  $p_B(f) \in \Delta_B X$ ,  $p_B(f) \leq f$  and the map  $p_B : \Delta_B X \rightarrow \Delta_B X$  is non-expanding. For all  $n \geq 1$ ,  $p_B(f) \leq q^n(f)$  and hence  $p_B(f)^* \geq q^n(f)^*$ , so

$$0 \leq p_B(f) - p_B(f)^* \leq q^n(f) - q^n(f)^* = 2(q^n(f) - q^{n+1}(f)).$$

As  $n \rightarrow \infty$ , the last term converges pointwise to 0, thus  $p_B(f)^* = p_B(f)$  and therefore,  $p_B(f) \in E_B(X)$  by Claim 5.6.  $\square$

Finally, define a non-expanding map  $p_A : \Delta_A X \rightarrow E_B X$  as  $p_A = p_B \circ q_A$ , where  $q_A, p_B$  are the non-expanding maps from Claims 5.5 and 5.7, respectively. The properties of these maps ensure that  $p_A(f) = p_B(q_A(f)) \leq q_A(f) \leq f$  for every  $f \in \Delta_A X$ .  $\square$

An isometry  $i : X \rightarrow Y$  between distance spaces is called *Isbell-injective* if for any isometry  $j : X \rightarrow Z$  to a distance space  $Z$ , there exists a non-expanding map  $f : Z \rightarrow Y$  such that  $i = f \circ j$ .

**Lemma 5.8.** *For any distance space  $X$  the isometry  $e_X : X \rightarrow \Delta X$  is Isbell-injective.*

*Proof.* Given any isometry  $j : X \rightarrow Z$  to a distance space  $Z$ , consider the map  $f : Z \rightarrow \Delta X$  assigning to each point  $z \in Z$  the function  $f_z : X \rightarrow \overline{\mathbb{R}}$  defined by  $f_z(x) = d_Z(z, j(x))$  for  $x \in X$ . Observe that for every  $x, y \in X$  we have

$$f_z(x) + f_z(y) = d_Z(z, j(x)) + d_Z(z, j(y)) \geq d_Z(j(x), j(y)) = d_X(x, y),$$

and hence  $f_z \in \Delta X$ .

For any  $x \in X$ , the function  $f_{j(x)}$  coincides with the function  $d_x$ . Indeed, for any  $y \in X$  we have  $f_{j(x)}(y) = d_Z(j(x), j(y)) = d_X(x, y) = d_x(y)$ . Consequently,  $f \circ j = \mathbf{e}_X : X \rightarrow EX$ .

For any points  $y, z \in Z$  and any  $x \in X$  the triangle inequality for the distance  $d_Z$  ensures that

$$d_{\overline{\mathbb{R}}}(f_y(x), f_z(x)) = |d_Z(y, j(x)) - d_Z(z, j(x))| \leq d_Z(y, z),$$

and hence

$$d_{\overline{\mathbb{R}^X}}(f_y, f_x) = \sup_{x \in X} d_{\overline{\mathbb{R}}}(f_y(x), f_x(x)) \leq d_Z(y, z),$$

which means that the map  $f : Z \rightarrow \Delta X$  is non-expanding.  $\square$

**Lemma 5.9.** *For any distance space  $X$ , the isometry  $\mathbf{e}_X : X \rightarrow EX$  is Isbell-injective.*

*Proof.* By Lemma 5.3, there exists a non-expanding map  $p : \Delta X \rightarrow EX$  such that  $p(f) = f$  for any  $f \in EX$ . Consequently,  $p \circ \mathbf{e}_X = \mathbf{e}_X$ .

To show that the isometry  $\mathbf{e}_X : X \rightarrow EX$  is Isbell-injective, take any isometry  $j : X \rightarrow Z$  to a distance space  $Z$ . By Lemma 5.8, there exists a non-expanding map  $f : Z \rightarrow \Delta X$  such that  $f \circ j = \mathbf{e}_X$ . Then  $g = p \circ f : Z \rightarrow EX$  is a non-expanding map such that  $g \circ j = p \circ f \circ j = p \circ \mathbf{e}_X = \mathbf{e}_X$ .  $\square$

**Lemma 5.10.** *Let  $X$  be a distance space and  $f : EX \rightarrow EX$  be a non-expanding map such that  $f \circ \mathbf{e}_X = \mathbf{e}_X$ . Then  $f$  is the identity map of  $EX$ .*

*Proof.* Given any function  $\alpha \in EX$  consider the function  $\beta = f(\alpha) \in EX$ . Consider the partition  $\{O(x, \infty) : x \in X\}$  into pseudometric components. Claim 5.5 implies that there are pseudometric components  $A, B$  of  $X$  such that  $\infty \notin \alpha[A] \cup \beta[B]$  and  $\alpha[X \setminus A] = \beta[X \setminus B] = \{\infty\}$ . We claim that  $A = B$ . Take any point  $a \in A$ . By Claim 5.6,

$$\infty > \alpha(a) = \sup_{x \in A} (d_a(x) - \alpha(x)) = d_{EX}(d_a, \alpha) \geq d_{EX}(f(d_a), f(\alpha)) = d_{EX}(d_a, \beta) \geq \beta(a)$$

and hence  $a \in B$  and  $A = B$ . Then  $\alpha \upharpoonright X \setminus A = \beta \upharpoonright X \setminus A$ . On the other hand, for every  $a \in A$ , by Claim 5.6,

$$\beta(a) = \sup_{x \in A} (d_a(x) - \beta(x)) = d_{EX}(d_a, \beta) = d_{EX}(f(d_a), f(\alpha)) \leq d_{EX}(d_a, \alpha) \leq \alpha(a),$$

so  $\beta = \alpha$  by the minimality of  $\alpha \in EX$ .  $\square$

An isometry  $i : X \rightarrow Y$  between distance spaces is called *tight* if any non-expanding map  $f : Y \rightarrow Z$  to a distance space  $Z$  is an isometry iff  $f \circ i$  is an isometry.

**Lemma 5.11.** *For any distance space  $X$ , the isometry  $\mathbf{e}_X : X \rightarrow EX$  is tight.*

*Proof.* Let  $f : EX \rightarrow Z$  be a non-expanding map to a distance space  $Z$  such that  $f \circ \mathbf{e}_X$  is an isometry. By Lemma 5.9, there exists a non-expanding map  $g : Z \rightarrow EX$  such that  $g \circ f \circ \mathbf{e}_X = \mathbf{e}_X$ . By Lemma 5.10,  $g \circ f$  is the identity map of  $EX$ . Then for any  $\alpha, \beta \in EX$  we have

$$d_{EX}(\alpha, \beta) = d_{EX}(g(f(\alpha)), g(f(\beta))) \leq d_Z(f(\alpha), f(\beta)) \leq d_{EX}(\alpha, \beta)$$

and hence  $d_Z(f(\alpha), f(\beta)) = d_{EX}(\alpha, \beta)$ , which means that  $f$  is an isometry.  $\square$

An isometry  $i : X \rightarrow Y$  between two distance spaces is called an *injective envelope* of  $X$  if it is tight and Isbell-injective.

Lemmas 5.9 and 5.11 imply the following main result of this section.

**Theorem 5.12.** *For any distance space  $X$  the map  $e_X : X \rightarrow EX$  is an injective envelope of  $X$ .*

The following proposition says that the injective envelope is unique up to an isometry.

**Proposition 5.13.** *For any injective envelope  $i : X \rightarrow Y$  of a distance space  $X$ , there exists a surjective isometry  $f : Y \rightarrow EX$  such that  $f \circ i = e_X$ .*

*Proof.* Since both isometries  $e_X : X \rightarrow EX$  and  $i : X \rightarrow Y$  are Isbell-injective, there are nonexpanding maps  $f : Y \rightarrow EX$  and  $g : EX \rightarrow Y$  such that  $f \circ i = e_X$  and  $g \circ e_X = i$ . Since the isometries  $i$  and  $e_X$  are tight, the non-expanding maps  $f, g$  are isometries. Observe that  $f \circ g \circ e_X = f \circ i = e_X$ . By Lemma 5.10,  $f \circ g$  is the identity map of  $EX$ . Then the isometry map  $f$  is surjective and  $g$  is injective.  $\square$

**Proposition 5.14.** *For any isometry  $i : X \rightarrow Y$  between distance spaces, there exists an isometry  $f : EX \rightarrow EY$  such that  $f \circ e_X = e_Y \circ i$ .*

*Proof.* By Lemma 5.9, there exists a non-expanding map  $f : EX \rightarrow EY$  such that  $f \circ e_X = e_Y \circ i$ . By Lemma 5.9, there exists a non-expanding map  $g : EY \rightarrow EX$  such that  $g \circ e_Y \circ i = e_X$ . Since  $g \circ f \circ e_X = g \circ e_Y \circ i = e_X$ , we can apply Lemma 5.10 and conclude that  $g \circ f$  is the identity map of  $EX$ . Then the non-expanding map  $f$  is an isometry.  $\square$

A distance space  $X$  is defined to be *injective* if for any isometry  $i : X \rightarrow Y$  to a distance space  $Y$  there exists a non-expanding map  $f : Y \rightarrow X$  such that  $f \circ i$  is the identity map of  $X$ .

**Proposition 5.15.** *A distance space  $X$  is injective if and only if there exists a non-expanding map  $r : EX \rightarrow X$  such that  $r \circ e_X$  is the identity map of  $X$ .*

*Proof.* The “only if” part follows from the definition of an injective distance space and Lemma 5.2.

To prove the “if” part, assume that  $X$  admits a non-expanding map  $r : EX \rightarrow X$  such that  $r \circ e_X$  is the identity map of  $X$ . Given any isometry  $i : X \rightarrow Y$ , apply Lemma 5.9 and find a non-expanding map  $g : Y \rightarrow EX$  such that  $g \circ i = e_X$ . Then the map  $f = r \circ g : Y \rightarrow X$  is non-expanding and for every  $x \in X$

$$f \circ i(x) = r \circ g \circ i(x) = r \circ e_X(x) = x.$$

$\square$

**Corollary 5.16.** *For any distance space  $X$  the  $\infty$ -metric space  $EX$  is injective.*

*Proof.* By Lemma 5.2, the maps  $e_X : X \rightarrow EX$  and  $e_{EX} : EX \rightarrow EEX$  are isometries. By Lemma 5.9, there exists a non-expanding map  $f : EEX \rightarrow EX$  such that  $f \circ e_{EX} \circ e_X = e_X$ . By Lemma 5.10, the composition  $f \circ e_{EX}$  is the identity map of  $EX$ . By Proposition 5.15, the distance space  $EX$  is injective.  $\square$

**Proposition 5.17.** *For any Lipschitz map  $f : X \rightarrow Y$  between distance spaces, there exists a Lipschitz map  $\tilde{f} : EX \rightarrow EY$  between their injective envelopes such that  $\tilde{f} \circ e_X = e_Y \circ f$  and  $\text{Lip}(\tilde{f}) \leq \text{Lip}(f)$ .*

*Proof.* By Proposition 4.2, there exists an isometry  $i : Y \rightarrow \bar{\mathbb{R}}^Y$ . By Proposition 4.3, for the Lipschitz map  $i \circ f : X \rightarrow \bar{\mathbb{R}}^Y$  and the isometry  $e_X : X \rightarrow EX$ , there exists a Lipschitz map  $g : EX \rightarrow \bar{\mathbb{R}}^Y$  such that  $g \circ e_X = i$  and  $\text{Lip}(g) \leq \text{Lip}(i \circ f) = \text{Lip}(f)$ . By Lemma 5.9, for the isometry  $i : Y \rightarrow \bar{\mathbb{R}}^Y$ , there exists a non-expanding map  $h : \bar{\mathbb{R}}^Y \rightarrow EY$  such that  $h \circ i = e_Y$ . Then  $\bar{f} = h \circ g : EX \rightarrow EY$  is a Lipschitz map with Lipschitz constant  $\text{Lip}(\bar{f}) \leq \text{Lip}(h) \cdot \text{Lip}(g) = \text{Lip}(g) \leq \text{Lip}(f)$  such that

$$\bar{f} \circ e_X = h \circ g \circ e_X = h \circ i = e_Y.$$

□

## 6. PRELIMINARIES ON FUNCTORS

All functors considered in this paper are covariant.

We say that a functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  has *finite supports* (or else,  $F$  is *finitely supported*) if for any set  $X$  and element  $a \in FX$  there exists a finite subset  $A \subseteq X$  such that  $a \in F[A; X]$  where  $F[A; X] := Fi_{A,X}[FA] \subseteq FX$  and  $i_{A,X} : A \rightarrow X$  is the identity embedding.

For a set  $X$  and an element  $a \in FX$  its *support* is defined as

$$\text{supp}(a) = \bigcap \{A \subseteq X : a \in F[A; X]\}.$$

The following helpful result is proved in [6].

**Proposition 6.1.** *If  $\text{supp}(a)$  is not empty, then  $a \in F[\text{supp}(a); X]$ . If  $\text{supp}(a)$  is empty, then  $a \in F[A; X]$  for any non-empty subset  $A \subseteq X$ .*

**Corollary 6.2.** *For any function  $f : X \rightarrow Y$  between sets with  $|f[X]| > 1$  and any  $a \in FX$  we have  $\text{supp}(Ff(a)) \subseteq f(\text{supp}(a))$ .*

*Proof.* If the set  $S := \text{supp}(a)$  is not empty, then  $a \in F[S; X]$  and there exists  $a' \in FS$  such that  $a = Fi_{S,X}(a')$ . Then  $Ff(a) = Ff \circ Fi_{S,X}(a') = F(f \circ i_{S,X})(a') \in F[f(S); X]$  and hence  $\text{supp}(Ff(a)) \subseteq f(S) = f(\text{supp}(a))$ .

Now assume that  $\text{supp}(a) = \emptyset$ . Since  $|f[X]| > 1$ , we can find two non-empty sets  $A, B \subseteq X$  with  $f[A] \cap f[B] = \emptyset$ . By Proposition 6.1,  $a \in F[A; X] \cap F[B; X]$ , which implies  $Fa \in F[f[A]; Y] \cap F[f[B]; Y]$ . Then  $\text{supp}(Ff(a)) \subseteq f[A] \cap f[B] = \emptyset = f(\text{supp}(a))$ . □

For a set  $X$  and a cardinal  $n \leq \omega$ , put  $F_n X = \{a \in FX : |\text{supp}(a)| \leq n\}$ . If  $FX = F_n X$  for any set  $X$ , then we say that the functor  $F$  has degree  $\leq n$ . The smallest number  $n$  with this property is called *the degree of the functor  $F$*  and is denoted by  $\text{deg}(F)$ . If the cardinal  $\text{deg}(F)$  is finite, then  $F$  is called a *functor of finite degree*. More information on functors (of finite degree) can be found in [32].

We will say that a functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$

- *preserves supports* if for any function  $f : X \rightarrow Y$  and any element  $a \in FX$  we get  $\text{supp}(Ff(a)) = f(\text{supp}(a))$ ;
- is *finitary* if for any finite set  $X$  the set  $FX$  is finite;
- *preserves singletons* if for the singleton  $1 = \{0\}$  the set  $F(1)$  is a singleton.

7. INTRODUCING THE DISTANCE  $d_{FX}^p$ 

Let  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  be a functor with finite supports. Fix a number  $p \in [1, \infty]$ .

For each distance  $d_X$  on a set  $X$  we define the distance  $d_{FX}^p$  on  $FX$  as follows. Let  $X^{<\omega} = \bigcup_{n \in \omega} X^n$ . The elements of  $X^{<\omega}$  are functions  $f : n \rightarrow X$  defined on finite ordinals  $n = \{0, \dots, n-1\}$ . On each finite power  $X^n$  we consider the  $\ell^p$ -distance  $d_{X^n}^p$  defined by the formula

$$d_{X^n}^p(f, g) = \begin{cases} \left( \sum_{i \in n} d_X(f(i), g(i))^p \right)^{\frac{1}{p}} & \text{if } p < \infty; \\ \max_{i \in n} d_X(f(i), g(i)) & \text{if } p = \infty. \end{cases}$$

The distances  $d_{X^n}^p$ ,  $n \in \omega$ , determine the distance  $d_{X^{<\omega}}^p$  on the set  $X^{<\omega} = \bigcup_{n \in \omega} X^n$ , defined by the formula

$$d_{X^{<\omega}}^p(f, g) = \begin{cases} d_{X^n}^p(f, g) & \text{if } f, g \in X^n \text{ for some } n \in \omega, \\ \infty, & \text{otherwise.} \end{cases}$$

For two elements  $a, b \in FX$  let  $L_{FX}(a, b)$  be the set of all finite sequences  $((a_0, f_0, g_0), \dots, (a_l, f_l, g_l))$  of triples

$$(a_0, f_0, g_0), \dots, (a_l, f_l, g_l) \in \bigcup_{n \in \omega} Fn \times X^n \times X^n$$

such that  $a = Ff_0(a_0)$ ,  $b = Fg_l(a_l)$  and  $Fg_i(a_i) = Ff_{i+1}(a_{i+1})$  for all  $0 \leq i < l$ . Elements of the set  $L_{FX}(a, b)$  are called  $(a, b)$ -linking chains.

For an  $(a, b)$ -linking chain  $s = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  put

$$\Sigma d_X^p(s) := \sum_{i=0}^l d_{X^{<\omega}}^p(f_i, g_i).$$

Define the distance  $d_{FX}^p : FX \times FX \rightarrow [0, \infty]$  letting

$$d_{FX}^p(a, b) = \inf (\{\infty\} \cup \{\Sigma d_X^p(s) : s \in L_{FX}(a, b)\}).$$

The correspondence  $F^p : (X, d_X) \mapsto (FX, d_{FX}^p)$  determines a functor  $F^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$  in the category  $\mathbf{Dist}$  of distance spaces, which is a lifting of the functor  $F$  with respect to the forgetful functor  $\mathbf{Dist} \rightarrow \mathbf{Set}$ . This lifting will be called the  $\ell^p$ -metrization of the functor  $F$ . In the remaining part of this paper we study the properties of the functor  $F^p$  and its modification  $\check{F}^p$ .

From now on, we assume that  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  is a functor with finite supports,  $X = (X, d_X)$  is a distance space, and  $p \in [1, \infty]$ .

8. THE DISTANCE  $d_{FX}^p$  BETWEEN ELEMENTS WITH SMALL SUPPORTS

In this section we evaluate the distance between elements of  $FX$ , supported by the empty set.

**Theorem 8.1.** *For any elements  $a, b \in FX$  with  $|\text{supp}(a) \cup \text{supp}(b)| \leq 1$ , we get*

$$d_{FX}^p(a, b) = \begin{cases} 0, & \text{if } a = b, \\ \infty, & \text{if } a \neq b. \end{cases}$$

*Proof.* It is clear that  $d_{FX}^p(a, b) = 0$  if  $a = b$ . So, we assume that  $a \neq b$ . In this case we prove that  $d_{FX}^p(a, b) = \infty$ . Separately we consider the cases of empty and non-empty set  $X$ .

First we consider the case of  $X = \emptyset$ . Assuming that  $d_{FX}^p(a, b) < \infty$ , we can find an  $(a, b)$ -linking chain  $s = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\Sigma d_X^p(s) < \infty$ . For every  $i \in \{0, \dots, l\}$  find a finite ordinal  $n_i$  such that  $f_i, g_i \in X^{n_i}$ . Then  $X = \emptyset \neq X^{n_i}$  implies that  $n_i = 0$  for all  $i \in \{0, \dots, l\}$  and hence  $f_i = g_i$  (as the set  $X^0$  is a singleton containing the unique function  $\emptyset \rightarrow \emptyset = X$ ). Then

$$a = Ff_0(a_0) = Fg_0(a_0) = Ff_1(a_1) = Fg_1(a_1) = \dots = Ff_l(a_l) = Fg_l(a_l) = b,$$

which is a desired contradiction.

Now we assume that the set  $X$  is not empty. In this case we can find a singleton  $S \subseteq X$  containing the union  $\text{supp}(a) \cup \text{supp}(b)$ . By Proposition 6.1,  $a, b \in F[S; X]$ . So, we can find (distinct) elements  $a', b' \in FS$  such that  $a = Fi_{S,X}(a')$  and  $b = Fi_{S,X}(b')$ . Here by  $i_{S,X} : S \rightarrow X$  we denote the identity embedding of  $S$  into  $X$ . Let  $f : X \rightarrow S$  be the unique constant map and observe that  $f \circ i_{S,X}$  is the identity map of  $S$ . Consequently,  $a' = Ff \circ Fi_{S,X}(a') = Ff(a)$  and  $b' = Ff(b)$ .

Assuming that  $d_{FX}^p(a, b) < \infty$ , we can find an  $(a, b)$ -linking chain  $s = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\Sigma d_X^p(s) < \infty$ . Applying to this chain the map  $f : X \rightarrow S$ , we obtain the  $(Ff(a), Ff(b))$ -linking chain  $((a_i, f \circ f_i, f \circ g_i))_{i=0}^l \in L_{FS}(Ff(a), Ff(b))$ . Since  $S$  is a singleton,  $f \circ f_i = f \circ g_i$  for all  $i \in \{0, \dots, l\}$  and hence

$$a' = Ff(a) = Ff \circ f_0(a_0) = Ff \circ g_0(a_0) = Ff \circ f_1(a_1) = \dots = Ff \circ g_l(a_l) = Ff(b) = b'.$$

Then  $a = Fi_{S,X}(a') = Fi_{S,X}(b') = b$ , which contradicts our assumption.  $\square$

**Corollary 8.2.** *If  $|X| \leq 1$ , then for any elements  $a, b \in FX$  we get*

$$d_{FX}^p(a, b) = \begin{cases} 0, & \text{if } a = b, \\ \infty, & \text{if } a \neq b. \end{cases}$$

## 9. SHORT LINKING CHAINS

Let  $a, b \in FX$  be two elements. An  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  is called

- *short* if for any distinct numbers  $i, j \in \{0, \dots, l\}$  with  $\text{dom}(f_i) = \text{dom}(f_j)$  we have  $a_i \neq a_j$ ;
- *very short* if it is short and  $\text{dom}(f_i) = \text{dom}(g_i) = \max\{1, \text{deg}(F)\}$  for every  $i \in \{0, \dots, l\}$ .

Observe that the length of any very short  $(a, b)$ -linking sequence does not exceed  $|Fn|$  where  $n = \max\{1, \text{deg}(F)\}$ .

By  $L_{FX}^s(a, b)$  (resp.  $L_{FX}^{vs}(a, b)$ ) we denote the subset of  $L_{FX}(a, b)$  consisting of (very) short  $(a, b)$ -linking chains.

**Theorem 9.1.** *For any  $a, b \in FX$  the following equality holds:*

$$d_{FX}^p(a, b) = \inf(\{\infty\} \cup \{\Sigma d_X^p(w) : w \in L_{FX}^s(a, b)\}).$$

*Proof.* Let  $D := \inf(\{\infty\} \cup \{\Sigma d_X^p(w) : w \in L_{FX}^s(a, b)\})$ . The inclusion  $L_{FX}^s(a, b) \subseteq L_{FX}(a, b)$  implies the inequality  $d_{FX}^p(a, b) \leq D$ . To derive a contradiction, assume that this inequality is strict. Then there exists an  $(a, b)$ -linking chain  $w = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\Sigma d_X^p(w) < D$ . We can assume that the length  $(l + 1)$  of this chain is the smallest possible.

We claim that in this case the chain  $w$  is short. Indeed, if for some  $i < j$  we have  $\text{dom}(f_i) = \text{dom}(f_j)$  and  $a_i = a_j$ , then the chain  $w$  can be replaced by the shorter  $(a, b)$ -linking chain

$$v := ((a_0, f_0, g_0), \dots, (a_i, f_i, g_j), (a_{j+1}, f_{j+1}, g_{j+1}), \dots, (a_l, f_l, g_l)),$$

which has

$$\begin{aligned} \Sigma d_X^p(v) &= \sum_{i=0}^{i-1} d_{X^{<\omega}}^p(f_i, g_i) + d_{X^{<\omega}}^p(f_i, g_j) + \sum_{i=j+1}^l d_{X^{<\omega}}^p(f_i, g_i) \leq \\ &\leq \sum_{i=0}^{i-1} d_{X^{<\omega}}^p(f_i, g_i) + \sum_{i=i}^j d_{X^{<\omega}}^p(f_i, g_i) + \sum_{i=j+1}^l d_{X^{<\omega}}^p(f_i, g_i) = \Sigma d_X^p(w) < D, \end{aligned}$$

but this contradicts the choice of  $w$  as the shortest chain with  $\Sigma d_X^p(w) < D$ .  $\square$

**Theorem 9.2.** *If the functor  $F$  has finite degree, then for any  $a, b \in FX$*

$$d_{FX}^p(a, b) = \inf(\{\infty\} \cup \{\Sigma d_X^p(w) : w \in L_{FX}^{vs}(a, b)\}).$$

*Proof.* Let  $D := \inf(\{\infty\} \cup \{\Sigma d_X^p(w) : w \in L_{FX}^{vs}(a, b)\})$ . The inclusion  $L_{FX}^{vs}(a, b) \subseteq L_{FX}(a, b)$  implies the inequality  $d_{FX}^p(a, b) \leq D$ . To derive a contradiction, assume that this inequality is strict.

Then  $a \neq b$  and there exists an  $(a, b)$ -linking chain  $w = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\Sigma d_X^p(w) < D$ . We can assume that the length  $(l+1)$  of this chain is the smallest possible. The minimality of  $l$  implies that  $f_i \neq g_i$  and hence  $n_i \neq 0$  for all  $i \in \{0, \dots, l\}$ .

Repeating the argument of the proof of Theorem 9.1, we can show that the chain  $w$  is short. By our assumption, the cardinal  $n = \text{deg}(F)$  is finite. Taking into account that  $a \neq b$  and  $d_{FX}^p(a, b) < \infty$ , we can apply Theorem 8.1 and conclude that  $n = \text{deg}(F) > 0$ . For every  $j \in \{0, \dots, l\}$  consider the cardinal  $n_j = \text{dom}(f_j) = \text{dom}(g_j) > 0$ . By Proposition 6.1, there exists a non-empty set  $S_j \subseteq n_j$  of cardinality  $|S_j| \leq n$  such that  $a_i \in F[S_j; n_j]$ . Then  $a_j = Fj_{S_j, n_j}(a'_j)$  for some element  $a'_j \in FS_j$ . Let  $h_j : |S_j| \rightarrow S_j$  be any bijection. Then  $Fh_j$  is a bijection too. Hence, there exists an element  $a''_j \in F|S_j|$  such that  $a'_j = Fh_j(a''_j)$ .

By definition, the cardinal  $|S_j|$  coincides with the subset  $\{0, \dots, |S_j| - 1\}$  of the cardinal  $n = \{0, \dots, n - 1\}$ . For every  $j \in \{0, \dots, k\}$ , let  $b_j = Fi_{|S_j|, n}(a''_j) \in Fn$  and choose functions  $f'_j, g'_j : n \rightarrow X$  such that  $f'_j \circ i_{|S_j|, n_i} = f_j \circ h_j$ ,  $g'_j \circ i_{|S_j|, n_i} = g_j \circ h_j$ , and  $f'_j \upharpoonright_{n \setminus |S_j|} = g'_j \upharpoonright_{n \setminus |S_j|}$ . It is easy to see that  $d_{X^{<\omega}}^p(f'_j, g'_j) = d_{X^{<\omega}}^p(f_j, g_j)$ .

Observe that

$$Ff'_j(b_j) = Ff_j \circ Fi_{|S_j|, n}(a''_j) = Ff_j(a_j)$$

and similarly  $Fg'_j(b_j) = Fg_j(a_j)$ . This implies that  $w' = ((b_j, f_j, g'_j))_{j=0}^l$  is a very short  $(a, b)$ -linking chain with  $\Sigma_X^p(w') = \Sigma_X^p(w) < D$ , which contradicts the definition of  $D$ .  $\square$

**Lemma 9.3.** *If the functor  $F$  is finitary and has finite degree, then for any  $\delta \in (0, \infty)$ , set  $A \subseteq X$  and elements  $a, b \in FX$  with  $d_{FX}^\infty(a, b) < \delta$ , there exists a function  $r : X \rightarrow X$  such that  $r \circ r = r$ ,  $Fr(a) = Fr(b)$ ,  $r(A) \subseteq A$  and*

$$\sup_{x \in X} d_X(x, r(x)) < n \cdot |Fn| \cdot \delta,$$

where  $n = \max\{1, \text{deg}(F)\}$ .



*Proof.* By Theorem 9.2, there exist  $l \leq |Fn|$  and a short  $(a, b)$ -linking chain  $w = ((a_i, f_i, g_i))_{i \in l} \in (Fn \times X^n \times X^n)^l$  such that

$$\varepsilon := \Sigma d_X^\infty(w) < \delta.$$

Let  $J = \{(f_i(k), g_i(k)) : i \in l, k \in n\}$  and  $\tilde{J}$  be the smallest equivalence relation on  $X$  containing the set  $J$ . For a point  $x \in X$  let  $\tilde{J}(x) := \{y \in X : (x, y) \in \tilde{J}\}$  be its equivalence class. If  $y \in \tilde{J}(x) \setminus \{x\}$ , then we can find a sequence of pairwise distinct points  $x_0, \dots, x_m$  in  $X$  such that  $x_0 = x$ ,  $x_m = y$  and  $(x_{i-1}, x_i) \in J \cup J^{-1}$  for every  $i \in \{1, \dots, m\}$ . Then  $d_X(x, y) \leq m \cdot \varepsilon \leq |J| \cdot \varepsilon \leq nl \cdot \varepsilon$  and hence  $\text{diam}(\tilde{J}(x)) \leq nl\varepsilon$  for every  $x \in X$ . Choose a set  $S \subseteq X$  such that

- $S \cap \tilde{J}(x)$  is a singleton for every  $x \in X$ ;
- $S \cap \tilde{J}(x) \subseteq A$  for every  $x \in A$ .

Let  $r : X \rightarrow S$  be the function assigning to each point  $x \in X$  the unique point of the singleton  $S \cap \tilde{J}(x)$ . Observe that for every  $x \in X$  we have  $d_X(x, r(x)) \leq \text{diam} \tilde{J}(x) \leq nl\varepsilon < n \cdot |Fn| \cdot \delta$ . The definition of  $r$  implies  $r \circ r = r$ .

We claim that  $Fr(a) = Fr(b)$ . For this observe that  $F(a) = Ff_i(a_0)$  and  $F(b) = Fg_l(a_l)$ . For every  $i \in l$  and  $k \in n$  we have  $(f_i(k), g_i(k)) \in J \subseteq \tilde{J}$  and hence  $r \circ f_i(k) = h \circ g_i(k)$ . Then  $r \circ f_i = r \circ g_i$  and  $Fr \circ Ff_i(a_i) = Fh \circ Fg_i(a_i)$ . Finally,

$$\begin{aligned} Fr(a) &= Fh \circ Ff_0(a_0) = Fr \circ Fg_0(a_0) = Fr \circ Ff_1(a_1) = Fr \circ Fg_1(a_1) = \dots \\ &= Fr \circ Ff_l(a_l) = Fr \circ Fg_l(a_l) = Fr(b). \end{aligned}$$

□

## 10. EQUIVALENT DEFINITIONS OF THE DISTANCE $d_{FX}^p$

The distance  $d_{FX}^p$  can be equivalently defined as follows.

**Theorem 10.1.** *The distance  $d_{FX}^p$  is equal to the largest distance on  $FX$  such that for every  $n \in \mathbb{N}$  and  $a \in Fn$  the map  $\xi_X^a : X^n \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding with respect to the  $\ell^p$ -distance  $d_{X^n}^p$  on  $X^n$ .*

*Proof.* The definition of the distance  $d_{FX}^p$  implies that for every  $n \in \mathbb{N}$  and  $a \in Fn$  the map  $\xi_X^a : X^n \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding.

Next, we show that  $d_{FX}^p \geq \rho$  for any distance  $\rho$  on  $FX$  such that for every  $n \in \mathbb{N}$  and  $a \in Fn$  the map  $\xi_X^a : (X^n, d_{X^n}^p) \rightarrow (FX, \rho)$  is non-expanding. Assuming that  $d_{FX}^p \not\geq \rho$ , we can find two elements  $a, b \in FX$  such that  $d_{FX}^p(a, b) < \rho(a, b)$ . The definition of the distance  $d_{FX}^p(a, b)$  yields an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\sum_{i=0}^l d_{X^{n_i}}^p(f_i, g_i) < \rho(a, b)$ .

For every  $i \in \{0, \dots, l\}$  find a number  $n_i \in \omega$  such that  $f_i, g_i \in X^{n_i}$ . The non-expanding property of the map  $\xi_X^{a_i} : X^{n_i} \rightarrow FX$  implies that

$$\rho(Ff_i(a_i), Fg_i(a_i)) = \rho(\xi_X^{a_i}(f_i), \xi_X^{a_i}(g_i)) \leq d_{X^{n_i}}^p(f_i, g_i) = d_{X^{<\omega}}^p(f_i, g_i).$$

Applying the triangle inequality, we conclude that

$$\rho(a, b) = \rho(Ff_0(a_0), Fg_l(a_l)) \leq \sum_{i=0}^l \rho(Ff_i(a_i), Fg_i(a_i)) \leq \sum_{i=0}^l d_{X^{<\omega}}^p(f_i, g_i) < \rho(a, b),$$

which is a desired contradiction proving the inequality  $\rho \leq d_{FX}^p$ . □

If the functor  $F$  has finite degree, then the distance  $d_{FX}^p$  can be equivalently defined as follows.

**Theorem 10.2.** *If the functor  $F$  has finite degree  $m = \deg(F)$ , then the distance  $d_{FX}^p$  is equal to the largest distance on  $FX$  such that for every  $a \in Fm$  the map  $\xi_X^a : X^m \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding with respect to the  $\ell^p$ -distance  $d_{X^m}^p$  on  $X^m$ .*

*Proof.* The definition of the distance  $d_{FX}^p$  implies that for every  $a \in Fm$  the map  $\xi_X^a : X^m \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding.

Next, we show that  $d_{FX}^p \geq \rho$  for any distance  $\rho$  on  $FX$  such that for every  $a \in Fm$  the map  $\xi_X^a : (X^m, d_{X^m}^p) \rightarrow (FX, \rho)$  is non-expanding. Assuming that  $d_{FX}^p \not\geq \rho$ , we can find two elements  $a, b \in FX$  such that  $d_{FX}^p(a, b) < \rho(a, b)$ . Then  $a \neq b$  and  $d_{FX}^p(a, b) < \infty$ . The definition of the distance  $d_{FX}^p(a, b)$  yields an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\sum_{i=0}^l d_{X^m}^p(f_i, g_i) < \rho(a, b)$ . We can assume that  $f_i \neq g_i$  for all  $i \in \{0, \dots, l\}$ .

Let us show that  $m > 0$ . Assuming that  $m = 0$ , we conclude that  $\text{supp}(a) = \text{supp}(b) = \emptyset$ . By Theorem 8.1,  $d_{FX}^p(a, b) = \infty$ , which contradicts  $d_{FX}^p(a, b) < \rho(a, b)$ . So,  $m \in \mathbb{N}$ .

For every  $i \in \{0, \dots, l\}$  find a number  $n_i \in \omega$  such that  $f_i, g_i \in X^{n_i}$ . Our assumption  $f_i \neq g_i$  ensures that  $n_i \neq 0$ . Since the functor  $F$  has finite degree  $\leq m$ , the support  $\text{supp}(a_i) \subseteq n_i$  of the element  $a_i$  has cardinality  $|\text{supp}(a_i)| \leq m$ . So, we can find a map  $\gamma_i : m \rightarrow n_i$  such that  $\text{supp}(a_i) \subseteq \gamma_i[m]$  and the image  $S_i := \gamma_i[m]$  has cardinality  $|S_i| = \max\{1, |\text{supp}(a_i)|\}$ . Since  $m > 0$ , we can choose a function  $s_i : n_i \rightarrow m$  such that  $\gamma_i \circ s_i(y) = y$  for any  $y \in \gamma_i[m]$ . Let  $b_i = Fs_i(a_i) \in Fm$ . Choose functions  $f'_i, g'_i : m \rightarrow X$  such that  $f'_i(x) = g'_i(x)$  for every  $x \in m \setminus s_i[S_i]$  and  $f'_i(x) = f_i \circ \gamma_i(x)$ ,  $g'_i(x) = g_i \circ \gamma_i(x)$  for any  $x \in s_i[S_i]$ .

We claim that  $Ff'_i(b_i) = Ff_i(a_i)$  and  $Fg'_i(b_i) = Fg_i(a_i)$ . By Proposition 6.1,  $a_i \in F[S_i; n_i]$ , so there exists an element  $a'_i \in FS_i$  such that  $a_i = Fi_{S_i, n_i}(a'_i)$ . Taking into account that  $\gamma_i \circ s_i \circ i_{S_i, n_i} = i_{S_i, n_i}$  and

$$f'_i \circ s_i \circ i_{S_i, n_i} = f_i \circ \gamma_i \circ s_i \circ i_{S_i, n_i} = f_i \circ i_{S_i, n_i},$$

we conclude that

$$Ff_i(a_i) = Ff_i \circ Fi_{S_i, n_i}(a'_i) = Ff'_i \circ Fs_i \circ Fi_{S_i, n_i}(a'_i) = Ff'_i \circ Fs_i(a_i) = Ff'_i(b_i).$$

By analogy we can prove that  $Fg'_i(b_i) = Fg_i(a_i)$ . So,  $((b_0, f'_0, g'_0), \dots, (b_l, f'_l, g'_l)) \in (Fm \times X^m \times X^m)^{l+1}$  is an  $(a, b)$ -linking chain as  $Ff'_0(b_0) = Ff_0(a_0) = a$ ,  $Fg'_l(b_l) = Fg_l(a_l) = b$ , and  $Fg'_i(b_i) = Fg_i(a_i) = Ff_{i+1}(a_{i+1}) = Ff'_{i+1}(b_{i+1})$  for all  $i < l$ . Now the choice of the distance  $\rho$  and the definition of the distance  $d_{FX}^p$  ensure that

$$\begin{aligned} \rho(a, b) &= \rho(Ff'_0(b_0), Fg'_l(b_l)) \leq \sum_{i=0}^l \rho(Ff'_i(b_i), Fg'_i(b_i)) = \sum_{i=0}^l \rho(\xi_X^{b_i}(f'_i), \xi_X^{b_i}(g'_i)) \leq \\ &\leq \sum_{i=0}^l d_{X^m}^p(f'_i, g'_i) = \sum_{i=0}^l \left( \sum_{x \in m} d_X(f'_i(x), g'_i(x))^p \right)^{\frac{1}{p}} = \sum_{i=0}^l \left( \sum_{x \in s_i[S_i]} d_X(f'_i(x), g'_i(x))^p \right)^{\frac{1}{p}} = \\ &= \sum_{i=0}^l \left( \sum_{x \in s_i[S_i]} d_X(f_i \circ \gamma_i(x), g_i \circ \gamma_i(x))^p \right)^{\frac{1}{p}} = \sum_{i=0}^l \left( \sum_{y \in S_i} d_X(f_i(y), g_i(y))^p \right)^{\frac{1}{p}} \leq \\ &\leq \sum_{i=0}^l \left( \sum_{y \in n_i} d_X(f_i(y), g_i(y))^p \right)^{\frac{1}{p}} = \sum_{i=0}^l d_{X^{n_i}}^p(f_i, g_i) < \rho(a, b), \end{aligned}$$

which is a desired contradiction completing the proof of the inequality  $\rho \leq d_{FX}^p$ .  $\square$

**Problem 10.3.** *Is there any efficient algorithm for calculating the distance  $d_{FX}^p$  (with arbitrary precision)?*

### 11. THE RELATION BETWEEN THE DISTANCES $d_{FX}^p$ AND $d_{FX}^q$

In this section we establish the relation between the distances  $d_{FX}^p$  for various numbers  $p \in [1, \infty]$ . For this we will use the following known relation between the distances  $d_{X^n}^p$  and  $d_{X^n}^q$  on the set  $X^n$ .

**Lemma 11.1.** *For any  $n \in \omega$  and any numbers  $p, q \in [1, \infty]$  with  $p \leq q$  the following inequalities hold:*

$$d_{X^n}^q \leq d_{X^n}^p \leq n^{\frac{q-p}{qp}} \cdot d_{X^n}^q.$$

*Proof.* If  $n = 0$ , then  $X^0$  is a singleton and

$$d_{X^n}^q = d_{X^n}^p = n^{\frac{q-p}{qp}} \cdot d_{X^n}^q = 0.$$

So, we assume that  $n > 0$ . Choose any distinct points  $x, y \in X^n$  and consider the function  $t : n \rightarrow [0, \infty]$ ,  $t : i \mapsto d_X(x(i), y(i))$ . If  $t(i) = \infty$  for some  $i \in n$ , then

$$d_{X^n}^q(x, y) = \infty = d_{X^n}^p(x, y) = n^{\frac{q-p}{qp}} \cdot d_{X^n}^q(x, y)$$

and we are done. So, assume that  $t(i) \in \mathbb{R}$  for all  $i \in n$ .

In this case  $d_{X^n}^p(x, y) = \|t\|_p$  where  $\|t\|_p = \left(\sum_{i \in n} t(i)^p\right)^{\frac{1}{p}}$ . If  $p \leq q$ , then the well-known inequality [17, Ex.1.8]  $\|t\|_q \leq \|t\|_p$  implies that  $d_{X^n}^q(x, y) \leq d_{X^n}^p(x, y)$ .

To prove that  $d_{X^n}^p(x, y) \leq n^{\frac{q-p}{qp}} d_{X^n}^q(x, y)$ , we will use the Hölder inequality [17, 1.5] for the functions  $t^p : n \rightarrow \mathbb{R}$ ,  $t^p : i \mapsto t(i)^p$  and  $\mathbf{1} : n \rightarrow \{1\} \subset \mathbb{R}$ :

$$\sum_{i \in n} t(i)^p \cdot 1 \leq \|t^p\|_{\frac{q}{p}} \cdot \|\mathbf{1}\|_{\frac{1}{1-\frac{p}{q}}} = \left(\sum_{i \in n} t^{\frac{q}{p}}\right)^{\frac{p}{q}} \cdot n^{1-\frac{p}{q}} = (\|t\|_q)^p \cdot n^{\frac{q-p}{q}}.$$

Taking this inequality to the power  $\frac{1}{p}$ , we obtain the required inequality

$$d_{X^n}^p(x, y) = \|t\|_p \leq \|t\|_q \cdot n^{\frac{q-p}{pq}} = d_{X^n}^q(x, y) \cdot n^{\frac{q-p}{qp}}.$$

$\square$

Lemma 11.1 and Theorems 10.2, 10.1 imply

**Theorem 11.2.** *For any numbers  $p, q \in [1, \infty]$  the inequality  $p \leq q$  implies*

$$d_{FX}^\infty \leq d_{FX}^q \leq d_{FX}^p \leq d_{FX}^1 \text{ and } d_{FX}^p \leq m^{\frac{q-p}{qp}} \cdot d_{FX}^q$$

where  $m = \deg(F)$ .

*Proof.* Assume that  $1 \leq p \leq q \leq \infty$ . By Theorem 10.1, for every  $n \in \mathbb{N}$  and every  $a \in Fn$  the function  $\xi_X^a : (X^n, \ell_{X^n}^q) \rightarrow (FX, d_{FX}^q)$  is non-expanding. By Lemma 11.1, the identity function  $(X^n, d_{X^n}^p) \rightarrow (X^n, d_{X^n}^q)$  is non-expanding. Then the map  $\xi_X^a : (X^n, d_{X^n}^p) \rightarrow (FX, d_{FX}^q)$  is non-expanding (as a composition of two non-expanding maps). Applying Theorem 10.1, we conclude that  $d_{FX}^p \leq d_{FX}^q$ .

Now assume that the functor  $F$  has degree  $m = \deg(F)$ . By Lemma 11.1,  $d_{X^m}^p \leq m^{\frac{q-p}{qp}} d_{X^m}^q$ . Then for the distance  $\rho_X := m^{\frac{q-p}{qp}} \cdot d_X$  on  $X$ , the identity map  $(X^m, \rho_{X^m}^q) \rightarrow (X^m, d_{X^m}^p)$  is

non-expanding. Looking at the definitions of the distances  $d_{FX}^q$  and  $\rho_{FX}^q$ , we can see that  $\rho_{FX}^q = m^{\frac{q-p}{qp}} \cdot d_{FX}^q$ .

By Theorem 10.1, for every  $a \in F_m$  the map  $\xi_X^a : (X^m, d_{X^m}^p) \rightarrow (FX, d_{FX}^p)$  is non-expanding and so is the map  $\xi_X^a : (X^m, \rho_{X^m}^q) \rightarrow (FX, d_{FX}^p)$ . Applying Theorem 10.2, we conclude that  $d_{FX}^p \leq \rho_{FX}^q = m^{\frac{q-p}{qp}} \cdot d_{FX}^q$ .  $\square$

## 12. PRESERVING LIPSCHITZ FUNCTIONS BY THE FUNCTOR $F^p$

In this section we show that the  $\ell^p$ -metrization  $F^p$  of the functor  $F$  preserves Lipschitz functions between distance spaces and does not increase their Lipschitz constant.

**Theorem 12.1.** *For any Lipschitz function  $f : (X, d_X) \rightarrow (Y, d_Y)$  between distance spaces, the function  $Ff : (FX, d_{FX}^p) \rightarrow (FY, d_{FY}^p)$  is Lipschitz with Lipschitz constant  $\text{Lip}(Ff) \leq \text{Lip}(f)$ .*

*Proof.* We need to check that  $d_{FY}^p(Ff(a), Ff(b)) \leq \text{Lip}(f) \cdot d_{FX}^p(a, b)$  for any points  $a, b \in FX$ . This inequality is trivial if  $d_{FX}^p(a, b) = \infty$ . So, we assume that  $d_{FX}^p(a, b) < \infty$ . In this case it suffices to prove that  $d_{FY}^p(Ff(a), Ff(b)) \leq \text{Lip}(f) \cdot (d_{FX}^p(a, b) + \varepsilon)$  for every  $\varepsilon > 0$ . By the definition of the distance  $d_{FX}^p(a, b)$ , there exists an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\sum_{i=0}^l d_{X^{\leftarrow \omega}}^p(f_i, g_i) < d_{FX}^p(a, b) + \varepsilon$ . Applying to this chain the function  $f : X \rightarrow Y$ , we obtain an  $(Ff(a), Ff(b))$ -linking chain  $(a_i, f \circ f_i, f \circ g_i)_{i=0}^l$  such that

$$\begin{aligned} d_{FY}^p(Ff(a), Ff(b)) &\leq \sum_{i=0}^l d_{Y^{\leftarrow \omega}}^p(f \circ f_i, f \circ g_i) = \sum_{i=0}^l \left( \sum_{x \in n} d_Y(f \circ f_i(x), f \circ g_i(x))^p \right)^{\frac{1}{p}} \leq \\ &\leq \sum_{i=0}^l \left( \sum_{x \in n_i} (\text{Lip}(f) \cdot d_X(f_i(x), g_i(x)))^p \right)^{\frac{1}{p}} = \text{Lip}(f) \cdot \sum_{i=0}^l \left( \sum_{x \in n_i} d_X(f_i(x), g_i(x))^p \right)^{\frac{1}{p}} = \\ &= \text{Lip}(f) \cdot \sum_{i=0}^l d_{X^{\leftarrow \omega}}^p(f_i, g_i) \leq \text{Lip}(f) \cdot (d_{FX}^p(a, b) + \varepsilon), \end{aligned}$$

where  $n_i := \text{dom}(f_i) = \text{dom}(g_i)$  for  $i \in \{0, \dots, l\}$ .  $\square$

## 13. PRESERVING ISOMETRIES BY THE FUNCTOR $F^p$

**Lemma 13.1.** *If  $f : X \rightarrow Y$  is a surjective isometry of distance spaces, then the map  $F^p f : F^p X \rightarrow F^p Y$  is an isometry.*

*Proof.* By the surjectivity of  $f$ , there exists a function  $g : Y \rightarrow X$  such that  $f \circ g$  is the identity map of  $Y$ . Then  $Ff \circ Fg$  is the identity map of  $FY$ , which implies that  $Ff$  is surjective.

Taking into account that  $f$  is an isometry, we conclude that for any  $y, y' \in Y$ ,

$$d_X(g(y), g(y')) = d_Y(f \circ g(y), f \circ g(y')) = d_Y(y, y'),$$

which means that  $g$  is an isometry. Consider the map  $r = g \circ f : X \rightarrow X$  and observe that for every  $x \in X$  we have

$$d_X(x, r(x)) = d_X(x, g \circ f(x)) = d_Y(f(x), f \circ g \circ f(x)) = d_Y(f(x), f(x)) = 0.$$

We claim that for any  $a \in FY$  the distance  $d_{FX}^p(a, Fr(b)) = 0$ . Since the functor  $F$  has finite supports, there exist  $n_0 \in \omega$ , map  $f_0 : n_0 \rightarrow X$  and  $a_0 \in Fn_0$  such that  $a = Ff_0(a_0)$ . Put  $g_0 = r \circ f_0$  and observe that  $((a_0, f_0, g_0))$  is an  $(a, Fr(b))$ -linking chain witnessing that

$$d_{FX}^p(a, Fr(a)) \leq d_{X^{<\omega}}^p(f_0, g_0) = \left( \sum_{i \in n_0} d_X(f_0(i), r \circ f_0(i))^p \right)^{\frac{1}{p}} = 0.$$

Now take any elements  $a, b \in FX$  and applying Theorem 12.1, conclude that

$$\begin{aligned} d_{FX}^p(a, b) &\leq d_{FX}^p(a, Fr(a)) + d_{FX}^p(Fr(a), Fr(b)) + d_{FX}^p(Fr(b), b) = \\ &= 0 + d_{FX}^p(Fg \circ Ff(a), Fg \circ Ff(b)) + 0 \leq d_{FY}^p(Ff(a), Ff(b)) \leq d_{FX}^p(a, b), \end{aligned}$$

which implies that all inequalities in this chain are equalities. In particular,

$$d_{FY}^p(Ff(a), Ff(b)) = d_{FX}^p(a, b),$$

which means that the map  $Ff$  is an isometry.  $\square$

We will say that a subset  $A \subseteq X$  of a distance space  $(X, d_X)$  is *dense* in  $X$  if for any  $\varepsilon > 0$  and  $x \in X$  there exists  $a \in A$  such that  $d_X(a, x) < \varepsilon$ .

**Lemma 13.2.** *If  $f : X \rightarrow Y$  is an injective isometry of distance spaces and the set  $f(X)$  is dense in  $Y$ , then the map  $F^p f : F^p X \rightarrow F^p Y$  is an injective isometry and the set  $F^p f[F^p X] = Ff[FX]$  is dense in  $F^p Y$ .*

*Proof.* If  $X = \emptyset$ , then  $Y = \emptyset$  by the density of the set  $f[X] = \emptyset$  in  $Y$ . In this case the map  $f : X \rightarrow Y$  is the identity map of  $\emptyset$  and  $F^p f$  is the identity map (and hence a bijective isometry) of the distance space  $F^p X = F^p Y = F^p \emptyset$ .

So, we assume that the set  $X$  is not empty. Then  $Y$  is not empty, too. By Theorem 12.1, the map  $F^p f : F^p X \rightarrow F^p Y$  is non-expanding. Assuming that  $F^p f$  is not an isometry, we can find two elements  $a, b \in F^p X$  such that  $d_{FY}^p(Ff(a), Ff(b)) < d_{FX}^p(a, b)$ . By the definition of the distance  $d_{FY}^p$ , there exists an  $(Ff(a), Ff(b))$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in LFY(Ff(a), Ff(b))$  such that  $\sum_{i=0}^l d_{Y^{<\omega}}^p(f_i, g_i) < d_{FX}^p(a, b)$ . For every  $i \in \{0, \dots, l\}$  let  $n_i := \text{dom}(f_i) = \text{dom}(g_i)$ . Choose  $\varepsilon > 0$  so small that

$$\sum_{i=0}^l (2\varepsilon \cdot \sqrt[p]{n_i} + d_{Y^{<\omega}}^p(f_i, g_i)) < d_{FX}^p(a, b).$$

Using the injectivity of the function  $f$  and the density of the set  $f[X]$  in  $Y$ , choose a function  $r : Y \rightarrow X$  such that  $r \circ f(x) = x$  for all  $x \in X$ , and  $d_Y(y, f \circ r(y)) < \varepsilon$  for all  $y \in Y \setminus f[X]$ . Applying the functor  $F$  to the identity map  $r \circ f$  of  $X$ , we conclude that  $Fr \circ Ff$  is the identity map of  $FX$ , which implies that the map  $Ff : FX \rightarrow FY$  is injective.

Consider the maps  $f'_i = r \circ f_i : n_i \rightarrow X$  and  $g'_i = r \circ g_i : n_i \rightarrow X$ . Taking into account that the map  $f$  is an isometry, we conclude that

$$\begin{aligned} d_{X^{<\omega}}^p(f'_i, g'_i) &= d_{Y^{<\omega}}^p(f \circ f'_i, f \circ g'_i) = d_{Y^{<\omega}}^p(f \circ r \circ f_i, f \circ r \circ g_i) \leq \\ &\leq d_{Y^{<\omega}}^p(f \circ r \circ f_i, f_i) + d_{Y^{<\omega}}^p(f_i, g_i) + d_{Y^{<\omega}}^p(g_i, f \circ r \circ g_i) < \varepsilon \sqrt[p]{n_i} + d_{Y^{<\omega}}^p(f_i, g_i) + \varepsilon \sqrt[p]{n_i}. \end{aligned}$$

We claim that  $Ff'_0(a_0) = a$  and  $Fg'_k(a_k) = b$ . It follows that  $\text{supp}(Ff(a)) \subseteq f[X]$  so we can find a finite set  $S \subseteq f[X]$  of cardinality  $|S_a| = \max\{1, |\text{supp}(Ff(a))|\}$  containing the support  $\text{supp}(Ff(a))$  of  $Ff(a)$ . By Proposition 6.1,  $Ff(a) \subseteq F[S; Y]$ . So, we can find an element

$a' \in FS$  such that  $Ff(a) = Fi_{S,Y}(a')$ . The definition of  $r$  and the inclusion  $S \subseteq f[X]$  imply  $f \circ r \circ i_{S,Y} = i_{S,Y}$ . Then

$$\begin{aligned} Ff(a) &= Fi_{S,Y}(a') = Ff \circ Fr \circ Fi_{S,Y}(a') = Ff \circ Fr \circ Ff(a) = \\ &= Ff \circ Fr \circ Ff_0(a_0) = Ff \circ Ff'_0(a_0) \end{aligned}$$

and hence  $a = Ff'_0(a_0)$  by the injectivity of the map  $Ff$ .

By analogy, we can prove that  $Fg'_l(a_l) = b$ . Consequently,  $((a_0, f'_0, g'_0), \dots, (a_l, f'_l, g'_l))$  is an  $(a, b)$ -linking chain and

$$d_{FX}^p(a, b) \leq \sum_{i=0}^l d_{X^{<\omega}}^p(f'_i, g'_i) < \sum_{i=0}^l (\varepsilon \sqrt[p]{n_i} + d_{X^{<\omega}}^p(f_i, g_i) + \varepsilon \sqrt[p]{n_i}) < d_{FX}^p(a, b),$$

which is a desired contradiction proving that  $d_{FY}^p(Ff(a), Ff(b)) = d_{FX}^p(a, b)$ .

Finally, we prove that the image  $Ff[FX]$  is dense in  $F^dY$ . Given any  $b \in FY$  and  $\varepsilon > 0$ , choose a finite set  $S \subseteq Y$  of cardinality  $|S| = \max\{1, |\text{supp}(b)|\}$  containing  $\text{supp}(b)$ . By Proposition 6.1,  $b \in F[S; Y]$ .

Let  $n_0 = |S|$ ,  $\beta : n_0 \rightarrow S$  be any bijection, and  $f_0 = i_{S,Y} \circ \beta : n_0 \rightarrow Y$ . Then  $F\beta : Fn_0 \rightarrow FS$  is a bijection and hence  $b \in F[S; Y] = Fi_{S,Y}[FS] = Fi_{S,Y}[F\beta[Fn_0]] = Ff_0[Fn_0]$ . Consequently, there exists an element  $b' \in Fn_0$  such that  $b = Ff_0(b')$ .

Since the set  $f[X]$  is dense in  $Y$ , there exists a function  $g_0 : n_0 \rightarrow f[X] \subseteq Y$  such that  $d_{Y^{<\omega}}^p(f_0, g_0) < \varepsilon$ . It follows that the map  $\alpha = f^{-1} \circ g_0 : n_i \rightarrow X$  is well-defined and  $f \circ \alpha = g_0$ . Let  $a := F\alpha(b') \in FX$  and observe that  $Ff(a) = Ff \circ F\alpha(b') = Fg_0(b')$  and

$$d_{FY}^p(b, Ff(a)) = d_{FX}^p(Ff_0(b'), Fg_0(b')) \leq d_{Y^{<\omega}}^p(f_0, g_0) < \varepsilon,$$

which means that the set  $Ff[FX]$  is dense in  $F^pY$ .  $\square$

Lemmas 13.1 and 13.2 imply

**Theorem 13.3.** *If  $f : X \rightarrow Y$  is an isometry of distance spaces and the set  $f[X]$  is dense in  $Y$ , then the map  $F^p f : F^p X \rightarrow F^p Y$  is an isometry and the set  $F^p f[F^p X] = Ff[FX]$  is dense in  $F^p Y$ .*

*Proof.* The isometry  $f$  can be written as the composition  $f = h \circ g$  of a surjective isometry  $g : X \rightarrow Z$  and an injective isometry  $h : Z \rightarrow Y$ . For example, we can take  $Z = f[X]$ ,  $g = f$  and  $h = i_{Z,Y}$ . By Lemma 13.1, the map  $Ff : F^p X \rightarrow F^p Z$  is a surjective isometry. By the density of the set  $f[X] = h[g[X]] = h[Z]$  in  $Y$  and Lemma 13.2, the map  $Fg : F^p X \rightarrow F^p Y$  is an isometry with dense image  $Fg[FX]$  in  $F^p Y$ . Then the function  $Ff = Fh \circ Fg$  is an isometry as the composition of two isometries. Moreover, the image  $Ff[FX] = Fh[Fg[FX]] = Fh[FZ]$  is dense in  $F^p Y$  by the density of  $Fg[FZ]$  in  $F^p Y$ .  $\square$

**Example 13.4.** In general, the functor  $F^p$  does not preserve isometries. To construct a counterexample, consider the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  assigning to each set  $X$  the family  $FX$  of all subsets  $A \subseteq X$  of cardinality  $0 < |A| \leq 2$ . To every map  $f : X \rightarrow Y$  between sets the functor  $F$  assigns the map  $Ff : FX \rightarrow FY$ ,  $Ff : a \mapsto f[a]$ . The functor  $F$  has finite degree 2, preserves supports and singletons (see Section 30 for more information on this functor).

Now consider the following finite subsets  $X \subseteq Y$  of the complex plane  $\mathbb{C}$ :

$$X := \{5 + \sqrt{3}i, 5 - \sqrt{3}i, -5 + \sqrt{3}i, -5 - \sqrt{3}i\} \quad \text{and} \quad Y = X \cup \{-4, 4\}.$$

In the set  $FX \subseteq FY$  consider the elements  $a = \{-5 + \sqrt{3}i, -5 - \sqrt{3}i\}$  and  $b = \{5 + \sqrt{3}i, 5 - \sqrt{3}i\}$ .

It can be shown that for every  $p \in [1, \infty]$

$$d_{FX}^p(a, b) = 10 + 4\sqrt{3} > 8 + 4\sqrt[3]{2} = d_{FY}^p(a, b).$$

Yet, we have the following partial positive result.

**Proposition 13.5.** *If the functor  $F$  is finitary and has finite degree, then for any isometry  $f : X \rightarrow Y$  between nonempty distance spaces and any elements  $a, b \in F^p X$ , we have*

$$d_{FX}^\infty(a, b) \leq 2n \cdot |Fn| \cdot d_{FY}^\infty(Ff(a), Ff(b)),$$

where  $n = \max\{1, \deg(F)\}$ .

*Proof.* To derive a contradiction, assume that  $d_{FX}^\infty(a, b) > 2n \cdot |Fn| \cdot d_{FY}^\infty(Ff(a), Ff(b))$ . Using Lemma 9.3, find a function  $r : Y \rightarrow Y$  such that  $Fr(Ff(a)) = Fr(Ff(b))$ ,  $r[f[X]] \subseteq f[X]$  and

$$\sup_{y \in Y} d_Y(y, r(y)) < n \cdot |Fn| \cdot \frac{1}{2n \cdot |Fn|} \cdot d_{FY}^\infty(a, b) = \frac{1}{2} d_{FX}^\infty(a, b).$$

Let  $s : f[X] \rightarrow X$  be any function such that  $s(y) \in f^{-1}(y)$  and hence  $f \circ s(y) = y$  for every  $y \in f[X]$ . Using Proposition 6.1, find two functions  $f_0, g_3 \in X^n$  and elements  $a_0, a_3 \in Fn$  such that  $a = Ff_0(a_0)$  and  $b = Fg_3(a_3)$ . Now consider the chain

$$w = ((a_0, f_0, g_0), (a_1, f_1, g_1), (a_2, f_2, g_2), (a_3, f_3, g_3)) \in (Fn \times X^n \times X^n)^4,$$

where

$$\begin{aligned} g_0 &= s \circ f \circ f_0, & a_1 &= a_0, & f_1 &= g_0, & g_1 &= s \circ r \circ f \circ f_0, \\ a_2 &= a_3, & f_2 &= s \circ r \circ f \circ g_3, & g_2 &= s \circ f \circ g_3, & f_3 &= g_2. \end{aligned}$$

It is clear that  $Fg_0(a_0) = Ff_1(a_1)$  and  $Fg_2(a_2) = Ff_3(a_3)$ . Also

$$\begin{aligned} Fg_1(a_1) &= Fs \circ Fr \circ Ff \circ Ff_0(a_0) = Fs \circ Fr \circ Ff(a) = \\ &Fs \circ Fr \circ Ff(b) = Fs \circ Fr \circ Ff \circ Fg_3(a_3) = Ff_2(a_2). \end{aligned}$$

Therefore, the chain  $w$  is  $(a, b)$ -linking.

Taking into account that  $f$  is an isometry and  $f \circ s(y) = y$  for  $y \in f[X]$ , we conclude that for every  $k \in n$ ,

$$\begin{aligned} d_X(f_0(k), g_0(k)) &= d_X(f_0(k), s \circ f \circ f_0(k)) = d_Y(f \circ f_0(k), f \circ s \circ f \circ f_0(k)) = \\ &d_Y(f \circ f_0(k), f \circ f_0(k)) = 0. \end{aligned}$$

Then  $d_{X^n}^\infty(f_0, g_0) = 0$ . By analogy we can show that  $d_{X^n}^\infty(f_3, g_3) = 0$ .

Also for any  $k \in n$  we have

$$\begin{aligned} d_X(f_1(k), g_1(k)) &= d_X(s \circ f \circ f_0(k), s \circ r \circ f \circ f_0(k)) = \\ &d_Y(f \circ s \circ f \circ f_0(k), f \circ s \circ r \circ f \circ f_0(k)) = d_Y(f \circ f_0(k), r \circ f \circ f_0(k)) \leq \\ &\sup_{y \in Y} d_Y(y, r(y)) < \frac{1}{2} d_{FX}^\infty(a, b) \end{aligned}$$

and hence  $d_{X^n}^\infty(f_1, g_1) < \frac{1}{2} d_{FX}^\infty(a, b)$ . By analogy we can prove that  $d_{X^n}^\infty(f_2, g_2) < \frac{1}{2} d_{FX}^\infty(a, b)$ . Then

$$d_{FX}^\infty(a, b) \leq d_X^\infty(w) = \sum_{i=0}^3 d_X^\infty(f_i, g_i) < 0 + \frac{1}{2} d_{FX}^\infty(a, b) + \frac{1}{2} d_{FX}^\infty(a, b) + 0 = d_{FX}^\infty(a, b),$$

which is a desired contradiction.  $\square$

#### 14. THE FUNCTOR $F^p f$ AND THE CONTINUITY MODULUS OF FUNCTIONS

Let us recall that for a function  $f : X \rightarrow Y$  between distance spaces, its continuity modulus  $\omega_f : (0, \infty) \rightarrow [0, \infty]$  is defined by the formula

$$\omega_f(\varepsilon) = \sup\{d_Y(f(x), f(y)) : x, y \in X \wedge d_X(x, y) < \varepsilon\} \text{ for } \varepsilon \in (0, \infty].$$

**Theorem 14.1.** *If the functor  $F$  has finite degree  $n = \deg(F)$ , then for any function  $f : X \rightarrow Y$  between distance spaces, the function  $F^p f : F^p X \rightarrow F^p Y$  has continuity modulus  $\omega_{F^p f} \leq |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f$ .*

*Proof.* Assuming that  $\omega_{F^p f} \not\leq |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f$ , we can find  $\varepsilon \in (0, \infty]$  and elements  $a, b \in F^p X$  such that  $d_{F^p X}^p(a, b) < \varepsilon$  but  $d_{F^p Y}^p(Ff(a), Ff(b)) > |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f(\varepsilon)$ . Then  $a \neq b$  and we can apply Theorem 8.1 to conclude that  $|\text{supp}(a) \cup \text{supp}(b)| > 1$  and hence  $n = \deg(F) > 0$ . By Theorem 9.2, there exists a very short  $(a, b)$ -linking chain  $w = ((a_i, f_i, g_i))_{i \in I} \in (Fn \times X^n \times X^n)^l$  such that  $\sum d_X^p(w) = \sum_{i \in I} d_{X^n}^p(f_i, g_i) < \varepsilon$ . Since the chain  $w$  is short, the sequence  $(Ff_i(a_i))_{i \in I}$  consists of pairwise distinct elements of  $F^p X$  and hence the elements  $a_i \in Fn$  are also pairwise distinct. Then  $k \leq |Fn|$ . Observe that  $((a_i, f \circ f_i, f \circ g_i))_{i \in I}$  is an  $(Ff(a), Ff(b))$ -linking chain and hence

$$\begin{aligned} |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f(\varepsilon) &< d_{F^p Y}^p(Ff(a), Ff(b)) \leq \sum_{i=0}^l d_{F^p Y}^p(f \circ f_i, f \circ g_i) = \\ &\sum_{i \in I} \left( \sum_{x \in n} d_Y(f(f_i(x)), f(g_i(x)))^p \right)^{\frac{1}{p}} \leq \sum_{i \in I} \left( \sum_{x \in n} \omega_f(d_X(f_i(x), g_i(x)))^p \right)^{\frac{1}{p}} \leq \\ &\sum_{i \in I} \left( \sum_{x \in n} \omega_f(\varepsilon)^p \right)^{\frac{1}{p}} = l \cdot n^{\frac{1}{p}} \cdot \omega_f(\varepsilon) \leq |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f(\varepsilon), \end{aligned}$$

which is a desired contradiction completing the proof of the inequality  $\omega_{F^p f} \leq |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f$ .  $\square$

**Corollary 14.2.** *Assume that the functor  $F$  is finitary and has finite degree. If a function  $f : X \rightarrow Y$  between distance spaces is asymptotically Lipschitz (resp. microform, macroform, duoform), then so is the function  $F^p f : F^p X \rightarrow F^p Y$ .*

**Lemma 14.3.** *For any nonempty set  $Z$  and functions  $f, g : Z \rightarrow X$  we have*

$$\sup_{a \in FZ} d_{F^p X}^p(Ff(a), Fg(a)) \leq n^{\frac{1}{p}} \cdot \sup_{z \in Z} d_X(f(z), g(z))$$

where  $n = \max\{1, \deg(F)\}$ .

*Proof.* To derive a contradiction, assume that  $d_{F^p X}^p(Ff(a), Fg(b)) > n^{\frac{1}{p}} \cdot d_{X^Z}(f, g)$  for some  $a \in FZ$ .

Given any  $a \in FZ$ , use Proposition 6.1 and find a nonempty finite subset  $S \subseteq X$  of cardinality  $|S| \leq n$  and an element  $a' \in FS$  such that  $a = Fi_{S,X}(a')$ . Let  $k = |S|$  and  $h : k \rightarrow S$  be any bijection. Then  $Fh$  is a bijection, too, and we can find an element  $a_0 \in Fk$  such that  $a' = Fh(a_0)$ . Consider the maps  $f_0 = f \circ i_{S,X} \circ h : k \rightarrow X$  and  $g_0 = g \circ i_{S,X} \circ h : k \rightarrow X$  and observe that

$$Ff_0(b) = Ff \circ Fi_{S,X} \circ Fh(a_0) = Ff \circ Fi_{S,X}(a') = Ff(a)$$



and similarly,  $Fg_0(b) = Fg(a)$ . Then  $((a_0, f_0, g_0))$  is an  $(Ff(a), Ff(b))$ -linking chain and hence

$$d_{FX}^p(Ff(a), Fg(a)) \leq \left( \sum_{i \in k} d_X(f_0(i), g_0(i))^p \right)^{\frac{1}{p}} \leq k^{\frac{1}{p}} \cdot d_{XZ}(f, g) \leq n^{\frac{1}{p}} \cdot d_{XZ}(f, g),$$

which contradicts the choice of  $a$ .  $\square$

**Corollary 14.4.** *Assume that the functor  $F$  is finitary and has finite degree. If a function  $f : X \rightarrow Y$  between distance spaces is a coarse equivalence (resp. quasi-isometry), then so is the function  $F^p f : F^p X \rightarrow F^p Y$ .*

*Proof.* If  $f$  is a coarse equivalence, then  $f$  is macroform and admits a macroform map  $g : Y \rightarrow X$  such that

$$d_{XX}(1_X, g \circ f) = \sup_{x \in X} d_X(x, g \circ f(x)) < \infty$$

and

$$d_{YY}(1_Y, f \circ g) = \sup_{y \in Y} d_Y(y, f \circ g(y)) < \infty.$$

By Corollary 14.2, the maps  $F^p f : F^p X \rightarrow F^p Y$  and  $F^p g : F^p Y \rightarrow F^p X$  are macroform and by Lemma 14.3,

$$\sup_{a \in F^p X} d_{FX}^p(a, F^p g \circ F^p f(a)) \leq \max\{1, \deg(F)\}^{\frac{1}{p}} \cdot d_{XX}(1_X, g \circ f) < \infty$$

and

$$\sup_{b \in F^p Y} d_{FY}^p(b, F^p f \circ F^p g(b)) \leq \max\{1, \deg(F)\}^{\frac{1}{p}} \cdot d_{YY}(1_Y, f \circ g) < \infty.$$

This means that the map  $F^p$  is a coarse equivalence.

By analogy we can prove that the map  $F^p f$  is a quasi-isometry if so is the map  $f$ .  $\square$

## 15. UPPER BOUNDS FOR $d_{FX}^p$

In this section we prove some upper bounds for the distance  $d_{FX}^p$ . In particular, we establish conditions implying that  $d_{FX}^p$  is a pseudometric. First we show that the functor  $F^p$  preserves the  $\{0, \infty\}$ -valuedness of distances.

A distance  $d$  on a set  $Y$  is called  $\{0, \infty\}$ -valued if  $d(Y \times Y) = \{0, \infty\}$ .

**Lemma 15.1.** *If the distance  $d_X$  in  $\{0, \infty\}$ -valued (and  $\infty$ -metric), then the distance  $d_{FX}^p$  on  $FX$  also is  $\{0, \infty\}$ -valued (and  $\infty$ -metric).*

*Proof.* To show that  $d_{FX}^p$  is  $\{0, \infty\}$ -valued (and  $\infty$ -metric), it suffices to check that  $d_{FX}^p(a, b) = 0$  (and  $a = b$ ) for any elements  $a, b \in FX$  with  $d_{FX}^p(a, b) < \infty$ . Using the definition of the distance  $d_{FX}^p$ , find an  $(a, b)$ -linking chain  $s = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\sum_{i=0}^l d_{X^{<\omega}}^p(f_i, g_i) < \infty$ . If  $d_X$  is  $\{0, \infty\}$ -valued, then so is the distance  $d_{X^{<\omega}}^p$ . Then for every  $i \leq l$  the inequality  $d_{X^{<\omega}}^p(f_i, g_i) < \infty$  is equivalent to  $d_{X^{<\omega}}^p(f_i, g_i) = 0$ , and finally

$$d_{FX}^p(a, b) \leq \sum_{i=0}^l d_{X^{<\omega}}^p(f_i, g_i) = 0.$$

Now assuming that  $d_X$  is an  $\infty$ -metric, we can show that the distance  $d_{X^{<\omega}}^p$  is an  $\infty$ -metric and hence the equalities  $d_{X^{<\omega}}^p(f_i, g_i) = 0$  imply  $f_i = g_i$  for all  $i \in \{0, \dots, l\}$ . Then

$$a = Ff_0(a_0) = Fg_0(a_0) = Ff_1(a_1) = Fg_1(a_1) = \dots = Fg_l(a_l) = b.$$

□

The distance  $d_X$  on  $X$  induces the equivalence relation  $\sim$  on  $X$ , defined by  $x \sim y$  iff  $d_X(x, y) < \infty$ . For an element  $x \in X$  let  $O(x, \infty) := \{y \in X : d_X(x, y) < \infty\}$  be the equivalence class of  $x$  under the equivalence relation  $\sim$ . Let  $X/\sim = \{O(x, \infty) : x \in X\}$  be the quotient set of  $X$  by the relation  $\sim$  and  $q : X \rightarrow X/\sim$  be the quotient map.

For two finite sets  $A, B \subseteq X$  let  $\underline{d}_X(A, B) = \inf\{d_X(x, y) : x \in A, y \in B\}$ . If one of the sets  $A$  or  $B$  is empty, then we put  $\underline{d}_X(A, B) = 0$  (by definition). Let

$$\underline{d}_X^p(A, B) = \left( \sum_{E \in X/\sim} \underline{d}_X(A \cap E, B \cap E)^p \right)^{\frac{1}{p}}$$

and observe that  $\underline{d}_X^p(A, B)$  is finite (in contrast to  $\underline{d}_X(A, B)$  which can be infinite).

For a subset  $S \subseteq X$  by

$$\bar{d}_X(S) = \sup(\{0\} \cup \{d_X(x, y) : x, y \in S, d_X(x, y) < \infty\})$$

we denote the real diameter of  $S$ .

**Theorem 15.2.** *Let  $q : X \rightarrow X/\sim$  be the quotient map defined above. For any two elements  $a, b \in FX$  the following conditions are equivalent:*

- 1)  $d_{FX}^p(a, b) < \infty$ ,
- 2)  $Fq(a) = Fq(b)$ ,
- 3)  $d_{FX}^p(a, b) \leq |S_a|^{\frac{1}{p}} \cdot \bar{d}_X(S_a) + \underline{d}_X^p(S_a, S_b) + |S_b|^{\frac{1}{p}} \cdot \bar{d}_X(S_b)$ ,
- 4)  $d_{FX}^p(a, b) \leq \min\{|S_a|^{\frac{1}{p}} \bar{d}_X(S_a) + |S_b|^{\frac{1}{p}} \bar{d}_X(S), |S_a|^{\frac{1}{p}} \bar{d}_X(S) + |S_b|^{\frac{1}{p}} \bar{d}_X(S_b)\}$ ,
- 5)  $d_{FX}^p(a, b) \leq (|S_a|^{\frac{1}{p}} + |S_b|^{\frac{1}{p}}) \cdot \bar{d}_X(S)$ ,

where  $S_a = \text{supp}(a)$ ,  $S_b = \text{supp}(b)$  and  $S = \text{supp}(a) \cup \text{supp}(b)$ .

*Proof.* Endow the set  $Y = X/\sim$  with a unique  $\{0, \infty\}$ -valued  $\infty$ -distance  $d_Y$ , and observe that the quotient map  $q : X \rightarrow Y$  is Lipschitz with Lipschitz constant  $\text{Lip}(q) = 0$  (here we assume that  $0 \cdot \infty = \infty$ ). By Theorem 12.1,  $\text{Lip}(Fq) = 0$ . By Lemma 15.1, the distance  $d_{FY}^p$  is a  $\{0, \infty\}$ -valued  $\infty$ -metric. Now we are ready to prove that (1)  $\Rightarrow$  (2)  $\Rightarrow$  (3)  $\Rightarrow$  (1) and (2)  $\Rightarrow$  (4)  $\Rightarrow$  (5)  $\Rightarrow$  (1). The implications (3)  $\Rightarrow$  (1)  $\Leftarrow$  (5)  $\Leftarrow$  (4) are trivial.

(1)  $\Rightarrow$  (2) If  $d_{FX}^p(a, b) < \infty$ , then  $d_{FY}^p(Fq(a), Fq(b)) \leq \text{Lip}(Fq) \cdot d_{FX}^p(a, b) = 0$  and finally  $Fq(a) = Fq(b)$  (because the distance  $d_{FY}^p$  is an  $\infty$ -metric).

(2)  $\Rightarrow$  (3, 4). Assume that  $Fq(a) = Fq(b)$  and put  $c = Ff(a) = Fg(b)$ . Let  $S_a := \text{supp}(a)$  and  $S_b := \text{supp}(b)$ . For every equivalence class  $y \in Y = X/\sim$  with  $S_a \cap y \neq \emptyset \neq S_b \cap y$  choose two points  $\alpha_{y,a} \in S_a \cap y$  and  $\beta_{y,b} \in S_b \cap y$  such that  $d_X(s_{y,a}, s_{y,b}) = \underline{d}_X(S_a \cap y, S_b \cap y)$ . Fix two maps  $\alpha, \beta : Y \rightarrow X$  such that for every  $y \in Y$  the following conditions hold:

- $\alpha(y) = \alpha_{y,a}$  and  $\beta(y) = \beta_{y,a}$  if  $S_a \cap y \neq \emptyset \neq S_b \cap y$ ;
- $\alpha(y) = \beta(y) \in S_a$  if  $S_a \cap y \neq \emptyset = S_b \cap y$ ;
- $\alpha(y) = \beta(y) \in S_b$  if  $S_a \cap y = \emptyset = S_b \cap y$ .

It can be shown that

$$\left( \sum_{y \in Y} d_X(\alpha(y), \beta(y))^p \right)^{\frac{1}{p}} = \underline{d}_X^p(S_a, S_b).$$

Choose a set  $\dot{S}_c \subseteq Y$  of cardinality  $|\dot{S}_c| = \max\{1, |\text{supp}(c)|\}$  such that  $\text{supp}(c) \subseteq \dot{S}_c$ . Next, choose a set  $\dot{S}_a \subseteq X$  of cardinality  $|\dot{S}_a| = \max\{1, |S_a|\}$  such that  $S_a \subseteq \dot{S}_a$ , and  $\dot{S}_a \subseteq \alpha(Y)$  if

$S_a = \emptyset$ . Also choose a set  $\dot{S}_b \subseteq X$  of cardinality  $|\dot{S}_b| = \max\{1, |S_b|\}$  such that  $S_b \subseteq \dot{S}_b$ , and  $\dot{S}_b \subseteq \beta(Y)$  if  $S_b = \emptyset$ .

Let  $n_0 = |\dot{S}_a|$ ,  $n_1 = |\dot{S}_c|$ ,  $n_2 = |\dot{S}_b|$ , and  $f_0 : n_0 \rightarrow \dot{S}_a$ ,  $\gamma : n_2 \rightarrow \dot{S}_c$ ,  $g_2 : n_2 \rightarrow \dot{S}_b$  be any bijective maps. By Proposition 6.1,  $a \in F[\dot{S}_a; X]$ ,  $b \in F[\dot{S}_b; X]$  and  $c \in F[\dot{S}_c; Y]$ . This allows us to find elements  $a_0 \in Fn_0$ ,  $a_1 \in Fn_1$ ,  $a_2 \in Fn_2$  such that  $a = Ff_0(a_0)$ ,  $c = F\gamma(a_1)$ ,  $b = Fg_2(a_2)$ . Let  $g_0 = \alpha \circ q \circ f_0$ ,  $f_1 = \alpha \circ \gamma$ ,  $g_1 = \beta \circ \gamma$ ,  $f_2 = \beta \circ q \circ g_2$ . To see that  $((a_0, f_0, g_0), (a_1, f_1, g_1), (a_2, f_2, g_2))$  is an  $(a, b)$ -linking chain, we need to check that  $Fg_0(a_0) = Ff_1(a_1)$  and  $Fg_1(a_1) = Ff_2(a_2)$ .

For the first equality, observe that

$$Fg_0(a_0) = F\alpha \circ Fq \circ Ff_0(a_0) = F\alpha \circ Fq(a) = F\alpha(c) = F\alpha(F\gamma(a_1)) = Ff_1(a_1).$$

By analogy we can check that  $Fg_1(a_1) = Ff_2(a_2)$ . Then  $((a_i, f_i, g_i))_{i=0}^k \in L_{FX}(a, b)$  and

$$\begin{aligned} d_{FX}^p(a, b) &\leq d_{X^{<\omega}}^p(f_0, g_0) + d_{X^{<\omega}}^p(f_1, g_1) + d_{X^{<\omega}}^p(f_2, g_2) = \\ &= d_{X^{<\omega}}^p(f_0, \alpha \circ q \circ f_0) + d_{X^{<\omega}}^p(\alpha \circ \gamma, \beta \circ \gamma) + d_{X^{<\omega}}^p(\beta \circ q \circ g_2, g_2) \leq \\ &\leq |S_a|^{\frac{1}{p}} \cdot \bar{d}_X(S_a) + \underline{d}_X(S_a, S_b) + |S_b|^{\frac{1}{p}} \cdot \bar{d}_X(S_b). \end{aligned}$$

Considering the  $(a, b)$ -linking chain  $((a_0, f_1, g_1), (a_2, f'_2, g_2))$  with  $f'_2 = \alpha \circ q \circ g_2$ , we can see that

$$\begin{aligned} d_{FX}^p(a, b) &\leq d_{X^{<\omega}}^p(f_0, g_0) + d_{X^{<\omega}}^p(f'_2, g_2) = \\ &= d_{X^{<\omega}}^p(f_0, \alpha \circ q \circ f_0) + d_{X^{<\omega}}^p(\alpha \circ q \circ g_2, g_2) \leq \\ &\leq |S_a|^{\frac{1}{p}} \cdot \bar{d}_X(S_a) + |S_b|^{\frac{1}{p}} \cdot \bar{d}_X(S), \end{aligned}$$

where  $S = S_a \cup S_b$ .

Considering the  $(a, b)$ -linking chain  $((a_0, f_1, g'_1), (a_2, f_2, g_2))$  with  $g'_1 = \beta \circ q \circ g_2$ , we can see that

$$\begin{aligned} d_{FX}^p(a, b) &\leq d_{X^{<\omega}}^p(f_0, g'_1) + d_{X^{<\omega}}^p(f_2, g_2) = \\ &= d_{X^{<\omega}}^p(f_0, \beta \circ q \circ f_0) + d_{X^{<\omega}}^p(\beta \circ q \circ g_2, g_2) \leq \\ &\leq |S_a|^{\frac{1}{p}} \cdot \bar{d}_X(S) + |S_b|^{\frac{1}{p}} \cdot \bar{d}_X(S_b), \end{aligned}$$

where  $S = S_a \cup S_b$ . □

**Corollary 15.3.** *Assume that the distance  $d_X$  is a pseudometric and let  $q : X \rightarrow 1$  be the constant map. Then for any  $a, b \in FX$  the following statements hold.*

- 1) *If  $Fq(a) \neq Fq(b)$ , then  $d_{FX}^p(a, b) = \infty$ .*
- 2) *If  $Fq(a) = Fq(b)$ , then  $d_{FX}^p(a, b) \leq (|S_a|^{\frac{1}{p}} + |S_b|^{\frac{1}{p}}) \cdot \bar{d}_X(S_a \cup S_b)$ , where  $S_a = \text{supp}(a)$ ,  $S_b = \text{supp}(b)$ .*

**Corollary 15.4.** *If  $F$  is a functor of finite degree  $n = \deg(F)$ , then the distance space  $F^p X$  has real diameter*

$$\bar{d}_{F^p X}^p(FX) \leq 2 \cdot |n|^{\frac{1}{p}} \cdot \bar{d}_X(X).$$

## 16. LOWER BOUNDS ON THE DISTANCE $d_{FX}^p$

In this section we find some lower bounds on the distance  $d_{FX}^p$ . We recall that for a subset  $A \subseteq X$  its separatedness number  $\underline{d}_X(A)$  is defined as

$$\underline{d}_X(A) := \inf(\{\infty\} \cup \{d_X(x, y) : x, y \in A, x \neq y\}).$$

**Proposition 16.1.** *The distance space  $(FX, d_{FX}^p)$  has the separatedness number  $\underline{d}_{FX}^p(FX) \geq \underline{d}_X(X)$ .*

*Proof.* Assuming that  $\underline{d}_{FX}^p(FX) < \underline{d}_X(X)$ , we can find two distinct elements  $a, b \in FX$  with  $d_{FX}^p(a, b) < \underline{d}_X(X)$  and then find an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^k \in L_{FX}(a, b)$  such that  $\sum_{i=0}^k d_{X^{\leftarrow}}^p(f_i, g_i) < \underline{d}_X(X)$ . The definition of the separatedness number  $\underline{d}_X(X)$  ensures that  $f_i = g_i$  for all  $i \in \{0, \dots, k\}$ . Then

$$a = Ff_0(a_0) = Fg_0(a_0) = Ff_1(a_1) = Fg_1(a_1) = \dots = Fg_k(a_k) = b,$$

which contradicts the choice of the distinct elements  $a, b$ .  $\square$

We say that a distance space  $(Y, d_Y)$  is *separated* if it has non-zero separatedness number  $\underline{d}_Y(Y)$ . In this case its distance  $d_Y$  also is called *separated*. Proposition 16.1 implies:

**Corollary 16.2.** *If the distance space  $(X, d_X)$  is separated, then so is the distance space  $(FX, d_{FX}^p)$ .*

We will say that a distance space  $(X, d_X)$  is *Lipschitz disconnected* if for any distinct points  $x, y \in X$  there exists a Lipschitz map  $f : X \rightarrow \{0, 1\} \subset \mathbb{R}$  such that  $f(x) = 0$  and  $f(y) = 1$ . The Lipschitz property of  $f$  implies that  $1 = d_{\mathbb{R}}(f(x), f(y)) \leq \text{Lip}(f) \cdot d_X(x, y)$  and hence  $d_X(x, y) > 0$ , witnessing that *each Lipschitz disconnected distance is an  $\infty$ -metric*. On the other hand, *each separated distance space is Lipschitz disconnected*.

**Theorem 16.3.** *If the distance space  $(X, d_X)$  is Lipschitz disconnected, then the distance  $d_{FX}^p$  is an  $\infty$ -metric.*

*Proof.* Given two distinct points  $a, b \in FX$ , we should prove that  $d_{FX}^p(a, b) > 0$ . This inequality follows from Theorem 8.1 if the set  $S = \text{supp}(a) \cup \text{supp}(b)$  contains at most one point. So, we can assume that  $|S| > 1$ . In this case  $a, b \in F[S; X]$  by Proposition 6.1 and we can find elements  $a', b' \in FS$  such that  $a = Fi_{S, X}(a')$  and  $b = Fi_{S, X}(b')$ . Since  $a \neq b$ , the elements  $a', b'$  are distinct. Since the set  $S$  is finite, the  $\infty$ -metric  $d_S = d_X \upharpoonright S \times S$  is separated. By Corollary 16.2, the distance  $d_{FS}^p$  is separated and hence is an  $\infty$ -metric. Consequently,  $d_{FS}^p(a', b') > 0$ .

Using the Lipschitz disconnectedness of  $X$ , we can construct a Lipschitz map  $f : X \rightarrow \{0, 1\}^m \subset \mathbb{R}^m$  for a suitable  $m$  such that  $f \upharpoonright S$  is injective. Take any function  $g : \{0, 1\}^m \rightarrow S$  such that  $g \circ f(x) = x$  for every  $x \in S$ , and observe that the function  $r = g \circ f : X \rightarrow S$  is Lipschitz and  $r \circ i_{S, X}$  is the identity map of  $S$ . Then  $a' = Fr \circ Fi_{S, X}(a') = Fr(a)$  and  $b' = Fr(b)$ . By Theorem 12.1, the function  $F^p r : F^p X \rightarrow F^p S$  is Lipschitz. Taking into account that

$$0 < d_{FS}^p(a', b') = d_{FS}^p(Fr(a), Fr(b)) \leq \text{Lip}(F^p r) \cdot d_{FX}^p(a, b),$$

we conclude that  $d_{FX}^p(a, b) > 0$ , which means that the distance  $d_{FX}^p$  is an  $\infty$ -metric.  $\square$

Less trivial conditions implying that  $d_{FX}^p$  is an  $\infty$ -metric are given in the following theorems.

**Theorem 16.4.** *For any distinct elements  $a, b \in FX$  we have*

$$d_{FX}^1(a, b) \geq \underline{d}_X(\text{supp}(a) \cup \text{supp}(b)).$$

*This implies that  $d_{FX}^1$  is an  $\infty$ -metric if  $d_X$  is an  $\infty$ -metric.*

*Proof.* We need to prove that  $d_{FX}^\infty(a, b) \geq \underline{d}_X(S)$  where  $S := \text{supp}(a) \cup \text{supp}(b)$ . To derive a contradiction, assume that  $d_{FX}^1(a, b) < \underline{d}_X(S)$  and find an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^k \in L_{FX}(a, b)$  such that  $\sum_{i=0}^k d_{X^{<\omega}}^1(f_i, g_i) < \underline{d}_X(S)$ .

Let  $\sim$  be the smallest equivalence relation on  $X$  containing all pairs  $(f_i(z), g_i(z))$ , where  $i \leq k$  and  $z \in \text{dom}(f_i) = \text{dom}(g_i)$ . We claim that  $x \not\sim y$  for any distinct points  $x, y \in S$ . Assuming that  $x \sim y$ , we could find a sequence  $x = x_0, \dots, x_l = y$  of pairwise distinct points of  $X$  such that for every positive  $i \leq l$  there exist a number  $j_i \in \{0, \dots, k\}$  and a point  $z_i \in \text{dom}(f_{j_i}) = \text{dom}(g_{j_i})$  such that  $\{x_{i-1}, x_i\} = \{f_{j_i}(z_i), g_{j_i}(z_i)\}$ .

We claim that for any  $j \in \{0, \dots, k\}$  and  $z \in \text{dom}(f_j)$  the set  $I_{j,z} = \{i \in \{0, \dots, l\} : (j_i, z_i) = (j, z)\}$  contains at most one number. Otherwise, we could find two distinct numbers  $i, \tilde{i} \in I_{j,z}$  and conclude that

$$\{x_{i-1}, x_i\} = \{f_{j_i}(z_i), g_{j_i}(z_i)\} = \{f_j(z), g_j(z)\} = \{f_{j_i}(z_{\tilde{i}}), g_{j_i}(z_{\tilde{i}})\} = \{x_{\tilde{i}-1}, x_{\tilde{i}}\},$$

which is not possible as the points  $x_0, \dots, x_l$  are pairwise distinct.

Then

$$\begin{aligned} d_X(x, y) &\leq \sum_{i=1}^l d_X(x_{i-1}, x_i) = \sum_{i=1}^l d_X(f_{j_i}(z_i), g_{j_i}(z_i)) = \sum_{j=0}^k \sum_{z \in \text{dom}(f_j)} \sum_{i \in I_{j,z}} d_X(f_{j_i}(z_i), g_{j_i}(z_i)) = \\ &\sum_{j=0}^k \sum_{z \in \text{dom}(f_j)} |I_{j,z}| \cdot d_X(f_j(z), g_j(z)) \leq \sum_{j=0}^k \sum_{z \in \text{dom}(f_j)} d_X(f_j(z), g_j(z)) = \sum_{j=0}^k d_{X^{<\omega}}^1(f_j, g_j) < \underline{d}_X(S), \end{aligned}$$

which contradicts the definition of  $\underline{d}_X(S)$ . This contradiction completes the proof of non-equivalence  $x \not\sim y$  for any distinct points  $x, y \in S$ .

Now choose any function  $r : X \rightarrow S \subseteq X$  such that

- $r(x) = r(y)$  for any points  $x, y \in X$  with  $x \sim y$  and
- $r(x) = x$  for any point  $x \in S$ .

The choice of the function  $r$  is always possible since distinct points of the set  $S$  are not equivalent.

The definition of the equivalence relation  $\sim$  guarantees that  $r \circ f_i = r \circ g_i$  for any  $i \leq k$ . Consequently,

$$F(r \circ f_0)(a_0) = F(r \circ g_0)(a_0) = F(r \circ f_1)(a_1) = \dots = F(r \circ g_k)(a_k).$$

We claim that  $F(r \circ f_0)(a_0) = Ff_0(a_0) = a$ . Proposition 6.1 guarantees that  $a \in F[S; X]$ , so we can find an element  $a' \in FS$  such that  $a = Fi_{S,X}(a')$  where  $i_{S,X} : S \rightarrow X$  denotes the identity inclusion. The equality  $r \circ i_{S,X} = i_{S,X}$  implies that  $Fr \circ Fi_{S,X} = Fi_{S,X}$  and hence

$$a = Fi_{S,X}(a') = Fr \circ Fi_{S,X}(a') = Fr(a) = Fr(Ff_0(a_0)) = F(r \circ f_0)(a_0).$$

By analogy we can prove that  $F(r \circ g_k)(b_k) = Fr(b) = b$ . Then

$$a = F(r \circ f_0)(a_0) = F(r \circ g_k)(b_k) = b,$$

which contradicts the choice of the (distinct) points  $a, b$ . □

Theorems 16.4 and 11.2 imply the following corollary.

**Corollary 16.5.** *If  $d_X$  is an  $\infty$ -metric and the functor  $F$  has finite degree  $m = \deg(F)$ , then for every  $p \in [1, \infty]$  the distance  $d_{FX}^p$  is an  $\infty$ -metric. Moreover, for any distinct elements  $a, b \in FX$  either  $d_{FX}^p(a, b) = \infty$  or*

$$d_{FX}^p(a, b) \geq m^{\frac{1-p}{p}} \cdot d_{FX}^1(a, b) \geq m^{\frac{1-p}{p}} \cdot \underline{d}_X(\text{supp}(a) \cup \text{supp}(b)) > 0.$$

Another condition guaranteeing that the distance  $d_{FX}^p$  is an  $\infty$ -metric was found by Shukel in [29]. Theorem 16.6 below is a modification of Theorem 1 of [29].

**Theorem 16.6.** *If the functor  $F$  preserves supports, then for any distinct elements  $a, b \in FX$  we have*

$$d_{FX}^p(a, b) \geq d_{FX}^\infty(a, b) \geq \max \{d_{HX}(\text{supp}(a), \text{supp}(b)), \frac{1}{3}\underline{d}_X(\text{supp}(a) \cup \text{supp}(b))\}.$$

*Proof.* The inequality  $d_{FX}^p(a, b) \geq d_{FX}^\infty(a, b)$  follows from Theorem 11.2. It remains to prove that  $d_{FX}^\infty(a, b) \geq \varepsilon$  where

$$\varepsilon := \max \{d_{HX}(S_a, S_b), \frac{1}{3}\underline{d}_X(S)\},$$

$S_a := \text{supp}(a)$ ,  $S_b = \text{supp}(b)$ , and  $S = S_a \cup S_b$ .

To derive a contradiction, assume that  $d_{FX}^\infty(a, b) < \varepsilon$  and find an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^k \in L_{FX}(a, b)$  such that  $\sum_{i=0}^k d_{X^{<\omega}}^\infty(f_i, g_i) < \varepsilon$ . We can assume that the length  $k$  of the chain is the smallest possible.

For every  $i \in \{0, \dots, k\}$ , consider the element  $b_i = Ff_i(a_i) \in FX$  and put  $b_{k+1} = Fg_k(a_k)$ . It follows that  $b_0 = a$ ,  $b_{k+1} = b$ , and  $b_i = Ff_i(a_i) = Fg_{i-1}(a_{i-1})$  for any  $i \in \{1, \dots, k\}$ . The minimality of  $k$  ensures that  $b_i \neq b_{i+1}$  for all  $i \in \{0, \dots, k\}$ .

Taking into account that the functor  $F$  preserves supports, we conclude that  $\text{supp}(b_i) = f_i[\text{supp}(a_i)]$  for every  $i \in \{0, \dots, k\}$ . We claim that for every  $i \in \{0, \dots, k\}$  the support  $S_i = \text{supp}(a_i)$  is not empty. Assuming that  $S_i = \emptyset$ , we conclude that  $\text{supp}(b_i) = f_i(\text{supp}(a_i)) = \emptyset = g_i(\text{supp}(a_i)) = \text{supp}(b_{i+1})$ . Since  $b_i \neq b_{i+1}$ , Theorem 8.1 ensures that  $d_{FX}^\infty(b_i, b_{i+1}) = \infty$ . On the other hand, the definition of the distance  $d_{FX}^\infty$  yields  $d_{FX}^\infty(b_i, b_{i+1}) = d_{FX}^\infty(Ff_i(a_i), Fg_i(a_i)) \leq d_{X^{<\omega}}^\infty(f_i, g_i) < \varepsilon \leq \infty$ . This contradiction shows that  $S_i \neq \emptyset$  and hence  $\text{supp}(b_i) = f(\text{supp}(a_i))$  is not empty, too.

The chain  $a = b_0, \dots, b_{k+1} = b$  of elements of  $FX$  induces the chain of finite non-empty subsets

$$S_a = \text{supp}(b_0), \dots, \text{supp}(b_{k+1}) = S_b$$

in  $X$ . Then by the triangle inequality for the Hausdorff distance  $d_{HX}$ , we obtain

$$\begin{aligned} d_{HX}(S_a, S_b) &\leq \sum_{i=0}^k d_{HX}(\text{supp}(b_i), \text{supp}(b_{i+1})) = \sum_{i=0}^k d_{HX}(f_i[\text{supp}(a_i)], g_i[\text{supp}(a_i)]) \leq \\ &\leq \sum_{i=0}^k d_{X^{<\omega}}^\infty(f_i, g_i) < \varepsilon = \max \{d_{HX}(S_a, S_b), \frac{1}{3}\underline{d}_S(S)\}, \end{aligned}$$

which implies that  $d_{HX}(S_a, S_b) < \varepsilon = \frac{1}{3}\underline{d}_X(S)$  and  $S_a = S_b = S$  by the definition of  $d_{HX}(S_a, S_b)$  and  $\underline{d}_X(S)$ .

Let  $O[S; \varepsilon] = \{x \in X : \exists s \in S \ d_X(s, x) < \varepsilon\}$  be the open  $\varepsilon$ -neighborhood of the finite non-empty set  $S$  in the distance space  $(X, d_X)$ .

Let  $r : X \rightarrow S \subseteq X$  be any function such that  $r(x, y) = x$  for any  $x \in S$  and  $y \in X$  with  $d(y, x) < \varepsilon$ . We claim that  $r \circ f_i \upharpoonright S_i = r \circ g_i \upharpoonright S_i$  for any  $i \in \{0, \dots, k\}$ . Indeed,

$$d_{HX}(S, \text{supp}(b_i)) \leq \sum_{j=0}^{i-1} d_{HX}(\text{supp}(b_j), \text{supp}(b_{j+1})) \leq \sum_{j=0}^{i-1} d_{X^{<\omega}}^\infty(f_j, g_j) < \varepsilon$$

and hence  $f_i(\text{supp}(a_i)) = \text{supp}(b_i) \subseteq O[S; \varepsilon]$  and  $g_i(\text{supp}(a_i)) = \text{supp}(Fg_i(a_i)) = \text{supp}(b_{i+1}) \subseteq O[S; \varepsilon]$ . As we already know, for every  $i \in \{0, \dots, k\}$  the support  $S_i = \text{supp}(a_i)$  is a non-empty set of the finite ordinal  $n_i := \text{dom}(f_i) = \text{dom}(g_i) \in \mathbb{N}$ .

The inclusions  $f_i[\text{supp}(a_i)] \cup g_i[\text{supp}(a_i)] \subseteq O[S; \varepsilon]$  and the inequality  $d_{X^{<\omega}}^\infty(f_i, g_i) < \varepsilon = \frac{1}{3} \underline{d}_X(S)$  imply that for any  $x \in S_i$  there exists a point  $s \in S$  with  $f_i(x), g_i(x) \in O[s; \varepsilon]$ . Consequently,  $r \circ f_i(x) = s = r \circ g_i(x)$ .

Proposition 6.1 guarantees that  $a_i \in F[\text{supp}(a_i); X]$  and hence  $a_i = Fi_{S_i, n_i}(a'_i)$  for some  $a'_i \in F(S_i)$ , where  $i_{S_i, n_i} : S_i \rightarrow n_i$  denotes the identity embedding of  $S_i$  into  $n_i$ . It follows from  $r \circ f_i \upharpoonright S_i = r \circ g_i \upharpoonright S_i$  that  $r \circ f_i \circ i_{S_i, n_i} = r \circ g_i \circ i_{S_i, n_i}$  and hence

$$\begin{aligned} (1) \quad Fr \circ Ff_i(a_i) &= Fr \circ Ff_i \circ Fi_{S_i, n_i}(a'_i) = \\ &= Fr \circ Fg_i \circ Fi_{S_i, n_i}(a'_i) = Fr \circ Fg_i(a_i) = Fr \circ Ff_{i+1}(a_{i+1}). \end{aligned}$$

Since  $\text{supp}(a) \subseteq S$ , by Proposition 6.1, there exists an element  $a' \in FS$  such that  $a = Fi_{S, X}(a')$ . Applying the functor  $F$  to the equality  $r \circ i_{S, X} = i_{S, X}$ , we conclude that  $Fr(a) = Fr \circ Fi_{S, X}(a') = Fi_{S, X}(a') = a$ . By analogy we can prove that  $Fr(b) = b$ . Taking into account the equality (1), we finally conclude that

$$\begin{aligned} a = Fr(a) &= Fr \circ Ff_0(a_0) = Fr \circ Fg_0(a_0) = Fr \circ Ff_1(a_1) = Fr \circ Fg_1(a_1) = \dots \\ &\dots = Fr \circ Fg_k(a_k) = Fr(b) = b, \end{aligned}$$

which contradicts the choice of the (distinct) points  $a, b$ .  $\square$

## 17. PRESERVATION OF CONTINUOUS FUNCTIONS BY THE FUNCTOR $F^p$

**Theorem 17.1.** *If the functor  $F$  is finitary, has finite degree and preserves supports, then for any continuous function  $f : X \rightarrow Y$  between distance spaces, the function  $F^p f : F^p X \rightarrow F^p Y$  is continuous.*

*Proof.* Let  $n = \max\{1, \deg(F)\}$ . To check the continuity of  $F^p f$ , fix any any real number  $\varepsilon \in (0, \infty)$  and any element  $a \in FX$ . By the continuity of  $f$ , there exists  $\delta \in (0, \infty)$  such that  $f(O(x, \delta)) \subseteq O(f(x), \varepsilon/\sqrt[n]{n})$  for all  $x \in \text{supp}(a)$ . Choose  $\delta' \in (0, \infty)$  such that

$$\delta' < \frac{1}{2n \cdot |Fn|} \min\{\delta, \underline{d}_X(\text{supp}(a))\}.$$

Given any element  $b \in FX$  with  $d_{FX}^p(a, b) < \delta'$ , we will prove that  $d_{FY}^p(Ff(a), Ff(b)) < \varepsilon$ . This inequality is trivially true if  $a = b$ . So we assume that  $a \neq b$ . By Theorem 8.1,  $d_{FX}^p(a, b) < \delta' < \infty$  implies that  $|S_a \cup S_b| > 1$ , where  $S_a := \text{supp}(a)$  and  $S_b := \text{supp}(b)$ .

By Lemma 9.3, there exists a function  $r : X \rightarrow X$  such that  $Fr(b) = Fr(a)$ ,  $r[S_a] \subseteq S_a$ , and  $\sup_{x \in X} d_X(r(x), x) < n \cdot |Fn| \cdot \delta' < \frac{1}{2} \underline{d}_X(S_a)$ . The latter strict inequality and the inclusion  $r[S_a] \subseteq S_a$  imply that  $r(x) = x$  for every  $x \in S_a$ .

Since  $F$  preserves supports,

$$r[\text{supp}(b)] = \text{supp}(Fr(b)) = \text{supp}(Fr(a)) = r[\text{supp}(a)] = \text{supp}(a)$$

and hence both sets  $S_a$  and  $S_b$  are not empty (because  $|S_a \cup S_b| > 1$ ). By Proposition 6.1, there are functions  $g, h \in X^n$  and elements  $a' \in Fn$  and  $b' \in Fn$  such that  $a = Fg(a')$ ,  $g[n] = S_a$ , and  $b = Fh(b')$ ,  $h[n] = S_b$ . Taking into account that  $r \circ g = g$ , we conclude that

$$F(r \circ h)(b') = Fr(Fh(b')) = Fr(b) = Fr(a) = Fr \circ Fg(a') = F(r \circ g)(a') = Fg(a') = a.$$

Then  $((b', r \circ h, h))$  is an  $(a, b)$ -linking chain and hence  $((b', f \circ r \circ h, f \circ h))$  is an  $(Ff(a), Ff(b))$ -linking chain.

For every  $x \in S_b$ , we have  $r(x) \in S_a$  and  $d_X(r(x), x) < n \cdot |Fn| \cdot \delta \leq \delta'$ , which implies  $d_Y(f(r(x)), f(x)) < \varepsilon/\sqrt[n]{n}$  by the choice of  $\delta'$ . Then

$$d_{FY}^p(Ff(a), Ff(b)) \leq d_{Yn}^p(f \circ r \circ h, f \circ h) = \left( \sum_{k \in n} d_Y(f \circ r \circ h(k), f \circ h(k)) \right) < \sqrt[n]{n} \cdot \frac{\varepsilon}{\sqrt[n]{n}} = \varepsilon.$$

□

**Example 17.2.** Consider the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  assigning to each set  $X$  the set  $FX = \{\emptyset\} \cup \{\{x, y\} : x, y \in X, x \neq y\}$  and to each function  $f : X \rightarrow Y$  the function  $Ff : FX \rightarrow FY$  such that  $Ff(\emptyset) = \emptyset$  and  $Ff(\{x, y\}) = \{f(x)\} \triangle \{f(y)\}$  for any  $\{x, y\} \in FX \setminus \{\emptyset\}$ . It is clear that  $F$  is finitary and has finite degree  $\deg(F) = 2$ , but  $F$  does not preserve supports. It can be shown that for the continuous map  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : x \mapsto x^3$ , the map  $F^p f : F^p \mathbb{R} \rightarrow F^p \mathbb{R}$  is not continuous. This example shows that the preservation of supports in Theorem 17.1 is essential.

## 18. METRIC PROPERTIES OF THE MAPS $\xi_X^a : X \rightarrow FX$ .

Given any element  $a \in F1$  consider the map  $\xi_X^a : X^1 \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ . If the set  $X$  is not empty, the the power  $X^1$  can be identified with  $X$  by identifying each point  $x \in X$  with the function  $\bar{x}_X : 1 \rightarrow \{x\} \subseteq X$ . In this case  $\xi_X^a(x) = F\bar{x}_X(a)$ . If  $X$  is empty, then  $X^1$  is empty and hence is equal to  $X$ . In both cases we can consider the map  $\xi_X^a$  as a function defined on  $X$ .

The following proposition shows that  $\xi^a$  is a natural transformation of the identity functor into the functor  $F$ .

**Proposition 18.1.** *For any  $a \in F1$  and any function  $f : X \rightarrow Y$  between sets, we have  $Ff \circ \xi_X^a = \xi_Y^a \circ f$ .*

*Proof.* For every  $x \in X$  and the map  $\bar{x} : 1 \rightarrow \{x\} \subseteq X$ , we have

$$Ff \circ \xi_X^a(x) = Ff \circ F\bar{x}_X(a) = F(f \circ \bar{x}_X)(a) = \xi_Y^a(f(x)) = \xi_Y^a \circ f(x).$$

□

By Theorem 10.1, for every  $a \in F1$  the map  $\xi_X^a : (X, d_X) \rightarrow (FX, d_{FX}^p)$  is non-expanding. In this section we find conditions guaranteeing that this map is an injective isometry.

**Lemma 18.2.** *Let  $a \in F1$  and assume that  $|X| \geq 2$ .*

- 1) *If the map  $\xi_2^a : 2 \rightarrow F2$  is not injective, then*
  - 1a) *the map  $\xi_X^a : X \rightarrow FX$  is constant;*
  - 1b)  *$\text{supp}(\xi_X^a(x)) = \emptyset$  for every  $x \in X$ .*
- 2) *If the map  $\xi_2^a : 2 \rightarrow F2$  is injective, then*
  - 2a)  *$a \notin F[\emptyset; 1] \subseteq F1$ ;*
  - 2b)  *$\xi_2^a(0) \notin F[\{1\}, 2]$ ;*
  - 2c)  *$\xi_X^a(x) \notin F[X \setminus \{x\}; X]$  for every  $x \in X$ ;*



- 2d)  $\text{supp}(\xi_X^a(x)) = \{x\}$  for any  $x \in X$ ;  
 2e) the map  $\xi_X^a : X \rightarrow FX$  is injective.

*Proof.* 1. Assume that the map  $\xi_2^a : 2 \rightarrow F2$  is not injective. Then  $F\bar{0}_2(a) = \xi_2^a(0) = \xi_2^a(1) = F\bar{1}_2(a)$ .

1a. To prove that the map  $\xi_X^a : X \rightarrow FX$  is constant, choose any distinct points  $x, y \in X$ . Let  $f : 2 \rightarrow X$  be the map defined by  $f(0) = x, f(1) = y$ . It follows that  $\bar{x}_X = f \circ \bar{0}_2$  and  $\bar{y}_X = f \circ \bar{1}_2$ . Then

$$\begin{aligned} \xi_X^a(x) = F\bar{x}_X(a) &= Ff \circ F\bar{0}_2(a) = Ff(\xi_2^a(0)) = Ff(\xi_2^a(1)) = \\ &= Ff \circ F\bar{1}_2(a) = F\bar{y}_X(a) = \xi_X^a(y). \end{aligned}$$

1b. Therefore, for any points  $x, y \in X$  we have  $\xi_X^a(x) = \xi_X^a(y) = F\bar{y}_X(a) \in F[\{y\}; X]$  and hence  $\text{supp}(\xi_X^a(x)) \subseteq \bigcap_{y \in X} \{y\} = \emptyset$ .

2. Assume that the map  $\xi_2^a : 2 \rightarrow F2$  is injective.

2a. First we show that  $a \notin F[\emptyset; 1]$ . To derive a contradiction, assume that  $a \in F[\emptyset; 1]$ . In this case we can find an element  $a' \in F\emptyset$  such that  $a = Fi_{\emptyset,1}(a')$ . It follows that for every  $z \in 2$

$$\xi_2^a(z) = F\bar{z}_2(a) = F\bar{z}_2(Fi_{\emptyset,1}(a')) = F(\bar{z}_2 \circ i_{\emptyset,1})(a') = Fi_{\emptyset,2}(a')$$

and hence the map  $\xi_2^a : 2 \rightarrow \{Fi_{\emptyset,2}(a')\} \subseteq F2$  is constant, which contradicts our assumption. This contradiction shows that  $a \notin F[\emptyset; 1]$ .

2b. Now we prove that  $\xi_2^a(0) \notin F[\{1\}; 2]$ . To derive a contradiction, assume that  $\xi_2^a(0) \in F[\{1\}; 2] = Fi_{\{1\},2}[F\{1\}]$ . Then we can find an element  $a' \in F\{1\}$  such that  $\xi_2^a(0) = Fi_{\{1\},2}(a')$ . Observe that  $\bar{1}_2 = i_{\{1\},2} \circ g$  where  $g : 1 \rightarrow \{1\}$  is the unique bijection. Since  $Fg : F1 \rightarrow F\{1\}$  is a bijection, there exists an element  $a'' \in F1$  such that  $a' = Fg(a'')$ . It follows that

$$F\bar{0}_2(a) = \xi_2^a(0) = Fi_{\{1\},2}(a') = Fi_{\{1\},2}(Fg(a'')) = F(i_{\{1\},2} \circ g)(a'') = F\bar{1}_2(a'').$$

Now consider the constant function  $\gamma : 2 \rightarrow 1$  and observe that  $\gamma \circ \bar{0}_2 = \gamma \circ \bar{1}_2$  is the identity map of the singleton 1. Then  $F\gamma \circ F\bar{0}_2 = F\gamma \circ F\bar{1}_2$  is the identity map of  $F1$ . Consequently,

$$a'' = F\gamma \circ F\bar{1}_2(a'') = F\gamma \circ F\bar{0}_2(a) = a$$

and then  $\xi_2^a(0) = F\bar{0}_2(a) = F\bar{1}_2(a'') = F\bar{1}_2(a) = \xi_2^a(1)$ , which contradicts the injectivity of the map  $\xi_2^a : 2 \rightarrow F2$ .

2c. Given any  $x \in X$  we will show that  $\xi_X^a(x) \notin F[X \setminus \{x\}; X]$ . To derive a contradiction, assume that  $\xi_X^a(x) \in F(A; X)$  where  $A := X \setminus \{x\}$ . Then  $\xi_X^a(x) = Fi_{A,X}(a')$  for some  $a' \in FA$ . Let  $\phi : A \rightarrow \{1\}$  be the constant function and  $\chi : X \rightarrow 2$  be the function defined by  $\chi(x) = 0$  and  $\chi(a) = 1$  for any  $a \in A$ . It follows that  $\bar{0}_2 = \chi \circ \bar{x}_X$ . Then

$$\begin{aligned} \xi_2^a(0) = F\bar{0}_2(a) &= F\chi \circ F\bar{x}_X(a) = F\chi(\xi_X^a(x)) = F\chi \circ Fi_{A,X}(a') = \\ &= Fi_{\{1\},2} \circ F\phi(a') \in Fi_{\{1\},2}[F\{1\}] \subseteq F[\{1\}, 2], \end{aligned}$$

which contradicts (2b).

2d. Now we are able to prove that  $\text{supp}(\xi_X^a(x)) = \{x\}$  for every  $x \in X$ . It follows from

$$\xi_X^a(x) = F\bar{x}_X(a) = Fi_{\{x\},X} \circ F\bar{x}_{\{x\}}(a) \in Fi_{\{x\},X}[F\{x\}] = F[\{x\}; X]$$

that  $\text{supp}(\xi_X^a(x)) \subseteq \{x\}$ .

Assuming that  $\text{supp}(\xi_X^a(x)) \neq \{x\}$ , we conclude that  $\text{supp}(\xi_X^a(x)) = \emptyset$  and hence  $\xi_X^a(x) \in F[X \setminus \{x\}; X]$ , which contradicts the statement (2c).

2e. By the statement (2d), for any distinct points  $x, y \in X$  the elements  $\xi_X^a(x)$  and  $\xi_X^a(y)$  are distinct as they have distinct supports. So, the function  $\xi_X^a : X \rightarrow FX$  is injective.  $\square$

**Lemma 18.3.** *For any element  $a \in F1$ , the map  $\xi_X^a : (X, d_X) \rightarrow (FX, d_{FX}^p)$  is an injective isometry if the map  $\xi_2^a : 2 \rightarrow F2$  is injective, the distance  $d_X$  is an  $\infty$ -metric and either  $p = 1$  or  $F$  preserves supports.*

*Proof.* Assume that the map  $\xi_2^a : 2 \rightarrow F2$  is injective,  $d_X$  is an  $\infty$ -metric and either  $p = 1$  or  $F$  preserves supports. By Lemma 18.2 and Theorem 10.1, the map  $\xi_X^a : (X, d_X) \rightarrow (FX, d_{FX}^p)$  is injective and non-expanding. To show that  $\xi_X^a$  is an isometry, fix any distinct elements  $x, y \in X$ . We need to show that  $d_{FX}^p(\xi_X^a(x), \xi_X^a(y)) \geq d_X(x, y)$ . By Lemma 18.2(2d),  $\text{supp}(\xi_X^a(z)) = \{z\}$  for any  $z \in X$ .

If  $p = 1$ , then by Theorem 16.4,

$$\begin{aligned} d_{FX}^p(\xi_X^a(x), \xi_X^a(y)) &\geq \min\{d_X(u, v) : u, v \in \text{supp}(\xi_X^a(x)) \cup \text{supp}(\xi_X^a(y)), u \neq v\} = \\ &= \min\{d_X(u, v) : u, v \in \{x\} \cup \{y\}, u \neq v\} = d_X(x, y). \end{aligned}$$

Next, we consider the case of functor  $F$  that preserves supports. To derive a contradiction, assume that  $d_{FX}^p(\xi_X^a(x), \xi_X^a(y)) < d_X(x, y)$  and hence

$$d_{FX}^\infty(\xi_X^a(x), \xi_X^a(y)) \leq d_{FX}^p(\xi_X^a(x), \xi_X^a(y)) < d_X(x, y)$$

according to Lemma 11.1.

By the definition of the distance  $d_{FX}^\infty$ , there exists a  $(\xi_X^a(x), \xi_X^a(y))$ -linking chain  $((a_i, f_i, g_i))_{i=0}^k \in L_{FX}(a, b)$  such that  $\sum_{i=0}^k d_{X^{\leftarrow \omega}}^\infty(f_i, g_i) < d_X(x, y)$ . Let  $b_0 := \xi_X^a(x) = Ff_0(a_0)$ ,  $b_{k+1} := \xi_X^a(y) = Fg_k(a_k)$  and  $b_i = Ff_i(a_i) = Fg_{i-1}(a_{i-1})$  for  $i \in \{1, \dots, k\}$ . Since the functor  $F$  preserves supports,

$$\text{supp}(b_i) = f_i[\text{supp}(a_i)] \quad \text{and} \quad \text{supp}(b_{i+1}) = g_i[\text{supp}(a_i)]$$

for all  $i \in \{0, \dots, k\}$ . Consequently,  $d_{HX}(\text{supp}(b_i), \text{supp}(b_{i+1})) \leq d_{X^{\leftarrow \omega}}^\infty(f_i, g_i)$  for every  $i \in \{0, \dots, k\}$ . Here by  $d_{HX}$  we denote the Hausdorff distance on the hyperspace  $[X]^{\leftarrow \omega}$  of all finite subsets of  $X$ . It follows that

$$\begin{aligned} d_X(x, y) &= d_{HX}(\{x\}, \{y\}) = d_{HX}(\text{supp}(b_0), \text{supp}(b_{k+1})) \leq \\ &\leq \sum_{i=0}^k d_{HX}(\text{supp}(b_i), \text{supp}(b_{i+1})) \leq \sum_{i=0}^k d_{X^{\leftarrow \omega}}^\infty(f_i, g_i) < d_X(x, y), \end{aligned}$$

which is a desired contradiction completing the proof of the inequality

$$d_{FX}^p(\xi_X^a(x), \xi_X^a(y)) \geq d_X(x, y). \quad \square$$

Now we can prove the main result of this section.

**Theorem 18.4.** *For an element  $a \in F1$ , the map  $\xi_X^a : (X, d_X) \rightarrow (FX, d_{FX}^p)$  is an injective isometry if the map  $\xi_2^a : 2 \rightarrow F2$  is injective and either  $p = 1$  or  $F$  preserves supports.*

*Proof.* Assume that the map  $\xi_2^a : 2 \rightarrow F2$  is injective and either  $p = 1$  or  $F$  preserves supports. By Lemma 18.2 and Theorem 10.1, the map  $\xi_X^a : (X, d_X) \rightarrow (FX, d_{FX}^p)$  is injective and non-expanding. To show that  $\xi_X^a$  is an isometry, it remains to check that  $d_{FX}^p(\xi_X^a(x), \xi_X^a(y)) \geq d_X(x, y)$  for any points  $x, y \in X$ .

On the set  $X$  consider the equivalence relation  $\approx$  defined by  $x \approx y$  iff  $d_X(x, y) = 0$ . For any  $x \in X$  let  $[x] := \{y \in X : d_X(x, y) = 0\}$  be its equivalence class. The quotient set  $\tilde{X} = \{[x] : x \in X\}$  of  $X$  by the equivalence relation  $\approx$  carries a unique  $\infty$ -metric  $d_{\tilde{X}}$  such that the quotient map  $q : X \rightarrow \tilde{X}$ ,  $q : x \mapsto [x]$ , is an isometry. Since  $d_{\tilde{X}}$  is an  $\infty$ -metric, we can apply Lemma 18.3 and conclude that for any points  $x, y \in \tilde{X}$  we have the equality  $d_{F\tilde{X}}^p(\xi_{\tilde{X}}^a(x), \xi_{\tilde{X}}^a(y)) = d_{\tilde{X}}(x, y)$ .

By Theorem 12.1, the map  $Fq : (FX, d_{FX}^p) \rightarrow (F\tilde{X}, d_{F\tilde{X}}^p)$  is non-expanding. Consequently, for any points  $x, y \in X$  we obtain the desired inequality:

$$\begin{aligned} d_{FX}^p(\xi^a(x), \xi^a(y)) &\geq d_{F\tilde{X}}^p(Fq \circ \xi_X^a(x), Fq \circ \xi_X^a(y)) = \\ &= d_{F\tilde{X}}^p(\xi_{\tilde{X}}^a(q(x)), \xi_{\tilde{X}}^a(q(y))) = d_{\tilde{X}}(q(x), q(y)) = d_X(x, y), \end{aligned}$$

witnessing that the map  $\xi_X^a : X \rightarrow F^p X$  is an isometry.  $\square$

## 19. PRESERVING THE COMPACTNESS BY THE FUNCTOR $F^p$

A distance space  $(X, d_X)$  is called *compact* if each sequence in  $X$  contains a convergent subsequence. It is standard to show that a distance space  $X$  is compact if and only if it is compact in the topological sense (i.e., each open cover of  $X$  has a finite subcover). Here on  $X$  we consider the topology consisting of sets  $U \subseteq X$  such that for every  $x \in X$  there exists  $\varepsilon > 0$  such that the  $\varepsilon$ -ball  $O(x; \varepsilon) = \{y \in X : d_X(y, x) < \varepsilon\}$  is contained in  $U$ . Discussing topological properties of distance spaces we have in mind this topology generated by the distance. The topology of a distance space  $(X, d_X)$  is Hausdorff if and only if the distance  $d_X$  is an  $\infty$ -metric.

**Theorem 19.1.** *If the functor  $F$  is finitary and has finite degree, then:*

- 1) *For any compact distance space  $(X, d_X)$  the distance space  $(FX, d_{FX})$  is compact;*
- 2) *For any continuous map  $f : (X, d_X) \rightarrow (Y, d_Y)$  between compact distance spaces the map  $Ff : (FX, d_{FX}^p) \rightarrow (FY, d_{FY}^p)$  is continuous.*

*Proof.* 1. Let  $(X, d_X)$  be a compact distance space. If  $X = \emptyset$ , then  $FX = F\emptyset$  is finite and hence compact, by the finitariness of the functor  $F$ . So, we assume that  $X \neq \emptyset$ .

Since the functor  $F$  has finite degree, there exists  $n \in \mathbb{N}$  such that the map  $\xi_X : Fn \times X^n \rightarrow FX$ ,  $\xi_X : (a, f) \mapsto Ff(a)$ , is surjective. The finitary property of the functor  $F$  guarantees that the space  $Fn$  is finite. Endow the space  $Fn$  with the discrete topology.

By Theorem 10.1, for every  $a \in Fn$  the map  $\xi_X^a : (X^n, d_{X^{<\omega}}^n) \rightarrow (FX, d_{FX}^p)$  is non-expanding and hence continuous. Then the distance space  $(FX, d_{FX}^p)$  is compact, being a continuous image of the compact topological space  $Fn \times X^n$ .

2. Let  $f : X \rightarrow Y$  be any continuous map between compact distance spaces  $(X, d_X)$  and  $(Y, d_Y)$ . By the compactness of  $X$ , the map  $f$  is uniformly continuous. By Corollary 14.2, the map  $F^p f : F^p X \rightarrow F^p Y$  is uniformly continuous and hence continuous.  $\square$

## 20. PRESERVING THE COMPLETENESS BY THE FUNCTOR $F^p$

In this section we address the problem of preservation of the completeness of distance spaces by the functor  $F^p$ . The completeness of distance spaces is defined by analogy with the completeness of metric spaces.

A sequence  $\{x_n\}_{n \in \omega} \subseteq X$  in a distance space  $(X, d_X)$  is *Cauchy* if for every  $\varepsilon \in (0, +\infty)$  there exists  $n \in \omega$  such that  $d_X(x_m, x_k) < \varepsilon$  for any  $m, k \geq n$ .

A distance space  $(X, d_X)$  is *complete* if every Cauchy sequence  $\{x_n\}_{n \in \omega} \subseteq X$  converges to some point  $x_\infty \in X$ . The latter means that for every  $\varepsilon \in (0, +\infty)$  there exists  $n \in \omega$  such that  $d_X(x_m, x_\infty) < \varepsilon$  for any  $m \geq n$ .

The following theorem is the principal result of this section.

**Theorem 20.1.** *Assume that the functor  $F$  is finitary and has finite degree. If a distance space  $(X, d_X)$  is complete, then so is the distance space  $F^p X$ .*

The proof of this theorem is preceded by several definitions and lemmas. Let  $\Omega$  be an infinite subset of  $\omega$ . A sequence  $\{x_n\}_{n \in \Omega} \subseteq X$  is called

- *convergent* if there exists  $x \in X$  such that  $\lim_{\Omega \ni i \rightarrow \infty} d_X(x_i, x) = 0$ , i.e.,  
 $\exists x \in X \forall \varepsilon \in (0, +\infty) \exists F \in [\Omega]^{<\omega} \forall i \in \Omega \setminus F \ d_X(x_i, x) < \varepsilon$ ;
- *Cauchy* if  $\lim_{\Omega \ni i, j \rightarrow \infty} d_X(x_i, x_j) = 0$ , i.e.,  
 $\forall \varepsilon \in (0, \infty) \exists F \in [\Omega]^{<\omega} \forall i, j \in \Omega \setminus F \ d_X(x_i, x_j) < \varepsilon$ ;
- $\varepsilon$ -*separated* for some  $\varepsilon \in (0, \infty)$  if  $d_X(x_i, x_j) \geq \varepsilon$  for any distinct numbers  $i, j \in \Omega$ ;
- *separated* if it is  $\varepsilon$ -separated for some  $\varepsilon > 0$ .

The following lemma is well-known for metric spaces and can be proved by analogy for distance spaces.

**Lemma 20.2.** *A Cauchy sequence  $(x_n)_{n \in \omega}$  in a distance space is convergent if and only if contains a convergent subsequence  $(x_n)_{n \in \Omega}$ .*

For an infinite subset  $\Omega$  of  $\omega$  we put  $[\Omega]^2 := \{(i, j) \in \Omega \times \Omega : i < j\}$ .

**Lemma 20.3.** *For every  $n \in \mathbb{N}$ , and every sequence  $\{f_i\}_{i \in \omega} \subseteq X^n$  there exists an infinite subset  $\Omega \subseteq \omega$  such that for any  $k, m \in n$  either*

$$\lim_{\Omega \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(m)) = 0 \quad \text{or} \quad \inf_{(i, j) \in [\Omega]^2} d_X(f_i(k), f_j(m)) > 0.$$

*Proof.* To prove the lemma for  $n = 1$ , it suffices to check that each sequence  $\{x_i\}_{i \in \omega} \subseteq X$  contains a subsequence  $(x_i)_{i \in \Omega}$ , which is either separated or Cauchy. Assuming that  $(x_n)_{n \in \omega}$  contains no separated subsequences, we will construct a Cauchy subsequence in  $(x_n)_{n \in \omega}$ .

For every  $k \in \omega$ , consider the function  $\chi_k : [\omega]^2 \rightarrow \{0, 1\}$  defined by

$$\chi_k(i, j) = \begin{cases} 0, & \text{if } d_X(x_i, x_j) < 2^{-k} \\ 1, & \text{otherwise.} \end{cases}$$

Applying the Ramsey Theorem [31, 1.3], find an infinite subset  $\Omega_0$  of  $\omega$  such that  $\chi_0[[\Omega_0]^2]$  is a singleton. If  $\chi_0[[\Omega_0]^2] = \{1\}$ , then  $(x_n)_{n \in \Omega_0}$  is a 1-separated subsequence of  $(x_n)_{n \in \omega}$ , which is forbidden by our assumption. Therefore,  $\chi_0[[\Omega_0]^2] = \{0\}$ . Applying the Ramsey Theorem inductively, we can construct a decreasing sequence of infinite sets  $(\Omega_k)_{k \in \omega}$  such that  $\chi_k[[\Omega_k]^2] = \{0\}$  for every  $k \in \omega$ . After completing the inductive construction, choose an infinite set  $\Omega \subseteq \omega$  such that  $\Omega \setminus \Omega_k$  is finite for every  $k \in \omega$ , and observe that the sequence  $(x_n)_{n \in \Omega}$  is Cauchy. This completes the proof of the lemma for  $n = 1$ .

Now assume that for some  $n \geq 2$  we have proved that for every  $l < n$ , every sequence  $\{f_i\}_{i \in \omega} \subseteq X^l$  contains a subsequence  $(f_i)_{i \in \Omega}$  such that for every  $k, m \in l$ , either

$$\lim_{\Omega \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(m)) = 0 \quad \text{or} \quad \inf_{(i, j) \in [\Omega]^2} d_X(f_i(k), f_j(m)) > 0.$$

Fix any sequence  $\{f_i\}_{i \in \omega} \subseteq X^n$ . By the inductive assumption, there exists an infinite subset  $\Omega' \subseteq \omega$  such that for every  $k, m \in n-1$ , either

$$\lim_{\Omega' \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(m)) = 0 \quad \text{or} \quad \varepsilon' := \inf_{(i, j) \in [\Omega']^2} d_X(f_i(k), f_j(m)) > 0.$$

Applying the inductive assumption once more, find an infinite set  $\Omega \subseteq \Omega'$  such that

$$\lim_{\Omega \ni i, j \rightarrow \infty} d_X(f_i(n-1), f_j(n-1)) = 0 \quad \text{or} \quad \inf_{(i, j) \in [\Omega]^2} d_X(f_i(n-1), f_j(n-1)) \geq \varepsilon$$

for some  $\varepsilon \in (0, \varepsilon']$ . Let  $C$  be the set of all number  $k \in n$  such that the sequence  $(f_i(k))_{i \in \Omega}$  is Cauchy.

Two cases are possible.

1)  $n-1 \in C$ . This case has two subcases.

1a) For some infinite subset  $\Lambda \subseteq \Omega$  and some  $l \in n-1$  we have

$$\lim_{\Lambda \ni i \rightarrow \infty} d_X(f_i(n-1), f_i(l)) = 0.$$

Replacing  $\Lambda$  by a smaller infinite subset, we can additionally assume that

$$\sup_{j \in \Lambda} d_X(f_j(n-1), f_j(l)) < \frac{1}{2}\varepsilon.$$

We claim that for any  $k, m \in n$  with  $\lim_{\Lambda \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(m)) \neq 0$  we have  $\inf_{(i, j) \in [\Lambda]^2} d_X(f_i(k), f_j(m)) \geq \frac{1}{2}\varepsilon$ . If  $k, m \in n-1$ , then this follows from the definition of  $\varepsilon' \geq \varepsilon > \frac{1}{2}\varepsilon$ . Now assume that  $k$  or  $m$  does not belong to  $n-1$ . The case  $k = m = n-1$  is impossible as  $n-1 \in C$ . So,  $k \neq m$  and we lose no generality assuming that  $k < m = n-1$ . It follows from

$$\lim_{\Lambda \ni i \rightarrow \infty} d_X(f_i(k), f_i(n-1)) \neq 0 = \lim_{\Lambda \ni i \rightarrow \infty} d_X(f_i(n-1), f_i(l))$$

that  $\lim_{\Lambda \ni i \rightarrow \infty} d_X(f_i(k), f_j(l)) \neq 0$  and then

$$\inf_{(i, j) \in [\Lambda]^2} d_X(f_i(k), f_j(n-1)) \geq \inf_{(i, j) \in [\Lambda]^2} d_X(f_i(k), f_j(l)) - \sup_{j \in \Lambda} d_X(f_j(n-1), f_j(l)) \geq \varepsilon' - \frac{1}{2}\varepsilon \geq \frac{1}{2}\varepsilon.$$

1b) For every infinite set  $\Lambda \subseteq \Omega$  and every  $l \in n-1$  we have

$\lim_{\Lambda \ni i \rightarrow \infty} d_X(f_i(n-1), f_i(l)) \neq 0$ . In this case, we can find an infinite subset  $\Lambda' \subseteq \Omega$  and a number  $\delta \in (0, \frac{1}{4}\varepsilon]$  such that  $\inf_{k \in n-1} \inf_{i \in \Lambda'} d_X(f_i(k), f_i(n-1)) \geq 2\delta$ . Find an infinite subset  $\Lambda'' \subseteq \Lambda'$  such that  $\sup_{i, j \in \Lambda''} d_X(f_i(k), f_j(k)) < \delta$  for every  $k \in C$ . By our assumption,  $n-1 \in C$  and for every  $k \in n \setminus C$  the sequence  $(f_i(k))_{i \in \Omega'}$  is  $\varepsilon'$ -separated. Then for the smallest number  $\lambda$  of the set  $\Lambda''$ , we can find an infinite subset  $\Lambda \subseteq \Lambda''$  such that

$$\min_{k \in n \setminus C} \inf_{i \in \Lambda} d_X(f_i(k), f_\lambda(n-1)) \geq \frac{1}{2}\varepsilon'.$$

We claim that for any  $k, m \in n$ ,

$$\lim_{\Lambda \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(m)) \neq 0 \Rightarrow \inf_{(i, j) \in [\Lambda]^2} d_X(f_i(k), f_j(m)) \geq \delta.$$

If  $k, m \in n-1$ , then this follows from the definition of  $\varepsilon' \geq \varepsilon \geq 4\delta$ . Now assume that  $k$  or  $m$  does not belong to  $n-1$ . We lose no generality assuming that  $k \leq m = n-1$ . If  $k \in C$ , then

$\lim_{\Lambda \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(n-1)) \neq 0$  implies  $\lim_{\Lambda \ni i \rightarrow \infty} d_X(f_i(k), f_i(n-1)) \neq 0$  and hence

$$\inf_{(i,j) \in [\Lambda]^2} d_X(f_i(k), f_j(n-1)) \geq \inf_{i \in \Lambda} d_X(f_i(k), f_i(n-1)) - \sup_{j \in \Lambda} d_X(f_j(n-1), f_i(n-1)) \geq 2\delta - \delta.$$

If  $k \in n \setminus C$ , then

$$\inf_{(i,j) \in [\Lambda]^2} d_X(f_i(k), f_j(n-1)) \geq \inf_{i \in \Lambda} d_X(f_i(k), f_\lambda(n-1)) - \sup_{j \in \Lambda} d_X(f_\lambda(n-1), f_j(n-1)) \geq \frac{1}{2}\varepsilon' - \delta.$$

2) Now consider the second case:  $n-1 \notin C$ . In this case the sequence  $(f_i(n-1))_{i \in \Omega}$  is  $\varepsilon$ -separated by the choice of  $\varepsilon$ . Find an element  $\lambda_0 \in \Omega$  such that

$$\sup\{d_X(f_i(k), f_j(k)) : i, j \in \Omega \cap [\lambda_0, \infty)\} < \frac{1}{4}\varepsilon \quad \text{for every } k \in C.$$

Using the  $\varepsilon$ -separatedness of the sequences  $(f_i(l))_{i \in \Omega}$  for  $l \in n \setminus C$ , we can inductively construct an increasing sequence  $\{\lambda_k\}_{k \in \omega} \subseteq \Omega$  such that for every  $k \in \omega$  the following condition holds:

$$(*) \text{ for any } i \leq \lambda_k, j \in \Omega \cap [\lambda_{k+1}, \infty), m \in n \text{ and } l \in n \setminus C \text{ we have } d_X(f_j(l), f_i(m)) \geq \frac{1}{2}\varepsilon.$$

Let  $\Lambda = \{\lambda_i : i \in \mathbb{N}\}$ . We claim that for any  $k, m \in n$ ,

$$\lim_{\Lambda \ni i, j \rightarrow \infty} d_X(f_i(k), f_j(m)) \neq 0 \Rightarrow \inf_{(i,j) \in [\Lambda]^2} d_X(f_i(k), f_j(m)) \geq \frac{1}{4}\varepsilon.$$

If  $k, m \in n-1$ , then this follows from the definition of  $\varepsilon' \geq \frac{1}{4}\varepsilon$ . Now assume that  $k$  or  $m$  does not belong to  $n-1$ . We lose no generality assuming that  $k \leq m = n-1$ . Choose any distinct numbers  $\lambda_i, \lambda_j \in \Lambda$ . If  $k \in C$ , then

$$d_X(f_{\lambda_i}(k), f_{\lambda_j}(n-1)) \geq d_X(f_{\lambda_0}(k), f_{\lambda_j}(n-1)) - d_X(f_{\lambda_0}(k), f_{\lambda_i}(k)) \geq \frac{1}{2}\varepsilon - \frac{1}{4}\varepsilon = \frac{1}{4}\varepsilon,$$

by  $(*)$  and the choice of  $\lambda_0$ . If  $k \notin C$ , then  $d_X(f_{\lambda_i}(k), f_{\lambda_j}(n-1)) \geq \frac{1}{2}\varepsilon$  by  $(*)$ .  $\square$

*Proof of Theorem 20.1.* Given any Cauchy sequence  $\{b_i\}_{i \in \omega} \subseteq F^p X$ , we should prove that it converges to some element  $b_\infty \in FX$ . Let  $s$  be the smallest number for which there exists an infinite set  $\Omega' \subseteq \omega$  and a sequence  $\{b'_i\}_{i \in \omega} \subseteq FX$  such that

- (a)  $\lim_{\Omega' \ni i \rightarrow \infty} d_{FX}^p(b_i, b'_i) = 0$ ;
- (b)  $d_{FX}^p(b'_i, b'_j) < \infty$  for all  $i, j \in \Omega'$ ;
- (c)  $|\text{supp}(b'_i)| = s$  for all  $i \in \Omega'$ .

Since the functor  $F$  has finite degree, the number  $s$  is well-defined and  $s \leq \text{deg}(F)$ . Fix an infinite set  $\Omega' \subseteq \omega$  satisfying the conditions (a)–(c). The condition (a) implies that the sequence  $(b'_i)_{i \in \Omega'}$  is Cauchy. If  $s = 0$ , then Theorem 8.1 and the condition (b) imply  $b'_i = b'_j$  for all  $i, j \in \Omega'$ . In this case the sequence  $(b'_i)_{i \in \Omega'}$  is convergent and so is the sequence  $(b_i)_{i \in \omega}$  (to the same limit point).

So, we assume that  $s > 0$ . Using Proposition 6.1, for every  $i \in \Omega'$ , find a bijective map  $h_i : s \rightarrow \text{supp}(b'_i) \subseteq X$  and an element  $a_i \in Fs$  such that  $b'_i = Fh_i(a_i)$ . Since  $F$  is finitary, the set  $Fs$  is finite. Then by the Pigeonhole Principle, there exists an element  $a \in Fs$  such that the set  $\Omega'' = \{i \in \Omega' : a_i = a\}$  is infinite. By Lemma 20.3, there exists an infinite set  $\Omega \subseteq \Omega''$  and  $\varepsilon \in (0, \infty)$  such that for any  $k, m \in s$  either

$$\lim_{\Omega \ni i, j \rightarrow \infty} d_X(h_i(k), h_j(m)) = 0 \quad \text{or} \quad \inf_{(i,j) \in [\Omega]^2} d_X(h_i(k), h_j(m)) \geq \varepsilon.$$

Replacing  $\Omega$  by a smaller infinite subset of  $\Omega$ , we can additionally assume that for any  $k, m \in s$ ,

$$\lim_{\Omega \ni i, j \rightarrow \infty} d_X(h_i(k), h_j(m)) = 0 \Rightarrow \sup_{i, j \in \Omega} d_X(h_i(k), h_j(m)) < \frac{1}{4}\varepsilon.$$

Let  $C$  be the set of all numbers  $k \in s$  such that the sequence  $(h_i(k))_{i \in \Omega}$  is Cauchy and hence converges to some point  $h(k) \in X$  of the complete distance space  $(X, d_X)$ . We can assume that  $h(k) = h(m)$  for any  $k, m \in C$  with  $d_X(h(k), h(m)) = 0$ . The choice of  $\varepsilon$  guarantees that

$$d_X(h(k), h(m)) \geq \inf_{(i,j) \in [\Omega]^2} d_X(h_i(k), h_j(m)) \geq \varepsilon$$

for any  $k, m \in C$  with  $d_X(h(k), h(m)) > 0$ . Consequently, the set  $h[C] := \{h(k) : k \in C\}$  is  $\varepsilon$ -separated. We claim that  $C = s$ . To derive a contradiction, assume that  $C \neq s$ .

Consider the (unique) map  $r : X \rightarrow X$  such that

$$r(x) = \begin{cases} y & \text{if } d_X(x, y) < \frac{1}{4}\varepsilon \text{ for some } y \in h[C]; \\ x & \text{otherwise.} \end{cases}$$

Observe that for every  $k \in C$ ,  $m \in s \setminus C$ , and distinct numbers  $i, j \in \Lambda$ , we have

$$d_X(r \circ h_i(k), r \circ h_j(m)) = d_X(h(k), h_j(m)) \geq d_X(h_i(k), h_j(m)) - d_X(h_i(k), h(k)) \geq \varepsilon - \frac{1}{4}\varepsilon.$$

For every  $i \in \Omega$ , consider the element  $b_i'' = Fr \circ Fh_i(b)$  and observe that

$$d_{FX}^p(b_i'', b_i') \leq d_{X^s}^p(r \circ h_i, h_i) \leq \left( \sum_{k \in S'} d_X(h_i(k), h(k))^p \right)^{\frac{1}{p}}.$$

The latter inequality implies that

$$\lim_{\Omega_s \ni i \rightarrow \infty} d_{FX}^p(b_i, b_i'') \leq \lim_{\Omega_s \ni i \rightarrow \infty} d_{FX}^p(b_i, b_i') + \lim_{\Omega_s \ni i \rightarrow \infty} d_{FX}^p(b_i', b_i'') = 0 + 0 = 0$$

and hence the sequence  $(b_i'')_{i \in \Omega_s}$  is Cauchy. Observe that for every  $i \in \Omega_s$  we have  $\text{supp}(b_i'') \subseteq r \circ h_i[s]$  and hence  $|\text{supp}(b_i'')| \leq |s| = s$ . Now the choice of  $s$  ensures that the set  $\{i \in \Omega : |\text{supp}(b_i'')| < s\}$  is finite. Since  $(b_i'')_{i \in \Omega}$  is Cauchy, we can find an infinite set  $\Lambda \subseteq \Omega$  such that

- (d)  $\text{supp}(b_i'') = r \circ h_i[s]$  for all  $i \in \Lambda$ ;
- (e)  $d_{FX}^1(b_i'', b_j'') \leq n^{1-\frac{1}{p}} d_{FX}^p(b_i'', b_j'') < \frac{1}{2}\varepsilon$  for all  $i, j \in \Lambda$ .

The first inequality in (e) follows Theorem 11.2.

Fix any distinct numbers  $i, j \in \Lambda$ . If  $C = \emptyset$ , then

$$\begin{aligned} \underline{d}_X(\text{supp}(b_i'') \cup \text{supp}(b_j'')) &= \underline{d}_X(r \circ h_i[s] \cup r \circ h_j[s]) = \underline{d}_X(h_i[s], h_j[s]) = \\ &= \min_{k, m \in s} d_X(h_i(k), h_j(m)) \geq \varepsilon. \end{aligned}$$

If  $h[C]$  is a singleton, then

$$\begin{aligned} \underline{d}_X(\text{supp}(b_i'') \cup \text{supp}(b_j'')) &= \underline{d}_X(r \circ h_i[s] \cup r \circ h_j[s]) = \\ &= \min\{\underline{d}_X(h_i[s \setminus C], h_j[s \setminus C]), \underline{d}_X(h[C], h_j[s \setminus C]), \underline{d}_X(h_i[s \setminus C], h[C])\} \geq \\ &= \min\{\varepsilon, \frac{3}{4}\varepsilon, \frac{3}{4}\varepsilon\} = \frac{3}{4}\varepsilon. \end{aligned}$$

If  $|h[C]| > 1$ , then

$$\begin{aligned} \underline{d}_X(\text{supp}(b_i'') \cup \text{supp}(b_j'')) &= \underline{d}_X(r \circ h_i[s] \cup r \circ h_j[s]) = \\ &= \min\{\underline{d}_X(h[C]), \underline{d}_X(h_i[s \setminus C], h_j[s \setminus C]), \underline{d}_X(h[C], h_j[s \setminus C]), \underline{d}_X(h_i[s \setminus C], h[C])\} \geq \\ &= \min\{\varepsilon, \varepsilon, \frac{3}{4}\varepsilon, \frac{3}{4}\varepsilon\} = \frac{3}{4}\varepsilon. \end{aligned}$$

In any case we obtain  $\underline{d}_X(\text{supp}(b'_i) \cup \text{supp}(b''_i)) \geq \frac{3}{4}\varepsilon > \frac{1}{2}\varepsilon$ . The last inequality, condition (e) and Theorem 16.4 imply that  $b''_i = b'_i$ . Then  $\text{supp}(b''_i) = \text{supp}(b'_i)$  and the symmetric difference  $\text{supp}(b'_i)\Delta\text{supp}(b''_i)$  is empty. On the other hand,

$$\text{supp}(b''_i)\Delta\text{supp}(b'_i) = (r \circ h_i[s])\Delta(r \circ h_j[s]) = h_i[s \setminus C] \cup h_j[s \setminus C] \neq \emptyset.$$

This contradiction shows that  $C = s$ . Then then for the element  $Fh(b) \in FX$  we have

$$\lim_{\Omega \ni i \rightarrow \infty} d_{FX}^p(b'_i, Fh(b)) = \lim_{\Omega \ni i \rightarrow \infty} d_{FX}^p(Fh_i(b), Fh(b)) \leq \lim_{\Omega \ni i \rightarrow \infty} d_{X^s}^p(h_i, h) = 0,$$

which means that the sequence  $(b'_i)_{i \in \Omega}$  converges to  $Fh(b)$  and so does the Cauchy sequence  $(b_i)_{i \in \omega}$  (as  $\lim_{\Omega \ni i \rightarrow \infty} d_{FX}^p(b_i, b'_i) = 0$ ).  $\square$

By a *completion* of a distance space  $X$  we understand a pair  $(Y, i)$  consisting of a complete distance space  $(Y, d_Y)$  and an injective isometry  $i : X \rightarrow Y$  such that the set  $i(X)$  is dense in  $Y$  and  $d_Y(y, y') > 0$  for every distinct points  $y, y' \in Y \setminus i(X)$ .

It can be shown that each distance space  $X$  has a completion and this completion is unique in the sense that for any completions  $(Y_1, i_1)$  and  $(Y_2, i_2)$  of  $X$  there exists a unique bijective isometry  $j : Y_1 \rightarrow Y_2$  such that  $j \circ i_1 = i_2$ .

**Problem 20.4.** *Given a distance space  $X$ , a finitely supported functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  and number  $p \in [1, \infty]$ , describe the completion of the distance space  $F^p X$ .*

**Remark 20.5.** Let  $H : \mathbf{Set} \rightarrow \mathbf{Set}$  be the functor assigning to each set  $X$  the set  $HX$  of nonempty finite subsets of  $X$  and to each function  $f : X \rightarrow Y$  between sets the function  $Hf : HX \rightarrow HY$ ,  $Hf : A \mapsto f[A]$ . For any complete metric space  $X$  the completion of the metric space  $H^\infty X$  can be identified with the space of all nonempty compact subsets of  $X$ , endowed with the Hausdorff metric. On the other hand, the completion of the metric space  $H^1 X$  can be identified with the space of all nonempty compact subsets of zero length in  $X$ , see [5].

## 21. THE FUNCTOR $F^p$ AND THE METRIC ENTROPY OF DISTANCE SPACES

For a distance space  $(X, d_X)$  and an  $\varepsilon \in (0, \infty)$ , let  $\mathbb{E}_\varepsilon(X, d_X)$  be the smallest cardinality of a cover of  $X$  by subsets of diameter  $\leq \varepsilon$ . If the distance  $d_X$  is clear from the context, then we write  $\mathbb{E}_\varepsilon(X)$  instead of  $\mathbb{E}_\varepsilon(X, d_X)$ .

For two numbers  $\varepsilon, \delta \in (0, \infty)$  let

$$\mathbb{E}_{\varepsilon, \delta}(X) = \sup\{\mathbb{E}_\varepsilon(A) : A \subseteq X \wedge \text{diam}(A) < \delta\}.$$

The functions  $\mathbb{E}_\varepsilon(X)$  and  $\mathbb{E}_{\varepsilon, \delta}(X)$  (of the variables  $\varepsilon, \delta$ ) are called *the metric entropy function* of the distance space  $X$ .

**Theorem 21.1.** *If the functor  $F$  has finite degree, then for every  $\varepsilon \in [0, \infty)$ ,*

$$\mathbb{E}_\varepsilon(F^p X) \leq |Fn| \cdot \mathbb{E}_\varepsilon(X^n, d_{X^n}^p) \leq |Fn| \cdot (\mathbb{E}_{\varepsilon/\sqrt[p]{n}}(X))^n,$$

where  $n = \max\{1, \deg(F)\}$ .

*Proof.* By Theorem 10.2, for every  $a \in Fn$  the map  $\xi^a : (X^n, d_{X^n}^p) \rightarrow (FX, d_{FX}^p)$  is non-expanding. By the definition of the cardinal  $\mathbb{E}_\varepsilon(X^n, d_{X^n}^p)$  there exists a cover of  $\mathcal{V}$  of  $X^n$  such that  $|\mathcal{V}| = \mathbb{E}_\varepsilon(X^n)$  and  $\text{mesh}(\mathcal{V}) \leq \varepsilon$ . Taking into account that  $FX = \bigcup_{a \in Fn} \xi^a[X^n]$ , we conclude that  $\mathcal{W} = \{\xi^a[V] : a \in Fn, V \in \mathcal{V}\}$  is a cover of  $F^p X$  of cardinality  $|\mathcal{W}| \leq |Fn| \cdot |\mathcal{V}| =$



$|Fn| \cdot \mathbb{E}_{X^n}(\varepsilon)$ . The non-expanding property of the maps  $\xi^a : X^n \rightarrow F^p X$  ensures that  $\text{mesh}(\mathcal{W}) \leq \text{mesh}(\mathcal{V}) \leq \varepsilon$ , witnessing that  $\mathbb{E}_\varepsilon(F^p X) \leq |\mathcal{W}| \leq |Fn| \cdot |\mathcal{V}| = |Fn| \cdot \mathbb{E}_\varepsilon(X^n, d_{X^n}^p)$ .

It remains to prove the inequality  $\mathbb{E}_\varepsilon(X^n, d_{X^n}^p) \leq (\mathbb{E}_{\varepsilon/\sqrt[p]{n}}(X))^n$ . By the definition of the cardinal  $\mathbb{E}_{\varepsilon/\sqrt[p]{n}}(X)$ , there exists a cover  $\mathcal{U}$  of  $X$  such that  $|\mathcal{U}| = \mathbb{E}_{\varepsilon/\sqrt[p]{n}}(X)$  and  $\text{mesh}(\mathcal{U}) \leq \varepsilon/\sqrt[p]{n}$ . For any sets  $U_0, \dots, U_{n-1} \in \mathcal{U}$  consider their product  $\prod_{i \in n} U_i \subseteq X^n$  and observe that

$$\text{diam}\left(\prod_{i \in n} U_i\right) \leq \left(\sum_{i \in n} \text{diam}(U_i)^p\right)^{\frac{1}{p}} \leq n^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}) \leq \varepsilon.$$

Then  $\mathcal{U}' = \{\prod_{i \in n} U_i : U_0, \dots, U_{n-1} \in \mathcal{U}\}$  is a cover of  $X^n$  witnessing that

$$\mathbb{E}_\varepsilon(X^n, d_{X^n}^p) \leq |\mathcal{U}'| \leq |\mathcal{U}|^n = (\mathbb{E}_{\varepsilon/\sqrt[p]{n}}(X))^n.$$

□

Now we deduce an upper bound for the metric entropy function  $\mathbb{E}_{\varepsilon, \delta}(F^p X)$  of the distance space  $F^p X$ .

**Theorem 21.2.** *Assume that the functor  $F$  has finite degree and preserves supports. Then for any  $\varepsilon, \delta \in [0, \infty)$  we have*

$$\mathbb{E}_{\varepsilon, \delta}(F^p X) \leq n^n \cdot |Fn| \cdot \mathbb{E}_{\varepsilon, \delta'}(X^n, d_{X^n}^p) \leq n^n \cdot |Fn| \cdot (\mathbb{E}_{\varepsilon', \delta'}(X))^n,$$

where  $n = \max\{1, \deg(F)\}$ ,  $\varepsilon' = \varepsilon/\sqrt[p]{n}$  and  $\delta' = 2\delta\sqrt[p]{n}$ .

*Proof.* Given any nonempty subset  $A \subseteq F^p X$  of  $\text{diam}(A) < \delta < \infty$ , we will prove that

$$\mathbb{E}_\varepsilon(A) \leq n^n \cdot |Fn| \cdot \mathbb{E}_{\varepsilon, \delta'}(X^n, d_{X^n}^p).$$

If  $\bigcup_{a \in A} \text{supp}(a) = \emptyset$ , then Theorem 8.1 implies that  $A$  is a singleton and hence  $\mathbb{E}_\varepsilon(A) = |A| = 1$ . So, suppose that  $\bigcup_{a \in A} \text{supp}(a) \neq \emptyset$  and choose an element  $a \in A$  with nonempty support  $S = \text{supp}(a)$ . Since the functor  $F$  has finite degree,  $|S| \leq n$ .

For every function  $g \in S^n$ , consider the ball

$$O_\infty(g; \delta) := \{f \in X^n : d_{X^n}^\infty(f, g) < \delta\}$$

in the distance space  $(X^n, d_{X^n}^\infty)$ .

**Claim 21.3.**  $A \subseteq \bigcup_{c \in Fn} \bigcup_{g \in S^n} \xi^c[O_\infty(g; \delta)]$ .

*Proof.* Given any element  $b \in A$ , apply Theorem 16.6 to conclude that

$$d_{HX}(\text{supp}(a), \text{supp}(b)) \leq d_{FX}^p(a, b) < \delta < \infty.$$

Then  $\text{supp}(b) \neq \emptyset$  and we can apply Proposition 6.1 to find a function  $f : n \rightarrow \text{supp}(b)$  and an element  $c \in Fn$  such that  $b = Ff(c)$ . Since  $d_{HX}(\text{supp}(b), S) < \delta$ , there exists a function  $r : \text{supp}(b) \rightarrow S$  such that  $d_X(r(x), x) < \delta$  for every  $x \in \text{supp}(b)$ . Consider the map  $g = r \circ f : n \rightarrow S$  and observe that  $d_{X^n}^\infty(f, g) < \delta$  and hence  $f \in O_\infty(g; \delta)$ . Then  $b = Ff(c) = \xi^c(f) \in \xi^c[O_\infty(g; \delta)]$ . □

By Theorem 11.2,  $O_\infty(g; \delta) \subseteq O_p(g; n^{\frac{1}{p}}\delta) := \{f \in X^n : d_{X^n}^p(f, g) < \delta\}$ . Taking into account that for every  $c \in Fn$ , the map  $\xi^c : X^n \rightarrow F^p X$  is non-expanding (for the distance

$d_{X^n}^p$  on  $X$ ), we conclude that

$$\begin{aligned} \mathbb{E}_\varepsilon(A) &\leq \sum_{c \in Fn} \sum_{g \in S^n} \mathbb{E}_\varepsilon(\xi^c[O_\infty(g; \delta)]) \leq \sum_{c \in Fn} \sum_{g \in S^n} \mathbb{E}_\varepsilon(\xi^c[O_p(g; \delta \sqrt[p]{n})]) \leq \\ &\sum_{c \in Fn} \sum_{g \in S^n} \mathbb{E}_\varepsilon(O_p(g; \delta \sqrt[p]{n})) \leq \sum_{c \in Fn} \sum_{g \in S^n} \mathbb{E}_{\varepsilon, 2\delta \sqrt[p]{n}}(X^n, d_{X^n}^p) \leq |Fn| \cdot n^n \cdot \mathbb{E}_{\varepsilon, \delta'}(X^n, d_{X^n}^p). \end{aligned}$$

The inequality  $\mathbb{E}_{\varepsilon, \delta'}(X^n, d_{X^n}^p) \leq (\mathbb{E}_{\varepsilon', \delta'}(X))^n$  can be verified by analogy with the inequality  $\mathbb{E}_\varepsilon(X^n, d_{X^n}^p) \leq (\mathbb{E}_\varepsilon(X))^n$ , proved in Theorem 21.1.  $\square$

The metric entropy functions allow us to define five *cardinal entropy characteristics* of a distance space  $X$ :

$$\begin{aligned} \mathbb{E}_0(X) &= \sup_{\varepsilon \in (0, \infty)} \mathbb{E}_\varepsilon(X)^+, & \mathbb{E}_{0, \infty}(X) &= \sup_{\varepsilon, \delta \in (0, \infty)} \mathbb{E}_{\varepsilon, \delta}(X)^+, & \mathbb{E}_\infty(X) &= \min_{\varepsilon \in (0, \infty)} \mathbb{E}_\varepsilon(X)^+, \\ \mathbb{E}_{0, 0}(X) &= \min_{\delta \in (0, \infty)} \sup_{\varepsilon \in (0, \infty)} \mathbb{E}_{\varepsilon, \delta}(X)^+, & \mathbb{E}_{\infty, \infty}(X) &= \min_{\varepsilon \in (0, \infty)} \sup_{\delta \in (0, \infty)} \mathbb{E}_{\varepsilon, \delta}(X)^+. \end{aligned}$$

In the above definitions  $\kappa^+$  stands for the successor of a cardinal  $\kappa$ . For example,  $\omega_1 = \omega^+$ .

It is easy to see that for any distance space  $X$  we have the inequalities:

$$\mathbb{E}_{0, 0}(X) \leq \mathbb{E}_{0, \infty}(X) \leq \mathbb{E}_0(X) \geq \mathbb{E}_\infty(X) \geq \mathbb{E}_{\infty, \infty}(X).$$

Many (known) properties of a distance space are equivalent to the countability of suitable cardinal entropy characteristics.

**Definition 21.4.** A distance space  $(X, d_X)$  is defined to be

- (0)-*finitary* (or else *totally bounded*) if  $\mathbb{E}_0(X) \leq \omega$  (i.e., for any  $\varepsilon \in (0, \infty]$  the entropy  $\mathbb{E}_\varepsilon(X)$  is finite);
- ( $\infty$ )-*finitary* if  $\mathbb{E}_\infty(X) \leq \omega$  (i.e.,  $X$  has finitely many pseudometric components);
- (0, 0)-*finitary* (or else *Assouad-bounded*) if  $\mathbb{E}_{0, 0}(X) \leq \omega$  (i.e., there exists  $\delta \in (0, \infty)$  such that for every  $\varepsilon \in (0, \delta]$  the entropy  $\mathbb{E}_{\varepsilon, \delta}(X)$  is finite);
- (0,  $\infty$ )-*finitary* if  $\mathbb{E}_{0, \infty}(X) \leq \omega$  (i.e., for any  $\varepsilon, \delta \in (0, \infty)$  the entropy  $\mathbb{E}_{\varepsilon, \delta}(X)$  is finite);
- ( $\infty, \infty$ )-*finitary* (or else *of bounded geometry*) if  $\mathbb{E}_{\infty, \infty}(X) \leq \omega$  (i.e., there exists  $\varepsilon \in (0, \infty)$  such that for every  $\delta \in (0, \infty)$  the cardinal  $\mathbb{E}_{\varepsilon, \delta}(X)$  is finite).

For any distance space these finitariness properties relate as follows:

$$\begin{array}{ccccc} (0, 0)\text{-finitary} & \iff & (0, \infty)\text{-finitary} & \iff & (0)\text{-finitary} & \implies & (\infty)\text{-finitary} & \implies & (\infty, \infty)\text{-finitary} \\ \parallel & & \parallel & & \parallel & & \parallel & & \parallel \\ \text{Assouad} & & \text{totally} & & & & \text{of bounded} & & \\ \text{bounded} & & \text{bounded} & & & & \text{geometry} & & \end{array}$$

Assouad bounded metric spaces were studied by Assouad in [2].

Theorems 21.1, 21.2 imply the following upper bounds for the cardinal entropy characteristics of the distance space  $F^p X$ .

**Corollary 21.5.** *If the functor  $F$  has finite degree, then for  $n := \max\{1, \deg(F)\}$ , the distance space  $F^p X$  has the following cardinal entropy characteristics.*

- 1)  $\mathbb{E}_0(F^p X) \leq |Fn| \cdot (\mathbb{E}_0(X))^n$ ;
- 2)  $\mathbb{E}_\infty(F^p X) \leq |Fn| \cdot (\mathbb{E}_\infty(X))^n$ .

If, in addition, the functor  $F$  preserves supports, then

- 3)  $\mathbb{E}_{0,0}(F^p X) \leq n^n \cdot |Fn| \cdot (\mathbb{E}_{0,0}(X))^n$ ;
- 4)  $\mathbb{E}_{0,\infty}(F^p X) \leq n^n \cdot |Fn| \cdot (\mathbb{E}_{0,\infty}(X))^n$ ;
- 5)  $\mathbb{E}_{\infty,\infty}(F^p X) \leq n^n \cdot |Fn| \cdot (\mathbb{E}_{\infty,\infty}(X))^n$ .

Corollary 21.5(1,5) has two important special cases:

**Corollary 21.6.** *Assume that the functor  $F$  is finitary and has finite degree. If a distance space  $(X, d_X)$  is totally bounded, then so is the distance space  $F^p X$ .*

**Corollary 21.7.** *Assume that the functor  $F$  is finitary, has finite degree and preserves supports. If a distance space  $X$  has bounded geometry (or is Assouad-bounded), then so is the distance space  $F^p X$ .*

**Example 21.8.** Consider the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  assigning to each set  $X$  the set  $FX = \{\emptyset\} \cup \{\{x, y\} : x, y \in X, x \neq y\}$  and to each function  $f : X \rightarrow Y$  the function  $Ff : FX \rightarrow FY$  such that  $Ff(\emptyset) = \emptyset$  and  $Ff(\{x, y\}) = \{f(x)\} \triangle \{f(y)\}$  for any  $\{x, y\} \in FX \setminus \{\emptyset\}$ . It is clear that  $F$  is finitary and has finite degree  $\deg(F) = 2$ , but  $F$  does not preserve supports. For the space  $X = \mathbb{R}$  we have  $\mathbb{E}_{0,0}(X) = \mathbb{E}_{0,\infty}(X) = \mathcal{E}_{\infty,\infty}(X) = \omega$  and  $\mathbb{E}_{0,0}(F^p X) = \mathbb{E}_{0,\infty}(F^p X) = \mathcal{E}_{\infty,\infty}(F^p X) = \omega_1$ . This example shows that the preservation of supports in Corollary 21.7 (and also in the statements (3)–(5) of Corollary 21.5) is essential.

Observe that a distance space  $X$  is *separable* (= has a countable dense subset) if and only if  $\mathbb{E}_0(X) \leq \omega_1$ . By Corollary 21.5(1), the distance space  $F^p$  is separable if  $X$  is separable and the functor  $F$  is finitary and has finite degree. In fact, this implication holds in a more general situation.

**Proposition 21.9.** *If for every  $n \in \mathbb{N}$  the set  $Fn$  is at most countable, then for any separable distance space  $X$ , the distance space  $F^p X$  is separable.*

*Proof.* By Proposition 6.1,  $FX = \bigcup_{n \in \mathbb{N}} \bigcup_{a \in Fn} \xi^a(X^n)$ . For every  $n \in \mathbb{N}$  the separability of the distance space  $X$  implies the separability of the power  $X^n$  endowed with the distance  $d_{X^n}^p$ . By Theorem 10.1, for every  $a \in Fn$ , the map  $\xi^a : X^n \rightarrow FX$  is non-expanding, which implies that the space  $\xi^a[X^n]$  is separable and then  $FX$  is separable, being a countable union of separable spaces.  $\square$

## 22. THE FUNCTOR $F^p$ AND THE ENTROPY DIMENSIONS OF DISTANCE SPACES

In this section we study the dependence between various entropy dimensions of distance spaces  $X$  and  $F^p X$ .

We will consider eight entropy dimensions divided into four pairs. The first pair form two well-known box-counting dimensions, defined for any totally bounded distance space. In our terminology totally bounded spaces are called (0)-finitary.

For a totally bounded distance space  $X$ , its upper and lower box-counting dimensions are defined by the formulas

$$\overline{\dim}_B X = \limsup_{\varepsilon \rightarrow +0} \frac{\ln(\mathbb{E}_\varepsilon(X))}{\ln(1/\varepsilon)} \quad \text{and} \quad \underline{\dim}_B X = \liminf_{\varepsilon \rightarrow +0} \frac{\ln(\mathbb{E}_\varepsilon(X))}{\ln(1/\varepsilon)}.$$

The box-counting dimensions play an important role in the Theory of Fractals, see [18, Ch.3].

**Example 22.1.** The Cantor ternary set

$$C = \left\{ \sum_{n=0}^{\infty} \frac{x_n}{3^n} : (x_n)_{n \in \omega} \in \{0, 1\}^{\omega} \right\} \subset \mathbb{R}$$

is totally bounded and has the box-counting dimensions

$$\overline{\dim}_B C = \underline{\dim}_B C = \frac{\ln 2}{\ln 3}.$$

The definition of the (0)-entropy dimensions and Theorem 21.1, imply the following corollary.

**Theorem 22.2.** *Assume that the functor  $F$  is finitary and has finite degree  $n = \deg(F)$ . If a distance space  $X$  is totally bounded, then the distance space  $F^p X$  is totally bounded and has upper and lower box-counting dimensions*

$$\overline{\dim}_B F^p X \leq n \cdot \overline{\dim}_B X \quad \text{and} \quad \underline{\dim}_B F^p X \leq n \cdot \underline{\dim}_B X.$$

A micro-version of the (upper and lower) box-counting dimensions are the (upper and lower) (0, 0)-entropy dimensions defined for any (0, 0)-finitary distance space  $X$  by the formulas

$$\overline{\dim}_{(0,0)} X = \lim_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \frac{\ln(\mathbb{E}_{\varepsilon, \delta}(X))}{\ln(\delta/\varepsilon)} \quad \text{and} \quad \underline{\dim}_{(0,0)} X = \lim_{\delta \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} \frac{\ln(\mathbb{E}_{\varepsilon, \delta}(X))}{\ln(\delta/\varepsilon)}.$$

Large-scale counterparts of the (upper and lower) (0, 0)-entropy dimensions are the (upper and lower)  $(\infty, \infty)$ -entropy dimensions, defined for any  $(\infty, \infty)$ -finitary distance space  $X$  by the formulas

$$\overline{\dim}_{(\infty, \infty)} X = \lim_{\varepsilon \rightarrow \infty} \limsup_{\delta \rightarrow \infty} \frac{\ln(\mathbb{E}_{\varepsilon, \delta}(X))}{\ln(\delta/\varepsilon)} \quad \text{and} \quad \underline{\dim}_{(\infty, \infty)} X = \lim_{\varepsilon \rightarrow \infty} \liminf_{\delta \rightarrow \infty} \frac{\ln(\mathbb{E}_{\varepsilon, \delta}(X))}{\ln(\delta/\varepsilon)}.$$

Finally, for any (0,  $\infty$ )-finitary distance space  $X$ , let us consider the (upper and lower) (0,  $\infty$ )-entropy dimensions:

$$\overline{\dim}_{(0, \infty)} X = \limsup_{(\varepsilon, \delta) \rightarrow (0, \infty)} \frac{\ln(\mathbb{E}_{\varepsilon, \delta}(X))}{\ln(\delta/\varepsilon)} \quad \text{and} \quad \underline{\dim}_{(0, \infty)} X = \liminf_{(\varepsilon, \delta) \rightarrow (0, \infty)} \frac{\ln(\mathbb{E}_{\varepsilon, \delta}(X))}{\ln(\delta/\varepsilon)}.$$

The upper (0,  $\infty$ )-entropy dimension  $\overline{\dim}_{(0, \infty)}$  was introduced and studied by Assouad in [2].

The relations between the six entropy dimensions are described by the following diagram in which an arrow  $a \rightarrow b$  indicates that  $a \leq b$ .

$$\begin{array}{ccccc} \overline{\dim}_{(0,0)}(X) & \longrightarrow & \overline{\lim}_{(0,\infty)}(X) & \longleftarrow & \overline{\dim}_{(\infty,\infty)}(X) \\ \uparrow & & \uparrow & & \uparrow \\ \underline{\dim}_{(0,0)}(X) & \longrightarrow & \underline{\dim}_{(0,\infty)}(X) & \longleftarrow & \underline{\dim}_{(\infty,\infty)}(X) \end{array}$$

**Example 22.3.** The extended Cantor set

$$C = \left\{ \sum_{n=m}^{\infty} \frac{x_n}{3^n} : m \in \mathbb{Z} \wedge (x_n)_{n \in \mathbb{Z}} \in \{0, 1\}^{\mathbb{Z}} \right\} \subset \mathbb{R}$$

is (0,  $\infty$ )-finitary and has the entropy dimensions

$$\underline{\dim}_{(0,0)} C = \overline{\dim}_{(0,0)} C = \underline{\dim}_{(0,\infty)} C = \overline{\dim}_{(0,\infty)} C = \underline{\dim}_{(\infty,\infty)} C = \overline{\dim}_{(\infty,\infty)} C = \frac{\ln 2}{\ln 3}.$$

For characterizations of the extended Cantor set in various categories, see [7].

The definition of the entropy dimensions and Theorem 21.1, imply the following corollary.

**Theorem 22.4.** *Assume that the functor  $F$  is finitary, preserves supports and has finite degree  $n = \deg(F)$ . Let  $(\varepsilon, \delta) \in \{(0, 0), (0, \infty), (\infty, \infty)\}$ . If a distance space  $X$  is  $(\varepsilon, \delta)$ -finitary, then the distance space  $F^p X$  is  $(\varepsilon, \delta)$ -finitary and has upper and lower  $(\varepsilon, \delta)$ -entropy dimensions:*

$$\overline{\dim}_{(\varepsilon, \delta)} F^p X \leq n \cdot \overline{\dim}_{(\varepsilon, \delta)} X, \quad \underline{\dim}_{(\varepsilon, \delta)} F^p X \leq n \cdot \underline{\dim}_{(\varepsilon, \delta)} X.$$

### 23. THE FUNCTOR $F^p$ AND THE HAUSDORFF DIMENSION OF DISTANCE SPACES

In this section we study the dependence between the Hausdorff dimensions of the distance spaces  $X$  and  $F^p X$ .

The Hausdorff dimension of distance spaces is defined by analogy with the Hausdorff dimension of metric spaces, see [18, Ch.2].

Let  $X$  be a nonempty distance space. For any  $\varepsilon \in (0, \infty]$ , let  $\text{cov}_\varepsilon(X)$  be the family of all countable covers of  $X$  by sets of diameter  $\leq \varepsilon$ . For a cover  $\mathcal{U} \in \text{cov}_\varepsilon(X)$  and a positive real number  $s$ , let

$$\mathcal{H}^d(\mathcal{U}) = \sum_{U \in \mathcal{U}} \text{diam}(U)^s := \sup_{\mathcal{F} \in [\mathcal{U}]^{<\omega}} \sum_{F \in \mathcal{F}} \text{diam}(F)^s \in [0, \infty],$$

and

$$\mathcal{H}_\varepsilon^s(X) = \inf(\{\infty\} \cup \{\mathcal{H}^d(\mathcal{U}) : \mathcal{U} \in \text{cov}_\varepsilon(X)\}).$$

It is clear that  $\mathcal{H}_{\varepsilon'}^s(X) \leq \mathcal{H}_\varepsilon^s(X)$  for any  $\varepsilon \leq \varepsilon'$ . Consequently, there exists the (finite or infinite) limit

$$\mathcal{H}^s(X) = \lim_{\varepsilon \rightarrow +0} \mathcal{H}_\varepsilon^s(X) = \sup_{\varepsilon \in (0, \infty)} \mathcal{H}_\varepsilon^s(X),$$

called the *s-dimensional Hausdorff measure* of the distance space  $X$ . It can be shown that if  $\mathcal{H}^s(X) < \infty$  for some  $s \in (0, \infty)$ , then  $\mathcal{H}^t(X) = 0$  for every  $t > s$ .

The (finite or infinite) number

$$\dim_H X = \inf(\{\infty\} \cup \{s \in (0, \infty) : \mathcal{H}^s(X) = 0\})$$

is called the *Hausdorff dimension* of the distance space  $X$ .

The following properties of the Hausdorff dimension of distance spaces can be proved by analogy with the corresponding properties of the Hausdorff dimension for metric spaces, see [18, §2.2].

**Proposition 23.1.** *Let  $X$  be a distance space.*

- 1) *Each subspace  $A \subseteq X$  has  $\dim_H A \leq \dim_H X$ .*
- 2) *For any countable cover  $\{X_i\}_{i \in \omega}$  of  $X$  we have  $\dim_H X = \sup_{i \in \omega} \dim_H X_i$ .*
- 3) *For any surjective Lipschitz map  $f : X \rightarrow Y$  to a distance space  $Y$  we have  $\dim_H Y \leq \dim_H X$ .*

**Theorem 23.2.** *Assume that the functor  $F$  has finite degree  $n = \deg(X)$  and the set  $F^n$  is at most countable. Then*

$$\dim_H F^p X \leq \dim_H X^n,$$

where  $X^n$  is endowed with the distance  $d_{X^n}^p$ .

*Proof.* The inequality is trivial if  $\dim_H X^n = \infty$  or  $\dim_H F^p X = 0$ . So, we assume that  $\dim_H F^p X > 0$  and  $\dim_H(X^n) < \infty$ . If  $n = \deg X = 0$ , then we can take any map  $f : 1 \rightarrow X$  and applying Theorem 8.1, conclude that  $FX = Ff[F1]$  and hence  $|F^p X| \leq |F1| \leq |Fn| \leq \omega$  and  $\dim_H F^p X = 0$  by Proposition 23.1(1). But this contradicts our assumption. Therefore,  $n = \deg(F) > 0$ . In this case, by Proposition 6.1,  $F^p X = \bigcup_{a \in F^n} \xi^a[X^n]$ . By Theorem 10.1, the function  $\xi^a : X^n \rightarrow F^p X$  is non-expanding and hence

$$\dim_H F^p X = \sup_{a \in F^n} \dim_H \xi^a[X^n] \leq \sup_{a \in F^n} \dim_H X^n = \dim_H X^n.$$

□

**Remark 23.3.** The Hausdorff dimension does not obey the logarithmic law. By Example 7.8 in [18], the real line contains two Borel subsets  $Y, Z$  such that  $\dim_H Y = \dim_H Z = 0$  but  $\dim_H(Y \times Z) \geq 1$ . Moreover, by Product Formulas 7.2 and 7.3 in [18], for any Borel subsets  $Y, Z$  in a Euclidean space  $\mathbb{R}^n$  we have

$$\dim_H(Y \times Z) \geq \dim_H Y + \dim_H Z \quad \text{and} \quad \dim_H(Y \times Z) \leq \dim_H Y + \overline{\dim}_B Z.$$

#### 24. THE FUNCTOR $F^p$ AND THE CONTROLLED DIMENSIONS OF DISTANCE SPACES

In this section we study the interplay between the controlled dimensions of the distance spaces  $X$  and  $F^p X$ . The controlled dimensions of distance spaces are generalizations of the covering dimension of Lebesgue and asymptotic dimension of Gromov. They are introduced as follows.

Let  $\mathcal{D}$  be a nonempty family of non-decreasing functions from  $(0, \infty)$  to  $(0, \infty]$ . The family  $\mathcal{D}$  will be called a *dimension controlling class* and its elements *dimension controlling functions*.

Let  $D : (0, \infty) \rightarrow (0, \infty]$  be a dimension controlling function and  $n$  be a cardinal. We say that a distance space  $(X, d_X)$  has  *$D$ -controlled dimension*  $\dim_D X \leq n$  if for every  $\varepsilon \in (0, +\infty)$  there exists a cover  $\mathcal{U} = \bigcup_{i \in 1+n} \mathcal{U}_i$  of  $X$  such that

$$\text{mesh}(\mathcal{U}) \leq D(\varepsilon) \quad \text{and} \quad \inf_{i \in 1+n} \underline{d}_X(\mathcal{U}_i) \geq \varepsilon,$$

where

$$\underline{d}_X(\mathcal{U}_i) = \inf(\{\infty\} \cup \{\underline{d}_X(U, V) : U, V \in \mathcal{U}_i, U \neq V\}).$$

The smallest cardinal  $n$  such that  $\dim_D X \leq n$  is called the  *$D$ -dimension* of the distance space  $X$  and is denoted by  $\dim_D X$ .

Given a dimension controlling class  $\mathcal{D}$ , the cardinal

$$\dim_{\mathcal{D}} X := \min_{D \in \mathcal{D}} \dim_D X$$

is called the  *$\mathcal{D}$ -controlled dimension* of the distance space  $X$ .

**Definition 24.1.** A dimension controlling class  $\mathcal{D}$  is called a *dimension controlling scale* if it satisfies the following conditions:

- the identity function  $(0, \infty) \rightarrow (0, \infty)$  belongs to  $\mathcal{D}$ ;
- for any functions  $f, g \in \mathcal{D}$  the functions  $f + g$  and  $f \circ g$  belong to  $\mathcal{D}$ ;
- for any function  $f \in \mathcal{D}$ , any non-decreasing function  $g \leq f$  belongs to  $\mathcal{D}$ .

Each dimension controlling class generates a unique dimension controlling scale (equal to the intersection of all dimension controlling scales that contain this class).

The main result of this section is the following theorem generalizing topological results of Basmanov [8, 9, 10] and “coarse” results of Radul, Shukel, Zarichnyi [27, 29, 30].

**Theorem 24.2.** *If the functor  $F$  is finitary and has finite degree, then for any dimensional controlling scale  $\mathcal{D}$  and any distance space  $X$ , the distance space  $F^p X = (FX, d_{FX}^p)$  has  $\mathcal{D}$ -controlled dimension*

$$\dim_{\mathcal{D}} F^p X \leq \deg(F) \cdot \dim_{\mathcal{D}} X.$$

This theorem follows from Theorems 24.11 and 24.13, treating the cases of finite and infinite dimension  $\dim_{\mathcal{D}} X$ , respectively. The proofs of Theorems 24.11, 24.13, is preceded by some lemmas.

For two families of subsets  $\mathcal{V}, \mathcal{W}$  of a distance space  $(X, d_X)$ , a positive real number  $\varepsilon$ , and a subset  $W \subseteq X$ , let

$$W \cup_{\varepsilon} \mathcal{V} := W \cup \bigcup \{V \in \mathcal{V} : \underline{d}_X(V, W) < \varepsilon\}$$

and let

$$\mathcal{W} \cup_{\varepsilon} \mathcal{V} = \{W \cup_{\varepsilon} \mathcal{V} : W \in \mathcal{W}\} \cup \{V \in \mathcal{V} : \underline{d}_X(V, \bigcup \mathcal{W}) \geq \varepsilon\}.$$

The following lemma easily follows from the definition of the family  $\mathcal{W} \cup_{\varepsilon} \mathcal{V}$ .

**Lemma 24.3.** *Let  $\varepsilon$  be a positive real number and  $\mathcal{V}, \mathcal{W}$  be two families of sets in a distance space  $(X, d_X)$ . If  $\underline{d}_X(\mathcal{V}) \geq \varepsilon$  and  $\underline{d}_X(\mathcal{W}) \geq 2\varepsilon + \text{mesh}(\mathcal{V})$ , then*

$$\underline{d}_X(\mathcal{W} \cup_{\varepsilon} \mathcal{V}) \geq \varepsilon \quad \text{and} \quad \text{mesh}(\mathcal{W} \cup_{\varepsilon} \mathcal{V}) \leq 2\varepsilon + 2 \cdot \text{mesh}(\mathcal{V}) + \text{mesh}(\mathcal{W}).$$

The following lemma is crucial for our proof. It is a slightly modified version of Lemma 31 of Bell and Dranishnikov [11] (who call it the Kolmogorov trick).

**Lemma 24.4.** *Let  $k \leq m$  be two nonzero finite cardinals,  $\varepsilon$  be a positive real number, and  $\mathcal{U} = \bigcup_{i \in m} \mathcal{U}_i$  be a cover of a distance space  $X$  such that  $\inf_{i \in m} \underline{d}_X(\mathcal{U}_i) \geq \varepsilon$ , and  $\bigcup_{i \in S} \bigcup \mathcal{U}_i = X$  for every  $S \in [m]^k$ . Then there exists a cover  $\mathcal{V} = \bigcup_{i \in m+1} \mathcal{V}_i$  of  $X$  such that*

- 1)  $\inf_{i \in m+1} \underline{d}_X(\mathcal{V}_i) \geq \frac{1}{3}\varepsilon$ ;
- 2)  $\text{mesh}(\mathcal{V}) \leq \text{mesh}(\mathcal{U}) + \frac{2}{3}\varepsilon$ ;
- 3)  $X = \bigcup_{i \in S} \bigcup \mathcal{V}_i$  for every  $S \in [m+1]^k$ .

*Proof.* For every  $i \in m$  and  $U \in \mathcal{U}_i$ , consider the family  $\mathcal{V}_i = \{O[U; \frac{1}{3}\varepsilon] : U \in \mathcal{U}_i\}$  of  $\frac{1}{3}\varepsilon$ -neighborhoods of the sets  $U \in \mathcal{U}_i$  in the distance space  $(X, d_X)$ . Also let

$$\mathcal{V}_m = \left\{ \bigcap_{i \in T} U_i \setminus \bigcup_{j \in m \setminus T} \mathcal{V}_j : T \in [m]^{k-m+1} \wedge (U_i)_{i \in T} \in \prod_{i \in T} \mathcal{U}_i \right\}.$$

We claim that the cover  $\mathcal{V} = \bigcup_{i \in m+1} \mathcal{V}_i$  has the properties (1)–(3). The properties (1),(2) will follow from the definitions of the families  $\mathcal{V}_i$  as soon as we check that  $\underline{d}_X(\mathcal{V}_m) \geq \frac{1}{3}\varepsilon$ . Assuming that  $\underline{d}_X(\mathcal{V}_m) < \frac{1}{3}\varepsilon$ , we could find two distinct sets  $V, V' \in \mathcal{V}_m$  and points  $v \in V, v' \in V'$  such that  $d_X(v, v') < \frac{1}{3}\varepsilon$ . Find two sets  $T, T' \in [m]^{m-k+1}$  and families of sets  $(U_i)_{i \in T} \in \prod_{i \in T} \mathcal{U}_i, (U'_i)_{i \in T'} \in \prod_{i \in T'} \mathcal{U}_i$  such that  $V = \bigcap_{i \in T} U_i \setminus \bigcup_{i \in m \setminus T} \mathcal{V}_i$  and  $V' = \bigcap_{i \in T'} U'_i \setminus \bigcup_{i \in m \setminus T'} \mathcal{V}_i$ . If  $T = T'$ , then there exists  $i \in T$  such that  $U_i \neq U'_i$ . Then

$$\frac{1}{3}\varepsilon > d_X(v, v') \geq \underline{d}_X(U_i, U'_i) \geq \varepsilon,$$

which is a contradiction showing that  $T \neq T'$ . Then there exists  $i \in T \setminus T'$ . The inequality  $d_X(v', v) < \frac{1}{3}\varepsilon$  implies  $v' \in O(v; \frac{1}{3}\varepsilon) \subseteq \bigcup \mathcal{V}_i$  and hence  $v' \notin V' \subseteq X \setminus \bigcup \mathcal{V}_i$ . This contradiction shows that the condition (2) is satisfied.

To verify the condition (3), assume that for some set  $S \in [m+1]^k$ , the family  $\bigcup_{i \in S} \bigcup \mathcal{V}_i$  does not contain some point  $x \in X$ . Then also  $x \notin \bigcup_{i \in S \cap m} \bigcup \mathcal{V}_i$ . The choice of the family

$\mathcal{U}$  guarantees that  $|S \cap m| < k$  and hence  $m \in S$  and  $S \cap m \in [m]^{k-1}$ . Consider the set  $T := m \setminus S \in [m]^{m-k+1}$  and observe that for every  $i \in T$  the set  $(S \cap m) \cup \{i\}$  has cardinality  $k$  and hence  $x \in U_i$  for some  $U_i \in \mathcal{U}_i$ . Then  $x \in \prod_{i \in T} U_i \setminus \bigcup_{m \setminus T} \mathcal{V}_i \subseteq \bigcup \mathcal{V}_m \subseteq \bigcup_{i \in S} \bigcup \mathcal{V}_i$ , which contradicts the choice of  $x$  and completes the proof of the lemma.  $\square$

**Lemma 24.5.** *Let  $n, m$  be two nonzero finite cardinals,  $\varepsilon$  be a positive real number, and  $\mathcal{U} = \bigcup_{i \in m+1} \mathcal{U}_i$  be a cover of a distance space  $X$  such that  $\inf_{i \in m+1} \underline{d}_X(\mathcal{U}_i) \geq 3^{nm-m+1}\varepsilon$ . Then there exists a cover  $\mathcal{V} = \bigcup_{i \in nm+1} \mathcal{V}_i$  of  $X$  such that*

- 1)  $\inf_{i \in m+1} \underline{d}_X(\mathcal{V}_i) \geq \varepsilon$ ;
- 2)  $\text{mesh}(\mathcal{V}) \leq \text{mesh}(\mathcal{U}) + 3^{nm-m+1}\varepsilon$ ;
- 3) for any  $T \in [X]^n$  there exists  $i \in nm+1$  such that  $T \subseteq \bigcup \mathcal{V}_i$ .

*Proof.* Let  $\delta = 3^{nm-m+1}\varepsilon$  and  $\mathcal{W}_1 = \bigcup_{i \in m+1} \mathcal{W}_{1,i}$  where  $\mathcal{W}_{1,i} = \mathcal{U}_i$  for  $i \in m+1$ . Observe that for every set  $S \in [m+1]^{1+m}$  we have  $\bigcup_{i \in S} \bigcup \mathcal{W}_{1,i} = X$  and  $\inf_{i \in m+1} \underline{d}_X(\mathcal{W}_{1,i}) \geq \delta$ . Applying Lemma 24.4 inductively, for every  $k \in \mathbb{N}$  construct a cover  $\mathcal{W}_k = \bigcup_{i \in m+k} \mathcal{W}_{k,i}$  satisfying the following conditions:

- $\inf_{i \in m+k} \underline{d}_X(\mathcal{W}_{k,i}) \geq \frac{1}{3^k}\delta$ ;
- $\text{mesh}(\mathcal{W}_k) \leq \text{mesh}(\mathcal{U}) + \sum_{i=1}^k \left(\frac{2}{3}\right)^i \delta \leq \text{mesh}(\mathcal{U}) + \delta$ ;
- $X = \bigcup_{i \in T} \bigcup \mathcal{W}_{k,i}$  for every  $T \in [m+k]^{1+m}$ .

Then the cover  $\mathcal{V} = \bigcup_{i \in nm+1} \mathcal{V}_i$ , where  $\mathcal{V}_i = \mathcal{W}_{nm-m+1,i}$  for  $i \in nm+1$ , has the properties:

- $\inf_{i \in nm+1} \underline{d}_X(\mathcal{V}_i) \geq \frac{1}{3^{nm-m+1}}\delta = \varepsilon$ ;
- $\text{mesh}(\mathcal{V}) \leq \text{mesh}(\mathcal{U}) + \sum_{i=1}^{nm-m+1} \left(\frac{2}{3}\right)^k \delta \leq \text{mesh}(\mathcal{U}) + \delta = \text{mesh}(\mathcal{U}) + 3^{nm-m+1}\varepsilon$ ;
- $X = \bigcup_{i \in S} \bigcup \mathcal{V}_i$  for every  $S \in [nm+1]^{1+m}$ .

The last property implies that for every  $x \in X$ , the set  $S_x = \{i \in nm+1 : x \notin \bigcup \mathcal{V}_i\}$  has cardinality  $|S_x| \leq m$ . Then for every set  $T \in [X]^n$ , there exists an index  $i \in (nm+1) \setminus \bigcup_{x \in T} S_x$ . For this index we have  $T \subseteq \bigcup \mathcal{V}_i$ , which means that the condition (3) of the lemma is satisfied.  $\square$

Given a finite cardinal  $k$ , consider the subspace

$$F_k^p X = \{a \in F^p X : |\text{supp}(a)| \leq k\}$$

of the distance space  $F^p X = (FX, d_{FX}^p)$ .

**Lemma 24.6.** *Assume that the functor  $F$  is finitary and has finite degree. Let  $n := \max\{1, \deg(F)\}$  and  $k, m$  be finite cardinals such that  $k \leq n$ . Let  $D, D', D''$  be dimension controlling functions such that for any  $\varepsilon \in (0, \infty)$  and the numbers*

$$\varepsilon' = 2n \cdot |Fn| \cdot \varepsilon \quad \text{and} \quad \delta = D(3^{km-m+1}\varepsilon') + 3^{km-m+1}\varepsilon',$$

we have

$$D''(\varepsilon) \geq 2\varepsilon + 2k^{\frac{1}{p}}\delta + D'(2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}}\delta).$$

Then for any distance space  $X$  with  $\dim_D X \leq m$ , the distance space  $F_k^p X$  has  $D''$ -controlled dimension

$$\dim_{D''}(F_k^p X) \leq \max\{\dim_{D'}(F_{k-1}^p X), k \cdot \dim_D(X)\}.$$



*Proof.* By definition of the cardinal  $m' = \dim_{D'}(F_{k-1}^p X)$ , there exists a cover  $\mathcal{V} = \bigcup_{i \in 1+m'} \mathcal{V}_i$  of  $F_{k-1}^p X$  such that

$$\inf_{i \in 1+m'} \underline{d}_{F_{k-1}^p X}(\mathcal{V}_i) \geq 2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}}\delta \quad \text{and} \quad \text{mesh}(\mathcal{V}_i) \leq D'(2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}}\delta).$$

Let  $\tilde{\mathcal{V}} = \bigcup_{i \in 1+m'} \tilde{\mathcal{V}}_i$  where  $\tilde{\mathcal{V}}_i = \{O[V; \varepsilon'] : V \in \mathcal{V}_i\}$  for  $i \in 1 + m'$ .

Consider the subspace  $W = F_k^p X \setminus \bigcup \tilde{\mathcal{V}}$ .

**Claim 24.7.** *For every  $a \in W$  we have  $|\text{supp}(a)| = k$  and  $\underline{d}_X(\text{supp}(a)) \geq \varepsilon'$ .*

*Proof.* The equality  $|\text{supp}(a)| = k$  follows from the inclusion  $W \subseteq F_k^p X \setminus F_{k-1}^p X$ . Assuming that  $\underline{d}_X(\text{supp}(a)) < \varepsilon'$ , find two distinct points  $x, y \in \text{supp}(a)$  with  $d_X(x, y) < \varepsilon'$ . Using Proposition 6.1, find a function  $f : k \rightarrow X$  and an element  $b \in Fk$  such that  $f[k] = \text{supp}(a)$  and  $Ff(b) = a$ . Let  $j \in k$  be the unique number such that  $f(j) = x$ , and  $g : k \rightarrow X$  be the function such that  $g(j) = y$  and  $g(i) = f(i)$  for every  $i \in k \setminus \{j\}$ . Then the element  $c = Fg(b)$  has support  $\text{supp}(c) \subseteq g[k] = \text{supp}(a) \setminus \{x\}$  and hence  $c \in F_{k-1} X$ . Then the inequality

$$d_{F_k^p X}^p(a, c) \leq d_{F_k^p X}^p(f, g) = d_X(f(i), g(i)) = d_X(x, y) < \varepsilon'$$

implies that  $a \in \bigcup \tilde{\mathcal{V}}$ , which contradicts the inclusion  $a \in W$ .  $\square$

Since  $\dim_D X \leq m$ , we can apply Lemma 24.5 and find a cover  $\mathcal{U} = \bigcup_{i \in km+1} \mathcal{U}_i$  such that  $\text{mesh}(\mathcal{U}) \leq D(3^{km-m+1}\varepsilon') + 3^{km-m+1}\varepsilon' = \delta$ ,  $\inf_{i \in km+1} \underline{d}_X(\mathcal{U}_i) \geq \varepsilon'$  and each set  $T \in [km+1]^k$  is contained in some set  $\bigcup \mathcal{U}_i$ . For every  $i \in km+1$ , consider the sets  $X_i = \bigcup \mathcal{U}_i \subseteq X$  and

$$W_i = \{a \in W : \text{supp}(a) \subseteq X_i\} \subseteq F_k^p X.$$

For every  $a \in W_i$  let  $\mathcal{W}_i(a)$  be the set of all elements  $b \in W_i$  for which there exists a sequence of elements  $a_0, \dots, a_l \in W_i$  such that  $a = a_0$ ,  $a_l = b$  and  $d_{F_k^p X}^p(a_{j-1}, a_j) < \varepsilon$  for every  $j \in \{1, \dots, l\}$ . Let  $\mathcal{W}_i := \{\mathcal{W}_i(a) : a \in W_i\}$ . It is clear that  $\mathcal{W}_i$  is a cover of the set  $W_i$  and  $\underline{d}_{F_k^p X}(\mathcal{W}_i) \geq \varepsilon$ .

**Claim 24.8.**  $\text{mesh}(\mathcal{W}_i) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}_i) \leq k^{\frac{1}{p}} \cdot \delta$ .

*Proof.* It suffices to show that for any distinct elements  $a, b \in W_i$  with  $\mathcal{W}_i(a) = \mathcal{W}_i(b)$ , we have  $d_{F_k^p X}^p(a, b) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}_i)$ . Since  $\mathcal{W}_i(a) = \mathcal{W}_i(b)$ , there exists a sequence of elements  $b_0, \dots, b_l \in W_i$  such that  $a = b_0$ ,  $b_l = b$ , and  $d_{F_k^p X}^p(b_{j-1}, b_j) < \varepsilon$  for every  $j \in \{1, \dots, l\}$ . By Lemma 9.3, for every  $j \in \{1, \dots, l\}$  there exists a function  $r_j : X \rightarrow X$  such that  $Fr_j(b_{j-1}) = Fr_j(b_j)$ ,  $r_j \circ r_j = r_j$ ,  $r_j[\text{supp}(b_{j-1}) \cup \text{supp}(b_j)] \subseteq \text{supp}(b_{j-1}) \cup \text{supp}(b_j)$ , and

$$\sup_{x \in X} d_X(x, r_j(x)) < n \cdot |Fn| \cdot \varepsilon = \frac{1}{2}\varepsilon'.$$

The latter inequality and Claim 24.7 imply that the restrictions  $r_j \upharpoonright \text{supp}(b_{j-1})$  and  $r_j \upharpoonright \text{supp}(b_j)$  are injective maps. Then

$$r_j[\text{supp}(b_{j-1})] = \text{supp}(Fr_j(b_{j-1})) = \text{supp}(Fr_j(b_j)) = r_j[\text{supp}(b_j)],$$

and we can define a function  $h_j : \text{supp}(b_{j-1}) \rightarrow \text{supp}(b_j)$  assigning to each point  $x \in \text{supp}(b_{j-1})$  a unique point  $y \in \text{supp}(b_j)$  such that  $r_j(x) = r_j(y)$ . Observe that

$$d_X(x, h_j(x)) \leq d_X(x, r_j(x)) + d_X(r_j(x), y) < \frac{1}{2}\varepsilon' + \frac{1}{2}\varepsilon = \varepsilon'.$$

Since  $r_j = r_j \circ r_j$ , we can extend the function  $h_j$  to a functions  $\bar{h}_j : X \rightarrow X$  such that  $r_j = r_j \circ \bar{h}_j$ .

**Claim 24.9.**  $F\bar{h}_j(b_{j-1}) = b_j$ .

*Proof.* Let  $S = \text{supp}(b_j)$  and  $e : S \rightarrow X_i$  be the identity embedding. The injectivity of the map  $r_j \circ e : S \rightarrow X$  implies the injectivity of the map  $Fr_j \circ Fe : FS \rightarrow FX$ . Then the maps  $Fe$  and  $Fr_j|_{F[S; X]}$  are injective. The equality  $r_j = r_j \circ \bar{h}_j$  implies  $Fr_j(b_j) = Fr_j(b_{j-1}) = Fr_j \circ F\bar{h}_j(b_{j-1})$ . Since  $b_j, F\bar{h}_j(b_{j-1}) \in F[S; X]$ , the injectivity of the restriction  $Fr_j|_{F[S; X]}$  ensures that  $b_j = F\bar{h}_j(b_{j-1})$ .  $\square$

Then for the bijective map  $h = h_l \circ \dots \circ h_1 : \text{supp}(b_0) \rightarrow \text{supp}(b_l)$  we get  $Fh(a) = Fh(b_0) = b_l = b$ .

For every  $j \in \{0, \dots, l\}$  consider the map  $u_j : \text{supp}(b_j) \rightarrow \mathcal{U}_i$  assigning to each point  $x \in \text{supp}(b_j)$  a unique set  $u_j(x) \in \mathcal{U}_i$  containing  $x$ . Taking into account that  $\underline{d}_X(\mathcal{U}_i) \geq \varepsilon' > d_X(x, h_j(x))$ , we conclude that  $u_{j-1} = u_j \circ h_j$  for every  $j \in \{1, \dots, l\}$ . Consequently,

$$u_0 = u_1 \circ h_1 = u_2 \circ h_2 \circ h_1 = \dots = u_l \circ h_l \circ \dots \circ h_1 = u_l \circ h$$

and for every  $x \in \text{supp}(a)$ ,

$$d_X(x, h(x)) \leq \text{diam}(u_0(x)) \leq \text{mesh}(\mathcal{U}_i).$$

Since  $a \in F_k X$ , there exists an element  $a' \in Fk$  and a function  $f : k \rightarrow \text{supp}(a)$  such that  $a = Ff(a')$ . Then  $b = Fh(a) = F(h \circ f)(a')$ . Now the definition of the metric  $d_{FX}^p$  ensures that

$$d_{FX}^p(a, b) \leq d_{Xk}^p(f, h \circ f) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}_i) \leq k^{\frac{1}{p}} \delta.$$

$\square$

**Claim 24.10.**  $\mathcal{W} = \bigcup_{i \in km+1} \mathcal{W}_i$  is a cover of the set  $W$  such that

$$\inf_{i \in km+1} \mathcal{W}_i \geq \varepsilon, \quad \text{mesh}(\mathcal{W}) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}) \leq k^{\frac{1}{p}} \cdot \delta \quad \text{and} \quad 2\varepsilon + \text{mesh}(\mathcal{W}) \leq \inf_{i \in 1+m'} \underline{d}_{FX}(\tilde{\mathcal{V}}_i).$$

*Proof.* The inequality  $\varepsilon \leq \inf_{i \in km+1} \mathcal{W}_i \geq \varepsilon$  follows from the definition of the families  $\mathcal{W}_i$ ; the upper bound  $\text{mesh}(\mathcal{W}) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}) \leq k^{\frac{1}{p}} \cdot \delta$  is proved in Claim 24.8.

The choice of the covers  $\mathcal{V}$  and  $\tilde{\mathcal{V}}$  guarantees that for every  $i \in 1+m'$  we have

$$\underline{d}_{FX}(\tilde{\mathcal{V}}_i) \geq \underline{d}_{FX}(\mathcal{V}_i) - 2\varepsilon' \geq 2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}} \delta - 2\varepsilon' = 2\varepsilon + k^{\frac{1}{p}} \delta \geq 2\varepsilon + \text{mesh}(\mathcal{W}).$$

$\square$

Let  $m'' = \max\{m', km - m\}$  and choose any surjective maps  $\alpha : 1+m'' \rightarrow 1+m'$  and  $\beta : 1+m'' \rightarrow nm - n + 1$ . For every  $i \in 1+m''$ , consider the family  $\tilde{\mathcal{W}}_i = \tilde{\mathcal{V}}_{\alpha(i)} \cup_{\varepsilon} \mathcal{W}_{\beta(i)}$ , and the cover  $\tilde{\mathcal{W}} = \bigcup_{i \in 1+m''} \tilde{\mathcal{W}}_i$  of the distance space  $F_k^p X$ . Claim 24.10 and Lemma 24.3 imply that  $\inf_{i \in 1+m''} \underline{d}_{FX}(\tilde{\mathcal{W}}_i) \geq \varepsilon$  and

$$\text{mesh}(\tilde{\mathcal{W}}) \leq 2\varepsilon + 2 \cdot \text{mesh}(\mathcal{W}) + \text{mesh}(\tilde{\mathcal{V}}) \leq 2\varepsilon + 2k^{\frac{1}{p}} \delta + D'(2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}} \delta) \leq D''(\varepsilon),$$

witnessing that

$$\dim_{D''} F_k^p X \leq m'' = \max\{\dim_{D'} F_{k-1}^p X, k \cdot \dim_D X\}.$$

$\square$

**Theorem 24.11.** *Assume that the functor  $F$  is finitary and has finitary degree. Let  $n = \max\{1, \deg(F)\}$ . Assume that for some dimension controlling function  $D$ , the distance space  $X$  has finite  $D$ -controlled dimension  $m = \dim_D X$ . Consider the sequence of dimension controlling functions  $(D_k)_{k \in \omega}$  defined by the recursive formula:  $D_0(\varepsilon) = 0$  and*

$$D_k(\varepsilon) = 2\varepsilon + 2k^{\frac{1}{p}}(D(2n \cdot |Fn| \cdot 3^{km-m+1}\varepsilon) + 2n \cdot |Fn| \cdot 3^{km-m+1}\varepsilon) + \\ D_{k-1}(2\varepsilon(1 + n \cdot |Fn|) + k^{\frac{1}{p}}(D(2n \cdot |Fn| \cdot 3^{km-m+1}\varepsilon) + 2n \cdot |Fn| \cdot 3^{km-m+1}\varepsilon))$$

for every  $\varepsilon \in (0, \infty)$ . Then  $\dim_{D_k} F_k^p X \leq k \cdot \dim_D X$  for every  $k \leq \deg(F)$ . In particular,  $\dim_{D_n} F^n X \leq \deg(F) \cdot \dim_D X$ .

*Proof.* By Theorem 8.1, for any distinct points  $a, b \in F_0 X$  we have  $d_{F_0 X}^p(a, b) = \infty$ , which implies that  $\dim_{D_0} F_0^p X = 0 \leq 0 \cdot \dim_D X$ . Now assume that for positive number  $k \leq \deg(F)$  we proved that  $\dim_{D_{k-1}} F_{k-1} X \leq (k-1) \cdot \dim_{D_{k-1}} X$ . Applying Lemma 24.6, we conclude that

$$\dim_{D_k} F_k^p X \leq \max\{\dim_{D_{k-1}} F_{k-1}^p X, k \cdot \dim_D X\} \leq \\ \max\{(k-1) \cdot \dim_D X, k \cdot \dim_D X\} = k \cdot \dim_D X. \quad \square$$

Next, we consider the case of infinite dimension  $\dim_D X$ .

**Lemma 24.12.** *Assume that the functor  $F$  is finitary and has finite degree. Let  $n := \max\{1, \deg(F)\}$  and  $k \leq n$ . Let  $D, D', D''$  be dimension controlling functions such that for any  $\varepsilon \in (0, \infty)$  and numbers*

$$\varepsilon' = 2n \cdot |Fn| \cdot \varepsilon \quad \text{and} \quad \delta = D(3^{k^2-2k+2}\varepsilon') + 3^{k^2-2k+2}\varepsilon',$$

we have

$$D''(\varepsilon) \geq 2\varepsilon + 2k^{\frac{1}{p}}\delta + D'(2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}}\delta).$$

Then for any distance space  $X$  with  $\dim_D X \geq \omega$ , the distance space  $F_k^p X$  has  $D''$ -controlled dimension

$$\dim_{D''}(F_k^p X) \leq \max\{\dim_{D'}(F_{k-1}^p X), \dim_D(X)\}.$$

*Proof.* By definition of the cardinal  $m' := \dim_{D'}(F_{k-1}^p X)$ , there exists a cover  $\mathcal{V} = \bigcup_{i \in 1+m'} \mathcal{V}_i$  of  $F_{k-1}^p X$  such that

$$\inf_{i \in 1+m'} \underline{d}_{F^p X}(\mathcal{V}_i) \geq 2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}}\delta \quad \text{and} \quad \text{mesh}(\mathcal{V}_i) \leq D'(2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}}\delta).$$

Let  $\tilde{\mathcal{V}} = \bigcup_{i \in 1+m'} \tilde{\mathcal{V}}_i$  where  $\tilde{\mathcal{V}}_i = \{O[V; \varepsilon'] : V \in \mathcal{V}_i\}$  for  $i \in 1+m'$ .

Consider the subspace  $W = F_k^p X \setminus \bigcup \tilde{\mathcal{V}}$ . By analogy with Claim 24.7, we can prove that for every  $a \in W$  we have  $|\text{supp}(a)| = k$  and  $\underline{d}_X(\text{supp}(a)) \geq \varepsilon'$ .

By definition of the (infinite) cardinal  $m := \dim_D X \geq \omega$ , there exists a cover  $\mathcal{U} = \bigcup_{i \in 1+m} \mathcal{U}_i$  of  $X$  such that  $\text{mesh}(\mathcal{U}) \leq D(3^{k^2-2k+2}\varepsilon')$  and  $\inf_{i \in 1+m} \underline{d}_X(\mathcal{U}_i) \geq 3^{k^2-2k+2}\varepsilon'$ . For any set  $\alpha \in [1+m]^k$ , consider the cover  $\bigcup_{i \in \alpha} \mathcal{U}_i$  of the set  $X_\alpha = \bigcup \mathcal{U}_\alpha$ . Observe that for every (in fact, unique) set  $S \in [\alpha]^k$ , we have  $X_\alpha = \bigcup_{i \in S} \mathcal{U}_i = \bigcup_{i \in \alpha} \mathcal{U}_i$  and

$$\inf_{i \in \alpha} \underline{d}_X(\mathcal{U}_i) \geq \inf_{i \in 1+m} \underline{d}_X(\mathcal{U}_i) \geq 3^{k^2-2k+2}\varepsilon' = 3^{k(k-1)-(k-1)+1}\varepsilon'.$$

By Lemma 24.5, there exists a cover  $\mathcal{U}_\alpha = \bigcup_{i \in k^2-k+1} \mathcal{U}_{(\alpha, i)}$  such that

- (a)  $\inf_{i \in k^2 - k + 1} \underline{d}_X(\mathcal{U}_{(\alpha, i)}) \geq \varepsilon'$ ;
- (b)  $\text{mesh}(\mathcal{U}_\alpha) \leq \text{mesh}(\mathcal{U}) + 3^{k^2 - 2k + 2} \varepsilon' \leq D(3^{k^2 - 2k + 2} \varepsilon') + 3^{k^2 - 2k + 2} \varepsilon' = \delta$ ;
- (c) for any  $T \in [X_\alpha]^k$  there exists  $i \in k^2 - k + 1$  such that  $T \subseteq \bigcup \mathcal{U}_{(\alpha, i)}$ .

For every  $\alpha \in [1 + m]^k$  and  $i \in k^2 - k + 1$ , consider the sets  $X_{(\alpha, i)} = \bigcup \mathcal{U}_{(\alpha, i)} \subseteq X$  and

$$W_{(\alpha, i)} = \{a \in W : \text{supp}(a) \subseteq X_{(\alpha, i)}\} \subseteq F_k^p X.$$

Let us show that

$$W = \bigcup_{\alpha \in [1+m]^k} \bigcup_{i \in k^2 - k + 1} W_{\alpha, i}.$$

Given any element  $a \in W \subseteq F_k X \setminus F_{k-1} X$ , consider its support  $\text{supp}(a) \in [X]^k$  and find a set  $\alpha \in [1 + m]^k$  such that  $\text{supp}(a) \subseteq \bigcup_{i \in \alpha} \mathcal{U}_i = X_\alpha$ . By the condition (c), there exists  $i \in k^2 - k + 1$  such that  $\text{supp}(a) \subseteq \bigcup \mathcal{U}_{(\alpha, i)} = X_{(\alpha, i)}$ . Then  $a \in W_{(\alpha, i)}$  by the definition of  $W_{(\alpha, i)}$ .

For every  $a \in W_{(\alpha, i)}$  let  $\mathcal{W}_{(\alpha, i)}(a)$  be the set of all elements  $b \in W_{(\alpha, i)}$  for which there exists a sequence of elements  $a_0, \dots, a_l \in W_{(\alpha, i)}$  such that  $a = a_0$ ,  $a_l = b$  and  $d_{F_k X}^p(a_{j-1} a_j) < \varepsilon$  for every  $j \in \{1, \dots, l\}$ . Let  $\mathcal{W}_{(\alpha, i)} := \{\mathcal{W}_{(\alpha, i)}(a) : a \in W_{(\alpha, i)}\}$ . It is clear that  $\mathcal{W}_{(\alpha, i)}$  is a cover of the set  $W_{(\alpha, i)}$  and  $\underline{d}_{F_k X}(\mathcal{W}_{(\alpha, i)}) \geq \varepsilon$ .

By analogy with Claim 24.8, we can prove that

$$\text{mesh}(\mathcal{W}_{(\alpha, i)}) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U}_i) \leq k^{\frac{1}{p}} \cdot \delta,$$

which implies that  $\mathcal{W} = \bigcup_{\alpha \in [1+m]^k} \bigcup_{i \in k^2 - k + 1} \mathcal{W}_{(\alpha, i)}$  is a cover of the set  $W$  such that

$\inf_{\alpha \in [1+m]^k} \inf_{i \in k^2 - k + 1} \underline{d}_X(\mathcal{W}_{(\alpha, i)}) \geq \varepsilon$  and  $\text{mesh}(\mathcal{W}) \leq k^{\frac{1}{p}} \cdot \text{mesh}(\mathcal{U})$ . By analogy with Claim 24.10, we can show that  $2\varepsilon + \text{mesh}(\mathcal{W}) \leq \inf_{i \in [1+m]'} \underline{d}_{F_k X}(\tilde{\mathcal{V}}_i)$ .

Let  $m'' = \max\{m', m\}$  and choose any surjective maps  $\alpha : 1 + m'' \rightarrow 1 + m'$  and  $\beta : 1 + m'' \rightarrow [1 + m]^k \times (k^2 - k + 1)$  (the map  $\beta$  exists since the cardinal  $m$  is infinite). For every  $i \in 1 + m''$ , consider the family  $\mathcal{U}_i = \tilde{\mathcal{V}}_{\alpha(i)} \cup_\varepsilon \mathcal{W}_{\beta(i)}$ , and the cover  $\mathcal{U} = \bigcup_{i \in [1+m]''} \mathcal{U}_i$  of the distance space  $F_k^p X$ . Using Lemma 24.3, we can show that  $\inf_{i \in [1+m]''} \underline{d}_{F_k X}(\mathcal{U}_i) \geq \varepsilon$  and

$$\text{mesh}(\mathcal{U}) \leq 2\varepsilon + 2 \cdot \text{mesh}(\mathcal{W}) + \text{mesh}(\tilde{\mathcal{V}}) \leq 2\varepsilon + 2k^{\frac{1}{p}} \delta + D'(2\varepsilon + 2\varepsilon' + k^{\frac{1}{p}} \delta) \leq D''(\varepsilon),$$

witnessing that

$$\dim_{D''} F_k^p X \leq m'' = \max\{\dim_{D'} F_{k-1}^p X, \dim_D X\}.$$

□

Lemma 24.12 implies the following infinite version of Theorem 24.11.

**Theorem 24.13.** *Assume that the functor  $F$  is finitary and has finitary degree. Let  $n = \max\{1, \deg(F)\}$ . Assume that for some dimension controlling function  $D$ , the distance space  $X$  has infinite  $D$ -controlled dimension  $\dim_D X$ . Consider the sequence of dimension controlling functions  $(D_k)_{k \in \omega}$  defined by the recursive formula:  $D_0(\varepsilon) = 0$  and*

$$\begin{aligned} D_k(\varepsilon) &= 2\varepsilon + 2k^{\frac{1}{p}} (D(2n \cdot |Fn| \cdot 3^{k^2 - 2k + 2} \varepsilon) + 2n \cdot |Fn| \cdot 3^{k^2 - 2k + 2} \varepsilon) + \\ &D_{k-1}(2\varepsilon(1 + n \cdot |Fn|) + k^{\frac{1}{p}} (D(2n \cdot |Fn| \cdot 3^{k^2 - 2k + 2} \varepsilon) + 2n \cdot |Fn| \cdot 3^{k^2 - 2k + 2} \varepsilon)) \end{aligned}$$

for every  $\varepsilon \in (0, \infty)$ . Then  $\dim_{D_k} F_k^p X \leq \dim_D X$  for every  $k \leq \deg(F)$ . In particular,  $\dim_{D_n} F^p X \leq \dim_D X$ .

Theorems 24.11, 24.13 imply Theorem 24.2 because for any dimension controlling scale  $\mathcal{D}$  and any function  $D \in \mathcal{D}$ , the dimension controlling functions  $D_k$  defined by the recursive formulas in Theorems 24.11, 24.13 belong to the scale  $\mathcal{D}$ .

Now we consider three important examples of dimension controlling scales.

**Example 24.14.** Consider the family  $\mathcal{AN}$  of all non-decreasing functions  $f : (0, \infty) \rightarrow (0, \infty)$  such that

$$\sup_{x \in (0, \infty)} \frac{f(x)}{x} < \infty.$$

It is easy to see that  $\mathcal{AN}$  is the smallest dimension controlling scale. For metric spaces, the  $\mathcal{AN}$ -controlled dimension  $\dim_{\mathcal{AN}}$  is known as the *Assouad-Nagata dimension*, see [25] and [15]. For this dimension, Theorem 24.2 implies the formula

$$\dim_{\mathcal{AN}} F^p X \leq \deg(F) \cdot \dim_{\mathcal{AN}} X$$

holding for any  $p \in [1, \infty]$ , distance space  $X$  and finitary functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  of finite degree.

**Example 24.15.** Consider the dimension controlling scale  $\mathcal{AS}$  consisting of all non-decreasing functions  $f : (0, \infty) \rightarrow (0, \infty)$ . For metric spaces, the  $\mathcal{AS}$ -controlled dimension  $\dim_{\mathcal{AS}}$  is known as the *asymptotic dimension of Gromov*, see [21]. For the asymptotic dimension, Theorem 24.2 implies the formula

$$\dim_{\mathcal{AS}} F^p X \leq \deg(F) \cdot \dim_{\mathcal{AS}} X$$

holding for any  $p \in [1, \infty]$ , distance space  $X$  and finitary functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  of finite degree. This formula has been announced by Radul and Shukel in [27], but their proof works only for support preserving functors.

**Example 24.16.** Consider the dimension controlling scale  $\mathcal{U}$  consisting of all non-decreasing functions  $f : (0, \infty) \rightarrow (0, \infty]$  such that

$$\lim_{x \rightarrow +0} f(x) = 0.$$

For metric spaces, the  $\mathcal{U}$ -controlled dimension  $\dim_{\mathcal{U}}$  is known as the *uniform covering dimension of Isbell*, see [23, Ch.5]. For the uniform dimension, Theorem 24.2 implies the formula

$$\dim_{\mathcal{U}} F^p X \leq \deg(F) \cdot \dim_{\mathcal{U}} X$$

holding for any  $p \in [1, \infty]$ , distance space  $X$  and finitary functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  of finite degree. For a compact metric space  $X$ , its uniform covering dimension  $\dim_{\mathcal{U}} X$  coincides with its topological dimension  $\dim X$ . In this case the above formula turns into the formula

$$\dim FX \leq \deg(F) \cdot \dim X,$$

proved for normal functors in the category of compact Hausdorff spaces by Basmanov, see [8, 9, 10].

25. CONCLUDING THEOREMS ON THE PROPERTIES OF THE DISTANCE  $d_{FX}^p$ 

In the following two theorems we unify the main results of the preceding sections, describing the properties of the functor  $F^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$ .

**Theorem 25.1.** *For any  $p \in [1, \infty]$ , and a nonempty distance space  $(X, d_X)$ , the  $\ell^p$ -metrization  $F^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$  of the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  has the following properties:*

- 1)  $d_{FX}^p$  is the largest distance on  $FX$  such that for every  $n \in \mathbb{N}$  and  $a \in Fn$ , the map  $\xi_X^a : (X^n, d_{X^n}^p) \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding.
- 2) if  $F$  has finite degree  $n = \deg(F)$ , then  $d_{FX}^p$  is the largest distance on  $FX$  such that for every  $a \in Fn$ , the map  $\xi_X^a : (X^n, d_{X^n}^p) \rightarrow FX$ ,  $\xi_X^a : f \mapsto Ff(a)$ , is non-expanding.
- 3)  $d_{FX}^p(a, b) = \infty$  for any distinct points  $a, b \in FX$  with  $|\text{supp}(a) \cup \text{supp}(b)| \leq 1$ .
- 4)  $d_{FX}^\infty \leq d_{FX}^p \leq d_{FX}^1$  and  $d_{FX}^p \leq n^{\frac{1}{p}} \cdot d_{FX}^\infty$  where  $n = \max\{1, \deg(F)\}$ .
- 5) For every Lipschitz map  $f$  between distance spaces, the map  $F^p f$  is Lipschitz with Lipschitz constant  $\text{Lip}(F^p f) \leq \text{Lip}(f)$ .
- 6) If  $f : X \rightarrow Y$  is an injective isometry of distance spaces and the image  $f[X]$  is dense in  $Y$ , then the map  $F^p f$  is an isometry.
- 7) If  $f : X \rightarrow Y$  is an isometry between distance spaces and the functor  $F$  is finitary and has finite degree, then for every  $a, b \in FX$ ,  $d_{FX}^\infty(a, b) \leq 2n \cdot |Fn| \cdot d_{FY}^\infty(Ff(a), Ff(b))$  where  $n = \max\{1, \deg(F)\}$ .
- 8) For any  $a, b \in FX$  with  $d_{FX}^p(a, b) < \infty$  we have  $d_{FX}^p(a, b) \leq (|S_a|^{\frac{1}{p}} + |S_b|^{\frac{1}{p}}) \cdot \bar{d}_X(S_a \cup S_b)$  and  $d_{FX}^p(a, b) \leq |S_a|^{\frac{1}{p}} \cdot \bar{d}_X(S_a) + \underline{d}_X(S_a, S_b) + |S_b|^{\frac{1}{p}} \cdot \bar{d}_X(S_b)$ , where  $S_a = \text{supp}(a)$  and  $S_b = \text{supp}(b)$ .
- 9) If the functor  $F$  has finite degree  $\leq n$ , then  $\bar{d}_{FX}^p(FX) \leq 2n^{\frac{1}{p}} \cdot \bar{d}_X(X)$ .
- 10)  $\underline{d}_{FX}^p(FX) \geq \underline{d}_X(X)$ ; moreover  $d_{FX}^1(a, b) \geq \underline{d}_X(\text{supp}(a) \cup \text{supp}(b))$  for any distinct elements  $a, b \in FX$ .
- 11) If  $F$  preserves supports, then for any distinct elements  $a, b \in FX$  we have  $d_{FX}^p(a, b) \geq d_{FX}^\infty(a, b) \geq \max\{d_{HX}(\text{supp}(a), \text{supp}(b)), \frac{1}{3}\underline{d}_X(\text{supp}(a) \cup \text{supp}(b))\}$ .
- 12) If  $F1$  is a singleton, and  $d_X$  is a pseudometric, then the distance  $d_{FX}^p$  is a pseudometric.
- 13) The distance  $d_{FX}^p$  is an  $\infty$ -metric if  $d_X$  is an  $\infty$ -metric and one of the following conditions holds: (i) the  $\infty$ -metric space  $(X, d_X)$  is Lipschitz disconnected, (ii)  $p = 1$ , (iii) the functor  $F$  has finite degree, (iv) the functor  $F$  preserves supports.
- 14) The distance  $d_{FX}^p$  is a metric if  $d_X$  is a metric,  $F1$  is a singleton and one of the following conditions holds: (i) the metric space  $(X, d_X)$  is Lipschitz disconnected, (ii)  $p = 1$ , (iii) the functor  $F$  has finite degree, (iv) the functor  $F$  preserves supports.
- 15) If for some  $a \in F1$  the map  $\xi_2^a : 2 \rightarrow F2$  is injective and  $p = 1$  or  $F$  preserves supports, then the function  $\xi_X^a : X \rightarrow F^p X$  is an injective isometry.
- 16) If  $d_X$  is  $\{0, \infty\}$ -valued (and  $\infty$ -metric) on  $X$ , then  $d_{FX}^p$  is  $\{0, \infty\}$ -valued (and  $\infty$ -metric) on  $FX$ .

*Proof.* 1. See Theorem 10.1.

2. See Theorem 10.2.

3. See Theorem 8.1.
4. See Theorem 11.2.
5. See Theorem 12.1.
6. See Theorem 13.3.
7. See Proposition 13.5.
8. See Theorem 15.2.
9. See Corollary 15.4.
10. See Proposition 16.1 and Theorem 16.4.
11. See Theorem 16.6.
12. See Corollary 15.3.
13. Apply Theorems 16.3, 16.4, 11.2, and 16.6.
14. Apply the statements (11) and (12) of this theorem.
15. See Theorem 18.4.
16. See Lemma 15.1. □

For finitary functors of finite degree we have some additional results.

**Theorem 25.2.** *If the functor  $F$  is finitary and has finite degree  $n = \deg(F) > 1$ , then for any  $p \in [1, \infty]$ , and any nonempty distance space  $(X, d_X)$ , the  $\ell^p$ -metrization  $F^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$  of the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  has the following properties:*

- 1) *For any function  $f : X \rightarrow Y$  to a distance space  $Y$ , the function  $F^p f : F^p X \rightarrow F^p Y$  has continuity modulus  $\omega_{F^p f} \leq |Fn| \cdot n^{\frac{1}{p}} \cdot \omega_f$ .*
- 2) *If a function  $f : X \rightarrow Y$  to a distance space is uniformly continuous (resp. asymptotically Lipschitz, macro-uniform, a quasi-isometry, a coarse equivalence), then so is the function  $F^p f$ .*
- 3) *If the functor  $F$  preserves supports, then for any continuous function  $f : X \rightarrow Y$  between distance spaces, the function  $F^p f : F^p X \rightarrow F^p Y$  is continuous.*
- 4) *If the distance space  $X$  is complete (and compact), then so is the space  $F^p X$ .*
- 5) *The distance space  $F^p X$  has metric entropy  $\mathbb{E}_\varepsilon(F^p X) \leq |Fn| \cdot \mathbb{E}_\varepsilon(X^n, d_{X^n}^p) \leq |Fn| \cdot (\mathbb{E}_{\varepsilon/\sqrt[p]{n}}(X))^n$  for every  $\varepsilon \in (0, \infty)$ .*
- 6) *The distance space  $F^p X$  has cardinal entropy characteristics  $\mathbb{E}_0(F^p X) \leq |Fn| \cdot \mathbb{E}_0(X)^n$  and  $\mathbb{E}_\infty(F^p X) \leq |Fn| \cdot \mathbb{E}_\infty(X)^n$ .*
- 7) *If the distance space  $X$  is totally bounded, then the distance space  $F^p X$  is totally bounded and has upper and lower box-counting dimensions  $\overline{\dim}_B F^p X \leq \deg(F) \cdot \overline{\dim}_B X$  and  $\underline{\dim}_B F^p X \leq \deg(F) \cdot \underline{\dim}_B X$ .*
- 8) *If the functor  $F$  preserves supports, then the distance space  $F^p X$  has metric entropy  $\mathbb{E}_{\varepsilon, \delta}(F^p X) \leq n^n \cdot |Fn| \cdot \mathbb{E}_{\varepsilon, 2\delta\sqrt[p]{n}}(X^n, d_{X^n}^p) \leq n^n \cdot |Fn| \cdot (\mathbb{E}_{\varepsilon/\sqrt[p]{n}, 2\delta\sqrt[p]{n}}(X))^n$  for any  $\varepsilon, \delta \in (0, \infty)$ .*
- 9) *If the functor  $F$  preserves supports, then the distance space  $F^p X$  has the cardinal entropy characteristics  $\mathbb{E}_{(\varepsilon, \delta)}(F^p X) \leq n^n \cdot |Fn| \cdot \mathbb{E}_{(\varepsilon, \delta)}(X)^n$  for any  $(\varepsilon, \delta) \in \{(0, 0), (0, \infty), (\infty, \infty)\}$ .*
- 10) *If for some  $(\varepsilon, \delta) \in \{(0, 0), (0, \infty), (\infty, \infty)\}$  a distance space  $X$  is  $(\varepsilon, \delta)$ -finitary, then the distance space  $F^p X$  is  $(\varepsilon, \delta)$ -finitary and has  $(\varepsilon, \delta)$ -entropy dimensions  $\overline{\dim}_{(\varepsilon, \delta)} F^p X \leq \deg(F) \cdot \overline{\dim}_{(\varepsilon, \delta)} X$  and  $\underline{\dim}_{(\varepsilon, \delta)} F^p X \leq \deg(F) \cdot \underline{\dim}_{(\varepsilon, \delta)} X$ .*
- 11) *The distance space  $F^p X$  has Hausdorff dimension  $\dim_H F^p X \leq \dim_H X^n$ .*
- 12) *For any dimension controlling scale  $\mathcal{D}$ , the distance space  $F^p X$  has  $\mathcal{D}$ -controlled dimension  $\dim_{\mathcal{D}} F^p X \leq \deg(F) \cdot \dim_{\mathcal{D}} X$ .*

*Proof.* 1. See Theorem 14.1.

2. See Corollaries 14.2 and 14.4.

3. See Theorem 17.1.

4. See Theorems 20.1 and 19.1.

5. See Theorem 21.1.

6. See Corollary 21.5(1,2).

7. See Corollary 21.6 and Theorem 22.2.

8. See Theorem 21.2.

9. See Corollary 21.5(3,4,5).

10. See Theorem 22.4.

11. See Theorem 23.2.

12. See Theorem 24.2. □

## 26. THE $\check{\ell}^p$ -METRIZATION $\check{F}^p$ OF THE FUNCTOR $F$

In this section we introduce a lifting  $\check{F}^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$  of the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$ , called the  $\check{\ell}^p$ -metrization of  $F$ . Comparing to the  $\ell_p$ -metrization  $F^p$ , the  $\check{\ell}^p$ -metrization  $\check{F}^p$  has some additional properties, in particular, it preserves isometries between distance spaces. The  $\check{\ell}^p$ -metrization is defined with the help of injective envelopes which were discussed in Section 5.

For any distance space  $(X, d_X)$ , the distance  $\check{d}_{FX}^p : FX \times FX \rightarrow [0, \infty]$  is defined by the formula

$$\check{d}_{FX}^p(a, b) = d_{FEX}^p(Fe_X(a), Fe_X(b)) \quad \text{for } a, b \in FX.$$

Here  $e_X : X \rightarrow EX$  is the canonical isometry of  $X$  into its injective envelope  $EX$ .

The  $\check{\ell}^p$ -metrization of the functor  $F$  is the functor  $\check{F}^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$  assigning to each distance space  $\mathbf{X} = (X, d_X)$  the distance space  $\check{F}^p\mathbf{X} = (FX, \check{d}_{FX}^p)$  and to each function  $f : (X, d_X) \rightarrow (Y, d_Y)$  between distance spaces the function  $\check{F}^p f := Ff$ .

It is clear that  $U \circ \check{F}^p = F \circ U$ , where  $U : \mathbf{Dist} \rightarrow \mathbf{Set}$ ,  $U : (X, d_X) \mapsto X$ , is the forgetful functor. So,  $\check{F}^p$  is a lifting of the functor  $F$  to the category  $\mathbf{Dist}$ . In the following theorem we establish some properties of this lifting.

**Theorem 26.1.** *The  $\check{\ell}^p$ -metrization  $\check{F}^p : \mathbf{Dist} \rightarrow \mathbf{Dist}$  of the functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  has the following properties for any nonempty distance space  $(X, d_X)$ :*

- 1)  $\check{d}^p(a, b) = \infty$  if  $a, b \in FX$  are distinct elements with  $|\text{supp}(a) \cup \text{supp}(b)| \leq 1$ .
- 2)  $\check{d}_{FX}^\infty \leq \check{d}_{FX}^p \leq \check{d}_{FX}^1$  and  $\check{d}_{FX}^p \leq n^{\frac{1}{p}} \cdot \check{d}_{FX}^\infty$ , where  $n = \max\{1, \deg(F)\}$ .
- 3)  $\check{d}_{FX}^p \leq d_{FX}^p$  and  $d_{FX}^\infty \leq n \cdot |Fn| \cdot \check{d}_{FX}^\infty$ , where  $n = \max\{1, \deg(F)\}$ ;  
moreover,  $\check{d}_{FX}^p = d_{FX}^p$  if the distance space  $(X, d_X)$  is injective.
- 4) For every Lipschitz map  $f$  between distance spaces, the map  $\check{F}^p f$  is Lipschitz with Lipschitz constant  $\text{Lip}(\check{F}^p f) \leq \text{Lip}(f)$ .
- 5) For any isometry  $f$  between distance spaces, the map  $\check{F}^p f$  is an isometry.
- 6)  $\check{d}_{FX}^p$  is a pseudometric if  $d_X$  is a pseudometric and  $F1$  is a singleton.
- 7)  $\check{d}_{FX}^1(a, b) \geq \underline{d}_X(\text{supp}(a) \cup \text{supp}(b))$  for any distinct elements  $a, b \in FX$ .
- 8) If  $F$  preserves supports, then for any distinct elements  $a, b \in FX$  we have  
 $\check{d}_{FX}^p(a, b) \geq \check{d}_{FX}^\infty(a, b) \geq \max\{d_{HX}(\text{supp}(a), \text{supp}(b)), \frac{1}{3}\underline{d}_X(\text{supp}(a) \cup \text{supp}(b))\}$ .
- 9)  $\check{d}_{FX}^p$  is an  $\infty$ -metric if the distances  $d_X$  and  $d_{FEX}^p$  are  $\infty$ -metric.



- 10)  $\check{d}_{FX}^p$  is an  $\infty$ -metric if  $d_X$  is an  $\infty$ -metric and one of the following conditions is satisfied: (i)  $p = 1$ , (ii)  $F$  has finite degree, (iii)  $F$  preserves supports.
- 11)  $\check{d}_{FX}^p$  is a metric if  $d_X$  is a metric,  $F1$  is a singleton, and one of the following conditions is satisfied: (i)  $p = 1$ , (ii)  $F$  has finite degree, (iii)  $F$  preserves supports.
- 12) If for some  $a \in F1$  the map  $\xi_2^a : 2 \rightarrow F2$  is injective and  $p = 1$  or  $F$  preserves supports, then the function  $\xi_X^a : X \rightarrow \check{F}^p X$  is an injective isometry.
- 16) If  $d_X$  is the  $\{0, \infty\}$ -valued  $\infty$ -metric on  $X$ , then  $\check{d}_{FX}^p$  is the  $\{0, \infty\}$ -valued  $\infty$ -metric on  $FX$ .

*Proof.* 1. Assume that  $a, b$  are two distinct elements of  $FX$  with  $|\text{supp}(a) \cup \text{supp}(b)| \leq 1$ . Take any singleton  $S \subseteq X$  containing the set  $\text{supp}(a) \cup \text{supp}(b)$ , and let  $r : X \rightarrow S$  be the constant map. By Proposition 6.1,  $a, b \in Fi_{S,X}[S; X]$  and hence  $a = Fi_{S,X}(a')$ ,  $b = Fi_{S,X}(b')$  for some distinct elements  $a', b' \in SX$ . Since the set  $ES$  is a singleton (containing the constant function  $S \rightarrow \{0\}$ ), the map  $e_S : S \rightarrow ES$  is bijective and so is the map  $Fe_X : FS \rightarrow FES$ . Let  $\bar{r} : EX \rightarrow ES$  be the unique (constant) map. The equality  $\bar{r} \circ e_X \circ i_{S,X} = e_S$  implies  $F\bar{r} \circ Fe_X \circ Fi_{S,X} = Fe_S$ . Then

$$F\bar{r} \circ Fe_X(a) = F\bar{r} \circ Fe_X \circ Fi_{S,X}(a') = Fe_S(a') \neq Fe_S(b') = F\bar{r} \circ Fe_X \circ Fi_{S,X}(b') = F\bar{r} \circ Fe_X(b)$$

and hence  $Fe_X(a) \neq Fe_X(b)$ . It follows from  $Fe_X(a), Fe_X(b) \in F[e_X[S]; EX]$  that  $\text{supp}(Fe_X(a)) \cup \text{supp}(Fe_X(b)) \subseteq e_X[S]$ . Applying Theorem 8.1, we conclude that

$$\check{d}_{FX}^p(a, b) = d_{FEX}^p(Fe_X(a), Fe_X(b)) = \infty.$$

2. The second statement follows from the definition of the distance  $\check{d}_{FX}$  and Theorem 25.1(4).

3. The inequality  $\check{d}_{FX}^p \leq d_{FX}^p$  follows from the definition of the distance  $\check{d}_{FX}$  and Theorem 25.1(5).

Next, we prove that  $d_{FX}^\infty \leq 2n \cdot |Fn| \cdot \check{d}_{FX}^\infty$  where  $n = \max\{1, \deg(F)\}$ . To derive a contradiction, assume that  $d_{FX}^\infty(a, b) > 2n \cdot |Fn| \cdot \check{d}_{FX}^\infty(a, b)$  for some elements  $a, b \in FX$ . In this case,  $2n \cdot |Fn| < \infty$ . Since  $d_{FEX}^\infty(Fe_X(a), Fe_X(b)) = \check{d}_{FX}^\infty(a, b) < \frac{1}{2n \cdot |Fn|} d_{FX}^\infty(a, b)$ , we can apply Lemma 9.3 and find a function  $r : EX \rightarrow EX$  such that  $Fr \circ Fe_X(a) = F \circ Fe_X(b)$ ,  $r[e_X[X]] \subseteq e_X[X]$ , and  $\sup_{y \in EX} d_{EX}(y, r(y)) < \frac{1}{2} d^\infty(a, b)$ .

Let  $s : X \rightarrow X$  be any function such that  $s(y) \in e_X^{-1}(y)$  for every  $y \in r[EX] \cap e_X[X]$ . Consider the function  $h = s \circ r \circ e_X : X \rightarrow X$  and observe that for every  $x \in X$  we have

$$d_X(x, h(x)) = d_{EX}(e_X(x), e_X \circ s \circ r \circ e_X(x)) = d_{EX}(e_X(x), r \circ e_X(x)) < \frac{1}{2} d^\infty(a, b).$$

Using Proposition 6.1, find a function  $f : n \rightarrow X$  and an element  $a' \in Fn$  such that  $a = Ff(a')$ . By the definition of the distance  $d_{FX}^\infty$ , we have

$$d_{FX}^\infty(a, Fh(a)) = d_{FX}^\infty(Ff(a'), F(h \circ f)(a')) \leq d_{X^n}^\infty(f, h \circ f) < \frac{1}{2} d_{FX}^\infty(a, b).$$

By analogy, we can prove that  $d_{FX}^\infty(b, Fh(b)) < \frac{1}{2} d_{FX}^\infty(a, b)$ .

Observe that  $Fh(a) = Fs \circ Fr \circ Fe_X(a) = Fs \circ Fr \circ Fe_X(b) = Fh(b)$ . Finally,

$$d_{FX}^\infty(a, b) \leq d_{FX}^\infty(a, Fh(a)) + d_{FX}^\infty(Fh(b), b) < \frac{1}{2} d_{FX}^\infty(a, b) + \frac{1}{2} d_{FX}^\infty(a, b) = d_{FX}^\infty(a, b),$$

which is a contradiction that completes the proof of the inequality  $d_{FX}^\infty \leq 2n \cdot |Fn| \cdot \check{d}_{FX}^\infty$ .

Now assume that the distance space  $(X, d_X)$  is injective. By Proposition 5.15, there exists a non-expanding map  $r : EX \rightarrow X$  such that  $r \circ e_X$  is the identity map of  $X$ . By Theorem 25.1(5), the maps  $Fr : FEX \rightarrow FX$  and  $Fe_X : FX \rightarrow FEX$  are non-expanding. Since

$Fr \circ Fe_X$  is the identity map of  $FX$ , the non-expanding map  $Fe_X : FX \rightarrow FEX$  is an isometry and hence  $d_{FX}^p(a, b) = d_{FEX}^p(Fe_X(a), Fe_X(b)) = d_{FX}^p(a, b)$  for any  $a, b \in FX$ .

4. Let  $f : X \rightarrow Y$  be a Lipschitz map between distance spaces. By Proposition 5.17, there exists a Lipschitz map  $\bar{f} : EX \rightarrow EY$  such that  $\bar{f} \circ e_X = e_Y \circ f$  and  $\text{Lip}(\bar{f}) \leq \text{Lip}(f)$ . By Theorem 25.1(5), the map  $F^p \bar{f} = F\bar{f} : FEX \rightarrow FEY$  is Lipschitz with Lipschitz constant  $\text{Lip}(F^p \bar{f}) \leq \text{Lip}(\bar{f}) \leq \text{Lip}(f)$ . Then for any  $a, b \in FX$  we have

$$\begin{aligned} d_{F^p Y}^p(Ff(a), Ff(b)) &= d_{FEY}^p(Fe_Y \circ Ff(a), Fe_Y \circ Ff(b)) = \\ &= d_{FEY}^p(F\bar{f} \circ Fe_X(a), F\bar{f} \circ Fe_X(b)) \leq d_{FEX}^p(Fe_X(a), Fe_X(b)) = d_{FX}^p(a, b), \end{aligned}$$

which means that the map  $\tilde{F}^p f = Ff : \tilde{F}^p X \rightarrow \tilde{F}^p Y$  is Lipschitz with Lipschitz constant  $\text{Lip}(\tilde{F}^p) \leq \text{Lip}(f)$ .

5. Let  $f : X \rightarrow Y$  be an isometry between distance spaces. We need to show that the map  $\tilde{F}^p f$  is an isometry. By Proposition 5.14, there exists an isometry  $\bar{f} : EX \rightarrow EY$  such that  $\bar{f} \circ e_X = e_Y \circ f$  and hence  $F\bar{f} \circ Fe_X = Fe_Y \circ Ff$ . By Lemma 5.9, there exists a non-expanding map  $g : EY \rightarrow EX$  such that  $g \circ e_Y \circ f = e_X$ . Observe that  $g \circ \bar{f} \circ e_X = g \circ e_Y \circ f = e_X$  and by Lemma 5.10,  $g \circ \bar{f}$  is the identity map of  $EX$ . Then  $Fg \circ F\bar{f}$  is the identity map of  $FEX$ . By Theorem 25.1(5), the maps  $F^p g$  and  $F^p \bar{f}$  are non-expanding. Since  $F^p g \circ F^p \bar{f} = \text{id}_{FEX}$ , the map  $F^p f$  is an isometry. Now take any elements  $a, b \in FX$  and observe that

$$\begin{aligned} d_{F^p X}^p(a, b) &= d_{FEX}^p(Fe_X(a), Fe_X(b)) = d_{FEY}^p(F\bar{f} \circ Fe_X(a), F\bar{f} \circ Fe_X(b)) = \\ &= d_{FEY}^p(Fe_Y \circ Ff(a), Fe_Y \circ Ff(b)) = d_{F^p Y}^p(Ff(a), Ff(b)), \end{aligned}$$

which means that the map  $\tilde{F}^p f : \tilde{F}^p X \rightarrow \tilde{F}^p Y$  is an isometry.

6. Assume that  $F1$  is a singleton and  $d_X$  is a pseudometric. By Theorem 15.2, the distance  $d_{FX}^p$  is a pseudometric. By Theorem 26.1(3),  $\check{d}_{FX}^p \leq d_{FX}^p$  and hence  $\check{d}_{FX}^p$  is a pseudometric.

7. Fix any distinct elements  $a, b \in FX$ . The inequality  $\check{d}_{FX}^1(a, b) \geq \underline{d}_X(\text{supp}(a) \cup \text{supp}(b))$  is trivial if  $\underline{d}_X(\text{supp}(a) \cup \text{supp}(b)) = 0$ . So, we assume that  $\underline{d}_X(\text{supp}(a) \cup \text{supp}(b)) > 0$ , which implies that the set  $S = \text{supp}(a) \cup \text{supp}(b)$  has cardinality  $|S| \geq 2$  and the restriction  $d_X|_{S \times S}$  is an  $\infty$ -metric. Taking into account that  $e_X : X \rightarrow EX$  is an isometry, we conclude that the map  $e_X|_S$  is injective and  $\underline{d}_{EX}(e_X[S]) = \underline{d}_X(S) > 0$ . Since  $|e_X[X]| \geq |e_X[S]| = |S| \geq 2$ , we can apply Corollary 6.2, and conclude that  $\text{supp}(Fe_X(a)) \subseteq e_X[\text{supp}(a)]$  and  $\text{supp}(Fe_X(b)) \subseteq e_X[\text{supp}(b)]$ . Since  $e_X$  is an isometry,

$$\begin{aligned} \underline{d}_{EX}(\text{supp}(Fe_X(a)) \cup \text{supp}(Fe_X(b))) &\geq \\ &= \underline{d}_{EX}(e_X[\text{supp}(a)] \cup e_X[\text{supp}(b)]) = \underline{d}_X(\text{supp}(a) \cup \text{supp}(b)). \end{aligned}$$

Now Theorem 25.1(10) and the definition of the distances  $\check{d}_{FX}^p$  and  $\check{d}_{FX}^1$  imply

$$\begin{aligned} \check{d}_{FX}^1(a, b) &= d_{FEX}^1(Fe_X(a), Fe_X(b)) \geq \\ &\geq \underline{d}_{EX}(\text{supp}(Fe_X(a)) \cup \text{supp}(Fe_X(b))) \geq \underline{d}_X(\text{supp}(a) \cup \text{supp}(b)). \end{aligned}$$

8. Assume that the functor  $F$  preserves supports and take any distinct points  $a, b \in FX$ . By Theorem 25.1(11),

$$\begin{aligned} \check{d}_{FX}^\infty(a, b) &= d_{FEX}^\infty(Fe_X(a), Fe_X(b)) \geq d_{HEX}(\text{supp}(Fe_X(a)), \text{supp}(Fe_X(b))) = \\ &= d_{HEX}(e_X[\text{supp}(a)], e_X[\text{supp}(b)]) = d_{HX}(\text{supp}(a), \text{supp}(b)). \end{aligned}$$

Next, we prove that  $\check{d}_{FX}^\infty(a, b) \geq \frac{1}{3}\underline{d}_X(S)$  where  $S = \text{supp}(s) \cup \text{supp}(b)$ . This inequality is trivially true if  $\underline{d}_X(S) = 0$ . So, we assume that  $\underline{d}_X(S) > 0$ . In this case the restriction  $d_X \upharpoonright S \times S$  is an  $\infty$ -metric, the isometry  $e_X \upharpoonright S$  is injective, and  $\underline{d}_{EX}(e_X[S]) = \underline{d}_X(S)$ . Applying Theorem 25.1(11) and taking into account that  $F$  preserves supports, we conclude that

$$\begin{aligned} \check{d}_{FX}^\infty(a, b) &= d_{FEX}^\infty(Fe_X(a), Fe_X(b)) \geq \frac{1}{3}\underline{d}_{EX}(\text{supp}(Fe_X(a)) \cup \text{supp}(Fe_X(b))) = \\ &= \frac{1}{3}\underline{d}_{EX}(e_X[\text{supp}(a)] \cup e_X[\text{supp}(b)]) = \frac{1}{3}\underline{d}_{EX}(e_X[S]) = \frac{1}{3}\underline{d}_X(S). \end{aligned}$$

The inequality  $\check{d}_{FX}^p \geq \check{d}_{FX}^\infty$  has been proved in Theorem 26.1(2).

9. Assume that the distances  $d_X$  and  $d_{FEX}^p$  are  $\infty$ -metrics. We should show that  $\check{d}_{FX}^p(a, b) > 0$  for any distinct elements  $a, b \in FX$ . Since  $d_X$  is an  $\infty$ -metric, the isometry  $e_X : X \rightarrow EX$  is injective. Then we can choose a function  $\gamma : EX \rightarrow X$  such that  $\gamma \circ e_X$  is the identity map of  $X$  and conclude that  $F\gamma \circ Fe_X$  is the identity map of  $FX$ , which implies that the map  $Fe_X$  is injective and hence  $\check{d}_{FX}^p(a, b) = d_{FEX}^p(Fe_X(a), Fe_X(b)) > 0$ .

10. The statement (10) follows from the statement (9) and Theorem 25.1(13).

11. The statement (11) follows from the statements (6) and (10) of this theorem.

12. Assume that for some  $a \in F1$  the map  $\xi_2^a : 2 \rightarrow F2$  is injective. By Theorems 10.1 and 26.1(3), the map  $\xi_X^a : (X, d) \rightarrow (FX, \check{d}_{FX}^p)$  is not expanding.

Now assume that  $p = 1$  or  $F$  preserves supports. In this case we will prove that the map  $\xi_X^a : X \rightarrow \check{F}^p X$  is an isometry. By Corollary 5.16, the distance space  $EX$  is injective and by Theorem 26.1(3),  $\check{d}_{FEX}^p = d_{FEX}^p$ . By Theorem 18.4, the map  $\xi_{EX}^a : EX \rightarrow F^p EX = \check{F}^p EX$  is an isometry. By Theorem 26.1(4), the map  $Fe_X : \check{F}^p X \rightarrow \check{F}^p EX = F^p EX$  is non-expanding. By Proposition 18.1, the following diagram commutes.

$$\begin{array}{ccc} X & \xrightarrow{e_X} & EX \\ \xi_X^a \downarrow & & \downarrow \xi_{EX}^a \\ \check{F}^p X & \xrightarrow{Fe_X} & \check{F}^p EX = F^p EX \end{array}$$

Now the isometry property of the map  $\xi_{EX}^a \circ e_X$  and the non-expanding property of  $Fe_X : \check{F}^p X \rightarrow \check{F}^p EX = F^p EX$  imply that the non-expanding map  $\xi_X^a : X \rightarrow \check{F}^p X$  is an isometry.

13. If  $d_X$  is the  $\{0, \infty\}$ -valued  $\infty$ -metric on  $X$ , then the  $\infty$ -metric of the space  $EX$  is  $\{0, \infty\}$ -valued (by the definition of  $EX$ ). By Theorem 25.1(16), the distance  $d_{FEX}^p$  is an  $\{0, \infty\}$ -valued  $\infty$ -metric. Then the distance  $\check{d}_{FX}^p$  is  $\{0, \infty\}$ -valued and by Theorem 26.1(9),  $\check{d}_{FX}^p$  is an  $\infty$ -metric.  $\square$

**Remark 26.2.** In general  $\check{d}_{FX}^p \neq d_{FX}^p$  even for support-preserving functors of finite degree. Indeed, it can be shown that for the functor  $H_2$ , the metric space  $X \subseteq \mathbb{C}$  and elements  $a, b \in H_2 X$  from Example 13.4 we have

$$\check{d}_{H_2 X}^p(a, b) = 4 + 4\sqrt[p]{2} < 10 + 4\sqrt{2} = d_{H_2}^p(a, b).$$

**Remark 26.3.** If the functor  $F$  is finitary and has finite degree, then by Theorem 26.1(3), the identity map  $F^p X \rightarrow \check{F}^p X$  is a Lipschitz isomorphism. In this case, Theorem 26.1 (with suitable modifications of the statements (1),(5),(8)) hold also for the functor  $\check{F}^p$ .

**Problem 26.4.** *Is there any efficient algorithm for calculating the distance  $\check{d}_{FX}^p$  (with arbitrary precision)?*

27. THE  $\ell^p$ -METRIZATION AND NATURAL TRANSFORMATIONS OF FUNCTORS

In this section we investigate the relation between the  $\ell^p$ -metrizations of functors  $F, G : \mathbf{Set} \rightarrow \mathbf{Set}$ , connected by a natural transformation  $\eta : F \rightarrow G$ . The natural transformation  $\eta$  assigns to each set  $X$  a function  $\eta_X : FX \rightarrow GX$  such that for any function  $f : X \rightarrow Y$  between sets the diagram

$$\begin{array}{ccc} FX & \xrightarrow{\eta_X} & GX \\ Ff \downarrow & & \downarrow Gf \\ FY & \xrightarrow{\eta_Y} & GY \end{array}$$

is commutative.

If for each  $X$  the function  $\eta_X : FX \rightarrow GX$  is injective (resp. bijective), then we say that  $\eta : F \rightarrow G$  is an *embedding* (resp. an *isomorphism*) of the functors  $F$  and  $G$ .

**Proposition 27.1.** *If  $\eta : F \rightarrow G$  is an embedding of functors  $F, G : \mathbf{Set} \rightarrow \mathbf{Set}$  and the functor  $G$  has finite supports, then  $F$  has finite supports, too.*

*Proof.* Given any set  $X$  and element  $a \in FX$ , we should find a finite set  $A \subseteq X$  such that  $a \in F[A; X] := Fi_{A,X}[FA]$ . If  $X$  is finite, then we put  $A = X$ . So, we assume that  $X$  is infinite and hence not empty. Since the functor  $G$  has finite supports, for the element  $b := \eta_X(a) \in \eta_X[FX] \subseteq GX$  there exists a non-empty finite set  $A \subseteq X$  such that  $b = Gi_{A,X}(b')$  for some  $b' \in GA$ . Since  $A$  is not empty, there exists a function  $r : X \rightarrow A$  such that  $r \circ i_{A,X}$  is the identity map of  $A$ . Then  $Fr \circ Fi_{A,X}$  is the identity map of  $FA$  and  $Gr \circ Gi_{A,X}$  is the identity map of  $GA \subseteq GX$ . Consider the element  $a' = Fr(a) \in FA$  and observe that

$$\begin{aligned} \eta_X \circ Fi_{A,X}(a') &= \eta_X \circ Fi_{A,X} \circ Fr(a) = Gi_{A,X} \circ \eta_A \circ Fr(a) = Gi_{A,X} \circ Gr \circ \eta_X(a) = \\ &= Gi_{A,X} \circ Gr(b) = Gi_{A,X} \circ Gr \circ Gi_{A,X}(b') = Gi_{A,X}(b') = b = \eta_X(a). \end{aligned}$$

Now the injectivity of  $\eta_X$  ensures that  $a = Fi_{A,X}(a') \in F[A; X]$ .  $\square$

**Theorem 27.2.** *Let  $F, G : \mathbf{Set} \rightarrow \mathbf{Set}$  be two functors with finite supports and  $\eta : F \rightarrow G$  be a natural transformation. For any distance space  $(X, d_X)$  and any  $p \in [1, \infty]$  the map  $\eta_X : (FX, d_{FX}^p) \rightarrow (GX, d_{GX}^p)$  is non-expanding and so is the map  $\eta_X : (FX, \check{d}_{FX}^p) \rightarrow (GX, \check{d}_{GX}^p)$ .*

*Proof.* Given two elements  $a, b \in FX$ , we should prove that

$$(2) \quad d_{GX}^p(\eta_X(a), \eta_X(b)) \leq d_{FX}^p(a, b).$$

To derive a contradiction, assume that  $d_{FX}^p(a, b) < d_{GX}^p(\eta_X(a), \eta_X(b))$  and find an  $(a, b)$ -linking chain  $w = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  with  $\Sigma d_X^p(w) < d_{GX}^p(\eta_X(a), \eta_X(b))$ . For every  $i \in \{0, \dots, l\}$  put  $n_i := \text{dom}(f_i) = \text{dom}(g_i)$  and  $b_i = \eta_{n_i}(a_i) \in Gn_i$ . Observe that  $\eta_X(a) = \eta_X(Ff_0(a_0)) = Gf_0(\eta_{m_0}(a_0)) = Gf_0(b_0)$ ,  $\eta_X(b) = \eta_X(Fg_l(a_l)) = Gg_l(\eta_{n_l}(a_l)) = Gg_l(b_l)$  and for every  $i \in \{1, \dots, l\}$

$$Gf_i(b_i) = Gf_i(\eta_{n_i}(a_i)) = \eta_X(Ff_i(a_i)) = \eta_X(Fg_{i-1}(a_{i-1})) = Gg_{i-1}(\eta_{n_{i-1}}(a_{i-1})) = Gg_{i-1}(b_{i-1}),$$

which means that  $\tilde{w} = ((b_i, f_i, g_i))_{i=0}^l$  is an  $(\eta_X(a), \eta_X(b))$ -connecting chain and hence

$$d_{GX}^p(\eta_X(a), \eta_X(b)) \leq \Sigma d_X^p(\tilde{w}) = \Sigma d_X^p(w) < d_{GX}^p(\eta_X(a), \eta_X(b)),$$

which is a desired contradiction, completing the proof of the inequality (2).

Next, observe that

$$\begin{aligned} \check{d}_{GX}^p(\eta_X(a), \eta_X(b)) &= d_{GEX}^p(\text{Ge}_X(\eta_X(a)), \text{Ge}_X(\eta_X(b))) = \\ &= d_{GEX}^p(\eta_{EX}(\text{Fe}_X(a)), \eta_{EX}(\text{Fe}_X(b))) \leq d_{FEX}^p(\text{Fe}_X(a), \text{Fe}_X(b)) = \check{d}_{FX}^p(a, b), \end{aligned}$$

which means that the map  $\eta_X : (FX, \check{d}_{FX}^p) \rightarrow (GX, \check{d}_{GX}^p)$  is non-expanding.  $\square$

A functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  is defined to be a *subfunctor* of a functor  $G : \mathbf{Set} \rightarrow \mathbf{Set}$  if for every set  $X$  we have  $FX \subseteq GX$  and for every function  $f : X \rightarrow Y$  between sets,  $Ff = Gf \upharpoonright FX$ . The latter condition means that the identity inclusion  $i_{FX, GX} : FX \rightarrow GX$  is a component of a natural transformation  $i : F \rightarrow G$ , which is a functor embedding.

A subfunctor  $F$  of a functor  $G : \mathbf{Set} \rightarrow \mathbf{Set}$  is defined to be *1-determined* if for any set  $X$  and the constant map  $\gamma : X \rightarrow 1$  we have  $FX = (G\gamma)^{-1}[F1] := \{a \in GX : G\gamma(a) \in F1 \subseteq G1\}$ .

**Theorem 27.3.** *Let  $G : \mathbf{Set} \rightarrow \mathbf{Set}$  be a functor with finite supports and  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  be a subfunctor of  $G$ . For any distance space  $(X, d_X)$ ,  $p \in [1, \infty]$  and any  $a, b \in FX \subseteq GX$  the following statements hold.*

- 1)  $d_{GX}^p(a, b) \leq d_{FX}^p(a, b)$  and  $\check{d}_{GX}^p(a, b) \leq \check{d}_{FX}^p(a, b)$ .
- 2) If  $F$  is 1-determined, then  $d_{GX}^p(a, b) = d_{FGX}^p(a, b)$  and  $\check{d}_{GX}^p(a, b) = \check{d}_{FX}^p(a, b)$ .

*Proof.* 1. The first statement follows from Proposition 27.1 and Theorem 27.2.

2. Now assuming that the subfunctor  $F$  of  $G$  is 1-determined, we will prove that  $d_{GX}^p(a, b) = d_{FX}^p(a, b)$ . By the preceding statement, it suffices to check that  $d_{GX}^p(a, b) \geq d_{FX}^p(a, b)$ . To derive a contradiction, assume that  $d_{GX}^p(a, b) < d_{FX}^p(a, b)$  and find an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{GX}(a, b)$  such that

$$\sum_{i=0}^l d_{X^{\leftarrow}}^p(f_i, g_i) < d_{FX}^p(a, b).$$

The definition of  $L_{GX}(a, b)$  ensures that  $a = Gf_0(a_0)$ ,  $b = Gg_l(a_l)$  and  $Gg_{i-1}(a_{i-1}) = Gf_i(a_i)$  for all  $i \in \{1, \dots, l\}$ . For every  $i \in \{0, \dots, l\}$  let  $n_i := \text{dom}(f_i) = \text{dom}(g_i) \in \omega$ .

Let  $\gamma : X \rightarrow 1$  and  $\gamma_i : n_i \rightarrow 1$  for  $i \in \{0, \dots, l\}$  be the constant maps. Taking into account that  $F$  is a 1-determined subfunctor of  $G$ , we conclude that the equality  $G\gamma_0(a_0) = G\gamma \circ Gf_0(a_0) = G\gamma(a) = F\gamma(a) \in F1$  implies that  $a_0 \in (G\gamma_0)^{-1}[F1] = Fn_0$ . By induction on  $i \in \{0, \dots, k\}$  we will prove that  $a_i \in Fn_i$ . For  $i = 0$  this has already been done. Assume that for some  $i \in \{1, \dots, k\}$  we have proved that  $a_{i-1} \in Fn_{i-1}$ . Then

$G\gamma_i(a_i) = G\gamma \circ Gf_i(a_i) = G\gamma \circ Gg_{i-1}(a_{i-1}) = G\gamma_{i-1}(a_{i-1}) = F\gamma_{i-1}(a_{i-1}) \in F\gamma_{i-1}[Fn_{i-1}] \subseteq F1$  and hence  $a_i \in (G\gamma_i)^{-1}[F1] = Fn_i$ .

After completing the inductive construction, we see that

$$((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b) \subseteq L_{GX}(a, b)$$

and thus

$$d_{FX}^p(a, b) \leq \sum_{i=0}^l d_{X^{\leftarrow}}^p(f_i, g_i) < d_{FX}^p(a, b),$$

which is a desired contradiction showing that  $d_{GX}^p(a, b) = d_{FX}^p(a, b)$ .

Then  $\check{d}_{GX}^p(a, b) = d_{GEX}^p(\text{Ge}_X(a), \text{Ge}_X(b)) = d_{FEX}^p(\text{Fe}_X(a), \text{Fe}_X(b)) = \check{d}_{FX}^p(a, b)$ .  $\square$

28. THE METRIZATIONS OF THE FUNCTOR OF  $n$ -TH POWER

Let  $n \in \omega$  and  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  be the functor assigning to each set  $X$  its  $n$ -th power  $X^n$  and to each map  $f : X \rightarrow Y$  the map  $Ff : FX \rightarrow FY$ ,  $Ff : \alpha \mapsto f \circ \alpha$ . It is clear that  $F$  is a functor of finite degree  $\leq n$ ,  $F$  preserves supports and singletons.

**Theorem 28.1.** *For every distance  $d_X$  on a set  $X$  we have the inequalities*

$$d_{X^n}^\infty \leq \check{d}_{FX}^p \leq d_{FX}^p \leq d_{X^n}^p \leq n^{\frac{1}{p}} \cdot d_{X^n}^\infty.$$

*In particular,  $\check{d}_{FX}^\infty = d_{FX}^\infty = d_{X^n}^\infty$ . If  $d_X[X \times X] \neq \{0\}$ ,  $p < \infty$  and  $n \geq 2$ , then  $d_{FX}^p \neq d_{X^n}^p$ .*

*Proof.* By Theorem 10.2, the pseudometric  $d_{FX}^p$  coincides with the largest pseudometric on  $FX$  such that for every  $a \in Fn = n^n$  the map  $\xi_X^a : X^n \rightarrow FX$ ,  $\xi_X^a : f \mapsto f \circ a$ , is non-expanding with respect to the  $\ell^p$ -metric  $d_{X^n}^p$  on  $X^n$ . The identity map  $a : n \rightarrow n$  induces the identity non-expanding map  $\xi_X^a : X^n \rightarrow FX$ , witnessing that  $d_{FX}^p \leq d_{X^n}^p$ . The inequality  $\check{d}_{FX}^p \leq d_{FX}^p$  follows from Theorem 26.1(3). The inequality  $d_{X^n}^p \leq n^{\frac{1}{p}} \cdot d_{X^n}^\infty$  follows from Lemma 11.1.

To see that  $d_{X^n}^\infty \leq d_{FX}^p$ , observe that for every  $a \in Fn = n^n$  and functions  $f, g \in Fn$ , we get the inequalities  $d_{X^n}^\infty(\xi_X^a(f), \xi_X^a(g)) = d_{X^n}^\infty(f \circ a, g \circ a) \leq d_{X^n}^\infty(f, g) \leq d_{X^n}^p(f, g)$ , witnessing that the map  $\xi_X^a : (X^n, d_{X^n}^p) \rightarrow (FX, d_{FX}^p)$  is non-expanding. By the maximality of  $d_{FX}^p$ , we have  $d_{X^n}^\infty \leq d_{FX}^p$ . Therefore, we obtain the chain of inequalities  $d_{X^n}^\infty \leq d_{FX}^p \leq d_{X^n}^p \leq n^{\frac{1}{p}} \cdot d_{X^n}^\infty$  which transforms into a chain of equalities if  $p = \infty$ .

Now we prove that  $d_{X^n}^\infty(a, b) \leq \check{d}_{FX}^p(a, b)$  for any  $a, b \in FX = X^n$ . Indeed,

$$\begin{aligned} \check{d}_{FX}^p(a, b) &= d_{FEX}^p(Fe_X(a), Fe_X(b)) \geq d_{FEX}^\infty(Fe_X(a), Fe_X(b)) = d_{(EX)^n}^\infty(e_X \circ a, e_X \circ b) = \\ &= \max_{i \in n} d_{EX}(e_X \circ a(i), e_X \circ b(i)) = \max_{i \in n} d_X(a(i), b(i)) = d_{X^n}^\infty(a, b). \end{aligned}$$

Now assume that  $d_X[X \times X] \neq \{0\}$ ,  $p < \infty$  and  $n \geq 2$ . Since  $d_X[X \times X] \neq \{0\}$ , there exists functions  $f, g \in X^n$  such that

$$d_X(f(0), g(0)) > 0 = d_X(f(1), g(1)) = \dots = d_X(f(n-1), g(n-1)).$$

Let  $a : n \rightarrow \{0\} \subseteq n$  be the constant map. Since the map  $\xi_X^a : (X^n, d_{X^n}^p) \rightarrow (FX, d_{FX}^p)$ ,  $\xi_X^a : \varphi \mapsto \varphi \circ a$ , is non-expanding,

$$d_{FX}^p(f \circ a, g \circ a) \leq d_{X^n}^p(f, g) = d_X(f(0), g(0)) < n^{\frac{1}{p}} \cdot d_X(f(0), g(0)) = d_{X^n}^p(f \circ a, g \circ a),$$

which implies that  $d_{FX}^p \neq d_{X^n}^p$ .  $\square$

For the functor  $F$  of  $n$ -th power we can repeat a special instance of Problems 10.3 and 26.4.

**Problem 28.2.** *Is there any efficient algorithm for calculating the distances  $d_{FX}^p$  and  $\check{d}_{FX}^p$  for the functor  $F$  of the  $n$ -th power?*

29. METRIZATIONS OF THE FUNCTOR OF  $M$ -VALUED FINITARY FUNCTIONS

Let  $(M, +)$  be a commutative monoid whose neutral element is denoted by 0. By  $M_0$  we denote the subgroup of invertible elements of  $M$ .

For every set  $X$  consider the set  $F(X, M)$  of all functions  $\varphi : X \rightarrow M$  with finite support  $\text{supp}(\varphi) := \{x \in X : \varphi(x) \neq 0\}$ . The set  $F(X, M)$  carries an obvious structure of a commutative monoid, whose neutral element is the constant function  $o : X \rightarrow \{0\} \subseteq M$  with empty support  $\text{supp}(o) = \emptyset$ .

For every  $x \in X$  and  $g \in M$  let  $\delta_x^g : X \rightarrow M$  be the function with  $\text{supp}(\delta_x^g) \subseteq \{x\}$  and  $\delta_x^g(x) = g$ . It is easy to see that each function  $\varphi \in F(X, M)$  can be written as the sum

$$\varphi = \sum_{i=1}^k \delta_{x_i}^{g_i}$$

for some points  $x_1, \dots, x_k \in X$  and elements  $g_1, \dots, g_k \in G$ .

For a subset  $A \subseteq X$  and a function  $\varphi \in F(X, M)$  by  $\sum_{x \in A} \varphi(x)$  we understand the finite sum  $\sum_{x \in A \cap \text{supp}(\varphi)} \varphi(x)$  in the commutative monoid  $M$ .

The subset  $F_0(X, M) = \{\varphi \in F(X, M) : \sum_{x \in X} \varphi(x) = 0\}$  is a subgroup in  $F(X, M)$  that coincides with the subgroup  $F_0(X, M_0)$ .

If  $M$  is a group, then the commutative monoids  $F_0(X, M) \subseteq F(X, M)$  are groups. Moreover, if  $M$  is the group  $\mathbb{Z}$  of integers, then  $F(X, M)$  can be identified with the free Abelian group over  $X$ . If  $M$  is a two-element group, then  $F(X, M)$  is a free Boolean group over  $X$ .

For any function  $f : X \rightarrow Y$  between sets let  $Ff : F(X, M) \rightarrow F(Y, M)$  be the function assigning to each function  $\varphi \in F(X, M)$  the (finitely supported) function  $F\varphi : Y \rightarrow M$ ,  $F\varphi : y \mapsto \sum_{x \in f^{-1}(y)} \varphi(x)$ . It follows that  $Ff(F_0(X, M)) \subseteq F_0(Y, M)$ .

This construction defines a functor  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  assigning to each set  $X$  the set  $FX := F(X, M)$  and to each function  $f : X \rightarrow Y$  between sets the function  $Ff : FX \rightarrow FY$ . It can be shown that  $Ff$  is a homomorphism between the commutative monoids  $FX$  and  $FY$ , so actually  $F$  is a functor from the category  $\mathbf{Set}$  to the category  $\mathbf{Mon}$  of commutative monoids and their homomorphisms. The functor  $F$  is called *the functor of  $M$ -valued finitary functions*.

Moreover,  $F_0$  is a 1-determined subfunctor of the functor  $F$  (for the definition of 1-determined subfunctors see Section 27).

Now let us consider the  $\ell^p$ - and  $\tilde{\ell}^p$ -metrizations of the functor  $F$ . Fix  $p \in [1, \infty]$  and a distance space  $(X, d_X)$ . The distance  $d_X$  induces the distances  $d_{FX}^p$  and  $\tilde{d}_{FX}^p$  on the set  $FX$ . These distances have all the properties described in Theorems 25.1 and 26.1. Moreover, these distances have nice algebraic properties.

**Proposition 29.1.** *The distance  $d_{FX}^p$  is subinvariant in the sense that  $d_{FX}^p(a + c, b + c) \leq d_{FX}^p(a, b)$  for any elements  $a, b, c \in FX$  of the commutative monoid  $FX$ .*

*Proof.* To derive a contradiction, assume that  $d_{FX}^p(a, b) < d_{FX}^p(a+c, b+c)$  and find an  $(a, b)$ -linking chain  $w = ((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\Sigma d_X^p(w) < d_{FX}^p(a+c, b+c)$ . Let  $m = |\text{supp}(c)| \in \omega$ .

For every  $i \in \{0, \dots, l\}$  let  $n_i = \text{dom}(f_i) = \text{dom}(g_i)$  and  $m_i := n_i + m$ . Let  $\gamma_i : m_i \setminus n_i \rightarrow \text{supp}(c)$  be any bijection. Let  $\tilde{a}_i : m_i \rightarrow M$  be the function such that  $\tilde{a}_i(x) = a_i(x)$  for  $x \in n_i$  and  $\tilde{a}_i(x) = c(\gamma_i(x))$  for  $x \in m_i \setminus n_i$ . Consider the functions  $\tilde{f}_i, \tilde{g}_i : m_i \rightarrow X$  defined by  $\tilde{f}_i|_{n_i} = f_i$ ,  $\tilde{g}_i|_{n_i} = g_i$  and  $\tilde{f}_i|_{m_i \setminus n_i} = \gamma = \tilde{g}_i|_{m_i \setminus n_i}$ . It can be shown that  $((\tilde{a}_i, \tilde{f}_i, \tilde{g}_i))_{i=0}^l \in L_{FX}(a+c, b+c)$  and

$$d_{FX}^p(a+c, b+c) \leq \sum_{i=0}^l d_{X^{\leftarrow \omega}}^p(\tilde{f}_i, \tilde{g}_i) = \sum_{i=0}^l d_{X^{\leftarrow \omega}}^p(f_i, g_i) < d_{FX}^p(a+c, b+c),$$

which is a desired contradiction, completing the proof.  $\square$

**Corollary 29.2.** *The distance  $\check{d}_{FX}^p$  is subinvariant in the sense that*

$$\check{d}_{FX}^p(a + c, b + c) \leq \check{d}_{FX}^p(a, b)$$

for any elements  $a, b, c \in FX$  of the commutative monoid  $FX$ .

*Proof.* By Proposition 29.1,

$$\begin{aligned} \check{d}_{FX}^p(a + c, b + c) &= d_{FEX}^p(\mathbf{Fe}_X(a + c), \mathbf{Fe}_X(b + c)) = \\ &= d_{FEX}^p(\mathbf{Fe}_X(a) + \mathbf{Fe}_X(c), \mathbf{Fe}_X(b) + \mathbf{Fe}_X(c)) \leq d_{FEX}^p(\mathbf{Fe}_X(a), \mathbf{Fe}_X(b)) = \check{d}_{FX}^p(a, b). \end{aligned}$$

$\square$

Since  $F_0$  is a 1-determined subfunctor of  $F$ , we can apply Theorem 27.3, Proposition 29.1 and Corollary 29.2 to deduce the following corollary.

**Corollary 29.3.**  *$d_{F_0X}^p = d_{FX}^p \upharpoonright F_0X \times F_0X$  and  $\check{d}_{F_0X}^p = \check{d}_{FX}^p \upharpoonright F_0X \times F_0X$ . Moreover, the distances  $d_{F_0X}^p$  and  $\check{d}_{F_0X}^p$  are invariant in the sense that*

$$d_{F_0X}^p(a + c, b + c) = d_{F_0X}^p(a, b) \quad \text{and} \quad \check{d}_{F_0X}^p(a + c, b + c) = \check{d}_{F_0X}^p(a, b)$$

for any elements  $a, b, c \in F_0X$  of the commutative group  $F_0X$ .

Let  $\sim$  be the equivalence relation on  $X$  defined by  $x \sim y$  iff  $d_X(x, y) < \infty$ . For each point  $x \in X$  its equivalence class coincides with the ball  $O(x; \infty) = \{y \in X : d_X(x, y) < \infty\}$ . Let  $X/\sim = \{O(x, \infty) : x \in X\}$  be the quotient set of all such  $\infty$ -balls.

In the group  $F_0X$  consider the subgroup  $F_{00}X$  consisting of finitely supported functions  $\varphi : X \rightarrow M_0$  such that  $\sum_{x \in E} \varphi(x) = 0$  for any equivalence class  $E \in X/\sim$ .

Now we are going to show that the distance  $\check{d}_{F_0X}^p$  is trivial on the subgroup  $F_{00}X$  whenever  $p > 1$  and the distance space is *Lipschitz-geodesic* in the sense that for any points  $x, y \in X$  with  $d_X(x, y) < \infty$  there exists a Lipschitz map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ . Here  $[0, 1] \subseteq \mathbb{R}$  is the closed unit interval on the real line with the standard Euclidean distance. It is easy to see that each injective metric space is Lipschitz-geodesic.

**Theorem 29.4.** *If  $p > 1$  and  $(X, d_X)$  is a Lipschitz-geodesic distance space, then  $d_{F_0X}^p(a, b) = 0$  for any elements  $a, b \in F_{00}X \subseteq F_0X$ .*

*Proof.* Observe that the invariant distance  $d_{F_0X}^p \upharpoonright F_0X \times F_0X$  on  $F_0X$  is uniquely determined by the norm  $\|\cdot\| : F_0X \rightarrow [0, \infty]$  assigning to each function  $\varphi \in F_0X$  the distance  $d_{F_0X}^p(\varphi, o)$  from  $\varphi$  to the zero function  $o : X \rightarrow \{0\} \subseteq G$ , which is the neutral element of the group  $F_0X$ .

The equality  $d_{F_0X}^p(a, b) = 0$  for any  $a, b \in F_{00}X$  will follow as soon as we prove that  $\|\varphi\| = 0$  for any  $\varphi \in F_{00}X$ .

Choose any function  $r : X \rightarrow X$  such that for each  $\infty$ -ball  $E \in X/\sim$  its image  $r(E)$  is a singleton in  $E$ . For every  $E \in X/\sim$  we have  $\sum_{x \in E} \varphi(x) = 0$  (by the definition of the group  $F_{00}M$ ). This implies that  $\varphi = \sum_{x \in \text{supp}(\varphi)} (\delta_x^{\varphi(x)} - \delta_{r(x)}^{\varphi(x)})$ . The triangle inequality will imply that  $\|\varphi\| = 0$  as soon as we prove that  $\|\delta_x^g - \delta_{r(x)}^g\| = 0$  for every  $x \in X$  and  $g \in M_0$ .

Since  $d_X(x, r(x)) < \infty$ , we can apply the Lipschitz-geodesic property of  $(X, d_X)$  and find a Lipschitz map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = r(x)$ . Now given any  $\varepsilon > 0$ , we will prove that  $\|\delta_x^g - \delta_{r(x)}^g\| < \varepsilon$ .



Find an even number  $n$  so large that  $\text{Lip}(\gamma) \cdot (n-1)^{\frac{1-p}{p}} < \varepsilon$  (such number  $n$  exists as  $p > 1$ ). Next, consider the function  $a_0 : n \rightarrow \{-g, g\} \subseteq M_0 \subseteq M$  such that  $a_0(i) = (-1)^i \cdot g$  for  $i \in n$ . Now consider the function  $f_0 : n \rightarrow X$  such that  $f_0(i) = f_0(i+1) = \gamma(\frac{i}{n-1})$  for every even number  $i \in n$ . Observe that  $Ff_0(a_0) = o$ . Next, let  $g_0 : n \rightarrow X$  be the function such that  $g_0(0) = x$ ,  $g_0(n-1) = r(x)$  and  $g_0(i-1) = g_0(i) = \gamma(\frac{i-1}{n-1})$  for every non-zero even number  $i \in n$ . It is easy to see that  $Fg_0(a_0) = \delta_x^g - \delta_{r(x)}^g$ . Then  $((a_0, f_0, g_0))$  is a  $(o, \delta_x^g - \delta_{r(x)}^g)$ -linking chain and hence

$$\begin{aligned} \|\delta_x^g - \delta_{r(x)}^g\| &= d_{FX}^p(o, \delta_x^g - \delta_{r(x)}^g) \leq d_{X^{\leftarrow \omega}}^p(f_0, g_0) = \left( \sum_{i \in n} d_X(f_0(i), g_0(i))^p \right)^{\frac{1}{p}} \leq \\ &\leq \left( 0 + (n-2) \left( \text{Lip}(\gamma) \frac{1}{n-1} \right)^p + 0 \right)^{\frac{1}{p}} < \text{Lip}(\gamma) \cdot (n-1)^{\frac{1-p}{p}} < \varepsilon, \end{aligned}$$

witnessing that  $\|\delta_x^g - \delta_{r(x)}^g\| = 0$ .  $\square$

**Corollary 29.5.** *If  $p > 1$  and the distance space  $(X, d_X)$  is Lipschitz-geodesic, then for any  $a, b \in F_0X$*

$$d_{FX}^p(a, b) = \begin{cases} 0 & \text{if } a - b \in F_{00}X; \\ \infty & \text{otherwise.} \end{cases}$$

*Proof.* Let  $q : X \rightarrow Y$ ,  $q : x \mapsto O(x, \infty)$  be the quotient map to the quotient space  $Y = X/\sim = \{O(x, \infty), x \in X\}$  of all  $\infty$ -balls in the distance space  $(X, d_X)$ .

Take any elements  $a, b$  of the group  $F_0X$ . If  $a - b \in F_{00}X$ , then by Corollary 29.3 and Theorem 29.4,

$$d_{FX}^p(a, b) = d_{FX}^p(a - b, o) = 0.$$

If  $a - b \notin F_{00}X$ , then  $\sum_{x \in E} (a(x) - b(x)) \neq 0$  for some  $\infty$ -ball  $E \in X/\sim$ , which implies that  $\sum_{x \in E} a(x) \neq \sum_{x \in E} b(x)$  and hence  $Fq(a) \neq Fq(b)$ . By Theorem 15.2,  $d_{FX}^p(a, b) = \infty$ .  $\square$

**Theorem 29.6.** *If  $p > 1$ , then for any  $a, b \in F_0X$  we have*

$$\check{d}_{FX}^p(a, b) = \begin{cases} 0 & \text{if } a - b \in F_{00}X; \\ \infty & \text{otherwise.} \end{cases}$$

*Proof.* By Corollary 5.16, the distance space  $EX$  is injective and hence Lipschitz-geodesic. Applying Corollary 29.5 to the Lipschitz-geodesic distance space  $EX$ , we conclude that

$$\check{d}_{FX}^p(a, b) = d_{FEX}^p(Fe_X(a), Fe_X(b)) = \begin{cases} 0 & \text{iff } Fe_X(a) - Fe_X(b) \in F_{00}EX \text{ iff } a - b \in F_{00}X; \\ \infty & \text{iff } Fe_X(a) - Fe_X(b) \notin F_{00}EX \text{ iff } a - b \notin F_{00}X. \end{cases}$$

$\square$

Theorem 29.6 and Corollary 29.5) yield simple formulas for calculating the distances  $\check{d}_{F_0X}^p$  and  $d_{F_0X}^p$  if  $p > 1$ . Now we present a simple formula for calculating the distance  $d_{F_0X}^1$ .

For an element  $\varphi \in FX$  let  $R_X(\varphi)$  be the set of all sequences

$$s = ((g_0, x_0, y_0), \dots, (g_l, x_l, y_l)) \in (M_0 \times X \times X)^{<\omega}$$

such that  $\varphi := \sum_{i=0}^l (\delta_{x_i}^{g_i} - \delta_{y_i}^{g_i})$  is equal to  $\varphi$ . Sequences  $s$  that belong to the set  $R_X(\varphi)$  will be called  $\varphi$ -representing sequences. For each sequence  $s = ((g_0, x_0, y_0), \dots, (g_l, x_l, y_l)) \in R_X(\varphi)$  put  $\Sigma d_X^1(s) := \sum_{i=0}^l d_X(x_i, y_i)$ .

In the family  $R_X(\varphi)$  consider two subfamilies:

$$R_X^{\subseteq}(\varphi) := \left\{ ((g_i, x_i, y_i))_{i=0}^l \in R_X(\varphi) : \bigcup_{i=0}^l \{x_i, y_i\} \subseteq \text{supp}(\varphi) \right\}$$

and

$$R_X^{\cap}(\varphi) := \left\{ ((g_i, x_i, y_i))_{i=0}^l \in R_X(\varphi) : \{x_i, y_i\} \cap \{x_j, y_j\} = \emptyset \text{ for any distinct } i, j \right\}.$$

It is easy to see that

$$R_X^{\cap}(\varphi) \subseteq R_X^{\subseteq}(\varphi) \subseteq R_X(\varphi)$$

and the families  $R_X^{\subseteq}(\varphi) \subseteq R_X(\varphi)$  are not empty if and only if  $\varphi \in F_0X$  (the latter fact can be proved by induction on  $|\text{supp}(\varphi)|$ ).

It will be convenient to assume that the zero function  $o \in F_0X$  is represented by the empty sequence  $() \in R_X^{\cap}(\varphi) \subseteq R_X^{\subseteq}(\varphi) \subseteq R_X(\varphi)$ .

For any  $\varphi \in F_0X$  consider the (finite or infinite) numbers

$$\|\varphi\|_{FX} := \inf(\{\Sigma d_X^1(s) : s \in R_X(\varphi)\} \cup \{\infty\}),$$

$$\|\varphi\|_{FX}^{\subseteq} := \inf(\{\Sigma d_X^1(s) : s \in R_X^{\subseteq}(\varphi)\} \cup \{\infty\}),$$

$$\|\varphi\|_{FX}^{\cap} := \inf(\{\Sigma d_X^1(s) : s \in R_X^{\cap}(\varphi)\} \cup \{\infty\}).$$

The inclusions  $R_X^{\cap}(\varphi) \subseteq R_X^{\subseteq}(\varphi) \subseteq R_X(\varphi)$  imply the inequalities

$$\|\varphi\| \leq \|\varphi\|^{\subseteq} \leq \|\varphi\|^{\cap}.$$

Also for any  $a, b \in FX$ , in the family  $L_{FX}(a, b)$  of  $(a, b)$ -connecting chains, consider the subfamily  $L_{FX}^{\subseteq}(a, b)$  consisting of  $(a, b)$ -connecting chains  $s = ((a_i, f_0, g_i))_{i=0}^k$  such that  $\bigcup_{i=0}^l f_i[\text{dom}(f_i)] \cup g_i[\text{dom}(g_i)] \subseteq \text{supp}(a) \cup \text{supp}(b)$ . For such chain  $s$  we put

$$\Sigma d_X^1(s) = \sum_{i=0}^l d_{X^{\leftarrow \omega}}^1(f_i, g_i) = \sum_{i=0}^l \sum_{x \in \text{dom}(f_i)} d_X(f_i(x), g_i(x)).$$

For every  $a, b \in FX$  consider the number

$$d_{FX}^{\subseteq}(a, b) = \inf(\{\Sigma d_X^1(s) : s \in L_{FX}^{\subseteq}(a, b)\} \cup \{\infty\})$$

and observe that  $d_{FX}^1(a, b) \leq d_{FX}^{\subseteq}(a, b)$  for any  $a, b \in FX$ .

**Theorem 29.7.** *For any elements  $a, b \in F_0X$*

- 1)  $d_{FX}^1(a, b) = \|a - b\|_{FX} \leq d_{FX}^{\subseteq}(a, b) \leq \|a - b\|_{FX}^{\subseteq}$ ;
- 2) if  $|M_0| = 2$ , then

$$\tilde{d}_{FX}^1(a, b) = d_{FX}^1(a, b) = d_{FX}^{\subseteq}(a, b) = \|a - b\|_{FX} = \|a - b\|_{FX}^{\subseteq} = \|a - b\|_{FX}^{\cap}.$$

*Proof.* 1. First we show that  $d_{FX}^1(a, b) \leq \|a - b\|_{FX}$  and  $d^{\subseteq}(a, b) \leq \|a - b\|_{FX}^{\subseteq}$ . To derive a contradiction, assume that  $\|a - b\|_{FX} < d_{FX}^1(a, b)$  and find an  $(a - b)$ -representing sequence  $((g_i, x_i, y_i))_{i=0}^l \in R_X(a - b)$  such that  $\sum_{i=0}^l d_X(x_i, y_i) < d_{FX}^1(a, b)$ . Let  $n_0 := 2l + 2$ ,  $a_0 := \sum_{i=0}^l (\delta_{2i}^{g_i} - \delta_{2i+1}^{g_i}) \in F_0(n_0)$ , and  $f_0, g_0 : 2n \rightarrow X$  be two functions defined by  $f_0(2i) = f_0(2i + 1) = x_i = g_0(2i)$  and  $g_0(2i + 1) = y_i$  for every  $i \in \{0, \dots, l\}$ . It is easy to see that

$Ff_0(a_0) = o$  and  $Fg_0(a_0) = \sum_{i=0}^l (\delta_{x_i}^{g_i} - \delta_{y_i}^{g_i}) = a - b$ . Then  $((a_0, f_0, g_0)) \in L_{FX}(o, a - b)$  is an  $(o, a - b)$ -linking chain, so

$$\begin{aligned} d_{FX}^1(a, b) &= d_{FX}^1(a - b, o) \leq d_{X^{\leftarrow \omega}}^1(f_0, g_0) = \sum_{j \in 2l+2} d_X(f_0(j), g_0(j)) = \\ &= \sum_{i=0}^l (d_X(f_0(2i), g_0(2i)) + d_X(f_0(2i+1), g_0(2i+1))) = \\ &= \sum_{i=0}^l (d_X(x_i, x_i) + d_X(x_i, y_i)) < d_{FX}^1(a, b), \end{aligned}$$

which is a desired contradiction showing that  $d_{FX}^1(a, b) \leq \|a - b\|_{FX}$ . Observe that the inclusion  $((g_i, x_i, y_i))_{i=0}^l \in R_X^{\subset}(a - b)$  implies that  $((a_0, f_0, g_0)) \in L_{FX}^{\subset}(a, b)$ , which yields the inequality  $d_{FX}^{\subset}(a, b) \leq \|a - b\|_{FX}^{\subset}$ . The inequality  $d_{FX}^1(a, b) \leq d_{FX}^{\subset}(a, b)$  follows from the inclusion  $L_{FX}^{\subset}(a, b) \subseteq L_{FX}(a, b)$ .

Next, we prove that  $\|a - b\|_{FX} \leq d_{FX}^1(a, b)$ . To derive a contradiction, assume that  $d_{FX}^1(a, b) < \|a - b\|_{FX}$  and find an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{FX}(a, b)$  such that  $\sum_{i=1}^l d_{X^{\leftarrow \omega}}^1(f_i, g_i) < \|a - b\|_{FX}$ . For every  $i \in \{0, \dots, l\}$  let  $n_i := \text{dom}(f_i) = \text{dom}(g_i)$ . Consider the sequence  $(m_i)_{i=0}^{l+1}$  of numbers  $m_i = \sum_{j < i} n_j$  and put  $m := m_{l+1}$ . For every number  $j < m$ , find (unique) numbers  $i_j \in \{0, \dots, l\}$  and  $r_j \in n_{i_j}$  such that  $j = m_{i_j} + r_j$ . Then put  $(g_j, x_j, y_j) := (a_{i_j}(r_j), f_{i_j}(r_j), g_{i_j}(r_j))$ . We claim that the sequence  $((g_j, x_j, y_j))_{j=0}^{m-1}$  is  $(a - b)$ -representing.

Indeed,

$$\begin{aligned} \sum_{j \in m} (\delta_{x_j}^{g_j} - \delta_{y_j}^{g_j}) &= \sum_{i=0}^l \sum_{r \in n_i} (\delta_{f_i(r)}^{a_i(r)} - \delta_{g_i(r)}^{a_i(r)}) = \sum_{i=0}^l (Ff_i(a_i) - Fg_i(a_i)) = \\ &= Ff_0(a_0) - Fg_l(a_l) + \sum_{i=1}^l (Ff_i(a_i) - Fg_{i-1}(a_{i-1})) = a - b + 0. \end{aligned}$$

Then

$$\|a - b\|_{FX} \leq \sum_{j \in m} d_X(x_j, y_j) = \sum_{i=0}^l \sum_{r \in n_i} d_X(f_i(r), g_i(r)) = \sum_{i=0}^l d_{X^{\leftarrow \omega}}^1(f_i, g_i) < \|a - b\|_{FX},$$

which is a desired contradiction that completes the proof of  $d_{FX}^1(a, b) = \|a - b\|_{FX} \leq d_{FX}^{\subset}(a, b) \leq \|a - b\|_{FX}^{\subset}$ .

2. Now assume that  $|M_0| = 2$  and let  $g$  be the unique non-zero element of the group  $M_0$ . For every  $\varphi \in F_0X$ , the inclusions  $R_X^{\cap}(\varphi) \subseteq R_X^{\subset}(\varphi) \subseteq R_X(\varphi)$  imply that  $\|\varphi\|_{FX} \leq \|\varphi\|_{FX}^{\subset} \leq \|\varphi\|_{FX}^{\cap}$ . Assuming that  $\|\varphi\|_{FX} < \|\varphi\|_{FX}^{\cap}$ , find a sequence  $s = ((g_i, x_i, y_i))_{i=0}^l \in R_X(\varphi)$  such that  $\sum_{i=0}^l d_X(x_i, y_i) < \|\varphi\|_{FX}^{\cap}$ . We can assume that the length  $k$  of this sequence is the smallest possible. It follows from  $\|\varphi\|_{FX} < \|\varphi\|_{FX}^{\cap}$  that  $a \neq b$  and hence  $k \geq 0$ . The minimality of  $l$  ensures that for every  $i \leq l$  the element  $g_i$  is non-zero and hence is equal to the unique non-zero element  $g$  of the 2-element group  $M_0$ . Also the minimality of  $l$  implies that  $\{x_i, y_i\} \neq \{x_j, y_j\}$  for any  $i < j$ . Indeed, assuming that  $\{x_i, y_i\} = \{x_j, y_j\}$  for some numbers  $i < j \leq l$ , we can see that  $\chi_{x_i, y_i}^{g_i} = \chi_{x_j, y_j}^{g_j}$  and hence  $\chi_{x_i, y_i}^{g_i} + \chi_{x_j, y_j}^{g_j} = o$ ,

which allows us to remove the triples  $(g_i, x_i, y_i)$  and  $(g_j, x_j, y_j)$  for the sequence  $s$  without breaking the  $\varphi$ -representation property. But this contradicts the minimality of  $l$ . If for some distinct  $i, j$  the doubletons  $\{x_i, y_i\}$  and  $\{x_j, y_j\}$  have a common point, then we can replace two triples  $(g_i, x_i, y_i)$  and  $(g_j, x_j, y_j)$  in the sequence  $s$  by one triple  $(g, x, y)$  where  $\{x, y\}$  coincides with the symmetric difference of the sets  $\{x_i, y_i\}$  and  $\{x_j, y_j\}$ . Here we observe that  $\chi_{x_i, y_j}^{g_i} + \chi_{x_j, y_j}^{g_j} = \chi_{x, y}^g$  and  $d_X(x, y) \leq d_X(x_i, y_i) + d_X(x_j, y_j)$ . So, we can make the sequence  $s$  shorter, which contradicts the minimality of  $l$ . Now we see that the sequence  $s$  belongs to the family  $R_X^\cap(\varphi)$ , which implies that  $\|\varphi\|_{FX}^\cap \leq \sum_{i=0}^l d_X(x_i, y_i) < \|\varphi\|_{FX}^\cap$ , and is a desired contradiction showing that  $\|\varphi\|_{FX} = \|\varphi\|_{FX}^\subseteq = \|\varphi\|_{FX}^\cap$ . Combining these equalities with the (in)equality  $d_{FX}^1(a, b) = \|a - b\|_{FX} \leq d_{FX}^\subseteq(a, b) \leq \|a - b\|_{FX}^\subseteq$ , proved in the first part of the theorem and the inequality  $\check{d}_{FX}^1(a, b) \leq d_{FX}^1(a, b)$  proved in Theorem 26.1(3), we conclude that

$$\check{d}_{FX}^1(a, b) \leq d_{FX}^1(a, b) = d_{FX}^\subseteq(a, b) = \|a - b\|_{FX} = \|a - b\|_{FX}^\subseteq = \|a - b\|_{FX}^\cap.$$

To complete the proof of the theorem, it remains to show that  $\|a - b\|_{FX} \leq \check{d}_{FX}^1(a, b)$ , which will follow from the invariantness of the distance  $\check{d}_{FX}^1$  as soon as we check that  $\|\varphi\|_{FX} \leq \check{d}_{FX}^1(\varphi, o)$  for every  $\varphi \in F_0X$ .

Assuming that the latter inequality does not hold, we conclude that the set

$$\Phi := \{\varphi \in F_0X : \check{d}_{FX}^1(\varphi, o) < \|\varphi\|_{FX}\}$$

is not empty. Let  $\varphi \in \Phi$  be an element whose support  $S = \text{supp}(\varphi)$  has the smallest possible cardinality. Since  $\varphi \in F_0X$ , the cardinality  $|S|$  is even (and non-zero as  $\|\varphi\|_{FX} > 0$ ). We claim that the restriction  $d_X \upharpoonright S$  is an  $\infty$ -metric. Indeed, assuming that  $S$  contains two distinct points  $x, y$  with  $d_X(x, y) = 0$ , we can consider the function  $\delta_x^g - \delta_y^g \in F_0X$  and conclude that  $0 \leq \check{d}_{FX}^1(\delta_x^g, \delta_y^g) \leq d_{FX}^1(\delta_x^g, \delta_y^g) \leq d_X(x, y) = 0$ . Then the function  $\psi = \varphi + \delta_x^g - \delta_y^g$  has support  $\text{supp}(\psi) = S \setminus \{x, y\}$  of cardinality  $|\text{supp}(\psi)| < |S| = |\text{supp}(\varphi)|$  and belongs to the set  $\Phi$  since

$$\check{d}_{FX}^1(\psi, o) \leq \check{d}_{FX}^1(\delta_x^g - \delta_y^g, o) + \check{d}_{FX}^1(\varphi, o) = \check{d}_{FX}^1(\varphi, o) < \|\varphi\|_{FX} \leq \|\psi\|_{FX} + \|\delta_x^g - \delta_y^g\|_{FX} = \|\psi\|_{FX},$$

but this contradicts the choice of  $\varphi$ . This contradiction shows that the restriction  $d_X \upharpoonright S \times S$  is an  $\infty$ -metric.

As we already know,

$$\|F\mathbf{e}_X(\varphi)\|_{FEX}^\subseteq = d_{FEX}^1(F\mathbf{e}_X(\varphi), o) = \check{d}_{FX}^1(\varphi, o) < d_{FX}^1(\varphi, o) = \|\varphi\|_{FX}.$$

By the definition of  $\|F\mathbf{e}_X(\varphi)\|_{FEX}^\subseteq$ , there is a sequence  $((g_i, x_i, y_i))_{i=0}^l \in R_{FEX}^\subseteq(F\mathbf{e}_X(\varphi))$  such that  $\sum_{i=0}^l (\delta_{x_i}^{g_i} - \delta_{y_i}^{g_i}) = F\mathbf{e}_X(\varphi)$  and  $\sum_{i=0}^l d_{EX}(x_i, y_i) < \|\varphi\|_{FX}$ . The definition of  $R_{EX}^\subseteq(F\mathbf{e}_X(\varphi))$  ensures that

$$\bigcup_{i=0}^l \{x_i, y_i\} \subseteq \text{supp}(F\mathbf{e}_X(\varphi)) \subseteq \mathbf{e}_X[\text{supp}(\varphi)] = \mathbf{e}_X[S].$$

For every  $i \in \{0, \dots, l\}$  choose points  $x'_i, y'_i \in S$  such that  $\mathbf{e}_X(x'_i) = x_i$  and  $\mathbf{e}_X(y'_i) = y_i$ . The injectivity of  $\mathbf{e}_X$  on the set  $S = \text{supp}(\varphi)$  and the equality  $F\mathbf{e}_X(\varphi) = \sum_{i=0}^l (\delta_{x_i}^{g_i} - \delta_{y_i}^{g_i})$  imply that  $\varphi = \sum_{i=0}^l (\delta_{x'_i}^{g_i} - \delta_{y'_i}^{g_i})$ . Then  $((g_i, x_i, y_i))_{i=0}^l \in R_{FX}(\varphi)$  and

$$\|\varphi\|_{FX} \leq \sum_{i=0}^l d_X(x'_i, y'_i) = \sum_{i=0}^l d_{EX}(x_i, y_i) < \|\varphi\|_{FX},$$

which is a desired contradiction that completes the proof.  $\square$

It should be mentioned that the assumption  $|M_0| = 2$  in Theorem 29.7 is essential and cannot be replaced by any weaker assumption.

**Example 29.8.** Let  $\{z_1, z_2, z_3\}$  be an equilateral triangle with unit sides in the Euclidean plane and  $z_0$  its center. Consider the distance space  $X = \{z_0, z_1, z_2, z_3\}$  endowed with the metric  $d_X$ , inherited from the plane. If  $|M_0| > 2$ , then there exists a function  $\varphi \in F_0X$  such that  $\varphi(z_0) = 0$  and  $\varphi(z_i) \neq 0$  for all  $i \in \{1, 2, 3\}$ . The function has  $\text{supp}(\varphi) = \{z_1, z_2, z_3\}$ . It can be shown that

$$d_{FX}^1(\varphi, 0) = \|\varphi\|_{FX} = \sqrt{3} < 2 = d_{FX}^C(\varphi, 0) = \|\varphi\|_{FX}^C < \|\varphi\|_{FX}^\cap = \infty.$$

**Remark 29.9.** If the monoid  $M$  is a cyclic group of order  $n \leq \omega$  with generator  $g$ , then the group  $FX = F(X, M)$  can be identified with the free Abelian group of exponent  $n$  over  $X$ . Each distance  $d_X$  on  $X$  induces the *Graev distance*  $d_{FX}^g$  on  $FX$ , defined by

$$d_{FX}^g(a, b) = \inf \left( \left\{ \sum_{i=0}^l d_X(x_i, y_i) : x_0, y_0, \dots, x_l, y_l \in X, a - b = \sum_{i=0}^l (\delta_{x_i}^g - \delta_{y_i}^g) \right\} \cup \{\infty\} \right).$$

Theorem 29.7 implies that  $d_{FX}^g = d_{FX}^1$  if and only if  $n \leq 3$ . More information on Graev distances on free Abelian (Boolean) groups can be found in [1, §7.2], [20] (and [28, §4]).

**Remark 29.10.** Theorem 29.7(2) suggests an algorithm for calculating the distances  $\check{d}_{FX}^1 = d_{FX}^1$  in case of  $|G| = 2$ . However, for calculating the norm  $\|\varphi\|_{FX}$  of for an element  $\varphi \in F_0X$  with support of cardinality  $|\text{supp}(\varphi)| = 2n$ , the brute force algorithm requires checking  $\frac{(2n)!}{2^n n!}$  cases, which yields superexponential algorithmic complexity.

**Problem 29.11.** *Is there a simpler (desirably, polynomial time) algorithm for calculating the distances  $\check{d}_{FX}^1 = d_{FX}^1$  on the group  $FX$  in case of the two-element group  $|G| = 2$ ?*

**Problem 29.12.** *Is there any reasonable algorithm for calculating the distances  $\check{d}_{FX}^1$  and  $d_{FX}^1$  on  $FX$  for arbitrary finite group  $G$  and arbitrary metric space  $(X, d_X)$  with computable metric?*

### 30. THE METRIZATIONS OF THE HYPERSPACE FUNCTOR

If  $M$  is a two-element semilattice, then the functor  $F = F(-, M)$ , considered in the preceding section, is isomorphic to the hyperspace functor  $H$  assigning to each set  $X$  the semilattice  $[X]^{<\omega}$  of finite subsets of  $X$  and to each function  $f : X \rightarrow Y$  between sets the function  $Hf : HX \rightarrow HY$ ,  $Hf : a \mapsto f[a]$ .

We recall that a *semilattice* is a commutative semigroup  $S$  whose any element  $x \in S$  is *idempotent* (i.e.  $x + x = x$ ). Each 2-element semilattice is isomorphic to the semilattice  $2 = \{0, 1\}$  endowed with the operation of maximum. The hyperspace  $HX$  is endowed with the semilattice operation of union.

Identifying each finite set  $A \subseteq X$  with its characteristic function  $\chi_A : X \rightarrow 2 = \{0, 1\}$ , we can identify the semilattice  $HX$  with the monoid  $F(X, 2)$ . This determines an isomorphism of the functors  $H$  and  $F(-, 2)$ .

It is easy to see that the functor  $H$  preserves supports; moreover,  $\text{supp}(a) = a$  for every set  $X$  and element  $a \in HX$ .

The hyperspace functor  $H$  has a metrization by the Hausdorff distance  $d_{HX}$ , defined in Section 6. The interplay between the distances  $d_{HX}$ ,  $d_{HX}^p$  and  $\check{d}_{HX}^p$  is described in the following theorem.

From now on, we assume that  $p \in [1, \infty]$  and  $(X, d_X)$  is a distance space.

**Theorem 30.1.**  $d_{HX} = \check{d}_{HX}^\infty = d_{HX}^\infty \leq \check{d}_{HX}^p \leq d_{HX}^p$ .

Moreover, the distances  $d_{HX}^p$  and  $\check{d}_{FX}^p$  are subinvariant in the sense that

$$d_{HX}^p(a \cup c, b \cup c) \leq d_{HX}^p(a, b) \text{ and } \check{d}_{HX}^p(a \cup c, b \cup c) \leq \check{d}_{HX}^p(a, b)$$

for every  $a, b, c \in HX$ .

*Proof.* The subinvariance of the distances  $d_{HX}$  and  $\check{d}_{HX}^p$  follows from Proposition 29.1 and Corollary 29.2.

By Theorem 16.6,  $d_{HX} \leq d_{HX}^\infty$ . Next, we prove that  $d_{HX}^\infty(a, b) \leq d_{HX}(a, b)$  for any  $a, b \in HX$ . This inequality is trivial if  $a = b$  or  $d_{HX}(a, b) = \infty$ . So, assume that  $a \neq b$  and  $d_{HX}(a, b) < \infty$ . In this case  $a, b$  are non-empty finite subsets of  $X$ . By the definition of  $d_H(a, b)$ , there are two functions  $p : a \rightarrow b$  and  $q : b \rightarrow a$  such that

$$\max\{\max_{x \in a} d_X(x, p(x)), \max_{y \in b} d_X(y, q(y))\} = d_{HX}(a, b).$$

Fix any bijective functions  $\alpha : |a| \rightarrow a$  and  $\beta : |b| \rightarrow b$ . Let  $n_0 = |a| + |b|$  and consider the functions  $f_0 : n_0 \rightarrow a$  and  $g_0 : n_0 \rightarrow b$  defined by the formulas

$$f_0(i) = \begin{cases} \alpha(i), & \text{if } 0 \leq i < |a|; \\ q \circ \beta(i - |a|), & \text{if } |a| \leq i < n_0; \end{cases} \quad \text{and} \quad g_0(i) = \begin{cases} p \circ \alpha(i), & \text{if } 0 \leq i < |a|; \\ \beta(i - |a|), & \text{if } |a| \leq i < n_0. \end{cases}$$

For the element  $a_0 := n_0 \in Hn_0$ , we get  $Hf_0(a_0) = f_0[n_0] = a$  and  $Hg_0(a_0) = g_0[n_0] = b$ , which means that  $((a_0, f_0, g_0)) \in L_{HX}(a, b)$  is an  $a, b$ -linking chain. Then

$$\begin{aligned} d_{HX}^\infty(a, b) &\leq d_{X^\omega}^\infty(f_0, g_0) = \max_{x \in n_0} d_X(f_0(x), g_0(x)) = \\ &= \max\{\max_{0 \leq i < |a|} d_X(\alpha(i), p \circ \alpha(i)), \max_{|a| \leq i < n_0} d_X(q \circ \beta(i), \beta(i))\} \leq \\ &\leq \max\{\max_{x \in a} d_X(x, p(x)), \max_{y \in b} d_X(q(y), y)\} = d_{HX}(a, b). \end{aligned}$$

Now we see that  $d_{HX} = d_{HX}^\infty$  (for any distance space  $(X, d_X)$ ). In particular,  $d_{HEX} = d_{HEX}^\infty$ .

Next, we prove that  $d_{HX}(a, b) = \check{d}_{HX}^\infty(a, b) = d_{HX}^\infty(a, b)$  for any elements  $a, b \in HX$ . Looking at the definition of the Hausdorff distance, we can see that

$$d_{HX}(a, b) = d_{HEX}(He_X(a), He_X(b)).$$

Now the definition of the distance  $\check{d}_{HX}$  and Theorem 26.1(3) ensure that

$$\begin{aligned} d_{HX}^\infty(a, b) &= d_{HX}(a, b) = d_{HEX}(He_X(a), He_X(b)) = \\ &= d_{HEX}^\infty(He_X(a), He_X(b)) = \check{d}_{HX}^\infty(a, b) \leq d_{HX}^\infty(a, b), \end{aligned}$$

which implies the desired equalities  $d_{HX}(a, b) = \check{d}_{HX}^\infty(a, b) = d_{HX}^\infty(a, b)$ . Applying Theorem 26.1(2,3), we conclude that  $d_{HX}(a, b) = \check{d}_{HX}^\infty(a, b) = d_{HX}^\infty(a, b) \leq \check{d}_{HX}^p(a, b) \leq d_{HX}^p(a, b)$ .  $\square$

For every  $n \in \mathbb{N}$  consider the subfunctor  $H_n$  of the functor  $H$ , assigning to each set  $X$  the subfamily  $H_n X := \{a \in HX : |a| \leq n\}$ . It is easy to see that the functor  $H_n$  has degree  $\leq n$  and preserves supports. The properties of its metizations are described in the following proposition.

**Proposition 30.2.**  $d_{HX} \leq \check{d}_{H_n X}^p \leq d_{H_n X}^p \leq n^{\frac{1}{p}} \cdot d_{H_n X}^\infty \leq 3 \cdot n^{\frac{1}{p}} \cdot d_{HX}$ .

*Proof.* The inequality  $d_{HX} \leq \check{d}_{H_n}^p$  follows from Theorem 30.1 and 27.3. The inequalities  $\check{d}_{H_n X}^p \leq d_{H_n X}^p \leq n^{\frac{1}{p}} \cdot d_{H_n X}^\infty$  follow from Theorems 26.1(3) and 25.1(4). It remains to prove that  $d_{H_n X}^\infty(a, b) \leq 3 \cdot d_{HX}(a, b)$  for any sets  $a, b \in H_n X$ . This inequality holds trivially if  $a = b$  or  $d_{HX}(a, b) = \infty$ . So, assume that  $a \neq b$  and  $d_{HX}(a, b) < \infty$ . In this case the sets  $a, b$  are non-empty. Let  $\gamma : a' \rightarrow b'$  be a maximal bijective function between subsets  $a' \subseteq a$  and  $b' \subseteq b$  such that  $d_X(x, \gamma(x)) \leq d_{HX}(a, b)$  for every  $x \in a'$ . By the maximality, the function  $\gamma$  has an extension  $p : a \rightarrow b'$  such that  $\max_{x \in a} d_X(x, p(x)) \leq d_{HX}(a, b)$ , and  $\gamma^{-1}$  has an extension  $q : b \rightarrow a'$  such that  $\max_{y \in b} d_X(y, q(y)) \leq d_{HX}(a, b)$ .

Now we define an  $(a, b)$ -linking chain  $((a_0, f_0, g_0), (a_1, f_1, g_1), (a_2, f_2, g_2)) \in L_{H_n X}(a, b)$  such that  $d_{H_n}^\infty(a, b) \leq \sum_{i=0}^2 d_{X^{\leq \omega}}^\infty(f_i, g_i) \leq 3 \cdot d_{HX}(a, b)$ .

Let  $n_0 = |a|$ ,  $a_0 = n_0 \in H_n n_0$ ,  $f_0 : |a| \rightarrow a$  be any bijective function and  $g_0 = p \circ f_0$ .

Let  $n_1 = |a'| = |b'|$ ,  $a_1 = n_1 \in H_n n_1$ ,  $f_1 : n_1 \rightarrow b'$  be any bijection and  $g_1 = \gamma^{-1} \circ f_1$ .

Let  $n_2 = |b|$ ,  $a_2 = n_2 \in H_n n_2$ ,  $g_2 : n_2 \rightarrow b$  be any bijection and  $f_2 = q \circ g_2$ .

Observe that  $a = f_0[n_0]$ ,  $g_0[n_0] = p[a] = b' = f_1[n_1]$ ,  $g_1[n_1] = a' = q[b] = f_2[n_2]$ ,  $g_2[n_2] = b$ , which means that  $((a_i, f_i, g_i))_{i=0}^2 \in L_{H_n X}(a, b)$  and then

$$d_{H_n X}^\infty(a, b) \leq \sum_{i=0}^2 d_{X^{\leq \omega}}^\infty(f_i, g_i) \leq 3 \cdot d_{HX}(a, b). \quad \square$$

**Problem 30.3.** *Can the constant 3 in Proposition 30.2 be improved?*

The distance  $d_{HX}^1$  admits an equivalent graph-theoretic description, similar to the description of the metric  $d_{FX}^1$  in Theorem 29.7. It is defined with the help of graphs, see [3], [4].

By a *graph* we understand a pair  $\Gamma = (V, E)$  consisting of a finite set  $V$  of vertices and a finite set  $E$  of edges. Each edge  $e \in E$  is a non-empty subset of  $V$  of cardinality  $|e| \leq 2$ . For a graph  $\Gamma = (V, E)$ , the *connected component* of a vertex  $v \in V$  is the set  $\Gamma_v$  of all vertices  $u \in V$  for which there exists a sequence  $v = v_0, \dots, v_n = u$  such that  $\{v_{i-1}, v_i\} \in E$  for every  $i \in \{1, \dots, n\}$ .

By a *graph in the distance space*  $(X, d_X)$  we understand any graph  $\Gamma = (V, E)$  with  $V \subseteq X$ . In this case we can define the length  $\ell(\Gamma)$  of  $\Gamma$  by the formula

$$\ell(\Gamma) = \sum_{e \in E} \bar{d}_X(e)$$

where  $\bar{d}_X(e) = \max\{d_X(x, y) : x, y \in e\}$ .

For two finite sets  $a, b \subseteq X$  let  $\mathbf{\Gamma}(a, b)$  be the family of graphs  $\Gamma = (V, E)$  in  $X$  such that  $a \cup b \subseteq V$  and each connected component of  $\Gamma$  intersects both sets  $a$  and  $b$ . Observe that the family  $\mathbf{\Gamma}(a, b)$  contains the complete graph on the set  $a \cup b$  and hence is not empty.

**Theorem 30.4.** *For any elements  $a, b \in HX$  we have*

$$d_{HX}(a, b) \leq d_{HX}^1(a, b) = \inf(\{\infty\} \cup \{\ell(\Gamma) : \Gamma \in \mathbf{\Gamma}(a, b)\}).$$

*Proof.* The inequality  $d_{HX} \leq d_{HX}^1$  was proved in Proposition 30.1.

Next, we prove that  $\hat{d}_{HX}(a, b) \leq d_{HX}^1(a, b)$ , where

$$\hat{d}_{HX}(a, b) := \inf(\{\infty\} \cup \{\ell(\Gamma) : \Gamma \in \mathbf{\Gamma}(a, b)\}).$$

To derive a contradiction, assume that  $d_{HX}^1(a, b) < \hat{d}_{HX}(a, b)$  and find an  $(a, b)$ -linking chain  $((a_i, f_i, g_i))_{i=0}^l \in L_{HX}(a, b)$  such that  $\sum_{i=0}^l d_{X^{\leq \omega}}^1(f_i, g_i) < \hat{d}_{HX}(a, b)$ . For every  $i \in \{0, \dots, l\}$

let  $n_i = \text{dom}(f_i) = \text{dom}(g_i)$ . Consider the graph  $\Gamma = (V, E)$  with

$$V = \bigcup_{i=0}^l (f_i[a_i] \cup g_i[a_i]) \quad \text{and} \quad E = \bigcup_{i=0}^l \{\{f_i(x), g_i(x)\} : x \in a_i\}.$$

Observe that

$$\ell(\Gamma) \leq \sum_{i=0}^l \sum_{x \in n_i} d_X(f_i(x), g_i(x)) = \sum_{i=0}^l d_{X^{\omega}}^1(f_i, g_i) < \hat{d}_{HX}(a, b).$$

We claim that  $\Gamma \in \mathbf{\Gamma}(a, b)$ . Observe that  $a \cup b = f_0[a_0] \cup g_l[a_l] \subseteq V$ . Next, we check that for every  $x \in V$  the connected component  $\Gamma_x$  of the graph  $\Gamma$  intersects the sets  $a$  and  $b$ . Find  $i \in \{0, \dots, l\}$  such that  $x \in f_i[a_i] \cup g_i[a_i]$ . Then  $x \in \{f_i(t_i), g_i(t_i)\}$  for some  $t_i \in a_i$ . For the point  $x_i := g_i(t_i) \in g_i[a_i] = f_{i+1}[a_{i+1}]$  we can find a point  $t_{i+1} \in a_{i+1}$  with  $x_i = f_{i+1}(t_{i+1})$  and consider the point  $x_{i+1} = g_{i+1}(t_{i+1})$ . Continuing by induction, we can construct sequences of points  $(x_j)_{j=x}^{l+1}$  and  $(t_j)_{j=i}^l \in \prod_{j=i}^l n_j$  such that  $\{x, x_i\} \in E$ ,  $x_i = f_i(t_i)$ ,  $x_{l+1} = g_l(t_l) \in g_l[a_l] = b$  and  $g_{j-1}(t_{j-1}) = x_j = f_j(t_j)$  for every  $j \in \{i+1, \dots, l\}$ . By the definition of the set  $E$ , for every  $j \in \{i, \dots, l\}$  the doubleton  $\{x_j, x_{j+1}\} = \{f_j(t_j), g_j(t_j)\}$  is an edge of the graph  $\Gamma$ . Then the chain  $x, x_i, \dots, x_{l+1}$  witnesses that  $x_{l+1} \in \Gamma_x \cap b$ . By analogy we can prove that the intersection  $\Gamma_x \cap a$  is not empty. Therefore,  $\Gamma \in \mathbf{\Gamma}(a, b)$  and  $\hat{d}_{HX}(a, b) \leq \ell(\Gamma) < \hat{d}_{HX}(a, b)$ . This contradiction shows that  $\hat{d}_{HX}(a, b) \leq d_{HX}^1(a, b)$ .

Next, we prove that  $d_{HX}^1(a, b) \leq \hat{d}_{HX}(a, b)$ . Assuming that  $\hat{d}_{HX}(a, b) < d_{HX}^1(a, b)$ , we can find a graph  $\Gamma \in \mathbf{\Gamma}(a, b)$  such that  $\ell(\Gamma) < d_{HX}^1(a, b)$ . By Lemma 30.5 (proved below), there exists an  $(a, b)$ -linking chain  $s \in L_{FX}(a, b)$  such that  $\Sigma d_X^1(s) \leq \ell(\Gamma)$ . Then  $d_{GX}^1(a, b) \leq \Sigma d_X^1(s) \leq \ell(\Gamma) < d_{HX}^1(a, b)$ , which is a desired contradiction showing that  $d_{HX}^1 = \hat{d}_{HX}$ .  $\square$

**Lemma 30.5.** *For any finite sets  $a, b \subseteq X$  and graph  $\Gamma \in \mathbf{\Gamma}(a, b)$ , there exists an  $(a, b)$ -linking chain  $s \in L_{FX}(a, b)$  such that  $\Sigma d_X^1(s) \leq \ell(\Gamma)$ .*

*Proof.* It suffices to prove that for every  $n \in \mathbb{N}$  the following statement  $(*_n)$  holds:

$(*_n)$  *for any finite sets  $a, b \subseteq X$  and graph  $\Gamma = (V, E) \in \mathbf{\Gamma}(a, b)$  with  $|V| \leq n$ , there exists an  $(a, b)$ -linking chain  $s \in L_{FX}(a, b)$  such that  $\Sigma d_X^1(s) \leq \ell(\Gamma)$ .*

To see that  $(*_1)$  is true, take any finite sets  $a, b \subseteq X$  and any graph  $\Gamma = \mathbf{\Gamma}(a, b)$  with  $|V| = 1$ . In this case  $\mathbf{\Gamma}(a, b) \neq \emptyset$  implies that either  $a, b$  are both empty or  $a = b = V$ . In both cases we get  $a = b$ . So, we can find an  $(a, b)$ -linking chain  $s \in L_{FX}(a, b)$  such that  $\Sigma d_X^1(s) = 0 \leq \ell(\Gamma)$ .

Now assume that for some  $n \in \mathbb{N}$  the statement  $(*_n)$  is true. Take any finite sets  $a, b \in \mathbb{N}$  and a graph  $\tilde{\Gamma} = (\tilde{V}, \tilde{E}) \in \mathbf{\Gamma}(a, b)$  with  $|\tilde{V}| = n+1$ . If  $a = b$ , then we can find an  $(a, b)$ -linking chain  $s \in L_{FX}(a, b)$  with  $\Sigma d_X^1(s) = 0 \leq \ell(\tilde{\Gamma})$  and finish the proof. So, we assume that  $a \neq b$ .

Let  $\Gamma = (V, E)$  be any minimal subgraph of the graph  $\tilde{\Gamma}$  such that  $\Gamma \in \mathbf{\Gamma}(a, b)$ . If  $|V| \leq n$ , then by the inductive assumption, there exists an  $(a, b)$ -linking chain  $s \in L_{FX}(a, b)$  such that  $\Sigma d_X^1(s) \leq \ell(\Gamma) \leq \ell(\tilde{\Gamma})$  and we are done. So, we assume that  $|\Gamma| = n+1$ . By the minimality of  $\Gamma$ , each connected component of  $\Gamma$  is a tree whose pendant vertices belong to the set  $a \cup b$ . Since  $\Gamma \in \mathbf{\Gamma}(a, b)$  and  $a \neq b$ , the graph  $\Gamma$  has some non-trivial connected component. Take any pendant vertex  $v$  of this non-trivial connected component and find a (unique) vertex  $u \in V \setminus \{v\}$  such that  $\{v, u\} \in E$ . Consider the subgraph  $\Gamma' = (V', E')$  of  $\Gamma$  with  $V' := V \setminus \{v\}$  and  $E' = \{e \in E : v \notin e\}$ . It follows that  $|V'| = n$ .

Three cases are possible.



1) If  $v \in b \setminus a$ , then let  $b' = (b \setminus \{v\}) \cup \{u\}$ . Observe that every path connecting the vertex  $v$  with another vertex of  $\Gamma$  includes the vertex  $u$ . Consequently,  $\Gamma' \in \mathbf{\Gamma}(a, b')$ . By the inductive assumption, there exists an  $(a, b')$ -linking chain  $s' = ((a_i, f_i, g_i))_{i=0}^{k-1} \in L_{FX}(a, b')$  such that  $\Sigma d_X^1(s') \leq \ell(\Gamma')$ . For this chain we have  $f_i[a_i] = a$  and  $g_{k-1}[a_i] = b'$ . Let  $n_k = |b|$ . It follows that  $b = b' \cup \{v\}$  or  $b = (b' \setminus \{u\}) \cup \{v\}$ .

1a) If  $b = b' \cup \{v\}$ , then choose any bijective map  $f_k : n_k \rightarrow b$  such that  $f_k(n_k - 2) = u$  and  $f_k(n_k - 1) = v$ . Next, consider the surjective map  $g_k : n_k \rightarrow b'$  such that  $g_k(i) = f_k(i)$  for all  $i \leq n_k - 2$  and  $g_k(n_k - 1) = u$ . Let  $a_k = n_k \in Hn_k$ . Observe that  $Hf_k(a_k) = f_k[n_k] = b'$  and  $Hg_k(a_k) = g_k[n_k] = b$ . Then  $s = ((a_i, f_i, g_i))_{i=0}^k$  is an  $(a, b)$ -linking chain such that  $\Sigma d_X^1(s) = \Sigma d^1(s') + d_X(u, v) \leq \ell(\Gamma') + d_X(u, v) = \ell(\Gamma)$ .

1b) If  $b = (b' \setminus \{u\}) \cup \{v\}$ , then choose any bijective map  $f_k : n_k \rightarrow b$  such that  $f_k(n_k - 1) = v$ . Next, consider the bijective map  $g_k : n_k \rightarrow b'$  such that  $g_k(i) = f_k(i)$  for all  $i \leq n_k - 2$  and  $g_k(n_k - 1) = u$ . Let  $a_k = n_k \in Hn_k$ . Observe that  $Hf_k(a_k) = f_k[n_k] = b'$  and  $Hg_k(a_k) = g_k[n_k] = b$ . Then  $s = ((a_i, f_i, g_i))_{i=0}^k$  is an  $(a, b)$ -linking chain such that  $\Sigma d_X^1(s) = \Sigma d^1(s') + d_X(u, v) \leq \ell(\Gamma') + d_X(u, v) = \ell(\Gamma)$ .

2) The case  $v \in a \setminus b$  can be considered by analogy with the case 1).

3) It remains to consider the third case  $v \in a \cap b$ . Let  $\Gamma_v = (V_v, E_v)$  be the connected component of the graph  $\Gamma$  that contains the point  $v$ . The third case has 4 subcases.

3a) The sets  $V_v \cap a \setminus \{v\}$  and  $V_v \cap b \setminus \{v\}$  are not empty. In this case put  $a' = a \setminus \{v\}$  and  $b' = b \setminus \{v\}$  and observe that the graph  $\Gamma'$  belongs to the family  $\mathbf{\Gamma}(a', b')$ . Applying the statement  $(*_n)$  to the graph  $\Gamma'$ , we can find an  $(a', b')$ -linking chain  $((a'_i, f_i, g_i))_{i \in k} \in L_{FX}(a', b')$  such that  $\Sigma d_X^1(s) \leq \ell(\Gamma')$ . For every  $i \in k$  let  $n'_i = \text{dom}(f'_i) = \text{dom}(g'_i)$ ,  $n_i := n'_i + 1$ ,  $a_i := a'_i \cup \{n'_i\} \subseteq n_i$  and  $f_i, g_i : n_i \rightarrow X$  be two maps such that  $f_i \upharpoonright n'_i = f'_i$ ,  $g_i \upharpoonright n'_i = g'_i$ , and  $f_i(n'_i) = g'_i(n'_i) = v$ . Then  $s = ((a_i, f_i, g_i))_{i \in k} \in L_{FX}(a, b)$  is an  $(a, b)$ -linking chain such that  $\Sigma d_X^1(s) = \Sigma d_X^1(s') \leq \ell(\Gamma') \leq \ell(\Gamma)$ .

3b) The sets  $V_v \cap a \setminus \{v\}$  and  $V_v \cap b \setminus \{v\}$  are empty. In this case  $a \cap V_v = \{v\} = b \cap V_v$ . Consider the subgraph  $\Gamma'' = (V'', E'')$  of  $\Gamma$  such that  $V'' = V \setminus V_v$  and  $E'' = \{e \in E : e \subseteq V''\}$  and let  $a' = a \setminus \{v\}$ ,  $b' = b \setminus \{v\}$ . Applying the inductive assumption to the graph  $\Gamma''$ , we can find an  $(a', b')$ -linking chain  $s' = ((a'_i, f_i, g'_i))_{i \in k} \in L_{FX}(a', b')$  such that  $\Sigma d_X^1(s') \leq \ell(\Gamma'')$ . For  $i \in k$  let  $n'_i = \text{dom}(f'_i) = \text{dom}(g'_i)$ ,  $n_i := n'_i + 1$ ,  $a_i := a'_i \cup \{n'_i\} \subseteq n_i$  and  $f_i, g_i : n_i \rightarrow X$  be two maps such that  $f_i \upharpoonright n'_i = f'_i$ ,  $g_i \upharpoonright n'_i = g'_i$ , and  $f_i(n'_i) = g'_i(n'_i) = v$ . Then  $s = ((a_i, f_i, g_i))_{i \in k} \in L_{FX}(a, b)$  is an  $(a, b)$ -linking chain such that  $\Sigma d_X^1(s) = \Sigma d_X^1(s') \leq \ell(\Gamma'') \leq \ell(\Gamma)$ .

3c) The set  $V_v \cap a \setminus \{v\}$  is not empty and  $V_v \cap b \setminus \{v\}$  is empty. Then for the sets  $a' = a \setminus \{v\}$  and  $b' = (b \setminus \{v\}) \cup \{u\}$ , the connected graph  $\Gamma'$  belongs to the family  $\mathbf{\Gamma}(a', b')$ . By the inductive assumption, there exists an  $(a', b')$ -linking chain  $s' = ((a'_i, f'_i, g'_i))_{i \in k} \in L_{FX}(a', b')$  such that  $\Sigma d_X^1(s') \leq \ell(\Gamma')$ . For every  $i \in k$  let  $n'_i = \text{dom}(f'_i) = \text{dom}(g'_i)$  and put  $n_i := n'_i + 1$ ,  $a_i := a'_i \cup \{n'_i\} \in Hn_i$ . Let  $f_i, g_i : n_i \rightarrow X$  be the functions such that  $f_i \upharpoonright n'_i = f'_i$ ,  $g_i \upharpoonright n'_i = g'_i$ , and  $f_i(n'_i) = v$ ,  $g_i(n'_i) = v$ . Let  $n_k = |b'| = |b|$ ,  $a_k = n_k$ , and  $f_k : n_k \rightarrow b'$  be any bijection such that  $f_k(n_k - 1) = u$ . Let  $g_k : n_k \rightarrow b$  be the bijective function such that  $g_k(i) = f_k(i)$  for  $i < n_k - 1$  and  $g_k(n_k - 1) = v$ . It is easy to see that  $s := ((a_i, f_i, g_i))_{i=0}^k$  is an  $(a, b)$ -linking chain such that  $\Sigma d_{FX}^1(s) = \Sigma d_{FX}^1(s') + d_X(u, v) \leq \ell(\Gamma') + d_X(u, v) = \ell(\Gamma)$ .

3d) The case  $V_v \cap a \setminus \{v\} = \emptyset \neq V_v \cap b \setminus \{v\}$  can be considered by analogy with the case 3c).

Thus we have handled all possible cases and proved the statement  $(*)_{n+1}$ .  $\square$

**Remark 30.6.** Let  $M$  be a commutative monoid,  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  be the functor of  $M$ -valued finitely supported functions, and  $(X, d_X)$  be a distance space.

- (1) If  $M$  is a two-element group, then  $\check{d}_{FX}^1 = d_{FX}^1$  on the group  $FX = F(X, M)$ , see Theorem 29.7(2).
- (2) If  $M$  is a two-element semilattice, then  $\check{d}_{FX}^\infty = d_{FX}^\infty$  on the semilattice  $FX = F(X, M)$ , see Theorem 30.1(2).

**Problem 30.7.** Find more functors  $F : \mathbf{Set} \rightarrow \mathbf{Set}$  such that  $\check{d}_{FX}^p = d_{FX}^p$  for some  $p \in [1, \infty]$  and all distance spaces  $(X, d_X)$ .

**Problem 30.8.** Are there (efficient) algorithms for calculation of the distances  $d_{HX}^p$  and  $\check{d}_{HX}^p$ .

**Remark 30.9.** The answer to Problem 30.8 strongly depends on the geometry of the distance space  $(X, d_X)$ . For subsets of the Euclidean plane, the problem of calculating the distance  $\hat{d}_{HX}$  is related to the classical Steiner's problem [14] of finding a tree of the smallest length that contains a given finite set. This problem is known [13] to be computationally very difficult. On the other hand, for finite subsets of the real line, there exists an algorithm [4] of complexity  $O(n \ln n)$ , calculating the distance  $d_{HX}^1(a, b)$  between two sets  $a, b \subset \mathbb{R}$  of cardinality  $|a| + |b| \leq n$ . Also there exists an algorithm of the same complexity  $O(n \ln n)$  calculating the Hausdorff distance  $d_{HX}(a, b)$  between two subsets of the real line. Finally, let us remark that the evident brute force algorithm for calculating the Hausdorff distance  $d_{HX}(a, b)$  between finite subsets of an arbitrary distance space  $(X, d_X)$  has complexity  $O(|a| \cdot |b|)$ .

#### REFERENCES

- [1] A. Arhangelskii, M. Tkachenko, *Topological groups and related structures*, Atlantis Press, Paris; World Sci. Publ., Hackensack, NJ, 2008. xiv+781 pp.
- [2] P. Assouad, *Étude d'une dimension métrique liée à la possibilité de plongements dans  $\mathbf{R}^n$* , C. R. Acad. Sci. Paris Sr. A-B **288**:15 (1979), A731–A734.
- [3] T. Banakh, *A new  $\ell^p$ -metric on the hyperspace of finite sets?*, (<https://mathoverflow.net/questions/277604>), 30.07.2017.
- [4] T. Banakh, *A quick algorithm for calculating the 1-distance between two finite sets on the real line?*, (<https://mathoverflow.net/a/277928/61536>), 04.08.2017.
- [5] T. Banakh, J. Garbulińska-Węgrzyn, *The completion of the hyperspace of finite subsets, endowed with the  $\ell^1$ -metric*, preprint.
- [6] T. Banakh, M. Martynenko, M. Zarichnyi, *On monomorphic topological functors with finite supports*, Carpathian Math. Publ. **4**:1 (2012) 4–12 (<http://arxiv.org/abs/1004.0457>).
- [7] T. Banakh, I. Zarichnyi, *Characterizing the Cantor bi-cube in asymptotic categories*, Groups, Geometry, and Dynamics, **5**:4 (2011) 691–728.
- [8] V.N. Basmanov, *Dimension and some functors with finite support*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. **6** (1981) 48–50.
- [9] V.N. Basmanov, *Covariant functors, retracts, and dimension*, Dokl. Akad. Nauk SSSR **271**:5 (1983), 1033–1036.
- [10] V.N. Basmanov, *Covariant functors of finite degrees on categories of Hausdorff compact spaces*, Fundam. Prikl. Mat. **2**:3 (1996), 637–654.
- [11] G. Bell, A. Dranishnikov, *Asymptotic dimension*, Topology Appl. **155**:12 (2008), 1265–1296.
- [12] Y. Benyamini, J. Lindenstrauss, *Geometric nonlinear functional analysis*, Amer. Math. Soc., Providence, RI, 2000. xii+488 pp.
- [13] M.W. Bern, R. Graham, *The shortest-network problem*, Scientific American. **260**:1 (1989) 84–89.
- [14] M. Brazil, R. Graham, D.A. Thomas, M. Zachariasen, *On the history of the Euclidean Steiner tree problem*, Arch. Hist. Exact Sci. **68**:3 (2014), 327–354.

- [15] S.V. Buyalo, N.D. Lebedeva, *Dimensions of locally and asymptotically self-similar spaces*, St. Petersburg Math. J. **19**:1 (2008), 45–65.
- [16] A. Dress, *Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: a note on combinatorial properties of metric spaces*, Adv. in Math. **53**:3 (1984), 321–402.
- [17] M. Fabian, P. Habala, P. Hájek, V.S. Montesinos, J. Pelant, V. Zizler, *Functional analysis and infinite-dimensional geometry*, Springer-Verlag, New York, 2001. x+451 pp.
- [18] K. Falconer, *Fractal Geometry. Mathematical foundations and applications*, John Wiley & Sons, Ltd., Chichester, 2003.
- [19] V.V. Fedorchuk, *Triples of infinite iterations of metrizable functors*, Izv. AN SSSR. Ser. Mat. **54**:3 (1990), 396–417.
- [20] M.I. Graev, *Free topological groups*, Izvestiya Akad. Nauk SSSR. Ser. Mat. **12** (1948) 279–324.
- [21] M. Gromov, *Asymptotic invariants of infinite groups*, *Geometric group theory*, Vol. 2 (Sussex, 1991), London Math. Soc. Lecture Note Ser., vol. 182, Cambridge Univ. Press, Cambridge, 1993, pp. 1295.
- [22] H. Herrlich, *Hyperconvex hulls of metric spaces*, Topology Appl. **44**:1-3 (1992), 181–187
- [23] J.R. Isbell, *Uniform spaces*, Mathematical Surveys, No. 12 Amer. Math. Soc., Providence, R.I. 1964.
- [24] J.R. Isbell, *Six theorems about injective metric spaces*, Comment. Math. Helv. **39** (1964), 6576.
- [25] U. Lang, T. Schlichenmaier, *Nagata dimension, quasisymmetric embeddings, and Lipschitz extensions*, International Mathematics Research Notices, **58** (2005), 3625–3655.
- [26] U. Lang, *Injective hulls of certain discrete metric spaces and groups*, J. Topol. Anal. **5**:3 (2013), 297–331.
- [27] T. Radul. O. Shukel, *Functors of finite degree and asymptotic dimension*, Mat. Stud. **29**:2 (2009), 204–206.
- [28] O. Sipacheva, *Free Boolean topological groups*, Axioms **4**:4 (2015), 492–517;  
(<https://arxiv.org/pdf/1612.04878.pdf>).
- [29] O. Shukel', *Functors of finite degree and asymptotic dimension zero*, Mat. Stud. **29**:1 (2008), 101–107; corrections in Mat. Stud. **32**:1 (2009), 112.
- [30] O. Shukel, M. Zarichnyi, *Asymptotic dimension of symmetric powers*, Math. Bull. Shevchenko Sci. Soc. **5** (2008), 304–311.
- [31] S. Todorčević, *Introduction to Ramsey spaces*, Princeton Univ. Press, 2010.
- [32] A. Teleiko, M. Zarichnyi, *Categorical topology of compact Hausdorff spaces*, VNTL Publishers, Lviv, (1999), 263 pp.

IVAN FRANKO NATIONAL UNIVERSITY OF LVIV AND JAN KOCHANOWSKI UNIVERSITY IN KIELCE  
*E-mail address:* `t.o.banakh@gmail.com`

IVAN FRANKO NATIONAL UNIVERSITY OF LVIV, UNIVERSYTETSKA 1, LVIV, 79000, UKRAINE  
*E-mail address:* `v.frider@yahoo.com`

IVAN FRANKO NATIONAL UNIVERSITY OF LVIV, UNIVERSYTETSKA 1, LVIV, 79000, UKRAINE  
*E-mail address:* `lesia.karчевska@gmail.com`

IVAN FRANKO NATIONAL UNIVERSITY OF LVIV AND UNIWERSYTET RZESZOWSKI  
*E-mail address:* `zarichnyi@yahoo.com`