

# Bending of light in novel 4D Gauss-Bonnet-de Sitter black holes by Rindler-Ishak method

Mohaddese Heydari-Fard<sup>1</sup>\*, Malihe Heydari-Fard<sup>2†</sup> and Hamid Reza Sepangi<sup>1‡</sup>

<sup>1</sup> *Department of Physics, Shahid Beheshti University, G. C., Evin, Tehran 19839, Iran*

<sup>2</sup> *Department of Physics, The University of Qom, 3716146611, Qom, Iran*

May 18, 2020

## Abstract

We study the bending of light in the space-time of black holes in the novel four-dimensional Einstein-Gauss-Bonnet theory of gravity, recently proposed by Glavan and Lin [26]. Using Rindler-Ishak method, the effect of Gauss-Bonnet coupling on the bending angle is studied. We show that a positive Gauss-Bonnet coupling gives a negative contribution to the Schwarzschild-de Sitter deflection angle, as one would expect.

**PACS numbers:** 04.50.Kd, 04.70. Bw, 04.80. Cc

## 1 Introduction

Many efforts have gone into understanding the nature of dark energy as being responsible of accelerated expansion of the observed universe. As is well known, one of the prime candidates for dark energy is the cosmological constant  $\Lambda$  [1]–[6]. Study of the effect of cosmological constant on local phenomena such as null geodesics, time delay of light [7]–[8] and the perihelion precession [9]–[12], has attracted considerable attention in the last decades. Many authors have investigated the effects of cosmological constant on the bending of light. Islam was the first to mention that the null geodesic equation in a spherically symmetric space-time does not involve  $\Lambda$  term and concluded that the cosmological constant does not affect the bending of light [13]. However, later Rindler and Ishak [14], by considering the intrinsic properties of the Schwarzschild-de Sitter space-time proposed a new method for calculating the deflection angle of light. This approach gives a new insight into the effect of cosmological constant on bending of light and, as they put it, “ $\Lambda$  does contribute to the observed bending of light.” Since the cosmological constant effectively counteracts gravity, one may intuitively expect that a positive  $\Lambda$  would diminish the bending of light. Different aspects of the method such as integration of the gravitational potentials and Fermats principle have been studied in [17]. The studies of light bending in Kerr-de Sitter and Reissner-Nordstrom-de Sitter space-times, using Rindler-Ishak method, have been carried out in [15] and [16], respectively. Also the method has been applied to galactic halos, of which an excellent example is the Mannheim-Kazanas solution of conformal Weyl gravity [18]–[20]. For further information on the use of Rindler-Ishak method, see [21]–[25].

The low energy limit of string theory contains quadratic and higher-order curvature invariants in addition to the usual scalar curvature in the Lagrangian. One specific combination of higher-order

---

\*Electronic address: m\_heydarifard@sbu.ac.ir

†Electronic address: heydarifard@qom.ac.ir

‡Electronic address: hr-sepangi@sbu.ac.ir

curvature invariants is the Gauss-Bonnet term which is a natural extension of Einsteins general relativity in a  $D$ -dimensional space-time with  $D - 4$  extra dimensions. For  $D = 4$  the Gauss-Bonnet (GB) term is a topological invariant and does not contribute to the gravitational field equations. However, recently a novel four-dimensional Einstein-Gauss-Bonnet (EGB) gravity is proposed by Glavan and Lin [26] where, by re-scaling the coupling constant  $\alpha \rightarrow \frac{\alpha}{D-4}$ , and in the limit  $D \rightarrow 4$ , the GB term does contribute to the gravitational dynamics, thus circumventing the conditions of Lovelocks theorem. The theory preserves the number of degrees of freedom and avoids the Ostrogradsky instability. Also, in contrast to general relativity, this theory is free from singularity problems. It should be noted at this point that although in the framework of semi-classical gravity with conformal anomaly a  $4D$  black hole solution similar to that of the EGB was initially found in [27], the EGB theory is a classical modified gravity on equal footing with general relativity. Shortly afterwards, black hole solutions and their extensions to charged and rotating black holes were studied and carried out in [28] and [29], respectively. In addition, a general study of solutions in Lovelock gravity has been presented in [30]–[31]. Also, an exact charged black hole solution of the theory surrounded by clouds of strings was obtained in [32]. The non-static solutions of radiating black holes as well as the Hayward and bardeen black hole solutions were constructed in [33]–[36], respectively. Exact spherically symmetric wormhole solutions for an isotropic and anisotropic matter sources, and thin-shell wormhole solutions are constructed, respectively in [37] and [38]. The authors in [39] have investigated the structure of relativistic stars in this theory and discussed observational constraints on the GB coupling  $\alpha$ . The study of geodesic motion and the effect of GB coupling on the shadow cast by black holes in four-dimensional Einstein-Gauss-Bonnet (EGB) gravity have been considered in [40] and [41]. Also, quasinormal modes of black holes in the framework of EGB gravity have been studied in [42]. In the strong deflection limit, gravitational lensing by static and spherically symmetric black holes of the theory have been investigated in [43]. The analysis of gravitational instability of asymptotically flat, de-Sitter and anti-de Sitter black holes in four-dimensional EGB gravity were carried out in [44]. Also, phase transition and thermodynamical behavior of such black holes have been investigated in [45]–[47]. For other studies on four-dimensional EGB gravity see [48]–[56].

$$\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \quad (1)$$

The purpose of this note is to investigate the problem of light bending in EGB black holes by Rindler-Ishak method. We investigate possible corrections resulting from GB coupling  $\alpha$  on the deflection angle of light. Conclusions are drawn in section 3.

## 2 Bending angle in $4D$ Einstein-Gauss-Bonnet-de Sitter black hole

Bending of light in the vicinity of compact gravitational objects is one of the most studied problems in general relativity and was first confirmed by Eddington and Dyson in 1919 [57]. The action of  $4D$  EGB gravity is given by

$$S = \frac{1}{16\pi G} \int \sqrt{-g} d^4x [R + \alpha (R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\lambda\rho}R^{\mu\nu\lambda\rho})], \quad (2)$$

where  $\alpha$  is the GB coupling with dimension of length squared. The static and spherically symmetric solution of the theory is given by [26]

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (3)$$

where

$$f(r) = 1 + \frac{r^2}{2\alpha} \left( 1 \pm \sqrt{1 + \frac{8\alpha M}{r^3}} \right), \quad (4)$$

and  $M$  denotes mass of the black hole. As is mentioned in [26], when  $\alpha$  takes negative values there is no real solution at short radial distances for which  $r^3 < -8\alpha M$ . However, for positive values of  $\alpha$  there are two branch of solutions. The negative branch asymptotically behaves as Schwarzschild solution with positive mass sign, whereas the positive branch behaves as Schwarzschild-de Sitter solution with a negative gravitational mass. Since the universe is in an accelerating expansion phase and the cosmological constant is the simplest candidate for its explanation, it would be interesting to study bending of light in the presence of a  $\Lambda$ -term in the theory. In this case the metric function  $f(r)$  reads [28]

$$f(r) = 1 + \frac{r^2}{2\alpha} \left( 1 \pm \sqrt{1 + \frac{8\alpha M}{r^3} + \frac{4\alpha\Lambda}{3}} \right), \quad (5)$$

where the negative branch asymptotically goes to Schwarzschild-de Sitter solution.

Now, let us obtain the deflection angle of light in EGB gravity. The standard approach for calculating the bending angle is [58]

$$\Delta\phi = 2|\phi(\infty) - \phi(r_0)| - \pi, \quad (6)$$

where  $r_0$  is the closest distance to the black hole. However, the space-time we consider here is not asymptotically flat and  $r \rightarrow \infty$  does not make sense. Therefore, we use Rindler-Ishak method proposed in [14] to obtain the deflection angle in an asymptotically non-flat space-time. Although the null geodesic equation for Schwarzschild-de Sitter space-time does not involve a  $\Lambda$  term, Rindler and Ishak have shown that by considering the effects of cosmological constant on the geometry of space-time one can obtain the contribution of  $\Lambda$  to the bending angle. Using the Euler-Lagrange equations for null geodesics in equatorial plan,  $\theta = \frac{\pi}{2}$ , we obtain the following equation

$$\frac{d^2u}{d\phi^2} + u = \frac{3Mu^2}{\sqrt{1 + 8\alpha Mu^3 + \frac{4\alpha\Lambda}{3}}}, \quad (7)$$

where the variable  $u$  is defined as  $u(\phi) = \frac{1}{r}$ . For small  $\alpha$ , by expanding the right-hand side of the above equation to first-order in parameters  $\alpha$  and  $\Lambda$ <sup>1</sup>, we find

$$\frac{d^2u}{d\phi^2} + u \approx 3Mu^2 - 2M\alpha\Lambda u^2 - 12\alpha M^2 u^5. \quad (8)$$

As one can see, the second term in the above equation involves both the cosmological constant and GB coupling, that is, in this theory the orbital equation of light contains the cosmological constant in contrast to Schwarzschild-de Sitter space-time in general relativity. We solve equation (8) using a perturbative method up to the third order and consider a solution as

$$u = u_0 + \delta u_1 + \delta u_2 + \delta u_3 + \dots, \quad (9)$$

where  $u_0 = \frac{\sin\phi}{R}$  is the undeflected straight line path and  $R$  is the closest distance to the mass, see figure 1. The corrections  $\delta u_1$ ,  $\delta u_2$  and  $\delta u_3$  satisfy the following equations

$$\frac{d^2(\delta u_1)}{d\phi^2} + \delta u_1 = 3mu_0^2, \quad (10)$$

$$\frac{d^2(\delta u_2)}{d\phi^2} + \delta u_2 = 6mu_0u_1, \quad (11)$$

$$\frac{d^2(\delta u_3)}{d\phi^2} + \delta u_3 = 6mu_0u_2 + 3mu_1^2 - 2m\alpha\Lambda u_0^2 - 12\alpha m^2 u_0^5. \quad (12)$$



As one can see, the deflection angle is modified by new terms containing the GB coupling and cosmological constant. The first term is the standard one for bending of light in a Schwarzschild field. In addition to the term  $-\frac{\Lambda R^3}{6M}$  representing the Schwarzschild-de Sitter space-time contribution and initially obtained in [14], the above expression contains the term  $-\frac{71R\Lambda M}{96}$  which is similar to that obtained in [59] and couples the cosmological constant to mass of the lens and has its origin in higher order corrections. Also, we obtain a local term  $\frac{25\pi\Lambda M^2}{512}$  which does not depend on  $R$ . Since all terms in the second bracket are positive, we find that for a positive  $\alpha$  the effects of the GB coupling is to diminish the deflection angle.

The effect of the cosmological constant on deflection angle at small scales such as the solar system is expected to be negligible. Therefore, by setting  $\Lambda = 0$  in equation (19) and keeping the linear term in the expansion, we find

$$2\epsilon \simeq \frac{4M}{R} \left( 1 - \frac{15\pi\alpha M}{16R^3} \right), \quad (20)$$

which only gives the corrections of the GB term to the deflection angle. The observational data on light deflection by the sun, from long baseline radio interferometry [60], gives  $\delta\phi_{LD} = \delta\phi_{LD}^{(GR)}(1 + \Delta_{LD})$ , with  $\Delta_{LD} \leq 0.0002 \pm 0.0008$ , where  $\delta\phi_{LD}^{(GR)} = 1.7510$  arcsec. By assuming that  $\Delta_{LD}$  is entirely due to the geometric effects of the GB term, the observational results constrain the coupling constant  $\alpha$  as  $|\alpha| \leq \frac{16R^3\Delta_{LD}}{15\pi M}$ . Taking for  $R$  and  $M$  values of the radius and mass of the sun,  $R_{\odot} = 6.955 \times 10^8 m$  and  $M_{\odot} = 1.989 \times 10^{30} kg$ , we obtain the constraint  $|\alpha| \leq (1.55 \pm 6.19) \times 10^{19} m^2$ . This result is compatible with that obtained in [61] by solar system measurements on GB gravity.

### 3 Conclusions

In this paper we have studied the bending of light in the space-time of a four-dimensional EGB-de Sitter black hole using the method of Rindler and Ishak. We have obtained an expression for the bending angle which contains the effects of the GB coupling and cosmological constant. We found that for a positive coupling constant  $\alpha$  the effect of GB term is to diminish the bending angle in the novel four-dimensional EGB gravity even up to third order in  $M$ . This is an interesting result in that the GB term as a candidate for dark energy, counteracts gravity. Also, using the observational data on bending of light by the sun, we have constrained the value of the GB coupling.

### References

- [1] S. Weinberg, *Rev. Mod. Phys* **61** (1989) 1.
- [2] M. S. Turner, *Phys. Rep* **619** (2000) 333.
- [3] V. Sahni and A. Starobinsky, *Int. J. Mod. Phys D* **09** (2000) 373.
- [4] T. Padmanabhan, *Phys. Rep*, **380** (2003) 235.
- [5] P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys* **75** (2003) 559.
- [6] A. Upadhye, M. Ishak, and P. J. Steinhardt, *Phys. Rev. D* **72** (2005) 063501.
- [7] B. Chen, R. Kantowski, and X. Dai, *Phys. Rev. D* **82** (2010) 043005.
- [8] K. E. Boudjemaa, M. Guenouche, and S. R. Zouzou, *Gen. Rel. Grav* **43** (2011) 1707.
- [9] R. A. Alpher, *Am. J. Phys* **35** (1967) 771.

- [10] A.W. Kerr, J. C. Hauck, and B. Mashhoon, *Class. Quant. Grav* **20** (2003) 2727.
- [11] N. Cruz, M. Olivares, and J. R. Villanueva, *Class. Quant. Grav* **22** (2005) 1167.
- [12] E. Hackmann and C. Lammerzahl, *Phys. Rev. Lett* **100** (2008) 171101.
- [13] N. J. Islam, *Phys. Lett. A* **97** (1983) 239.
- [14] W. Rindler and M. Ishak, *Phys. Rev. D* **76** (2007) 043006.
- [15] J. Sultana, *Phys. Rev. D* **88** (2013) 042003.
- [16] M. Heydari-Fard, S. Mojahed and S. Y. Rokni *Astrophys Space Sci* **351** (2014) 251.
- [17] M. Ishak, *Phys. Rev. D* **78** (2008) 103006.
- [18] A. Bhattacharya, R. Isaev, M. Scalia, C. Cattani and K. K. Nandi, *JCAP* **09** (2010) 004.
- [19] A. Bhattacharya, G. M. Garipova, E. Laserra, A. Bhadra and K. K. Nandia, *JCAP* **02** (2011) 028.
- [20] C. Cattani, M. Scalia, E. Laserra, I. Bochicchio and K. K. Nandi, *Phys. Rev. D* **87** (2013) 047503.
- [21] G. Farrugia, J. L. Said and M. L. Ruggiero, *Phys. Rev. D* **93** (2016) 104034.
- [22] A. Mishra and S. Chakraborty, *Eur. Phys. J. C* **78** (2018) 374.
- [23] M. Guenouche and S. R. Zouou, *Phys. Rev. D* **98** (2018) 123508.
- [24] M. S. Ali and S. Bhattacharya, *Phys. Rev. D* **97** (2018) 024029.
- [25] M. Haluk Secuk and O. Delice, arXiv: 1908.00509 [gr-qc]
- [26] D. Glavan and C. Lin, *Phys. Rev. Lett* **124** (2020) 081301.
- [27] R. G. Cai, L. M. Cao and N. Ohta, *JHEP* **1004** (2010) 082; R. G. Cai, *Phys. Lett. B*, **733** (2014) 183.
- [28] P. G. S. Fernandes, arXiv:2003.05491 [gr-qc].
- [29] R. Kumar and S. G. Ghosh, arXiv:2003.08927 [gr-qc].
- [30] R. A. Konoplya and A. Zhidenko, *Phys. Rev. D* **101** (2020) 084038.
- [31] A. Casalino, A. Colléaux, M. Rinaldi and S. Vicentini, arXiv:2003.07068 [gr-qc].
- [32] D. V. Singh, S. G. Ghosh and S. D. Maharaj, arXiv:2003.14136 [gr-qc].
- [33] S. G. Ghosh and S. D. Maharaj, arXiv:2003.09841 [gr-qc].
- [34] S. G. Ghosh and R. Kumar, arXiv:2003.12291 [gr-qc].
- [35] A. Kumar and S. G. Ghosh, arXiv:2004.01131 [gr-qc].
- [36] A. Kumar and R. Kumar, arXiv:2003.13104 [gr-qc].
- [37] K. Jusufi, A. Banerjee and S. G. Ghosh, arXiv:2004.10750 [gr-qc].
- [38] P. Liu, C. Niu, X. Wang, C. Y. Zhang, arXiv:2004.14267 [gr-qc].
- [39] D. D. Doneva and S. S. Yazadjiev, arXiv:2003.10284 [gr-qc].

- [40] M. Guo and P. C. Li, arXiv:2003.02523 [gr-qc].
- [41] S. W. Wei and Y. X. Liu, arXiv:2003.07769 [gr-qc]; R. Roy and S. Chakrabarti, arXiv:2003.14107 [gr-qc]
- [42] R. A. Konoplya and A. F. Zinhailo, arXiv:2003.01188 [gr-qc]; M. S. Churilova, arXiv:2004.00513 [gr-qc]; A. Aragón, R. Bécar, P. A. González and Y. Vásquez, 2004.05632 [gr-qc]; S. Devi, R. Roy and S. Chakrabarti, 2004.14935 [gr-qc].
- [43] S. U. Islam, R. Kumar and S. G. Ghosh, arXiv: 2004.01038 [gr-qc]; R. Kumar, S. U. Islam and S. G. Ghosh, arXiv:2004.012970 [gr-qc]; X. H. Jin, Y. X. Gao and D. J. Liu, arXiv: 2004.02261 [gr-qc].
- [44] R. A. Konoplya and A. Zhidenko, arXiv:2003.12492 [gr-qc].
- [45] S. A. Hosseini Mansoori, arXiv: 2003.13382 [gr-qc].
- [46] K. Hegde, A. N. Kumara, C. L. A. Rizwan, K. M. Ajith and M. S. Ali, arXiv:2003.08778 [gr-qc].
- [47] D. V. Singh and S. Siwach, arXiv:2003.11754 [gr-qc].
- [48] C. Y. Zhang, P. C. Li and M. Guo, arXiv:2003.13068 [gr-qc].
- [49] R. A. Konoplya and A. Zhidenko, arXiv:2003.12171 [gr-qc].
- [50] Y. P. Zhang, S. W. Wei and Y. X. Liu, arXiv:2003.10960 [gr-qc].
- [51] H. Lü and Y. Pang, arXiv:2003.11552 [gr-qc].
- [52] C. Liu, T. Zhu and Q. Wu, arXiv:2004.01662 [gr-qc].
- [53] B. Eslam Panah and Kh. Jafarzade, arXiv:2004.04058 [gr-qc].
- [54] X. H. Ge and S. J. Sin, arXiv:2004.12191 [hep-th].
- [55] S. Ying, arXiv:2004.09480 [gr-qc].
- [56] G. Narain and H. Q. Zhang, arXiv:2005.05183 [gr-qc]
- [57] F. W. Dyson, A. S. Eddington and C. Davidson, *philosophical transactions of the royal society of London. series A, containing papers of a mathematical or physical character*, **332** (1920) 291.
- [58] S. Weinberg, *Gravitation and Cosmology*, (John Wiley and Sons, 1972)
- [59] M. Sereno, *Phys. Rev. D* **77** (2008) 043004.
- [60] D. S. Robertson, W. E. Carter, W. H. Dillinger, *Nature* **349** (1991) 768; D.E. Lebach, B.E. Corey, I.I. Shapiro, M.I. Ratner, J.C. Webber, A.E.E. Rogers, J.L. Davis, T.A. Herring, *Phys. Rev. Lett* **75** (1995) 1439.
- [61] L. Amendola, C. Charmousis and S. Davis, *JCAP* **0710** (2007) 004.