

Detecting fibered strongly quasi-positive links

ALBERTO CAVALLO

Max Planck Institute for Mathematics,
Bonn 53111, Germany

cavallo@mpim-bonn.mpg.de

Abstract

We prove that an n -component fibered link L in S^3 is strongly quasi-positive if and only if $\tau(L) = g_3(L) + n - 1$, where $g_3(L)$ denotes the Seifert genus and τ is the Ozsváth-Szabó concordance invariant. We also provide a table which contains a list of some fibered prime links with at most 9 crossings; and we explicitly determine the ones that are strongly quasi-positive and their maximal self-linking number.

1 Introduction

It is a very known result in low dimensional topology that every contact 3-manifold can be presented by an open book decomposition; this is a pair (L, π) where L is a smooth link in M and $\pi : M \setminus L \rightarrow S^1$ is a locally trivial fibration whose fiber's closure is a compact surface F with $\partial F = L$. Such an L is then called a fibered link in M and F is a fibered surface for L .

In this paper we are only interested in the case where the ambient manifold M is the 3-sphere. Fibered links in S^3 possess special properties; in fact, those links have unique fibered surfaces up to isotopy and, once such a surface is fixed, the fibration π is also completely determined. It follows that in S^3 we can define fibered links as the links whose complement fibers over the circle.

Since an open book decomposition (L, π) always determines a contact structure, we obtain a partition of fibered links in S^3 by saying that L carries the structure ξ_L . In particular, it is a consequence of classical results of Harer and Stallings [9, 22], and the Giroux correspondence [8], that the subset of fibered links, inducing the unique tight structure ξ_{st} on S^3 , coincides with the set of strongly quasi-positive links. We recall that a link is said strongly quasi-positive if it can be written as closure of the composition of d -braids of the form

$$(\sigma_i \cdots \sigma_{j-2})\sigma_{j-1}(\sigma_i \cdots \sigma_{j-2})^{-1} \quad \text{for some} \quad d \geq j \geq i + 2 \geq 3$$

or

$$\sigma_i \quad \text{for} \quad i = 1, \dots, d - 1,$$

where $\sigma_1, \dots, \sigma_{d-1}$ are the Artin generators of the d -braids group.

Remark 1.1. Fibered and strongly quasi-positive links in S^3 are connected transverse $\mathbb{C}$ -links in the sense of [3].

The main result of our paper is a criterion for detecting exactly which fibered links in S^3 are strongly quasi-positive, and then carry the contact structure ξ_{st} , generalizing results of Hedden [11] (for knots) and Boileau, Boyer and Gordon [1] (for non-split alternating links). We recall that Ozsváth, Stipsicz

and Szabó defined the τ -set (of 2^{n-1} integers) of an n -component link in [17], using the link Floer homology group $cHFL^-$, and they denote with $\tau_{\max}$ and $\tau_{\min}$ its maximum and minimum. Moreover, the τ -set contains τ , the concordance invariant defined in [2] by the author.

Theorem 1.2. *A fibered link L in S^3 is strongly quasi-positive if and only if $\tau(L) = g_3(L) + n - 1$ and, in this case, one also has $\tau(L) = \tau_{\max}(L)$.*

Combining Theorem 1.2 with [6, Theorem 1.1] and [3, Theorem 1.4] we easily obtain the following corollary.

Corollary 1.3. *For a fibered link L the fact that $\tau(L) = g_3(L) + n - 1$ immediately implies $SL(L) = 2\tau(L) - n$, where $SL(L)$ denotes the maximal self-linking number of L .*

Furthermore, we give a table where we list all prime alternating fibered links with crossing number at most 9, together with the prime non-alternating fibered ones with crossing number at most 7, and we denote whether they are strongly quasi-positive. We plan to expand the table in the future, but unfortunately at the moment we do not have knowledge of any program to compute τ for links.

The paper is organized as follows: in Section 2 we recall the basics properties of link Floer homology and fibered surfaces in the 3-sphere; moreover, we prove Theorem 1.2. Finally, in Section 3 we present our table of small prime fibered links and we explain how to read it.

Funding: The author has a post-doctoral fellowship at the Max Planck Institute for Mathematics in Bonn.

2 Proof of the main theorem

2.1 Link Floer homology

In [21] Ozsváth and Szabó describe how to construct the chain complex $(\widehat{CFL}_*(\mathcal{D}), \hat{\partial})$ from a Heegaard diagram $\mathcal{D} = (\Sigma, \alpha, \beta, \mathbf{w}, \mathbf{z})$, where the sets of basepoints both contain n elements. They show that, under some conditions, the diagram $\mathcal{D}$ represents an n -component link L .

If we ignore the information given by $\mathbf{z}$ then $\mathcal{D}$ is just a (multi-pointed) Heegaard diagram for S^3 and, using the same procedure, we obtain the complex $(\widehat{CF}_*(\mathcal{D}), \hat{\partial})$, see [18]. When $n = 1$ the homology of the latter complex is denoted by $\widehat{HF}_*(S^3) \cong \mathbb{F}_0$, where $\mathbb{F}$ is the field with two elements. In general, we have that

$$H_*(\widehat{CF}(\mathcal{D})) \cong \widehat{HF}_*(S^3) \otimes (\mathbb{F}_{-1} \oplus \mathbb{F}_0)^{\otimes n-1} ;$$

hence, if $\mathcal{D}_1$ and $\mathcal{D}_2$ have the same number of basepoints then $\widehat{CF}_*(\mathcal{D}_1)$ is chain homotopy equivalent to $\widehat{CF}_*(\mathcal{D}_2)$.

The homology of $\widehat{CFL}_*(\mathcal{D})$, following the notation of the author in [2], is denoted by $\widehat{\mathcal{HFL}}_*(L)$, but for what we said before this group is graded isomorphic to $H_*(\widehat{CF}(\mathcal{D}))$. The difference, in the case of links, is given by the fact that the basepoints in $\mathbf{z}$ define the Alexander (collapsed) increasing filtration $\mathcal{A}^s \widehat{CFL}_*(\mathcal{D})$ with $s \in \mathbb{Z}$. Such a filtration descends into homology in the following way: consider the quotient projection $\pi_d : \text{Ker } \hat{\partial}_d \rightarrow \widehat{\mathcal{HFL}}_d(L)$, where $\hat{\partial}_d : \widehat{CFL}_d(\mathcal{D}) \rightarrow \widehat{CFL}_{d-1}(\mathcal{D})$ is the restriction of $\hat{\partial}$ to $\widehat{CFL}_d(\mathcal{D})$, and define

$$\text{Ker } \hat{\partial}_{d,s} = \text{Ker } \hat{\partial}_d \cap \mathcal{A}^s \widehat{CFL}_d(\mathcal{D}) .$$

Then we say that

$$\mathcal{A}^s \widehat{\mathcal{HFL}}_d(L) = \pi_d(\text{Ker } \hat{\partial}_{d,s}) \subset \widehat{\mathcal{HFL}}_d(L)$$

for any $d \in \mathbb{Z}$. It is important to observe that $\mathcal{A}^s \widehat{\mathcal{HFL}}_*(L)$ is not the same homology group obtained by just taking the homology of the Alexander level s of $\widehat{CFL}(\mathcal{D})$, which is instead denoted with $H_*\left(\mathcal{A}^s \widehat{CFL}(\mathcal{D})\right)$; more specifically, the latter group is obtained by first restricting $\widehat{CFL}_d(\mathcal{D})$ and $\widehat{\partial}_d$ to the Alexander level s and then extracting the homology. In particular, note that $\mathcal{A}^s \widehat{\mathcal{HFL}}_*(L)$ is, by definition, always a subgroup of $\widehat{\mathcal{HFL}}_*(L)$, while this needs not be true for $H_*\left(\mathcal{A}^s \widehat{CFL}(\mathcal{D})\right)$.

In addition, we write $\widehat{HFL}_{*,*}(L)$ for the homology of the graded object associated to $\widehat{ACFL}_*(\mathcal{D})$, which is the bigraded complex $(\text{gr}_{*,*}(\mathcal{D}), \text{gr}(\widehat{\partial}))$ defined as follows, see [17]: we have that $\text{gr}_{d,s}(\mathcal{D})$ is the subspace of $\mathcal{A}^s \widehat{CFL}_d(\mathcal{D})$ spanned by generators that are not in $\mathcal{A}^{s-1} \widehat{CFL}_d(\mathcal{D})$, while $\text{gr}(\widehat{\partial})$ is the component of $\widehat{\partial}$ which preserves the Alexander grading, so that $\text{gr}_{d,s}(\widehat{\partial}) := \text{gr}(\widehat{\partial})|_{\text{gr}_{d,s}(\mathcal{D})}: \text{gr}_{d,s}(\mathcal{D}) \rightarrow \text{gr}_{d-1,s}(\mathcal{D})$.

We conclude this subsection by recalling that in [2, Theorem 1.3] the invariant $\tau_{\min}(L)$ was shown by the author to coincide with the minimal integer s such that $\mathcal{A}^s \widehat{\mathcal{HFL}}_*(L)$ is non-zero.

2.2 Fibered surfaces and the Thurston norm

It is a result of Ghiggini [7] and Ni [15] that link Floer homology detects fibered links in the 3-sphere.

Theorem 2.1 (Ghiggini-Ni). *A link $L \hookrightarrow S^3$ with n -components is fibered if and only if $\dim \widehat{HFL}_{*,s_{\text{top}}}(L) = 1$, where s_{top} is the maximal $s \in \mathbb{Z}$ such that the group $\widehat{HFL}_{*,s}(L)$ is non-zero.*

We recall that $\|L\|_T$ denotes the evaluation of the Thurston semi-norm [23] at the homology class represented by the Seifert surfaces of L ; and it is defined as

$$\|L\|_T = o(L) - \max \{ \chi(\Sigma) \},$$

where Σ is a compact and oriented surface in S^3 , such that $\partial\Sigma = L$, and $o(L)$ is the number of unknotted unknots in L .

Hence, it follows from [16] that, for an L as in Theorem 2.1, one has $2s_{\text{top}} = \|L\|_T + n$, assuming that L is not the unknot. Moreover, we have the following result.

Proposition 2.2. *If L is a fibered n -component link in S^3 then $s_{\text{top}} = g_3(L) + n - 1$.*

Proof. The statement is clearly true for the unknot and it is a classical result, see [13, Chapters 4 and 5], that fibered surfaces are always connected and minimize the Thurston norm of L . Therefore, we have

$$s_{\text{top}} = \frac{-\chi(F) + n}{2} = \frac{-(2 - 2g(F) - n) + n}{2} = g(F) + n - 1,$$

where F is a fibered surface for L .

Now, if L admits a Seifert surface T , with $g(T) < g(F)$, then this contradicts the fact that F minimizes $\|L\|_T$ and the claim follows. $\square$

2.3 The contact invariant $c(L)$

In order to prove Theorem 1.2, we need to recall the original definition of the contact invariant $\widehat{c}(\xi) \in \widehat{HF}(S^3)$ from [19] and adapt it to our setting. In [19] Ozsváth and Szabó, for a given fibered knot $K \hookrightarrow S^3$, constructed a Heegaard diagram $\mathcal{D}_K$, representing the mirror image of the knot K , with a distinguished intersection point $c(K)$ such that

$$A(c(K)) = -g_3(K) \quad \text{and} \quad A(y) > -g_3(K)$$

for every other intersection point y in $\mathcal{D}_K$. In particular, this gives that

$$H_* \left(\mathcal{A}^{-g_3(K)} \widehat{CFK}(\mathcal{D}_K) \right) \cong \mathbb{F}$$

and

$$H_* \left(\mathcal{A}^s \widehat{CFK}(\mathcal{D}_K) \right) \cong \{0\} \quad \text{for any } s < -g_3(K).$$

This construction works exactly in the same way for an n -component fibered link L : only that now one has $A(c(L)) = -g$ with $g \in \mathbb{Z}$ and $c(L)$ is still the unique minimal Alexander grading intersection point in $\mathcal{D}_L$.

Lemma 2.3. *Suppose that $L, c(L)$ and g are as before. Then we have that $g = g_3(L) + n - 1$.*

Proof. The element $c(L)$ is necessarily a cycle and $[c(L)]$ has to be non-zero in $\widehat{HFL}(L^*)$; hence, we have that $A(c(L))$ is the minimal integer s such that $\widehat{HFL}_{*,s}(L^*)$ is non-zero. Because of the symmetries of $\widehat{HFL}(L)$, see [20], and Proposition 2.2 this means that $A(c(L)) = -g_3(L) - n + 1 = -g$. $\square$

If we consider the inclusion $i^{L,-g} : \mathcal{A}^{-g} \widehat{CFL}(\mathcal{D}_L) \hookrightarrow \widehat{CF}(\mathcal{D}_L)$ then we can see $c(L)$ as a cycle in $\widehat{CF}(\mathcal{D}_L)$ by dropping the information about the Alexander grading.

The element $c(L)$ can then be identified with the contact element defined in [4], using multi-pointed Legendrian Heegaard diagrams, generalizing the presentation of Honda, Kazez and Matić, see [12]. This means that

$$[c(L)] = \widehat{c}(\xi_L) \otimes \mathbf{e}_{1-n} \in \widehat{HF}(S^3) \otimes (\mathbb{F}_{-1} \oplus \mathbb{F}_0)^{\otimes n-1}, \quad (2.1)$$

where $\mathbf{e}_{1-n}$ is the unique generator of $(\mathbb{F}_{-1} \oplus \mathbb{F}_0)^{\otimes n-1}$ in grading $1 - n$. In particular, we can state the following lemma.

Lemma 2.4. *Suppose that L is a fibered link and ξ_L is the contact structure on S^3 carried by L . Then $c(L)$ is zero in homology whenever ξ_L is overtwisted.*

Proof. From [19] one has $\widehat{c}(\xi_L) = [0]$ when ξ_L is overtwisted. Hence, the homology class $[c(L)]$ is equal to zero for Equation (2.1). $\square$

From the definition of $c(L)$ we observe that $c(L)$ is non-zero in homology if and only if $-g$ is the minimal t such that the map

$$i_*^{L,t} : H_* \left(\mathcal{A}^t \widehat{CFL}(\mathcal{D}_L) \right) \longrightarrow \widehat{\mathcal{HFL}}_*(L^*) \cong \widehat{HF}(S^3) \otimes (\mathbb{F}_{-1} \oplus \mathbb{F}_0)^{\otimes n-1} \quad (2.2)$$

induced by the inclusion $i^{L,t}$ is non-trivial.

Proof of Theorem 1.2. If L is strongly quasi-positive then from [3, Theorem 1.4] one has $\tau(L) = \tau_{\max}(L) = g_3(L) + n - 1$. Suppose now that $\tau_{\max}(L) = g_3(L) + n - 1 = g$, see Lemma 2.3; we want to prove that L is strongly quasi-positive.

We have that

$$-g = -\tau_{\max}(L) = \tau_{\min}(L^*) = \min_{s \in \mathbb{Z}} \left\{ \mathcal{A}^s \widehat{\mathcal{HFL}}(L^*) \text{ is non-zero} \right\} = \min_{s \in \mathbb{Z}} \{ i_*^{L,s} \text{ is non-trivial} \} \quad (2.3)$$

from the definition of τ -set given in [2]. Since L is fibered, the combination of Equations (2.2) and (2.3) tells us precisely that $c(L)$ is non-zero in homology and, applying Lemma 2.4, that $\xi_L = \xi_{\text{st}}$.

At this point, we conclude by recalling that a fibered link L carries the tight structure on S^3 if and only if it is strongly quasi-positive, as we observed in the introduction. $\square$

3 Table of small prime fibered strongly quasi-positive links

Using Theorem 1.2 and some computation we can detect exactly which prime links are fibered and, among these, which ones are strongly quasi-positive. Those data are enclosed in Table 1. We denote the links following the notation on Linkinfo database [14]. Therefore, the number inside brackets identifies the relative orientation of the link. Clearly, being fibered and strongly quasi-positive does depend on relative orientations. Moreover, a link L is fibered if and only if L^* is fibered; while when in Table 1 we write that L is strongly quasi-positive, we mean that at least one between L and L^* is. In fact, the following result holds.

Theorem 3.1. *Suppose that L is a strongly quasi-positive link in S^3 . Then its mirror image L^* is also strongly quasi-positive if and only if L is an unlink.*

Theorem 3.1 follows from a more general result of Hayden [10], but we can prove it directly using only link Floer homology. Let us start from the following lemma.

Lemma 3.2. *If L is strongly quasi-positive then the same is true for L' , where $L = L' \sqcup \bigcirc_{o(L)}$ and $\bigcirc_\ell$ is the ℓ -unlink. Furthermore, one has $\|L\|_T = \|L'\|_T$.*

Proof. The link L bounds a quasi-positive surface Σ . It is known that quasi-positive surfaces always minimize the Thurston norm; hence, since one has $\|L\|_T = \|L'\|_T$ from the additivity of $\|\cdot\|_T$, we have that Σ is the disjoint union of Σ' with some disks, where $\partial\Sigma' = L'$. This implies that L' is strongly quasi-positive. $\square$

At this point, in order to prove Theorem 3.1 we only need Lemma 3.3, which solves Exercise 8.4.9 (a) in [17].

Lemma 3.3. *For every n -component link L we have*

$$0 \leq \tau_{\max}(L) - \tau_{\min}(L) \leq n - 1.$$

Proof. It follows from [5] because $\tau_{\min}(L) = -\tau_{\max}(L^*)$, for the symmetries of $cHFL^-(L)$, and $\tau_{\max}(L)$ is a slice-torus concordance invariant. To show this we can apply [5, Lemma 2.1]; then the only non-trivial fact to prove is that $\tau_{\max}$ coincides with τ for torus links, but this follows from [3, Theorem 1.4]. $\square$

In particular, this result implies that $\tau(L) + \tau(L^*) \leq n - 1$.

Proof of Theorem 3.1. From Lemma 3.2 we can suppose that $o(L) = 0$. By using [3, Theorem 1.4] again, the fact that L and L^* are both strongly quasi-positive gives $2\tau(L) - n = 2\tau(L^*) - n$, since $\|L\|_T = \|L^*\|_T$. Therefore, from Lemma 3.3 one has

$$2\tau(L) \leq n - 1 \quad \text{and then} \quad \|L\|_T \leq -1,$$

but this is impossible. $\square$

The following easier criterion can be used for alternating links. We recall that fibered links are always non-split and we denote by $\sigma(L)$ the signature and by ∇_L the Conway polynomial of L .

Proposition 3.4. *A quasi-alternating n -component link is fibered and strongly quasi-positive if and only if $\nabla_L(z) = z^{-\sigma(L)} + f(z)$, where $f(z)$ has degree strictly smaller than $-\sigma(L)$.*

Proof. It follows from Theorem 2.1 and the facts that $\widehat{HFL}(L)$ categorifies the polynomial $\left(t^{\frac{1}{2}} - t^{-\frac{1}{2}}\right)^{n-1} \cdot \nabla_L\left(t^{\frac{1}{2}} - t^{-\frac{1}{2}}\right)$ and quasi-alternating links are $\widehat{HFL}$ -thin, see [17]. $\square$

This result agrees with [1], where is shown that, for a non-split alternating link L , being strongly quasi-positive is equivalent to being positive and this happens if and only if $2g_3(L) = 1 - n - \sigma(L)$. In fact, we also recall that for such a link L one has $2\tau(L) = n - 1 - \sigma(L)$.

name	strongly QP (SL)	name	strongly QP (SL)	name	strongly QP (SL)
L2a1{0}	Y (0)	L8a10{1}	N	L9a41{0}	N
L2a1{1}	Y (0)	L8a14{0}	Y (6)	L9a42{0}	N
L4a1{1}	Y (2)	L8a15{0,0}	N	L9a42{1}	N
L5a1{0}	N	L8a16{0,0}	N	L9a43{1,0}	N
L5a1{1}	N	L8a16{1,1}	N	L9a43{0,1}	N
L6a1{0}	N	L8a17{1,1}	N	L9a43{1,1}	N
L6a3{0}	Y (4)	L8a18{0,0}	N	L9a44{0,0}	N
L6a4{0,0}	N	L8a18{1,1}	N	L9a44{1,0}	N
L6a4{1,0}	N	L8a19{0,0}	N	L9a44{0,1}	N
L6a4{0,1}	N	L8a19{1,1}	N	L9a46{0,0}	N
L6a4{1,1}	N	L8a20{0,0}	N	L9a46{1,0}	N
L6a5{1,0}	N	L8a20{1,0}	N	L9a46{0,1}	N
L6a5{0,1}	N	L8a21{1,0,0}	N	L9a46{1,1}	N
L6a5{1,1}	N	L8a21{0,1,0}	N	L9a47{1,0}	N
L6n1{0,0}	N	L8a21{0,0,1}	N	L9a47{0,1}	N
L6n1{1,0}	N	L8a21{1,0,1}	N	L9a47{1,1}	N
L6n1{0,1}	Y (3)	L8a21{0,1,1}	N	L9a48{0,0}	N
L6n1{1,1}	N	L8a21{1,1,1}	N	L9a49{1,0}	N
L7a1{0}	N	L9a2{0}	N	L9a49{0,1}	N
L7a1{1}	N	L9a2{1}	N	L9a50{0,0}	N
L7a2{1}	N	L9a5{0}	N	L9a50{1,0}	N
L7a3{0}	N	L9a6{1}	N	L9a50{1,1}	N
L7a3{1}	N	L9a8{0}	N	L9a51{0,0}	N
L7a5{1}	N	L9a8{1}	N	L9a51{1,0}	N
L7a6{0}	N	L9a9{0}	N	L9a51{1,1}	N
L7a7{0,0}	N	L9a9{1}	N	L9a52{1,0}	N
L7a7{1,0}	N	L9a11{0}	N	L9a52{1,1}	N
L7a7{0,1}	N	L9a12{1}	N	L9a53{0,0}	N
L7n1{0}	Y (4)	L9a14{0}	N	L9a53{1,0}	N
L7n1{1}	N	L9a14{1}	N	L9a53{0,1}	N
L7n2{0}	N	L9a16{0}	N	L9a53{1,1}	N
L7n2{1}	N	L9a20{0}	N	L9a54{0,0}	N
L8a1{0}	N	L9a21{0}	N	L9a54{1,0}	N
L8a1{1}	N	L9a22{0}	N	L9a54{0,1}	N
L8a2{0}	N	L9a24{1}	N	L9a54{1,1}	N
L8a2{1}	N	L9a26{1}	N	L9a55{0,0,0}	N
L8a3{1}	N	L9a27{0}	N	L9a55{0,1,0}	N
L8a4{0}	N	L9a28{0}	N	L9a55{0,0,1}	N
L8a4{1}	N	L9a29{0}	N	L9a55{1,0,1}	N
L8a5{0}	N	L9a31{0}	N		
L8a7{1}	N	L9a32{1}	N		
L8a8{0}	N	L9a33{0}	N		
L8a8{1}	N	L9a36{0}	N		
L8a9{0}	N	L9a38{0}	N		
L8a9{1}	N	L9a39{0}	N		

Table 1: Some fibered prime links with 9 or fewer crossings. In the second column we write whether or not the corresponding fibered link (or its mirror image) is strongly quasi-positive. In the latter case, the maximal self-linking number is denoted between brackets.

References

- [1] M. Boileau, S. Boyer and C. Gordon, *Branched covers of quasipositive links and L -spaces*, J. of Topology, **12** (2019), no. 2, pp. 536–576.
- [2] A. Cavallo, *The concordance invariant tau in link grid homology*, Algebr. Geom. Topol., **18** (2018), no. 4, pp. 1917–1951.
- [3] A. Cavallo, *On Bennequin-type inequalities for links in tight contact 3-manifolds*, J. Knot Theory Ramifications, **29** (2020), no. 8, 2050055.
- [4] A. Cavallo, *An invariant of Legendrian and transverse links from open book decompositions of contact 3-manifolds*, Glasgow Math. J., **63** (2021), no. 2, pp. 451–483.
- [5] A. Cavallo and C. Collari, *Slice-torus concordance invariants and Whitehead doubles of links*, Canad. J. Math., **72** (2020), no. 6, pp. 1423–1462.
- [6] J. Etnyre and J. Van Horn-Morris, *Fibered transverse knots and the Bennequin bound*, Int. Math. Res. Not. IMRN, (2011), no. 7, pp. 1483–1509.
- [7] P. Ghiggini, *Knot Floer homology detects genus-one fibred knots*, Amer. J. Math., **130** (2008), no. 5, pp. 1151–1169.
- [8] E. Giroux, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, Proceedings of the International Congress of Mathematicians, Vol. II, Higher Ed. Press, Beijing, 2002, pp. 405–414.
- [9] J. Harer, *How to construct all fibered knots and links*, Topology, **21** (1982), no. 3, pp. 263–280.
- [10] K. Hayden, *Minimal braid representatives of quasipositive links*, Pacific J. Math., **295** (2018), no. 2, pp. 421–427.
- [11] M. Hedden, *Notions of positivity and the Ozsváth-Szabó concordance invariant*, J. Knot Theory Ramifications, **19** (2010), no. 5, pp. 617–629.
- [12] K. Honda, W. Kazez and G. Matić, *On the contact class in Heegaard Floer homology*, J. Differential Geom., **83** (2009), no. 2, pp. 289–311.
- [13] A. Kawauchi, *A survey of knot theory*, Birkhäuser Verlag, Basel, 1996, pp. xxii and 420.
- [14] C. Livingston and A. H. Moore, *KnotInfo: Table of Link Invariants*, <http://www.indiana.edu/~knotinfo>, December 27, 2019.
- [15] Y. Ni, *Knot Floer homology detects fibred knots*, Invent. Math., **170** (2007), no. 3, pp. 577–608.
- [16] Y. Ni, *Link Floer homology detects the Thurston norm*, Geom. Topol., **13** (2009), no. 5, pp. 2991–3019.
- [17] P. Ozsváth, A. Stipsicz and Z. Szabó, *Grid homology for knots and links*, AMS, volume 208 of Mathematical Surveys and Monographs, 2015.
- [18] P. Ozsváth and Z. Szabó, *Holomorphic disks and topological invariants of closed three-manifolds*, Ann. of Math. (2), **159** (2004), no. 3, pp. 1159–1245.
- [19] P. Ozsváth and Z. Szabó, *Heegaard Floer homology and contact structures*, Duke Math. J., **129** (2005), no. 1, pp. 39–61.

- [20] P. Ozsváth and Z. Szabó, *Heegaard diagrams and Floer homology*, International Congress of Mathematicians Vol. II, pp. 1083–1099, Eur. Math. Soc., Zürich, 2006.
- [21] P. Ozsváth and Z. Szabó, *Holomorphic disks, link invariants and the multi-variable Alexander polynomial*, *Algebr. Geom. Topol.*, **8** (2008), no. 2, pp. 615–692.
- [22] J. Stallings, *Constructions of fibred knots and links*, Algebraic and geometric topology, Part 2, pp. 55–60, Proc. Sympos. Pure Math., XXXII, Amer. Math. Soc., Providence, R.I., 1978.
- [23] W. Thurston, *A norm for the homology of 3-manifolds*, *Mem. Amer. Math. Soc.*, **59** (1986), no. 339, pp. i-vi and 99–130.