Green's function for cut points of chordal SLE attached with boundary arcs

Dapeng Zhan*

September 16, 2022

Abstract

A technique of two-curve Green's function is used to study the Green's function of cut points of chordal SLE_{κ} for $\kappa \in (4, 8)$. In order to apply the technique, we take the union of the SLE curve with two open boundary arcs, which share two boundary points other than the endpoints of the SLE curve. The Green's function of interest is, for any z_0 in the domain, the limit as $r \downarrow 0$ of the $r^{-\alpha}$ times the probability that $\{|z - z_0| < r\}$ contains a cut point of the above union, where $\alpha = \frac{3}{8}\kappa - 1$. We prove the limit exists, obtain an exact formula up to a multiplicative constant, and derive a rate of convergence.

1 Introduction and Main Result

The Schramm-Loewner evolution (SLE), first introduced by Oded Schramm in 1999 ([16]), is a one-parameter ($\kappa \in (0, \infty)$) family of measures on plane curves. The SLE with different parameters have been proved to be scaling limits of various lattice models. We refer the reader to Lawler's textbook [5] for basic properties of SLE.

An SLE_{κ} curve is simple for $\kappa \in (0, 4]$, space-filling for $\kappa \in [8, \infty)$, and non-simple and nonspace-filling for $\kappa \in (4, 8)$ (cf. [14]). The Hausdorff dimension of an SLE_{κ} curve is min $\{1 + \frac{\kappa}{8}, 2\}$ (cf. [14, 2]). Suppose $\kappa \in (0, 8)$. Then the probability that a chordal SLE_{κ} curve η exactly passes through any fixed interior point z_0 in the domain is zero. People were interested in the decay rate of the probability $\mathbb{P}[\operatorname{dist}(z_0, \eta) < r]$ as $r \downarrow 0$. Specifically, it is about the limit $\lim_{r\downarrow 0} r^{-\alpha} \mathbb{P}[\operatorname{dist}(z_0, \eta) < r]$, where $\alpha = 1 - \frac{\kappa}{8} > 0$ equals 2 minus the Hausdorff dimension. The limit is called the Green's function for SLE.

A Green's function problem usually includes two parts: proving the existence of the limit and finding the exact formula of the Green's function. The exact formula of the Green's function of chordal SLE was predicted in [14], and the convergence and the formula were proved later in [6]. The paper [6] also proves the convergence of two-point Green's function, i.e., the limit

^{*}Michigan State University, zhan@math.msu.edu

of the renormalized probability that an SLE_{κ} curve gets close to two distinct points, and uses the one- and two- point Green's functions to prove the existence of Minkowski content of SLE.

Miller and Wu studied (cf. [11]) the Hausdorff dimensions of the sets of double points and cut points of SLE. Among other results, they proved that the Hausdorff dimension of the set of cut points of SLE_{κ} is $3 - \frac{3}{8}\kappa$ for $\kappa \in (4, 8)$. They derived a Green's function type estimate: for a chordal SLE_{κ} curve η and a point z_0 in the same domain,

 $\mathbb{P}[\eta \text{ has a cut point contained in } \{|z - z_0| < r\}] = r^{\alpha_0 + o(1)}, \quad \text{as } r \downarrow 0, \tag{1.1}$

where the exponent α_0 given by

$$\alpha_0 = \frac{3}{8}\kappa - 1 \tag{1.2}$$

equals 2 minus the Hausdorff dimension of the cut-set, and $o(1) \rightarrow 0$ as $r \downarrow 0$.

The intention of the current paper is to improve the estimate (1.1). Let η be a chordal SLE_{κ} curve in a simply connected domain from a_1 to a_2 . We expect that the limit

$$G_{D;a_1,a_2}(z_0) := \lim_{r \downarrow 0} r^{-\alpha_0} \mathbb{P}[\eta \text{ has a cut point contained in } \{|z - z_0| < r\}]$$
(1.3)

should converge to a finite positive number. This is inspired by [4], which studies the one-point and two-point Green's functions for cut points of Brownian motion, and uses them to prove the existence of Minkowski content for the set of cut points.

A new technique was introduced in [20], which studies the Green's function of 2-SLE_{κ} for $\kappa \in (0,8)$. A 2-SLE_{κ} configuration is a pair of random curves (η_1, η_2) in a simply connected domain D connecting four marked boundary points such that when any curve is given, the other curve is a chordal SLE_{κ} in a complement domain of the given curve. The Green's function for such 2-SLE_{κ} at $z_0 \in D$ is the limit $\lim_{r\downarrow 0} r^{-\alpha} \mathbb{P}[\eta_j \cap \{|z - z_0| < r\} \neq \emptyset, j = 1, 2]$, where $\alpha > 0$ is given by $\frac{(12-\kappa)(\kappa+4)}{8\kappa}$. Such a limit turns out to converge to a positive number, whose value can be described using a hypergeometric function.

The main idea in [20] is the following. Assume that $D = \mathbb{D} := \{|z| < 1\}$ and $z_0 = 0$ by conformal invariance. Suppose η_j connects a_j with b_j , j = 1, 2, and a_1, b_1, a_2, b_2 are oriented clockwise. After a reduction we may assume that $b_1 = 1$ and $b_2 = -1$. Then we simultaneously grow η_1 and η_2 respectively from a_1 and a_2 with some random speeds such that at any time t before the process stops,

- (C1) the conformal radius of the remaining domain viewed from 0 is e^{-t} , and
- (C2) 1 and -1 equally split the harmonic measure of the remaining domain viewed from 0.

The process stops at some time T when the two curves together disconnect 0 from any of b_1, b_2 . By Koebe's 1/4 theorem and Beurling's estimate, we then know that for any $t \in [0, T)$, max{dist $(0, \eta_j[0, t]), j = 1, 2$ } $\approx e^{-t}$. After the time T, the two curves will not both get closer to 0, and so max{dist $(0, \eta_j), j = 1, 2$ $\approx e^{-T}$.

Thus, the original limit problem is transformed into finding the decay rate of $\mathbb{P}[T > s]$ as $s = -\log(r) \to \infty$. For any $t \in [0, T)$, one may find a conformal map, which maps the

remaining domain at t conformally onto \mathbb{D} , and fixes 0, 1, -1. Then $\eta_1(t)$ and $\eta_2(t)$ are mapped to $e^{iZ_1(t)}$ and $-e^{iZ_2(t)}$ for some $Z_1(t), Z_2(t) \in (0, \pi)$. The (Z_1, Z_2) is a two-dimensional diffusion process with lifetime T. The explicit transition density of this process was derived in [20] using Itô's calculus and orthogonal polynomials of two variables, which implies the key estimate that there is a function G on $(0, \pi)^2$ such that if (Z_1, Z_2) starts from (z_1, z_2) , then P[T > s] = $G(z_1, z_2)e^{-\alpha_1 s}(1 + O(e^{-\beta s}))$ for some constant $\beta > 0$. Such an estimate was then used to prove the convergence of the Green's function.

The two-curve technique in [20] was later used in [19] to study the existence of Green's function for 2-SLE_{κ} at a boundary point. The results of [20, 19] may serve as the first few steps in proving the existence of Minkowski content of double points of SLE_{κ}.

In the current paper, we are going to apply the two-curve technique to study the cut point Green's function. Let D be a Jordan domain, and $a_1 \neq a_2 \in \partial D$. Let $\kappa \in (4, 8)$, and γ_1 be a chordal SLE_{κ} curve in D from a_1 to a_2 . In order to apply the technique, we introduce two other points $b_1, b_2 \in \partial D$ such that b_j lies on the open boundary arc of D from a_j to a_{3-j} in the clockwise direction. Let A_j be the open boundary arc of D with end points b_1, b_2 that contains $a_j, j = 1, 2$. Instead of considering the Green's function for cut points of γ_1 , we study the Green's function for the cut points of the union $\gamma_1 \cup A_1 \cup A_2$. This is a compromise we have to make now in order to apply the two-curve technique in [20]. Our main theorem is the following.

Theorem 1.1. Let $z_0 \in D$ and $R = \text{dist}(z_0, D^c)$. Let f be a conformal map from D onto \mathbb{D} such that $f(z_0) = 0$. Let $w_j, v_j \in \mathbb{R}$ be such that $e^{iw_j} = f(a_j)$ and $e^{iv_j} = f(b_j)$, j = 1, 2. Let α_0 be as in (1.2). Define $G_{D;a_1,b_1,a_2,b_2}(z_0) = |f'(z_0)|^{\alpha_0} \widetilde{G}(w_1, v_1, w_2, v_2)$, where

$$\widetilde{G}(w_1, v_1, w_2, v_2) := \left| \sin\left(\frac{w_1 - w_2}{2}\right) \right|^{\frac{8}{\kappa} - 1} \left| \sin\left(\frac{v_1 - v_2}{2}\right) \right|^{\frac{(\kappa - 4)^2}{2\kappa}} \prod_{j=1}^2 \prod_{s=1}^2 \left| \sin\left(\frac{w_j - v_s}{2}\right) \right|^{1 - \frac{4}{\kappa}}.$$
(1.4)

Let $\beta_0 = 1 - \frac{8}{5\kappa}$. Then there is a positive constant C_0 depending only on κ such that

 $\mathbb{P}[\gamma_1 \cup A_1 \cup A_2 \text{ has a cut point contained in } \{|z - z_0| < r\}]$

$$= C_0 G_{D;a_1,b_1,a_2,b_2}(z_0) r^{\alpha_0} (1 + O(r/R)^{\beta_0}), \quad \text{as } r/R \downarrow 0.$$

Here the implicit constants in the $O(\cdot)$ symbol depend only on κ .

We will basically follow the approach in [20]. First suppose $D = \mathbb{D}$, $z_0 = 0$, $b_1 = 1$, and $b_2 = -1$. Let γ_2 be a time-reversal of γ_1 , which by reversibility of SLE_{κ} (cf. [9]) is a chordal SLE_{κ} in \mathbb{D} from a_2 to a_1 . We then grow γ_1 and γ_2 simultaneously respectively from a_1 and a_2 with random speeds such that at any time t before the first time T that $A_1 \cup \eta_1[0,T]$ intersects $A_2 \cup \eta_2[0,T]$, Conditions (C1) and (C2) are both satisfied. By mapping the remaining domain conformally onto \mathbb{D} with 0, 1, -1 fixed, we obtain a two-dimensional diffusion process $(Z_1(t), Z_2(t))$ taking values in $(0, \pi)^2$. We then use orthogonal polynomials to derive the transition density of (Z_1, Z_2) , and use it to prove Theorem 1.1. This paper can be viewed as the first step of proving the existence of Minkowski content of cut points of SLE. For that purpose, following the approach of [6], we will also need the ordered two-point Green's function, i.e., for $z_1 \neq z_2 \in D$, the limit

$$\lim_{r_1, r_2 \downarrow 0} r_1^{-\alpha_0} r_2^{-\alpha_0} \mathbb{P}[\gamma_1 \text{ first visits a cut point of } \gamma_1 \cup A_1 \cup A_2 \text{ in } \{|z - z_1| < r_1\}$$

and then visits a cut point of $\gamma_1 \cup A_1 \cup A_2$ in $\{|z - z_2| < r_2\}$.

The first result on two-point Green's functions was derived in [8], where the ordered twopoint Green's function for a chordal SLE_{κ} curve γ_1 in D from a_1 to a_2 is expressed as

$$G_{D;a_1,a_2}(z_1) \cdot \mathbb{E}[G_{D_{\gamma_*};z_1,a_2}(z_2)]_{\gamma_*}$$

where $G_{D;a_1,a_2}$ is the one-point Green's function for γ_1 , D_{γ_*} is the complement domain of a random curve γ_* in D whose boundary contains a_2 , where γ_* is the first arm of a two-sided radial SLE_{κ} curve in D from a_1 to a_2 via z_1 , which is also a radial SLE_{κ}(2) curve in D from a_1 to z_1 with the force point at a_2 , and the expectation is with respect to the randomness of γ_* . Here the SLE_{κ}(2) curve γ_* can be intuitively viewed as the chordal SLE_{κ} curve γ_1 conditioned on the singular event that it passes through z_1 , and z_1 is a boundary point of D_{γ_*} .

Thus, we expect that the ordered two-point Green's function for cut points of $\gamma_1 \cup A_1 \cup A_2$ can be expressed as the product

$$G_{D;a_1,b_1,a_2,b_2}(z_1) \cdot \mathbb{E}[G_{D_{\gamma_*};z_1,b_1,a_2,b_2}(z_2)],$$

where $G_{D;a_1,b_1,a_2,b_2}$ is the (one-point) Green's function as in Theorem 1.1, D_{γ_*} is the complement domain of a random curve γ_* in D whose boundary contains a_2 , where γ_* is an $\text{SLE}_{\kappa}(2, \kappa - 4, \kappa - 4)$ curve in D from a_1 to z_1 with force points at a_2, b_1, b_2 . We will see later in Section 3.2.3 that such an $\text{SLE}_{\kappa}(2, \kappa - 4, \kappa - 4)$ curve γ_* can be intuitively viewed as the chordal SLE_{κ} curve γ_1 conditioned on the singular event that $\gamma_1 \cup A_1 \cup A_2$ has a cut point at z_1 .

One may also consider the boundary Green's function for cut points of SLE. This is the limit of (1.3) in the case that z_0 is a boundary point. Here α_0 should be replaced by another exponent α_1 , and ∂D is assumed to be analytic near z_0 . Similarly as in [8], the boundary Green's function for cut points may be used to study the two-point cut point Green's function. While the existence of the boundary Green's function for cut points is beyond reach now, the technique of this paper may be used to prove the convergence of the limit:

$$\lim_{r \downarrow 0} r^{-1} \mathbb{P}[\gamma_1 \cup A_1 \cup A_2 \text{ has a cut point contained in } \{|z - b_1| < r\}],$$

where b_1, b_2, A_1, A_2 are as before. This means that we set $b_1 = z_0$, and take $\alpha_1 = 1$.

Theorem 1.1 does not immediately imply the convergence of the original limit (1.3), which motivated this paper. We made the problem easier by adding two marked boundary points. One approach to attack the original problem is to add two more marked boundary points, which means that the attached boundary arcs do not share end points. In that case the formula of the Green's function is expected to be a multivariable special function.

We believe that the limit (1.3) does converge. We may consider the simple case that $D = \mathbb{H} = \{ \operatorname{Im} z > 0 \}, a_1 = 0 \text{ and } a_2 = \infty$. By the scaling property and reflection symmetry of SLE, there is a positive function $S(\theta)$ on $(0,\pi)$, which is symmetric about $\pi/2$, such that $G_{\mathbb{H};0,\infty}(z) = |z|^{-\alpha_0} S(\arg z)$. To get the explicit formula of G, we need to find out the formula of S. The boundary exponent $\alpha_1 = 1$ suggests that $\lim_{\theta \downarrow 0} S(\theta)/\theta$ is finite and positive.

Below is a sketch of the rest of the paper. In Section 2, we review Loewner equations and two-time-parameter stochastic processes. In Section 3, we recall the setup and results from [20, Sections 3,4,5] with suitable modifications to fit the setting here. We then prove Theorem 1.1 in Section 4. In the appendix, we give a detailed proof of a two directional domain Markov property for chordal SLE_{κ} which satisfies reversibility.

2 Preliminary

Let $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ and $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$. Let J(z) = -1/z. By $f : D \xrightarrow{\text{Conf}} E$ we mean that f maps D conformally onto E. Let $\cot_2, \sin_2, \cos_2, \tan_2$ denote the functions $\cot(\cdot/2), \sin(\cdot/2), \cos(\cdot/2), \tan(\cdot/2), \operatorname{respectively}$.

2.1 Prime ends

For the sake of rigor, we will use the notion of prime ends (cf. [1]). We prefer the description used in [25]. Let D be a simply connected domain. Consider the set of pairs (f,ζ) , where $f: D \xrightarrow{\text{Conf}} \mathbb{D}$, and $\zeta \in \overline{\mathbb{D}}$. Such a set is nonempty by Riemann mapping theorem. Define an equivalence relation "~" on this set such that $(f_1,\zeta_1) \sim (f_2,\zeta_2)$ if (the continuation of) $f_2 \circ f_1^{-1}$ maps ζ_1 to ζ_2 . The set of (f,ζ) modulo the equivalence relation is called the prime end closure of D, which is equipped with a unique metrizable topology such that for any $f: D \xrightarrow{\text{Conf}} \mathbb{D}$, the map $[(f,\zeta)] \mapsto \zeta$ is a homeomorphism from \widehat{D} onto $\overline{\mathbb{D}}$. Note that $z \mapsto [(f,f(z))]$ is an embedding from D into \widehat{D} . We identify D as an open subset of \widehat{D} by identifying $z \in D$ with [(f,f(z))], which does not depend on the choice of f. We call $\widehat{\partial}D := \widehat{D} \setminus D$ the prime end boundary of D, and every $p \in \widehat{\partial}D$ a prime end of D. If $g: D_1 \xrightarrow{\text{Conf}} D_2$, then g induces a homeomorphism $g: [(f,\zeta)] \mapsto [(f \circ g^{-1},\zeta)]$ from \widehat{D}_1 onto \widehat{D}_2 , which maps $\widehat{\partial}D_1$ onto $\widehat{\partial}D_2$.

Let $\overline{D}^{\#}$ be the closure of D in the Riemann sphere $\widehat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$, and $\partial^{\#}D = \overline{D}^{\#} \setminus D$. Let $\zeta \in \partial^{\#}D$ (a point in $\widehat{\mathbb{C}}$) and $p \in \widehat{\partial}D$ (a prime end of D). If $z \in D$ and $z \to p$ (w.r.t. the topology of \widehat{D}) implies that $z \to \zeta$ (w.r.t. the metric of $\widehat{\mathbb{C}}$), then we say that ζ is determined by p. On the other hand, if $z \in D$ and $z \to \zeta$ implies that $z \to \gamma$, then we say that ζ determines p. If p and ζ determine each other, then we do not distinguish them. For example, if D is a Jordan domain in $\widehat{\mathbb{D}}$ such as \mathbb{D} or \mathbb{H} , then every $p \in \widehat{D}$ is identified with a point $\zeta \in \partial^{\#}D$.

Suppose $D \subset D$ are two simply connected domains. Let p be a prime end of D. If D is a neighborhood of p in D, i.e., the intersection of D and a neighborhood of p in \hat{D} , then by

Schewarz reflection principle, there is a unique prime end \tilde{p} of \tilde{D} such that for $z \in \tilde{D}, z \to p$ in D iff $z \to \tilde{p}$ in \tilde{D} . When this happens, we do not distinguish \tilde{p} from p, and say that D and \tilde{D} share the prime end p.

2.2 Chordal Loewner equations

A relatively closed subset K of \mathbb{H} is called an \mathbb{H} -hull if K is bounded and $\mathbb{H} \setminus K$ is a simply connected domain, and $\mathbb{H} \setminus K$ is called an \mathbb{H} -domain. For an \mathbb{H} -hull K, there is a unique $g_K : \mathbb{H} \setminus K \xrightarrow{\text{Conf}} \mathbb{H}$ such that $g_K(z) = z + \frac{c}{z} + O(1/z^2)$ as $z \to \infty$ for some $c \ge 0$. The constant c, denoted by hcap(K), is called the \mathbb{H} -capacity of K, which is zero iff $K = \emptyset$. We write hcap₂ for hcap $(\cdot/2)$. For a set $S \subset \mathbb{C}$, if there is an \mathbb{H} -hull K such that $\mathbb{H} \setminus K$ is the unbounded connected component of $\mathbb{H} \setminus \overline{S}$, then we say that K is the \mathbb{H} -hull generated by S, and write K = Hull(S). In this case we write hcap(S) and hcap₂(S) respectively for hcap(K) and hcap₂(K).

Let $\widehat{w} \in C([0,T),\mathbb{R})$ for some $T \in (0,\infty]$. The chordal Loewner equation driven by \widehat{w} is

$$\partial_t g_t(z) = \frac{2}{g_t(z) - \widehat{w}(t)}, \quad 0 \le t < T; \quad g_0(z) = z.$$

For a fixed $z \in \mathbb{C}$, this is an ODE in t with initial value at 0. Let $\tau_z \in (0, \infty]$ be such that $[0, \tau_z)$ is the maximal interval of the solution. For $t \in [0, T)$, let $K_t = \{z \in \mathbb{H} : \tau_z \leq t\}$. It turns out that each K_t is an \mathbb{H} -hull with hcap₂ $(K_t) = t$, and each g_t agrees with the g_{K_t} associated with K_t . We call g_t and K_t the chordal Loewner maps and hulls, respectively, driven by \hat{w} .

If for every $t \in [0,T)$, g_t^{-1} extends continuously to $\overline{\mathbb{H}}$, and $\eta(t) := g_t^{-1}(\widehat{w}(t))$, $0 \leq t < T$, is a continuous curve, then we say that η is the chordal Loewner curve driven by \widehat{w} . Such η may not exist in general. When it exists, we have $\eta(0) = \widehat{w}(0) \in \mathbb{R}$, $\eta(t) \in \overline{K_t} \subset \overline{\mathbb{H}}$ and $K_t = \operatorname{Hull}(\eta[0,t])$ for all $0 \leq t < T$. On the other hand, if there is a continuous curve η such that $K_t = \operatorname{Hull}(\eta[0,t])$ for each $0 \leq t < T$, then η is the chordal Loewner curve driven by \widehat{w} . For every $t_0 \in [0,T)$, the set $S_{t_0} := (t_0,T) \cap \eta^{-1}(\mathbb{H} \setminus K_{t_0})$ is dense in $[t_0,T)$, and as $t \to t_0$ along $S_{t_0}, g_{t_0}(\eta(t)) \to \widehat{w}(t_0)$, and so $\eta(t)$ tends to the prime end $g_t^{-1}(\widehat{w}(t_0))$ of $\mathbb{H} \setminus K_{t_0}$. When there is no ambiguity, we also use $\eta(t)$ to denote the prime end $g_t^{-1}(\widehat{w}(t))$ of $\mathbb{H} \setminus K_t$.

For $\kappa \in (0, \infty)$, chordal SLE_{κ} is defined by solving the chordal Loewner equation with the driving function being $\hat{w}(t) = \sqrt{\kappa}B(t)$, $0 \leq t < \infty$, where *B* is a standard linear Brownian motion. It is known (cf. [14, 7]) that a.s. \hat{w} generates a chordal Loewner curve η , which satisfies $\eta(0) = \hat{w}(0) = 0$ and $\lim_{t\to\infty} \eta(t) = \infty$. Such η is called a chordal SLE_{κ} curve in \mathbb{H} from 0 to ∞ , or a standard chordal SLE_{κ}. If *D* is a simply connected domain with two distinct prime ends *a*, *b*, then there is a conformal map *f* from \mathbb{H} onto *D* such that f(0) = a and $f(\infty) = b$. Then the *f*-image of a standard chordal SLE_{κ}, which is a continuous curve in the prime end closure of *D*, is called an SLE_{κ} curve in *D* from *a* to *b*. Whenever $\partial^{\#}D$ is locally connected, the chordal SLE_{κ} is also a curve in the spherical closure of *D*.

Chordal SLE_{κ} satisfies the following DMP (domain Markov property, cf. [5]). Let η be a chordal SLE_{κ} curve in D from a to b. Let \mathcal{A} be a σ -algebra independent of η , and $\mathcal{F}^{\mathcal{A}}$ be the usual (complete and right continuous) augmentation of the filtration $(\sigma(\eta_1(s):s \leq t) \lor \mathcal{A})_{t\geq 0}$.

Then for any $\mathcal{F}^{\mathcal{A}}$ -stopping time T, conditionally on $\mathcal{F}_{T}^{\mathcal{A}}$ and the event $\{T < \infty\}$, $\eta(T + \cdot)$ is a chordal SLE_{κ} curve from (the prime end) $\eta(T)$ to b in the connected component of $D \setminus \eta[0,T]$ which shares the prime end b with D. This follows easily from the Markov property of the (rescaled) Brownian motion $\sqrt{\kappa}B_t$.

For $\kappa \in (0, 8]$, chordal SLE_{κ} satisfies reversibility (cf. [7] for $\kappa = 8$, [24] for $\kappa \in (0, 4]$, and [9] for $\kappa \in (4, 8)$). This means that, if η is a chordal SLE_{κ} curve in D from a to b, then any time-reversal of η is a time-change of a chordal SLE_{κ} curve in D from b to a.

2.3 Radial Loewner equations

A relatively closed subset K of \mathbb{D} is called a \mathbb{D} -hull if $\mathbb{D} \setminus K$ is a simply connected domain that contains 0. For an \mathbb{D} -hull K, there is a unique conformal map g_K from $\mathbb{D} \setminus K$ onto \mathbb{D} such that $g_K(0) = 0$ and $g'_K(0) \ge 1$. The constant $\log |g'_K(0)| \ge 0$ is called the \mathbb{D} -capacity of K and is denoted by dcap(K). For a set $S \subset \mathbb{C}$, if there is a \mathbb{D} -hull K such that $D \setminus K$ is the connected component of $\mathbb{D} \setminus \overline{S}$ that contains 0, then we say that K is the \mathbb{D} -hull generated by S, and write $K = \operatorname{Hull}(S)$, dcap $(S) = \operatorname{dcap}(K)$, and $g_S = g_K$.

Let $\widehat{w} \in C([0,T),\mathbb{R})$ for some $T \in (0,\infty]$. The radial Loewner equation driven by \widehat{w} is

$$\partial_t g_t(z) = g_t(z) \frac{e^{i\widehat{w}(t)} + g_t(z)}{e^{i\widehat{w}(t)} - g_t(z)}, \quad 0 \le t < T; \quad g_0(z) = z.$$

For $z \in \mathbb{C}$, let $\tau_z \in (0, \infty]$ be such that $[0, \tau_z)$ is the maximal interval of the solution. For $t \in [0, T)$, let $K_t = \{z \in \mathbb{D} : \tau_z \leq t\}$. It turns out that each K_t is an \mathbb{D} -hull with dcap $(K_t) = t$, and each g_t agrees with the g_{K_t} associated with K_t . We call g_t and K_t the radial Loewner maps and hulls, respectively, driven by \hat{w} . By the Schwarz lemma and Koebe's 1/4 theorem, we have $e^{-t}/4 \leq \operatorname{dist}(0, K_t) \leq e^{-t}$ for 0 < t < T.

If for every $t \in [0,T)$, g_t^{-1} extends continuously to $\overline{\mathbb{D}}$, and $\eta(t) := g_t^{-1}(e^{i\widehat{w}(t)})$, $0 \le t < T$, is a continuous curve, then we say that η is the radial Loewner curve driven by \widehat{w} . Such η may not exist in general. When it exists, we have $\eta(0) = e^{i\widehat{w}(0)} \in \partial \mathbb{D}$, $\eta(t) \in \overline{K_t} \subset \overline{\mathbb{D}}$, and $K_t = \operatorname{Hull}(\eta[0,t])$ for all $0 \le t < T$. On the other hand, if there is a continuous curve η such that $K_t = \operatorname{Hull}(\eta[0,t])$ for each $t \in [0,T)$, then η is the radial Loewner curve driven by \widehat{w} . When there is no ambiguity, we also use $\eta(t)$ to denote the prime end $g_t^{-1}(e^{i\widehat{w}(t)})$ of $\mathbb{D} \setminus K_t$.

We will use covering radial Loewner equations. Let e^i denote the map $z \mapsto e^{iz}$, which maps \mathbb{H} onto $\mathbb{D} \setminus \{0\}$. Let $\widehat{w} \in C([0,T), \mathbb{R})$. The covering radial Loewner equation driven by \widehat{w} is

$$\partial_t \widetilde{g}_t(z) = \cot_2(\widetilde{g}_t(z) - \widehat{w}(t)), \quad \widetilde{g}_0(z) = z.$$

For each $0 \leq t < T$, let $\widetilde{K}_t = \{z \in \mathbb{H} : \widetilde{\tau}_z \leq t\}$, where $\widetilde{\tau}_z$ is such that $[0, \widetilde{\tau}_z)$ is the maximal interval of the solution. We call \widetilde{g}_t and \widetilde{K}_t respectively the covering radial Loewner maps and hulls driven by \widehat{w} . If g_t and K_t are respectively the radial Loewner maps and hulls driven by \widetilde{w}_t , then $\widetilde{K}_t = (e^i)^{-1}(K_t)$ and $e^i \circ \widetilde{g}_t = g_t \circ e^i$ for all $0 \leq t < T$.

If there is a continuous and strictly increasing function u on [0, T) with u(0) = 0, such that $K_{u^{-1}(t)}$ and $g_{u^{-1}(t)}$, $0 \le t < u(T)$, are respectively the radial Loewner hulls and maps driven by

 $\widehat{w} \circ u^{-1}$, then we say that K_t and g_t are respectively radial Loewner hulls and maps with speed du (understood as a measure) driven by \widehat{w} . If u is absolutely continuous and u' = q, we also say that the speed is q. We similarly define radial Loewner curve and covering radial Loewner hulls and maps with some speed. If η is a radial Loewner curve with some speed, then we also understand $\eta(t)$ as the prime end $g_{\eta[0,t]}^{-1}(\widehat{w}(t))$ of $\mathbb{H} \setminus \text{Hull}(\eta[0,t])$.

We now recall radial $\text{SLE}_{\kappa}(\underline{\rho})$ processes, where $\kappa > 0$ and $\underline{\rho} = (\rho_1, \ldots, \rho_m) \in \mathbb{R}^m$ for some $m \in \mathbb{N} \cup \{0\}$. Let $e^{iw}, e^{iv_1}, \ldots, e^{iv_m}$ be distinct points on $\partial \mathbb{D}$. A radial $\text{SLE}_{\kappa}(\underline{\rho})$ curve in \mathbb{D} started from e^{iw} aimed at 0 with force points $e^{iv_1}, \ldots, e^{iv_m}$ is the radial Loewner curve driven by $\hat{w}(t), 0 \leq t < \hat{T}$, which solves the SDE:

$$d\widehat{w}(t) = \sqrt{\kappa} dB_t + \sum_{j=1}^m \frac{\rho_j}{2} \cot_2(\widehat{w}(t) - \widetilde{g}_t^{\widehat{w}}(v_j)) dt, \quad \widehat{w}(0) = w.$$

where *B* is a standard Brownian motion, and $\tilde{g}_t^{\widehat{w}}$ are covering radial Loewner maps driven by \widehat{w} . The process stops whenever the solution $t \mapsto \tilde{g}_t^{\widehat{w}}(v_j)$ blows up for any *j*. The radial $\text{SLE}_{\kappa}(\underline{\rho})$ curve, whose existence follows from the Girsanov Theorem (cf. [15]), starts from e^{iw} , and may or may not end at 0 (depending on κ and ρ). The following proposition is [20, Lemma 3.4].

Proposition 2.1. Let $\kappa > 0$, $n \in \mathbb{N}$. Suppose $\underline{\rho} = (\rho_1, \ldots, \rho_n) \in \mathbb{R}^n$ satisfies $\rho_1, \rho_n \geq \frac{\kappa}{2} - 2$ and $\rho_k \geq 0$, $1 \leq k \leq n$. Let $e^{iw}, e^{iv_1}, \ldots, e^{iv_n}$ be distinct points on \mathbb{T} such that $w > v_1 > \cdots > v_n > w - 2\pi$. Let $\eta(t), 0 \leq t < T$, be a radial $SLE_{\kappa}(\underline{\rho})$ curve in \mathbb{D} started from e^{iw} aimed at 0 with force points $e^{iv_1}, \ldots, e^{iv_n}$. Then a.s. $T = \infty$, 0 is a subsequential limit of $\eta(t)$ as $t \to \infty$, and η does not hit the arc $J := \{e^{i\theta} : v_1 \geq \theta \geq v_n\}$.

2.4 Two-parameter stochastic processes

Now we recall the results in [20, Section 2.3]. We assign a partial order \leq to $\mathbb{R}^2_+ = [0, \infty)^2$ such that $\underline{t} = (t_1, t_2) \leq (s_1, s_2) = \underline{s}$ iff $t_j \leq s_j$, j = 1, 2. The minimal element of \mathbb{R}^2_+ is $\underline{0} = (0, 0)$. We write $\underline{t} < \underline{s}$ if $t_j < s_j$, j = 1, 2. We define $\underline{t} \wedge \underline{s} = (t_1 \wedge s_1, t_2 \wedge s_2)$. Given $\underline{t}, \underline{s} \in \mathbb{R}^2_+$, we define $[\underline{t}, \underline{s}] = \{\underline{r} \in \mathbb{R}^2_+ : \underline{t} \leq \underline{r} \leq \underline{s}\}$. Let $\underline{e}_1 = (1, 0)$ and $\underline{e}_2 = (0, 1)$. So $(t_1, t_2) = t_1\underline{e}_1 + t_2\underline{e}_2$. For a function X defined on a subset U of \mathbb{R}^2_+ , $j \in \{1, 2\}$, and $t_j \in \mathbb{R}_+$, we use $X|_{t_j}^j(t)$ to denote the function $t \mapsto X(t_j\underline{e}_j + t\underline{e}_{3-j})$, whose domain is those $t \in \mathbb{R}_+$ such that $t_j\underline{e}_j + t\underline{e}_{3-j} \in U$.

Definition 2.2. Let $\mathcal{F}_{\underline{t}}, \underline{t} \in \mathbb{R}^2_+$, be a family of σ -algebras on a measurable space Ω such that $\mathcal{F}_{\underline{t}} \subset \mathcal{F}_{\underline{s}}$ whenever $\underline{t} \leq \underline{s}$. Then we call $(\mathcal{F}_{\underline{t}})_{\underline{t} \in \mathbb{R}^2_+}$ an \mathbb{R}^2_+ -indexed filtration on Ω , and simply denote it by \mathcal{F} .

From now on, we let \mathcal{F} be an \mathbb{R}^2_+ -indexed filtration on Ω , and let $\mathcal{F}_{\underline{\infty}} = \vee_{\underline{t} \in \mathbb{R}^2_+} \mathcal{F}_{\underline{t}}$.

Definition 2.3. A random map $\underline{T} : \Omega \to [0,\infty]^2$ is called an \mathcal{F} -stopping time if for any deterministic $\underline{t} \in \mathbb{R}^2_+$, $\{\underline{T} \leq \underline{t}\} \in \mathcal{F}_{\underline{t}}$. We call \underline{T} finite if it takes values in \mathbb{R}^2_+ , and bounded if there is a deterministic $\underline{t} \in \mathbb{R}^2_+$ such that $\underline{T} \leq \underline{t}$. For an \mathcal{F} -stopping time \underline{T} , we define $\mathcal{F}_{\underline{T}} = \{A \in \mathcal{F}_{\underline{\infty}} : A \cap \{\underline{T} \leq \underline{t}\} \in \mathcal{F}_{\underline{t}}, \forall \underline{t} \in \mathbb{R}^2_+\}.$

Definition 2.4. A relatively open subset \mathcal{R} of \mathbb{R}^2_+ is called a history complete region, or simply an HC region, if for any $\underline{t} \in \mathcal{R}$, we have $[\underline{0}, \underline{t}] \subset \mathcal{R}$. Given an HC region \mathcal{R} , for $j \in \{1, 2\}$, define $T_j^{\mathcal{R}} : \mathbb{R}_+ \to \mathbb{R}_+ \cup \{\infty\}$ by $T_j^{\mathcal{R}}(t) = \sup\{s \ge 0 : s\underline{e}_j + t\underline{e}_{3-j} \in \mathcal{R}\}$, where we set $\sup \emptyset = 0$.

The space of HC regions is naturally equipped with a σ -algebra, which is generated by the family $\{\mathcal{R} : \underline{t} \in \mathcal{R}\}, \underline{t} \in \mathbb{R}^2_+$. A map \mathcal{D} from Ω into the space of HC regions is called an \mathcal{F} -stopping region if for any $\underline{t} \in \mathbb{R}^2_+$, $\{\omega \in \Omega : \underline{t} \in \mathcal{D}(\omega)\} \in \mathcal{F}_{\underline{t}}$. Such a map is clearly measurable. A random function $X(\underline{t})$ defined on Ω with a random domain \mathcal{D} is called an \mathcal{F} adapted HC process if \mathcal{D} is an \mathcal{F} -stopping region, and for every $\underline{t} \in \mathbb{R}^2_+$, $X_{\underline{t}}$ restricted to $\{\underline{t} \in \mathcal{D}\}$ is \mathcal{F}_t -measurable.

The following proposition is [20, Lemma 2.7].

Proposition 2.5. Let \underline{T} and \underline{S} be two \mathcal{F} -stopping times. Then (i) $\{\underline{T} \leq \underline{S}\} \in \mathcal{F}_{\underline{S}}$; (ii) if \underline{S} is a constant $\underline{s} \in \mathbb{R}^2_+$, then $\{\underline{T} \leq \underline{S}\} \in \mathcal{F}_{\underline{T}}$; and (iii) if f is an $\mathcal{F}_{\underline{T}}$ -measurable function, then $\mathbf{1}_{\{T \leq S\}}f$ is $\mathcal{F}_{\underline{S}}$ -measurable. In particular, if $\underline{T} \leq \underline{S}$, then $\mathcal{F}_{\underline{T}} \subset \mathcal{F}_{\underline{S}}$.

Definition 2.6. Suppose that there are two \mathbb{R}_+ -indexed filtrations \mathcal{F}^1 and \mathcal{F}^2 such that $\mathcal{F}_{(t_1,t_2)} = \mathcal{F}_{t_1}^1 \vee \mathcal{F}_{t_2}^2$, $(t_1,t_2) \in \mathbb{R}_+^2$. Then we say that \mathcal{F} is a separable filtration generated by \mathcal{F}^1 and \mathcal{F}^2 . For such a separable filtration, if T_j is an \mathcal{F}^j -stopping time, j = 1, 2, then $\underline{T} := (T_1, T_2)$ is called a separable stopping time (w.r.t. \mathcal{F}^1 and \mathcal{F}^2).

The following proposition is [20, Lemma 2.13].

Proposition 2.7. Suppose \mathcal{F} is separable. Let \underline{T} and \underline{S} be two \mathcal{F} -stopping times, where \underline{S} is separable. Then $\{\underline{T} \leq \underline{S}\} \in \mathcal{F}_{\underline{T}}$.

Definition 2.8. The right-continuous augmentation of \mathcal{F} is another \mathbb{R}^2_+ -indexed filtration $\overline{\mathcal{F}}$ defined by $\overline{\mathcal{F}}_{\underline{t}} = \bigcap_{\underline{s} > \underline{t}} \mathcal{F}_{\underline{s}}, \underline{t} \in \mathbb{R}^2_+$. We say that \mathcal{F} is right-continuous if $\overline{\mathcal{F}} = \mathcal{F}$.

The following proposition follows from a standard argument.

Proposition 2.9. The right-continuous augmentation $\overline{\mathcal{F}}$ of \mathcal{F} is right-continuous. An $[0, \infty]^2$ -valued map \underline{T} on Ω is an $\overline{\mathcal{F}}$ -stopping time if and only if $\{\underline{T} < \underline{t}\} \in \mathcal{F}_{\underline{t}}$ for any $\underline{t} \in \mathbb{R}^2_+$; and for such \underline{T} , $A \in \overline{\mathcal{F}}_{\underline{T}}$ if and only if $A \cap \{\underline{T} < \underline{t}\} \in \mathcal{F}_{\underline{t}}$ for any $\underline{t} \in \mathbb{R}^2_+$. If \mathcal{F} is right-continuous, and if $(\underline{T}^n)_{n \in \mathbb{N}}$ is a decreasing sequence of \mathcal{F} -stopping times, then $\underline{T} := \inf_n \underline{T}^n$ is also an \mathcal{F} -stopping time, and $\mathcal{F}_{\underline{T}} = \bigcap_n \mathcal{F}_{\underline{T}^n}$.

Now we fix a probability measure \mathbb{P} on $(\Omega, \mathcal{F}_{\underline{\infty}})$, and let $\mathbb{E}[\cdot|\mathcal{F}_t]$ denote the corresponding conditional expectations.

Definition 2.10. An \mathcal{F} -adapted process $(X_{\underline{t}})_{\underline{t}\in\mathbb{R}^2_+}$ is called an \mathcal{F} -martingale (w.r.t. \mathbb{P}) if for any $\underline{s} \leq \underline{t} \in \mathbb{R}^2_+$, a.s. $\mathbb{E}[X_{\underline{t}}|\mathcal{F}_{\underline{s}}] = X_{\underline{s}}$. If there is $\zeta \in L^1(\Omega, \mathcal{F}_{\underline{\infty}}, \mathbb{P})$ such that a.s. $X_{\underline{t}} = \mathbb{E}[\zeta|\mathcal{F}_{\underline{t}}]$ for all $\underline{t} \in \mathbb{R}^2_+$, then we say that X is closed by ζ .

The following is [20, Lemma 2.11].

Proposition 2.11 (Optional Stopping Theorem). Suppose X is a continuous \mathcal{F} -martingale. Then the following are true. (i) If X is closed by ζ , then for any finite \mathcal{F} -stopping time \underline{T} , $X_T = \mathbb{E}[\zeta|\mathcal{F}_{\underline{T}}]$. (ii) If $\underline{T} \leq \underline{S}$ are two bounded \mathcal{F} -stopping times, then $\mathbb{E}[X_{\underline{S}}|\mathcal{F}_{\underline{T}}] = X_{\underline{T}}$.

3 Ensemble of Two Radial Loewner Chains

3.1 Deterministic ensemble

Let $m \in \mathbb{N} \cup \{0\}$. Let $w_1, w_2, v_1, \ldots, v_m \in \mathbb{R}$ be such that $e^{iw_1}, e^{iw_2}, e^{iv_1}, \ldots, e^{iv_m}$ are pairwise distinct. For j = 1, 2, let $\hat{w}_j \in C([0, \hat{T}_j), \mathbb{R})$ be a radial Loewner driving function with $\hat{w}_j(0) = w_j$. Suppose \hat{w}_j generates radial Loewner hulls $K_j(t)$, radial Loewner maps $g_j(t, \cdot)$, $0 \leq t < \hat{T}_j$. Let \mathcal{D} denote the set of $(t_1, t_2) \in [0, \hat{T}_1) \times [0, \hat{T}_2)$ such that $\overline{K_1(t_1)} \cap \overline{K_2(t_2)} = \emptyset$ and $e^{iv_1}, \ldots, e^{iv_m} \notin \overline{K_1(t_1)} \cup \overline{K_2(t_2)}$. Then \mathcal{D} is an HC region as in Definition 2.4, and we may define functions $T_1^{\mathcal{D}}$ and $T_2^{\mathcal{D}}$. For $(t_1, t_2) \in \mathcal{D}$, let $K(t_1, t_2) = K_1(t_1) \cup K_2(t_2)$. Then $K(t_1, t_2)$ is also an \mathbb{D} -hull. Let $g((t_1, t_2), \cdot) = g_{K(t_1, t_2)}$, and $m(t_1, t_2) = dcap(K(t_1, t_2))$. For $(t_1, t_2) \in \mathcal{D}$ and $j \neq k \in \{1, 2\}$, let $K_{j,t_k}(t_j) = g_k(t_k, K_j(t_j))$, and $g_{j,t_k}(t_j, \cdot) = g_{K_{j,t_k}(t_j)}$. Let $\tilde{K}(t_1, t_2), \tilde{K}_{j,t_k}(t_j) \subset \mathbb{H}$ be the preimages of $K(t_1, t_2), K_{j,t_k}(t_j)$, respectively, under the map e^i . Let $\tilde{g}(\underline{t}, \cdot), \underline{t} \in \mathcal{D}$, be the unique family of maps, such that $\tilde{g}(\underline{t}, z)$ is jointly continuous in $\underline{t}, z, \tilde{g}(\underline{0}, \cdot) = id$, and for each $\underline{t} \in \mathcal{D}$, similarly. Fix $j \neq k \in \{1, 2\}$ and $s \in \{1, \ldots, m\}$. Let $(t_1, t_2) \in \mathcal{D}$. We define the following real valued functions on \mathcal{D} :

$$V_{s}(t_{1}, t_{2}) = \tilde{g}((t_{1}, t_{2}), v_{s}), \quad V_{s,1}(t_{1}, t_{2}) = \tilde{g}'((t_{1}, t_{2}), v_{s});$$
$$W_{j}(t_{1}, t_{2}) = \tilde{g}_{k,t_{j}}(t_{k}, \hat{w}_{j}(t_{j})), \quad W_{j,h}(t_{1}, t_{2}) = \tilde{g}_{k,t_{j}}^{(h)}(t_{k}, \hat{w}_{j}(t_{j}));$$
$$W_{j,S}(t_{1}, t_{2}) = \frac{W_{j,3}(t_{1}, t_{2})}{W_{j,1}(t_{1}, t_{2})} - \frac{3}{2} \left(\frac{W_{j,2}(t_{1}, t_{2})}{W_{j,1}(t_{1}, t_{2})}\right)^{2}.$$

Here the prime and the superscript (h) are respectively the partial derivative and h-th partial derivative w.r.t. the space variable: v_s or $\hat{w}_i(t_i)$.

Let $j \neq k \in \{1, 2\}$ and $s \in \{1, \ldots, m\}$. By [20, Section 3.1], for any $t_k \geq 0$, $K_{j,t_k}(t_j)$ and $g_{j,t_k}(t_j, \cdot), 0 \leq t_j < T_j^{\mathcal{D}}(t_k)$, are radial Loewner hulls and maps, respectively, driven by $W_j|_{t_k}^k$ with speed $(W_{j,1}|_{t_k}^k)^2$. Moreover, we have the following formulas.

$$\partial_{t_j} \mathbf{m} = W_{j,1}^2 \partial t_j; \tag{3.1}$$

$$\partial_{t_j} W_k = W_{j,1}^2 \cot_2(W_k - W_j) \partial t_j, \quad \partial_{t_j} V_s = W_{j,1}^2 \cot_2(V_s - W_j) \partial t_j; \tag{3.2}$$

$$\frac{\partial_{t_j} W_{k,1}}{W_{k,1}} = W_{j,1}^2 \cot_2' (W_k - W_j) \partial t_j, \quad \frac{\partial_{t_j} V_{s,1}}{V_{s,1}} = W_{j,1}^2 \cot_2' (V_s - W_j) \partial t_j; \quad (3.3)$$

$$\partial_{t_j} W_{k,S} = W_{j,1}^2 W_{k,1}^2 \cot_2^{\prime\prime\prime} (W_k - W_j) \partial t_j;$$
(3.4)

$$\partial_{t_j}\widetilde{g}_{k,t_j}(t_k,\widehat{z})|_{\widehat{z}=\widehat{w}_j(t_j)} = -3\widetilde{g}_{k,t_j}''(t_k,\widehat{w}_j(t_j)) = -3W_{j,2};$$
(3.5)

$$\frac{\partial_{t_j} \widetilde{g}'_{k,t_j}(t_k,\widehat{z})|_{\widehat{z}=\widehat{w}_j(t_j)}}{\widetilde{g}'_{k,t_j}(t_k,\widehat{z})|_{\widehat{z}=\widehat{w}_j(t_j)}} = \frac{1}{2} \left(\frac{W_{j,2}}{W_{j,1}}\right)^2 - \frac{4}{3} \frac{W_{j,3}}{W_{j,1}} - \frac{1}{6} (W_{j,1}^2 - 1).$$
(3.6)

Finally, suppose that \widehat{w}_1 and \widehat{w}_2 generate radial Loewner curves η_1 and η_2 , respectively. For each $j \neq k \in (1,2)$ and $\underline{t} = (t_1, t_2) \in \mathcal{D}$, we define $\eta_j^{t_k}(t_j) = g_k(t_k, \eta_j(t_j))$. Then we have $K_{j,t_k}(t_j) = \operatorname{Hull}(\eta_j^{t_k}[0, t_j])$. So for any $t_k \geq 0$, $\eta_j^{t_k}(t_j)$, $0 \leq t_j < T_j^{\mathcal{D}}(t_k)$, is a radial Loewner curve driven by $W_j|_{t_k}^k$ with speed $(W_{j,1}|_{t_k}^k)^2$.

3.2 Commutation couplings

3.2.1 The statement

We use the setup in the previous subsection. We view $(\widehat{w}_1(t))_{0 \leq t < \widehat{T}_1}$ and $(\widehat{w}_2(t))_{0 \leq t < \widehat{T}_2}$ as elements in $\Sigma := \bigcup_{0 < T \leq \infty} C([0,T),\mathbb{R})$. Now suppose \widehat{w}_1 and \widehat{w}_2 are random. Then their laws are probability measures on Σ . Let \mathcal{F}^1 and \mathcal{F}^2 be the \mathbb{R}_+ -indexed filtrations respectively generated by \widehat{w}_1 and \widehat{w}_2 . More specifically, for j = 1, 2 and $t \in \mathbb{R}_+$, \mathcal{F}_t^j is the σ -algebra generated by $\{s < \widehat{T}_i\} \cap \{\widehat{w}(s) \in U\}, 0 \leq s \leq t, U \in \mathcal{B}(\mathbb{R})$. We are going to prove the following.

Proposition 3.1. Let $\kappa \in (0, \infty)$ and $\underline{\rho} = (\rho_1, \ldots, \rho_m) \in \mathbb{R}^m$. We write $e^{i\underline{\nu}}$ for $(e^{i\nu_1}, \ldots, e^{i\nu_m})$. There is a coupling of two random radial Loewner curves $\eta_j(t)$, $0 \leq t < \widehat{T}_j$, j = 1, 2, with radial Loewner driving functions \widehat{w}_j and maps $g_j(t, \cdot)$ such that $\widehat{w}_j(0) = w_j$ and the following holds. For j = 1, 2, η_j is a radial $SLE_{\kappa}(2, \underline{\rho})$ curve in \mathbb{D} started from e^{iw_1} aimed at 0 with force points $(e^{iw_2}, e^{i\underline{\nu}})$. For any $j \neq k \in \{1, 2\}$ and any \mathcal{F}^k -stopping time τ_k with $\tau_k < \widehat{T}_k$, conditionally on $\mathcal{F}^k_{\tau_k}$, the $\eta^{\tau_k}_j(t_j) = g_k(\tau_k, \eta_j(t_j))$, $0 \leq t_j < T^{\mathcal{D}}_j(\tau_k)$, has the law of a timechange of a radial $SLE_{\kappa}(2, \underline{\rho})$ curve in \mathbb{D} started from $g_k(\tau_k, e^{iw_j})$ aimed at 0 with force points $(e^{i\widehat{w}_k(\tau_k)}, g_k(\tau_k, e^{iv_1}), \ldots, g_k(\tau_k, e^{iv_m}))$ stopped at some stopping time.

The η_1 and η_2 in Proposition 3.1 are said to commute with each other. The idea of the commutation relation between SLE curves is generated in [3]. The chordal counterpart of this proposition, with $e^{iw_1}, e^{iw_2}, e^{i\underline{v}}, 0$ replaced by $w_1, w_2, \underline{v}, \infty$, follows easily from the imaginary geometry developed in [10]. In that case, one may construct the two chordal $\text{SLE}_{\kappa}(2, \underline{\rho})$ curves as two flow lines of a GFF in \mathbb{H} with piecewise constant boundary data changing values at $w_1, w_2, v_1, \ldots, v_m$, where the constants are determined by $\kappa, \underline{\rho}$. Similarly, Proposition 3.1 could be probably proved by developing a radial version of imaginary geometry theory. We will not develop such theory here. Instead, we give another proof, which provides us some information of the coupling measure, e.g., Lemma 3.8, that will be needed later. The proof in the special case that m = 2 and $\rho_1 = \rho_2 = 2$ were given in [20, Section 3]. Here we are interested in the case that m = 2 and $\rho_1 = \rho_2 = \kappa - 4$. For the reader's convenience and future reference, we provide a complete proof of Proposition 3.1 here.

For j = 1, 2, let Ξ_j denote the space of simple crosscuts of \mathbb{D} that separate e^{iw_j} from $e^{iw_{3-j}}, e^{iv_1}, \ldots, e^{iv_m}, 0$. For j = 1, 2 and $\xi_j \in \Xi_j$, let $\tau_{\xi_j}^j$ be the first time that η_j hits the closure of ξ_j . Let $\Xi = \{(\xi_1, \xi_2) \in \Xi_1 \times \Xi_2, \operatorname{dist}(\xi_1, \xi_2) > 0\}$. For $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, let $\tau_{\underline{\xi}} = (\tau_{\xi_1}^1, \tau_{\xi_2}^2)$. We may pick a countable subset Ξ^* of Ξ such that for every $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, there is $\underline{\xi}^* = (\xi_1^*, \xi_2^*)$ such that ξ_j^* disconnects ξ_j from 0, j = 1, 2. The significance of Ξ and Ξ^* is that $[\underline{0}, \tau_{\underline{\xi}}] \subset \mathcal{D}$ for any $\underline{\xi} \in \Xi$, and $\mathcal{D} = \bigcup_{\underline{\xi} \in \Xi^*} [\underline{0}, \tau_{\underline{\xi}}]$.

We call the joint law of \hat{w}_1 and \hat{w}_2 in Proposition 3.1 a global commutation coupling. Such a measure is unique, if it exists, because the stated property determines the marginal law of η_2 (taking $\tau_1 = 0$) and the conditional law of η_1 given the part of η_2 up to any \mathcal{F}^2 -stopping time τ_2 that happens before η_2 ends. Let $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$. If η_1, η_2 satisfy the properties in Proposition 3.1 with the following modifications: (i) τ_k is required to be $\leq \tau_{\xi_k}$, and (ii) the time that η_j hits $\eta_k[0, \tau_k]$ is replaced by τ_{ξ_j} , then we call the joint law of the driving functions for η_1, η_2 a local commutation coupling within $\underline{\xi}$. The global commutation coupling is automatically a local commutation coupling within $\underline{\xi}$ for every $\underline{\xi} \in \Xi$. A local commutation coupling within $\underline{\xi}$, when restricted to $\mathcal{F}^1_{\tau_{\xi_1}} \vee \mathcal{F}^2_{\tau_{\xi_2}^2}$, is unique.

3.2.2 Two-variable local martingale

For j = 1, 2, let \mathbb{P}_*^j denote the law of the driving function of a radial $\mathrm{SLE}_{\kappa}(2, \underline{\rho})$ curve in \mathbb{D} started from e^{iw_j} aimed at 0 with force points $(e^{iw_{3-j}}, e^{i\underline{v}})$, which is a probability measure on Σ . Let $\mathbb{P}_*^i = \mathbb{P}_*^1 \times \mathbb{P}_*^2$. The superscript i means "independence". We may prove the existence of the global commutation coupling using the stochastic coupling technique developed in [24, 23]. It suffices to prove that there is a positive continuous process \widehat{M}_* defined on \mathcal{D} with the following properties.

- $\widehat{M}_*(\cdot, 0) = \widehat{M}_*(0, \cdot) \equiv 1.$
- For any $\xi \in \Xi$, $\log(\widehat{M}_*)$ is uniformly bounded on $[\underline{0}, \tau_{\xi}]$.
- For any $\underline{\xi} \in \Xi$, $\mathbb{R}^2_+ \ni \underline{t} \mapsto \widehat{M}_*(\underline{t} \wedge \tau_{\xi})$ is an \mathbb{R}^2_+ -indexed martingale under \mathbb{P}^i_* .
- For any $\underline{\xi} \in \Xi$, the probability measure $\mathbb{P}^{\underline{\xi}}_*$ on Σ^2 defined by $d\mathbb{P}^{\underline{\xi}}_* = \widehat{M}_*(\tau_{\underline{\xi}})d\mathbb{P}^i_*$ is a local commutation coupling within ξ .

We have the following proposition (cf. [24, Sections 6 and 7]).

Proposition 3.2. Whenever the above \widehat{M}_* exists, there exists a global commutation coupling, denoted by \mathbb{P}^c_* , which agrees with $\mathbb{P}^{\underline{\xi}}_*$ on $\mathcal{F}^1_{\tau^1_{\xi_1}} \vee \mathcal{F}^2_{\tau^2_{\xi_2}}$ for any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$. In particular, for any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, \mathbb{P}^c_* is absolutely continuous w.r.t. \mathbb{P}^i_* on $\mathcal{F}^1_{\tau^1_{\xi_1}} \vee \mathcal{F}^2_{\tau^2_{\xi_2}}$, and the Radon-Nikodym derivative is $\widehat{M}_*(\tau_{\xi})$.

For j = 1, 2, let \mathbb{P}_B^j denote the law of the process $w_j + \sqrt{\kappa}B(t)$, $0 \leq t < \infty$, where B is a standard linear Brownian motion. Let $\mathbb{P}_B^i = \mathbb{P}_B^1 \times \mathbb{P}_B^2$. In order to construct \widehat{M}_* , it suffices to find another positive continuous process M_* on \mathcal{D} with the following properties.

- For any $\xi \in \Xi$, $\log(M_*)$ is uniformly bounded on $[\underline{0}, \tau_{\xi}]$.
- For any $\underline{\xi} \in \Xi$, $\mathbb{R}^2_+ \ni \underline{t} \mapsto M_*(\underline{t} \wedge \tau_{\underline{\xi}})$ is an \mathbb{R}^2_+ -indexed martingale under \mathbb{P}^i_B .

• For any $\underline{\xi} \in \Xi$, the probability measure $\mathbb{P}^{\underline{\xi}}_*$ on Σ^2 defined by $d\mathbb{P}^{\underline{\xi}}_* = (M_*(\tau_{\underline{\xi}})/M_*(\underline{0}))d\mathbb{P}^{\mathrm{i}}_B$ is a local commutation coupling within ξ .

If such M_* exists, then for any $j \in \{1,2\}$ and $\xi_j \in \Xi_j$, \mathbb{P}^j_* is absolutely continuous w.r.t. \mathbb{P}^j_B on $\mathcal{F}^j_{\tau^j_{\xi_j}}$, and the Radon-Nikodym derivative is $M_*|_0^{3-j}(\tau^j_{\xi_j})/M_*(\underline{0})$. So for any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, \mathbb{P}^i_* is absolutely continuous w.r.t. \mathbb{P}^i_B on $\mathcal{F}^1_{\tau^1_{\xi_1}} \vee \mathcal{F}^2_{\tau^2_{\xi_2}}$, and the Radon-Nikodym derivative is $\frac{M_*(\tau^1_{\xi_1}, 0)M_*(0, \tau^2_{\xi_2})}{M_*(0, 0)M_*(0, 0)}$. Thus, the \widehat{M}_* defined by $\widehat{M}_*(t_1, t_2) = \frac{M_*(t_1, t_2)M_*(0, 0)}{M_*(t_1, 0)M_*(0, t_2)}$ satisfies the properties stated in the previous paragraph. Moreover, the global commutation coupling \mathbb{P}^c_* is absolutely continuous w.r.t. \mathbb{P}^i_B on $\mathcal{F}^1_{\tau^1_{\xi_1}} \vee \mathcal{F}^2_{\tau^2_{\xi_2}}$ for any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, and the Radon-Nikodym derivative is $M_*(\tau_{\underline{\xi}})/M_*(\underline{0})$. From now on, we are going to construct such M_* .

First suppose (\hat{w}_1, \hat{w}_2) follows the law \mathbb{P}_B^i . Then for two independent Brownian motions B_1 and B_2 , we have $\hat{w}_j(t) = w_j + \sqrt{\kappa}B_j(t), t \ge 0, j = 1, 2$.

Recall the boundary scaling exponent b and central charge c in the literature:

$$\mathbf{b} = \frac{6-\kappa}{2\kappa}, \quad \mathbf{c} = \frac{(3\kappa - 8)(6-\kappa)}{2\kappa}$$

Fix $j \neq k \in \{1,2\}$. Let τ_k be an \mathcal{F}^k -stopping time with $\tau_k < \widehat{T}_k$. Let $\mathcal{F}^{(j,\infty)}$ denote usual augmentation of the \mathbb{R}_+ -indexed filtration $(\mathcal{F}^j_t \vee \mathcal{F}^k_\infty)_{t\geq 0}$. By independence, B_j is an $\mathcal{F}^{(j,\infty)}$ -Brownian motion. From now on, we will repeatedly apply Itô's formula (cf. [15]), where all SDE are $\mathcal{F}^{(j,\infty)}$ -adapted. By (3.5), we have

$$dW_j|_{\tau_k}^k(t) = W_{j,1}|_{\tau_k}^k(t)\sqrt{\kappa}dB_j(t) - \kappa \,\mathrm{b}\,W_{j,2}|_{\tau_k}^k(t)dt_j.$$

To make the symbols less heavy, we write the SDE as

$$\partial_j W_j = W_{j,1} \sqrt{\kappa} \partial B_j - \kappa \, \mathrm{b} \, W_{j,2} \partial t_j. \tag{3.7}$$

The symbol " ∂_j " in the SDE could be understood as the partial derivative w.r.t. the *j*-th variable. We keep in mind that the *k*-th variable is fixed to be τ_k . Using (3.6), we get

$$\frac{\partial_{j} W_{j,1}^{\rm b}}{W_{j,1}^{\rm b}} = \mathrm{b} \, \frac{W_{j,2}}{W_{j,1}} \sqrt{\kappa} \partial B_j + \frac{\mathrm{c}}{6} W_{j,S} \partial t_j - \frac{\mathrm{b}}{6} (W_{j,1}^2 - 1) \partial t_j.$$
(3.8)

Using (3.2,3.3) we get for $r \neq s \in \{1, ..., m\}$,

$$\frac{\partial_j \sin_2(W_k - V_r)^{\frac{\rho_r}{\kappa}}}{\sin_2(W_k - V_r)^{\frac{\rho_r}{\kappa}}} = -\frac{\rho_r}{2\kappa} [1 + \cot_2(W_j - W_k) \cot_2(W_j - V_r)] W_{j,1}^2 \partial t_j;$$
(3.9)

$$\frac{\partial_j \sin_2(V_r - V_s)^{\frac{\rho_r \rho_s}{2\kappa}}}{\sin_2(V_r - V_s)^{\frac{\rho_r \rho_s}{2\kappa}}} = -\frac{\rho_r \rho_s}{4\kappa} [1 + \cot_2(W_j - V_r) \cot_2(W_j - V_s)] W_{j,1}^2 \partial t_j;$$
(3.10)

$$\frac{\partial_j W_{k,1}^{\rm b}}{W_{k,1}^{\rm b}} = -\frac{{\rm b}}{2} [1 + \cot_2(W_j - W_k) \cot_2(W_j - W_k)] W_{j,1}^2 \partial t_j;$$
(3.11)

$$\frac{\partial_j V_{r,1}^{\frac{\rho_r(\rho_r+4-\kappa)}{4\kappa}}}{V_{r,1}^{\frac{\rho_r(\rho_r+4-\kappa)}{4\kappa}}} = -\frac{\rho_r(\rho_r+4-\kappa)}{8\kappa} [1 + \cot_2(W_j - V_r) \cot_2(W_j - V_r)] W_{j,1}^2 \partial t_j;$$
(3.12)

$$\begin{aligned} \frac{\partial_j \sin_2(W_j - W_k)}{\sin_2(W_j - W_k)} &= \frac{\sqrt{\kappa}}{2} \cot_2(W_j - W_k) W_{j,1} \partial B_{\tau_k}^k - \frac{\kappa}{2} \operatorname{b} \cot_2(W_j - W_k) W_{j,2} \partial t_j \\ &\quad + \frac{1}{2} \cot_2(W_j - W_k)^2 W_{j,1}^2 \partial t_j - \frac{\kappa}{8} W_{j,1}^2 \partial t_j; \\ \frac{\partial_j \sin_2(W_j - V_r)}{\sin_2(W_j - V_r)} &= \frac{\sqrt{\kappa}}{2} \cot_2(W_j - V_r) W_{j,1} \partial B_j - \frac{\kappa}{2} \operatorname{b} \cot_2(W_j - V_r) W_{j,2} \partial t_j \\ &\quad + \frac{1}{2} \cot_2(W_j - V_r)^2 W_{j,1}^2 \partial t_j - \frac{\kappa}{8} W_{j,1}^2 \partial t_j. \end{aligned}$$

The last two formulas further imply that

$$\frac{\partial_{j} \sin_{2}(W_{j} - W_{k})^{\frac{\rho_{0}}{\kappa}}}{\sin_{2}(W_{j} - W_{k})^{\frac{\rho_{0}}{\kappa}}} = \frac{\rho_{0}}{2} \cot_{2}(W_{j} - W_{k})W_{j,1}\frac{\partial B_{j}}{\sqrt{\kappa}} - \frac{\rho_{0}}{2} \operatorname{b} \cot_{2}(W_{j} - W_{k})W_{j,2}\partial t_{j} + \frac{\rho_{0}(\rho_{0} + 4 - \kappa)}{8\kappa} \cot_{2}(W_{j} - W_{k})^{2}W_{j,1}^{2}\partial t_{j} - \frac{\rho_{0}}{8}W_{j,1}^{2}\partial t_{j}, \quad \rho_{0} \in \mathbb{R};$$
(3.13)
$$\frac{\partial_{j} \sin_{2}(W_{j} - V_{r})^{\frac{\rho_{r}}{\kappa}}}{\sin_{2}(W_{j} - V_{r})^{\frac{\rho_{r}}{\kappa}}} = \frac{\rho_{r}}{2} \cot_{2}(W_{j} - V_{r})W_{j,1}\frac{\partial B_{j}}{\sqrt{\kappa}} - \frac{\rho_{r}}{2} \operatorname{b} \cot_{2}(W_{j} - V_{r})W_{j,2}\partial t_{j} + \frac{\rho_{r}(\rho_{r} + 4 - \kappa)}{8\kappa} \cot_{2}(W_{j} - V_{r})^{2}W_{j,1}^{2}\partial t_{j} - \frac{\rho_{r}}{8}W_{j,1}^{2}\partial t_{j}.$$
(3.14)

Define a function I on ${\mathcal D}$ by

$$I(t_1, t_2) = \exp\Big(\int_0^{t_1} \int_0^{t_2} W_{1,1}(s_1, s_2)^2 W_{2,1}(s_1, s_2)^2 \cot_2'''(W_1(s_1, s_2) - W_2(s_1, s_2)) ds_2 ds_1\Big).$$

By (3.4) and the fact that $W_{j,S}|_0^k \equiv 0$, we get

$$\frac{\partial_j I^{-\frac{c}{6}}}{I^{-\frac{c}{6}}} = -\frac{c}{6} W_{j,S} \partial t_j.$$
(3.15)

Define $\widehat{\mathbf{m}}$ on \mathcal{D} by $\widehat{\mathbf{m}}(t_1, t_2) = \mathbf{m}(t_1, t_2) - t_1 - t_2$. By (3.1),

$$\partial_j \widehat{\mathbf{m}} = (W_{j,1}^2 - 1)\partial t_j. \tag{3.16}$$

Let $\rho_{\Sigma} = \sum_{s} \rho_{s}$. Define a positive continuous process M_{*} on \mathcal{D} by

$$M_{*} = e^{\frac{(\rho_{\Sigma}+2)(\rho_{\Sigma}+6)}{8\kappa}m}e^{\frac{b}{6}\widehat{m}}W_{1,1}^{b}W_{2,1}^{b}I^{-\frac{c}{6}}|\sin_{2}(W_{1}-W_{2})|^{\frac{2}{\kappa}}\prod_{1\leq r< s\leq m}|\sin_{2}(V_{r}-V_{s})|^{\frac{\rho_{r}\rho_{s}}{2\kappa}} \times \prod_{s\in\{1,...,m\}} \left(V_{s,1}^{\frac{\rho_{s}(\rho_{s}+4-\kappa)}{4\kappa}}\prod_{j\in\{1,2\}}|\sin_{2}(W_{j}-V_{s})|^{\frac{\rho_{s}}{\kappa}}\right).$$
(3.17)

Combining (3.1, 3.8-3.16), where ρ_0 is set to be 2 in (3.13), we get

$$\frac{\partial_j M_*}{M_*} = \mathbf{b} \frac{W_{j,2}}{W_{j,1}} \sqrt{\kappa} \partial B_j + \cot_2(W_j - W_k) W_{j,1} \frac{\partial B_j}{\sqrt{\kappa}} + \sum_r \frac{\rho_r}{2} \cot_2(W_j - V_r) W_{j,1} \frac{\partial B_j}{\sqrt{\kappa}}.$$
 (3.18)

This means that $M_*|_{\mathcal{T}_{\mathbf{r}}}^k(t)$ is a continuous $\mathcal{F}^{(j,\infty)}$ -local martingale.

Lemma 3.3. For any $\underline{\xi} \in \Xi$, $|\log(M_*)|$ is uniformly bounded on $[\underline{0}, \tau_{\underline{\xi}}]$ by a constant depending only on $\kappa, \xi, e^{iw_1}, e^{iw_2}, e^{iv_1}, \dots, e^{iv_m}$.

Proof. It suffices to show that m, $\log |\sin_2(W_1 - W_2)|$, $\log |\sin_2(W_j - V_s)|$, $\log |\sin_2(V_r - V_s)|$, $\log |S_{s,1}|$, $\log |W_{j,1}|$, $j = 1, 2, r \neq s \in \{1, \ldots, m\}$, are all bounded in absolute value on $[\underline{0}, \tau_{\underline{\xi}}]$ by constants depending only on $\underline{\xi}, e^{iw_1}, e^{iw_2}, e^{iv_1}, \ldots, e^{iv_m}$. These statements were all proved in the proof of Lemma 3.1 of [20].

Let $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$. Let $j \neq k \in \{1, 2\}$. Let $\tau_k \leq \tau'_k$ be two \mathcal{F}^k -stopping times such that $\tau_k \leq \tau^k_{\xi_k}$. By Lemma 3.3 and (3.18), $M|_{\tau_k}^k(\cdot \wedge \tau_{\xi_j})$ under \mathbb{P}^i_B is an $\mathcal{F}^{(j,\infty)}$ -martingale closed by $M|_{\tau_k}^k(\tau^j_{\xi_j})$. The filtration $\mathcal{F}^{(j,\infty)}$ in the statement can be replaced by $(\mathcal{F}^j_t \vee \mathcal{F}^k_{\tau'_k})_{t\geq 0}$ since $M|_{\tau_k}^k(\cdot \wedge \tau_{\xi_j})$ is adapted to it. Let \mathcal{F} denote the \mathbb{R}^2_+ -indexed filtration generated by \mathcal{F}^1 and \mathcal{F}^2 . Applying the above result twice: first to k = 1 and $\tau_1 = \tau'_1 = \tau^1_{\xi_1}$, and then to k = 2, $\tau_2 = t_2 \wedge \tau^2_{\xi_2}$ and $\tau'_2 = t_2$, we get the following lemma (cf. [20, Corollary 3.2]).

Lemma 3.4. For any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, $\mathbb{R}^2_+ \ni \underline{t} \mapsto M_*(\underline{t} \wedge \tau_{\underline{\xi}})$ is an \mathcal{F} -martingale under \mathbb{P}^i_B closed by $M_*(\tau_{\xi})$. In particular, we have $\mathbb{E}^i_B[M_*(\tau_{\xi})] = M_*(\underline{0})$.

3.2.3 Construction of the coupling measure

Fix $\underline{\xi} \in \Xi$. We may now define a new probability measure $\widetilde{\mathbb{P}}_*$ by $d\widetilde{\mathbb{P}}_*/d\mathbb{P}_B^{\mathbf{i}} = M_*(\tau_{\underline{\xi}})/M_*(\underline{0})$. Fix $j \neq k \in \{1, 2\}$ and an \mathcal{F}^k -stopping time τ_k with $\tau_k \leq \tau_{\xi_k}^k$. Define

$$\widetilde{B}_{j}(t) = B_{j}(t) - \int_{0}^{t \wedge \tau_{\xi_{j}}^{j}} \frac{2}{2\sqrt{\kappa}} \cot_{2}(W_{j}|_{\tau_{k}}^{k}(s) - W_{k}|_{\tau_{k}}^{k}(s))W_{j,1}|_{\tau_{k}}^{k}(s)^{2}ds$$
$$-\sum_{l=1}^{m} \int_{0}^{t \wedge \tau_{\xi_{j}}^{j}} \frac{\rho_{s}}{2\sqrt{\kappa}} \cot_{2}(W_{j}|_{\tau_{k}}^{k}(s) - V_{l}|_{\tau_{k}}^{k}(s))W_{j,1}|_{\tau_{k}}^{k}(s)^{2}ds, \quad t \ge 0.$$

By (3.18) and the Girsanov Theorem, \widetilde{B}_j is an $\mathcal{F}^{(j,\infty)}$ -Brownian motion under $\widetilde{\mathbb{P}}_*$. By (3.7) and the definition of \widetilde{B} , we find that the W_j (with t_k fixed to be τ_k) satisfies the SDE

$$\partial_j W_j = W_{j,1} \sqrt{\kappa} \partial \widetilde{B} + \cot_2 (W_j - W_k) W_{j,1}^2 \partial t_j + \sum_{s=1}^m \frac{\rho_s}{2} \cot_2 (W_j - V_s) W_{j,1}^2 \partial t_j, \quad 0 \le t_j \le \tau_{\xi_j}^j.$$

Since $\eta_j^{\tau_k}(t_j)$, $0 \leq t_j < T_j^{\mathcal{D}}(\tau_k)$, is a radial Loewner curve with speed $(W_{j,1}|_{\tau_k}^k)^2$ driven by $W_j|_{\tau_k}^k$, the SDE implies that, under $\widetilde{\mathbb{P}}_*$, conditionally on $\mathcal{F}_{\tau_k}^k$, the $\eta_j^{\tau_k}$ up to $\tau_{\xi_j}^j$ has the law of a timechange of a radial $\mathrm{SLE}_{\kappa}(2,\underline{\rho})$ curve in \mathbb{D} started from $g_k(\tau_k, e^{iw_j})$ aimed at 0 with force points $e^{iw_k(\tau_k)}, g_k(\tau_k, e^{i\underline{v}})$. This means that $\widetilde{\mathbb{P}}_*$ is a local commutation coupling within $\underline{\xi}$. Thus, the M_* satisfies all required properties. By Proposition 3.2, we then complete the proof of Proposition 3.1. Moreover, we get the following proposition.

Proposition 3.5. For any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, the global coupling measure in Proposition 3.1 is absolutely continuous w.r.t. \mathbb{P}^1_B on $\mathcal{F}^1_{\tau^1_{\xi_1}} \vee \mathcal{F}^2_{\tau^2_{\xi_2}}$, and the Radon-Nikodym derivative is $M_*(\tau_{\xi})/M_*(\underline{0})$, where M_* is defined by (3.17).

Lemma 3.6. Suppose $\kappa > 4$, m = 2, $\rho_1 = \rho_2 = \kappa - 4$, and $w_1 > v_1 > w_2 > v_2 > w_1 - 2\pi$. Let η_1 and η_2 be given by Proposition 3.1. Let $\mathcal{D} = \{(t_1, t_2) \in \mathbb{R}^2_+ : \eta_1[0, t_1] \cap \eta_2[0, t_2] = \emptyset\}$. Then a.s. $\mathcal{D} = \mathbb{R}^2_+$.

Proof. Let \widehat{w}_j be the driving function of η_j , j = 1, 2. Since the force values $\rho_1 = \rho_2 = \kappa - 4 > \frac{\kappa}{2} - 2$, and 2 > 0, by Proposition 2.1, a.s. η_j does not intersect A_{3-j} , and so has lifetime ∞ . Fix a deterministic time $t_2 \in \mathbb{N}$. Conditionally on $\mathcal{F}_{t_2}^2$, the $g_2(t_2, \cdot)$ -image of the part of η_1 before hitting $\eta_2[0, t_2]$ has the law of a time-change of a radial SLE_{κ} $(2, \kappa - 4, \kappa - 4)$ curve in \mathbb{D} started from $g_2(t_2, e^{iw_1})$ aimed at 0 with force points $e^{i\widehat{w}_2(t_2)}, g_2(t_2, e^{iv_1}), g_2(t_2, e^{iv_2})$. If η_1 does hit $\eta_2[0, t_2]$, then the (time-changed) radial SLE_{κ} $(2, \kappa - 4, \kappa - 4)$ curve $\eta_1^{t_2}$ hits the arc on $\partial \mathbb{D}$ with endpoints $g_2(t_2, e^{iv_1}), g_2(t_2, e^{iv_2})$ that contains $e^{i\widehat{w}_2(t_2)}$. By Proposition 2.1 again, the latter event has probability zero. Thus, \mathbb{P}^c_* -a.s. η_1 does not intersect $\eta_2[0, t_2]$, which implies that $\mathbb{R}_+ \times [0, t_2] \subset \mathcal{D}$. Since this holds for every $t_2 \in \mathbb{N}$, we get the conclusion.

We use \mathbb{P}^c_* to denote the joint law of the driving functions \hat{w}_1 and \hat{w}_2 of the η_1 and η_2 in Lemma 3.6. The superscript c stands for coupling, and the subscript * means that both curves end at 0.

For the w_j and v_j as above, we will need a different kind of commutation coupling as follows. Define M_c on \mathcal{D} by

$$M_c = e^{\frac{(\kappa-6)(\kappa-2)}{8\kappa}} {}^{\mathrm{m}} e^{\frac{b}{6}\widehat{\mathrm{m}}} W^{\mathrm{b}}_{1,1} W^{\mathrm{b}}_{2,1} I^{-\frac{c}{6}} |\sin_2(W_1 - W_2)|^{\frac{\kappa-6}{\kappa}}.$$
(3.19)

Let Ξ be as usual for e^{iw_j} , e^{iv_j} , j = 1, 2. Then Lemma 3.3 holds for M_c . Fix $j \neq k \in \{1, 2\}$. Let τ_k be an \mathcal{F}^k -stopping time. Using (3.1,3.8,3.11,3.13 (for $\rho_0 = \kappa - 6$),3.15,3.16), we see that M_c , with t_k fixed to be τ_k , is a continuous local martingale under \mathbb{P}^i_B satisfying the following SDE:

$$\frac{\partial_j M_c}{M_c} = \mathbf{b} \frac{W_{j,2}}{W_{j,1}} \sqrt{\kappa} \partial B_j + \frac{\kappa - 6}{2} \cot_2(W_j - W_k) W_{j,1} \frac{\partial B_j}{\sqrt{\kappa}}$$

So Lemma 3.4 also holds for M_c .

Fix $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$. One may define a probability measure $\widetilde{\mathbb{P}}_c$ by $d\widetilde{\mathbb{P}}_c/d\mathbb{P}_B^i = M_c(\tau_{\underline{\xi}})/M_c(\underline{0})$. Fix $j \neq k \in \{1, 2\}$ and an \mathcal{F}^k -stopping time τ_k with $\tau_k \leq \tau_{\xi_k}^k$. Using the Girsanov Theorem, one can show that, under $\widetilde{\mathbb{P}}_c$, conditionally on $\mathcal{F}_{\tau_k}^k$, the $g_k(\tau_k, \cdot)$ image of the part of η_j up to $\tau_{\xi_j}^j$ has the law of a time-change of a radial $\mathrm{SLE}_\kappa(\kappa - 6, 0, 0)$ curve in \mathbb{D} started from $g_k(\tau_k, e^{iw_j})$ aimed at 0 with force points $e^{iw_k(\tau_k)}, g_k(\tau_k, e^{iv_1}), g_k(\tau_k, e^{iv_2})$. So we get the following proposition.

Proposition 3.7. Let $\kappa \in (0,\infty)$. There is a coupling of two random radial Loewner curves $\eta_j(t), 0 \leq t < \hat{T}_j, j = 1, 2$, with radial Loewner driving functions \hat{w}_j and maps $g_j(t, \cdot)$ such that $\hat{w}_j(0) = w_j$ and the following holds. For $j = 1, 2, \eta_j$ is a radial $SLE_{\kappa}(\kappa - 6, 0, 0)$ curve in \mathbb{D} started from e^{iw_j} aimed at 0 with force points $e^{iw_{3-j}}, e^{iv_1}, e^{iv_2}$. For any $j \neq k \in \{1, 2\}$ and any \mathcal{F}^k -stopping time τ_k with $\tau_k < \hat{T}_k$, conditionally on $\mathcal{F}^k_{\tau_k}$, the $\eta_j^{\tau_k}(t_j), 0 \leq t_j < T_j^{\mathcal{D}}(\tau_k)$, has the law of a time-change of a radial $SLE_{\kappa}(\kappa - 6, 0, 0)$ curve in \mathbb{D} started from $g_k(\tau_k, e^{iw_j})$ aimed at 0 with force points $e^{i\widehat{w}_k(\tau_k)}, g_k(\tau_k, e^{iv_1}), g_k(\tau_k, e^{iv_2})$ stopped at the first time that it separates 0 from any of the three force points in \mathbb{D} . Moreover, for any $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$, the joint law of \widehat{w}_1 and \widehat{w}_2 is absolutely continuous w.r.t. \mathbb{P}^i_B on $\mathcal{F}^1_{\tau_{\xi_1}} \vee \mathcal{F}^2_{\tau_{\xi_2}}$, and the Radon-Nikodym derivative is $M_c(\tau_{\xi})/M_c(\underline{0})$, where M_c is defined by (3.19).

By [17], for j = 1, 2, the η_j in Proposition 3.7 is a time-change of a chordal SLE_{κ} in \mathbb{D} from e^{iw_j} to $e^{iw_{3-j}}$ stopped at the first time that it separates 0 from any of $e^{iw_{3-j}}, e^{iv_1}, e^{iv_2}$ in \mathbb{D} . When $\kappa \in (0, 8]$, the coupling of η_1 and η_2 in Proposition 3.7 can be easily constructed using the DMP and reversibility of chordal SLE_{κ} . For the construction, we start with a chordal SLE_{κ} in \mathbb{D} from e^{iw_1} to e^{iw_2} , let η_1 and η_2 be respectively the part of η and the reversal of η up to the first time that it separates 0 from any of the three force points in \mathbb{D} . But such a construction does not provide us the Radon-Nikodym derivative process M_c .

Let \mathbb{P}_c^c denote the joint law of the driving functions \hat{w}_1 and \hat{w}_2 of the η_1 and η_2 obtained from Proposition 3.7. The superscript and subscript respectively stand for coupling and chordal.

We are mostly interested in the case $\kappa \in (4, 8)$. In that case, under \mathbb{P}_c^c the η_1 and η_2 are parts of the same chordal $\operatorname{SLE}_{\kappa}$ curve in \mathbb{D} from e^{iw_1} to e^{iw_2} . Under \mathbb{P}_*^c the η_1 and η_2 both end at 0, and $\eta_1 \cup A_1$ is disjoint from $\eta_2 \cup A_2$, where for $j = 1, 2, A_j$ is the connected component of $\mathbb{D} \setminus \{e^{iv_1}, e^{iv_2}\}$ that contains e^{iw_j} . Thus, under \mathbb{P}_*^c , 0 is a cut point of $\eta_1 \cup \eta_2 \cup A_1 \cup A_2 \cup \{0\}$. Intuitively, we may understand \mathbb{P}_*^c as the chordal $\operatorname{SLE}_{\kappa}$ curve in \mathbb{D} from e^{iw_1} to e^{iw_2} conditioned on the singular event that 0 is a cut point of $\eta \cup A_1 \cup A_2$. See Figure 3.2.3.

Let $M_{*\to c} = M_c/M_*$, which can be expressed in terms of the α_0 and G defined in (1.2,1.4):

$$M_{*\to c} = e^{-\alpha_0 \,\mathrm{m}} \widetilde{G}(W_1, W_2, V_1, V_2)^{-1}.$$
(3.20)

Lemma 3.8. Suppose $\kappa \in (4,8)$. For any \mathcal{F} -stopping time \underline{T} , \mathbb{P}_c^c restricted to $\mathcal{F}_{\underline{T}} \cap \{\underline{T} \in \mathcal{D}\}$ is absolutely continuous w.r.t. \mathbb{P}_*^c , and the RN derivative is $M_{*\to c}(\underline{T})/M_{*\to c}(\underline{0})$. In other words, if $A \in \mathcal{F}_{\underline{T}}$ and $A \subset \{\underline{T} \in \mathcal{D}\}$, then $\mathbb{P}_c^c[A] = \mathbb{E}_*^c[\mathbf{1}_A M_{*\to c}(\underline{T})/M_{*\to c}(\underline{0})]$.

Proof. Let $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$ and $\underline{t} = (t_1, t_2) \in \mathbb{R}^2_+$. By Lemma 3.4, Proposition 3.5, and the facts



Figure 1: The above figure is a sketch of the left boundary (red) and right boundary (blue) of a chordal SLE_{κ} curve γ ($\kappa \in (4, 8)$) in \mathbb{D} connecting a_1 with a_2 , where each boundary is a boundary-touching simple curve composed of countably many crosscuts of \mathbb{D} , exactly one of which connects the upper semicircle A_1 with the lower semicircle A_2 . The p_1, \ldots, p_6 illustrated are intersections of the two boundaries, and are cut points of γ , but only the intersection p_4 of the two crosscuts connecting A_1 and A_2 is a cut point of $\gamma \cup A_1 \cup A_2$. The small circle has radius r. If one conditions γ on the event that $\gamma \cup A_1 \cup A_2$ has a cut point in $\{|z| < r\}$, and lets $r \to 0$, then the result is a curve from a_1 to a_2 passing through its cut point 0, and the two sub-curves (respectively from a_1, a_2 to 0) jointly follow the law \mathbb{P}^c_* .

$$\begin{split} \mathcal{F}_{t_1 \wedge \tau_{\xi_1}^1}^1 & \vee \mathcal{F}_{t_2 \wedge \tau_{\xi_2}^2}^2 \subset \mathcal{F}_{\underline{t}} \text{ and that } M_*(\underline{t} \wedge \tau_{\underline{\xi}}) \text{ is } \mathcal{F}_{t_1 \wedge \tau_{\xi_1}^1}^1 \vee \mathcal{F}_{t_2 \wedge \tau_{\xi_2}^2}^2 \text{-measurable, we see that} \\ & \frac{d\mathbb{P}_*^c | \mathcal{F}_{t_1 \wedge \tau_{\xi_1}^1}^1 \vee \mathcal{F}_{t_2 \wedge \tau_{\xi_2}^2}^2}{d\mathbb{P}_B^i | \mathcal{F}_{t_1 \wedge \tau_{\xi_1}^1}^1 \vee \mathcal{F}_{t_2 \wedge \tau_{\xi_2}^2}^2} = \frac{\mathbb{E}\Big[M_*(\tau_{\underline{\xi}}) \Big| \mathcal{F}_{t_1 \wedge \tau_{\xi_1}^1}^1 \vee \mathcal{F}_{t_2 \wedge \tau_{\xi_2}^2}^2\Big]}{M_*(\underline{0})} = \frac{M_*(\underline{t} \wedge \tau_{\underline{\xi}})}{M_*(\underline{0})}. \end{split}$$

A similar formula holds with \mathbb{P}_c^c and M_c respectively in place of \mathbb{P}_*^c and M_* . So we get

$$\frac{d\mathbb{P}_{c}^{c}|\mathcal{F}_{t_{1}\wedge\tau_{\xi_{1}}^{1}}^{1}\vee\mathcal{F}_{t_{2}\wedge\tau_{\xi_{2}}^{2}}^{2}}{d\mathbb{P}_{*}^{c}|\mathcal{F}_{t_{1}\wedge\tau_{\xi_{1}}^{1}}^{1}\vee\mathcal{F}_{t_{2}\wedge\tau_{\xi_{2}}^{2}}^{2}} = \frac{M_{*\to c}(\underline{t}\wedge\tau_{\underline{\xi}})}{M_{*\to c}(\underline{0})}.$$
(3.21)

Let $\underline{T} = (T_1, T_2)$ be an \mathcal{F} -stopping time. Let $\underline{\xi} \in \Xi^*$. Fix $A \in \mathcal{F}_{\underline{T}}$ with $A \subset \{\underline{T} < \tau_{\underline{\xi}}\}$. For $n \in \mathbb{N}$, let $\underline{T}^n = (T_1^n, T_2^n)$, where $T_j^n = \frac{[2^n T]}{2^n}$, j = 1, 2. It is easy to see that each \underline{T}^n is an \mathcal{F} -stopping time, and $\underline{T}^n \downarrow \underline{T}$. Let $A_n = A \cap \{\underline{T}^n < \tau_{\underline{\xi}}\}$, $n \in \mathbb{N}$. Then $A_n \uparrow A$. Fix $n \in \mathbb{N}$. By

Proposition 2.7, $A_n \in \mathcal{F}_{\underline{T}^n}$. We know that \underline{T}^n takes values in $\frac{1}{2^n}\mathbb{Z}^2_+$. Let $\underline{s} = (s_1, s_2) \in \frac{1}{2^n}\mathbb{Z}^2_+$ and $A_{n,\underline{s}} = A_n \cap \{\underline{T}^n = \underline{s}\}$. By Proposition 2.5, $A_{n,\underline{s}} \in \mathcal{F}_{\underline{s}}$. Since $\mathcal{F}_{s_j}^j \cap \{s_j \le s_j \land \tau_{\xi_j}^j\} \subset \mathcal{F}_{s_j \land \tau_{\xi_j}^j}^j$, j = 1, 2, by a monotone class argument we get $\mathcal{F}_{\underline{s}} \cap \{\underline{s} \le \underline{s} \land \tau_{\underline{\xi}}\} \subset \mathcal{F}_{t_1 \land \tau_{\xi_1}^1}^1 \lor \mathcal{F}_{t_2 \land \tau_{\xi_2}^2}^2$. Since $A_{n,\underline{s}} \in \mathcal{F}_{\underline{s}} \cap \{\underline{s} < \tau_{\underline{\xi}}\}$, we see that $A_{n,\underline{s}} \in \mathcal{F}_{t_1 \land \tau_{\xi_1}^1}^1 \lor \mathcal{F}_{t_2 \land \tau_{\xi_2}^2}^2$. By (3.21),

$$\mathbb{P}_{c}^{c}[A_{n,\underline{s}}] = \mathbb{E}_{*}^{c}[\mathbf{1}_{A_{n,\underline{s}}}M_{*\to c}(\underline{s}\wedge\tau_{\underline{\xi}})/M_{*\to c}(\underline{0})] = \mathbb{E}_{*}^{c}[\mathbf{1}_{A_{n,\underline{s}}}M_{*\to c}(\underline{T}^{n})/M_{*\to c}(\underline{0})],$$

where the last equality holds because $\underline{T}^n = \underline{s} \leq \tau_{\underline{\xi}}$ on $A_{n,\underline{s}}$. Summing up over $\underline{s} \in \frac{1}{2^n} \mathbb{Z}^2_+$, we get $\mathbb{P}^c_c[A_n] = \mathbb{E}^c_*[\mathbf{1}_{A_n}M_{*\to c}(\underline{T}^n)/M_{*\to c}(\underline{0})]$. Sending $n \to \infty$ and using dominated convergence theorem, we get $\mathbb{P}^c_c[A] = \mathbb{E}^c_*[\mathbf{1}_A M_{*\to c}(\underline{T})/M_{*\to c}(\underline{0})]$. Here we use the facts that $\underline{T}^n < \tau_{\underline{\xi}}$ on A_n and that log $M_{*\to c}$ is uniformly bounded on $[0, \tau_{\underline{\xi}}]$. This means that, for any $\xi \in \Xi^*$, \mathbb{P}^c_c restricted to $\mathcal{F}_{\underline{T}} \cap \{\underline{T} < \tau_{\underline{\xi}}\}$ is absolutely continuous w.r.t. \mathbb{P}^c_* , and the RN derivative is $M_{*\to c}(\underline{T})/M_{*\to c}(\underline{0})$. The conclusion of the lemma then follows since $\bigcup_{\xi\in\Xi^*}\{\underline{T} < \tau_{\underline{\xi}}\} = \{\underline{T} \in \mathcal{D}\}$.

3.3 A time curve in the time region

Suppose η_1 and η_2 are random radial Loewner curves driven by \hat{w}_1 and \hat{w}_2 , which jointly follow one of the three laws \mathbb{P}^i_B , \mathbb{P}^c_* , \mathbb{P}^c_c . Let $\theta = V_1 - V_2$ and $Z_j = W_j - V_j$, j = 1, 2. By [20, (4.1)],

$$\partial_j \theta = \frac{-W_{j,1}^2 \sin_2(\theta)}{\sin_2(W_j - V_1) \sin_2(W_j - V_2)} \partial t_j, \quad j = 1, 2.$$

When j = 2, using $0 > \sin_2(W_j - V_1) \sin_2(W_j - V_2) \ge -\sin(\theta/4)^2$ and (3.1), we get

$$\partial_2 \theta \ge 2W_{2,1}^2 \cot(\theta/4) \partial t_j = 2 \cot(\theta/4) \partial_2 \mathbf{m} \,. \tag{3.22}$$

Recall that most of the processes we have encountered are defined on the time region $\mathcal{D} \subset \mathbb{R}^2_+$. From now on, suppose $v_1 - v_2 = \pi$. Then $\theta(\underline{0}) = \pi$. By [20, Section 4], there exists a continuous increasing curve $\underline{u} = (u_1, u_2) : \mathbb{R}_+ \to \overline{\mathcal{D}}$ with $\underline{u}(0) = \underline{0}$, and an extended number $T^u \in (0, \infty]$, such that \underline{u} is strictly increasing and takes values in \mathcal{D} on $[0, T^u)$, takes constant values in ∂D on $[T^u, \infty)$, and for any $t \in [0, T^u)$, $m(\underline{u}(t)) = t$ and $\theta(\underline{u}(t)) = \pi$. We use such curve \underline{u} to obtain a one-time-parameter process $X^u := X \circ \underline{u}$ from any two-time-parameter process X on \mathcal{D} . We have the following facts.

- For $0 \le t < T^u$, $m^u(t) = t$, $\theta^u(t) = \pi$, and $Z_j^u \in (0, \pi)$, j = 1, 2.
- For any $j \in \{1, 2\}$ and $t \ge 0, u_j(t) \le t$.
- \underline{u} is differentiable with positive derivatives on $[0, T^u)$ that satisfy

$$(W_{j,1}^u)^2 u'_j = \frac{\sin(Z_j^u)}{\sin(Z_1^u) + \sin(Z_2^u)}, \quad \text{on } [0, T^u), \quad j = 1, 2.$$
(3.23)

• For any deterministic time $t \in \mathbb{R}_+$, $\underline{u}(t)$ is an \mathcal{F} -stopping time.

Using the last property, we define an \mathbb{R}_+ -indexed filtration \mathcal{F}^u by $\mathcal{F}^u_t = \mathcal{F}_{\underline{u}(t)}, t \ge 0$. For $\xi = (\xi_1, \xi_2) \in \Xi$, let $\tau^u_{\underline{\xi}}$ denote the first $t \ge 0$ such that $u_1(t) = \tau^1_{\xi_1}$ or $u_2(t) = \tau^2_{\xi_2}$, whichever comes first. Then for any $\xi \in \Xi, \tau^u_{\underline{\xi}}$ is an \mathcal{F}^u -stopping time; $\underline{u}(\tau^u_{\underline{\xi}})$ is an \mathcal{F} -stopping time; and for every deterministic $t \in \mathbb{R}_+, \underline{u}(t \land \tau^u_{\underline{\xi}})$ is an \mathcal{F} -stopping time.

First, suppose (\hat{w}_1, \hat{w}_2) follows the law \mathbb{P}_B^i . Then there are independent Brownian motions B_1 and B_2 such that $\hat{w}_j(t_j) = w_j + \sqrt{\kappa}B_j(t_j), t_j \ge 0, j = 1, 2$. So we get five continuous \mathcal{F} -martingales: $\hat{w}_j(t_j), \hat{w}_j(t_j)^2 - \kappa t_j, j = 1, 2$, and $\hat{w}_1(t_1)\hat{w}_2(t_2)$. Using Proposition 2.11 and the fact that for any $t \in \mathbb{R}_+, \underline{u}(t)$ is an \mathcal{F} -stopping time bounded by (t, t), we see that $\hat{w}_j^u(t), \hat{w}_j^u(t)^2 - \kappa u_j(t), j = 1, 2$, and $\hat{w}_1^u(t)\hat{w}^u(t)$ are all \mathcal{F}^u -martingales. Note that $\hat{w}_j^u(t) = \hat{w}_j(u_j(t))$. So we get quadratic variation and covariation for $\hat{w}_j^u, j = 1, 2$:

$$\langle \widehat{w}_j^u \rangle_t = \kappa u_j(t), \quad j = 1, 2, \quad \langle \widehat{w}_1^u, \widehat{w}_2^u \rangle \equiv 0.$$
 (3.24)

Fix $\underline{\xi} = (\xi_1, \xi_2) \in \Xi$. Let $M \in \{M_*, M_c\}$. We have known that $M(\cdot \wedge \tau_{\underline{\xi}})$ is a bounded \mathcal{F} -martingale (under \mathbb{P}^i_B). From Proposition 2.11, $M(\underline{u}(\cdot) \wedge \tau_{\underline{\xi}})$ is an \mathcal{F}^u -martingale. Since $\tau_{\underline{\xi}}^u$ is an \mathcal{F}^u -stopping time, we see that $M(\underline{u}(\cdot \wedge \tau_{\underline{\xi}}^u) \wedge \tau_{\underline{\xi}}) = M(\underline{u}(\cdot \wedge \tau_{\underline{\xi}}^u)) = M^u(\cdot \wedge \tau_{\underline{\xi}}^u)$ is an \mathcal{F}^u -martingale. Here the first equality holds because $\underline{u}(t \wedge \tau_{\underline{\xi}}^u) \leq \tau_{\underline{\xi}}$. Since $[0, T^u) = \bigcup_{\xi \in \Xi^*} [0, \tau_{\underline{\xi}}^u]$ and Ξ^* is countable, we conclude that M^u_* and M^u_c are \mathcal{F}^u -local martingales with lifetime T^u under the probability measure \mathbb{P}^i_B .

Next, we compute the SDE for M^u_* in terms of \widehat{w}^u_1 and \widehat{w}^u_2 . By (3.17) we may express M^u_* as a product of several factors. Among these factors, $\sin_2(W^u_1 - W^u_2)^{\frac{2}{\kappa}}$, $(W^u_{j,1})^b$ and $|\sin_2(W^u_j - V^u_s)|^{1-\frac{4}{\kappa}}$, $j, s \in \{1, 2\}$, contribute the diffusion part of M^u_* , and other factors are differentiable in t. For $j \neq k \in \{1, 2\}$ and $s \in \{1, 2\}$, using (3.2,3.5,3.6), we get the following SDEs:

$$dW_{j}^{u} = W_{j,1}^{u} d\widehat{w}_{j}^{u} - \kappa \, \mathrm{b} \, W_{j,2} u_{j}' dt + \cot_{2} (W_{j}^{u} - W_{k}^{u}) (W_{k,1}^{u})^{2} u_{k}' dt, \qquad (3.25)$$

$$\frac{d(W_{j,1}^{u})^{\mathrm{b}}}{(W_{j,1}^{u})^{\mathrm{b}}} = \mathrm{b} \, \frac{W_{j,2}^{u}}{W_{j,1}^{u}} \, d\widehat{w}_{j}^{u} + \mathrm{drift \ terms},$$

$$\frac{d\sin_{2} (W_{j}^{u} - V_{s}^{u})^{1 - \frac{4}{\kappa}}}{\sin_{2} (W_{j}^{u} - V_{s}^{u})^{1 - \frac{4}{\kappa}}} = \frac{\kappa - 4}{2\kappa} \cot_{2} (W_{j}^{u} - V_{s}^{u}) W_{j,1}^{u} d\widehat{w}_{j}^{u} + \mathrm{drift \ terms},$$

$$\frac{d\sin_{2} (W_{1}^{u} - W_{2}^{u})^{\frac{2}{\kappa}}}{\sin_{2} (W_{1}^{u} - W_{2}^{u})^{\frac{2}{\kappa}}} = \frac{1}{\kappa} \cot_{2} (W_{j}^{u} - W_{k}^{u}) [W_{j,1}^{u} d\widehat{w}_{j}^{u} - W_{k,1}^{u} d\widehat{w}_{k}^{u}] + \mathrm{drift \ terms}.$$

Since we already know that M^u_* , \hat{w}^u_1 and \hat{w}^u_2 are \mathcal{F}^u -local martingales, we get the SDE:

$$\frac{dM^u_*}{M^u_*} = \sum_{j=1}^2 \left[b \frac{W^u_{j,2}}{W^u_{j,1}} + \frac{1}{\kappa} \cot_2(W^u_j - W^u_{3-j}) W^u_{j,1} + \sum_{s=1}^2 \frac{\kappa - 4}{2\kappa} \cot_2(W^u_j - V^u_s) W^u_{j,1} \right] d\hat{w}^u_j. \quad (3.26)$$

Lemma 3.9. Under \mathbb{P}^c_* , for j = 1, 2, \widehat{w}^u_j satisfies the SDE

$$d\widehat{w}_{j}^{u} = \sqrt{\kappa u_{j}^{\prime}} dB_{j}^{u} + \left[\kappa \operatorname{b} \frac{W_{j,2}^{u}}{W_{j,1}^{u}} + \operatorname{cot}_{2}(W_{j}^{u} - W_{3-j}^{u})W_{j,1}^{u} + \sum_{l=1}^{2} \frac{\kappa - 4}{2} \operatorname{cot}_{2}(W_{j}^{u} - V_{l}^{u})W_{j,1}^{u}\right] u_{j}^{\prime} dt,$$
(3.27)

where $B_{i}^{u}(t)$, j = 1, 2, are two independent Brownian motions.

Proof. This lemma is similar to [20, Lemma 4.3], and the proof uses a Girsanov-type argument. Here is a sketch. We first define \tilde{w}_{j}^{u} , j = 1, 2, such that

$$\widetilde{w}_{j}^{u}(t) = \widehat{w}_{j}^{u}(t) - \int_{0}^{t} \left[\kappa \operatorname{b} \frac{W_{j,2}^{u}(s)}{W_{j,1}^{u}(s)} + \operatorname{cot}_{2}(W_{j}^{u}(s) - W_{3-j}^{u}(s))W_{j,1}^{u}(s) + \sum_{l=1}^{2} \frac{\kappa - 4}{2} \operatorname{cot}_{2}(W_{j}^{u}(s) - V_{l}^{u}(s))W_{j,1}^{u}(s) \right] u_{j}'(s)ds.$$
(3.28)

By Lévy's characterization of Brownian motion, it suffices to show that \widetilde{w}_1^u and \widetilde{w}_2^u are local martingals under \mathbb{P}^c_* , $d\langle \widetilde{w}_i^u \rangle = du_j$, j = 1, 2, and $\langle \widetilde{w}_1^u, \widetilde{w}_2^u \rangle \equiv 0$.

Using (3.26,3.28) and Itô's formula, it is straightforward to calculate that $\widetilde{w}_j^u M_*^u$, j = 1, 2, are \mathcal{F}^u -local martingales under \mathbb{P}_B^i . Fix $j \in \{1, 2\}$. Now we prove that \widetilde{w}_j^u is a local martingale under \mathbb{P}_*^c . It suffices to prove that, for any $\underline{\xi} \in \Xi$, $\widetilde{w}_j^u(\cdot \wedge \tau_{\underline{\xi}}^u)$ is an \mathcal{F}^u -martingale under \mathbb{P}_*^c . Fix $\underline{\xi} \in \Xi$. Using an argument similar to the proof of Lemma 3.3, we can show that $|\widetilde{w}_j^u|$ is bounded on $[0, \tau_{\underline{\xi}}^u]$ by a constant depending only on $\kappa, \underline{\xi}, w_1, w_2, v_1, v_2$. So we see that $\widetilde{w}_j^u(\cdot \wedge \tau_{\underline{\xi}}^u)M_*^u(\cdot \wedge \tau_{\underline{\xi}}^u)$, j = 1, 2, are bounded \mathcal{F}^u -martingales under \mathbb{P}_B^i .

To prove that $\widetilde{w}_{j}^{u}(\cdot \wedge \tau_{\underline{\xi}}^{u})$ is a martingale under \mathbb{P}_{*}^{c} , we fix $t > s \geq 0$ and $A \in \mathcal{F}_{s}^{u}$, and need to prove that

$$\mathbb{E}^{c}_{*}[\mathbf{1}_{A}\widetilde{w}^{u}_{j}(t\wedge\tau^{u}_{\underline{\xi}})] = \mathbb{E}^{c}_{*}[\mathbf{1}_{A}\widetilde{w}^{u}_{j}(s\wedge\tau^{u}_{\underline{\xi}})].$$
(3.29)

Let $A_1 = A \cap \{\tau_{\underline{\xi}}^u \le s\}$ and $A_2 = A \cap \{\tau_{\underline{\xi}}^u > s\}$. Then (3.29) would follow from

$$\mathbb{E}^{c}_{*}[\mathbf{1}_{A_{1}}\widetilde{w}^{u}_{j}(t\wedge\tau^{u}_{\underline{\xi}})] = \mathbb{E}^{c}_{*}[\mathbf{1}_{A_{1}}\widetilde{w}^{u}_{j}(s\wedge\tau^{u}_{\underline{\xi}})]; \qquad (3.30)$$

$$\mathbb{E}^{c}_{*}[\mathbf{1}_{A_{2}}\widetilde{w}^{u}_{j}(t\wedge\tau^{u}_{\underline{\xi}})] = \mathbb{E}^{c}_{*}[\mathbf{1}_{A_{2}}\widetilde{w}^{u}_{j}(s\wedge\tau^{u}_{\underline{\xi}})].$$
(3.31)

Equality (3.30) is trivially true because $\widetilde{w}_{j}^{u}(t \wedge \tau_{\underline{\xi}}^{u}) = \widetilde{w}_{u}^{j}(\tau_{\underline{\xi}}^{u}) = \widetilde{w}_{j}^{u}(s \wedge \tau_{\underline{\xi}}^{u})$ on A_{1} .

Since $\widetilde{w}_j^u(\cdot \wedge \tau_{\xi}^u) M^u_*(\cdot \wedge \tau_{\xi}^u)$ is an \mathcal{F}^u -martingale under \mathbb{P}_B^i and $A_2 \in \mathcal{F}_s^u$, we get

$$\mathbb{E}^{\mathbf{i}}_{B}[\mathbf{1}_{A_{2}}\widetilde{w}^{u}_{j}(t\wedge\tau^{u}_{\underline{\xi}})M^{u}_{*}(t\wedge\tau^{u}_{\underline{\xi}})] = \mathbb{E}^{\mathbf{i}}_{B}[\mathbf{1}_{A_{2}}\widetilde{w}^{u}_{j}(s\wedge\tau^{u}_{\underline{\xi}})M^{u}_{*}(s\wedge\tau^{u}_{\underline{\xi}})].$$

Then (3.31) would follow from the above equality and the equalities

$$\mathbb{E}_{B}^{i}[\mathbf{1}_{A_{2}}\widetilde{w}_{j}^{u}(r\wedge\tau_{\underline{\xi}}^{u})M_{*}^{u}(r\wedge\tau_{\underline{\xi}}^{u})] = M_{*}(\underline{0})\mathbb{E}_{*}^{c}[\mathbf{1}_{A_{2}}\widetilde{w}_{j}^{u}(r\wedge\tau_{\underline{\xi}}^{u})], \quad r\in\{s,t\}.$$
(3.32)

We now prove (3.32) in the case r = t. The case r = s is similar. By Proposition 3.5, $\mathbb{P}^c_* | \mathcal{F}_{\tau_{\underline{\xi}}}$ is absolutely continuous w.r.t. $\mathbb{P}^i_B | \mathcal{F}_{\tau_{\underline{\xi}}}$, and the RN derivative is $M_*(\tau_{\underline{\xi}})/M_*(\underline{0})$. Since $M_*(\cdot \wedge \tau_{\underline{\xi}})$ is a martingale, for any two-dimensional stopping time $\underline{T} \leq \tau_{\underline{\xi}}$, $\mathbb{P}^c_* | \mathcal{F}_{\underline{T}}$ is absolutely continuous w.r.t. $\mathbb{P}^i_B | \mathcal{F}_{\underline{T}}$, and the RN derivative is $M_*(\underline{T})/M_*(\underline{0})$. Let $\underline{T} = \underline{u}(t \wedge \tau_{\underline{\xi}})$. Then \underline{T} is a stopping time $\leq \tau_{\underline{\xi}}$. Since $\{\tau_{\underline{\xi}}^u > s\} \subset \{\underline{u}(s) \leq \underline{u}(s \wedge \tau_{\underline{\xi}}^u)\}$ and $A \in \mathcal{F}^u_s = \mathcal{F}_{\underline{u}(s)}$, we get $A_2 \in \mathcal{F}_{\underline{u}(s \wedge \tau_{\underline{\xi}}^u)} \subset \mathcal{F}_{\underline{u}(t \wedge \tau_{\underline{\xi}})} = \mathcal{F}_{\underline{T}}$. Thus, $\mathbf{1}_{A_2} \widetilde{w}^u_j(t \wedge \tau_{\underline{\xi}}^u)$ is $\mathcal{F}_{\underline{T}}$ -measurable. So we get (3.32) in the case r = t using the RN derivative between $\mathbb{P}^c_* | \mathcal{F}_{\underline{T}}$ and $\mathbb{P}^i_B | \mathcal{F}_{\underline{T}}$.

Now we have proved that \widetilde{w}_{j}^{u} , j = 1, 2, are \mathcal{F}^{u} -martingales under \mathbb{P}_{*}^{c} . Since (3.24) holds for \widehat{w}_{j}^{u} under \mathbb{P}_{B}^{i} , it also holds for \widetilde{w}_{j}^{u} under \mathbb{P}_{B}^{i} . For any $\xi \in \Xi$, using the local absolute continuity between $\mathbb{P}_{*}^{c} | \mathcal{F}_{\tau_{\underline{\xi}}}$ and $\mathbb{P}_{B}^{i} | \mathcal{F}_{\tau_{\underline{\xi}}}$, we can then conclude that (3.24) also holds for \widetilde{w}_{j}^{u} under \mathbb{P}_{*}^{c} up to $\tau_{\underline{\xi}}^{u}$. Since this holds for any $\xi \in \Xi$, we get $d\langle \widetilde{w}_{j}^{u} \rangle = du_{j}, j = 1, 2$, and $\langle \widetilde{w}_{1}^{u}, \widetilde{w}_{2}^{u} \rangle \equiv 0$ throughout. The proof is then complete.

Now we suppose (\hat{w}_1, \hat{w}_2) follows the law \mathbb{P}^c_* , and let B^u_1 and B^u_2 be the independent Brownian motions from Lemma 3.9. Combining (3.23,3.25,3.27), we get, for j = 1, 2,

$$dW_j^u = W_{j,1}^u \sqrt{\kappa u_j'} dB_j^u + \cot_2(W_j^u - W_{3-j}^u) dt + \sum_{s=1}^2 \frac{\kappa - 4}{2} \cot_2(W_j^u - V_s^u) (W_{j,1}^u)^2 u_j' dt.$$

Using (3.2) we get, for j = 1, 2,

$$dV_j^u = -\cot_2(W_j^u - V_j^u)(W_{j,1}^u)^2 u_j' dt - \cot_2(W_{3-j}^u - V_j^u)(W_{3-j,1}^u)^2 u_{3-j}' dt.$$

Since $Z_j^u = W_j^u - V_j^u$, $W_{3-j}^u - V_j^u = W_{3-j}^u - V_{3-j}^u \pm \pi = Z_{3-j}^u \pm \pi$, $W_j^u - W_{3-j}^u = Z_j^u - Z_{3-j}^u \mp \pi$, and $W_j^u - V_{3-j}^u = Z_j^u \mp \pi$, combining the above two displayed formulas with (3.23), we get, for j = 1, 2,

$$dZ_j^u = \sqrt{\frac{\kappa \sin(Z_j^u)}{\sin(Z_1^u) + \sin(Z_2^u)}} dB_j^u + \frac{(\kappa - 2)\cos(Z_j^u)}{\sin(Z_1^u) + \sin(Z_2^u)} dt.$$
(3.33)

Remark 3.10. If the two force values $\kappa - 4$ and $\kappa - 4$ at e^{iv_1} and e^{iv_2} are respectively replaced by $\rho_1, \rho_2 \in \mathbb{R}$, then $Z_j^u, j = 1, 2$, satisfy the SDE:

$$dZ_j^u = \sqrt{\frac{\kappa \sin(Z_j^u)}{\sin(Z_1^u) + \sin(Z_2^u)}} dB_j^u + \frac{(2 + (\rho_1 + \rho_2)/2)\cos(Z_j^u)}{\sin(Z_1^u) + \sin(Z_2^u)} dt + \frac{(\rho_j - \rho_{3-j})/2}{\sin(Z_1^u) + \sin(Z_2^u)} dt.$$

3.4 Transition density

We are going to derive the transition density of (Z_1^u, Z_2^u) under \mathbb{P}_*^c or \mathbb{P}_c^c following the approach of [20, Section 5]. The idea, originated in [21, Appendix B], is to solve a PDE eigenvalue problem using orthogonal polynomials. First suppose that the underlying probability measure is \mathbb{P}^c_* . Recall that (Z_1^u, Z_2^u) satisfy (3.33), where B_1^u and B_2^u are independent Brownian motions by Lemma 3.9. Define B^u_+ and B^u_- by

$$B_{\pm}^{u}(t) = \int_{0}^{t} \sqrt{\frac{\sin(Z_{1}^{u}(s))}{\sin(Z_{1}^{u}(s)) + \sin(Z_{2}^{u}(s))}} dB_{1}^{u}(s) \pm \int_{0}^{t} \sqrt{\frac{\sin(Z_{2}^{u}(s))}{\sin(Z_{1}^{u}(s)) + \sin(Z_{2}^{u}(s))}} dB_{1}^{u}(s)$$

Then B^u_+ and B^u_- are standard linear Brownian motions with quadratic covariation

$$d\langle B_{+}^{u}, B_{-}^{u} \rangle_{t} = \cot_{2}(Z_{1}^{u} + Z_{2}^{u}) \tan_{2}(Z_{1}^{u} - Z_{2}^{u}) dt.$$
(3.34)

Define $Z_{\pm}^u = (Z_1^u \pm Z_2^u)/2$. Then $Z_{\pm}^u \in (0,\pi), Z_{\pm}^u \in (-\pi/2,\pi/2)$, and they satisfy the SDEs:

$$dZ_{+}^{u} = \frac{\sqrt{\kappa}}{2} dB_{+}^{u} + \frac{\kappa - 2}{2} \cot(Z_{+}^{u}) dt;$$
$$dZ_{-}^{u} = \frac{\sqrt{\kappa}}{2} dB_{-}^{u} - \frac{\kappa - 2}{2} \tan(Z_{-}^{u}) dt.$$

Let $X = \cos(Z_+^u)$ and $Y = \sin(Z_-^u)$. Then $X, Y \in (-1, 1)$, and satisfy the SDEs

$$dX = -\frac{\sqrt{\kappa}}{2}\sqrt{1 - X^2}dB_+^u - \left(\frac{\kappa - 2}{2} + \frac{\kappa}{8}\right)Xdt;$$

$$dY = +\frac{\sqrt{\kappa}}{2}\sqrt{1 - Y^2}dB_-^u - \left(\frac{\kappa - 2}{2} + \frac{\kappa}{8}\right)Ydt.$$

From (3.34) we have

$$d\langle X,Y\rangle_t = -\frac{\kappa}{4}XYdt.$$

We have $(X, Y) \in \mathbb{D}$ because $X^2 + Y^2 = 1 - \sin(Z_1^u) \sin(Z_2^u) < 1$.

Define a second order differential operator \mathcal{L} by

$$\mathcal{L} := \frac{\kappa}{8}(1-x^2)\partial_x^2 + \frac{\kappa}{8}(1-y^2)\partial_y^2 - \frac{\kappa}{4}xy\partial_x\partial_y - (\frac{\kappa-2}{2} + \frac{\kappa}{8})x\partial_x - (\frac{\kappa-2}{2} + \frac{\kappa}{8})y\partial_y.$$

Using orthogonal polynomials, we may find eigenvectors and eigenvalues of \mathcal{L} . Define

$$\lambda_s = -\frac{\kappa}{8}s(s+4-\frac{8}{\kappa}), \quad s \in \mathbb{Z}_+ = \{0, 1, 2, 3, \dots\}.$$

A straightforward calculation shows that, for any $n, m \in \mathbb{Z}_+$, $\mathcal{L}(x^n y^m)$ equals $\lambda_{n+m} x^n y^m$ plus a polynomial in x, y of degree less than n + m. Hence, for each $n, m \in \mathbb{Z}_+$, there is a polynomial $P_{(n,m)}(x, y)$ of degree n + m, which equals $x^n y^m$ plus a polynomial in x, y of degree less than n + m, such that $\mathcal{L}P_{(n,m)} = \lambda_{n+m}P_{(n,m)}$.

Let $\Psi(x,y) = (1-x^2-y^2)^{1-\frac{4}{\kappa}}$, and define $\langle f,g \rangle_{\Psi} := \iint_{\mathbb{D}} f(x,y)g(x,y)\Psi(x,y)dxdy$. Since $\Psi \equiv 0$ on \mathbb{T} , using integration by parts, we can show that for smooth functions f and g on $\overline{\mathbb{D}}$,

 $\langle \mathcal{L}f,g \rangle_{\Psi} = \langle f,\mathcal{L}g \rangle_{\Psi}.$ In fact, if we let $a_{xx} = \frac{\kappa}{8}(1-x^2), a_{yy} = \frac{\kappa}{8}(1-y^2), a_{xy} = a_{yx} = -\frac{\kappa}{8}xy, b_x = -(\frac{\kappa-2}{2} + \frac{\kappa}{8})x$, and $b_y = -(\frac{\kappa-2}{2} + \frac{\kappa}{8})y$, then both $\langle \mathcal{L}f,g \rangle_{\Psi}$ and $\langle f,\mathcal{L}g \rangle_{\Psi}$ equal

$$-\int\!\!\int_{\mathbb{D}} [(\partial_x f) a_{xx} \Psi(\partial_x g) + (\partial_x f) a_{xy} \Psi(\partial_y g) + (\partial_y f) a_{yx} \Psi(\partial_x g) + (\partial_y f) a_{yy} \Psi(\partial_y g)] dxdy.$$

Here we use $\partial_x(a_{xx}\Psi) + \partial_y(a_{xy}\Psi) = b_x\Psi$ and $\partial_x(a_{yx}\Psi) + \partial_y(a_{yy}\Psi) = b_y\Psi$.

Thus, the eigenvectors of \mathcal{L} with different eigenvalues are orthogonal w.r.t. $\langle \cdot \rangle_{\Psi}$, and we may use $P_{(n,m)}$, $n, m \in \mathbb{Z}_+$, to construct an $\langle \cdot \rangle_{\Psi}$ -orthonormal family of polynomials $v_{(n,s)}$, $n \in \mathbb{Z}_+$, $s \in \{0, 1, \ldots, n\}$, such that $v_{(n,s)}$ is of degree n and $\mathcal{L}v_{(n,s)} = \lambda_n v_{(n,s)}$. From [18, Section 1.2.2], we may choose $v_{(n,s)}$ such that for each $n \geq 0$, $v_{(n,0)}, v_{(n,1)}, \ldots, v_{(n,n)}$ are given by

$$v_{n,j,1} = h_{n,j,1} P_j^{(1-\frac{4}{\kappa}, n-2j)} (2r^2 - 1) r^{n-2j} \cos((n-2j)\theta), \quad 0 \le j \le \lfloor n/2 \rfloor,$$
$$v_{n,j,2} = h_{n,j,2} P_j^{(1-\frac{4}{\kappa}, n-2j)} (2r^2 - 1) r^{n-2j} \sin((n-2j)\theta), \quad 0 \le j \le \lfloor (n-1)/2 \rfloor,$$

where $P_j^{(1-\frac{4}{\kappa},n-2j)}$ are Jacobi polynomials of index $(1-\frac{4}{\kappa},n-2j)$, (r,θ) is the polar coordinate of (x,y): $x = r \cos \theta$ and $y = r \sin \theta$, and $h_{n,j,i} > 0$ are normalization constants. Using the polar integration and Formula [12, Table 18.3.1]:

$$\int_{-1}^{1} P_{j}^{(\alpha,\beta)}(x)^{2} (1-x)^{\alpha} (1+x)^{\beta} dx = \frac{2^{\alpha+\beta+1} \Gamma(j+\alpha+1) \Gamma(j+\beta+1)}{j! (2j+\alpha+\beta+1) \Gamma(j+\alpha+\beta+1)}$$

with $\alpha = 1 - \frac{4}{\kappa}$ and $\beta = n - 2j$, we compute

$$h_{n,j} := h_{n,j,1} = h_{n,j,2} = \sqrt{\frac{1 + \mathbf{1}_{n \neq 2j}}{\pi}} \cdot \frac{j!(n + 2 - \frac{4}{\kappa})\Gamma(n - j + 2 - \frac{4}{\kappa})}{\Gamma(j + 2 - \frac{4}{\kappa})\Gamma(n - j + 1)}$$

Using the supremum norm (over [-1, 1]) of $P_j^{(\alpha,\beta)}$ ([12, 18.14.1, 18.14.2]):

$$\|P_j^{(\alpha,\beta)}\|_{\infty} = \frac{\Gamma(\max\{\alpha,\beta\}+j+1)}{j!\Gamma(\max\{\alpha,\beta\}+1)}, \quad \text{if } \max\{\alpha,\beta\} \ge -1/2 \text{ and } \min\{\alpha,\beta\} > -1,$$

we get

$$\|v_{n,j,1}\|_{\infty} = \|v_{n,j,2}\|_{\infty} = h_{n,j} \max\left\{\frac{\Gamma(2-\frac{4}{\kappa}+j)}{j!\Gamma(2-\frac{4}{\kappa})}, \frac{\Gamma(n-j+1)}{j!\Gamma(n-2j+1)}\right\}.$$
(3.35)

For t > 0, $(x, y), (x^*, y^*) \in \mathbb{D}$, we define

$$p_t((x,y),(x^*,y^*)) = \sum_{n=0}^{\infty} \sum_{s=0}^n \Psi(x^*,y^*) v_{(n,s)}(x,y) v_{(n,s)}(x^*,y^*) e^{\lambda_n t},$$
(3.36)

and $p_{\infty}(x^*, y^*) = \frac{1}{\pi}(2 - \frac{4}{\kappa})\Psi(x^*, y^*)$, which is the term in the summation for n = s = 0. The following propositions are similar to Lemma 5.1, Lemma 5.2 and Corollary 5.3 of [20]. The proofs use the estimate (3.35) and the orthogonality between $v_{n,s}$ w.r.t. $\langle \cdot \rangle_{\Psi}$. The exponent $1 - \frac{5}{8}\kappa$ in Lemma 3.11 is the λ_1 here.

Lemma 3.11. For any $t_0 > 0$, the series in (3.36) converges uniformly on $[t_0, \infty) \times \overline{\mathbb{D}} \times \overline{\mathbb{D}}$, and there is $C_{t_0} \in (0, \infty)$ depending only on κ and t_0 such that

$$|p_t((x,y),(x^*,y^*)) - p_{\infty}(x^*,y^*)| \le C_{t_0} e^{(1-\frac{5}{8}\kappa)t} p_{\infty}(x^*,y^*), \quad t \ge t_0, \quad (x,y), (x^*,y^*) \in \overline{\mathbb{D}}.$$

Moreover, for any t > 0 and $(x^*, y^*) \in \overline{\mathbb{D}}$,

$$p_{\infty}(x^*, y^*) = \iint_{\mathbb{D}} p_{\infty}(x, y) p_t((x, y), (x^*, y^*)) dx dy.$$

Lemma 3.12. (X,Y) (under \mathbb{P}^c_*) has transition density p_t and invariant density p_{∞} .

Corollary 3.13. (Z_1^u, Z_2^u) under \mathbb{P}^c_* has transition density

$$p_t^Z(\underline{z}, \underline{z}^*) := p_t((\cos_2(z_1 + z_2), \sin_2(z_1 - z_2)), (\cos_2(z_1^* + z_2^*), \sin_2(z_1^* - z_2^*))) \frac{\sin z_1^* + \sin z_2^*}{4}$$

and invariant density $p_{\infty}^{Z}(\underline{z}^{*}) := p_{\infty}(\cos_{2}(z_{1}^{*}+z_{2}^{*}), \sin_{2}(z_{1}^{*}-z_{2}^{*}))\frac{\sin z_{1}^{*}+\sin z_{2}^{*}}{4}$.

We may then use Lemma 3.8 to derive the transition density for (Z_1^u, Z_2^u) under the law \mathbb{P}_c^c . Let $t \in \mathbb{R}_+$. Since $\underline{u}(t)$ is an \mathcal{F} -stopping time, and $\underline{u}(t) \in \mathcal{D}$ iff $T^u > t$, by Lemma 3.8,

$$\frac{d\mathbb{P}_c^c|\mathcal{F}_t^u \cap \{T^u > t\}}{d\mathbb{P}_*^c|\mathcal{F}_t^u \cap \{T^u > t\}} = \frac{M_{*\to c}^u(t)}{M_{*\to c}^u(0)}.$$
(3.37)

Let $\widetilde{G}^u(z_1, z_2) := \widetilde{G}(\pi + z_1, \pi, z_2, 0)$. By (3.20), we get $M^u_{* \to c}(t) = e^{-\alpha_0 t} \widetilde{G}^u(Z^u_1(t), Z^u_2(t))^{-1}$. Combining this with (3.37) and Corollary 3.13, we get the transition density for (Z^u_1, Z^u_2) under the law \mathbb{P}^c_c in the lemma below, which resembles [20, Lemma 5.4].

Lemma 3.14. Under \mathbb{P}^c_c , $\underline{Z}^u(t) := (Z^u_1(t), Z^u_2(t)), 0 \le t < T^u$, has transition density

$$\widetilde{p}_t^Z(\underline{z},\underline{z}^*) := e^{-\alpha_0 t} p_t^Z(\underline{z},\underline{z}^*) \frac{\widetilde{G}^u(\underline{z})}{\widetilde{G}^u(\underline{z}^*)}$$

Here the meaning of the transition density is: if \underline{Z}^u starts with $\underline{z} \in (0,\pi)^2$, then for any t > 0and any bounded measurable function f on $(0,\pi)^2$,

$$\mathbb{E}_{c}^{c}[\mathbf{1}_{\{T^{u}>t\}}f(\underline{Z}^{u}(t))] = \int_{(0,\pi)^{2}} \widetilde{p}_{t}^{Z}(\underline{z},\underline{z}^{*})f(\underline{z}^{*})d\underline{z}^{*}.$$

We compute that $\frac{p_{\infty}^{z}(z_1,z_2)}{\tilde{G}^{u}(z_1,z_2)} = C \cos_2(z_1-z_2)^{2-\frac{8}{\kappa}} \sin_2(z_1+z_2)$ for some constant $C \in (0,\infty)$ depending only on κ . Since $2-\frac{8}{\kappa} > 0$, we may define

$$\mathcal{Z} = \int_{(0,\pi)^2} \frac{p_{\infty}^Z(\underline{z})}{\widetilde{G}^u(\underline{z})} d\underline{z} \in (0,\infty), \qquad (3.38)$$

$$\widetilde{p}_{\infty}^{Z}(\underline{z}) = \frac{1}{\mathcal{Z}} \frac{p_{\infty}^{Z}(\underline{z})}{\widetilde{G}^{u}(\underline{z})}, \quad \underline{z} \in (0, \pi)^{2}.$$
(3.39)

The following lemma resembles [20, Lemma 5.5].

Lemma 3.15. (i) For any t > 0 and $\underline{z}^* \in (0, \pi)^2$,

$$\int_{(0,\pi)^2} \widetilde{p}_{\infty}^Z(\underline{z}) \widetilde{p}_t^Z(\underline{z}, \underline{z}^*) d\underline{z} = \widetilde{p}_{\infty}^Z(\underline{z}^*) e^{-\alpha_0 t}.$$
(3.40)

(ii) If \underline{Z}^u starts from $\underline{z} \in (0, \pi)^2$, then as $t \to \infty$,

$$\mathbb{P}_c^c[T^u > t] = \mathcal{Z}\widetilde{G}^u(\underline{z})e^{-\alpha_0 t}(1 + O(e^{(1 - \frac{5}{8}\kappa)t})),$$

$$\widetilde{p}_t^Z(\underline{z}, \underline{z}^*) = \mathbb{P}_c^c[T^u > t]\widetilde{p}_{\infty}^Z(\underline{z}^*)(1 + O(e^{(1 - \frac{5}{8}\kappa)t})),$$

where the implicit constant depends only on κ . The two formulas together imply that

$$\widetilde{p}_t^Z(\underline{z},\underline{z}^*) = \mathcal{Z}\widetilde{G}^u(\underline{z})\widetilde{p}_{\infty}^Z(\underline{z}^*)e^{-\alpha_0 t}(1 + O(e^{(1-\frac{5}{8}\kappa)t})), \quad \underline{z},\underline{z}^* \in (0,\pi)^2.$$
(3.41)

4 Proof of the Main Theorem

We now start the proof of Theorem 1.1. By conformal invariance, Koebe's distortion theorem and reflection symmetry, we may assume that $D = \mathbb{D}$, $z_0 = 0$, and $a_j = e^{iw_j}$ and $b_j = e^{iv_j}$, j = 1, 2, where $w_1 > v_1 > w_2 > v_2 > w_1 - 2\pi$. Then we have $G_{D;a_1,b_1,a_2,b_2}(0) = \widetilde{G}(w_1, v_1, w_2, v_2)$. Let γ_1 be as in the theorem, and let γ_2 be a time-reversal of γ_1 . By reversibility, γ_2 is a chordal SLE_{κ} in \mathbb{D} from e^{iw_2} to e^{iw_1} . We may assume that for $j = 1, 2, \gamma_j$ is parametrized by some capacity viewed from a_{3-j} . This means that there is a conformal map $f_j : \mathbb{D} \xrightarrow{\text{Conf}} \mathbb{H}$, which sends a_{3-j} to ∞ such that hcap₂ $f_j(\gamma_j[0,t]) = t$ for $t \ge 0$. For $r \in (0,1)$, let E(r) denote the event that $\gamma \cup A_1 \cup A_2$ has a cut point that lies in $\{|z| < r\}$, which also depends on w_1, v_1, w_2, v_2 , and let $P(w_1, v_2, w_2, v_2; r) = \mathbb{P}[E(r)]$. We need to show that there is a constant $C_0 > 0$ depending only on κ such that

$$P(w_1, v_1, w_2, v_2; r) = C_0 \widetilde{G}^u(w_1, v_1, w_2, v_2) r^{\alpha_0} (1 + O(r^{\beta_0})), \quad \text{as } r \to 0.$$
(4.1)

For j = 1, 2, let \widehat{S}_j be the first time that γ_j separates 0 from any of $e^{iw_{3-j}}, e^{iv_1}, e^{iv_2}$ in \mathbb{D} . Let $v_j(t) = \operatorname{dcap}(\gamma_j[0,t]), \ 0 \leq t < \widehat{S}_j, \ \widehat{T}_j = v_j(\widehat{S}_j), \ u_j = v_j^{-1} : [0,\widehat{T}_j) \to [0,\widehat{S}_j)$, and $\eta_j = \gamma_j \circ u_j$. By [17], η_j is a radial SLE_{κ}($\kappa - 6, 0, 0$) curve in \mathbb{D} started from e^{iw_j} with force points $e^{iw_{3-j}}, e^{iv_1}, e^{iv_2}$. Let \widehat{w}_j be a radial Loewner driving function of η_j with $\widehat{w}_j(0) = w_j$. We will use the notation in Section 3.1 for the radial Loewner curves η_1, η_2 in this section.

For j = 1, 2, let \mathcal{G}^j and \mathcal{F}^j be respectively the filtrations generated by γ_j and η_j . Then \widehat{S}_j is an \mathcal{G}^j -stopping time. Let $\overline{\mathcal{F}}^j$ and $\overline{\mathcal{G}}^j$ be respectively the right-continuous augmentation of \mathcal{F}^j and \mathcal{G}^j . We extend u_j to $[0, \infty]$ such that $u_j \equiv \infty$ on $[\widehat{T}_j, \infty]$. The following is a well-known proposition about time-change.

Proposition 4.1. For $j \in \{1, 2\}$, if τ is an $\overline{\mathcal{F}}^j$ -stopping time, then $u_j(\tau)$ is an $\overline{\mathcal{G}}^j$ -stopping time, and $\overline{\mathcal{F}}^j_{\tau} \subset \overline{\mathcal{G}}^j_{u_j(\tau)}$.

Fix $j \neq k \in \{1, 2\}$. Let τ_k be an \mathcal{F}^k -stopping time less than \widehat{T}_k and let $\sigma_k = u_k(\tau_k)$, which by Proposition 4.1 is an $\overline{\mathcal{G}}_{\sigma_k}^k$ -stopping time. By DMP of chordal SLE_{κ}, conditionally on $\overline{\mathcal{G}}_{\sigma_k}^k$, the part of γ_k after σ_k is a chordal SLE_{κ} in $\mathbb{D} \setminus K_k(\tau_k)$ from $\gamma_k(\sigma_k) = \eta_k(\tau_k)$ to e^{iw_j} . Since γ_j is a time-reversal of γ_k , by reversibility of chordal SLE_{κ}, conditionally on $\overline{\mathcal{G}}_{\sigma_k}^k$, the part of γ_j up to the time that it hits $\eta_k(\tau_k)$ is a chordal SLE_{κ} in $\mathbb{D} \setminus K_k(\tau_k)$ from e^{iw_j} to $\eta_k(\tau_k)$. Recall that η_j is a time-change of the part of γ_j up to the first time that it separates 0 from any of $e^{iw_k}, e^{iv_1}, e^{iv_2}$ in \mathbb{D} . So the part of η_j up to the first time that it hits $\eta_k[0, \tau_k]$ is a time-change of the part of γ_j up to the first time that it separates 0 from any of $e^{iw_k}, e^{iv_1}, e^{iv_2}$ in $\mathbb{D} \setminus K_k(\tau_k)$.

Since $g_k(\tau_k, \cdot)$ maps $\mathbb{D}\setminus K_k(\tau_k)$ conformally onto \mathbb{D} , sends $\eta_k(\tau_k)$ to $e^{i\widehat{w}_k(\tau_k)}$, and is measurable w.r.t. $\mathcal{F}^k_{\tau_k} \subset \overline{\mathcal{G}}^k_{\sigma_k}$, by conformal invariance of chordal SLE, we see that, conditionally on $\overline{\mathcal{G}}^k_{\sigma_k}$, the $g_k(\tau_k, \cdot)$ -image of the part of η_j up to the first time that it hits $\eta_k[0, \tau_k]$ is a time-change of a chordal SLE_{κ} in \mathbb{D} from $g_k(\tau_k, e^{iw_j})$ to $e^{i\widehat{w}_k(\tau_k)}$ up to the first time that it separates 0 from any of $e^{i\widehat{w}_k(\tau_k)}, g_k(\tau_k, e^{iv_1}), g_k(\tau_k, e^{iv_2})$ in \mathbb{D} , which by [17] is a time-change of a radial SLE_{κ}($\kappa - 6, 0, 0$) curve in \mathbb{D} started from $g_k(\tau_k, e^{iw_j})$ with force points $e^{i\widehat{w}_k(\tau_k)}, g_k(\tau_k, e^{iv_1}), g_k(\tau_k, e^{iv_2})$. Since $g^k_{\tau_k}$ and $e^{i\widehat{w}_k(\tau_k)}$ are $\mathcal{F}^k_{\tau_k}$ -measurable, the above statement holds with $\mathcal{F}^k_{\tau_k}$ in place of $\overline{\mathcal{G}}^k_{\sigma_k}$. So we obtain the commutation coupling in Proposition 3.7, and the law of $(\widehat{w}_1, \widehat{w}_2)$ is \mathbb{P}^c_c .

Let \mathcal{F} and \mathcal{G} be respectively the separable \mathbb{R}^2_+ -indexed filtration generated by $\mathcal{F}^1, \mathcal{F}^2$ and $\mathcal{G}^1, \mathcal{G}^2$, and let $\overline{\mathcal{F}}$ and $\overline{\mathcal{G}}$ be their right-continuous augmentation. Let $\underline{\widehat{T}} = (\widehat{T}_1, \widehat{T}_2)$ and $\underline{\widehat{S}} = (\widehat{S}_1, \widehat{S}_2)$. Define u_{\otimes} on $[0, \infty]^2$ such that if $(t_1, t_2) \in [0, \widehat{T}_1) \times [0, \widehat{T}_2), u_{\otimes}(t_1, t_2) = (u_1(t_1), u_2(t_2)),$ and otherwise $u_{\otimes}(t_1, t_2) = (\infty, \infty)$.

Lemma 4.2. For any $\overline{\mathcal{F}}$ -stopping time \underline{T} , $u_{\otimes}(\underline{T})$ is an $\overline{\mathcal{G}}$ -stopping time and $\overline{\mathcal{F}}_{\underline{T}} \subset \overline{\mathcal{G}}_{u_{\otimes}(\underline{T})}$. *Proof.* Let $A \in \overline{\mathcal{F}}_T$ and $\underline{s} = (s_1, s_2) \in \mathbb{R}^2_+$. We have

$$A \cap \{u_{\otimes}(\underline{T}) < \underline{s}\} = \bigcup_{\underline{p} \in \mathbb{Q}^2_+} (A \cap \{\underline{T} < \underline{p}\} \cap \{u_{\otimes}(\underline{p}) < \underline{s}\}).$$
(4.2)

By Proposition 2.9, $A \cap \{\underline{T} < \underline{p}\} \in \mathcal{F}_{\underline{p}}$. For any j = 1, 2 and $B_j \in \mathcal{F}_{p_j}^j$, by Proposition 4.1, $B_j \cap \{u_j(p_j) < s_j\} \in \mathcal{G}_{s_j}^j$. By a monotone class argument, we get $B \cap \{u_{\otimes}(\underline{p}) < \underline{s}\} \in \mathcal{G}_{\underline{s}}$ for any $B \in \mathcal{F}_{\underline{p}}$. So $A \cap \{\underline{T} < \underline{p}\} \cap \{u_{\otimes}(\underline{p}) < \underline{s}\} \in \mathcal{G}_{\underline{s}}$. By (4.2) we get $A \cap \{u_{\otimes}(\underline{T}) < \underline{s}\} \in \mathcal{G}_{\underline{s}}$. Since this holds for all $A \in \overline{\mathcal{F}}_{\underline{T}}$ and $\underline{s} \in \mathbb{R}^2_+$, by Proposition 2.9, we see that $u_{\otimes}(\underline{T})$ is an $\overline{\mathcal{G}}$ -stopping time and $\overline{\mathcal{F}}_{\underline{T}} \subset \overline{\mathcal{G}}_{u_{\otimes}(\underline{T})}$.

Lemma 4.3. Let 0 < r < 1/4. Let $\underline{T} = (T_1, T_2)$ be an $\overline{\mathcal{F}}$ -stopping time such that $T_j < -\log(4r)$, j = 1, 2, and when $\underline{T} \in \mathcal{D}$, $m(\underline{T}) < -\log(4r)$. On the event $\{\underline{T} \in \mathcal{D}\}$, let $R_+ > R_- \in (0, 1)$ be such that $\frac{R_{\pm}}{(1\pm R_{\pm})^2} = e^{m(\underline{T})}r$, which exist uniquely because $e^{m(\underline{T})}r < 1/4$. Then

$$\mathbb{P}[E(r)|\overline{\mathcal{F}}_{\underline{T}}] \leq \mathbf{1}_{\{\underline{T}\in\mathcal{D}\}} \cdot P(W_1(\underline{T}), V_1(\underline{T}), W_2(\underline{T}), V_2(\underline{T}); R_{\pm}).$$

$$(4.3)$$

Here and below the symbol \leq means that when we choose + and - in the \pm on the RHS, the inequality holds with \leq and \geq , respectively.

Proof. For j = 1, 2, let τ_r^j and $\tilde{\tau}_r^j$ be respectively the first time that η_j and γ_j visits $\{|z| \leq r\}$. First, suppose the event E(r) happens. Then γ_1 and γ_2 do reach $\{|z| \leq r\}$, i.e., $\tilde{\tau}_r^j < \infty, j = 1, 2$. By duality of SLE (cf. [10, Theorem 1.4] and [11, the discussion before Theorem 1.5]), the left boundary and right boundary of γ_1 are two (simple) flow lines, whose intersection is the cut-set of η_1 . In fact, the two simple flow lines are $\operatorname{SLE}_{\kappa'}(\kappa'-4,\frac{\kappa'}{2}-2)$ and $\operatorname{SLE}_{\kappa'}(\frac{\kappa'}{2}-2,\kappa'-4)$ curves, where $\kappa' = 16/\kappa$, starting from $\eta_1(0)$ with the force points being $\eta_1(0)^+$ and $\eta_1(0)^-$. We do not need the laws of these curves. Thus, for $j = 1, 2, \gamma_j[0, \tilde{\tau}_r^j]$ does not intersect A_{3-j} or disconnect 0 from $\partial \mathbb{D}$; and $\gamma_1[0, \tilde{\tau}_r^1] \cap \gamma_2[0, \tilde{\tau}_r^2] = \emptyset$. The former condition implies that $\tilde{\tau}_r^j < \hat{S}_j$, and so $\tau_r^j = v_j(\tilde{\tau}_r^j) < \hat{T}_j$; and the latter implies that $\eta_1[0, \tau_r^1] \cap \eta_2[0, \tau_r^2] = \emptyset$. So we get $(\tau_r^1, \tau_r^2) \in \mathcal{D}$. Since $\mathbb{D} \setminus K_j(\tau_r^j) \supset \{|z| < r\}$, by Koebe's 1/4 theorem, $\tau_r^j \ge -\log(4r) \ge T_j, j = 1, 2$. Since $(\tau_r^1, \tau_r^2) \in \mathcal{D}$ and \mathcal{D} is an HC region, we get $\underline{T} \in \mathcal{D}$. Thus, $E(r) \subset \{\underline{T} \in \mathcal{D}\}$.

Now we assume that $\{\underline{T} \in \mathcal{D}\}$ (instead of E(r)) happens. Let $\tilde{\gamma}_1$ denote the part of γ_1 between $\eta_1(T_1)$ and $\eta_2(T_2)$. Let $\tilde{A}_j := A_j \cup \eta[0, T_j], j = 1, 2$. Then $\gamma_1 \cup A_1 \cup A_2 = \tilde{\gamma}_1 \cup \tilde{A}_1 \cup \tilde{A}_2$. By Lemma 4.2, $u_{\otimes}(\underline{T})$ is an $\overline{\mathcal{G}}$ -stopping time. By Theorem A.1, conditionally on $\overline{\mathcal{G}}_{u_{\otimes}(\underline{T})}$ and the event $\{\underline{T} \in \mathcal{D}\}, \tilde{\gamma}$ is a time-change of a chordal SLE_{κ} in $\mathbb{D} \setminus K(\underline{T})$ from $\eta_1(T_1) = \gamma_1(u_1(T_1))$ to $\eta_2(T_2) = \gamma_2(u_2(T_2))$. Since the event $\{\underline{T} \in \mathcal{D}\}$ and the random elements $\mathbb{D} \setminus K(\underline{T})$ and $\eta_j(T_j), j = 1, 2$, are all $\overline{\mathcal{F}}_{\underline{T}}$ -measurable, the above statement holds with $\overline{\mathcal{F}}_{\underline{T}}$ in place of $\overline{\mathcal{G}}_{u_{\otimes}(\underline{T})}$. Since $g(\underline{T}, \cdot)$ is $\overline{\mathcal{F}}_{\underline{T}}$ -measurable, and sends $\eta_1(T_1)$ and $\eta_2(T_2)$ respectively to $e^{iW_1(\underline{T})}$ and $e^{iW_2(\underline{T})}$, we conclude that, conditionally on $\overline{\mathcal{F}}_{\underline{T}}$ and the event $\{\underline{T} \in \mathcal{D}\}, g(\underline{T}, \tilde{\gamma})$ is a time-change of a chordal SLE_{κ} in \mathbb{D} from $e^{iW_1(\underline{T})}$ to $e^{iW_2(\underline{T})}$.

For j = 1, 2, let $A_j(\underline{T})$ be the boundary arc of \mathbb{D} connecting $e^{iV_1(\underline{T})}$ and $e^{iV_2(\underline{T})}$ which contains $e^{iW_j(\underline{T})}$. Since $g(\underline{T}, \cdot)$ sends e^{iv_1} and e^{iv_2} respectively to $e^{iV_1(\underline{T})}$ and $e^{iV_2(\underline{T})}$, that $\tilde{\gamma}_1 \cup \tilde{A}_1 \cup \tilde{A}_2$ contains a cut point in $\{|z| < r\}$ is then equivalent to that $g(\underline{T}, \tilde{\gamma}_1) \cup A_1(\underline{T}) \cup A_2(\underline{T})$ has a cut point contained in $g(\underline{T}, \{|z| < r\})$. Recall that $g(\underline{T}, \cdot)$ maps $\mathbb{D} \setminus K(\underline{T})$ conformally onto \mathbb{D} , fixes 0, and has derivative $e^{m(\underline{T})}$ at 0. By Koebe's distortion theorem applied to $e^{m(\underline{T})}g(\underline{T}, \cdot)^{-1}$,

$$\Big\{|z| < \frac{e^{-\mathbf{m}(\underline{T})}R}{(1+R)^2}\Big\} \subset g(\underline{T}, \cdot)^{-1}(\{|z| < R\}) \subset \Big\{|z| < \frac{e^{-\mathbf{m}(\underline{T})}R}{(1-R)^2}\Big\}, \quad \forall R \in (0,1).$$

Taking $R = R_{\pm}$, we get $\{|z| < R_{-}\} \subset g(\underline{T}, \{|z| < r\}) \subset \{|z| < R_{+}\}$. Then (4.3) follows immediately.

For $z_1, z_2 \in (0, \pi)$, we write $P^u(z_1, z_r; r)$ for $P(\pi + z_1, \pi, z_2, 0; r)$. By rotation symmetry, if $v_1 - v_2 = \pi$, then $P(w_1, v_1, w_2, v_2; r) = P^u(w_1 - v_1, w_2 - v_2; r)$.

Lemma 4.4. Let $\widetilde{p}_t^Z(\underline{z},\underline{z}^*)$, $t \in (0,\infty)$, $\underline{z}, \underline{z}^* \in (0,\pi)^2$, be the transition density given by Lemma 3.14. Let $r \in (0,1/4)$ and $t_0 \in (0, -\log(4r))$. Let $R_+, R_- \in (0,1)$ be such that $\frac{R_{\pm}}{(1\pm R_{\pm})^2} = e^{t_0}r$. Then for any $\underline{z} \in (0,\pi)^2$,

$$P^{u}(\underline{z};r) \leq \int_{(0,\pi)^{2}} P^{u}(\underline{z}^{*},R_{\pm}) \widetilde{p}_{t_{0}}^{Z}(\underline{z},\underline{z}^{*}) d\underline{z}^{*}.$$
(4.4)

Proof. Fix $\underline{z} = (z_1, z_2) \in (0, \pi)^2$. Let $w_1 = \pi + z_1$, $v_1 = \pi$, $w_2 = z_2$, $v_2 = 0$. Then $\mathbb{P}[E(r)] = P^u(\underline{z}, r)$. Since $v_1 - v_2 = \pi$, we may define the curve $\underline{u} = (u_1, u_2)$ as in Section 3.3. If $\underline{u}(t_0) \in \mathcal{D}$, then $u_j(t_0) \leq \mathrm{m}(\underline{u}(t_0)) = t_0$, j = 1, 2. If $\underline{u}(t_0) \notin \mathcal{D}$, then $t_0 \geq T^u$ and $\underline{u}(t_0) = \underline{u}(T^u) = \lim_{t \uparrow T^u} \underline{u}(t) \leq T^u \leq t_0$ because for $0 \leq t < T^u$, we have $\underline{u}(t) \in \mathcal{D}$, and so $u_j(t) \leq t \leq T^u$, j = 1, 2. Thus, in any case we have $u_j(t_0) \leq t_0$, j = 1, 2.

Apply Lemma 4.3 to the \mathcal{F} -stopping time $\underline{u}(t_0)$. Since $m(\underline{u}(t_0)) = t_0$ when $\underline{u}(t_0) \in \mathcal{D}$, the R_{\pm} here agree with the R_{\pm} in Lemma 4.3. From $V_1^u(t_0) - V_2^u(t_0) = \pi$, we then get

$$\mathbb{P}[E(r)|\mathcal{F}_{t_0}^u] \leq \mathbf{1}_{\underline{u}(t_0)\in\mathcal{D}}P^u(Z_1^u(t_0), Z_2^u(t_0); R_{\pm}).$$

By integration we get

$$P^{u}(\underline{z};r) = \mathbb{P}[E(r)] \leq \mathbb{E}[\mathbf{1}_{\underline{u}(t_{0})\in\mathcal{D}}P^{u}(\underline{Z}^{u}(t_{0});R_{\pm})].$$

$$(4.5)$$

Since $(\widehat{w}_1, \widehat{w}_2)$ follows the law $\mathbb{P}_c^c, \widetilde{p}_t^Z(\underline{z}, \underline{z}^*)$ is the transition density of the process $(\underline{Z}^u(t))$. So the density of $\underline{Z}^u(t_0)$ is $\widetilde{p}_{t_0}^Z(\underline{z}, \cdot)$, and the RHS of (4.5) agrees with the RHS of (4.4).

Let $\widetilde{p}_{\infty}^{Z}$ be given by (3.39). Define $\overline{P}(r) = \int_{(0,\pi)^2} P^u(\underline{z},r) \widetilde{p}_{\infty}^{Z}(\underline{z}) d\underline{z}, r \in (0,1).$

Lemma 4.5. We have $\overline{P}(r) = \widehat{c}r^{\alpha_0}(1 + O(r))$ as $r \downarrow 0$, where $\widehat{c} \ge 0$ is a constant depending only on κ , and the implicit constants in O(r) depend only on κ .

Proof. Let r, t_0, R_{\pm} be as in Lemma 4.4. By integrating both sides of (4.4) against $\widetilde{p}_{\infty}^Z(\underline{z})$ and using (3.40), we get $\overline{P}(r) \leq e^{-\alpha_0 t_0} \overline{P}(R_{\pm})$. Let $Q(r) = r^{-\alpha_0} \overline{P}(r)$. Since $\frac{R_{\pm}}{(1 \pm R_{\pm})^2} = e^{t_0} r$, we get

$$Q(r) \leq (1 \pm R_{\pm})^{2\alpha_0} Q(R_{\pm}).$$
 (4.6)

Let $R \in (0, \frac{1}{6}]$ and $r \in (0, \frac{R}{2}]$. Then $r < \frac{R}{(1+R)^2} < \frac{R}{(1-R)^2} < \frac{1}{4}$. So there are $t_+, t_- \in (0, -\log(4r))$ such that $\frac{R}{(1\pm R)^2} = e^{t\pm}r$. By (4.6) (which does not contain t_0),

$$Q(r) \leq (1 \pm R)^{2\alpha_0} Q(R), \quad \text{if } 0 < r \leq R/2 < R \leq 1/6.$$
 (4.7)

Thus, for any $a \in (0, \frac{1}{12}]$ and $r_1, r_2 \in (0, a]$, we have $(1 + 2a)^{-2\alpha_0}Q(r_1) \leq (1 - 2a)^{-2\alpha_0}Q(r_2)$. This implies that $\lim_{r \downarrow 0} Q(r)$ converges. Let the limit be denoted by \hat{c} , which is nonnegative and depends only on κ . Letting $r \downarrow 0$ in (4.7), we get $Q(R) \geq \hat{c}(1 \pm R)^{-2\alpha_0}$, if $R \leq 1/6$. This immediately implies the conclusion.

Recall the \mathcal{Z} defined in (3.38). Let $C_0 = \widehat{c}\mathcal{Z}$. Then C_0 is a nonnegative constant depending only on κ . We are now ready to prove that (4.1) holds for such C_0 . First suppose $v_1 - v_2 = \pi$. Then $P(w_1, v_1, w_2, v_2; r) = P^u(z_1, z_2; r)$ and $\widetilde{G}(w_1, v_1, w_2, v_2) = \widetilde{G}^u(z_1, z_2)$, where $z_j = w_j - v_j$. Recall that $\beta_0 = 1 - \frac{8}{5\kappa}$. Let $R = r^{\beta_0}$. When r is small enough (depending on β_0), we may choose $t_+, t_- \in (0, -\log(4r))$ such that $\frac{R}{(1\pm R)^2} = e^{t\pm}r$. By (3.41) and Lemmas 4.4 and 4.5,

$$P^{u}(\underline{z};r) \stackrel{\leq}{\leq} \int_{(0,\pi)^{2}} P^{u}(\underline{z}^{*},R) \widetilde{p}^{Z}_{t\pm}(\underline{z},\underline{z}^{*}) d\underline{z}^{*} = \mathcal{Z} \widetilde{G}^{u}(\underline{z}) e^{-\alpha_{0}t_{\pm}} \overline{P}(R) (1 + O(e^{(1-\frac{5}{8}\kappa)t_{\pm}}))$$

$$= C_0 \widetilde{G}^u(\underline{z}) e^{-\alpha_0 t_{\pm}} R^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}})) = C_0 \widetilde{G}^u(\underline{z}) r^{\alpha_0} (1 + O(R) + O(e^{(1 - \frac{5}{8}\kappa)t_{\pm}}))$$

Since $O(R) = O(e^{(1-\frac{5}{8}\kappa)t_{\pm}}) = O(r^{\beta_0})$, the above formula implies that

$$P^{u}(\underline{z};r) = C_0 \tilde{G}^{u}(\underline{z}) r^{\alpha_0} (1 + O(r^{\beta_0})), \quad \text{as } r \downarrow 0.$$

$$(4.8)$$

So we have finished the proof of (4.1) in the case $v_1 - v_2 = \pi$.

Now suppose $\theta(\underline{0}) = v_1 - v_2 < \pi$. By (3.22), θ is increasing in t_2 . Let τ_2 be the first $t_2 \in (0, \hat{T}_2)$ such that $\theta(0, \tau_2) = \pi$, if this time exists; otherwise let $\tau_2 = \hat{T}_2$. From (3.22) and that m(0,t) = t, we see that $\cos(\theta(0,t)/4) \leq e^{-t/2} \cos(\theta(\underline{0})/4) \leq e^{-t/2}$, $0 \leq t < \hat{T}_2$. Then for $0 \leq t < \tau_2$, $t \leq -2\log\cos(\theta(0,t)/4) \leq \log(2)$, which implies that $\tau_2 \leq \log(2)$. Suppose $r \in (0,1/8)$. Then $\tau_2 \leq -\log(4r)$. Let $R_{\pm} \in (0,1)$ be such that $\frac{R_{\pm}}{(1\pm R_{\pm})^2} = e^{\tau_2}r$. Then $R_{\pm} = e^{\tau_2}r(1+O(r))$ as $r \downarrow 0$. We now apply Lemma 4.3 to $\underline{T} = (0,\tau_2)$. Since $\theta(0,\tau_2) = \pi$ when $(0,\tau_2) \in \mathcal{D}$, using (4.8), we get, as $r \downarrow 0$,

$$\mathbb{P}[E(r)|\mathcal{F}_{(0,\tau_2)}] \leq \mathbf{1}_{\{(0,\tau_2)\in\mathcal{D}\}} P^u(\underline{Z}(0,\tau_2); R_{\pm}) = \mathbf{1}_{\{(0,\tau_2)\in\mathcal{D}\}} P^u(\underline{Z}(0,\tau_2); e^{\tau_2}r(1+O(r)))$$
$$= C_0 \mathbf{1}_{\{(0,\tau_2)\in\mathcal{D}\}} e^{\alpha_0\tau_2} \widetilde{G}^u(\underline{Z}(0,\tau_2)) r^{\alpha_0}(1+O(r^{1-\frac{8}{5\kappa}})),$$

which together with (3.20) implies that

$$\mathbb{P}[E(r)] = C_0 r^{\alpha_0} (1 + O(r^{1 - \frac{8}{5\kappa}})) \mathbb{E}[\mathbf{1}_{\{(0,\tau_2) \in \mathcal{D}\}} e^{\alpha_0 \operatorname{m}(0,\tau_2)} \widetilde{G}^u(\underline{Z}(0,\tau_2))]$$
$$= C_0 r^{\alpha_0} (1 + O(r^{1 - \frac{8}{5\kappa}})) \mathbb{E}[\mathbf{1}_{\{(0,\tau_2) \in \mathcal{D}\}} M_{* \to c}(0,\tau_2)^{-1}]$$

Since $\mathbb{E} = \mathbb{E}_c^c$, by Lemmas 3.6 and 3.8,

$$\mathbb{E}[\mathbf{1}_{\{(0,\tau_2)\in\mathcal{D}\}}M_{*\to c}(0,\tau_2)^{-1}] = M_{*\to c}(\underline{0})^{-1}\mathbb{P}^c_*[(0,\tau)\in\mathcal{D}] = M_{*\to c}(\underline{0})^{-1} = \widetilde{G}(w_1,v_1,w_2,v_2).$$

The last two displayed formulas together imply that (4.1) holds in the case $v_1 - v_2 < \pi$.

The case that $v_1 - v_2 > \pi$ can be handled in a similar way. By (3.22), θ is decreasing in t_1 . We may apply Lemma 4.3 to $\underline{T} = (\tau_1, 0)$, where τ_1 is the first $t_1 \in [0, \hat{T}_1)$ such that $\theta(t_1, 0) = \pi$, if this time exists; and $\tau_1 = \hat{T}_1$ otherwise. Thus, (4.1) holds in all cases.

It remains to show that $C_0 > 0$. Suppose $C_0 = 0$. Then for any $z_0 \in D$, we have $\mathbb{P}[E_{z_0}(r)] = 0$ when r > 0 is small enough, which implies that a.s. $\gamma_1 \cup A_1 \cup A_2$ does not have a cut point. However, since $\kappa \in (4, 8)$, γ_1 does have a cut point, and when this happens, one can always find b_1 and b_2 in some deterministic countable dense subset of $\partial D \setminus \{a_1, a_2\}$ such that $\gamma_1 \cup A_1 \cup A_2$ contains a cut point, where for $j = 1, 2, A_j$ is the connected component of $\partial \mathbb{D} \setminus \{b_1, b_2\}$ that contains a_j . This implies that for some choice of b_1, b_2 , the probability that $\gamma_1 \cup A_1 \cup A_2$ has a cut point is positive. The contradiction shows that $C_0 > 0$. The proof is now complete.

Appendices

A Domain Markov Property in Two Directions

Let $\kappa \in (0, 8]$. In this appendix, we combine the reversibility of chordal SLE_{κ} with the usual (one-directional) DMP to derive a two-directional DMP.

Let D be a simply connected domain with locally connected boundary. Let a_1, a_2 be distinct prime ends of D. For j = 1, 2, let γ_j be a chordal SLE_{κ} curve in D from a_j to a_{3-j} , parametrized by the capacity viewed from a_{3-j} , such that γ_1 and γ_2 are time-reversal of each other. Such γ_1, γ_2 exist by reversibility of SLE_{κ}. Let ϕ_2 be the decreasing auto-homeomorphism of $[0, \infty]$ such that $\gamma_1 \circ \phi_2 = \gamma_2$. For j = 1, 2 and $t \ge 0$, let D_t^j be the connected component of $D \setminus \gamma_j[0, t]$, which shares the prime end a_{3-j} with D. We may view $\gamma_j(t)$ as a prime end of D_t^j . For j = 1, 2, let $\mathcal{D}_j = \{(t_1, t_2) \in \mathbb{R}^2_+ : \gamma_j(t_j) \notin \gamma_{3-j}[0, t_{3-j}]\}$ and $\mathcal{D}_{\cap} = \mathcal{D}_1 \cap \mathcal{D}_2$. When $t_1 < \phi_2(t_2)$, let D_t denote the connected component of $D \setminus \bigcup_{j=1,2} \gamma_j[0, t_j]$ whose closure contains $\gamma_1[t_1, \phi_2(t_2)]$. If $\underline{t} \in \mathcal{D}_{\cap}, D_{\underline{t}}$ shares the prime end $\gamma_j(t_j)$ with $D_{t_j}^j$, j = 1, 2. For j = 1, 2, let \mathcal{G}^j be the \mathbb{R}_+ -indexed filtration generated by γ_j . Let \mathcal{G} be the separable \mathbb{R}^2_+ -indexed filtration generated by \mathcal{G}^1 and \mathcal{G}^2 , and $\overline{\mathcal{G}}$ be the right-continuous augmentation of \mathcal{G} . We are going to prove the following theorem, which has been used in Lemma 4.3.

Theorem A.1. If $\underline{T} = (T_1, T_2)$ is an $\overline{\mathcal{G}}$ -stopping time, then conditionally on $\overline{\mathcal{G}}_{\underline{T}}$ and the event $\{\underline{T} \in \mathcal{D}_{\cap}\}, \gamma_1|_{[T_1,\phi_2(T_2)]}$ is a time-change of a chordal SLE_{κ} curve from the prime end $\gamma_1(T_1)$ to the prime end $\gamma_2(T_2)$ in $D_{\underline{T}}$.

Remark A.2. A more specific statement of the theorem is the following. The event $\{\underline{T} \in \mathcal{D}_{\cap}\}$ is $\overline{\mathcal{G}}_T$ -measurable, and on this event, there are

- an $\overline{\mathcal{G}}_{\underline{T}}$ -measurable conformal map $h_{\underline{T}}$ from \mathbb{H} onto $D_{\underline{T}}$, which has continuation to $\overline{\mathbb{H}}^{\#}$, and sends 0 and ∞ respectively to the prime ends $\gamma_1(\overline{T}_1)$ and $\gamma_2(T_2)$,
- a standard chordal SLE_{κ} curve $\beta_1^{\underline{T}}$ conditionally independent of $\overline{\mathcal{G}}_{\underline{T}}$, and
- an increasing homeomorphism u_1^T from $[0,\infty]$ onto $[T_1,\phi_2(T_2)]$,

such that $\gamma_1 \circ u_1^{\underline{T}} = h_{\underline{T}} \circ \beta_1^{\underline{T}}$ a.s. on $\{\underline{T} \in \mathcal{D}_{\cap}\}$.

The random curve $\beta_1^{\underline{T}}$ is only defined on the event $\{\underline{T} \in \mathcal{D}_{\cap}\}$. By saying that $\beta_1^{\underline{T}}$ is a standard chordal SLE_{κ} curve conditionally independent of $\overline{\mathcal{G}}_{\underline{T}}$, we mean that under the new probability measure $\mathbb{P}[\cdot|\{\underline{T} \in \mathcal{D}_{\cap}\}]$, where \mathbb{P} is the original underlying probability measure, $\beta_1^{\underline{T}}$ has the law of a standard chordal SLE_{κ} curve and is independent of $\overline{\mathcal{G}}_{\underline{T}}$.

Remark A.3. In the statement of the theorem, we condition on the σ -algebra $\overline{\mathcal{G}}_{\underline{T}}$ and an event $\{\underline{T} \in \mathcal{D}_{\cap}\}$, which is $\overline{\mathcal{G}}_{\underline{T}}$ -measurable. The bigger the event, the stronger the statement is. For the application of Theorem A.1 in Lemma 4.3, we only need a weaker result, in which \mathcal{D}_{\cap} is replaced by $\mathcal{D}_0 = \{(t_1, t_2) \in \mathbb{R}^2_+ : \gamma_1[0, t_1] \cap \gamma_2[0, t_2] = \emptyset\}$, which is a proper subset of \mathcal{D}_{\cap} in the case $\kappa \in (4, 8]$. The strongest possible form of the theorem for $\kappa \in (4, 8)$ is when \mathcal{D}_{\cap} is replaced by the bigger set $\{(t_1, t_2) \in \mathbb{R}^2_+ : t_1 < \phi_2(t_2)\}$. In that case, the statement has to be modified since $D_{\underline{T}}$ may not share the prime end $\gamma_j(T_j)$ with $D^j_{T_j}$, and so we may not view $\gamma_j(T_j)$ as a prime end of D_T . The proof of such a strong statement remains open to the author.

Remark A.4. Theorem A.1 is intuitively correct, but still requires a proof, which we could not find in the literature. We make a complete (nontrivial) proof of the theorem here because it is a solid step in the development of the multi-time-parameter framework.

The essential property of the chordal $\operatorname{SLE}_{\kappa}$ curve for $\kappa \in (0, 8)$ used in the proof is the fact that such a curve commutes with its time reversal, which is also a chordal $\operatorname{SLE}_{\kappa}$ curve. Thus, the argument can be easily extended to any commuting pair of $\operatorname{SLE-type}$ curves. Taking the η_1 and η_2 in Proposition 3.1 for example, we have the following result. Let \mathcal{D} be as defined in Section 3.1. Let \mathcal{F}^j be the \mathbb{R}_+ -indexed filtration generated by η_j , j = 1, 2, and let $\overline{\mathcal{F}}$ be the right-continuous augmentation of the separable \mathbb{R}^2_+ -indexed filtration generated by \mathcal{F}^1 and \mathcal{F}^2 . Then for any $\overline{\mathcal{F}}$ -stopping time $\underline{T} = (T_1, T_2)$, conditionally on $\overline{\mathcal{F}}_{\underline{T}}$ and $\{\underline{T} \in \mathcal{D}\}$, for j = 1, 2, the $g(\underline{T}, \cdot)$ -images of $\eta_j(T_j + \cdot)$ is a time-change of a radial $\operatorname{SLE}_{\kappa}(2, \underline{\rho})$ curve in \mathbb{D} started from $e^{iW_j(\underline{T})}$ aimed at 0 with force points $e^{iW_{3-j}(\underline{T})}, e^{iV_1(\underline{T})}, \dots, e^{iV_m(\underline{T})}$. In addition, the two new curves $\widehat{\eta}_1$ and $\widehat{\eta}_2$ commute in the same way as in η_1 and η_2 do. A similar result has been used in [19, Lemma 4.1].

We now give a sketch of the proof. Recall that the one-directional DMP follows from the strong Markov property of Brownian motion and the definition of chordal Loewner equation. The standard way to prove the strong Markov property of Brownian motion at an arbitrary stopping time T is to approximate T by a decreasing sequence of stopping times (T_n) taking countably many values and apply the Markov property of Brownian motion at deterministic times to prove that the strong Markov property holds for each T_n .

To prove Theorem A.1, we could also approximate \underline{T} using a decreasing sequence of stopping times (\underline{T}^n) such that each \underline{T}^n takes countably many values. Using the one-directional DMP and reversibility of chordal SLE_{κ}, it is easy to prove that the theorem holds for a deterministic times $\underline{t} = (t_1, t_2)$, which means that, conditionally on $\overline{\mathcal{G}}_{\underline{t}}$ and the event $\{\underline{t} \in \mathcal{D}_{\cap}\}, \gamma_1|_{[t_1,\phi_2(t_2)]}$ is a time-change of a chordal SLE_{κ} curve from $\gamma_1(t_1)$ to $\gamma_2(t_2)$ in the domain $D_{\underline{t}}$. Since each \underline{T}^n takes countably many values, the statement also holds for \underline{T}^n .

Extending the results from \underline{T}^n to \underline{T} takes significant amount of work. We observe that, for each $\underline{t} = (t_1, t_2) \in \mathcal{D}_{\cap}$, there exists at least one conformal map from \mathbb{H} onto $D_{\underline{t}}$, which sends 0 and ∞ respectively to the prime ends $\gamma_1(t_1)$ and $\gamma_2(t_2)$, but such conformal maps are not unique. We will prove Lemma A.13, which says that we may choose the conformal maps $f_{\underline{t}}^1, \underline{t} \in \mathcal{D}_{\cap}$, such that $f_{\underline{t}}^1(z)$ is jointly continuous in \underline{t} and z. The continuity fact implies that $f_T^1 = \lim_{n \to \infty} f_{T^n}^1$ is $\overline{\mathcal{G}}_{\underline{T}}$ -measurable. Fix $\underline{t} \in \mathcal{D}_{\cap}$. By DMP and reversibility of chordal SLE_{κ}, there is a random Loewner curve $\eta_{\underline{t}}$ in \mathbb{H} , which is a time-change of a standard SLE_{κ} such that $f_{\underline{t}}^1 \circ \eta_{\underline{t}} = \eta_1|_{[t_1,\phi_2(t_2)]}$. Suppose $\eta_{\underline{t}}$ is driven by $\widehat{w}_{\underline{t}}$ with speed $u_{\underline{t}}$. Let $\widehat{v}_{\underline{t}} = \widehat{w}_{\underline{t}} \circ u_{\underline{t}}^{-1}$. Then $(\widehat{v}_{\underline{t}})$ has the law of $(\sqrt{\kappa}B_t)$. Since \underline{T}^n takes countably many values, $(\widehat{v}_{\underline{T}^n}(t))$ also has the law of $(\sqrt{\kappa}B_t)$. We will prove Lemma A.12, which will imply that $\widehat{w}_{\underline{t}}(t)$ and $u_{\underline{t}}(t)$ both have versions that are jointly continuous in \underline{t} and t, and so the same is true for $\widehat{v}_{\underline{t}}(t)$, which then implies that $\widehat{v}_{\underline{T}}(t) = \lim_{n \to \infty} \widehat{v}_{\underline{T}^n}(t)$ has the law of $(\sqrt{\kappa}B_t)$. This fact shows that $\eta_{\underline{T}}(t)$ is a time-change of a standard SLE_{κ}. Since $f_{\underline{t}}^1 \circ \eta_{\underline{t}} = \eta_1|_{[t_1,\phi_2(t_2)]}$ and $f_{\underline{T}}^1$ is $\overline{\mathcal{G}}_{\underline{T}}$ -measurable, we arrive at the conclusion.

The proofs of Lemmas A.12 and A.13 use the Carathéodory topology for the convergence of domains developed in [25, Sections 5.1-5.2], whose definition and basic properties will be recalled in Definition A.6 and Propositions A.7 and A.9. We will further develop this topology to include one prime end and derive some of its properties in Lemma A.10.

The above description of the proof gives the basic idea but is very sketchy and not precise. Here are some examples. We will not go directly from \underline{T}^n to \underline{T} . Instead, we will first approximate $\underline{T} = (T_1, T_2)$ using (T_1^n, T_2) , where T_1^n takes countably many values (but (T_1^n, T_2) does not), and then approximate each (T_1^n, T_2) using stopping times taking countably many values. Lemma A.12 will appear before Lemma A.13 and be used in the proof of the latter lemma. The precise statement about the law of $(\widehat{v}_{\underline{T}^n}(t))$ should be: conditionally on $\overline{\mathcal{G}}_{\underline{T}^n}$ and the event $\{\underline{T}^n \in \mathcal{D}_{\cap}\}, (\widehat{v}_{\underline{T}^n}(t))$ has the law of $(\sqrt{\kappa}B_t)$.

Two groups of intertwined factors contribute to the significant length of the proof of Theorem A.1. The factors on the Probability side is the treatment of two-time-variable filtration, stopping times and stochastic processes; the factors on the Complex Analysis side is the argument involving Loewner's equation, prime ends, Carathéodory topology and extremal length. In addition, we have to deal with the annoying random time-changes and the fact that many random elements are not defined on the whole probability space.

Now we start the proof of Theorem A.1. By conformal invariance, it suffices to work on standard chordal $\operatorname{SLE}_{\kappa}$ curves. Let J(z) = -1/z. By reversibility of chordal $\operatorname{SLE}_{\kappa}$, there are two standard chordal $\operatorname{SLE}_{\kappa}$ curves η_1 and η_2 , and a decreasing auto-homeomorphism ϕ_2 of $[0,\infty]$ such that $\eta_1 \circ \phi_2 = J \circ \eta_2$. For j = 1, 2, let \hat{w}_j be the chordal Loewner driving function for η_j , K_t^j , $0 \leq t < \infty$, be the chordal Loewner hulls driven by \hat{w}_j , $\eta_j^J = J \circ \eta_j$, and $K_t^{j,J} = J(K_t^j)$. We also view $\eta_j(t)$ and $\eta_j^J(t)$ as prime ends of $\mathbb{H} \setminus K_t^j$ and $\mathbb{H} \setminus K_t^{j,J}$, respectively. For j = 1, 2, let \mathcal{F}^j be the filtration generated by \hat{w}_j . Let \mathcal{F} be the separable \mathbb{R}^2_+ -indexed filtration generated by \mathcal{F}^1 and \mathcal{F}^2 , and $\overline{\mathcal{F}}$ be the right-continuous augmentation of \mathcal{F} . Let $\mathcal{D}_j = \{(t_1, t_2) \in \mathbb{R}^2_+ : \eta_j^J(t_j) \notin \eta_{3-j}[0, t_{3-j}]\}, j = 1, 2$, and $\mathcal{D}_{\cap} = \mathcal{D}_1 \cap \mathcal{D}_2$. For $\underline{t} \in \mathcal{D}_{\cap}$, there is a unique connected component of $\mathbb{H} \setminus (\eta_1[0, t_1] \cup \eta_2^J[0, t_2])$, which shares the prime end $\eta_1(t_1)$ with $\mathbb{H} \setminus K_{t_1}^1$ and the prime end $\eta_2^J(t_2)$ with $\mathbb{H} \setminus K_{t_2}^{2,J}$. Let this component be denoted by $H_{\underline{t}}^1$. Theorem A.1 is equivalent to the following theorem.

Lemma A.5. Let $\underline{T} = (T_1, T_2)$ be an $\overline{\mathcal{F}}$ -stopping time. Then the event $\{\underline{T} \in \mathcal{D}_{\cap}\} \in \overline{\mathcal{F}}_{\underline{T}}$, and on this event there are

• an $\mathcal{F}_{\underline{T}}$ -measurable random conformal map $f_{\underline{T}}$ from \mathbb{H} onto H^1_T , which sends 0 and ∞ to

the prime ends $\eta_1(T_1)$ and $\eta_2^J(T_2)$, respectively,

- a standard chordal SLE_{κ} curve $\zeta_1^{\underline{T}}$ conditionally independent of $\overline{\mathcal{F}}_{\underline{T}}$, and
- an increasing homeomorphism u_1^T from $[0,\infty]$ onto $[T_1,\phi_2(T_2)]$,

such that $\eta_1 \circ u_1^{\underline{T}} = f_{\underline{T}} \circ \zeta_1^{\underline{T}}$ a.s. on $\{\underline{T} \in \mathcal{D}_{\cap}\}$.

The rest of the appendix is devoted to the proof of Lemma A.5. We first review some revised Carathéodory topology introduced in [25], which is similar to but different from the Carathéodory kernel convergence in [13, Section 1.4].

Definition A.6. Let $(D_n)_{n \in \mathbb{N}}$ and D be domains in \mathbb{C} . We say that (D_n) converges to D in the Carathéodory topology, and write $D_n \xrightarrow{\text{Cara}} D$, if

- (i) for every compact set $K \subset D$, there exists $n_0 \in \mathbb{N}$ such that $K \subset D_n$ if $n \ge n_0$;
- (ii) for every $z_0 \in \partial D$ there exists $z_n \in \partial D_n$ for each n such that $z_n \to z_0$.

Let (D_n) and D be as in the definition. Suppose $f_n : D_n \to \mathbb{C}$ and $f : D \to \mathbb{C}$. By $f_n \xrightarrow{\text{l.u.}} f$ in D we mean that f_n converges to f locally uniformly in D, i.e., uniformly on every compact subset of D. The following is [25, Lemma 5.1].

Proposition A.7. Let D_n , $n \in \mathbb{N}$, and D be domains in \mathbb{C} such that $D_n \xrightarrow{\text{Cara}} D$. Let $f_n : D_n \xrightarrow{\text{Conf}} E_n$ for some domain $E_n \subset \mathbb{C}$, $n \in \mathbb{N}$. Suppose $f_n \xrightarrow{\text{Lu.}} f$ in D for some nonconstant $f: D \to \mathbb{C}$. Then f is a conformal map, $E_n \xrightarrow{\text{Cara}} E := f(D)$, and $f_n^{-1} \xrightarrow{\text{Lu.}} f^{-1}$ in E.

The following proposition is well known (cf. [19, Proposition 2.3]).

Proposition A.8. If K is an \mathbb{H} -hull with $K \subset \{|z - x_0| \leq r\}$ for some $x_0 \in \mathbb{R}$ and r > 0, then $|g_K(z) - z| \leq 3r$ for any $z \in \mathbb{C} \setminus K^{\text{doub}}$.

For a nonempty \mathbb{H} -hull K, we write $a_K = \min(\overline{K} \cap \mathbb{R})$, $b_K = \max(\overline{K} \cap \mathbb{R})$, $B_K = [a_K, b_K]$, and $K^{\text{doub}} = K \cup \{\overline{z} : z \in K\} \cup B_K$. For $K \subset L \in \mathcal{H}$ with $L \neq \emptyset$, by Schwarz reflection principle, g_K extends to a conformal map from $\mathbb{C} \setminus (K^{\text{doub}} \cup B_L)$ conformally onto $\mathbb{C} \setminus S_K^L$ for some compact interval S_K^L . We write S_K for S_K^K . Let $c_K^L < d_K^L$ and $c_K < d_K$ be such that $S_K^L = [c_K^L, d_K^L]$ and $S_K = [c_K, d_K]$. Then g_K maps $(-\infty, a_K)$ and (b_K, ∞) respectively onto $(-\infty, c_K)$ and (d_K, ∞) , and satisfies that

$$g_K(x) < x, \quad \forall x \in (-\infty, a_K); \quad g_K(x) > x, \quad \forall x \in (b_K, \infty).$$
 (A.1)

Let $f_K = g_K^{-1}$. Then $f_K : \mathbb{H} \xrightarrow{\text{Conf}} \mathbb{H} \setminus K$ and $f_K : \mathbb{C} \setminus S_K \xrightarrow{\text{Conf}} \mathbb{C} \setminus K^{\text{doub}}$. Let \mathcal{H} denote the space of \mathbb{H} -hulls. There is a metric $d_{\mathcal{H}}$ on \mathcal{H} such that $K_n \to K$ w.r.t. $d_{\mathcal{H}}$

Let \mathcal{H} denote the space of \mathbb{H} -hulls. There is a metric $d_{\mathcal{H}}$ on \mathcal{H} such that $K_n \to K$ w.r.t. $d_{\mathcal{H}}$ iff $f_{K_n} \xrightarrow{\text{l.u.}} f_K$ in \mathbb{H} . By Proposition A.7, this implies that $\mathbb{H} \setminus K_n \xrightarrow{\text{Cara}} \mathbb{H} \setminus K$. But the converse is not true. For $K \in \mathcal{H}$, let $\mathcal{H}(K) = \{L \in \mathcal{H} : L \subset K\}$. The following is [25, Lemma 5.4 (i)]. **Proposition A.9.** For any $K \in \mathcal{H}$ with $K \neq \emptyset$, $\mathcal{H}(K)$ is compact w.r.t. $d_{\mathcal{H}}$. This means that every sequence (K_n) in $\mathcal{H}(K)$ contains a convergent subsequence w.r.t. $d_{\mathcal{H}}$ with limit in $\mathcal{H}(K)$. Moreover, if $K_n \to K_0$ in $\mathcal{H}(K)$, then $f_{K_n} \xrightarrow{1.u.} f_{K_0}$ in $\mathbb{C} \setminus S_K$, $\mathbb{C} \setminus (K_n^{\text{doub}} \cup B_K) \xrightarrow{\text{Cara}} \mathbb{C} \setminus (K_0^{\text{doub}} \cup B_K)$, and $g_{K_n} \xrightarrow{1.u.} g_{K_0}$ in $\mathbb{C} \setminus (K_0^{\text{doub}} \cup B_K)$.

Let \mathcal{P} denote the space of pairs (K, p), where K is an \mathbb{H} -hull, and p is a prime end of $\mathbb{H} \setminus K$ other than ∞ . Note that $g_K(p) \in \mathbb{R}$ for $(K, p) \in \mathcal{P}$. Equip \mathcal{P} with the metric

$$d_{\mathcal{P}}((K_1, p_1), (K_2, p_2)) = d_{\mathcal{H}}(K_1, K_2) + |g_{K_1}(p_1) - g_{K_2}(p_2)|.$$

For $(K,p) \in \mathcal{P}$, let $g_{K,p} = g_K - g_K(p)$ and $f_{K,p} = g_{K,p}^{-1}$. If $(K_n, p_n) \to (K, p)$ w.r.t. $d_{\mathcal{P}}$, then $g_{K_n,p_n} \xrightarrow{\text{l.u.}} g_{K,p}$ in $\mathbb{H} \setminus K$ and $f_{K_n,p_n} \xrightarrow{\text{l.u.}} f_{K,p}$ in \mathbb{H} . For a nonempty \mathbb{H} -hull L, let $\mathcal{P}(L)$ denote the set of $(K,p) \in \mathcal{P}$ such that $K \in \mathcal{H}(L)$, and $p \notin (-\infty, a_L) \cup (b_L, \infty)$. Then $g_K(p) \in S_K^L$. Let $S_{K,p}^L = S_K^L - g_K(p)$. Then $0 \in S_{K,p}^L$ and $g_{K,p} : \mathbb{C} \setminus (K^{\text{doub}} \cup B_L) \xrightarrow{\text{Conf}} \mathbb{C} \setminus S_{K,p}^L$.

Subset of $(H,p) \in \mathcal{F}$ back that $H \in \mathcal{F}(L)$, and $p \notin (- \operatorname{oc}, a_L) \in (\mathfrak{o}_L, \mathfrak{o}_L)$. Then $g_K(p) \in \mathcal{S}_K^*$. Let $S_{K,p}^L = S_K^L - g_K(p)$. Then $0 \in S_{K,p}^L$ and $g_{K,p} : \mathbb{C} \setminus (K^{\operatorname{doub}} \cup B_L) \xrightarrow{\operatorname{Conf}} \mathbb{C} \setminus S_{K,p}^L$. The following example is important. Suppose K_t , $0 \leq t < T$, are chordal Loewner hulls driven by \hat{w} . For $0 \leq t < T$, let p_t be the prime end $g_{K_t}^{-1}(\hat{w}(t))$ of $\mathbb{H} \setminus K_t$. Then (K_t, p_t) , $0 \leq t < T$, is a continuous curve in \mathcal{P} . The corresponding maps g_{K_t,p_t} and f_{K_t,p_t} are called centered Loewner maps in the literature.

Let Ξ denote the set of pairs $(L^1, L^2) \in \mathcal{H}^2$ such that L^j contains a neighborhood of 0 in \mathbb{H} , j = 1, 2, and $\operatorname{dist}((L^1)^{\operatorname{doub}}, J((L^2)^{\operatorname{doub}})) > 0$. For each $\underline{L} = (L^1, L^2) \in \Xi$, let $\mathcal{PP}(\underline{L}) = \mathcal{P}(L^1) \times \mathcal{P}(L^2)$. Let $\mathcal{PP} = \bigcup_{\underline{L} \in \Xi} \mathcal{PP}(\underline{L})$. Suppose $(K_1, p_1; K_2, p_2) \in \mathcal{PP}$. Define $H^1_{K_1; K_2} = \mathbb{H} \setminus (K_1 \cup J(K_2))$. Then $H^1_{K_1; K_2}$ is simply connected and shares the prime ends p_1 and $J(p_2)$ respectively with $\mathbb{H} \setminus K_1$ and $\mathbb{H} \setminus J(K_2)$. Let $g^J_{K_2, p_2} = J \circ g_{K_2, p_2} \circ J$ and $f^J_{K_2, p_2} = (g^J_{K_2, p_2})^{-1}$. Let $\widetilde{K}_1, \widetilde{p}_1$ be respectively the $g^J_{K_2, p_2}$ -images of K_1, p_1 . Since $g^J_{K_2, p_2} : H^1_{K_1; K_2} \xrightarrow{\operatorname{Conf}} \mathbb{H} \setminus \widetilde{K}_1$, we see that $\mathbb{H} \setminus \widetilde{K}_1$ is also simply connected. Since $J(p_2)$ is a prime end of $H^1_{K_1, K_2}$ bounded away from K_1 , and is sent to ∞ by $g^J_{K_2, p_2}$, we see that $\mathbb{H} \setminus \widetilde{K}_1$ shares the prime end ∞ with \mathbb{H} . Thus, $\widetilde{K}_1 \in \mathcal{H}$. Since p_1 and $J(p_2)$ are distinct prime ends of \mathbb{H}_{K_1, K_2} , we have $\widetilde{p}_1 = g^J_{K_2, p_2}(p_1) \neq g^J_{K_2, p_2}(J(p_2)) = \infty$ in terms of prime ends. So $(\widetilde{K}_1, \widetilde{p}_1) \in \mathcal{P}$.

Lemma A.10. Let $(L^1, L^2) \in \Xi$. For j = 1, 2, let (K_n^j, p_n^j) , $n \in \mathbb{N}$, be a convergent sequence in $\mathcal{P}(L^j)$ with limit $(K_{\infty}^j, p_{\infty}^j) \in \mathcal{P}(L_j)$. For $n \in \overline{\mathbb{N}} := \mathbb{N} \cup \{\infty\}$, let $\widetilde{K}_n^1, \widetilde{p}_n^1$ be respectively the $g_{K_n^2, p_n^2}^J$ -images of K_n^1, p_n^1 . Then $(\widetilde{K}_n^1, \widetilde{p}_n^1) \to (\widetilde{K}_{\infty}^1, \widetilde{p}_{\infty}^1)$ w.r.t. $d_{\mathcal{P}}, g_{\widetilde{K}_n^1} \xrightarrow{\text{lus}} g_{\widetilde{K}_{\infty}^1}$ in $\mathbb{C} \setminus ((\widetilde{K}_{\infty}^1)^{\text{doub}} \cup g_{K_{\infty}^2, p_{\infty}^2}^J(B_{L^1}))$, and $g_{\widetilde{K}_n^1}^{-1} \xrightarrow{\text{lus}} g_{\widetilde{K}_n^1}^{-1}$ in $\mathbb{C} \setminus (S_{\widetilde{K}_n^1} \cup g_{\widetilde{K}_\infty^1} \circ g_{\widetilde{K}_\infty^2, p_{\infty}^2}^J(B_{L^1}))$.

Proof. From $K_n^j \to K_\infty^j$, j = 1, 2, we know that $H_{K_n^1;K_n^2}^1 \xrightarrow{\text{Cara}} H_{K_\infty^1;K_\infty^2}^1$. Since $g_{K_n^2,p_n^2}^J \xrightarrow{\text{l.u.}} g_{K_\infty^2,p_\infty^2}^J$ in $H_{K_\infty^1,K_\infty^2}^1$, by Proposition A.7, $\mathbb{H} \setminus \widetilde{K}_n^1 \xrightarrow{\text{Cara}} \mathbb{H} \setminus \widetilde{K}_\infty^1$. Let $\Lambda = \Lambda(L^1,L^2) \in (0,\infty)$ be the extremal distance (cf. [1]) between $(L^2)^{\text{doub}}$ and $J((L^1)^{\text{doub}})$. By comparison principle, for any $n \in \mathbb{N}$, the extremal distance between $(K_n^2)^{\text{doub}} \cup B_{L^2}$ and $J((K_n^1)^{\text{doub}} \cup B_{L^1})$ is at least Λ . By conformal invariance, the extremal distance between $S_{K_n^2,p_n^2}^{L^2}$ and $g_{K_n^2,p_n^2} \circ J((K_n^1)^{\text{doub}})$ is at least A. Since $S_{K_n^2,p_n^2}^{L^2}$ contains 0 and has length $\geq b_{L^2} - a_{L^2} > 0$ by (A.1), there is $r = r(L^1, L^2) > 0$ such that dist $(0, g_{K_n^2,p_n^2} \circ J(K_n^1)) \geq r$, which implies that $\widetilde{K}_n^1 \subset \{|z| \leq 1/r\}$. By Proposition A.9, $\{\widetilde{K}_n^1\}$ is pre-compact. Since $\mathbb{H} \setminus \widetilde{K}_n^1 \stackrel{\text{Cara}}{\longrightarrow} \mathbb{H} \setminus \widetilde{K}_\infty^1$, we get $\widetilde{K}_n^1 \to \widetilde{K}_\infty^1$ w.r.t. $d_{\mathcal{H}}$. For each $n \in \overline{\mathbb{N}}, g_n := g_{\widetilde{K}_n^1} \circ g_{K_n^2,p_n^2}^J \circ f_{K_n^1}$ is a conformal map from $\mathbb{C} \setminus (S_{K_n^1} \cup g_{K_n^1} \circ J((K_n^2)^{\text{doub}}))$

For each $n \in \overline{\mathbb{N}}$, $g_n := g_{\widetilde{K}_n^1} \circ g_{K_n^2, p_n^2}^{-2} \circ f_{K_n^1}$ is a conformal map from $\mathbb{C} \setminus (S_{K_n^1} \cup g_{K_n^1} \circ J((K_n^2)^{\text{doub}}))$ onto $\mathbb{C} \setminus (S_{\widetilde{K}_n^1} \cup g_{\widetilde{K}_n^1} \circ J(S_{K_n^2}))$, and maps a neighborhood of $S_{K_n^1}$ to a neighborhood of $S_{\widetilde{K}_n^1}$. By Schwarz reflection principle, it extends to a conformal map on $\mathbb{C} \setminus g_{K_n^1} \circ J((K_n^2)^{\text{doub}})$, and maps $S_{K_n^1}$ onto $S_{\widetilde{K}_n^1}$. From $K_n^1 \to K_\infty^1$, $\widetilde{K}_n^1 \to \widetilde{K}_\infty^1$ and $(K_n^2, p_n^2) \to (K_\infty^2, p_\infty^2)$, we know that $g_n \xrightarrow{\text{l.u.}} g_0$ in $\mathbb{C} \setminus (S_{K_n^1}^{L_1} \cup g_{K_\infty^1} \circ J((K_\infty^2)^{\text{doub}}))$. By maximum principle, we get $g_n \xrightarrow{\text{l.u.}} g_0$ in $\mathbb{C} \setminus g_{K_n^1} \circ J((K_\infty^2)^{\text{doub}})$. Since $g_{\widetilde{K}_n^1}(\widetilde{p}_n^1) = g_n \circ g_{K_n^1}(p_n^1)$ for $n \in \overline{\mathbb{N}}$, and $g_{K_n^1}(p_n^1) \to g_{K_\infty^1}(p_\infty^1) \in$ $\mathbb{C} \setminus g_{K_\infty^1} \circ J((K_\infty^2)^{\text{doub}})$, we get $g_{\widetilde{K}_n^1}(\widetilde{p}_n^1) \to g_{\widetilde{K}_\infty^1}(\widetilde{p}_\infty^1)$. Thus, $(\widetilde{K}_n^1, \widetilde{p}_n^1) \to (\widetilde{K}_\infty^1, \widetilde{p}_\infty^1)$.

Since $K_n^1 \to K_\infty^1$ in $\mathcal{H}(L^1)$, we get $\mathbb{C} \setminus ((K_n^1)^{\text{doub}} \cup B_{L^1}) \xrightarrow{\text{Cara}} \mathbb{C} \setminus ((K_\infty^1)^{\text{doub}} \cup B_{L^1})$. By Proposition A.7, we get $\mathbb{C} \setminus ((\widetilde{K}_n^1)^{\text{doub}} \cup g_{K_n^2, p_n^2}^J(B_{L^2})) \xrightarrow{\text{Cara}} \mathbb{C} \setminus ((\widetilde{K}_\infty^1)^{\text{doub}} \cup g_{K_\infty^2, p_\infty^2}^J(B_{L^1}))$. From the first paragraph of the proof, there is $R \in (0, \infty)$ such that $\widetilde{K}_n^1 \subset \{|z| < R\}$ for all $n \in \overline{\mathbb{N}}$. By Proposition A.8, $|g_{\widetilde{K}_n^1}(z) - z| \leq 3R$ for all $n \in \overline{\mathbb{N}}$ and $z \in \mathbb{C} \setminus (\widetilde{K}_n^1)^{\text{doub}}$. Thus, any subsequence of $\{g_{\widetilde{K}_n^1}\}$ contains a further subsequence, which converges locally uniformly in $\mathbb{C} \setminus ((\widetilde{K}_\infty^1)^{\text{doub}} \cup g_{K_\infty^2, p_\infty^2}^J(B_{L^1}))$. The limit must be $g_{\widetilde{K}_\infty^1}$ since we already know that $g_{\widetilde{K}_n^1} \xrightarrow{\text{l.u.}} g_{\widetilde{K}_n^1}$ in $\mathbb{H} \setminus \widetilde{K}_\infty^1$. So we get $g_{\widetilde{K}_n^1} \xrightarrow{\text{l.u.}} g_{\widetilde{K}_\infty^1}$ in $\mathbb{C} \setminus ((\widetilde{K}_\infty^1)^{\text{doub}} \cup g_{K_\infty^2, p_\infty^2}^J(B_{L^1}))$. By Proposition A.7, $g_{\widetilde{K}_n^1}^{-1} \xrightarrow{\text{l.u.}} g_{\widetilde{K}_\infty^1}^{-1}$ in $g_{\widetilde{K}_\infty^1}(\mathbb{C} \setminus ((\widetilde{K}_\infty^1)^{\text{doub}} \cup g_{K_\infty^2, p_\infty^2}^J(B_{L^1}))) = \mathbb{C} \setminus (S_{\widetilde{K}_\infty^1} \cup g_{\widetilde{K}_\infty^1} \circ g_{K_\infty^2, p_\infty^2}^J(B_{L^1})))$.

If $\eta(t)$, $0 \leq t < T$, is a continuous curve in $\overline{\mathbb{H}}$, and there are a continuous function \widehat{w} on [0,T) and a continuous and strictly increasing function u on [0,T) with u(0) = 0, such that $\eta \circ u^{-1}(t)$, $0 \leq t < u(T)$, is the chordal Loewner curve driven by $\widehat{w} \circ u^{-1}$, then we say that η is a chordal Loewner curve with speed du driven by \widehat{w} . For each $0 \leq t < T$, we also understand $\eta(t)$ as the prime end $g_{\eta[0,t]}^{-1}(\widehat{w}(t))$ of $\mathbb{H} \setminus \operatorname{Hull}(\eta[0,t])$. The following proposition is a well-known result in the Loewner theory, and so we omit its proof.

Proposition A.11. Suppose $\eta(t)$, $0 \le t < T$, is a chordal Loewner curve. Let I be an open real interval, which contains $\eta \cap \mathbb{R}$. Let U be a subdomain of \mathbb{H} , which is a neighborhood of I in \mathbb{H} , and contains $\bigcup_{0 \le t < T} \operatorname{Hull}(\eta[0, t])$. Let ψ be a conformal map from U into \mathbb{H} , which extends continuously to $U \cup I$, and maps I into \mathbb{R} . Then $\psi \circ \eta(t)$, $0 \le t < T$, is a chordal Loewner curve with some speed, and for each $t \in [0, T)$, ψ sends the prime end $\eta(t)$ of $\mathbb{H} \setminus \operatorname{Hull}(\eta[0, t])$ to the prime end $\psi \circ \eta(t)$ of $\mathbb{H} \setminus \operatorname{Hull}(\psi \circ \eta[0, t])$.

Now we come back to the proof of Lemma A.5. For j = 1, 2 and $t \ge 0$, let g_t^j denote the centered Loewner map for η_j at time t, i.e., $g_t^j = g_{K_t^j} - \hat{w}_j(t)$. Let $f_t^j = (g_t^j)^{-1}$ and $f_t^{j,J} = J \circ f_t^j \circ J$. Fix $t_2 \ge 0$. By DMP of SLE, conditionally on $\mathcal{F}_{t_2}^2$, $\eta_2(t_2 + \cdot)$ is a chordal SLE_{κ} in $\mathbb{H} \setminus K_{t_2}^2$ from $\eta_2(t_2)$ to ∞ . Since f_t^2 maps \mathbb{H} conformally onto $\mathbb{H} \setminus K_t^2$, extends continuously to $\overline{\mathbb{H}}^{\#}$, and sends 0 and ∞ respectively to $\eta_2(t)$ and ∞ , there is a standard chordal SLE_{κ} curve $\eta_2^{t_2+}$ independent of $\mathcal{F}_{t_2}^2$ such that a.s. $\eta_2(t_2+\cdot) = f_{t_2}^2 \circ \eta_2^{t_2+}$. The $\eta_2^{t_2+}$ is the chordal Loewner curve driven by $\widehat{w}_2^{t_2+} := \widehat{w}_2(t_2+\cdot) - \widehat{w}_2(t_2)$. Let $K_t^{2,t_2+} = \operatorname{Hull}(\eta_2^{t_2+}[0,t])$ and g_t^{2,t_2+} be the centered Loewner map for $\eta_2^{t_2+}$, i.e., $g_{K_t^{2,t_2+}} - \widehat{w}_2^{t_2+}(t)$. Then $f_{t_2}^2 = f_{t_2+t}^2 \circ g_t^{2,t_2+}$, $t \ge 0$.

Since $\eta_1(t)$, $0 \le t \le \phi_2(t_2)$, is a time-reversal of $\eta_2^J(t)$, $t_2 \le t \le \infty$, by reversibility of SLE_{κ}, conditionally on $\mathcal{F}_{t_2}^2$, $\eta_1(t)$, $0 \le t \le \phi_2(t_2)$, is a time-change of a chordal SLE_{κ} in $\mathbb{H} \setminus K_{t_2}^{2,J}$ from 0 to $\eta_2^J(t_2)$. Since $f_{t_2}^{2,J}$: $(\mathbb{H}; 0, \infty) \xrightarrow{\text{Conf}} (\mathbb{H} \setminus K_{t_2}^{2,J}; 0, \eta_2^J(t_2))$, there is a standard chordal SLE_{κ} curve $\zeta_1^{t_2}$ independent of $\mathcal{F}_{t_2}^2$, and an increasing homeomorphism $u_1^{t_2}$ from $[0, \phi_2(t_2)]$ onto $[0, \infty]$, such that a.s. $\eta_1 = f_{t_2}^{2,J} \circ \zeta_1^{t_2} \circ u_1^{t_2}$ on $[0, \phi_2(t_2)]$. Let $\hat{v}_1^{t_2}$ be the driving function for $\zeta_1^{t_2}$. Let $\eta_1^{t_2} = \zeta_1^{t_2} \circ u_1^{t_2}$ and $\hat{w}_1^{t_2} = \hat{v}_1^{t_2} \circ u_1^{t_2}$. Then $\eta_1^{t_2}$ is the chordal Loewner curve with speed $du_1^{t_2}$ driven by $\hat{w}_1^{t_2}$, and a.s. $\eta_1 = f_{t_2}^{2,J} \circ \eta_1^{t_2}$ on $[0, \phi_2(t_2)]$.

Lemma A.12. The functions $\eta_1^{t_2}(t_1), u_1^{t_2}(t_1), \widehat{w}_1^{t_2}(t_1)$ all have continuous versions on $\mathcal{D}_2 = \{(t_1, t_2) : \eta_2^1(t_2) \notin \eta_1[0, t_1]\}$. More specifically, there are continuous processes $\eta_1^* : \mathcal{D}_2 \to \overline{\mathbb{H}}, u_1^* : \mathcal{D}_2 \to \mathbb{R}_+$ and $\widehat{w}_1^* : \mathcal{D}_2 \to \mathbb{R}$ such that for any fixed $t_2 \in \mathbb{R}_+, \eta_1^*(\cdot, t_2)$ is the chordal Loewner curve with speed $du_1^*(\cdot, t_2)$ driven by $\widehat{w}_1^*(\cdot, t_2)$; and a.s. $\eta_1^*(\cdot, t_2), u_1^*(\cdot, t_2)$ and $\widehat{w}_1^*(\cdot, t_2)$ are respectively equal to $\eta_1^{t_2}, u_1^{t_2}$ and $\widehat{w}_1^{t_2}$, which further implies that a.s. $\eta_1|_{[0,\phi_2(t_2))} = f_{t_2}^{2,J} \circ \eta_1^*(\cdot, t_2)$. Here for two functions to be equal, they must have the same domain.

Proof. Let \mathcal{A} denote the set of triples $(t_1^0; t_2^0, t_1^1)$, where $t_1^0, t_2^0, t_1^1 \in \mathbb{Q}_+$ and $t_2^0 < t_2^1$. For each $\underline{v} = (t_1^0; t_2^0, t_2^1) \in \mathcal{A}$, define $R_{\underline{v}} = [0, t_1^0] \times [t_2^0, t_2^1]$ and $E_{\underline{v}} = \{R_{\underline{v}} \subset \mathcal{D}_2\}$. We make the following claim: for any $\underline{v} = (t_1^0; t_2^0, t_2^1) \in \mathcal{A}$, on the event $E_{\underline{v}}$, there are continuous processes $\eta_1^* : R_{\underline{v}} \to \overline{\mathbb{H}}$, $u_1^* : R_{\underline{v}} \to \mathbb{R}_+$ and $\widehat{w}_1^* : R_{\underline{v}} \to \mathbb{R}$ such that for any $t_2 \in [t_2^0, t_2^1]$, $\eta_1^*(\cdot, t_2)$ is the chordal Loewner curve with speed $du_1^*(\cdot, t_2)$ driven by $\widehat{w}_1^*(\cdot, t_2)$; and a.s. $\eta_1^*(\cdot, t_2), u_1^*(\cdot, t_2)$ and $\widehat{w}_1^*(\cdot, t_2)$ are respectively equal to the restrictions of $\eta_1^{t_2}, u_1^{t_2}$ and $\widehat{w}_1^{t_2}$ to $[0, t_1^0]$.

Loewner curve with speed $du_1^{(\cdot)} \cdot t_2^{\underline{v}}$ driven by $\widehat{w}_1^{(\cdot)}(\cdot, t_2)$; and a.s. $\eta_1^{(\cdot)}(\cdot, t_2)$, $u_1^{(\cdot)}(\cdot, t_2)$ and $\widehat{w}_1^{(\cdot)}(\cdot, t_2)$ are respectively equal to the restrictions of $\eta_1^{t_2}$, $u_1^{t_2}$ and $\widehat{w}_1^{t_2}$ to $[0, t_1^0]$. Fix $\underline{v} = (t_1^0; t_2^0, t_2^1) \in \mathcal{A}$. Let $\Delta t_2 = t_2^1 - t_2^0$. Suppose $E_{\underline{v}}$ happens. Then $\eta_2^J[t_2^0, t_2^1] \cap \eta_1[0, t_1^0] = \emptyset$. Since $\eta_2^J[t_2^0, t_2^1] = f_{t_2^0}^{2,J} \circ J \circ \eta_2^{t_2^0+}[0, \Delta t_2]$ and $\eta^1[0, t_1^0] = f_{t_2^0}^{2,J} \circ \eta_1^{t_2^0}[0, t_1^0]$, we get $(J \circ \eta_2^{t_2^0+}[0, \Delta t_2]) \cap \eta_1^{t_2^0}[0, t_1^0] = \emptyset$. Since both $\eta_1^{t_2}$ and $\eta_2^{t_2^0+}$ start from 0, we can find a (random) pair $\underline{L} = (L^1, L^2) \in \Xi$ such that $K_{t_1^0}^{1,t_2^0} \subset L^1$ and $K_{\Delta t_2}^{2,t_2^0+} \subset L^2$. Then $(K_{t_1}^{1,t_2^0}, \eta_1^{t_2^0}(t_1)), 0 \le t_1 \le t_1^0$, and $(K_{t_2}^{2,t_2^0+}, \eta_2^{t_2^0+}(t_2)), 0 \le t_2 \le \Delta t_2$, are respectively continuous curves in $\mathcal{P}(L^1)$ and $\mathcal{P}(L^2)$. Let $g_t^{2,t_2^0+,J} = J \circ g_t^{2,t_2^0+} \circ J$. Let $t_2 \in [t_2^0, t_2^1]$. We have a.s., for all $t_1 \in [0, t_1^0]$,

$$f_{t_2}^{2,J} \circ \eta_1^{t_2}(t_1) = \eta_1(t_1) = f_{t_2^0}^{2,J} \circ \eta_1^{t_2^0}(t_1) = f_{t_2}^{2,J} \circ g_{t_2-t_2^0}^{2,t_2^0+,J} \circ \eta_1^{t_2^0}(t_1),$$

which implies that a.s. for all $t_1 \in S_1 := [0, t_1^0] \cap (\eta_1^{t_2^0})^{-1}(\mathbb{H}), \ \eta_1^{t_2}(t_1) = g_{t_2-t_2^0}^{2, t_2^0+, J} \circ \eta_1^{t_2^0}(t_1).$ Since S_1 is dense in $[0, t_1^0]$, both $\eta_1^{t_2}$ and $\eta_1^{t_2^0}$ are continuous on $[0, t_1^0]$, and $g_{t_2-t_2^0}^{2, t_2^0+, J}$ is continuous on $\overline{\mathbb{H}} \setminus J(\overline{K_{t_2-t_2^0}^{2, t_2^0+, J}}) \supset \overline{\mathbb{H}} \setminus J(\overline{L^2}) \supset \overline{L^1} \supset \eta_1^{t_2^0}[0, t_1^0]$, we get a.s. $\eta_1^{t_2} = g_{t_2-t_2^0}^{2, t_2^0+, J} \circ \eta_1^{t_2^0}$ on $[0, t_1^0]$.

Define η_1^* and u_1^* by $\eta_1^*(t_1, t_2) = g_{t_2-t_2^0}^{2,t_2^0+,J} \circ \eta_1^{t_2^0}(t_1)$ and $u_1^*(t_1, t_2) = \text{hcap}_2(\eta_1^*([0, t_1], t_2))$ for $(t_1, t_2) \in R_{\underline{v}}$. Then η_1^* and u_1^* are continuous on $R_{\underline{v}}$ by Proposition A.9. By the last paragraph, for any $t_2 \in [t_2^0, t_2^1]$, a.s. $\eta_1^*(\cdot, t_2) = \eta_1^{t_2}$ and $\eta_1 = f_{t_2}^{2,J} \circ \eta_1^*(\cdot, t_2)$ on $[0, t_1^0]$. So η_1^* and u_1^* are continuous versions of $\eta_1^{t_2}(t_1)$ and $u_1^{t_2}(t_1)$ on $R_{\underline{v}}$. By the continuity of η_1, η_1^* , and $f_{t_2}^{2,J}$, we then get a.s. $\eta_1(t_1) = f_{t_2}^{2,J} \circ \eta_1^*(t_1, t_2)$ for any $(t_1, t_2) \in R_{\underline{v}}$. By Proposition A.11, $\eta_1^*(\cdot, t_2)$ is a chordal Loewner curve with speed $du_1^*(\cdot, t_2)$. Define $\hat{w}_1^* : R_{\underline{v}} \to \mathbb{R}$ such that for every $t_2 \in [t_2^0, t_2^1]$, $\hat{w}_1^*(\cdot, t_2)$ is the driving function for $\eta_1^*(\cdot, t_2)$. For any $t_2 \in [t_2^0, t_2^1]$, since a.s. $\eta_1^*(\cdot, t_2) = \eta_1^{t_2}$ on $[0, t_1^0]$, and $\hat{w}_1^{t_2}$ is the driving function for $\eta_1^{t_2}$, we find that a.s. $\hat{w}_1^*(\cdot, t_2) = \hat{w}_1^{t_2}$ on $[0, t_1^0]$. To prove the claim, it remains to prove that \hat{w}_1^* is a.s. continuous on $R_{\underline{v}}$.

 $\begin{bmatrix} [0, t_1^r], \text{ and } w_1^{-r} \text{ is the driving function for } y_1^{-r}, \dots \\ \\ \text{ From } \eta_1 = f_{t_2}^{2,J}(\eta_1^*(\cdot, t_2)) = f_{t_2}^{2,J}(\eta_1^*(\cdot, t_2^0)), \text{ we get } \eta_1^*(\cdot, t_2) = g_{t_2-t_2}^{2,t_2^0+,J} \circ \eta_1^*(\cdot, t_2^0), t_2 \geq t_2^0. \text{ For } \\ \\ \underline{t} = (t_1, t_2) \in R_{\underline{v}}, \text{ let } K_{t_1}^{1,t_2} = \text{Hull}(\eta_1^*([0, t_1], t_2)). \text{ Then we have } K_{t_1}^{1,t_2} = g_{t_2-t_2^0}^{2,t_2^0+,J}(K_{t_1}^{1,t_2}) \text{ and } \\ \\ \widehat{w}_1^*(\underline{t}) = g_{K_{t_1}^{1,t_2}}(\eta_1^*(\underline{t})). \text{ Define } G_{\underline{t}} = g_{K_{t_1}^{1,t_2}} \circ g_{t_2-t_2^0}^{2,t_2^0+,J} \circ g_{K_{t_1}^{-1,t_2}}^{-1}, \text{ which maps } \mathbb{H} \setminus (g_{K_{t_1}^{1,t_2}} \circ J(K_{t_2-t_2^0}^{2,t_2^0+,J})) \\ \\ \text{ conformally onto } \mathbb{H}, \text{ and extends to a conformal map from } \mathbb{C} \setminus (g_{K_{t_1}^{1,t_2^0}} \circ J((K_{t_2-t_2^0}^{2,t_2^0+,J}))), \text{ which } \\ \\ \text{ contains } \widehat{w}_1^*(t_1^0, t_2), \text{ into } \mathbb{C}. \text{ From } \widehat{w}_1^*(\underline{t}) = g_{K_{t_1}^{1,t_2}}(\eta_1^*(\underline{t})), \ \widehat{w}_1^*(t_1, t_2^0) = g_{K_{t_1}^{1,t_2}}(\eta_1^*(t_1, t_2^0)), \text{ and } \\ \\ \\ \eta_1^*(\cdot, t_2) = g_{t_2-t_2^0}^{2,t_2^0+,J} \circ \eta_1^*(\cdot, t_2^0), \text{ we get } w_1^*(\underline{t}) = G_{\underline{t}}(w_1^*(t_1, t_2^0)). \text{ Since } w_1^*(\cdot, t_2^0) = w_1^{t_2^0}, \text{ which is } \\ \\ \text{ continuous in } t_1, \text{ to show that } w_1^* \text{ is continuous in } \underline{t}, \text{ it suffices to show that } G_{\underline{t}}(x) \text{ is jointly } \\ \\ \text{ continuous in } \underline{t} \text{ and } x \text{ on a domain that contains } \{(\underline{t}, x) : \underline{t} \in R_{\underline{v}}, x \in S_{K_{t_1}^{1,t_2}}\}. \end{cases}$

Since $G_{\underline{t}}$ is a conformal map for each $\underline{t} \in \underline{v}$, to prove the joint continuity, by maximum principle, it suffices to show that, for any fixed $\underline{t}^* = (t_1^*, t_2^*) \in R_{\underline{v}}$, there is a Jordan curve ξ , whose interior domain contains $S_{K_{t_1}^{1,t_2^0}}$, and is contained in $\mathbb{C} \setminus (g_{K_{t_1}^{1,t_2^0}} \circ J((K_{t_2-t_2^0}^{2,t_2^0+}))))$, such that as $R_{\underline{v}} \ni \underline{t}^n \to \underline{t}^*$, $G_{\underline{t}^n} \to G_{\underline{t}^*}$ uniformly on ξ . To choose such ξ , we may first find a Jordan curve ξ' , which separates $(L^1)^{\text{doub}}$ from $J((L^2)^{\text{doub}})$, and then let $\xi = g_{K_{t_1}^{1,t_2^0}}(\xi')$. Then we use

Lemma A.10 to conclude that $G_{\underline{t}^n} \to G_{\underline{t}^*}$ uniformly on ξ if $R_{\underline{v}} \ni \underline{t}^n \to \underline{t}^*$.

Now we have proved the claim. Since \mathcal{D}_2 is the union of $R_{\underline{v}}$ over those $\underline{v} \in \mathcal{A}$ such that $R_{\underline{v}} \subset \mathcal{D}_2$, from the claim we conclude the existence of continuous maps η_1^* , u_1^* and \widehat{w}_1^* defined on \mathcal{D}_2 with all properties described in the statement except that now we can only say that a.s. $\eta_1^*(\cdot, t_2)$, $u_1^*(\cdot, t_2)$ and $\widehat{w}_1^*(\cdot, t_2)$ are respectively equal to the restrictions of $\eta_1^{t_2}$, $u_1^{t_2}$ and $\widehat{w}_1^{t_2}$ to $[0, T_2(t_2))$, where $T_2(t_2) := \inf\{t_1 : \eta_1(t_1) = \eta_2^J(t_2)\}$. From $\eta_1|_{[0,\phi_2(t_2))} = f_{t_2}^{2,J} \circ \eta_1^{t_2}$ we know that a.s. $f_{t_2}^{2,J} \circ \eta_1^*(\cdot, t_2)$ equals $\eta_1|_{[0,T_2(t_2))}$. We know that $\eta_1^{t_2}$, $u_1^{t_2}$ and $\widehat{w}_1^{t_2}$ are defined on $[0, \phi_2(t_2))$. To conclude the proof, it suffices to show that a.s. $T_2(t_2) = \phi_2(t_2)$. Since $\eta_1|_{[0,\phi_2(t_2)]}$ is a time-change of a chordal SLE_ κ in $\mathbb{H} \setminus J(K_{t_2}^{2,J})$ from 0 to the prime end

Since $\eta_1|_{[0,\phi_2(t_2)]}$ is a time-change of a chordal SLE_{κ} in $\mathbb{H} \setminus J(K_{t_2}^{2,J})$ from 0 to the prime end $\eta_2^J(t_2)$, η_1 first visits the prime end $\eta_2^J(t_2)$ at the time $\phi_2(t_2)$. We claim that a.s. $\eta_2^J(t_2)$ is the only prime end of $\mathbb{H} \setminus J(K_{t_2}^{2,J})$, which determines the boundary point $\eta_2^J(t_2)$. When $\kappa \leq 4$ this is true since η_2^J is simple. When $\kappa > 4$, this follows from [22, Theorem 6.1]. Thus, a.s. η_1 first

visits the point $\eta_2^J(t_2)$ at the time $\phi_2(t_2)$. So we get a.s. $T_2(t_2) = \phi_2(t_2)$, as desired.

By discarding a null event we may assume that $\eta_1(t_1) = f_{t_2}^{2,J} \circ \eta_1^*(t_1, t_2)$ for every $(t_1, t_2) \in \mathcal{D}_2$. For $\underline{t} = (t_1, t_2) \in \mathcal{D}_2$, let $K_{t_1}^{1,t_2} = \operatorname{Hull}(\eta_1^*([0, t_1], t_2))$ and $f_{t_1}^{1,t_2}$ be the centered Loewner map $f_{K_{t_1}^{1,t_2},\eta_1^*(t_1, t_2)}$. Let $f_{\underline{t}}^1 = f_{t_2}^{2,J} \circ f_{t_1}^{1,t_2}$. It is clear that $\{\underline{t} \in \mathcal{D}_2\} = \{\eta_2^J(t_2) \notin \eta_1[0, t_1]\} \in \mathcal{F}_{\underline{t}}$, and $f_{t_2}^{2,J}$ is $\mathcal{F}_{t_2}^2$ -measurable. Since $\eta_1^*(t, t_2), 0 \leq t \leq t_1$, are determined by $\eta_1|_{[0,t_1]}$ and $\eta_2|_{[0,t_2]}$, $f_{t_1}^{1,t_2}$ defined on the event $\{\underline{t} \in \mathcal{D}_2\}$ is $\mathcal{F}_{\underline{t}}$ -measurable, and so is $f_{\underline{t}}^1 = f_{t_2}^{2,J} \circ f_{t_1}^{1,t_2}$. Since η_1^* is continuous on \mathcal{D}_2 , by Proposition A.9, $(t_1, t_2; z) \mapsto f_{t_1}^{1,t_2}(z)$ is jointly continuous on $\mathcal{D}_2 \times \mathbb{H}$.

- **Lemma A.13.** (i) For every $\underline{t} = (t_1, t_2) \in \mathcal{D}_2$, $f_{\underline{t}}^1$ is a conformal map on \mathbb{H} , has continuation on $\overline{\mathbb{H}}^{\#}$, and maps 0 and ∞ respectively to the points $\eta_1(t_1)$ and $\eta_2^J(t_2)$. Moreover, $f_{\underline{t}}^1(\mathbb{H})$ is the connected component of $\mathbb{H} \setminus (\eta_1[0, t_1] \cup \eta_2^J[0, t_2])$ which shares the prime end $\eta_2^J(t_2)$ with $\mathbb{H} \setminus K_{t_2}^{2,J}$; and f_t^1 sends the prime end ∞ to the prime end $\eta_2^J(t_2)$.
- (ii) If $\underline{t} = (t_1, t_2) \in \mathcal{D}_{\cap}$, then besides the results in (i), $f_{\underline{t}}^1(\mathbb{H})$ also shares the prime end $\eta_1(t_1)$ with $\mathbb{H} \setminus K_{t_1}^1$, and $f_{\underline{t}}^1$ sends the prime end 0 to the prime end $\eta_1(t_1)$.

Proof. (i) Let $\underline{t} = (t_1, t_2) \in \mathcal{D}_2$. Since $f_{t_2}^{2,J}$ and $f_{t_1}^{1,t_2}$ are both conformal maps from \mathbb{H} into \mathbb{H} , and have continuation on $\overline{\mathbb{H}}^{\#}$, the same is true for $f_{\underline{t}}^1 = f_{t_2}^{2,J} \circ f_{t_1}^{1,t_2}$. Since $f_{t_1}^{1,t_2}(0) = \eta_1^*(\underline{t})$, we get $f_{\underline{t}}^1(0) = f_{t_2}^{2,J} \circ \eta_1^*(\underline{t}) = \eta_1(t_1)$. Since $f_{t_1}^{1,t_2}(\mathbb{H}) = \mathbb{H} \setminus K_{t_1}^{1,t_2}$ is a connected component of $\mathbb{H} \setminus \eta_1^*([0,t_1],t_2), f_{\underline{t}}^1(\mathbb{H})$ is a connected component of $f_{t_2}^{2,J}(\mathbb{H}) \setminus f_{t_2}^{2,J} \circ \eta_1^*([0,t_1],t_2) = \mathbb{H} \setminus (K_{t_2}^{2,J} \cup \eta_1[0,t_1])$, which is also a connected component of $\mathbb{H} \setminus (\eta_2^J[0,t_2] \cup \eta_1[0,t_1])$. Since $f_{t_1}^{1,t_2}(\mathbb{H}) = \mathbb{H} \setminus K_{t_1}^{1,t_2}$ is the connected component of $\mathbb{H} \setminus \eta_1^*([0,t_1],t_2)$ that shares the

Since $f_{t_1}^{1,t_2}(\mathbb{H}) = \mathbb{H} \setminus K_{t_1}^{1,t_2}$ is the connected component of $\mathbb{H} \setminus \eta_1^*([0,t_1],t_2)$ that shares the prime end ∞ with \mathbb{H} , we see that $f_{\underline{t}}^1(\mathbb{H}) = f_{t_2}^{2,J}(\mathbb{H} \setminus K_{t_1}^{1,t_2})$ is contained in and shares the prime end $f_{t_2}^{2,J}(\infty) = \eta_2^J(t_2)$ with $f_{t_2}^{2,J}(\mathbb{H}) = \mathbb{H} \setminus K_{t_2}^{2,J}$. Moreover, in terms of prime ends, we have

$$f_{\underline{t}}^{1}(\infty) = J \circ f_{t_{2}}^{2} \circ J \circ f_{t_{1}}^{1,t_{2}}(\infty) = J \circ f_{t_{2}}^{2} \circ J(\infty) = J \circ f_{t_{2}}^{2}(0) = \eta^{J}(t_{2}).$$

(ii) Suppose now $\underline{t} \in \mathcal{D}_{\cap}$. Then $\eta_1(t_1) \notin \eta_2^J[0, t_2]$. So there is a connected component of $\mathbb{H} \setminus (\eta_1[0, t_1] \cup \eta_2^J[0, t_2])$, which is contained in and shares the prime end $\eta_1(t_1)$ with $\mathbb{H} \setminus K_{t_1}^1$. Let the domain be denoted by $D_{\underline{t}}^1$. There is $\delta > 0$ such that $\eta_1[t_1, t_1 + \delta] \cap \eta_2^J[0, t_2] = \emptyset$. Let $S = \{t \in [t_1, t_1 + \delta] : \eta_1(t) \in \mathbb{H} \setminus K_{t_1}^1\}$. Then $\eta_1(S) \subset D_{\underline{t}}^1$, and S is dense in $[t_1, t_1 + \delta]$. Since $\eta_1 = f_{t_2}^{2,J} \circ \eta_1^*(\cdot, t_2)$, we get $\eta_1^*(S, t_2) \subset \mathbb{H} \setminus \eta_1^*([0, t_1], t_2)$, which implies that $\eta_1^*(S, t_2) \subset \mathbb{H} \setminus K_{t_1}^{1,t_2} = f_{t_1}^{1,t_2}(\mathbb{H})$. So $\eta_1(S) \subset f_{t_2}^{2,J} \circ f_{t_1}^{1,t_2}(\mathbb{H}) = f_{\underline{t}}^1(\mathbb{H})$. Now $f_{\underline{t}}^1(\mathbb{H})$ and $D_{\underline{t}}^1$ are both some connected component of $\mathbb{H} \setminus (\eta_1[0, t_1] \cup \eta_2^J[0, t_2])$, and have nonempty intersection: $\eta_1(S)$. So they must agree. Thus, $f_t^1(\mathbb{H})$ is contained in and shares the prime end $\eta_1(t_1)$ with $\mathbb{H} \setminus K_{t_1}^1$.

As $t \downarrow t_1$ along S, $\eta_1^*(\bar{t}, t_2)$ and $\eta_1(t) = f_{t_2}^{2,J} \circ \eta_1^*(t, t_2)$ respectively approach the prime ends $\eta_1^*(t_1, t_2)$ of $\mathbb{H} \setminus K_{t_1}^{1,t_2}$ and $\eta_1(t_1)$ of $D_{\underline{t}}^1$. So $f_{t_2}^{2,J}$ sends the prime end $\eta_1^*(\underline{t})$ to the prime end $\eta_1(t_1)$. Thus, in terms of prime ends, $f_{\underline{t}}^1(0) = f_{t_2}^{2,J} \circ f_{t_1}^{1,t_2}(0) = f_{t_2}^{2,J}(\eta_1^*(\underline{t})) = \eta_1(t_1)$.

Lemma A.14. For any $\overline{\mathcal{F}}$ -stopping time \underline{T} , the event $\{\underline{T} \in \mathcal{D}_2\}$ is $\overline{\mathcal{F}}_{\underline{T}}$ -measurable, and $f_{\underline{T}}^1$ is an $\overline{\mathcal{F}}_{\underline{T}}$ -measurable random conformal map defined on the event $\{\underline{T} \in \mathcal{D}_2\}$.

Proof. Let $\underline{T} = (T_1, T_2)$ be an $\overline{\mathcal{F}}$ -stopping time. For each $n \in \mathbb{N}$ and j = 1, 2, define $T_j^{\lfloor n \rfloor} = 2^{-n} \lfloor 2^n T_j \rfloor$ and $T_j^n = T_j^{\lfloor n \rfloor} + 2^{-n}$. Let $\underline{T}^{\lfloor n \rfloor} = (T_1^{\lfloor n \rfloor}, T_2^{\lfloor n \rfloor})$ and $\underline{T}^n = (T_1^n, T_2^n)$. Then $\underline{T}^{\lfloor n \rfloor} \leq T \leq \underline{T}^n, \underline{T}^n$ is an \mathcal{F} -stopping time, $\underline{T}^{\lfloor n \rfloor} \uparrow \underline{T}$, and $\underline{T}^n \downarrow \underline{T}$. For each $n \in \mathbb{N}$ and $\underline{t} \in (2^{-n} \mathbb{N})^2$, by Proposition 2.5

$$\{[\underline{T}^{\lfloor n \rfloor}, \underline{T}^n] \subset \mathcal{D}_2\} \cap \{\underline{T}^n = \underline{t}\} = \{[\underline{t} - (2^{-n}, 2^{-n}), \underline{t}] \subset \mathcal{D}_2\} \cap \{\underline{T}^n = \underline{t}\}$$
$$= \{\eta_2^J[t_2 - 2^{-n}, t_2] \cap \eta_1[0, t_1] = \emptyset\} \in \mathcal{F}_{\underline{t}} \cap \{\underline{T}^n = \underline{t}\} \subset \mathcal{F}_{\underline{T}^n}.$$

Thus, $\{[\underline{T}^{\lfloor n \rfloor}, \underline{T}^n] \subset \mathcal{D}_2\} = \bigcup_{\underline{t} \in (2^{-n} \mathbb{N})^2} (\{[\underline{T}^{\lfloor n \rfloor}, \underline{T}^n] \subset \mathcal{D}_2\} \cap \{\underline{T}^n = \underline{t}\}) \in \mathcal{F}_{\underline{T}^n}$. Similarly, for each $\underline{t} \in (2^{-n} \mathbb{N})^2$, $f_{\underline{T}^n}^1$ restricted to $\{\underline{T}^n \in \mathcal{D}_2\} \cap \{\underline{T}^n = \underline{t}\}$ is measurable w.r.t. $\mathcal{F}_{\underline{t}} \cap \{\underline{T}^n = \underline{t}\} \subset \mathcal{F}_{\underline{T}^n}$, and so $f_{\underline{T}^n}^1$ is $\mathcal{F}_{\underline{T}^n}$ -measurable on $\{\underline{T}^n \in \mathcal{D}_2\}$.

Since \mathcal{D}_2 is open, by Proposition 2.9, $\{\underline{T} \in \mathcal{D}_2\} = \bigcup_{m \ge l} \bigcap_{n \ge m} \{[\underline{T}^{\lfloor n \rfloor}, \underline{T}^n] \subset \mathcal{D}_2\} \in \mathcal{F}_{\underline{T}^l}$ for any $l \in \mathbb{N}$. Thus, $\{\underline{T} \in \mathcal{D}_2\} \in \bigcap_l \mathcal{F}_{\underline{T}^l} = \overline{\mathcal{F}}_{\underline{T}}$. Since $f_{\underline{t}}^1(z)$ is jointly continuous on $\mathcal{D}_2 \times \mathbb{H}$, $f_{\underline{T}^n}^1 \xrightarrow{\text{l.u.}} f_{\underline{T}}^1$ in \mathbb{H} on $\{\underline{T} \in \mathcal{D}_2\}$. By Proposition 2.9, $f_{\underline{T}}^1$ is $\overline{\mathcal{F}}_{\underline{T}}$ -measurable.

Lemma A.15. Let \mathcal{F}_n , $n \in \mathbb{N}$, and \mathcal{F}_0 be sub- σ -algebras of a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\mathcal{F}_0 \subset \bigcap_n \mathcal{F}_n$. Let $E_n \in \mathcal{F}_n$, $n \in \mathbb{N} \cup \{0\}$, be such that $E_0 \subset \liminf E_n$. For $n \in \mathbb{N} \cup \{0\}$, let $X_n(t)$, $t \ge 0$, be a continuous process defined on the event E_n such that a.s. on E_0 , $X_n \to X_0$ pointwise on \mathbb{R}_+ . Let μ be a probability measure on $C(\mathbb{R}_+)$. Suppose for each $n \in \mathbb{N}$, the conditional law of X_n given \mathcal{F}_n and E_n is μ . Then the conditional law of X_0 given \mathcal{F}_0 and E_0 is also μ .

Proof. Let $S = \{t_1 < \cdots < t_m\}$ be a nonempty finite subset of \mathbb{R}_+ , and μ_S be the finite dimensional distribution for μ on S. Then for any $n \in \mathbb{N}$, the conditional law of $(X_n(t_1), \ldots, X_n(t_m))$ given \mathcal{F}_n and E_n is μ_S . It suffices to show that the same is true for X_0 , \mathcal{F}_0 and E_0 . Let $f \in C_b(\mathbb{R}^m, \mathbb{R})$ and $A_0 \in \mathcal{F}_0 \cap E_0$. Then for every $n \in \mathbb{N}$, $A_0 \cap E_n \in \mathcal{F}_n \cap E_n$, and so

Since $E_0 \subset \liminf E_n$, we have $\mathbf{1}_{A_0 \cap E_n} \to \mathbf{1}_{A_0}$. By the continuity of f and that $X_n \to X$, we get $f(X_n(t_1), \ldots, X_n(t_m)) \to f(X_0(t_1), \ldots, X_0(t_m))$. Letting $n \to \infty$ and applying the dominated convergence theorem, we get $\mathbb{E}[\mathbf{1}_{A_0}f(X_0(t_1), \ldots, X_0(t_m))] = \mathbb{P}[A_0] \int f d\mu_S$. Since this holds for any $f \in C_b(\mathbb{R}^m)$, the conditional law of $(X_0(t_1), \ldots, X_0(t_m))$ given \mathcal{F}_0 and E_0 is μ_S .

Proof of Lemma A.5. By Lemma A.14, $\{\underline{T} \in \mathcal{D}_2\} \in \overline{\mathcal{F}}_{\underline{T}}$ and $f_{\underline{T}}^1$ is $\overline{\mathcal{F}}_{\underline{T}}$ -measurable. Symmetrically, $\{\underline{T} \in \mathcal{D}_1\} \in \overline{\mathcal{F}}_{\underline{T}}$. So $\{\underline{T} \in \mathcal{D}_{\cap}\} \in \overline{\mathcal{F}}_{\underline{T}}$. We are going to show that Lemma A.5 holds with $f_{\underline{T}} := f_{\underline{T}}^1 = f_{T_2}^{2,J} \circ f_{T_1}^{1,T_2}$. By Lemma A.13, on the event $\{\underline{T} \in \mathcal{D}_{\cap}\}, f_{\underline{T}}^1$ maps \mathbb{H} conformally onto H_T^1 , and sends 0 and ∞ respectively to $\eta_1(t_1)$ and $\eta_2^J(t_2)$.

We first consider four special cases. In the first three cases we prove a stronger result, where $\{\underline{T} \in \mathcal{D}_{\cap}\}$ is replaced by a bigger event $\{\underline{T} \in \mathcal{D}_{2}\}$.

<u>Case 1.</u> The <u>T</u> is a deterministic point $\underline{t} = (t_1, t_2) \in \mathbb{R}^2_+$. This is not completely trivial because we have to handle the right-continuous augmentation. We prove Case 1 in two steps.

<u>Step 1.</u> We prove that the statement holds with $\overline{\mathcal{F}}$ replaced by \mathcal{F} (without right-continuous augmentation). We give a complete proof for this simpler statement, which also prepares us for the proof in Step 2. We write E for the event $\{\underline{t} \in \mathcal{D}_2\}$.

Recall that $\zeta_1^{t_2}$ is a standard chordal SLE_{κ} curve independent of $\mathcal{F}_{t_2}^1$, $u_1^{t_2}$ is an increasing homeomorphism from $[0, \phi_2(t_2)]$ onto $[0, \infty]$, and $\eta_1 = f_{t_2}^{2,J} \circ \zeta_{t_2}^1 \circ u_{t_2}^1$ on $[0, \phi_2(t_2)]$. Let $\mathcal{F}_t^{1,t_2} = \sigma(\sigma(\zeta_1^{t_2}(s):s \leq t) \cup \mathcal{F}_{t_2}^2), t \geq 0$. We extend $u_1^{t_2}$ from $[0, \phi_2(t_2))$ to $[0, \infty]$ such that it equals ∞ on $[\phi_2(t_2), \infty]$. For any $a \geq 0$, $\{u_1^{t_2}(t_1) \leq a\} = \{\operatorname{hcap}_2(f_{t_2}^{2,J} \circ \zeta_1^{t_2}[0,a]) \geq t\} \in \mathcal{F}_a^{1,t_2}$. Thus, $u_1^{t_2}(t_1)$ is an \mathcal{F}^{1,t_2} -stopping time. Applying the usual DMP to the standard chordal SLE_{κ} curve $\zeta_1^{t_1}$ at the time $u_1^{t_2}(t_1)$, we get a standard chordal SLE_{$\kappa} curve \zeta_1^t$ defined on the event E, which is conditionally independent of $\mathcal{F}_{u_1^{t_2}(t_1)}^{1,t_2}$, such that $\zeta_1^{t_2}(u_1^{t_2}(t_1) + \cdot) = f_{t_1}^{1,t_2} \circ \zeta_1^t$ as on E. Let $u_1^t = (u_1^{t_2})^{-1}(u_1^{t_2}(t_1) + \cdot)$. Then u_1^t is an increasing homeomorphism from $[0, \infty]$ onto $[t_1, \phi_2(t_2)]$. Combining $f_{\underline{t}}^1 = f_{t_2}^{2,J} \circ f_{t_1}^{1,t_2}$, $\eta_1 = f_{t_2}^{2,J} \circ \zeta_1^{t_2} \circ u_1^{t_2}$ and $\zeta_1^{t_2}(u_1^{t_2}(t_1) + \cdot) = f_{t_1}^{1,t_2} \circ \zeta_{\underline{t}}^t$, we get $\eta_1 \circ u_{\underline{t}}^t = f_{t_2}^{1,0} \circ f_{t_1}^{1,t_2}$, $\eta_1 = f_{t_2}^{2,J} \circ \zeta_1^{t_2} \circ u_1^{t_2}$ and $\zeta_1^{t_2}(u_1^{t_2}(t_1) + \cdot) = f_{t_1}^{1,t_2} \circ \zeta_{\underline{t}}^t$, we get $\eta_1 \circ u_{\underline{t}}^t = f_{t_2}^t \circ f_{\underline{t}}^t$ as. on E.</sub>

 $\eta_1 \circ u_1^{\underline{t}} = f_{\underline{t}}^1 \circ \zeta_1^{\underline{t}} \text{ a.s. on } E.$ Since $\mathcal{F}_{t_2}^2 \subset \mathcal{F}_{u_1^{t_2}(t_1)}^{1,t_2}$, and on the event E, for $0 \leq t \leq t_1$, $\eta_1(t) = f_{t_2}^{2,J} \circ \zeta_1^{t_2} \circ u_1^{t_2}(t)$, we have $\mathcal{F}_{\underline{t}} \cap E \subset \mathcal{F}_{u_1^{t_2}(t_1)}^{1,t_2}$. Since given E, $\zeta_1^{\underline{t}}$ is conditionally independent of $\mathcal{F}_{u_1^{t_2}(t_1)}^{1,t_2}$, it is also conditionally independent of $\mathcal{F}_{\underline{t}}$. So we have finished Step 1 of Case 1.

<u>Step 2.</u> We prove that the $\zeta_1^{\underline{t}}$ in Step 1 is conditionally independent of $\overline{\mathcal{F}}_{\underline{t}}$. When this is done, the proof of Case 1 is complete.

Let $\Omega_2 = \{(u_1^*(s_1, s_2), s_2) : (s_1, s_2) \in \mathcal{D}_2\}$. Define $u_1^{*-1}, \zeta_1^*, \widehat{v}_1^*$ on Ω_2 such that for any $s_2 \in \mathbb{R}_+, u_1^{*-1}(\cdot, s_2) = u_1^*(\cdot, s_2)^{-1}, \zeta_1^*(\cdot, s_2) = \eta_1^*(u_1^{*-1}(\cdot, s_2), s_2), \text{ and } \widehat{v}_1^*(\cdot, s_2) = \widehat{w}_1^*(u_1^{*-1}(\cdot, s_2), s_2).$ Then ζ_1^* and \widehat{v}_1^* are continuous on Ω_2 . For any $s_2 \ge 0$, $\zeta_1^*(\cdot, s_2)$ is the chordal Loewner curve driven by $\widehat{v}_1^*(\cdot, s_2)$, and $\zeta_1^*(\cdot, s_2)$ and $\widehat{v}_1^*(\cdot, s_2)$ are respectively equal to $\zeta_1^{s_2}$ and $\widehat{v}_1^{s_2}$.

From the construction of $\zeta_1^t(t)$, $t \ge 0$, its driving function, say $\hat{v}_1^t(t)$, is given by $\hat{v}_1^t(t) := \hat{v}_1^{t_2}(u_1^{t_2}(t_1) + t) - \hat{v}_1^{t_2}(u_1^{t_2}(t_1))$, $t \ge 0$. By discarding a null event, we may assume that $\hat{v}_1^t = \hat{v}_1^*(u_1^*(t_1, t_2) + \cdot, t_2) - \hat{v}_1^*(u_1^*(t_1, t_2), t_2)$. Similarly, we may assume that $u_1^t = u_1^{*-1}(u_1^*(t_1, t_2) + \cdot, t_2)$. Then \hat{v}_1^t and u_1^t are both continuous in \underline{t} by continuity of $\hat{v}_1^*, u_1^*, u_1^{*-1}$.

Choose $\underline{t}^n \downarrow \underline{t}$. Then $\overline{\mathcal{F}}_{\underline{t}} \subset \bigcap_n \mathcal{F}_{\underline{t}^n}$. Let $E_n = \{\underline{t}^n \in \mathcal{D}_2\} \in \mathcal{F}_{\underline{t}^n}$. Since \mathcal{D}_2 is open, we have $E \subset \liminf E_n$. For any $n \in \mathbb{N}$, by the result of Step 1, conditionally on $\mathcal{F}_{\underline{t}^n}$ and E_n , $\widehat{v}_1^{\underline{t}^n}/\sqrt{\kappa}$ has the law of a standard Brownian motion. By the continuity of $\widehat{v}_1^{\underline{t}}$ in \underline{t} , $\widehat{v}_1^{\underline{t}^n} \to \widehat{v}^{\underline{t}}$ pointwise. By Lemma A.15, $\widehat{v}_1^{\underline{t}}$ is conditionally independent of $\overline{\mathcal{F}}_{\underline{t}}$ given E. The same is true for $\zeta_1^{\underline{t}}$ since it is the chordal Loewner curve determined by $\widehat{v}_1^{\underline{t}}$. This finishes Step 2.

<u>Case 2.</u> T takes values in a countable set S. This follows easily from Case 1.

<u>Case 3.</u> T_2 takes values in a countable set S_2 . For $n \in \mathbb{N}$, define $\underline{T}^n = (T_1^n, T_2)$ such that $T_1^n = 2^{-n} \lceil 2^n T_1 \rceil$. Then each \underline{T}^n is an $\overline{\mathcal{F}}$ -stopping time taking values in a countable set. By the

results of Cases 1 and 2, for each n, on the event $E_n := \{\underline{T}^n \in \mathcal{D}_2\}$ there are a chordal Loewner curve $\zeta_1^{\underline{T}^n}$, whose driving function is given by $\hat{v}_1^{\underline{T}^n} := \hat{v}_1^*(u_1^*(\underline{T}^n) + \cdot, T_2) - \hat{v}_1^*(u_1^*(\underline{T}^n), T_2)$, and an increasing homeomorphism $u_1^{\underline{T}^n} := u_1^{*-1}(u_1^*(\underline{T}^n) + \cdot, T_2)$ from $[0, \infty]$ onto $[T_1^n, \phi_2(T_2)]$, such that a.s. $\eta_1 \circ u_1^{\underline{T}^n} = f_{\underline{T}^n}^1 \circ \zeta_1^{\underline{T}^n}$ on E_n , and $\hat{v}_1^{\underline{T}^n}/\sqrt{\kappa}$ is a standard Brownian motion conditionally independent of $\overline{\mathcal{F}}_{\underline{T}^n}$ given the event E_n . Using the same argument as in Step 2 of Case 1, we know that $\hat{v}_1^{\underline{T}} := \hat{v}_1^*(u_1^*(\underline{T}) + \cdot, T_2) - \hat{v}_1^*(u_1^*(\underline{T}), T_2)$ is conditionally independent of $\overline{\mathcal{F}}_{\underline{T}}$ with the law of $\sqrt{\kappa}B$ given $E := \{\underline{T} \in \mathcal{D}_2\} = \bigcup_n E_n$. Thus, $\hat{v}_1^{\underline{T}}$ a.s. generates a chordal Loewner curve $\zeta_1^{\underline{T}}$, which has the law of a standard chordal SLE_{κ}, and is conditionally independent of $\overline{\mathcal{F}}_{\underline{T}}$ given E. Since $\hat{v}_1^*(\cdot, T_2)$ is the driving function for $\zeta_1^*(\cdot, T_2)$, from the definitions of $\hat{v}_1^{\underline{T}}$ and $f_{T_1}^{1,T_2}$ we know that $\zeta_1^*(u_1^*(\underline{T}) + \cdot, T_2) = f_{T_1}^{1,T_2} \circ \zeta_1^{\underline{T}}$. Combining this with $\eta_1^*(\cdot, T_2) = \zeta_1^*(\cdot, T_2) \circ u_1^*(\cdot, T_2)$, $\eta_1 = f_{T_2}^{2,J} \circ \eta_1^*(\cdot, T_2), u_1^{\underline{T}} = u_1^{*-1}(u_1^*(\underline{T}) + t, T_2)$, and $f_{\underline{T}}^1 = f_{T_2}^{2,J} \circ f_{T_1}^{1,T_2}$, we get $\eta_1 \circ u_1^{\underline{T}} = f_{\underline{T}}^1 \circ \zeta_1^{\underline{T}}$ a.s. on $\{\underline{T} \in \mathcal{D}_2\}$.

<u>Case 4.</u> T_1 takes values in a countable set S_1 . Swapping indices "1" and "2" and using the result of Case 3, we know that on the event $\{\underline{T} \in \mathcal{D}_1\}$, there are

- an $\overline{\mathcal{F}}_{\underline{T}}$ -measurable conformal map $f_{\underline{T}}^2$ on \mathbb{H} with continuation on $\overline{\mathbb{H}}^{\#}$,
- a standard chordal SLE_{κ} curve $\zeta_2^{\underline{T}}$ conditionally independent of $\overline{\mathcal{F}}_{\underline{T}}$, and
- an increasing homeomorphism u_2^T from $[0, \infty]$ onto $[T_2, \phi_1(T_1)]$,

such that $\eta_2 \circ u_{\overline{2}}^{\underline{T}} = f_{\underline{T}}^2 \circ \zeta_{\overline{2}}^{\underline{T}}$ a.s. on $\{\underline{T} \in \mathcal{D}_1\}$. Moreover, on the event $E_{\cap} := \{\underline{T} \in \mathcal{D}_{\cap}\}, f_{\underline{T}}^2$ maps \mathbb{H} conformally onto the connected component $D_{\underline{T}}^2$ of $\mathbb{H} \setminus (\eta_2[0, T_2] \cup J \circ \eta_1[0, T_1])$ which shares the prime ends $\eta_2(T_2)$ and $J \circ \eta_1(T_1)$ respectively with $\mathbb{H} \setminus K_{T_2}^2$ and $\mathbb{H} \setminus J(K_{T_1}^1)$, and sends 0 and ∞ respectively to the prime ends $\eta_2(T_2)$ and $J \circ \eta_1(T_1)$.

Suppose E_{\cap} happens. We have $J(D_{\underline{T}}^2) = D_{\underline{T}}^1$ since they are both a connected component of $\mathbb{H} \setminus (\eta_1[0,T_1] \cup \eta_2^J[0,T_2])$ that shares the prime ends $\eta_1(T_1)$ and $\eta_2^J(T_2)$ respectively with $\mathbb{H} \setminus K_{T_1}^1$ and $\mathbb{H} \setminus K_{T_2}^{2,J}$. Let $f_{\underline{T}}^{2,J} = J \circ f_{\underline{T}}^2 \circ J$. Then both $f_{\underline{T}}^{2,J}$ and $f_{\underline{T}}^1$ map \mathbb{H} conformally onto $J(D_{\underline{T}}^2) = D_{\underline{T}}^1$, send 0 and ∞ respectively to the same prime ends $\eta_1(T_1)$ and $\eta_2^J(T_2)$, and are $\overline{\mathcal{F}}_{\underline{T}}$ -measurable. So there is an $\overline{\mathcal{F}}_{\underline{T}}$ -measurable random variable a > 0 such that $f_{\underline{T}}^{2,J} = f_{\underline{T}}^1(\cdot/a)$.

Applying the reversibility of SLE to ζ_2^T , we get another standard chordal SLE_{κ} curve $\tilde{\zeta}_1^T$, which is also conditionally independent of $\overline{\mathcal{F}}_T$, and a decreasing auto-homeomorphism $\hat{\phi}_2$ of $[0, \infty]$, such that a.s. $J \circ \zeta_2^T = \tilde{\zeta}_1^T \circ \hat{\phi}_2$. Let $u_1^T(t) = \phi_2 \circ u_2^T \circ \hat{\phi}_2^{-1}(a^2t)$. Then u_1^T is an increasing homeomorphism from $[0, \infty]$ onto $[T_1, \phi_2(T_2)]$. Let $\zeta_1^T(t) = \tilde{\zeta}_1^T(a^2t)/a$. Since a is $\overline{\mathcal{F}}_T$ -measurable, by the scaling property of chordal SLE_{κ}, ζ_1^T is also a standard chordal SLE_{κ} curve conditionally independent of $\overline{\mathcal{F}}_T$. Moreover, on $\{\underline{T} \in \mathcal{D}_{\cap}\}$, a.s.

$$J \circ \eta_1 \circ u_1^{\underline{T}} = \eta_2 \circ \phi_2^{-1} \circ u_1^{\underline{T}} = \eta_2 \circ u_2^{\underline{T}} \circ \widehat{\phi}_2^{-1}(a^2 \cdot) = f_{\underline{T}}^2 \circ \zeta_2^{\underline{T}} \circ \widehat{\phi}_2^{-1}(a^2 \cdot)$$
$$= f_{\underline{T}}^2 \circ J \circ \widetilde{\zeta}_1^{\underline{T}}(a^2 \cdot) = f_{\underline{T}}^2 \circ J(a\zeta_1^{\underline{T}}) = J \circ f_{\underline{T}}^{2,J}(a\zeta_1^{\underline{T}}) = J \circ f_{\underline{T}}^1 \circ \zeta_1^{\underline{T}},$$

which implies that $\eta_1 \circ u_1^T = f_T^1 \circ \zeta_1^T$. This completes Case 4.

<u>General Case.</u> For $n \in \mathbb{N}$, define $\underline{T}^n = (T_1^n, T_2)$ such that $T_1^n = 2^{-n} \lceil 2^n T_1 \rceil$. Then each \underline{T}^n is an $\overline{\mathcal{F}}$ -stopping time with the first variable taking values in a countable set. Let $E_0 = \{\underline{T} \in \mathcal{D}_{\cap}\}$ and $E_n = \{\underline{T}^n \in \mathcal{D}_{\cap}\}, n \in \mathbb{N}$. By Case 4, for each n, on the event E_n , there are a standard chordal SLE_{κ} curve $\zeta_1^{\underline{T}^n}$, which is conditionally independent of $\overline{\mathcal{F}}_{\underline{T}^n}$ given E_n , and an increasing homeomorphism $u_1^{\underline{T}^n}$ from $[0, \infty]$ onto $[T_1^n, \phi_2(T_2)]$, such that $\eta_1 \circ u_1^{\underline{T}^n} = f_{\underline{T}^n}^1 \circ \zeta_1^{\underline{T}^n}$ a.s. on E_n .

Suppose E_n happens. Then $T_1^n < \phi_2(T_2)$. Let $u_1^n = (u_1^{\underline{T}^n})^{-1}(\cdot + T_1^n)$, which is an increasing homeomorphism from $[0, \phi_2(T_2) - T_1^n]$ onto $[0, \infty]$. Let $\widehat{v_1}^{\underline{T}^n}$ be the driving function for $\zeta_1^{\underline{T}^n}$. Let $\eta_1^{\underline{T}^n} = \zeta_1^{\underline{T}^n} \circ u_1^n$ and $\widehat{w_1}^{\underline{T}^n} = \widehat{v_1}^{\underline{T}^n} \circ u_1^n$. Then $\eta_1^{\underline{T}^n}$ is the chordal Loewner curve with speed du_1^n driven by $\widehat{w_1}^{\underline{T}^n}$. From a.s. $\eta_1 \circ u_1^{\underline{T}^n} = f_{\underline{T}^n} \circ \zeta_1^{\underline{T}^n}$ we get a.s. $\eta_1(T_1^n + \cdot) = f_{\underline{T}^n}^1 \circ \eta_1^{\underline{T}^n}$.

Define $\eta_1^{T_2,n}, u_1^{T_2,n}, \hat{w}_1^{T_2,n}$ on $[0, \phi_2(T_2))$ such that, for $0 \le t_1 < T_1^n, \eta_1^{T_2,n}(t_1) = \eta_1^*(t_1, T_2), u_1^{T_2,n}(t_1) = \hat{w}_1^*(t_1, T_2);$ and for $T_1^n \le t_1 < \phi_2(T_2), \eta_1^{T_2,n}(t_1) = f_{T_1^{n}}^{T_2,n}(t_1) = u_1^*(T_1^n) + u_1^n(t_1 - T_1^n), u_1^{T_2,n}(t_1) = \hat{w}_1^*(T_1^n) + \hat{w}_1^{T^n}(t_1 - T_1^n).$ Since $\eta_1^*(\cdot, t_2)$ is the chordal Loewner curve with speed $du_1^*(\cdot, t_2)$ driven by $\hat{w}_1^*(\cdot, t_2), f_{T_1^n}^{T_1,T_2}$ is the centered Loewner map for $\eta_1^*(\cdot, t_2)$ at the time T_1^n , and $\eta_1^{T^n}$ is the chordal Loewner curve with speed $du_1^{T_2,n}$ is the chordal Loewner curve with speed $du_1^{T_2,n}$ and $f_1^{T_2,n}$, we get $\eta_1(t) = f_{T_2}^{2,J} \circ \eta_1^{T_2,n}(t), 0 \le t < \phi_2(T_2), \eta_1(T_1^n + \cdot) = f_{T_1^n} \circ \eta_1^{T^n}, and f_{T_1^n} = f_{T_2}^{2,J} \circ f_{T_1^n}^{1,T_2}, we get <math>\eta_1(t) = f_{T_2}^{2,J} \circ \eta_1^{T_2,n}(t), 0 \le t < \phi_2(T_2), a.s.$ on E_n .

driven by $w_1^{-2,n}$. From a.s. $\eta_1 = f_{T_2}^{-1} \circ \eta_1^*(\cdot, T_2)$, $\eta_1(T_1^n + \cdot) = f_{\underline{T}^n} \circ \eta_{\underline{1}}^{-1}$, and $f_{\underline{T}^n} = f_{T_2}^{-1} \circ f_{T_1}^{-n}^{-2}$, we get $\eta_1(t) = f_{T_2}^{2,J} \circ \eta_1^{T_2,n}(t)$, $0 \le t < \phi_2(T_2)$, a.s. on E_n . Let $\zeta_1^{T_2,n} = \eta_1^{T_2,n} \circ (u_1^{T_2,n})^{-1}$ and $\hat{v}_1^{T_2,n} = \hat{w}_1^{T_2,n} \circ (u_1^{T_2,n})^{-1}$. Then $\zeta_1^{T_2,n}$ is the chordal Loewner curve driven by $\hat{v}_1^{T_2,n}$, and $\eta_1|_{[0,\phi_2(T_2)]} = f_{T_2}^{2,J} \circ \zeta_1^{T_2,n} \circ u_1^{T_2,n}$ a.s. on E_n . The last equality implies that a.s. $\zeta_1^{T_2,n} = \zeta_1^{T_2,n}$ and $u_1^{T_2,n} = u_1^{T_2,m}$ on $E_n \cap E_m$ for any $n, m \in \mathbb{N}$. Since $E_0 \subset \lim \inf E_n$, we may define a Loewner curve $\zeta_1^{T_2}$ and a time-change function $u_1^{T_2}$ on E_0 such that for every $n \in \mathbb{N}, \zeta_1^{T_2} = \zeta_1^{T_2,n}$ and $u_1^{T_2} = u_1^{T_2,n}$ a.s. on $E_0 \cap E_n$. Then we have $\eta_1|_{[0,\phi_2(T_2))} = f_{T_2}^{2,J} \circ \zeta_1^{T_2} \circ u_1^{T_2}$ a.s. on E_0 . Let $\hat{v}_1^{T_2}$ be the driving function for $\zeta_1^{T_2}$. Then $v_1^{T_2} = v_1^{T_2,n}$ a.s. on $E_0 \cap E_n$. By the definition of $u_1^{T_2,n}, (u_1^{T_2,n})^{-1}(u_1^*(\underline{T}^n) + t_1) = T_1^n + (u_1^n)^{-1}(t_1)$. By the definition of $\hat{w}_1^{T_2,n}, \hat{w}_1^{T_2,n}(T_1^n + (u_1^n)^{-1}(t_1)) = \hat{w}_1^*(\underline{T}^n) + \hat{w}_1^{\overline{T}^n} \circ (u_1^n)^{-1}(t_1)$. By these two equalities and the definition of $\hat{v}_1^{T_2,n}$ and that $\hat{w}_1^{\overline{T}^n} = \hat{v}_1^{\overline{T}^n} \circ u_1^n$ we get $\hat{v}_1^{T_n} = \hat{v}_1^{T_2,n}(u_1^*(\underline{T}^n) + \cdot) - v_1^{T_2,n}(u_1^*(\underline{T}^n))$. From the independence property of $\zeta_1^{\overline{T}^n}$, we see that, conditionally on $\overline{\mathcal{F}}_{\underline{T}^n}$ and the event $E_n, \hat{v}_1^{\overline{T}^n}$ has the law of $(\sqrt{\kappa} B_4)$. It is clear that as on $E_0, \hat{v}_1^{\overline{T}^n} \to \hat{v}_1^{\overline{T}} = \hat{v}_1^{T_2}(u_1^*(T) + \cdot) - \hat{v}_1^{T_2}(u_2^*(T))$ pointwise

By the definition of $u_1^{T_2,n}$, $(u_1^{T_2,n})^{-1}(u_1^*(\underline{T}^n) + t_1) = T_1^n + (u_1^n)^{-1}(t_1)$. By the definition of $\hat{w}_1^{T_2,n}$, $\hat{w}_1^{T_2,n}(T_1^n + (u_1^n)^{-1}(t_1)) = \hat{w}_1^*(\underline{T}^n) + \hat{w}_1^{\overline{T}^n} \circ (u_1^n)^{-1}(t_1)$. By these two equalities and the definition of $\hat{v}_1^{T_2,n}$ and that $\hat{w}_1^{\overline{T}^n} = \hat{v}_1^{\overline{T}^n} \circ u_1^n$ we get $\hat{v}_1^{\overline{T}^n} = \hat{v}_1^{T_2,n}(u_1^*(\underline{T}^n) + \cdot) - v_1^{T_2,n}(u_1^*(\underline{T}^n))$. From the independence property of $\zeta_1^{\overline{T}^n}$, we see that, conditionally on $\overline{\mathcal{F}}_{\underline{T}^n}$ and the event E_n , $\hat{v}_1^{\underline{T}^n}$ has the law of $(\sqrt{\kappa}B_t)$. It is clear that, a.s. on E_0 , $\hat{v}_1^{\overline{T}^n} \to \hat{v}_1^{\overline{T}} := \hat{v}_1^{T_2}(u_1^*(\underline{T}) + \cdot) - \hat{v}_1^{T_2}(u_1^*(\underline{T}))$ pointwise on \mathbb{R}_+ since $\hat{v}_1^{T_2,n} = \hat{v}_1^{T_2}$ and $u_1^{T_2,n} = \hat{u}_1^{T_2}$ for n big enough. By Lemma A.15, conditionally on $\overline{\mathcal{F}}_{\underline{T}}$ and the event E_0 , $\hat{v}_1^{\overline{T}}$ also has the law of $(\sqrt{\kappa}B_t)$. Thus, \hat{v}_1^T generates a chordal Loewner curve $\zeta_1^{\overline{T}}$ defined on E_0 , which has the law of a standard chordal SLE_{κ} conditionally on $\overline{\mathcal{F}}_{\underline{T}}$ and E_0 . Since $(f_t^{1,T_2})_{t\geq 0}$ are the centered Loewner maps for $\eta_1^{T_2} = \zeta_1^{T_2} \circ u_1^*(\cdot,T_2)$, we have $\zeta_1^{T_2}(u_1^*(\underline{T}) + \cdot) = f_{T_1}^{1,T_2} \circ \zeta_1^{\overline{T}}$. Let $u_1^T(s) = (u_1^{T_2})^{-1}(s + u_1^*(\underline{T}))$, $s \geq 0$. Then u_1^T is an increasing homeomorphism from $[0,\infty]$ onto $[T_1,\phi_2(T_2)]$. Since $\eta_1 = f_{T_2}^{2,J} \circ \zeta_1^{T_2} \circ u_1^{T_2}$ a.s. on E_0 and $f_{\underline{T}} = f_{T_2}^{2,J} \circ \zeta_1^{T_2} \circ u_1^{T_2}$ a.s. on E_0 and $f_{\underline{T}} = f_{\underline{T}}^2 \circ \zeta_1^T$ a.s. on E_0 , as desired.

References

- Lars V. Ahlfors. Conformal invariants: topics in geometric function theory. McGraw-Hill Book Co., New York, 1973.
- [2] V. Beffara. The dimension of SLE curves, Ann. Probab., **36**:1421-1452, 2008.
- [3] Julien Dubédat. Commutation relations for SLE, Comm. Pure Applied Math., 60(12):1792-1847, 2007.
- [4] Nina Holden, Gregory F. Lawler, Xinyi Li and Xin Sun. Minkowski content of Brownian cut points. In preprint, arXiv:1803.10613.
- [5] Gregory Lawler. Conformally Invariant Processes in the Plane, Amer. Math. Soc, 2005.
- [6] Gregory F. Lawler and Mohammad A. Rezaei. Minkowski content and natural parametrization for the Schramm-Loewner evolution. Ann. Probab., 43(3):1082-1120, 2015.
- [7] G. Lawler. O. Schramm, and W. Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees, Ann. Probab., 32:939–995, 2004.
- [8] G. Lawler and B. Werness. Multi-point Green's function for SLE and an estimate of Beffara, Annals of Prob. 41:1513-1555, 2013.
- [9] Jason Miller and Scott Sheffield. Imaginary Geometry III: reversibility of SLE_{κ} for $\kappa \in (4, 8)$. Ann. Math., 184(2):455-486, 2016.
- [10] Jason Miller and Scott Sheffield. Imaginary Geometry I: intersecting SLEs. Probab. Theory Relat. Fields, 164(3):553-705, 2016.
- [11] Jason Miller and Hao Wu. Intersections of SLE Paths: the double and cut point dimension of SLE. Prob. Theory Relat. Fields, 167(1-2):45-105, 2017.
- [12] NIST Digital Library of Mathematical Functions. http://dlmf.nist.gov/18, Release 1.0.6 of 2013-05-06.
- [13] Pommerenke Ch. Boundary behaviour of conformal maps. Springer-Verlag, Berlin, Heidelberg, 1992
- [14] Steffen Rohde and Oded Schramm. Basic properties of SLE. Ann. Math., 161:879-920, 2005.
- [15] Daniel Revuz and Marc Yor. Continuous Martingales and Brownian Motion. Springer, Berlin, 1991.
- [16] O. Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. Israel J. Math., 118:221-288, 2000.

- [17] Oded Schramm and David B. Wilson. SLE coordinate changes. New York J. Math., 11:659– 669, 2005.
- [18] Yuan Xu. Lecture notes on orthogonal polynomials of several variables. Inzell Lectures on Orthogonal Polynomials. W. zu Castell, F. Filbir, B. Forster (eds.). Advances in the Theory of Special Functions and Orthogonal Polynomials. Nova Science Publishers Volume 2, 2004, Pages 135-188.
- [19] Dapeng Zhan. Two-curve Green's function for 2-SLE: the boundary case, *Electron. J. Probab.*, 26, article no. 32:1-58, 2021.
- [20] Dapeng Zhan. Two-curve Green's function for 2-SLE: the interior case. Comm. Math. Phys., **375**:1-40, 2020.
- [21] Dapeng Zhan. Ergodicity of the tip of an SLE curve. *Prob. Theory Relat. Fields*, **164**(1):333-360, 2016.
- [22] Dapeng Zhan. Duality of chordal SLE, II. Ann. I. H. Poincare-Pr., 46(3):740-759, 2010.
- [23] Dapeng Zhan. Duality of chordal SLE. Invent. Math., **174**(2):309-353, 2008.
- [24] Dapeng Zhan. Reversibility of chordal SLE. Ann. Probab., 36(4):1472-1494, 2008.
- [25] Dapeng Zhan. The Scaling Limits of Planar LERW in Finitely Connected Domains. Ann. Probab. 36:467-529, 2008.

This figure "cut-png.png" is available in "png" format from:

http://arxiv.org/ps/2004.02242v2