EXTERNAL TRIANGULATION OF THE HOMOTOPY CATEGORY OF EXACT QUASI-CATEGORY

HIROYUKI NAKAOKA AND YANN PALU

ABSTRACT. Extriangulated categories axiomatize extension-closed subcategories of triangulated categories. We show that the homotopy category of an exact quasi-category can be equipped with a natural extriangulated structure.

Contents

1. Introduction	1
2. Notations and terminology	3
2.1. Squares, rectangles and cubes	3
3. Exact sequences in exact quasi-categories	10
3.1. Morphisms of exact sequences	11
3.2. Decompositions of morphisms	15
3.3. Splitting exact sequences	20
3.4. Equivalence of exact sequences	24
3.5. (Co)products of exact sequences	25
4. Extriangulated structure on the homotopy category	28
4.1. Construction of the functor \mathbb{E}	29
4.2. Biadditivity of \mathbb{E}	31
4.3. External triangulation of $h\mathscr{C}$	34
4.4. Compatibility with the triangulation in stable case	39
References	41

1. Introduction

For triangulated categories, several possible higher-categorical enhancements are known, one of which is the celebrated notion of a $stable \infty$ -category by Lurie ([L2]). Recently, as a wider class of ∞ -categories, the notion of an $exact \infty$ -category has appeared in Barwick's [B1], where it is used in order to prove that the K-theory of an idempotent complete stable quasi-category endowed with a bounded t-structure is weakly equivalent to the K-theory of its heart. A non-additive version of exact quasi-categories also appeared in [DK] as a source of examples of 2-Segal spaces, via Waldhausen's S-construction. Instances of exact quasi-categories are given by

• the nerve of any ordinary exact category,

1

The first author is supported by JSPS KAKENHI Grant Numbers JP19K03468, JP20K03532. Both authors were partially supported by the French ANR grant SC3A (15 \times CE40 0004 01). Both authors wish to thank Gustavo Jasso for his interest, and for introducing them a possible relation with a future work of his student.

 \bullet any extension-closed full additive subcategory of a stable ∞ -category as stated in [B1, Examples 3.3 and 3.5].

Regarding its importance and naturality, it should be natural to expect that the homotopy category of an exact ∞ -category would have a nice structure very close to triangulation. In this article, we give a positive answer to this expectation, with the notion of an extriangulated category which has been introduced as a unification of exact categories and triangulated categories in our previous article [NP]. It is defined as an additive category equipped with a structure called an external triangulation, which is more flexible than triangulations in the sense that it is naturally inherited by relative theories, extension-closed subcategories, and ideal quotients by some projective-injectives ([HLN, Section 3.2], [NP, Remark 2.18, Proposition 3.30]). In particular, extension-closed subcategories of triangulated categories are typical examples of extriangulated categories, which dovetails the above-mentioned feature of exact ∞ -categories.

Let $\mathscr C$ be an exact quasi-category (the definition is recalled in Section 3). We say that two exact sequences in $\mathscr C$

$$\mathbf{S} = \bigvee_{Q \longrightarrow C}^{i} B \qquad \qquad A \longrightarrow B'$$

$$\mathbf{S}' = \bigvee_{Q' \longrightarrow C} \bigvee_{Q' \longrightarrow C}$$

are equivalent if there is a cube in ${\mathcal C}$ from ${\bf S}$ to ${\bf S}'$

$$\mathbf{C} = \begin{bmatrix} A & \longrightarrow B \\ & & O & \longrightarrow C \\ & & & \downarrow C \\ & & & & \downarrow C \\ & & & & \downarrow C \\ & \\ \\ & \downarrow C \\ \\ \\ &$$

where $\overline{a} = \mathrm{id}_A$ and $\overline{c} = \mathrm{id}_C$ in the homotopy category $h\mathscr{C}$. This is an equivalence relation (Proposition 3.28), and we define $\mathbb{E}(C,A)$ to be the set of equivalence classes of exact sequences with fixed end terms A and C. Then $\mathbb{E}: (h\mathscr{C})^{\mathrm{op}} \times (h\mathscr{C}) \to Ab$ is an additive bifunctor (Corollary 4.19). We also define an additive realization \mathfrak{s} (Lemma 4.21) by sending the equivalence class of \mathbf{S} to the equivalence class (defined similarly as for short exact sequences) of $A \xrightarrow{\overline{i}} B \xrightarrow{\overline{p}} C$ in $h\mathscr{C}$.

Theorem 1.1 (Theorem 4.22 and Proposition 4.28). Let $\mathscr C$ be an exact quasicategory. Then

- (1) its homotopy category $(h\mathscr{C}, \mathbb{E}, \mathfrak{s})$ is extriangulated.
- (2) Moreover, the functor $\mathscr{C} \to h\mathscr{C}$ sends exact sequences \mathbf{S} as above to extriangles $A \xrightarrow{\overline{\imath}} B \xrightarrow{\overline{p}} C \xrightarrow{\mathbf{S}}$.
- (3) If $F: \mathcal{C} \to \mathcal{D}$ is an exact functor of exact quasi-categories, then the induced functor $hF: h\mathcal{C} \to h\mathcal{D}$ is an exact functor of extriangulated categories.

When $\mathscr C$ is stable, this recovers the usual triangulated structure on $h\mathscr C$ (Proposition 4.30).

In Section 2, we organize notations and terminology which will be used in the proceeding sections. In Section 3 we remind the definition of an exact ∞ -category

in [B1], and summarize properties of exact sequences. In Section 4, after a brief introduction of the definition of an extriangulated category, we proceed to construct an external triangulation of the homotopy category. As the main theorem (Theorem 4.22), indeed we will show that the homotopy category of any exact ∞ -category becomes an extriangulated category. In the last Subsection 4.4, in the stable case we will also see that this extriangulated structure is compatible with the triangulation given in [L2].

This work came out of our exploration for an existing higher-categorical notion appropriate to enhance extriangulated categories. We hope that the existence of such a notion in turn guarantees the validity and naturality of the notion of an extriangulated category.

2. Notations and terminology

Basically we follow the notations and terminology in [L1] and [B1], while we use the term quasi-category as in Joyal's [J], instead of ∞ -category. Besides, we will also employ some ad hoc ones which we will introduce in this section, to facilitate the later argument dealing with exact sequences.

Throughout this article, $\mathscr C$ denotes an essentially small quasi-category ([B2, Notation 1.3]). By definition it is a simplicial set which admits every inner horn extensions ([L1, Definition 1.1.2.4.]). As being a simplicial set, it has the face maps $d_k \colon \mathscr C_{n+1} \to \mathscr C_n$ $(k=0,1,\ldots,n+1)$, and the degeneracy maps $s_k \colon \mathscr C_n \to \mathscr C_{n+1}$ $(k=0,1,\ldots,n)$ for any $n \geq 0$, where each $\mathscr C_n$ denotes the set of its n-simplices. In particular 0-simplices and 1-simplices in $\mathscr C$ are called objects and morphisms. For any pair of objects $A, B \in \mathscr C_0$, we write $f \in \mathscr C_1(A,B)$ to indicate that f is a morphism satisfying $d_1(f) = A$ and $d_0(f) = B$. For an object $A \in \mathscr C_0$, we abbreviate $s_0(A) \in \mathscr C_1(A,A)$ to 1_A for simplicity. The homotopy category of $\mathscr C$ will be denoted by $h\mathscr C$. For a morphism $a \in \mathscr C_1(A,B)$, its image in the homotopy category will be denoted by $\overline{a} \in (h\mathscr C)(A,B)$. We use the symbol $\mathrm{id}_A = \overline{1}_A$ exclusively for the identity morphisms in the homotopy category.

2.1. Squares, rectangles and cubes. As in [L1, Section 4.4.2], a square in $\mathscr C$ is a map $\mathbf S\colon \Delta^1\times\Delta^1\to\mathscr C$, namely an object in the quasi-category of functors $\operatorname{Fun}(\Delta^1\times\Delta^1,\mathscr C)$ (see [L1, Notation 1.2.7.2, Proposition 1.2.7.3]). We will often denote $\mathbf S$ as a diagram in $\mathscr C$

$$\mathbf{S}(0,0) \to \mathbf{S}(0,1)$$

$$\downarrow \qquad \qquad \downarrow$$

$$\mathbf{S}(1,0) \to \mathbf{S}(1,1)$$

or by labelling some of its simplices

(2.1)
$$A \xrightarrow{x} B \qquad A \xrightarrow$$

It can be also identified with a pair of 2-simplices ξ and η satisfying $d_1(\xi) = d_1(\eta)$, which correspond to

$$\xi = \mathbf{S}|_{(\{0\} \times \Delta^1) * \{(1,1)\}}$$
 and $\eta = \mathbf{S}|_{\{(0,0)\} * (\{1\} \times \Delta^1)}$.

In this article we will also express a square as above by $\mathbf{S} = (\eta \setminus \xi)$. For a square $\mathbf{S} = (\eta \setminus \xi)$, its transpose $\mathbf{S}^t = (\xi \setminus \eta)$ is given by the below.

$$\begin{array}{c|c}
A & \xrightarrow{i} D \\
x & \xrightarrow{z} & \downarrow^{j} \\
B & \xrightarrow{y} & C
\end{array}$$

If we fix a map $F : \Lambda_0^2 \to \mathscr{C}$ denoted by a diagram

$$\begin{array}{ccc}
A & \xrightarrow{x} & B \\
\downarrow & & \downarrow \\
D & & D
\end{array}$$

in \mathscr{C} , then squares

$$\mathbf{S} = i \int_{0}^{1} \int_{0}^{1} \mathbf{S} d\mathbf{s} d\mathbf{s} d\mathbf{s} = i \int_{0}^{1} \int_{0}^{1} \mathbf{S} d\mathbf{s} d\mathbf{s} d\mathbf{s} d\mathbf{s} = i \int_{0}^{1} \int_{0}^{1} \mathbf{S} d\mathbf{s} d\mathbf$$

can be regarded as objects in the quasi-category under (2.2) ([L1, Remark 1.2.9.5.]). In this undercategory, a morphism from \mathbf{S} to \mathbf{S}' is nothing but a pair of 3-simplices φ, ψ in $\mathscr C$ satisfying

$$d_1(\varphi) = d_1(\psi), d_2(\varphi) = \eta', d_2(\psi) = \xi', d_3(\varphi) = \eta, d_3(\psi) = \xi.$$

In this article, we denote this morphism by $\lceil \varphi | \psi \rceil \colon \mathbf{S} \to \mathbf{S}'$. We refer to such a pair simply as a morphism in "the" undercategory, when (2.2) is clear from the context.

A rectangle in this article means a map $\mathbf{R} \colon \Delta^1 \times \Delta^2 \to \mathscr{C}$ which will be denoted as a diagram

$$\mathbf{R}(0,0) \Rightarrow \mathbf{R}(0,1) \Rightarrow \mathbf{R}(0,2)$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$\mathbf{R}(1,0) \Rightarrow \mathbf{R}(1,1) \Rightarrow \mathbf{R}(1,2)$$

often by labelling some of its simplices similarly as for squares. It can be identified with a triplet of 3-simplices $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$ satisfying $d_1(\mathcal{X}) = d_1(\mathcal{Y})$ and $d_2(\mathcal{Y}) = d_2(\mathcal{Z})$, which correspond to

$$\mathcal{X} = \mathbf{R}|_{\{(0,0)\}*(\{1\}\times\Delta^2)}, \quad \mathcal{Y} = \mathbf{R}|_{(\{0\}\times[0,1])*(\{1\}\times[1,2])}, \quad \mathcal{Z} = \mathbf{R}|_{(\{0\}\times\Delta^2)*\{(1,2)\}}.$$

In this article, we express such a rectangle by $\mathbf{R} = \begin{bmatrix} \mathcal{X} & \mathcal{Y} & \mathcal{Z} \end{bmatrix}$. The left, right, and outer squares in \mathbf{R} will be denoted by $\mathbf{R}_{\text{left}} = \mathbf{R}|_{\Delta^1 \times [0,1]}$, $\mathbf{R}_{\text{right}} = \mathbf{R}|_{\Delta^1 \times [1,2]}$ and $\mathbf{R}_{\text{out}} = \mathbf{R}|_{\Delta^1 \times \Lambda_1^2}$, respectively. If \mathbf{R}_{left} is a push-out, then \mathbf{R}_{out} is a push-out if and only if $\mathbf{R}_{\text{right}}$ is a push-out ([L1, Lemma 4.4.2.1]). Dually for pull-backs.

Remark that a square (2.1) can be thought of as a morphism in the quasi-category $\operatorname{Fun}(\Delta^1, \mathscr{C})$ from $A \xrightarrow{x} B$ to $D \xrightarrow{j} C$. In this view, a consecutive pair of squares

can be filled into a 2-simplex in Fun(Δ^1, \mathscr{C}). Slightly more strictly, we may also designate the 2-simplices in \mathscr{C} used as guides for this pasting operation, as follows.

Proposition 2.1. Let

$$\mathbf{S} = \begin{array}{cccc} A & \xrightarrow{x} & B \\ & \downarrow & \downarrow & \\ & \downarrow & \downarrow & \\ & A' & \xrightarrow{x'} & B' \\ & & A'' & \xrightarrow{x''} & B'' \\ & & & & A'' & \xrightarrow{x''} & B'' \\ \end{array}$$

be any pair of squares such that $\mathbf{S}|_{\{1\}\times\Delta^1}=\mathbf{S}'|_{\{0\}\times\Delta^1}=x'$, and let

be arbitrarily taken pair of 2-simplices. Then there exists a rectangle

$$\mathbf{R} = x \begin{vmatrix} a'' \\ a \\ A' & a' \end{vmatrix} A''$$

$$B \Rightarrow B' \Rightarrow B' \Rightarrow B''$$

such that $\mathbf{R}_{\text{left}} = \mathbf{S}^t$, $\mathbf{R}_{\text{right}} = \mathbf{S}'^t$ and $\mathbf{R}|_{\{0\} \times \Delta^2} = \alpha$, $\mathbf{R}|_{\{1\} \times \Delta^2} = \beta$. Moreover, if $\mathbf{S} = (s_0(x') \setminus \xi)$ and $\alpha = s_0(a')$ (hence consequently $A' = A, a = 1_A, z = x', a'' = a'$), then \mathbf{R} can be chosen to be of the form $\mathbf{R} = [\mathcal{X} \mid s_0(\xi') \mid s_0(\eta')]$.

Proof. This is just rephrasing the fact that $(\Delta^1 \times \Lambda_1^2) \cup (\partial \Delta^1 \times \Delta^2) \hookrightarrow \Delta^1 \times \Delta^2$ is an inner anodyne map. ([J]. See also [L1, Proposition 2.3.2.1], [C, Proposition 3.2.3]). We also remark that as for the latter part, we only have to take a 3-simplex

$$\mathcal{X} = B \xrightarrow{\begin{array}{c} x \\ z \\ b \end{array}} B''$$

$$B'$$

such that $d_0(\mathcal{X}) = \beta$, $d_1(\mathcal{X}) = \xi'$, $d_3(\mathcal{X}) = \xi$.

If we are given a rectangle $\mathbf{R} \colon \Delta^1 \times \Delta^2 \to \mathscr{C}$ and a morphism $f \in \mathscr{C}_1(\mathbf{R}((1,2)), C)$ to some $C \in \mathscr{C}_0$, then we can extend them to obtain a map $F \colon (\Delta^1 \times \Delta^2)^{\triangleright} \to \mathscr{C}$. More strictly, we may designate some of the simplices involved, as below. This ad hoc lemma will be used later in the proof of Lemma 3.10.

Lemma 2.2. Let

$$\mathbf{R} = \left[\begin{array}{c|c} \psi & \varphi & \theta \end{array}\right] = x \downarrow x' \downarrow x' \downarrow x'' \downarrow x' \downarrow x$$

be any rectangle, let

$$\lceil \zeta | \zeta' \rceil \colon \mathbf{R}_{\text{out}} \to \mathbf{S} = \begin{cases} A \xrightarrow{a''} A'' \\ x \downarrow \\ B \xrightarrow{a''} C \end{cases}$$

be any morphism in the undercategory, and let

$$\wp = A' \xrightarrow{a'} A''$$

be any 3-simplex in $\mathscr C$ satisfying $d_1(\wp) = d_2(\zeta')$ and $d_3(\wp) = d_3(\theta)$. Then we can extend this to a map $(\Delta^1 \times \Delta^2)^{\triangleright} \to \mathscr C$. Namely, there is a triplet of 4-simplices

$$\Psi = \begin{pmatrix} A & A & A \\ A & A & A \\ B' & B' & B'' \end{pmatrix}, \quad \Phi = \begin{pmatrix} A' & A & A \\ A' & A' & A' \\ B' & B'' \end{pmatrix}, \quad \Theta = \begin{pmatrix} A' & A' & A' \\ A'' & A'' & A'' \\ A'' & A'' & A'' \end{pmatrix}$$

such that $d_1(\Psi) = d_1(\Phi)$, $d_2(\Phi) = d_2(\Theta)$, $\lceil \zeta | \zeta' \rceil = \lceil d_2(\Psi) | d_1(\Theta) \rceil$, $d_3(\Theta) = \wp$, and $\lceil \psi \mid \varphi \mid \theta \rceil = \lceil d_4(\Phi) \mid d_4(\Psi) \mid d_4(\Theta) \rceil$.

Proof. Take a 2-simplex

$$B'$$

$$B'' \longrightarrow C$$

such that $d_0(\varpi) = d_0 d_1(\zeta) = d_0 d_1(\zeta')$ and $d_2(\varpi) = b'$. Take 3-simplices

$$\alpha = A'' \xrightarrow{A'} C, \ \beta = B' \xrightarrow{A} C, \ \gamma = B' \xrightarrow{A'} C, \ \delta = B' \xrightarrow{B''} C$$

such that

$$\begin{split} d_0(\alpha) &= d_0(\zeta'), \ d_2(\alpha) = d_0(\wp), \ d_3(\alpha) = d_0(\theta), \\ d_0(\beta) &= \varpi, \ d_1(\beta) = d_1(\zeta), \ d_3(\beta) = d_1(\varphi) = d_1(\psi), \\ d_0(\gamma) &= \varpi, \ d_1(\gamma) = d_1(\alpha), \ d_3(\gamma) = d_0(\varphi), \\ d_0(\delta) &= \varpi, \ d_1(\delta) = d_0(\zeta), \ d_3(\delta) = d_0(\psi). \end{split}$$

Then we can take Θ, Φ, Ψ so that

$$d_0(\Theta) = \alpha, \ d_1(\Theta) = \zeta', \ d_3(\Theta) = \wp, \ d_4(\Theta) = \theta,$$

$$d_0(\Phi) = \gamma, \ d_1(\Phi) = \beta, \ d_2(\Phi) = d_2(\Theta), \ d_4(\Phi) = \varphi,$$

$$d_0(\Psi) = \delta, \ d_1(\Psi) = \beta, \ d_2(\Psi) = \zeta, \ d_4(\Psi) = \psi$$

hold, as desired.

A cube in \mathscr{C} is a map \mathbb{C} : $\Delta^1 \times \Delta^1 \times \Delta^1 \to \mathscr{C}$, namely an object in Fun($\Delta^1 \times \Delta^1 \times \Delta^1, \mathscr{C}$), which we denote by a diagram

$$\mathbf{C}(0,0,0) \longrightarrow \mathbf{C}(0,1,0)$$

$$\downarrow \mathbf{C}(1,0,0) \longrightarrow \mathbf{C}(1,1,0)$$

$$\mathbf{C}(0,0,1) \longrightarrow \mathbf{C}(0,1,1)$$

$$\mathbf{C}(1,0,1) \longrightarrow \mathbf{C}(1,1,1)$$

in \mathscr{C} or by labelling some of its simplices for example as follows,

similarly as for squares. We may identify it with a pair of rectangles

$$\mathbf{R}_{\mathbf{f}} = A \xrightarrow{x} B \xrightarrow{y} C$$

$$\downarrow a \downarrow b \downarrow b \downarrow c \text{ and } \mathbf{R}_{\mathbf{b}} = A \xrightarrow{i} D \xrightarrow{j} C$$

$$\downarrow a \downarrow d \downarrow d \downarrow c$$

$$\downarrow a \downarrow d \downarrow d \downarrow d$$

$$\downarrow a \downarrow d \downarrow d \downarrow d$$

$$\downarrow a \downarrow$$

sharing the outer square in common. Each of these rectangles is expressed as $\mathbf{R}_f = \left[\begin{array}{c|c} \mathcal{X}_f & \mathcal{Y}_f & \mathcal{Z}_f \end{array} \right]$ and $\mathbf{R}_b = \left[\begin{array}{c|c} \mathcal{X}_b & \mathcal{Y}_b & \mathcal{Z}_b \end{array} \right]$. In summary, a cube \mathbf{C} can be expressed by a 6-tuple of 3-simplices in \mathscr{C}

(2.4)
$$\mathbf{C} = \begin{bmatrix} \mathcal{X}_{f} & \mathcal{Y}_{f} & \mathcal{Z}_{f} \\ \mathcal{X}_{b} & \mathcal{Y}_{b} & \mathcal{Z}_{b} \end{bmatrix}$$

satisfying
$$d_1(\mathcal{X}_f) = d_1(\mathcal{Y}_f), d_2(\mathcal{Y}_f) = d_2(\mathcal{Z}_f), d_2(\mathcal{X}_f) = d_2(\mathcal{X}_b), d_1(\mathcal{Z}_f) = d_1(\mathcal{Z}_b), d_1(\mathcal{X}_b) = d_1(\mathcal{Y}_b), d_2(\mathcal{Y}_b) = d_2(\mathcal{Z}_b).$$

Remark 2.3. From a cube C as above, we may obtain cubes

by transpositions, and also
$$\left[\begin{array}{c|c|c} \mathcal{X}_b & \mathcal{Y}_b & \mathcal{Z}_b \\ \mathcal{X}_f & \mathcal{Y}_f & \mathcal{Z}_f \end{array} \right], \left[\begin{array}{c|c|c} \mathcal{Y}_f & \mathcal{X}_f & \mathcal{X}_b \\ \mathcal{Z}_f & \mathcal{Z}_b & \mathcal{Y}_b \end{array} \right], \left[\begin{array}{c|c|c} \mathcal{Z}_b & \mathcal{Z}_f & \mathcal{Y}_f \\ \mathcal{Y}_b & \mathcal{X}_b & \mathcal{X}_f \end{array} \right].$$

We often regard a cube (2.3) as a morphism from $\mathbf{C}|_{\Delta^1 \times \Delta^1 \times \{0\}}$ to $\mathbf{C}|_{\Delta^1 \times \Delta^1 \times \{1\}}$ in Fun($\Delta^1 \times \Delta^1, \mathscr{C}$). In this view, we have the following.

Example 2.4. Let

$$\mathbf{S} = i \bigvee_{j}^{A} \bigvee_{i}^{x} \bigvee_{j}^{B} V$$

be any square.

(1) There is a cube consisting of degenerated 3-simplices

$$\mathbf{I_S} = \begin{bmatrix} s_0(\xi) & s_1(\xi) & s_2(\xi) \\ s_0(\eta) & s_1(\eta) & s_2(\eta) \end{bmatrix}$$

such that $\mathbf{I}_{\mathbf{S}}|_{\Delta^1 \times \Delta^1 \times \{0\}} = \mathbf{I}_{\mathbf{S}}|_{\Delta^1 \times \Delta^1 \times \{1\}} = \mathbf{S}$. This is nothing but $1_{\mathbf{S}} = s_0(\mathbf{S})$ in $\operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})$.

(2) Suppose that \mathscr{C} is pointed ([L2, Definition 2.1]). Then any square

$$\mathbf{O} = \begin{array}{c|c} O_A \xrightarrow{x_o} O_B \\ \vdots & z_o & y_o \\ O_D \xrightarrow{j_o} O_C \end{array}$$

in which O_A, O_B, O_C, O_D are zero objects in $\mathscr C$, gives a zero object in $\operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr C)$. In particular, there exists a cube $\mathbf C$ such that $\mathbf C|_{\Delta^1 \times \Delta^1 \times \{0\}} = \mathbf S$ and $\mathbf C|_{\Delta^1 \times \Delta^1 \times \{1\}} = \mathbf O$. Indeed, by using the definition of zero objects, such a cube can be taken compatibly with arbitrarily given 5-tuple of squares

$$A \xrightarrow{x} B \qquad A \xrightarrow{i} D \qquad B \xrightarrow{y} C \qquad D \xrightarrow{j} C \qquad A \xrightarrow{z} C$$

$$\downarrow a \downarrow \qquad \downarrow b \qquad , \qquad \downarrow d \qquad , \qquad \downarrow b \downarrow \qquad \downarrow c \qquad , \qquad \downarrow c \qquad , \qquad \downarrow d \qquad , \qquad \downarrow c \qquad \downarrow c$$

$$O_{A \xrightarrow{z}} O_{B} \qquad O_{A \xrightarrow{z}} O_{D} \qquad O_{B \xrightarrow{z}} O_{C} \qquad O_{D \xrightarrow{z}} O_{C} \qquad O_{A \xrightarrow{z}} O_{C}$$

sharing the morphisms with the same labels in common. Similarly for the existence of a cube \mathbf{C}' with $\mathbf{C}'|_{\Delta^1 \times \Delta^1 \times \{0\}} = \mathbf{O}$ and $\mathbf{C}'|_{\Delta^1 \times \Delta^1 \times \{1\}} = \mathbf{S}$.

Similarly as in Proposition 2.1, any consecutive pair of cubes can be filled into a 2-simplex in Fun($\Delta^1 \times \Delta^1, \mathscr{C}$). More strictly, the following holds.

Proposition 2.5. Let

be a pair of cubes satisfying $\mathbf{C}|_{\Delta^1 \times \Delta^1 \times \{1\}} = \mathbf{C}'|_{\Delta^1 \times \Delta^1 \times \{0\}}$, and let

be arbitrarily taken 2-simplices. Then there exists $F: \Delta^1 \times \Delta^1 \times \Delta^2 \to \mathscr{C}$ such that $F|_{\Delta^1 \times \Delta^1 \times [0,1]} = \mathbf{C}, F|_{\Delta^1 \times \Delta^1 \times [1,2]} = \mathbf{C}'$ and

$$F|_{\{(0,0)\}\times\Delta^2} = \alpha, F|_{\{(0,1)\}\times\Delta^2} = \beta, F|_{\{(1,1)\}\times\Delta^2} = \gamma, F|_{\{(1,0)\}\times\Delta^2} = \delta.$$

Especially we obtain a cube

$$F|_{\Delta^{1}\times\Delta^{1}\times\Lambda_{1}^{2}} = \begin{array}{c} A \xrightarrow{x} B & y \\ \downarrow & D \xrightarrow{j} C \\ A'' \xrightarrow{x''} & B'' & y'' \\ \downarrow & B'' & y'' \\ \downarrow & B'' & y'' \end{array}$$

compatible with the given data.

Proof. Similarly as Proposition 2.1, this is just rephrasing the fact that $((\Delta^1 \times \Delta^1) \times \Lambda_1^2) \cup (\{(0,0),(0,1),(1,0),(1,1)\} \times \Delta^2) \hookrightarrow \Delta^1 \times \Delta^1 \times \Delta^2$ is an inner anodyne map ([J], [L1, Proposition 2.3.2.1], [C, Proposition 3.2.3]).

Corollary 2.6. Let S, S' be any pair of squares such that S((0,0)) = S'((0,0)) = A and S((1,1)) = S'((1,1)) = C. Assume that there is a cube C such that

$$\mathbf{C}|_{\Lambda^1 \times \Lambda^1 \times \{0\}} = \mathbf{S}, \ \mathbf{C}|_{\Lambda^1 \times \Lambda^1 \times \{1\}} = \mathbf{S}'$$

and that $a = \mathbf{C}|_{\{(0,0)\}\times\Delta^1}$ and $c = \mathbf{C}|_{\{(1,1)\}\times\Delta^1}$ satisfies $\overline{a} = \mathrm{id}_A$ and $\overline{c} = \mathrm{id}_C$. Then there exists a cube \mathbf{C}' such that

$$\mathbf{C}'|_{\Delta^1 \times \Delta^1 \times \{0\}} = \mathbf{S}, \ \mathbf{C}'|_{\Delta^1 \times \Delta^1 \times \{1\}} = \mathbf{S}'$$

and $\mathbf{C}'|_{\{(0,0)\}\times\Delta^1}=1_A$, $\mathbf{C}'|_{\{(0,1)\}\times\Delta^1}=\mathbf{C}|_{\{(0,1)\}\times\Delta^1}$, $\mathbf{C}'|_{\{(1,0)\}\times\Delta^1}=\mathbf{C}|_{\{(1,0)\}\times\Delta^1}$, $\mathbf{C}'|_{\{(1,1)\}\times\Delta^1}=1_C$.

Proof. This follows from Proposition 2.5 applied to the cubes C, $I_{S'}$ and 2-simplices of the following form,

where we denote $\mathbf{C}|_{\{(0,1)\}\times\Delta^1}$ and $\mathbf{C}|_{\{(1,0)\}\times\Delta^1}$ by $B\stackrel{b}{\longrightarrow} B'$ and $D\stackrel{d}{\longrightarrow} D'$, respectively.

3. Exact sequences in exact quasi-categories

First we recall the definition of an exact quasi-category. For the detail, see [B1],[B2]. A quasi-category $\mathscr C$ is called *additive* ([B1, Definition 2.2]) if it is pointed, has all finite products and finite coproducts, and moreover if the homotopy category $h\mathscr C$ is additive as an ordinary category. In particular, for any pair of objects $X_1, X_2 \in h\mathscr C$, the unique morphism $\mathfrak v \in (h\mathscr C)(X_1 \amalg X_2, X_1 \times X_2)$ which satisfies

$$p_{k'} \circ \mathfrak{v} \circ i_k \colon X_k \to X_{k'} = \begin{cases} \mathrm{id} & \mathrm{if } k = k' \\ 0 & \mathrm{if } k \neq k' \end{cases}$$

should become an isomorphism. Here, $X_1 \stackrel{p_1}{\longleftarrow} X_1 \times X_2 \stackrel{p_2}{\longrightarrow} X_2$ and $X_1 \stackrel{i_1}{\longrightarrow} X_1 \coprod X_2 \stackrel{i_2}{\longleftarrow} X_2$ are product and coproduct of X_1, X_2 in $h\mathscr{C}$, respectively.

Let \mathscr{C} be a quasi-category, and let $\mathscr{C}_{\dagger}, \mathscr{C}^{\dagger}$ be two subcategories of \mathscr{C} containing all homotopy equivalences. A morphism in \mathscr{C}_{\dagger} is called *ingressive*, and a morphism in \mathscr{C}^{\dagger} is called *egressive*. A square

$$\begin{array}{ccc}
A & \xrightarrow{x} & B \\
\downarrow & & \downarrow y \\
D & \xrightarrow{j} & C
\end{array}$$

is called an *ambigressive pull-back* if it is a pull-back square in which j is ingressive and y is egressive. Dually, (3.1) is an *ambigressive push-out* if it is a push-out square in which x is ingressive and i is egressive.

Definition 3.1. ([B1, Definition 3.1]) The triplet $(\mathscr{C}, \mathscr{C}_{\dagger}, \mathscr{C}^{\dagger})$ is an exact quasicategory (= exact ∞ -category) if it satisfies the following conditions.

- (Ex0) \mathscr{C} is additive.
- (Ex1) If $O \in \mathscr{C}_0$ is a zero object, then any morphism $O \to X$ in \mathscr{C} is ingressive. Dually, any morphism $X \to O$ in \mathscr{C} is egressive.
- (Ex2) Push-outs of ingressive morphisms exist and are ingressive. Dually, pull-backs of egressive morphisms exist and are egressive.
- (Ex3) A square in $\mathscr C$ is an ambigressive pull-back if and only if it is an ambigressive push-out.

In the rest of this article, let $(\mathscr{C}, \mathscr{C}_{\dagger}, \mathscr{C}^{\dagger})$ be an exact quasi-category defined as above. We also simply say that \mathscr{C} is an exact quasi-category.

Definition 3.2. ([B1, Definition 3.1])

(1) An exact sequence starting from A and ending in C is an ambigressive pull-back (hence an ambigressive push-out)

(3.2)
$$\mathbf{S} = i \bigvee_{j}^{A} \bigvee_{i}^{x} B \bigvee_{j}^{y} C$$

in which O is a zero object in \mathscr{C} . When we emphasize the end-objects A and C, we write ${}_{A}\mathbf{S}_{C}$ in this article.

Example 3.3. If a morphism $x \in \mathscr{C}_1(X, X)$ satisfies $\overline{x} = \mathrm{id}_X$, then any square

in which O, O' are zero objects, is an exact sequence.

3.1. Morphisms of exact sequences.

Definition 3.4. Let $\mathscr{E} \subseteq \operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})$ be the full subcategory spanned by exact sequences. In particular, for a pair of exact sequences

(3.3)
$$\mathbf{S} = i \bigvee_{j}^{A} \sum_{\xi}^{x} \bigvee_{y} \text{ and } \mathbf{S}' = i \bigvee_{j}^{x'} \sum_{\xi'}^{x'} \bigvee_{y'}^{y'},$$

a morphism of exact sequences from S to S' means a cube

(3.4)
$$\mathbf{C} = \begin{bmatrix} A & \xrightarrow{x} & B & y \\ & & O & \downarrow & C \\ & & & \downarrow & \downarrow & C \\ & & & & \downarrow & \downarrow & C \\ & & & & & \downarrow & C' \\ & & & & & & \downarrow & C' \end{bmatrix}$$

which satisfies $\mathbf{C}|_{\Delta^1 \times \Delta^1 \times \{0\}} = \mathbf{S}$ and $\mathbf{C}|_{\Delta^1 \times \Delta^1 \times \{1\}} = \mathbf{S}'$. We abbreviate this to $\mathbf{C} \colon \mathbf{S} \to \mathbf{S}'$. When we indicate morphisms a and c, we will also write ${}_a\mathbf{C}_c \colon \mathbf{S} \to \mathbf{S}'$ in this article.

Proposition 3.5. Let ${}_{A}\mathbf{S}_{C}$, ${}_{A}\mathbf{S}'_{C}$, be any pair of exact sequences starting from A and ending in C, and let ${}_{a}\mathbf{C}_{c} \colon \mathbf{S} \to \mathbf{S}'$ be a morphism such that $\overline{a} = \mathrm{id}_{A}$ and $\overline{c} = \mathrm{id}_{C}$, as depicted in (3.5) below. Suppose that there also exists a morphism in the opposite direction ${}_{a'}\mathbf{C}'_{c'} \colon \mathbf{S}' \to \mathbf{S}$ as below, such that $\overline{a'} = \mathrm{id}_{A}$ and $\overline{c'} = \mathrm{id}_{C}$.

Then b is a homotopy equivalence.

(3.5)
$$\mathbf{C} = \begin{bmatrix} A & \xrightarrow{x} & B & y \\ & & & & \\ & & & \\ A & \xrightarrow{x'} & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\$$

Proof. It suffices to show that \bar{b} is an isomorphism in $h\mathscr{C}$. Remark that \mathbf{C} and \mathbf{C}' yield commutative diagrams

$$\begin{array}{cccc}
A & \overline{x} & B & \overline{y} & C & A & \overline{x'} & B' & \overline{y'} & C \\
\parallel & \Diamond & \downarrow_{\overline{b}} & \Diamond & \parallel & , & \parallel & \Diamond & \downarrow_{\overline{b'}} & \Diamond & \parallel \\
A & \longrightarrow & B' & \longrightarrow & C & A & \longrightarrow & B & \longrightarrow & C
\end{array}$$

in $h\mathscr{C}$, whose rows are both weak kernel and weak cokernel. For $\mathfrak{b} = \overline{b'} \circ \overline{b} \in (h\mathscr{C})(B,B)$, since we have $(\mathrm{id}_B - \mathfrak{b}) \circ \overline{x} = 0$ and $\overline{y} \circ (\mathrm{id}_B - \mathfrak{b}) = 0$, there are $\mathfrak{d} \in (h\mathscr{C})(B,A)$, $\mathfrak{e} \in (h\mathscr{C})(C,B)$ such that $\overline{x} \circ \mathfrak{d} = \mathrm{id}_B - \mathfrak{b}$ and $\mathfrak{e} \circ \overline{y} = \mathrm{id}_B - \mathfrak{b}$. It follows that $(\mathrm{id}_B - \mathfrak{b}) \circ (\mathrm{id}_B - \mathfrak{b}) = \mathfrak{e} \circ \overline{y} \circ \overline{x} \circ \mathfrak{d} = 0$, equivalently $\mathfrak{b} \circ (2 \cdot \mathrm{id}_B - \mathfrak{b}) = (2 \cdot \mathrm{id}_B - \mathfrak{b}) \circ \mathfrak{b} = \mathrm{id}_B$. In particular $\mathfrak{b} = \overline{b'} \circ \overline{b}$ is an isomorphism. The same argument shows that $\overline{b} \circ \overline{b'}$ is also an isomorphism. Hence so is \overline{b} .

Remark 3.6. Later in Proposition 3.26 we will see (in combination with Corollary 2.6) that the existence of \mathbf{C}' is not necessary to assume, and that it always holds automatically.

Proposition 3.7. Let $\mathbf{S} = (\eta \setminus \xi), \mathbf{S}' = (\eta' \setminus \xi')$ be any pair of exact sequences as in (3.3), and let

(3.6)
$$\mathbf{T} = \begin{bmatrix} A & \xrightarrow{x} & B \\ \downarrow b \\ A' & \xrightarrow{x'} & B' \end{bmatrix}$$

be any square. Then there exists a morphism of exact sequences $\mathbf{C} \colon \mathbf{S} \to \mathbf{S}'$ such that $\mathbf{C}|_{\Delta^1 \times \{0\} \times \Delta^1} = \mathbf{T}$. More precisely, the following holds.

(1) There exist rectangles of the form

$$\mathbf{R}' = \begin{bmatrix} \mathcal{X}' \mid \mathcal{Y}' \mid \mathcal{Z}' \end{bmatrix} = \begin{bmatrix} A & \xrightarrow{a} & A' & \xrightarrow{i'} & O' \\ x & & & & \downarrow j' \\ B & \xrightarrow{b} & B' & \xrightarrow{y'} & C' \end{bmatrix}$$

sharing some common morphism $i_p \in \mathcal{C}_1(A, O')$, such that $\mathbf{S} = (\mathbf{R}_{left})^t$, $\mathbf{S}' = (\mathbf{R}'_{right})^t$ and $\mathbf{T} = (\mathbf{R}'_{left})^t$.


Moreover, if A = A', $a = 1_A$ and $\mathbf{T} = (s_0(x') \setminus \xi)$, then we may take $i_p = i'$, and \mathbf{R}' can be chosen to be of the form $\begin{bmatrix} \mathcal{X}' \mid s_0(\xi') \mid s_0(\eta') \end{bmatrix}$.

(2) For any $\mathbf{R} = [s_2(\xi) \mid s_2(\eta) \mid \mathcal{Z}]$ and $\mathbf{R}' = [\mathcal{X}' \mid \mathcal{Y}' \mid \mathcal{Z}']$ as in (1), there exist three 3-simplices Φ, Ψ_f, Ψ_b which gives a morphism


$$\mathbf{C} = \left[egin{array}{c|c} \mathcal{Y}' & \mathcal{X}' & \Psi_{\mathrm{f}} \ \mathcal{Z}' & \Phi & \Psi_{\mathrm{b}} \end{array}
ight] : \mathbf{S}
ightarrow \mathbf{S}'$$

satisfying $d_3(\Phi) = d_3(\mathcal{Z})$.

Proof. (1) By using the definition of a zero object, we may take a 3-simplex



with arbitrarily chosen morphisms $A \xrightarrow{i_p} O'$, $O \to O'$, $O' \to C$, such that $d_2(\mathcal{Z}) = \eta$. This gives **R** as desired. By the definition of a zero object, there also exists a 2-simplex α as below. Also, we may take a 2-simplex β arbitrarily as below, with some $y_p \in \mathscr{C}_1(B, C')$.



Then the existence of \mathbf{R}' follows from Proposition 2.1 applied to $\mathbf{T}, \mathbf{S}', \alpha, \beta$. The latter part is also obvious from the construction, in which case we use $\alpha = s_0(i')$.


(2) We may take a 3-simplex

$$\Phi = O \xrightarrow{i \qquad i_p} C'$$

such that $d_1(\Phi) = d_1(\mathcal{Z}')$ and $d_3(\Phi) = d_3(\mathcal{Z})$, with arbitrarily taken compatible $d_0(\Phi)$ by using the definition of a zero object. Put $\mathbf{S}^p = (\mathbf{R}_{\text{out}})^t = (d_1(\mathcal{Z}) \setminus \xi)$ and

$$\mathbf{S}^{q} = (d_{1}(\mathcal{Z}') \backslash d_{2}(\mathcal{X}')) = \begin{cases} A \xrightarrow{x} B \\ i_{p} \downarrow \\ O' \xrightarrow{j'} C' \end{cases}.$$

Since $(\mathbf{R}_{\mathrm{left}})^t$, $(\mathbf{R}_{\mathrm{right}})^t$ are exact sequences, so is \mathbf{S}^p . Thus by the definition of push-out, there exists a morphism $\lceil \Gamma | \Psi_{\mathrm{f}} \rceil$ from \mathbf{S}^p to \mathbf{S}^q in the undercategory, namely a pair of 3-simplices Ψ_{f} , Γ in $\mathscr E$ satisfying $d_2(\Psi_{\mathrm{f}}) = d_2(\mathcal X')$, $d_3(\Psi_{\mathrm{f}}) = \xi$, $d_2(\Gamma) = d_1(\mathcal Z')$, $d_3(\Gamma) = d_1(\mathcal Z)$, and $d_1(\Gamma) = d_1(\Psi_{\mathrm{f}})$. We may take a 3-simplex ν such that $d_0(\nu) = d_0(\Gamma)$, $d_2(\nu) = d_0(\Phi)$, $d_3(\nu) = d_0(\mathcal Z)$ and then a 4-simplex



such that $d_0(\Theta) = \nu$, $d_1(\Theta) = \Gamma$, $d_3(\Theta) = \Phi$, $d_4(\Theta) = \mathcal{Z}$. If we put $\Psi_b = d_2(\Theta)$, then Φ , Ψ_f , Ψ_b satisfy the required conditions.

The following will be used later to replace the zero objects appearing in exact sequences.

Lemma 3.8. Let

$$\nu = O \xrightarrow{A} C \quad \text{and} \quad \nu' = O \xrightarrow{A} C$$

be any pair of 3-simplices satisfying $d_2(\nu)=d_2(\nu')$, in which O,O',O'' are zero objects. Let $o\in \mathscr{C}_1(O'',O')$ be arbitrarily taken morphism. Then there exists a 3-simplex

$$\mu = O'' \xrightarrow{A} C$$

such that $d_1(\mu) = d_1(\nu)$ and $d_2(\mu) = d_1(\nu')$.

Proof. We can show the existence of a 4-simplex

$$\Phi = O \xrightarrow{A} C$$

$$O'' \xrightarrow{o} O'$$

such that $d_2(\Phi) = \nu$ and $d_3(\Phi) = \nu'$, with arbitrarily chosen compatible 3-simplices $d_0(\Phi)$ and $d_4(\Phi)$ by using the definition of zero objects. This gives $\mu = d_1(\Phi)$ as desired.

Proposition 3.9. Let

$${}_{A}\mathbf{S}_{C} = (\eta \backslash \xi) = i \bigvee_{j}^{A} \bigvee_{z \in \mathbb{Z}}^{\xi} \bigvee_{y} \quad and \quad {}_{A}\mathbf{S}'_{C} = (\eta' \backslash \xi) = i \bigvee_{j'}^{A} \bigvee_{z \in \mathbb{Z}}^{z} \bigvee_{y'}^{z} \bigvee_{j'}^{z} C$$

be any pair of squares sharing common ξ in the upper triangles. Suppose that there is a 3-simplex ν such that $d_1(\nu) = \eta'$ and $d_2(\nu) = \eta$. Then the following holds.

- (1) There exists a 3-simplex ν' such that $d_1(\nu') = \eta$ and $d_2(\nu') = \eta'$.
- (2) S is an exact sequence if and only if S' is an exact sequence.

We also remark that in case (2), the given ν and ν' obtained in (1) are nothing but morphisms of exact sequences

$$\begin{bmatrix} s_0(\xi) & s_1(\xi) & s_2(\xi) \\ s_0(\eta') & \nu & s_2(\eta) \end{bmatrix} : \mathbf{S} \to \mathbf{S}' \quad and \quad \begin{bmatrix} s_0(\xi) & s_1(\xi) & s_2(\xi) \\ s_0(\eta) & \nu' & s_2(\eta') \end{bmatrix} : \mathbf{S}' \to \mathbf{S}.$$

Proof. (1) is immediate from Lemma 3.8, applied to ν and $s_1(\eta)$. (2) follows from the existence of a rectangle $\mathbf{R} = \begin{bmatrix} \nu \mid s_0(\eta') \mid s_0(\xi) \end{bmatrix}$, in which \mathbf{R}_{left} is an exact sequence and $\mathbf{R}_{\text{out}} = \mathbf{S}$, $\mathbf{R}_{\text{right}} = \mathbf{S}'$.

3.2. Decompositions of morphisms.

Lemma 3.10. Let ${}_{A}\mathbf{S}_{C}=(\eta\backslash\xi)$ be any exact sequence as in (3.2), and let $a\in\mathscr{C}_{1}(A,A')$ be any morphism. Let

(3.7)
$$\mathbf{P} = \begin{bmatrix} A & \xrightarrow{x} B \\ \downarrow b_0 \\ A' & \xrightarrow{x_0} B_0 \end{bmatrix}$$

be arbitrarily chosen push-out square, which always exists by the definition of exact quasi-category. Then, we have the following.

(1) There exists an exact sequence

(3.8)
$$\mathbf{S}_{0} = i_{0} \begin{vmatrix} A' & x_{0} \\ & & \downarrow \\ & z_{0} \\ & & \downarrow \\ & O & y_{0} \end{vmatrix} y_{0}$$

and a morphisms of exact sequences $\mathbf{D} \colon \mathbf{S} \to \mathbf{S}_0$ of the form

$$\mathbf{D} = \begin{bmatrix} \varphi & \psi & s_2(\xi) \\ \theta & s_1(\eta) & s_2(\eta) \end{bmatrix} = \begin{bmatrix} A & \xrightarrow{x} & B & y \\ \downarrow & O & \downarrow & C \\ \downarrow & \downarrow & \downarrow & \downarrow \\ A' & \xrightarrow{x_0} & \downarrow & B_0 & y_0 \\ \downarrow & \downarrow & \downarrow & C \end{bmatrix}$$

such that $\mathbf{D}|_{\Delta^1 \times \{0\} \times \Delta^1} = \mathbf{P}$. We denote any such square \mathbf{S}_0 admitting \mathbf{D} by $a_*\mathbf{S}$.

(2) Let $\mathbf{D} \colon \mathbf{S} \to \mathbf{S}_0 = a_* \mathbf{S}$ be arbitrarily as in (1). For any morphism of exact sequences

$${}_{a}\mathbf{C}_{c} = \left[\begin{array}{c|c} \mathcal{X}_{\mathrm{f}} & \mathcal{Y}_{\mathrm{f}} & \mathcal{Z}_{\mathrm{f}} \\ \mathcal{X}_{\mathrm{b}} & \mathcal{Y}_{\mathrm{b}} & \mathcal{Z}_{\mathrm{b}} \end{array} \right] : {}_{A}\mathbf{S}_{C} \rightarrow {}_{A'}\mathbf{S}_{C'}' = (\eta' \backslash \xi')$$

as in (3.4), there exists a morphism of exact sequences $a_{A'} \mathbf{C}'_c \colon a_* \mathbf{S} \to \mathbf{S}'$ of the form

for some $\mathcal{Y}_{\mathrm{f}}', \mathcal{Y}_{\mathrm{b}}', \mathcal{Z}_{\mathrm{f}}', \mathcal{Z}_{\mathrm{b}}'$, such that $\overline{b} = \overline{b'} \circ \overline{b}_0$ holds in $h\mathscr{C}$.

Dually, for any ${}_{A}\mathbf{S}_{C}=(\eta\backslash\xi)$ as above and any $c\in\mathscr{C}(C',C)$, we may obtain a morphism of exact sequences of the form

$$\mathbf{1}_{A}\mathbf{D}_{c}' = \left[\begin{array}{c|c} s_{0}(\xi) & s_{1}(\xi) \\ s_{0}(\eta) & \mathcal{Y}_{\mathrm{b}} \end{array} \right] \mathcal{Z}_{\mathrm{f}} \\ \mathcal{Z}_{\mathrm{b}} \end{array} \right] : \mathbf{S}_{0}' \to \mathbf{S}$$

in which $\mathbf{D}'|_{\Delta^1 \times \{1\} \times \Delta^1}$ is a pull-back. In this case, we denote such \mathbf{S}'_0 by $c^*\mathbf{S}$. It has a property dual to (2).

Proof. (1) By using the definition of a zero object, we can take a 3-simplex

$$\theta = A \xrightarrow{i} C$$


$$A' \xrightarrow{i_0} C$$

such that $d_1(\theta) = \eta$, with some morphisms $i_0 \in \mathcal{C}_1(A', O)$ and $z_0 \in \mathcal{C}_1(A', C)$. Put $\nu = d_2(\theta)$. Since **P** is a push-out, we obtain a morphism

$$[\varphi|\psi] \colon \mathbf{P} \to \begin{bmatrix} A & \xrightarrow{x} & B \\ & \downarrow & \downarrow \\ A' & \xrightarrow{z_0} & C \end{bmatrix}$$

in the undercategory. Then this gives a rectangle $\mathbf{R} = \begin{bmatrix} \psi & \varphi & \theta \end{bmatrix}$ with $\mathbf{R}_{\text{left}} = \mathbf{P}^t$, $\mathbf{R}_{\text{out}} = \mathbf{S}^t$ and $\mathbf{R}_{\text{right}} = (\mathbf{S}_0)^t$. Since \mathbf{P} , \mathbf{S}^t are push-outs, it follows that x_0 is ingressive and that $\mathbf{S}_0 = (\mathbf{R}_{\text{right}})^t$ is also a push-out. Thus \mathbf{S}_0 is an exact sequence, and $\mathbf{D} = \begin{bmatrix} \varphi & \psi & s_2(\xi) \\ \theta & s_1(\eta) & s_2(\eta) \end{bmatrix} : \mathbf{S} \to \mathbf{S}_0$ indeed gives a morphism with the stated property.

(2) We may take a 4-simplex



such that $d_1(\Omega) = \mathcal{Y}_b$ and $d_2(\Omega) = \mathcal{X}_b$, for arbitrarily chosen compatible $d_0(\Omega)$ and $d_4(\Omega)$ satisfying $d_3d_4(\Omega) = d_3(\theta)$ using the definition of zero object. Applying Lemma 2.2 to

$$\mathbf{R} = \left[\begin{array}{c|c} \psi & \varphi & \theta \end{array} \right], \quad \left[\mathcal{Z}_f | \mathcal{Z}_b \right] \colon \mathbf{R}_{out} \to x \\ & \downarrow \\ B \longrightarrow C' \end{array}, \quad \wp = d_3(\Omega),$$

we obtain 4-simplices Ψ, Φ, Θ such that $d_1(\Psi) = d_1(\Phi), d_2(\Phi) = d_2(\Theta), \lceil \mathcal{Z}_f | \mathcal{Z}_b \rceil = \lceil d_2(\Psi) | d_1(\Theta) \rceil, d_3(\Theta) = \wp$, and $\lceil \psi \mid \varphi \mid \theta \rceil = \lceil d_4(\Phi) \mid d_4(\Psi) \mid d_4(\Theta) \rceil$. If we put

$$\mathbf{T} = \mathbf{C}|_{\{0\} \times \Delta^1 \times \Delta^1} = (d_3(\mathcal{X}_f) \setminus d_3(\mathcal{Y}_f)),$$

$$\mathbf{T}' = (d_2(\mathcal{X}_h) \setminus d_2(\mathcal{Z}_f)) = (d_2(\mathcal{X}_f) \setminus d_2(\mathcal{Y}_f)),$$

Then T, T' and P can be regarded as objects in the quasi-category under

$$A \xrightarrow{x} B$$

$$\downarrow a \qquad \qquad A'$$

and there is a morphism $\lceil \mathcal{X}_f | \mathcal{Y}_f \rceil \colon \mathbf{T} \to \mathbf{T}'$. By the construction so far, we also have $\lceil d_3(\Phi) | d_3(\Psi) \rceil \colon \mathbf{P} \to \mathbf{T}'$ in this undercategory. Since \mathbf{P} is a push-out, we obtain another morphism $\lceil \kappa | \lambda \rceil \colon \mathbf{P} \to \mathbf{T}$. Then, again by the definition of push-out, we obtain a 2-simplex in this undercategory, namely a pair of 4-simplices in \mathscr{C}

$$\Gamma = A' \xrightarrow{a} A C'$$

$$B_0 \xrightarrow{B_0} B'$$

$$B_0 \xrightarrow{B'} B'$$

such that

$$d_1(\Gamma) = d_1(\Xi), d_2(\Gamma) = \mathcal{X}_f, d_2(\Xi) = \mathcal{Y}_f,$$

$$d_3(\Gamma) = d_3(\Phi), d_3(\Xi) = d_3(\Psi), d_4(\Gamma) = \kappa, d_4(\Xi) = \lambda.$$

Now

$$\mathbf{C}' = \left[\begin{array}{c|c} s_0(\xi') & d_0(\Gamma) & d_0(\Phi) \\ s_0(\eta') & d_0(\Omega) & d_0(\Theta) \end{array} \right] : \mathbf{S}_0 \to \mathbf{S}'$$

gives a morphism of exact sequences with the stated properties.

Lemma 3.11. Let ${}_{A}\mathbf{S}_{C}, {}_{A'}\mathbf{S}'_{C'}$ be two exact sequences as in (3.3), and let

$${}_{a}\mathbf{C}_{c} = \left[\begin{array}{c|c} \mathcal{X}_{\mathrm{f}} & \mathcal{Y}_{\mathrm{f}} & \mathcal{Z}_{\mathrm{f}} \\ \mathcal{X}_{\mathrm{b}} & \mathcal{Y}_{\mathrm{b}} & \mathcal{Z}_{\mathrm{b}} \end{array} \right] : \mathbf{S} \rightarrow \mathbf{S}'$$

be a morphism as in (3.4). If C = C' and $\overline{c} = \mathrm{id}_C$, then the following holds.

(1) The square

(3.10)
$$\mathbf{C}|_{\{0\}\times\Delta^{1}\times\Delta^{1}} = \begin{array}{c} A \xrightarrow{x} B \\ \downarrow b \\ A' \xrightarrow{x'} B' \end{array}$$

is a pull-back.

(2) If moreover b is egressive, then (3.10) is an ambigressive push-out.

Proof. (1) \mathbf{C} contains rectangles

$$\mathbf{R} = \left[\begin{array}{c|c} \mathcal{Y}_{f} & \mathcal{X}_{f} & \mathcal{X}_{b} \end{array} \right] = \begin{array}{c} A \xrightarrow{a} A' \xrightarrow{i'} O' \\ x \downarrow & x' \downarrow & \downarrow j' \\ B \xrightarrow{b} B' \xrightarrow{u'} C \end{array}$$

$$\mathbf{R}' = \begin{bmatrix} \mathcal{Z}_{\mathbf{f}} \mid \mathcal{Z}_{\mathbf{b}} \mid \mathcal{Y}_{\mathbf{b}} \end{bmatrix} = \begin{array}{c} A \xrightarrow{i} O \xrightarrow{o} O' \\ x \downarrow & j \downarrow \\ B \xrightarrow{y} C \xrightarrow{c} C \end{array}$$

which satisfies $(\mathbf{R}_{\text{right}})^t = \mathbf{S}'$, $\mathbf{R}_{\text{out}} = \mathbf{R}'_{\text{out}}$ and $(\mathbf{R}'_{\text{left}})^t = \mathbf{S}$. Since $(\mathbf{R}_{\text{right}})^t$ is an exact sequence as in Example 3.3, it follows that $\mathbf{C}|_{\{0\}\times\Delta^1\times\Delta^1}=(\mathbf{R}_{\text{left}})^t$ is a pull-back. (2) is immediate from the definition of exact quasi-category.

Proposition 3.12. Let

$$\mathbf{S} = (\eta \backslash \xi) = i \bigvee_{j}^{A} \bigvee_{i}^{\xi} \bigvee_{j}^{\xi} y \quad and \quad \mathbf{S}' = (\eta' \backslash \xi') = i \bigvee_{j}^{A} \bigvee_{i}^{\xi'} \bigvee_{j}^{\xi'} \bigvee_{j}^{\eta'} C$$

be two exact sequences starting from A and ending in C. Let

$$(3.11) 1_{A}\mathbf{C}_{1_{C}} = \begin{bmatrix} \mathcal{X}_{f} & \mathcal{Y}_{f} & \mathcal{Z}_{f} \\ \mathcal{X}_{b} & \mathcal{Y}_{b} & \mathcal{Z}_{b} \end{bmatrix} = \begin{bmatrix} A & \xrightarrow{x} & B & y \\ & & \downarrow & & \downarrow & \\ & A & \xrightarrow{x'} & \downarrow & B' & y' \\ & & & \downarrow & C \end{bmatrix} : \mathbf{S} \to \mathbf{S}'$$

be any morphism.

If moreover $b = \mathbf{C}|_{\{(0,1)\} \times \Delta^1}$ is a homotopy equivalence in \mathscr{C} , then there also exists a morphism in the opposite direction $_{1_A}(\mathbf{C}')_{1_C} \colon \mathbf{S}' \to \mathbf{S}$. In addition, $b' = \mathbf{C}'|_{\{(0,1)\} \times \Delta^1}$ becomes a homotopy equivalence in \mathscr{C} .

Proof. Put $\xi^{(1)} = d_2(\mathcal{Y}_f), \eta^{(1)} = d_2(\mathcal{Y}_b), \eta^{(2)} = d_1(\mathcal{X}_b), \text{ and put}$

$$\mathbf{S}^{(1)} = (\eta^{(1)} \backslash \xi^{(1)}) = i \bigvee_{O \longrightarrow C} A \xrightarrow{x} B \\ O \xrightarrow{x} C \qquad \mathbf{S}^{(2)} = (\eta^{(2)} \backslash \xi^{(1)}) = \bigvee_{O' \xrightarrow{i'} C} A \xrightarrow{x} B \\ O' \xrightarrow{i'} C$$

In the following rectangle, since $(\mathbf{R}_{left})^t = \mathbf{S}$ and $(\mathbf{R}_{right})^t$ are exact sequences,

$$\mathbf{R} = \begin{bmatrix} \mathcal{Z}_{f} \mid \mathcal{Z}_{b} \mid s_{1}(\eta^{(1)}) \end{bmatrix} = \begin{cases} A \xrightarrow{i} O \xrightarrow{1_{O}} O \\ \downarrow \downarrow \downarrow \downarrow \\ B \xrightarrow{y} C \xrightarrow{1_{C}} C \end{cases}$$

so is $\mathbf{S}^{(1)} = (\mathbf{R}_{\text{out}})^t$. Moreover,

$$\mathbf{1}_{A}\mathbf{C}_{1_{C}}^{(1)} = \begin{bmatrix} s_{0}(\xi^{(1)}) & s_{1}(\xi^{(1)}) & \mathcal{Z}_{\mathrm{f}} \\ s_{0}(\eta^{(1)}) & s_{1}(\eta^{(1)}) & \mathcal{Z}_{\mathrm{b}} \end{bmatrix} : {}_{A}\mathbf{S}_{C} \to {}_{A}\mathbf{S}_{C}^{(1)}$$

is a morphism of exact sequences. By Proposition 3.9, the existence of the 3-simplex \mathcal{Y}_b shows that $\mathbf{S}^{(2)}$ is also an exact sequence, and that

$${}_{1_{A}}\mathbf{C}_{1_{C}}^{(2)} = \left[\begin{array}{c|c} s_{0}(\xi^{(1)}) & s_{1}(\xi^{(1)}) \\ s_{0}(\eta^{(2)}) & \mathcal{Y}_{b} & s_{2}(\eta^{(1)}) \end{array} \right] : {}_{A}\mathbf{S}_{C}^{(1)} \to {}_{A}\mathbf{S}_{C}^{(2)}$$

is a morphisms of exact sequences. Besides, there is another morphism

$$\mathbf{1}_{A}\mathbf{C}_{1_{C}}^{(3)} = \left[egin{array}{c|c} \mathcal{X}_{\mathrm{f}} & \mathcal{Y}_{\mathrm{f}} & s_{2}(\xi^{(1)}) \ s_{1}(\eta^{(2)}) & s_{2}(\eta^{(2)}) \end{array}
ight] : {}_{A}\mathbf{S}_{C}^{(2)}
ightarrow {}_{A}\mathbf{S}_{C}^{\prime},$$

hence we have a sequence of morphisms $\mathbf{S} \xrightarrow{\mathbf{C}^{(1)}} \mathbf{S}^{(1)} \xrightarrow{\mathbf{C}^{(2)}} \mathbf{S}^{(2)} \xrightarrow{\mathbf{C}^{(3)}} \mathbf{S}'$.

By Lemma 3.11, it follows that $\mathbf{C}^{(3)}|_{\{0\}\times\Delta^1\times\Delta^1}$ is an ambigressive push-out. Thus $\mathbf{C}^{(3)}$ satisfies the conditions required in Lemma 3.10 (1). Thus by (2) of the same lemma applied to $\mathbf{C}^{(3)}$ and $\mathbf{I}_{\mathbf{S}^{(2)}}\colon \mathbf{S}^{(2)}\to \mathbf{S}^{(2)}$, we obtain a morphism $\mathbf{I}_A(\mathbf{C}^{(3)\prime})_{\mathbf{I}_C}\colon \mathbf{S}'\to \mathbf{S}^{(2)}$. Similarly, by using the duals of Lemmas 3.10 and 3.11, we obtain a morphism $\mathbf{I}_A(\mathbf{C}^{(1)\prime})_{\mathbf{I}_C}\colon \mathbf{S}^{(1)}\to \mathbf{S}$. By Proposition 3.9, we also have a morphism $\mathbf{I}_A(\mathbf{C}^{(2)\prime})_{\mathbf{I}_C}\colon \mathbf{S}^{(2)}\to \mathbf{S}^{(1)}$. Now by Proposition 2.5 applied iteratively to the sequence of morphisms

$$\mathbf{S} \overset{\mathbf{C}^{(1)'}}{\longleftarrow} \mathbf{S}^{(1)} \overset{\mathbf{C}^{(2)'}}{\longleftarrow} \mathbf{S}^{(2)} \overset{\mathbf{C}^{(3)'}}{\longleftarrow} \mathbf{S}'.$$

we obtain a morphism $\mathbf{C}' \colon \mathbf{S}' \to \mathbf{S}$ as desired. The last assertion for b' follows from Proposition 3.5.

Remark 3.13. Proposition 3.12 also follows immediately from [C, Corollary 3.5.12]. In fact, the morphism $\mathbf{C} \in \operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})(\mathbf{S}, \mathbf{S}')$ satisfying the assumptions in Proposition 3.12 becomes a homotopy equivalence in $\operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})(\mathbf{S}, \mathbf{S}')$ (hence in \mathscr{E}).

Remark 3.14. The assumption on b in Proposition 3.12 is redundant. Indeed in Proposition 3.26 we will see that it holds automatically.

3.3. Splitting exact sequences.

Definition 3.15. Let $A, C \in \mathscr{C}_0$ be any pair of objects, and let

$$(3.12) A C$$

be any 2-simplex in \mathscr{C} , in which $O \in \mathscr{C}_0$ is a zero object. Let $A \stackrel{p_A}{\longleftarrow} A \times C \stackrel{p_C}{\longrightarrow} C$ and $A \stackrel{i_A}{\longrightarrow} A \coprod C \stackrel{i_C}{\longleftarrow} C$ be a product and a coproduct of A and C, respectively.

(1) By the definition of product, we obtain a pair of 2-simplices as below.

$$(3.13) \qquad A \qquad \downarrow u_A \qquad z \\ A \stackrel{\mu_{A,C}}{\longleftarrow} A \times C \stackrel{\pi_{A,C}}{\longrightarrow} C$$

(2) Dually, we obtain a pair of 2-simplices as below.

$$A \xrightarrow{i_{A}} A \coprod C \xrightarrow{i_{C}} C$$

$$\downarrow^{i_{A,C}} \downarrow^{j_{C}} \downarrow^{j_{C}} \downarrow^{j_{C}} \downarrow^{j_{C}}$$

We will use these symbols in the rest, often abbreviating $\pi_{A,C}$ to π and so on.

Proposition 3.16. Let (3.12) be any 2-simplex, in which $O \in \mathcal{C}_0$ is a zero object. Then the squares

(3.14)
$$A\mathbf{N}_{C} = i \int_{j}^{u_{A}} A \times C \qquad A\mathbf{N}'_{C} = i \int_{j}^{i_{A}} A \coprod C \qquad J_{j} C$$

 $are\ exact\ sequences.$

Proof. By using the definition of a zero object we may obtain a rectangle ${\bf R}$ of the following form

$$\begin{array}{c|c}
A & & & & \\
 & & & & \\
\downarrow & & & \\
\downarrow & & & & \\
\downarrow & & \\
\downarrow & & & \\
\downarrow &$$

in which $\mathbf{R}_{\text{left}} = \mathbf{N}$, and $\mathbf{R}_{\text{right}}$ is a pull-back. As in Example 3.3 we know that $\mathbf{R}_{\text{right}}$ is an exact sequence, thus it follows that \mathbf{N} is also a pull-back and p_C is egressive. This shows that \mathbf{N} is an exact sequence. Dually for \mathbf{N}' .

Remark 3.17. The above definition of ${}_{A}\mathbf{N}_{C}$ and ${}_{A}\mathbf{N}'_{C}$ depends on the choice of (3.12) and simplices taken. Later we will introduce an equivalence relation for exact sequences in Section 3.4, to show that their equivalence classes ${}_{A}\underline{\mathbf{N}}_{C}$, ${}_{A}\underline{\mathbf{N}}'_{C}$ do not depend on these choices, and moreover satisfy ${}_{A}\underline{\mathbf{N}}_{C} = {}_{A}\underline{\mathbf{N}}'_{C}$.

Lemma 3.18. Let ${}_{A}\mathbf{S}_{C}=(\eta\backslash\xi)$ be any exact sequence as in (3.2). Let ${}_{A}\mathbf{N}_{C}=(\eta\backslash\pi),{}_{A}\mathbf{N}_{C}'=(\eta\backslash\iota)$ be as in Proposition 3.16, obtained by using the 2-simplex η . The following holds.

(1) If \overline{x} is a split monomorphism in $h\mathscr{C}$, then there is a 3-simplex

$$\varphi = X \xrightarrow{B} Y$$

$$A \xrightarrow{y} C$$

$$A \times C$$

such that $d_1(\varphi) = \pi_{A,C}$ and $d_2(\varphi) = \xi$.

(2) Dually, if \overline{y} is a split epimorphism in $h\mathscr{C}$, then there is a 3-simplex ψ such that $d_1(\psi) = \xi$ and $d_2(\psi) = \iota_{A,C}$.

Proof. Since (2) can be shown dually, it is enough to show (1). Suppose that \overline{x} is a split monomorphism in $h\mathscr{C}$. This means that there is a 2-simplex ω of the following form.



Then by the definition of product, we obtain a pair of 2-simplices as below.

$$(3.15) \qquad A \xrightarrow{r} A \times C \xrightarrow{g} C$$

Also, ω and ξ gives the following pair of 2-simplices.

$$(3.16) A \xrightarrow{1_A} A \xrightarrow{x} z$$

$$A \xrightarrow{x} B \xrightarrow{x} C$$

By the definition of product, (3.15) and (3.16) give a pair of 3-simplices as below,

$$\varphi' = A \xrightarrow{1_A} A \xrightarrow{r} B , \quad \varphi = A \xrightarrow{x} A \xrightarrow{z} C$$

$$A \xrightarrow{p_A} A \xrightarrow{u_A} b \qquad A \times C$$

$$A \times C \qquad A \times C$$

such that $d_3(\varphi) = d_3(\varphi')$ holds and the pairs $(d_k(\varphi'), d_0(\varphi))$ (k = 0, 1, 2) agree with (3.15), (3.13), (3.16), respectively. This φ satisfies the stated properties.

Proposition 3.19. Let (3.12) be any 2-simplex in which $O \in \mathcal{C}_0$ is a zero object, and let ${}_{A}\mathbf{N}_{C}$, ${}_{A}\mathbf{N}_{C}'$ be as in Proposition 3.16. Then there exists a 3-simplex


$$\aleph_{A,C} = \begin{array}{c} A \coprod C \\ i_A \\ v_{A,C} \\ A \end{array} \begin{array}{c} j_C \\ \downarrow \\ p_C \\ A \times C \end{array}$$

such that $d_1(\aleph_{A,C}) = \pi_{A,C}$, $d_2(\aleph_{A,C}) = \iota_{A,C}$ and that $\overline{\upsilon_{A,C}}$ is an isomorphism in $h\mathscr{C}$.

Proof. Take any 2-simplex as below.



By the definition of a coproduct, we obtain a pair of 2-simplices



similarly as in Definition 3.15 (2). Then we have $\overline{j_A} \circ \overline{i_A} = \mathrm{id}_A$ in $h\mathscr{C}$, hence by Lemma 3.18 we obtain a 2-simplex $\aleph_{A,C}$ with $d_1(\aleph_{A,C}) = \pi_{A,C}, d_2(\aleph_{A,C}) = \iota_{A,C}$. As in the proof of Lemma 3.18, morphism $\upsilon_{A,C} = d_0 d_3(\aleph_{A,C})$ is obtained as a morphism appearing in the following

$$A \coprod C$$

$$j_{A} \downarrow v_{A,C} \downarrow j_{C}$$

$$A \times C \xrightarrow{p_{C}} C$$

by construction. In the homotopy category $h\mathscr{C}$, morphism $\mathfrak{v} = \overline{v_{A,C}}$ is the unique one which makes

(3.17)
$$A \coprod C$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

commutative, since the bottom row gives a product of A and C in $h\mathscr{C}$. By the additivity of $h\mathscr{C}$, this forces \mathfrak{v} to be an isomorphism.

Remark 3.20. In the sequel we will continue to use the symbol $v_{A,C}$ for a morphism which gives the unique (iso)morphism $\mathfrak{v} = \overline{v_{A,C}}$ in $h\mathscr{C}$ making (3.17) commutative.

Corollary 3.21. Let ${}_{A}\mathbf{S}_{C}=(\eta\backslash\xi), {}_{A}\mathbf{N}_{C}=(\eta\backslash\pi)$ and ${}_{A}\mathbf{N}_{C}'=(\eta\backslash\iota)$ be as in Lemma 3.18. Then, the following are equivalent.

(i) There exists a morphism

(3.18)
$$a\mathbf{C}_{c} = \begin{bmatrix} A & \xrightarrow{x} & B & y \\ & & O & \downarrow & C \\ & & & \downarrow & b & \downarrow c \\ A & \xrightarrow{u_{A}} & A \times C & p_{C} & \downarrow c \\ & & & & \downarrow & C \end{bmatrix} : \mathbf{S} \to \mathbf{N}$$

such that $\overline{a} = id_A$, $\overline{c} = id_C$.

- (i)' There exists a morphism $a' \mathbf{C}'_{c'} \colon \mathbf{N}' \to \mathbf{S}$ such that $\overline{a'} = \mathrm{id}_A$, $\overline{c'} = \mathrm{id}_C$.
- (ii) There exists a morphism $_{1_A}\mathbf{C}_{1_C}\colon \mathbf{S}\to \mathbf{N}$ in which $b=\mathbf{C}|_{\{(0,1)\}\times\Delta^1}$ is a homotopy equivalence.
- (ii)' There exists a morphism $_{1_A}\mathbf{C}'_{1_C} \colon \mathbf{N}' \to \mathbf{S}$ in which $b' = \mathbf{C}'|_{\{(0,1)\} \times \Delta^1}$ is a homotopy equivalence.
- (iii) \overline{x} is a split monomorphism in $h\mathscr{C}$.
- (iii)' \overline{y} is a split epimorphism in hC.

Remark that the conditions (iii) and (iii)' are independent of the choices of N and N'.

Proof. Equivalence (ii) \Leftrightarrow (ii)' follows from Proposition 2.5, Propositions 3.12 and 3.19. Since (i)' \Leftrightarrow (ii)' \Leftrightarrow (iii)' can be shown dually, it is enough to show (i) \Leftrightarrow (ii) \Leftrightarrow (iii). Remark that (ii) \Rightarrow (i) is trivial, and that (iii) \Rightarrow (i) follows from in Lemma 3.18. Also, (i) \Rightarrow (iii) is immediate. Indeed if there is a morphism (3.18) as in (i), then we have $\overline{p_A} \circ \overline{b} \circ \overline{x} = \overline{p_A} \circ \overline{u_A} \circ \overline{a} = \mathrm{id}_A$, which in particular means that \overline{x} is a split monomorphism.

It remains to show (i) \Rightarrow (ii). Suppose that we have a morphism (3.18) satisfying $\overline{a} = \mathrm{id}_A$ and $\overline{c} = \mathrm{id}_C$. By Corollary 2.6, we may assume $a = 1_A$ and $c = 1_C$ from the beginning. Let us show that \overline{b} becomes an isomorphism in $h\mathscr{C}$. Remark that the 2-simplex $\pi_{A,C}$ in (3.13) induces a split short exact sequence

$$A \xrightarrow{\overline{u_A}} A \times C \xrightarrow{\overline{p_C}} C$$

in $h\mathscr{C}$. Replacing it by the isomorphic one $A \xrightarrow{\left[\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}\right]} A \oplus C \xrightarrow{\left[\begin{smallmatrix} 0 & 1 \end{smallmatrix}\right]} C$, we see that **C** induces the following commutative diagram in $h\mathscr{C}$,

(3.19)
$$A \xrightarrow{\overline{x}} B \xrightarrow{\overline{y}} C$$

$$\downarrow C \downarrow \begin{bmatrix} \overline{v} \\ \overline{y} \end{bmatrix} \circlearrowleft \downarrow C$$

$$A \xrightarrow{\lceil \frac{1}{0} \rceil} A \oplus C \xrightarrow{\lceil 0 \ 1 \rceil} C$$

where we put $\mathfrak{r} = \overline{p_A} \circ \overline{b} \in (h\mathscr{C})(B,A)$. It suffices to show that $\left[\frac{\mathfrak{r}}{y}\right]$ is an isomorphism. Since the top row of (3.19) is a weak cokernel sequence, there is some $\mathfrak{w} \in (h\mathscr{C})(C,B)$ such that $\mathrm{id}_B - \overline{x} \circ \mathfrak{r} = \mathfrak{w} \circ \overline{y}$. This shows that $\left[\overline{x} \ \mathfrak{w}\right] : A \oplus C \to B$

gives a left inverse of $\left\lceil \frac{\mathfrak{r}}{y} \right\rceil$. On the other hand, since the right square of (3.19) is a weak push-out by the dual of Lemma 3.11 (1), it can be verified that $\left\lceil \frac{\mathfrak{r}}{y} \right\rceil$ is an epimorphism, by a straightforward argument in the ordinary additive category $h\mathscr{C}$. Thus $\left\lceil \frac{\mathfrak{r}}{y} \right\rceil$ is an isomorphism.

Definition 3.22. An exact sequence **S** is said to *split*, if it satisfies condition (iii) (or other equivalent conditions) in Corollary 3.21.

Corollary 3.23. Let $A, C \in \mathcal{C}_0$ be any pair of objects, and let ${}_{A}\mathbf{S}_{C}, {}_{A}\mathbf{S}'_{C}$ be exact sequences starting from A and ending in C. Suppose that there exists a morphism ${}_{1_{A}}\mathbf{C}_{1_{C}} \colon \mathbf{S} \to \mathbf{S}'$. If one of ${}_{A}\mathbf{S}_{C}, {}_{A}\mathbf{S}'_{C}$ splits, then so does the other.

Proof. This can be easily checked by using (iii) and (iii)' in Corollary 3.21. \Box

3.4. Equivalence of exact sequences.

Lemma 3.24. Let ${}_{A}\mathbf{S}_{C}$ be any exact sequence as in (3.2). Then as $x_{*}\mathbf{S}$ of Lemma 3.10, we may take a splitting one. Dually, $y^{*}\mathbf{S}$ can be chosen to split.

Proof. By [B1, Lemma 4.6], we have a pull-back of the following form.

$$\begin{array}{ccc}
A \times B \xrightarrow{p_B} B \\
\downarrow & & \downarrow y \\
B \xrightarrow{y} C
\end{array}$$

If we use this pull-back, the resulting y^*S becomes of the form

$$A \longrightarrow A \times B$$

$$\downarrow \qquad \qquad \downarrow^{p_B},$$

$$O \longrightarrow B$$

which obviously satisfies (iii)' of Corollary 3.21. Dually for x_*S .

Remark 3.25. Later in Corollary 4.6 (1) we will see that all $x_*\mathbf{S}$ become equivalent, and hence in particular always split, independently of the choices made. Similarly for $y^*\mathbf{S}$.

Proposition 3.26. For any morphism of exact sequences $_{1_A}\mathbf{C}_{1_C}\colon \mathbf{S}\to \mathbf{S}',$ the morphism $b=\mathbf{C}|_{\{(0,1)\}\times\Delta^1}$ becomes a homotopy equivalence.

Thus the assumption on b in Proposition 3.12 is always satisfied, hence there exists a morphism in the opposite direction ${}_{1_A}\mathbf{C}'_{1_C}\colon \mathbf{S}'\to \mathbf{S}$.

Proof. Label C as in (3.11). Take $x_*\mathbf{S}$ and $x_*\mathbf{S}'$. By Lemma 3.24, we may choose $x_*\mathbf{S}$ to split. There are morphisms

$$_{x}\mathbf{D}_{1_{C}}\colon \mathbf{S} \to x_{*}\mathbf{S}$$
 and $_{x}\mathbf{D}_{1_{C}}'\colon \mathbf{S}' \to x_{*}\mathbf{S}'$

as in Lemma 3.10 (1). By Proposition 2.5 applied to $\mathbf{S} \xrightarrow{^{1}_{A}\mathbf{C}_{1_{C}}} \mathbf{S}' \xrightarrow{^{x}(\mathbf{D}')_{1_{C}}} x_{*}\mathbf{S}'$, we obtain a morphism $_{x}\mathbf{C}'_{1_{C}} \colon \mathbf{S} \to x_{*}\mathbf{S}'$. Thus by Lemma 3.10 (2), we obtain a morphism $_{1_{B}}\mathbf{C}''_{1_{C}} \colon x_{*}\mathbf{S} \to x_{*}\mathbf{S}'$. Since $x_{*}\mathbf{S}$ splits, it follows that $x_{*}\mathbf{S}'$ also splits by Corollary 3.23.

Let us show that \overline{b} is an isomorphism in $h\mathscr{C}$. If we label simplices of \mathbf{D}' as below,

$$\begin{array}{c|c}
A \xrightarrow{x'} B' & y' \\
\downarrow i' & O' \xrightarrow{1} C \\
\downarrow 1_{O'} & \downarrow b_M \\
B \xrightarrow{x_M} & M & y_M \\
\downarrow O' \xrightarrow{j'} C
\end{array}$$

then we have the following commutative diagram

$$A \xrightarrow{\overline{x'}} B' \xrightarrow{\overline{y'}} C$$

$$\overline{x} \downarrow \quad \Diamond \quad \downarrow \overline{b_M} \Diamond \quad \parallel$$

$$B \xrightarrow{\overline{x_M}} M \xrightarrow{\overline{y_M}} C$$

in $h\mathscr{C}$. Since x_*S' splits, there is some $\mathfrak{r} \in (h\mathscr{C})(M,B)$ such that $\mathfrak{r} \circ \overline{x_M} = \mathrm{id}_B$. Since $A \xrightarrow{\overline{x}} B \xrightarrow{\overline{y}} C$ is a weak cokernel sequence, there exists $\mathfrak{d} \in (h\mathscr{C})(C,B)$ such that $\mathrm{id}_B - \mathfrak{r} \circ \overline{b_M} \circ \overline{b} = \mathfrak{d} \circ \overline{y}$. If we put $\mathfrak{b} = \mathfrak{r} \circ \overline{b_M} + \mathfrak{d} \circ \overline{y'} \in (h\mathscr{C})(B',B)$, then it satisfies

$$\mathfrak{b} \circ \overline{b} = \mathfrak{r} \circ \overline{b_M} \circ \overline{b} + \mathfrak{d} \circ \overline{y'} \circ \overline{b} = \mathrm{id}_B,$$

which shows that \mathfrak{b} is a left inverse of \overline{b} .

A dual argument using $y'^*\mathbf{S}$ and $y'^*\mathbf{S}'$ shows that \overline{b} has also a right inverse, and thus it is an isomorphism.

Definition 3.27. Let $A, C \in \mathcal{C}_0$ be any pair of objects. Let

$$(3.20) A \xrightarrow{x} B A \xrightarrow{x'} B'$$

$$Q \xrightarrow{j} C Q' \xrightarrow{j'} C$$

be two exact sequences starting from A and ending in C. We say \mathbf{S} is equivalent to \mathbf{S}' if there is a morphism of exact sequences ${}_{1_A}\mathbf{C}_{1_C}\colon \mathbf{S}\to \mathbf{S}'$. In this case we write $\mathbf{S}\sim \mathbf{S}'$. By Corollary 2.6, this is equivalent to the existence of a morphism ${}_a\mathbf{C}'_c\colon \mathbf{S}\to \mathbf{S}'$ satisfying $\overline{a}=\mathrm{id}_A$ and $\overline{c}=\mathrm{id}_C$.

Proposition 3.28. For any $A, C \in \mathcal{C}_0$, relation \sim is an equivalence relation.


Proof. Reflexive law is obvious, since there is always $I_S: S \to S$ for any exact sequence S as in Example 2.4. Symmetric law and transitive law follow from Propositions 3.26 and 2.5, respectively.

3.5. (Co)products of exact sequences. We begin with the following consequence of [L1, Corollary 5.1.2.3]. Let

(3.21)
$$\mathbf{S}_{k} = \begin{bmatrix} A_{k} & x_{k} \\ \vdots & B_{k} \\ z_{k} & y_{k} \\ O_{k} & x_{k} \\ \vdots & \vdots \\ C_{k} & x_{k$$

be any pair of squares. By [L1, Corollary 5.1.2.3], a product $\mathbf{S}_1 \times \mathbf{S}_2$ of $\mathbf{S}_1, \mathbf{S}_2$ in $\operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})$ can be obtained by taking products on each vertex of $\Delta^1 \times \Delta^1$ as follows,

together with projections $\mathbf{Q}_k \in \operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})_1(\mathbf{S}_1 \times \mathbf{S}_2, \mathbf{S}_k)$ (k = 1, 2). In particular, for any square \mathbf{S} and any pair of morphisms $\mathbf{C}_k \in \operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})_1(\mathbf{S}, \mathbf{S}_k)$ (k = 1, 2), we obtain a morphism $(\mathbf{C}_1, \mathbf{C}_2)$ and a pair of 2-simplices



in Fun($\Delta^1 \times \Delta^1, \mathscr{C}$). Dually, we have a coproduct

$$\mathbf{S}_1 \coprod \mathbf{S}_2 = i_1 \coprod i_2 \underbrace{ \begin{matrix} x_1 \coprod x_2 \\ \xi_1 \coprod \xi_2 \end{matrix}}_{j_1 \coprod j_2} B_1 \coprod B_2$$


$$O_1 \coprod O_2 \xrightarrow[j_1 \coprod j_2]{} C_1 \coprod C_2$$

For any square **S** and any $\mathbf{C}_k \in \operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})_1(\mathbf{S}_k, \mathbf{S})$ (k = 1, 2), we have $\mathbf{C}_1 \cup \mathbf{C}_2 \in \operatorname{Fun}(\Delta^1 \times \Delta^1, \mathscr{C})_1(\mathbf{S}_1 \coprod \mathbf{S}_2, \mathbf{S})$ with a property dual to the above.

Again by [L1, Corollary 5.1.2.3], if S_1, S_2 are pull-backs, then so is $S_1 \times S_2$. Similarly, if S_1, S_2 are push-outs, then so is $S_1 \coprod S_2$.


Proposition 3.29. If S_1, S_2 are exact sequences, then so are $S_1 \times S_2$ and $S_1 \coprod S_2$.

Proof. Let $\mathbf{S}_1, \mathbf{S}_2$ be as in (3.21). Since egressive morphisms are stable under taking pull-backs, it follows that $y_1 \times 1_{A_2}$ and $1_{B_1} \times y_2$ are egressive morphisms. Since there is a 2-simplex



in \mathscr{C} , it follows that $y_1 \times y_2$ is also egressive. Since $j_1 \times j_2$ in (3.22) is ingressive because $O_1 \times O_2$ is a zero object, we see that $\mathbf{S}_1 \times \mathbf{S}_2$ is an exact sequence. Dually for $\mathbf{S}_1 \coprod \mathbf{S}_2$.

Example 3.30. Let ${}_{A}\mathbf{S}_{C}$ be any exact sequence as in (3.2). From $\mathbf{I}_{\mathbf{S}} \colon \mathbf{S} \to \mathbf{S}$, we obtain the following morphisms of exact sequences, which we denote by $\nabla_{\mathbf{S}} = \mathbf{I}_{\mathbf{S}} \cup \mathbf{I}_{\mathbf{S}} \colon \mathbf{S} \coprod \mathbf{S} \to \mathbf{S}$ and $\Delta_{\mathbf{S}} = (\mathbf{I}_{\mathbf{S}}, \mathbf{I}_{\mathbf{S}}) \colon \mathbf{S} \to \mathbf{S} \times \mathbf{S}$, respectively.



Proposition 3.31. Let

$$\mathbf{C}_{k} = \begin{pmatrix} A_{k} & X_{k} & B_{k} & y_{k} \\ \downarrow^{i_{k}} & O_{k} & \downarrow^{j_{k}} & C_{k} \\ A_{k}' & X_{k}' & \downarrow^{j_{k}} & B_{k}' & y_{k}' \\ \downarrow^{i_{k}} & O_{k}' & X_{k}' & C_{k}' \end{pmatrix} : \mathbf{S}_{k} \to \mathbf{S}_{k}' \quad (k = 1, 2)$$

be any pair of morphisms of exact sequences. Then we have the following morphism of exact sequences $S_1 \coprod S_2 \to S_1' \times S_2'$

(3.23)
$$A_{1} \coprod A_{2} \xrightarrow{x_{1} \coprod x_{2}} B_{1} \coprod B_{2}$$

$$V_{a_{1},a_{2}} \downarrow O_{1} \coprod O_{2} \xrightarrow{j_{1} \coprod j_{2}} C_{1} \coprod C_{2}$$

$$V_{o_{1},o_{2}} \downarrow V_{o_{1},o_{2}} \downarrow V_{o_{1},b_{2}} \downarrow V_{o_{1},b_{2}} \downarrow V_{o_{1},c_{2}}$$

$$A'_{1} \times A'_{2} \xrightarrow{x'_{1} \times x'_{2}} B'_{1} \times B'_{2} \xrightarrow{y'_{1} \times y'_{2}} V_{c_{1},c_{2}}$$

$$i'_{1} \times i'_{2} O'_{1} \times O'_{2} \xrightarrow{j'_{1} \times j'_{2}} C'_{1} \times C'_{2}$$

in which v_{a_1,a_2} gives the unique morphism $\overline{v_{a_1,a_2}} = \overline{a_1} \oplus \overline{a_2} \in (h\mathscr{C})(A_1 \oplus A_2, A_1' \oplus A_2')$ in the homotopy category, and similarly for $v_{b_1,b_2}, v_{c_1,c_2}, v_{o_1,o_2}$.

Proof. As in Example 2.4, we have morphisms

$$\mathbf{I}_k' = \mathbf{I}_{\mathbf{S}_k'} \colon \mathbf{S}_k' \to \mathbf{S}_k', \quad \ \mathbf{C}_k' \colon \mathbf{S}_k' \to \mathbf{O}, \quad \mathbf{C}_k'' \colon \mathbf{O} \to \mathbf{S}_k'.$$

By Proposition 2.5 applied to $\mathbf{S}_1' \xrightarrow{\mathbf{C}_1'} \mathbf{O} \xrightarrow{\mathbf{C}_2''} \mathbf{S}_2'$, we obtain a morphism $\mathbf{Z}_1' \colon \mathbf{S}_1' \to \mathbf{S}_2'$. Similarly for $\mathbf{Z}_2' \colon \mathbf{S}_2' \to \mathbf{S}_1'$.

Then we have morphisms

$$(\mathbf{I}_1', \mathbf{Z}_1') \colon \mathbf{S}_1' \to \mathbf{S}_1' \times \mathbf{S}_2'$$
 and $(\mathbf{Z}_2', \mathbf{I}_2') \colon \mathbf{S}_2' \to \mathbf{S}_1' \times \mathbf{S}_2'$

By Proposition 2.5 applied to $\mathbf{S}_1 \xrightarrow{\mathbf{C}_1} \mathbf{S}_1' \xrightarrow{(\mathbf{I}_1', \mathbf{Z}_1')} \mathbf{S}_1' \times \mathbf{S}_2'$ and $\mathbf{S}_2 \xrightarrow{\mathbf{C}_2} \mathbf{S}_2' \xrightarrow{(\mathbf{Z}_2', \mathbf{I}_2')} \mathbf{S}_1' \times \mathbf{S}_2'$ respectively, we obtain morphisms $\mathbf{F}_1 \colon \mathbf{S}_1 \to \mathbf{S}_1' \times \mathbf{S}_2'$ and $\mathbf{F}_2 \colon \mathbf{S}_2 \to \mathbf{S}_1' \times \mathbf{S}_2'$. Then we obtain a morphism $\mathbf{F}_1 \cup \mathbf{F}_2 \colon \mathbf{S}_1 \coprod \mathbf{S}_2 \to \mathbf{S}_1' \times \mathbf{S}_2'$ as in (3.23) with the stated property.

Corollary 3.32. For any pair of squares S_k (k = 1, 2) as in (3.21), we obtain the following morphism $S_1 \coprod S_2 \to S_1 \times S_2$

$$\begin{array}{c|c} A_1 \coprod A_2 \xrightarrow[i_1 \coprod i_2]{} B_1 \coprod B_2 \\ v_{A_1,A_2} & O_1 \coprod O_2 \xrightarrow[j_1 \coprod j_2]{} C_1 \coprod C_2 \\ v_{O_1,O_2} & v_{B_1,B_2} \\ A_1 \times A_2 \xrightarrow[i_1 \times i_2]{} B_1 \times B_2 \\ & O_1 \times O_2 \xrightarrow[j_1 \times j_2]{} C_1 \times C_2 \end{array}$$

in which, the vertical morphisms are homotopy equivalences as in Remark 3.20.

Proof. This follows from Proposition 3.31 applied to $I_{S_1}: S_1 \to S_1$ and $I_{S_2}: S_2 \to S_2$.

4. Extriangulated structure on the homotopy category

By definition, an extriangulated category is an additive category equipped with a structure $(\mathbb{E}, \mathfrak{s})$ satisfying some conditions. For the convenience of the reader, we write them down in the case of the homotopy category $h\mathscr{C}$. For the detail, see [NP].

Definition 4.1. ([NP, Definition 2.7]) For a pair of objects $A, C \in h\mathscr{C}$, two consecutive morphisms $A \xrightarrow{\mathfrak{r}} B \xrightarrow{\mathfrak{y}} C$ and $A \xrightarrow{\mathfrak{r}'} B' \xrightarrow{\mathfrak{y}'} C$ are said to be *equivalent* if there is an isomorphism $\mathfrak{b} \in (h\mathscr{C})(B, B')$ such that $\mathfrak{b} \circ \mathfrak{x} = \mathfrak{x}'$ and $\mathfrak{y}' \circ \mathfrak{b} = \mathfrak{y}$.

In the sequel, the equivalence class to which a sequence $A \xrightarrow{\mathfrak{x}} B \xrightarrow{\mathfrak{y}} C$ belongs will be denoted by $[A \xrightarrow{\mathfrak{x}} B \xrightarrow{\mathfrak{y}} C]$.

Definition 4.2. ([NP, Definitions 2.9 and 2.10]) Suppose that there is a biadditive functor \mathbb{E} : $(h\mathscr{C})^{\mathrm{op}} \times (h\mathscr{C}) \to Ab$ to the category of abelian groups Ab. Let \mathfrak{s} be a correspondence which associates an equivalence class $\mathfrak{s}(\delta) = [A \xrightarrow{\mathfrak{r}} B \xrightarrow{\mathfrak{y}} C]$ to each element $\delta \in \mathbb{E}(C,A)$. Such \mathfrak{s} is called an *additive realization of* \mathbb{E} if it satisfies the following conditions.

(i) Let $\delta \in \mathbb{E}(C, A)$ and $\delta' \in \mathbb{E}(C', A')$ be any pair of elements, with

(4.1)
$$\mathfrak{s}(\delta) = [A \xrightarrow{\mathfrak{r}} B \xrightarrow{\mathfrak{y}} C], \quad \mathfrak{s}(\delta') = [A' \xrightarrow{\mathfrak{r}'} B' \xrightarrow{\mathfrak{y}'} C'].$$

Then, for any pair of morphisms $\mathfrak{a} \in (h\mathscr{C})(A, A')$, $\mathfrak{c} \in (h\mathscr{C})(C, C')$ satisfying $\mathbb{E}(C, \mathfrak{a})(\delta) = \mathbb{E}(\mathfrak{c}, A')(\delta')$, there exists a morphism $\mathfrak{b} \in (h\mathscr{C})(B, B')$ such that $\mathfrak{b} \circ \mathfrak{x} = \mathfrak{x}' \circ \mathfrak{a}$ and $\mathfrak{y}' \circ \mathfrak{b} = \mathfrak{c} \circ \mathfrak{y}$.

(ii) For any $A, C \in h\mathscr{C}$, the zero element $0 \in \mathbb{E}(C, A)$ satisfies

$$\mathfrak{s}(0) = [A \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} A \oplus C \xrightarrow{[0 \ 1]} C].$$

(iii) For any pair of elements $\delta \in \mathbb{E}(C,A)$ and $\delta' \in \mathbb{E}(C',A')$ with (4.1), we have

$$\mathfrak{s}(\delta \oplus \delta') = [A \oplus A' \xrightarrow{\mathfrak{x} \oplus \mathfrak{x}'} B \oplus B' \xrightarrow{\mathfrak{y} \oplus \mathfrak{y}'} C \oplus C'].$$

Here $\delta \oplus \delta' \in \mathbb{E}(C \oplus C', A \oplus A')$ denotes the element which corresponds to $(\delta, 0, 0, \delta')$ through the natural isomorphism

$$\mathbb{E}(C \oplus C', A \oplus A') \cong \mathbb{E}(C, A) \oplus \mathbb{E}(C, A') \oplus \mathbb{E}(C', A) \oplus \mathbb{E}(C', A')$$

induced from the biadditivity of \mathbb{E} .

If $\mathfrak s$ is an additive realization, any pair $(A \xrightarrow{\mathfrak r} B \xrightarrow{\mathfrak v} C, \delta)$ of a sequence $A \xrightarrow{\mathfrak r} B \xrightarrow{\mathfrak v} C$ and an element $\delta \in \mathbb E(C,A)$ satisfying $\mathfrak s(\delta) = [A \xrightarrow{\mathfrak r} B \xrightarrow{\mathfrak v} C]$ is called an $\mathfrak s$ -triangle (or simply an extriangle if $\mathfrak s$ is obvious from the context), and abbreviately expressed by $A \xrightarrow{\mathfrak r} B \xrightarrow{\mathfrak v} C \xrightarrow{\delta}$.

Definition 4.3. ([NP, Definition 2.12]) A triplet $(h\mathscr{C}, \mathbb{E}, \mathfrak{s})$ is called an *extriangulated category* if it satisfies the following conditions. In this case, the pair $(\mathbb{E}, \mathfrak{s})$ is called an *external triangulation* of $h\mathscr{C}$.

- (ET1) $\mathbb{E}: (h\mathscr{C})^{\mathrm{op}} \times (h\mathscr{C}) \to Ab$ is a biadditive functor.
- (ET2) \mathfrak{s} is an additive realization of \mathbb{E} .
- (ET3) Let $\delta \in \mathbb{E}(C, A)$ and $\delta' \in \mathbb{E}(C', A')$ be any pair of elements with (4.1). Then, for any commutative square

$$(4.2) A \xrightarrow{\mathfrak{x}} B \xrightarrow{\mathfrak{y}} C$$

$$\downarrow \mathfrak{b}$$

$$A' \xrightarrow{\mathfrak{x}'} B' \xrightarrow{\mathfrak{y}'} C'$$

in $h\mathscr{C}$, there exists $\mathfrak{c} \in (h\mathscr{C})(C, C')$ such that $\mathfrak{c} \circ \mathfrak{y} = \mathfrak{y}' \circ \mathfrak{b}$ and $\mathbb{E}(C, \mathfrak{a})(\delta) = \mathbb{E}(\mathfrak{c}, A')(\delta')$.

(ET3)^{op} Dual of (ET3).

(ET4) Let $\delta \in \mathbb{E}(D,A)$ and $\delta' \in \mathbb{E}(F,B)$ be any pair of elements, with

$$\mathfrak{s}(\delta) = [A \xrightarrow{\mathfrak{f}} B \xrightarrow{\mathfrak{f}'} D] \text{ and } \mathfrak{s}(\delta') = [B \xrightarrow{\mathfrak{g}} C \xrightarrow{\mathfrak{g}'} F].$$

Then there exist an object $E \in h\mathscr{C}$, a commutative diagram

$$\begin{array}{cccc}
A & \xrightarrow{\mathfrak{f}} B & \xrightarrow{\mathfrak{f}'} D \\
\parallel & \circlearrowleft & \mathfrak{g} & \circlearrowleft & \flat & \mathfrak{d} \\
A & \xrightarrow{\mathfrak{h}} C & \xrightarrow{\mathfrak{h}'} E \\
& & & & & & & & \downarrow \mathfrak{c} \\
& & & & & & & \downarrow \mathfrak{c} \\
& & & & & & & & \downarrow \mathfrak{c}
\end{array}$$

in $h\mathscr{C}$, and an element $\delta'' \in \mathbb{E}(E,A)$ such that $\mathfrak{s}(\delta'') = [A \xrightarrow{\mathfrak{h}} C \xrightarrow{\mathfrak{h}'} E]$, which satisfy the following compatibilities.

- (i) $\mathfrak{s}(\mathbb{E}(F,\mathfrak{f}')(\delta')) = [D \xrightarrow{\mathfrak{d}} E \xrightarrow{\mathfrak{e}} F],$
- (ii) $\mathbb{E}(\mathfrak{d}, A)(\delta'') = \delta$,
- (iii) $\mathbb{E}(E,\mathfrak{f})(\delta'') = \mathbb{E}(\mathfrak{e},B)(\delta').$

(ET4)^{op} Dual of (ET4).

In the rest of this section, we will show that $h\mathscr{C}$ can be equipped with an external triangulation $(\mathbb{E}, \mathfrak{s})$, for any exact quasi-category \mathscr{C} .

4.1. Construction of the functor \mathbb{E} . Firstly, in this subsection we construct a functor \mathbb{E} : $(h\mathscr{C})^{\mathrm{op}} \times (h\mathscr{C}) \to Set$ to the category Set of sets.

Definition 4.4. Let $A, C \in \mathcal{C}_0$. Define $\mathbb{E}(C, A)$ to be the set of equivalence classes of exact sequences **S** starting from A and ending in C, modulo the equivalence relation \sim defined in Definition 3.27.

For each exact sequence ${}_{A}\mathbf{S}_{C}$, let $\delta = \underline{\mathbf{S}}$ denote its equivalence class with respect to \sim . When we emphasize the end-objects A and C, we write ${}_{A}\delta_{C} = {}_{A}\underline{\mathbf{S}}_{C}$.

We make use of the notation $a_*\mathbf{S}$ from Lemma 3.10.

Proposition 4.5. Suppose that we have ${}_{A}\mathbf{S}_{C} \sim {}_{A}\mathbf{S}_{C}'$, and let



be any 2-simplex in \mathscr{C} . Then we have $a_{2*}a_{1*}\mathbf{S}' \sim a_{3*}\mathbf{S}$.

Proof. By the assumption, there is a morphism $_{1_A}\mathbf{C}_{1_C} \colon \mathbf{S} \to \mathbf{S}'$. By the definition of $a_{1*}\mathbf{S}'$, $a_{2*}a_{1*}\mathbf{S}'$ and $a_{3*}\mathbf{S}$, there are morphisms of exact sequences

$$a_1 \mathbf{D}_{1_C}^{(1)} \colon \mathbf{S}' \to a_{1*} \mathbf{S}', \ a_2 \mathbf{D}_{1_C}^{(2)} \colon a_{1*} \mathbf{S}' \to a_{2*} a_{1*} \mathbf{S}', \ a_3 \mathbf{D}_{1_C}^{(3)} \colon \mathbf{S} \to a_{3*} \mathbf{S}'$$

in which $\mathbf{D}^{(k)}|_{\{0\}\times\Delta^1\times\Delta^1}$ (k=1,2,3) are push-outs. By Proposition 2.5 applied iteratively to $\mathbf{S} \stackrel{\mathbf{C}}{\longrightarrow} \mathbf{S}' \stackrel{\mathbf{D}^{(1)}}{\longrightarrow} a_{1*}\mathbf{S}' \stackrel{\mathbf{D}^{(2)}}{\longrightarrow} a_{2*}a_{1*}\mathbf{S}'$, we obtain a morphism ${}_{a_3}\mathbf{C}'_{1_C} \colon \mathbf{S} \to a_{2*}a_{1*}\mathbf{S}'$. Then by Lemma 3.10 (2), we obtain a morphism ${}_{1_{A''}}\mathbf{C}''_{1_C} \colon a_{3*}\mathbf{S} \to a_{2*}a_{1*}\mathbf{S}'$. This shows $a_{3*}\mathbf{S} \sim a_{2*}a_{1*}\mathbf{S}'$.

Corollary 4.6. We have the following.

(1) If morphisms $a, a' \in \mathcal{C}_1(A, A')$ and exact sequences ${}_{A}\mathbf{S}_{C}$, ${}_{A}\mathbf{S}_{C}$ satisfy $\overline{a} = \overline{a'}$ and $\mathbf{S} \sim \mathbf{S}'$, then $a_*\mathbf{S} \sim a_*'\mathbf{S}'$. Thus for any $\mathfrak{a} = \overline{a} \in (h\mathcal{C})(A, A')$ and any $C \in \mathcal{C}_0$, the map

$$\mathfrak{a}_* \colon \mathbb{E}(C, A) \to \mathbb{E}(C, A') \; ; \; \delta = \underline{\mathbf{S}} \mapsto \mathfrak{a}_* \delta = a_* \mathbf{S}$$

is well-defined.

- (2) The maps obtained in (1) satisfy the following.
 - (i) For any $A, C \in \mathscr{C}_0$, we have $(\mathrm{id}_A)_* = \mathrm{id}_{\mathbb{E}(C,A)} \colon \mathbb{E}(C,A) \to \mathbb{E}(C,A)$.
 - (ii) For any $\mathfrak{a} \in (h\mathscr{C})(A, A')$, $\mathfrak{a}' \in (h\mathscr{C})(A', A'')$ and any $C \in \mathscr{C}_0$, we have $(\mathfrak{a}' \circ \mathfrak{a})_* = \mathfrak{a}'_* \circ \mathfrak{a}_* \colon \mathbb{E}(C, A) \to \mathbb{E}(C, A'')$.

Proof. (1) and (ii) of (2) are immediate from Proposition 4.5. (i) of (2) is also obvious by the existence of $\mathbf{I}_{\mathbf{S}}$ for any ${}_{A}\mathbf{S}_{C}$.

Lemma 4.7. Let ${}_{A}\mathbf{S}_{C}$, ${}_{A'}\mathbf{S}'_{C'}$ be any pair of exact sequences, and put $\delta = \underline{\mathbf{S}}$, $\delta' = \underline{\mathbf{S}'}$. Let $\mathfrak{a} \in (h\mathscr{C})(A, A')$ and $\mathfrak{c} \in (h\mathscr{C})(C, C')$ be any pair of morphisms in the homotopy category. The following are equivalent.

- (1) $\mathfrak{a}_*\delta = \mathfrak{c}^*\delta'$.
- (2) There exists a morphism ${}_{a}\mathbf{C}_{c} \colon \mathbf{S} \to \mathbf{S}'$ such that $\overline{a} = \mathfrak{a}$ and $\overline{c} = \mathfrak{c}$.

Proof. (1) \Rightarrow (2). Take any $a \in \mathscr{C}_1(A, A')$ and $c \in \mathscr{C}_1(C, C')$ such that $\mathfrak{a} = \overline{a}$ and $\mathfrak{c} = \overline{c}$. By definition, there exist morphisms $\mathbf{S} \xrightarrow{a\mathbf{D}_{1_C}} a_*\mathbf{S} \xrightarrow{1_{A'}\mathbf{C}_{1_C}^{(1)}} c^*\mathbf{S}' \xrightarrow{1_{A'}\mathbf{D}_{c}'} \mathbf{S}'$ of exact sequences. By Proposition 2.5, we obtain a morphism ${}_a\mathbf{C}_c \colon \mathbf{S} \to \mathbf{S}'$ as desired.

 $(2) \Rightarrow (1)$. Let ${}_{a}\mathbf{C}_{c} \colon \mathbf{S} \to \mathbf{S}'$ be a morphism as in (3.4) such that $\overline{a} = \mathfrak{a}$, $\overline{c} = \mathfrak{c}$. By Lemma 3.10, we obtain a morphism ${}_{1_{A'}}\mathbf{C}'_{c} \colon a_{*}\mathbf{S} \to \mathbf{S}'$. Then by the dual of Lemma 3.10 applied to \mathbf{C}' and a pull-back along c, we obtain a morphism ${}_{1_{A'}}\mathbf{C}''_{1_{C}} \colon a_{*}\mathbf{S} \to c^{*}\mathbf{S}'$. Thus $a_{*}\mathbf{S} \sim c^{*}\mathbf{S}'$ follows.

Proposition 4.8. For any $\mathfrak{a} \in (h\mathscr{C})(A, A')$ and $\mathfrak{c} \in (h\mathscr{C})(C', C)$, we have

$$\mathfrak{a}_*\mathfrak{c}^* = \mathfrak{c}^*\mathfrak{a}_* \colon \mathbb{E}(C,A) \to \mathbb{E}(C',A').$$

Proof. Take any $a \in \mathcal{C}_1(A, A')$ and $c \in \mathcal{C}_1(C', C)$ so that $\mathfrak{a} = \overline{a}$ and $\mathfrak{c} = \overline{c}$ hold. Let ${}_{A}\mathbf{S}_{C}$ be any exact sequence starting from A and ending in C. By Lemma 3.10 (1) and its dual, we obtain morphisms of exact sequences

$$c^*\mathbf{S} \xrightarrow{\mathbf{1}_A \mathbf{D}_c'} \mathbf{S} \xrightarrow{a \mathbf{D}_{\mathbf{1}_C}} a_*\mathbf{S}.$$

By Proposition 2.5, we obtain a morphism ${}_{a}\mathbf{C}''_{c} \colon c^{*}\mathbf{S} \to a_{*}\mathbf{S}$. Lemma 4.7 shows $\mathfrak{a}_{*}(\mathfrak{c}^{*}\underline{\mathbf{S}}) = \mathfrak{c}^{*}(\mathfrak{a}_{*}\underline{\mathbf{S}})$.

The argument so far gives the following functor.

Definition 4.9. Functor $\mathbb{E}: (h\mathscr{C})^{\mathrm{op}} \times h\mathscr{C} \to Set$ is defined by the following.

- For an object $(C, A) \in (h\mathscr{C})^{\mathrm{op}} \times (h\mathscr{C})$, the set $\mathbb{E}(C, A)$ is the one defined in Definition 4.4.
- For a morphism $(\mathfrak{c},\mathfrak{a}) \in ((h\mathscr{C})^{\mathrm{op}} \times (h\mathscr{C}))((C,A),(C',A'))$, the map

$$\mathbb{E}(\mathfrak{c},\mathfrak{a})\colon \mathbb{E}(C,A)\to \mathbb{E}(C',A')$$

is the one defined by $\mathbb{E}(\mathfrak{c},\mathfrak{a}) = \mathfrak{c}^*\mathfrak{a}_* = \mathfrak{a}_*\mathfrak{c}^*$.

4.2. **Biadditivity of** \mathbb{E} . In this subsection we will show that the functor \mathbb{E} obtained in Definition 4.9 indeed factors through Ab.

Definition 4.10. By Corollary 3.23, the equivalence class of the splitting exact sequences forms one element in $\mathbb{E}(C,A)$ for each $A,C \in h\mathscr{C}$. We denote this particular element by ${}_{A}0_{C}$. Especially we have ${}_{A}0_{C} = {}_{A}\underline{\mathbf{N}}_{C} = {}_{A}\underline{\mathbf{N}}_{C}'$.

Remark 4.11. It is not hard to see that $\mathbb{E}(O,O)$ consists of only one object for each zero object $O \in \mathscr{C}_0$, which necessarily agrees with ${}_O O_O$. Remark that a morphism $a \in \mathscr{C}_1(A,A')$ satisfies $\overline{a}=0$ in $h\mathscr{C}$ if and only if it factors through some (equivalently, any) zero object. For any $\delta \in \mathbb{E}(C,A)$ and any $a \in \mathscr{C}_1(A,A')$ such that $\overline{a}=0$, we have $a_*\delta = {}_{A'}O_C$ by Lemma 3.24. Similarly we have $0^*\delta = {}_A O_{C'}$ for the zero morphism $0 \in (h\mathscr{C})(C',C)$.

Proposition 4.12. The following holds.

(1) For any $A, A' \in \mathcal{C}_0$, there is a natural isomorphism

$$(v_{A,A'})_* \colon \mathbb{E}(-, A \coprod A') \stackrel{\cong}{\Longrightarrow} \mathbb{E}(-, A \times A')$$

of functors $(h\mathscr{C})^{\mathrm{op}} \to Set$.

(2) For any $C, C' \in \mathcal{C}_0$, there is a natural isomorphism

$$(v_{C,C'})^* \colon \mathbb{E}(C \times C', -) \stackrel{\cong}{\Longrightarrow} \mathbb{E}(C \coprod C', -)$$

of functors $h\mathscr{C} \to Set$.

(3) For any $A, A', C, C' \in \mathcal{C}_0$, there is an isomorphism

$$\vartheta = \vartheta_{A',C'}^{A,C} \colon \mathbb{E}(C \coprod C', A \coprod A') \stackrel{\cong}{\Longrightarrow} \mathbb{E}(C \times C', A \times A')$$

given by $\vartheta_{A',C'}^{A,C} = ((\upsilon_{C,C'})^*)^{-1} \circ (\upsilon_{A,A'})_*$. This sends $\delta \coprod \delta' = \underline{\mathbf{S} \coprod \mathbf{S}'}$ to $\delta \times \delta' = \underline{\mathbf{S} \times \mathbf{S}'}$ for any $\delta = \underline{\mathbf{S}} \in \mathbb{E}(C,A)$ and $\delta' = \underline{\mathbf{S}'} \in \mathbb{E}(C',A')$.

Proof. Since $\overline{v_{A,A'}}$ is an isomorphism in $h\mathscr{C}$, (1) is a formal consequence of the functoriality of \mathbb{E} . Dually for (2).

The former part of (3) is immediate from (1) and (2). The latter part follows from Corollary 3.32, since it implies $(v_{A,A'})_*(\mathbf{S} \coprod \mathbf{S'}) = (v_{C,C'})^*(\mathbf{S} \times \mathbf{S'})$ by Lemma 4.7.

Remark 4.13. In the rest we will identify $A \oplus A' = A \coprod A'$ with $A \times A'$ through the isomorphism $\overline{v_{A,A'}} \colon A \oplus A' \xrightarrow{\cong} A \times A'$ in $h\mathscr{C}$.

Also we put $\delta \oplus \delta' = \delta \coprod \delta' \in \mathbb{E}(C \oplus C', A \oplus A')$. By the above identification and Proposition 4.12 (3), we will often identify $\delta \oplus \delta'$ with $\delta \times \delta' = \vartheta(\delta \oplus \delta')$. This agrees with the one in (iii) of Definition 4.2 via the above identification.

Lemma 4.14. We have the following.

(1) For any $\delta \in \mathbb{E}(C, A)$, we have

$$(4.4) (\nabla_A)_*(\delta \oplus \delta) = (\nabla_C)^*\delta \text{ and } (\Delta_A)_*\delta = (\Delta_C)^*(\delta \oplus \delta).$$

(2) For any $\delta_k \in \mathbb{E}(C_k, A_k)$, $\mathfrak{a}_k \in (h\mathscr{C})(A_k, A'_k)$, $\mathfrak{c}_k \in (h\mathscr{C})(C_k, C'_k)$ (k = 1, 2), we have

$$(\mathfrak{a}_1\oplus\mathfrak{a}_2)_*(\delta_1\oplus\delta_2)=\mathfrak{a}_{1*}\delta_1\oplus\mathfrak{a}_{2*}\delta_2\ \ \mathrm{and}\ \ (\mathfrak{c}_1\oplus\mathfrak{c}_2)^*(\delta_1\oplus\delta_2)=\mathfrak{c}_1^*\delta_1\oplus\mathfrak{c}_2^*\delta_2.$$

Proof. (1) This follows from Lemma 4.7 applied to the morphisms in Example 3.30.

(2) Take representatives $\delta_k = \underline{\mathbf{S}_k}$, $\mathfrak{a}_k = \overline{a_k}$ (k = 1, 2). By the definition of $a_{k*}\mathbf{S}_k$, there are morphisms of exact sequences

$$a_k(\mathbf{D}_k)_{1_{C_k}}: A_k(\mathbf{S}_k)_{C_k} \to A'_k(a_{k*}\mathbf{S}_k)_{C_k}$$

for k=1,2. By Proposition 3.31, we have a morphism ${}_{v}\mathbf{C}_{v'}\colon \mathbf{S}_1 \coprod \mathbf{S}_2 \to \mathbf{S}_1 \times \mathbf{S}_2$ in which, under the identifications in Remark 4.13, the morphisms $v \in \mathscr{C}_1(A_1 \coprod A_2, A'_1 \times A'_2)$ and $v' \in \mathscr{C}_1(C_1 \coprod C_2, C_1 \times C_2)$ induce

$$\overline{v} = \overline{a_1} \oplus \overline{a_2} = \mathfrak{a}_1 \oplus \mathfrak{a}_2 \in (h\mathscr{C})(A_1 \oplus A_2, A_1' \oplus A_2'),$$

$$\overline{v'} = \overline{1_{C_1}} \oplus \overline{1_{C_2}} = \operatorname{id}_{C_1 \oplus C_2} \in (h\mathscr{C})(C_1 \oplus C_2, C_1 \oplus C_2)$$

in the homotopy category, thus implies

$$(\mathfrak{a}_1 \oplus \mathfrak{a}_2)_*(\delta_1 \oplus \delta_2) = \underline{\upsilon_*(\mathbf{S}_1 \coprod \mathbf{S}_2)} = \underline{\upsilon'^*(a_{1*}\mathbf{S}_1 \times a_{2*}\mathbf{S}_2)} = \mathfrak{a}_{1*}\delta_1 \oplus \mathfrak{a}_{2*}\delta_2$$
 by Lemma 4.7. Similarly for $(\mathfrak{c}_1 \oplus \mathfrak{c}_2)^*(\delta_1 \oplus \delta_2) = \mathfrak{c}_1^*\delta_1 \oplus \mathfrak{c}_2^*\delta_2$.

Definition 4.15. For any $\delta_1, \delta_2 \in \mathbb{E}(C, A)$, define $\delta_1 + \delta_2 \in \mathbb{E}(C, A)$ by

$$\delta_1 + \delta_2 = (\nabla_A)_* (\Delta_C)^* (\delta_1 \oplus \delta_2).$$

Obviously this operation is commutative.

Proposition 4.16. Let $\delta \in \mathbb{E}(C, A)$ be any element.

- (1) For any pair of endomorphisms $\mathfrak{a}, \mathfrak{a}' \in \operatorname{End}_{h\mathscr{C}}(A)$ of A in $h\mathscr{C}$, we have $(\mathfrak{a} + \mathfrak{a}')_* \delta = \mathfrak{a}_* \delta + \mathfrak{a}'_* \delta$.
- (2) For any pair of endomorphisms $\mathfrak{c}, \mathfrak{c}' \in \operatorname{End}_{h\mathscr{C}}(C)$ of C in $h\mathscr{C}$, we have $(\mathfrak{c} + \mathfrak{c}')^*\delta = \mathfrak{c}^*\delta + \mathfrak{c}'^*\delta$.

Proof. (1) By Lemma 4.14, we have

$$\mathfrak{a}_*\delta + \mathfrak{a}'_*\delta = (\nabla_A)_*(\Delta_C)^*(\mathfrak{a}_*\delta \oplus \mathfrak{a}'_*\delta) = (\nabla_A)_*(\Delta_C)^*(\mathfrak{a} \oplus \mathfrak{a}')_*(\delta \oplus \delta)
= (\nabla_A)_*(\mathfrak{a} \oplus \mathfrak{a}')_*(\Delta_C)^*(\delta \oplus \delta) = (\nabla_A)_*(\mathfrak{a} \oplus \mathfrak{a}')_*(\Delta_A)_*\delta
= (\nabla_A \circ (\mathfrak{a} \oplus \mathfrak{a}') \circ \Delta_A)_*\delta = (\mathfrak{a} + \mathfrak{a}')_*\delta.$$

Similarly for
$$(2)$$
.

In the following corollary, remark that $h\mathscr{C}$ is an additive category by definition, hence there are $-\mathrm{id}_A \in (h\mathscr{C})(A,A)$ and $-\mathrm{id}_C \in (h\mathscr{C})(C,C)$.

Corollary 4.17. For any $\delta \in \mathbb{E}(C, A)$, the following holds in $\mathbb{E}(C, A)$.

- (1) $\delta + {}_A 0_C = \delta$.
- (2) $\delta + (-\mathrm{id}_C)^* \delta = \delta + (-\mathrm{id}_A)_* \delta = {}_A 0_C.$

In particular it follows that $(-id_C)^*\delta = (-id_A)_*\delta$ gives the additive inverse of δ .

Proof. This follows from Remark 4.11 and Proposition 4.16.

Proposition 4.18. The following holds.

(1) For any $A_1, A_2, C \in \mathscr{C}_0$,

(4.5)
$$([1 \ 0]_*, [0 \ 1]_*) \colon \mathbb{E}(C, A_1 \oplus A_2) \to \mathbb{E}(C, A_1) \times \mathbb{E}(C, A_2)$$

is bijective. (2) For any $A, C_1, C_2 \in \mathscr{C}_0$,

$$(\begin{bmatrix} 1\\0 \end{bmatrix}^*, \begin{bmatrix} 0\\1 \end{bmatrix}^*) \colon \mathbb{E}(C_1 \oplus C_2, A) \to \mathbb{E}(C_1, A) \times \mathbb{E}(C_2, A)$$

is bijective.

Proof. (1) Let us denote the map (4.5) by F for simplicity. It suffices to show that

$$G \colon \mathbb{E}(C, A_1) \times \mathbb{E}(C, A_2) \to \mathbb{E}(C, A_1 \oplus A_2) \; ; \; (\delta_1, \delta_2) \mapsto \begin{bmatrix} 1 \\ 0 \end{bmatrix}_* \delta_1 + \begin{bmatrix} 0 \\ 1 \end{bmatrix}_* \delta_2$$

gives its inverse. $G \circ F = \text{id}$ follows from Proposition 4.16, since we have

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix}_{*} \begin{bmatrix} 1 \\ 0 \end{bmatrix}_{*} \delta + \begin{bmatrix} 0 \\ 1 \end{bmatrix}_{*} \begin{bmatrix} 0 \\ 1 \end{bmatrix}_{*} \delta = (\begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix})_{*} \delta = (\operatorname{id}_{A_{1} \oplus A_{2}})_{*} \delta = \delta$$

for any $\delta \in \mathbb{E}(C, A_1 \oplus A_2)$.

For any $(\delta_1, \delta_2) \in \mathbb{E}(C, A_1) \times \mathbb{E}(C, A_2)$, we have

$$[1 \ 0]_*(\delta_1 \oplus \delta_2) = [1 \ 0]^*\delta_1$$

by Proposition 3.31 applied to $I: \mathbf{S}_1 \to \mathbf{S}_1$ and $\mathbf{S}_2 \to \mathbf{O}$. Thus by Lemma 4.14, we obtain

$$[1 \ 0]_*(\begin{bmatrix} 1 \ 0 \end{bmatrix}_* \delta_1 + \begin{bmatrix} 0 \ 1 \end{bmatrix}_* \delta_2) = [1 \ 0]_*(\nabla_{A_1 \oplus A_2})_*(\Delta_C)^*(\begin{bmatrix} 1 \ 0 \end{bmatrix}_* \delta_1 \oplus \begin{bmatrix} 0 \ 1 \end{bmatrix}_* \delta_2)$$

$$= [1 \ 0]_*(\nabla_{A_1 \oplus A_2})_*(\Delta_C)^*(\begin{bmatrix} 1 \ 0 \end{bmatrix} \oplus \begin{bmatrix} 0 \ 1 \end{bmatrix})_*(\delta_1 \oplus \delta_2)$$

$$= (\Delta_C)^*([1 \ 0] \circ \nabla_{A_1 \oplus A_2} \circ (\begin{bmatrix} 1 \ 0 \end{bmatrix} \oplus \begin{bmatrix} 0 \ 1 \end{bmatrix})_*(\delta_1 \oplus \delta_2)$$

$$= (\Delta_C)^*[1 \ 0]_*(\delta_1 \oplus \delta_2) = (\Delta_C)^*[1 \ 0]^*\delta_1 = \delta_1.$$

Similarly we have $[0\ 1]_*(\begin{bmatrix}1\\0\end{bmatrix}_*\delta_1 + \begin{bmatrix}0\\1\end{bmatrix}_*\delta_2) = \delta_2$, and thus $F \circ G((\delta_1, \delta_2)) = (\delta_1, \delta_2)$.

Corollary 4.19. $\mathbb{E}(C, A)$ has a natural structure of additive group, whose addition is given by Definition 4.15, with the zero element ${}_{A}0_{C}$. Moreover,

$$\mathbb{E} \colon (h\mathscr{C})^{\mathrm{op}} \times h\mathscr{C} \to Ab$$

becomes a biadditive functor.

Proof. This is a formal consequence of the functoriality of \mathbb{E} : $(h\mathscr{C})^{\mathrm{op}} \times h\mathscr{C} \to Set$, Corollary 4.17 and Proposition 4.18.

4.3. External triangulation of $h\mathscr{C}$.

Definition 4.20. Let $A, C \in \mathcal{C}_0$ be any pair of objects. For each $\delta \in \mathbb{E}(C, A)$, take any

$$\mathbf{S} = i \bigvee_{j}^{A} \xrightarrow{x} B \\ O \xrightarrow{j} C$$

such that $\delta = \underline{\mathbf{S}}$, and put $\mathfrak{s}(\delta) = [A \xrightarrow{\overline{x}} B \xrightarrow{\overline{y}} C]$, where the equivalence class of the sequences is the one in Definition 4.1. Well-definedness follows from Proposition 3.26.

Lemma 4.21. \mathfrak{s} is an additive realization of \mathbb{E} .

Proof. Let us confirm the conditions (i),(ii),(iii) in Definition 4.2.

- (i) This follows from Lemma 4.7.
- (ii) Let $A, C \in \mathcal{C}_0$ be any pair of objects. Since ${}_A 0_C = {}_A \underline{\mathbf{N}'}_C$, we have

$$\mathfrak{s}({}_{A}0{}_{C}) = [A \xrightarrow{\overline{\imath}_{A}} A \coprod C \xrightarrow{\overline{\jmath}_{C}} C] = [A \xrightarrow{\left[\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}\right]} A \coprod C \xrightarrow{\left[\begin{smallmatrix} 0 & 1 \end{smallmatrix}\right]} C].$$

(iii) Let $\delta = \underline{\mathbf{S}} \in \mathbb{E}(C, A), \delta' = \underline{\mathbf{S}'} \in \mathbb{E}(C', A')$ be any pair of elements, with

$$\mathbf{S} = i \downarrow \begin{matrix} A \xrightarrow{x} B \\ \downarrow y \quad \text{and} \quad \mathbf{S}' = i' \downarrow \begin{matrix} x' \\ \downarrow y' \\ O' \xrightarrow{j'} C' \end{matrix}$$

By definition we have

$$\mathfrak{s}(\delta) = [A \xrightarrow{\mathfrak{r}} B \xrightarrow{\mathfrak{y}} C], \quad \mathfrak{s}(\delta') = [A' \xrightarrow{\mathfrak{r}'} B' \xrightarrow{\mathfrak{y}'} C'],$$

and $\delta \oplus \delta' = \underline{\mathbf{S} \coprod \mathbf{S'}}$ as in Remark 4.13. It follows

$$\mathfrak{s}(\delta\oplus\delta')=[A\amalg A'\xrightarrow{\overline{x\amalg x'}}B\amalg B'\xrightarrow{\overline{y\amalg y'}}C\amalg C']=[A\oplus A'\xrightarrow{\overline{x}\oplus\overline{x'}}B\oplus B'\xrightarrow{\overline{y}\oplus\overline{y'}}C\oplus C'].$$

Theorem 4.22. $(h\mathscr{C}, \mathbb{E}, \mathfrak{s})$ is an extriangulated category.

Proof. (ET1) and (ET2) are shown in Corollary 4.19 and Lemma 4.21, respectively. (ET3) follows from Proposition 3.7 and Lemma 4.7. Dually for (ET3) op . It remains to show (ET4) and (ET4) op .

Since (ET4)^{op} can be shown dually, it is enough to show (ET4). Let $\delta = \underline{\mathbf{S}} \in \mathbb{E}(D,A)$ and $\delta' = \underline{\mathbf{S}'} \in \mathbb{E}(F,B)$ be any pair of elements given by the following exact sequences.

$$\mathbf{S} = i \bigvee_{j}^{A} \underbrace{\begin{matrix} f \\ \xi \end{matrix} }_{j}^{B} D \qquad B \underbrace{\begin{matrix} g \\ \zeta \end{matrix} }_{j'}^{C} C \\ g' \underbrace{\begin{matrix} f' \\ z' \end{matrix} }_{j'} F$$

By definition, we have $\mathfrak{s}(\delta) = [A \xrightarrow{\overline{f}} B \xrightarrow{\overline{f'}} D]$ and $\mathfrak{s}(\delta') = [B \xrightarrow{\overline{g}} C \xrightarrow{\overline{g'}} F]$. Since f, g are ingressive, we may take a 2-simplex

$$A \xrightarrow{f} B$$

in which h is ingressive by the definition of exact quasi-category. As in [B1, Remark 3.2], h should appear as the fiber of some exact sequence as below.

$$\mathbf{S}'' = i'' \bigvee_{j''}^{A} \bigvee_{j''}^{h'} \bigvee_{k}^{C}$$

Put $\delta'' = \underline{\mathbf{S}''}$. We have $\mathfrak{s}(\delta'') = [A \xrightarrow{\overline{h}} C \xrightarrow{\overline{h'}} E]$ by definition. Applying Proposition 3.7 to \mathbf{S}, \mathbf{S}'' and

$$\mathbf{T} = (s_0(h)\backslash \chi) = \begin{array}{c} A \xrightarrow{f} B \\ \downarrow A \xrightarrow{s_0(h)} Q \end{array}$$

we obtain a morphism of the form

$${}_{1_{A}}\mathbf{C}_{d} = \begin{bmatrix} s_{0}(\xi'') & \mathcal{Y}_{f} & \mathcal{Z}_{f} \\ s_{0}(\eta'') & \mathcal{Y}_{b} & \mathcal{Z}_{b} \end{bmatrix} = \begin{bmatrix} A & \xrightarrow{f} & B & f' \\ \downarrow i & O & \downarrow & \downarrow \\ \downarrow i & O & \downarrow g & \downarrow d \\ A & \xrightarrow{h} & \downarrow C & h' \\ \downarrow i'' & O'' & \xrightarrow{j''} & E \end{bmatrix} : \mathbf{S} \to \mathbf{S}''.$$

By the dual of Lemma 3.11 (1), the square

$$\mathbf{P} = \left(\mathbf{C}|_{\Delta^{1} \times \{1\} \times \Delta^{1}}\right)^{t} = \begin{cases} B \xrightarrow{g} C \\ f' \downarrow & \downarrow h' \\ D \xrightarrow{d} E \end{cases}$$

is a push-out. By Lemma 3.10, using this push-out we obtain a morphism of exact sequences

$$f'\mathbf{D}_{1_{F}} = \begin{bmatrix} \mathcal{X}'_{\mathbf{f}} & \mathcal{Y}'_{\mathbf{f}} & s_{2}(\xi') \\ \mathcal{X}'_{\mathbf{b}} & s_{1}(\eta') & s_{2}(\eta') \end{bmatrix} = \begin{cases} B \xrightarrow{g} C & g' \\ \downarrow & \downarrow & \downarrow \\ I_{O'} & \downarrow & \downarrow \\ D \xrightarrow{d} & \downarrow & E & e \\ \downarrow & \downarrow & \downarrow \\ O' & \xrightarrow{j'} & F \end{cases} : \mathbf{S}' \to f'_{*}\mathbf{S}'.$$

By the construction so far, we have a commutative diagram in $h\mathscr{C}$

$$\begin{array}{c|c}
A & \overline{f} \\
 & \nearrow B & \overline{f'} \\
 & & \searrow \overline{g} \\
 & & \searrow \overline{g} \\
 & & & \searrow \overline{g} \\
 & & & \searrow \overline{g'} \\
 & & & & \searrow \overline{g'} \\
 & & & & & & F \\
\hline$$

satisfying $\mathfrak{s}(\overline{f'}_*\delta') = [D \xrightarrow{\overline{d}} E \xrightarrow{\overline{e}} F]$. Besides, the existence of \mathbf{C} shows $\delta = \overline{d}^*\delta''$ by Lemma 4.7.

It remains to show $\overline{f}_*\delta'' = \overline{e}^*\delta'$. By Lemma 4.7, it suffices to show the existence of a morphism ${}_f\mathbf{C}'_e \colon \mathbf{S}'' \to \mathbf{S}'$. Remark that we have cubes

$$\begin{bmatrix} \mathcal{Y}_{f} & s_{0}(\xi'') & s_{0}(\eta'') \\ \mathcal{Z}_{f} & \mathcal{Z}_{b} & \mathcal{Y}_{b} \end{bmatrix} = \begin{cases} A \xrightarrow{1_{A}} A & i'' \\ \downarrow & O \xrightarrow{l} O'' \\ \downarrow & J \\ \downarrow$$

as transpositions of C and D. As in Proposition 2.5, we obtain a map $\Delta^1 \times \Delta^1 \times \Delta^2 \to \mathscr{C}$

$$A \xrightarrow{1_A} A \xrightarrow{i''} O \xrightarrow{j} O''$$

$$A \xrightarrow{j} A \xrightarrow{i''} O''$$

$$A \xrightarrow{j} O'' \xrightarrow{j} A \xrightarrow{j} O''$$

$$A \xrightarrow{j} O' \xrightarrow{j} O' \xrightarrow{j} A \xrightarrow{j} O''$$

$$A \xrightarrow{j} A \xrightarrow{j'} O''$$

$$A \xrightarrow{j} O'' O' \xrightarrow{j'} A \xrightarrow{j} O''$$

$$A \xrightarrow{j'} O'' \xrightarrow{j'} A \xrightarrow{j''} O''$$

$$A \xrightarrow{j'} O'' \xrightarrow{j'} O''$$

$$A \xrightarrow{j'} O \xrightarrow{j'} O' \xrightarrow{j'} O'$$

$$A \xrightarrow{j'} O \xrightarrow{j'} O' \xrightarrow{j'} O'$$

compatibly with these cubes. Especially it contains 4-simplices

$$\Theta =
\begin{array}{c}
B \\
\downarrow \\
C \\
h'
\end{array}$$

$$E =
\begin{array}{c}
O \\
\downarrow \\
D \\
\downarrow \\
O'
\end{array}$$

$$A \\
D \\
\downarrow \\
O'$$

$$A \\
F \\
F \\
O''$$

$$A \\
F \\
O''$$

$$A \\
F \\
O''$$

$$O'' \\
O'' \\
O''$$

$$E =
\begin{array}{c}
O \\
\downarrow \\
O \\
O''
\end{array}$$

such that $\mathbf{S}'' = (d_1 d_4(\Psi) \setminus d_1 d_4(\Theta)), \mathbf{S}' = (d_0 d_2(\Phi) \setminus d_0 d_3(\Theta))$ and

$$d_2d_3(\Theta) = d_2d_2(\Phi), \ d_1d_1(\Theta) = d_1d_1(\Psi), \ d_1d_2(\Xi) = d_1d_2(\Phi).$$

Applying Lemma 3.8 to $d_2(\Xi)$ and $d_3(\Psi)$, we obtain a 3-simplex μ such that $d_1(\mu) = d_1d_2(\Xi)$ and $d_2(\mu) = d_1d_3(\Psi)$. If we put

$$\mathbf{C}' = \left[\begin{array}{c|c} d_3(\Theta) & s_1 d_1 d_3(\Theta) & d_1(\Theta) \\ d_2(\Phi) & \mu & d_1(\Psi) \end{array} \right],$$

then this give a morphism ${}_{f}\mathbf{C}'_{e} \colon \mathbf{S}'' \to \mathbf{S}'$ as desired.

Definition 4.23. Inspired from [S], we call *topological* any extriangulated category which is equivalent to the homotopy category of an exact quasi-category.

Lemma 4.24. Let \mathscr{C} be an exact quasi-category, and let \mathscr{D} be a full subcategory of \mathscr{C} closed by homotopy equivalences of objects in \mathscr{C} . Assume that \mathscr{D} is extension-closed in \mathscr{C} , meaning that, \mathscr{D} contains the zero objects of \mathscr{C} and that, for any exact sequence ${}_{A}\mathbf{S}_{C}$ in \mathscr{C} with $A, C \in \mathscr{D}_{0}$, the middle term B also belongs to \mathscr{D}_{0} . Define $(\mathscr{D}_{\dagger})_{1}$ by considering those morphisms in \mathscr{D} that are ingressive in \mathscr{C} and for which some cofiber belongs to \mathscr{D} . (For $n \geq 2$, we define $(\mathscr{D}_{\dagger})_{n} \subseteq (\mathscr{C}_{\dagger})_{n} \cap \mathscr{D}_{n}$ to be the subset consisting of n-simplices whose edges are in $(\mathscr{D}_{\dagger})_{1}$. The face and degeneracy

maps in \mathcal{Q}_{\dagger} are restrictions of those in \mathcal{Q} .) Define \mathcal{Q}^{\dagger} dually. Then $(\mathcal{Q}, \mathcal{Q}_{\dagger}, \mathcal{Q}^{\dagger})$ is an exact quasi-category.

Proof. Stability under (split) extensions implies that \mathcal{D} is additive. To see that \mathcal{D}_{\dagger} is a subcategory, let



any 2-simplex in \mathscr{D} in which f, g belong to $(\mathscr{D}_{\dagger})_1$. Consider the diagram of ambigressive push-outs

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$O \longrightarrow Z' \longrightarrow Y'$$

$$\downarrow \qquad \qquad \downarrow$$

$$O' \longrightarrow X'$$

where Z' and X' belong to \mathscr{D} by assumption. The bottom square is an exact sequence, hence Y' also belongs to \mathscr{D} . The outer upper rectangle shows that $X \stackrel{h}{\longrightarrow} Z$ belongs to \mathscr{D}_{\dagger} . Dually, \mathscr{D}^{\dagger} is a subcategory of \mathscr{D} . The remaining axioms readily follow from those for \mathscr{C} .

Proposition 4.25. Every extension-closed full subcategory, containing zero, of a topological extriangulated category is topological.

Proof. Let \mathscr{C} be an exact quasi-category and let \mathcal{D} be a full subcategory of $h\mathscr{C}$ that we assume extension-closed. By [NP, Remark 2.18], \mathcal{D} inherits an extriangulated structure from that of $h\mathscr{C}$. Define \mathscr{D} to be the full subcategory of \mathscr{C} given by \mathcal{D} . More explicitly, \mathscr{D} is given by the pull-back, in the category of simplicial sets:

Then \mathscr{D} is extension-closed in \mathscr{C} . Indeed, any exact sequence $X \to Y \to Z$ in \mathscr{C} with $X, Z \in \mathscr{D}_0$ induce, by Theorem 4.22, an extriangle $X \rightarrowtail Y \twoheadrightarrow Z \dashrightarrow$ in $h\mathscr{C}$. By assumption, we have $Y \in \mathcal{D}$. We can thus apply Lemma 4.24: \mathscr{D} is an exact quasi-category. The claim follows since $h\mathscr{D} = \mathcal{D}$.

We plan to investigate the following

Question 4.26. Is every topological extriangulated category equivalent to some extension-closed full subcategory of a triangulated category?

Our next result is an elementary tool that might help answering Question 4.26. Let $F \colon \mathscr{C} \to \mathscr{D}$ be an exact functor of exact quasi-categories [B1, Definition 4.1]. In other words, F is a map of simplicial sets that preserves zero objects, ingressive morphisms, egressive morphisms, push-outs along ingressive morphisms and pull-backs along egressive morphisms.

Because F is a map of simplicial sets, it commutes to faces and degeneracies and thus induces a functor $hF: h\mathscr{C} \to h\mathscr{D}$ which coincide with F on objects. It is defined on morphisms by $hF \overline{a} = \overline{Fa}$.

Definition 4.27 ([B-TS, Definition 2.31]). A functor $G: (\mathcal{C}, \mathbb{E}, \mathfrak{s}) \to (\mathcal{D}, \mathbb{E}', \mathfrak{s}')$ between extriangulated categories is *exact* (or *extriangulated*) if it is additive, and if there is a natural transformation $\Gamma: \mathbb{E} \Rightarrow \mathbb{E}'(G^{\mathrm{op}}-, G-)$ such that, for any \mathfrak{s} -triangle $A \xrightarrow{i} B \xrightarrow{p} C \xrightarrow{-i}$ in \mathcal{C} , its image $GA \xrightarrow{Gi} GB \xrightarrow{Gp} GC \xrightarrow{\Gamma_{C,A}(\delta)}$ is an \mathfrak{s}' -triangle in \mathcal{D} .

Proposition 4.28. Let $F: \mathcal{C} \to \mathcal{D}$ be an exact functor of exact quasi-categories. Then $hF: h\mathcal{C} \to h\mathcal{D}$ is an exact functor of extriangulated categories.

Proof. Because F preserves push-outs of ingressive morphisms, it preserves coproducts, and similarly for products. It follows that hF is additive. Define Γ as follows: For any $A, C \in \mathscr{C}_0$, and any $\delta = \underline{\mathbf{S}} \in \mathbb{E}(C,A)$, let $\Gamma_{C,A}(\delta) = \underline{F}\underline{\mathbf{S}}$. The map $\Gamma_{C,A}$ is well-defined since if ${}_a\mathbf{C}_c$ is a cube in \mathscr{C} from \mathbf{S} to \mathbf{S}' with $\overline{a} = \mathrm{id}_A$ and $\overline{c} = \mathrm{id}_C$ in $h\mathscr{C}$, then ${}_{Fa}\mathbf{C}'_{Fc} = F\mathbf{C}$ is a cube in \mathscr{D} from $F\mathbf{S}$ to $F\mathbf{S}'$ with $\overline{Fa} = \mathrm{id}_{FA}$ and $\overline{Fc} = \mathrm{id}_{FC}$ in $h\mathscr{D}$. It remains to show that $\Gamma = (\Gamma_{C,A})$ is natural. Let $\delta = \underline{\mathbf{S}} \in \mathbb{E}(C,A)$ and let $a \in \mathscr{C}_1(A,A')$. Consider a cube

$$\mathbf{D} = \begin{bmatrix} A & \xrightarrow{x} & B & y \\ & i & O & \xrightarrow{j} & C \\ & A' & \xrightarrow{x_0} & \downarrow b_0 & \downarrow b_0 \\ & A' & \xrightarrow{x_0} & \downarrow b_0 & \downarrow b_0 \\ & & & \downarrow b_0 & \downarrow b_0 \\ & \downarrow b_0 & \downarrow b_0 \\$$

defining $a_*\mathbf{S}$ as in Lemma 3.10. In particular, the back square is a push-out, with x ingressive. Since F preserves push-outs along ingressive morphisms, the cube $F\mathbf{D}$ witnesses the fact that $F(a_*\mathbf{S}) = (Fa)_*F\mathbf{S}$. Dually, for any $C' \stackrel{c}{\longrightarrow} C$, we have $F(c^*\mathbf{S}) = (Fc)^*F\mathbf{S}$. Whence $\Gamma_{C',A'} \circ \overline{c}^*\overline{a}_* = (hF\overline{c})^*(hF\overline{a})_* \circ \Gamma_{C,A}$.

4.4. Compatibility with the triangulation in stable case. In this subsection, let $\mathscr C$ be a stable quasi-category, which is a particular case of an exact quasi-category, as in [B1, Example 3.3]. In this case it is known that the homotopy category $h\mathscr C$ can be equipped with a structure of a triangulated category ([L2]). Let us show that the structure of an extriangulated category on $h\mathscr C$ obtained in Theorem 4.22 is compatible with it.

First let us briefly review the construction of the shift functor. For any object $A \in \mathcal{C}_0$, we choose an exact sequence

$$\mathbf{U}_{A} = (\eta_{A} \backslash \zeta_{A}) = \bigvee_{O'_{A} \to \Sigma A} \mathcal{O}_{A} \qquad \left(\text{respectively, } \mathbf{V}_{A} = \bigvee_{O''_{A} \to A} \mathcal{O}''_{A}\right)$$

in which O_A, O'_A (resp. O''_A, O'''_A) are zero objects.

If we put $\mu_A = \underline{\mathbf{U}_A} \in \mathbb{E}(\Sigma A, A)$ (resp. $\nu_A = \underline{\mathbf{V}_A} \in \mathbb{E}(A, \Omega A)$), then we have an \mathfrak{s} -triangle

$$(4.6) A \to 0 \to \Sigma A \xrightarrow{\mu_A}, (resp. \ \Omega A \to 0 \to A \xrightarrow{\nu_A}).$$

Especially, as in [NP, Proposition 3.3] we obtain natural isomorphisms of functors induced by the Yoneda's lemma

$$(\mu_A)_{\sharp} \colon (h\mathscr{C})(-,\Sigma A) \stackrel{\cong}{\Longrightarrow} \mathbb{E}(-,A)$$

which assigns $(\mu_A)_{\sharp}(\mathfrak{c}) = \mathfrak{c}^*(\mu_A)$ to each $\mathfrak{c} \in (h\mathscr{C})(C, \Sigma A)$. Thus for any $\mathfrak{a} \in (h\mathscr{C})(A, A')$, there exists a morphism $\Sigma \mathfrak{a} \in (h\mathscr{C})(\Sigma A, \Sigma A')$ uniquely so that

$$(h\mathscr{C})(-,\Sigma A) \xrightarrow{(\mu_A)_{\sharp}} \mathbb{E}(-,A)$$

$$(\Sigma \mathfrak{a}) \circ - \bigvee \qquad \Diamond \qquad \bigvee \mathfrak{a}_*$$

$$(h\mathscr{C})(-,\Sigma A') \xrightarrow{(\mu_{A'})_{\sharp}} \mathbb{E}(-,A')$$

becomes commutative. Then the correspondence

- for any object $A \in h\mathscr{C}$, associate $\Sigma A \in h\mathscr{C}$ using the chosen \mathbf{U}_A ,
- for any morphism $\mathfrak{a} \in (h\mathscr{C})(A, A')$, associate the morphism $\Sigma \mathfrak{a} \in (h\mathscr{C})(\Sigma A, \Sigma A')$ by the above

gives the endofunctor $\Sigma \colon h\mathscr{C} \to h\mathscr{C}$. By Lemma 4.7, morphism $\Sigma \overline{a}$ for $a \in \mathscr{C}_1(A, A')$ is equal to the one given by $\Sigma \overline{a} = \overline{s}$ where $s \in \mathscr{C}_1(\Sigma A, \Sigma A')$ admits some morphism ${}_a\mathbf{C}_s \colon \mathbf{U}_A \to \mathbf{U}_{A'}$ of exact sequences. Similarly for the functor $\Omega \colon h\mathscr{C} \to h\mathscr{C}$, which gives a quasi-inverse of Σ .

Using these structures, we can complete any morphism $\overline{z} \in (h\mathscr{C})(C, \Sigma A)$ into a triangle diagram of the form $A \to B \to C \xrightarrow{\overline{z}} \Sigma A$ compatibly with the external triangulation in the following way. Indeed, this is how to relate triangulations with external triangulations ([NP, Proposition 3.22]).

• For each $\overline{z} \in (h\mathscr{C})(C, \Sigma A)$, take

(4.7)
$$\mathfrak{s}((\mu_A)_{\sharp}(\overline{z})) = [A \xrightarrow{\overline{x}} B \xrightarrow{\overline{y}} C]$$
 to obtain $A \xrightarrow{\overline{x}} B \xrightarrow{\overline{y}} C \xrightarrow{\overline{z}} \Sigma A$.

Remark 4.29. In fact, the existence of \mathfrak{s} -triangles of the form (4.6) is equivalent to that the extriangulated category $(h\mathscr{C},\mathbb{E},\mathfrak{s})$ admits a compatible triangulation. Indeed, existence of such \mathfrak{s} -triangles is equivalent to that $(h\mathscr{C},\mathbb{E},\mathfrak{s})$ is Frobenius with trivial projective-injectives in the sense of [NP, Definition 7.1] and [ZZ, Subsection 3.3], and thus $(h\mathscr{C},\Sigma,\Delta)$ becomes a triangulated category by [NP, Corollary 7.6] or [ZZ, Example 3.15], in which Δ is defined to be the class of triangles isomorphic to those $A \xrightarrow{\overline{x}} B \xrightarrow{\overline{y}} C \xrightarrow{\overline{z}} \Sigma A$ obtained in the above way.

On the other hand, the distinguished triangles in the triangulation given in [L2] are those isomorphic to $A \xrightarrow{\overline{x'}} B' \xrightarrow{\overline{y'}} C \xrightarrow{\overline{z}} \Sigma A$ which fits in some rectangle

(4.8)
$$\mathbf{R} = \begin{bmatrix} \mathcal{X} \mid \mathcal{Y} \mid \mathcal{Z} \end{bmatrix} = \begin{pmatrix} A \xrightarrow{x'} B' \longrightarrow O_A \\ \downarrow & \downarrow \\ O'_A \longrightarrow C \xrightarrow{z} \Sigma A \end{pmatrix}$$

such that $\mathbf{R}_{\text{out}} = \mathbf{U}_A$ and that $\mathbf{R}_{\text{right}}$ is a pull-back. Thus to see the compatibility, it is enough to show the following.

Proposition 4.30. For any morphism $z \in \mathscr{C}_1(C, \Sigma A)$ and any rectangle (4.8) in which $\mathbf{R}_{\text{out}} = \mathbf{U}_A$ holds and $\mathbf{R}_{\text{right}}$ is a pull-back, we have $\mathfrak{s}((\mu_A)_{\sharp}(\overline{z})) = [A \xrightarrow{\overline{x'}} B' \xrightarrow{\overline{y'}} C]$.

Proof. In the rectangle \mathbf{R} , we see that $\mathbf{S}' = \mathbf{R}_{\text{left}}$ is an exact sequence, and satisfies $\mathfrak{s}(\underline{\mathbf{S}'}) = [A \xrightarrow{\overline{x'}} B' \xrightarrow{\overline{y'}} C]$ by definition. Moreover, \mathbf{R} gives a morphism of exact sequences

$${}_{1_A}\mathbf{C}_z = \left[\begin{array}{c|c} s_0(\zeta_A) & \mathcal{Z} & \mathcal{Y} \\ s_0(\eta_A) & s_1(\eta_A) & \mathcal{X} \end{array} \right] \colon \mathbf{S}' \to \mathbf{U}_A,$$
 which shows $(\mu_A)_{\sharp}(\overline{z}) = (\overline{z})^*\mathbf{U}_A = \underline{\mathbf{S}'}$ by Lemma 4.7.

References

- [B1] Barwick, C.: On exact ∞-categories and the theorem of the heart. Compos. Math. 151 (2015), no. 11, 2160–2186.
- [B2] Barwick, C.: On the algebraic K-theory of higher categories. J. Topol. 9 (2016), no. 1, 245–347.
- [B-TS] Bennett-Tennenhaus, R.; Shah, A.: Transport of structure in higher homological algebra. arXiv:2003.02254
- [C] Cisinski, D-C.: Higher categories and homotopical algebra. Cambridge Studies in Advanced Mathematics, 180. Cambridge University Press, Cambridge, 2019. xviii+430 pp.
- [DK] Dyckerhoff, T.; Kapranov, M.: Higher Segal spaces. Lecture Notes in Mathematics, 2244. Springer, Cham, 2019. xv+218 pp.
- $[HLN] \ \ Herschend, \ H.; \ Liu, \ L.; \ Nakaoka, \ H.: \ n-exangulated \ categories. \ arXiv:1709.06689v2.$
- [J] Joyal, A.: Notes on quasi-categories.
- [L1] Lurie, J.: Higher topos theory. Annals of Mathematics Studies, 170. Princeton University Press, Princeton, NJ, 2009. xviii+925 pp.
- [L2] Lurie, J.: Stable infinity categories. arXiv:math/0608228.
- [NP] Nakaoka, H.; Palu, Y.: Extriangulated categories, Hovey twin cotorsion pairs and model structures. Cah. Topol. Géom. Différ. Catég. 60 (2019), no. 2, 117–193.
- [S] Schwede, S.: Algebraic versus topological triangulated categories. In Cambridge Univ. Press, Cambridge, Triangulated categories, London Math. Soc. Lecture Note Ser. 375 (2010), 389-407.
- [ZZ] Zhou, P.; Zhu, B.: Triangulated quotient categories revisited. J. Algebra 502 (2018), 196– 232

Graduate School of Mathematics, Nagoya University, Furocho, Chikusaku, Nagoya 464-8602, Japan

E-mail address: nakaoka.hiroyuki@math.nagoya-u.ac.jp

LAMFA, UNIVERSITÉ DE PICARDIE JULES VERNE, 33 RUE SAINT-LEU, AMIENS, FRANCE E-mail address: yann.palu@u-picardie.fr
URL: http://www.lamfa.u-picardie.fr/palu/