

Casimir elements and Sugawara operators for Takiff algebras

A. I. Molev*

School of Mathematics and Statistics
University of Sydney, NSW 2006, Australia

Abstract

For every simple Lie algebra \mathfrak{g} we consider the associated Takiff algebra \mathfrak{g}_ℓ defined as the truncated polynomial current Lie algebra with coefficients in \mathfrak{g} . We use a matrix presentation of \mathfrak{g}_ℓ to give a uniform construction of algebraically independent generators of the center of the universal enveloping algebra $U(\mathfrak{g}_\ell)$. A similar matrix presentation for the affine Kac–Moody algebra $\widehat{\mathfrak{g}}_\ell$ is then used to prove an analogue of the Feigin–Frenkel theorem describing the center of the corresponding affine vertex algebra at the critical level. The proof relies on an explicit construction of a complete set of Segal–Sugawara vectors for the Lie algebra \mathfrak{g}_ℓ .

1 Introduction

For each simple finite-dimensional Lie algebra \mathfrak{g} over \mathbb{C} and any positive integer ℓ consider the truncated polynomial current Lie algebra \mathfrak{g}_ℓ which is defined as the quotient of $\mathfrak{g} \otimes \mathbb{C}[v]$ by the ideal $\mathfrak{g} \otimes \mathbb{C}[v]v^{\ell+1}$. The Lie algebra \mathfrak{g}_ℓ is also called the *generalized* or *ℓ -th Takiff algebra* following the pioneering work [20], where such algebras were studied in the case $\ell = 1$. As shown in that paper, the subalgebra of \mathfrak{g}_ℓ -invariants in the symmetric algebra $S(\mathfrak{g}_\ell)$ is an algebra of polynomials. This result was extended by Raïs and Tauvel [18] to all values of ℓ . More recently, Macedo and Savage [12] proved its multi-parameter generalization, while Panyushev and Yakimova [17] showed that this generalization remains valid for a wide class of Lie algebras \mathfrak{g} beyond simple Lie algebras.

The results of Raïs and Tauvel were used by Geoffriau [8] to describe properties of an analogue of the Harish-Chandra homomorphism for the center of the universal enveloping algebra $U(\mathfrak{g}_\ell)$. Explicit generators of the center in type A were given in [13]. In this case the ℓ -th Takiff algebra associated with \mathfrak{gl}_n coincides with the centralizer of a certain nilpotent element e in $\mathfrak{gl}_{n(\ell+1)}$; namely, e is the direct sum of n Jordan blocks of size $\ell + 1$. The construction of central elements was extended by Brown and Brundan [4] to arbitrary nilpotents.

Here we give a uniform explicit construction of algebraically independent generators of the center of $U(\mathfrak{g}_\ell)$ for all simple Lie algebras \mathfrak{g} and all $\ell \geq 1$.

*alexander.molev@sydney.edu.au

Then we equip \mathfrak{g}_ℓ with an invariant symmetric bilinear form by extending a standard normalized Killing form on \mathfrak{g} . The corresponding affine Kac–Moody algebra $\widehat{\mathfrak{g}}_\ell$ is defined as a central extension of the Lie algebra of Laurent polynomials $\mathfrak{g}_\ell[t, t^{-1}]$. The vacuum module over $\widehat{\mathfrak{g}}_\ell$ is a vertex algebra whose center $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ is a commutative associative algebra. In the case $\ell = 0$ the structure of the center $\mathfrak{z}(\widehat{\mathfrak{g}})$ at the critical level was described by a celebrated theorem of Feigin and Frenkel [6] (see also [7]), which states that $\mathfrak{z}(\widehat{\mathfrak{g}})$ is an algebra of polynomials in infinitely many variables. We show that this property is shared by the center $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ for all $\ell \geq 1$.

Our arguments rely on matrix presentations of the Lie algebras \mathfrak{g}_ℓ and $\widehat{\mathfrak{g}}_\ell$. Such presentations of the classical Lie algebras and the exceptional Lie algebra of type G_2 played a key role in the constructions of generators of the Feigin–Frenkel center $\mathfrak{z}(\widehat{\mathfrak{g}})$ in [14] and [16]; see also [22] for a different approach. A recent work by Wendlandt [21] provides a significant extension of the matrix techniques by giving a presentation of $U(\mathfrak{g})$ for any simple Lie algebra \mathfrak{g} associated with its arbitrary faithful representation. We recall some results from that paper below as they will be needed for our calculations. To show that our central elements are free generators of $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ we use the classical limit and employ the Macedo–Savage theorem [12] in the particular case of ‘double’ Takiff algebras.

An analogue of the Feigin–Frenkel theorem for Takiff algebras in type A was already proved by Arakawa and Premet [1] as a particular case of a more general theorem describing the centers at the critical level of the affine vertex algebras associated with centralizers of nilpotent elements in simple Lie algebras. Explicit generators of the center in type A were produced in [15].

As in [1] and [15], our generators of the center $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ can be used to produce generators of quantum shift of argument subalgebras of $U(\mathfrak{g}_\ell)$. We expect that under certain regularity conditions they will be ‘quantizations’ of the Mishchenko–Fomenko subalgebras of $S(\mathfrak{g}_\ell)$ thus yielding a solution of Vinberg’s quantization problem for Takiff algebras.

2 Matrix presentations

We start by recalling some standard tensor product notation. Any $N \times N$ matrix $X = [X_{ij}]$ with entries in an associative algebra \mathcal{A} will be regarded as the element

$$X = \sum_{i,j=1}^N X_{ij} \otimes e_{ij} \in \mathcal{A} \otimes \text{End } \mathbb{C}^N, \quad (2.1)$$

where the $e_{ij} \in \text{End } \mathbb{C}^N$ denote the standard matrix units. We will need tensor product algebras of the form $\mathcal{A} \otimes \text{End } (\mathbb{C}^N)^{\otimes m}$. For any $a \in \{1, \dots, m\}$ we will denote by X_a the element (2.1) associated with the a -th copy of $\text{End } \mathbb{C}^N$ so that

$$X_a = \sum_{i,j=1}^N X_{ij} \otimes 1^{\otimes(a-1)} \otimes e_{ij} \otimes 1^{\otimes(m-a)} \in \mathcal{A} \otimes \text{End } (\mathbb{C}^N)^{\otimes m}.$$

Given any element

$$C = \sum_{i,j,k,l=1}^N c_{ijkl} e_{ij} \otimes e_{kl} \in \text{End } \mathbb{C}^N \otimes \text{End } \mathbb{C}^N,$$

for any two indices $a, b \in \{1, \dots, m\}$ such that $a < b$, we set

$$C_{ab} = \sum_{i,j,k,l=1}^N c_{ijkl} 1^{\otimes(a-1)} \otimes e_{ij} \otimes 1^{\otimes(b-a-1)} \otimes e_{kl} \otimes 1^{\otimes(m-b)} \in \text{End}(\mathbb{C}^N)^{\otimes m}.$$

Sometimes an additional copy of the endomorphism algebra $\text{End } \mathbb{C}^N$ labelled by 0 will be used so that the notation extends accordingly to that case.

For any $a \in \{1, \dots, m\}$ the *partial trace* tr_a will be understood as the linear map

$$\text{tr}_a : \text{End}(\mathbb{C}^N)^{\otimes m} \rightarrow \text{End}(\mathbb{C}^N)^{\otimes(m-1)}$$

which acts as the usual trace map on the a -th copy of $\text{End } \mathbb{C}^N$ and is the identity map on all the remaining copies. Similarly, the *partial transposition* t_a is the linear map on $\text{End}(\mathbb{C}^N)^{\otimes m}$ which acts as the usual transposition $t : e_{ij} \mapsto e_{ji}$ on the a -th copy of $\text{End } \mathbb{C}^N$ and is the identity map on all the remaining copies.

For a given simple Lie algebra \mathfrak{g} over \mathbb{C} introduce a symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$ as the normalized Killing form

$$\langle X, Y \rangle = \frac{1}{2h^\vee} \text{tr}(\text{ad } X \text{ ad } Y), \quad (2.2)$$

where h^\vee is the dual Coxeter number for \mathfrak{g} . Fix a basis J^1, \dots, J^d of \mathfrak{g} and let J_1, \dots, J_d be the basis dual to J^1, \dots, J^d with respect to the form (2.2). Define Casimir elements by

$$\overline{\Omega} = \sum_{i=1}^d J_i \otimes J^i \in \text{U}(\mathfrak{g}) \otimes \text{U}(\mathfrak{g}) \quad \text{and} \quad \omega = \sum_{i=1}^d J_i J^i \in \text{U}(\mathfrak{g}). \quad (2.3)$$

With the chosen normalization of the Killing form, the eigenvalue of ω in the adjoint representation equals $2h^\vee$ which coincides with the value of the parameter $c_{\mathfrak{g}}$ in [21]. Now let $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a faithful representation of \mathfrak{g} of dimension $\dim V = N$. Identify the vector space V with \mathbb{C}^N by choosing a basis and set

$$\Omega = (\rho \otimes \rho)(\overline{\Omega}) \in \text{End } \mathbb{C}^N \otimes \text{End } \mathbb{C}^N.$$

Furthermore, define the matrix

$$F = \sum_{i,j=1}^N F_{ij} \otimes e_{ij} \in \text{U}(\mathfrak{g}) \otimes \text{End } \mathbb{C}^N, \quad (2.4)$$

by setting $F = -(1 \otimes \rho)(\overline{\Omega})$.

The following presentation of $\text{U}(\mathfrak{g})$ is due to Wendlandt [21, Proposition 4.4].

Proposition 2.1. *The algebra $\text{U}(\mathfrak{g})$ is generated by the elements F_{ij} with $1 \leq i, j \leq N$ subject only to the relations*

$$F_1 F_2 - F_2 F_1 = \Omega F_2 - F_2 \Omega \quad (2.5)$$

in $\text{U}(\mathfrak{g}) \otimes \text{End } \mathbb{C}^N \otimes \text{End } \mathbb{C}^N$, and

$$F^2 - \left((F^t)^2\right)^t = h^\vee F. \quad (2.6)$$

It is easy to see that $[F_1 + F_2, \Omega] = 0$ so that relation (2.5) can be written in the equivalent form

$$F_1 F_2 - F_2 F_1 = F_1 \Omega - \Omega F_1. \quad (2.7)$$

Remark 2.2. Let $P \in \text{End } \mathbb{C}^N \otimes \text{End } \mathbb{C}^N$ be the operator permuting the tensor factors (see also (4.9) below). Proposition 2.1 will hold if (2.6) is replaced by a system of linear relations on the entries of the matrix F given by

$$\text{tr}_2(\Omega F_2 - F_2 \Omega)P = h^\vee F_1. \quad (2.8)$$

If ρ is the vector representation for the classical types, then in types B, C and D this is equivalent to the skew-symmetry condition $F + F' = 0$ with respect to the bilinear form defining the orthogonal or symplectic Lie algebra. In type A the relation poses no extra conditions on the generator matrix. An explicit form of the symmetry conditions (2.8) for the 7-dimensional representation in type G_2 can be found in [16].

One consequence of relation (2.5) is the following well-known property of the powers of the generator matrix; cf. [14, Proposition 4.2.1].

Corollary 2.3. *All elements $\text{tr } F^m$ with $m \geq 0$ belong to the center of $U(\mathfrak{g})$.*

It is also known that by taking ρ to be the lowest-dimension representation of \mathfrak{g} , one can choose algebraically independent generators of the center of $U(\mathfrak{g})$ among the Casimir elements $\text{tr } F^m$ (with the exception of type D , where a Pfaffian-type element $\text{Pf } F$ has to be added). The required values of m coincide with the degrees of basic invariants of the symmetric algebra $S(\mathfrak{g})$ as given in Table 1.

Type of \mathfrak{g}	Degrees of generators
A_n	$2, 3, \dots, n+1$
B_n	$2, 4, \dots, 2n$
C_n	$2, 4, \dots, 2n$
D_n	$2, 4, \dots, 2n-2, n$
E_6	$2, 5, 6, 8, 9, 12$
E_7	$2, 6, 8, 10, 12, 14, 18$
E_8	$2, 8, 12, 14, 18, 20, 24, 30$
F_4	$2, 6, 8, 12$
G_2	$2, 6$

Table 1: Degrees of basic invariants

More precisely, the following holds.

Corollary 2.4. *Except for type D_n , the elements $\text{tr } F^m$ with m running over the values specified in Table 1 are algebraically independent generators of the center of the algebra $U(\mathfrak{g})$.*

In type D_n the elements $\text{tr } F^m$ with $m = 2, 4, \dots, 2n-2$ and $\text{Pf } F$ are algebraically independent generators of the center of the algebra $U(\mathfrak{g})$.

Corollary 2.4 is a classical result for types A, B, C, D , but it appears to be less known for the exceptional types. In those cases, to prove that the elements $\text{tr } F^m$ are algebraically independent generators, one only needs to verify that their top degree components are basic \mathfrak{g} -invariants of the symmetric algebra $S(\mathfrak{g})$. The latter property goes back to Kuin' [10], [11], with some cases previously considered by Coxeter [5] (E_6) and Takeuchi [19] (F_4). A direct claim about these Casimir elements was made in [2], [3].

To give more details for the exceptional types, recall that the theorem of Kuin', whose proof is given in [11, §2.2], reads as follows. Suppose that $\Lambda_1, \dots, \Lambda_N$ are the weights of the lowest-dimension representation (V, ρ) of \mathfrak{g} . The weights are understood as elements of the Cartan subalgebra \mathfrak{h} of \mathfrak{g} , where \mathfrak{h}^* and \mathfrak{h} are identified via the form (2.2). That is, V has a basis v_1, \dots, v_N such that any element $X \in \mathfrak{h}$ acts by

$$X v_a = \langle \Lambda_a, X \rangle v_a, \quad a = 1, \dots, N.$$

The theorem states that the power sums

$$P_m = \sum_{a=1}^N \Lambda_a^m \quad (2.9)$$

with m running over the respective degrees in Table 1, are algebraically independent generators of the subalgebra of W -invariants $S(\mathfrak{h})^W$ in $S(\mathfrak{h})$, where W denotes the Weyl group of \mathfrak{g} .

Returning to Corollary 2.4, choose a special form of the Casimir element $\overline{\Omega}$ in (2.3) by taking a basis J^1, \dots, J^d of \mathfrak{g} such that J^1, \dots, J^n form a basis of the Cartan subalgebra \mathfrak{h} , while the remaining J^i 's are root vectors. Then the vectors J_1, \dots, J_n of the dual basis will also belong to \mathfrak{h} . Consider the entries F_{ij} of the matrix F defined in (2.4) as elements of the symmetric algebra $S(\mathfrak{g})$. A nonzero contribution to the image of the element $\text{tr } F^m \in S(\mathfrak{g})^{\mathfrak{g}}$ under the Chevalley isomorphism $S(\mathfrak{g})^{\mathfrak{g}} \rightarrow S(\mathfrak{h})^W$ will only come from its part $\text{tr } \overline{F}^m$, where

$$\overline{F} = - \sum_{i=1}^n J_i \otimes \rho(J^i).$$

By our choice of parameters, $\rho(J^i)$ is a diagonal matrix of the form

$$\rho(J^i) = \sum_{a=1}^N \langle \Lambda_a, J^i \rangle e_{aa}.$$

Hence, the Chevalley image of $\text{tr } F^m$ equals

$$\sum_{a=1}^N \left(- \sum_{i=1}^n \langle \Lambda_a, J^i \rangle J_i \right)^m = (-1)^m \sum_{a=1}^N \Lambda_a^m = (-1)^m P_m$$

which thus coincides with the element (2.9), up to a sign.

3 Casimir elements for Takiff algebras

We will use the presentation of \mathfrak{g} associated with an arbitrary faithful representation ρ , as given in Proposition 2.1. Introduce elements of the Takiff algebra \mathfrak{g}_ℓ as defined in the Introduction, by $F_{ij}^{(r)} = F_{ij} v^r$ for $r = 0, 1, \dots, \ell$. We combine them into the respective matrices

$$F^{(r)} = \sum_{i,j=1}^N F_{ij}^{(r)} \otimes e_{ij} \in U(\mathfrak{g}_\ell) \otimes \text{End } \mathbb{C}^N.$$

For a variable u set

$$F(u) = F^{(0)} + F^{(1)}u + \dots + F^{(\ell)}u^\ell.$$

The traces of powers of this matrix are polynomials in u of the form

$$\text{tr } F(u)^m = \sum_r \theta_m^{(r)} u^r, \quad \theta_m^{(r)} \in U(\mathfrak{g}_\ell).$$

Proposition 3.1. *All elements $\theta_m^{(r)}$ with $m \geq 1$ and $r = m\ell, m\ell - 1, \dots, m\ell - \ell$ belong to the center of the algebra $U(\mathfrak{g}_\ell)$.*

Proof. Since $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}]$, the Lie algebra \mathfrak{g}_ℓ is generated by the elements $F_{ij}^{(0)}$ and $F_{ij}^{(1)}$ with $1 \leq i, j \leq N$. Hence, it is sufficient to verify that all commutators

$$[F_{ij}^{(0)}, \text{tr } F(u)^m] \quad \text{and} \quad [F_{ij}^{(1)}, \text{tr } F(u)^m]$$

are polynomials in u whose degrees are less than $m\ell - \ell$. We will use the matrix notation of Section 2 and consider the algebra $U(\mathfrak{g}_\ell) \otimes \text{End } \mathbb{C}^N \otimes \text{End } \mathbb{C}^N$ with the tensor factors $\text{End } \mathbb{C}^N$ labelled by 0 and 1. Relation (2.5) implies

$$\begin{aligned} [F_0^{(0)}, F(u)_1^m] &= \sum_{i=1}^m F(u)_1^{i-1} [F_0^{(0)}, F(u)_1] F(u)_1^{m-i} \\ &= \sum_{i=1}^m F(u)_1^{i-1} [\Omega_{01}, F(u)_1] F(u)_1^{m-i} = [\Omega_{01}, F(u)_1^m]. \end{aligned}$$

Therefore, taking the partial trace tr_1 on both sides and using its cyclic property we can conclude that $[F_0^{(0)}, \text{tr}_1 F(u)_1^m] = 0$. We also have

$$u [F_0^{(1)}, F(u)_1] = [\Omega_{01}, F(u)_1 - F_1^{(0)}].$$

Hence, a similar calculation gives

$$u [F_0^{(1)}, \text{tr}_1 F(u)_1^m] = - \sum_{i=1}^m \text{tr}_1 F(u)_1^{i-1} [\Omega_{01}, F_1^{(0)}] F(u)_1^{m-i}.$$

However, the degree of the polynomial in u on the right hand side is $m\ell - \ell$ so that the commutator $[F_0^{(1)}, \text{tr}_1 F(u)_1^m]$ is a polynomial of degree less than $m\ell - \ell$, as required. \square

Remark 3.2. Proposition 3.1 and its proof extend to the reductive Lie algebra \mathfrak{gl}_N , where the F_{ij} should be replaced by the standard basis elements E_{ij} of \mathfrak{gl}_N . If ρ is the vector representation of \mathfrak{gl}_N , the Casimir elements provided by the proposition are closely related to one of the families produced in [13]. \square

Now suppose that $\mathfrak{g} = \mathfrak{o}_{2n}$ is the orthogonal Lie algebra (of type D_n). We will use its presentation where the elements of \mathfrak{g} are skew-symmetric $2n \times 2n$ matrices with the usual matrix commutator. In relation (2.5) we have

$$\Omega = \sum_{i,j=1}^{2n} (e_{ij} \otimes e_{ji} - e_{ij} \otimes e_{ij}),$$

while (2.8) is equivalent to $F + F^t = 0$. Define the *Pfaffian* of the matrix $F(u)$ by the formula

$$\text{Pf } F(u) = \sum_{\sigma} \text{sgn } \sigma \cdot F_{\sigma(1)\sigma(2)}(u) \cdots F_{\sigma(2n-1)\sigma(2n)}(u), \quad (3.1)$$

summed over the elements σ of the subset $\mathcal{A}_{2n} \subset \mathfrak{S}_{2n}$ of the symmetric group \mathfrak{S}_{2n} which consists of the permutations with the properties $\sigma(2k-1) < \sigma(2k)$ for all $k = 1, \dots, n$ and $\sigma(1) < \sigma(3) < \cdots < \sigma(2n-1)$. Introduce the coefficients of the polynomial $\text{Pf } F(u)$ by

$$\text{Pf } F(u) = \sum_r \pi^{(r)} u^r.$$

Proposition 3.3. *All coefficients $\pi^{(r)}$ with $r = n\ell, n\ell - 1, \dots, n\ell - \ell$ belong to the center of the algebra $U(\mathfrak{g}_\ell)$.*

Proof. It is enough to prove that the commutators

$$[F_{ij}^{(0)}, \text{Pf } F(u)] \quad \text{and} \quad [F_{ij}^{(1)}, \text{Pf } F(u)] \quad (3.2)$$

are polynomials in u of degree less than $n\ell - \ell$. Note that for any permutation $\pi \in \mathfrak{S}_{2n}$ the mapping $F_{ij}^{(r)} \mapsto F_{\pi(i)\pi(j)}^{(r)}$ defines an automorphism of the Lie algebra \mathfrak{g}_ℓ . Hence, it is sufficient to verify the required properties of the commutators in (3.2) for $i = 1$ and $j = 2$. This follows by a straightforward calculation with the use of the commutation relations

$$[F_{ij}^{(0)}, F_{kl}(u)] = \delta_{kj} F_{il}(u) - \delta_{il} F_{kj}(u) - \delta_{ki} F_{jl}(u) + \delta_{jl} F_{ki}(u)$$

and

$$u[F_{ij}^{(1)}, F_{kl}(u)] = [F_{ij}^{(0)}, F_{kl}(u) - F_{kl}^{(0)}]$$

and arguing as in the proof of Proposition 3.1. \square

Remark 3.4. Analogous expressions for the Pfaffian-type central elements can also be written for the presentations of the orthogonal Lie algebra associated with arbitrary non-degenerate symmetric bilinear forms on \mathbb{C}^{2n} ; cf. [14, Sec. 8]. \square

Return to an arbitrary simple Lie algebra \mathfrak{g} and suppose now that ρ is the lowest-dimension representation of \mathfrak{g} .

Theorem 3.5. *Except for type D_n , the elements $\theta_m^{(r)}$ with m running over the values specified in Table 1 and $r = m\ell, m\ell - 1, \dots, m\ell - \ell$, are algebraically independent generators of the center of the algebra $U(\mathfrak{g}_\ell)$.*

In type D_n the elements $\theta_m^{(r)}$ with $m = 2, 4, \dots, 2n - 2$ and $r = m\ell, m\ell - 1, \dots, m\ell - \ell$ together with $\pi^{(r)}$ with $r = n\ell, n\ell - 1, \dots, n\ell - \ell$, are algebraically independent generators of the center of the algebra $U(\mathfrak{g}_\ell)$.

Proof. All these elements belong to the center of $U(\mathfrak{g}_\ell)$ by Propositions 3.1 and 3.3. Their symbols in the symmetric algebra $S(\mathfrak{g}_\ell)$ are \mathfrak{g}_ℓ -invariants. Moreover, these invariants are associated with basic \mathfrak{g} -invariants in $S(\mathfrak{g})$ in the way described in [18, Sec. 3.1]. Therefore, applying [18, Théorème 4.5], we can conclude that the symbols are algebraically independent generators of the subalgebra of \mathfrak{g}_ℓ -invariants in $S(\mathfrak{g}_\ell)$. This implies the desired property of the central elements in $U(\mathfrak{g}_\ell)$. \square

4 Segal–Sugawara vectors

We will identify the Lie algebra \mathfrak{g} with a subalgebra of \mathfrak{g}_ℓ via the embedding $F_{ij} \mapsto F_{ij}^{(0)}$. Extend the form (2.2) defined on this subalgebra to the Lie algebra \mathfrak{g}_ℓ by positing that all elements $F_{ij}^{(r)}$ with $r = 1, \dots, \ell$ belong to its kernel. This defines a symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$ on \mathfrak{g}_ℓ . The corresponding affine Kac–Moody algebra $\widehat{\mathfrak{g}}_\ell$ is the central extension

$$\widehat{\mathfrak{g}}_\ell = \mathfrak{g}_\ell[t, t^{-1}] \oplus \mathbb{C}K,$$

where $\mathfrak{g}_\ell[t, t^{-1}]$ is the Lie algebra of Laurent polynomials in t with coefficients in \mathfrak{g}_ℓ . For any $r \in \mathbb{Z}$ and $X \in \mathfrak{g}_\ell$ we will write $X[r] = X t^r$. The commutation relations of the Lie algebra $\widehat{\mathfrak{g}}_\ell$ have the form

$$[X[r], Y[s]] = [X, Y][r + s] + r \delta_{r, -s} \langle X, Y \rangle K, \quad X, Y \in \mathfrak{g}_\ell, \quad (4.1)$$

and the element K is central in $\widehat{\mathfrak{g}}_\ell$.

The *vacuum module at the level $k \in \mathbb{C}$* over $\widehat{\mathfrak{g}}_\ell$ is the quotient $V_k(\mathfrak{g}_\ell) = U(\widehat{\mathfrak{g}}_\ell)/I$, where I is the left ideal of $U(\widehat{\mathfrak{g}}_\ell)$ generated by $\mathfrak{g}_\ell[t]$ and the element $K - k$. The Poincaré–Birkhoff–Witt theorem implies that this quotient is isomorphic to the universal enveloping algebra $U(t^{-1}\mathfrak{g}_\ell[t^{-1}])$, as a vector space. The vacuum module is equipped with a vertex algebra structure; see e.g. [7], [9]. We will call the level $k = -(\ell + 1)h^\vee$ *critical*, as the vacuum module $V_{\text{cri}}(\mathfrak{g}_\ell)$ at this level turns out to exhibit similar properties to its counterpart for $\ell = 0$. We will denote by $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ the center of the vertex algebra $V_{\text{cri}}(\mathfrak{g}_\ell)$ which is defined as the subspace

$$\mathfrak{z}(\widehat{\mathfrak{g}}_\ell) = \{v \in V_{\text{cri}}(\mathfrak{g}_\ell) \mid \mathfrak{g}_\ell[t]v = 0\}.$$

It follows from the axioms of vertex algebra that $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ is a unital commutative associative algebra which can be regarded as a subalgebra of $U(t^{-1}\mathfrak{g}_\ell[t^{-1}])$. This subalgebra is invariant with respect

to the *translation operator* T which is the derivation of the algebra $U(t^{-1}\mathfrak{g}_\ell[t^{-1}])$ whose action on the generators is given by

$$T : X[r] \mapsto -rX[r-1], \quad X \in \mathfrak{g}_\ell, \quad r < 0. \quad (4.2)$$

Any element of $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ is called a *Segal–Sugawara vector*. By the Feigin–Frenkel theorem [6], [7], in the case $\ell = 0$ the center $\mathfrak{z}(\widehat{\mathfrak{g}})$ contains a *complete set of Segal–Sugawara vectors* S_1, \dots, S_n , which means that the translations $T^r S_p$ with $r \geq 0$ and $p = 1, \dots, n (= \text{rank } \mathfrak{g})$ are algebraically independent generators of the algebra $\mathfrak{z}(\widehat{\mathfrak{g}})$.

Our goal is to prove that this property is shared by the algebras $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ for all $\ell \geq 1$. Note that this has already been proved in type A by Arakawa and Premet [1] as a particular case of a more general theorem on affine vertex algebras associated with centralizers of nilpotent elements in \mathfrak{g} ; see also [15] for an explicit construction of a complete set of Segal–Sugawara vectors.

We begin by producing some families of Segal–Sugawara vectors for \mathfrak{g}_ℓ and then will show how to choose a complete set of such vectors. As in Section 3, we will use the presentation of \mathfrak{g} associated with an arbitrary faithful representation ρ , given in Proposition 2.1. Introduce polynomials in u of the form

$$\mathcal{F}(u) = F^{(0)}[-1] + F^{(1)}[-1]u + \dots + F^{(\ell)}[-1]u^\ell,$$

where $F^{(r)}[p]$ denotes the matrix

$$F^{(r)}[p] = \sum_{i,j=1}^N F_{ij}^{(r)}[p] \otimes e_{ij}, \quad p \in \mathbb{Z}. \quad (4.3)$$

Define elements $\Theta_m^{(r)} \in V_{\text{cri}}(\mathfrak{g}_\ell) \cong U(t^{-1}\mathfrak{g}_\ell[t^{-1}])$ as the coefficients of the polynomial

$$\text{tr } \mathcal{F}(u)^m = \sum_r \Theta_m^{(r)} u^r.$$

Proposition 4.1. *Suppose that $\ell \geq 1$. Then all coefficients $\Theta_m^{(r)}$ with the parameters $m \geq 1$ and $r = m\ell, m\ell - 1, \dots, m\ell - \ell$ are Segal–Sugawara vectors for \mathfrak{g}_ℓ .*

Proof. Since $F_{ij}^{(0)}[0], F_{ij}^{(0)}[1]$ and $F_{ij}^{(1)}[0]$ with $1 \leq i, j \leq N$ are generators of the Lie algebra $\mathfrak{g}_\ell[t]$, it is sufficient to verify that they annihilate the elements $\Theta_m^{(r)}$. Using the matrix notation as in the proof of Proposition 3.1, we derive from (2.5) that

$$[F^{(0)}[0]_0, \mathcal{F}(u)_1] = [\Omega_{01}, \mathcal{F}(u)_1].$$

Hence, $F^{(0)}[0]_0 \text{tr}_1 \mathcal{F}(u)_1^m = 0$ in $V_{\text{cri}}(\mathfrak{g}_\ell)$ which follows in the same way as in the proof of Proposition 3.1. As in that proof, we also have

$$u [F^{(1)}[0]_0, \mathcal{F}(u)_1] = [\Omega_{01}, \mathcal{F}(u)_1 - F^{(0)}[-1]_1],$$

which implies that the degree of the polynomial $F^{(1)}[0]_0 \text{tr}_1 \mathcal{F}(u)_1^m$ in u is less than $m\ell - \ell$.

Furthermore, as an immediate consequence of (2.7) and (4.1), we obtain

$$\left[F^{(0)}[r]_0, F^{(0)}[s]_1 \right] = \left[F^{(0)}[r+s]_0, \Omega_{01} \right] + r \delta_{r,-s} \Omega_{01} K.$$

Hence, for the remaining generators of $\mathfrak{g}_\ell[t]$ we have

$$\left[F^{(0)}[1]_0, \mathcal{F}(u)_1 \right] = \left[\mathcal{F}(0, u)_0, \Omega_{01} \right] + \Omega_{01} K,$$

where we used the notation

$$\mathcal{F}(0, u) = F^{(0)}[0] + F^{(1)}[0]u + \cdots + F^{(\ell)}[0]u^\ell. \quad (4.4)$$

Calculating in the vacuum module we then find

$$F^{(0)}[1]_0 \operatorname{tr}_1 \mathcal{F}(u)_1^m = \sum_{i=1}^m \operatorname{tr}_1 \mathcal{F}(u)_1^{i-1} \left(-\Omega_{01} \mathcal{F}(0, u)_0 + \mathcal{F}(0, u)_0 \Omega_{01} + \Omega_{01} K \right) \mathcal{F}(u)_1^{m-i}.$$

Observe that modulo a polynomial in u of degree less than ℓ , we can write

$$\left[\mathcal{F}(0, u)_0, \mathcal{F}(u)_1 \right] \equiv (\ell + 1) \left[\Omega_{01}, F^{(\ell)}[-1]_1 u^\ell \right] \equiv (\ell + 1) \left[\Omega_{01}, \mathcal{F}(u)_1 \right]. \quad (4.5)$$

Therefore,

$$\mathcal{F}(0, u)_0 \mathcal{F}(u)_1^{m-i} \equiv (\ell + 1) \left[\Omega_{01}, \mathcal{F}(u)_1^{m-i} \right] \quad (4.6)$$

modulo a polynomial in u of degree less than $(m - i)\ell$.

Now use the following general property of the partial transposition t_1 :

$$\operatorname{tr}_1 XY = \operatorname{tr}_1 X^{t_1} Y^{t_1}. \quad (4.7)$$

Taking $X = \mathcal{F}(u)_1^{i-1} \mathcal{F}(0, u)_0$ and $Y = \Omega_{01} \mathcal{F}(u)_1^{m-i}$ we obtain

$$\operatorname{tr}_1 \mathcal{F}(u)_1^{i-1} \mathcal{F}(0, u)_0 \Omega_{01} \mathcal{F}(u)_1^{m-i} = \operatorname{tr}_1 \left(\mathcal{F}(u)_1^{i-1} \right)_1^t \mathcal{F}(0, u)_0 \left(\mathcal{F}(u)_1^{m-i} \right)_1^t (\Omega_{01})^{t_1}.$$

Applying the transposition to both sides of (4.6) we get

$$\mathcal{F}(0, u)_0 \left(\mathcal{F}(u)_1^{m-i} \right)_1^t \equiv (\ell + 1) \left[\left(\mathcal{F}(u)_1^{m-i} \right)_1^t, (\Omega_{01})^{t_1} \right].$$

Thus, bringing the calculations together, we obtain the following relation in $V_{\text{cri}}(\mathfrak{g}_\ell)$ modulo a polynomial in u of degree less than $m\ell - \ell$:

$$\begin{aligned} F^{(0)}[1]_0 \operatorname{tr}_1 \mathcal{F}(u)_1^m &\equiv K \sum_{i=1}^m \operatorname{tr}_1 \mathcal{F}(u)_1^{i-1} \Omega_{01} \mathcal{F}(u)_1^{m-i} \\ &\quad + (\ell + 1) \sum_{i=1}^m \operatorname{tr}_1 \left(\mathcal{F}(u)_1^{i-1} \Omega_{01} \mathcal{F}(u)_1^{m-i} \Omega_{01} - \mathcal{F}(u)_1^{i-1} \Omega_{01}^2 \mathcal{F}(u)_1^{m-i} \right. \\ &\quad \left. - \left(\mathcal{F}(u)_1^{i-1} \right)_1^t (\Omega_{01})^{t_1} \left(\mathcal{F}(u)_1^{m-i} \right)_1^t (\Omega_{01})^{t_1} + \left(\mathcal{F}(u)_1^{i-1} \right)_1^t \left(\mathcal{F}(u)_1^{m-i} \right)_1^t \left((\Omega_{01})^{t_1} \right)^2 \right). \end{aligned}$$

The application of (4.7) to the terms in the last line brings this expression to the form

$$\begin{aligned} F^{(0)}[1]_0 \operatorname{tr}_1 \mathcal{F}(u)_1^m &\equiv \sum_{i=1}^m \operatorname{tr}_1 \mathcal{F}(u)_1^{i-1} \left(K \Omega_{01} + (\ell+1) \left(((\Omega_{01})^{t_1})^2 \right)^{t_1} - (\ell+1) \Omega_{01}^2 \right) \mathcal{F}(u)_1^{m-i} \\ &+ (\ell+1) \sum_{i=1}^m \operatorname{tr}_1 \left(\mathcal{F}(u)_1^{i-1} \Omega_{01} \mathcal{F}(u)_1^{m-i} \Omega_{01} - \Omega_{01} \mathcal{F}(u)_1^{i-1} \Omega_{01} \mathcal{F}(u)_1^{m-i} \right). \end{aligned}$$

Recall that $K = -(\ell+1)h^\vee$ at the critical level and so the first sum is zero. This follows from the identity

$$\Omega_{01}^2 - \left(((\Omega_{01})^{t_1})^2 \right)^{t_1} + h^\vee \Omega_{01} = 0$$

which is a consequence of relation (2.6); we just need to apply $\rho \otimes 1$ to its both sides and note that $\Omega_{01} = -(\rho \otimes 1)(F)$.

The second sum is a polynomial in u of degree at most $m\ell - \ell$. It remains to verify that the coefficient of $u^{m\ell - \ell}$ in the sum is zero. This coefficient equals

$$\sum_{i=1}^m \operatorname{tr}_1 \left(\Phi_1^{i-1} \Omega_{01} \Phi_1^{m-i} \Omega_{01} - \Omega_{01} \Phi_1^{i-1} \Omega_{01} \Phi_1^{m-i} \right), \quad (4.8)$$

where we set $\Phi = F^{(\ell)}[-1]$ for brevity. In the algebra

$$\operatorname{U}(\widehat{\mathfrak{g}}_\ell) \otimes \operatorname{End} \mathbb{C}^N \otimes \operatorname{End} \mathbb{C}^N \otimes \operatorname{End} \mathbb{C}^N$$

with the tensor factors $\operatorname{End} \mathbb{C}^N$ labelled by 0, 1 and 2 we can write

$$\operatorname{tr}_1 \Omega_{01} \Phi_1^{i-1} \Omega_{01} \Phi_1^{m-i} = \operatorname{tr}_{1,2} \Omega_{01} \Phi_1^{i-1} \Omega_{01} \Phi_2^{m-i} P_{12},$$

where

$$P_{12} = \sum_{i,j=1}^N e_{ij} \otimes e_{ji} \quad (4.9)$$

is the permutation operator. We have used the relation $\Phi_2^{m-i} P_{12} = P_{12} \Phi_1^{m-i}$ and observed that $\operatorname{tr}_2 P_{12} = 1$. Since $\ell \geq 1$, the matrix elements of the matrix Φ pairwise commute and so

$$\operatorname{tr}_{1,2} \Omega_{01} \Phi_1^{i-1} \Omega_{01} \Phi_2^{m-i} P_{12} = \operatorname{tr}_{1,2} \Phi_2^{m-i} \Omega_{01} \Phi_1^{i-1} \Omega_{01} P_{12} = \operatorname{tr}_{1,2} \Phi_2^{m-i} P_{12} \Omega_{02} \Phi_2^{i-1} \Omega_{02}$$

which equals

$$\operatorname{tr}_2 \Phi_2^{m-i} \Omega_{02} \Phi_2^{i-1} \Omega_{02} = \operatorname{tr}_1 \Phi_1^{m-i} \Omega_{01} \Phi_1^{i-1} \Omega_{01}.$$

Thus, the expression (4.8) is zero so that $F^{(0)}[1]_0 \operatorname{tr}_1 \mathcal{F}(u)_1^m$ is a polynomial in u of degree less than $m\ell - \ell$. \square

Remark 4.2. Proposition 4.1 does not hold under the assumption $\ell = 0$; see e.g. [14, Secs 7.1, 8.2 and 8.4] for counterexamples in classical types. \square

Consider now the orthogonal Lie algebra $\mathfrak{g} = \mathfrak{o}_{2n}$ with its presentation used in Section 3. Define elements $\Pi^{(r)} \in V_{\text{cri}}(\mathfrak{g}_\ell)$ as the coefficients of the *Pfaffian* of the matrix $\mathcal{F}(u)$

$$\text{Pf } \mathcal{F}(u) = \sum_r \Pi^{(r)} u^r,$$

where $\text{Pf } \mathcal{F}(u)$ is given by formula (3.1) applied to the matrix $\mathcal{F}(u)$.

Proposition 4.3. *All coefficients $\Pi^{(r)}$ with $r = n\ell, n\ell-1, \dots, n\ell-\ell$ are Segal–Sugawara vectors for \mathfrak{g}_ℓ .*

Proof. Given any permutation $\pi \in \mathfrak{S}_{2n}$, the mapping

$$F_{ij}^{(r)}[p] \mapsto F_{\pi(i)\pi(j)}^{(r)}[p], \quad K \mapsto K$$

defines an automorphism of the Lie algebra $\widehat{\mathfrak{g}}_\ell$. Therefore, it is sufficient to verify that the elements $F_{12}^{(0)}[0], F_{12}^{(1)}[0]$ and $F_{12}^{(0)}[1]$ acting in the vacuum module $V_{\text{cri}}(\mathfrak{g}_\ell)$ annihilate the given coefficients $\Pi^{(r)}$. The argument for the first two elements is a straightforward calculation with the use of the commutation relations involving the entries of the matrix $\mathcal{F}(u) = [\mathcal{F}_{ij}(u)]$,

$$[F_{ij}^{(0)}[0], \mathcal{F}_{kl}(u)] = \delta_{kj} \mathcal{F}_{il}(u) - \delta_{il} \mathcal{F}_{kj}(u) - \delta_{ki} \mathcal{F}_{jl}(u) + \delta_{jl} \mathcal{F}_{ki}(u)$$

and

$$u[F_{ij}^{(1)}[0], \mathcal{F}_{kl}(u)] = [\mathcal{F}_{ij}^{(0)}, \mathcal{F}_{kl}(u) - F_{kl}^{(0)}[-1]];$$

cf. the case $\ell = 0$ in [14, Proposition 8.1.4]. For the remaining element we use the relations

$$\begin{aligned} [F_{ij}^{(0)}[1], \mathcal{F}_{kl}(u)] &= \delta_{kj} \mathcal{F}_{il}(0, u) - \delta_{il} \mathcal{F}_{kj}(0, u) - \delta_{ki} \mathcal{F}_{jl}(0, u) + \delta_{jl} \mathcal{F}_{ki}(0, u) \\ &\quad + K(\delta_{kj} \delta_{il} - \delta_{ki} \delta_{jl}) \end{aligned}$$

involving the entries of the matrix (4.4). Consider first the summands in formula (3.1) for the matrix $\mathcal{F}(u)$ with $\sigma(1) = 1$ and $\sigma(2) = 2$. In the vacuum module we have

$$F_{12}^{(0)}[1] \mathcal{F}_{12}(u) \mathcal{F}_{\sigma(3)\sigma(4)}(u) \dots \mathcal{F}_{\sigma(2n-1)\sigma(2n)}(u) = -K \mathcal{F}_{\sigma(3)\sigma(4)}(u) \dots \mathcal{F}_{\sigma(2n-1)\sigma(2n)}(u).$$

Furthermore, let $\tau \in \mathcal{A}_{2n}$ with $\tau(2) > 2$. Then $\tau(3) = 2$ and $\tau(4) > 2$ and so

$$F_{12}^{(0)}[1] \mathcal{F}_{1\tau(2)}(u) \mathcal{F}_{2\tau(4)}(u) \dots \mathcal{F}_{\tau(2n-1)\tau(2n)}(u) = -\mathcal{F}_{2\tau(2)}(0, u) \mathcal{F}_{2\tau(4)}(u) \dots \mathcal{F}_{\tau(2n-1)\tau(2n)}(u).$$

Considering this expression modulo a polynomial in u of degree less than $n\ell - \ell$, we may apply relations (4.5) to conclude that the expression coincides with

$$(\ell + 1) \mathcal{F}_{\tau(2)\tau(4)}(u) \dots \mathcal{F}_{\tau(2n-1)\tau(2n)}(u).$$

Suppose now that $\sigma \in \mathcal{A}_{2n}$ is an element with the fixed values $\sigma(1) = 1$ and $\sigma(2) = 2$, and calculate the coefficient of the monomial

$$\mathcal{F}_{\sigma(3)\sigma(4)}(u) \dots \mathcal{F}_{\sigma(2n-1)\sigma(2n)}(u)$$

in the expansion of $F_{12}^{(0)}[1] \text{Pf } \mathcal{F}(u)$ modulo a polynomial in u of degree less than $n\ell - \ell$. Essentially the same calculation was already performed in the case $\ell = 0$ in [14, Proposition 8.1.4] showing that the coefficient equals $-K - (\ell + 1)(2n - 2)$, which is zero at the critical level since $h^\vee = 2n - 2$. \square

Now let \mathfrak{g} be an arbitrary simple Lie algebra and suppose that ρ is the lowest-dimension representation of \mathfrak{g} . The next theorem shows that a natural extension of the Feigin–Frenkel theorem [6] holds for the Lie algebra \mathfrak{g}_ℓ with $\ell \geq 1$; that is, the center $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$ is an algebra of polynomials.

Theorem 4.4. *Except for type D_n , the elements $\Theta_m^{(r)}$ with m running over the values specified in Table 1 and $r = m\ell, m\ell - 1, \dots, m\ell - \ell$, form a complete set of Segal–Sugawara vectors for the Lie algebra \mathfrak{g}_ℓ .*

In type D_n the elements $\Theta_m^{(r)}$ with $m = 2, 4, \dots, 2n - 2$ and $r = m\ell, m\ell - 1, \dots, m\ell - \ell$ together with $\Pi^{(r)}$ with $r = n\ell, n\ell - 1, \dots, n\ell - \ell$, form a complete set of Segal–Sugawara vectors for the Lie algebra \mathfrak{g}_ℓ .

Proof. All elements are Segal–Sugawara vectors by Propositions 4.1 and 4.3. We need to show that all the shifted vectors $T^s \Theta_m^{(r)}$ (together with $T^s \Pi^{(r)}$ in type D_n) with $s \geq 0$ are algebraically independent generators of the algebra $\mathfrak{z}(\widehat{\mathfrak{g}}_\ell)$. We will follow the approach which was used for the case $\ell = 0$; cf. [7, Secs 3.3 and 3.4] and [14, Sec. 6.3].

Regard $V_{\text{cri}}(\mathfrak{g}_\ell)$ as a $\mathfrak{g}_\ell[t]$ -module obtained by restriction of the action of $\widehat{\mathfrak{g}}_\ell$ to the subalgebra $\mathfrak{g}_\ell[t]$. By identifying the vector space $V_{\text{cri}}(\mathfrak{g}_\ell)$ with the algebra $U(t^{-1}\mathfrak{g}_\ell[t^{-1}])$ and using its canonical filtration, equip the associated graded space $\text{gr } V_{\text{cri}}(\mathfrak{g}_\ell)$ with the structure of a $\mathfrak{g}_\ell[t]$ -module. As a vector space, $\text{gr } V_{\text{cri}}(\mathfrak{g}_\ell)$ will be identified with the symmetric algebra $S(t^{-1}\mathfrak{g}_\ell[t^{-1}])$. The action of $\mathfrak{g}_\ell[t]$ on $S(t^{-1}\mathfrak{g}_\ell[t^{-1}])$ is obtained by extending the adjoint representation of $\mathfrak{g}_\ell[t]$ on $\mathfrak{g}_\ell[t, t^{-1}]/\mathfrak{g}_\ell[t] \cong t^{-1}\mathfrak{g}_\ell[t^{-1}]$ to the symmetric algebra.

The symbols $\overline{\Theta}_m^{(r)}$ of the Segal–Sugawara vectors $\Theta_m^{(r)}$ (and the symbols $\overline{\Pi}^{(r)}$ of the Segal–Sugawara vectors $\Pi^{(r)}$ in type D_n) belong to the subalgebra $S(t^{-1}\mathfrak{g}_\ell[t^{-1}])^{\mathfrak{g}_\ell[t]}$ of $\mathfrak{g}_\ell[t]$ -invariants in $S(t^{-1}\mathfrak{g}_\ell[t^{-1}])$. The translation operator defined in (4.2) induces a derivation of the symmetric algebra which we will also denote by T . We only need to verify that all the shifts $T^s \overline{\Theta}_m^{(r)}$ (together with $T^s \overline{\Pi}^{(r)}$ in type D_n) with $s \geq 0$ are algebraically independent generators of the algebra of invariants.

For the rest of the proof we will work with the symmetric algebra and so we will keep the same notation $F_{ij}^{(r)}[-p-1]$ with $p \geq 0$ for the images of these elements of the enveloping algebra in the associated graded algebra $S(t^{-1}\mathfrak{g}_\ell[t^{-1}])$. Furthermore, we will only consider the elements $\overline{\Theta}_m^{(r)}$; the extension to the Pfaffian-type invariants in type D_n will then be obvious.

Using matrices (4.3), introduce power series in a variable z by

$$\mathcal{F}^{(r)}(z) = \sum_{p=0}^{\infty} F^{(r)}[-p-1] z^p \quad (4.10)$$

and write the series

$$\overline{\Theta}_m(z, u) = \text{tr} \left(\mathcal{F}^{(0)}(z) + \dots + \mathcal{F}^{(\ell)}(z) u^\ell \right)^m \quad (4.11)$$

as a polynomial in u ,

$$\overline{\Theta}_m(z, u) = \sum_r \overline{\Theta}_m^{(r)}(z) u^r.$$

The element $T^s \overline{\Theta}_m^{(r)}$ then equals $s!$ times the coefficient of z^s in the series $\overline{\Theta}_m^{(r)}(z)$.

On the other hand, we can use the non-degenerate invariant symmetric bilinear form on the Lie algebra \mathfrak{g}_ℓ defined by

$$\langle X_0 + X_1 v + \cdots + X_\ell v^\ell, Y_0 + Y_1 v + \cdots + Y_\ell v^\ell \rangle_\ell = \langle X_0, Y_\ell \rangle + \cdots + \langle X_\ell, Y_0 \rangle$$

to identify the $\mathfrak{g}_\ell[t]$ -module $t^{-1}\mathfrak{g}_\ell[t^{-1}]$ with the restricted dual $\mathfrak{g}_\ell[t]^*$. Namely, we will use the isomorphism such that for any $X, Y \in \mathfrak{g}_\ell$ and all values $p, s \geq 0$ the image of the element $X[-p-1] \in t^{-1}\mathfrak{g}_\ell[t^{-1}]$ is the functional whose value on $Y[s] \in \mathfrak{g}_\ell[t]$ equals $\delta_{ps} \langle X, Y \rangle_\ell$. Denote the image of the matrix $F^{(r)}[-p-1]$ under this isomorphism by $\mathcal{X}^{(\ell-r)}[p]$, and the corresponding image of the series (4.10) by $\mathcal{X}^{(\ell-r)}(z)$. Accordingly, the image of the series (4.11) will then be written as

$$\mathcal{T}_m(z, u) = \text{tr} \left(\mathcal{X}^{(\ell)}(z) + \cdots + \mathcal{X}^{(0)}(z) u^\ell \right)^m$$

so that

$$u^{-m\ell} \mathcal{T}_m(z, u) = \text{tr} \left(\mathcal{X}^{(0)}(z) + \cdots + \mathcal{X}^{(\ell)}(z) u^{-\ell} \right)^m.$$

Expand this as a polynomial in u^{-1}

$$\text{tr} \left(\mathcal{X}^{(0)}(z) + \cdots + \mathcal{X}^{(\ell)}(z) u^{-\ell} \right)^m = \sum_{r \geq 0} \mathcal{T}_m^{(r)}(z) u^{-r}$$

thus defining the power series

$$\mathcal{T}_m^{(r)}(z) = \sum_{s=0}^{\infty} \mathcal{T}_m^{(r)}[s] z^s.$$

By identifying $S(\mathfrak{g}_\ell[t]^*)$ with the algebra of polynomial functions $\text{Fun } \mathfrak{g}_\ell[t]$ on $\mathfrak{g}_\ell[t]$ we come to proving that all coefficients of the series $\mathcal{T}_m^{(r)}(z)$ with m running over the values specified in Table 1 and $r = 0, 1, \dots, \ell$ are algebraically independent generators of the algebra of $\mathfrak{g}_\ell[t]$ -invariants of $\text{Fun } \mathfrak{g}_\ell[t]$. It is sufficient to establish the corresponding property for each truncated algebra $\mathfrak{g}_{\ell, \ell'}$ defined as the quotient of $\mathfrak{g}_\ell[t]$ by the ideal $t^{\ell'+1}\mathfrak{g}_\ell[t]$. More precisely, we need to show that for any nonnegative integer ℓ' , the coefficients $\mathcal{T}_m^{(r)}[s]$ with the same conditions on m and r as above, and with s taking the values $0, 1, \dots, \ell'$, are algebraically independent generators of the algebra of $\mathfrak{g}_{\ell, \ell'}$ -invariants of $\text{Fun } \mathfrak{g}_{\ell, \ell'}$. Writing the definition of the coefficients $\mathcal{T}_m^{(r)}[s]$ in the form

$$\text{tr} \left(\sum_{r=0}^{\ell} \sum_{p=0}^{\ell'} \mathcal{X}^{(r)}[p] z^p u^{-r} \right)^m = \sum_{r, s \geq 0} \mathcal{T}_m^{(r)}[s] z^s u^{-r},$$

observe that the required property of these coefficients holds by [12, Theorem 5.4(b)]. This follows since the traces $\text{tr } \mathcal{X}^m$ with m running over the values specified in Table 1, are algebraically independent generators of the algebra of \mathfrak{g} -invariants of $\text{Fun } \mathfrak{g}$, where \mathcal{X} denotes the image of the matrix F under the isomorphism $S(\mathfrak{g}) \cong \text{Fun } \mathfrak{g}$ defined by the form (2.2). \square

Acknowledgements

I am grateful to Oksana Yakimova for providing very useful answers to my questions. The support of the Australian Research Council, grant DP180101825 is acknowledged.

Data Availability

The data that supports the findings of this study are available within the article.

References

- [1] T. Arakawa and A. Premet, *Quantizing Mishchenko–Fomenko subalgebras for centralizers via affine W -algebras*, Trans. Mosc. Math. Soc. **78** (2017), 217–234.
- [2] F. Berdjis, *A criterion for completeness of Casimir operators*, J. Math. Phys. **22** (1981), 1851–1856.
- [3] F. Berdjis and E. Beslmüller, *Casimir operators for F_4 , E_6 , E_7 , and E_8* , J. Math. Phys. **22** (1981), 1857–1860.
- [4] J. Brown and J. Brundan, *Elementary invariants for centralizers of nilpotent matrices*, J. Aust. Math. Soc. **86** (2009), 1–15.
- [5] H. S. M. Coxeter, *The product of the generators of a finite group generated by reflections*, Duke Math. J. **18** (1951), 765–782.
- [6] B. Feigin and E. Frenkel, *Affine Kac–Moody algebras at the critical level and Gelfand–Dikii algebras*, Int. J. Mod. Phys. A **7**, Suppl. 1A (1992), 197–215.
- [7] E. Frenkel, *Langlands correspondence for loop groups*, Cambridge Studies in Advanced Mathematics, 103. Cambridge University Press, Cambridge, 2007.
- [8] F. Geoffriau, *Homomorphisme de Harish-Chandra pour les algèbres de Takiff généralisées*, J. Algebra **171** (1995), 444–456.
- [9] V. Kac, *Vertex algebras for beginners*, University Lecture Series, 10. American Mathematical Society, Providence, RI, 1997.
- [10] D. Kuin', *Poincaré polynomials of certain compact homogeneous spaces*, Soviet Math. Dokl. **6** (1965), 115–119.
- [11] D. Kuin', *The Poincaré polynomials of compact homogeneous Riemannian spaces with irreducible stationary group* (Russian), Trudy Semin. Vektor. Tenzor. Analizu **14** (1968), 33–93.
- [12] T. Macedo and A. Savage, *Invariant polynomials on truncated multicurrent algebras*, J. Pure Appl. Algebra **223** (2019), 349–368.
- [13] A. I. Molev, *Casimir elements for certain polynomial current Lie algebras*, in “Group 21, Physical Applications and Mathematical Aspects of Geometry, Groups, and Algebras,” Vol. 1, (H.-D. Doebner, W. Scherer, P. Nattermann, Eds). World Scientific, Singapore, 1997, 172–176.

- [14] A. Molev, *Sugawara operators for classical Lie algebras*. Mathematical Surveys and Monographs 229. AMS, Providence, RI, 2018.
- [15] A. Molev, *Center at the critical level for centralizers in type A*, J. Algebra **566** (2021), 163–186.
- [16] A. I. Molev, E. Ragoucy and N. Rozhkovskaya, *Segal–Sugawara vectors for the Lie algebra of type G_2* , J. Algebra **455** (2016), 386–401.
- [17] D. Panyushev and O. Yakimova, *Takiff algebras with polynomial rings of symmetric invariants*, Transform. Groups **25** (2020), 609–624.
- [18] M. Raïs and P. Tauvel, *Indice et polynômes invariants pour certaines algèbres de Lie*, J. Reine Angew. Math. **425** (1992), 123–140.
- [19] M. Takeuchi, *On Pontrjagin classes of compact symmetric spaces*, J. Fac. Sci. Univ. Tokyo Sect. I **9** (1962), 313–328.
- [20] S. J. Takiff, *Rings of invariant polynomials for a class of Lie algebras*, Trans. Amer. Math. Soc. **160** (1971), 249–262.
- [21] C. Wendlandt, *The R-matrix presentation for the Yangian of a simple Lie algebra*, Comm. Math. Phys. **363** (2018), 289–332.
- [22] O. Yakimova, *Symmetrisation and the Feigin-Frenkel centre*, [arXiv:1910.10204](https://arxiv.org/abs/1910.10204).