

# HEINZ-TYPE MEAN CURVATURE ESTIMATES IN LORENTZ-MINKOWSKI SPACE

ATSUFUMI HONDA, YU KAWAKAMI, MIYUKI KOISO, AND SYUNSUKE TORI

*Dedicated to Professor Atsushi Kasue on the occasion of his 65th birthday*

ABSTRACT. We provide a unified description of Heinz-type mean curvature estimates under an assumption on the gradient bound for space-like graphs and time-like graphs in the Lorentz-Minkowski space. As a corollary, we give a unified vanishing theorem of mean curvature for these entire graphs of constant mean curvature.

## 1. INTRODUCTION

The celebrated theorem of Bernstein [6] (as for a simple proof, see [23]), stating that the only entire minimal graph in the Euclidean 3-space  $\mathbf{E}^3$  is a plane, has played a seminal role in the evolution of surface theory. For instance, by using this theorem, we can show the uniqueness theorem for an entire graph of constant mean curvature in  $\mathbf{E}^3$ . Indeed Heinz [15] proved that if  $\varphi(u^1, u^2)$  is a  $C^2$ -differentiable function defined on the open disk  $B^2(R)$  with center at the origin and radius  $R(> 0)$  in the  $(u^1, u^2)$ -plane, and the mean curvature  $H$  of the graph  $\Gamma_\varphi := \{(u^1, u^2, \varphi(u^1, u^2)) \in \mathbf{E}^3; (u^1, u^2) \in B^2(R)\}$  of  $\varphi$  satisfies  $|H| \geq \alpha > 0$ , where  $\alpha$  is constant, then  $R \leq 1/\alpha$ . As a corollary, we have a vanishing theorem of mean curvature for entire graphs of constant mean curvature, that is, if  $\Gamma_\psi$  is an entire graph of constant mean curvature, then  $H \equiv 0$ . Combining these results, we obtain that the only entire graph of constant mean curvature in  $\mathbf{E}^3$  is a plane (see [18, Section 2.1]). The Heinz result was extended to the case of graphic hypersurfaces in the Euclidean  $(n+1)$ -space  $\mathbf{E}^{n+1}$  by Chern [11] and Flanders [14].

We will study this subject for graphic hypersurfaces in the Lorentz-Minkowski space. Here we recall some basic definitions and fundamental facts. We denote by  $\mathbf{R}_1^{n+1} = (\mathbf{R}^{n+1}, \langle \cdot, \cdot \rangle_L)$  the Lorentz-Minkowski  $(n+1)$ -space with the Lorentz metric

$$\langle (x^1, \dots, x^n, x^{n+1}), (y^1, \dots, y^n, y^{n+1}) \rangle_L := x^1 y^1 + \dots + x^n y^n - x^{n+1} y^{n+1},$$

where  $(x^1, \dots, x^n, x^{n+1}), (y^1, \dots, y^n, y^{n+1}) \in \mathbf{R}^{n+1}$ . Let  $M^n$  be a smooth orientable  $n$ -manifold. An immersion  $f: M^n \rightarrow \mathbf{R}_1^{n+1}$  is called *space-like* if the induced metric  $g := f^* \langle \cdot, \cdot \rangle_L$  on  $M^n$  is Riemannian, and is called *time-like* if  $g$  on  $M^n$  is Lorentzian. A

---

2020 *Mathematics Subject Classification*. Primary 53A10; Secondary 53B30, 53C24, 53C42.

*Key words and phrases*. Heinz-type mean curvature estimate, Bernstein-type theorem, space-like graph, time-like graph, constant mean curvature.

The authors were partially supported by JSPS KAKENHI Grant Number JP18H04487, JP19K03463, JP19K14526, JP20H04642 and JP20H01801.

space-like (resp. time-like) immersion of constant mean curvature is called a *space-like* (resp. *time-like*) *constant mean curvature immersion*. In particular, a space-like immersion with vanishing mean curvature is called a *space-like maximal immersion* and a time-like immersion with vanishing mean curvature is called a *time-like minimal immersion*.

Let  $\psi: \Omega \rightarrow \mathbf{R}$  be a  $C^2$ -differentiable function defined on a domain  $\Omega$  in  $\mathbf{R}^n$ . We consider the graph  $\Gamma_\psi := \{(u^1, \dots, u^n, \psi(u^1, \dots, u^n)) \in \mathbf{R}_1^{n+1}; (u^1, \dots, u^n) \in \Omega\}$  of  $\psi$ . If  $\Omega = \mathbf{R}^n$ , then  $\Gamma_\psi$  is called an *entire graph* of  $\psi$  in  $\mathbf{R}_1^{n+1}$ . If the graph  $\Gamma_\psi$  of  $\psi$  is space-like, the gradient  $\nabla\psi := (\psi_{u^1}, \dots, \psi_{u^n})$  of  $\psi$  satisfies  $|\nabla\psi| := \sqrt{(\psi_{u^1})^2 + \dots + (\psi_{u^n})^2} < 1$  on  $\Omega$ , and

$$(1) \quad nH = \sum_{i=1}^n \frac{\partial}{\partial u^i} \left( \frac{\psi_{u^i}}{\sqrt{1 - |\nabla\psi|^2}} \right) = \operatorname{div} \left( \frac{\nabla\psi}{\sqrt{1 - |\nabla\psi|^2}} \right),$$

that is,

$$(1 - |\nabla\psi|^2) \sum_{i=1}^n \frac{\partial^2 \psi}{(\partial u^i)^2} + \sum_{i,j=1}^n \frac{\partial \psi}{\partial u^i} \frac{\partial \psi}{\partial u^j} \frac{\partial^2 \psi}{\partial u^i \partial u^j} = nH(1 - |\nabla\psi|^2)^{3/2}$$

holds. Here  $H$  is the mean curvature of  $\Gamma_\psi$  and  $\psi_{u^i} := \partial\psi/\partial u^i$ . If the graph  $\Gamma_\psi$  of  $\psi$  is time-like, its gradient  $\nabla\psi$  satisfies  $|\nabla\psi| > 1$  on  $\Omega$ , and

$$(2) \quad nH = \sum_{i=1}^n \frac{\partial}{\partial u^i} \left( \frac{\psi_{u^i}}{\sqrt{|\nabla\psi|^2 - 1}} \right) = \operatorname{div} \left( \frac{\nabla\psi}{\sqrt{|\nabla\psi|^2 - 1}} \right)$$

holds.

For entire space-like maximal graphs in  $\mathbf{R}_1^{n+1}$ , the following uniqueness theorem, called the Calabi-Bernstein theorem is well-known. This result was proved by Calabi [8] for  $n \leq 4$  and Cheng-Yau [10] for all  $n$ .

**FACT 1.1** (The Calabi-Bernstein theorem). *Any entire space-like maximal graph is a hyperplane in  $\mathbf{R}_1^{n+1}$ .*

We remark that an improvement of this theorem and a fluid mechanical interpretation of the duality between minimal graphs in  $\mathbf{E}^3$  and space-like maximal graphs in  $\mathbf{R}_1^3$  are discussed in [2] (cf. [7]).

On the other hand, for entire space-like constant mean curvature graphs, the uniqueness theorem does not hold in general. In fact, the graph of the function

$$(3) \quad \psi(u^1, \dots, u^n) = \left( (u^1)^2 + \dots + (u^n)^2 + \frac{1}{H^2} \right)^{1/2} \quad (H > 0)$$

is an entire space-like constant mean curvature  $H$  graph which is not a hyperplane. This graph is a hyperboloid. Many other entire space-like constant mean curvature graphs are constructed by Treibergs [27]. Hence the following questions naturally arise. What are the situations where Heinz-type mean curvature estimate and Bernstein-type uniqueness

theorem hold for space-like graphs in  $\mathbf{R}_1^{n+1}$ ? What are these situations for time-like graphs in  $\mathbf{R}_1^{n+1}$ ?

In this paper, we perform a systematic study for these questions. In particular, we give a unified description of Heinz-type mean curvature estimates under an assumption on the gradient bound for space-like graphs and time-like graphs in  $\mathbf{R}_1^{n+1}$  (Theorem 2.1). As a corollary, we obtain a unified vanishing theorem of mean curvature for entire graphs of constant mean curvature with a gradient bound (Corollary 2.4). In Section 3, by applying these results, we answer the questions mentioned above. Moreover we give a short commentary about the recent study by the authors for entire constant mean curvature graphs in  $\mathbf{R}_1^{n+1}$ .

Finally, the authors gratefully acknowledge the useful comments of Shintaro Akamine and Atsushi Kasue during the preparation of this paper.

## 2. MAIN RESULTS

The main theorem is stated as follows:

**THEOREM 2.1.** *Let  $B^n(R)$  be the open  $n$ -ball with center at the origin and radius  $R(> 0)$  in  $\mathbf{R}^n$ , and  $\psi(u^1, \dots, u^n)$  a  $C^2$ -differentiable function on  $B^n(R)$ . Suppose that there exist constants  $M > 0$  and  $k \in \mathbf{R}$  such that*

$$(4) \quad \frac{|\nabla\psi|}{\sqrt{|1 - |\nabla\psi|^2|}} \leq M \left( (u^1)^2 + \dots + (u^n)^2 \right)^k$$

*on  $B^n(R)$ . Assume that  $\Gamma_\psi$  is a space-like or time-like graph of  $\psi$  in  $\mathbf{R}_1^{n+1}$ . If the mean curvature  $H$  of  $\Gamma_\psi$  satisfies the inequality*

$$|H| \geq \alpha > 0,$$

*where  $\alpha$  is constant, then the following inequality holds:*

$$(5) \quad \alpha \leq MR^{2k-1}.$$

**PROOF.** Let  $\omega$  be the  $(n-1)$ -form given by

$$(6) \quad \omega = \sum_{i=1}^n (-1)^{i-1} \left( \frac{\psi_{u^i}}{\sqrt{|1 - |\nabla\psi|^2|}} \right) du^1 \wedge \dots \wedge \widehat{du^i} \wedge \dots \wedge du^n,$$

where  $\widehat{du^i}$  means that  $du^i$  is omitted from the wedge product. Then we can obtain  $d\omega = nH du^1 \wedge \dots \wedge du^n$  by (1) and (2). Take any positive number  $R'$  satisfying  $0 < R' < R$ . The Stokes theorem yields the following equality:

$$(7) \quad \int_{\overline{B^n(R')}} d\omega = \int_{\partial B^n(R')} \omega,$$

where  $\overline{B^n(R')}$  is the closure and  $\partial B^n(R')$  is the boundary of  $B^n(R')$ .

We may assume  $H \geq \alpha > 0$  by changing the direction of the normal vector if necessary. Let  $V_n$  be the volume of the unit closed  $n$ -ball  $\overline{B}^n(1)$  and  $A_{n-1}$  the volume of the unit  $(n-1)$ -sphere  $\partial B^n(1)$ . Then, by radial integration,  $nV_n = A_{n-1}$  holds. The absolute value of the left-hand side of (7) becomes

$$\begin{aligned} \left| \int_{\overline{B}^n(R')} d\omega \right| &= n \int_{\overline{B}^n(R')} H du^1 \cdots du^n \\ &\geq n\alpha \int_{\overline{B}^n(R')} du^1 \cdots du^n = n\alpha(R')^n V_n. \end{aligned}$$

On the other hand, by using the Cauchy-Schwarz inequality, the absolute value of the right-hand side of (7) becomes

$$\begin{aligned} \left| \int_{\partial B^n(R')} \omega \right| &= \left| \int_{\partial B^n(R')} \sum_{i=1}^n (-1)^{i-1} \left( \frac{\psi_{u^i}}{\sqrt{|1 - |\nabla\psi|^2|}} \right) du^1 \cdots \widehat{du^i} \cdots du^n \right| \\ &\leq \int_{\partial B^n(R')} \frac{|\nabla\psi|}{\sqrt{|1 - |\nabla\psi|^2|}} \left( \sum_{i=1}^n \left( du^1 \cdots \widehat{du^i} \cdots du^n \right)^2 \right)^{1/2}. \end{aligned}$$

By (4), we have

$$\begin{aligned} \left| \int_{\partial B^n(R')} \omega \right| &\leq M(R')^{2k} \int_{\partial B^n(R')} \left( \sum_{i=1}^n \left( du^1 \cdots \widehat{du^i} \cdots du^n \right)^2 \right)^{1/2} \\ &= M(R')^{n+2k-1} A_{n-1}. \end{aligned}$$

We thus obtain  $n\alpha(R')^n V_n \leq M(R')^{n+2k-1} A_{n-1}$ , that is,  $\alpha \leq M(R')^{2k-1}$ . The proof is completed by letting  $R' \rightarrow R$ .  $\square$

From the above argument, we give the following generalization of the result which was obtained by Salavessa [26, Theorem 1.5] for space-like graphs in  $\mathbf{R}_1^{n+1}$ .

**PROPOSITION 2.2.** *Let  $D$  be a relatively compact domain (i.e. its closure  $\overline{D}$  is compact) of  $\mathbf{R}^n$  with smooth boundary  $\partial D$ . Assume that  $\Gamma_\psi \subset \mathbf{R}_1^{n+1}$  is a space-like or time-like graph of a  $C^2$ -differentiable function  $\psi$ . Set  $m_D := \max_{\overline{D}} |\nabla\psi|$  if  $\Gamma_\psi$  is a space-like graph in  $\mathbf{R}_1^{n+1}$  and  $m_D := \min_{\overline{D}} |\nabla\psi|$  if  $\Gamma_\psi$  is a time-like graph in  $\mathbf{R}_1^{n+1}$ . For the mean curvature  $H$  of  $\Gamma_\psi$ , we have*

$$(8) \quad \min_{\overline{D}} |H| \leq \frac{1}{n} \frac{m_D}{\sqrt{|1 - (m_D)^2|}} \frac{A(\partial D)}{V(\overline{D})}.$$

Here  $V(\overline{D})$  (resp.  $A(\partial D)$ ) is the volume of  $\overline{D}$  (resp.  $\partial D$ ).

REMARK 2.3. To be precise, in [26, Theorem 1.5] Salavessa obtained this inequality for space-like graphs of  $M \times N$ , where  $M$  is a Riemannian  $n$ -manifold and  $N$  is an oriented Lorentzian 1-manifold.

Indeed, from the Stokes theorem for the  $(n-1)$ -form  $\omega$  given by (6), we obtain

$$(9) \quad \int_{\overline{D}} d\omega = \int_{\partial D} \omega.$$

Then the absolute value of the left-hand side of (9) becomes

$$\left| \int_{\overline{D}} d\omega \right| \geq n \min_{\overline{D}} |H| V(\overline{D}).$$

Since the function  $t/\sqrt{1-t^2}$  is monotone increasing on  $0 < t < 1$  and  $t/\sqrt{t^2-1}$  is monotone decreasing on  $t > 1$ , we have

$$\left| \int_{\partial D} \omega \right| \leq \frac{m_D}{\sqrt{|1-(m_D)^2|}} A(\partial D).$$

We have thus proved the inequality (8).

As a corollary of Theorem 2.1, we give the following unified vanishing theorem of mean curvature for entire space-like graphs and time-like graphs of constant mean curvature in  $\mathbf{R}_1^{n+1}$ .

COROLLARY 2.4. *Assume that  $\Gamma_\psi$  is an entire space-like or time-like graph of a  $C^2$ -differentiable function  $\psi(u^1, \dots, u^n)$  in  $\mathbf{R}_1^{n+1}$ . If the entire graph  $\Gamma_\psi$  has constant mean curvature and there exist constants  $M > 0$  and  $\varepsilon > 0$  such that*

$$(10) \quad \frac{|\nabla\psi|}{\sqrt{|1-|\nabla\psi|^2|}} \leq M \left( (u^1)^2 + \dots + (u^n)^2 \right)^{(1/2)-\varepsilon}$$

on  $\mathbf{R}^n$ , then its mean curvature must vanish everywhere.

PROOF. By Theorem 2.1, the mean curvature  $H$  of  $\Gamma_\psi$  satisfies

$$|H| \leq \frac{M}{R^{2\varepsilon}}$$

on  $B^n(R)$ . We obtain  $H \equiv 0$  by letting  $R \rightarrow +\infty$ . □

### 3. APPLICATIONS

3.1. **Space-like case.** Then  $|\nabla\psi| < 1$  holds. We give a geometric interpretation for  $|\nabla\psi|/\sqrt{1-|\nabla\psi|^2}$ . When the graph  $\Gamma_\psi$  is space-like,

$$(11) \quad \nu(= \nu(u^1, \dots, u^n)) = \frac{1}{\sqrt{1-|\nabla\psi|^2}}(\psi_{u^1}, \dots, \psi_{u^n}, 1)$$

is the time-like unit normal vector field  $\nu$  of  $\Gamma_\psi$ . Since  $e_{n+1} = (0, \dots, 0, 1) \in \mathbf{R}_1^{n+1}$  is also time-like, there exists a unique real-valued function  $\theta(= \theta(u^1, \dots, u^n)) \geq 0$  such that

$\langle \nu, e_{n+1} \rangle_L = -\cosh \theta$ . This function  $\theta$  is called the *hyperbolic angle* between  $\nu$  and  $e_{n+1}$  (see [22]). By simple calculation, we have

$$(12) \quad \sinh \theta = \frac{|\nabla \psi|}{\sqrt{1 - |\nabla \psi|^2}}.$$

We remark that López [21] studies the function  $|\nabla \psi|/\sqrt{1 - |\nabla \psi|^2}$  from the viewpoint of the Dirichlet problem.

From Theorem 2.1, we obtain the following Heinz-type mean curvature estimate.

**COROLLARY 3.1.** *Let  $\Gamma_\psi$  be a space-like graph of a  $C^2$ -differentiable function  $\psi(u^1, \dots, u^n)$  on  $B^n(R)$  in  $\mathbf{R}_1^{n+1}$  and  $\theta$  the hyperbolic angle between  $\nu$  and  $e_{n+1}$ . Suppose that there exist  $M > 0$  and  $k \in \mathbf{R}$  such that*

$$(13) \quad \sinh \theta \leq M \left( (u^1)^2 + \dots + (u^n)^2 \right)^k$$

*on  $B^n(R)$ . If the mean curvature  $H$  of  $\Gamma_\psi$  satisfies the inequality  $|H| \geq \alpha > 0$ , where  $\alpha$  is constant, then  $\alpha \leq MR^{2k-1}$  holds.*

By virtue of Fact 1.1 and Corollary 2.4, we obtain the following Bernstein-type theorem for entire space-like constant mean curvature graphs in  $\mathbf{R}_1^{n+1}$ .

**COROLLARY 3.2.** *If an entire space-like graph  $\Gamma_\psi$  in  $\mathbf{R}_1^{n+1}$  has constant mean curvature and there exist constants  $M > 0$  and  $\varepsilon > 0$  such that*

$$(14) \quad \sinh \theta \leq M \left( (u^1)^2 + \dots + (u^n)^2 \right)^{(1/2)-\varepsilon}$$

*on  $\mathbf{R}^n$ , then it must be a space-like hyperplane. Here  $\theta$  is the hyperbolic angle between  $\nu$  and  $e_{n+1}$ .*

Corollary 3.2 is optimal because there exists an example which is not congruent to a space-like hyperplane and satisfies (14) for  $\varepsilon = 0$ . In fact, the function  $\psi$  given by (3) is not linear and its graph has constant mean curvature. Moreover it satisfies

$$\sinh \theta = H \sqrt{(u^1)^2 + \dots + (u^n)^2},$$

which is the equality in (14) for  $\varepsilon = 0$ .

From the case where  $\varepsilon = 1/2$  in Corollary 3.2, we obtain the following Liouville-type result.

**COROLLARY 3.3.** *Let  $\Gamma_\psi$  be an entire space-like constant mean curvature graph of a  $C^2$ -differentiable function  $\psi(u^1, \dots, u^n)$  in  $\mathbf{R}_1^{n+1}$  and  $\theta$  the hyperbolic angle between  $\nu$  and  $e_{n+1}$ . If  $\sinh \theta$  is bounded on  $\mathbf{R}^n$ , then  $\Gamma_\psi$  must be a space-like plane.*

The following result is equivalent to Corollary 3.3.

COROLLARY 3.4. [4, Corollary 9.5] *If an entire space-like graph  $\Gamma_\psi$  in  $\mathbf{R}_1^{n+1}$  has constant mean curvature and there exists some constant  $C(< 1)$  such that  $|\nabla\psi| \leq C$  on  $\mathbf{R}^n$ , then it must be a space-like hyperplane.*

Indeed, if  $|\nabla\psi| \leq C$  holds on  $\mathbf{R}^n$ , then we obtain

$$\frac{|\nabla\psi|}{\sqrt{1 - |\nabla\psi|^2}} \leq \frac{C}{\sqrt{1 - C^2}}$$

on  $\mathbf{R}^n$ , that is,  $\sinh \theta$  is bounded because the function  $t/\sqrt{1 - t^2}$  is monotone increasing on  $0 < t < 1$ . By the same argument, the converse also holds, that is, if  $\sinh \theta$  is bounded on  $\mathbf{R}^n$ , then there exists some constant  $C(< 1)$  such that  $|\nabla\psi| \leq C$  on  $\mathbf{R}^n$ .

As described in [4, Section 9.3], Corollary 3.4 is a case of nonparametric form of the following value-distribution-theoretical property of the Gauss map.

FACT 3.5 ([1, 29], [24] for a first weaker version). *Let  $f: M^n \rightarrow \mathbf{R}_1^{n+1}$  be a complete space-like constant mean curvature immersion. If the image of the time-like unit normal vector field (i.e., Gauss map)  $\nu(M^n)$  is contained in a geodesic ball in the hyperbolic  $n$ -space  $\mathbf{H}^n(-1)$ , then  $f(M^n)$  is a hyperplane in  $\mathbf{R}_1^{n+1}$ .*

In fact, every entire space-like constant mean curvature graph in  $\mathbf{R}_1^{n+1}$  is complete (see [10], note that the converse statement is also true, that is, every complete space-like hypersurface in  $\mathbf{R}_1^{n+1}$  is an entire graph ([5])) and the image  $\nu(\mathbf{R}^n)$  is contained in a geodesic ball in  $\mathbf{H}^n(-1)$  if and only if  $\sqrt{1 - |\nabla\psi|^2}$  is bounded away from zero on  $\mathbf{R}^n$ . However the proof of Fact 3.5 is not as simple as our argument in this paper because it is obtained via an application of the Omori-Yau maximum principle ([9], [25], [30]). See the book [4] for details.

For a space-like graph in  $\mathbf{R}_1^{n+1}$ ,  $\sinh \theta$  can be replaced with  $1/\sqrt{1 - |\nabla\psi|^2}$  in (13) and (14). In fact,

$$\sinh \theta = \frac{|\nabla\psi|}{\sqrt{1 - |\nabla\psi|^2}} < \frac{1}{\sqrt{1 - |\nabla\psi|^2}}$$

holds. Thus we have the following result.

COROLLARY 3.6. *If an entire space-like graph  $\Gamma_\psi$  in  $\mathbf{R}_1^{n+1}$  has constant mean curvature and there exist constants  $M > 0$  and  $\varepsilon > 0$  such that*

$$\frac{1}{\sqrt{1 - |\nabla\psi|^2}} \leq M \left( (u^1)^2 + \cdots + (u^n)^2 \right)^{(1/2) - \varepsilon}$$

on  $\mathbf{R}^n$ , then it must be a space-like hyperplane.

From Corollary 3.6, we can show the following proposition which was given by Dong.

PROPOSITION 3.7 ([12]). Let  $\Gamma_\psi$  be an entire space-like constant mean curvature graph of a  $C^2$ -differentiable function  $\psi(u^1, \dots, u^n)$  in  $\mathbf{R}_1^{n+1}$ . If the function  $\psi$  satisfies

$$\frac{1}{\sqrt{1 - |\nabla\psi|^2}} = o(r) \quad \text{as } r \rightarrow +\infty,$$

where  $r = \sqrt{(u^1)^2 + \dots + (u^n)^2}$ , then  $\Gamma_\psi$  must be a space-like hyperplane.

REMARK 3.8. In [12], Dong showed a generalization of this result for an entire space-like graph of parallel mean curvature in the pseudo-Euclidean  $(m+n)$ -space  $\mathbf{R}_n^{m+n}$  of index  $n$ .

3.2. **Time-like case.** Then  $|\nabla\psi| > 1$  holds. By Theorem 2.1, we obtain the following Heinz-type estimate for the mean curvature of time-like graphs in  $\mathbf{R}_1^{n+1}$ .

COROLLARY 3.9. Let  $\Gamma_\psi$  be a time-like graph of a  $C^2$ -differentiable function  $\psi(u^1, \dots, u^n)$  on  $B^n(R)$  in  $\mathbf{R}_1^{n+1}$ . Suppose that there exist constants  $M > 0$  and  $k \in \mathbf{R}$  such that

$$(15) \quad \frac{|\nabla\psi|}{\sqrt{|\nabla\psi|^2 - 1}} \leq M \left( (u^1)^2 + \dots + (u^n)^2 \right)^k$$

on  $B^n(R)$ . If the mean curvature  $H$  of  $\Gamma_\psi$  satisfies the inequality

$$|H| \geq \alpha > 0,$$

where  $\alpha$  is constant, then the following inequality holds:

$$(16) \quad \alpha \leq MR^{2k-1}.$$

Moreover, we obtain the following result from Corollary 2.4.

COROLLARY 3.10. If an entire time-like graph  $\Gamma_\psi$  in  $\mathbf{R}_1^{n+1}$  has constant mean curvature  $H$  and there exist constants  $M > 0$  and  $\varepsilon > 0$  such that

$$(17) \quad \frac{|\nabla\psi|}{\sqrt{|\nabla\psi|^2 - 1}} \leq M \left( (u^1)^2 + \dots + (u^n)^2 \right)^{(1/2)-\varepsilon}$$

on  $\mathbf{R}^n$ , then it must be minimal (i.e.  $H \equiv 0$ ).

We note that there exist many entire time-like minimal graphs in  $\mathbf{R}_1^{n+1}$ . In fact, for a  $C^2$ -differentiable function  $h(t)$  of one variable which satisfies  $h'(t) > 0$  for all  $t \in \mathbf{R}$ ,  $\psi(u^1, \dots, u^n) := u^n + h(u^1)$  gives an entire time-like minimal graph  $\Gamma_\psi$  in  $\mathbf{R}_1^{n+1}$ .

We do not know whether Corollary 3.10 is optimal or not. However, for example, from the case where  $\varepsilon = 1/2$  and  $n = 2$  in Corollary 3.10, we provide the following result.

COROLLARY 3.11. Let  $\Gamma_\psi$  be an entire time-like constant mean curvature graph of a  $C^2$ -differentiable function  $\psi(u^1, u^2)$  in  $\mathbf{R}_1^3$ . If  $|\nabla\psi|/\sqrt{|\nabla\psi|^2 - 1}$  is bounded on  $\mathbf{R}^2$ , then  $\Gamma_\psi$  is a translation surface, that is,  $\psi(u^1, u^2) = h(u^1) + k(u^2)$ , where  $h(u^1)$  is a  $C^2$ -differentiable function of  $u^1$  and  $k(u^2)$  is a  $C^2$ -differentiable function of  $u^2$ .



The latter half of the statement of Corollary 3.11 follows from the following fact.

FACT 3.12 ([19, 20]). *Every entire time-like minimal graph in  $\mathbf{R}_1^3$  is a translation surface.*

**3.3. Closing Remark.** This paper is focused on the study of entire space-like graphs and time-like graphs in  $\mathbf{R}_1^{n+1}$ . Here we explain the recent development of the study of entire constant mean curvature graphs in  $\mathbf{R}_1^{n+1}$ . Let  $\psi: \Omega \rightarrow \mathbf{R}$  be a  $C^2$ -differentiable function defined on a domain  $\Omega$  in  $\mathbf{R}^n$ . A point where  $|\nabla\psi| > 1$  (resp.  $|\nabla\psi| < 1$ ,  $|\nabla\psi| = 1$ ) is said to be *space-like* (resp. *time-like*, *light-like*). We remark that the first fundamental form  $g$  of the graph  $\Gamma_\psi$  in  $\mathbf{R}_1^{n+1}$  is degenerate at each light-like point. If  $\Omega$  contains space-like points and time-like points, then the graph  $\Gamma_\psi$  is called of *mixed type*. Akamine, the first author, Umehara and Yamada proved the following fact.

FACT 3.13. [3, Corollary C] *An entire real analytic constant mean curvature graph in  $\mathbf{R}_1^{n+1}$  which has no time-like points and does have a light-like point must be a light-like hyperplane.*

See [3] (also [28]) for the precise definition of constant mean curvature graph which does not depend on its causal type. From this result, we know that any entire space-like constant mean curvature graph in  $\mathbf{R}_1^{n+1}$  with light-like points is a light-like hyperplane although there exist many nontrivial examples of entire space-like constant mean curvature graphs which have no light-like points. Moreover, the first and third authors, Kokubu, Umehara and Yamada [16, Remark 2.2] showed that there does not exist an entire real analytic nonzero constant mean curvature graph of mixed type in  $\mathbf{R}_1^{n+1}$ .

REMARK 3.14. Several entire zero mean curvature graphs of mixed type in  $\mathbf{R}_1^3$  were found in recent years. A systematic construction of this class is given in [13]. See [17] for details about mixed type surfaces in Lorentzian 3-manifolds.

## REFERENCES

- [1] R. AIYAMA, On the Gauss map of complete space-like hypersurfaces of constant mean curvature in Minkowski space, Tsukuba J. Math. **16** (1992), 353–361.
- [2] S. AKAMINE, M. UMEHARA AND K. YAMADA, Improvement of the Bernstein-type theorem for space-like zero mean curvature graphs in Lorentz-Minkowski space using fluid mechanical duality, Proc. Amer. Math. Soc. Ser. B **7** (2020), 17–27.
- [3] S. AKAMINE, A. HONDA, M. UMEHARA AND K. YAMADA, Bernstein-type theorem for zero mean curvature hypersurfaces without time-like points in Lorentz-Minkowski space, to appear in Bull. Braz. Math. Soc., New Series, <https://doi.org/10.1007/s00574-020-00196-8>.
- [4] L. J. ALÍAS, P. MASTROLIA, M. RIGOLI, Maximum principles and geometric applications, Springer Monographs in Mathematics, Springer, 2016.
- [5] L. J. ALÍAS, A. ROMERO, M. SÁNCHEZ, Uniqueness of complete spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes, Gen. Relativ. Gravit. **27**, 71–84 (1995).

- [6] S. BERNSTEIN, Sur un théorème de géométrie et ses applications aux équations aux dérivées partielles du type elliptique, *Comm. de la Soc. Math. de Kharkov (2éme sér.)* **15**, 38–45 (1915–1917).
- [7] L. BERS, *Mathematical aspects of subsonic and transonic gas dynamics*, John Wiley and Sons, 1958.
- [8] E. CALABI, Examples of Bernstein problem for some nonlinear equations, *Proc Symp. Pure Appl. Math.* **15** (1968), 223–230.
- [9] Q. CHEN, X. L. XIN, A generalized maximum principle and its applications in geometry, *Amer. J. Math.* **114** (1992), 355–366.
- [10] S.-Y. CHENG, S.-T. YAU, Maximal spacelike hypersurfaces in the Lorentz-Minkowski spaces, *Ann. of Math.* **104** (1976), 407–419.
- [11] S.-S. CHERN, On the curvatures of a piece of hypersurface in euclidean space, *Abh. Math. Sem. Univ. Hamburg*, **29** (1965), 77–91.
- [12] Y. DONG, Bernstein theorems for space-like graphs with parallel mean curvature and controlled growth, *J. Geom. Phys.* **58** (2008), 324–333.
- [13] S. FUJIMORI, Y. KAWAKAMI, M. KOKUBU, W. ROSSMAN, M. UMEHARA, K. YAMADA, Entire zero-mean curvature graphs of mixed type in Lorentz-Minkowski 3-space, *Q. J. Math.* **67** (2016), 801–837.
- [14] H. FLANDERS, Remark on mean curvature, *J. London Math. Soc.* **41** (1966), 364–366.
- [15] E. HEINZ, Über Flächen mit eineindeutiger Projektion auf eine Ebene, deren Krümmungen durch Ungleichungen eingeschränkt sind, *Math. Ann.* **129** (1955), 451–454.
- [16] A. HONDA, M. KOISO, M. KOKUBU, M. UMEHARA, K. YAMADA, Mixed type surfaces with bounded mean curvature in 3-dimensional space-times, *Diff. Geom. and its Appl.* **52** (2017), 64–77.
- [17] A. HONDA, K. SAJI, K. TERAMOTO, Mixed type surfaces with bounded Gaussian curvature in three-dimensional Lorentzian manifolds, *Adv. Math.* **365** (2020), to appear, <https://doi.org/10.1016/j.aim.2020.107036>.
- [18] K. KENMOTSU, Surfaces with constant mean curvature, Translated from the 2000 Japanese original by Katsuhiko Moriya and revised by the author, *Translations of Mathematical Monographs*, Vol. 221, Am. Math. Soc., Providence (2003).
- [19] M. A. MAGID, The Bernstein problem for timelike surfaces, *Yokohama Math. J.* **37** (1990), 125–137.
- [20] T. K. MILNOR, A conformal analog of Bernstein’s theorem for timelike surfaces in Minkowski 3-space, in: *The Legacy of Sonya Kovalevskaya*, in: *Contemp. Math.* **64** (1987), 123–130.
- [21] R. LÓPEZ, The Dirichlet problem of the constant mean curvature equation in Lorentz-Minkowski space and in Euclidean space, *Mathematics special volume Differential Geometry: Theory and Applications*, to appear, arXiv: 1912.07866.
- [22] B. O’NEILL, *Semi-Riemannian geometry with applications to relativity*, Academic Press, London (1983).
- [23] J. C. C. NITSCHKE, Elementary proof of Bernstein’s theorem on minimal surfaces, *Ann. of Math.* **66** (1957), 543–544.
- [24] B. PALMER, The Gauss map of a spacelike constant mean curvature hypersurface of Minkowski space, *Comment. Math. Helv.* **65** (1990), 52–57.
- [25] H. OMORI, Isometric immersions of Riemannian manifolds, *J. Math. Soc. Japan* **95** (1967), 205–214.
- [26] I. M. C. SALAVESSA, Spacelike graphs with parallel mean curvature, *Bull. Belg. Math. Soc. Simon Stevin* **15** (2008), 65–76.
- [27] A. TREIBERGS, Entire spacelike hypersurfaces of constant mean curvature in Minkowski space, *Invent. Math.* **66** (1982), 39–56.

- [28] M. UMEHARA, K. YAMADA, Hypersurfaces with light-like points in a Lorentzian manifold, *J. Geom. Anal.* **29** (2019), 3405–3437.
- [29] Y. L. XIN, On the Gauss image of a spacelike hypersurface with constant mean curvature in Minkowski space, *Comment. Math. Helv.* **66** (1991), 590–598.
- [30] S. -T. YAU, Harmonic functions on complete Riemannian manifolds, *Comm. Pure Appl. Math.* **28** (1975), 201–228.

DEPARTMENT OF APPLIED MATHEMATICS,  
FACULTY OF ENGINEERING,  
YOKOHAMA NATIONAL UNIVERSITY,  
79-5, TOKIWADAI, HODOGATA-KU, YOKOHAMA, 240-8501, JAPAN  
*E-mail address:* honda-atsufumi-kp@ynu.ac.jp

FACULTY OF MATHEMATICS AND PHYSICS,  
INSTITUTE OF SCIENCE AND ENGINEERING,  
KANAZAWA UNIVERSITY,  
KANAZAWA, 920-1192, JAPAN  
*E-mail address:* y-kwkami@se.kanazawa-u.ac.jp

INSTITUTE OF MATHEMATICS FOR INDUSTRY,  
KYUSHU UNIVERSITY,  
744, MOTOOKA NISHI-KU, FUKUOKA, 819-0395, JAPAN  
*E-mail address:* koiso@math.kyushu-u.ac.jp

GRADUATE SCHOOL OF NATURAL SCIENCE AND TECHNOLOGY,  
KANAZAWA UNIVERSITY,  
KANAZAWA, 920-1192, JAPAN  
*E-mail address:* shunsuke-1350@stu.kanazawa-u.ac.jp