

CIRCUM- AND INCONIC INVARIANTS OF 3-PERIODICS IN THE ELLIPTIC BILLIARD

DAN REZNIK AND RONALDO GARCIA

ABSTRACT. A Circumconic passes through a triangle's vertices; an Inconic is tangent to the sidelines. We study the variable geometry of certain conics derived from the 1d family of 3-periodics in the Elliptic Billiard. Some display intriguing invariances such as aspect ratio and pairwise ratio of focal lengths.

Keywords elliptic billiard, periodic trajectories, triangle center, circumconic, circumellipse, circumhyperbola, conservation, invariance, invariant.

MSC 51M04 and 37D50 and 51N20 and 51N35 and 68T20

1. INTRODUCTION

Given a triangle, a *circumconic* passes through its three vertices and satisfies two additional constraints, e.g., center or perspector¹ location. An *Inconic* touches each side and is centered at a specified location. Both these objects are associated² with simple lines on the plane [29, Circumconic, Inconic] and therefore lend themselves to agile algebraic manipulation.

We study properties and invariants of such conics derived from a 1d family of triangles: 3-periodics in an Elliptic Billiard (EB): these are triangles whose bisectors coincide with normals to the boundary (bounces are elastic), see Figure 1.

Amongst all planar curves, the EB is uniquely integrable [15]. It can be regarded as a special case of Poncelet's Porism [6]. These two properties imply two classic invariances: N -periodics have constant perimeter and envelop a confocal Caustic. The seminal work is [27] and more recent treatments include [18, 25].

We have shown 3-periodics also conserve the Inradius-to-Circumradius³ ratio which implies an invariant sum of cosines, and that their *Mittenpunkt*⁴ is stationary at the EB center [24]. Indeed many such invariants have been effectively generalized for $N > 3$ [1, 3].

We have also studied the loci of 3-periodic Triangle Centers over the family: out of the first 100 listed in [16], 29 sweep out ellipses (a remarkable fact on its own) with the remainder sweeping out higher-order curves [9]. Related is the study of loci described by the Triangle Centers of the Poristic Triangle family [20]. We have also showed [22] the aspect ratio of the *Circumbilliard*

¹Where reference and polar triangles are perspective [29].

²The Isogonal and Isotomic conjugation of such conics are lines [29, Perspector].

³As does the Poristic Family [7].

⁴Where lines drawn from each Excenter thru sides' midpoints meet.




FIGURE 1. 3-periodics (blue) in the Elliptic Billiard (EB, black): normals to the the boundary at vertices (black arrows) are bisectors. The family is constant-perimeter and envelopes a confocal Caustic (brown). This family conserves the ratio inradius-to-circumradius and has a stationary Mittenpunkt at the EB center. **Video:** [21, PL#01].

(a circumellipse which is an EB for a generic triangle) is invariant over the 1d family of *Poristic*, triangles, with fixed Incircle and Circumcircle [28, 20]. See [21, PL#07].

Main Results:

- Theorem 1 in Section 2: The ratio of semi-axis of the X_1 -centered Circumellipse is invariant over the 3-periodic family. We conjecture this to be the case for a 1d-family of circumellipses.
- Theorem 2 in Section 3: The focal lengths of two special circum-hyperbola (Feuerbach and Excentral Jerabek) is constant over the 3-periodic family.
- Theorems 4 and 3 and in Section 4 show the aspect ratios of two important Excentral Inconics are invariant and that one of the Inconics is a 90° -rotated copy of the X_1 -centered Circumellipse.

A reference table with all Triangle Centers, Lines, and Symbols appears in Appendix D. Videos of many of the experiments are assembled on Table 2 in Section 5.

2. INVARIANTS IN CIRCUMELLIPSES

Let the boundary of the EB satisfy:

$$(1) \quad f(x, y) = \left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1.$$

Where $a > b > 0$ denote the EB semi-axes throughout the paper. Below we use *aspect ratio* as the ratio of an ellipse's semi-axes. When referring to Triangle Centers we adopt Kimberling X_i notation [16], e.g., X_1 for the Incenter, X_2 for the Barycenter, etc., see Table 3 in Appendix D.

The following five-parameter equation is assumed for all circumconics not passing through $(0, 0)$.

$$(2) \quad 1 + c_1x + c_2y + c_3xy + c_4x^2 + c_5y^2 = 0$$




FIGURE 2. A reference triangle is shown (blue) as well as its Medial (red). The latter sides divide the plane into 7 regions, including the Medial's interior. When a Circumconic center lies on any of the shaded regions (resp. unshaded) it is an Ellipse (resp. Hyperbola). Parabolas have centers at infinity. For illustration, the X_1 and X_9 -centered Circumellipses and the X_{11} -centered Feuerbach Hyperbola are shown. Note that over the family of 3-periodics, a given Circumconic may alternate between Ellipse and Hyperbola, e.g., when centered on X_4, X_5, X_6 , etc.

In Appendix A we provide a method to compute the c_i given 3 points a conic must pass through as well as its center.

The Medial Triangle divides the plane in 7 regions, see Figure 2. The following is a known fact [2, 13]:

Remark 1. *If the center of a Circumconic lies within 4 of these (resp. the remainder 3), the conic will be an Ellipse (resp. Hyperbola).*

Centers X_1, X_2 , and X_9 are always interior to the Medial Triangle [16], so the Circumconics $E_i, i = 1, 2, 9$ centered on them will ellipses, Figure 3. E_2 is the *Steiner Circumellipse*, least-area over all possible Circumellipses [29], and E_9 is the Elliptic Billiard.

It is known that E_1 intersects the EB and the Circumcircle at X_{100} , the Anticomplement of the Feuerbach Point. Also that E_2 intersects E_9 at X_{190} , the Yff Parabolic Point [14, 16]. These two ellipses intersect at X_{664} ⁵ [19].

Given a generic triangle T :

Proposition 1. *The axes of E_1 are parallel to E_9 's.*

The proof is in Appendix B.

Theorem 1. *Over the 3-periodic family, the major η_1' (resp. minor η_1) semiaxis lengths of E_1 are variable, though their ratio is invariant. These are given by:*

⁵This is the isogonal conjugate of X_{663} , i.e., \mathcal{L}_{663} mentioned before is coincidentally its *Trilinear Polar* [29].




FIGURE 3. E_1 (green) and E_2 (brown) are Circumellipses to the orbit (blue) centered on X_1 (green) and X_2 (brown). The former (resp. latter) intersect the EB at X_{100} (resp. X_{190}). They intersect at the vertices and at X_{664} . E_1 axes remain parallel to the EB over the orbit family, and the ratio of their lengths is constant. The axes of E_2 are only parallel to the EB's when the orbit is an isosceles. **Video:** [21, PL#08].

$$\begin{aligned}\eta'_1 &= R + d, & \eta_1 &= R - d \\ \frac{\eta'_1}{\eta_1} &= \frac{R + d}{R - d} = \frac{1 + \sqrt{1 - 2\rho}}{\rho} - 1 > 1\end{aligned}$$

Where $d = |X_1 X_3| = \sqrt{R(R - 2r)}$.

Proof. Calculate the ratio using vertex locations (see [10]) for an isosceles orbit, and then verify with a Computer Algebra System (CAS) the expression holds over the entire family. A related proof appears in [23]. \square

Note: experimentally η'_1 is maximal (resp. minimal) when the 3-periodic is an isosceles with axis of symmetry parallel to the EB's minor (resp. major) axis.

Proposition 2. *The axes of E_2 are only parallel to E_9 if T is isosceles.*

See Appendix B.

2.1. Parallel-Axis Pencil. The Feuerbach Circumhyperbola of a Triangle is a rectangular hyperbola⁶ centered on X_{11} [29]. Peter Moses has contributed a stronger generalization [19]:

Remark 2. *The pencil of Circumconics whose centers lie on the Feuerbach Circumhyperbola F_{med} of the Medial Triangle have mutually-parallel axes.*

The complement⁷ of X_{11} is X_{3035} [16] so F_{med} is centered there, see Figure 4. The following is a list of Circumellipses whose centers lie on F_{med} [19]: X_i , $i=1, 3, 9, 10^8, 119, 142, 214, 442, 600, 1145, 2092, 3126, 3307$,

⁶Since it passes through the Orthocenter X_4 [29].

⁷The 2:1 reflection of a point about X_2 .

⁸Notice X_{10} is the Incenter of the Medial. Interestingly, X_8 , the Incenter of the ACT, does not belong to this select group.




FIGURE 4. An $a/b = 1.5$ EB is shown (black) centered on X_9 as well as a sample 3-periodic (blue). Also shown are Circumellipses centered on X_i , $i = 1, 3, 10, 142$, whose centers lie on the Feuerbach Circumhyperbola of the Medial Triangle (both shown red), centered on X_{3035} , the complement of X_{11} . Notice all conics drawn (including the Circumhyp) have axes parallel to the EB and all Circumellipses pass through X_{100} . Note: the Circumellipse centered on X_3 is the Circumcircle, its axes, drawn diagonally, are immaterial.

3308, 3647, 5507, 6184, 6260, 6594, 6600, 10427, 10472, 11517, 11530, 12631, 12639, 12640, 12864, 13089, 15346, 15347, 15348, 17057, 17060, 18258, 18642, 19557, 19584, 22754, 34261, 35204.

Proposition 3. *A circumellipse has center on F_{med} iff it passes through X_{100} .*

A proof appears in Appendix C. The following has been observed experimentally:

Conjecture 1. *Over the family of 3-periodics, all Circumellipses in Moses' pencil conserve the ratio of their axes.*

3. A SPECIAL PAIR OF CIRCUMHYPERBOLAE

Here we study invariants of two well-known Circumhyperbolae⁹: the Feuerbach and Jerabek Hyperbolas F and J [29, Jerabek Hyperbola]. Both are rectangular since they contain X_4 [29]. The former is centered on X_{11} and the latter on X_{125} . With respect to 3-periodics no invariants have been detected for J . However, the Jerabek J_{exc} of the Excentral Triangle, which

⁹Its centers lie in the unshaded regions in Figure 2.

passes through the Excenters and is centered on X_{100} ¹⁰, does produce interesting invariants¹¹. F is known to pass through X_1 and X_9 of its reference triangle. Interestingly J_{exc} also passes through X_1 and X_9 . This stems from the fact that J passes through X_4 and X_6 . Since the Excentral Triangle is always acute [5], its X_4 is X_1 . Likewise, the excentral X_6 is X_9 .

The Isogonal Conjugate of a Circumconic is a line [29, Circumconic]. Remarkably:

Remark 3. *The Isogonal conjugate of F with respect to a reference triangle and that of J_{exc} with respect to the Excentral one is line $X_1X_3 = \mathcal{L}_{650}$.*

The first part is well-known [29, Feuerbach Hyperbola]. For the second part, consider that J is the Isogonal Conjugate of the Euler Line [29, Jerabek Hyperbola]. The Euler Line of the Excentral Triangle passes through its X_4 and X_5 which are X_1 and X_3 in the reference 3-periodic.

Referring to Figure 5:

Proposition 4. *J_{exc} intersects E_9 in exactly two locations.*

Proof. Let $s_i, i = 1, 2, 3$ refer to 3-periodic sidelengths. The perspector of J_{exc} is $X_{649} = s_1(s_2 - s_3) :: (\text{cyclical})$ [16]. Therefore the trilinears $x : y : z$ of J_{exc} satisfy [30]:

$$J_{exc} : s_1(s_2 - s_3)x^2 + s_2(s_3 - s_1)y^2 + s_3(s_1 - s_2)z^2 = 0.$$

Notice the above is satisfied for the Excenters $[1 : 1 : -1]$, $[1 : -1 : 1]$ and $[-1 : 1 : 1]$. As $X_1 = [1 : 1 : 1]$ and $X_9 = s_2 + s_3 - s_1 :: (\text{cyclical})$ it follows that $J_{exc}(X_1) = J_{exc}(X_9) = 0$.

Eliminating variable x , the intersection of $J_{exc} = 0$ and $E_9 = 0$ is given by the quartic:

$$\begin{aligned} & s_2(s_1 - s_3)k_1y^4 + 2s_2(s_1 - s_3)k_1k_2y^3z \\ & + 2s_3(s_1 - s_2)k_1k_2yz^3 + s_3(s_1 - s_2)k_1z^4 = 0 \end{aligned}$$

With $k_1 = (s_1 + s_2 - s_3)^2$ and $k_2 = s_1 + s_3 - s_2$. The discriminant of the above equation is:

$$-432[(s_2 - s_3)(s_1 - s_3)(s_1 - s_2)(s_1 + s_3 - s_2)^2(s_1 - s_2 - s_3)^2(s_1 + s_2 - s_3)^2(s_1s_2s_3)]^2$$

Since it is negative, there will be two real and two complex solutions [4]. \square

Proposition 5. *F intersects the X_9 -centered Circumellipse at X_{1156} .*

Proof. The perspector of X_9 is X_1 and that of X_{11} is $X_{650} = (s_3 - s_3)(s_3 + s_3 - s_1) :: (\text{cyclical})$. Therefore, the trilinears $x : y : z$ of F and E_9 satisfy:

¹⁰The Excentral's X_{125} [29].

¹¹The Feuerbach Hyperbola F_{exc} has not yet yielded any detectable invariants over the 3-periodic family.

$$\begin{aligned}
F &: (s_2 - s_3)(s_2 + s_3 - s_1)/x + \\
& (s_3 - s_1)(s_3 + s_1 - s_2)/y + \\
& (s_1 - s_2)(s_1 + s_2 - s_3)/z = 0 \\
E_9 &: 1/x + 1/y + 1/z = 0.
\end{aligned}$$

X_{1156} is given by $1/[(s_2 - s_3)^2 + s_1(s_2 + s_3 - 2s_1)] ::$ (cyclic). This point can be readily checked to satisfy both of the above. \square

Given a generic triangle T , the following two claims are known:

Proposition 6. *The asymptotes of both F and J_{exc} are parallel to the X_9 -centered circumconic, i.e., c_4 and c_5 in (2) vanish.*

Proof. To see the first part, consider that since the Caustic is centered on X_9 and tangent to the 3-periodics, it is the (stationary) Mandart Inellipse I_9 of the family [29]. This inconic is known to have axes parallel to the asymptotes of F [11]. Since the Caustic is confocal with the EB, F asymptotes must be parallel to the EB axes.

Secondly, 3-periodics are the Orthic Triangles of the Excentrals, therefore the EB is the (stationary) Excentral's Orthic Inconic [29]. The latter's axes are known to be parallel to the asymptotes of the Jerabek hyperbola. [29, Orthic Inconic].

An alternate, algebraic proof appears in Appendix B. \square

Let λ (resp. λ') be the focal length of F (resp. J_{exc}).

Remark 4. *Isosceles 3-periodics have $\lambda' = \lambda = 0$.*

To see this consider the sideways isosceles 3-periodic with $P_1 = (a, 0)$. P_2 and P_3 will lie on the 2nd and 3rd quadrants at $(-a_c, \pm y')$, where $a_c = a(\delta - b^2)/(a^2 - b^2)$ is the length of the Caustic major semi-axis [9]. X_1 and X_4 will lie along the 3-periodic's axis of symmetry, i.e., the x-axis. To pass through all 5 points, F degenerates to a pair of orthogonal lines: the x-axis and the vertical line $x = -a_c$. The foci will collapse to the point $(-a_c, 0)$. A similar degeneracy occurs for the upright isosceles, i.e., when $P_1 = (0, b)$, namely, the foci collapse to $(0, -b_c)$, where $b_c = b(a^2 - \delta)/(a^2 - b^2)$ is the Caustic minor semi-axis length.

Theorem 2. *For all non-isosceles 3-periodics, λ'/λ is invariant and given by:*

$$(3) \quad \frac{\lambda'}{\lambda} = \sqrt{2/\rho} > 2$$

Proof. Assume the EB is in the form of (1). Let the 3-periodic be given by $P_i = (x_i, y_i), i = 1, 2, 3$. F passes through the P_i, X_1 and $X_9 = (0, 0)$. The asymptotes of F are parallel to the EB axes, therefore this hyperbola is

given by $c_1x + c_2y + c_3xy = 0$ and $\lambda^2 = |8c_1c_2/c_3^2|$, where:

$$\begin{aligned}
c_1 &= y_2y_3(x_2 - x_3)x_1^2 + (x_2^2y_3 - x_3^2y_2 - y_2^2y_3 + y_2y_3^2)x_1y_1 \\
&\quad + y_2y_3(x_2 - x_3)y_1^2 - (x_2y_3 - x_3y_2)(x_2x_3 + y_2y_3)y_1 \\
(4) \quad c_2 &= x_2x_3(y_2 - y_3)x_1^2 + (x_2x_3^2 - x_2^2x_3 - x_2y_3^2 + x_3y_2^2)x_1y_1 \\
&\quad + (x_2y_3 - x_3y_2)(x_2x_3 + y_2y_3)x_1 - x_2x_3(y_2 - y_3)y_1^2 \\
c_3 &= (x_2y_3 - x_3y_2)x_1^2 + (x_3^2y_2 - x_2^2y_3 + y_2^2y_3 - y_2y_3^2)x_1 \\
&\quad + (x_3y_2 - x_2y_3)y_1^2 + (x_2^2x_3 - x_2x_3^2 + x_2y_3^2 - x_3y_2^2)y_1
\end{aligned}$$

Let $P'_i = (x'_i, y'_i)$, $i = 1, 2, 3$ be the Excenters. They are given by

$$\begin{aligned}
P'_1 &= \left(\frac{-x_1s_1 + x_2s_2 + x_3s_3}{s_2 + s_3 - s_1}, \frac{-y_1s_1 + y_2s_2 + y_3s_3}{s_2 + s_3 - s_1} \right) \\
(5) \quad P'_2 &= \left(\frac{x_1s_1 - x_2s_2 + x_3s_3}{s_3 + s_1 - s_2}, \frac{y_1s_1 - y_2s_2 + y_3s_3}{s_3 + s_1 - s_2} \right) \\
P'_3 &= \left(\frac{x_1s_1 + x_2s_2 - x_3s_3}{s_1 + s_2 - s_3}, \frac{y_1s_1 + y_2s_2 - y_3s_3}{s_1 + s_2 - s_3} \right)
\end{aligned}$$

Here, $s_1 = |P_2 - P_3|$, $s_2 = |P_1 - P_3|$ and $s_3 = |P_1 - P_2|$.

Since J_{exc} is also centered on the origin and has horizontal/vertical asymptotes, J_{exc} is given by $c'_1x + c'_2y + c'_3xy = 0$, and $(\lambda')^2 = |8c'_1c'_2/(c'_3)^2|$, where c'_i are constructed as (4) replacing (x_i, y_i) with (x'_i, y'_i) .

Consider a right-triangle¹² 3-periodic, e.g., with $P_1(x^\perp, y^\perp)$ [22, Proposition 5]:

$$x^\perp = \frac{a^2\sqrt{a^4 + 3b^4 - 4b^2\delta}}{c^3}, \quad y^\perp = \frac{b^2\sqrt{-b^4 - 3a^4 + 4a^2\delta}}{c^3}$$

P_2 and P_3 are obtained explicitly [8]. From these obtain c_i using (4). Using (5) obtain P'_i and the c'_i . Finally, obtain a symbolic expression for λ'/λ . After some manipulation and simplification with a Computer Algebra System (CAS), we obtain (3) which we call a *candidate*.

Parametrize the 3-periodic family with $P_1(t) = (a \cos t, b \sin t)$ and using the sequence above arrive at an expression for λ'/λ in terms of t . Subtract that from the right-triangle candidate. After some algebraic manipulation and CAS simplification verify the subtraction vanishes, i.e., λ'/λ is independent of t . \square

3.1. Focal Length Extrema. Let $P_1(t) = (a \cos t, b \sin t)$. While their ratio is constant, λ and λ' undergo three simultaneous maxima in $t \in (0, \pi/2)$, see Figure 6. In fact, the following additional properties occur at configurations of maximal focal length (we omit the rather long algebraic proofs), see Figure 7:

- F' is tangent to the Caustic at $\pm X_{11}$.
- J'_{exc} is tangent to the EB at $\pm X_{100}$, i.e., at $\mp X_{1156}$ (see below).

Remark 5. Like F , J'_{exc} intersects the EB at X_{1156} .

¹²We found this to best simplify the algebra.




FIGURE 5. An $a/b = 1.5$ EB is shown (black) as well as a sample 3-periodic (blue), the confocal Caustic (brown), and the Excentral Triangle (green). The 3-periodic's Feuerbach Circumhyperbola F (orange) passes through its three vertices as well as X_1 , X_9 , and X_4 . The Excentral's Jerabek Circumhyperbola J_{exc} (purple) passes through the three Excenters, as well as X_1 , X_9 and X_{40} (not shown). Two invariants have been detected over the orbit family: (i) the asymptotes (dashed) of both F and J_{exc} stay parallel to the EB axes, (ii) the ratio of focal lengths is constant (focal axis appears dashed). F intersects the Billiard at X_{1156} .

This happens because X_{1156} is the reflection of X_{100} about X_9 . If the latter is placed on the origin, then $X_{1156} = -X_{100}$, and J'_{exc} passes through $\pm X_{100}$.

Let F' and J'_{exc} be copies of F and J_{exc} translated by $-X_{11}$ and $-X_{100}$ respectively, i.e., they become concentric with the EB (focal lengths are unchanged). Since their asymptotes are parallel to the EB axes and centered on the origin, their equations will be of the form:

$$F' : xy = k'_F, \quad J'_{exc} : xy = k'_J$$

Remark 6. $\lambda = 2\sqrt{2k'_F}$, $\lambda' = 2\sqrt{2k'_J}$, $\lambda'/\lambda = \sqrt{k'_J/k'_F} = \sqrt{2/\rho}$.

4. INCONIC INVARIANTS

A triangle's *Inconic* touches its three sides while satisfying two other constraints, e.g., the location of its center. Similar to Circumconics, if the latter is interior to the 4 shaded regions in Figure 2 it is an ellipse, else it is a hyperbola. Lines drawn from each vertex to the Inconic tangency points concur at the perspector or Brianchon Point B [29].




FIGURE 6. Focal lengths λ, λ' of F, J_{exc} vs the parameter t in $P_1(t) = (a \cos t, b \sin t)$ are shown red and green. The solid (resp. dashed) curves correspond to $a/b = 1.5$ (resp. $a/b = 1.3$). In the first quadrant there are 3 maxima. λ'/λ (blue) remain constant for the whole interval.




FIGURE 7. Two snapshots of J and F_{exc} drawn solid blue and solid green, respectively, for $a/b = 1.5$. Also shown (dashed) are copies F' and J'_{exc} of both hyperbolas translated so they are dynamically concentric with the EB (translate J by $-X_{11}$ and F_{exc} by $-X_{100}$). Their focal lengths λ, λ' are identical to the original ones; their focal axes are collinear and shown the dashed diagonal through the EB center. Notice that like F, J' also intersects the EB at X_{1156} . **Left:** $t = 10.1^\circ$, showing an intermediate value of ether focal length. **Right:** $t = 6.2^\circ$, focal lengths are at a maximum. When this happens, the translated copy of F (resp. J_{exc}) is tangent to the Caustic (resp. EB) at X_{11} (resp. X_{1156}). **Video:** [21, PL#09,10]

Let the Inconic center C be specified by Barycentrics¹³ $g(s_1, s_2, s_3)$ (cyclic), then B is given by $1/[g(s_2, s_3, s_1) + g(s_3, s_1, s_2) - g(s_1, s_2, s_3)]$ (cyclic) [26]. For example, consider the Inconic centered on X_1 (the Incircle), i.e., $g = s_2 s_3$ (cyclic). Then $B = 1/(s_1 s_3 + s_1 s_2 - s_2 s_3)$. Dividing by the product $s_1 s_2 s_3$ obtain $B = 1/(s_2 + s_3 - s_1)$ (cyclic), confirming that the perspector of the Incircle is the Gergonne Point X_7 [29, Perspector]. The contact points are the vertices of the Cevian triangle through B .

Above we identified the confocal Caustic with the Mandart Inellipse I_9 [29] of the 3-periodic family, i.e., it is a stationary ellipse centered on X_9 and axis-aligned with the EB, Figure 1. Its semi-axes a_c, b_c are given by [8]:

¹³Barycentrics g can be easily converted to Trilinears f via: $f(s_1, s_2, s_3) = g(s_1, s_2, s_3)/s_1$ (cyclic) [30].

$$a_c = \frac{a(\delta - b^2)}{a^2 - b^2}, \quad b_c = \frac{b(a^2 - \delta)}{a^2 - b^2}.$$

Similarly, the X_9 -centered Inconic of the family of Excentral Triangles is the stationary EB, i.e., the EB is the Orthic Inconic [29] of the Excentrals.

4.1. Excentral X_3 -Centered Inconic. No particular invariants have yet been found for any of the Inconics to 3-periodics with centers in X_i , $i = 3, 4, \dots, 12$. Let I_3 denote the X_3 -centered inconic. Its Brianchon Point is X_{69} and its foci aren't named centers [19].

Remark 7. I_3 is always an ellipse.

To see this consider the Circumcenter of an acute, right, or obtuse triangle lies inside, on a vertex, or opposite to the interior of the Medial, respectively, i.e., within one of the 4 shaded regions in Figure 2.

Let I'_3 denote the X_3 -centered Inconic of the Excentral Triangle. Its Brianchon Point is X_{69} of the Excentral, i.e., X_{2951} of the reference 3-periodic [19], Figure 8(left). Let μ'_3, μ_3 denote I'_3 major and minor semi-axes. In [23] we show:

Proposition 7. *Let T be a triangle and T' its excentral. The Inconic I'_3 to T' on its Circumcenter (X_{40} of T) is a 90° -rotated copy of the E_1 , the X_1 -centered Circumconic, i.e.:*

$$(6) \quad \mu'_3 = \eta'_1, \quad \mu_3 = \eta_1$$

Remark 8. *As E_1 , I'_3 is axis-aligned with the EB and contains X_{100} , Figure 8 (left).*

Theorem 3. *Over 3-periodics, μ'_3 and μ_3 are variable though their ratio is invariant. These are given by:*

$$\begin{aligned} \mu'_3 &= R + d, & \mu_3 &= R - d \\ \frac{\mu'_3}{\mu_3} &= \frac{R + d}{R - d} = \frac{1 + \sqrt{1 - 2\rho}}{\rho} - 1 > 1 \end{aligned}$$

As above, $d = \sqrt{R(R - 2r)}$, and $\rho = r/R$.

Proof. The above was initially a candidate an isosceles 3-periodic and then verified it held any 3-periodic configuration with a CAS. A related proof appears in [23]. \square

4.2. Excentral X_5 -Centered (MacBeath) Inconic. Above the locus of the Excenters is identified with the Excentral MacBeath Circumconic E'_6 . The MacBeath Inconic I_5 of a triangle is centered on X_5 and has foci on X_4 and X_3 . Its Brianchon Point is X_{264} , and it can be both an ellipse or a hyperbola [29, MacBeath Inconic]. No invariants have yet been found for I_5 .

Consider I'_5 , the MacBeath Inconic of the Excentral Triangle, Figure 8(right). The center and foci of I'_5 with respect to the reference triangle are X_3 , X_1 , and X_{40} , respectively, and its Brianchon is X_{1742} [19]. Unlike I'_3 , the axes of I'_5 are askew with respect to the EB.




FIGURE 8. **Left:** The X_3 -centered Inconic I'_3 (red) of the Excentral Triangle (green), whose center is X_{40} in terms of the 3-periodic (blue). Its Brianchon Point is X_{2951} [19]. Its axes are aligned with the EB which it intersects at X_{100} , and its aspect ratio is invariant. The X_1 -centered Circumconic E_1 (green) is a 90° -rotated copy of I'_3 . **Right:** The MacBeath Inconic I'_5 of the Excentral (red) is centered on the latter's X_5 and has foci on X_4 and X_3 . These correspond to X_3 , X_1 , and X_{40} of the reference triangle (blue). Its Brianchon Point is X_{1742} [19]. Though its axes are askew with respect to the EB, its aspect ratio is invariant over the 3-periodic family. **Video:** [21, PL#11].

Remark 9. I'_5 is always an ellipse.

This is due to the fact that the Excentral is acute as is its homothetic Medial. Since X_5 is the latter's Circumcenter, it must lie inside it. Let μ'_5, μ_5 denote I'_5 major and minor semi-axes.

Theorem 4. Over the 3-periodic family μ'_5 and μ_5 are variable and their ratio is invariant. These are given by:

$$\mu'_5 = R, \quad \mu_5 = \sqrt{R^2 - d^2}$$

$$\frac{\mu'_5}{\mu_5} = \frac{R}{\sqrt{R^2 - d^2}} = \frac{1}{\sqrt{2\rho}}$$

The above was derived for an isosceles 3-periodic and shown to work for any a, b . A related proof is given in [23].

5. CONCLUSION

The various quantities and invariants described above are summarized on Table 1.

Length	Meaning	Value	Invariant
η'_1	Major semiaxis of E_1	$R + d$	–
η_1	Minor semiaxis of E_1	$R - d$	–
η'_1/η_1	E_1 aspect ratio	$(1 - \rho + \sqrt{1 - 2\rho})/\rho$	yes
λ'	J_{exc} focal length	–	–
λ	F focal length	–	–
λ'/λ	Ratio of focal lengths	$\sqrt{2/\rho}$	yes
μ'_3	Major semiaxis of I'_3	$= \eta'_1$	–
μ_3	Minor semiaxis of I'_3	$= \eta_1$	–
μ'_3/μ_3	I'_3 aspect ratio	$= \eta'_1/\eta_1$	yes
μ'_5	Major semiaxis of I'_5	R	–
μ_5	Minor semiaxis of I'_5	$\sqrt{R^2 - d^2}$	–
μ'_5/μ_5	I'_5 aspect ratio	$1/\sqrt{2\rho}$	yes

TABLE 1. Summary of Circumconic and Inconic Quantities. $\rho = r/R$ and $d = \sqrt{R(R - 2r)}$.

Videos mentioned above have been placed on a [playlist](#) [21]. Table 2 contains quick-reference links to all videos mentioned, with column “PL#” providing video number within the playlist.

Additionally to Conjecture 1 we submit the following questions to the reader:

- Can expressions be derived for λ' and λ (we only provide one for their ratio)?
- Can alternate proofs be found for Theorems 1 and 2 with tools of algebraic and/or projective geometry?
- Are there other notable circumconic pairs which exhibit interesting invariants?
- Can any of the invariants cited above be generalized to N -periodics?
- Are there other Inconic invariants over 3-periodics and/or their derived triangles?
- Are there interesting properties for the loci of the *foci* of Feuerbach and/or Jerabek Circumhyperbolas?
- The Yff Circumparabola whose focus is on X_{190} is shown in [21, PL#12] over 3-periodics. Are there interesting invariants?
- Does any Triangle Circumcubic display interesting invariants over 3-periodics? We found none for the Thomson Cubic shown in [21, PL#13]. How about the Darboux, Neuberg, Lucas, and myriad others catalogued in [12].

ACKNOWLEDGMENTS

We would like to thank Peter Moses and Clark Kimberling, for their prompt help with dozens of questions. We would like to thank Boris Odehnal for his help with some proofs. A warm thanks goes out to Profs. Jair Koiller and Daniel Jaud who provided critical editorial help.

PL#	Title	Section
01	Mittenpunkt stationary at EB center	1
07	Invariant Aspect Ratio of Circumbilliard of Poristic Family	1
08	The X_1 - and X_2 -centered Circumellipses	2
09	Orbit Feuerbach and Excentral Jerabek Circumhyperbolas	3
10	Invariant Focal Length Ratio for F and J_{exc}	3
11	Excentral MacBeath and X_3 -Centered Inconics: Invariant Aspect Ratio	4
12	The Yff Circumparabola of 3-Periodics	5
13	The Thomson Cubic of 3-Periodics	5

TABLE 2. Videos mentioned in the paper. Column “PL#” indicates the entry within the playlist [21]

The second author is fellow of CNPq and coordinator of Project PRONEX/CNPq/ FAPEG 2017 10 26 7000 508.

REFERENCES

- [1] Akopyan, A., Schwartz, R., Tabachnikov, S.: Billiards in ellipses revisited (2020). URL <https://arxiv.org/abs/2001.02934> 1
- [2] Akopyan, A., Zaslavsky, A.: Geometry of Conics. Amer. Math. Soc., Providence, RI (2007) 3
- [3] Bialy, M., Tabachnikov, S.: Dan Reznik’s identities and more (2020). URL <https://arxiv.org/abs/2001.08469> 1
- [4] Burnside, W.S., Panton, A.W.: The theory of equations: With an introduction to the theory of binary algebraic forms. 2 volumes. Dover Publications, Inc., New York (1960) 6
- [5] Coxeter, H.S.M., Greitzer, S.L.: Geometry Revisited, *New Mathematical Library*, vol. 19. Random House, Inc., New York (1967) 6
- [6] Dragović, V., Radnović, M.: Poncelet Porisms and Beyond: Integrable Billiards, Hyperelliptic Jacobians and Pencils of Quadrics. *Frontiers in Mathematics*. Springer, Basel (2011). URL <https://books.google.com.br/books?id=QcOmDAEACAAJ> 1
- [7] Gallatly, W.: The modern geometry of the triangle. Francis Hodgson (1914) 1
- [8] Garcia, R.: Elliptic billiards and ellipses associated to the 3-periodic orbits. *American Mathematical Monthly* **126**(06), 491–504 (2019). URL <https://doi.org/10.1080/00029890.2019.1593087> 8, 10
- [9] Garcia, R., Reznik, D., Koiller, J.: Loci of 3-periodics in an elliptic billiard: why so many ellipses? (2020). URL <https://arxiv.org/abs/2001.08041> 1, 7, 19
- [10] Garcia, R., Reznik, D., Koiller, J.: New properties of triangular orbits in elliptic billiards (2020). URL <https://arxiv.org/abs/2001.08054> 4
- [11] Gibert, B.: Generalized Mandart conics. *Forum Geom.* **4**, 177–198 (2004) 7
- [12] Gibert, B.: Cubics in the triangle plane (2020). URL <https://bernard-gibert.pagesperso-orange.fr/> 13
- [13] Glaeser, G., Stachel, H., Odehnal, B.: The universe of conics. Springer Spektrum, Berlin (2016). DOI 10.1007/978-3-662-45450-3. URL <https://doi.org/10.1007/978-3-662-45450-3>. From the ancient Greeks to 21st century developments 3
- [14] Grozdev, S., Dekov, D.: The computer program “Discoverer” as a tool of mathematical investigation. *International Journal of Computer Discovered Mathematics (IJCDM)* (2014). URL <http://www.ddekov.eu/j/2014/JCGM201405.pdf> 3

- [15] Kaloshin, V., Sorrentino, A.: On the integrability of Birkhoff billiards. *Phil. Trans. R. Soc. A* (376) (2018). DOI <https://doi.org/10.1098/rsta.2017.0419> 1
- [16] Kimberling, C.: Encyclopedia of triangle centers (2019). URL <https://faculty.evansville.edu/ck6/encyclopedia/ETC.html> 1, 2, 3, 4, 6, 16
- [17] Kimberling, C.: Central lines of triangle centers (2020). URL <https://faculty.evansville.edu/ck6/encyclopedia/CentralLines.html> 19
- [18] Lynch, P.: Integrable elliptic billiards and ballyards. *European Journal of Physics* **41**(1), 015,005 (2019). DOI 10.1088/1361-6404/ab4748. URL <https://arxiv.org/abs/1907.09295> 1
- [19] Moses, P.: Various questions regarding centers and perspectors of circumconics and inconics. Private Communication (March, 2020) 3, 4, 11, 12, 19, 20
- [20] Odehnal, B.: Poristic loci of triangle centers. *J. Geom. Graph.* **15**(1), 45–67 (2011) 1, 2
- [21] Reznik, D.: Playlist for “Circumphenomena of 3-Periodics in Elliptic Billiards” (2020). URL <https://bit.ly/2WDPeVk> 2, 4, 10, 12, 13, 14
- [22] Reznik, D., Garcia, R.: The circumbilliard: an elliptic billiard for a generic triangle (2020). In prepatation 1, 8
- [23] Reznik, D., Garcia, R.: New invariants for the poristic triangle family (2020). In preparation 4, 11, 12
- [24] Reznik, D., Garcia, R., Koiller, J.: Can the elliptic billiard still surprise us? *Math Intelligencer* **42** (2019). DOI 10.1007/s00283-019-09951-2. URL <https://rdcu.be/b2cg1> 1
- [25] Rozikov, U.A.: *An Introduction To Mathematical Billiards*. World Scientific Publishing Company (2018) 1
- [26] Stothers, W.: Further notes on circumconics and inconics (2001). URL <http://www.maths.gla.ac.uk/wws/cabripages/misc/candi.htm> 10
- [27] Tabachnikov, S.: *Geometry and Billiards*, *Student Mathematical Library*, vol. 30. American Mathematical Society, Providence, RI (2005). DOI 10.1090/stml/030. URL <http://www.personal.psu.edu/sot2/books/billiardsgeometry.pdf>. Mathematics Advanced Study Semesters, University Park, PA 1
- [28] Weaver, J.H.: Invariants of a poristic system of triangles. *Bull. Amer. Math. Soc.* **33**(2), 235–240 (1927). DOI 10.1090/S0002-9904-1927-04367-1. URL <https://doi.org/10.1090/S0002-9904-1927-04367-1> 2
- [29] Weisstein, E.: Mathworld (2019). URL <http://mathworld.wolfram.com> 1, 3, 4, 5, 6, 7, 9, 10, 11
- [30] Yiu, P.: A tour of triangle geometry (2003). URL <http://math.fau.edu/Yiu/TourOfTriangleGeometry/MAAFlorida37040428.pdf> 6, 10

APPENDIX A. COMPUTING A CIRCUMCONIC

Let a Circumconic have center $M = (x_m, y_m)$ Equation (2) is subject to the following 5 constraints¹⁴: it must be satisfied for vertices P_1, P_2, P_3 , and its gradient must vanish at M :

$$f(P_i) = 0, \quad i = 1, 2, 3$$

$$\frac{dg}{dx}(x_m, y_m) = c_1 + c_3 y_m + 2c_4 x_m = 0$$

$$\frac{dg}{dy}(x_m, y_m) = c_2 + c_3 x_m + 2c_5 y_m = 0$$

Written as a linear system:

¹⁴If M is set to X_9 one obtains the Circumbilliard.

$$\begin{bmatrix} x_1 & y_1 & x_1 y_1 & x_1^2 & y_1^2 \\ x_2 & y_2 & x_2 y_2 & x_2^2 & y_2^2 \\ x_3 & y_3 & x_3 y_3 & x_3^2 & y_3^2 \\ 1 & 0 & y_m & 2x_m & 0 \\ 0 & 1 & x_m & 0 & 2y_m \end{bmatrix} \cdot \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \\ c_5 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ -1 \\ 0 \\ 0 \end{bmatrix}$$

Given sidelengths s_1, s_2, s_3 , the coordinates of $X_9 = (x_m, y_m)$ can be obtained by converting its Trilinears ($s_2 + s_3 - s_1 :: \dots$) to Cartesians [16].

Principal axes' directions are given by the eigenvectors of the Hessian matrix H (the jacobian of the gradient), whose entries only depend on c_3, c_4 , and c_5 :

$$(7) \quad H = J(\nabla g) = \begin{bmatrix} 2c_4 & c_3 \\ c_3 & 2c_5 \end{bmatrix}$$

The ratio of semiaxes' lengths is given by the square root of the ratio of H 's eigenvalues:

$$(8) \quad a/b = \sqrt{\lambda_2/\lambda_1}$$

Let $U = (x_u, y_u)$ be an eigenvector of H . The length of the semiaxis along u is given by the distance t which satisfies:

$$g(M + tU) = 0$$

This yields a two-parameter quadratic $d_0 + d_2 t^2$, where:

$$\begin{aligned} d_0 &= 1 + c_1 x_m + c_4 x_m^2 + c_2 y_m + c_3 x_m y_m + c_5 y_m^2 \\ d_2 &= c_4 x_u^2 + c_3 x_u y_u + c_5 y_u^2 \end{aligned}$$

The length of the semi-axis associated with U is then $t = \sqrt{-d_0/d_2}$. The other axis can be computed via (8).

The eigenvectors (axes of the conic) of H are given by the zeros of the quadratic form

$$q(x, y) = c_3(y^2 - x^2) + 2(c_2 - c_5)xy$$

APPENDIX B. CIRCUMELLIPSES OF ELEMENTARY TRIANGLE

Let a triangle T have vertices $P_1 = (0, 0)$, $P_2 = (1, 0)$ and $P_3 = (u, v)$ and sidelengths s_1, s_2, s_3 . Using the linear system in Appendix A, one can obtain implicit equations for the circumellipses E_9, E_1, E_2 centered on T 's Mittenpunkt X_9 , Incenter X_1 , and Barycenter X_2 , respectively:

$$\begin{aligned}
E_9(x, y) &= v^2 x^2 - v(s_1 - s_2 - 1 + 2u)xy + ((s_1 - s_2 - 1)u + u^2 + s_2)y^2 \\
&\quad - v^2 x + ((s_1 - s_2 - 1)u + u^2 + s_2)y^2 + v(u - s_2)y = 0 \\
E_1(x, y) &= (L - 2)v^2 x^2 + (L - 2s_2 - 2u)(L - 2)vxy \\
&\quad + (-L^2 u + (2u + 1)Ls_2 + (u^2 + 2u)L - 2s_2^2 - 4us_2 - 2u^2)y^2 \\
&\quad - (L - 2)v^2 x - v(Ls_2 - uL - 2s_2^2 + 2u)y \\
E_2(x, y) &= v^2 x^2 + v(1 - 2u)xy + (u^2 - u + 1)y^2 - v^2 x + v(u - 1)y = 0 \\
s_1 &= \sqrt{(u - 1)^2 + v^2}, \quad s_2 = \sqrt{u^2 + v^2}, \quad L = s_1 + s_2 + 1
\end{aligned}$$

Consider the quadratic forms

$$\begin{aligned}
q_9(x, y) &= v(s_1 - s_2 + 2u - 1)x^2 + 2((s_2 - s_1)u - s_2 - u^2 + u + v^2)xy \\
&\quad + v(1 - 2u - s_1 + s_2)y^2 \\
q_1(x, y) &= -v(L - 2)(L - 2s_2 - 2u)x^2 + v(L - 2)(L - 2s_2 - 2u)y^2 \\
&\quad + 2(L^2 u - (2u + 1)Ls_2 + (v^2 - u^2 - 2u)L + 4s_2 u + 4u^2)xy \\
q_2(x, y) &= v(2u + 1)x^2 + 2(-u^2 + v^2 + u - 1)xy + v(1 - 2u)y^2
\end{aligned}$$

The axes of E_9 (resp. E_1) are defined by the zeros of q_9 (resp. q_1). Using the above equations it is straightforward to show that the axes of E_1 and E_9 are parallel.

The axes of E_2 and E_9 are parallel if and only if $(u - 1)^2 + v^2 = 1$ or $u^2 + v^2 = 1$; this means that the triangle is isosceles.

The implicit equations of the circumhyperbolas F passing through the vertices of the orbit centered on X_{11} and J_{exc} passing through the vertices of the excentral triangle and centered on X_{100} are:

$$\begin{aligned}
F(x, y) &= v^3(2u - 1)(x^2 - y^2) + v^3(1 - 2u)x \\
&\quad + [(s_2^3 + (u - 1)s_2^2 - us_2)s_1 + (2u - 1)s_2^2 - us_1^2 s_2 - u^4 - 4u^2 v^2 + v^4 + 2uv^2]xy \\
&\quad + [s_1^2 u^2 s_2 + (-us_2^3 - u(u - 1)s_2^2 + s_2 u^2)s_1 + us_2^4 - v^2 s_2^2 - u^3(2u - 1)]y = 0 \\
J_{exc}(x, y) &= 4v^3(2u - 1)(x^2 - y^2) \\
&\quad + [(4s_2^3 + 4(u - 1)s_2^2 - 4us_2)s_1 - 4us_1^2 s_2 - 4s_2^2 \\
&\quad - 4u^4 - 16u^2 v^2 + 4v^4 + 8u^3 + 16uv^2]xy \\
&\quad + [(2(-s_1 s_2^3 + ((1 - s_1)u - v^2 + s_1)s_2^2 + us_1(s_1 + 1)s_2 + u(u - 2)s_1^2))v]x \\
&\quad + [(4 - 4u)s_2((u^2 - (\frac{1}{2}s_1 + 1)u + \frac{1}{2}s_1 + \frac{1}{2})s_2 + \frac{1}{2}us_1(s_1 + 1) - \frac{1}{2}s_2^2(s_2 + s_1))]y \\
&\quad - us_1^2 v s_2 + (vs_2^3 + (-1 + u)vs_2^2 - uvs_2)s_1 + vs_2^4 - (2u^2 - 2u + 1)vs_2^2 = 0
\end{aligned}$$

Using the above equations it is straightforward to show that the axes of E_9 and asymptotes of F and J_{exc} are parallel.

APPENDIX C. CIRCUMELLIPSES WITH PARALLEL AXES

Consider a triangle with vertices $A = (u, v)$, $B = (-1, 0)$ and $C = (1, 0)$. Let $s_2 = |A - B|$ and $s_3 = |B - C|$. The equation of F_{med} is given by:

$$\begin{aligned}
(9) \quad F_{med}(x, y) = & 4uv^3(x^2 - y^2)(2(u-1)s_3s_2^2 - (2(u-1)s_3^2 - 2(u^2 + v^2 - 1)s_3)s_2 \\
& 2(u^2 - 1)^2 + 2v^2(4u^2 - v^2))xy \\
& + (-((u^2 + v^2 - 1)s_3 + s_3^2(u-1))vs_2 + s_2^2(u+1)vs_3 \\
& - v(v^4 + (u^2 - 1)^2)x \\
& + (2u^2 + 2v^2 - 2)uv^2 y = 0
\end{aligned}$$

A one parameter family of circumellipses passing through A , B , C and X_{100} is given by:

$$\begin{aligned}
(10) \quad E_b(x, y) = & uv^3(bxy - 4x^2) \\
& + ((b(u^2 + v^2 - 1)s_3 + b(u-1)s_3^2)s_2 - b(u+1)s_2^2s_3 \\
& - v^2(b(4u^2 - v^2) + 4uv) - b(u^2 - 1)^2)y^2 \\
& - v((b(u^2 + v^2 - 1)s_2 - b(u+1)s_2^2)s_3 + b(u-1)s_3^2s_2 \\
& - b(u^2 + v^2 - 1)(u^2 - v^2 + uv - 1))y + 4uv^3 = 0
\end{aligned}$$

It is straightforward to verify that the family E_b is axis aligned (independent of b). Denoting the center of E_b by (x_c, y_c) it follows, using CAS, that $F_{med}(x_c, y_c) = 0$.

The reciprocal follows similarly.

APPENDIX D. TABLE OF SYMBOLS

Tables 3 and 4 lists most Triangle Centers and symbols mentioned in the paper.

Center	Meaning	Note
X_1	Incenter	Locus is Ellipse
X_2	Barycenter	Perspector of Steiner Circum/Inellipses
X_3	Circumcenter	Locus is Ellipse, Perspector of M
X_4	Orthocenter	Exterior to EB when 3-periodic is obtuse
X_5	Center of the 9-Point Circle	
X_6	Symmedian Point	Locus is Quartic [9]
X_7	Gergonne Point	Perspector of Incircle
X_8	Nagel Point	Perspector of I_9 , X_1 of ACT Incircle
X_9	Mittenpunkt	Center of (Circum)billiard
X_{10}	Spieler Point	Incenter of Medial
X_{11}	Feuerbach Point	on confocal Caustic
X_{40}	Bevan Point	X_3 of Excentral
X_{69}	X_6 of the ACT	Perspector of I_3
X_{100}	Anticomplement of X_{11}	On Circumcircle and EB, J_{exc} center
X_{125}	Center of Jerabek Hyperbola J	
X_{142}	X_9 of Medial	Midpoint of X_9X_7 , lies on $L(2, 7)$
X_{144}	Anticomplement of X_7	Perspector of ACT and its Intouch Triangle
X_{168}	X_9 of the Excentral Triangle	Non-elliptic Locus
X_{190}	Focus of the Yff Parabola	Intersection of E_2 and the EB
X_{264}	Isotomic Conjugate of X_3	Perspector of I_5
X_{649}	Cross-difference of X_1, X_2	Perspector of J_{exc}
X_{664}	Trilinear Pole of $L(2, 7)$	Intersection of E_1 and E_2 [19]
X_{650}	Cross-difference of X_1, X_3	Perspector of F
X_{1156}	Isogonal Conjugate of Schröder Point X_{1155}	intesection of F with EB
X_{1742}	Mimosa Transform of X_{212}	Perspector of I'_5
X_{2951}	Excentral-Isogonal Conjugate of X_{57}	Perspector of I'_3
X_{3035}	Complement of X_{11}	Center of F_{med}
X_{3659}	X_{11} of Excentral Triangle	Center of F_{exc}
$L(2, 7)$	ACT-Medial Mittenpunkt Axis	Line \mathcal{L}_{663} [17]
$L(1, 3)$	Isogonal Conjugate of F and J_{exc}	Line \mathcal{L}_{650} [17]

TABLE 3. Kimberling Centers and Central Lines mentioned in the paper.

DAN REZNIK, DATA SCIENCE CONSULTING,, RIO DE JANEIRO, RJ, BRAZIL
E-mail address: dan@dat-sci.com

RONALDO GARCIA, INST. DE MATEMÁTICA E ESTATÍSTICA,, UNIV. FEDERAL DE
 GOIÁS,, GOIÂNIA, GO, BRAZIL
E-mail address: ragarcia@ufg.br

Symbol	Meaning	Note
a, b	EB semi-axes	$a > b > 0$
P_i, s_i	Vertices and sidelengths of 3-periodic	invariant $\sum s_i$
P'_i	Vertices of the Excentral Triangle	
a_c, b_c	Semi-axes of confocal Caustic	
r, R, ρ	Inradius, Circumradius, r/R	ρ is invariant
δ	Oft-used constant	$\sqrt{a^4 - a^2b^2 + b^4}$
d	Distance $ X_1X_3 $	$\sqrt{R(R - 2r)}$
P^\perp	Obtuse 3-periodic limits on EB	
F, J	Feuerbach, Jerabek Hyperbola	Centers X_{11}, X_{125} , Perspectors X_{650}, X_{647}
F_{exc}	F of Excentral Triangle	Center X_{3659} [19]
J_{exc}	J of Excentral Triangle	Center X_{100} , Perspector X_{649}
F', J'_{exc}	F, J_{exc} translated by $-X_{11}, -X_{100}$	Origin-centered
F_{med}	F of Medial	Center X_{3035} [19]
E_i	Circumellipse centered on X_i	Axes parallel to E_9 if X_i on F_{med}
E'_6	Excentral MacBeath Circumellipse	Center X_9 , Perspector X_{40}
I_3, I_5	Inellipses Centered on X_3, X_5	Perspectors X_{69}, X_{264}
I_9	Mandart Inellipse	Perspector X_8
I'_3	Excentral I_3	Center X_{40} , Perspector X_{2951} [19]
I'_5	Excentral I_5	Center X_3 , Perspector X_{1742} [19]
η'_i, η_i	Major and minor semiaxis of E_i	Invariant ratio if X_i on F_{med}
μ'_i, μ_i	Major and minor semiaxis of I'_i	Invariant ratio for $i = 3, 5$
λ', λ	Focal lengths of J_{exc}, F (and J'_{exc}, F')	Invariant ratio

TABLE 4. Symbols mentioned in the paper.