

Geodesic motion in Bogoslovsky-Finsler Spacetimes

M. Elbistan^{1*}, P. M. Zhang^{1†}, N. Dimakis^{2‡}, G. W. Gibbons^{3§}, P. A. Horvathy^{4¶}

¹ *School of Physics and Astronomy,
Sun Yat-sen University, Zhuhai, China*

² *Center for Theoretical Physics,
College of Physical Science and Technology,
Sichuan University,
Chengdu 610065, China*

³*D.A.M.T.P., Cambridge University,
Wilberforce Road,
Cambridge CB3 0WA, United Kingdom*

⁴*Institut Denis Poisson,
Tours University – Orléans University,
UMR 7013 (France)*

(Dated: July 6, 2020)

* email:elbistan@impcas.ac.cn.

† corresponding author. email:zhangpm5@mail.sysu.edu.cn

‡ email:nsdimakis@gmail.com

§ email:G.W.Gibbons@damtp.cam.ac.uk

¶ email:horvathy@lmpt.univ-tours.fr

Abstract

We study the free motion of a massive particle moving in the background of a Finslerian deformation of a plane gravitational wave in Einstein's general relativity. The deformation is a curved version of a one-parameter family of relativistic Finsler structures introduced by Bogoslovsky, which are invariant under a certain deformation of Cohen and Glashow's very special relativity group ISIM(2). The partially broken Carroll symmetry we derive using Baldwin-Jeffery-Rosen coordinates allows us to integrate the geodesics equations. The transverse coordinates of timelike Finsler-geodesics are identical to those of the underlying plane gravitational wave for any value of the Bogoslovsky-Finsler parameter b . We then replace the underlying plane gravitational wave with a homogenous pp-wave solution of the Einstein-Maxwell equations. We conclude by extending the theory to the Finsler-Friedmann-Lemaitre model.

Phys. Rev. D 102, 024014. DOI: 10.1103/PhysRevD.102.024014

PACS numbers: 04.20.-q Classical general relativity;

04.30.-w Gravitational waves

02.20.Sv Lie algebras of Lie groups;

Contents

I. Introduction	3
II. Finsler spaces	5
III. Bogoslovsky-Finsler metrics	9
IV. Geodesics	12
V. Partially broken Carroll symmetry	17
VI. Prolongation vectors and symmetries	21
VII. An Einstein-Maxwell example	23
VIII. Bogoslovsky-Finsler-Friedmann-Lemaître Model	24
A. The Λ CDM model	25
B. The choice of null vector field	26
C. Hubble friction	26
D. Conformal flatness	27
E. Redshifting	28
F. A possible choice for $f(\cdot)$	29
IX. Bogoslovsky-Finsler-Friedmann-Lemaître Geodesics	30
X. Conclusion	34
Acknowledgments	36
References	36

I. INTRODUCTION

Our present fundamental physical theories are based on local Lorentz invariance and hence on local isotropy. This leads naturally to the introduction of pseudo-Riemannian geometry and its associated metric tensor. It has long been known, however that a “principle of

relativity” can be made compatible with anisotropy by deforming the Lorentz group by the inclusion of dilations [1] (although the experiments of Hughes, and Drever indicate that the anisotropy must be very weak [2]).

Currently there is also a great deal of activity exploring the astrophysics and cosmology of alternative gravitational theories based on standard Lorentzian geometry. Laboratory tests of local Lorentz invariance are very well developed and have reached impressive levels of precision.

Riemann himself envisaged more general geometries. An elegant construction combining these ideas was provided some time ago by Bogoslovsky [3, 4] (for more recent accounts, see [5]). In what is now known as Finsler geometry, the line element is a general homogeneous function of degree 1 in displacements, rather than the square root of a quadratic form.

The theory proposed by Bogoslovsky, which is the main subject of interest of this paper, has turned out to be relevant for attempts to accommodate a proposal of Cohen and Glashow [1], accounting for weak CP violation in the standard model of particle physics, in the gravitational background [6, 7].

The first significant application of Finsler geometry to physics is due to Randers [8] who pointed out that the world line of a particle of mass m and electric charge e extremizes the action

$$S_0 = \int \mathcal{L}_0 d\tau = - \int m \sqrt{-g_{\mu\nu} dx^\mu dx^\nu} + e A_\mu dx^\mu ; \quad (\text{I.1})$$

where τ is an arbitrary parameter and A_μ is the electromagnetic potential. Randers applied this idea to Kaluza-Klein theory. Further studies followed [9–12] ; it has also been applied to the gravitomagnetic effects occurring in stationary spacetimes [13]. For more recent work on Finsler spaces, see [14–24]. Null geodesics and causality are considered in particular in [24].

The aim of the present paper is to contribute to the physical applications of the Finslerian generalization of general relativity by exploring the motion of freely moving massive particles in the background of Bogoslovsky-Finsler deformations of plane gravitational waves and spatially flat Friedmann-Lemaître cosmologies.

II. FINSLER SPACES

In this section, we shall briefly summarize some earlier results [9–12]. An excellent general reference to Finsler geometry used by these authors is [25] to which we refer the reader for more details of the general theory. If $F(x^\mu; y^\nu)$ is a Finsler function ¹, then it is assumed that $F^2(x; y)$ may be written such that [9]

$$F^2(x; y) = \mathcal{F}(x; y)g_{\mu\nu}y^\mu y^\nu; \quad (\text{II.1})$$

where $\mathcal{F}(x; y)$ is a positive function which is positively homogeneous in the velocities. Moreover [9], if $H(x)_{\alpha_1, \alpha_2, \dots, \alpha_N}$ is a totally symmetric tensor of rank N which is covariantly constant with respect to the Levi-Civita covariant derivative of the metric $g_{\mu\nu}$, then if

$$! = g(x)_{\mu\nu}y^\mu y^\nu = (H_{\alpha_1, \alpha_2, \dots, \alpha_N} y^{\alpha_1} y^{\alpha_2} \dots y^{\alpha_N})^{2/N} \quad \text{and} \quad \mathcal{F}(x; y) = \mathcal{F}(!) \quad (\text{II.2})$$

then the set of Finsler geodesics of F and the set of standard Riemannian geodesics of $g_{\mu\nu}$ coincide [see Eq. (20) in Ref. [9].] Further aspects are considered in Refs. [22, 23].

The case of Finsler pp-waves [15, 16] occurs when H is a covariantly constant null covector and $g_{\mu\nu}$ is the metric of a pp-wave, a special case of which is a plane gravitational wave. This was in effect pointed out by Tavakol and Van der Bergh [12] in 1986 and elaborated and extended by Roxburgh in 1991. Bogoslovky's original flat Finsler metric [3, 4] is a special case of their work but no mention of Bogoslovsky is made in Refs. [9, 11, 12], and so one assumes that they were unaware of it.

We next recall some basic definitions and notation used in Refs. [9, 11, 12]. Given a Finsler function $F(x; y)$, one may define the *Finsler metric tensor*

$$f_{\mu\nu}(x; y) = \frac{1}{2} \frac{\partial^2 F^2(x; y)}{\partial y^\mu \partial y^\nu} \quad (\text{II.3})$$

which is homogeneous of degree 0 in y^μ . That is, $f_{\mu\nu}(x; y)$ depends only upon the *direction*. Differentiating the identity $F^2(x^\alpha; y^\mu) = F^2(x^\alpha; y^\mu)$ twice with respect to y^μ implies that

$$f_{\mu\nu}y^\mu y^\nu = F^2(x; y); \quad (\text{II.4})$$

¹ See Refs. [9, 11, 12, 25]; y^μ is a four-velocity and (x^μ, y^μ) are local coordinates on TM , the tangent bundle of the spacetime manifold M .

The Finsler line element or arc length ds along a curve with tangent vector $y^\mu = \frac{dx^\mu}{d\lambda}$ is given by

$$ds^2 = F^2(x^\mu; dx^\mu) = f_{\mu\nu}(x; y) dx^\mu dx^\nu; \quad (\text{II.5})$$

and a Finsler geodesic is one for which $\int_\gamma F(x; dx^\mu) = \int_\gamma ds = 0$: The Euler-Lagrange equations are

$$\frac{d^2 x^\mu}{ds^2} + \frac{\mu}{\alpha\beta} \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds} = 0; \quad (\text{II.6})$$

where

$$\mu_{\nu\kappa} = \frac{1}{2} \left(\frac{\partial f_{\kappa\nu}}{\partial x^\mu} + \frac{\partial f_{\mu\kappa}}{\partial x^\nu} - \frac{\partial f_{\mu\nu}}{\partial x^\kappa} \right); \quad \frac{\mu}{\nu\kappa} = f^{\mu\sigma} \nu_{\sigma\kappa} \quad (\text{II.7})$$

are the analogues of *Christoffel symbols of the first and second kind*, respectively. In deriving the Euler-Lagrange equations, one uses the fact that $y^\kappa \frac{\partial f_{\alpha\beta}}{\partial y^\kappa} = 0$ because $f_{\mu\nu}$ is homogeneous of degree 0 in y^μ . Evidently, under a change of parameter $s \rightarrow \lambda = \lambda(s)$ we have $\frac{d}{ds} = \lambda' \frac{d}{d\lambda}$; $f_{\mu\nu} \rightarrow f_{\mu\nu}$ since $f_{\mu\nu}$ is homogenous of degree 0 in velocities. Thus, as in the standard Lorentzian situation,

$$\frac{d^2 x^\mu}{d\lambda^2} + \frac{\mu}{\alpha\beta} \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} = - \frac{\lambda''}{\lambda'} \frac{dx^\mu}{d\lambda}; \quad (\text{II.8})$$

If $\lambda'' = 0$, λ is called an affine parameter, and in what follows, unless otherwise stated, λ will denote an affine parameter.

In Refs. [9, 11, 12] the quantities

$$G^\mu = \frac{1}{2} \frac{\mu}{\nu\kappa} y^\nu y^\kappa \quad G^\mu_{\nu\kappa} = \frac{\partial^2 G^\mu}{\partial y^\nu \partial y^\kappa} \quad (\text{II.9})$$

are introduced. Although in general $G^\mu_{\nu\kappa} \neq \frac{\mu}{\nu\kappa}$; by virtue of the homogeneity of degree 0 of $\frac{\mu}{\nu\kappa}$ in y^μ , one has

$$G^\mu_{\nu\kappa} y^\nu y^\kappa = \frac{\mu}{\nu\kappa} y^\nu y^\kappa; \quad (\text{II.10})$$

and therefore Euler-Lagrangian equations may be rewritten as

$$\frac{d^2 x^\mu}{d\lambda^2} + G^\mu_{\nu\kappa} \frac{dx^\nu}{d\lambda} \frac{dx^\kappa}{d\lambda} = 0; \quad (\text{II.11})$$

In general $G^\mu_{\nu\kappa}$ depends upon the direction [12]; a *Berwald-Finsler manifold* is one for which $G^\mu_{\nu\kappa}$ is independent of the direction, –i.e.,

$$G^\mu_{\nu\kappa} = G^\mu_{\nu\kappa}(x); \quad (\text{II.12})$$

The motivation for Refs. [9, 11, 12] came from a classic paper of Ehlers, Pirani, and Schild [26] examining the fundamental assumptions justifying the use of pseudo-Riemannian geometry adapted in Einstein's general relativity. Roughly speaking the idea was that

- *The Principle of Universality of free fall* endows spacetime \mathcal{M} with a *projective structure*, that is, an equivalence class of curves, $\gamma : \mathbb{R} \rightarrow \mathcal{M}$ up to reparametrization.
- *The Principle of Einstein Causality* endows spacetime with a causal structure such that light rays are determined by some connection.

They conjectured that the only way of achieving this was that freely falling particles and null rays follow the geodesics of a pseudo-Riemannian metric, and in the case of particles that the curves carry a privileged parametrization given by proper time with respect to the pseudo-Riemannian metric along their paths in spacetime, whether freely falling or not.

In Refs. [11, 12] Tavakol and Van den Bergh sought to show that one could pass to a Finsler structure as well, provided one assumes

- {Ai}

$$F^2(\mathbf{x}; \mathbf{y}) = e^{2\sigma(\mathbf{x}, \mathbf{y})} \mathbf{g}_{\mu\nu}(\mathbf{x}) y^\mu y^\nu ; \quad (\text{II.13})$$

where $\mathbf{g}_{\mu\nu}(\mathbf{x})$ is a Lorentzian metric and $(\mathbf{x}^\alpha; \mathbf{y}^\mu)$ is homogeneous degree 0 in \mathbf{y}^μ .

This condition ensures that the conformal structures of the Finsler metric and the Lorentzian metric agree locally, in the spirit of Ref. [26]. It is pointed out in Ref. [12] that Eq. (II.13) is not equivalent to

$$\mathbf{f}_{\mu\nu} = e^{2\sigma(\mathbf{x}, \mathbf{y})} \mathbf{g}_{\mu\nu} \quad (\text{II.14})$$

because if this were true then $\mathbf{f}_{\mu\nu}$ would only depend upon \mathbf{x} and hence $\mathbf{f}_{\mu\nu}$ and $\mathbf{g}_{\mu\nu}$ would be conformally related, citing Ref. [25].²

- {Aii}

$$G_{\nu\kappa}^\mu = \left\{ \begin{matrix} \mu \\ \nu\kappa \end{matrix} \right\} \quad (\text{II.15})$$

where $\left\{ \begin{matrix} \mu \\ \nu\kappa \end{matrix} \right\}$ are the Christoffel symbols of the Lorentzian metric $\mathbf{g}_{\mu\nu}$. This condition ensures that the projective structures of the Finsler structure $F(\mathbf{x}; \mathbf{y})$ and the Lorentzian structure $\mathbf{g}_{\mu\nu}$ agree locally, again in the spirit of Ref. [26].

² In fact, Eq. (II.13) is obviously equivalent to Eq. (II.1) which is the form used by Ref. [9] (who appears to regard it as always true, although $g_{\mu\nu}$ is not necessarily unique).

Tavakol and Van den Bergh [12] claimed that the necessary and sufficient condition on $(\mathbf{x}; \mathbf{y})$ is

$$\frac{\partial}{\partial \mathbf{x}^\mu} - \mathbf{y}^\nu \frac{\partial}{\partial \mathbf{y}^\kappa} \left\{ \begin{matrix} \kappa \\ \mu\nu \end{matrix} \right\} = 0: \quad (\text{II.16})$$

and referred to it as the *metricity condition*. The name originates in the theory of the so-called *Cartan connection*. One defines

$$\mathbf{C}_{\mu\nu\kappa} = \frac{1}{2} \frac{\partial \mathbf{f}_{\mu\nu}}{\partial \mathbf{y}^\kappa} \quad (\text{II.17})$$

which is from Eq. (II.3), as totally symmetric in μ, ν, κ . Then one defines

$$\Gamma_{\mu\nu\kappa} = \mathbf{C}_{\mu\nu\kappa} - \left(\mathbf{C}_{\sigma\kappa\nu} \frac{\partial \mathbf{G}^\sigma}{\partial \mathbf{y}^\mu} + \mathbf{C}_{\sigma\kappa\mu} \frac{\partial \mathbf{G}^\sigma}{\partial \mathbf{y}^\nu} - \mathbf{C}_{\mu\sigma\nu} \frac{\partial \mathbf{G}^\sigma}{\partial \mathbf{y}^\kappa} \right): \quad (\text{II.18})$$

Acting on a vector $W^\mu(\mathbf{x}; \mathbf{y})$ the Cartan covariant derivative is defined by

$$\nabla_{\mathbf{x}^\kappa}^{\text{Cartan}} W^\mu = \frac{\partial W^\mu}{\partial \mathbf{x}^\kappa} + \mathbf{y}^\sigma \Gamma_{\sigma\kappa}^\lambda \frac{\partial W^\mu}{\partial \mathbf{y}^\kappa} + \Gamma_{\nu\sigma}^\mu W^\sigma \quad (\text{II.19})$$

and extended to tensors of arbitrary valence in the obvious way. The Cartan connection satisfies

$$\nabla_{\mathbf{x}^\kappa}^{\text{Cartan}} \mathbf{f}_{\mu\nu} = 0: \quad (\text{II.20})$$

This is equivalent to

$$\frac{\partial \mathbf{F}^2}{\partial \mathbf{x}^\kappa} - \frac{\partial \mathbf{F}^2}{\partial \mathbf{y}^\sigma} \frac{\partial \mathbf{G}^\sigma}{\partial \mathbf{y}^\kappa} = 0 \quad (\text{II.21})$$

and ensures that the norms of the vector remain constant under parallel transport along different routes. In Ref. [9] it is written as

$$\frac{\partial \mathbf{F}}{\partial \mathbf{x}^\kappa} - \frac{\partial \mathbf{F}}{\partial \mathbf{y}^\sigma} \frac{\partial \mathbf{G}^\sigma}{\partial \mathbf{y}^\kappa} = 0: \quad (\text{II.22})$$

In Ref. [12] it was suggested that

$$\mathbf{g}_{\mu\nu} \mathbf{x}^\mu \mathbf{d}\mathbf{x}^\nu = -2\mathbf{d}u\mathbf{d}v + (\mathbf{u})\mathbf{d}\mathbf{x}^2 + (\mathbf{u})\mathbf{d}\mathbf{y}^2 \quad (\text{II.23})$$

with $(\mathbf{x}^1; \mathbf{x}^2; \mathbf{x}^3; \mathbf{x}^4) = (\mathbf{x}; \mathbf{y}; \mathbf{u}; \mathbf{v})$, which they call a plane wave, might lead to a solution. They find [in their Eq. (34)] that

$$= \left(\frac{\dot{\mathbf{x}}^2 + \dot{\mathbf{y}}^2 - 2\dot{\mathbf{u}}\dot{\mathbf{v}}}{\dot{\mathbf{u}}^2} \right) \quad (\text{II.24})$$

and they claim that it is indeed a solution.

The treatment of Ref. [9] starts with the helpful observation that the sums, products, and ratios of solutions are again solutions. Roxburgh investigated Lorentzian metrics with covariantly constant vector fields and pointed out that pp-waves are a special case.

III. BOGOSLOVSKY-FINSLER METRICS

Bogoslovsky's theory [3, 4] was based on the Finsler line element such that the proper time along a future-directed *timelike* world line $x^\mu(\cdot)$ in flat Minkowski spacetime is obtained by combining the Minkowski line element with what we call here the Bogoslovsky factor,

$$d\tau = (-g_{\mu\nu}dx^\mu dx^\nu)^{\frac{1-b}{2}} (-g_{\mu\nu}l^\mu dx^\nu)^b; \quad (\text{III.1})$$

where $0 \leq b < 1$ is a dimensionless constant, $g_{\mu\nu}$ is the flat Minkowski metric tensor (with mainly positive signature), and l^μ , is a constant future directed null vector – i.e.

$$\partial_\mu l^\nu = 0; \quad g_{\mu\nu}l^\mu l^\nu = 0; \quad l^0 > 0; \quad (\text{III.2})$$

where $\partial_\mu = \frac{\partial}{\partial x^\mu}$. The Bogoslovsky factor makes (III.1) homogenous of degree 1 – i.e., Finslerian.

The parameter b introduces spatial anisotropy which might be relevant at the early stages of the Universe [27]. The constant b is very small by Hughes-Drever-type experiments [2]; Bogoslovsky argues that $b < 10^{-10}$ [28]. For $b = 0$ we recover the Minkowski proper time element, cf. Eq. (I.1) with $A_\mu = 0$.

Bogoslovsky's Finsler line element has an obvious generalization: in Eq. (III.1), one replaces $g_{\mu\nu}$ with a curved pseudo-Riemannian metric $g_{\mu\nu}(x)$ and l^μ with a future-directed null vector such that

$$\nabla_\mu l^\nu = 0; \quad g_{\mu\nu}l^\mu l^\nu = 0; \quad (\text{III.3})$$

where ∇_μ is the Levi-Civita connection of the Lorentzian metric $g_{\mu\nu}$. This idea has recently been explored in Refs. [15–17, 21], where such spacetimes are called “Finsler pp-waves”.

Such spacetimes are also referred to as Brinkman [29] or Bargmann [30] spacetimes. They admit Brinkmann coordinates $X^\mu = (V; U; X^i)$ such that

$$g_{\mu\nu}dX^\mu dX^\nu = 2dV dU + dX^i dX^i - 2H(X^i; U)dU^2; \quad (\text{III.4})$$

where the spacetime dimension is $d + 1$ and $i = 1; 2; \dots; d - 1$. $H(X^i; U)$ is an arbitrary not

identically vanishing function of its arguments ³. $U;V$ may be written as

$$V = X^- = \frac{1}{\sqrt{2}}(X^d - X^0); U = X^+ = \frac{1}{\sqrt{2}}(X^d + X^0); \quad (\text{III.5})$$

We have $l^\mu @_\mu = -@_V$ so that $-g_{\mu\nu} l^\mu dX^\nu = dU$: Then the Finsler pp-line element is

$$d = (-g_{\mu\nu} dX^\mu dX^\nu)^{\frac{1-b}{2}} (-g_{\mu\nu} l^\mu dX^\nu)^b = (-g_{\mu\nu} dX^\mu dX^\nu)^{\frac{1-b}{2}} (dU)^b; \quad (\text{III.6})$$

where $g_{\mu\nu}$ is the pp-wave metric.

Returning to the pp-waves, we recall that the metric is Ricci flat if and only if $H(X^i; U)$ is a harmonic function of the coordinates X^i ; it may, however, have arbitrary dependence upon U . It then represents a left-moving (i.e., in the negative X^d direction) gravitational wave such that X^i 's are transverse to the direction of motion. The wave fronts $U = \text{constant}$ are null hypersurfaces, and the covariantly constant and hence Killing null vector field $@_V$ lies in the wave fronts.

If, in addition, $H(X^i; U)$ is quadratic in the transverse coordinates then we have a *plane gravitational wave*. If $d = 3$, which we assume from now on, then

$$-2H = \mathcal{A}_+(U)(X_1^2 - X_2^2) + \mathcal{A}_\times(U)2X_1X_2 = K_{ij}(U)X^iX^j; \quad (\text{III.7})$$

where $\mathcal{A}_+(U)$ and $\mathcal{A}_\times(U)$ are the amplitudes of the two plane polarization states.

For general $\mathcal{A}_+(U)$ and $\mathcal{A}_\times(U)$ there is a five-dimensional isometry group G_5 which acts multiply transitively on the three-dimensional wave fronts $U = \text{constant}$ [31, 32]. This group is a subgroup of the six-dimensional Carroll group $\text{Carr}(2)$ in three spacetime dimensions [33, 34] in which the $SO(2)$ subgroup is omitted [36].

The Carroll group $\text{Carr}(2)$ may be regarded as a subgroup of the Poincaré group $ISO(3; 1)$ defined by freezing out U -translations [34]; it acts on the null hyperplanes $U = \text{constant}$. If we label the Killing vector fields of the Poincaré group as

$$P_\mu = \frac{@}{@X^\mu}; \quad L_{\mu\nu} = X_\mu P_\nu - X_\nu P_\mu; \quad (\text{III.8})$$

³ Our choice of sign for g_{UV} has the advantage that raising and lowering of indices entails no minus signs, merely swapping U and V and is consistent with our previous papers. It has however the consequence that if we choose a time orientation such that U increases to the future then V decreases to future. In other words $\frac{\partial}{\partial U}$ is a future directed null vector field and $\frac{\partial}{\partial V}$ is a past directed vector field.

then the Carroll group is generated by

$$P_- = \frac{\partial}{\partial V}; \quad P_i = \frac{\partial}{\partial X^i}; \quad (\text{III.9a})$$

$$L_{ij} = X_i P_j - X_j P_i; \quad L_{-i} = X_- P_i - X_i P_- = U P_i - X_i P_-; \quad (\text{III.9b})$$

$i = 1; 2$, and each hyperplane $U = \text{const.}$, is left invariant. The generators in Eq. (III.9a) are translations, whereas those in Eq. (III.9b) are planar rotation and boosts. Since $d = 3$, we may relabel the generators $L_{ij} = J$, and the U-V boost

$$N_0 = L_{+-} = X_+ P_- - X_- P_+ = V P_- - U_- P_+ \quad \text{where} \quad P_+ = \frac{\partial}{\partial U} \quad (\text{III.10})$$

and find that the four generators $N_0; J; L_{-i}$ generate a group which is abstractly isomorphic to the group SIM(2), the group of similarities, that is dilations, rotations, and translations of the Euclidean plane \mathbb{E}^2 . SIM(2) is the largest proper subgroup of the Lorentz group SO(3; 1). Adjoining the generators $P_+; P_-; P_i$ gives rise to the eight generators of ISIM(2), which is a subgroup of the Poincaré group. The group ISIM(2) acts multiply transitively on Minkowski spacetime $\mathbb{E}^{3,1}$.

It was suggested by Cohen and Glashow [1] that ISIM(2), which may be thought of as the subgroup of ISO(3; 1) leaving invariant a null direction, could explain weak CP violation while being compatible with tests of Lorentz invariance, since it would rule out *spurions*, that is, tensor vacuum expectation values.

In Ref. [37] it was pointed out that Ricci flat pp-waves are *strongly universal*. In particular, they have nonvanishing scalar invariants constructed from the Riemann tensor and as a consequence satisfy almost any set of covariant field equations. Quantum corrections to the metric vanish. Thus this property may be thought of as the analogue for the proposed curved Bogoslovsky-Finsler structures with a Ricci flat metric $g_{\mu\nu}$ of Cohen and Glashow's "no spurions" condition.

In Ref. [6], an attempt was made to find a link with general relativity in which Minkowski spacetime $\mathbb{E}^{3,1}$ may be regarded as the coset ISO(3; 1)=SO(3; 1). The only two deformations of the Poincaré group led to the two de Sitter groups SO(4; 1) and SO(3; 2) for which translations act in a noncommutative fashion on the cosets' de Sitter spacetime $dS_4 = \text{SO}(4; 1)=\text{SO}(3; 1)$ and anti-de Sitter spacetime $\text{AdS}_4 = \text{SO}(3; 2)=\text{SO}(3; 1)$.

They therefore investigated the deformations of ISIM(2) and found that there exists a family of deformations depending upon two dimensionless parameters a and b . However for

all \mathbf{a} and \mathbf{b} the translations $P_+; P_-; P_i$ failed to commute. In general, the rotation J became a noncompact generator unless $\mathbf{a} = 0$, leaving $\text{DISIM}_b(2)$ depending on a dimensionless parameter \mathbf{b} . They then observed that this is precisely the symmetry of Bogoslovsky's Finsler metric [Eq. (III.1)]. For a review of these ideas and their relation to the much earlier work of Voigt [38], the reader is directed to the recent review in Ref. [39]. For a recent discussion of Bogoslovsky-Finsler deformations in the light of the ideas of Segal, see Ref. [40].

In a recent paper [15], the authors have shown, among other things, that the Bogoslovsky-Finsler pp-waves enjoy the same universal properties with respect to generalizations of the Einstein equations to Finsler-Einstein equations as those in the pseudo-Riemannian case discussed in Ref. [37].

IV. GEODESICS

The geodesics of a Finsler metric with Finsler function $F(x^\mu; \dot{x}^\mu)$, where F is homogeneous of degree 1 in \dot{x}^μ , are extrema of

$$I = \int F(x^\mu; \frac{dx^\mu}{d}) d \quad ; \quad (\text{IV.1})$$

In the case we are considering, we restrict our attention to future-directed timelike curves for which both $g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu$ and $-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu$ are strictly positive in order to ensure that F is real. For a particle of mass m , the action with respect to Lagrangians is

$$S_b = -m \int F d \quad ; \quad (\text{IV.2})$$

where F is the Bogoslovsky-Finsler line element [Eq. (III.6)]. The integral is independent of the parameter λ . Therefore if $p_\mu = \frac{\partial(-mF)}{\partial \dot{x}^\mu}$, then $\mathcal{H} = p_\mu \dot{x}^\mu + mF$ is a constant of the motion. This is indeed true, but because $F(x^\mu; \dot{x}^\mu)$ is homogeneous of degree 1 in \dot{x}^μ one has $\dot{x}^\mu \frac{\partial F}{\partial \dot{x}^\mu} = F$ and consequently the constant vanishes identically. Standard Riemannian or Lorentzian metrics are, of course, a special case of this general fact.

Now we analyze the motion along geodesics in Bogoslovsky-Finsler plane gravitational waves adapting the discussion for the standard Einstein case given in Refs. [36, 41]. First, we find it convenient to pass to Baldwin-Jeffery-Rosen (BJR) coordinates $x^\mu = (v; u; x^i)$, defined by

$$X^i = P_{ij} x^j \quad ; \quad U = u \quad ; \quad V = v - \frac{1}{4} \frac{da_{ij}}{du} x^i x^j \quad ; \quad (\text{IV.3})$$

where $\mathbf{a} \equiv (a_{ij}) = \mathbf{P}^t \mathbf{P}$; and the matrix \mathbf{P} satisfies the matrix Sturm-Liouville equation

$$\frac{d^2 \mathbf{P}}{du^2} = \mathbf{K} \mathbf{P}; \quad \mathbf{P}^t \frac{d\mathbf{P}}{du} = \frac{d\mathbf{P}^t}{du} \mathbf{P}; \quad (\text{IV.4})$$

where $\mathbf{K} = (K_{ij})$ is the profile in Brinkmann coordinates; see Eq. (III.7).

In BJR coordinates we have

$$g_{\mu\nu} dx^\mu dx^\nu = 2du dv + a_{ij}(u) \dot{x}^i \dot{x}^j; \quad l^\mu \frac{\partial}{\partial x^\mu} = -\frac{\partial}{\partial v}; \quad (\text{IV.5})$$

Here $\dot{x}^\mu = \frac{dx^\mu}{d\lambda}$ where λ is an arbitrary parameter. The Lagrangian is proportional to the Bogoslovsky-Finsler function,

$$\mathcal{L}_b = -mF \quad \text{where} \quad F = \left(-2\dot{u}\dot{v} - a_{ij}(u) \dot{x}^i \dot{x}^j \right)^{\frac{1}{2}(1-b)} (\dot{u})^b \quad (\text{IV.6})$$

For the curve to be timelike we must have $\dot{u}\dot{v} < 0$ and $\dot{u} > 0$. Since the integral (IV.1) is independent of the choice of the parameter λ , we are entitled to make the choice $\lambda = u$ and extremize

$$\int \left(-2 \frac{dv}{du} - a_{ij}(u) \frac{dx^i}{du} \frac{dx^j}{du} \right)^{\frac{1}{2}(1-b)} du; \quad (\text{IV.7})$$

With this choice of parametrization the integrand of (IV.7) is now no longer homogeneous in the velocities $\frac{dv}{du}$ and $\frac{dx^i}{du}$ but because a_{ij} depends on the ‘‘time’’ u , there is no conserved analogue of the quantity \mathcal{H} . The symmetry aspects will be further investigated in Sec.V.

Before analyzing the general case we recall, for later comparison, some aspects of the geodesics of a pp-wave described by the square-root Lagrangian [Eq. (I.1)].

Geodesics in a pp-wave

Let us thus first consider a pp-wave written in BJR coordinates, whose geodesics are described by Eq. (I.1) with $\mathbf{A}_\mu = 0$:

$$\mathcal{L}_0 = -m \sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu} = -m \sqrt{-a_{ij}(u) \dot{x}^i \dot{x}^j - 2\dot{u}\dot{v}}; \quad (\text{IV.8})$$

The canonical momenta $\mathbf{p}_\mu = \frac{\partial \mathcal{L}_0}{\partial \dot{x}^\mu}$ are

$$\mathbf{p}_u = \frac{m\dot{v}}{\sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}}; \quad \mathbf{p}_i = \frac{ma_{ij} \dot{x}^j}{\sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}}; \quad \mathbf{p}_v = \frac{m\dot{u}}{\sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}} \quad (\text{IV.9})$$

of which \mathbf{p}_i and \mathbf{p}_v are constants of the motion since $a_{ij} = a_{ij}(u)$. For a u -dependent profile \mathbf{p}_u is not conserved, though. The geodesic equations of motion are

$$\ddot{u} = \dot{u} \frac{d}{d} \ln (\sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}); \quad (\text{IV.10a})$$

$$\ddot{x}^i + \dot{u} a^{ij} a'_{jk} \dot{x}^k = \dot{x}^i \frac{d}{d} \ln (\sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}); \quad (\text{IV.10b})$$

$$\ddot{v} - \frac{1}{2} a'_{ij} \dot{x}^i \dot{x}^j = \dot{v} \frac{d}{d} \ln (\sqrt{-g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu}); \quad (\text{IV.10c})$$

where $a'_{ij} = \frac{da_{ij}}{du}$. Using the first equation, the two remaining ones simplify to ⁴

$$\ddot{x}^i + \dot{u} a^{ij} a'_{jk} \dot{x}^k = \dot{x}^i \frac{\ddot{u}}{\dot{u}}; \quad (\text{IV.11a})$$

$$\ddot{v} - \frac{1}{2} a'_{ij} \dot{x}^i \dot{x}^j = \dot{v} \frac{\ddot{u}}{\dot{u}}; \quad (\text{IV.11b})$$

An ingenious way to solve these equations is to use the conserved quantities. We first define the constants of the motion by setting

$$P_i = \frac{p_i}{p_v} = \frac{a_{ij} \dot{x}^j}{\dot{u}}; \quad (\text{IV.12})$$

The resulting first-order differential equation for x^i is solved at once as

$$x^i(u) = S^{ij}(u) P_j + x_0^i; \quad (\text{IV.13})$$

where $S \equiv S^{ij}$ is the Souriau matrix [36], defined by

$$\frac{dS(u)}{du} = a^{-1}(u); \quad (\text{IV.14})$$

p_v in Eq. (IV.9) provides us in turn with a first-order equation for v :

$$\dot{v} = -\frac{1}{2} a^{ij} P_i P_j \dot{u} - \frac{1}{2} \frac{2}{0} \dot{u} \quad \text{where} \quad \frac{0}{0} = \frac{m}{p_v}; \quad (\text{IV.15})$$

This equation is then solved as

$$v = -\frac{1}{2} P_i P_j S^{ij}(u) - \frac{1}{2} \frac{2}{0} u + v_0; \quad (\text{IV.16})$$

The transverse motion (IV.13) is the same for all values of the mass m , which enters only the v motion (IV.16) by a shift which is linear in u and proportional to the mass-quotient term $\frac{0}{0}$ in Eq. (IV.15), familiar from Ref. [42].

Finsler Geodesics.

⁴ Choosing the affine parameter $\lambda = u$, the rhs would vanish.

Let us now consider the Bogoslovsky-Finsler Lagrangian \mathcal{L}_b in Eq. (IV.6). Its canonical momenta are

$$\mathbf{p}_u = m(\dot{u})^{b-1} (-2\dot{u}\dot{v} - a_{ij}\dot{x}^i\dot{x}^j)^{-\frac{1+b}{2}} ((1+b)\dot{u}\dot{v} + b a_{ij}\dot{x}^i\dot{x}^j); \quad (\text{IV.17a})$$

$$\mathbf{p}_i = m(1-b)(a_{ij}\dot{x}^j)\dot{u}^b (-2\dot{u}\dot{v} - a_{ij}\dot{x}^i\dot{x}^j)^{-\frac{1+b}{2}}; \quad (\text{IV.17b})$$

$$\mathbf{p}_v = m(1-b)\dot{u}^{b+1} (-2\dot{u}\dot{v} - a_{ij}\dot{x}^i\dot{x}^j)^{-\frac{1+b}{2}}; \quad (\text{IV.17c})$$

\mathbf{p}_i and \mathbf{p}_v are constants of the motion as before and we have the dispersion relation in Eq. (18) of Ref. [6]:

$$\mathbf{p}^2 \equiv g^{\mu\nu}\mathbf{p}_\mu\mathbf{p}_\nu = -m^2(1-b^2)\dot{u}^{2b} \left(-2\dot{v} - a_{ij}\dot{x}^i\dot{x}^j \right)^{-b}; \quad (\text{IV.18})$$

The geodesic equations,

$$(b+1)\ddot{u} = \dot{u} \frac{d}{d} \ln \left(-2\dot{u}\dot{v} - a_{ij}\dot{x}^i\dot{x}^j \right)^{\frac{1+b}{2}}; \quad (\text{IV.19a})$$

$$\ddot{x}^i + \dot{u} a^{ij} a'_{jk} \dot{x}^k + b \frac{\ddot{u}}{\dot{u}} \dot{x}^i = \dot{x}^i \frac{d}{d} \ln \left(-2\dot{u}\dot{v} - a_{ij}\dot{x}^i\dot{x}^j \right)^{\frac{1+b}{2}}; \quad (\text{IV.19b})$$

$$\dot{v} + \frac{3b-1}{2(1+b)} a'_{ij} \dot{x}^i \dot{x}^j + \frac{2b}{\dot{u}(1+b)} a_{ij} \dot{x}^i \dot{x}^j = \frac{1}{b+1} \left(\dot{v} + \frac{2b}{1+b} a_{ij} \dot{x}^i \dot{x}^j \right) \left(\frac{d}{d} \ln \left(-2\dot{u}\dot{v} - a_{ij}\dot{x}^i\dot{x}^j \right)^{\frac{1+b}{2}} \right); \quad (\text{IV.19c})$$

reduce to Eq. (IV.10) when $b = 0$.

The remarkable fact is that using Eq. (IV.19a), the two remaining equations become the *same* [Eq. (IV.11)], as for the square root Lagrangian [Eq. (IV.8)].

This does *not* imply identical solutions, though, as seen by solving the geodesics equations along the same lines as before. Setting once again $\mathbf{P}_i = \frac{p_i}{p_v}$ provides us with the transverse motion

$$\boxed{\mathbf{x}^i(\cdot) = S^{ij}(u(\cdot)) \mathbf{P}_j + \mathbf{x}_0^i} \quad (\text{IV.20})$$

which is again Eq. (IV.13). Then, from Eq. (IV.17b), we infer that

$$\dot{v} = -\frac{1}{2} (a^{ij} \mathbf{P}_i \mathbf{P}_j + \frac{2}{b}) \dot{u}; \quad \text{where} \quad b = \left(\frac{m}{p_v} (1-b) \right)^{\frac{1}{1+b}}; \quad (\text{IV.21})$$

whose integration yields

$$\boxed{\mathbf{v} = -\frac{1}{2} S^{ij}(u) \mathbf{P}_i \mathbf{P}_j - \frac{1}{2} \frac{2}{b} \mathbf{u} + v_0;} \quad (\text{IV.22})$$

Let us observe that this takes the same form as for Eq. (IV.16), however, with a new, \mathbf{b} -dependent mass-quotient term, $\frac{m}{\dot{u}}$. For $\mathbf{b} = 0$ the latter reduces to $\frac{m}{\dot{u}}$, and the massive Eq. (IV.16) is recovered.

The family of pp-wave geodesics are given by (IV.13) and (IV.16) and are labeled by the constants of integration $P_i; \mathbf{x}_0^i; v_0$ and \dot{u}_0 . The Finsler geodesics are given by Eqs. (IV.20) and (IV.22) and are labeled by the constants of integration $P_i; \mathbf{x}_0^i; v_0$ and \dot{u}_0 . It is clear that the two sets of geodesics are identical up to a \mathbf{b} -dependent relabeling of the last constant of integration.

In the massless case $m = 0$ (photons) that, the \mathbf{b} -dependent term drops out from Eq. (IV.22). Letting $\mathbf{b} \rightarrow 1$ turns off the mass-quotient term, $\frac{m}{\dot{u}} \rightarrow 0$, and all geodesics behave as if they were massless, consistently with Eq. (IV.18). See Fig. 1 in Sec. VII for an illustration.

Another way to see the surprising identity of the geodesics is to consider the Euler-Lagrange equations

$$E_\mu = \frac{\partial \mathcal{L}_0}{\partial \dot{\mathbf{x}}^\mu} - \frac{d}{dt} \left(\frac{\partial \mathcal{L}_0}{\partial \dot{\mathbf{x}}^\mu} \right) = 0 \quad \text{and} \quad \tilde{E}_\mu = \frac{\partial \mathcal{L}_b}{\partial \dot{\mathbf{x}}^\mu} - \frac{d}{dt} \left(\frac{\partial \mathcal{L}_b}{\partial \dot{\mathbf{x}}^\mu} \right) = 0 \quad (\text{IV.23})$$

of the two Lagrangians \mathcal{L}_0 and \mathcal{L}_b in Eqs. (IV.8) and (IV.6), respectively.

Both systems can be described by three independent equations, since the following identities hold: $\dot{\mathbf{x}}^\mu E_\mu \equiv 0; \dot{\mathbf{x}}^\mu \tilde{E}_\mu \equiv 0$: Then the combinations are as follows:

1. For the first system :

$$\frac{(-g_{\mu\nu} \dot{\mathbf{x}}^\mu \dot{\mathbf{x}}^\nu)^{1/2}}{m} \left(\frac{2\dot{v}}{\dot{u}} E_v + \frac{\dot{\mathbf{x}}^i}{\dot{u}} E_i \right) = 0 \quad (\text{IV.24})$$

$$\frac{(-g_{\mu\nu} \dot{\mathbf{x}}^\mu \dot{\mathbf{x}}^\nu)^{1/2}}{m} \left(\frac{\dot{\mathbf{x}}^i}{\dot{u}} E_v - a^{ij} E_j \right) = 0: \quad (\text{IV.25})$$

2. For the second system:

$$\frac{(-g_{\mu\nu} \dot{\mathbf{x}}^\mu \dot{\mathbf{x}}^\nu)^{\frac{1+b}{2}}}{m} \left[\frac{1}{(1-b^2)\dot{u}^{b+2}} (2\dot{u}\dot{v} - ba_{ij}\dot{\mathbf{x}}^i\dot{\mathbf{x}}^j) \tilde{E}_v + \frac{\dot{\mathbf{x}}^i}{(1-b)\dot{u}^{b+1}} \tilde{E}_i \right] = 0 \quad (\text{IV.26})$$

$$\frac{(-g_{\mu\nu} \dot{\mathbf{x}}^\mu \dot{\mathbf{x}}^\nu)^{\frac{1+b}{2}}}{m} \left[\frac{\dot{\mathbf{x}}^i}{(1-b)\dot{u}^{b+1}} \tilde{E}_v + \frac{a^{ij}}{(b-1)\dot{u}^b} \tilde{E}_j \right] = 0 \quad (\text{IV.27})$$

(where $i; j = 1; 2$) yield both parts of Eq. (IV.11).

V. PARTIALLY BROKEN CARROLL SYMMETRY

Generic plane gravitational waves are invariant under the same five-parameter group we denote by G_5 [31, 32]. Expressed in BJR coordinates, G_5 is implemented as [36],

$$\mathbf{u} \rightarrow \mathbf{u}; \quad \mathbf{x} \rightarrow \mathbf{x} + \mathbf{S}(\mathbf{u})\mathbf{b} + \mathbf{c}; \quad \mathbf{v} \rightarrow \mathbf{v} - \mathbf{b} \cdot \mathbf{x} - \frac{1}{2}\mathbf{b} \cdot \mathbf{S}(\mathbf{u})\mathbf{b} + \mathbf{f}; \quad (\text{V.1})$$

where \mathbf{S} is Souriau's matrix [Eq. (IV.14)]. The 2-vectors \mathbf{b} and \mathbf{c} and \mathbf{f} are constants, interpreted as boosts, and as transverse and vertical translations. These same transformations are isometries also for the Bogoslovsky-Finsler metric [Eq. (III.6)] because \mathbf{u} is fixed, and the pp isometries leave the pp-wave metric – and hence their powers – invariant. The transformations in Eq. (V.1) are generated by the vector fields

$$\mathbf{B}_i = \mathbf{S}_{ij}(\mathbf{u})\partial_j - \mathbf{x}_i\partial_v; \quad \partial_i \quad \text{and} \quad \partial_v; \quad (\text{V.2})$$

respectively. The only nonvanishing Lie bracket is

$$[\partial_i; \mathbf{B}_j] = -\partial_j\partial_v; \quad (\text{V.3})$$

Rotations, generated by $\mathbf{L}_{ij} = \mathbf{x}_i\partial_j - \mathbf{x}_j\partial_i$, are not symmetries in general.

The restriction of a pp-wave to the $\mathbf{u} = 0$ hypersurface \mathcal{C}_0 carries a Carroll structure. The “vertical” coordinate \mathbf{v} is interpreted as “Carrollian time” [33–35]. \mathcal{C}_0 is left invariant by the action (V.1) and the generators then satisfy the Carroll algebra in two space dimensions with rotations omitted [36]. Then Eq. (V.1) tells us how the Carroll group is implemented on any hypersurface $\mathbf{u} = \mathbf{u}_0 = \text{const}$:

In the flat case, $\mathbf{a}_{ij} = \delta_{ij}$, we have further symmetries. In particular, adding the vector fields $\partial_u; \mathbf{L}_{ij}$ and $\mathbf{L}_{+-} = \mathbf{v}\partial_v - \mathbf{u}\partial_u$ yields the Lie algebra of an eight-parameter subgroup of the Poincaré group. We now have

$$[\mathbf{v}\partial_v - \mathbf{u}\partial_u; \partial_v] = -\partial_v; \quad (\text{V.4})$$

and so the *direction* of the null Killing vector field ∂_v is preserved. The eight-dimensional group they generate is ISIM(2). Omitting the translations $\partial_v; \partial_u; \partial_i$ gives SIM(2), the largest proper subgroup of the Lorentz group $\text{SO}(3;1)$. This is the symmetry of Cohen and Glashow's *very special relativity* [1].

Returning to the case of general pp-waves and their Bogoslovsky-Finsler version [Eq. (III.6)], we emphasize that the BJR matrix $\mathbf{a} = (a_{ij})$ and thus the Souriau matrix \mathbf{S} ,

depends on the pp-wave metric only, but *not* on the deformation parameter \mathbf{b} . Therefore, the isometries in Eq. (V.1) act, for Eq. (III.6), exactly as for standard plane waves.

The invariance of the Bogoslovsky-Finsler model can be confirmed with respect to the partially broken Carroll group. The infinitesimal version of Eq. (V.1) is Y_{iso} in Eq. (V.2). The linear momenta in Eq. (IV.17) are readily recovered ; using Eq. (V.2) for boosts, we get in turn,

$$\mathbf{k}^i = \mathbf{p}_v \mathbf{x}^i - S_{ij} \mathbf{p}_j; \quad (\text{V.5})$$

– just as for a gravitational wave [36]. Its conservation follows from Noether’s theorem, and can also be confirmed by a direct calculation. The dependence on \mathbf{b} is hidden in the momenta in Eq. (IV.17). The initial position \mathbf{x}_0^i in Eq. (IV.13) is the conserved value of \mathbf{k}^i .

For $\mathbf{b} = 0$ the flat Bogoslovsky-Finsler model has one more isometry, identified with the U-V boost $\mathbf{N}_0 = \mathbf{L}_{+-}$ in Eq. (III.10). For $\mathbf{b} \neq 0$, this generator is broken but not entirely lost : Let us explain how this comes about.

As said above, the (rotationless) Carroll isometry group \mathbf{G}_5 in Eq. (V.1) of the initial pp-wave remains a symmetry with identical generators for its Bogoslovsky-Finsler extension.

To see what happens to U-V boosts we start with the Minkowski metric, ${}_{\mu\nu}d\mathbf{x}^\mu d\mathbf{x}^\nu = {}_{ij}d\mathbf{x}^i d\mathbf{x}^j + 2d\mathbf{u}d\mathbf{v}$. A U-V boost, implemented as

$$\mathbf{u} \rightarrow \lambda^{-1}\mathbf{u}; \quad \mathbf{x}^i \rightarrow \mathbf{x}^i; \quad \mathbf{v} \rightarrow \lambda \mathbf{v} \quad (\text{V.6})$$

where $\lambda = \text{const} > 0$ is an isometry. Moreover, its \mathbf{b} -dependent deformation of Eq. (V.6),

$$\mathbf{u} \rightarrow \lambda^{b-1}\mathbf{u}; \quad \mathbf{x}^i \rightarrow \lambda^b \mathbf{x}^i; \quad \mathbf{v} \rightarrow \lambda^{b+1}\mathbf{v} \quad (\text{V.7})$$

is readily seen to leave the Bogoslovsky-Finsler line element (III.1) invariant – although for $\mathbf{b} \neq 0$ it is only a conformal transformation for the Minkowski metric, ${}_{\mu\nu}d\mathbf{x}^\mu d\mathbf{x}^\nu \rightarrow \lambda^{2b} {}_{\mu\nu}d\mathbf{x}^\mu d\mathbf{x}^\nu$, and not an isometry ⁵. We record for later use that the \mathbf{b} -deformed boost [Eq. (V.7)] is generated by

$$\mathbf{N}_b = (\mathbf{b} - 1)\mathbf{u}_{@_u} + (\mathbf{b} + 1)\mathbf{v}_{@_v} + \mathbf{b}\mathbf{x}^i_{@_i}; \quad (\text{V.8})$$

Both Eqs. (V.6) and (V.7) leave the hypersurface $\mathbf{u} = 0$ invariant, and extend the Carroll action (V.1). We note that the restriction to \mathcal{C}_0 of the deformed U-V boost [Eq. (V.7)] scales

⁵ Noting that $\lambda^{b-1} = (\lambda^b)^{\frac{b-1}{b}}$ shows that (V.7) has dynamical exponent $z = 1 - \frac{1}{b} < 0$ which corresponds to the conformal Galilei algebra labeled by z [45, 46].

the Carrollian time, $v \rightarrow {}^{b+1}v$. Therefore it is only the *direction* of $@_v$ (and not $@_v$ itself) which is preserved the isometry [Eq. (III.1)] is “chronoprojective” [41, 43]:

$$@_v \rightarrow {}^{-1-b}@_v : \quad (\text{V.9})$$

In the flat case, two more isometries — namely \mathbf{u} -translations and rotations complete the algebra to one with eight-parameter. With some abuse, we will still refer to \mathbf{G}_5 extended with \mathbf{U} -translations (but with no rotations) as “Carroll” for simplicity and denote it by \mathbf{G}_6 . Its further extension by $\mathbf{U} - \mathbf{V}$ boosts will be called chrono-Carroll [41] and denoted by \mathbf{G}_7 .

The Lie algebra structure is most easily checked by taking the commutators of the vector fields in Eqs. (III.9) and (III.10) and comparing with those given in Eq. (9) of Ref. [6], which gives the structure constants of the deformed group $\text{DISIM}_b(2)$; those of $\text{ISIM}(2)$ are obtained by setting $\mathbf{b} = 0$.

Further insight is gained by decomposing the deformed \mathbf{U} - \mathbf{V} boost generator \mathbf{N}_b in Eq. (V.8) into the sum of the undeformed expression $\mathbf{N}_0 = \mathbf{L}_{+-}$ and a relativistic dilation, \mathbf{D} :

$$\mathbf{N}_b = v@_v - \mathbf{u}@_u + \mathbf{b}(\mathbf{u}@_u + v@_v + \mathbf{x}^i@_i) = \mathbf{N}_0 + \mathbf{bD} : \quad (\text{V.10})$$

For $\mathbf{b} \neq 0$, \mathbf{N}_0 is broken, and it is only the above combination of \mathbf{U} - \mathbf{V} boosts and dilations which is a symmetry — a situation familiar from gravitational plane waves [36, 41, 44].

It is instructive to see how this comes about. In the flat Minkowski case $\mathbf{a}_{ij} = \delta_{ij}$ and Eq. (IV.17) yield

$$\mathbf{p}_i = (\dot{\mathbf{x}}_i = \dot{\mathbf{u}})\mathbf{p}_v \quad \text{and} \quad \mathbf{p}_u = \frac{(1 + \mathbf{b})\dot{\mathbf{u}}\dot{v} - \mathbf{b}\dot{\mathbf{x}}^i\dot{\mathbf{x}}_i}{(1 - \mathbf{b})\dot{\mathbf{u}}^2}\mathbf{p}_v :$$

Then for $\mathcal{D} = \mathbf{D}^\mu\mathbf{p}_\mu$ and $\mathcal{N}_0 = \mathbf{N}_0^\mu\mathbf{p}_\mu$ we have

$$\dot{\mathcal{D}} = \frac{(2\dot{\mathbf{u}}\dot{v} + \dot{\mathbf{x}}^i\dot{\mathbf{x}}_i)}{(1 - \mathbf{b})\dot{\mathbf{u}}}\mathbf{p}_v \quad \text{and} \quad \dot{\mathcal{N}}_0 = -\mathbf{b}\frac{(2\dot{\mathbf{u}}\dot{v} + \dot{\mathbf{x}}^i\dot{\mathbf{x}}_i)}{(1 - \mathbf{b})\dot{\mathbf{u}}}\mathbf{p}_v = -\mathbf{b}\dot{\mathcal{D}} ; \quad (\text{V.11})$$

so that the combination of the two expressions is conserved :

$$\dot{\mathcal{N}}_b = 0 \quad \text{for} \quad \mathcal{N}_b = \mathcal{N}_0 + \mathbf{bD} : \quad (\text{V.12})$$

Now we turn to the curved case. Let us consider a conformal transformation \mathbf{f} of a pp wave with metric $\mathbf{g}_{\mu\nu}$:

$$\mathbf{f}_*\mathbf{g}_{\mu\nu} = \Omega^2\mathbf{g}_{\mu\nu} ; \quad (\text{V.13})$$

where \mathbf{f}_* is the pullback map. This changes the “pp factor” in Eq. (IV.5), as

$$(\mathbf{g}_{\mu\nu}d\mathbf{x}^\mu d\mathbf{x}^\nu)^{\frac{1}{2}(1-b)} \rightarrow \Omega^{1-b}(\mathbf{g}_{\mu\nu}d\mathbf{x}^\mu d\mathbf{x}^\nu)^{\frac{1}{2}(1-b)} ; \quad (\text{V.14})$$

The change can be compensated by the ‘‘B-F factor’’, though. Assuming that $\mathbf{f}_* \mathbf{l}_\mu = \Omega^a \mathbf{l}_\mu$ for some constant \mathbf{a} yields

$$(\mathbf{l}_\mu d\mathbf{x}^\mu)^b (\mathbf{g}_{\mu\nu} d\mathbf{x}^\mu d\mathbf{x}^\nu)^{\frac{1}{2}(1-b)} \rightarrow \Omega^{ab+1-b} (\mathbf{l}_\mu d\mathbf{x}^\mu)^b (\mathbf{g}_{\mu\nu} d\mathbf{x}^\mu d\mathbf{x}^\nu)^{\frac{1}{2}(1-b)}: \quad (\text{V.15})$$

In the undeformed case $\mathbf{b} = 0$ this is a conformal transformation with conformal factor Ω ; obtaining an isometry requires \mathbf{f} to be an isometry for the pp-wave. This is consistent with our findings in the flat case for the U-V boost [Eq. (V.6)].

For $\mathbf{b} \neq 0$ we have another option. If the exponent of Ω in Eq. (V.15) vanishes,

$$ab = \mathbf{b} - 1; \quad (\text{V.16})$$

then we do get an isometry again. For the \mathbf{b} -deformed U-V boost in Eq. (V.7), we have $\Omega = \Omega^{\frac{1}{\mathbf{b}}}$, consistently with Eq. (V.16).

In the case of plane gravitational waves, one drops the angular momentum \mathbf{J} and $@_U$ and then the issue is what to do about \mathbf{N}_b . Our only certainty so far is that the flat-space implementation [Eq. (V.8)] does not work.

The symmetry of the Bogoslovsky-Finsler model is in fact of the very special relativity (VSR) type — more precisely, a subgroup of the eight-parameter $\text{DISIM}_b(2)$, where $0 < \mathbf{b} < 1$ is a deformation parameter [7, 45]. $\text{DISIM}_b(2)$ is isomorphic to the conformal Galilei group with dynamical exponent [46]

$$z = 1 - \frac{1}{\mathbf{b}}: \quad (\text{V.17})$$

For $\mathbf{b} \neq 0$, U-V boosts (which are isometries for the Minkowski case) are deformed to Eq. (V.8), a combination of U-V boosts and relativistic dilations.

One can be puzzled about whether the ‘‘deformation trick’’ can work also for a nontrivial profile. The answer is that it *might* work for a particular profile. Let us consider, for example, a pp wave [Eq. (III.4)] written in Brinkmann coordinates with the (singular) profile

$$2H(\mathbf{X}^i; \mathbf{U}) = -\frac{\mathbf{K}_{ij}^0}{\mathbf{U}^2} \mathbf{X}^i \mathbf{X}^j: \quad \mathbf{K}_{ij}^0 = \text{const}: \quad (\text{V.18})$$

This wave has a six-parameter isometry group [32, 41, 44, 47]. It is, in particular, invariant under a U-V boost [Eq. (V.6)]. Then we find that the deformed U-V boost,

$$\mathbf{U} \rightarrow \Lambda \mathbf{U}; \quad \mathbf{X} \rightarrow \Lambda^{\frac{\mathbf{b}}{\mathbf{b}-1}} \mathbf{X}; \quad \mathbf{V} \rightarrow \Lambda^{\frac{\mathbf{b}+1}{\mathbf{b}-1}} \mathbf{V} \quad (\text{V.19})$$

leaves the Bogoslovsky-Finsler line element

$$ds_{BF} = \left(-2dUdV - d\mathbf{X}^2 - \frac{K_{ij}^0}{U^2} X^i X^j dU^2 \right)^{\frac{1-b}{2}} (dU)^b \quad (\text{V.20})$$

invariant. The usual U-V boost is recovered for $\mathbf{b} = 0$. Writing $\Lambda = \Lambda_{\frac{b}{b-1}}$ shows, moreover, that when $\mathbf{b} \neq 0$, the dynamical exponent is $z = -1 + \frac{1}{b}$, minus that in Eq. (V.17). Note that Eq. (V.19) is, once again, a conformal transformation of the pp-wave metric [Eqs. (III.4)-(V.18)] with conformal factor $\Omega^2 = \Lambda^{\frac{2b}{b-1}}$.

VI. PROLONGATION VECTORS AND SYMMETRIES

The connection of the aforementioned symmetries to integrals of the motion is established through Noether's first theorem [48]: each generator of any finite-dimensional Lie group of transformations which leaves the action form-invariant up to a surface term [49] is associated with a conserved quantity.

Consider, for example, a dynamical system with dependent and independent variables $x^\mu(\lambda)$ and λ , respectively. The most general point transformation one can have is,

$$\Upsilon = \left(\lambda; \mathbf{x} \right)_{\lambda}^{\textcircled{}} + Y^\mu \left(\lambda; \mathbf{x} \right)_{x^\mu}^{\textcircled{}}; \quad (\text{VI.1})$$

where the coefficient $\left(\lambda; \mathbf{x} \right)_{\lambda}^{\textcircled{}}$ accounts for transformations which might also involve the parameter λ . This vector can be extended to the space of the first derivatives $\dot{x}^\mu = \frac{dx^\mu}{d\lambda}$, i.e., we can consider the *first prolongation* of Υ , defined as [50, 51]

$$\text{pr}^{(1)}\Upsilon = \Upsilon + \Phi^\mu \frac{\textcircled{}}{\dot{x}^\mu}; \quad \text{where} \quad \Phi^\mu = \frac{dY^\mu}{d\lambda} - \dot{x}^\mu \frac{d}{d\lambda}; \quad (\text{VI.2})$$

The coefficient Φ^μ here is to guarantee the correct transformation law for the derivatives. Given, for example, the generator (VI.1), up to first order in the transformation parameter (say ϵ) we may write

$$\tilde{\lambda} \sim \lambda + \epsilon \left(\lambda; \mathbf{x} \right)_{\lambda}^{\textcircled{}} \quad \tilde{x}^\mu \sim x^\mu + \epsilon Y^\mu \left(\lambda; \mathbf{x} \right)_{x^\mu}^{\textcircled{}} \quad (\text{VI.3})$$

which furthermore implies

$$\frac{d\tilde{x}^\mu}{d\tilde{\lambda}} \sim \frac{d(x^\mu + \epsilon Y^\mu)}{d(\lambda + \epsilon \left(\lambda; \mathbf{x} \right)_{\lambda}^{\textcircled{}})} \sim \left(\frac{dx^\mu}{d\lambda} + \frac{dY^\mu}{d\lambda} \right) \left(1 - \epsilon \frac{d}{d\lambda} \right) \simeq \dot{x}^\mu + \epsilon \Phi^\mu; \quad (\text{VI.4})$$

With the use of the extended vector $\text{pr}^{(1)}\Upsilon$, the initial requirement of Noether's theorem written as

$$(\text{Ld}) = d\Sigma \quad (\text{VI.5})$$

where $\Sigma = \Sigma(\ ; \mathbf{x})$ is some function can be cast in infinitesimal form as

$$\text{pr}^{(1)}\Upsilon(\text{L}) + \text{L} \frac{d}{d} = \frac{d\Sigma}{d}: \quad (\text{VI.6})$$

To an appropriate generator Υ and a function Σ satisfying the above relation there corresponds a conserved quantity :

$$\text{I} = \Upsilon^\mu \frac{\partial \text{L}}{\partial \dot{x}^\mu} - \left(\dot{x}^\alpha \frac{\partial \text{L}}{\partial \dot{x}^\alpha} - \text{L} \right) - \Sigma: \quad (\text{VI.7})$$

The geodesic system is invariant under arbitrary changes of the parameter $\ ;$ therefore the inclusion of the coefficient into Eq. (VI.1) does not contribute in the conservation law. As can be seen using Eq. (VI.7), essentially multiplies the Hamiltonian, which is identically zero for Lagrangians which are homogeneous functions of degree 1 in the velocities. The coefficient plays a role instead in *Noether's second theorem* and an identity among the Euler-Lagrange equations of motion [52]. As a result, we may restrict ourselves to consider pure spacetime transformations generated by vectors $\Upsilon = \Upsilon^\alpha(\mathbf{x})\partial_\alpha$. Then the first prolongation becomes

$$\text{pr}^{(1)}\Upsilon = \Upsilon + \frac{d\Upsilon^\alpha}{d} \frac{\partial}{\partial \dot{x}^\alpha} = \Upsilon^\alpha(\mathbf{x}) \frac{\partial}{\partial x^\alpha} + \frac{\partial \Upsilon^\alpha}{\partial x^\beta} \dot{x}^\beta \frac{\partial}{\partial \dot{x}^\alpha}: \quad (\text{VI.8})$$

and Eqs. (VI.6) and (VI.7) reduce to

$$\text{pr}^{(1)}\Upsilon(\text{L}) = \frac{d\Sigma}{d}; \quad \text{I} = \Upsilon^\mu \frac{\partial \text{L}}{\partial \dot{x}^\mu} - \Sigma: \quad (\text{VI.9})$$

If for a given spacetime vector Υ , the relation $\text{pr}^{(1)}\Upsilon(\text{L}) = 0$ is satisfied (as for isometries of the geodesic system), then Σ is just a constant and can be omitted, thus having $\tilde{\text{I}} = \text{I} + \Sigma = \Upsilon^\mu \frac{\partial \text{L}}{\partial \dot{x}^\mu} = \text{const}$:

To illustrate the prolongation technique, we note that for a system in the background $\mathfrak{g}_{\mu\nu}$ [Eq. (IV.5)] whose Lagrangian is L , the first prolongation of the isometries in Eq. (V.2),

$$\Upsilon_{iso} = (\text{S}^{ij} \partial_j + \partial_i) \partial_i + (- \partial_i \text{X}^i + \partial') \partial_v;$$

is

$$\text{pr}^{(1)}\Upsilon_{iso}(\text{L}) = \left(\Upsilon_{iso}^\alpha \partial_\alpha + \frac{\partial \Upsilon_{iso}^\alpha}{\partial x^\beta} \dot{x}^\beta \frac{\partial}{\partial \dot{x}^\alpha} \right) (\text{L}): \quad (\text{VI.10})$$

If the rhs is a total derivative, then we have a symmetry for the system.

Applying Eq. (VI.10) first to the pp-wave Lagrangian

$$\mathcal{L}_{pp} = \dot{u}\dot{v} + \frac{1}{2}a_{ij}\dot{x}^i\dot{x}^j \quad (\text{VI.11})$$

confirms that Y_{iso} is a symmetry for the pp-wave.

Next, for the Bogoslovsky-Finsler Lagrangian \mathcal{L}_b in Eq. (IV.6), we find that the rhs of Eq. (VI.10) vanishes:

$$\text{pr}^{(1)}Y_{iso}(\mathcal{L}_b) = \left(m(1-b)(-a_{ij}\dot{x}^i\dot{x}^j - 2\dot{u}\dot{v})^{-\frac{1+b}{2}}\dot{u}^b \right) \text{pr}^{(1)}Y_{iso}(\mathcal{L}_{pp}) = 0; \quad (\text{VI.12})$$

proving that the Carroll group (with broken rotations) generates symmetries also for the Bogoslovsky-Finsler metric. The conserved quantities listed in Sec.V are recovered using Eq. (VI.9).

Turning now to U-V boosts we check first that for the *flat Minkowski metric* the prolongation of the deformed boost N_b in Eq. (V.8) vanishes:

$$\text{pr}^{(1)}N_b(\mathcal{L}_0) = 0 \quad (\text{VI.13})$$

and thus generates the constant of the motion \mathcal{N}_b in Eq. (V.12).

However, the same calculation carried out in the *curved* background $g_{\mu\nu}$ [Eq. (IV.5)] yields instead

$$\text{pr}^{(1)}N(\mathcal{L}_b) = m\dot{u}\dot{v}^b(b-1)^2 \left(\frac{da_{ij}}{du}\dot{x}^i\dot{x}^j \right) (-a_{ij}\dot{x}^i\dot{x}^j - 2\dot{u}\dot{v})^{\frac{-1-b}{2}}; \quad (\text{VI.14})$$

Consistently with what we said before, this vanishes for the flat metric $g_{\mu\nu}$. However, it is manifestly *not* a total derivative in general whenever $\mathbf{a} = (a_{ij})$ is not a constant matrix.

VII. AN EINSTEIN-MAXWELL EXAMPLE

In this section, we treat the motion in the Bogoslovsky-Finsler deformation of a pp-wave which is not Ricci flat. It is

$$ds^2 = (dX^1)^2 + (dX^2)^2 + 2dUdV - \frac{1}{4} \left((X^1)^2 + (X^2)^2 \right) dU^2; \quad (\text{VII.1})$$

From Eq. (24.5) on p. 385 of Ref. [32], one learns that it belongs to a class first considered by Baldwin and Jeffery [53]. It is conformally flat and is an Einstein-Maxwell solution with

a covariantly constant null Maxwell field. From the Bargmann point of view, this metric describes an isotropic harmonic oscillator in the plane with frequency \dagger [30]. The kinematic group arising from the null reduction is the Newton-Hooke group [54]. Because the metric (VII.1) is of the form of Eq. (III.4) with

$$-2H = K_{ij}X^iX^j \quad (\text{VII.2})$$

where K_{ij} is nondegenerate and independent of U , it is also a Cahen-Wallach symmetric space [55–58]. Following the procedure outlined in Sec. IV, the metric (VII.1) can be presented in the BJR form. We set $\mathbf{a} = P^tP$, where

$$P = \begin{bmatrix} \left(1 - \sin(\dagger u)\right)^{1/2} \cos & -\left(1 + \sin(\dagger u)\right)^{1/2} \sin \\ \left(1 - \sin(\dagger u)\right)^{1/2} \sin & \left(1 + \sin(\dagger u)\right)^{1/2} \cos \end{bmatrix} \quad (\text{VII.3})$$

is a solution of the Sturm-Liouville equation (IV.4) with diagonal profile $K = -\frac{\omega^2}{4}\text{Id}$. Then, using Eq. (IV.3), we end up with

$$ds^2 = (1 - \sin(\dagger u))dx^2 + (1 + \sin(\dagger u))dy^2 + 2dudv \quad (\text{VII.4})$$

which has $\mathbf{a} = P^tP = \text{diag}(1 - \sin(\dagger u); 1 + \sin(\dagger u))$. On p. 386 of Ref. [32], this result is ascribed to Brdička [59]. Equation (VII.4) shows that the U-V boost symmetry is manifestly broken.

The Souriau matrix is found by integrating the inverse of (\mathbf{a}_{ij}) , cf. Eq. (IV.14),

$$S(u) = \frac{1}{\dagger} \begin{bmatrix} \tan\left(\frac{\omega u}{2} + \frac{\pi}{4}\right) + C_1 & 0 \\ 0 & \tan\left(\frac{\omega u}{2} - \frac{\pi}{4}\right) + C_2 \end{bmatrix}; \quad (\text{VII.5})$$

where $C_{1,2}$ are integration constants. Choosing $u_0 = 0$ yields $C_1 = -1$ and $C_2 = 1$. The trajectories (IV.13)-(IV.22) for different values of \mathbf{b} are depicted in Fig. 1.

We mention that the profile of the metric (VII.1) is U -independent and therefore U -translation, $U \rightarrow U +$ is an additional isometry. This carries over trivially to its Finslerized line element [Eq. (III.6)], since both the pp-wave metric and the ‘‘Bogoslovsky-Finsler factor’’ are invariant.

VIII. BOGOSLOVSKY-FINSLER-FRIEDMANN-LEMAÎTRE MODEL

In this section we shall describe a simple extension of Bogoslovsky’s theory to take into account the expansion of the Universe. For some previous work, see Refs. [60–63]. In




FIG. 1: Consistently with (IV.13), the Bogoslovsky-Finsler geodesics project to the same curve in 2D transverse space for all values of the parameter b while their v coordinates differ, according to (IV.22), in a b -dependent term, which is linear in retarded time, u . Experiments indicate that the anisotropy and hence b is very small. When $b \rightarrow 1$ the trajectory approaches to the massless one (in heavy black), consistently with (IV.22).

contrast to our work, these authors consider only Finsler metrics which share the isotropy and spatial homogeneity of Friedmann-Lemaître models. This necessarily excludes the use of a null vector field.

A. The Λ CDM model

The simplest standard model consistent with current observational data is the spatially flat Friedmann-Lemaître model with metric

$$g_{\mu\nu}dx^\mu dx^\nu = -dt^2 + a^2 dx^2 \quad (\text{VIII.1})$$

where $a = a(t)$ and $\mathbf{x} = (x; y; z)$. The “scale factor” $a(t)$ is determined by the Einstein equations once the matter content has been specified. The favored Λ CDM model has

$$a(t) = \sinh^{\frac{2}{3}}\left(\frac{\sqrt{3\Lambda}}{4} t\right); \quad (\text{VIII.2})$$

which enjoys the remarkable property that the jerk equals 1,

$$j = a^2 \left(\frac{da}{dt} \right)^{-3} \frac{d^3 a}{dt^3} = 1 : \quad (\text{VIII.3})$$

See Ref. [64] for details and original references.

Here we shall leave the precise form of $a(t)$ unspecified. The coordinate t is called *cosmic time*. The spatial coordinate \mathbf{x} is usually said to be *comoving* since the world lines of the cosmic fluid have constant \mathbf{x} . Two events simultaneous with respect to constant time, i.e. with $\mathbf{x}_1^\mu = (0; \mathbf{x}_1)$ and $\mathbf{x}_2^\mu = (0; \mathbf{x}_2)$ have a time-dependent proper separation $a(t)(\mathbf{x}_1 - \mathbf{x}_2)$.

B. The choice of null vector field

The vector field

$$l^\mu \frac{\partial}{\partial x^\mu} = g(t) \frac{1}{\sqrt{2}} \left(a \frac{\partial}{\partial t} - \frac{\partial}{\partial z} \right) \quad (\text{VIII.4})$$

where $g(t)$ is a nonvanishing arbitrary function is past directed and null but is neither covariantly constant nor Killing, as it can be checked by a tedious calculation. The associated one-form is

$$l_\mu dx^\mu = -a^2 g \frac{1}{\sqrt{2}} \left(\frac{1}{a} dt + dz \right) : \quad (\text{VIII.5})$$

If $\dot{x}^\mu = \frac{dx^\mu}{d\lambda}$ then

$$\mathcal{L} = -m(a^2 g)^b \left(\frac{1}{\sqrt{2}} (a(t)^{-1} \dot{t} + \dot{z}) \right)^b (\dot{t}^2 - a^2 \dot{\mathbf{x}}^2)^{\frac{1}{2}(1-b)} \quad (\text{VIII.6})$$

is a possible Bogoslovsky-Finsler-type Lagrangian for a particle of mass m . It admits three commuting symmetries generated by $\frac{\partial}{\partial \mathbf{x}}$ and hence three conserved momenta $\mathbf{p} = \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{x}}}$.

If $b = 0$, then Eq. (VIII.6) is the standard action for a freely moving particle in a flat isotropic Friedmann-Lemaître universe.

C. Hubble friction

A notable feature of the free motion of a massive particle moving in a Friedmann-Lemaître universe is *Hubble friction*. The conserved momenta are

$$\mathbf{p} = ma^2 \frac{d\mathbf{x}}{d} \quad \text{where} \quad d = \sqrt{1 - \left(a \frac{d\mathbf{x}}{dt} \right)^2} dt : \quad (\text{VIII.7})$$

Here d is the increment of proper time along the world line of a particle. The four-velocity of the particle with respect to the local inertial reference frame $\frac{\partial}{\partial t}; \frac{\partial}{a(t)\partial \mathbf{x}}$ is

$$\mathbf{u} = a(t) \frac{d\mathbf{x}}{d} \quad \text{whence} \quad \mathbf{u} = \frac{\mathbf{p}}{ma(t)} : \quad (\text{VIII.8})$$

One may also define a velocity \mathbf{v} measured in units of cosmic time t , $\mathbf{v} = a(t) \frac{d\mathbf{x}}{dt}$; so that

$$d = \sqrt{1 - \mathbf{v}^2} dt; \quad \mathbf{u} = \frac{\mathbf{v}}{\sqrt{1 - \mathbf{v}^2}} \quad \text{and} \quad \mathbf{v} = \frac{\mathbf{p}}{m} \frac{1}{\sqrt{a^2 + \frac{\mathbf{p}^2}{m^2}}} : \quad (\text{VIII.9})$$

Hence,

$$\boxed{\frac{d\mathbf{x}}{dt} = \frac{\mathbf{p}}{a \sqrt{m^2 a^2 + \mathbf{p}^2}}} : \quad (\text{VIII.10})$$

Thus, in an expanding phase in which $a(t)$ increases with time, both \mathbf{v} and \mathbf{u} decrease with time. However, as a consequence of isotropy, their directions remain constant. The fact that we have three conserved momenta and the constraint

$$\left(\frac{dt}{d}\right)^2 = 1 + a^2 \left(\frac{d\mathbf{x}}{d}\right)^2 \quad (\text{VIII.11})$$

implies that the system of geodesics is completely integrable. In fact,

$$d\mathbf{x} = \frac{\mathbf{p}}{ma^2} \frac{dt}{\sqrt{1 + \left(\frac{\mathbf{p}}{ma}\right)^2}} \quad \text{and} \quad d = \frac{dt}{\sqrt{1 + \left(\frac{\mathbf{p}}{am}\right)^2}} : \quad (\text{VIII.12})$$

D. Conformal flatness

Before proceeding further we recall that the Friedmann-Lemaître metric (VIII.1) is conformally flat, as becomes clear if we define *conformal time* by

$$(t) = \int^t \frac{d\tilde{t}}{a(\tilde{t})} \quad (\text{VIII.13})$$

where the lower limit is left unspecified for the time being. In terms of cosmic time, the Friedmann-Lemaître metric (VIII.1) becomes

$$g_{\mu\nu} dx^\mu dx^\nu = a^2 \left\{ -d^2 + dz + dx^i dx^i \right\} = a^2 \left\{ 2dudv + dx^i dx^i \right\} \quad (\text{VIII.14})$$

where a^2 is regarded as a function of conformal time (t) , and we introduce the light-cone coordinates

$$u = \frac{z +}{\sqrt{2}}; \quad v = \frac{z -}{\sqrt{2}} : \quad (\text{VIII.15})$$

From Eq. (VIII.5), we learn that $l_\mu dx^\mu = -a^2 g du$ whence our Bogoslovsky-Finsler-ized line element is

$$ds = a^{1+b} g^b \left(-2dudv - dx^i dx^i \right)^{\frac{1}{2}(1-b)} (du)^b; \quad (\text{VIII.16})$$

This Lagrangian would yield Bogoslovsky's original flat spacetime model provided we choose $g(t)$ such that

$$f = a^{1+b} g^b = 1; \quad (\text{VIII.17})$$

The only freedom with this model would be to introduce an arbitrary factor

$$\mathcal{L}_f = f(\cdot) \left(-2\dot{u}\dot{v} - \dot{x}^i \dot{x}^i \right)^{\frac{1}{2}(1-b)} (\dot{u})^b; \quad (\text{VIII.18})$$

which amounts to saying that the mass depends upon cosmic time.

This situation is the same as in the ordinary spatially flat Friedmann-Lemaître cosmology for which $b = 0$. We can either say the Universe is expanding, but our rulers, are constructed from massive particles, all of whose masses are constant in cosmic time t , or that the Universe is time independent, but the rulers all change with the same time dependence. In that case, the phenomenon of Hubble friction would be ascribed not to the expansion of the Universe but to masses are getting heavier.

E. Redshifting

If we adopt (VIII.18) then light rays move along straight lines in $(t; \mathbf{x})$ coordinates. Emitters and observers (e.g. galaxies and astronomers) are usually held to be at rest in these coordinates.

Suppose the observer is at the origin at $(t_0; 0; 0; 0)$ and receives light rays from a galaxy at $(t_e; x_e; y_e; z_e)$ so that the duration of emission in conformal time is d_e and the duration of the corresponding observation is d_0 , then

$$d_e = d_0; \quad (\text{VIII.19})$$

Then the emitted and observed proper times are $d_e = f(t_e) d_e; d_0 = f(t_0) d_0$ and so the redshift is

$$1 + z = d_0/d_e = f(t_0)/f(t_e); \quad (\text{VIII.20})$$

Thus, if the universe is expanding — that is, if $f' > 0$ — then the signal received is redshifted, and contrariwise if the Universe is contracting — that is, if $f' < 0$.

Note that under these assumptions, the emitted light from all galaxies at the same conformal time will be redshifted in the same way. That is, *the redshift should be isotropic*.

F. A possible choice for $f(\eta)$

As mentioned earlier, our observed Universe is well described by a scale factor $a(t)$ given by Eq. (VIII.2). Applying the Einstein equations to the Friedmann-Lemaître metric [Eq. (VIII.1)], one finds that it is supported by a pressure-free fluid (some of it visible and some of it not – so-called dark matter) and a positive cosmological constant term Λ often called dark energy. Near the big bang — i.e., for small t — $a(t) \propto t^{2/3}$, because the Λ term is negligible. This is the Einstein-de Sitter model. At late times, $a(t) \propto \exp \sqrt{\Lambda=3} t$, which exhibits cosmic acceleration. This is de Sitter spacetime.

From Eq. (VIII.13), choosing Eq. (VIII.2) and setting $ag = 1$ in Eq. (VIII.16), we have

$$t = \int_0^t \sinh^{-\frac{2}{3}} \left(\frac{\sqrt{3\Lambda}}{4} \tilde{t} \right) d\tilde{t} \quad (\text{VIII.21a})$$

$$f(\eta) = a(t) = \sinh^{\frac{2}{3}} \left(\frac{\sqrt{3\Lambda}}{4} t \right); \quad (\text{VIII.21b})$$

This step depends only on the scale factor a in (VIII.1) and does not involve the deformation




FIG. 2: The conformal factor (VIII.21b) of the Friedmann-Lemaître model (VIII.1), expressed as function of the conformal time, η , obtained by numerical integration of (VIII.21).

parameter b . See Fig.2. It is worth noting that conformal time as a function of cosmic time is bounded from above — as happens for de Sitter space, to which our spacetime tends when $t \rightarrow \infty$.

IX. BOGOSLOVSKY-FINSLER-FRIEDMANN-LEMAÎTRE GEODESICS

Written in coordinates $(\eta; \mathbf{x}; \mathbf{y}; \mathbf{z})$, the Lagrangian (VIII.18) is

$$\mathcal{L}_f = -mf(\eta) \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^b (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{\frac{1}{2}(1-b)}; \quad (\text{IX.1})$$

providing us with the momenta

$$\mathbf{p}_x = m(1-b)f(\eta) \dot{\mathbf{x}} \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^b (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{-\frac{1}{2}(1+b)} \quad (\text{IX.2a})$$

$$\mathbf{p}_y = m(1-b)f(\eta) \dot{\mathbf{y}} \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^b (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{-\frac{1}{2}(1+b)} \quad (\text{IX.2b})$$

$$\begin{aligned} \mathbf{p}_z &= m(1-b)f(\eta) \dot{\mathbf{z}} \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^b (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{-\frac{1}{2}(1+b)} \\ &\quad - \frac{b}{\sqrt{2}} mf(\eta) \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^{-1+b} (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{\frac{1}{2}(1-b)} \end{aligned} \quad (\text{IX.2c})$$

$$\begin{aligned} \mathbf{p}_\eta &= -mf(\eta) \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^{b-1} (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{-\frac{1}{2}(1+b)} \\ &\quad \left((1-b) \dot{\eta} \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right) + \frac{b}{\sqrt{2}} (\dot{\eta}^2 - \dot{\mathbf{x}}^2 + \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2) \right) \end{aligned} \quad (\text{IX.2d})$$

Evidently the three momenta $\mathbf{p}_x, \mathbf{p}_y, \mathbf{p}_z$ are conserved. Moreover, since

$$\frac{\mathbf{p}_y}{\mathbf{p}_x} = \frac{dy}{dx} \quad (\text{IX.3})$$

the projections of the geodesics onto the transverse $\mathbf{x} - \mathbf{y}$ plane are straight lines. Choosing the proper time as parameter, $\eta = \tau$, one has the constraint

$$f(\tau) \left(\frac{\dot{\eta} + \dot{\mathbf{z}}}{\sqrt{2}} \right)^b (\dot{\eta}^2 - \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2)^{\frac{1}{2}(1-b)} = 1 \quad (\text{IX.4})$$

which we may rewrite in terms of conformal time, $\eta = \tau$, as an equation for τ as

$$\tau' = f(\tau) \left(\frac{1 + \mathbf{z}'}{\sqrt{2}} \right)^b (1 - (\mathbf{x}')^2 - (\mathbf{y}')^2 - (\mathbf{z}')^2)^{\frac{1}{2}(1-b)}; \quad (\text{IX.5})$$

where $(\mathbf{x}'; \mathbf{y}'; \mathbf{z}') = \left(\frac{dx}{d\eta}, \frac{dy}{d\eta}, \frac{dz}{d\eta} \right)$.

If $f' = 0$, then $\mathbf{p}_\mu = f(\eta)$ is independent of η , leaving us with the same straight-line motion at constant velocity as for the flat Bogoslovsky spacetime.

As we have seen above, even if $f' \neq 0$, the projections of the motion on the $\mathbf{x} - \mathbf{y}$ plane are straight lines, although not with constant speed with respect to the conformal time τ . The

speeds of the projections onto the $\mathbf{x} - \mathbf{z}$ and $\mathbf{y} - \mathbf{z}$ planes are also not at constant speed, but they are not straight lines either. Over conformal times that are short compared with $\frac{f}{f'}$ they are approximately straight lines with slopes given by $\frac{p_x}{mf(\eta)}$ but over longer time periods, the speeds and directions change reflecting precisely the effects of Hubble friction.

The geodesics are conveniently studied by switching to conformal time, τ . Introducing

$$\dot{\mathbf{u}} = \frac{\dot{\mathbf{x}} + \dot{\mathbf{z}}}{\sqrt{2}} \quad \text{and} \quad \dot{w} = \dot{\mathbf{x}}^2 - \dot{\mathbf{y}}^2 - \dot{\mathbf{z}}^2 \quad (\text{IX.6})$$

in place of $\dot{\mathbf{x}}$ and $\dot{\mathbf{z}}$, Eqs. (IX.2a), (IX.2b), and (IX.2c) become

$$m(b-1)f(\tau)\dot{\mathbf{x}}\dot{u}^b + p_x \dot{w}^{\frac{b+1}{2}} = 0 \quad (\text{IX.7a})$$

$$m(b-1)f(\tau)\dot{\mathbf{y}}\dot{u}^b + p_y \dot{w}^{\frac{b+1}{2}} = 0 \quad (\text{IX.7b})$$

$$\frac{mf(\tau)}{2\sqrt{2}}\dot{u}^{b-1}\dot{w}^{\frac{1}{2}(-b-1)}(2(b-1)\dot{u}^2 + (b+1)\dot{w} - (b-1)(\dot{\mathbf{x}}^2 + \dot{\mathbf{y}}^2)) + p_z = 0: \quad (\text{IX.7c})$$

The first two equations imply identical evolution:

$$\dot{\mathbf{x}} = \frac{\dot{u}^{-b}\dot{w}^{\frac{b+1}{2}}}{m(1-b)f(\tau)} p_x; \quad \dot{\mathbf{y}} = \frac{\dot{u}^{-b}\dot{w}^{\frac{b+1}{2}}}{m(1-b)f(\tau)} p_y \quad (\text{IX.8})$$

which confirms once again that the transverse projection is a straight line, owing to $\dot{\mathbf{x}}=\dot{\mathbf{y}} = p_x=p_y = \text{const}$: With their help, Eq. (IX.7c) becomes

$$-\sqrt{2}(b-1)m^2f(\tau)^2\dot{u}^{2b}(2(b-1)\dot{u}^2 + (b+1)\dot{w}) - 4(b-1)mp_zf(\tau)\dot{u}^{b+1}\dot{w}^{\frac{b+1}{2}} + \sqrt{2}(p_x^2 + p_y^2)\dot{w}^{b+1} = 0: \quad (\text{IX.9})$$

By reparametrizing $w(\tau)$ as

$$w(\tau) = \int (\tau')^2 \dot{u}(\tau')^2 d\tau'; \quad (\text{IX.10})$$

where τ' is a new function that we introduce, Eq. (IX.9) reduces from a differential one to an algebraic

$$-\sqrt{2}(b-1)m^2f(\tau')^2((b+1)(\tau')^2 + 2(b-1)) - 4(b-1)mf(\tau')p_z(\tau')^{b+1} + \sqrt{2}(p_x^2 + p_y^2)(\tau')^{2(b+1)} = 0: \quad (\text{IX.11})$$

For $b = 0$ this is simply quadratic in τ' , but for $b \neq 0$ it is not trivial to solve it for τ' , cf. Fig.3(a). However, as it is quadratic in $f(\tau')$, the inverse problem [which amounts to choosing τ' to find the corresponding $f(\tau')$] still works. The functions τ' and $w(\tau')$ are plotted in Fig.3. Using Eqs. (IX.6), (IX.10) and (IX.8) we get




FIG. 3: (a) $\sigma(\eta)$ and (b) $w(\eta)$ in (IX.10) plotted for $\mathbf{b} = \mathbf{0}$ and for $\mathbf{b} = \mathbf{0.5}$.

$$\dot{\cdot} = \frac{\dot{u}}{2\sqrt{2}} \left(2 + \frac{(p_x^2 + p_y^2)^{2(b+1)}}{(b-1)^2 m^2 f(\cdot)^2} \right) \quad (\text{IX.12a})$$

$$\dot{x} = \dot{u} \frac{b+1}{(1-b)m f(\cdot)} p_x \quad (\text{IX.12b})$$

$$\dot{y} = \dot{u} \frac{b+1}{(1-b)m f(\cdot)} p_y \quad (\text{IX.12c})$$

$$\dot{z} = \frac{\dot{u}}{2\sqrt{2}} \left(2 - \frac{(p_x^2 + p_y^2)^{2(b+1)}}{(b-1)^2 m^2 f(\cdot)^2} \right) \quad (\text{IX.12d})$$

together with the algebraic constraint between (\cdot) and $f(\cdot)$ in Eq. (IX.11).

The joint system can be shown to satisfy the Euler-Lagrange equations.

The $u(\cdot)$ that remains unspecified in (IX.12) and disappears from Eq. (IX.11) serves as a gauge parameter [by seeing the ratios of derivatives that are being formed in Eq. (IX.12)], for which we can simply set $u(\cdot) = \cdot$. So in this time-gauge $(\cdot) + z(\cdot) = \sqrt{2}$, which is compatible with Eqs. (IX.12a) and (IX.12d), as seen above. We thus have, in the

“conformal-time gauge,”

$$\frac{dx}{d} = \frac{2\sqrt{2}(1-b)m^{b+1}f(\)}{(1-b)^2m^2(\ ^2+2)f(\)^2 + (\mathbf{p}_x^2 + \mathbf{p}_y^2)^{2(b+1)}} \mathbf{p}_x \quad (\text{IX.13a})$$

$$\frac{dy}{d} = \frac{2\sqrt{2}(1-b)m^{b+1}f(\)}{(1-b)^2m^2(\ ^2+2)f(\)^2 + (\mathbf{p}_x^2 + \mathbf{p}_y^2)^{2(b+1)}} \mathbf{p}_y \quad (\text{IX.13b})$$

$$\frac{dz}{d} = \frac{2 - \ ^2 - \frac{(\mathbf{p}_x^2 + \mathbf{p}_y^2)^{2(b+1)}}{(1-b)^2m^2f(\)^2}}{2 + \ ^2 + \frac{(\mathbf{p}_x^2 + \mathbf{p}_y^2)^{2(b+1)}}{(1-b)^2m^2f(\)^2}}; \quad (\text{IX.13c})$$

The Friedmann-Lemaître case $b = 0$

If $b = 0$, the algebraic relation (IX.11) is quadratic and can be solved for $\$:

$$\ (\) = \pm \frac{\sqrt{2}mf(\)}{\sqrt{m^2f(\)^2 + \mathbf{p}^2 \pm p_z}}; \quad (\text{IX.14})$$

shown by the blue line in Fig. 3(a) for the upper sign⁶. In terms of conformal time

$$\boxed{\frac{d\mathbf{x}}{d} = \pm \frac{\mathbf{p}}{\sqrt{m^2f(\)^2 + \mathbf{p}^2}}}; \quad (\text{IX.15})$$

to be compared with Eq. (VIII.10). The consistency with the equations in Sec. VIII C follows from

$$d = \frac{mf^2d}{\sqrt{m^2f^2 + \mathbf{p}^2}}; \quad d\mathbf{x} = \frac{d}{\sqrt{m^2f^2 + \mathbf{p}^2}} \mathbf{p} \Rightarrow \boxed{\frac{d\mathbf{x}}{d} = \frac{1}{mf^2} \mathbf{p}}; \quad (\text{IX.16})$$

We could not obtain analytical expressions; however, using the numerically calculated values of $f(\)$ (see Fig. 2) allows us to plot $\mathbf{x}(\)$ by solving Eq. (IX.15), as shown in Fig. 4⁷.

The $b = 0$ case nicely illustrates *Hubble friction*: all trajectories slow down and ultimately come to rest. For $b \neq 0$ it seems that this happens only for the transverse motion but not for the motion in the z direction, see Fig. 6. The slowing down in the transverse case is plausible from Eqs. (IX.12b) and (IX.12c).

⁶ Choosing the lower sign would amount to an overall sign change when that of p_z is also reversed.

⁷ The two signs in (IX.14) can be compensated by $\mathbf{p} \rightarrow -\mathbf{p}$ implying an overall sign change. In what follows the upper sign will be chosen.




FIG. 4: For $\mathbf{b} = \mathbf{0}$ all trajectories follow straight lines and have identical evolution. For $\mathbf{b} = \mathbf{0.5}$ $z(\eta)$ become different from the transverse trajectories $(x(\eta), y(\eta))$ consistently with (IX.13), as shown in Fig.5.




FIG. 5: For $\mathbf{b} > \mathbf{0}$ the motion in the $x - z$ plane is not more along a straight line (as it is for $\mathbf{b} = \mathbf{0}$). The Hubble friction slows down the z motion for $\mathbf{b} = \mathbf{0}$ but not when $\mathbf{b} > \mathbf{0}$.

X. CONCLUSION

In this paper, motivated by work by Bogoslovsky [3–5, 7], and by that of Tavakol and Van den Bergh, and Roxburgh [9, 11, 12], and by Cohen and Glashow [1], and more recently by others [14–21], we have studied the free motion of a massive particle moving in a one-parameter family of Finslerian deformations of a plane gravitational wave. By free motion, we mean that it extremizes the proper time along its timelike world line. Finslerian proper




FIG. 6: For the Friedmann-Lemaître model for $\mathbf{b} = \mathbf{0}$ the 3D trajectory is a straight line. For the Bogoslovsky-Finsler modification $\mathbf{b} = \mathbf{0.5}$ however, while the projection to the $x - y$ plane is still along a straight line, the z -component becomes curved, consistently with Fig.5.

time is measured by replacing the usual square-root integrand

$$\sqrt{-g_{\mu\nu} \frac{dx^\mu}{d} \frac{dx^\nu}{d}} \quad \text{with} \quad \left(-g_{\mu\nu} l^\mu \frac{dx^\nu}{d}\right)^b \left(-g_{\mu\nu} \frac{dx^\mu}{d} \frac{dx^\nu}{d}\right)^{\frac{1}{2}(1-b)};$$

where l^μ is a null vector field and b is a dimensionless constant.

In earlier work, we have shown that because of the five-dimensional isometry group of plane gravitational waves, the motion of the usual timelike geodesics is completely integrable.

In the present paper, we have shown that the five-dimensional partially broken Carroll symmetry group G_5 remains a symmetry of our Finslerian line element provided we choose the null vector field l^μ to be the covariantly constant null vector of the underlying gravitational wave. As a consequence, we find that that not only is the free motion completely integrable, but it differs only in that the “vertical” coordinate v involves in turn a b -dependent term, which is linear in the retarded time coordinate u . The motion in the transverse directions is unchanged. The situation is analogous to what happens for massive vs massless geodesics in a pp-wave [42].

The symmetry of the Bogoslovsky-Finsler model is in fact that of the very special rela-

tivity (VRS) type ; in the Minkowski case it is the eight-parameter $\text{DISIM}_b(2)$ [7, 40, 45]. The clue is to deform a U-V boost \mathbf{N}_0 to \mathbf{N}_b as in Eq. (V.8). The trick works for certain nontrivial profiles, as for the \mathbf{U}^{-2} discussed at the end of Sec.V.

We have also examined the free motion of a Finslerian deformation of a homogeneous pp-wave which is an Einstein-Maxwell solution. The resulting spacetime is a Cahen-Wallach symmetric space [55] and arises in a wide variety of physical applications and whose null reduction in the fashion of Eisenhart and Duval et al. [30] is a simple harmonic oscillator with a Newton-Hooke-type symmetry. Here again, the free motion is qualitatively independent of the deformation parameter b .

We have also studied a simple anisotropic cosmological model based on that of Friedmann and Lemaître with vanishing spatial curvature. Because the latter is conformally flat, the motion of massive particles is equivalent to motion in flat Bogoslovsky spacetime except that all masses become time dependent with identical time dependence.

Although our present Universe shows little sign of anisotropy of the sort that arises in Bogoslovsky-Finsler metrics, that may not have been true earlier in the history of the Universe since the absence of anisotropy now is usually ascribed to a rapid phase of inflation during which the scale factor of the Universe increased by a factor of perhaps 60 e-folds. It is of interest, therefore to study geodesics in Bogoslovsky-Finsler deformations of Friedmann-Lemaître metrics.

Acknowledgments

We would like to thank Ettore Minguzzi for correspondence. This work was partially supported by the National Key Research and Development Program of China (No. 2016YFE0130800) and the National Natural Science Foundation of China (Grant No. 11975320).

-
- [1] A. G. Cohen and S. L. Glashow, “Very special relativity”, *Phys. Rev. Lett.* **97** (2006) 021601 doi:10.1103/PhysRevLett.97.021601 [hep-ph/0601236].
 - [2] V W Hughes, H G Robinson and V Betran-Lopez, “Upper limit for the anisotropy of inertial mass from nuclear resonance experiments,” *Phys. Rev. Lett.* **4** (1961) 342-4 ; R W P Drever,

- “A search for anisotropy of inertial mass using a free precession method, ” *Phil. Mag.* **6** (1961) 683-7.
- [3] G. Y. Bogoslovsky, “A Special-Relativistic Theory of the locally Anisotropic Spacetime: I The metric and Group of motions of the Anisotropic Space of Events,” *Il Nuovo Cimento* **40 B** (1977) 99-115
- [4] G.Y. Bogoslovsky, “A Special-Relativistic Theory of the Locally Anisotropic Spacetime. II: Mechanics and Electrodynamics in the Anisotropic Space,” “ *Il Nuovo Cimento* **40 B** (1977) 116-134.
- [5] V. Balan, G. Y. Bogoslovsky, S. S. Kokarev, D. G. Pavlov, S. V. Siparov and N. Voicu, “Geometrical Models of the Locally Anisotropic Space-Time,” *J. Mod. Phys.* **3** (2012) 1314 doi:10.4236/jmp.2012.329170 [arXiv:1111.4346 [gr-qc]].
- [6] G. W. Gibbons, J. Gomis and C. N. Pope, “General very special relativity is Finsler geometry,” *Phys. Rev. D* **76** (2007) 081701 doi:10.1103/PhysRevD.76.081701 [arXiv:0707.2174 [hep-th]].
- [7] G. Y. Bogoslovsky, “DISIMb(2) Local Relativistic Symmetry and Finslerian Extension of the Theory of Relativity,” arXiv:2002.02843 [gr-qc].
- [8] G. Randers, “On an Asymmetrical Metric in the Four-Space of General Relativity,” *Phys. Rev.* **59** (1941) no.2, 195. doi:10.1103/PhysRev.59.195
- [9] Ian W. Roxburgh, “Finsler Spaces with Riemannian Geodesics,” *Gen. Rel. Grav.* **23** 1071 (1991)
- [10] Ian W. Roxburgh, R. K. Tavakol and N. Van den Bergh “The geometry of space time and Berwald spaces. A review of results on no-Riemannian Berwald spaces. *Tensor (N.S.)* **51** (1992) 72.
- [11] R. K. Tavakol and N. Van den Bergh, “Finsler spaces and the underlying geometry of space-time,” *Phys. Lett. A* **112**, 23, (1985)
- [12] R. K. Tavakol and N. Van den Bergh, “Viability Criteria for the Theories of Gravity and Finsler Spaces,” *Gen. Rel. Grav.* **18** 849 (1986)
- [13] G. W. Gibbons, C. A. R. Herdeiro, C. M. Warnick and M. C. Werner, “Stationary Metrics and Optical Zermelo-Randers-Finsler Geometry,” *Phys. Rev. D* **79** (2009) 044022 doi:10.1103/PhysRevD.79.044022 [arXiv:0811.2877 [gr-qc]].
- [14] A. Kostelecky, “Riemann-Finsler geometry and Lorentz-violating kinematics,” *Phys. Lett. B* **701** (2011), 137-143 doi:10.1016/j.physletb.2011.05.041 [arXiv:1104.5488 [hep-th]].

- [15] A. Fuster and C. Pabst, “Finsler pp-waves,” *Phys. Rev. D* **94** (2016) no.10, 104072 doi:10.1103/PhysRevD.94.104072 [arXiv:1510.03058 [gr-qc]].
- [16] A. Fuster, C. Pabst and C. Pfeifer, “Berwald spacetimes and very special relativity,” *Phys. Rev. D* **98** (2018) no.8, 084062 doi:10.1103/PhysRevD.98.084062 [arXiv:1804.09727 [gr-qc]].
- [17] A. García-Parrado Gómez-Lobo and E. Minguzzi, “Pseudo-Finsler spaces modeled on a pseudo-Minkowski space,” *Rept. Math. Phys.* **82** (2018), 29-42. doi:10.1016/S0034-4877(18)30069-7 [arXiv:1612.00829 [math.DG]].
- [18] C. Lämmerzahl and V. Perlick, “Finsler geometry as a model for relativistic gravity,” *Int. J. Geom. Meth. Mod. Phys.* **15** (2018) no.supp01, 1850166 doi:10.1142/S0219887818501669 [arXiv:1802.10043 [gr-qc]].
- [19] B. R. Edwards and V. A. Kostelecky, “RiemannFinsler geometry and Lorentz-violating scalar fields,” *Phys. Lett. B* **786** (2018), 319-326 doi:10.1016/j.physletb.2018.10.011 [arXiv:1809.05535 [hep-th]].
- [20] A. Fuster, S. Heefer, C. Pfeifer and N. Voicu, “On the non metrizable of Berwald Finsler spacetimes,” [arXiv:2003.02300 [math.DG]].
- [21] A. Bernal, M. A. Javaloyes and M. Sánchez, “Foundations of Finsler spacetimes from the Observers’ Viewpoint,” arXiv:2003.00455 [gr-qc].
- [22] M. Hohmann, C. Pfeifer and N. Voicu, “Finsler gravity action from variational completion,” *Phys. Rev. D* **100** (2019) no.6, 064035 doi:10.1103/PhysRevD.100.064035 [arXiv:1812.11161 [gr-qc]].
- [23] E. Caponio and A. Masiello, “On the analyticity of static solutions of a field equation in Finsler gravity,” *Universe* **6** (2020) no.4, 59 doi:10.3390/universe6040059 [arXiv:2004.10613 [math.DG]].
- [24] E. Caponio, M. A. Javaloyes and A. Masiello, “On the energy functional on Finsler manifolds and applications to stationary spacetimes,” *Math. Ann.* **351** (2011), 365-392 doi:10.1007/s00208-010-0602-7 [arXiv:math/0702323 [math.DG]].
- [25] H. Rund, *The Differential Geometry of Finsler spaces*. Springer Verlag (1953)
- [26] J. E. Ehlers, F. A. Pirani and A. Schild, ”The geometry of free fall and light propagation,” in *General Relativity*. L. O. Raifeartaigh ed. Oxford University Press, London (1972).
- [27] G. Y. Bogoslovsky and H. F. Goenner, “Finslerian spaces possessing local relativistic symmetry,” *Gen. Rel. Grav.* **31** (1999) 1565 doi:10.1023/A:1026786505326 [gr-qc/9904081].

- [28] G. Y. Bogoslovsky, “Some physical displays of the space anisotropy relevant to the feasibility of its being detected at a laboratory,” arXiv:0706.2621 [gr-qc].
- [29] M. W. Brinkmann, “Einstein spaces which are mapped conformally on each other,” *Math. Ann.* **94** (1925) 119–145.
- [30] C. Duval, G. Burdet, H. Kunzle, M. Perrin, “Bargmann structures and Newton-Cartan theory,” *Phys. Rev. D* **31** (1985) 1841; C. Duval, G.W. Gibbons, P. Horvathy, “Celestial mechanics, conformal structures and gravitational waves,” *Phys. Rev.* **D43** (1991) 3907. [hep-th/0512188].
- [31] H. Bondi, F. A. E. Pirani and I. Robinson, “Gravitational waves in general relativity. 3. Exact plane waves,” *Proc. Roy. Soc. Lond. A* **251** (1959) 519. doi:10.1098/rspa.1959.0124
- [32] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers and E. Herlt, “Exact solutions of Einstein’s field equations,” Cambridge Univ. Press (2003). doi:10.1017/CBO9780511535185
- [33] J. M. Lévy-Leblond, “Une nouvelle limite non-relativiste du group de Poincaré,” *Ann. Inst. H Poincaré* **3** (1965) 1;
- [34] C. Duval, G. W. Gibbons, P. A. Horvathy and P. M. Zhang, “Carroll versus Newton and Galilei: two dual non-Einsteinian concepts of time,” *Class. Quant. Grav.* **31** (2014) 085016 [arXiv:1402.5894 [gr-qc]].
- [35] E. Bergshoeff, J. Gomis and G. Longhi, “Dynamics of Carroll Particles,” *Class. Quant. Grav.* **31** (2014) no.20, 205009 doi:10.1088/0264-9381/31/20/205009 [arXiv:1405.2264 [hep-th]].
- [36] C. Duval, G. W. Gibbons, P. A. Horvathy and P.-M. Zhang, “Carroll symmetry of plane gravitational waves,” *Class. Quant. Grav.* **34** (2017) no.17, 175003 doi:10.1088/1361-6382/aa7f62 [arXiv:1702.08284 [gr-qc]]; J.-M. Souriau, “Ondes et radiations gravitationnelles,” *Colloques Internationaux du CNRS No 220*, pp. 243-256. Paris (1973).
- [37] A. A. Coley, G. W. Gibbons, S. Hervik and C. N. Pope, “Metrics With Vanishing Quantum Corrections,” *Class. Quant. Grav.* **25** (2008) 145017 doi:10.1088/0264-9381/25/14/145017 [arXiv:0803.2438 [hep-th]].
- [38] W. Voigt, “Über das Doppler’sche Prinzip”, *Gott. Nachr.* 2 41-51 (1887)
- [39] O. Chashchina, N. Dudisheva and Z. K. Silagadze, “Voigt transformations in retrospect: missed opportunities? ”, *Annales Fond. Broglie* 44 (2019) 39-110. arXiv:1609.08647 [physics.hist-ph].
- [40] S. Dhasmana and Z. Silagadze, “Finsler space-time in light of Segal’s principle,” *Mod. Phys.*

- Lett. A **35** (2019) no.06, 2050019 doi:10.1142/S0217732320500194 [arXiv:1911.04233 [gr-qc]].
- [41] P.-M. Zhang, M. Cariglia, M. Elbistan and P. A. Horvathy, “Scaling and conformal symmetries for plane gravitational waves,” J. Math. Phys. **61** (2020) no.2, 022502 doi:10.1063/1.5136078, 10.1063/1.513607 [arXiv:1905.08661 [gr-qc]].
- [42] K. Andrzejewski, N. Dimakis, M. Elbistan, P. A. Horvathy, P. Kosinski and P.-M. Zhang, “Conformal symmetries and integrals of the motion in pp waves with external electromagnetic fields,” arXiv:2003.07649 [gr-qc].
- [43] M. Perrin, G. Burdet and C. Duval, “Chrono-projective Invariance of the Five-dimensional Schrödinger Formalism,” Class. Quant. Grav. **3** (1986) 461. doi:10.1088/0264-9381/3/3/020
- [44] A. Ilderton, “Screw-symmetric gravitational waves: a double copy of the vortex,” Phys. Lett. B **782** (2018), 22-27 doi:10.1016/j.physletb.2018.04.069 [arXiv:1804.07290 [gr-qc]].
- [45] G. Gibbons, J. Gomis and C. Pope, “Deforming the Maxwell-Sim Algebra,” Phys. Rev. D **82** (2010), 065002 doi:10.1103/PhysRevD.82.065002 [arXiv:0910.3220 [hep-th]].
- [46] J. Negro, M. A. del Olmo, An A. Rodríguez-Marco, “Nonrelativistic conformal groups,” J. Math. Phys. **38** (1997), 3786, and “Nonrelativistic conformal groups. II. Further developments and physical applications,” *ibid.* 3810. M. Henkel, “Local Scale Invariance and Strongly Anisotropic Equilibrium Critical Systems,” Phys. Rev. Lett. **78** (1997), 1940-1943 doi:10.1103/PhysRevLett.78.1940 [arXiv:cond-mat/9610174 [cond-mat.stat-mech]]. J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, “Exotic Galilean conformal symmetry and its dynamical realizations,” Phys. Lett. A **357** (2006), 1. [arXiv:hep-th/0511259]. A. Bagchi, and R. Gopakumar, “Galilean Conformal Algebras and AdS/CFT,” arXiv:0902.1385 [hep-th]. C. Duval and P. A. Horvathy, “Non-relativistic conformal symmetries and Newton-Cartan structures,” J. Phys. A **42** (2009) 465206 doi:10.1088/1751-8113/42/46/465206. [arXiv:0904.0531 [math-ph]].
- [47] K. Andrzejewski and S. Prencel, “Memory effect, conformal symmetry and gravitational plane waves,” Phys. Lett. B **782** (2018), 421-426 doi:10.1016/j.physletb.2018.05.072 [arXiv:1804.10979 [gr-qc]].
- [48] E. Noether, “Invariante Variationsproblem”, Gott. Nachr. 1918 (1918) 235; [English translation M. A. Tavel: E. Noether, “Invariant Variation Problems” Transp. Theory Statist. Phys. 1 (1971) 186]
- [49] E. Bessel-Hagen “Über die Erhaltungssätze der Elektrodynamik, Math. Ann. **84**, 258-276

(1921)

- [50] P. J. Olver, *Applications of Lie Groups to Differential Equations*, 2nd edn, Berlin, Springer, (2000).
- [51] T. Christodoulakis, N. Dimakis and Petros A. Terzis, “Lie point and variational symmetries in minisuperspace Einstein gravity,” *J. Phys. A* **47** (2014) 095202
- [52] K. Sundermeyer, *Constrained Dynamics*, Springer Verlag, Berlin, Heidelberg, New York (1982)
- [53] O. R. Baldwin and G. B. Jeffery, “The Relativity Theory of Plane Waves” *Proc. Roy Soc. A.* **111** (1926) 95-104
- [54] G. W. Gibbons and C. E. Patricot, “Newton-Hooke space-times, Hpp waves and the cosmological constant,” *Class. Quant. Grav.* **20** (2003) 5225 doi:10.1088/0264-9381/20/23/016 [hep-th/0308200].
- [55] M. Cahen and N. Wallach, “Lorentzian Symmetric Spaces,” *Bull. Am. Math. Soc.* **76** (1970) 585-591.
- [56] J. M. Figueroa-O’Farrill and G. Papadopoulos, “Homogeneous fluxes, branes and a maximally supersymmetric solution of M theory,” *JHEP* **0108** (2001) 036 doi:10.1088/1126-6708/2001/08/036 [hep-th/0105308].
- [57] M. Blau and M. O’Loughlin, “Homogeneous plane waves,” *Nucl. Phys. B* **654** (2003) 135 doi:10.1016/S0550-3213(03)00055-5 [hep-th/0212135].
- [58] M. Blau, P. Meessen and M. O’Loughlin, “Gödel, Penrose, anti-Mach: Extra supersymmetries of time dependent plane waves,” *JHEP* **0309** (2003) 072 doi:10.1088/1126-6708/2003/09/072 [hep-th/0306161]
- [59] M. Brdička, “On Gravitational Waves” *Proc. Roy. Irish. Acad. A* **54** (1951) 137
- [60] I. W. Roxburgh and R. Tavakol, “The Gravitational Theories of Poincaré and Milne and non-Riemannian Kinematic models of the Universe”, *Mon. Not. R. Astr. Soc.* **170** (1975) 599-610
- [61] J. D. Bekenstein, “The Relation between physical and gravitational geometry,” *Phys. Rev. D* **48** (1993) 3641 doi:10.1103/PhysRevD.48.3641 [gr-qc/9211017].
- [62] A. P. Kouretsis, M. Stathakopoulos and P. C. Stavrinou, “The General Very Special Relativity in Finsler Cosmology,” *Phys. Rev. D* **79** (2009) 104011 doi:10.1103/PhysRevD.79.104011 [arXiv:0810.3267 [gr-qc]].
- [63] M. Hohmann and C. Pfeifer, “Geodesics and the magnitude-redshift relation on

cosmologically symmetric Finsler spacetimes,” *Phys. Rev. D* **95** (2017) 104021
doi:10.1103/PhysRevD.95.104021 [arXiv:1612.08187 [gr-qc]].

- [64] M. Dunajski and G. Gibbons, “Cosmic Jerk, Snap and Beyond,” *Class. Quant. Grav.* **25**
(2008) 235012 doi:10.1088/0264-9381/25/23/235012 [arXiv:0807.0207 [gr-qc]].