

A probabilistic proof of the finite geometric series

Raju Dey and Suchandan Kayal*

Department of Mathematics, National Institute of Technology Rourkela, Rourkela-769008, India

Abstract

In this note, we present a probabilistic proof of the well-known finite geometric series. The proof follows by taking the moments of the sum and the difference of two independent exponentially distributed random variables.

1 Introduction

There are various popular series in the literature related to infinite series. One of these is the geometric series. Each term of the geometric series is a constant multiple of the previous term. For example, if a is the first entry of the series, then the second entry is a constant (a real number, say $\rho \neq 0$) multiple of a , the third entry is (same) constant multiple of $a\rho$, and so on. The real number ρ is known as the common ratio of the series. The infinite geometric series is

$$a + a\rho + a\rho^2 + a\rho^3 + \cdots . \quad (1.1)$$

Without loss of generality, here, we assume that $a = 1$. Thus, the infinite geometric series becomes

$$1 + \rho + \rho^2 + \rho^3 + \cdots . \quad (1.2)$$

The partial sum of the first n terms of the infinite geometric series given by (1.2) is

$$S_n = \sum_{k=0}^{n-1} \rho^k = \frac{1 - \rho^n}{1 - \rho}. \quad (1.3)$$

Note that the formula given in (1.3) is also known as the geometric identity for finite number of terms. In this letter, we establish the identity given by (1.3) from a probabilistic point of view, which is provided in the next section.

*Email address (corresponding author): kayals@nitrrkl.ac.in, suchandan.kayal@gmail.com

2 Proof of (1.3)

To establish the sum formula for the finite geometric series, the following lemma is useful. Let X be an exponentially distributed random variable with probability density function

$$f_X(x|\lambda) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x > 0, \\ 0, & \text{otherwise,} \end{cases} \quad (2.1)$$

where $\lambda > 0$. If X has the density function given by (2.1), then for convenience, we denote $X \sim \text{Exp}(\lambda)$.

Lemma 2.1. *Let X and Y be two independent random variables such that $X \sim \text{Exp}(\lambda)$ and $Y \sim \text{Exp}(\mu)$, where $\lambda \neq \mu$. Further, let $U = X + Y$ and $V = X - Y$. Then,*

$$f_U(u|\lambda, \mu) = \begin{cases} \frac{\lambda\mu}{\lambda - \mu} (e^{-\mu u} - e^{-\lambda u}), & \text{if } u > 0, \\ 0, & \text{otherwise} \end{cases} \quad (2.2)$$

and

$$f_V(v|\lambda, \mu) = \begin{cases} \frac{\lambda\mu}{\lambda + \mu} e^{-\lambda v}, & \text{if } v > 0, \\ \frac{\lambda\mu}{\lambda + \mu} e^{\mu v}, & \text{otherwise.} \end{cases} \quad (2.3)$$

Proof. Consider the transformations $U = X + Y$ and $U_1 = X$. Then, the joint probability density function of U and U_1 is obtained as

$$f_{U,U_1}(u, u_1) = \lambda\mu e^{-(\lambda+\mu)u_1} e^{-\mu(u-u_1)}, \quad 0 < u < \infty, \quad 0 < u_1 < u. \quad (2.4)$$

Thus, the probability density function of U given by (2.2) can be obtained after integrating (2.4) with respect to u_1 . Similarly, the density function in (2.3) can be derived if we take the transformations $V = X - Y$ and $V_1 = X$. $\square$

Now, we present the proof of (1.3). Note that the k th order moments of X and Y about the origin are respectively given by $E(X^k) = k!/\lambda^k$ and $E(Y^k) = k!/\mu^k$, where $k = 0, 1, 2, \dots$ (see Rohatgi and Saleh, 2015). Further, using (2.2), it can be shown that

$$\begin{aligned} E(X + Y)^{n-1} &= \int_0^\infty u^{n-1} \left(\frac{\lambda\mu}{\lambda - \mu} \right) (e^{-\mu u} - e^{-\lambda u}) du \\ &= \frac{(n-1)!(\lambda\mu)}{\lambda - \mu} \left(\frac{1}{\mu^n} - \frac{1}{\lambda^n} \right). \end{aligned} \quad (2.5)$$

The binomial theorem in (2.5) yields

$$\begin{aligned}
E\left(\sum_{k=0}^{n-1} \binom{n-1}{k} X^k Y^{n-1-k}\right) &= \frac{(n-1)!(\lambda\mu)}{\lambda-\mu} \left(\frac{1}{\mu^n} - \frac{1}{\lambda^n}\right) \\
\Rightarrow \sum_{k=0}^{n-1} \binom{n-1}{k} E(X^k) E(Y^{n-1-k}) &= \frac{(n-1)!(\lambda\mu)}{\lambda-\mu} \left(\frac{1}{\mu^n} - \frac{1}{\lambda^n}\right) \\
\Rightarrow \sum_{k=0}^{n-1} \binom{n-1}{k} \frac{k!}{\lambda^k} \frac{(n-1-k)!}{\mu^{n-1-k}} &= \frac{(n-1)!(\lambda\mu)}{\lambda-\mu} \left(\frac{1}{\mu^n} - \frac{1}{\lambda^n}\right) \\
&\Rightarrow \sum_{k=0}^{n-1} \left(\frac{\mu}{\lambda}\right)^k = \frac{\lambda\mu^n}{\lambda-\mu} \left(\frac{1}{\mu^n} - \frac{1}{\lambda^n}\right) \\
&= \frac{1}{1-\frac{\mu}{\lambda}} \left(1 - \left(\frac{\mu}{\lambda}\right)^n\right) \tag{2.6}
\end{aligned}$$

Thus, the identity given by (1.3) is established when $\rho = \mu/\lambda$ is a strictly positive real number. To prove the identity when $\rho = \mu/\lambda < 0$, we consider the expectation of $(X-Y)^{n-1}$, where the probability density function of $V = X - Y$ is given by (2.3). Now,

$$E(X - Y)^{n-1} = \int_0^\infty u^{n-1} \frac{\lambda\mu}{\lambda+\mu} e^{-\lambda u} du + \int_{-\infty}^0 u^{n-1} \frac{\lambda\mu}{\lambda+\mu} e^{\mu u} du \tag{2.7}$$

$$= \frac{(n-1)!\lambda\mu}{\lambda+\mu} \left(\frac{1}{\lambda^n} + (-1)^{n-1} \frac{1}{\mu^n}\right). \tag{2.8}$$

Using similar arguments to (2.6), we obtain

$$\sum_{k=0}^{n-1} \left(-\frac{\mu}{\lambda}\right)^k = \frac{1 - \left(-\frac{\mu}{\lambda}\right)^n}{1 - \left(-\frac{\mu}{\lambda}\right)}. \tag{2.9}$$

Thus, the desired identity is proved when $\rho = -\mu/\lambda$ is strictly negative. Combining (2.6) and (2.9), the identity given by (1.3) is established for any nonzero real number ρ .

Remark 2.1. Taking limit $n \rightarrow \infty$ in (1.3), one can prove that $\sum_{k=0}^\infty \rho^k = \frac{1}{1-\rho}$ for $|\rho| < 1$.

References

Rohatgi, V. K. and Saleh, A. M. E. (2015). An introduction to probability and statistics, John Wiley & Sons.