

# EXISTENCE AND MULTIPLICITY OF POSITIVE SOLUTIONS FOR THE FRACTIONAL LAPLACIAN UNDER SUBCRITICAL OR CRITICAL GROWTH

SILVIA FRASSU, ANTONIO IANNIZZOTTO

ABSTRACT. We study a Dirichlet type problem for an equation involving the fractional Laplacian and a reaction term subject to either subcritical or critical growth conditions, depending on a positive parameter. Applying a critical point result of Bonanno, we prove existence of one or two positive solutions as soon as the parameter lies under a (explicitly determined) threshold. As an application, we find two positive solutions for a fractional Brezis-Nirenberg problem.

Version of April 7, 2020

## 1. INTRODUCTION

This paper is devoted to the following Dirichlet problem for a pseudo-differential equation of fractional order:

$$(1.1) \quad \begin{cases} (-\Delta)^s u = \lambda f(u) & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{in } \Omega^c. \end{cases}$$

Here  $s \in (0, 1)$ ,  $\Omega \subset \mathbb{R}^N$  ( $N > 2s$ ) is a bounded domain with  $C^{1,1}$  boundary, and the leading operator is the fractional Laplacian defined for all  $u \in \mathcal{S}(\mathbb{R}^N)$  by

$$(1.2) \quad (-\Delta)^s u(x) = 2 \text{P.V.} \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy.$$

The autonomous reaction  $f \in C(\mathbb{R})$  is assumed to be non-negative and dominated at infinity by a power of  $u$ , namely, for all  $t \in \mathbb{R}$

$$(1.3) \quad 0 \leq f(t) \leq a_0(1 + |t|^{q-1}) \quad (a_0 > 0, q \leq 2_s^*),$$

where  $2_s^* = 2N/(N - 2s)$  denotes the critical exponent for the fractional Sobolev space  $H^s(\mathbb{R}^N)$  (see [18]). Finally,  $\lambda > 0$  is a parameter.

Problem (1.1) admits a variational formulation by means of the energy functional

$$J_\lambda(u) = \frac{[u]_s^2}{2} - \lambda \int_\Omega F(u) dx,$$

where  $[\cdot]_s$  denotes the Gagliardo seminorm and  $F$  is the primitive of  $f$ , i.e., weak solutions of (1.1) coincide with critical points of  $J_\lambda$  in a convenient subspace of  $H^s(\mathbb{R}^N)$  (see Section 2 below for details). We note that, for  $\lambda = 1$ , problem (1.1) embraces the following Dirichlet problem with pure power nonlinearities:

$$(1.4) \quad \begin{cases} (-\Delta)^s u = \mu u^{p-1} + u^{q-1} & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{in } \Omega^c, \end{cases}$$

with  $1 < p < q \leq 2_s^*$  and  $\mu > 0$ .

For a general introduction to the fractional Laplacian we refer to [13–15, 18]. The study of (1.1) (or closely related problems) via variational methods started from the work of Servadei and Valdinoci [29, 30]. Here we distinguish between the *subcritical* ( $q < 2_s^*$  in (1.3)) and *critical* ( $q = 2_s^*$ ) cases. In the subcritical case, we mention for instance the contributions of [5, 16, 19–21, 24, 27, 32] and the monograph [25].

2010 *Mathematics Subject Classification.* 35R11, 35S15, 35A15.

*Key words and phrases.* Fractional Laplacian, Critical growth, Variational methods.

In the critical case, the main difficulty lies in the fact that  $J_\lambda$  does not satisfy the (usual in variational methods) Palais-Smale compactness condition. In particular, problem (1.4) with  $q = 2_s^*$  represents a fractional counterpart of the famous Brezis-Nirenberg problem [11]. Again, the first result in this direction is due to Servadei and Valdinoci [31] (see also [2, 3, 26]). Later, Barrios *et al.* [4] studied (1.4) with  $1 < p < 2 < q = 2_s^*$  (concave case) proving that, for  $\mu > 0$  small enough, such problem has at least two positive solutions  $u_\mu < w_\mu$ , employing both topological (sub-supersolutions) and variational methods, in a way that was first introduced in [1] for elliptic PDE's.

Our approach to problem (1.1) is purely variational, mainly based on a critical point theorem of Bonanno [6] and some of its consequences, presented in [7–9]. The main feature of such method is a strategy to find a local minimizer of a  $J_\lambda$ -type functional, which only requires a *local* Palais-Smale condition. Our results are the following:

- (a) In the subcritical case ( $q < 2_s^*$ ) we apply an abstract result of [7] and explicitly compute a real number  $\lambda^* > 0$  s.t. problem (1.1) admits at least two positive solutions  $u_\lambda, v_\lambda$  for all  $\lambda \in (0, \lambda^*)$ .
- (b) In the critical case ( $q = 2_s^*$ ) we first study a generalization of problem (1.4), explicitly determining a real number  $\mu^* > 0$  s.t. there exist at least one positive solution  $u_\mu$  for all  $\mu \in (0, \mu^*)$ . Then, we focus on (1.4) with  $q = 2_s^*$  and, applying the mountain pass theorem, we produce a second positive solution  $w_\mu > u_\mu$  for all  $\mu \in (0, \mu^*)$  (here we mainly follow [9]).

To our knowledge, this is the first application of the ideas of [6] in the field of fractional Laplacian equations. A noteworthy difference with respect to the classical elliptic case is the following: in this approach, it is essential to explicitly compute  $J_\lambda(\bar{u})$  at some Sobolev-type function  $\bar{u} : \Omega \rightarrow \mathbb{R}$ , which is usually chosen in such a way to have a piecewise constant  $|\nabla \bar{u}|$ . In the fractional framework, functions may have no gradient at all, and the computation of the Gagliardo seminorm is often prohibitive, so  $\bar{u}$  will be chosen as (a multiple of) the solution of a fractional torsion equation in a ball (see (2.3)).

We also remark that our main result in part (b) is formally equivalent to the main result of [4], but with two substantial differences: the first solution  $u_\mu$  is found as a local minimizer of  $J_\lambda$  (instead of being detected via sub-supersolutions, and *a posteriori* proved to be a minimizer), and moreover the interval  $(0, \mu^*)$  is *explicitly* determined (though possibly not optimal).

The paper has the following structure: in Section 2 we collect the necessary preliminaries; in Section 3 we develop part (a) of our study; in Sections 4 and 5 we focus on part (b).

**Notation:** Throughout the paper, for any  $A \subset \mathbb{R}^N$  we shall set  $A^c = \mathbb{R}^N \setminus A$ . By  $|A|$  we will denote either the  $N$ -dimensional Lebesgue measure or the  $(N - 1)$ -dimensional Hausdorff measure of  $A$ , which will be clear from the context. For any two measurable functions  $u, v$ ,  $u = v$  in  $A$  will stand for  $u(x) = v(x)$  for a.e.  $x \in A$  (and similar expressions). We will often write  $t^\nu = |t|^{\nu-1}t$  for  $t \in \mathbb{R}$ ,  $\nu > 1$ . For any  $t \in \mathbb{R}$  we set  $t^\pm = \max\{\pm t, 0\}$ . By  $B_r(x)$  we denote the open ball centered at  $x \in \mathbb{R}^N$  of radius  $r > 0$ . For all  $\nu \in [1, \infty]$ ,  $\|\cdot\|_\nu$  denotes the standard norm of  $L^\nu(\Omega)$  (or  $L^\nu(\mathbb{R}^N)$ , which will be clear from the context). Every function  $u$  defined in  $\Omega$  will be identified with its 0-extension to  $\mathbb{R}^N$ . Moreover,  $C$  will denote a positive constant (whose value may change line by line).

## 2. PRELIMINARIES

We begin by recalling some basic notions about fractional Sobolev spaces (for details we refer to [18]). We define the Gagliardo seminorm by setting for all measurable  $u : \mathbb{R}^N \rightarrow \mathbb{R}$

$$[u]_s = \left[ \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{(u(x) - u(y))^2}{|x - y|^{N+2s}} dx dy \right]^{\frac{1}{2}}.$$

Accordingly, we define the space

$$H^s(\mathbb{R}^N) = \{u \in L^2(\mathbb{R}^N) : [u]_s < \infty\}.$$

The embedding  $H^s(\mathbb{R}^N) \hookrightarrow L^{2_s^*}(\mathbb{R}^N)$  is continuous, and the fractional Talenti constant is given by the following lemma (see [17, Theorem 1.1] and [18, Proposition 3.6]):

**Lemma 2.1.** *We have*

$$T(N, s) = \max_{u \in H^s(\mathbb{R}^N) \setminus \{0\}} \frac{\|u\|_{2_s^*}}{[u]_s} = \frac{s^{\frac{1}{2}} \Gamma(\frac{N-2s}{2})^{\frac{1}{2}} \Gamma(N)^{\frac{s}{N}}}{2^{\frac{1}{2}} \pi^{\frac{N+2s}{4}} \Gamma(1-s)^{\frac{1}{2}} \Gamma(\frac{N}{2})^{\frac{s}{N}}} > 0,$$

the maximum being attained at the functions

$$u(x) = \frac{a}{(b + |x - x_0|^2)^{\frac{N-2s}{2}}} \quad (a, b > 0, x_0 \in \mathbb{R}^N).$$

Now we establish a variational formulation for (1.1), following [30] (see also [22]). Set

$$H_0^s(\Omega) = \{u \in H^s(\mathbb{R}^N) : u = 0 \text{ in } \Omega^c\},$$

a Hilbert space under the inner product

$$\langle u, v \rangle = \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{N+2s}} dx dy$$

and the corresponding norm  $\|u\| = [u]_s$  (see [30, Lemma 7]). The dual space of  $H_0^s(\Omega)$  is denoted  $H^{-s}(\Omega)$ . By Lemma 2.1 and Hölder's inequality, for any  $\nu \in [1, 2_s^*]$  the embedding  $H_0^s(\Omega) \hookrightarrow L^\nu(\Omega)$  is continuous and for all  $u \in H_0^s(\Omega)$  we have

$$(2.1) \quad \|u\|_\nu \leq T(N, s) |\Omega|^{\frac{2_s^* - \nu}{2_s^*}} \|u\|.$$

Further, the embedding is compact iff  $\nu < 2_s^*$  (see [30, Lemma 8]).

In order to deal with problem (1.1) variationally, we assume the following hypotheses on the reaction  $f$ :

- H<sub>0</sub>**  $f \in C(\mathbb{R})$ ,  $F(t) = \int_0^t f(\tau) d\tau$ , and
- (i)  $f(t) \geq 0$  for all  $t \in \mathbb{R}$ ;
  - (ii)  $f(t) \leq a_0(1 + |t|^{2_s^* - 1})$  for all  $t \in \mathbb{R}$  ( $a_0 > 0$ ).

We set for all  $u \in H_0^s(\Omega)$ ,  $\lambda > 0$

$$\Phi(u) = \frac{\|u\|^2}{2}, \quad \Psi(u) = \int_\Omega F(u) dx, \quad J_\lambda(u) = \Phi(u) - \lambda\Psi(u)$$

( $\Psi$  is well defined by virtue of hypothesis **H<sub>0</sub>** (i) (ii)). Then  $\Phi, \Psi, J_\lambda \in C^1(H_0^s(\Omega))$  with

$$\langle J'_\lambda(u), \varphi \rangle = \langle u, \varphi \rangle - \lambda \int_\Omega f(u)\varphi dx$$

for all  $u, \varphi \in H_0^s(\Omega)$ . We say that  $u$  is a (weak) solution of problem (1.1) if  $J'_\lambda(u) = 0$  in  $H^{-s}(\Omega)$ , that is, for all  $\varphi \in H_0^s(\Omega)$  we have

$$(2.2) \quad \langle u, \varphi \rangle = \lambda \int_\Omega f(u)\varphi dx.$$

The regularity theory for fractional Dirichlet problems was essentially developed in [28] (see also [4, 22]). While smooth in  $\Omega$ , solutions are in general singular on  $\partial\Omega$ , so the best global regularity we can expect is weighted Hölder continuity, in the following sense. Set for all  $x \in \overline{\Omega}$

$$d_\Omega(x) = \text{dist}(x, \Omega^c),$$

then define the spaces

$$C_s^0(\overline{\Omega}) = \left\{ u \in C^0(\overline{\Omega}) : \frac{u}{d_\Omega^s} \in C^0(\overline{\Omega}) \right\}, \quad \|u\|_{0,s} = \left\| \frac{u}{d_\Omega^s} \right\|_\infty,$$

and for any  $\alpha \in (0, 1)$

$$C_s^\alpha(\overline{\Omega}) = \left\{ u \in C^0(\overline{\Omega}) : \frac{u}{d_\Omega^s} \in C^\alpha(\overline{\Omega}) \right\}, \quad \|u\|_{\alpha,s} = \|u\|_{0,s} + \sup_{x \neq y} \frac{|u(x)/d_\Omega^s(x) - u(y)/d_\Omega^s(y)|}{|x - y|^\alpha}.$$

The positive order cone of  $C_s^0(\overline{\Omega})$  has a nonempty interior given by

$$\text{int}(C_s^0(\overline{\Omega})_+) = \left\{ u \in C_s^0(\overline{\Omega}) : \frac{u}{d_\Omega^s} > 0 \text{ in } \overline{\Omega} \right\}.$$

For the reader's convenience we recall from [22, Theorems 2.3, 3.2 and Lemma 2.7] the main properties of weak solutions:

**Proposition 2.2.** *Let **H<sub>0</sub>** hold,  $u \in H_0^s(\Omega)$  be a weak solution of (1.1). Then:*

- (i) (a priori bound)  $u \in L^\infty(\Omega)$ ;
- (ii) (regularity)  $u \in C_s^\alpha(\overline{\Omega})$  with  $\alpha \in (0, s]$  depending only on  $s$  and  $\Omega$ ;

(iii) (Hopf's lemma) if  $u \neq 0$ , then  $u \in \text{int}(C_s^0(\overline{\Omega})_+)$ .

By Proposition 2.2 (iii) we see that, whenever  $u \in H_0^s(\Omega) \setminus \{0\}$  satisfies (2.2), then in particular  $u > 0$  in  $\Omega$ . Moreover, assuming further that  $f$  is locally Lipschitz in  $\mathbb{R}$ , from [28, Corollary 1.6] we deduce that  $u \in C^\beta(\Omega)$  for any  $\beta \in [1, 1 + 2s)$ , which along with Proposition 2.2 (ii) implies that for all  $x \in \mathbb{R}^N$  the mapping

$$x \mapsto \frac{u(x) - u(y)}{|x - y|^{N+2s}}$$

lies in  $L^1(\mathbb{R}^N)$ . Then, testing (2.2) with any  $\varphi \in C_c^\infty(\Omega)$  and applying (1.2), we have

$$\int_{\Omega} (-\Delta)^s u \varphi \, dx = \langle u, \varphi \rangle = \int_{\Omega} f(u) \varphi \, dx,$$

i.e.,  $u$  solves (1.1) pointwisely.

We also recall the following result, relating the local minimizers of the energy functional  $J_\lambda$  in  $H_0^s(\Omega)$  and in  $C_s^0(\overline{\Omega})$ , respectively (see [22, Theorem 1.1], [4, Proposition 2.5], and [23, Theorem 1.1] for a nonlinear extension), namely an analog for the fractional case of the main result of [12]:

**Proposition 2.3.** *Let  $\mathbf{H}_0$  hold,  $u \in H_0^s(\Omega)$ . Then, the following are equivalent:*

- (i) *there exists  $\rho > 0$  s.t.  $J_\lambda(u + v) \geq J_\lambda(u)$  for all  $v \in H_0^s(\Omega) \cap C_s^0(\overline{\Omega})$ ,  $\|v\|_{0,s} \leq \rho$ ;*
- (ii) *there exists  $\sigma > 0$  s.t.  $J_\lambda(u + v) \geq J_\lambda(u)$  for all  $v \in H_0^s(\Omega)$ ,  $\|v\| \leq \sigma$ .*

As pointed out in the Introduction, we will make use of the following fractional torsion equation on a ball:

$$(2.3) \quad \begin{cases} (-\Delta)^s u_R = 1 & \text{in } B_R(x_0) \\ u_R = 0 & \text{in } B_R(x_0)^c, \end{cases}$$

where  $x_0 \in \mathbb{R}^N$ ,  $R > 0$ . The solution of (2.3) (defined as in (2.2)) is unique, given by

$$u_R(x) = A(N, s)(R^2 - |x - x_0|^2)_+^s, \quad A(N, s) = \frac{s\Gamma(\frac{N}{2})}{2\pi^{\frac{N}{2}}\Gamma(1+s)\Gamma(1-s)}$$

(see [13, p. 33] or [28, equation (1.4)]). This simple example is popular in fractional regularity theory, as it shows that solutions of Dirichlet problems may be singular at the boundary. For future use we compute some norms of  $u_R$ :

**Lemma 2.4.** *For all  $x_0 \in \mathbb{R}^N$ ,  $R > 0$  we have*

- (i)  $\|u_R\|_\nu = A(N, s) \left[ \frac{\pi^{\frac{N}{2}} \Gamma(1 + \nu s) R^{N+2\nu s}}{\Gamma(\frac{N+2\nu s+2}{2})} \right]^{\frac{1}{\nu}}$  for all  $\nu \geq 1$ ;
- (ii)  $[u_R] = \left[ \frac{s\Gamma(\frac{N}{2}) R^{N+2s}}{2\Gamma(1-s)\Gamma(\frac{N+2s+2}{2})} \right]^{\frac{1}{2}}$ .

*Proof.* First we recall the well-known formulas

$$|\partial B_1(0)| = \frac{2\pi^{\frac{N}{2}}}{\Gamma(\frac{N}{2})}, \quad \int_0^1 (1 - \rho^2)^\alpha \rho^{N-1} \, d\rho = \frac{\Gamma(\frac{N}{2})\Gamma(1 + \alpha)}{2\Gamma(\frac{N+2\alpha+2}{2})} \quad (\alpha > 0),$$

then for all  $\nu \geq 1$  we compute

$$\begin{aligned} \int_{B_R(x_0)} u_R^\nu(x) \, dx &= A(N, s)^\nu \int_{B_R(x_0)} (R^2 - |x - x_0|^2)^{\nu s} \, dx \\ &= A(N, s)^\nu R^{N+2\nu s} |\partial B_1(0)| \int_0^1 (1 - \rho^2)^{\nu s} \rho^{N-1} \, d\rho \\ &= A(N, s)^\nu \frac{\pi^{\frac{N}{2}} \Gamma(1 + \nu s) R^{N+2\nu s}}{\Gamma(\frac{N+2\nu s+2}{2})}, \end{aligned}$$

which implies (i). Further, testing (2.3) with  $u_R \in H_0^s(B_R(x_0))$  and applying (i) with  $\nu = 1$ , we have

$$\begin{aligned} [u_R]_s^2 &= \int_{B_R(x_0)} u_R dx \\ &= A(N, s) \frac{\pi^{\frac{N}{2}} \Gamma(1+s) R^{N+2s}}{\Gamma(\frac{N+2s+2}{2})} \\ &= \frac{s \Gamma(\frac{N}{2}) R^{N+2s}}{2 \Gamma(1-s) \Gamma(\frac{N+2s+2}{2})}, \end{aligned}$$

which gives (ii).  $\square$

**Remark 2.5.** We note that some results here are affected by the definition (1.2), which is the same adopted in [4]. Other works on the subject, for instance [13, 17, 18], define the fractional Laplacian as

$$(-\Delta)^s u(x) = C(N, s) \text{P.V.} \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x-y|^{N+2s}} dy, \quad C(N, s) = \frac{2^{2s} s \Gamma(\frac{N+2s}{2})}{\pi^{\frac{N}{2}} \Gamma(1-s)} > 0,$$

where the multiplicative constant is required to equivalently define  $(-\Delta)^s$  by means of the Fourier transform. In this paper, explicit constants are one of the main issues, so we decide to follow the standard of [4] in order to easily compare similar results.

### 3. TWO POSITIVE SOLUTIONS UNDER SUBCRITICAL GROWTH

In this section, following [7] as a model, we study (1.1) under the following hypotheses:

- H<sub>1</sub>**  $f \in C(\mathbb{R})$ ,  $F(t) = \int_0^t f(\tau) d\tau$  satisfy
- (i)  $f(t) \geq 0$  for all  $t \in \mathbb{R}$ ;
  - (ii)  $f(t) \leq a_p |t|^{p-1} + a_q |t|^{q-1}$  for all  $t \in \mathbb{R}$  ( $1 \leq p < 2 < q < 2_s^*$ ,  $a_p, a_q > 0$ );
  - (iii)  $\lim_{t \rightarrow 0^+} \frac{F(t)}{t^2} = \infty$
  - (iv)  $0 < \rho F(t) \leq f(t)t$  for all  $t \geq M$  ( $\rho > 2$ ,  $M > 0$ ).

Hypotheses **H<sub>1</sub>** conjure for  $f$  a subcritical, superlinear growth at infinity, as well as a sublinear growth near the origin, while (iv) is an Ambrosetti-Rabinowitz condition.

First, we recall the classical Palais-Smale condition at level  $c \in \mathbb{R}$ , for a functional  $J \in C^1(X)$  on a Banach space  $X$ :

$(PS)_c$  Every sequence  $(u_n)$  in  $X$ , s.t.  $J(u_n) \rightarrow c$  and  $J'(u_n) \rightarrow 0$  in  $X^*$ , has a convergent subsequence.

We say that  $J$  satisfies  $(PS)$ , if  $J$  satisfies  $(PS)_c$  for any  $c \in \mathbb{R}$ .

We will apply the following abstract result, slightly rephrased from [7, Theorem 2.1]:

**Theorem A.** Let  $X$  be a Banach space,  $\Phi, \Psi \in C^1(X)$ ,  $J_\lambda = \Phi - \lambda \Psi$  ( $\lambda > 0$ ),  $r \in \mathbb{R}$ ,  $\bar{u} \in X$  satisfy

- (A<sub>1</sub>)  $\inf_{u \in X} \Phi(u) = \Phi(0) = \Psi(0) = 0$ ;
- (A<sub>2</sub>)  $0 < \Phi(\bar{u}) < r$ ;
- (A<sub>3</sub>)  $\sup_{\Phi(u) \leq r} \frac{\Psi(u)}{r} < \frac{\Psi(\bar{u})}{\Phi(\bar{u})}$ ;
- (A<sub>4</sub>)  $\inf_{u \in X} J_\lambda(u) = -\infty$  for all  $\lambda \in I_r = \left( \frac{\Phi(\bar{u})}{\Psi(\bar{u})}, \left[ \sup_{\Phi(u) \leq r} \frac{\Psi(u)}{r} \right]^{-1} \right)$ .

Then, for all  $\lambda \in I_r$  for which  $J_\lambda$  satisfies  $(PS)$ , there exist  $u_\lambda, v_\lambda \in X$  s.t.

$$J'_\lambda(u_\lambda) = J'_\lambda(v_\lambda) = 0, \quad J_\lambda(u_\lambda) < 0 < J_\lambda(v_\lambda).$$

Let  $T(N, s) > 0$  be defined by Lemma 2.1, set

$$(3.1) \quad \lambda^* = \frac{1}{2T(N, s)^2 |\Omega|^{\frac{2_s^*-2}{2_s^*}}} \left( \frac{a_p}{p} \right)^{\frac{2-p}{q-p}} \left( \frac{a_q}{q} \right)^{\frac{p-2}{q-p}} \left( \frac{2-p}{q-2} \right)^{\frac{2-p}{q-p}} \frac{q-2}{q-p} > 0.$$

We have the following multiplicity result:

**Theorem 3.1.** *Let  $\mathbf{H}_1$  hold,  $\lambda^* > 0$  be defined by (3.1). Then, for all  $\lambda \in (0, \lambda^*)$ , (1.1) has at least two solutions  $u_\lambda, v_\lambda \in \text{int}(C_s^0(\bar{\Omega})_+)$ .*

*Proof.* Without loss of generality we may assume  $f(t) = 0$  for all  $t \leq 0$ . We are going to apply Theorem A. Set  $X = H_0^s(\Omega)$  and define  $\Phi, \Psi, J_\lambda$  as in Section 2, then clearly  $\Phi, \Psi \in C^1(H_0^s(\Omega))$  and

$$\inf_{u \in H_0^s(\Omega)} \Phi(u) = \Phi(0) = \Psi(0) = 0,$$

hence hypothesis (A<sub>1</sub>) holds. Set

$$(3.2) \quad r = \frac{|\Omega|^{\frac{2}{2s}}}{2T(N, s)^2} \left[ \frac{a_p q (2-p)}{a_q p (q-2)} \right]^{\frac{2}{q-p}} > 0.$$

For all  $u \in H_0^s(\Omega)$ ,  $\Phi(u) \leq r$ , we have  $\|u\| \leq (2r)^{\frac{1}{2}}$ . So, by hypotheses  $\mathbf{H}_1$  (i) (ii), along with (2.1), (3.1) and (3.2), we obtain

$$\begin{aligned} \frac{\Psi(u)}{r} &\leq \frac{a_p}{pr} \|u\|_p^p + \frac{a_q}{qr} \|u\|_q^q \\ &\leq \frac{a_p}{pr} T(N, s)^p |\Omega|^{\frac{2s-p}{2s}} (2r)^{\frac{p}{2}} + \frac{a_q}{qr} T(N, s)^q |\Omega|^{\frac{2s-q}{2s}} (2r)^{\frac{q}{2}} \\ &= 2T(N, s)^2 |\Omega|^{\frac{2s-2}{2s}} \left( \frac{a_p}{p} \right)^{\frac{q-2}{q-p}} \left( \frac{a_q}{q} \right)^{\frac{2-p}{q-p}} \left( \frac{2-p}{q-2} \right)^{\frac{p-2}{q-p}} \\ &\quad + 2T(N, s)^2 |\Omega|^{\frac{2s-2}{2s}} \left( \frac{a_p}{p} \right)^{\frac{q-2}{q-p}} \left( \frac{a_q}{q} \right)^{\frac{2-p}{q-p}} \left( \frac{2-p}{q-2} \right)^{\frac{q-2}{q-p}} \\ &= 2T(N, s)^2 |\Omega|^{\frac{2s-2}{2s}} \left( \frac{a_p}{p} \right)^{\frac{q-2}{q-p}} \left( \frac{a_q}{q} \right)^{\frac{2-p}{q-p}} \left( \frac{2-p}{q-2} \right)^{\frac{p-2}{q-p}} \frac{q-p}{q-2} = \frac{1}{\lambda^*}. \end{aligned}$$

Summarizing,

$$(3.3) \quad \sup_{\Phi(u) \leq r} \frac{\Psi(u)}{r} \leq \frac{1}{\lambda^*}.$$

Now fix  $\lambda \in (0, \lambda^*)$ . Since  $\partial\Omega$  is  $C^{1,1}$ , we can find  $x_0 \in \mathbb{R}^N$ ,  $R > 0$  largest s.t.  $B_R(x_0) \subseteq \Omega$ . Let  $K > 0$  be s.t.

$$(3.4) \quad K \frac{s\Gamma(\frac{N}{2})\Gamma(1+2s)\Gamma(\frac{N+2s+2}{2})R^{2s}}{\pi^{\frac{N}{2}}\Gamma(1+s)^2\Gamma(1-s)\Gamma(\frac{N+4s+2}{2})} > \frac{1}{\lambda}.$$

By  $\mathbf{H}_1$  (iii), we can find  $\varepsilon > 0$  s.t. for all  $t \in [0, \varepsilon]$

$$(3.5) \quad F(t) \geq Kt^2.$$

Finally, fix

$$(3.6) \quad 0 < \delta < \min \left\{ \left[ \frac{4\Gamma(1-s)\Gamma(\frac{N+2s+2}{2})r}{s\Gamma(\frac{N}{2})R^{N+2s}} \right]^{\frac{1}{2}}, \frac{2\pi^{\frac{N}{2}}\Gamma(1+s)\Gamma(1-s)\varepsilon}{s\Gamma(\frac{N}{2})R^{2s}} \right\}.$$

Now let  $u_R$  be the solution of (2.3) in  $B_R(x_0)$ , and set  $\bar{u} = \delta u_R \in H_0^s(\Omega)$ . Then we have by Lemma 2.4 (ii) and (3.6)

$$\Phi(\bar{u}) = \frac{s\Gamma(\frac{N}{2})R^{N+2s}\delta^2}{4\Gamma(1-s)\Gamma(\frac{N+2s+2}{2})} < r,$$

which implies (A<sub>2</sub>). Besides, by (3.6) we have for all  $x \in \Omega$

$$0 \leq \bar{u}(x) \leq \frac{s\Gamma(\frac{N}{2})R^{2s}\delta}{2\pi^{\frac{N}{2}}\Gamma(1+s)\Gamma(1-s)} < \varepsilon,$$

hence by (3.5) and Lemma 2.4 (i)

$$\Psi(\bar{u}) \geq \int_{\Omega} K\bar{u}^2 dx = K\delta^2 \|u_R\|_2^2 = K \frac{s^2\Gamma(\frac{N}{2})^2\Gamma(1+2s)R^{N+4s}}{4\pi^{\frac{N}{2}}\Gamma(1+s)^2\Gamma(1-s)^2\Gamma(\frac{N+4s+2}{2})} \delta^2.$$

The relations above and (3.4) imply

$$\frac{\Psi(\bar{u})}{\Phi(\bar{u})} \geq K \frac{s\Gamma(\frac{N}{2})\Gamma(1+2s)\Gamma(\frac{N+2s+2}{2})R^{2s}}{\pi^{\frac{N}{2}}\Gamma(1+s)^2\Gamma(1-s)\Gamma(\frac{N+4s+2}{2})} > \frac{1}{\lambda}.$$

Recalling that  $\lambda < \lambda^*$ , by (3.3) we have

$$\sup_{\Phi(u) \leq r} \frac{\Psi(u)}{r} < \frac{1}{\lambda} < \frac{\Psi(\bar{u})}{\Phi(\bar{u})},$$

which yields at once (A<sub>3</sub>) and  $\lambda \in I_r$ . By **H**<sub>1</sub> (iv) we can find  $C > 0$  s.t. for all  $t \geq M$

$$(3.7) \quad F(t) \geq Ct^\rho.$$

Now pick  $w \in C_c^\infty(\Omega) \setminus \{0\}$ . By (3.7), and recalling that  $F(t) \geq 0$  for all  $t \in \mathbb{R}$ , we have for all  $\tau > 0$

$$\begin{aligned} J_\lambda(\tau w) &\leq \frac{\|w\|^2}{2}\tau^2 - \lambda \int_{\{w \leq M/\tau\}} F(\tau w) dx - \lambda \int_{\{w > M/\tau\}} C(\tau w)^\rho dx \\ &\leq \frac{\|w\|^2}{2}\tau^2 - \lambda \int_\Omega C(\tau w)^\rho dx + \lambda \int_{\{w \leq M/\tau\}} CM^\rho dx \\ &\leq \frac{\|w\|^2}{2}\tau^2 - \lambda C\|w\|_\infty^\rho |\Omega| \tau^\rho + \lambda CM^\rho |\Omega| \end{aligned}$$

and the latter tends to  $-\infty$  as  $\tau \rightarrow \infty$  (since  $\rho > 2$ ). So we see that (A<sub>4</sub>) holds as well.

Finally, we prove that  $J_\lambda$  satisfies (PS). Let  $(u_n)$  be a sequence in  $H_0^s(\Omega)$  s.t.  $|J_\lambda(u_n)| \leq C$ ,  $J'_\lambda(u_n) \rightarrow 0$  in  $H^{-s}(\Omega)$ . Then, for all  $n \in \mathbb{N}$  we have

$$(3.8) \quad \frac{\|u_n\|^2}{2} - \lambda \int_\Omega F(u_n) dx \leq C$$

and for all  $\varphi \in H_0^s(\Omega)$

$$(3.9) \quad \left| \langle u_n, \varphi \rangle - \lambda \int_\Omega f(u_n)\varphi dx \right| \leq \|J'_\lambda(u_n)\| \|\varphi\|$$

Multiplying (3.8) by  $\rho > 2$  (as in **H**<sub>1</sub> (iv)), testing (3.9) with  $u_n$ , and subtracting,

$$\begin{aligned} \frac{\rho-2}{2}\|u_n\|^2 &\leq \lambda \int_\Omega (\rho F(u_n) - f(u_n)u_n) dx + \|J'_\lambda(u_n)\| \|u_n\| + C \\ &\leq \lambda \int_{\{0 \leq u_n \leq M\}} C(|u_n|^\rho + |u_n|^q) dx + \|J'_\lambda(u_n)\| \|u_n\| + C \\ &\leq \lambda C(M^\rho + M^q)|\Omega| + \|J'_\lambda(u_n)\| \|u_n\| + C. \end{aligned}$$

So  $(u_n)$  is bounded in  $H_0^s(\Omega)$ . Passing to a subsequence, we have  $u_n \rightharpoonup u$  in  $H_0^s(\Omega)$ ,  $u_n \rightarrow u$  in  $L^p(\Omega)$ ,  $L^q(\Omega)$ , and  $u_n(x) \rightarrow u(x)$  for a.e.  $x \in \Omega$ . Testing (3.9) this time with  $u_n - u \in H_0^s(\Omega)$ , we have for all  $n \in \mathbb{N}$

$$\begin{aligned} \|u_n - u\|^2 &\leq \langle u, u_n - u \rangle + \lambda \int_\Omega (a_p|u_n|^{p-1} + a_q|u_n|^{q-1})|u_n - u| dx + \|J'_\lambda(u_n)\| \|u_n - u\| \\ &\leq \langle u, u_n - u \rangle + \lambda(a_p\|u_n\|_p^{p-1}\|u_n - u\|_p + a_q\|u_n\|_q^{q-1}\|u_n - u\|_q) + \|J'_\lambda(u_n)\| \|u_n - u\|, \end{aligned}$$

(where we used **H**<sub>1</sub> (ii) and Hölder's inequality), and the latter tends to 0 as  $n \rightarrow \infty$ . So,  $u_n \rightarrow u$  in  $H_0^s(\Omega)$ . (Note that we actually proved that  $J_\lambda$  is unbounded from below and satisfies (PS) for all  $\lambda > 0$ .)

By Theorem A, there exist  $u_\lambda, v_\lambda \in H_0^s(\Omega)$  s.t.

$$J'_\lambda(u_\lambda) = J'_\lambda(v_\lambda) = 0, \quad J_\lambda(u_\lambda) < 0 < J_\lambda(v_\lambda).$$

Therefore,  $u_\lambda, v_\lambda \neq 0$  solve (1.1). By **H**<sub>0</sub> (i) and Proposition 2.2, finally, we have  $u_\lambda, v_\lambda \in \text{int}(C_s^0(\bar{\Omega})_+)$ .  $\square$

We focus now on problem (1.4), with  $1 < p < 2 < q < 2_s^*$  (subcritical case) and  $\mu > 0$ . Set

$$(3.10) \quad \mu^* = \left[ 2T(N, s)^2 |\Omega|^{\frac{2_s^* - 2}{2s}} \right]^{\frac{p-q}{q-2}} p q^{\frac{2-p}{q-2}} \left( \frac{2-p}{q-2} \right)^{\frac{2-p}{q-2}} \left( \frac{q-2}{q-p} \right)^{\frac{q-p}{q-2}} > 0.$$

We have the following multiplicity result:

**Corollary 3.2.** *Let  $1 < p < 2 < q < 2_s^*$ ,  $\mu^* > 0$  be defined by (3.10). Then, for all  $\mu \in (0, \mu^*)$  (1.4) has at least two solutions  $u_\mu, v_\mu \in \text{int}(C_s^0(\overline{\Omega})_+)$ .*

*Proof.* Set for all  $t \in \mathbb{R}$ ,  $\mu \in (0, \mu^*)$

$$f(t) = \mu(t^+)^{p-1} + (t^+)^{q-1}.$$

Then  $f$  satisfies  $\mathbf{H}_1$  with  $a_p = \mu$ ,  $a_q = 1$ , and any  $\rho \in (2, q)$ . In view of (3.10), here (3.1) rephrases as

$$\lambda^* = \frac{1}{2T(N, s)^2 |\Omega|^{\frac{2_s^*-2}{2_s^*}}} p^{\frac{q-2}{q-p}} q^{\frac{2-p}{q-p}} \left( \frac{2-p}{q-2} \right)^{\frac{2-p}{q-p}} \frac{q-2}{q-p} \mu^{\frac{2-q}{q-p}} > 1.$$

Hence we can apply Theorem 3.1 with  $\lambda = 1$  and find  $u_\mu, v_\mu \in \text{int}(C_s^0(\overline{\Omega})_+)$  solutions to (1.4).  $\square$

We present an example for Corollary 3.2:

**Example 3.3.** Set  $s = \frac{1}{2}$ ,  $p = \frac{3}{2}$ ,  $q = 3$ ,  $N = 2$  and

$$\Omega = \left\{ (x, y) \in \mathbb{R}^2 : \frac{x^2}{4} + \frac{y^2}{9} \leq 1 \right\}.$$

Then we have  $2_{1/2}^* = 4 > 3$ ,  $|\Omega| = 6\pi$ , while Lemma 2.1 gives

$$T\left(2, \frac{1}{2}\right) = \frac{\left(\frac{1}{2}\right)^{\frac{1}{2}} \Gamma\left(\frac{1}{2}\right)^{\frac{1}{2}} \Gamma(2)^{\frac{1}{4}}}{2^{\frac{1}{2}} \pi^{\frac{3}{4}} \Gamma\left(\frac{1}{2}\right)^{\frac{1}{2}} \Gamma(1)^{\frac{1}{4}}} = \frac{1}{2\pi^{\frac{3}{4}}}.$$

Therefore, (3.10) becomes

$$\mu^* = \left[ 2 \left( \frac{1}{2\pi^{\frac{3}{4}}} \right)^2 (6\pi)^{\frac{1}{2}} \right]^{-\frac{3}{2}} 3^{\frac{1}{2}} \left( \frac{1}{2} \right)^{\frac{1}{2}} \left( \frac{2}{3} \right)^{\frac{3}{2}} = \frac{2^{\frac{3}{4}} \pi^{\frac{3}{2}}}{3^{\frac{3}{4}}}.$$

By Corollary 3.2, for all  $\mu \in (0, \mu^*)$  (1.4) has at least two positive solutions.

#### 4. ONE POSITIVE SOLUTION UNDER CRITICAL GROWTH

In this section, we study the following slight generalization of problem (1.4):

$$(4.1) \quad \begin{cases} (-\Delta)^s u = \mu g(u) + u^{2_s^*-1} & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{in } \Omega^c, \end{cases}$$

with  $\mu > 0$  and assuming the following hypotheses on  $g$ :

$\mathbf{H}_2$   $g \in C(\mathbb{R})$ ,  $G(t) = \int_0^t g(\tau) d\tau$  satisfy

- (i)  $g(t) \geq 0$  for all  $t \in \mathbb{R}$ ;
- (ii)  $g(t) \leq a_p |t|^{p-1}$  for all  $t \in \mathbb{R}$  ( $p \in (1, 2_s^*)$ ,  $a_p > 0$ );
- (iii)  $\lim_{t \rightarrow 0^+} \frac{G(t)}{t^2} = \infty$ .

Note that, due to hypothesis  $\mathbf{H}_1$  (iii), problem (4.1) reduces to (1.4) with  $g(t) = t^{p-1}$  only for  $p \in (1, 2)$  (concave case). Although, the results of this section also embrace the case  $p \in [2, 2_s^*)$  (linear/convex case).

Due to the presence of the critical term  $u^{2_s^*-1}$  in (4.1), we cannot apply Theorem A, as the associated energy functional does not satisfy (PS) in general. So we introduce the following local Palais-Smale condition for functionals of the type  $J_\lambda = \Phi - \lambda\Psi$ , with  $\Phi, \Psi \in C^1(X)$ ,  $\lambda > 0$ , defined on a Banach space  $X$ , and  $r > 0$ :

(PS)<sup>r</sup> Every sequence  $(u_n)$  in  $X$ , s.t.  $(J_\lambda(u_n))$  is bounded in  $\mathbb{R}$ ,  $J'(u_n) \rightarrow 0$  in  $X^*$ , and  $\Phi(u_n) \leq r$  for all  $n \in \mathbb{N}$ , has a convergent subsequence.

In this case, our main tool is the following local minimum result, slightly rephrased from [9, Theorem 3.3]:

**Theorem B.** *Let  $X$  be a Banach space,  $\Phi, \Psi \in C^1(X)$ ,  $J_\lambda = \Phi - \lambda\Psi$  ( $\lambda > 0$ ),  $r \in \mathbb{R}$ ;  $\bar{u} \in X$  satisfy*

- (B<sub>1</sub>)  $\inf_{u \in X} \Phi(u) = \Phi(0) = \Psi(0) = 0$ ;
- (B<sub>2</sub>)  $0 < \Phi(\bar{u}) < r$ ;
- (B<sub>3</sub>)  $\sup_{\Phi(u) \leq r} \frac{\Psi(u)}{r} < \frac{\Psi(\bar{u})}{\Phi(\bar{u})}$ .

Let

$$I_r = \left( \frac{\Phi(\bar{u})}{\Psi(\bar{u})}, \left[ \sup_{\Phi(u) \leq r} \frac{\Psi(u)}{r} \right]^{-1} \right).$$

Then, for all  $\lambda \in I_r$  for which  $J_\lambda$  satisfies  $(PS)^r$ , there exists  $u_\lambda \in X$  s.t.

$$0 < \Phi(u_\lambda) < r, \quad J_\lambda(u_\lambda) = \min_{0 < \Phi(u) < r} J_\lambda(u).$$

Set for all  $\mu > 0, t \in \mathbb{R}$

$$f(t) = \mu g(t) + (t^+)^{2^*-1}, \quad F(t) = \int_0^t f(\tau) d\tau,$$

then define  $\Phi, \Psi \in C^1(H_0^s(\Omega))$  as in Section 2. Further, for all  $\lambda > 0$  set  $J_\lambda = \Phi - \lambda\Psi$ . Set for all  $r, \mu > 0$

$$(4.2) \quad \lambda_r^* = \min \left\{ \left[ \frac{2^{\frac{2^*}{2}} T(N, s) 2_s^* r^{\frac{2^*-2}{2}}}{2_s^*} + \mu \frac{2^{\frac{p}{2}} a_p T(N, s)^p |\Omega|^{\frac{2^*-p}{2_s^*}} r^{\frac{p-2}{2}}}{p} \right]^{-1}, \frac{1}{T(N, s) 2_s^*} \left[ \frac{s}{2Nr} \right]^{\frac{2_s^*}{N-2_s^*}} \right\}$$

We prove now that  $J_\lambda$  satisfies  $(PS)^r$  for all  $r > 0$  and all  $\lambda > 0$  small enough:

**Lemma 4.1.** *Let  $r, \mu > 0, \lambda_r^* > 0$  be defined by (4.2). Then  $J_\lambda$  satisfies  $(PS)^r$  for all  $\lambda \in (0, \lambda_r^*)$ .*

*Proof.* Let  $(u_n)$  be a sequence in  $H_0^s(\Omega)$  s.t.  $(J_\lambda(u_n))$  is bounded,  $J'_\lambda(u_n) \rightarrow 0$  in  $H^{-s}(\Omega)$ , and  $\Phi(u_n) \leq r$  for all  $n \in \mathbb{N}$ . Then  $(u_n)$  is bounded in  $H_0^s(\Omega)$ , hence in  $L^{2_s^*}(\Omega)$  (Lemma 2.1). Passing to a subsequence we have  $u_n \rightharpoonup u$  in  $H_0^s(\Omega)$ ,  $L^{2_s^*}(\Omega)$ ,  $u_n \rightarrow u$  in  $L^p(\Omega)$ ,  $u_n(x) \rightarrow u(x)$  for a.e.  $x \in \Omega$ , and  $J_\lambda(u_n) \rightarrow c$ .

First we see that

$$(4.3) \quad J'_\lambda(u) = 0.$$

Indeed, since  $(u_n^{2_s^*-1})$  is bounded in  $L^{(2_s^*)'}(\Omega)$ , up to a further subsequence we have  $u_n^{2_s^*-1} \rightharpoonup u^{2_s^*-1}$  in  $L^{(2_s^*)'}(\Omega)$ , while by **H**<sub>2</sub> (i) (ii) we have  $g(u_n) \rightarrow g(u)$  in  $L^{p'}(\Omega)$ . So, for all  $\varphi \in H_0^s(\Omega)$  we have

$$\begin{aligned} \langle J'_\lambda(u_n), \varphi \rangle &= \langle u_n, \varphi \rangle - \lambda \int_\Omega u_n^{2_s^*-1} \varphi dx - \lambda \mu \int_\Omega g(u_n) \varphi dx \\ &\rightarrow \langle u, \varphi \rangle - \lambda \int_\Omega u^{2_s^*-1} \varphi dx - \lambda \mu \int_\Omega g(u) \varphi dx = \langle J'_\lambda(u), \varphi \rangle, \end{aligned}$$

which along with  $J'_\lambda(u_n) \rightarrow 0$  gives (4.3). Besides,

$$(4.4) \quad J_\lambda(u) > -r.$$

Indeed, since  $u_n \rightharpoonup u$  in  $H_0^s(\Omega)$  and  $\Phi$  is convex, we have  $\Phi(u) \leq r$ , i.e.,  $\|u\| \leq (2r)^{\frac{1}{2}}$ . So using Lemma 2.1, (2.1) with  $\nu = p$ , (4.2), and  $\lambda < \lambda_r^*$ , we have

$$\begin{aligned} J_\lambda(u) &\geq -\lambda\Psi(u) \\ &\geq -\frac{\lambda}{2_s^*} \|u\|_{2_s^*}^{2_s^*} - \frac{\lambda\mu a_p}{p} \|u\|_p^p \\ &\geq -\frac{\lambda}{2_s^*} T(N, s) 2_s^* (2r)^{\frac{2_s^*}{2}} - \frac{\lambda\mu a_p}{p} T(N, s)^p |\Omega|^{\frac{2_s^*-p}{2_s^*}} (2r)^{\frac{p}{2}} \\ &\geq -\lambda r \left[ \frac{2^{\frac{2^*}{2}} T(N, s) 2_s^* r^{\frac{2^*-2}{2}}}{2_s^*} + \mu \frac{2^{\frac{p}{2}} a_p T(N, s)^p |\Omega|^{\frac{2_s^*-p}{2_s^*}} r^{\frac{p-2}{2}}}{p} \right] \\ &\geq -\frac{\lambda r}{\lambda_r^*}, \end{aligned}$$

and the latter gives (4.4) since  $\lambda > \lambda_r^*$ . Now set  $v_n = u_n - u$ . We have

$$(4.5) \quad \lim_n \left[ \Phi(v_n) - \frac{\lambda}{2_s^*} \|v_n\|_{2_s^*}^{2_s^*} \right] = c - J_\lambda(u).$$

Indeed, since  $v_n \rightharpoonup 0$  in  $H_0^s(\Omega)$ , we have

$$\|v_n\|^2 = \|u_n\|^2 - 2\langle u_n, u \rangle + \|u\|^2 = \|u_n\|^2 - \|u\|^2 + o(1)$$

(as  $n \rightarrow \infty$ ). Since  $v_n \rightarrow 0$  in  $L^{2_s^*}(\Omega)$ , by the Brezis-Lieb Lemma [10, Theorem 1] we have

$$\|v_n\|_{2_s^*}^{2_s^*} = \|u_n\|_{2_s^*}^{2_s^*} - \|u\|_{2_s^*}^{2_s^*} + o(1).$$

Since  $u_n \rightarrow u$  in  $L^p(\Omega)$ , we have  $G(u_n) \rightarrow G(u)$  in  $L^1(\Omega)$ . So,

$$\begin{aligned} \Phi(v_n) - \frac{\lambda}{2_s^*} \|v_n\|_{2_s^*}^{2_s^*} &= [\Phi(u_n) - \Phi(u)] - \frac{\lambda}{2_s^*} [\|u_n\|_{2_s^*}^{2_s^*} - \|u\|_{2_s^*}^{2_s^*}] - \lambda\mu \int_{\Omega} [G(u_n) - G(u)] dx + o(1) \\ &= J_{\lambda}(u_n) - J_{\lambda}(u) + o(1) \rightarrow c - J_{\lambda}(u). \end{aligned}$$

On the other hand,

$$(4.6) \quad \lim_n [\|v_n\|^2 - \lambda \|v_n\|_{2_s^*}^{2_s^*}] = 0.$$

Indeed, arguing as above and recalling that  $g(u_n)u_n \rightarrow g(u)u$  in  $L^1(\Omega)$ , we have

$$\begin{aligned} \|v_n\|^2 - \lambda \|v_n\|_{2_s^*}^{2_s^*} &= [\|u_n\|^2 - \|u\|^2] - \lambda [\|u_n\|_{2_s^*}^{2_s^*} - \|u\|_{2_s^*}^{2_s^*}] - \lambda\mu \int_{\Omega} [g(u_n)u_n - g(u)u] dx + o(1) \\ &= \langle J'_{\lambda}(u_n), u_n \rangle - \langle J'_{\lambda}(u), u \rangle + o(1), \end{aligned}$$

and the latter tends to 0 as  $n \rightarrow \infty$ , by  $J'_{\lambda}(u_n) \rightarrow 0$ , boundedness of  $(u_n)$ , and (4.3). Recalling that  $(v_n)$  is bounded in  $H_0^s(\Omega)$ , up to a subsequence we have  $\|v_n\| \rightarrow \beta \geq 0$ . We prove that

$$(4.7) \quad \beta = 0,$$

arguing by contradiction. Assume  $\beta > 0$ . Then, by (4.6) we have

$$\beta^2 = \lim_n \lambda \|v_n\|_{2_s^*}^{2_s^*} \leq \lambda T(N, s)^{2_s^*} \beta^{2_s^*},$$

hence

$$\beta \geq \left[ \frac{1}{\lambda T(N, s)^{2_s^*}} \right]^{\frac{1}{2_s^*-2}}.$$

By (4.4) and (4.5), we also have

$$\left( \frac{1}{2} - \frac{1}{2_s^*} \right) \beta^2 = c - J_{\lambda}(u) < 2r,$$

hence

$$\beta < \left[ \frac{2Nr}{s} \right]^{\frac{1}{2}}.$$

Comparing the last inequalities and recalling (4.2), we get

$$\lambda > \frac{1}{T(N, s)^{2_s^*}} \left[ \frac{s}{2Nr} \right]^{\frac{2_s^*}{N-2s}} \geq \lambda_r^*,$$

a contradiction. So (4.7) is proved, which means  $u_n \rightarrow u$  in  $H_0^s(\Omega)$ . Thus,  $J_{\lambda}$  satisfies  $(PS)^r$ .  $\square$

Set

$$(4.8) \quad \mu^* = \min \left\{ \left[ \frac{2_s^*}{2^{\frac{2_s^*+2}{2}} T(N, s)^{2_s^*}} \right]^{\frac{2}{2_s^*-2}}, \frac{s}{3NT(N, s)^{\frac{N}{s}}} \right\}^{\frac{2-p}{2}} \frac{p}{2^{\frac{p+2}{2}} a_p T(N, s)^p |\Omega|^{\frac{2_s^*-p}{2_s^*}}} > 0.$$

We have the following existence result for problem (4.1):

**Theorem 4.2.** *Let  $\mathbf{H}_2$  hold,  $\mu^* > 0$  be defined by (4.8). Then, for all  $\mu \in (0, \mu^*)$ , (4.1) has at least one solution  $u_{\mu} \in \text{int}(C_s^0(\bar{\Omega})_+)$ .*

*Proof.* Fix  $\mu \in (0, \mu^*)$  and set

$$(4.9) \quad r = \min \left\{ \left[ \frac{2_s^*}{2^{\frac{2_s^*+2}{2}} T(N, s)^{2_s^*}} \right]^{\frac{2}{2_s^*-2}}, \frac{s}{3NT(N, s)^{\frac{N}{s}}} \right\} > 0.$$

By (4.8), (4.9) we have

$$\frac{2^{\frac{2_s^*}{2}} T(N, s)^{2_s^*} r^{\frac{2_s^*-2}{2}}}{2_s^*} + \mu \frac{2^{\frac{p}{2}} a_p T(N, s)^p |\Omega|^{\frac{2_s^*-p}{2_s^*}} r^{\frac{p-2}{2}}}{p} \leq \frac{1}{2} + \frac{\mu}{2\mu^*} < 1,$$

as well as

$$\frac{1}{T(N, s)2_s^{2_s^*}} \left[ \frac{s}{2Nr} \right]^{\frac{2_s^*}{N-2_s^*}} \geq \frac{1}{T(N, s)2_s^{2_s^*}} \left[ \frac{s}{2N} \frac{3NT(N, s)^{\frac{N}{s}}}{s} \right]^{\frac{2_s^*}{N-2_s^*}} = \left( \frac{3}{2} \right)^{\frac{2_s^*}{N-2_s^*}} > 1,$$

hence by (4.2) we have  $\lambda_r^* > 1$ .

We intend to apply Theorem B. First, we see that hypothesis (B<sub>1</sub>) holds. Then, for all  $u \in H_0^s(\Omega)$ ,  $\Phi(u) \leq r$  we have by **H**<sub>2</sub> (i) (ii), Lemma 2.1, and (2.1)

$$\begin{aligned} \frac{\Psi(u)}{r} &\leq \frac{\|u\|_{2_s^{2_s^*}}^{2_s^*}}{2_s^* r} + \mu \frac{a_p \|u\|_p^p}{pr} \\ &\leq \frac{T(N, s)2_s^{2_s^*} (2r)^{\frac{2_s^*}{2}}}{2_s^* r} + \mu \frac{a_p T(N, s)^p |\Omega|^{\frac{2_s^* - p}{2_s^*}} (2r)^{\frac{p}{2}}}{pr} \\ &\leq \frac{1}{\lambda_r^*}. \end{aligned}$$

On the other hand, by **H**<sub>2</sub> (iii) we have

$$\lim_{t \rightarrow 0^+} \frac{F(t)}{t^2} = \infty.$$

So, arguing as in the proof of Theorem 3.1, we can find  $\bar{u} \in H_0^s(\Omega)$  s.t.

$$0 < \Phi(\bar{u}) < r, \quad \frac{\Psi(\bar{u})}{\Phi(\bar{u})} > \frac{1}{\lambda_r^*},$$

which ensures (B<sub>2</sub>) and (B<sub>3</sub>). Finally, since  $\lambda_r^* > 1$ , by Lemma 4.1 the functional  $J_1$  satisfies (PS)<sup>r</sup>.

Since  $1 \in I_r$ , from Theorem B we deduce the existence of a (relabelled) function  $u_\mu \in H_0^s(\Omega)$  s.t.

$$0 < \Phi(u_\mu) < r, \quad J_1(u_\mu) = \min_{0 < \Phi(u_\mu) < r} J_1(u).$$

In particular, we have  $J_1'(u_\mu) = 0$  in  $H^{-s}(\Omega)$ . Thus, by Proposition 2.2,  $u_\mu \in \text{int}(C_s^0(\bar{\Omega})_+)$  is a solution of (4.1).  $\square$

**Remark 4.3.** The proof of Theorem 4.2 gives additional information:  $u_\mu$  is a local minimizer of  $J_1$  in  $H_0^s(\Omega)$ , satisfies the bound  $\|u_\mu\| < (2r)^{\frac{1}{2}}$ , and the mapping  $\mu \mapsto J_1(u_\mu)$  is decreasing in  $(0, \mu^*)$ .

## 5. TWO POSITIVE SOLUTIONS UNDER CRITICAL GROWTH

Finally, we turn to problem (1.4) with  $q = 2_s^*$ , namely, the Brezis-Nirenberg problem for the fractional Laplacian:

$$(5.1) \quad \begin{cases} (-\Delta)^s u = \mu u^{p-1} + u^{2_s^*-1} & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{in } \Omega^c, \end{cases}$$

with  $p \in (1, 2)$ ,  $\mu > 0$ . This is a special case of (4.1) with  $g(t) = (t^+)^{p-1}$ , which satisfies **H**<sub>2</sub> with  $a_p = 1$ . We know from [4, Theorem 1.1] that (5.1) has at least two positive solutions for all  $\mu > 0$  small enough. Our last result yields an explicitly estimate of 'how small'  $\mu$  should be, given by (4.8) which in the present case rephrases as

$$(5.2) \quad \mu^* = \min \left\{ \left[ \frac{2_s^*}{2^{\frac{2_s^*+2}{2}} T(N, s) 2_s^{2_s^*}} \right]^{\frac{2}{2_s^*-2}}, \frac{s}{3NT(N, s)^{\frac{N}{s}}} \right\}^{\frac{2-p}{2}} \frac{p}{2^{\frac{p+2}{2}} T(N, s)^p |\Omega|^{\frac{2_s^*-p}{2_s^*}}} > 0.$$

Indeed, we have the following multiplicity result:

**Theorem 5.1.** *Let  $p \in (1, 2)$ ,  $\mu^* > 0$  be defined by (5.2). Then, for all  $\mu \in (0, \mu^*)$ , (5.1) has at least two solutions  $u_\mu, w_\mu \in \text{int}(C_s^0(\bar{\Omega})_+)$ ,  $u_\mu < w_\mu$  in  $\Omega$ .*

*Proof.* Fix  $\mu \in (0, \mu^*)$ , define  $f \in C(\mathbb{R})$ ,  $\Phi, \Psi \in C^1(H_0^s(\Omega))$  as in Section 4, and set for brevity  $J = J_1 = \Phi - \Psi$ . From Theorem 4.2 and Remark 4.3 we know that there exists  $u_\mu \in H_0^s(\Omega) \cap \text{int}(C_s^0(\overline{\Omega})_+)$  which solves (5.1) and is a local minimizer of  $J$ . Set for all  $(x, t) \in \Omega \times \mathbb{R}$

$$\begin{aligned}\tilde{f}(x, t) &= f(u_\mu(x) + t^+) - f(u_\mu(x)), \\ \tilde{F}(x, t) &= \int_0^t \tilde{f}(x, \tau) d\tau = F(u_\mu(x) + t^+) - F(u_\mu(x)) - f(u_\mu(x))t^+.\end{aligned}$$

For all  $v \in H_0^s(\Omega)$  set

$$\tilde{\Psi}(v) = \int_\Omega \tilde{F}(x, v) dx, \quad \tilde{J}(v) = \Phi(v) - \tilde{\Psi}(v).$$

As in Section 2, it is easily seen that  $\tilde{J} \in C^1(H_0^s(\Omega))$  and all its critical points solve the (nonautonomous) auxiliary problem

$$(5.3) \quad \begin{cases} (-\Delta)^s v = \tilde{f}(x, v) & \text{in } \Omega \\ v = 0 & \text{in } \Omega^c. \end{cases}$$

The functionals  $\tilde{J}$  and  $J$  are related to each other by the following inequality for all  $v \in H_0^s(\Omega)$ :

$$(5.4) \quad \tilde{J}(v) \geq J(u_\mu + v^+) - J(u_\mu) + \frac{\|v^-\|^2}{2}.$$

Indeed, we have  $v^\pm \in H_0^s(\Omega)$  and, setting

$$\Omega_+ = \{x \in \Omega : v(x) > 0\}, \quad \Omega_- = \Omega \setminus \Omega_+,$$

from  $v = v^+ - v^-$  we have

$$\begin{aligned}\|v\|^2 &= \|v^+\|^2 + \|v^-\|^2 - 2 \iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{(v^+(x) - v^+(y))(v^-(x) - v^-(y))}{|x - y|^{N+2s}} dx dy \\ &\geq \|v^+\|^2 + \|v^-\|^2,\end{aligned}$$

as the integrand vanishes everywhere but in  $\Omega_+ \times \Omega_-$  and in  $\Omega_- \times \Omega_+$ , where it is negative. So we have

$$\begin{aligned}\tilde{J}(v) &= \frac{\|v\|^2}{2} - \int_\Omega \tilde{F}(x, v) dx \\ &\geq \frac{\|v^+\|^2}{2} + \frac{\|v^-\|^2}{2} - \int_\Omega [F(u_\mu + v^+) - F(u_\mu) - f(u_\mu)v^+] dx \\ &= \frac{\|u_\mu + v^+\|^2}{2} - \frac{\|u_\mu\|^2}{2} - \langle u_\mu, v^+ \rangle + \frac{\|v^-\|^2}{2} - \int_\Omega [F(u_\mu + v^+) - F(u_\mu) - f(u_\mu)v^+] dx \\ &= J(u_\mu + v^+) - J(u_\mu) + \frac{\|v^-\|^2}{2}\end{aligned}$$

(where we used that  $u_\mu$  solves (5.1)).

We claim that 0 is a local minimizer of  $\tilde{J}$ . Indeed, by Proposition 2.3 there exists  $\rho > 0$  s.t. for all  $v \in H_0^s(\Omega) \cap C_s^0(\overline{\Omega})$ ,  $\|v\|_{0,s} \leq \rho$  we have  $J(u_\mu + v) \geq J(u_\mu)$ . Then, for any such  $v$  we have as well  $\|v^+\|_{0,s} \leq \rho$ , which along with (5.4) implies

$$\tilde{J}(v) \geq J(u_\mu + v^+) - J(u_\mu) + \frac{\|v^-\|^2}{2} \geq 0 = \tilde{J}(0).$$

So, 0 is a local minimizer of  $\tilde{J}$  in  $C_s^0(\overline{\Omega})$  and hence, by Proposition 2.3 again, it is such also in  $H_0^s(\Omega)$ . In particular,  $\tilde{J}'(0) = 0$  in  $H^{-s}(\Omega)$ , i.e., 0 solves (5.3).

From now on we closely follow [4]. Arguing by contradiction, assume that 0 is the *only* critical point of  $\tilde{J}$  in  $H_0^s(\Omega)$ . Under such assumption, by [4, Lemma 2.10]  $\tilde{J}$  satisfies  $(PS)_c$  at any level  $c < c^*$ , where

$$(5.5) \quad c^* = \frac{s}{NT(N, s)^{\frac{N}{s}}}.$$

Fix  $x_0 \in \Omega$ , and for all  $\varepsilon > 0$ , define  $v_\varepsilon \in H^s(\mathbb{R}^N)$  by setting for all  $x \in \mathbb{R}^N$

$$v_\varepsilon(x) = \frac{\varepsilon^{\frac{N-2s}{2}}}{(\varepsilon^2 + |x - x_0|^2)^{\frac{N-2s}{2}}}.$$

By Lemma 2.1 we have

$$(5.6) \quad \|v_\varepsilon\|_{2_s^*} = T(N, s)[v_\varepsilon]_s.$$

Now fix  $r > 0$  s.t.  $\overline{B}_r(x_0) \subset \Omega$ ,  $\eta \in C^\infty(\mathbb{R}^N)$  s.t.  $\eta = 1$  in  $B_{\frac{r}{2}}(x_0)$ ,  $\eta = 0$  in  $B_1^c(x_0)$ , and  $0 \leq \eta \leq 1$  in  $\mathbb{R}^N$ , then define  $w_\varepsilon \in H_0^s(\Omega)$  by setting for all  $x \in \mathbb{R}^N$

$$w_\varepsilon(x) = \frac{\eta(x)v_\varepsilon(x)}{\|\eta v_\varepsilon\|_{2_s^*}}.$$

Clearly  $\|w_\varepsilon\|_{2_s^*} = 1$ . Besides, we will prove that for all  $\varepsilon > 0$  small enough

$$(5.7) \quad \max_{\tau \geq 0} \tilde{J}(\tau w_\varepsilon) < c^*.$$

Assume  $N > 4s$ . Then, by [29, Propositions 21, 22] we find for all  $\varepsilon > 0$  small enough

$$\begin{aligned} \|w_\varepsilon\|^2 &\leq \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} \\ \|w_\varepsilon\|_2^2 &\geq C\varepsilon^{2s} - C\varepsilon^{N-2s} \end{aligned}$$

( $C > 0$  denotes several constants, independent of  $\varepsilon$ ). By convexity we have for all  $x \in \Omega$ ,  $t \geq 0$

$$\tilde{F}(x, t) \geq \frac{t^{2_s^*}}{2_s^*} + \frac{C}{2} u_\mu(x)^{2_s^*-2} t^2.$$

Using (5.6) and the relations above, we see that for all  $\varepsilon > 0$  small enough and all  $\tau \geq 0$

$$(5.8) \quad \begin{aligned} \tilde{J}(\tau w_\varepsilon) &\leq \frac{\tau^2}{2} \|w_\varepsilon\|^2 - \frac{\tau^{2_s^*}}{2_s^*} \|w_\varepsilon\|_{2_s^*}^{2_s^*} - \frac{C\tau^2}{2} \int_\Omega u_\mu^{2_s^*-2} w_\varepsilon^2 dx \\ &\leq \frac{\tau^2}{2} \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} - C'\varepsilon^{2s} \right] - \frac{\tau^{2_s^*}}{2_s^*} =: h_\varepsilon(\tau) \end{aligned}$$

( $C, C' > 0$  independent of  $\varepsilon$ ). Now we focus on the mapping  $h_\varepsilon \in C^1(\mathbb{R}_+)$ . First we note that

$$\lim_{\tau \rightarrow \infty} h_\varepsilon(\tau) = -\infty,$$

so there exists  $\tau_\varepsilon \geq 0$  s.t.

$$h_\varepsilon(\tau_\varepsilon) = \max_{\tau \geq 0} h_\varepsilon(\tau).$$

If  $\tau_\varepsilon = 0$ , from (5.8) we immediately deduce (5.7). So, let  $\tau_\varepsilon > 0$ . Differentiating  $h_\varepsilon$ , we get

$$\tau_\varepsilon = \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} - C'\varepsilon^{2s} \right]^{\frac{1}{2_s^*-2}},$$

which tends to  $T(N, s)^{-\frac{2}{2_s^*-2}} > 0$  as  $\varepsilon \rightarrow 0^+$ . So, taking  $\varepsilon > 0$  small enough, we have  $\tau_\varepsilon \geq \tau_0 > 0$ . Set

$$\tilde{\tau}_\varepsilon = \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} \right]^{\frac{1}{2_s^*-2}},$$

and note that the mapping

$$\tau \mapsto \frac{\tau^2}{2} \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} \right] - \frac{\tau^{2_s^*}}{2_s^*}$$

is increasing in  $[0, \tilde{\tau}_\varepsilon]$ . So we have

$$\begin{aligned} h_\varepsilon(\tau_\varepsilon) &= \frac{\tau_\varepsilon^2}{2} \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} \right] - \frac{\tau_\varepsilon^{2_s^*}}{2_s^*} - \frac{C'\varepsilon^{2s}\tau_\varepsilon^2}{2} \\ &\leq \frac{\tilde{\tau}_\varepsilon^2}{2} \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} \right] - \frac{\tilde{\tau}_\varepsilon^{2_s^*}}{2_s^*} - C''\varepsilon^{2s} \\ &= \frac{s}{N} \left[ \frac{1}{T(N, s)^2} + C\varepsilon^{N-2s} \right]^{\frac{N}{2s}} - C''\varepsilon^{2s}. \end{aligned}$$

Since  $N - 2s > 2s$ , for all  $\varepsilon > 0$  small enough we have by (5.5)

$$h_\varepsilon(\tau_\varepsilon) < \frac{s}{NT(N, s)^{\frac{N}{s}}} = c^*.$$

Then, by (5.8) we obtain (5.7). The cases  $2s < N \leq 4s$  are treated in similar ways, see [4, Lemma 2.11].

As a byproduct of (5.8) we have that  $\tilde{J}(\tau w_\varepsilon) \rightarrow -\infty$  as  $\tau \rightarrow \infty$ , so we can find  $\bar{\tau} > 0$  s.t.

$$\tilde{J}(\bar{\tau} w_\varepsilon) < 0.$$

Since  $\tilde{J}$  has a local minimum at 0 and no other critical point, we can find  $\sigma \in (0, \|\bar{\tau} w_\varepsilon\|)$  s.t.  $\tilde{J}(v) > 0$  for all  $v \in H_0^s(\Omega)$ ,  $\|v\| = \sigma$ . That is,  $\tilde{J}$  exhibits a mountain pass geometry around 0. Set

$$\Gamma = \{ \gamma \in C([0, 1], H_0^s(\Omega)) : \gamma(0) = 0, \gamma(1) = \bar{\tau} w_\varepsilon \}, \quad c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} \tilde{J}(\gamma(t)).$$

Clearly,  $\gamma(t) = t\bar{\tau} w_\varepsilon$  define a path of the family  $\Gamma$ , so by (5.7) we have

$$c \leq \max_{t \in [0, 1]} \tilde{J}(t\bar{\tau} w_\varepsilon) < c^*.$$

Thus,  $\tilde{J}$  satisfies  $(PS)_c$ . By the Mountain Pass Theorem, there exists  $v \in H_0^s(\Omega) \setminus \{0\}$  s.t.  $\tilde{J}'(v) = 0$  in  $H^{-s}(\Omega)$ , a contradiction.

So we have proved the existence of  $v_\mu \in H_0^s(\Omega) \setminus \{0\}$  s.t.  $\tilde{J}'(v_\mu) = 0$  in  $H^{-s}(\Omega)$ . Such  $v_\mu$  solves (5.3), and by monotonicity of  $f$  we have for a.e.  $x \in \Omega$

$$\tilde{f}(x, v_\mu(x)) = f(u_\mu(x) + v_\mu^+(x)) - f(u_\mu(x)) \geq 0,$$

so by the fractional Hopf lemma (see for instance [22, Lemma 2.7], as Proposition 2.2 here does not apply) we have  $v_\mu \in \text{int}(C_s^0(\bar{\Omega})_+)$ . Now set

$$w_\mu = u_\mu + v_\mu \in \text{int}(C_s^0(\bar{\Omega})_+).$$

Clearly  $w_\mu > u_\mu$  in  $\Omega$ , and for all  $\varphi \in H_0^s(\Omega)$  we have

$$\begin{aligned} \langle J'(w_\mu), \varphi \rangle &= \langle u_\mu + v_\mu, \varphi \rangle - \int_\Omega f(u_\mu + v_\mu) \varphi \, dx \\ &= \left[ \langle u_\mu, \varphi \rangle - \int_\Omega f(u_\mu) \varphi \, dx \right] + \left[ \langle v_\mu, \varphi \rangle - \int_\Omega \tilde{f}(x, v_\mu) \varphi \, dx \right] \\ &= \langle J'(u_\mu), \varphi \rangle + \langle \tilde{J}'(v_\mu), \varphi \rangle = 0, \end{aligned}$$

so  $w_\mu$  solves (5.1), which concludes the proof.  $\square$

**Acknowledgement.** Both authors are members of GNAMPA (Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni) of INdAM (Istituto Nazionale di Alta Matematica 'Francesco Severi') and supported by the research project *Integro-differential Equations and nonlocal Problems* funded by Fondazione di Sardegna (2017). A. Iannizzotto is also supported by the grant PRIN n. 2017AYM8XW: *Non-linear Differential Problems via Variational, Topological and Set-valued Methods*. We thank Prof. G. Bonanno for his precious suggestions.

## REFERENCES

- [1] A. AMBROSETTI, H. BREZIS, G. CERAMI, Combined effects of concave and convex nonlinearities in some elliptic problems, *J. Funct. Anal.* **122** (1994) 519–543.
- [2] G. AUTUORI, P. PUCCI, Elliptic problems involving the fractional Laplacian in  $\mathbb{R}^N$ , *J. Differential Equations* **255** (2013) 2340–2362.
- [3] B. BARRIOS, E. COLORADO, A. DE PABLO, U. SÁNCHEZ, On some critical problems for the fractional Laplacian operator, *J. Differential Equations* **252** (2012) 6133–6162.
- [4] B. BARRIOS, E. COLORADO, R. SERVADEI, F. SORIA, A critical fractional equation with concave-convex power nonlinearities, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **32** (2015) 875–900.
- [5] Z. BINLIN, G. MOLICA BISCI, R. SERVADEI, Superlinear nonlocal fractional problems with infinitely many solutions, *Nonlinearity* **28** (2015) 2247–2264.
- [6] G. BONANNO, A critical point theorem via the Ekeland variational principle, *Nonlinear Anal.* **75** (2012) 2992–3007.
- [7] G. BONANNO, G. D’AGUÌ, Two non-zero solutions for elliptic Dirichlet problems, *Zeitsch. Anal. Anwend.* **35** (2016) 449–465.
- [8] G. BONANNO, G. D’AGUÌ, Critical nonlinearities for elliptic Dirichlet problems, *Dynam. Systems Appl.* **22** (2013) 411–418.
- [9] G. BONANNO, G. D’AGUÌ, D. O’REGAN, A local minimum theorem and critical nonlinearities, *An. Șt. Univ. Ovidius Constanța* **24** (2016) 67–86.
- [10] H. BREZIS, E. LIEB, A relation between pointwise convergence of functions and convergence of functionals, *Proc. Amer. Math. Soc.* **88** (1983) 486–490.
- [11] H. BREZIS, L. NIRENBERG, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponent, *Comm. Pure Appl. Math.* **36** (1983) 437–477.
- [12] H. BREZIS, L. NIRENBERG,  $H^1$  versus  $C^1$  minimizers, *C. R. Acad. Sci. Paris* **317** (1993) 465–472.
- [13] C. BUCUR, E. VALDINOCI, Nonlocal diffusion and applications, Springer, New York (2016).
- [14] X. CABRÉ, Y. SIRE, Nonlinear equations for fractional Laplacians I: Regularity, maximum principles, and Hamiltonian estimates, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **31** (2014) 23–53.
- [15] X. CABRÉ, Y. SIRE, Nonlinear equations for fractional Laplacians II: Existence, uniqueness, and qualitative properties of solutions, *Trans. Amer. Math. Soc.* **367** (2015) 911–941.
- [16] X. CHANG, Z.Q. WANG, Nodal and multiple solutions of nonlinear problems involving the fractional Laplacian, *J. Differential Equations* **256** (2014) 2965–2992.
- [17] A. COTSIOLIS, N. K. TAVOULARIS, Best constants for Sobolev inequalities for higher order fractional derivatives, *J. Math. Anal. Appl.* **295** (2004) 225–236.
- [18] E. DI NEZZA, G. PALATUCCI, E. VALDINOCI, Hitchhiker’s guide to the fractional Sobolev spaces, *Bull. Sci. Math.* **136** (2012) 521–573.
- [19] F.G. DÜZGÜN, A. IANIZZOTTO, Three nontrivial solutions for nonlinear fractional Laplacian equations, *Adv. Nonlinear Anal.* **7** (2018) 211–226.
- [20] A. FISCELLA, Saddle point solutions for non-local elliptic operators, *Topol. Methods Nonlinear Anal.* **44** (2014) 527–538.
- [21] S. FRASSU, Nonlinear Dirichlet problem for the nonlocal anisotropic operator  $L_K$ , *Commun. Pure Appl. Anal.* **18** (2019) 1847–1867.
- [22] A. IANIZZOTTO, S. MOSCONI, M. SQUASSINA,  $H^s$  versus  $C^0$ -weighted minimizers, *Nonlinear Differ. Equ. Appl.* **22** (2015) 477–497.
- [23] A. IANIZZOTTO, S. MOSCONI, M. SQUASSINA, Sobolev versus Hölder minimizers for the degenerate fractional  $p$ -Laplacian, *Nonlinear Anal.* **191** (2020) DOI: 10.1016/j.na.2019.111635.
- [24] A. IANIZZOTTO, N.S. PAPAGEORGIOU, Existence and multiplicity results for resonant fractional boundary value problems, *Discrete Contin. Dyn. Syst. Ser. S* **11** (2018) 511–532.
- [25] G. MOLICA BISCI, V.D. RĂDULESCU, R. SERVADEI, Variational methods for nonlocal fractional problems, Cambridge University Press, Cambridge (2016).
- [26] G. MOLICA BISCI, R. SERVADEI, Lower semicontinuity of functionals of fractional type and applications to nonlocal equations with critical Sobolev exponent, *Adv. Differential Equations* **20** (2015) 635–660.
- [27] D. MUGNAI, D. PAGLIARDINI, Existence and multiplicity results for the fractional Laplacian in bounded domains, *Adv. Calc. Var.* **10** (2017) 111–124.
- [28] X. ROS-OTON, J. SERRA, The Dirichlet problem for the fractional Laplacian: regularity up to the boundary, *J. Math. Pures Appl.* **101** (2014) 275–302.
- [29] R. SERVADEI, E. VALDINOCI, Mountain pass solutions for non-local elliptic operators, *J. Math. Anal. Appl.* **389** (2012) 887–898.
- [30] R. SERVADEI, E. VALDINOCI, Variational methods for non-local operators of elliptic type, *Discrete Contin. Dyn. Syst.* **33** (2013) 2105–2137.
- [31] R. SERVADEI, E. VALDINOCI, The Brezis-Nirenberg result for the fractional Laplacian, *Trans. Amer. Math. Soc.* **367** (2015) 67–102.
- [32] Y. WEI, X. SU, Multiplicity of solutions for nonlocal elliptic equations driven by the fractional Laplacian, *Calc. Var.* **52** (2015) 95–124.

(S. Frassu, A. Iannizzotto) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE  
 UNIVERSITY OF CAGLIARI  
 VIA OSPEDALE 72, 09124 CAGLIARI, ITALY  
 E-mail address: [silvia.frassu@unica.it](mailto:silvia.frassu@unica.it), [antonio.iannizzotto@unica.it](mailto:antonio.iannizzotto@unica.it)