EXTENDED AFFINE LIE ALGEBRAS, VERTEX ALGEBRAS, AND REDUCTIVE GROUPS

FULIN CHEN¹, HAISHENG LI, SHAOBIN TAN², AND QING WANG³

ABSTRACT. In this paper, we explore natural connections among the representations of the extended affine Lie algebra $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ with $\mathbb{C}_q=\mathbb{C}_q[t_0^{\pm 1},t_1^{\pm 1}]$ an irrational quantum 2-torus, the simple affine vertex algebra $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ with ℓ a positive integer, and Levi subgroups $\mathrm{GL}_{\mathbf{I}}$ of $\mathrm{GL}_{\ell}(\mathbb{C})$. First, we give a canonical isomorphism between the category of integrable restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ and that of equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ -modules. Second, we classify irreducible \mathbb{N} -graded equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ -modules. Third, we establish a duality between irreducible \mathbb{N} -graded equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ -modules and irreducible regular $\mathrm{GL}_{\mathbf{I}}$ -modules on certain fermionic Fock spaces. Fourth, we obtain an explicit realization of every irreducible \mathbb{N} -graded equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ -module. Fifth, we completely determine the following branchings: (i) The branching from $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0) \otimes L_{\widehat{\mathfrak{sl}_\infty}}(\ell',0)$ to $L_{\widehat{\mathfrak{sl}_\infty}}(\ell+\ell',0)$ for quasi modules. (ii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its Levi subalgebras. (iii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its Levi subalgebras. (iii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its subalgebras $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$.

1. Introduction

This paper is to establish and explore a natural connection between the extended affine Lie algebras of type A_{N-1} coordinated by irrational quantum tori and vertex algebras. Extended affine Lie algebra, written as EALA in short, was first introduced by Hoegh-Krohn and Torresani in [H-KT] under the name of quasisimple Lie algebra, and since then it has been extensively studied in literature (see [AABGP, BGK, N] and the references therein). By definition, an EALA is a Lie algebra together with a finite-dimensional ad-diagonalizable subalgebra and a nondegenerate symmetric invariant bilinear form, satisfying a list of conditions. One of the conditions is that the group generated by the isotropic roots is a free abelian group of finite rank called the *nullity*. It is known that EALAs of nullity 0 are exactly finite-dimensional simple Lie algebras, while EALAs of nullity 1 are precisely affine Kac-Moody algebras (see [ABGP]).

²⁰¹⁰ Mathematics Subject Classification. 17B67 & 17B69.

¹Partially supported by China NSF grant (No.11971397) and the Fundamental Research Funds for the Central Universities (No.20720190069).

²Partially supported by China NSF grants (Nos.11531004, 11971397).

³Partially supported by China NSF grants (Nos.11531004, 11622107).

Note that affine Kac-Moody algebras were classified as untwisted affine Lie algebras and twisted affine Lie algebras (see [K]). It is known that untwisted affine Lie algebras through their highest weight modules can be naturally associated with vertex operator algebras and modules (see [FZ]), whereas twisted affine Lie algebras can be associated to twisted modules for the vertex operator algebras associated to the corresponding untwisted affine Lie algebras (see [FLM], [Li2]). These (affine) vertex operator algebras, being the major building blocks in vertex operator algebra theory, play an important role. An eminent problem is to establish and explore natural connections of vertex algebras with all EALAs.

The structure of a general EALA is now pretty well understood and the EALAs with positive nullity are like affine Kac-Moody algebras in many ways. Among these Lie algebras, toroidal extended affine Lie algebras are multi-loop generalizations of untwisted affine Lie algebras. It was proved (see [BGK, ABFP, N]) that every (maximal) EALA of positive nullity is isomorphic to a toroidal EALA, or the subalgebra of the fixed points in a toroidal EALA with respect to a finite abelian automorphism group, or an EALA of type A coordinated by an irrational quantum torus. Meanwhile, there have been several studies on EALAs and their related algebras where vertex algebras have played an important role. In [BBS], Berman, Billig, and Szmigielski established a natural connection between toroidal Lie algebras and affine vertex operator algebras. Later on, Billig studied in [B1] the structure of the vertex operator algebras associated to full toroidal Lie algebras. By using the vertex operator algebras constructed in [B1], some irreducible integrable modules for toroidal EALAs and their subalgebras of the fixed-points under certain automorphism groups were constructed in [B2, CLT] and [BL], respectively. As for the EALAs coordinated by irrational quantum tori, an explicit natural connection with vertex algebras is yet to be established. This is the main concern of this present paper.

In this paper, we concentrate ourselves to nullity 2 EALAs coordinated by irrational quantum tori. Let q be a generic complex number. By definition, the quantum 2-torus, denoted by $\mathbb{C}_q[t_0^{\pm 1}, t_1^{\pm 1}]$ or simply by \mathbb{C}_q , is the associative algebra with underlying space $\mathbb{C}[t_0^{\pm 1}, t_1^{\pm 1}]$ and with the basic commutation relation $t_1t_0 = qt_0t_1$. On the other hand, let N be a positive integer with $N \geq 2$. Denote by $\mathfrak{gl}_N(\mathbb{C}_q)$ the matrix Lie algebra over \mathbb{C}_q and set $\mathfrak{sl}_N(\mathbb{C}_q) = [\mathfrak{gl}_N(\mathbb{C}_q), \mathfrak{gl}_N(\mathbb{C}_q)]$, the derived subalgebra. Furthermore, let $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ denote the universal central extension of $\mathfrak{sl}_N(\mathbb{C}_q)$. The nullity 2 EALA $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ coordinated by \mathbb{C}_q by definition is the extension of $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ by the two canonical degree-zero derivations (see [BGK]). Representations of $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ and $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ have been extensively studied in literature (cf. [BS, BGT, G1, G2, G3, GZ, ER, CT]). In particular, a fermionic construction of the basic modules for $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ was obtained by Gao in [G3] and integrable highest weight modules for $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ were studied by Rao (see [ER]). As for this paper, our focus will be on the Lie algebra $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$.

Note that the EALAs coordinated by irrational quantum tori cannot be directly associated to modules or twisted modules for vertex algebras, due to the fact that their canonical generating functions are not "local." Then we come to a theory of what were called quasi modules for vertex algebras. The notion of quasi module was introduced in [Li3] to associate vertex algebras to a certain family of Lie algebras. Indeed, with this new theory a much wider variety of Lie algebras, including the quantum 2-torus Lie algebra, can be associated with vertex algebras. The theory of quasi modules was developed further in [Li4], where a notion of vertex Γ -algebra with Γ a group was introduced and an enhanced notion of quasi module, called equivariant quasi module for a vertex Γ -algebra, was introduced. The notion of quasi module from definition generalizes that of module. On the other hand, this notion also generalizes the notion of twisted module in a certain natural way (see [Li5]).

In this paper, we shall intensively study a natural connection of the Lie algebra $\mathfrak{sl}_N(\mathbb{C}_q)$ with vertex algebras and their equivariant quasi modules. As the first step, we relate $\mathfrak{sl}_N(\mathbb{C}_q)$ to a (general) affine Lie algebra. Let \mathfrak{gl}_{∞} be the Lie algebra of infinite order complex matrices with only finitely many nonzero entries and set $\mathfrak{sl}_{\infty} = [\mathfrak{gl}_{\infty}, \mathfrak{gl}_{\infty}]$ (the derived subalgebra). Equip \mathfrak{gl}_{∞} with a suitable nondegenerate symmetric invariant bilinear form. Then we have an affine Lie algebra $\widehat{\mathfrak{sl}_{\infty}} = \mathfrak{sl}_{\infty} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}\mathbf{k}$. (Note that $\widehat{\mathfrak{sl}_{\infty}}$ is different from the infinite rank affine Kac-Moody algebra $\overline{\mathfrak{sl}_{\infty}}$ which is a completion of \mathfrak{sl}_{∞} (see [K]).) For any complex number ℓ , we have a vertex algebra $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ and its simple quotient vertex algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. We show that $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ has an intrinsic connection with the affine Lie algebra \mathfrak{sl}_{∞} . As the first main result, using this intrinsic connection we establish a canonical category isomorphism between the categories of restricted $\mathfrak{sl}_N(\mathbb{C}_q)$ -modules of level ℓ and equivariant quasi modules for $V_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$. Furthermore, assuming that ℓ is a positive integer, we show that under the canonical category isomorphism, integrable and restricted $\mathfrak{sl}_N(\mathbb{C}_q)$ -modules of level ℓ correspond exactly to equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -modules.

In representation theory, one of the prominent notions is that of dual pair which was first studied by Howe for reductive groups (see [H1, H2]). In [F], Igor Frenkel discovered a duality for affine Lie algebras of type A, which is now commonly known as the level-rank duality. Wang in [W2] obtained certain dualities between a completed infinite rank affine Kac-Moody algebra and certain reductive Lie groups. An interesting level-rank duality was also found for vertex operator algebras of types A by Jiang and Lin in [JLin] and B, D by Jiang and Lam in [JLam]. In [VV1], the usual Schur ($\mathfrak{gl}_N(\mathbb{C}), S_\ell$)-duality was generalized to the two-parameters quantum toroidal algebra and the classical limit (see [VV2]) gives rise to a duality between $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ and a distinguished double affine Hecke algebra.

As the second main result of this paper, we obtain a level-rank type duality between irreducible integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules of level $\ell \in \mathbb{Z}_+$ and irreducible regular modules for Levi subgroups $\operatorname{GL}_{\mathbf{I}}$ of $\operatorname{GL}_{\ell}(\mathbb{C})$, where the parameter \mathbf{I} is a partition of $\{1,\ldots,\ell\}$. To a certain extent, this duality is analogous to the classical skew $(\mathfrak{gl}_N(\mathbb{C}),\operatorname{GL}_{\ell}(\mathbb{C}))$ -duality (see [H2]). By using this $(\widehat{\mathfrak{sl}}_N(\mathbb{C}_q),\operatorname{GL}_{\mathbf{I}})$ -duality, we furthermore obtain a fermionic realization of every irreducible integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module. Consequently, we obtain a realization of every irreducible \mathbb{N} -graded equivariant quasi $L_{\widehat{\mathfrak{sl}}_\infty}(\ell,0)$ -module.

Note that one of important applications of dual pair is to study branchings. For a pair $H \subset G$ of (complex) reductive groups, the branching law from G to H is a description of the irreducible H-modules and their multiplicities that occur in the decomposition of each irreducible regular G-module. Among the better known classical branchings are the branching from $H \times H$ to H and the branching from a group G to a Levi subgroup G. Analogously, the two better branchings in vertex operator algebra theory are the branchings from $L_{\widehat{\mathfrak{g}}}(\ell,0) \otimes L_{\widehat{\mathfrak{g}}}(\ell',0)$ to $L_{\widehat{\mathfrak{g}}}(\ell+\ell',0)$ and from $L_{\widehat{\mathfrak{g}}}(\ell,0)$ to $L_{\widehat{\mathfrak{f}}}(\ell,0)$, where ℓ,ℓ' are positive integers and \mathfrak{h} is a Levi subalgebra of \mathfrak{g} . On the other hand, Igor Frenkel studied in [F] the $\widehat{\mathfrak{g}} \to \widehat{\mathfrak{g}}^{(M)}$ branching, where $\widehat{\mathfrak{g}}^{(M)} = \mathfrak{g} \otimes \mathbb{C}[t^M, t^{-M}] \oplus \mathbb{C}\mathbf{k} \subset \widehat{\mathfrak{g}}$ with M a positive integer. Frenkel made a conjecture on the decomposition of the basic modules and he confirmed the conjecture for $\mathfrak{g} = \mathfrak{sl}_N$ by using the level-rank duality for $\widehat{\mathfrak{sl}}_N$.

As the third part of this paper, we study three analogous branchings. More specifically, we determine the following branchings: (i) The branching from $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$ to $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ on equivariant quasi modules. (ii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its Levi subalgebras. (iii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its subalgebra $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0},t_1^{\pm M_1}])$. It turns out that all the multiplicities in these branchings are *finite*, which can be calculated by the Littlewood-Richardson rule (cf. [GW]). This is quite different from the branching from $L_{\widehat{\mathfrak{g}}}(\ell,0) \otimes L_{\widehat{\mathfrak{g}}}(\ell',0)$ to $L_{\widehat{\mathfrak{g}}}(\ell+\ell',0)$ on ordinary modules with $\widehat{\mathfrak{g}}$ a finite rank affine Kac-Moody algebra, where the multiplicities are in general infinite and determining such branchings is an important and difficult problem (cf. [JLin, JLam, KMPX]).

Now, we start a more detailed introduction section by section. In Section 2, we recall the Lie algebra $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ and establish a canonical isomorphism between $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ and what was called the covariant algebra of the affine Lie algebra $\widehat{\mathfrak{sl}}_\infty$ (see [G-KK], [Li3]). Recall that associated to any Lie algebra \mathfrak{g} with a nondegenerate symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$ we have an affine Lie algebra $\widehat{\mathfrak{g}}$. Furthermore, let G be an automorphism group of \mathfrak{g} , which preserves the form $\langle \cdot, \cdot \rangle$, and let $\chi: G \to \mathbb{C}^\times$ be a linear character such that for any $a, b \in \mathfrak{g}$,

$$[ga, b] = 0$$
 and $\langle ga, b \rangle = 0$ for all but finitely many $g \in G$.

The covariant algebra $\widehat{\mathfrak{g}}[G]$ of the affine Lie algebra $\widehat{\mathfrak{g}}$ is defined to be the Lie algebra with $\widehat{\mathfrak{g}}$ as the generating space space, subject to the following relations

$$g(a \otimes t^n) = \chi(g)^n ga \otimes t^n,$$

$$[a \otimes t^m, b \otimes t^n] = \sum_{g \in G} \chi(g)^m \left([ga, b] \otimes t^{m+n} + m \delta_{m+n,0} \langle ga, b \rangle \mathbf{k} \right)$$

for $g \in G$, $a, b \in \mathfrak{g}$, $m, n \in \mathbb{Z}$, and $[\mathbf{k}, \widehat{\mathfrak{g}}] = 0$. For the Lie algebra \mathfrak{gl}_{∞} , take the automorphism group $\mathbb{Z} = \langle \sigma^N \rangle$, where $\sigma E_{m,n} = E_{m+1,n+1}$ for $m, n \in \mathbb{Z}$, together with the linear character χ_q defined by $\chi_q(n) = q^n$ for $n \in \mathbb{Z}$. It is proved that $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ is canonically isomorphic to the covariant algebra $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$.

In Section 3, we give a precise connection between $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ and equivariant quasi modules for vertex algebras $V_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ and $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$. We first recall the notion of quasi module for a vertex algebra V and some basic results. The notion of quasi module is defined by simply replacing the Jacobi identity in the definition of a module with the "quasi Jacobi identity" stating that for any $u, v \in V$, the usual Jacobi identity after multiplied by a nonzero polynomial depending on u, v holds. For the notion of vertex Γ -algebra, Γ is a group with a linear character $\chi: \Gamma \to \mathbb{C}^\times$, where a $vertex\ \Gamma$ -algebra is a vertex algebra V equipped with a representation R of Γ on V such that $R_q(1) = 1$ for $g \in \Gamma$ and

$$R_g Y(v, x) R_g^{-1} = Y(R_g v, \chi(g)^{-1} x)$$
 for $g \in \Gamma, v \in V$.

Note that Γ does not act on V as an automorphism group in general. On the other hand, suppose V is a \mathbb{Z} -graded vertex algebra in the sense that V is a vertex algebra with a \mathbb{Z} -grading $V = \bigoplus_{n \in \mathbb{Z}} V_{(n)}$ such that $\mathbf{1} \in V_{(0)}$ and

$$u_r V_{(n)} \subset V_{(m+n-r-1)}$$
 for $u \in V_{(m)}, m, r, n \in \mathbb{Z}$.

Then for any automorphism group Γ of V such that $gV_{(n)} \subset V_{(n)}$ for $g \in \Gamma$, $n \in \mathbb{Z}$ and for any linear character χ of Γ , V becomes a vertex Γ -algebra with $R_g = \chi(g)^{-L(0)}g$ for $g \in \Gamma$, where L(0) denotes the grading operator on V. For a vertex Γ -algebra V, an equivariant quasi module is a quasi module (W, Y_W) satisfying the conditions that

$$Y_W(R_q v, x) = Y_W(v, \chi(q)x)$$
 for $q \in \Gamma$, $v \in V$

and that for $u, v \in V$, there exists a nonzero polynomial f(z) whose roots are contained in $\chi(\Gamma)$ such that

$$f(x_1/x_2)[Y_W(u,x_1),Y_W(v,x_2)]=0.$$

For any complex number ℓ , we have a \mathbb{Z} -graded vertex algebra $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ and its simple quotient $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. On the other hand, the automorphism σ of the Lie algebra \mathfrak{sl}_{∞} induces an automorphism of $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ and $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. Take $\mathbb{Z}=\langle\sigma^N\rangle$ with the linear character χ_q . Then $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ and $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ are both vertex (\mathbb{Z},χ_q) -algebras. As the main results of this section, we establish a canonical category isomorphism between the categories of restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ and equivariant quasi modules for $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. Furthermore, assuming that ℓ is a positive integer, we prove that under the canonical category isomorphism, integrable

and restricted $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules of level ℓ correspond exactly to equivariant quasi $L_{\widehat{\mathfrak{sl}}_N}(\ell,0)$ -modules.

In Section 4, we classify irreducible N-graded equivariant quasi modules for $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ with ℓ any positive integer. First, we show that every irreducible N-graded equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -module is an irreducible integrable highest weight $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module of level ℓ . Then, slightly generalizing a result of Rao in [ER] we obtain a classification of irreducible integrable highest weight $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ , which are parameterized by a positive integer k and a pair

$$(\boldsymbol{\lambda}, \boldsymbol{c}) = ((\lambda_1, \dots, \lambda_k), (c_1, \dots, c_k)) \in P_+^k \times (\mathbb{C}^\times)^k$$

such that $\lambda_1 + \cdots + \lambda_k$ is of level ℓ , where P_+ denotes the set of dominant integral weights for the affine Lie algebra $\widehat{\mathfrak{sl}_N}$. From the canonical category isomorphism given in Section 3, we obtain an explicit classification of irreducible N-graded equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ -modules.

In Sections 5 and 6, we present a level-rank duality between irreducible integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules of level ℓ and finite-dimensional irreducible regular modules for an arbitrary Levi subgroup of the general linear group $\mathrm{GL}_\ell = \mathrm{GL}_\ell(\mathbb{C})$. As an application, we obtain an explicit realization of every irreducible N-graded equivariant quasi $L_{\widehat{\mathfrak{sl}}_\infty}(\ell,0)$ -module. More specifically, set

$$(\mathbb{C}^{\times})_q^{\ell} = \{(a_1, \dots, a_{\ell}) \in (\mathbb{C}^{\times})^{\ell} \mid \text{either } a_i = a_j \text{ or } a_i \notin a_j \Gamma_q \text{ for } 1 \leq i, j \in \ell\},$$

where $\Gamma_q = \{q^n \mid n \in \mathbb{Z}\}$. For any $\boldsymbol{a} \in (\mathbb{C}^\times)_q^\ell$, by using a result of Gao (see [G3]) and the Chari-Pressely evaluation module construction (see [CP]), we construct a fermionic vacuum module \mathcal{F}_N^a for $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ of level ℓ . On the other hand, for any partition \mathbf{I} of $\{1,\ldots,\ell\}$, denote by $\mathrm{GL}_{\mathbf{I}}$ the Levi subgroup of GL_ℓ associated to \mathbf{I} . From [FF], \mathcal{F}_N^a is naturally a locally regular $\mathrm{GL}_{\mathbf{I}}$ -module. Using Frenkel's level-rank duality, we prove that for any $\boldsymbol{a} \in (\mathbb{C}^\times)_q^\ell$, there is a partition \mathbf{I}_a of $\{1,\ldots,\ell\}$ associated to \boldsymbol{a} such that $(\widehat{\mathfrak{sl}}_N(\mathbb{C}_q),\mathrm{GL}_{\mathbf{I}_a})$ (and hence $(L_{\widehat{\mathfrak{sl}}_\infty}(\ell,0),\mathrm{GL}_{\mathbf{I}_a})$) form a dual pair on \mathcal{F}_N^a in the sense of Howe (see [H1]). We also determine the irreducible isotypic decomposition of \mathcal{F}_N^a as an $(\widehat{\mathfrak{sl}}_N(\mathbb{C}_q),\mathrm{GL}_{\mathbf{I}_a})$ -module. Furthermore, it is proved that every finite-dimensional irreducible regular module of every Levi subgroup of GL_ℓ occurs in \mathcal{F}_N^a and every irreducible integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module of level ℓ occurs in \mathcal{F}_N^a for a suitable choice of \boldsymbol{a} .

In Sections 7, 8, and 9, we study the following branchings respectively: (i) The branching from $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$ to $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$. (ii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its Levi subalgebras. (iii) The branching from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to its subalgebra $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$. Note that the first branching is about quasi modules and it amounts to the tensor product decomposition of irreducible integrable highest weight $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules. By employing the reciprocity laws associated to certain seesaw pairs related to the dual pair $(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}_a})$, we completely determine

these three branchings. More specifically, in Section 7, for every irreducible N-graded equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$ -module W, we show that W as a quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ -module is completely reducible and we determine the irreducible quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ -modules and their multiplicities in W. We also show that the branching from $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(1,0)$ to $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+1,0)$ is multiplicity free. Similarly, we completely determine the other two branchings.

For this paper, we use symbols \mathbb{Z}_+ and \mathbb{N} for the sets of positive integers and nonnegative integers, respectively. We work on the field \mathbb{C} of complex numbers.

2. Extended affine Lie algebras coordinated by quantum tori

In this section, we first recall the EALA (and its core) of type A_{N-1} coordinated by the 2-dimensional quantum torus and then we give a realization as a covariant Lie algebra of the affine Lie algebra of \mathfrak{sl}_{∞} .

2.1. The extended affine Lie algebra $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$. Let $N \geq 2$ be a positive integer. For any unital associative algebra \mathcal{A} (over \mathbb{C}), we denote by $\mathfrak{gl}_N(\mathcal{A})$ the associative algebra of all $N \times N$ matrices with entries in \mathcal{A} . Naturally, $\mathfrak{gl}_N(\mathcal{A})$ is a Lie algebra with commutator as its Lie bracket. Note that $\mathfrak{gl}_N(\mathcal{A})$ is naturally a (left) \mathcal{A} -module (with each $a \in \mathcal{A}$ as a scalar). Set

(2.1)
$$\mathfrak{sl}_N(\mathcal{A}) = [\mathfrak{gl}_N(\mathcal{A}), \mathfrak{gl}_N(\mathcal{A})],$$

the derived Lie subalgebra. For $1 \le i, j \le N$, $a \in \mathcal{A}$, we also write $E_{i,j}a = aE_{i,j}$, the matrix whose only nonzero entry is the (i, j)-entry which is a.

Let q be a nonzero complex number, which is fixed throughout this paper. Denote by $\mathbb{C}_q[t_0^{\pm 1}, t_1^{\pm 1}]$ the 2-dimensional quantum torus associated to q. By definition,

$$\mathbb{C}_q[t_0^{\pm 1}, t_1^{\pm 1}] = \mathbb{C}[t_0^{\pm 1}, t_1^{\pm 1}]$$

as a vector space, where

$$(t_0^{m_0}t_1^{m_1})(t_0^{n_0}t_1^{n_1})=q^{m_1n_0}t_0^{m_0+n_0}t_1^{m_1+n_1}$$

for $m_0, m_1, n_0, n_1 \in \mathbb{Z}$. From now on, we simply write \mathbb{C}_q for $\mathbb{C}_q[t_0^{\pm 1}, t_1^{\pm 1}]$. Throughout this paper, we assume that q is not a root of unity. Note that this assumption amounts to that \mathbb{C}_q is a simple associative algebra.

Consider the following two-dimensional central extension of Lie algebra $\mathfrak{gl}_N(\mathbb{C}_q)$:

(2.2)
$$\widehat{\mathfrak{gl}_N}(\mathbb{C}_q) = \mathfrak{gl}_N(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{k}_0 \oplus \mathbb{C}\mathbf{k}_1,$$

where $\mathbf{k}_0, \mathbf{k}_1$ are (linearly independent) central elements, and

(2.3)
$$[E_{i,j}t_0^{m_0}t_1^{m_1}, E_{k,l}t_0^{n_0}t_1^{n_1}]$$

$$= \delta_{j,k}q^{m_1n_0}E_{i,l}t_0^{m_0+n_0}t_1^{m_1+n_1} - \delta_{i,l}q^{n_1m_0}E_{k,j}t_0^{m_0+n_0}t_1^{m_1+n_1}$$

$$+ \delta_{j,k}\delta_{i,l}\delta_{m_0+n_0,0}\delta_{m_1+n_1,0}q^{m_1n_0}(m_0\mathbf{k}_0 + m_1\mathbf{k}_1)$$

for $1 \leq i, j, k, l \leq N$ and for $m_0, m_1, n_0, n_1 \in \mathbb{Z}$.

It is clear that $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ is a $\mathbb{Z} \times \mathbb{Z}$ -graded Lie algebra with

$$\deg \mathbf{k}_0 = (0,0), \ \deg \mathbf{k}_1 = (0,0), \ \deg (E_{i,j} t_0^{m_0} t_1^{m_1}) = (m_0, m_1)$$

for $1 \leq i, j \leq N$, $m_0, m_1 \in \mathbb{Z}$. Define degree-zero derivations $\mathbf{d}_0, \mathbf{d}_1$ of $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ by

(2.4)
$$\mathbf{d}_r(E_{i,j}t_0^{m_0}t_1^{m_1}) = m_r E_{i,j}t_0^{m_0}t_1^{m_1}$$

for r = 0, 1 and for $1 \leq i, j \leq N, m_0, m_1 \in \mathbb{Z}$. Adding derivations $\mathbf{d}_0, \mathbf{d}_1$ to $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$, we obtain a Lie algebra

(2.5)
$$\widetilde{\mathfrak{gl}_N}(\mathbb{C}_q) = \widehat{\mathfrak{gl}_N}(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{d}_0 \oplus \mathbb{C}\mathbf{d}_1.$$

Set

$$(2.6) \widehat{\mathfrak{sl}_N}(\mathbb{C}_q) = [\widehat{\mathfrak{gl}_N}(\mathbb{C}_q), \widehat{\mathfrak{gl}_N}(\mathbb{C}_q)] = \mathfrak{sl}_N(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{k}_0 \oplus \mathbb{C}\mathbf{k}_1,$$

the derived subalgebra of $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$. It is known (see [BGK]) that $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ is a universal central extension of $\mathfrak{sl}_N(\mathbb{C}_q)$ and

(2.7)
$$\widehat{\mathfrak{gl}_N}(\mathbb{C}_q) = \widehat{\mathfrak{sl}_N}(\mathbb{C}_q) \oplus \mathbb{C}I_N,$$

where $I_N = E_{1,1} + \cdots + E_{N,N}$ (the identity matrix). Set

(2.8)
$$\widetilde{\mathfrak{sl}_N}(\mathbb{C}_q) = \widehat{\mathfrak{sl}_N}(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{d}_0 \oplus \mathbb{C}\mathbf{d}_1 \subset \widetilde{\mathfrak{gl}_N}(\mathbb{C}_q).$$

The Lie algebra $\mathfrak{sl}_N(\mathbb{C}_q)$ is known to be a nullity-2 extended affine Lie algebra of type A_{N-1} , where $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ is called the core of $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$. We refer the reader to [BGK] or [AABGP] for details.

In this paper, we shall focus on the subalgebra $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$. From [BGK] we have:

Lemma 2.1. The following elements form a basis of $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$:

(2.9)
$$E_{i,j}t_0^{m_0}t_1^{m_1}, \quad E_{r,r}-E_{r+1,r+1}, \quad \mathbf{k}_0, \quad \mathbf{k}_1,$$

where $1 \le i, j \le N$, $m_0, m_1 \in \mathbb{Z}$ with $(i - j, m_0, m_1) \ne (0, 0, 0)$ and $1 \le r \le N - 1$.

2.2. The (\mathbb{Z}, χ_q) -covariant algebra $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$ of $\widehat{\mathfrak{sl}_{\infty}}$. We start with the definition of a covariant algebra of an affine Lie algebra. Let \mathfrak{g} be a (possibly infinite-dimensional) Lie algebra equipped with a symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$. Associated to the pair $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$, we have an affine Lie algebra

$$\widehat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}\mathbf{k},$$

where **k** is a (nonzero) central element, and for $a, b \in \mathfrak{g}$, $m, n \in \mathbb{Z}$,

$$[a \otimes t^m, b \otimes t^n] = [a, b] \otimes t^{m+n} + m\delta_{m+n,0}\langle a, b\rangle \mathbf{k}.$$

Assume that Γ is a group acting on \mathfrak{g} as an automorphism group preserving the bilinear form $\langle \cdot, \cdot \rangle$ such that for $a, b \in \mathfrak{g}$,

(2.10)
$$[ga, b] = 0$$
 and $\langle ga, b \rangle = 0$ for all but finitely many $g \in \Gamma$.

Let $\chi:\Gamma\to\mathbb{C}^{\times}$ be any linear character. We lift the Γ -action from \mathfrak{g} to $\widehat{\mathfrak{g}}$ by

(2.11)
$$g(a \otimes t^m + \beta \mathbf{k}) = \chi(g)^m (ga \otimes t^m) + \beta \mathbf{k}$$

for $g \in \Gamma$, $a \in \mathfrak{g}$, $m \in \mathbb{Z}$, $\beta \in \mathbb{C}$. From [Li4], the (Γ, χ) -covariant algebra of $\widehat{\mathfrak{g}}$ is the Lie algebra $\widehat{\mathfrak{g}}[\Gamma]$, where

(2.12)
$$\widehat{\mathfrak{g}}[\Gamma] = \widehat{\mathfrak{g}}/\operatorname{span}\{gu - u \mid g \in \Gamma, u \in \widehat{\mathfrak{g}}\}\$$

as a vector space and the Lie bracket is given by

$$[\overline{u}, \overline{v}] = \sum_{q \in \Gamma} \overline{[gu, v]} \text{ for } u, v \in \widehat{\mathfrak{g}}.$$

Here and below, for $u \in \widehat{\mathfrak{g}}$, \overline{u} stands for the image of u in $\widehat{\mathfrak{g}}[\Gamma]$ under the natural quotient map $\widehat{\mathfrak{g}} \to \widehat{\mathfrak{g}}[\Gamma]$. Note that for $a, b \in \mathfrak{g}$, $m, n \in \mathbb{Z}$ and $g \in \Gamma$, we have

$$(2.13) \overline{a \otimes t^m} = \chi(g)^m \overline{ga \otimes t^m},$$

$$(2.14) \qquad [\overline{a \otimes t^m}, \overline{b \otimes t^n}] = \sum_{g \in \Gamma} \chi(g)^m \left(\overline{[ga, b] \otimes t^{m+n}} + m \delta_{m+n,0} \langle ga, b \rangle \mathbf{k} \right),$$

where we still denote $\overline{\mathbf{k}}$ by \mathbf{k} for simplicity.

The following is straightforward:

Lemma 2.2. Let \mathfrak{g} be a Lie algebra, Γ a group as above. Suppose that \mathfrak{g}_0 is a subalgebra of \mathfrak{g} which is stable under the action of Γ . Then the embedding map of \mathfrak{g}_0 into \mathfrak{g} gives rise to a Lie algebra homomorphism $\psi: \widehat{\mathfrak{g}}_0[\Gamma] \to \widehat{\mathfrak{g}}[\Gamma]$.

We also formulate a simple lemma which we shall use to determine a basis of the covariant Lie algebra $\widehat{\mathfrak{g}}[\Gamma]$. Let G be a group and let U be a G-module. Define U/G to be the quotient space of U modulo the subspace spanned by $\{gu-u\mid g\in G,\ u\in U\}$.

Lemma 2.3. Let G be a group, U a G-module. (a) If $U = \bigoplus_{\alpha \in S} U_{\alpha}$ as a G-module, then $U/G \simeq \bigoplus_{\alpha \in S} U_{\alpha}/G$. (b) Suppose that U has a basis B with a subset B_0 satisfying the condition that for every $b \in B$, there exist unique $g \in G$, $b_0 \in B_0$ such that $b \in \mathbb{C}gb_0$. Then B_0 gives rise to a basis for the quotient space U/G.

Proof. Part (a) follows immediately from a standard result in linear algebra. For Part (b), from the assumption, $\{g(b_0) \mid g \in G, b_0 \in B_0\}$ is also a basis of U. It then follows that $B_0 \cup \{g(b_0) - b_0 \mid g \in G, g \neq e, b_0 \in B_0\}$ is also a basis of U. Note that $g(hb_0) - hb_0 = (ghb_0 - b_0) - (hb_0 - b_0)$ for $g, h \in G$, $b_0 \in B_0$. Using this and using the basis $\{g(b_0) \mid g \in G, b_0 \in B_0\}$ of U, we see that $\{g(b_0) - b_0 \mid g \in G, g \neq e, b_0 \in B_0\}$ is a basis of the span of $\{gu - u \mid g \in G, u \in U\}$. Consequently, B_0 gives rise to a basis for the quotient space U/G.

Let \mathfrak{gl}_{∞} be the associative algebra of all doubly infinite complex matrices with only finitely many nonzero entries, which is also naturally a Lie algebra. As before,

for $m, n \in \mathbb{Z}$, let $E_{m,n}$ denote the unit matrix whose only nonzero entry is the (m, n)-entry which is 1. Equip \mathfrak{gl}_{∞} with the bilinear form $\langle \cdot, \cdot \rangle$ defined by

$$\langle E_{i,j}, E_{k,l} \rangle = \delta_{j,k} \, \delta_{i,l} \quad \text{for } i, j, k, l \in \mathbb{Z},$$

which is nondegenerate, symmetric and (associative) invariant. Set

$$\mathfrak{sl}_{\infty} = [\mathfrak{gl}_{\infty}, \mathfrak{gl}_{\infty}],$$

the derived subalgebra of \mathfrak{gl}_{∞} . We see that $\langle \cdot, \cdot \rangle$ is also nondegenerate on \mathfrak{sl}_{∞} . Then we have the affine Lie algebra $\widehat{\mathfrak{sl}_{\infty}}$ associated to the pair $(\mathfrak{sl}_{\infty}, \langle \cdot, \cdot \rangle)$.

Definition 2.4. Let σ be the automorphism of the algebra \mathfrak{gl}_{∞} defined by

(2.16)
$$\sigma(E_{m,n}) = E_{m+1,n+1} \quad \text{for } m, n \in \mathbb{Z}.$$

The following is straightforward:

Lemma 2.5. The automorphism σ of \mathfrak{gl}_{∞} preserves the bilinear form $\langle \cdot, \cdot \rangle$ and the Lie subalgebra \mathfrak{sl}_{∞} . Furthermore, for any positive integer N, the map

(2.17)
$$\rho_N : \mathbb{Z} \to \operatorname{Aut}(\mathfrak{gl}_{\infty}) \text{ (resp. } \operatorname{Aut}(\mathfrak{sl}_{\infty})), \quad r \mapsto \sigma^{Nr} \quad (r \in \mathbb{Z})$$

is a one-to-one group homomorphism, and for any $a, b \in \mathfrak{gl}_{\infty}$ (resp. \mathfrak{sl}_{∞}),

$$[\sigma^r(a), b] = 0$$
 and $\langle \sigma^r(a), b \rangle = 0$ for all but finitely many $r \in \mathbb{Z}$.

From now on, we fix the group actions of \mathbb{Z} on \mathfrak{gl}_{∞} and \mathfrak{sl}_{∞} via the homomorphism ρ_N . Define a linear character $\chi_q: \mathbb{Z} \to \mathbb{C}^{\times}$ by

(2.18)
$$\chi_q(r) = q^r \quad \text{for } r \in \mathbb{Z}.$$

Then we have the (\mathbb{Z}, χ_q) -covariant algebras $\widehat{\mathfrak{gl}_{\infty}}[\mathbb{Z}]$ and $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$. In what follows, we shall show that Lie algebra $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ is isomorphic to $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$.

Define a linear map $\theta_{N,q}: \widehat{\mathfrak{gl}_N}(\mathbb{C}_q) \to \widehat{\mathfrak{gl}_\infty}[\mathbb{Z}]$ by

(2.19)
$$\theta_{N,q}(E_{i,j}t_0^{m_0}t_1^{m_1}) = \overline{E_{Nm_1+i,j} \otimes t^{m_0}}, \quad \theta_{N,q}(\mathbf{k}_0) = \mathbf{k}, \quad \theta_{N,q}(\mathbf{k}_1) = 0$$
 for $1 \le i, j \le N, m_0, m_1 \in \mathbb{Z}$. Then we have:

Proposition 2.6. The linear map $\theta_{N,q}: \widehat{\mathfrak{gl}_N}(\mathbb{C}_q) \to \widehat{\mathfrak{gl}_\infty}[\mathbb{Z}]$ is a surjective Lie homomorphism with $\ker(\theta_{N,q}) = \mathbb{C}\mathbf{k}_1$.

Proof. By Lemma 2.3, we see that the central element \mathbf{k} together with the elements

$$\overline{E_{Nm_1+i,j} \otimes t^{m_0}}$$
 for $1 \le i, j \le N, m_0, m_1 \in \mathbb{Z}$,

form a basis of $\widehat{\mathfrak{gl}_{\infty}}[\mathbb{Z}]$. Thus $\theta_{N,q}$ is a surjective linear map with $\ker(\theta_{N,q}) = \mathbb{C}\mathbf{k}_1$. Let $1 \leq i, j, k, l \leq N$ and $m_0, m_1, n_0, n_1 \in \mathbb{Z}$. By definition (see (2.14)) we have

$$\overline{\left[E_{Nm_1+i,j} \otimes t^{m_0}, \overline{E_{Nn_1+k,l} \otimes t^{n_0}}\right]} \\
= \sum_{r \in \mathbb{Z}} \chi_q(r)^{m_0} \left(\overline{\left[E_{N(m_1+r)+i,Nr+j} \otimes t^{m_0}, E_{Nn_1+k,l} \otimes t^{n_0}\right]}\right)$$

$$+ m_{0}\delta_{m_{0}+n_{0},0}\langle E_{N(m_{1}+r)+i,Nr+j}, E_{Nn_{1}+k,l}\rangle\mathbf{k})$$

$$= \delta_{j,k}q^{n_{1}m_{0}}\overline{E_{N(m_{1}+n_{1})+i,l}\otimes t^{m_{0}+n_{0}}} - \delta_{i,l}q^{-m_{1}m_{0}}\overline{E_{Nn_{1}+k,-Nm_{1}+j}\otimes t^{m_{0}+n_{0}}}$$

$$+ m_{0}q^{m_{0}n_{1}}\delta_{m_{0}+n_{0},0}\delta_{m_{1}+n_{1},0}\delta_{j,k}\delta_{i,l}\mathbf{k}$$

$$= \delta_{j,k}q^{n_{1}m_{0}}\overline{E_{N(m_{1}+n_{1})+i,l}\otimes t^{m_{0}+n_{0}}} - \delta_{i,l}q^{m_{1}n_{0}}\overline{E_{N(m_{1}+n_{1})+k,j}\otimes t^{m_{0}+n_{0}}}$$

$$+ m_{0}q^{m_{0}n_{1}}\delta_{m_{0}+n_{0},0}\delta_{m_{1}+n_{1},0}\delta_{j,k}\delta_{i,l}\mathbf{k},$$

noticing that (see (2.13) and (2.18))

$$\overline{E_{Nn_1+k,-Nm_1+j} \otimes t^{m_0+n_0}} = \chi_q(m_1)^{m_0+n_0} \overline{\sigma_{N,m_1}(E_{Nn_1+k,-Nm_1+j}) \otimes t^{m_0+n_0}}$$

$$= q^{(m_0+n_0)m_1} \overline{E_{N(n_1+m_1)+k,j} \otimes t^{m_0+n_0}}.$$

It then follows from (2.3) and (2.20) that $\theta_{N,q}$ is a Lie algebra homomorphism. \square

Now, we continue to study Lie algebra $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$. For $1 \leq i, j \leq N$, $m_0, m_1 \in \mathbb{Z}$ with $(i, m_0, m_1) \neq (j, 0, 0)$, set

$$e_{i,j}(m_0, m_1) = \overline{E_{Nm_1+i,j} \otimes t^{m_0}} \quad \text{if } (i, m_1) \neq (j, 0),$$

$$e_{i,i}(m_0, 0) = \frac{1}{1 - q^{-m_0}} \overline{(E_{i,i} - E_{N+i,N+i}) \otimes t^{m_0}} \quad \text{if } m_0 \neq 0.$$

On the other hand, set

$$\mathbf{k}' = \overline{E_{N+1,N+1} - E_{1,1}}$$
 and $\overline{h_r} = \overline{E_{r,r} - E_{r+1,r+1}}$ for $1 \le r \le N - 1$.

Then we have:

Lemma 2.7. The following relations hold for $1 \le i, j \le N$, $m_0, m_1, n_1 \in \mathbb{Z}$ with $(i, m_1) \ne (j, n_1)$:

(2.21)
$$\overline{E_{Nm_1+i,Nn_1+j} \otimes t^{m_0}} = q^{-m_0n_1}e_{i,j}(m_0, m_1 - n_1),$$

(2.22)
$$\overline{(E_{Nm_1+i,Nm_1+i} - E_{Nn_1+j,Nn_1+j}) \otimes t^{m_0}} = \begin{cases}
q^{-m_0m_1}e_{i,i}(m_0,0) - q^{-m_0n_1}e_{j,j}(m_0,0) & \text{if } m_0 \neq 0 \\
\overline{E_{i,i} - E_{j,j}} + (m_1 - n_1)\mathbf{k'} & \text{if } m_0 = 0.
\end{cases}$$

Furthermore, the vectors

(2.23)
$$e_{i,j}(m_0, m_1), \quad \overline{h_r}, \quad \mathbf{k}', \quad \mathbf{k},$$

where $1 \leq i, j \leq N$, $m_0, m_1 \in \mathbb{Z}$ with $(i - j, m_0, m_1) \neq (0, 0, 0)$ and $1 \leq r \leq N - 1$, form a basis of $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$.

Proof. From definition (see (2.13)) we get (2.21) as

$$\overline{E_{Nm_1+i,Nn_1+j} \otimes t^{m_0}} = \chi_q(-n_1)^{m_0} \overline{\sigma^{N(-n_1)}(E_{Nm_1+i,Nn_1+j})} \otimes t^{m_0} \\
= q^{-m_0n_1} \overline{E_{N(m_1-n_1)+i,j} \otimes t^{m_0}} = q^{-m_0n_1} e_{i,j}(m_0, m_1 - n_1).$$

Assume $m_0 \neq 0$. For convenience, define $e_{r,r}(m_0,0)$ for any $r \in \mathbb{Z}$ in the same way:

$$e_{r,r}(m_0,0) = \frac{1}{1 - q^{-m_0}} \overline{(E_{r,r} - E_{N+r,N+r}) \otimes t^{m_0}}.$$

Note that

$$e_{r+kN,r+kN}(m_0,0) = q^{-km_0}e_{r,r}(m_0,0)$$

for any $r, k \in \mathbb{Z}$. Using this we get

$$(1 - q^{-m_0})\overline{(E_{m,m} - E_{n,n}) \otimes t^{m_0}}$$

$$= \left(\overline{(E_{m,m} - E_{n,n}) \otimes t^{m_0}} - \overline{(E_{N+m,N+m} - E_{N+n,N+n}) \otimes t^{m_0}}\right)$$

$$= \overline{(E_{m,m} - E_{N+m,N+m}) \otimes t^{m_0} - (E_{n,n} - E_{N+n,N+n}) \otimes t^{m_0}}$$

$$= (1 - q^{-m_0}) \left(e_{m,m}(m_0, 0) - e_{n,n}(m_0, 0)\right),$$

which gives

$$\overline{(E_{m,m} - E_{n,n}) \otimes t^{m_0}} = e_{m,m}(m_0, 0) - e_{n,n}(m_0, 0)$$

for any $m, n \in \mathbb{Z}$. Then we have

$$\overline{(E_{Nm_1+i,Nm_1+i} - E_{Nn_1+j,Nn_1+j}) \otimes t^{m_0}}
= e_{Nm_1+i,Nm_1+i}(m_0,0) - e_{Nn_1+j,Nn_1+j}(m_0,0)
= q^{-m_1m_0}e_{i,i}(m_0,0) - q^{-n_1m_0}e_{j,j}(m_0,0).$$

Now, assume $m_0 = 0$. For any integer r, as

$$\overline{E_{N+1,N+1} - E_{1,1}} - \overline{E_{N+r,N+r} - E_{r,r}} = \overline{\sigma_{N,1}(E_{1,1} - E_{r,r}) - (E_{1,1} - E_{r,r})} = 0,$$

we have

$$\overline{E_{N+r,N+r} - E_{r,r}} = \overline{E_{N+1,N+1} - E_{1,1}} = \mathbf{k}'.$$

It follows that

$$\overline{E_{nN+i,nN+i} - E_{i,i}} = n \mathbf{k'} \quad \text{for all } 1 \le i \le N, \ n \in \mathbb{Z}.$$

Then we have

$$\overline{E_{Nm_1+i,Nm_1+i} - E_{Nn_1+j,Nn_1+j}}
= \overline{E_{Nm_1+i,Nm_1+i} - E_{i,i}} + \overline{E_{i,i} - E_{j,j}} + \overline{E_{j,j} - E_{Nn_1+j,Nn_1+j}}
= \overline{E_{i,i} - E_{j,j}} + (m_1 - n_1)\mathbf{k}'.$$

This proves the first assertion.

Now, we prove the basis assertion. Note that matrices $E_{m,n}$ and $E_{m,m}-E_{m+1,m+1}$ for $m,n\in\mathbb{Z}$ with $m\neq n$ form a basis of \mathfrak{sl}_{∞} . Then the elements

$$E_{m,n}\otimes t^{m_0}, \quad (E_{m,m}-E_{m+1,m+1})\otimes t^{m_0}$$

for $m, n, m_0 \in \mathbb{Z}$ with $m \neq n$ together with **k** form a basis of $\widehat{\mathfrak{sl}}_{\infty}$. Recall that for $r \in \mathbb{Z}$, $\rho_N(r) = (\sigma^N)^r$ on \mathfrak{sl}_{∞} is the restriction of the automorphism of \mathfrak{gl}_{∞} defined by $\sigma^{Nr}(E_{m,n}) = E_{m+rN,n+rN}$ for $m, n \in \mathbb{Z}$ and that

$$\rho_N(r)(E_{m,n}\otimes t^{m_0})=\chi_q(r)^{m_0}E_{m+rN,n+rN}\otimes t^{m_0}=q^{rm_0}E_{m+rN,n+rN}\otimes t^{m_0}$$

for $m, n, m_0 \in \mathbb{Z}$. It then follows from Lemma 2.3 that the elements

$$\overline{E_{mN+i,j}\otimes t^{m_0}}, \overline{(E_{i,i}-E_{i+1,i+1})\otimes t^{m_0}}$$

for $1 \leq i, j \leq N$, $m, m_0 \in \mathbb{Z}$ with $(i, m) \neq (j, 0)$ together with **k** form a basis of $\widehat{\mathfrak{sl}}_{\infty}[\mathbb{Z}]$. When $m_0 = 0$, it is clear that $\{\overline{h_1}, \ldots, \overline{h_{N-1}}, \mathbf{k'}\}$ is also a basis for the subspace spanned by $(E_{i,i} - E_{i+1,i+1}) \otimes t^{m_0}$ for $1 \leq i \leq N$.

Let m_0 be any nonzero integer. From the first assertion, the N vectors

(2.24)
$$\overline{(E_{i,i} - E_{N+i,N+i}) \otimes t^{m_0}} \quad \text{for } 1 \le i \le N$$

linearly span the N-dimensional subspace spanned by

(2.25)
$$\overline{(E_{i,i} - E_{i+1,i+1}) \otimes t^{m_0}}$$
 for $1 \le i \le N$.

Consequently, the N vectors in (2.24) also form a basis for the subspace spanned by the vectors in (2.25). Therefore,

$$\overline{E_{mN+i,j}\otimes t^{m_0}}, \quad \overline{(E_{i,i}-E_{N+i,N+i})\otimes t^{m_0}}$$

for $1 \leq i, j \leq N$, $m, m_0 \in \mathbb{Z}$ with $(i, m) \neq (j, 0)$ together with \mathbf{k}' and \mathbf{k} form a basis of $\widehat{\mathfrak{sl}}_{\infty}[\mathbb{Z}]$. Now, the proof is complete.

As the main result of this section, we have:

Theorem 2.8. The Lie algebra $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ is isomorphic to the covariant algebra $\widehat{\mathfrak{sl}_\infty}[\mathbb{Z}]$, where an isomorphism $\theta: \widehat{\mathfrak{sl}_N}(\mathbb{C}_q) \to \widehat{\mathfrak{sl}_\infty}[\mathbb{Z}]$ is given by

$$\theta(\mathbf{k}_0) = \mathbf{k}, \quad \theta(\mathbf{k}_1) = \mathbf{k}',$$

$$\theta(E_{i,i}t_0^{m_0}t_1^{m_1}) = e_{i,i}(m_0, m_1), \quad \theta(E_{r,r} - E_{r+1,r+1}) = \overline{E_{r,r} - E_{r+1,r+1}}$$

for
$$1 \le i, j \le N$$
, $m_0, m_1 \in \mathbb{Z}$ with $(i - j, m_0, m_1) \ne (0, 0, 0)$ and $1 \le r \le N - 1$.

Proof. In view of Lemma 2.7, the linear map θ is a linear isomorphism from $\mathfrak{sl}_N(\mathbb{C}_q)$ onto $\widehat{\mathfrak{sl}_\infty}[\mathbb{Z}]$, so it remains to prove that θ is a Lie algebra homomorphism. Recall from Proposition 2.6 that we have a Lie algebra homomorphism $\theta_{N,q}:\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)\to\widehat{\mathfrak{gl}_\infty}[\mathbb{Z}]$. This gives a Lie algebra homomorphism from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to $\widehat{\mathfrak{gl}_\infty}[\mathbb{Z}]$, also denoted by $\theta_{N,q}$, such that $\ker(\theta_{N,q}) = \mathbb{C}\mathbf{k}_1$.

On the other hand, by Lemma 2.3 the Lie algebra embedding of \mathfrak{sl}_{∞} into \mathfrak{gl}_{∞} gives rise to a Lie algebra homomorphism $\psi: \widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}] \to \widehat{\mathfrak{gl}_{\infty}}[\mathbb{Z}]$. Notice that the following relations hold in $\widehat{\mathfrak{gl}_{\infty}}[\mathbb{Z}]$:

$$\overline{(E_{i,i} - E_{N+i,N+i}) \otimes t^{m_0}} = (1 - q^{-m_0}) \overline{(E_{i,i} \otimes t^{m_0})},$$

$$\overline{E_{N+1,N+1} - E_{1,1}} = 0$$
 as $\overline{E_{N+j,N+j} \otimes t^{m_0}} = q^{-m_0} \overline{E_{j,j} \otimes t^{m_0}}$ for $1 \leq j \leq N$. Thus
$$\psi(e_{i,i}(m_0,0)) = \overline{E_{i,i} \otimes t^{m_0}} \quad \text{for } 1 \leq i \leq N, \ m_0 \neq 0$$

and $\psi(\mathbf{k}') = 0$. It follows that $\ker(\psi) = \mathbb{C}\mathbf{k}'$ and $\theta_{N,q}(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)) = \psi(\widehat{\mathfrak{sl}_\infty}[\mathbb{Z}])$. Consequently, θ reduces to a Lie algebra isomorphism from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)/\mathbb{C}\mathbf{k}_1$ onto $\widehat{\mathfrak{sl}_\infty}[\mathbb{Z}]/\mathbb{C}\mathbf{k}'$. To show that θ is a Lie algebra homomorphism, it suffices to check the central extensions involving \mathbf{k}_1 and \mathbf{k}' , respectively. From the Lie commutator relation (2.3), we only need to consider $[E_{i,j}t_0^{m_0}t_1^{m_1}, E_{k,l}t_0^{n_0}t_1^{n_1}]$ in $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ with $\delta_{j,k}\delta_{i,l}\delta_{m_0+n_0,0}\delta_{m_1+n_1,0} = 1$ and $m_1 \neq 0$.

Let $1 \leq i, j \leq N$, $m_0, m_1 \in \mathbb{Z}$ with $m_1 \neq 0$. From (2.3) we have

$$[E_{i,j}t_0^{m_0}t_1^{m_1}, E_{j,i}t_0^{-m_0}t_1^{-m_1}] = q^{-m_1m_0}(E_{i,i} - E_{j,j} + m_0\mathbf{k}_0 + m_1\mathbf{k}_1).$$

On the other hand, by definition (see (2.14)) we have

$$\begin{aligned} & [\overline{E_{Nm_1+i,j}} \otimes t^{m_0}, \overline{E_{-Nm_1+j,i}} \otimes t^{-m_0}] \\ &= \sum_{r \in \mathbb{Z}} \chi_q(r)^{m_0} \Big(\overline{[E_{N(m_1+r)+i,Nr+j}} \otimes t^{m_0}, E_{-Nm_1+j,i} \otimes t^{-m_0}] \\ & + m_0 \delta_{m_0+n_0,0} \langle E_{N(m_1+r)+i,Nr+j}, E_{-Nm_1+j,i} \rangle \mathbf{k} \Big) \\ &= \overline{q^{-m_1 m_0} E_{i,i}} - q^{-m_1 m_0} \overline{E_{-Nm_1+j,-Nm_1+j}} + m_0 q^{-m_0 m_1} \mathbf{k} \\ &= q^{-m_1 m_0} \left(\overline{E_{i,i}} - \overline{E_{j,j}} + m_1 \mathbf{k}' \right) + m_0 q^{-m_0 m_1} \mathbf{k}, \end{aligned}$$

where we are using (2.22) for the last equality. We see that the central extension by \mathbf{k}_1 matches the central extension by \mathbf{k}' . Therefore, θ is a Lie algebra isomorphism from $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ to $\widehat{\mathfrak{sl}_\infty}[\mathbb{Z}]$.

3. Equivariant quasi modules for simple vertex algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$

In this section, first we recall from [Li3] the notion of vertex Γ -algebra and the notion of equivariant quasi module for a vertex Γ -algebra. Then, as the main result of this section, we prove that for any nonnegative integer ℓ , the category of equivariant quasi modules for the simple vertex algebra $\widehat{\mathfrak{sl}_{\infty}}(\ell,0)$ associated to the affine Lie algebra $\widehat{\mathfrak{sl}_{\infty}}$ is canonically isomorphic to that of integrable restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ .

3.1. Vertex Γ -algebras and their equivariant quasi modules. We start with the notion of vertex Γ -algebra introduced in [Li3].

Definition 3.1. Let Γ be a group. A *vertex* Γ -algebra is a vertex algebra V equipped with two group homomorphisms

$$R: \Gamma \to \mathrm{GL}(V); g \mapsto R_g,$$

$$\chi:\Gamma\to\mathbb{C}^{\times}$$
,

satisfying the condition that $R_g(1) = 1$ and

(3.1)
$$R_q Y(v, x) R_q^{-1} = Y(R_q(v), \chi(g)^{-1} x) \text{ for } g \in \Gamma, \ v \in V.$$

Remark 3.2. Let V be a \mathbb{Z} -graded vertex algebra in the sense that V is a vertex algebra equipped with a \mathbb{Z} -grading $V = \bigoplus_{n \in \mathbb{Z}} V_{(n)}$ such that $\mathbf{1} \in V_{(0)}$ and

$$(3.2) u_m V_{(n)} \subset V_{(n+k-m-1)} \text{for } u \in V_{(k)}, m, n, k \in \mathbb{Z}.$$

Define a linear operator L(0) on V by L(0)v = nv for $v \in V_{(n)}$ with $n \in \mathbb{Z}$. Assume that Γ is an automorphism group of V as a \mathbb{Z} -graded vertex algebra and let $\chi : \Gamma \to \mathbb{C}^{\times}$ be any linear character. For $g \in \Gamma$, set $R_g = \chi(g)^{-L(0)}g$. Then V with group homomorphisms R and χ is a vertex Γ -algebra (see [Li3]).

Example 3.3. Here we recall the vertex algebras associated to affine Lie algebras. Let \mathfrak{g} be a Lie algebra equipped with a nondegenerate symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$. Equip the affine Lie algebra $\widehat{\mathfrak{g}}$ with a \mathbb{Z} -grading $\widehat{\mathfrak{g}} = \bigoplus_{n \in \mathbb{Z}} \widehat{\mathfrak{g}}_{(n)}$, where

(3.3)
$$\widehat{\mathfrak{g}}_{(0)} = \mathfrak{g} \oplus \mathbb{C} \mathbf{k} \text{ and } \widehat{\mathfrak{g}}_{(n)} = \mathfrak{g} \otimes \mathbb{C} t^{-n} \text{ for } n \neq 0.$$

Let ℓ be a complex number. View \mathbb{C} as a $(\mathfrak{g} \otimes \mathbb{C}[t] \oplus \mathbb{C}\mathbf{k})$ -module with $\mathfrak{g} \otimes \mathbb{C}[t]$ acting trivially and with \mathbf{k} acting as scalar ℓ . Form the induced $\widehat{\mathfrak{g}}$ -module

$$(3.4) V_{\widehat{\mathfrak{g}}}(\ell,0) = \mathcal{U}(\widehat{\mathfrak{g}}) \otimes_{\mathcal{U}(\mathfrak{g} \otimes \mathbb{C}[t] \oplus \mathbb{C}\mathbf{k})} \mathbb{C},$$

which is naturally N-graded by defining deg $\mathbb{C} = 0$. Set $\mathbf{1} = 1 \otimes 1 \in V_{\widehat{\mathfrak{g}}}(\ell, 0)$ and identify \mathfrak{g} as the degree-one subspace of $V_{\widehat{\mathfrak{g}}}(\ell, 0)$ through the linear map

$$a \mapsto a(-1)\mathbf{1} \in V_{\widehat{\mathfrak{g}}}(\ell,0) \quad \text{for } a \in \mathfrak{g}.$$

Following the standard practice, we alternatively write a(m) for $a \otimes t^m \in \widehat{\mathfrak{g}}$. It was known (cf. [FZ]) that there exists a vertex algebra structure on $V_{\widehat{\mathfrak{g}}}(\ell,0)$, which is uniquely determined by the condition that **1** is the vacuum vector and

$$Y(a,x) = a(x) := \sum_{m \in \mathbb{Z}} a(m)x^{-m-1}$$
 for $a \in \mathfrak{g}$.

Denote by $J_{\widehat{\mathfrak{g}}}(\ell,0)$ the unique maximal graded $\widehat{\mathfrak{g}}$ -submodule of $V_{\widehat{\mathfrak{g}}}(\ell,0)$. Then $J_{\widehat{\mathfrak{g}}}(\ell,0)$ is a (two-sided) ideal of vertex algebra $V_{\widehat{\mathfrak{g}}}(\ell,0)$. Define

(3.5)
$$L_{\widehat{\mathfrak{g}}}(\ell,0) = V_{\widehat{\mathfrak{g}}}(\ell,0)/J_{\widehat{\mathfrak{g}}}(\ell,0),$$

which is a simple \mathbb{Z} -graded vertex algebra. One can show that the vertex algebra $L_{\widehat{\mathfrak{g}}}(\ell,0)$ is simple. Assume that Γ is a group which acts on \mathfrak{g} as an automorphism group preserving the bilinear form. It is a simple fact that the action of Γ on \mathfrak{g} can be uniquely lifted to an action on $V_{\widehat{\mathfrak{g}}}(\ell,0)$ where Γ acts as an automorphism group preserving the \mathbb{Z} -grading. As $J_{\widehat{\mathfrak{g}}}(\ell,0)$ is Γ -stable, Γ naturally acts on $L_{\widehat{\mathfrak{g}}}(\ell,0)$. From Remark 3.2, for any linear character χ of Γ , $V_{\widehat{\mathfrak{g}}}(\ell,0)$ and $L_{\widehat{\mathfrak{g}}}(\ell,0)$ are vertex Γ -algebras with $R_g = \chi(g)^{-L(0)}g$ for $g \in \Gamma$.

The following notion was introduced in [Li4]:

Definition 3.4. Let V be a vertex Γ -algebra. An equivariant quasi V-module is a vector space W equipped with a linear map

$$Y_W(\cdot, x): V \to \operatorname{Hom}(W, W((x))) \subset (\operatorname{End}W)[[x, x^{-1}]]$$

 $v \mapsto Y_W(v, x),$

satisfying the conditions that

$$(3.6) Y_W(\mathbf{1}, x) = 1_W (the identity operator on W),$$

$$(3.7) Y_W(R_q v, x) = Y_W(v, \chi(g)x) \text{for } g \in \Gamma, \ v \in V$$

and that for $u, v \in V$, the quasi Jacobi identity

$$x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)p(x_1,x_2)Y_W(u,x_1)Y_W(v,x_2)$$

$$-x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)p(x_1,x_2)Y_W(v,x_2)Y_W(u,x_1)$$

$$= x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)p(x_1,x_2)Y_W(Y(u,x_0)v,x_2)$$
(3.8)

holds on W for some polynomial $p(x_1, x_2)$ of the form

(3.9)
$$p(x_1, x_2) = (x_1 - \chi(g_1)x_2) \cdots (x_1 - \chi(g_k)x_2),$$

where $g_1, \ldots, g_k \in \Gamma$.

To emphasize the dependence on the group Γ and the linear character χ , we shall also use the term " (Γ, χ) -equivariant quasi V-module." Assume that $V = \bigoplus_{n \in \mathbb{Z}} V_{(n)}$ is also \mathbb{Z} -graded. A \mathbb{Z} -graded equivariant quasi V-module is an equivariant quasi V-module W with a \mathbb{Z} -grading $W = \bigoplus_{n \in \mathbb{Z}} W_{(n)}$ such that

(3.10)
$$u_m W_{(n)} \subset W_{(n+k-m-1)} \text{ for } u \in V_{(k)}, \ m, n, k \in \mathbb{Z}.$$

Recall the (Γ, χ) -covariant algebra $\widehat{\mathfrak{g}}[\Gamma]$ of the affine Lie algebra $\widehat{\mathfrak{g}}$ from Section 2.2. The following result was obtained in [Li4]:

Proposition 3.5. Let \mathfrak{g}, Γ be given as in Example 3.3, let $\chi : \Gamma \to \mathbb{C}^{\times}$ be a one-to-one group homomorphism, and let $\ell \in \mathbb{C}$. Then for any restricted $\widehat{\mathfrak{g}}[\Gamma]$ -module W of level ℓ , there is a (Γ, χ) -equivariant quasi $V_{\widehat{\mathfrak{g}}}(\ell, 0)$ -module structure $Y_W(\cdot, x)$ which is uniquely determined by

$$Y_W(a,x) = \bar{a}(x) := \sum_{n \in \mathbb{Z}} \overline{a(n)} x^{-n-1}$$
 for $a \in \mathfrak{g}$.

On the other hand, any (Γ, χ) -equivariant quasi $V_{\widehat{\mathfrak{g}}}(\ell, 0)$ -module W is a restricted $\widehat{\mathfrak{g}}[\Gamma]$ -module of level ℓ with $\overline{a}(x) = Y_W(a, x)$ for $a \in \mathfrak{g}$.

Remark 3.6. Let V_1, \ldots, V_r be vertex Γ -algebras with the same linear character χ . It is straightforward to see that the tensor product vertex algebra $V_1 \otimes \cdots \otimes V_r$

(see [FHL]) is also a vertex Γ -algebra. On the other hand, let W_i be a (Γ, χ) -equivariant quasi V_i -module for $1 \leq i \leq r$. Then $W_1 \otimes \cdots \otimes W_r$ is an equivariant quasi $V_1 \otimes \cdots \otimes V_r$ -module with

$$Y_{W_1 \otimes \cdots \otimes W_r}(v_1 \otimes \cdots \otimes v_r, x) = Y_{W_1}(v_1, x) \otimes \cdots \otimes Y_{W_r}(v_r, x)$$
 for $v_i \in V_i$.

3.2. Equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -modules. Recall from Section 2.2 that $\mathbb Z$ acts on \mathfrak{sl}_{∞} via the representation ρ_N (see (2.17)) and χ_q is the linear character of $\mathbb Z$ defined by $\chi_q(n)=q^n$ for $n\in\mathbb Z$. From Example 3.3, for any complex number ℓ we have $\mathbb Z$ -graded vertex $\mathbb Z$ -algebras $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ and $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. To emphasize the dependence on the $\mathbb Z$ -action via ρ_N , we shall also denote the vertex $\mathbb Z$ -algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ by $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N)$. In this subsection, for any nonnegative integer ℓ , we give a characterization of $(\mathbb Z,\chi_q)$ -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -modules in terms of integrable restricted $\widehat{\mathfrak{sl}_N}(\mathbb C_q)$ -modules of level ℓ .

Definition 3.7. An $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module W is said to be *restricted* if for any $w \in W$ and for $1 \leq i, j \leq N, \ m_0, m_1 \in \mathbb{Z}$ with $(i - j, m_0, m_1) \neq (0, 0, 0)$,

$$(E_{i,j}t_0^{m_0}t_1^{m_1})w=0$$
 for m_0 sufficiently large,

and it is said to be of level $\ell \in \mathbb{C}$ if \mathbf{k}_0 acts as scalar ℓ .

For $1 \leq i, j \leq N$, $m \in \mathbb{Z}$ with $(i - j, m) \neq (0, 0)$, we form a generating function

(3.11)
$$(E_{i,j}t_1^m)(x) = \sum_{n \in \mathbb{Z}} (E_{i,j}t_0^n t_1^m) x^{-n-1} \in \widehat{\mathfrak{sl}_N}(\mathbb{C}_q)[[x, x^{-1}]],$$

and we form generating functions

(3.12)
$$H_i(x) = \sum_{n \in \mathbb{Z}} (E_{i,i} - E_{i+1,i+1}) t_0^n x^{-n-1} \quad \text{for } 1 \le i \le N-1,$$

(3.13)
$$H_N(x) = \sum_{n \in \mathbb{Z}} (E_{N,N} t_0^n - q^{-n} E_{1,1} t_0^n - \delta_{n,0} \mathbf{k}_1) x^{-n-1}.$$

It follows from Lemma 2.1 that all the components of these generating functions together with \mathbf{k}_0 form a basis of $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$.

The following result gives an isomorphism between the category of (\mathbb{Z}, χ_q) equivariant quasi $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -modules and the category of restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules
of level ℓ (cf. [Li4]):

Proposition 3.8. Let ℓ be any complex number. For any restricted $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ module W of level ℓ , there exists a (\mathbb{Z}, χ_q) -equivariant quasi $V_{\widehat{\mathfrak{sl}}_{\infty}}(\ell, 0)$ -module
structure $Y_W(\cdot, x)$ on W, which is uniquely determined by

$$Y_W(E_{mN+i,nN+j}, x) = q^n \left(E_{i,j} t_1^{m-n} \right) (q^n x),$$

$$Y_W(E_{nN+i,nN+i} - E_{nN+i+1,nN+i+1}, x) = q^n H_i(q^n x),$$

for $1 \leq i, j \leq N, m, n \in \mathbb{Z}$ with $(i, m) \neq (j, n)$. On the other hand, for any (\mathbb{Z}, χ_q) equivariant quasi $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module (W, Y_W) , W is a restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module of
level ℓ with the action uniquely determined by

$$(E_{i,j}t_1^m)(x) = Y_W(E_{mN+i,j}, x), \quad H_i(x) = Y_W(E_{i,i} - E_{i+1,i+1}, x)$$

for $1 \le i, j \le N, \ m \in \mathbb{Z}$ with $(i - j, m) \ne (0, 0)$.

Proof. Let W be a restricted $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module of level ℓ . It follows from Proposition 2.8 that W is a restricted $\widehat{\mathfrak{sl}}_{\infty}[\mathbb{Z}]$ -module of level ℓ on which

$$\frac{\overline{E_{mN+i,j} \otimes t^n} = E_{i,j} t_0^n t_1^m, \ \overline{(E_{k,k} - E_{k+1,k+1}) \otimes t^n} = (E_{k,k} - E_{k+1,k+1}) t_0^n,
\overline{(E_{N,N} - E_{N+1,N+1}) \otimes t^n} = E_{N,N} t_0^n - q^{-n} E_{1,1} t_0^n - \delta_{n,0} \mathbf{k}_1,$$

where $1 \leq i, j \leq N, 1 \leq k \leq N-1$ and $m, n \in \mathbb{Z}$ with $(i-j, m) \neq (0, 0)$. As χ_q is injective, from Proposition 3.5, the $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$ -module W is naturally a (\mathbb{Z}, χ_q) -equivariant quasi $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module with

(3.14)
$$Y_W(a,x) = \bar{a}(x) = \sum_{n \in \mathbb{Z}} \overline{a(n)} x^{-n-1} \quad \text{for } a \in \mathfrak{sl}_{\infty}.$$

Then (\mathbb{Z}, χ_q) -equivariant quasi $V_{\widehat{\mathfrak{sl}}_{\infty}}(\ell, 0)$ -module structure $Y_W(\cdot, x)$ on W is uniquely determined by

$$Y_{W}(E_{mN+i,nN+j},x) = \overline{E_{mN+i,nN+j}}(x) = q^{n} \overline{E_{(m-n)N+i,j}}(q^{n}x) = q^{n} \left(E_{i,j}t_{1}^{m-n}\right)(q^{n}x),$$

$$Y_{W}(E_{nN+i,nN+i} - E_{nN+i+1,nN+i+1},x) = q^{n} \overline{E_{i,i} - E_{i+1,i+1}}(q^{n}x) = q^{n} H_{i}(q^{n}x),$$
for $1 \le i \ne j \le N, \ m, n \in \mathbb{Z}$ with $(i,m) \ne (j,n)$.

On the other hand, let (W, Y_W) be a (\mathbb{Z}, χ_q) -equivariant quasi $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module. From Proposition 3.5 also, W is a restricted $\widehat{\mathfrak{sl}_{\infty}}[\mathbb{Z}]$ -module of level ℓ with $\bar{a}(x) = Y_W(a, x)$ for $a \in \mathfrak{sl}_{\infty}$. By Proposition 2.8, W becomes a restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module of level ℓ on which

$$(E_{i,j}t_1^m)(x) = \overline{E_{mN+i,j}}(x) = Y_W(E_{mN+i,j}, x),$$

$$H_i(x) = \overline{E_{i,i} - E_{i+1,i+1}}(x) = Y_W(E_{i,i} - E_{i+1,i+1}, x)$$

for $1 \le i, j \le N, m \in \mathbb{Z}$ with $(i - j, m) \ne (0, 0)$.

We fix a \mathbb{Z} -grading on $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ given by derivation $-\mathbf{d}_0$ (recall (2.4)), i.e.,

(3.15)
$$\deg(E_{i,j}t_0^n t_1^m) = -n \quad \text{for } 1 \le i, j \le N, \ m, n \in \mathbb{Z}.$$

Then $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q) = \bigoplus_{n \in \mathbb{Z}} \widehat{\mathfrak{sl}_N}(\mathbb{C}_q)_{(n)}$, where for nonzero $n \in \mathbb{Z}$,

(3.16)
$$\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)_{(n)} = \sum_{1 \le i,j \le N} \sum_{m \in \mathbb{Z}} \mathbb{C}E_{i,j} t_0^{-n} t_1^m$$

and

$$(3.17)$$
 $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)_{(0)}$

$$= \sum_{1 \le i, j \le N, m \in \mathbb{Z}, (i-j,m) \ne (0,0)} \mathbb{C} E_{i,j} t_1^m + \sum_{r=1}^{N-1} \mathbb{C} (E_{r,r} - E_{r+1,r+1}) + \mathbb{C} \mathbf{k}_0 + \mathbb{C} \mathbf{k}_1.$$

Definition 3.9. A \mathbb{C} -graded $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module is an $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module W equipped with a \mathbb{C} -grading $W = \bigoplus_{\alpha \in \mathbb{C}} W(\alpha)$ such that $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)_{(n)} W(\alpha) \subset W(n+\alpha)$ for $n \in \mathbb{Z}, \ \alpha \in \mathbb{C}$.

Note that that a \mathbb{C} -graded $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module is the same as an $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q) \rtimes \mathbb{C}\mathbf{d}_0$ -module on which \mathbf{d}_0 acts semisimply. The notions of homomorphism and isomorphism for \mathbb{C} -graded $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules are defined in the obvious way (which are required to preserve the \mathbb{C} -gradings).

Remark 3.10. It follows from Proposition 3.8 that for any complex number ℓ , a \mathbb{Z} -graded (\mathbb{Z}, χ_q) -equivariant quasi $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module structure on a vector space W amounts to a \mathbb{Z} -graded restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module structure of level ℓ .

Recall that $J_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ is the maximal $\widehat{\mathfrak{sl}_{\infty}}$ -submodule of $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. The following is a description of $J_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ for nonnegative integers ℓ .

Lemma 3.11. Let ℓ be a nonnegative integer. Then

(3.18)
$$E_{i,j}(-1)^{\ell+1}\mathbf{1} \in J_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \quad \text{for any } i \neq j \in \mathbb{Z}.$$

Furthermore, $J_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ as an $\widehat{\mathfrak{sl}_{\infty}}$ -module is generated by the vectors

(3.19)
$$(E_{mN+i,nN+j}(-1))^{\ell+1}\mathbf{1}$$
 for $1 \le i \ne j \le N, \ m, n \in \mathbb{Z}$.

Proof. Let $i, j \in \mathbb{Z}$ with $i \neq j$. We first show that

(3.20)
$$E_{m,n}(k)E_{i,j}(-1)^{\ell+1}\mathbf{1} = 0, \quad (E_{m,m} - E_{n,n})(k)E_{i,j}(-1)^{\ell+1}\mathbf{1} = 0$$

for all $m, n, k \in \mathbb{Z}$ with $m \neq n, k > 0$. Let $m, n \in \mathbb{Z}$ with $m \neq n$. Pick a finite interval I of \mathbb{Z} , containing m, n, i, j. Set

$$\mathfrak{sl}_I = \operatorname{Span}\{E_{i,j}, E_{i,i} - E_{j,j} \mid i, j \in I \text{ with } i \neq j\}.$$

It follows from the P-B-W theorem that the $\widehat{\mathfrak{sl}_I}$ -submodule $\mathcal{U}(\widehat{\mathfrak{sl}_I})\mathbf{1}$ of $V_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ is isomorphic to $V_{\widehat{\mathfrak{sl}_I}}(\ell,0)$. On the other hand, it was known (cf. [LL]) that

$$(\mathfrak{sl}_I \otimes t\mathbb{C}[t]) E_{i,j}(-1)^{\ell+1} \mathbf{1} = 0 \text{ in } V_{\widehat{\mathfrak{sl}_I}}(\ell, 0)$$

for all $i, j \in I$ with $i \neq j$. Thus (3.20) holds. Then it follows from the P-B-W theorem that $\mathcal{U}(\widehat{\mathfrak{sl}_{\infty}})E_{i,j}(-1)^{\ell+1}\mathbf{1}$ is a proper graded submodule of $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$, and hence a submodule of $J_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. This proves the first assertion.

For the second assertion, denote by J' the $\widehat{\mathfrak{sl}_{\infty}}$ -submodule of $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ generated by the vectors in (3.19). Set $V' = V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)/J'$ and $\mathbf{1}' = 1 + J' \in V'$. Then it suffices to show that V' is an irreducible $\widehat{\mathfrak{sl}_{\infty}}$ -module. Let v be any nonzero vector in V'.

Fix a finite interval I = [Nm + 1, Nn] of \mathbb{Z} with m < n such that $v \in \mathcal{U}(\widehat{\mathfrak{sl}_I})\mathbf{1}'$. Note that $E_{mN+1,nN}$ is a highest root vector in \mathfrak{sl}_I and that (see (3.19))

(3.21)
$$E_{mN+1,nN}(-1)^{\ell+1}\mathbf{1}' = 0.$$

Then the highest weight $\widehat{\mathfrak{sl}}_I$ -module $\mathcal{U}(\widehat{\mathfrak{sl}}_I)\mathbf{1}'$ is integrable and hence irreducible (see [K]), which implies $\mathcal{U}(\widehat{\mathfrak{sl}}_I)\mathbf{1}' = \mathcal{U}(\widehat{\mathfrak{sl}}_I)v$. Thus

$$\mathbf{1'} \in \mathcal{U}(\widehat{\mathfrak{sl}}_I)\mathbf{1'} = \mathcal{U}(\widehat{\mathfrak{sl}}_I)v \subset \mathcal{U}(\widehat{\mathfrak{sl}}_\infty)v.$$

As $V' = \mathcal{U}(\widehat{\mathfrak{sl}}_{\infty})\mathbf{1}'$, we get $V' = \mathcal{U}(\widehat{\mathfrak{sl}}_{\infty})v$. This proves that V' is an irreducible $\widehat{\mathfrak{sl}}_{\infty}$ -module, and hence $J' = J_{\widehat{\mathfrak{sl}}_{\infty}}(\ell,0)$. Now, the proof is complete.

The following notion (cf. [ER]) is similar to the notion of integrable module for affine Kac-Moody algebras:

Definition 3.12. An $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module W is said to be *integrable* if for any $1 \leq i \neq j \leq N, m_0, m_1 \in \mathbb{Z}, E_{i,j}t_0^{m_0}t_1^{m_1}$ acts locally nilpotently on W.

Notice that for any $1 \le i \ne j \le N$, $m, n \in \mathbb{Z}$, from (2.3) we have

$$(3.22) [(E_{i,j}t_1^m)(x_1), (E_{i,j}t_1^n)(x_2)] = 0 in \widehat{\mathfrak{gl}_N}(\mathbb{C}_q)[[x_1^{\pm 1}, x_2^{\pm 1}]]$$

and hence this is true in $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)[[x_1^{\pm 1},x_2^{\pm 1}]]$. The following is analogous to a result for affine Kac-Moody algebras:

Proposition 3.13. Let W be a restricted $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module of level $\ell \in \mathbb{C}$. Then W is integrable if and only if ℓ is a nonnegative integer and

$$(E_{i,j}t_1^m)(x)^{\ell+1} = 0$$
 on W

for all $1 \le i \ne j \le N$ and $m \in \mathbb{Z}$.

Proof. Let $\{e, h, f\}$ be the standard basis of \mathfrak{sl}_2 such that

$$[e, f] = h, \quad [h, e] = 2e, \quad [h, f] = -2f.$$

It was known (see [LP], [DLM]) that a restricted module U of level ℓ for the affine Lie algebra $\widehat{\mathfrak{sl}_2}$ is integrable if and only if ℓ is a nonnegative integer and

$$e(x)^{\ell+1} = 0 = f(x)^{\ell+1}$$
 on U ,

where $a(x) = \sum_{n \in \mathbb{Z}} (a \otimes t^n) x^{-n-1}$ for $a \in \mathfrak{sl}_2$. For $1 \leq i < j \leq N, \ m \in \mathbb{Z}$, set

$$\widehat{\mathcal{A}}_{i,j}(m) = \operatorname{Span}\{E_{i,j}t_0^n t_1^m, E_{j,i}t_0^n t_1^{-m}, q^{mn} E_{i,i}t_0^n - E_{j,j}t_0^n + m\delta_{n,0}\mathbf{k}_1, \mathbf{k}_0 \mid n \in \mathbb{Z}\}.$$

It is straightforward to show that $\widehat{\mathcal{A}}_{i,j}(m)$ is a subalgebra of $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ isomorphic to $\widehat{\mathfrak{sl}}_2$, where an isomorphism is given by

$$q^{mn}E_{i,j}t_0^nt_1^m \mapsto e \otimes t^n, \quad E_{j,i}t_0^nt_1^{-m} \mapsto f \otimes t^n,$$

$$q^{mn}E_{i,i}t_0^n - E_{j,j}t_0^n + m\delta_{n,0}\mathbf{k}_1 \mapsto h \otimes t^n, \quad \mathbf{k}_0 \mapsto \mathbf{k}$$

for $n \in \mathbb{Z}$. Then W is an integrable $\widehat{\mathcal{A}}_{i,j}(m)$ -module if and only if ℓ is a nonnegative integer and

$$(E_{i,j}t_1^m)(x)^{\ell+1} = (E_{j,i}t_1^{-m})(x)^{\ell+1} = 0$$
 on W .

On the other hand, from definition W is an integrable $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module if and only if for any $1 \leq i < j \leq N, \ m \in \mathbb{Z}, W$ is an integrable $\widehat{\mathcal{A}}_{i,j}(m)$ -module. Then it follows immediately.

The following is a special case of [LTW, Corollary 5.3]:

Lemma 3.14. Let V be a vertex Γ -algebra, let $a \in V$ such that $a_n a = 0$ for $n \geq 0$, and let ℓ be a nonnegative integer. Suppose that (W, Y_W) is an equivariant quasi V-module such that

$$[Y_W(a, x_1), Y_W(a, x_2)] = 0.$$

If $(a_{-1})^{\ell+1}\mathbf{1} = 0$ in V, then

$$(3.24) Y_W(a,x)^{\ell+1} = 0 on W.$$

On the other hand, the converse is also true if (W, Y_W) is faithful.

As $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ is a quotient algebra of $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$, in view of Proposition 3.8, (\mathbb{Z},χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -modules are naturally restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ . The following is the first main result of this paper:

Theorem 3.15. Let ℓ be a nonnegative integer. Then (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -modules exactly correspond to integrable restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ .

Proof. Let (W, Y_W) be a (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module. In view of Proposition 3.8, W is a restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module of level ℓ with

$$(E_{i,j}t_1^m)(x) = Y_W(E_{Nm+i,j}, x)$$
 for $1 \le i \ne j \le N, m \in \mathbb{Z}$.

Let $1 \le i \ne j \le N$, $m, n \in \mathbb{Z}$. Noticing that $Nm + i \ne Nn + j$, we have

$$[E_{Nm+i,Nn+j}(x_1), E_{Nm+i,Nn+j}(x_2)] = 0$$

in $\widehat{\mathfrak{sl}_{\infty}}$, which implies

$$(3.25) (E_{Nm+i,Nn+j})_k (E_{Nm+i,Nn+j}) = 0 \text{for } k \ge 0$$

in $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ and hence in $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$. From Lemma 3.11 we also have

$$E_{Nm+i,j}(-1)^{\ell+1}\mathbf{1} = 0$$
 in $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$.

On the other hand, as $i \neq j$, by (3.22) we have

$$[Y_W(E_{Nm+i,j},x_1),Y_W(E_{Nm+i,j},x_2)] = [(E_{i,j}t_1^m)(x_1),(E_{i,j}t_1^m)(x_2)] = 0.$$

Then by Lemma 3.14 we get

$$Y_W(E_{Nm+i,j}, x)^{\ell+1} = 0$$
 on W .

Thus

$$(E_{i,j}t_1^m)(x)^{\ell+1} = Y_W(E_{Nm+i,j}, x)^{\ell+1} = 0$$
 on W .

In view of Proposition 3.13, W is an integrable $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module.

On the other hand, let W be an integrable restricted $\mathfrak{sl}_N(\mathbb{C}_q)$ -module of level ℓ . By Propositions 3.8 and 3.13, there is a (\mathbb{Z}, χ_q) -equivariant quasi $V_{\widehat{\mathfrak{sl}_\infty}}(\ell, 0)$ -module structure $Y_W(\cdot, x)$ on W such that

$$Y_W(E_{Nm+i,Nn+j}, x) = q^n (E_{i,j} t_1^m) (q^n x)$$
 for $1 \le i \ne j \le N, \ m, n \in \mathbb{Z}$.

Then we have

$$(3.26) Y_W(E_{Nm+i,Nn+j},x)^{\ell+1} = q^{n(\ell+1)} (E_{i,j}t_1^m) (q^n x)^{\ell+1} = 0$$

for $1 \leq i \neq j \leq N$ and $m, n \in \mathbb{Z}$, noticing that

$$[Y_W(E_{Nm+i,Nn+j},x_1),Y_W(E_{Nm+i,Nn+j},x_1)]$$

$$=[q^n(E_{i,j}t_1^m)(q^nx_1),q^n(E_{i,j}t_1^m)(q^nx_2)]=0.$$

Note that W is naturally a faithful equivariant quasi module for $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)/\ker Y_W$. With (3.25), (3.27) and (3.26), it follows from Lemma 3.14 that

$$E_{Nm+i,Nn+j}(-1)^{\ell+1}\mathbf{1} = 0$$
 in $V_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)/\ker Y_W$

for all $1 \leq i \neq j \leq N$ and $m, n \in \mathbb{Z}$. Then by Lemma 3.11 $Y_W(\cdot, x)$ reduces to an equivariant quasi module structure $\overline{Y_W}(\cdot, x)$ for $L_{\widehat{\mathfrak{sl}_\infty}}(\ell, 0)$. Consequently, W is an equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell, 0)$ -module with the required property. \square

4. Classification of irreducible equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -modules

The main goal of this section is to classify irreducible N-graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_\infty}}(\ell,0)$ -modules for any nonnegative integer ℓ , or equivalently irreducible N-graded integrable $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -modules of level ℓ . Irreducible N-graded (\mathbb{Z},χ_q) -equivariant quasi modules for the tensor product vertex \mathbb{Z} -algebra $L_{\widehat{\mathfrak{sl}_\infty}}(\ell_1,0)\otimes\cdots\otimes L_{\widehat{\mathfrak{sl}_\infty}}(\ell_d,0)$ are also determined.

4.1. Integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules. In this subsection, we introduce a notion of highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module and classify all irreducible \mathbb{N} -graded integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules.

Note that $\mathfrak{sl}_N(\mathbb{C}[t_0, t_0^{-1}]) \oplus \mathbb{C}\mathbf{k}_0 \oplus \mathbb{C}\mathbf{d}_0$ is a subalgebra of $\widetilde{\mathfrak{sl}}_N(\mathbb{C}_q)$, which is isomorphic to the affine Kac-Moody algebra $\widetilde{\mathfrak{sl}}_N$ of type $A_{N-1}^{(1)}$. Here, we identify $\widetilde{\mathfrak{sl}}_N$ with this subalgebra of $\widetilde{\mathfrak{sl}}_N(\mathbb{C}_q)$:

(4.1)
$$\widetilde{\mathfrak{sl}_N} = \mathfrak{sl}_N(\mathbb{C}[t_0, t_0^{-1}]) \oplus \mathbb{C}\mathbf{k}_0 \oplus \mathbb{C}\mathbf{d}_0 \subset \widetilde{\mathfrak{sl}_N}(\mathbb{C}_q).$$

Set

(4.2)
$$\mathfrak{h} = \sum_{i=1}^{N-1} \mathbb{C}(E_{i,i} - E_{i+1,i+1}),$$

a Cartan subalgebra of $\mathfrak{sl}_N(=\mathfrak{sl}_N(\mathbb{C}))$, and set

$$(4.3) H = \mathfrak{h} \oplus \mathbb{C}\mathbf{k}_0 \oplus \mathbb{C}\mathbf{d}_0,$$

a Cartan subalgebra of the affine Kac-Moody algebra $\widehat{\mathfrak{sl}_N}$. We shall use the following triangular decomposition of $\widehat{\mathfrak{sl}_N}(\mathbb{C}_a)$:

$$\widehat{\mathfrak{sl}_N}(\mathbb{C}_q) = \widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+ \oplus \widehat{\mathcal{H}} \oplus \widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^-,$$

where

(4.4)
$$\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^{\pm} = \sum_{\pm m_0 \in \mathbb{Z}_+, m_1 \in \mathbb{Z}} \mathbb{C} E_{i,j} t_0^{m_0} t_1^{m_1} + \sum_{\pm (j-i) \in \mathbb{Z}_+, m \in \mathbb{Z}} \mathbb{C} E_{i,j} t_1^m,$$

and

(4.5)
$$\widehat{\mathcal{H}} = \sum_{1 \le i \le N, \ n \in \mathbb{Z}} \mathbb{C}h_{i,n} + \mathbb{C}\mathbf{k}_1,$$

where

$$h_{i,n} = (E_{i,i} - E_{i+1,i+1})t_1^n \quad \text{for } 1 \le i \le N-1, \ n \in \mathbb{Z},$$

$$(4.6) \quad h_{N,n} = -q^n E_{1,1}t_1^n + E_{N,N}t_1^n \quad \text{for } n \in \mathbb{Z} \setminus \{0\},$$

$$h_{N,0} = \mathbf{k}_0 - (E_{1,1} - E_{N,N}).$$

To emphasize the dependence on the positive integer N, we shall also alternatively denote the subalgebra $\widehat{\mathcal{H}}$ by $\widehat{\mathcal{H}}_N$. Note that $\{h_{i,n} \mid 1 \leq i \leq N, n \in \mathbb{Z}\} \cup \{\mathbf{k}_1\}$ is a basis of $\widehat{\mathcal{H}}$.

Definition 4.1. An $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module W is called a *highest weight module with highest weight* $\lambda \in \widehat{\mathcal{H}}^*$ if there exists a nonzero vector $v_\lambda \in W$, called a *highest weight vector*, such that $W = \mathcal{U}(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q))v_\lambda$, $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+v_\lambda = 0$, and

$$(4.7) hv_{\lambda} = \lambda(h)v_{\lambda} \text{for } h \in \widehat{\mathcal{H}}.$$

An \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module with highest weight $\lambda \in \widehat{\mathcal{H}}^*$ is a highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module W equipped with an \mathbb{N} -grading $W = \bigoplus_{n \in \mathbb{N}} W(n)$ such that the highest weight vector $v_{\lambda} \in W(0)$.

Remark 4.2. Recall that $H = \mathfrak{h} + \mathbb{C}\mathbf{k}_0 + \mathbb{C}\mathbf{d}_0$ is a Cartan subalgebra of $\widehat{\mathfrak{sl}}_N$. We see that an \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module W is an H-weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{d}_0$ -module with \mathbf{d}_0 acting trivially on the highest weight vector. Then W is a highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{d}_0$ -module defined in [ER] with respect to the usual partial order on H^* .

Note that the condition (4.7) implies that $\lambda(\mathbf{k}_1) = 0$. Now, let $\lambda \in \widehat{\mathcal{H}}^*$ such that $\lambda(\mathbf{k}_1) = 0$. Let $\mathbb{C}v_{\lambda}$ be the one-dimensional $(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+ + \widehat{\mathcal{H}})$ -module with

$$\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)^+ v_\lambda = 0$$
 and $hv_\lambda = \lambda(h)v_\lambda$ for $h \in \widehat{\mathcal{H}}$.

Then form an induced $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module

$$(4.8) V(\lambda) = \mathcal{U}(\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)) \otimes_{\mathcal{U}(\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)^+ + \widehat{\mathcal{H}})} \mathbb{C}v_{\lambda}.$$

It is clear that $V(\lambda)$ is an $(\mathfrak{h} + \mathbb{C}\mathbf{k}_0 + \mathbb{C}\mathbf{k}_1)$ -weight module. On the other hand, define $\deg v_{\lambda} = 0$ to make $V(\lambda)$ an \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module. It follows that $V(\lambda)$ has a unique maximal H-weight submodule, which we denote by J_{λ} . Set

$$(4.9) L(\lambda) = V(\lambda)/J_{\lambda},$$

which is a graded irreducible $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module. One can show that $L(\lambda)$ is also an irreducible $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module. Furthermore, it is straightforward to see that for any $\lambda, \mu \in \widehat{\mathcal{H}}^*$, $L(\lambda) \simeq L(\mu)$ as an N-graded $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module if and only if $\lambda = \mu$.

Remark 4.3. Note that for any $\lambda \in \widehat{\mathcal{H}}^*$ such that $\lambda(\mathbf{k}_1) = 0$, every \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module with highest weight λ is naturally a homomorphism image of $V(\lambda)$ and any such irreducible module is isomorphic to $L(\lambda)$.

Next, we identify integrable $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules among the irreducible highest weight modules $L(\lambda)$. We first define a class of linear functionals on $\widehat{\mathcal{H}}$. Let k be a positive integer. For any pair

(4.10)
$$(\boldsymbol{\lambda}, \boldsymbol{c}) \in (H^*)^k \times (\mathbb{C}^\times)^k$$

with $\lambda = (\lambda_1, \dots, \lambda_k)$, $\boldsymbol{c} = (c_1, \dots, c_k)$, we define a linear functional $\eta_{\lambda, \boldsymbol{c}}$ on $\widehat{\mathcal{H}}$ by $\eta_{\lambda, \boldsymbol{c}}(\mathbf{k}_1) = 0$ and

(4.11)
$$\eta_{\boldsymbol{\lambda},\boldsymbol{c}}(h_{j,n}) = \sum_{i=1}^{k} \lambda_i(h_{j,0}) c_i^n \quad \text{for } 1 \le j \le N, \ n \in \mathbb{Z}.$$

In particular, for any $\lambda \in H^*$, $c \in \mathbb{C}^{\times}$, we have a linear functional $\eta_{\lambda,c}$. Then

$$\eta_{\lambda,c} = \eta_{\lambda_1,c_1} + \cdots + \eta_{\lambda_k,c_k}.$$

Set

(4.12)
$$P_{+} = \{ \lambda \in H^* \mid \lambda(h_{i,0}) \in \mathbb{N}, \ \lambda(\mathbf{d}_0) = 0 \text{ for } 1 \le i \le N \},$$

the set of dominant integral weights of \mathfrak{sl}_N .

The following is a classification result (cf. [ER, CT]):

Proposition 4.4. Let $(\lambda, \mathbf{c}) \in (P_+)^k \times (\mathbb{C}^\times)^k$ with k a positive integer. Then the irreducible \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module $L(\eta_{\lambda,\mathbf{c}})$ is integrable. Furthermore, every irreducible integrable \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module is isomorphic to a module of this form.

Remark 4.5. Note that the first assertion of Proposition 4.4 was proved in [ER], while the second assertion was proved for Lie algebra $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q) \oplus \mathbb{C}\mathbf{d}_0$ under the assumption that the H-weight subspaces of W are finite-dimensional.

To prove Proposition 4.4, we shall need a result on level zero integrable modules for affine Lie algebra $\widehat{\mathfrak{sl}_2}$. Let $\{e,h,f\}$ be the standard basis of \mathfrak{sl}_2 and let $\chi:\mathbb{Z}\to\mathbb{C}$ be any function. An $\widehat{\mathfrak{sl}_2}$ -module W of level zero is called a *loop-highest weight module with loop-highest weight* χ if there is a nonzero vector w in W such that

$$(4.13) W = \mathcal{U}(\widehat{\mathfrak{sl}_2})w, (e \otimes t^m) w = 0, (h \otimes t^m) w = \chi(m)w for m \in \mathbb{Z}.$$

The notion "loop-highest weight module" (see [CG] for example) is designated to distinguish this from the standard highest weight module (see [K]).

Lemma 4.6. If there exists an integrable loop-highest weight $\widehat{\mathfrak{sl}}_2$ -module W with a nonzero loop-highest weight χ , then there exist finitely many positive integers p_1, \ldots, p_k and nonzero complex numbers b_1, \ldots, b_k such that

$$\chi(m) = \sum_{j=1}^{k} p_j b_j^m \quad \text{for } m \in \mathbb{Z}.$$

Proof. Recall from [BZ] that χ is called an exp-polynomial if there exist finitely many polynomials f_1, \ldots, f_k and nonzero complex numbers b_1, \ldots, b_k such that $\chi(m) = \sum_{j=1}^k f_j(m) b_j^m$ for all $m \in \mathbb{Z}$. A result of Kac-Jocobsen (see [JK, Proposition 6.2]) states that if χ is not an exp-polynomial, then the Verma type loophighest weight $\widehat{\mathfrak{sl}}_2$ -module with loop-highest weight χ is irreducible. On the other hand, every Verma type loop-highest weight $\widehat{\mathfrak{sl}}_2$ -module is not integrable. As W is assumed to be integrable, the Verma type loop-highest weight $\widehat{\mathfrak{sl}}_2$ -module with loop-highest weight χ must be reducible and hence χ is an exp-polynomial.

Let w be a (nonzero) loop-highest weight vector in W. Then hw = nw for some nonnegative integer n as w is a highest weight vector in the integrable \mathfrak{sl}_2 -module W. Noticing that $W = \mathcal{U}(\mathbb{C}f \otimes \mathbb{C}[t, t^{-1}])w$, we have

$$W = \bigoplus_{m \in \mathbb{Z}, m \le n} W_m$$
 with $W_n = \mathbb{C}w$,

where $W_m = \{w \in W \mid hw = mw\}$. Since W is an integrable \mathfrak{sl}_2 -module, we have $W = \bigoplus_{m \in \mathbb{Z}, |m| \leq n} W_m$. From [BZ], all the h-weight subspaces of the (unique) irreducible quotient module \overline{W} of W are finite-dimensional. Consequently, \overline{W} is finite-dimensional. Then this lemma follows from [CP, Proposition 2.1 (iii)]. \square

Proof of Proposition 4.4: The first assertion was proved in [ER]. (This also follows from an explicit realization of $L(\eta_{\lambda,c})$ in Section 6.) Now, let W be an irreducible integrable \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module with highest weight χ and highest weight vector v. For $1 \leq i \leq N-1$, set

$$\mathfrak{a}_i = \text{Span}\{E_{i,i+1}t_1^n, E_{i+1,i}t_1^n, h_{i,n}, \mathbf{k}_1 \mid n \in \mathbb{Z}\},\$$

and set

$$\mathfrak{a}_N = \operatorname{Span} \{ q^n E_{N,1} t_0 t_1^n, E_{1,N} t_0^{-1} t_1^n, h_{N,n}, \mathbf{k}_1 \mid n \in \mathbb{Z} \}.$$

It can be readily seen that \mathfrak{a}_i for $1 \leq i \leq N$ are subalgebras of $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$, which are isomorphic to $\widehat{\mathfrak{sl}_2}$. Then for every i, $\mathcal{U}(\mathfrak{a}_i)v$ is an integrable loop-highest weight $\widehat{\mathfrak{sl}_2}$ -module with loop-highest weight χ_i , where

$$\chi_i(n) = \chi(h_{i,n})$$
 for $n \in \mathbb{Z}$.

By Lemma 4.6, for each $1 \leq i \leq N$, there exist finitely many nonzero complex numbers $b_{i,1}, \ldots, b_{i,k_i}$ and nonnegative integers $p_{i,1}, \ldots, p_{i,k_i}$ such that

$$\chi_i(m) = \sum_{j=1}^{k_i} p_{i,j} b_{i,j}^m \quad \text{for } m \in \mathbb{Z}.$$

Let c_1, \ldots, c_r be all the distinct elements of $\{b_{i,j} \mid 1 \leq i \leq N, 1 \leq j \leq k_i\}$. Then

$$\chi_i(m) = \sum_{i=1}^r q_{i,j} c_j^m \quad \text{ for } 1 \le i \le N, \ m \in \mathbb{Z},$$

where $q_{i,j}$ are nonnegative integers which are independent of m. Define r linear functionals $\lambda_1, \ldots, \lambda_r$ on H by $\lambda_i(h_{j,0}) = q_{i,j}$ for $1 \leq i, j \leq N$ and $\lambda_i(\mathbf{d}_0) = 0$. We have $\lambda_i \in P_+$ and $\chi = \eta_{\lambda,\mathbf{c}}$. Then W is isomorphic to $L(\eta_{\lambda,\mathbf{c}})$, as desired.

Remark 4.7. Recall that a fundamental dominant weight is an element $\lambda \in P_+$ such that $\lambda(\mathbf{k}_0) = 1$. Let $(\lambda, \mathbf{c}) \in P_+^k \times (\mathbb{C}^\times)^k$ with $\lambda_i \neq 0$ for $1 \leq i \leq k$. Set $\ell_i = \lambda_i(\mathbf{k}_0) \in \mathbb{Z}_+$. Furthermore, set $\ell = \ell_1 + \cdots + \ell_k$. Note that $\eta_{\lambda, \mathbf{c}} + \eta_{\lambda', \mathbf{c}'} = \eta_{(\lambda, \lambda'), (\mathbf{c}, \mathbf{c}')}$ and $\eta_{\lambda + \mu, \mathbf{c}} = \eta_{(\lambda, \mu), (\mathbf{c}, \mathbf{c})}$. It then follows that there exists a pair $(\mu, \mathbf{d}) \in P_+^\ell \times (\mathbb{C}^\times)^\ell$ such that $\eta_{\lambda, \mathbf{c}} = \eta_{\mu, \mathbf{d}}$, where $\mu = (\mu_1, \dots, \mu_\ell)$ with μ_1, \dots, μ_ℓ fundamental dominant weights.

It is known (see [DLM]) that every irreducible integrable restricted module for any (untwisted) affine Kac-Moody Lie algebra is a highest weight module. For the extended affine Lie algebra $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$, we have the following result:

Proposition 4.8. Let $W = \bigoplus_{n \in \mathbb{N}} W(n)$ be a nonzero \mathbb{N} -graded integrable $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ module on which \mathbf{k}_0 and \mathbf{k}_1 act as scalars ℓ_0 and ℓ_1 , respectively. Then ℓ_0 is a
nonnegative integer, $\ell_1 = 0$, and Ω_W is a nonzero $\widehat{\mathcal{H}}$ -submodule of W, where

(4.14)
$$\Omega_W = \{ w \in W \mid \widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+ w = 0 \}.$$

Furthermore, every irreducible \mathbb{N} -graded integrable (restricted) $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module is a highest weight module.

Proof. For the first assertion, by Proposition 3.13, ℓ_0 is a nonnegative integer. We now prove $\ell_1 = 0$. Fix two integers i, j with $1 \le i < j \le N$. Let $m \in \mathbb{Z}$. Note that the correspondence

$$e \mapsto E_{i,j}t_1^m, \quad f \mapsto E_{j,i}t_1^{-m}, \quad h \mapsto E_{i,i} - E_{j,j} + m\mathbf{k}_1$$

gives rise to a Lie algebra embedding of \mathfrak{sl}_2 into $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$. Under this embedding, W becomes an \mathfrak{sl}_2 -module on which e and f act locally nilpotently. Then W is an integrable \mathfrak{sl}_2 -module and hence it is a direct sum of finite-dimensional irreducible \mathfrak{sl}_2 -modules. Let v be a (nonzero) highest weight vector of weight k. As

$$hv = (E_{i,i} - E_{j,j} + m\mathbf{k}_1)v = (k + m\ell_1)v,$$

 $k + m\ell_1$ must be a nonnegative integer. Thus $k + m\ell_1 \in \mathbb{N}$ for all $m \in \mathbb{Z}$, which implies $\ell_1 = 0$.

Now that \mathbf{k}_1 acts trivially, W is an integrable module for the loop algebra $\mathfrak{sl}_N(\mathbb{C}[t_1,t_1^{-1}])$. As W is an \mathbb{N} -graded module, there exists $n\in\mathbb{N}$ such that $W(n)\neq 0$ and W(m)=0 for all m< n. Then

$$(4.15) (E_{i,j}t_0^{m_0}t_1^{m_1})W(n) \subset W(n-m_0) = 0$$

for all $1 \leq i, j \leq N$, $m_0 \geq 1$, $m_1 \in \mathbb{Z}$. Notice that W(n) is an $\mathfrak{sl}_N(\mathbb{C}[t_1, t_1^{-1}])$ -submodule of W. As W is an integrable restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module, by Proposition 3.13 we have $(E_{i,j}t_1^m)(z)^{\ell_0+1}=0$ for $1\leq i\neq j\leq N$, $m\in\mathbb{Z}$. This together with (4.15) implies that $(E_{i,j}t_1^m)^{\ell_0+1}=0$ on W(n) for all $m\in\mathbb{Z}$. In particular, we have $E_{i,j}^{\ell+1}=0$ on W(n) for all $1\leq i\neq j\leq N$. Then W(n) is a direct sum of some finite-dimensional irreducible \mathfrak{sl}_N -modules $L(\lambda)$ where λ are dominant integral weights satisfying the condition $\lambda(\theta^\vee)\leq \ell$ (with θ denoting the highest positive root). As there are only finitely many such dominant integral weights, there exists $\lambda\in\mathfrak{h}^*$ such that $W(n)_\lambda\neq 0$ and $W(n)_{\lambda+\alpha}=0$ for any positive root α of \mathfrak{sl}_N . Then $(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+)W(n)_\lambda=0$. Thus we have $W(n)_\lambda\subset\Omega_W$, proving that $\Omega_W\neq 0$. As $[\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+,\widehat{\mathcal{H}}]\subset\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+$, it follows that Ω_W is an $\widehat{\mathcal{H}}$ -submodule of W.

Now, assume that W is an irreducible \mathbb{N} -graded integrable $\mathfrak{sl}_N(\mathbb{C}_q)$ -module. Then \mathbf{k}_0 and \mathbf{k}_1 act as scalars ℓ_0 and ℓ_1 on W, respectively. As \mathbf{k}_1 acts trivially on W by the first assertion, Ω_W becomes a module for the abelian Lie algebra $\widehat{\mathcal{H}}/\mathbb{C}\mathbf{k}_1$. Furthermore, the irreducibility of W implies that Ω_W is an irreducible $\widehat{\mathcal{H}}$ -module. Consequently, $\Omega_W = \mathbb{C}v_\lambda$ for some $v_\lambda \in \Omega_W \cap W_\lambda$. It then follows that W is a highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module with highest weight vector v_λ .

Combining Propositions 4.8 and 4.4 we immediately have:

Corollary 4.9. Every irreducible \mathbb{N} -graded integrable $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module of level $\ell \in \mathbb{N}$ is isomorphic to $L(\eta_{\lambda, \mathbf{c}})$ for some $\lambda = (\lambda_1, \dots, \lambda_k) \in (P_+)^k$ and $\mathbf{c} = (c_1, \dots, c_k) \in (\mathbb{C}^{\times})^k$ with k a positive integer such that $\sum_{i=1}^k \lambda_i(\mathbf{k}_0) = \ell$.

Combining Corollary 4.9 with Remark 3.10 and Theorem 3.15, we immediately have the main result of this section:

Theorem 4.10. For any $\ell \in \mathbb{N}$, irreducible \mathbb{N} -graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -modules up to isomorphism are exactly the irreducible integrable \mathbb{N} -graded highest weight $\widehat{\mathfrak{sl}_{N}}(\mathbb{C}_q)$ -modules $L(\eta_{\lambda, \mathbf{c}})$ for $\lambda = (\lambda_1, \ldots, \lambda_k) \in (P_+)^k$ and $\mathbf{c} = (c_1, \ldots, c_k) \in (\mathbb{C}^\times)^k$ with $k \in \mathbb{Z}_+$ such that $\sum_{i=1}^k \lambda_i(\mathbf{k}_0) = \ell$.

Furthermore, we have:

Lemma 4.11. Let $(\lambda, c) \in (H^*)^r \times (\mathbb{C}^{\times})^r$ with r a positive integer such that $\eta_{\lambda, c} \neq 0$. Then there exists

$$(\boldsymbol{\mu}, \boldsymbol{d}) = ((\mu_1, \dots, \mu_s), (d_1, \dots, d_s)) \in (H^*)^s \times (\mathbb{C}^\times)^s$$

for some positive integer s, satisfying the condition

(4.16)
$$\mu_1, \ldots, \mu_s$$
 are nonzero and d_1, \ldots, d_s are distinct,

such that $\eta_{\lambda,c} = \eta_{\mu,d}$. Furthermore, assume $(\mu', d') \in (H^*)^{s'} \times (\mathbb{C}^{\times})^{s'}$ is another pair satisfying the same condition above. Then $\eta_{\mu,d} = \eta_{\mu',d'}$ if and only if s = s' and there is a permutation $\tau \in S_s$ such that

$$\mu_{\tau(i)} = \mu'_i$$
 and $d_{\tau(i)} = d'_i$ for $i = 1, ..., s$.

Proof. Write $\lambda = (\lambda_1, \dots, \lambda_r) \in (H^*)^r$ and $\mathbf{c} = (c_1, \dots, c_r) \in (\mathbb{C}^\times)^r$. Let c'_1, \dots, c'_k be all the distinct numbers in $\{c_1, \dots, c_r\}$. Define $\lambda' = (\lambda'_1, \dots, \lambda'_k)$ with $\lambda'_j = \sum_{i, c_i = c'_j} \lambda_i \in H^*$ for $1 \leq j \leq k$. Then $\eta_{\lambda, \mathbf{c}} = \eta_{\lambda', \mathbf{c}'}$. As $\eta_{\lambda, \mathbf{c}} \neq 0$, we have $\lambda'_j \neq 0$ for some j. Let μ_1, \dots, μ_s be all the nonzero elements in the list $\lambda'_1, \dots, \lambda'_k$ and let d_1, \dots, d_s be the corresponding nonzero numbers in the list c'_1, \dots, c'_k . Then we have $\eta_{\lambda, \mathbf{c}} = \eta_{\mu, \mathbf{d}}$ with $\mu = (\mu_1, \dots, \mu_s)$ and $\mathbf{d} = (d_1, \dots, d_s)$, as desired.

Note that for any $(\lambda, c) \in (H^*)^r \times (\mathbb{C}^{\times})^r$ with c_1, \ldots, c_r distinct, $\eta_{\lambda, c} = 0$ if and only if $\lambda_i = 0$ for all i. Assume $\eta_{\mu, d} = \eta_{\mu', d'}$. Set

$$I = \{1 \le i \le s \mid d_i \ne d'_j \text{ for all } 1 \le j \le s'\}.$$

If $I \neq \emptyset$, the equality $\eta_{\boldsymbol{\mu},\boldsymbol{d}} = \eta_{\boldsymbol{\mu}',\boldsymbol{d}'}$ implies

$$\sum_{i \in I} \eta_{\mu_i, d_i} + \sum_{j=1}^{s'} \eta_{\nu_j, d'_j} = 0$$

for some $\nu_j \in \mathfrak{h}^*$. This is impossible as d_i with $i \in I$ and $d'_1, \ldots, d'_{s'}$ are all distinct and $\mu_i \neq 0$ for $i \in I$. Thus $I = \emptyset$, i.e., $\{d_1, \ldots, d_s\} \subset \{d'_1, \ldots, d'_{s'}\}$. Symmetrically, we have $\{d'_1, \ldots, d'_{s'}\} \subset \{d_1, \ldots, d_s\}$. Thus $\{d_1, \ldots, d_s\} = \{d'_1, \ldots, d'_{s'}\}$. In

particular, we have s = s' and there exists a permutation τ such that $d_{\tau(i)} = d'_i$ for i = 1, ..., s. Furthermore, we have

$$\eta_{\boldsymbol{\mu},\boldsymbol{d}} = \sum_{i=1}^{s} \eta_{\mu_{i},d_{i}} = \sum_{i=1}^{s} \eta_{\mu_{\tau(i)},d_{\tau(i)}} = \sum_{i=1}^{s} \eta_{\mu_{\tau(i)},d'_{i}} \text{ and } \eta_{\boldsymbol{\mu}',\boldsymbol{d}'} = \sum_{i=1}^{s} \eta_{\mu'_{i},d'_{i}},$$

which imply $\mu_{\tau(i)} = \mu'_i$ for $i = 1, \ldots, s$.

4.2. Irreducible equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_1,0)\otimes\cdots\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_d,0)$ -modules. Let d be a positive integer and let $\ell_1,\ldots,\ell_d,N_1,\ldots,N_d$ be positive integers with $N_i\geq 2$ for $1\leq i\leq d$. Set $\boldsymbol{\ell}=(\ell_1,\ldots,\ell_d)$. From Example 3.6, we have a simple \mathbb{Z} -graded vertex \mathbb{Z} -algebra

$$(4.17) L_{\widehat{\mathfrak{sl}_{\infty}}}(\boldsymbol{\ell},0) := (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_1,0),\rho_{N_1}) \otimes \cdots \otimes (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_r,0),\rho_{N_d}),$$

recalling that $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_i,0),\rho_{N_i})$ denotes the simple vertex \mathbb{Z} -algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_i,0)$ with the \mathbb{Z} -action on \mathfrak{sl}_{∞} given by ρ_{N_i} . For $1 \leq i \leq d$, we view $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_i,0),\rho_{N_i})$ as a vertex \mathbb{Z} -subalgebra of $L_{\widehat{\mathfrak{sl}_{\infty}}}(\boldsymbol{\ell},0)$ in the obvious way. The main goal of this section is to prove the following (cf. [FHL, Theorem 4.7.4]):

Proposition 4.12. Let W_i be an irreducible (\mathbb{Z}, χ_q) -equivariant quasi $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_i, 0), \rho_{N_i})$ module for $1 \leq i \leq d$. Then the tensor product equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module $W_1 \otimes \cdots \otimes W_d$ is irreducible. On the other hand, every irreducible \mathbb{N} -graded (\mathbb{Z}, χ_q) equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module is isomorphic to such a tensor product module.

Proof. Note that each W_i is of countable dimension over \mathbb{C} , so that the Schur lemma holds. Then the first assertion follows from the Jacobson density theorem just as in [FHL]. Now, we consider the second assertion. Let W be any irreducible \mathbb{N} -graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module. Then for each $1 \leq i \leq d$, W is naturally an \mathbb{N} -graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_i, 0)$ -module, and for $1 \leq i \neq j \leq d$, the actions of $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_i, 0)$ and $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell_j, 0)$ on W commute. By Theorem 3.15, W is naturally an \mathbb{N} -graded integrable and restricted $\widehat{\mathfrak{sl}_{N_i}}(\mathbb{C}_q)$ -module of level ℓ_i . Furthermore, the actions of $\widehat{\mathfrak{sl}_{N_i}}(\mathbb{C}_q)$ and $\widehat{\mathfrak{sl}_{N_j}}(\mathbb{C}_q)$ on W commute for $1 \leq i \neq j \leq d$. By Proposition 4.8, we have $\Omega_W^i \neq 0$, where

$$\Omega_W^i = \{ w \in W \mid \widehat{\mathfrak{sl}_{N_i}}(\mathbb{C}_q)^+ w = 0 \}.$$

For $1 \leq i \neq j \leq d$, as $[\widehat{\mathfrak{sl}_{N_i}}(\mathbb{C}_q), \widehat{\mathfrak{sl}_{N_j}}(\mathbb{C}_q)] = 0$ on W we have $\widehat{\mathfrak{sl}_{N_j}}(\mathbb{C}_q)\Omega_W^i \subset \Omega_W^i$. It follows (from the first part of Proposition 4.8) that $\bigcap_{i=1}^d \Omega_W^i \neq 0$. Set

$$\mathcal{K} = \widehat{\mathfrak{sl}_{N_1}}(\mathbb{C}_q) \oplus \cdots \oplus \widehat{\mathfrak{sl}_{N_d}}(\mathbb{C}_q).$$

As an irreducible N-graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module, W is also an irreducible N-graded K-module. From the proof of the second assertion of

Proposition 4.8, we conclude that $\bigcap_{i=1}^d \Omega_W^i = \mathbb{C}v$, where $v \in \bigcap_{i=1}^d \Omega_W^i$ and there exists a linear functional λ on $\widehat{\mathcal{H}}_{N_1} \oplus \cdots \oplus \widehat{\mathcal{H}}_{N_d}$ such that $W = U(\mathcal{K})v$ and

$$hv = \lambda(h)v \quad \text{for } h \in \widehat{\mathcal{H}}_{N_1} \oplus \cdots \oplus \widehat{\mathcal{H}}_{N_d}.$$

Given a linear functional λ on $\widehat{\mathcal{H}}_{N_1} \oplus \cdots \oplus \widehat{\mathcal{H}}_{N_d}$, we say that a \mathcal{K} -module M is a highest weight module of highest weight λ if there is a nonzero vector $w \in M$ such that $\mathcal{U}(\mathcal{K})w = M$, $(\widehat{\mathfrak{sl}}_{N_1}(\mathbb{C}_q)^+ \oplus \cdots \oplus \widehat{\mathfrak{sl}}_{N_d}(\mathbb{C}_q)^+)w = 0$ and

$$hw = \lambda(h)w$$
 for $h \in \widehat{\mathcal{H}}_{N_1} \oplus \cdots \oplus \widehat{\mathcal{H}}_{N_d}$.

That is, W is an \mathbb{N} -graded highest weight \mathcal{K} -module. On the other hand, we define Verma module $M_{\mathcal{K}}(\lambda)$ in the obvious way, which is a universal \mathbb{N} -graded highest weight \mathcal{K} -module. Note that $M_{\mathcal{K}}(\lambda)$ has a unique irreducible \mathbb{N} -graded quotient module. Consequently, irreducible \mathbb{N} -graded highest weight \mathcal{K} -modules of highest weight λ are unique up to isomorphism. It then follows that any irreducible \mathbb{N} -graded highest weight \mathcal{K} -module of highest weight λ is isomorphic to the tensor product \mathcal{K} -module

$$L(\lambda|_{\widehat{\mathcal{H}}_{N_1}}) \otimes \cdots \otimes L(\lambda|_{\widehat{\mathcal{H}}_{N_d}}),$$

where $L(\lambda|_{\widehat{\mathcal{H}}_{N_1}})$ is the irreducible highest weight $\widehat{\mathfrak{sl}_{N_i}}(\mathbb{C}_q)$ -module with highest weight $\lambda|_{\widehat{\mathcal{H}}_{N_i}} \in (\widehat{\mathcal{H}}_{N_i})^*$. This proves the second assertion.

5. Fermionic Fock modules and $(\widehat{\mathfrak{gl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}_a})$ -duality

In this section, we give a family of fermionic Fock modules for $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ and establish a duality between $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ and Levi subgroups of $\mathrm{GL}_\ell(\mathbb{C})$.

We start by recalling from [G3] a fermionic representation of $\mathfrak{gl}_N(\mathbb{C}_q)$. Let N and ℓ be positive integers with $N \geq 2$ as before. Define \mathcal{C}_N^{ℓ} to be the associative (Clifford) algebra with generators

$$\psi_i^{\mu}(m), \quad \bar{\psi}_i^{\mu}(m) \quad \text{(where } 1 \leq i \leq N, \ 1 \leq \mu \leq \ell, \ m \in \mathbb{Z}),$$

subject to relations

$$\psi_{i}^{\mu}(m)\psi_{j}^{\nu}(n) + \psi_{j}^{\nu}(n)\psi_{i}^{\mu}(m) = 0, \quad \bar{\psi}_{i}^{\mu}(m)\bar{\psi}_{j}^{\nu}(n) + \bar{\psi}_{j}^{\nu}(n)\bar{\psi}_{i}^{\mu}(m) = 0,$$

$$\psi_{i}^{\mu}(m)\bar{\psi}_{j}^{\nu}(n) + \bar{\psi}_{j}^{\nu}(n)\psi_{i}^{\mu}(m) = \delta_{i,j}\delta_{\mu,\nu}\delta_{m+n,0}1$$

for $1 \leq i, j \leq N, 1 \leq \mu, \nu \leq \ell, m, n \in \mathbb{Z}$. Form generating functions

$$\psi_i^{\mu}(x) = \sum_{m \in \mathbb{Z}} \psi_i^{\mu}(m) x^{-m}$$
 and $\bar{\psi}_i^{\mu}(x) = \sum_{m \in \mathbb{Z}} \bar{\psi}_i^{\mu}(m) x^{-m}$

for $1 \leq i \leq N$, $1 \leq \mu \leq \ell$. It is clear that \mathcal{C}_N^{ℓ} is a \mathbb{Z} -graded algebra with

(5.1)
$$\operatorname{deg} \psi_i^{\mu}(n) = \operatorname{deg} \bar{\psi}_i^{\mu}(n) = -n$$

for $1 \le i \le N$, $1 \le \mu \le \ell$. Set

$$(\mathcal{C}_{N}^{\ell})^{an} = \langle \psi_{i}^{\mu}(m+1), \ \bar{\psi}_{i}^{\mu}(m) \mid 1 \leq i \leq N, \ 1 \leq \mu \leq \ell, \ m \in \mathbb{N} \rangle,$$

$$(\mathcal{C}_{N}^{\ell})^{cr} = \langle \psi_{i}^{\mu}(-m), \ \bar{\psi}_{i}^{\mu}(-m-1) \mid 1 \leq i \leq N, \ 1 \leq \mu \leq \ell, \ m \in \mathbb{N} \rangle,$$

(the subalgebras of \mathcal{C}_N^{ℓ} generated by the indicated elements). We have

$$\mathcal{C}_N^{\ell} \simeq (\mathcal{C}_N^{\ell})^{cr} \otimes (\mathcal{C}_N^{\ell})^{an}$$

as a vector space.

Denote by \mathcal{F}_N^{ℓ} the \mathcal{C}_N^{ℓ} -module generated by vector $|0\rangle$, subject to relations

(5.2)
$$\psi_i^{\mu}(m+1)|0\rangle = \bar{\psi}_i^{\mu}(m)|0\rangle = 0 \text{ for } 1 \le i \le N, \ 1 \le \mu \le \ell, \ m \in \mathbb{N}.$$

Then $\mathcal{F}_N^{\ell} = (\mathcal{C}_N^{\ell})^{cr}$ as a $(\mathcal{C}_N^{\ell})^{cr}$ -module. It is well known that \mathcal{F}_N^{ℓ} is an irreducible \mathcal{C}_N^{ℓ} -module. Defining deg $|0\rangle = 0$, we make \mathcal{F}_N^{ℓ} an \mathbb{N} -graded \mathcal{C}_N^{ℓ} -module

$$\mathcal{F}_N^{\ell} = \bigoplus_{n \in \mathbb{N}} \mathcal{F}_N^{\ell}(n)$$

with $\mathcal{F}_N^{\ell}(n)$ denoting the homogeneous subspace of degree n. It is straightforward to show that $\mathcal{F}_N^{\ell}(n)$ is finite-dimensional for every $n \in \mathbb{N}$.

Following [G3], we introduce the following normal ordering:

(5.4)
$$: \psi_i^{\mu}(m)\bar{\psi}_j^{\nu}(n) := \begin{cases} \psi_i^{\mu}(m)\bar{\psi}_j^{\nu}(n) & \text{if } m \leq n, \\ -\bar{\psi}_i^{\nu}(n)\psi_i^{\mu}(m) & \text{if } m > n \end{cases}$$

for $1 \le i, j \le N, 1 \le \mu, \nu \le \ell, m, n \in \mathbb{Z}$.

Remark 5.1. For $1 \le i, j \le N$, $1 \le \mu, \nu \le \ell$, $m, n \in \mathbb{Z}$, it is straightforward to show that the normal ordering defined in (5.4) coincides with the one defined by

$$: \psi_i^{\mu}(m)\bar{\psi}_j^{\nu}(n) := \begin{cases} \psi_i^{\mu}(m)\bar{\psi}_j^{\nu}(n) & \text{if } n \ge 0, \\ -\bar{\psi}_j^{\nu}(n)\psi_i^{\mu}(m) & \text{if } n < 0. \end{cases}$$

For $1 \leq i, j \leq N$, $m \in \mathbb{Z}$ and $\boldsymbol{a} = (a_1, \dots, a_\ell) \in (\mathbb{C}^{\times})^{\ell}$, define a field

$$\Psi_{i,j}^{\mathbf{a}}(m,x) = \begin{cases} \sum_{p=1}^{\ell} : \psi_i^p(x) \bar{\psi}_j^p(x) : & \text{if } m = 0\\ \sum_{p=1}^{\ell} a_p^m \left(: \psi_i^p(x) \bar{\psi}_j^p(q^m x) : + \delta_{i,j} \frac{q^m}{1 - q^m} \right) & \text{if } m \neq 0. \end{cases}$$

Furthermore, use the expansion

(5.5)
$$\Psi_{i,j}^{\boldsymbol{a}}(m,x) = \sum_{n \in \mathbb{Z}} \Psi_{i,j}^{\boldsymbol{a}}(n,m)x^{-n}$$

to define operators $\Psi_{i,j}^{\boldsymbol{a}}(n,m) \in \text{End}(\mathcal{F}_N^{\ell})$. Explicitly, we have

(5.6)
$$\Psi_{i,j}^{\mathbf{a}}(n,0) = \sum_{p=1}^{\ell} \sum_{k \in \mathbb{Z}} : \psi_i^p(n-k) \bar{\psi}_j^p(k) :,$$

$$\Psi_{i,j}^{\mathbf{a}}(n,m) = \sum_{p=1}^{\ell} a_p^m \left(\sum_{k \in \mathbb{Z}} q^{-mk} : \psi_i^p(n-k) \bar{\psi}_j^p(k) : + \delta_{n,0} \delta_{i,j} \frac{q^m}{1-q^m} \right)$$

for $m \neq 0$. The following is a generalization of a result of [G3]:

Proposition 5.2. Let ℓ be a positive integer and let $\mathbf{a} \in (\mathbb{C}^{\times})^{\ell}$. Then \mathcal{F}_{N}^{ℓ} is a $\widehat{\mathfrak{gl}_{N}}(\mathbb{C}_{q})$ -module with $\mathbf{k}_{0} = \ell$, $\mathbf{k}_{1} = 0$, and

(5.7)
$$E_{i,j}t_0^{m_0}t_1^{m_1} = \Psi_{i,j}^{\boldsymbol{a}}(m_0, m_1) \quad \text{for } 1 \le i, j \le N, \ m_0, m_1 \in \mathbb{Z}.$$

Furthermore, \mathcal{F}_N^{ℓ} is an integrable and restricted $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module which is \mathbb{N} -graded with finite-dimensional homogeneous subspaces.

Proof. In case that $\ell = 1$ and $\boldsymbol{a} = a_1 = 1$, this was proved in [G3]. Denote this particular $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -module simply by \mathcal{F}_N . Now, we consider the general case. Associated to the ℓ -tuple \boldsymbol{a} , we have an evaluation $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -module $(\mathcal{F}_N)(a_1) \otimes \cdots \otimes (\mathcal{F}_N)(a_\ell)$, where

$$(\mathcal{F}_N)(a_1)\otimes\cdots\otimes(\mathcal{F}_N)(a_\ell)=\mathcal{F}_N^{\otimes\ell}$$

as a vector space and where $\mathbf{k}_0 = \ell$, $\mathbf{k}_1 = 0$, and

$$E_{i,j}t_0^{m_0}t_1^{m_1}(v_1 \otimes \cdots \otimes v_\ell) = \sum_{r=1}^{\ell} a_r^{m_1}(v_1 \otimes \cdots \otimes E_{i,j}t_0^{m_0}t_1^{m_1}v_r \otimes \cdots \otimes v_\ell)$$

for $1 \leq i, j \leq N$, $m_0, m_1 \in \mathbb{Z}$, and $v_r \in \mathcal{F}_N$. Then through the natural identification of \mathcal{F}_N^{ℓ} with $(\mathcal{F}_N)(a_1) \otimes \cdots \otimes (\mathcal{F}_N)(a_{\ell})$, we obtain a $\widehat{\mathfrak{gl}}_N(\mathbb{C}_q)$ -module structure on \mathcal{F}_N^{ℓ} with the action given by (5.7) as desired.

For the second assertion, using the fact that \mathcal{F}_N^{ℓ} is a tensor product of level 1 $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules, it suffices to deal with the case with $\ell=1$. It is clear from (5.6) that the $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module \mathcal{F}_N is restricted. For the integrability, in view of Proposition 3.13, it suffices to show that for $1 \leq i \neq j \leq N$ and $m_1 \in \mathbb{Z}$,

$$\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_1)\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_2) = \Psi_{i,j}^{\boldsymbol{a}}(m_1, x_2)\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_1)$$

and $\Psi_{i,j}^{\boldsymbol{a}}(m_1,x)^2=0$ on \mathcal{F}_N . Note that

$$\psi_i^1(x_1)\psi_i^1(x_2) = -\psi_i^1(x_2)\psi_i^1(x_1), \quad \bar{\psi}_i^1(x_1)\bar{\psi}_i^1(x_2) = -\bar{\psi}_i^1(x_2)\bar{\psi}_i^1(x_1),$$

which imply $\psi_i^1(x)^2 = 0$ and $\bar{\psi}_j^1(x_1)^2 = 0$. On the other hand, as $i \neq j$, we also have $\psi_i^1(x_1)\bar{\psi}_j^1(x_2) = -\bar{\psi}_j^1(x_2)\psi_i^1(x_1)$, which implies

$$: \psi_i^1(x)\bar{\psi}_j^1(q^{m_1}x) := \psi_i^1(x)\bar{\psi}_j^1(q^{m_1}x).$$

Then

$$\Psi_{i,j}^{\boldsymbol{a}}(m_1,x) = a_1^{m_1} : \psi_i^1(x)\bar{\psi}_i^1(q^{m_1}x) := a_1^{m_1}\psi_i^1(x)\bar{\psi}_i^1(q^{m_1}x).$$

It follows that

$$\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_1)\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_2) = \Psi_{i,j}^{\boldsymbol{a}}(m_1, x_2)\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_1)$$

and

$$\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_1)\Psi_{i,j}^{\boldsymbol{a}}(m_1, x_2) = -a_1^{2m_1}\psi_i^1(x_1)\psi_i^1(x_2)\bar{\psi}_j^1(q^{m_1}x_1)\bar{\psi}_j^1(q^{m_1}x_2),$$

which implies $\Psi_{i,j}^a(m_1,x)^2=0$. This completes the proof.

Definition 5.3. For $a \in (\mathbb{C}^{\times})^{\ell}$, denote by

(5.8)
$$\rho_{\mathbf{a}}: \widehat{\mathfrak{gl}_N}(\mathbb{C}_q) \longrightarrow \operatorname{End}(\mathcal{F}_N^{\ell})$$

the representation of $\widehat{\mathfrak{gl}}_N(\mathbb{C}_q)$ on \mathcal{F}_N^{ℓ} obtained in Proposition 5.2, and we also denote by \mathcal{F}_N^a the $\widehat{\mathfrak{gl}}_N(\mathbb{C}_q)$ -module.

Next, we present an analogue in the toroidal setting of the skew $(\mathfrak{gl}_N(\mathbb{C}), \operatorname{GL}_{\ell}(\mathbb{C}))$ duality. Let n be a positive integer. Set $\operatorname{GL}_n = \operatorname{GL}_n(\mathbb{C})$, let H_n be the subgroup
of GL_n , consisting of diagonal matrices, and let N_n^+ be the subgroup consisting of
upper-triangular unipotent matrices. Following $[\operatorname{GW}]$, set

(5.9)
$$\mathbb{Z}_{++}^{n} = \{ \boldsymbol{\mu} = (\mu_1, \dots, \mu_n) \in \mathbb{Z}^n \mid \mu_1 \ge \mu_2 \ge \dots \ge \mu_n \}$$

and view each $\boldsymbol{\mu} = (\mu_1, \dots, \mu_n) \in \mathbb{Z}_{++}^n$ as a dominant regular character of H_n by

$$\mu(h) = h^{\mu} = h_1^{\mu_1} \cdots h_n^{\mu_n}$$
 for $h = \text{diag}\{h_1, \dots, h_n\} \in H_n$.

For $\boldsymbol{\mu} \in \mathbb{Z}_{++}^n$, denote by $L_{\mathrm{GL}_n}(\boldsymbol{\mu})$ the irreducible highest weight GL_n -module of highest weight $\boldsymbol{\mu}$. A basic fact is that $L_{\mathrm{GL}_n}(\boldsymbol{\mu})$ is regular, finite-dimensional, and the space of N_n^+ -fixed vectors of weight $\boldsymbol{\mu}$ is 1-dimensional.

More generally, let ℓ be a positive integer and let **I** be a partition of $\{1, \ldots, \ell\}$ with the associated equivalence relation denoted by \sim . Define

(5.10)
$$GL_{\mathbf{I}} = \{ (c_{ij})_{i,j=1}^{\ell} \in GL_{\ell} \mid c_{ij} = 0 \text{ if } i \not\sim j \},$$

the Levi subgroup of GL_{ℓ} associated to I. Correspondingly, we have the following linear algebraic subgroups of $GL_{\mathbf{I}}$:

(5.11)
$$N_{\mathbf{I}}^{+} = N_{\ell}^{+} \cap GL_{\mathbf{I}} \quad \text{and} \quad H_{\mathbf{I}} = H_{\ell} \cap GL_{\mathbf{I}}.$$

Then we identify dominant regular characters of H_I with elements of

$$(5.12) \quad \mathbb{Z}_{++}^{\mathbf{I}} := \{ \boldsymbol{\mu} = (\mu_1, \dots, \mu_\ell) \in \mathbb{Z}^\ell \mid \mu_i \ge \mu_j \text{ for } 1 \le i, j \le \ell, \ i \sim j, \ i < j \}.$$

For each $\mu \in \mathbb{Z}_{++}^{\mathbf{I}}$, denote by $L_{\mathrm{GL}_{\mathbf{I}}}(\mu)$ the finite-dimensional irreducible highest weight $\mathrm{GL}_{\mathbf{I}}$ -module of highest weight μ .

Let W be any locally regular $GL_{\mathbf{I}}$ -module of countable dimension. Then W is a direct sum of finite-dimensional irreducible regular $GL_{\mathbf{I}}$ -modules (cf. [GW]):

$$W = \bigoplus_{\boldsymbol{\mu} \in P(W)} W^{\mathbf{N}_{\mathbf{I}}^+}(\boldsymbol{\mu}) \otimes L(\boldsymbol{\mu}),$$

where $W^{\mathbf{N_{I}^{+}}}(\boldsymbol{\mu})$ denotes the space of $\mathbf{N_{I}^{+}}$ -fixed vectors of weight $\boldsymbol{\mu}$ in W and $P(W) = \{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}} \mid W^{\mathbf{N_{I}^{+}}}(\boldsymbol{\mu}) \neq 0\}.$

Definition 5.4. Let \mathcal{L} be a Lie algebra and let V be a vertex Γ-algebra. We say that $(\mathcal{L}, \operatorname{GL}_{\mathbf{I}})$ (resp. $(V, \operatorname{GL}_{\mathbf{I}})$) is a (Howe) dual pair on a locally regular $\operatorname{GL}_{\mathbf{I}}$ -module W if (i) W is an \mathcal{L} -module (resp. equivariant quasi V-module) such that the actions of \mathcal{L} (resp. V) and $\operatorname{GL}_{\mathbf{I}}$ commute. (ii) For every $\boldsymbol{\mu} \in P(W)$, the \mathcal{L} -submodule (resp. equivariant quasi V-submodule) $W^{\mathrm{N}_{\mathbf{I}}^+}(\boldsymbol{\mu})$ of W is irreducible. (iii) For any $\boldsymbol{\mu}, \boldsymbol{\nu} \in P(W)$, $W^{\mathrm{N}_{\mathbf{I}}^+}(\boldsymbol{\mu}) \cong W^{\mathrm{N}_{\mathbf{I}}^+}(\boldsymbol{\nu})$ if and only if $\boldsymbol{\mu} = \boldsymbol{\nu}$.

Recall from [FF] that \mathcal{F}_N^{ℓ} is a module for the general Lie algebra \mathfrak{gl}_{ℓ} with

(5.13)
$$E_{r,s} = \sum_{i=1}^{N} \sum_{n \in \mathbb{Z}} : \psi_i^r(-n) \bar{\psi}_i^s(n) :$$

for $1 \leq r, s \leq \ell$. It is clear that $\mathcal{F}_N^{\ell} = \bigoplus_{n \in \mathbb{Z}} \mathcal{F}_N^{\ell}(n)$ (recall (5.3)) is a direct sum of finite-dimensional \mathfrak{gl}_{ℓ} -submodules. Then this \mathfrak{gl}_{ℓ} -module structure on \mathcal{F}_N^{ℓ} gives rise to a module structure for GL_{ℓ} . Furthermore, for any partition \mathbf{I} of $\{1, \ldots, \ell\}$, \mathcal{F}_N^{ℓ} is a locally regular $GL_{\mathbf{I}}$ -module.

Definition 5.5. For any positive integer ℓ , define

$$(\mathbb{C}^{\times})_q^{\ell} = \{(a_1, \dots, a_{\ell}) \in (\mathbb{C}^{\times})^{\ell} \mid \text{ either } a_i = a_j \text{ or } a_i \notin a_j \Gamma_q \text{ for } 1 \leq i, j \in \ell\},$$

where $\Gamma_q = \{q^n \mid n \in \mathbb{Z}\}$. Furthermore, for any $\boldsymbol{a} \in (\mathbb{C}^{\times})_q^{\ell}$, we define an equivalence relation $\sim_{\boldsymbol{a}}$ on $\{1, \ldots, \ell\}$ by $i \sim_{\boldsymbol{a}} j$ if and only if $a_i = a_j$ for $1 \leq i, j \leq \ell$, and denote by $\mathbf{I}_{\boldsymbol{a}}$ the partition of $\{1, \ldots, \ell\}$ associated to $\sim_{\boldsymbol{a}}$.

The following is the main result of this section:

Theorem 5.6. For any $\mathbf{a} \in (\mathbb{C}^{\times})_q^{\ell}$ with ℓ a positive integer, $(\widehat{\mathfrak{gl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}_a})$ is a dual pair on the Fock space \mathcal{F}_N^{ℓ} with the $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -module structure afforded by ρ_a .

The rest of this section is devoted to the proof of this theorem. We start with a duality result of [F]. Set

$$(5.14) \overline{\mathfrak{gl}}_{\infty} = \mathfrak{gl}_{\infty} \oplus \mathbb{C}\mathbf{k},$$

a one-dimensional central extension of Lie algebra \mathfrak{gl}_{∞} , where

$$[A, B] = AB - BA + \sum_{i \le 0} \operatorname{tr}([E_{i,i}, A]B)\mathbf{k}$$

for $A, B \in \mathfrak{gl}_{\infty}$ (see [DJKM]). It is known (see [F]) that \mathcal{F}_N^{ℓ} is a $\overline{\mathfrak{gl}}_{\infty}$ -module of level ℓ (with $\mathbf{k} = \ell$), where

(5.15)
$$E_{mN+i,nN+j} = \sum_{p=1}^{\ell} : \psi_i^p(-m)\bar{\psi}_j^p(n) :$$

for $1 \le i, j \le N$, $m, n \in \mathbb{Z}$. Furthermore, we have (see [F]; cf. [FKRW, W1]):

Proposition 5.7. Let N and ℓ be positive integers. Then $(\overline{\mathfrak{gl}}_{\infty}, \operatorname{GL}_{\ell})$ is a dual pair on \mathcal{F}_{N}^{ℓ} . Furthermore, for any partition $\mathbf{I} = \{S_{1}, \ldots, S_{s}\}$ of $\{1, \ldots, \ell\}$, $(\overline{\mathfrak{gl}}_{\infty}^{\oplus s}, \operatorname{GL}_{\mathbf{I}})$ is a dual pair on \mathcal{F}_{N}^{ℓ} .

Note that for Proposition 5.7 the $\overline{\mathfrak{gl}}_{\infty}^{\oplus s}$ -action on \mathcal{F}_{N}^{ℓ} is given by

(5.16)
$$\iota_r(\mathbf{k}) = |S_r|, \quad \iota_r(E_{Nm+i,Nn+j}) = \sum_{p \in S_r} : \psi_i^p(-m)\bar{\psi}_j^p(n) :$$

for $1 \leq r \leq s, \ 1 \leq i, j \leq N, \ m, n \in \mathbb{Z}$, where i_r denotes the identification map of $\overline{\mathfrak{gl}}_{\infty}$ with the r-th component of $\overline{\mathfrak{gl}}_{\infty}^{\oplus s}$.

Lemma 5.8. Let ℓ be a positive integer and let $\mathbf{a} \in (\mathbb{C}^{\times})_{q}^{\ell}$ with the associated partition $\mathbf{I}_{\mathbf{a}} = \{S_{1}, \ldots, S_{s}\}$. Then a subspace U of \mathcal{F}_{N}^{ℓ} is a $\widehat{\mathfrak{gl}_{N}}(\mathbb{C}_{q})$ -submodule through $\rho_{\mathbf{a}}$ if and only if U is a $\overline{\mathfrak{gl}_{\infty}^{\oplus s}}$ -submodule. Furthermore, for any $\widehat{\mathfrak{gl}_{N}}(\mathbb{C}_{q})$ -submodules V and W of \mathcal{F}_{N}^{ℓ} , a linear map f from V to W is a $\widehat{\mathfrak{gl}_{N}}(\mathbb{C}_{q})$ -module homomorphism if and only if f is a $\overline{\mathfrak{gl}_{\infty}^{\oplus s}}$ -module homomorphism.

Proof. For $1 \leq i, j \leq N, m_0, m_1 \in \mathbb{Z}$, set

$$E_{i,j}(m_0,0) = E_{i,j}t_0^{m_0},$$

$$E_{i,j}(m_0, m_1) = E_{i,j} t_0^{m_0} t_1^{m_1} + \delta_{i,j} \delta_{m_0,0} \frac{q^{m_1}}{q^{m_1} - 1} \left(\sum_{r=1}^{\ell} a_r^{m_1} \right) \frac{\mathbf{k}_0}{\ell} \in \widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$$

for $m_1 \neq 0$. Then a subspace U of \mathcal{F}_N^{ℓ} (= $\mathcal{F}_N^{\boldsymbol{a}}$) is a $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -submodule if and only if $E_{i,j}(m_0,m_1)U \subset U$ for all $1 \leq i,j \leq N, \ m_0,m_1 \in \mathbb{Z}$. On the other hand, from (5.6) and (5.7) we have

(5.17)
$$E_{i,j}(m_0, m_1)v = \sum_{k \in \mathbb{Z}} \sum_{r=1}^{s} (b_r q^{-k})^{m_1} \left(\sum_{p \in S_r} : \psi_i^p(m_0 - k) \bar{\psi}_j^p(k) : v \right)$$

for $v \in \mathcal{F}_N^{\boldsymbol{a}}$, where $b_r = a_p$ for $p \in S_r$. This together with (5.16) implies that every $\overline{\mathfrak{gl}}_{\infty}^{\oplus s}$ -submodule of $\mathcal{F}_N^{\boldsymbol{\ell}}$ is a $\widehat{\mathfrak{gl}}_N(\mathbb{C}_q)$ -submodule of $\mathcal{F}_N^{\boldsymbol{a}}$.

Now, let V be a $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -submodule of \mathcal{F}_N^a $(=\mathcal{F}_N^\ell)$. Let $1 \leq i, j \leq N, m_0 \in \mathbb{Z}$ and $v \in V$ be fixed. Then there is a finite subset K of \mathbb{Z} such that

$$\sum_{p \in S_r} : \psi_i^p(m_0 - k) \bar{\psi}_j^p(k) : v = 0 \quad \text{for all } 1 \le r \le s, \ k \notin K.$$

Using this we get

$$\sum_{k \in K} \sum_{r=1}^{s} (b_r q^{-k})^{m_1} \left(\sum_{p \in S_r} : \psi_i^p(m_0 - k) \bar{\psi}_j^p(k) : v \right) \in V$$

for $1 \le m_1 \le s|K|$. As q is not a root of unity and $b_i \notin b_j \Gamma_q$ for $1 \le i \ne j \le s$, we have that $b_r q^{-k}$ are distinct for all $1 \le r \le s$, $k \in \mathbb{Z}$. Then we obtain

$$\sum_{p \in S_r} : \psi_i^p(m_0 - k) \bar{\psi}_j^p(k) : v \in V \quad \text{for } 1 \le r \le s, \ k \in K.$$

Note that this is also true for $k \notin K$. Combining this with (5.16), we conclude that V is a $\overline{\mathfrak{gl}}_{\infty}^{\oplus s}$ -submodule of \mathcal{F}_N^{ℓ} . This proves the first assertion. The second assertion can be proved similarly.

Proof of Theorem 5.6: Now we can conclude the proof: From Proposition 5.7, the \mathfrak{gl}_{ℓ} -action given by (5.13) on \mathcal{F}_N^{ℓ} with

$$E_{r,s} = \sum_{i=1}^{N} \sum_{n \in \mathbb{Z}} : \psi_i^r(-n) \bar{\psi}_i^s(n) :$$

for $1 \leq r, s \leq \ell$ commutes with the action of $\overline{\mathfrak{gl}}_{\infty}$ with

$$E_{mN+i,nN+j} = \sum_{r=1}^{\ell} : \psi_i^r(-m)\bar{\psi}_j^r(n) :$$

for $m, n \in \mathbb{Z}$, $1 \leq i, j \leq N$. Then we see that the \mathfrak{gl}_{ℓ} -action given by (5.13) on \mathcal{F}_N^{ℓ} commutes with the $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -action given by (5.7) and (5.6) through ρ_a . Since $(\widehat{\mathfrak{gl}_{\infty}^{\ell}}, \operatorname{GL}_{\mathbf{I}_a})$ is a dual pair on \mathcal{F}_N^{ℓ} , it follows immediately from Lemma 5.8 that $(\widehat{\mathfrak{gl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}_a})$ is a dual pair on \mathcal{F}_N^{ℓ} .

Remark 5.9. With $\mathfrak{gl}_N(\mathbb{C})$ a subalgebra of $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$, the $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -module \mathcal{F}_N^a is naturally a $\mathfrak{gl}_N(\mathbb{C})$ -module. Note that the $\mathfrak{gl}_N(\mathbb{C})$ -action on \mathcal{F}_N^ℓ is independent of \boldsymbol{a} and the degree-zero subspace $\mathcal{F}_N^\ell(0)$ is a $\mathfrak{gl}_N(\mathbb{C})$ -submodule of \mathcal{F}_N^a . It was known (see [H2]) that $(\mathfrak{gl}_N(\mathbb{C}), \mathrm{GL}_\ell)$ is a dual pair on $\mathcal{F}_N^\ell(0)$. In view of this, Theorem 5.6 can be viewed as an extension of the skew $(\mathfrak{gl}_N(\mathbb{C}), \mathrm{GL}_\ell)$ -duality.

6.
$$(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\operatorname{GL}_{\mathbf{I}_{\boldsymbol{a}}})$$
-duality and Fermionic realization

Let ℓ be a positive integer as before. Recall from Lemma 5.2 that for every $\boldsymbol{a} \in (\mathbb{C}^{\times})^{\ell}$, we have an N-graded integrable restricted $\widehat{\mathfrak{sl}}_{N}(\mathbb{C}_{q})$ -module $\mathcal{F}_{N}^{\boldsymbol{a}} (= \mathcal{F}_{N}^{\ell})$ of level ℓ . Then by Theorem 3.15, $\mathcal{F}_{N}^{\boldsymbol{a}}$ is naturally an N-graded (\mathbb{Z}, χ_{q}) -equivariant quasi $L_{\widehat{\mathfrak{sl}}_{\infty}}(\ell, 0)$ -module. In this section, we continue to show that for any $\boldsymbol{a} \in (\mathbb{C}^{\times})_{q}^{\ell}$, $\mathcal{F}_{N}^{\boldsymbol{a}}$ is a direct sum of irreducible quasi $L_{\widehat{\mathfrak{sl}}_{\infty}}(\ell, 0)$ -submodules with finite

multiplicities. Furthermore, we prove that every irreducible \mathbb{N} -graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module is isomorphic to an irreducible submodule of \mathcal{F}_N^a for some $\mathbf{a} \in (\mathbb{C}^{\times})_q^{\ell}$.

We start with some notations. First, for $\mu \in \mathbb{Z}$, we define integers $\dot{\mu}$ and $\ddot{\mu}$ by

(6.1)
$$\mu = \dot{\mu}N + \ddot{\mu} \quad \text{with } 1 \le \ddot{\mu} \le N.$$

Definition 6.1. For any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_\ell) \in \mathbb{Z}^\ell$ and $\boldsymbol{a} = (a_1, \dots, a_\ell) \in (\mathbb{C}^\times)^\ell$, we define a linear functional $\eta_{\boldsymbol{\mu},\boldsymbol{a}}$ on $\widehat{\mathcal{H}}$ by $\eta_{\boldsymbol{\mu},\boldsymbol{a}}(\mathbf{k}_1) = 0$ and

(6.2)
$$\eta_{\boldsymbol{\mu},\boldsymbol{a}}(h_{i,n}) = \sum_{k=1}^{\ell} (a_k q^{-\dot{\mu}_k})^n \delta_{\ddot{\mu}_k,i}$$

for $1 \le i \le N$, $n \in \mathbb{Z}$.

Remark 6.2. For $1 \leq i \leq N$, define $\Lambda_i \in \mathcal{H}^*$ by

(6.3)
$$\Lambda_i(h_{j,0}) = \delta_{i,j} \quad \text{for } 1 \le j \le N.$$

For $\boldsymbol{\mu} \in \mathbb{Z}^{\ell}$ and $\boldsymbol{a} \in (\mathbb{C}^{\times})^{\ell}$, set $\boldsymbol{\lambda} = (\Lambda_{\ddot{\mu}_1}, \dots, \Lambda_{\ddot{\mu}_{\ell}})$ and $\boldsymbol{c} = (a_1 q^{-\dot{\mu}_1}, \dots, a_{\ell} q^{-\dot{\mu}_{\ell}})$. Then we have $\eta_{\boldsymbol{\mu},\boldsymbol{a}} = \eta_{\boldsymbol{\lambda},\boldsymbol{c}}$ which was defined in (4.11). Furthermore, let $\boldsymbol{\nu} \in \mathbb{Z}^{\ell}$ and $\boldsymbol{b} \in (\mathbb{C}^{\times})^{\ell}$. Then it follows from Lemma 4.11 that $\eta_{\boldsymbol{\mu},\boldsymbol{a}} = \eta_{\boldsymbol{\nu},\boldsymbol{b}}$ if and only if there is a permutation σ on $\{1,\dots,\ell\}$ and an $\boldsymbol{m} = (m_1,\dots,m_{\ell}) \in \mathbb{Z}^{\ell}$ such that

$$\sigma(\boldsymbol{\mu}) = \boldsymbol{\nu} + N\boldsymbol{m} = (\nu_1 + Nm_1, \dots, \nu_{\ell} + Nm_{\ell}),$$

and that

$$\sigma(\boldsymbol{a}) = q^{\boldsymbol{m}}\boldsymbol{b} := (q^{m_1}b_1, \dots, q^{m_\ell}b_\ell).$$

Remark 6.3. For $1 \le i \le N$, $\mu \in \mathbb{Z}_+$, set

(6.4)
$$J_{i,\mu} = \{ k \in \mathbb{Z}_{\geq 0} \mid kN + i \leq \mu \}.$$

From definition, for $1 \le i \le N-1$ we have

(6.5)
$$J_{i,\mu} = \begin{cases} J_{i+1,\mu} \cup \{\dot{\mu}\} & \text{if } i = \ddot{\mu} \\ J_{i,\mu} = J_{i+1,\mu} & \text{otherwise.} \end{cases}$$

We also have

(6.6)
$$J_{1,\mu} = \{0, 1, \dots, \dot{\mu}\} = \begin{cases} J_{N,\mu} & \text{if } \ddot{\mu} = N \\ J_{N,\mu} \cup \{\dot{\mu}\} & \text{otherwise.} \end{cases}$$

On the other hand, for a negative integer μ , we set

(6.7)
$$\bar{J}_{i,\mu} = \{k \in \mathbb{Z}_{\geq 0} \mid (k+1)N - i + 1 \leq -\mu\} = J_{N+1-i,-\mu}.$$

For $1 \leq i \leq N-1$, we have $\bar{J}_{i+1,\mu} = \bar{J}_{i,\mu} \cup \{-\dot{\mu}-1\}$ if $i = \ddot{\mu}$ and $\bar{J}_{i,\mu} = \bar{J}_{i+1,\mu}$ otherwise. We also have

(6.8)
$$\bar{J}_{N,\mu} = \begin{cases} \{0, 1, \dots, -\dot{\mu} - 2\} = \bar{J}_{1,\mu} & \text{if } \ddot{\mu} = N \\ \{0, 1, \dots, -\dot{\mu} - 1\} = \bar{J}_{1,\mu} \cup \{-\dot{\mu} - 1\} & \text{otherwise.} \end{cases}$$

Set $I_N = \{1, 2, ..., N\}$ for convenience. Define maps π and $\bar{\pi} : \mathbb{Z} \times I_N \to \mathbb{Z}$ by

(6.9)
$$\pi(n,i) = nN - i, \quad \bar{\pi}(n,i) = nN + i - 1 \quad \text{for } (n,i) \in \mathbb{Z} \times I_N.$$

Note that both π and $\bar{\pi}$ are bijections and

$$\pi(\mathbb{Z}_+ \times I_N) = \mathbb{Z}_{\geq 0}, \quad \pi(\mathbb{Z}_- \times I_N) = \mathbb{Z}_-, \quad \bar{\pi}(\mathbb{Z}_{\geq 0} \times I_N) = \mathbb{Z}_{\geq 0}, \quad \bar{\pi}(\mathbb{Z}_- \times I_N) = \mathbb{Z}_-.$$

For convenience, we define ϕ_m^p , $\bar{\phi}_m^p$ for $1 \leq p \leq \ell$, $m \in \mathbb{Z}$ by

$$\phi_{\pi(n,i)}^p = \psi_i^p(n), \quad \bar{\phi}_{\bar{\pi}(n,i)}^p = \bar{\psi}_i^p(n)$$

for $1 \le i \le N$, $n \in \mathbb{Z}$, i.e.,

(6.10)
$$\phi_{nN-i}^p = \psi_i^p(n), \qquad \bar{\phi}_{nN+i-1}^p = \bar{\psi}_i^p(n).$$

Then

(6.11)
$$\phi_m^p \phi_n^q + \phi_n^q \phi_m^p = 0, \quad \bar{\phi}_m^p \bar{\phi}_n^q + \bar{\phi}_n^q \bar{\phi}_m^p = 0,$$

(6.12)
$$\phi_m^p \bar{\phi}_n^q + \bar{\phi}_n^q \phi_m^p = \delta_{p,q} \delta_{m+n+1,0}$$

for $1 \leq p, q \leq \ell, \ m, n \in \mathbb{Z}$.

For $\mu \in \mathbb{Z}$, $1 \leq p \leq \ell$, we define an element $A^p(\mu)$ of the Clifford algebra \mathcal{C}_N^{ℓ} in terms of ϕ_n^p and $\bar{\phi}_n^p$ for $n \in \mathbb{Z}$ by

(6.13)
$$A^{p}(\mu) = \begin{cases} \phi_{-\mu}^{p} \cdots \phi_{-2}^{p} \phi_{-1}^{p} & \text{for } \mu \geq 1, \\ 1 & \text{for } \mu = 0, \\ \bar{\phi}_{\mu}^{p} \cdots \bar{\phi}_{-2}^{p} \bar{\phi}_{-1}^{p} & \text{for } \mu \leq -1. \end{cases}$$

Let μ be a positive integer. Then

(6.14)
$$\phi_n^p A^p(\mu)|0\rangle = 0 \quad \text{for } n \ge 0, \quad \bar{\phi}_n^p A^p(\mu)|0\rangle = 0 \quad \text{for } n \ge \mu,$$

(6.15)
$$\phi_{-n}^{p} A^{p}(\mu) = 0 \text{ for } 1 \le n \le \mu.$$

(Note that $\phi_n^p A^p(\mu) = (-1)^\mu A^p(\mu) \phi_n^p$ and $\phi_{-n}^p \phi_{-n}^p = 0$.) Similarly, we have

(6.16)
$$\bar{\phi}_n^p A^p(-\mu)|0\rangle = 0 \quad \text{for } n \ge 0, \quad \phi_n^p A^p(-\mu)|0\rangle = 0 \quad \text{for } n \ge \mu,$$

(6.17)
$$\bar{\phi}_{-n}^p A^p(-\mu) = 0 \text{ for } 1 \le n \le \mu.$$

We also have

(6.18)
$$A^{p}(\mu)A^{q}(\nu) = \pm A^{q}(\nu)A^{p}(\mu) \quad \text{for } 1 \le p \ne q \le \ell, \ \mu, \nu \in \mathbb{Z}.$$

Using these properties, we have (cf. [W1]):

Proposition 6.4. Let $a \in (\mathbb{C}^{\times})_q^{\ell}$ with ℓ a positive integer and let $\mu \in \mathbb{Z}_{++}^{\mathbf{I}_a}$. Set

(6.19)
$$v_{\boldsymbol{\mu}} = A(\boldsymbol{\mu})|0\rangle = A^{1}(\mu_{1}) \cdots A^{\ell}(\mu_{\ell})|0\rangle \in \mathcal{F}_{N}^{\ell}.$$

Then v_{μ} is a joint highest weight vector in $\mathcal{F}_{N}^{a} = \mathcal{F}_{N}^{\ell}$ in the sense that

(6.20)
$$\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+ v_{\mu} = 0, \quad hv_{\mu} = \eta_{\mu,a}(h)v_{\mu} \quad \text{for } h \in \widehat{\mathcal{H}}$$

and that

(6.21)
$$N_{\mathbf{I}_a}^+ v_{\mu} = v_{\mu}, \quad hv_{\mu} = \mu(h)v_{\mu} \quad \text{for } h \in \mathbf{H}_{\mathbf{I}_a}.$$

Proof. First, from (6.16)-(6.18) we have the following facts for $1 \le p \le \ell$, $n \in \mathbb{Z}$:

(6.22)
$$\phi_n^p A(\boldsymbol{\mu})|0\rangle = 0 \quad \text{if } n \ge -\mu_p,$$

(6.23)
$$\bar{\phi}_n^p A(\boldsymbol{\mu})|0\rangle = 0 \quad \text{if } n \ge \mu_p.$$

(A) Annihilation property $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)^+v_{\mu}=0.$

Let $1 \le i, j \le N$, $n \in \mathbb{Z}$ such that either n > 0 or n = 0 and i < j. We claim

$$(6.24) : \psi_i^p(n-k)\bar{\psi}_i^p(k) : A(\boldsymbol{\mu})|0\rangle = 0 \text{for all } 1 \le p \le \ell, \ k \in \mathbb{Z}.$$

(I) The case with n = 0 and i < j. Note that it is equivalent to prove

$$\sum_{k \in \mathbb{Z}} f(k) : \psi_i^p(-k) \bar{\psi}_j^p(k) : A(\boldsymbol{\mu})|0\rangle = 0$$

for any complex-valued function f on \mathbb{Z} . From definition, for $1 \leq p, q \leq \ell$ we have

$$(6.25) \qquad \sum_{k \in \mathbb{Z}} f(k) : \psi_{i}^{p}(-k) \bar{\psi}_{j}^{q}(k) :$$

$$= \sum_{k \geq 0} f(k) \psi_{i}^{p}(-k) \bar{\psi}_{j}^{q}(k) - \sum_{k \leq -1} f(k) \bar{\psi}_{j}^{q}(k) \psi_{i}^{p}(-k)$$

$$= \sum_{k \geq 0} f(k) \phi_{-kN-i}^{p} \bar{\phi}_{kN+j-1}^{q} - \sum_{k \leq -1} f(k) \bar{\phi}_{kN+j-1}^{q} \phi_{-kN-i}^{p}$$

$$= \sum_{k \geq 0} \left(f(k) \phi_{-kN-i}^{p} \bar{\phi}_{kN+j-1}^{q} - f(-k-1) \bar{\phi}_{-(k+1)N+j-1}^{q} \phi_{(k+1)N-i}^{p} \right).$$

Let $k \geq 0$. Assume $\mu_p \geq 0$. We have $\phi_{(k+1)N-i}^p A(\boldsymbol{\mu})|0\rangle = 0$ as $(k+1)N-i \geq 0$. One the other hand, if $kN+j > \mu_p$, then $\bar{\phi}_{kN+j-1}^p A(\boldsymbol{\mu})|0\rangle = 0$. If $kN+j \leq \mu_p$, then $kN+i < kN+j \leq \mu_p$. In this case, we have $\phi_{-kN-i}^p \bar{\phi}_{kN+j-1}^p A(\boldsymbol{\mu})|0\rangle = 0$ as

$$\phi^{p}_{-kN-i}\bar{\phi}^{p}_{kN+j-1}A(\boldsymbol{\mu}) = -\bar{\phi}^{p}_{kN+j-1}\phi^{p}_{-kN-i}A(\boldsymbol{\mu}) = 0.$$

Thus $\sum_{k\in\mathbb{Z}} f(k) : \psi_i^p(-k) \bar{\psi}_j^p(k) : A(\boldsymbol{\mu})|0\rangle = 0$ from (6.25). The case with $\mu_p < 0$ follows from a similar argument.

(II) The case with n > 0. Note that

$$: \psi_i^p(n-k)\bar{\psi}_i^p(k) := \psi_i^p(n-k)\bar{\psi}_i^p(k) = -\bar{\psi}_i^p(k)\psi_i^p(n-k).$$

We here prove this for the case $\mu_p > 0$ while the case $\mu_p < 0$ follows from a similar argument. If k < n, we have

$$\psi_i^p(n-k)A(\boldsymbol{\mu})|0\rangle = \pm A(\boldsymbol{\mu})\psi_i^p(n-k)|0\rangle = 0.$$

Assuming $k \geq n$, we have

$$\psi_i^p(n-k)\bar{\psi}_i^p(k)A(\boldsymbol{\mu})|0\rangle = \phi_{(n-k)N-i}^p\bar{\phi}_{kN+i-1}^pA(\boldsymbol{\mu})|0\rangle.$$

If $kN + j > \mu_p$, we have $\bar{\phi}_{kN+j-1}^p A(\boldsymbol{\mu})|0\rangle = 0$. If $kN + j \leq \mu_p$, we have

$$(n-k)N - i = -kN - j + (nN + j - i) > -\mu_p,$$

so that $\phi_{(n-k)N-i}^p A(\boldsymbol{\mu}) = 0$. Hence

$$\phi^{p}_{(n-k)N-i}\bar{\phi}^{p}_{kN+j-1}A(\mu)|0\rangle = -\bar{\phi}^{p}_{kN+j-1}\phi^{p}_{(n-k)N-i}A(\mu)|0\rangle = 0.$$

Therefore we have : $\psi_i^p(n-k)\overline{\psi}_j^p(k)$: $A(\boldsymbol{\mu})|0\rangle = 0$ for $\mu_p > 0$. Then it follows from (4.4) and (5.7) (and (5.6)) that $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)^+v_{\boldsymbol{\mu}} = 0$.

(B) Determination of the highest weight.

For $1 \le i \le N$, $n \in \mathbb{Z}$, from (5.6) and (5.7) we have

$$E_{i,i}t_1^n = \sum_{p=1}^{\ell} \sum_{k \in \mathbb{Z}} (a_p q^{-k})^n : \psi_i^p(-k) \bar{\psi}_i^p(k) : +(1 - \delta_{n,0}) \frac{a_p^n q^n}{1 - q^n}.$$

Then

$$h_{i,n} = (E_{i,i} - E_{i+1,i+1})t_1^n$$

$$= \sum_{p=1}^{\ell} \sum_{k \in \mathbb{Z}} (a_p q^{-k})^n \left(: \psi_i^p(-k) \bar{\psi}_i^p(k) : - : \psi_{i+1}^p(-k) \bar{\psi}_{i+1}^p(k) : \right)$$

for $1 \le i \le N - 1$, $n \in \mathbb{Z}$,

$$h_{N,0} = \mathbf{k}_0 - E_{1,1} + E_{N,N} = \ell + \sum_{p=1}^{\ell} \sum_{k \in \mathbb{Z}} \left(: \psi_N^p(-k) \bar{\psi}_N^p(k) : - : \psi_1^p(-k) \bar{\psi}_1^p(k) : \right),$$

$$h_{N,n} = -q^n E_{1,1} t_1^n + E_{N,N} t_1^n$$

$$= \sum_{p=1}^{\ell} \sum_{k \in \mathbb{Z}} (a_p q^{-k})^n \left(: \psi_N^p(-k) \bar{\psi}_N^p(k) : -q^n : \psi_1^p(-k) \bar{\psi}_1^p(k) : \right) + a_p^n q^n$$

for $n \neq 0$.

Claim B: For $1 \le p \le \ell$, we have

$$\sum_{k\in\mathbb{Z}} f(k) : \psi_i^p(-k)\bar{\psi}_i^p(k) : A(\boldsymbol{\mu})|0\rangle = \begin{cases} \left(\sum_{k\in J_{i,\mu_p}} f(k)\right) A(\boldsymbol{\mu})|0\rangle & \text{if } \mu_p > 0, \\ -\left(\sum_{k\in \bar{J}_{i,\mu_p}} f(k)\right) A(\boldsymbol{\mu})|0\rangle & \text{if } \mu_p < 0, \end{cases}$$

where f is any complex-valued function on \mathbb{Z} .

We here give a proof for the case $\mu_p > 0$ while the case $\mu_p < 0$ follows from a similar argument. Let $1 \le i \le N$, $1 \le p \le \ell$, $k \ge 0$. We have

$$\bar{\phi}_{-kN-i}^p \phi_{kN+i-1}^p A(\mu) |0\rangle = \pm \bar{\phi}_{-kN-i}^p A(\mu) \phi_{kN+i-1}^p |0\rangle = 0.$$

On the other hand, we have

$$\phi_{-kN-i}^p \bar{\phi}_{kN+i-1}^p A(\boldsymbol{\mu})|0\rangle = \begin{cases} 0 & \text{if } kN+i > \mu_p, \\ A(\boldsymbol{\mu})|0\rangle & \text{if } kN+i \leq \mu_p, \end{cases}$$

noticing that $\bar{\phi}_{kN+i-1}^p A(\boldsymbol{\mu})|0\rangle = \pm A(\boldsymbol{\mu})\bar{\phi}_{kN+i-1}^p|0\rangle = 0$ if $kN+i>\mu_p$ and

$$\phi^p_{-kN-i}\bar{\phi}^p_{kN+i-1}A(\boldsymbol{\mu})|0\rangle = -\bar{\phi}^p_{kN+i-1}\phi^p_{-kN-i}A(\boldsymbol{\mu})|0\rangle + A(\boldsymbol{\mu})|0\rangle = A(\boldsymbol{\mu})|0\rangle$$

if $kN + i \leq \mu_p$, as $\phi_{-kN-i}^p A(\boldsymbol{\mu}) = 0$. Then Claim B follows from (6.25) in this case. Now, using Claim B and Remark 6.3, we get $h_{i,n}A(\boldsymbol{\mu})|0\rangle = \eta_{\boldsymbol{\mu},\boldsymbol{a}}(h_{i,n})A(\boldsymbol{\mu})|0\rangle$ for $1 \leq i \leq N, \ n \in \mathbb{Z}$, as desired.

The other assertions can be proved similarly and we omit the details. \Box

As the main result of this section, we have:

Theorem 6.5. Let ℓ be a positive integer and let $\mathbf{a} \in (\mathbb{C}^{\times})_q^{\ell}$. Then both $(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}_{\mathbf{a}}})$ and $(L_{\widehat{\mathfrak{sl}_\infty}}(\ell, 0), \operatorname{GL}_{\mathbf{I}_{\mathbf{a}}})$ are dual pairs on $\mathcal{F}_N^{\mathbf{a}}$. Furthermore, we have

(6.26)
$$\mathcal{F}_{N}^{a} = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}}} L(\eta_{\boldsymbol{\mu},a}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\mu})$$

as an $(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}_q})$ -module.

Proof. We write I simply for I_a . From Theorem 5.6 we have

(6.27)
$$\mathcal{F}_{N}^{a} = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}}} (\mathcal{F}_{N}^{a})^{N_{\mathbf{I}}^{+}}(\boldsymbol{\mu}) \otimes L_{GL_{\mathbf{I}}}(\boldsymbol{\mu})$$

as a $(\widehat{\mathfrak{gl}_N}(\mathbb{C}_q), \operatorname{GL}_{\mathbf{I}})$ -module, where $(\mathcal{F}_N^a)^{\operatorname{N}_{\mathbf{I}}^+}(\boldsymbol{\mu})$ if nonzero is an irreducible $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -module. As $(\mathcal{F}_N^a)^{\operatorname{N}_{\mathbf{I}}^+}(\boldsymbol{\mu}) \neq 0$ for $\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}}$ by Proposition 6.4, it must be an irreducible $\widehat{\mathfrak{gl}_N}(\mathbb{C}_q)$ -module, so that the identity matrix I_N acts as a scalar. Then it follows from (2.7) that $(\mathcal{F}_N^a)^{\operatorname{N}_{\mathbf{I}}^+}(\boldsymbol{\mu})$ is an irreducible $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module. Also from Proposition 6.4, $v_{\boldsymbol{\mu}}$ is a (nonzero) highest weight vector in $(\mathcal{F}_N^a)^{\operatorname{N}_{\mathbf{I}}^+}(\boldsymbol{\mu})$ of weight $\eta_{\boldsymbol{\mu},\boldsymbol{a}}$. It follows that $(\mathcal{F}_N^a)^{\operatorname{N}_{\mathbf{I}}^+}(\boldsymbol{\mu})$ is a highest weight irreducible $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module isomorphic to $L(\eta_{\boldsymbol{\mu},\boldsymbol{a}})$. Thus we have the decomposition (6.26). Finally, the dual pair assertion follows from the fact that $L(\eta_{\boldsymbol{\mu},\boldsymbol{a}}) \cong L(\eta_{\boldsymbol{\nu},\boldsymbol{a}})$ if and only if $\eta_{\boldsymbol{\mu},\boldsymbol{a}} = \eta_{\boldsymbol{\nu},\boldsymbol{a}}$, which is equivalent to $\boldsymbol{\mu} = \boldsymbol{\nu}$ by Remark 6.2.

Remark 6.6. Note that for any partition **I** of $\{1,\ldots,\ell\}$, there exists $\boldsymbol{a} \in (\mathbb{C}^{\times})_q^{\ell}$ such that $\mathbf{I} = \mathbf{I}_a$. It then follows from Theorem 6.5 that every irreducible regular module for the Levi subgroup $\mathrm{GL}_{\mathbf{I}}$ of GL_{ℓ} occurs in the decomposition (6.27). On the other hand, every Levi subgroup of GL_{ℓ} is conjugate to $\mathrm{GL}_{\mathbf{I}}$ for some partition **I** of $\{1,\ldots,\ell\}$.

As the first application of Theorem 6.5, we have the following realization of irreducible N-graded integrable restricted $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules of level ℓ , or equivalently irreducible N-graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}}_\infty}(\ell, 0)$ -modules:

Theorem 6.7. Let ℓ be a positive integer. Then for any $\mathbf{a} \in (\mathbb{C}^{\times})_{q}^{\ell}$ and $\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}$, the space $(\mathcal{F}_{N}^{\mathbf{a}})^{\mathbb{N}_{\mathbf{I}_{a}}^{+}}(\boldsymbol{\mu})$ of the $\mathbb{N}_{\mathbf{I}_{a}}^{+}$ -fixed points of weight $\boldsymbol{\mu}$ in $\mathcal{F}_{N}^{\mathbf{a}}$ is an irreducible \mathbb{N} -graded (\mathbb{Z}, χ_{q}) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module. On the other hand, every irreducible \mathbb{N} -graded (\mathbb{Z}, χ_{q}) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)$ -module can be realized this way.

Proof. The first assertion follows from Theorem 6.5. As for the second assertion, recall from (the proof of) Theorem 6.5 that $(\mathcal{F}_N^a)^{\mathrm{N}_{\mathbf{I}_a}^+}(\boldsymbol{\mu}) \cong L(\eta_{\boldsymbol{\mu},a})$. Thus it suffices to show that for any pair $(\boldsymbol{\lambda}, \boldsymbol{c}) \in (P_+)^k \times (\mathbb{C}^\times)^k$ in Theorem 4.10, there exist $\boldsymbol{a} \in (\mathbb{C}^\times)_q^\ell$ and $\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_a}$ such that $\eta_{\boldsymbol{\mu},a} = \eta_{\boldsymbol{\lambda},c}$. In view of Remark 4.7, it suffices to consider the case with $k = \ell$ and $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_\ell)$ such that $\lambda_1, \dots, \lambda_\ell \in P_+$ are all fundamental.

Define an equivalence relation on $\{1,\ldots,\ell\}$ such that $i \sim j$ if and only if $c_i = q^n c_j$ for some $n \in \mathbb{Z}$. Let $\mathbf{I} = \{S_1,\ldots,S_s\}$ be the partition of $\{1,\ldots,\ell\}$ associated to \sim . For $1 \leq r \leq s$, pick a representative $b_r \in \{c_i \mid i \in S_r\}$. Define $\mathbf{a} = (a_1,\ldots,a_\ell) \in (\mathbb{C}^\times)^\ell$ where $a_k = b_r$ if $k \in S_r$. Then \mathbf{a} lies in $(\mathbb{C}^\times)^\ell_q$ and $\mathbf{I} = \mathbf{I}_{\mathbf{a}}$ is the partition of $\{1,\ldots,\ell\}$ associated to \mathbf{a} . As $\lambda_1,\ldots,\lambda_\ell \in P_+$ are all fundamental, for each $1 \leq i \leq \ell$ there is a unique $m_i \in \{1,\ldots,N\}$ such that $\lambda_i(h_{j,0}) = \delta_{j,m_i}$ for $1 \leq j \leq N$. On the other hand, for each $1 \leq i \leq \ell$, there exist unique $1 \leq r \leq s$ and $n_i \in \mathbb{Z}$ such that $c_i = b_r q^{-n_i}$. Then define $\mu_i = N n_i + m_i$. Since $\eta_{\tau \lambda, \tau c} = \eta_{\lambda, c}$ for any permutation τ on $\{1,\ldots,\ell\}$, we can assume $\mu_i \geq \mu_j$ for $i \sim j$ and i < j. Then the ℓ -tuple $\boldsymbol{\mu} := (\mu_1,\ldots,\mu_\ell)$ lies in $\mathbb{Z}^{\mathbf{I}}_{++}$, and we have $\eta_{\boldsymbol{\mu},\boldsymbol{a}} = \eta_{\lambda,c}$, as desired.

Remark 6.8. Note that although the \mathbb{Z} -graded vertex algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ has infinite-dimensional homogenous subspaces, it follows from Theorem 6.7 and Proposition 5.2 that all the homogenous subspaces of every irreducible \mathbb{N} -graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ -module are finite-dimensional.

7.
$$L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)\to L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$$
 branching

Recall that for any nonzero complex number ℓ , the \mathbb{Z} -graded vertex algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ contains \mathfrak{sl}_{∞} as its degree-one subspace which generates $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ as a vertex algebra. Let ℓ and ℓ' be positive integers which are fixed throughout this section. It is known that $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ as an $\widehat{\mathfrak{sl}_{\infty}}$ -module is isomorphic to the submodule generated by the vacuum vector $\mathbf{1}\otimes\mathbf{1}$ in $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$. This gives rise to a \mathbb{Z} -graded vertex algebra embedding of $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ into $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$, which is uniquely determined by

$$(7.1) u \mapsto u \otimes \mathbf{1} + \mathbf{1} \otimes u \quad \text{for } u \in \mathfrak{sl}_{\infty}.$$

It is clear that this is also a vertex \mathbb{Z} -algebra embedding with the particular \mathbb{Z} -module structures defined in Section 3.2. We then view $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ as a vertex \mathbb{Z} -subalgebra of $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$. In this section, we shall determine the $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)\to L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ branching rule for irreducible equivariant quasi modules.

In view of Proposition 4.12 and Theorem 6.7, every irreducible N-graded (\mathbb{Z}, χ_q) -equivariant quasi module for $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0) \otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell', 0)$ is isomorphic to

$$L(\eta_{\boldsymbol{\mu},\boldsymbol{a}})\otimes L(\eta_{\boldsymbol{\nu},\boldsymbol{b}})$$

for some pairs

$$(\boldsymbol{a}, \boldsymbol{\mu}) \in (\mathbb{C}^{\times})_{q}^{\ell} \times \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}}} \quad \text{and} \quad (\boldsymbol{b}, \boldsymbol{\nu}) \in (\mathbb{C}^{\times})_{q}^{\ell'} \times \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{b}}}.$$

For any $1 \leq i \leq \ell$, set $m_i = 0$ if $a_i \notin b_j \Gamma_q$ for any j and otherwise set m_i to be the unique integer such that $q^{m_i}a_i \in \{b_1, \ldots, b_{\ell'}\}$. Furthermore, set $\boldsymbol{m} = (m_1, \ldots, m_{\ell}) \in \mathbb{Z}^{\ell}$. Recall from Remark 6.2 that

$$\eta_{\boldsymbol{\mu},\boldsymbol{a}} = \eta_{\boldsymbol{\mu}+N\boldsymbol{m},q^{\boldsymbol{m}}\boldsymbol{a}}.$$

It is straightforward to see that $q^{\boldsymbol{m}}\boldsymbol{a} \in (\mathbb{C}^{\times})_q^{\ell}, \boldsymbol{\mu} + N\boldsymbol{m} \in \mathbb{Z}_{++}^{\mathbf{I}_q \boldsymbol{m}_{\boldsymbol{a}}}$, and

$$(q^{\boldsymbol{m}}\boldsymbol{a},\boldsymbol{b}) \in (\mathbb{C}^{\times})_{q}^{\ell+\ell'}.$$

In view of this, it suffices to consider $L(\eta_{\mu,a}) \otimes L(\eta_{\nu,b})$ with $(a,b) \in (\mathbb{C}^{\times})_q^{\ell+\ell'}$.

View $GL_{\ell} \times GL_{\ell'}$ as a subgroup of $GL_{\ell+\ell'}$ in the obvious way. Then we have a natural group embedding

$$\operatorname{GL}_{\mathbf{I}_{\boldsymbol{a}}} \times \operatorname{GL}_{\mathbf{I}_{\boldsymbol{b}}} (\subset \operatorname{GL}_{\ell} \times \operatorname{GL}_{\ell'}) \hookrightarrow \operatorname{GL}_{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}} (\subset \operatorname{GL}_{\ell+\ell'}).$$

Write the $\mathrm{GL}_{\mathbf{I}_{(a,b)}} \to \mathrm{GL}_{\mathbf{I}_a} \times \mathrm{GL}_{\mathbf{I}_b}$ branching as

(7.2)
$$\operatorname{Res}_{\operatorname{GL}_{\mathbf{I}_{a}} \times \operatorname{GL}_{\mathbf{I}_{b}}}^{\operatorname{GL}_{\mathbf{I}_{(a,b)}}} L_{\operatorname{GL}_{\mathbf{I}_{(a,b)}}}(\boldsymbol{\xi}) = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}, \ \boldsymbol{\nu} \in \mathbb{Z}_{++}^{\mathbf{I}_{b}}} D_{\boldsymbol{\mu},\boldsymbol{\nu}}^{\boldsymbol{\xi}} \left(L_{\operatorname{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\mu}) \otimes L_{\operatorname{GL}_{\mathbf{I}_{b}}}(\boldsymbol{\nu}) \right)$$

for $\boldsymbol{\xi} \in \mathbb{Z}_{++}^{\mathbf{I}_{(a,b)}}$, where $D_{\boldsymbol{\mu},\boldsymbol{\nu}}^{\boldsymbol{\xi}}$ denotes the (finite) multiplicity of the $\mathrm{GL}_{\mathbf{I}_a} \times \mathrm{GL}_{\mathbf{I}_b}$ module $L_{\mathrm{GL}_{\mathbf{I}_a}}(\boldsymbol{\mu}) \otimes L_{\mathrm{GL}_{\mathbf{I}_b}}(\boldsymbol{\nu})$ in $L_{\mathrm{GL}_{\mathbf{I}_{(a,b)}}}(\boldsymbol{\xi})$. Using this decomposition we have the following main result of this section:

Theorem 7.1. Let ℓ, ℓ' be positive integers and let

$$oldsymbol{a} \in (\mathbb{C}^{ imes})_q^\ell, \ oldsymbol{b} \in (\mathbb{C}^{ imes})_q^{\ell'}, \ oldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{oldsymbol{a}}}, \ oldsymbol{
u} \in \mathbb{Z}_{++}^{\mathbf{I}_{oldsymbol{b}}}$$

such that $(\boldsymbol{a}, \boldsymbol{b}) \in (\mathbb{C}^{\times})_{q}^{\ell+\ell'}$. Then $L(\eta_{\boldsymbol{\mu}, \boldsymbol{a}}) \otimes L(\eta_{\boldsymbol{\nu}, \boldsymbol{b}})$ as a quasi $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell', 0)$ -module decomposes into irreducible submodules as

(7.3)
$$\operatorname{Res}_{L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)}^{L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)} L(\eta_{\boldsymbol{\mu},\boldsymbol{a}}) \otimes L(\eta_{\boldsymbol{\nu},\boldsymbol{b}}) = \bigoplus_{\boldsymbol{\xi}\in\mathbb{Z}_{++}^{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}}} D_{\boldsymbol{\mu},\boldsymbol{\nu}}^{\boldsymbol{\xi}} L(\eta_{\boldsymbol{\xi},(\boldsymbol{a},\boldsymbol{b})}).$$

Proof. Identify the Fock space $\mathcal{F}_N^a \otimes \mathcal{F}_N^b$ (= $\mathcal{F}_N^{\ell} \otimes \mathcal{F}_N^{\ell'}$) with $\mathcal{F}_N^{(a,b)}$ (= $\mathcal{F}_N^{\ell+\ell'}$) canonically. By Theorem 6.5, we have the following two dual pairs

$$(7.4) \qquad (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0), \operatorname{GL}_{\mathbf{I}_{\boldsymbol{a}}} \times \operatorname{GL}_{\mathbf{I}_{\boldsymbol{b}}}) \quad \text{and} \quad (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0), \operatorname{GL}_{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}})$$

on $\mathcal{F}_N^a \otimes \mathcal{F}_N^b \ (= \mathcal{F}_N^{(a,b)})$. Recall the following canonical embeddings

$$L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0) \hookrightarrow L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0) \quad \text{and} \quad \mathrm{GL}_{\mathbf{I}_{a}} \times \mathrm{GL}_{\mathbf{I}_{b}} \hookrightarrow \mathrm{GL}_{\mathbf{I}_{(a,b)}}.$$

Notice that the $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)$ -action on $\mathcal{F}_N^{(a,b)}$ coincides with that on $\mathcal{F}_N^a\otimes\mathcal{F}_N^a$ via the embedding $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0)\hookrightarrow L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)\otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell',0)$, while the $\mathrm{GL}_{\mathbf{I}_a}\times\mathrm{GL}_{\mathbf{I}_b}$ -action on $\mathcal{F}_N^{(a,b)}$ via the embedding $\mathrm{GL}_{\mathbf{I}_a}\times\mathrm{GL}_{\mathbf{I}_b}\hookrightarrow\mathrm{GL}_{\mathbf{I}_{(a,b)}}$ coincides with that on $\mathcal{F}_N^a\otimes\mathcal{F}_N^a$. That is, the two dual pairs in (7.4) form a seesaw pair in the sense of [Ku]. The rest essentially follows from the reciprocity law attached to this seesaw pair as in the classical case (see [H2]).

First, by (6.26) we have

$$\mathcal{F}_N^{\boldsymbol{a}} = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}}}} L(\eta_{\boldsymbol{\mu},\boldsymbol{a}}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{\boldsymbol{a}}}}(\boldsymbol{\mu}) \quad \text{and} \quad \mathcal{F}_N^{\boldsymbol{b}} = \bigoplus_{\boldsymbol{\nu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{b}}}} L(\eta_{\boldsymbol{\nu},\boldsymbol{n}}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{\boldsymbol{b}}}}(\boldsymbol{\nu}).$$

Then

$$(7.5) \quad \mathcal{F}_{N}^{\boldsymbol{a}} \otimes \mathcal{F}_{N}^{\boldsymbol{b}} = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}}}, \ \boldsymbol{\nu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{b}}}} \left(L(\eta_{\boldsymbol{\mu},\boldsymbol{a}}) \otimes L(\eta_{\boldsymbol{\nu},\boldsymbol{a}}) \right) \otimes \left(L_{\mathrm{GL}_{\mathbf{I}_{\boldsymbol{a}}}}(\boldsymbol{\mu}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{\boldsymbol{b}}}}(\boldsymbol{\nu}) \right)$$

as an $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0),\operatorname{GL}_{\mathbf{I}_a}\times\operatorname{GL}_{\mathbf{I}_b})$ -module. On the other hand, by (6.26) we have

$$\mathcal{F}_{N}^{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}} = \bigoplus_{\boldsymbol{\xi} \in \mathbb{Z}_{++}^{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}}} L(\eta_{\boldsymbol{\xi},(\boldsymbol{a},\boldsymbol{b})}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}}}(\boldsymbol{\xi}).$$

Combining this with (7.2), we get

$$(7.6) \qquad \mathcal{F}_{N}^{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}} = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}}}, \ \boldsymbol{\nu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{b}}} \in \mathbb{Z}_{++}^{\mathbf{I}_{(\boldsymbol{a},\boldsymbol{b})}} } D_{\boldsymbol{\mu},\boldsymbol{\nu}}^{\boldsymbol{\xi}} L(\eta_{\boldsymbol{\xi},(\boldsymbol{a},\boldsymbol{b})}) \otimes (L_{\mathrm{GL}_{\mathbf{I}_{\boldsymbol{a}}}}(\boldsymbol{\mu}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{\boldsymbol{b}}}}(\boldsymbol{\nu}))$$

as an $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+\ell',0), \operatorname{GL}_{\mathbf{I}_a} \times \operatorname{GL}_{\mathbf{I}_b})$ -module. With the identification $\mathcal{F}_N^{(a,b)} = \mathcal{F}_N^a \otimes \mathcal{F}_N^b$, combining (7.5) with (7.6) we obtain (7.3) as desired.

It is known (cf. [GW]) that for any positive integer n, the branching from GL_{n+1} to $GL_n \times GL_1$ on every finite-dimensional irreducible module is multiplicity free. Then as an immediate consequence of Theorem 7.1 (with $\ell' = 1$) we have:

Corollary 7.2. The branching from $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(1,0)$ to $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+1,0)$ on every irreducible \mathbb{N} -graded (\mathbb{Z},χ_q) -equivariant quasi module is multiplicity free.

It is important to note that the branching from $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \otimes L_{\widehat{\mathfrak{sl}_{\infty}}}(1,0)$ to $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell+1,0)$ on irreducible N-graded modules are *not* multiplicity free.

Remark 7.3. In view of Theorem 3.15, Theorem 7.1 also gives an explicit tensor product decomposition for any two irreducible integrable highest weight $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules. Let \mathcal{O}_{int}^f be the category of $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -modules which are sums of finitely many irreducible integrable highest weight modules. By Theorem 7.1 we conclude that \mathcal{O}_{int}^f is a semisimple abelian tensor category. This can be viewed as a q-analogue of a result of Wang [W1, Theorem 5.2].

8.
$$L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \to L_{\widehat{\mathfrak{sl}_{\infty}}\mathbf{N}}(\ell,0)$$
 branching law

Let ℓ, d be positive integers and let $\mathbf{N} = (N_1, \dots, N_d) \in \mathbb{Z}_+^d$ with $N_1 \geq 2, \dots, N_d \geq 2$. Set

(8.1)
$$N_{(0)} = 0, N_{(r)} = N_1 + \dots + N_r \text{ for } 1 \le r \le d.$$

Especially, set

$$(8.2) N = N_{(d)} = N_1 + \dots + N_d.$$

For 1 < r < d, set

(8.3)
$$\mathfrak{sl}_{\infty,\mathbf{N}}^{(r)} = \langle E_{mN+i,nN+j} \mid m, n \in \mathbb{Z}, \ N_{(r-1)} + 1 \le i \ne j \le N_{(r)} \rangle,$$

a Lie subalgebra of \mathfrak{sl}_{∞} . Then set

(8.4)
$$\mathfrak{sl}_{\infty,\mathbf{N}} = \sum_{r=1}^{d} \mathfrak{sl}_{\infty,\mathbf{N}}^{(r)},$$

a subalgebra of \mathfrak{sl}_{∞} , which actually is a direct sum of the Lie algebras $\mathfrak{sl}_{\infty,\mathbf{N}}^{(r)}$. Notice that each subalgebra $\mathfrak{sl}_{\infty,\mathbf{N}}^{(r)}$ is isomorphic to \mathfrak{sl}_{∞} with an isomorphism π_r given by

(8.5)
$$\pi_r(E_{mN+i,nN+j}) = E_{mN_r + \bar{i}, nN_r + \bar{j}}$$

for $m, n \in \mathbb{Z}$, $N_{(r-1)} + 1 \leq i \neq j \leq N_{(r)}$, where $\bar{i} = i - N_{(r-1)}$, $\bar{j} = j - N_{(r-1)}$ (so that $1 \leq \bar{i} \neq \bar{j} \leq N_r$). Then $\mathfrak{sl}_{\infty,\mathbf{N}}$ is naturally isomorphic to $\mathfrak{sl}_{\infty}^{\oplus d}$ with an isomorphism

$$\pi = \pi_1 \times \pi_2 \times \cdots \times \pi_d.$$

Equip $\mathfrak{sl}_{\infty,\mathbf{N}}$ with the bilinear form inherited from \mathfrak{sl}_{∞} , and on the other hand, equip $\mathfrak{sl}_{\infty}^{\oplus d}$ with the canonical bilinear form associated to the bilinear form on \mathfrak{sl}_{∞} . It is straightforward to see that the isomorphism preserves the invariant bilinear forms. Consequently, the isomorphism π extends uniquely to an affine Lie algebra isomorphism $\hat{\pi}$ from $\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}$ to $\widehat{\mathfrak{sl}_{\infty}^{\oplus d}}$, preserving the canonical central elements. Furthermore, $\hat{\pi}$ gives rise to a vertex algebra isomorphism

(8.6)
$$\pi_{va}: L_{\widehat{\mathfrak{sl}_{\infty}}, \mathbf{N}}(\ell, 0) \longrightarrow L_{\widehat{\mathfrak{sl}_{\oplus}}^{\oplus d}}(\ell, 0) \cong L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell, 0)^{\otimes d}.$$

Recall from Definition 2.4 the automorphism σ of \mathfrak{gl}_{∞} , where

$$\sigma(E_{m,n}) = E_{m+1,n+1}$$
 for $m, n \in \mathbb{Z}$.

It can be readily seen that σ^N preserves the subalgebras $\mathfrak{sl}_{\infty,\mathbf{N}}^{(r)}$ for $1 \leq r \leq d$, hence the subalgebra $\mathfrak{sl}_{\infty,\mathbf{N}}$. We then view σ^N as an automorphism of the \mathbb{Z} -graded vertex algebra $L_{\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}}(\ell,0)$. Let R_N be the representation of \mathbb{Z} on $L_{\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}}(\ell,0)$ given by

(8.7)
$$R_N(n) = \sigma^{nN} \chi_q(n)^{-L(0)} = \sigma^{nN} q^{-nL(0)} \quad \text{for } n \in \mathbb{Z}.$$

Then $L_{\widehat{\mathfrak{sl}_{\infty},\mathbf{N}}}(\ell,0)$ is a \mathbb{Z} -graded vertex \mathbb{Z} -algebra. Note that $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ contains $L_{\widehat{\mathfrak{sl}_{\infty},\mathbf{N}}}(\ell,0)$ as a \mathbb{Z} -graded vertex subalgebra.

Notice that under the Lie algebra isomorphism π_r from $\mathfrak{sl}_{\infty,\mathbf{N}}^{(r)}$ to \mathfrak{sl}_{∞} (see (8.5)), the automorphism σ^N of $\mathfrak{sl}_{\infty,\mathbf{N}}^{(r)}$ corresponds to the automorphism σ^{N_r} of \mathfrak{sl}_{∞} .

Definition 8.1. For $1 \leq r \leq d$, with $n \in \mathbb{Z}$ acting as σ^{nN_r} on the \mathbb{Z} -graded vertex algebra $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$, we obtain a vertex \mathbb{Z} -algebra, which we denote by $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_r})$.

With this, the vertex algebra isomorphism π_{va} (see (8.6) is actually a vertex \mathbb{Z} -algebra isomorphism from $(L_{\widehat{\mathfrak{sl}_{o},\mathbf{N}}}(\ell,0),\rho_N)$ to the tensor product vertex \mathbb{Z} -algebra

$$(8.8) (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_1}) \otimes \cdots \otimes (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_d})$$

(cf. Example 3.6 and Lemma 3.11).

Our main goal of this section is to determine the $L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0) \to L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)$ branching law. Recall that equivariant quasi $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_r})$ -modules correspond to integrable and restricted $\widehat{\mathfrak{sl}_{N_r}}(\mathbb{C}_q)$ -modules of level ℓ for $1 \le r \le d$. Recall also from Definition 6.1 that the linear functional $\eta_{\mu,a}$ on $\widehat{\mathcal{H}}$ also depends on the fixed positive integer N. In the following, we shall need such linear functionals for various positive integers N. For this reason, we here shall denote the functional $\eta_{\mu,a}$ by $\eta_{\mu,a}^{(N)}$, to show its dependence on N.

Remark 8.2. Note that with the vertex \mathbb{Z} -algebra isomorphism π_{va} , it follows from Proposition 4.12 and Theorem 6.7 that each irreducible \mathbb{N} -graded (\mathbb{Z}, χ_q) -equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}, \mathbf{N}}}(\ell, 0)$ -module is of the form

(8.9)
$$L(\eta_{\boldsymbol{\mu}_1,\boldsymbol{a}_1}^{(N_1)}) \otimes \cdots \otimes L(\eta_{\boldsymbol{\mu}_d,\boldsymbol{a}_d}^{(N_d)}),$$

where $\boldsymbol{a}_r \in (\mathbb{C}^{\times})_q^{\ell}$, $\boldsymbol{\mu}_r \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}_r}}$ for $1 \leq r \leq d$.

Let **I** be a partition of $\{1, \ldots, \ell\}$. Set $\operatorname{GL}_{\mathbf{I}}^d = \operatorname{GL}_{\mathbf{I}} \times \cdots \times \operatorname{GL}_{\mathbf{I}}$ (*d*-times) and view $\operatorname{GL}_{\mathbf{I}}$ as the diagonal subgroup. Write the $\operatorname{GL}_{\mathbf{I}}^d \to \operatorname{GL}_{\mathbf{I}}$ branching (tensor product decomposition) as

(8.10)
$$\operatorname{Res}_{\operatorname{GL}_{\mathbf{I}}}^{\operatorname{GL}_{\mathbf{I}}^{d}} L(\boldsymbol{\mu}_{1}) \otimes \cdots \otimes L(\boldsymbol{\mu}_{d}) = \bigoplus_{\boldsymbol{\xi} \in \mathbb{Z}_{++}^{\mathbf{I}}} C_{\boldsymbol{\mu}_{1}, \dots, \boldsymbol{\mu}_{d}}^{\boldsymbol{\xi}} L(\boldsymbol{\xi}),$$

where $\mu_1, \ldots, \mu_d \in \mathbb{Z}_{++}^{\mathbf{I}}$ and $C_{\mu_1, \ldots, \mu_d}^{\boldsymbol{\xi}}$ denotes the indicated multiplicity.

Remark 8.3. Let $\mathbf{I} = \{S_1, \dots, S_s\}$ be a partition of $\{1, \dots, \ell\}$. Note that in case $s = \ell$, we have $\mathrm{GL}_{\mathbf{I}} = \mathrm{GL}_1^{\ell}$, so the tensor product decomposition above is always multiplicity free. On the other hand, if $\ell = 2$ and s = 1, we have $\mathrm{GL}_{\mathbf{I}} = \mathrm{GL}_2$. It is known that the tensor product decomposition above with d = 2 is also always multiplicity free.

As the main result of this section, we have:

Theorem 8.4. Let ℓ, d be positive integers and let $\mathbf{N} = (N_1, \dots, N_d) \in \mathbb{Z}_+^d$ with $N_i \geq 2$. Set $N = N_1 + \dots + N_d$. Then for any $\mathbf{a} \in (\mathbb{C}^{\times})_q^{\ell}$, $\boldsymbol{\xi} \in \mathbb{Z}_{++}^{\mathbf{I}_{\mathbf{a}}}$, $L(\eta_{\boldsymbol{\xi}, \mathbf{a}}^{(N)})$ decomposes into irreducible equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty}, \mathbf{N}}}(\ell, 0)$ -submodules with multiplicities as

$$(8.11) \operatorname{Res}_{L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)}^{L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0)} L(\eta_{\boldsymbol{\xi},\boldsymbol{a}}^{(N)}) = \bigoplus_{\boldsymbol{\mu}_{1},\dots,\boldsymbol{\mu}_{d} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}}}} C_{\boldsymbol{\mu}_{1},\dots,\boldsymbol{\mu}_{d}}^{\boldsymbol{\xi}} L(\eta_{\boldsymbol{\mu}_{1},\boldsymbol{a}}^{(N_{1})}) \otimes \cdots \otimes L^{d}(\eta_{\boldsymbol{\mu}_{d},\boldsymbol{a}}^{(N_{d})}),$$

where the multiplicities $C_{\mu^1,\dots,\mu_d}^{\xi}$ are the same as in (8.10).

Proof. Recall that \mathcal{F}_N^a is an equivariant quasi module for $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N)$ with $\mathcal{F}_N^a = \mathcal{F}_N^\ell$ as a vector space. Then we have an equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty,N}}}(\ell,0)$ -module \mathcal{F}_N^a via the natural embedding of $L_{\widehat{\mathfrak{sl}_{\infty,N}}}(\ell,0)$ into $(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N)$. On the other hand, we have an equivariant quasi $L_{\widehat{\mathfrak{sl}_{\infty,N}}}(\ell,0)$ -module $\mathcal{F}_{N_1}^a\otimes\cdots\otimes\mathcal{F}_{N_d}^a$ via the vertex algebra isomorphism

$$\pi_{va}: (L_{\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}}(\ell,0),\rho_N) \cong (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_1}) \otimes \cdots \otimes (L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_d}).$$

It is straightforward to show that there exists a (Clifford) algebra isomorphism

(8.12)
$$\theta_{\mathbf{N}}^{\ell}: \quad \mathcal{C}_{N_1}^{\ell} \otimes \cdots \otimes \mathcal{C}_{N_d}^{\ell} \longrightarrow \mathcal{C}_{N}^{\ell}$$

such that

$$\theta_{\mathbf{N}}^{\ell}(\psi_{i}^{p}(n)) = \psi_{i+N_{(r-1)}}^{p}(n), \quad \theta_{\mathbf{N}}^{\ell}(\bar{\psi}_{i}^{p}(n)) = \bar{\psi}_{i+N_{(r-1)}}^{p}(n)$$

for $1 \le r \le d$, $1 \le i \le N_r$, $1 \le p \le \ell$, $n \in \mathbb{Z}$. Furthermore, this gives rise to a Fock space identification

(8.13)
$$\Theta_{\mathbf{N}}^{\ell}: \quad \mathcal{F}_{N_1}^{\ell} \otimes \cdots \otimes \mathcal{F}_{N_d}^{\ell} \longrightarrow \mathcal{F}_{N}^{\ell}.$$

It is easy to see that $\Theta_{\mathbf{N}}^{\ell}$ is an isomorphism of quasi $L_{\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}}(\ell,0)$ -modules from $\mathcal{F}_{N_1}^{\boldsymbol{a}}\otimes\cdots\otimes\mathcal{F}_{N_d}^{\boldsymbol{a}}$ to \mathcal{F}_N^{ℓ} . It can be readily seen that $\Theta_{\mathbf{N}}^{\ell}$ is also a $\mathrm{GL}_{\mathbf{I}_{\boldsymbol{a}}}$ -module isomorphism. Now, by Theorem 6.5, we get a seesaw pair

$$((L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_1})\otimes\cdots\otimes(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_d}),\operatorname{GL}_{\mathbf{I}_{\boldsymbol{a}}}^d)$$
 and $((L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N),\operatorname{GL}_{\mathbf{I}_{\boldsymbol{a}}})$ on \mathcal{F}_N^ℓ with the embeddings

$$(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_1})\otimes\cdots\otimes(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_{N_d})\cong(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N)\hookrightarrow(L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N)$$

and $\operatorname{GL}_{\mathbf{I}_a} \hookrightarrow \operatorname{GL}_{\mathbf{I}_a}^d$. Using the first dual pair and (8.10), we obtain the following $(L_{\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}}(\ell,0),\operatorname{GL}_{\mathbf{I}_a})$ -module decomposition

$$\mathcal{F}_{N_{1}}^{a} \otimes \cdots \otimes \mathcal{F}_{N_{d}}^{a}
= \left(\bigoplus_{\boldsymbol{\mu}_{1} \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}} L(\eta_{\boldsymbol{\mu}_{1}, \boldsymbol{a}}^{(N_{1})}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\mu}_{1}) \right) \otimes \cdots \otimes \left(\bigoplus_{\boldsymbol{\mu}_{d} \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}} L(\eta_{\boldsymbol{\mu}_{d}, \boldsymbol{a}}^{(N_{d})}) \otimes L_{\mathrm{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\mu}_{d}) \right)
\cong \bigoplus_{\boldsymbol{\mu}^{1}, \dots, \boldsymbol{\mu}^{d} \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}} \left(L(\eta_{\boldsymbol{\mu}_{1}, \boldsymbol{a}}^{(N_{1})}) \otimes \cdots \otimes L(\eta_{\boldsymbol{\mu}_{d}, \boldsymbol{a}}^{(N_{d})}) \right) \otimes \left(L_{\mathrm{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\mu}_{1}) \otimes \cdots \otimes L_{\mathrm{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\mu}_{d}) \right)
= \bigoplus_{\boldsymbol{\xi}, \boldsymbol{\mu}_{1}, \dots, \boldsymbol{\mu}_{d} \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}} C_{\boldsymbol{\mu}_{1}, \dots, \boldsymbol{\mu}_{d}}^{\boldsymbol{\xi}} \left(L(\eta_{\boldsymbol{\mu}_{1}, \boldsymbol{a}}^{(N_{1})}) \otimes \cdots \otimes L^{d}(\eta_{\boldsymbol{\mu}_{d}, \boldsymbol{a}}^{(N_{d})}) \right) \otimes L_{\mathrm{GL}_{\mathbf{I}_{a}}}(\boldsymbol{\xi}).$$

Combining this with the decomposition for the dual pair $((L_{\widehat{\mathfrak{sl}_{\infty}}}(\ell,0),\rho_N),\mathrm{GL}_{\mathbf{I}_a})$ on \mathcal{F}_N^{ℓ} (see (6.26)) we obtain (8.11).

As immediate consequences of Theorem 8.4 and Remark 8.3 we have:

Corollary 8.5. The branching from $(L_{\widehat{\mathfrak{sl}_{\infty}}}(1,0), \rho_N)$ to $L_{\widehat{\mathfrak{sl}_{\infty,N}}}(1,0)$ in Theorem 8.4 on equivaraint quasi modules is multiplicity free.

Corollary 8.6. Assume $\ell = 2$ and $\mathbf{N} = (N_1, N_2)$ (with d = 2). Then the branching from $L_{\widehat{\mathfrak{sl}_{\infty},\mathbf{N}}}(2,0)$ to $L_{\widehat{\mathfrak{sl}_{\infty,\mathbf{N}}}}(2,0)$ in Theorem 8.4 is multiplicity free.

Remark 8.7. Here, we formulate a Lie algebra analogue of Theorem 8.4. For $1 \leq r \leq d$, let $\widehat{\mathfrak{sl}_{\mathbf{N}}}^{(r)}(\mathbb{C}_q)$ denote the subalgebra of $\widehat{\mathfrak{sl}_{N}}(\mathbb{C}_q)$, generated by the elements

$$E_{i,j}t_0^{m_0}t_1^{m_1}$$
 for $m_0, m_1 \in \mathbb{Z}$, $N_{(r-1)} + 1 \le i \ne j \le N_{(r)}$.

Set

$$\widehat{\mathfrak{sl}_{\mathbf{N}}}(\mathbb{C}_q) = \sum_{r=1}^d \widehat{\mathfrak{sl}_{\mathbf{N}}}^{(r)}(\mathbb{C}_q),$$

which is a subalgebra of $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$. It is easy to see that $\widehat{\mathfrak{sl}}_{\mathbf{N}}^{(r)}(\mathbb{C}_q)$ is isomorphic to $\widehat{\mathfrak{sl}}_{N_r}(\mathbb{C}_q)$ (as $N_r = N_{(r)} - N_{(r-1)}$) and for $1 \leq r \neq s \leq d$,

$$[\widehat{\mathfrak{sl}_{\mathbf{N}}}^{(r)}(\mathbb{C}_q), \widehat{\mathfrak{sl}_{\mathbf{N}}}^{(s)}(\mathbb{C}_q)] = 0.$$

Then it follows that every irreducible \mathbb{N} -graded integrable restricted $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ module has the form of (8.9). Under the setting of Theorem 8.4, we have the $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q) \to \widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ branching law:

$$\operatorname{Res}_{\widehat{\mathfrak{sl}_{\mathbf{N}}}(\mathbb{C}_q)}^{\widehat{\mathfrak{sl}_{\mathbf{N}}}(\mathbb{C}_q)} L(\eta_{\boldsymbol{\xi},\boldsymbol{a}}^{(N)}) = \bigoplus_{\boldsymbol{\mu}_1,\dots,\boldsymbol{\mu}_d \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{a}}}} C_{\boldsymbol{\mu}_1,\dots,\boldsymbol{\mu}_d}^{\boldsymbol{\xi}} L(\eta_{\boldsymbol{\mu}_1,\boldsymbol{a}}^{(N_1)}) \otimes \cdots \otimes L(\eta_{\boldsymbol{\mu}_d,\boldsymbol{a}}^{(N_d)}).$$

9.
$$\widehat{\mathfrak{sl}_N}(\mathbb{C}_q) \to \widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$$
 branching law

Let M_0 and M_1 be positive integers, which are fixed throughout this section. Denote by $\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}]$ the subalgebra of $\mathbb{C}_q = \mathbb{C}_q[t_0^{\pm 1}, t_1^{\pm 1}]$, generated by $t_0^{\pm M_0}$ and $t_1^{\pm M_1}$. Set

$$(9.1) \qquad \widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}]) = \mathfrak{sl}_N(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}]) \oplus \mathbb{C}\mathbf{k}_0 \oplus \mathbb{C}\mathbf{k}_1 \subset \widehat{\mathfrak{sl}_N}(\mathbb{C}_q).$$

Note that $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$ is simply the subalgebra of $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ generated by

$$E_{i,j}t_0^{m_0M_0}t_1^{m_1M_1}$$
 for $1 \le i \ne j \le N$, $m_0, m_1 \in \mathbb{Z}$.

The main goal of this section is to determine the $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q) \to \widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$ branching law.

First of all, it is straightforward to see that as an algebra $\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}]$ is isomorphic to $\mathbb{C}_{q^{M_0M_1}}[t_0^{\pm 1}, t_1^{\pm 1}]$ with $t_0^{n_0M_0}t_1^{n_1M_1}$ corresponding to $t_0^{n_0}t_1^{n_1}$ for $n_0, n_1 \in \mathbb{Z}$. Furthermore, we have the following straightforward result:

Lemma 9.1. There is a Lie algebra isomorphism

$$\pi_{M_0,M_1}:\ \widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0},t_1^{\pm M_1}]) \to \widehat{\mathfrak{sl}_N}(\mathbb{C}_{q^{M_0M_1}}),$$

which is uniquely determined by the assignment

(9.2)
$$E_{i,j}t_0^{m_0M_0}t_1^{m_1M_1} \mapsto E_{i,j}t_0^{m_0}t_1^{m_1}, \quad M_0\mathbf{k}_0 \mapsto \mathbf{k}_0, \quad M_1\mathbf{k}_1 \mapsto \mathbf{k}_1$$

for $1 \le i \ne j \le N, \ m_0, m_1 \in \mathbb{Z}$.

Let $\mathbf{I} = \{S_1, \dots, S_s\}$ be a partition of $\{1, 2, \dots, \ell\}$. Set

(9.3)
$$\mathbf{I}^{M_0} = \{k\ell + S_i \mid 0 \le k \le M_0 - 1, \ 1 \le i \le s\},\$$

where by definition $k\ell + S_i = \{k\ell + p \mid p \in S_i\}$. Then \mathbf{I}^{M_0} is a partition of $\{1, 2, \dots, M_0\ell\}$. For $\boldsymbol{a} = (a_1, \dots, a_\ell) \in (\mathbb{C}^{\times})^{\ell}$, we define $(\boldsymbol{a}q)_{M_0}^{M_1} \in (\mathbb{C}^{\times})^{M_0\ell}$ by

(9.4)
$$((\boldsymbol{a}q)_{M_0}^{M_1})_{k\ell+r} = (a_r q^{-k})^{M_1}$$

for $0 \le k \le M_0 - 1$, $1 \le r \le \ell$. Then we have:

Lemma 9.2. Let M_0, M_1 and ℓ be positive integers, let $\mathbf{a} \in (\mathbb{C}^{\times})_q^{\ell}$ and let $\mathbf{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_a}$. Then there exist $\mathbf{b} \in (\mathbb{C}^{\times})_q^{\ell}$ and $\mathbf{\nu} \in \mathbb{Z}_{++}^{\mathbf{I}_b}$ such that

$$\eta_{\boldsymbol{\mu}, \boldsymbol{a}} = \eta_{\boldsymbol{\nu}, \boldsymbol{b}}, \quad (\boldsymbol{b}q)_{M_0}^{M_1} \in (\mathbb{C}^{\times})_{q^{M_0M_1}}^{M_0\ell} \quad \text{and} \quad \mathbf{I}_{(\boldsymbol{b}q)_{M_0}^{M_1}} = \mathbf{I}_{\boldsymbol{b}}^{M_0}.$$

Proof. Define an equivalence relation \sim on $I:=\{1,2,\ldots,\ell\}$ by claiming $i\sim j$ if and only if $a_i^{M_1}=q^{nM_1}a_j^{M_1}$ for some $n\in\mathbb{Z}$. Let $\{I_1,\ldots,I_s\}$ be the partition of I, associated to this equivalence relation. For $1\leq i\leq \ell$, set $A_i=\{a_i\mid i\in I_i\}$ and choose a representative c_i from A_i . Then for each $j\in I$ there exist unique $1\leq i\leq s$ and $n_j\in\mathbb{Z}$ such that $q^{n_jM_1}a_j^{M_1}=c_i^{M_1}$. Set $\boldsymbol{n}=(n_1,\ldots,n_\ell)\in\mathbb{Z}^\ell$ and

$$\boldsymbol{b} = q^{\boldsymbol{n}} \boldsymbol{a} = (q^{n_1} a_1, \dots, q^{n_\ell} a_\ell) \in (\mathbb{C}^{\times})^{\ell},$$

$$\boldsymbol{\nu} = \boldsymbol{\mu} + \boldsymbol{n} N = (\mu_1 + n_1 N, \dots, \mu_{\ell} + n_{\ell} N) \in \mathbb{Z}^{\ell}.$$

It is straightforward to see that $\boldsymbol{b} \in (\mathbb{C}^{\times})_{q}^{\ell}$ (with $\mathbf{I}_{\boldsymbol{b}} = \mathbf{I}_{\boldsymbol{a}}$) and $\boldsymbol{\nu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\boldsymbol{b}}}$. Furthermore, for $1 \leq i, j \leq \ell$, we have either $b_i^{M_1} = b_j^{M_1}$ or $b_i^{M_1} \neq q^{nM_1}b_j^{M_1}$ for any $n \in \mathbb{Z}$. This implies that $(\boldsymbol{b}q)_{M_0}^{M_1}$ lies in $(\mathbb{C}^{\times})_{q^{M_1}}^{M_0\ell}$ (and hence in $(\mathbb{C}^{\times})_{q^{M_0M_1}}^{M_0\ell}$) and $\mathbf{I}_{(\boldsymbol{b}q)_{M_0}^{M_1}} = \mathbf{I}_{\boldsymbol{b}}^{M_0}$. Finally, it follows from Remark 6.2 that $\eta_{\boldsymbol{\mu},\boldsymbol{a}} = \eta_{\boldsymbol{\nu},\boldsymbol{b}}$, as desired. \square

Let I be a partition of $\{1, \ldots, \ell\}$. From definition, we have

$$\operatorname{GL}_{\mathbf{I}^{M_0}} \cong \operatorname{GL}_{\mathbf{I}} \times \cdots \times \operatorname{GL}_{\mathbf{I}} = (\operatorname{GL}_{\mathbf{I}})^{M_0}.$$

View $GL_{\mathbf{I}}$ as a subgroup of $GL_{\mathbf{I}^{M_0}}$ through the diagonal embedding. Write the $GL_{\mathbf{I}^{M_0}} \to GL_{\mathbf{I}}$ branching as

(9.5)
$$\operatorname{Res}_{\operatorname{GL}_{\mathbf{I}}}^{\operatorname{GL}_{\mathbf{I}^{M_0}}} L(\boldsymbol{\xi}) = \bigoplus_{\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}}} E_{\boldsymbol{\mu}}^{\boldsymbol{\xi}} L(\boldsymbol{\mu})$$

for $\boldsymbol{\xi} \in \mathbb{Z}_{++}^{\mathbf{I}^{M_0}}$, where $E_{\boldsymbol{\mu}}^{\boldsymbol{\xi}}$ denotes the multiplicity of the $\mathrm{GL}_{\mathbf{I}}$ -module $L(\boldsymbol{\mu})$ in the $\mathrm{GL}_{\mathbf{I}^{M_0}}$ -module $L(\boldsymbol{\xi})$.

The following is the main result of this section:

Theorem 9.3. Let ℓ , M_0 , M_1 be positive integers, let $\mathbf{a} \in (\mathbb{C}^{\times})_q^{\ell}$ such that $(\mathbf{a}q)_{M_0}^{M_1} \in (\mathbb{C}^{\times})_{q^{M_0M_1}}^{M_0\ell}$ and $\mathbf{I}_{(\mathbf{a}q)_{M_0}^{M_1}} = \mathbf{I}_{\mathbf{a}}^{M_0}$, and let $\boldsymbol{\mu} \in \mathbb{Z}_{++}^{\mathbf{I}_{\mathbf{a}}}$. Then the $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q)$ -module $L(\eta_{\boldsymbol{\mu},\mathbf{a}})$ viewed as an $\widehat{\mathfrak{sl}}_N(\mathbb{C}_q[t_0^{\pm M_0},t_1^{\pm M_1}])$ -module decomposes into irreducible submodules with multiplicities as

(9.6)
$$L(\eta_{\mu,a}) = \bigoplus_{\xi \in \mathbb{Z}_{++}^{\mathbf{I}_{a}}^{M_{0}}} E_{\mu}^{\xi} L(\eta_{\xi,(aq)_{M_{0}}^{M_{1}}}).$$

Proof. Recall that we have Clifford algebras $\mathcal{C}_N^{M_0\ell}$ and \mathcal{C}_N^{ℓ} . It is straightforward to show that the assignment

(9.7)
$$\psi_i^{k\ell+p}(n) \mapsto \psi_i^p(M_0n-k), \qquad \bar{\psi}_i^{k\ell+p}(n) \mapsto \bar{\psi}_i^p(M_0n+k)$$

for $1 \leq i \leq N$, $0 \leq k \leq M_0 - 1$, $1 \leq p \leq \ell$, $n \in \mathbb{Z}$ extends uniquely to an algebra isomorphism from $\mathcal{C}_N^{M_0\ell}$ to \mathcal{C}_N^ℓ , denoted by ϕ . (Note that every integer $1 \leq p' \leq M_0\ell$ can be written uniquely as $p' = k\ell + p$ with $1 \leq p \leq \ell$, $0 \leq k \leq M_0 - 1$.) We have

(9.8)
$$\phi\left(:\psi_i^{k\ell+p}(n)\bar{\psi}_{i'}^{k'\ell+p'}(n'):\right) =: \psi_i^p(M_0n-k)\bar{\psi}_{i'}^{p'}(M_0n'+k'):$$

for $1 \leq i, i' \leq N$, $1 \leq p, p' \leq \ell$, $0 \leq k, k' \leq M_0 - 1$ and $n, n' \in \mathbb{Z}$, recalling Remark 5.1 and noticing that $M_0n' + k' \geq 0$ if and only if $n' \geq 0$. Via this isomorphism ϕ , the \mathcal{C}_N^{ℓ} -module \mathcal{F}_N^{ℓ} becomes a $\mathcal{C}_N^{M_0\ell}$ -module, which is denoted by $(\mathcal{F}_N^{\ell})^{\phi}$. Notice that for $0 \leq k \leq M_0 - 1$, $n \in \mathbb{Z}$, $M_0n - k > 0$ if and only if n > 0, and $M_0n + k \geq 0$ if and only if $n \geq 0$. It then follows from the construction of $\mathcal{F}_N^{M_0\ell}$

that $(\mathcal{F}_N^\ell)^\phi \simeq \mathcal{F}_N^{M_0\ell}$ as a $\mathcal{C}_N^{M_0\ell}$ -module. That is, we have a linear isomorphism $\Phi: \mathcal{F}_N^{M_0\ell} \to \mathcal{F}_N^\ell$ such that $\Phi(|0\rangle) = |0\rangle$ and

$$\Phi(\psi_i^{k\ell+p}(n)w) = \psi_i^p(M_0n - k)\bar{\phi}(w), \quad \Phi(\bar{\psi}_i^{k\ell+p}(n)w) = \bar{\psi}_i^p(M_0n + k)\bar{\phi}(w)$$

for $1 \le i \le N$, $1 \le p \le \ell$, $0 \le k \le M_0 - 1$, $n \in \mathbb{Z}$, $w \in \mathcal{F}_N^{\ell}$.

Now, with $\mathbf{a} = (a_1, \dots, a_\ell) \in (\mathbb{C}^\times)^\ell$, we have an $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q)$ -module $\mathcal{F}_N^{\mathbf{a}}$, which is naturally an $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$ -module. For $1 \leq i \neq j \leq N$, $m_0, m_1 \in \mathbb{Z}$, recall from (5.17) that $E_{i,j}t_0^{M_0m_0}t_1^{M_1m_1}$ (in $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0}, t_1^{\pm M_1}])$) acts on $\mathcal{F}_N^{\mathbf{a}}$ as

$$\sum_{n\in\mathbb{Z}}\sum_{r=1}^{\ell} (a_r q^{-n})^{M_1 m_1} : \psi_i^r (m_0 M_0 - n) \bar{\psi}_j^r (n) :$$

$$= \sum_{n \in \mathbb{Z}} \sum_{r=1}^{\ell} \sum_{k=0}^{M_0-1} \left((a_r q^{-k})^{M_1} q^{-nM_0 M_1} \right)^{m_1} : \psi_i^r ((m_0 - n) M_0 - k) \bar{\psi}_j^r (n M_0 + k) :.$$

On the other hand, via the Lie algebra isomorphism π_{M_0,M_1} , the $\widehat{\mathfrak{sl}_N}(\mathbb{C}_{q^{M_0M_1}})$ module $\mathcal{F}_N^{(aq)_{M_0}^{M_1}}(=\mathcal{F}_N^{M_0\ell})$ becomes an $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0},t_1^{\pm M_1}])$ -module, where the element $E_{i,j}t_0^{M_0m_0}t_1^{M_1m_1}$ acts as

$$\sum_{n\in\mathbb{Z}}\sum_{k=0}^{M_0-1}\sum_{r=1}^{\ell}\left((a_rq^{-k})^{M_1}q^{-nM_0M_1}\right)^{m_1}:\psi_i^{k\ell+r}(m_0-n)\bar{\psi}_j^{k\ell+r}(n):.$$

It then follows from (9.8) that the Fock space identification

$$\Psi:~\mathcal{F}_N^{(oldsymbol{a}q)_{M_0}^{M_1}}=\mathcal{F}_N^{M_0\ell}\simeq^\Phi\mathcal{F}_N^\ell=\mathcal{F}_N^{oldsymbol{a}}$$

is an $\widehat{\mathfrak{sl}_N}(\mathbb{C}_q[t_0^{\pm M_0},t_1^{\pm M_1}])\text{-module}$ isomorphism.

Next, we show that Ψ is also a $\mathrm{GL}_{\mathbf{I}_a}$ -module isomorphism. Recall that the action of $\mathrm{GL}_{\mathbf{I}_a}$ on \mathcal{F}_N^{ℓ} is given by the action of the Lie algebra $\mathfrak{gl}_{\mathbf{I}_a}$ from (5.13), where

$$\mathfrak{gl}_{\mathbf{I}_a} = \operatorname{span}\{E_{p,p'} \mid 1 \leq p, p' \leq \ell, \ p \sim_a p'\} \subset \mathfrak{gl}_{\ell}.$$

Similarly, the action of $\mathrm{GL}_{\mathbf{I}^{M_0}}$ on $\mathcal{F}_N^{M_0\ell}$ is given by the action of the Lie algebra $\mathfrak{gl}_{\mathbf{I}_a}{}^{M_0}$ ($\subset \mathfrak{gl}_{M_0\ell}$). Furthermore, $\mathfrak{gl}_{\mathbf{I}_a}$ acts on $\mathcal{F}_N^{M_0\ell}$ through the embedding of $\mathfrak{gl}_{\mathbf{I}_a}$ into $\mathfrak{gl}_{\mathbf{I}_a}{}^{M_0}$ with

$$E_{p,p'} \mapsto \sum_{k=0}^{M_0 - 1} E_{k\ell + p, k\ell + p'}$$

for $1 \leq p, p' \leq \ell$ with $p \sim_a p'$. We see that the element $E_{p,p'}$ of the Lie algebra $\mathfrak{gl}_{\mathbf{I}_a}$ acts on $\mathcal{F}_N^{M_0\ell}$ as

$$\sum_{k=0}^{M_0-1} \sum_{i=1}^{N} : \psi_i^{k\ell+p}(-n) \bar{\psi}_i^{k\ell+p'}(n) : .$$

Then it follows from (9.8) that Ψ is a $\mathfrak{gl}_{\mathbf{I}_a}$ -module isomorphism. Thus Ψ is also a $\mathrm{GL}_{\mathbf{I}_a}$ -module isomorphism.

Finally, by using Ψ we transport the $\mathrm{GL}_{\mathbf{I}_a^{M_0}}$ -module structure from $\mathcal{F}_N^{(aq)_{M_0}^{M_1}}$ to \mathcal{F}_N^a , obtaining a seesaw pair

$$(\widehat{\mathfrak{sl}_N}(\mathbb{C}_q),\operatorname{GL}_{\mathbf{I}_{\boldsymbol{a}}})$$
 and $(\widehat{\mathfrak{sl}_N}(\mathbb{C}_{q^{M_0M_1}}),\operatorname{GL}_{\mathbf{I}_{\underline{a}}^{M_0}})$

on the Fock space \mathcal{F}_N^a . Then just as in the proof of Theorem 8.4, (9.6) follows from this seesaw pair.

REFERENCES

- [AABGP] B. Allison, S. Azam, S. Berman, Y. Gao, and A. Pianzola, Extended Affine Lie Algebras and Their Root Systems, *Memoirs Amer. Math. Soc.* **126**,1997.
- [ABFP] B. Allison, S. Berman, J. Faulkner and A. Pianzola, Multiloop realization of extended affine Lie algebras and Lie tori, *Trans. Amer. Math. Soc.* **361** (2009), 4807-4842.
- [ABGP] B. Allison, S. Berman, Y. Gao, and A. Pianzola, A characterization of affine Kac-Moody Lie algebras, Commun. Math. Phys. 185 (1997), 671-688.
- [BGK] S. Berman, Y. Gao, and Y. Krylyuk, Quantum tori and the structure of elliptic quasisimple Lie algebras, J. Funct. Anal. 135 (1996) 339-389.
- [BGT] S. Berman, Y. Gao, and S. Tan, A unified view of some vertex operator constructions, *Israel J. Math.* **134** (2003) 29-60.
- [BBS] S. Berman, Y. Billig and J. Szmigielski, Vertex operator algebras and the representation theory of toroidal algebras, in *Recent Developments in Infinite-Dimensional Lie Algebras and Conformal Field Theory (Charlottesville, VA, 2000)*, Contemporary Math. 297, Amer. Math. Soc., Providence, 2002, 1-26.
- [BS] S. Berman and J. Szmigielski, Principal realization for extended affine Lie algebra of type \mathfrak{sl}_2 with coordinates in a simple quantum torus with two variables, Contemporary Math., 248, (1999), 39-67.
- [B1] Y. Billig, A category of modules for the full toroidal Lie algebra, *Int. Math. Res. Not.* (2006), Art. ID 68395, 46 pp.
- [B2] Y. Billig, Representations of toroidal extended affine Lie algebras, J. Algebra 308 (2007) 252-269.
- [BL] Y. Billig and M. Lau, Irreducible modules for extended affine Lie algebras, J. Algebra 327 (2011) 208-235.
- [BZ] Y. Billig and K. Zhao, Weight modules over exp-polynomial Lie algebra, *J. Pure Appl. Algebra* **191** (2004) 23-42.
- [CG] V. Chari and J. Greenstein, Graded level zero integrable representations of affine Lie algebras, *Tran. Amer. Math. Soc.* Vol. **360** (2008) 2923-2940.
- [CP] V. Chari and A. Pressley, Weyl modules for classical and quantum affine algebras, Representation Theory 5 (2001) 191-223.
- [CLT] F. Chen, Z. Li and S. Tan, Integrable representations for toroidal extended affine Lie algebras, J. Algebra 519 (2019) 228-252.
- [CT] F. Chen and S. Tan, Integrable representations for extended affine Lie algebras coordinated by quantum tori, J. Lie Theory 23 (2) (2013) 383-405.
- [DJKM] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, Operator approach to the Kadomtsev-Petviashvili equation. Transformation groups for soliton equations III, J. Phys. Soc. Japan 50 (1981) 3806-3812.

- [DLM] C. Dong, H.-S. Li and G. Mason, Regularity of rational vertex operator algebras, *Advances in Math.* **132** (1997) 148-166.
- [ER] S. Eswara Rao, A class of integrable modules for the core of EALA coordinatized by quantum tori, J. Algebra 275 (2004) 59-74.
- [FF] A. Feingold, I. Frenkel, Classical affine algebras, Advances in Math. 56 (1985) 117-172.
- [FKRW] E. Frenkel, V. Kac, A. Radul, and W. Wang, $W_{1+\infty}$ and $W(gl_N)$ with central charge N, Commun. Math. Phys. **170** (1995) 337-357.
- [F] I. B. Frenkel, Representations of affine Lie algebras, Hecke modular forms and Korteweg-de Vires type equations, Lecture Notes in Mathematics, Vol. 933, Springer-Verlag, Berlin, (1982), 71-110.
- [FHL] I. B. Frenkel, Y. Huang and J. Lepowsky, On Axiomatic Approaches to Vertex Operator Algebras and Modules, *Memoirs Amer. Math. Soc.* **104**, 1993.
- [FLM] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Math., Vol. 134, Academic Press, Boston, 1988.
- [FZ] I. B. Frenkel and Y.-C. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992) 123-168.
- [G1] Y. Gao, Representations of extended affine Lie algebras coordinatized by certain quantum tori, *Compositio Math.* **123** (2000) 1-25.
- [G2] Y. Gao, Vertex operators arising from the homogeneous realization for \hat{gl}_N , Commun. Math. Phys. **211** (2000) 745-777.
- [G3] Y. Gao, Fermionic and bosonic representations of the extended affine Lie algebra $\widehat{gl_N(\mathbb{C}_q)}$, Canada Math. Bull. 45 (2002) 623-633.
- [GZ] Y. Gao and Z. Zeng, Hermitian representations of the extended affine Lie algebra $\widehat{gl_2(\mathbb{C}_q)}$, Advances in Math. **207** (2006) 244-265.
- [G-KK] M. Golenishcheva-Kutuzova and V. Kac, Γ-conformal algebras, J. Math. Phys. 39 (1998) 2290-2305.
- [GW] R. Goodman and R. Wallach, Symmetry, Representations, and Invariants, Graduate Texts in Mathematics, 255.
- [H-KT] R. Høegh-Krohn and B. Torresani, Classification and construction of quasisimple Lie algebras, J. Funct. Anal. 89 (1990) 106-136.
- [H1] R. Howe, Remarks on classical invariant theory, *Trans. Amer. Math. Soc.* **313** (1989) 539-570.
- [H2] R. Howe, Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond, Israel Math. Conf. Proc., 8 (1992), 1-182.
- [JK] H. Jakobsen and V. Kac, A new class of unitarizable highest weight representations of infinite dimensional Lie algebras, in: Nonlinear Equations in Classical and Quantum Field Theory, Meudon/Paris, 1983/1984, Springer, Berlin, (1985), 1-20.
- [JLin] C. Jiang and Z. Lin, Tensor Decomposition, Parafermions, Level-Rank Duality and Reciprocity Law for vertex operator algebras, arXiv:1406.4191.
- [JLam] C. Jiang and C.-H. Lam, Level-Rank Duality for Vertex Operator Algebras of types B and D, Bull. Inst. Math. Acad. Sin.(N.S.) 14 (2019) 31-54.
- [K] V. G. Kac, Infinite-dimensional Lie Algebras, 3rd ed., Cambridge Univ. Press, Cambridge, 1990.
- [KMPX] V. G. Kac, P. Möseneder Frajria, P. Papi, F. Xu, Conformal embeddings and simple current extensions, *Int. Mat. Res. Not.*, **14** (2015), 5229-5288.
- [KR] V. G. Kac and A. Radul, Quasifinite highest weight modules over the Lie algebra of differential operators on the circle, *Commun. Math. Phys.* **157** (1993) 429-457.

- [Ku] S. Kulda, Seesaw dual reductive pairs, in "Automorphic forms of several variables", Proceedings of the 1983 Taniguchi Symposium (I. Satake and Y. Morita. eds.), Birkhauser, Basel, (1984) 244-268.
- [LL] J. Lepowsky and H.-S. Li, Introduction to Vertex Operator Algebras and Their Representations, Progress in Math. 227, Birkhäuser, Boston, 2004.
- [LP] J. Lepowsky and M. Primc, Structure of the Standard Modules for the Affine Lie Algebra $A_1^{(1)}$, Contemporary Math. 46, Amer. Math. Soc., Providence, 1985.
- [Li1] H.-S. Li, Local systems of vertex operators, vertex superalgebras and modules, *J. Pure Appl. Algebra* **109** (1996) 143-195.
- [Li2] H.-S. Li, Local systems of twisted vertex operators, vertex superalgebras and twisted modules, Contemporary Math. 193, Amer. Math. Soc., Providence, 1996, 203-236.
- [Li3] H.-S. Li, A new construction of vertex algebras and quasi modules for vertex algebras, *Advances in Math.* **202** (2006) 232-286.
- [Li4] H.-S. Li, On certain generalizations of twisted affine Lie algebras and quasi modules for Γ-vertex algebras, J. Pure Appl. Algebra 209 (2007) 853-871.
- [Li5] H.-S. Li, Twisted modules and quasi-modules for vertex operator algebras, Contemporary Math. 422, Amer. Math. Soc., Providence, 2007, 389-400.
- [LTW] H.-S. Li, S. Tan and Q. Wang, Trigonometric Lie algebras, affine Lie algebras, and vertex algebras, Advances in Math. 363 (2020) 106985.
- [N] E. Neher, Extended affine Lie algebras, C. R. Math. Acad. Sci. Soc. R. Can. 26 (2004) 90-96.
- [VV1] M. Varagnolo and E. Vasserot, Schur duality in the toroidal setting, Commun. Math. Phys. 182 (1996) 469-484.
- [VV2] M. Varagnolo and E. Vasserot, Double-loop algebras and the Fock space, *Invent. Math.* 133 (1998) 133-159.
- [W1] W. Wang, Dual pairs and tensor categories of modules over the Lie algebras \widehat{gl}_{∞} and $W_{1+\infty}$, J. Algebra **216** (1999) 159-177.
- [W2] W. Wang, Duality in infinite dimensional Fock representations, Commun. Contem. Math. 1 (1999) 155-199.

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005 E-mail address: chenf@xmu.edu.cn

Department of Mathematical Sciences, Rutgers University, Camden, NJ 08102, USA

 $E ext{-}mail\ address: hli@camden.rutgers.edu}$

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005 $E\text{-}mail\ address:}$ tans@xmu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005 $E\text{-}mail\ address:}$ qingwang@xmu.edu.cn