

# Torsion functions and the Cheeger problem: a fractional approach

H. Bueno, G. Ercole, S.S. Macedo and G.A. Pereira\*

*Departamento de Matemática*

*Universidade Federal de Minas Gerais, Belo Horizonte, 31.270-901, Brazil*

*e-mail: hamilton@mat.ufmg.br, grey@mat.ufmg.br, sdasilvamacedo@gmail.com*

*and gilbertoapereira@yahoo.com.br*

**Abstract:** Let  $\Omega$  be a Lipschitz bounded domain of  $\mathbb{R}^N$ ,  $N \geq 2$ . The fractional Cheeger constant  $h_s(\Omega)$ ,  $0 < s < 1$ , is defined by  $h_s(\Omega) = \inf_{E \subset \Omega} \frac{P_s(E)}{|E|}$ , where  $P_s(E) = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\chi_E(x) - \chi_E(y)|}{|x-y|^{N+s}} dx dy$  with  $\chi_E$  denoting the characteristic function of the smooth subdomain  $E$ . The main purpose of this paper is to show that  $\lim_{p \rightarrow 1+} |\phi_p^s|_{L^\infty(\Omega)}^{1-p} = h_s(\Omega) = \lim_{p \rightarrow 1+} |\phi_p^s|_{L^1(\Omega)}^{1-p}$ , where  $\phi_p^s$  is the fractional  $(s, p)$ -torsion function of  $\Omega$ , that is, the solution of the Dirichlet problem for the fractional  $p$ -Laplacian:  $-(\Delta_p^s) u = 1$  in  $\Omega$ ,  $u = 0$  in  $\mathbb{R}^N \setminus \Omega$ . For this, we derive suitable bounds for the first eigenvalue  $\lambda_{1,p}^s(\Omega)$  of the fractional  $p$ -Laplacian operator in terms of  $\phi_p^s$ . We also show that  $\phi_p^s$  minimizes the  $(s, p)$ -Gagliardo seminorm in  $\mathbb{R}^N$ , among the functions normalized by the  $L^1$ -norm.

**Key words:** Fractional Cheeger problem, fractional  $p$ -Laplacian, fractional torsion functions.

**MSC 2010:** Primary 35P15, 35R11; Secondary 47A75.

## 1 Introduction

The Cheeger constant  $h(\Omega)$  of a bounded domain  $\Omega \subset \mathbb{R}^N$  ( $N > 1$ ) is defined by

$$h(\Omega) = \inf_{E \subset \Omega} \frac{P(E)}{|E|}, \quad (1.1)$$

where  $E$  is a smooth subset of  $\Omega$  and the nonnegative values  $P(E)$  and  $|E|$  denote, respectively, the distributional perimeter and the  $N$ -dimensional Lebesgue measure of  $E$ . A subset  $E$  that minimizes the quotient is a Cheeger set of  $\Omega$ .

In [7] Kawohl and Fridman proved that

$$h(\Omega) = \lim_{p \rightarrow 1+} \lambda_{1,p}(\Omega),$$

where  $\lambda_{1,p}(\Omega)$  is the first eigenvalue of the Dirichlet  $p$ -Laplacian operator, that is, the least real number  $\lambda$  such that the Dirichlet problem

$$\begin{cases} -\Delta_p u &= \lambda |u|^{p-2} u & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega, \end{cases}$$

has a nontrivial solution. (Let us recall that the  $p$ -Laplacian operator is defined by  $\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ ,  $p > 1$ .)

The first eigenvalue  $\lambda_{1,p}(\Omega)$  is also variationally characterized by

$$\lambda_{1,p}(\Omega) := \min \left\{ |\nabla u|_p^p : u \in W_0^{1,p}(\Omega), |u|_p = 1 \right\},$$

---

\*The authors acknowledge the support of CNPq-Brazil and FAPEMIG.

with  $|\cdot|_r$  standing for the usual norm of  $L^r(\Omega)$ ,  $1 \leq r \leq \infty$  (this notation will be adopted from now on).

In [2] a different characterization of the Cheeger constant of  $\Omega$  was obtained:

$$\lim_{p \rightarrow 1^+} \frac{1}{|\phi_p|_\infty^{p-1}} = h(\Omega) = \lim_{p \rightarrow 1^+} \frac{1}{|\phi_p|_1^{p-1}}$$

where  $\phi_p$  denotes the  $p$ -torsion function of  $\Omega$ , that is, the solution of the Dirichlet problem

$$\begin{cases} -\Delta_p u &= 1 & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega, \end{cases}$$

which is known as the  $p$ -torsional creep problem (see [6]).

We remark that

$$\frac{1}{|\phi_p|_1^{p-1}} = \min \left\{ |\nabla u|_p^p : u \in W_0^{1,p}(\mathbb{R}^N), |u|_1 = 1 \right\},$$

since the minimum is attained at  $\phi_p/|\phi_p|_1$ .

The fractional version of problem (1.1) consists in minimizing that quotient when  $P(E)$  is substituted by  $P_s(E)$ , given by

$$P_s(E) = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\chi_E(x) - \chi_E(y)|}{|x - y|^{N+s}} dx dy,$$

where  $\chi_E$  stands for the characteristic function of the smooth subdomain  $E$ . The value  $P_s(E)$  is called the (nonlocal)  $s$ -perimeter of  $E$ . So, the fractional Cheeger problem is the minimization problem

$$h_s(\Omega) = \inf_{E \subset \Omega} \frac{P_s(E)}{|E|},$$

and  $h_s(\Omega)$  is called the  $s$ -Cheeger constant of  $\Omega$ .

For  $1 < p < \infty$  and  $s \in (0, 1)$ , the fractional  $(s, p)$ -Laplacian  $(-\Delta)_p^s$  is the nonlinear nonlocal operator defined by

$$(-\Delta)_p^s u(x) = \lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}^N \setminus B_\epsilon(x)} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} dy.$$

This definition is consistent, up to a normalization constant depending on  $p$ ,  $s$  and  $N$ , with the usual  $p$ -Laplacian operator.

The first (fractional) eigenvalue  $\lambda_{1,p}^s(\Omega)$  of  $(-\Delta)_p^s$  is the least number  $\lambda$  such that the problem

$$\begin{cases} (-\Delta)_p^s u &= \lambda |u|^{p-2} u & \text{in } \Omega \\ u &= 0 & \text{on } \mathbb{R}^N \setminus \Omega, \end{cases}$$

has a nontrivial weak solution (see [8, 4]). Its variational characterization is given by (see [1]):

$$\lambda_{1,p}^s(\Omega) := \min \left\{ [u]_{s,p}^p : u \in W_0^{s,p}(\Omega), |u|_p = 1 \right\}, \quad (1.2)$$

where

$$[u]_{s,p} := \left( \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{1}{p}} \quad (1.3)$$

is the  $(s, p)$ -seminorm of Gagliardo in  $\mathbb{R}^N$  of a measurable function  $u$  and  $W_0^{s,p}(\Omega)$  is a suitable fractional Sobolev space defined in the sequel (see Definition 2.1).

In [1] Brasco, Lindgren and Parini proved the  $s$ -Cheeger version of the result originally obtained by Kawohl and Friedman [7] for the Cheeger problem:

$$h_s(\Omega) = \lim_{p \rightarrow 1^+} \lambda_{1,p}^s(\Omega). \quad (1.4)$$

In this paper, by assuming that  $\Omega$  is a Lipschitz bounded domain, we show, in the spirit of the paper [2], that the fractional version of the torsional creep problem,

$$\begin{cases} (-\Delta)_p^s u = 1 & \text{in } \Omega \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega, \end{cases}$$

is intrinsically connected to both the  $s$ -Cheeger problem and the first eigenproblem for the fractional Dirichlet  $p$ -Laplacian, as  $p$  goes to 1.

This connection will be developed in Section 2, where we introduce the  $(s, p)$ -torsion function of  $\Omega$ , that is, the weak solution  $\phi_p^s$  of (1). We will derive the estimates

$$\frac{1}{|\phi_p^s|_\infty^{p-1}} \leq \lambda_{1,p}^s(\Omega) \leq \left( \frac{|\Omega|}{|\phi_p^s|_1} \right)^{p-1} \quad (1.5)$$

and

$$\frac{|\phi_p^s|_\infty}{|\phi_p^s|_1} \leq \frac{1}{|B_1|} \left( \frac{sp + N(p-1)}{sp} \right)^{\frac{sp+N(p-1)}{sp}} \left( \frac{\lambda_{1,p}^s(\Omega)}{\lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp}}, \quad (1.6)$$

where  $B_1$  denotes the unit ball of  $\mathbb{R}^N$ .

Then, taking (1.4) into account, we will combine (1.5) with (1.6) in order to conclude the main result of this paper:

$$\lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_\infty^{p-1}} = h_s(\Omega) = \lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_1^{p-1}}.$$

Still in Section 2 we prove that  $\phi_p^s$  minimizes, in  $W_0^{s,p}(\Omega) \setminus \{0\}$ , the Rayleigh quotient  $[u]_{s,p}^p / |u|_1^p$ . As an immediate consequence of this fact, we show that  $\phi_p^s$  is a radial function when  $\Omega$  is a ball.

## 2 The main results

From now on  $\Omega$  denotes a Lipschitz bounded domain of  $\mathbb{R}^N$ ,  $N \geq 2$ , and  $0 < s < 1 < p < \frac{N}{s}$ .

**Definition 2.1** *The Sobolev space  $W_0^{s,p}(\Omega)$  is the closure of  $C_0^\infty(\Omega)$  with respect to the norm*

$$\|u\| := [u]_{s,p} + |u|_p, \quad (2.1)$$

where  $[u]_{s,p}$  is defined by (1.3).

Functions in  $W_0^{s,p}(\Omega)$  have a natural extension to  $\mathbb{R}^N$  and, although  $u = 0$  in  $\mathbb{R}^N \setminus \Omega$ , the identity

$$[u]_{s,p}^p = \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy + 2 \int_{\mathbb{R}^N \setminus \Omega} \int_{\Omega} \frac{|u(x)|^p}{|x - y|^{N+sp}} dx dy$$

shows dependence on values in  $\mathbb{R}^N \setminus \Omega$ .

It is worth mentioning that  $W_0^{s,p}(\Omega)$  is a reflexive Banach space and that this space coincides with the closure of  $C_0^\infty(\Omega)$  relative to the norm

$$u \mapsto \left( \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{1}{p}} + |u|_p,$$

if  $\partial\Omega$  is Lipschitz, (see [1, Proposition B.1]).

Moreover, thanks the fractional Poincaré inequality (see [1, Lemma 2.4])

$$|u|_p^p \leq C_{N,s,p,\Omega} [u]_{s,p}^p, \quad \forall u \in C_0^\infty(\Omega),$$

$[\cdot]_{s,p}$  is also a norm in  $W_0^{s,p}(\Omega)$ , equivalent to norm defined in (2.1).

We refer the reader to [3] for fractional Sobolev spaces.

**Definition 2.2** We say that a function  $u \in W_0^{s,p}(\Omega)$  is a weak solution of the fractional Dirichlet problem

$$\begin{cases} (-\Delta)_p^s u &= f & \text{in } \Omega \\ u &= 0 & \text{on } \mathbb{R}^N \setminus \Omega, \end{cases} \quad (2.2)$$

if

$$\langle (-\Delta)_p^s u, \varphi \rangle = \int_{\Omega} f \varphi \, dx, \quad \forall \varphi \in W_0^{s,p}(\Omega), \quad (2.3)$$

where

$$\langle (-\Delta)_p^s u, \varphi \rangle := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (\varphi(x) - \varphi(y))}{|x - y|^{N+sp}} \, dx dy,$$

a notation that will be used from now on.

Existence of a weak solution of (2.2), when  $f \in L^1(\Omega)$  follows from direct minimization in  $W_0^{s,p}(\Omega)$  of the functional

$$\frac{1}{p} [u]_{s,p}^p - \int_{\Omega} f(x) u(x) \, dx,$$

whereas uniqueness comes from, for instance, the comparison principle for the fractional  $p$ -Laplacian (see [8, Lemma 9]). The same principle shows that if  $f$  is nonnegative then the weak solution  $u$  is nonnegative as well. When  $f \in L^\infty(\Omega)$  and  $\Omega$  is sufficiently smooth, say with boundary at least of class  $C^{1,1}$ , the weak solutions are  $\alpha$ -Hölder continuous up to the boundary for some  $\alpha \in (0, 1)$ , see [5].

When  $f \equiv 1$  the Dirichlet problem (2.2) will be referred to as the  $(s, p)$ -fractional torsional creep problem and its unique weak solution will be called  $(s, p)$ -torsion function. Let us denote this function by  $\phi_p^s$ . We have  $\phi_p^s \geq 0$  and

$$\langle (-\Delta)_p^s \phi_p^s, \varphi \rangle = \int_{\Omega} \varphi \, dx, \quad \forall \varphi \in W_0^{s,p}(\Omega). \quad (2.4)$$

In particular, by taking  $\varphi = \phi_p^s$  we obtain

$$\langle (-\Delta)_p^s \phi_p^s, \phi_p^s \rangle = [\phi_p^s]_{s,p}^p = |\phi_p^s|_1.$$

**Theorem 2.1** We have

$$\frac{1}{|\phi_p^s|_1^{p-1}} = \min \left\{ [v]_{s,p}^p : v \in W_0^{s,p}(\Omega), |v|_1 = 1 \right\} = \left[ \frac{\phi_p^s}{|\phi_p^s|_1} \right]_{s,p}. \quad (2.5)$$

Moreover,  $\frac{\phi_p^s}{|\phi_p^s|_1}$  is the only nonnegative function attaining the minimum.

**Proof.** Since the functional  $v \mapsto |v|_1$  is not differentiable, we will first consider the minimization problem

$$m := \inf \left\{ [v]_{s,p}^p : v \in W_0^{s,p}(\Omega), \int_{\Omega} v \, dx = 1 \right\} \quad (2.6)$$

and show that it is uniquely solved by the positive function  $\frac{\phi_p^s}{|\phi_p^s|_1}$ .

Thus, let us take a sequence  $(u_n) \subset W_0^{s,p}(\Omega)$  such that

$$\int_{\Omega} u_n \, dx = 1 \quad \text{and} \quad [u_n]_{s,p}^p \rightarrow m.$$

We observe that the sequence  $(u_n)$  is bounded in  $L^1(\Omega)$ :

$$|u_n|_1^p \leq |\Omega|^{p-1} |u_n|_p^p \leq |\Omega|^{p-1} \lambda_{1,p}^s(\Omega)^{-1} [u_n]_{s,p}^p.$$

Since  $W_0^{s,p}(\Omega)$  is reflexive, the  $L^1$ -boundedness of  $(u_n)$  implies the existence of  $u \in W_0^{s,p}(\Omega)$  such that, up to a subsequence,  $u_n \rightharpoonup u$  (weak convergence) in  $W_0^{s,p}(\Omega)$  and  $u_n \rightarrow u$  in  $L^1(\Omega)$ . The convergence in  $L^1(\Omega)$  implies that  $\int_{\Omega} u \, dx = 1$ , so that  $m \leq [u]_{s,p}^p$ . On its turn, the weak convergence guarantees that

$$[u]_{s,p} + 1 = \|u\| \leq \liminf \|u_n\| = \liminf ([u_n]_{s,p} + 1) = m^{\frac{1}{p}} + 1.$$

It follows that  $m = [u]_{s,p}^p$ , so that the infimum in (2.6) is attained by the weak limit  $u$ .

By applying Lagrange multipliers, we infer the existence of a real number  $\lambda$  such that

$$\langle (-\Delta)_p^s u, \varphi \rangle = \lambda \int_{\Omega} \varphi \, dx, \quad \forall \varphi \in W_0^{s,p}(\Omega). \quad (2.7)$$

Taking  $\varphi = u$  we conclude that  $\lambda = m > 0$ , since  $m = [u]_{s,p}^p$  and  $\int_{\Omega} u \, dx = 1$ .

This fact and (2.7) imply that  $u$  is a weak solution of the Dirichlet problem

$$\begin{cases} (-\Delta)_p^s u &= m & \text{in } \Omega \\ u &= 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases}$$

By uniqueness, we have  $u = m^{\frac{1}{p-1}} \phi_p^s \geq 0$ . Since  $\int_{\Omega} u \, dx = 1$  we conclude that

$$m = \frac{1}{|\phi_p^s|_1^{p-1}} \quad \text{and} \quad u = \frac{\phi_p^s}{|\phi_p^s|_1}.$$

We remark that

$$\frac{1}{|\phi_p^s|_1^{p-1}} \leq [|v|]_{s,p}^p \leq [v]_{s,p}^p$$

for every  $v \in W_0^{s,p}(\Omega)$  such that  $|v|_1 = 1$ . This finishes the proof since

$$\frac{1}{|\phi_p^s|_1^{p-1}} = \left[ \frac{\phi_p^s}{|\phi_p^s|_1} \right]_{s,p} \quad \text{and} \quad \left| \frac{\phi_p^s}{|\phi_p^s|_1} \right|_1 = 1.$$

■

The next result recovers Lemma 4.1 of [5]:

**Corollary 2.1** *The  $(s,p)$ -torsion function is radial when  $\Omega$  is a ball.*

**Proof.** Let  $(\phi_p^s)^* \in W_0^{s,p}(\Omega)$  be the Schwarz symmetrization of  $\phi_p^s$ , that is, the radially decreasing function such that

$$\{\phi_p^s > t\}^* = \{(\phi_p^s)^* > t\}, \quad t > 0,$$

where, for any  $D \subset \mathbb{R}^N$ ,  $D^*$  stands for the  $N$ -dimensional ball with the same volume of  $D$ .

It is well-known that  $(\phi_p^s)^* \geq 0$ ,  $[(\phi_p^s)^*]_{s,p} \leq [\phi_p^s]_{s,p}$  and  $|(\phi_p^s)^*|_1 = |\phi_p^s|_1$ . Therefore,  $(\phi_p^s)^* / |(\phi_p^s)^*|_1$  attains the minimum in (2.5) and by uniqueness we have  $(\phi_p^s)^* = \phi_p^s$ . ■

It is also well-known that the first eigenfunctions of the fractional  $p$ -Laplacian belong to  $L^\infty(\Omega)$  and are either positive or negative almost everywhere in  $\Omega$ . Moreover, they are scalar multiple each other. So, let us denote by  $e_p^s$  the positive and  $L^\infty$ -normalized first eigenfunction. It follows that  $|e_p^s|_\infty = 1$  and

$$\begin{cases} (-\Delta)_p^s e_p^s &= \lambda_{1,p}^s(\Omega) (e_p^s)^{p-1} & \text{in } \Omega \\ e_p^s &= 0 & \text{on } \mathbb{R}^N \setminus \Omega, \end{cases}$$

meaning that

$$\langle (-\Delta)_p^s e_p^s, \varphi \rangle = \lambda_{1,p}^s(\Omega) \int_{\Omega} (e_p^s)^{p-1} \varphi \, dx, \quad \forall \varphi \in W_0^{s,p}(\Omega). \quad (2.8)$$

Of course, by taking  $\varphi = e_p^s$  in (2.8) we obtain

$$\langle (-\Delta)_p^s e_p^s, e_p^s \rangle = [e_p^s]_{s,p}^p = \lambda_{1,p}^s(\Omega) |e_p^s|_p^p.$$

As mentioned in the introduction,  $\lambda_{1,p}^s(\Omega)$  is variationally characterized by (1.2).

**Proposition 2.1** *Let  $u \in W_0^{s,p}(\Omega)$  be the weak solution of (2.2) with  $f \in L^\infty(\Omega) \setminus \{0\}$ . Then,*

$$|f|_\infty^{-\frac{1}{p-1}} u \leq \phi_p^s \quad \text{a.e. in } \Omega \quad (2.9)$$

and

$$\lambda_{1,p}^s(\Omega) \leq |f|_\infty \left( \frac{|\Omega|}{|u|_1} \right)^{p-1}. \quad (2.10)$$

**Proof.** Since  $u$  and  $\phi_p^s$  are both equal zero in  $\mathbb{R}^N \setminus \Omega$  and

$$\begin{aligned} \langle (-\Delta)_p^s (|f|_\infty^{-\frac{1}{p-1}} u), \varphi \rangle &= |f|_\infty^{-1} \langle (-\Delta)_p^s u, \varphi \rangle \\ &= |f|_\infty^{-1} \int_{\Omega} f \varphi \, dx \\ &\leq \int_{f \geq 0} \varphi \, dx \leq \int_{\Omega} \varphi \, dx \leq \langle (-\Delta)_p^s \phi_p^s, \varphi \rangle \end{aligned}$$

holds for every nonnegative  $\varphi \in W_0^{s,p}(\Omega)$ , (2.9) follows from the comparison principle (see [8, Lemma 9]).

In order to prove (2.10) we use (1.2) and Hölder inequality:

$$\begin{aligned} \lambda_{1,p}^s(\Omega) &\leq \frac{[u]_{s,p}^p}{|u|_p^p} \\ &= \frac{\int_{\Omega} f u \, dx}{|u|_p^p} \leq \frac{|f|_\infty |u|_1}{|u|_p^p} \leq \frac{|f|_\infty |u|_1}{|u|_1^p} |\Omega|^{p-1} = \|f\|_\infty \left( \frac{|\Omega|}{|u|_1} \right)^{p-1}. \end{aligned}$$

■

**Corollary 2.2** *It holds*

$$e_p^s \leq \lambda_{1,p}^s(\Omega)^{\frac{1}{p-1}} \phi_p^s \quad \text{a.e. in } \Omega \quad (2.11)$$

and

$$\frac{1}{|\phi_p^s|_\infty^{p-1}} \leq \lambda_{1,p}^s(\Omega) \leq \frac{|\Omega|^{p-1}}{|\phi_p^s|_1^{p-1}}. \quad (2.12)$$

**Proof.** Taking  $u = e_p^s$  and  $f = \lambda_{1,p}^s(\Omega)(e_p^s)^{p-1}$  in (2.9) we readily obtain (2.11). Hence, passing to maxima, we arrive at the first inequality in (2.12). The second inequality, on its turn, follows from (2.10) with  $u = \phi_p^s$  and  $f \equiv 1$ . ■

We would like to emphasize the following consequence of (2.11):  $\phi_p^s > 0$  almost everywhere in  $\Omega$ .

A Faber-Krahn inequality also holds true for the first fractional eigenvalue.

**Lemma 2.1 (Theorem 3.5 of [1])** Let  $p > 1$  and  $s \in (0, 1)$ . For every bounded domain  $D \subset \mathbb{R}^N$  we have

$$|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1) = |B|^{\frac{sp}{N}} \lambda_{1,p}^s(B) \leq |D|^{\frac{sp}{N}} \lambda_{1,p}^s(D) \quad (2.13)$$

where  $B$  is any  $N$ -dimensional ball and  $B_1$  denotes the unit ball of  $\mathbb{R}^N$ .

**Remark 2.1** Since  $h_s(D) = \lim_{p \rightarrow 1^+} \lambda_{1,p}^s(D)$  one has, immediately,

$$|B_1|^{\frac{s}{N}} h_s(B_1) = |B|^{\frac{s}{N}} h_s(B) \leq |D|^{\frac{s}{N}} h_s(D). \quad (2.14)$$

The next estimate is obtained by applying standard set-level techniques; however, the bounds obtained are adequate to study the asymptotic behavior as  $p \rightarrow 1^+$ .

**Proposition 2.2** Let  $u \in W_0^{1,p}(\Omega) \setminus \{0\}$  be a nonnegative, weak solution of (2.2) with  $f \in L^\infty(\Omega)$ . Then  $u \in L^\infty(\Omega)$  and

$$\frac{|u|_\infty}{|u|_1} \leq \frac{1}{|B_1|} \left( \frac{sp + N(p-1)}{sp} \right)^{\frac{sp + N(p-1)}{sp}} \left( \frac{|f|_\infty}{\lambda_{1,p}^s(B_1) |u|_\infty^{p-1}} \right)^{\frac{N}{sp}}. \quad (2.15)$$

**Proof.** For each  $k > 0$  we set

$$A_k = \{x \in \Omega : u(x) > k\}.$$

Since  $u \in W_0^{s,p}(\Omega)$  and  $u \geq 0$  in  $\Omega$ , the function

$$(u - k)^+ = \max\{u - k, 0\} = \begin{cases} u - k, & \text{if } u > k \\ 0, & \text{if } u \leq k \end{cases}$$

belongs to  $W_0^{s,p}(\Omega)$ . Therefore, choosing  $\varphi = (u - k)^+$  in (2.3) we obtain

$$\langle (-\Delta)_p^s u, (u - k)^+ \rangle = \int_{A_k} f(x)(u - k) \, dx.$$

It is not difficult to check that

$$[(u - k)^+]_{s,p}^p \leq \langle (-\Delta)_p^s u, (u - k)^+ \rangle.$$

Thus, we have

$$[(u - k)^+]_{s,p}^p \leq \int_{A_k} (u - k) f \, dx \leq |f|_\infty \int_{A_k} (u - k) \, dx. \quad (2.16)$$

We now consider  $k > 0$  such that  $|A_k| > 0$ . In order to estimate  $[(u - k)^+]_{s,p}^p$  from below, let us fix a ball  $B \subset \mathbb{R}^N$  and apply Lemma 2.1 to obtain

$$|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1) |A_k|^{-\frac{sp}{N}} \leq \lambda_{1,p}^s(A_k) \leq \frac{[(u - k)^+]_{s,p}^p}{\int_{A_k} (u - k)^p \, dx}.$$

Hence, Hölder's inequality yields

$$\left( \int_{A_k} (u - k) \, dx \right)^p \leq |A_k|^{p-1} \int_{A_k} (u - k)^p \, dx \leq |A_k|^{p-1} \frac{|A_k|^{\frac{sp}{N}} [(u - k)^+]_{s,p}^p}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)}.$$

Thus, it follows from (2.16) that

$$\begin{aligned} |B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1) |A_k|^{-\frac{sp + N(p-1)}{N}} \left( \int_{A_k} (u - k) \, dx \right)^p &\leq [(u - k)^+]_{s,p}^p \\ &\leq |f|_\infty \int_{A_k} (u - k) \, dx, \end{aligned}$$

what yields

$$\left( \int_{A_k} (u - k) \, dx \right)^{p-1} \leq \frac{|f|_\infty |A_k|^{\frac{sp+N(p-1)}{N}}}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)}$$

and so

$$\left( \int_{A_k} (u - k) \, dx \right)^{\frac{N(p-1)}{sp+N(p-1)}} \leq \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp+N(p-1)}} |A_k|. \quad (2.17)$$

Define

$$g(k) := \int_{A_k} (u - k) \, dx = \int_k^\infty |A_k| \, dt,$$

the last equality being a consequence of Cavalieri's principle. Combining the definition of  $g(k)$  with (2.17), we have

$$[g(k)]^{\frac{N(p-1)}{sp+N(p-1)}} \leq - \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp+N(p-1)}} g'(k).$$

Therefore,

$$1 \leq - \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp+N(p-1)}} [g(k)]^{-\frac{N(p-1)}{sp+N(p-1)}} g'(k). \quad (2.18)$$

Integration of (2.18) from 0 to  $k$  produces

$$\begin{aligned} k &\leq \left( \frac{sp + N(p-1)}{sp} \right) \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp+N(p-1)}} [g(0)^{\frac{sp}{sp+N(p-1)}} - g(k)^{\frac{sp}{sp+N(p-1)}}] \\ &\leq \left( \frac{sp + N(p-1)}{sp} \right) \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp+N(p-1)}} (|u|_1)^{\frac{sp}{sp+N(p-1)}}, \end{aligned}$$

since  $g(k) \geq 0$  and  $g(0) = |u|_1$ .

Let  $c$  denote, just for a moment, the right-hand side of the latter inequality. We have proved that  $k \leq c$  whenever  $|A_k| > 0$ . Since  $c$  does not depend on  $k$  this implies that  $|A_k| = 0$  for every  $k > c$ , thus allowing us to conclude that  $u \in L^\infty(\Omega)$  and also that  $|u|_\infty \leq c$ . So,

$$|u|_\infty \leq \left( \frac{sp + N(p-1)}{sp} \right) \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp+N(p-1)}} |u|_1^{\frac{sp}{sp+N(p-1)}}$$

or, what is the same,

$$|u|_\infty^{1+\frac{N(p-1)}{sp}} \leq \left( \frac{sp + N(p-1)}{sp} \right)^{\frac{sp+N(p-1)}{sp}} \left( \frac{|f|_\infty}{|B_1|^{\frac{sp}{N}} \lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp}} |u|_1,$$

from what follows (2.15). ■

**Corollary 2.3** *The  $(s, p)$ -torsion function  $\phi_p^s$  belongs to  $L^\infty(\Omega)$  and, in addition,*

$$\frac{1}{|\Omega|} \leq \frac{|\phi_p^s|_\infty}{|\phi_p^s|_1} \leq \frac{1}{|B_1|} \left( \frac{sp + N(p-1)}{sp} \right)^{\frac{sp+N(p-1)}{sp}} \left( \frac{\lambda_{1,p}^s(\Omega)}{\lambda_{1,p}^s(B_1)} \right)^{\frac{N}{sp}}. \quad (2.19)$$



**Proof.** The first inequality is obvious. Proposition 2.2 with  $u = \phi_p^s$  and  $f \equiv 1$  yields

$$\frac{|\phi_p^s|_\infty}{|\phi_p^s|_1} \leq \frac{1}{|B_1|} \left( \frac{sp + N(p-1)}{sp} \right)^{\frac{sp+N(p-1)}{sp}} \left( \frac{1}{\lambda_{1,p}^s(B_1) |\phi_p^s|_\infty^{p-1}} \right)^{\frac{N}{sp}}.$$

Now, the second inequality in (2.19) follows from the first inequality in (2.12). ■

**Theorem 2.2** *One has*

$$\lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_\infty^{p-1}} = h_s(\Omega) = \lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_1^{p-1}}.$$

**Proof.** Taking (1.4) into account, we have

$$\lim_{p \rightarrow 1^+} \frac{\lambda_{1,p}^s(\Omega)}{\lambda_{1,p}^s(B_1)} = \frac{h_s(\Omega)}{h_s(B_1)} \in (0, \infty).$$

Hence, it follows from (2.19) that

$$\lim_{p \rightarrow 1^+} \left( \frac{|\phi_p^s|_\infty}{|\phi_p^s|_1} \right)^{p-1} = 1.$$

Thus, by making  $p$  go to 1 in (2.12) we have

$$\begin{aligned} \lim_{p \rightarrow 1^+} \lambda_{1,p}^s(\Omega) &\leq \lim_{p \rightarrow 1^+} \frac{|\Omega|^{p-1}}{|\phi_p^s|_1^{p-1}} \\ &= \lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_1^{p-1}} \\ &= \lim_{p \rightarrow 1^+} \left( \frac{|\phi_p^s|_\infty}{|\phi_p^s|_1} \right)^{p-1} \lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_\infty^{p-1}} = \lim_{p \rightarrow 1^+} \frac{1}{|\phi_p^s|_\infty^{p-1}} \leq \lim_{p \rightarrow 1^+} \lambda_{1,p}^s(\Omega). \end{aligned}$$

Since  $\lim_{p \rightarrow 1^+} \lambda_{1,p}^s(\Omega) = h_s(\Omega)$ , we are done. ■

In [1], the authors also proved that

$$h_s(\Omega) = \inf \left\{ [v]_{s,1}^p : v \in W_0^{s,1}(\Omega), |v|_1 = 1 \right\}.$$

Since  $W_0^{s,1}(\Omega)$  is not reflexive, they were able to prove that the minimum  $h_s(\Omega)$  is attained on the larger Sobolev space

$$\mathcal{W}_0^{s,1}(\Omega) := \left\{ v \in L^1(\Omega) : [v]_{s,1}^p < \infty, u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega \right\}.$$

For completeness, we state the following result on the behavior of the  $L^1$ -normalized family  $\left\{ \frac{\phi_p^s}{|\phi_p^s|_1} \right\}$  as  $p \rightarrow 1$ .

It corresponds to [1, Theorem 7.2], which was proved for the family  $\left\{ \frac{e_p^s}{|e_p^s|_1} \right\}$ . Its proof follows the same script and will be omitted.

**Theorem 2.3** *Let  $u_p := \frac{\phi_p^s}{|\phi_p^s|_1}$ . There exists a sequence  $(p_n)$  such that  $p_n \rightarrow 1^+$  and  $u_{p_n} \rightarrow u$  in  $L^q(\Omega)$ , for every  $q < \infty$ . The limit function  $u$  is a solution of the minimization problem*

$$h_s(\Omega) = \min_{v \in \mathcal{W}_0^{s,1}(\Omega)} \left\{ [v]_{s,1} : u \geq 0, |u|_1 = 1 \right\}.$$

Moreover,  $u \in L^\infty(\Omega)$  and

$$\frac{1}{|\Omega|} \leq |u|_\infty \leq \frac{1}{|B_1|} \left( \frac{h_s(\Omega)}{h_s(B_1)} \right)^{\frac{N}{s}}. \quad (2.20)$$

The upper bound that appears in the statement of Theorem 7.2 of [1] is

$$\left[ \frac{|B|^{\frac{N-s}{N}}}{P_s(B)} \right]^{\frac{N}{s}} h_s(\Omega)^{\frac{N}{s}}.$$

However, it is very simple to check, by applying (2.14), that it is equal to the upper bound in (2.20).

We remark that, once obtained the convergence in  $L^q(\Omega)$  stated above, the upper bound in (2.20) follows from (2.19). Indeed, since

$$|u|_q = \lim_{n \rightarrow \infty} |u_{p_n}|_q \leq |\Omega|^{\frac{1}{q}} \lim_{n \rightarrow \infty} |u_{p_n}|_\infty$$

(2.19) implies that

$$|\Omega|^{-\frac{1}{q}} |u|_q \leq \lim_{n \rightarrow \infty} |u_{p_n}|_\infty \leq \frac{1}{|B_1|} \left( \frac{h_s(\Omega)}{h_s(B_1)} \right)^{\frac{N}{s}}.$$

Hence, the upper bound in (2.20) follows, since  $|u|_\infty = \lim_{q \rightarrow \infty} |\Omega|^{-\frac{1}{q}} |u|_q$ .

The lower bound in (2.20), which does not appear in the statement of Theorem 7.2 of [1], follows by taking  $q = 1$ , since

$$1 = \lim_{n \rightarrow \infty} |u_{p_n}|_1 = |u|_1 \leq |u|_\infty |\Omega|.$$

It is interesting to note that, as it happens with the standard  $p$ -torsion functions,  $|u|_\infty = |\Omega|^{-1}$  when  $\Omega$  is a ball. In fact, in this case (2.14) yields

$$\frac{1}{|B_1|} \left( \frac{h_s(\Omega)}{h_s(B_1)} \right)^{\frac{N}{s}} = \frac{1}{|\Omega|}.$$

## References

- [1] L. Brasco, E. Lindgren and E. Parini: *The fractional Cheeger problem*, Interfaces and Free Boundaries **16** (2014), 419–458.
- [2] H. Bueno and G. Ercole: *Solutions of the Cheeger problem via torsion functions*, J. Math. Anal. Appl. **381** (2011), 263–279.
- [3] R. Di Nezza, G. Palatucci and E. Valdinoci: *Hitchhikers guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (5) (2012), 521–573.
- [4] G. Franzina and G. Palatucci: *Fractional  $p$ -eigenvalue*, Riv. Mat. Univ. Parma, **5**, no. 5 (2014), 373–386.
- [5] A. Iannizzotto, S. Mosconi and M. Squassina: *Global Hölder regularity for the fractional  $p$ -laplacian*, preprint (2014) available at <http://arxiv.org/abs/1411.2956>.
- [6] B. Kawohl: *On a family of torsional creep problems*, J. Reine Angew. Math. **410** (1990), 1–22.
- [7] B. Kawohl and V. Fridman: *Isoperimetric estimates for the first eigenvalue of the  $p$ -Laplace operator and the Cheeger constant*, Comment Math. Univ. **44** (2003), 659–667.
- [8] E. Lindgren and P. Lindqvist: *Fractional Eigenvalues*, Calc. Var. Partial Differential Equations **49**, no. 1-2 (2014), 795–826.