

Twisted Alexander invariants of knot group representations II; computation and duality.

Takefumi Nosaka¹

Abstract

Given a homomorphism from a link group to a group, we introduce a K_1 -class, which is a generalization of the 1-variable Alexander polynomial. We compare the K_1 -class with K_1 -classes in [Nos] and with Reidemeister torsions. As a corollary, we show a relation to Reidemeister torsions of finite cyclic covering spaces, and show reciprocity in some senses.

Keywords

knot, Alexander polynomial, K_1 -group, Novikov ring

1 Introduction

Let $L \subset S^3$ be a knot in the 3-sphere. The Alexander polynomial of L has been studied in many ways, and its applications and beautiful properties are discovered, including, e.g., relations to Reidemeister torsions, Fox derivatives, and applications to cyclic coverings and slice knots. As a generalization, after twisted Alexander polynomials are introduced [Wada, Lin], similar applications and properties are discovered (however, such results frequently require some assumptions); see [FV, FKK] and references therein.

In the previous paper [Nos], which is inspired by [Mil1, Tur], given a homomorphism from a link group to a group, the author suggested elements of a K_1 -group, which are a generalization of the twisted Alexander polynomials. This paper gives several approaches to the K_1 -class in another way. In Sections 2 and 4, we give two other definitions as K_1 -classes from the viewpoint of the Fox derivatives or Reidemeister torsion over K_1 . Here, an advantage of the definitions is applicable to not only knots but also links; see Definition 2.2. Furthermore, in the knot case, we show the equivalence (up to some ambiguity) of the three definitions (Theorem 3.1). As a corollary of the equivalence, we compute the K_1 -classes of some 2-bridge knots; see Section 2.1.

In Sections 5–7, we will address some applications. First, under some conditions, we give a relation to (commutative) Reidemeister torsion of the m -fold cyclic covering space of $S^3 \setminus L$; see Section 5. Next, we will see that the conditions are suitable to group homomorphisms $f : \pi_1(S^3 \setminus L) \rightarrow \mathbb{Z} \rtimes G$, where G is a finite group. In this situation, we show (Theorem 6.1) a duality theorem of the K_1 -class, as in reciprocity of the (twisted) Alexander polynomials [FKK, Mil1, Tur]. In application, we give an estimate of sliceness of knots, which is a slight generalization of the works of Herald-Kirk-Livingston [HKL, KL]. In fact, we find that the conditions in Section 7 are applicable to the Casson-Gordan theory [CG].

Finally, under some conditions, we give a relation to the circle valued Morse theory (see Appendix A), and a reduction to the higher order Alexander polynomial [C, Har]; see Appendix B. These discussions rely on the works of [P, Fri2].

¹ E-mail address: nosaka@math.titech.ac.jp

Conventional notation. For a group G and a commutative ring A , we denote the group ring by $A[G]$, and the abelianization by G_{ab} . Every non-commutative ring R has always 1, and is assumed to satisfy that R^r and R^s with $r \neq s$ are not isomorphic as R -modules. We mean by R^\times the multiplicative group consisting of units in R .

Acknowledgments

The author expresses his gratitude to Takahiro Kitayama, Ryoto Tange for valuable comments.

2 Definition of the Alexander K_1 -class

We give the definition as a generalization of the Alexander polynomial; see Definition 2.2.

As a similar setting to [Nos], we set up algebraic terminologies and need an assumption. Let \mathcal{A} be a ring, which is possibly non-commutative, and take a ring isomorphism $\kappa : \mathcal{A} \rightarrow \mathcal{A}$. Then, we have the completed skew Laurent-polynomial ring $\mathcal{A}_\kappa((\tau))$. Namely, $\mathcal{A}_\kappa((\tau))$ is the set of formal power series $\sum_{i=-N}^{\infty} a_i \tau^i$ where $a_i \in \mathcal{A}$ and $\tau^n a = \kappa^n(a) \tau^n$; in other words, $\mathcal{A}_\kappa((\tau))$ is equal to $\mathcal{A}_\kappa[[\tau]][\tau^{-1}]$, which is also called the Novikov ring in [Fri1, PR]. Furthermore, let us fix a group G with presentation $\langle x_1, \dots, x_m | r_1, \dots, r_{m-1} \rangle$ of deficiency 1, and consider the following assumption throughout this paper.

Assumption (\dagger) Let \mathcal{A} be a ring and $\kappa : \mathcal{A} \rightarrow \mathcal{A}$ be as above. Suppose a ring homomorphism $\rho : \mathbb{Z}[G] \rightarrow \mathcal{A}_\kappa((\tau))$ such that, for any $i \leq m$, there is $w_i \in \mathcal{A}_\kappa((\tau))^\times$ such that $\rho(x_i) = w_i^{-1} \tau w_i$.

Example 2.1 (A special case of Novikov completion). Set a link L in the 3-sphere S^3 . Choose a link diagram D , and let m be the number of the arcs on D . Then, the Wirtinger presentation from D gives such a presentation $\pi_1(S^3 \setminus L) \cong \langle x_1, \dots, x_m | r_1, \dots, r_{m-1} \rangle$, possibly $r_j = 1$ (see, e.g., [Lic], [Wada, §5]).

Given a group homomorphism $h : \pi_1(S^3 \setminus L) \rightarrow H \rtimes \mathbb{Z}$ such that H is a group, $h(x_i) = (g_i, 1)$ for some $g_i \in G$. Let \mathcal{A} be the group ring $B[H]$ over a commutative ring B . If we replace $h(x_i)$ by $h(x_i)\tau$ and define $\kappa(a) := (0, 1)(a, 0)(0, -1)$ for $a \in H$ we have $\mathcal{A}_\kappa((\tau))$. This h canonically gives rise to $\rho : \mathbb{Z}[\pi_1(S^3 \setminus L)] \rightarrow \mathcal{A}_\kappa((\tau))$ satisfying (\dagger).

In general, we obtain such an h from any group G and any group homomorphism $f : \pi_1(S^3 \setminus L) \rightarrow G$, as follows. Let $\text{Ab} : \pi_1(S^3 \setminus L) \rightarrow \mathbb{Z}^{\#L}$ be an abelianization, and $\mu : \mathbb{Z}^{\#L} \rightarrow \mathbb{Z}$ be the multiplication such that $\mu \circ \text{Ab}(x_i) = 1$ for any i . Notice the isomorphism $\pi_1(S^3 \setminus L) \cong \text{Ker}(\mu \circ \text{Ab}) \rtimes \mathbb{Z}$. Let $H \subset G$ be the restricted image $f(\text{Ker}(\mu \circ \text{Ab}))$, and let $\mathbb{Z} = \{\tau^n\}_{n \in \mathbb{Z}}$ act on H by $g \cdot \tau^n := f(x_1)^n g f(x_1)^{-n}$. By the action, we have the semi-direct product $H \rtimes \mathbb{Z}$, and can define a homomorphism

$$h : \pi_1(S^3 \setminus L) = \text{Ker}(\mu \circ \text{Ab}) \rtimes \mathbb{Z} \rightarrow H \rtimes \mathbb{Z} \quad \text{by} \quad h(g, n) = (f(g), n),$$

which satisfy the condition in the previous paragraph.

Let us review up K_1 -groups. For a ring R with unit, let $GL_n(R)$ be the general linear group over R of size n . Since $GL_n(R)$ diagonally injects into $GL_{n+1}(R)$, we have the colimit $GL(R) = \lim GL_n(R)$. The K_1 -group, $K_1(R)$, is defined to be the abelianization $GL(R)_{\text{ab}}$. By abuse of notation, we often regard elements of $GL_n(R)$ as those of $K_1(R)$. When $R = \mathcal{A}_\kappa((\tau))$

as above, -1 , τ , and $\rho(g)$ are represented by the invertible (1×1) -matrices. Let $\pm\tau$ denote the subgroup of $K_1(\mathcal{A}_\kappa((\tau)))$ generated by -1 and τ , which is isomorphic to either \mathbb{Z} or $\mathbb{Z} \times \mathbb{Z}/2$. Similarly, let $\pm\rho(G)$ be the subgroup of $K_1(\mathcal{A}_\kappa((\tau)))$ generated by -1 and the image $\rho(G)$. We later use the quotient groups $K_1(\mathcal{A}_\kappa((\tau)))/\pm\tau$ and $K_1(\mathcal{A}_\kappa((\tau)))/\pm\rho(G)$.

Furthermore, we study the Fox derivative and a Jacobi matrix. Let \mathcal{F} be the free group with a basis, x_1, \dots, x_m . We define a \mathbb{Z} -linear map $\frac{\partial}{\partial x_i} : \mathbb{Z}[\mathcal{F}] \rightarrow \mathbb{Z}[\mathcal{F}]$ by the following identities:

$$\frac{\partial x_j}{\partial x_i} = \delta_{ij}, \quad \frac{\partial(hk)}{\partial x_i} = \frac{\partial h}{\partial x_i} + h \frac{\partial k}{\partial x_i} \quad (h, k \in F).$$

Consider the $(m-1) \times m$ matrix, A_ρ , over $\mathcal{A}_\kappa((\tau))$ whose (i, j) -th entry is $\rho(\frac{\partial r_i}{\partial x_j}) \in \mathcal{A}_\kappa((\tau))$. For $1 \leq k \leq m$, let us denote by $A_{\rho,k}$ by the $(m-1) \times (m-1)$ -matrix obtained from A_ρ by removing the k -th column.

Definition 2.2. Choose $k \in \mathbb{N}$ as above. Suppose that $A_{\rho,k}$ is an invertible matrix. We define the K_1 -Alexander class (with respect to ρ) to be the K_1 -class

$$[A_{\rho,k}] \in K_1(\mathcal{A}_\kappa((\tau)))/\pm\tau.$$

On the other hand, if $A_{\rho,k}$ is not invertible, we define the K_1 -class to be zero.

Theorem 2.3. Let $G = \pi_1(S^3 \setminus L)$ be a link group with a Wirtinger presentation, as in Example 2.1. Then, the K_1 -class $\Delta_\rho^{K_1}$ depends only on the homomorphism $\rho : \mathbb{Z}[\pi_1(S^3 \setminus L)] \rightarrow \mathcal{A}_\kappa((\tau))$.

We will give the proof in §2.2. Although we suppose invertibility of $A_{F,W}$, we give a criterion for the invertibility:

Proposition 2.4. Suppose $\#L = 1$, i.e., L is a knot. Then, L is fibered if and only if $A_{\rho,k}$ is invertible for any homomorphism ρ satisfying (\dagger) .

We conclude this section by explaining that the twisted Alexander polynomial of [Wada] can be formulated from our K_1 -Alexander class:

Example 2.5. Let R be a commutative ring. The paper [Wada, §5] defines a polynomial from a representation $\rho^{\text{pre}} : \pi_1(S^3 \setminus L) \rightarrow SL_n(R)$ as follows. For the formulation, let \mathcal{A} be the matrix ring $\text{Mat}(n \times n, R)$, and let κ be the identity $\text{id}_\mathcal{A}$. Formally, let τ be $\rho^{\text{pre}}(\mathbf{m})$ as a commutative indeterminate. Since there is a ring homomorphism $\mathbb{Z}[SL_n(R)] \rightarrow \text{Mat}(n \times n, R)$ which sends $\sum_g a_g g$ to $\sum_g a_g g$, the ρ^{pre} gives rise to a ring homomorphism $\rho : \mathbb{Z}[\pi_1(S^3 \setminus L)] \rightarrow \mathcal{A}_\kappa((\tau))$. Notice that the determinant $\text{Mat}(n \times n, R_\kappa((\tau))) \rightarrow R_\kappa((\tau))$ induces a homomorphism

$$\det : K_1(\mathcal{A}_\kappa((\tau)))/\pm\tau \longrightarrow R_\kappa((\tau))^\times/\pm\tau.$$

Then, if $\#L = 1$, checking [Wada, Corollary 5] carefully, we verify by construction that the determinant $\det(\Delta_\rho^{K_1}) \in R_\kappa((\tau))^\times/\pm\tau$ exactly coincides with the twisted Alexander polynomial of [Wada]. In other words, our K_1 -class $\Delta_\rho^{K_1}$ is a lift of the twisted Alexander polynomial.

2.1 Computation for some 2-bridge knots

We will compute the K_1 -classes with respect to every 2-bridge knots of Seifert genus 1.




Figure 1: The 2-bridge knot of genus 1.

For non-zero integers m and n , let $K(m, n)$ be the 2-bridge knot $S(4mn + 1, 2m)$ in Schubert's form; see Figure 1. We may suppose $m > 0$. As is known, every 2-bridge knot of genus 1 is represented as one of $K(m, n)$. Thanks to Proposition 2.1 in [HT], the knot group $\pi_1(S^3 \setminus K(m, n))$ has the presentation

$$\langle x, y | w^n x = y w^n \rangle, \quad \text{where } w = (xy^{-1})^m (x^{-1}y)^m,$$

where x and y are conjugate to a meridian. Moreover, as seen in the proof of the proposition, this presentation is strongly Tietze equivalent to a Wirtinger presentation.

Proposition 2.6. *Let $m > 0$. Consider the following element of $\mathcal{A}_\kappa((\tau))$:*

$$Z_{m,n} := \begin{cases} \rho(w^n) + (1 - \rho(y)) \frac{1 - \rho(w^n)}{1 - \rho(w)} (1 - \rho(wx^{-1})) \frac{1 - \rho(xy^{-1})^m}{1 - \rho(xy^{-1})} & (n > 0), \\ \rho(w^n) + (\rho(y) - 1) \rho(w^{-1}) \frac{1 - \rho(w^n)}{1 - \rho(w)} (1 - \rho(wx^{-1})) \frac{1 - \rho(xy^{-1})^m}{1 - \rho(xy^{-1})} & (n < 0). \end{cases}$$

The matrix $A_{F,W}$ is invertible if and only if $Z_{m,n}$ is an invertible element. If so, the K_1 -Alexander invariant is given by

$$\Phi_{\rho,k} = Z_{m,n} \in K_1(\mathcal{A}_\kappa((\tau))) / \pm \tau.$$

Proof. Let $r = w^n x w^{-n} y^{-1}$. It is sufficient to show $Z_{m,n} = \rho(\frac{\partial r}{\partial x})$. Notice that

$$\begin{aligned} \frac{\partial r}{\partial x} &= \frac{\partial w^n}{\partial x} + w^n \frac{\partial}{\partial x} (x w^{-n} y^{-1}) = \frac{\partial w^n}{\partial x} + w^n (1 + x \frac{\partial}{\partial x} (w^{-n})) \\ &= \begin{cases} (1 + w + \dots + w^{n-1}) \frac{\partial w}{\partial x} + w^n + y w^n (1 + w^{-1} + \dots + w^{1-n}) \frac{\partial w^{-1}}{\partial x} & (n > 0), \\ (1 + w^{-1} + \dots + w^{n+1}) \frac{\partial w}{\partial x} + w^n + y w^n (1 + w^{-1} + \dots + w^{-n-1}) \frac{\partial w^{-1}}{\partial x} & (n < 0), \end{cases} \\ &= \begin{cases} w^n + (1 - y)(1 + w + \dots + w^{n-1}) \frac{\partial w}{\partial x} & (n > 0), \\ w^n + (y - 1)(w^{-1} + \dots + w^n) \frac{\partial w}{\partial x} & (n < 0). \end{cases} \end{aligned}$$

Here, the last equality is due to $\frac{\partial w^{-1}}{\partial x} = -w^{-1} \frac{\partial w}{\partial x}$. We should observe

$$\begin{aligned} \frac{\partial w}{\partial x} &= (1 + xy^{-1} + \dots + (xy^{-1})^{m-1}) - (xy^{-1})^m (1 + x^{-1}y + \dots + (x^{-1}y)^{m-1}) x^{-1} \\ &= (1 - (xy^{-1})^m y^{-1} (yx^{-1})^m) (1 + xy^{-1} + \dots + (xy^{-1})^{m-1}) \\ &= (1 - wx^{-1}) (1 - (xy^{-1})^m) / (1 - xy^{-1}). \end{aligned}$$

Therefore, we have $Z_{m,n} = \rho(\frac{\partial r}{\partial x})$ as required. \square

The above computation is done as an element of $\mathcal{A}_\kappa((\tau))$; however, there are many examples such that $\Phi_{\rho,k}$ the K_1 -Alexander invariant $\Phi_{\rho,k}$ can not be represented by any 1×1 -matrix. e.g, L is a non-fibered pretzel knot. Indeed, even concerning the classical Alexander module, the minimal number of sizes of the matrix $\Phi_{\rho,k}$ is estimated by Nakanishi index.

2.2 Proof of Theorem 2.3

For the proof, we review the strongly Tietze transformations [Wada]. For a finite presentable group $G = \langle g_1, \dots, g_m | r_1, \dots, r_n \rangle$ and a word w of g_1, \dots, g_m , the transformation of the following types are called the *strongly Tietze transformations*:

- (Ia) To replace one of the relators r_i by its inverse r_i^{-1} .
- (Ib) To replace one of the relators r_i by its conjugate $wr_i^{-1}w^{-1}$.
- (Ic) To replace one of the relators r_i by $r_i r_j$ for any $j \neq i$.
- (II) To add a new generator y and a new relator yw^{-1} . In other words, the resulting presentation is given by $\langle g_1, \dots, g_m, y | r_1, \dots, r_n, yw^{-1} \rangle$.

Two presentations of G are said to be *strongly Tietze equivalent*, if they are related by a finite sequence of the operations of the above types and their inverse operations.

Proof of Theorem 2.3. It is shown [Wada, Lemma 6 in §5] that any two Wirtinger presentations for a given link L are strongly Tietze equivalent. Hence, for the proof of Theorem 2.3, it is enough to show the following proposition, which claims an invariance with respect to strongly Tietze equivalence: \square

Proposition 2.7 (cf. [Ki1, Theorem 4.5]). *The K_1 -class $[\Phi_{\rho,k}]$ in $K_1(\mathcal{A}_\kappa((\tau)))/\pm\tau$ does not depend on the choice of k . Moreover, if we change another presentation of G which is strongly Tietze equivalent to the above presentation, the associated K_1 -class $[\Phi_{\rho,k}]$ in $K_1(\mathcal{A}_\kappa((\tau)))/\pm\tau$ is invariant.*

Proof. The proof is essentially based on the proofs of [Wada, Lemma 2 in §3] and [Ki1, Theorem 4.5]. Following the proofs, we will check (i) the independence of the choice of k , and (ii) the invariance with respect to each transformations (Ia)(Ib)(Ic)(II).

To show (i), choose k' . Let $e_j(a)$ be the diagonal $(m-1)^2$ -matrix whose (j, j) -th entry is a and (k, k) -th entry is 1 where $j \neq k$. Then, according to the fundamental formula

$$\sum_{j=1}^m \frac{\partial r_i}{\partial x_j} (x_j - 1) = r_i - 1$$

(see [Fox, (2.3)]), we have

$$\begin{aligned} A_{\rho,k'} e_k(\rho(x_k) - 1) &= \left(\dots, \rho\left(\frac{\partial r_i}{\partial x_k}\right) \rho(x_k - 1), \dots \right) = \left(\dots, -\sum_{j \neq k} \rho\left(\frac{\partial r_i}{\partial x_j}\right) \rho(x_j - 1), \dots \right) \\ &= \left(\dots, -\rho\left(\frac{\partial r_i}{\partial x_{k'}}\right) \rho(x_{k'} - 1), \dots \right) = (-1)^{k-k'} A_{\rho,k} e_{k'}(\rho(x_{k'}) - 1) \in \text{Mat}(m \times m; \mathcal{A}_\kappa((\tau))). \end{aligned}$$

Notice that $e_k(\rho(x_k) - 1)$ is invertible, since $1 - \rho(x_k)$ is invertible because of $(1 - \rho(x_k))(1 + \sum_{j=1}^\infty w_k^{-1} \tau^j w_k) = 1$. Therefore, the invertibility of $A_{\rho,k}$ implies that $A_{\rho,k'}$ is also invertible. Moreover, since $\rho(x_k) - 1 = \rho(w_k^{-1})(\tau - 1)\rho(w_k) = \tau - 1$ in K_1 by Assumption (\dagger), we obtain $A_{\rho,k'} = A_{\rho,k}$ in the quotient $K_1(\mathcal{A}_\kappa((\tau)))/\pm\tau$.

Next, concerning (ii), we consider strongly Tietze transformations. Notice

$$\frac{\partial(r_i^{-1})}{\partial x_j} = -r_i^{-1} \frac{\partial r_i}{\partial x_j}, \quad \frac{\partial(wr_iw^{-1})}{\partial x_j} = w \frac{\partial r_i}{\partial x_j}, \quad \frac{\partial(r_i r_\ell)}{\partial x_j} = \frac{\partial r_i}{\partial x_j} + r_i \frac{\partial r_\ell}{\partial x_j}.$$

Therefore, the changes of the K_1 -classes with respect to Ia and Ib are $A_{\rho,k} \mapsto -A_{\rho,k}$, and $A_{\rho,k} \mapsto (-1)^k \rho(w) A_{\rho,k}$, respectively. Furthermore, recalling from Whitehead Lemma [Wei, Lemma III.1.3.3] that any elementary matrix is 1 in K_1 , we can easily verify that the changes of the K_1 -values with respect to Ic and II are $A_{\rho,k} \mapsto A_{\rho,k}$, and $A_{\rho,k} \mapsto \rho(w^{-1}) A_{\rho,k}$, respectively. Since $\rho(x_k) = \tau \in K_1(\mathcal{A}_\kappa((\tau)))$, we have $\rho(w) = \tau^{n_w} \in K_1(\mathcal{A}_\kappa((\tau)))$ for some $n_w \in \mathbb{Z}$ by assumption. Hence, the observation in the quotient $K_1(\mathcal{A}_\kappa((\tau)))/\pm \tau$ proves (ii). \square

3 Relation to the K_1 -class in [Nos]

In what follows, we let L be a knot embedded in the 3-sphere S^3 , i.e., $\#L = 1$, and choose a knot diagram on \mathbb{R}^2 .

In the paper [Nos], the author constructed another K_1 -class, where the construction is inspired by [Lin]. This section shows (Theorem 3.1) that this class and another $[\Phi_{\rho,k}]$ in §2 are almost equal. In this section, we assume that L is a knot, and choose a meridian $\mathbf{m} \in \pi_1(S^3 \setminus L)$ such that $\rho(\mathbf{m}) = \tau$.

We recall the presentation (1) and the definition of the K_1 -class below. We choose a Seifert surface F of genus g and a bouquet of circles $W \subset F$ such that W is a deformation retract of F and the inclusion $F \subset S^3$ is isotopic to the standard embedding $W \subset F$. Take a bicollar $F \times [-1, 1]$ of F such that $F \times \{0\} = F$. Let $\iota_\pm : F \rightarrow S^3 \setminus F$ be the embeddings whose images are $F \times \{\pm 1\}$. Take generating sets $W := \{u_1, \dots, u_{2g}\}$ of $\pi_1 F$ and $X := \{x_1, \dots, x_{2g}\}$ of $\pi_1(S^3 \setminus F)$, and set $y_i^\# := (\iota_+)_*(u_i)$ and $z_i = (\iota_-)_*(u_i)$; a von Kampen argument yields a presentation

$$\langle x_1, \dots, x_{2g}, \mathbf{m} \mid r_i^F := \mathbf{m}^{-1} y_i \mathbf{m} z_i^{-1} \quad (1 \leq i \leq 2g) \rangle, \quad (1)$$

of $\pi_1(S^3 \setminus L)$. Notice that

$$\frac{\partial r_i^F}{\partial x_j} = \mathbf{m} \frac{\partial y_i}{\partial x_j} - r_i^F \frac{\partial z_i}{\partial x_j},$$

and consider the square matrix of the form

$$A_{F,W} := \left\{ \rho\left(\frac{\partial r_i^F}{\partial x_j}\right) \right\}_{1 \leq i,j \leq 2g} = \left\{ \tau \rho\left(\frac{\partial y_j}{\partial x_i}\right) - \rho\left(\frac{\partial z_j}{\partial x_i}\right) \right\}_{1 \leq i,j \leq 2g} \in \text{Mat}(2g \times 2g, \mathcal{A}_\kappa((\tau))). \quad (2)$$

Then, the paper [Nos, §3] defined a quotient group

$$\mathcal{Q}_{\mathcal{A},\kappa} := K_1(\mathcal{A}_\kappa((\tau)))/K_1(\mathcal{A}), \quad (3)$$

and if $A_{F,W}$ is an invertible matrix, we define the K_1 -Alexander invariant (with respect to ρ) to be the K_1 -class

$$\Delta_\rho^{K_1} := [\tau^{-g} A_{F,W}] \in \mathcal{Q}_{\mathcal{A},\kappa}.$$

On the other hand, if $A_{F,W}$ is not invertible, we define $\Delta_\rho^{K_1}$ to be zero. The main theorem in [Nos] shows that this $\Delta_\rho^{K_1}$ does not depend on the choice of F, W , and X .

Then, the theorem of this section is as follows:

Theorem 3.1. *Let G be a knot group with a Wirtinger presentation, as in Example 2.1, and let $\rho : \mathbb{Z}[\pi_1(S^3 \setminus L)] \rightarrow \mathcal{A}_\kappa((\tau))$ satisfy (\dagger) .*

Then, $\Phi_{\rho,k}$ is invertible if and only if so is $A_{F,W}$. In addition, if so, the K_1 -class $[\tau^{-g} A_{F,W}]$ is equal to $[\Phi_{\rho,k}]$ in the quotient group $\mathcal{Q}_{\mathcal{A},\kappa}/\{\tau^n\}_{n \in \mathbb{Z}}$.

4 Review of the Reidemeister torsion in a K_1 -group

For the proof of Theorem 3.1, the Reidemeister torsion appearing in a K_1 -group plays a key rule. We will review the Reidemeister torsion. The definition and properties are based on [Mil1] or [Tur, §I.3].

Consider an exact sequence of length 3

$$C_* : 0 \rightarrow C_3 \xrightarrow{\partial_3} C_2 \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \rightarrow 0,$$

where C_* is a finitely generated free R -module (C_3 may be zero). Let us choose a basis $\mathcal{C}_i \subset C_i$ for all i with $C_i \neq 0$. Assume that $B_i = \text{Im}(\partial_{i+1}) \subset C_i$ is free, pick a basis \mathcal{B}_i of B_i and a lift $\tilde{\mathcal{B}}_i$ of \mathcal{B}_i to C_i . By $\mathcal{B}_i \tilde{\mathcal{B}}_{i-1}$ we mean the collection of elements given by \mathcal{B}_i and $\tilde{\mathcal{B}}_{i-1}$. Since C_* is exact, $\mathcal{B}_i \tilde{\mathcal{B}}_{i-1}$ is indeed a basis for C_i . For bases d, e of a complex, denote by $[d/e]$ the invertible matrix of a basis change, i.e., $[d/e] = (a_{ij})$ where $d_i = \sum_j a_{ij} e_j$. Then we define the Reidemeister torsion of the based acyclic complex (C_*, \mathcal{C}_i) to be

$$\mathcal{T}(C_*, \mathcal{C}_i) := [\tilde{\mathcal{B}}_2/\mathcal{C}_3][\mathcal{B}_2 \tilde{\mathcal{B}}_1/\mathcal{C}_2]^{-1}[\mathcal{B}_1 \tilde{\mathcal{B}}_0/\mathcal{C}_1][\mathcal{B}_0/\mathcal{C}_0]^{-1} \in K_1(R).$$

Meanwhile, in the case where the R -modules B_i are not free, Section 3 in [Tur] gives a definition of the torsion of C_* . Since this paper does not consider such a case in details, we omit the details.

We will study homology groups in local coefficients. Let X be a connected CW complex. Denote the universal covering space of X by \tilde{X} . We regard the chain complex of space, $C_*(\tilde{X})$, as a chain complex of right $\mathbb{Z}[\pi_1(X)]$ -modules, where the $\mathbb{Z}[\pi_1(X)]$ -module structure is defined via covering transformations. Given a ring homomorphism $\rho : \mathbb{Z}[\pi_1(X)] \rightarrow R$, we can therefore consider the chain complex $C_*(X; R) = C_*(\tilde{X}) \otimes_{\mathbb{Z}[\pi_1(X)]} R$. We denote its homology by $H_*(X; R)$.

We now suppose that X is of finite type and $\dim X \leq 3$. Then, if the homology $H_i(X; R)$ is not zero for some i , we write $\mathcal{T}(X, \rho) = 0$. Otherwise, denote the i -cells of X by $\sigma_i^1, \dots, \sigma_i^{r_i}$, and choose an orientation for each cell σ_i^j , and also pick a lift $\tilde{\sigma}_i^j$ for each cell σ_i^j to the universal cover \tilde{X} . Since the set $\{\tilde{\sigma}_i^1, \dots, \tilde{\sigma}_i^{r_i}\}$ makes a basis \mathcal{C}_i of $C_i(X; R)$, we can define

$$\mathcal{T}(C_i(X; R), \{\mathcal{C}_i\}) \in K_1(R). \quad (4)$$

Moreover, consider the quotient of $K_1(R)$ passage by the image $\{\pm \rho(g)\}_{g \in \pi_1(X)}$. Then, as is known, the class

$$\mathcal{T}(X, \rho) := [\mathcal{T}(C_i(X; R), \{\mathcal{C}_i\})] \in K_1(R) / \pm \rho(\pi_1(X)) \quad (5)$$

depends only on the simple homotopy type of X and the homomorphism $\rho : \pi_1(X) \rightarrow R$.

Next, we give a procedure for computing $\mathcal{T}(X, \rho)$, following [Tur, p. 8]. Take subsets, $\xi_i \subset \{1, 2, \dots, \text{rank}(C_i)\}$ so that $\xi_0 = \emptyset$, and denote $(\xi_0, \xi_1, \dots, \xi_m)$ by ξ . For i, j, k , define the matrix $\{a_{jk}^i\}_{j,k}$ by considering only the ξ_i -columns of A_i and with the ξ_{i-1} -rows removed. Such a matrix chain ξ is called a \mathcal{T} -chain if $A_1(\xi), \dots, A_m(\xi)$ are square matrices. The following is the generalization of Turaev's Theorem 2.2 to the noncommutative setting, and is stated in [Fri1, Theorem 2.1].

Proposition 4.1 (cf [Tur, Theorem 2.2]. See also [Fri1, Theorem 2.1]). *Let ξ be a \mathcal{T} -chain such that $A_i(\xi)$ is invertible for all odd i . Suppose that every B_i is free. Then, $A_i(\xi)$ is invertible for all even i if and only if $H_*(C_*(X; R)) = 0$. If $H_*(C_*(X; R)) = 0$, then*

$$\mathcal{T}(C_i(X; R), \{\mathcal{C}_i\}) = \varepsilon \prod_{i=0}^m A_i(\xi)^{(-1)^i} \in K_1(R) \quad \text{for some } \varepsilon \in \pm 1.$$

4.1 Proof of Theorem 3.1

Proof of Theorem 3.1. We first recall the facts on invariance of Reidemeister torsion. Let Y be a finite connected finite CW complex. According to [Tur, Theorem 9.1], if there is a cellular map $f : Y \rightarrow X$ which is homotopy equivalent, and $H_*(X; R) = 0$ and the Whitehead group of $\pi_1(Y)$ is zero, then $H_*(Y; R) = 0$ and the associated torsion $\mathcal{T}(Y, \rho')$ is equal to $\mathcal{T}(X, \rho)$ in $K_1(R)/\pm \rho(\pi_1(X))$. Here, $\rho' = \rho \circ f_*$. Furthermore, if X is homotopic to the knot complement $S^3 \setminus L$, then the Whitehead group of $\pi_1(X)$ vanishes by Waldhausen [Wal]. Therefore, it is enough for the proof to find X and Y whose homotopy type are $S^3 \setminus L$ such that $\mathcal{T}(X, \rho) \cdot (1 - \tau)$ equals $\Phi_{\rho, m}$ and $\mathcal{T}(Y, \rho)(1 - \tau)$ equals $[A_{F, W}]$

First, we take the CW-complex X corresponding with the Wirtinger presentation. Namely, X consists of a single vertex, m edges labeled by the generators x_1, \dots, x_m and $(m - 1)$ 2-cells attached by the relations r_1, \dots, r_{m-1} . As is known, we can easily verify that W is homotopic to $S^3 \setminus L$. We regards these x_1, \dots, x_m and r_1, \dots, r_{m-1} as cells of X . The cellular complex is written in

$$0 \rightarrow \bigoplus_{j=1}^{m-1} \mathcal{A}_\kappa((\tau))r_j \xrightarrow{\partial_2} \bigoplus_{i=1}^m \mathcal{A}_\kappa((\tau))x_i \xrightarrow{\partial_1} \mathcal{A}_\kappa((\tau)) \rightarrow 0, \quad (6)$$

as left $\mathcal{A}_\kappa((\tau))$ -modules where the boundary maps are given by

$$\partial_2(r_j) = \sum_i \rho\left(\frac{\partial r_j}{\partial x_i}\right)x_i, \quad \text{and} \quad \partial_1(x_i) = 1 - \rho(x_i).$$

Notice that the restriction of ∂_1 on $\mathcal{A}_\kappa((\tau))x_m$ is $1 - \rho(x_m) = 1 - w_m^{-1}\tau w_m$; hence, it is invertible. Therefore, the acyclicity of (6) is equivalent to the invertibility of $\Phi_{\rho, k}$. If we set $\xi_1 := \{1\}$ and $\xi_2 := \{1, \dots, m\}$ as in Proposition 4.1, the Reidemeister torsion $\mathcal{T}(X, \rho)$ is given by $\Phi_{\rho, m} \cdot (1 - \tau)^{-1}$, as required.

Next, we let Y be the CW complex corresponding with the presentation (1) in a similar way. Then, it is shown [Tro, Section 2.3] that this Y is known to be homeomorphic to $S^3 \setminus L$, and the chain complex is given by

$$0 \rightarrow \bigoplus_{j=1}^{2g} \mathcal{A}_\kappa((\tau))r_j^F \xrightarrow{\partial_2} \bigoplus_{i=1}^{2g} \mathcal{A}_\kappa((\tau))x_i \oplus \mathcal{A}_\kappa((\tau))\mathbf{m} \xrightarrow{\partial_1} \mathcal{A}_\kappa((\tau)) \rightarrow 0,$$

where the boundary maps are given by

$$\partial_2(r_j) = (1 - \rho(y_j))\mathbf{m} + \sum_i \rho\left(\frac{\partial r_j^F}{\partial x_i}\right)x_i, \quad \text{and} \quad \partial_1(x_i) = 1 - \rho(x_i), \quad \partial_1(\mathbf{m}) = 1 - \tau.$$

Notice that the restriction of ∂ on $\mathcal{A}_\kappa((\tau))\mathfrak{m}$ is $1 - \tau$ invertible. Therefore, similarly, Proposition 4.1 implies that the Reidemeister torsion $\mathcal{T}(Y, \rho)$ is $A_{F,W} \cdot (1 - \tau)^{-1}$ by definition, as required. This completes the proof. \square

Proof of Proposition 2.4. It is proven in the paper [Nos] that, K is fibered, if and only if the matrix $A_{F,W}$ is shown to be invertible for any ρ . Thus, Proposition 2.4 immediately deduces the proof. \square

5 Relation to the torsions of cyclic coverings

Throughout this section, we assume the existence of $m \in \mathbb{N}$ such that $\kappa^m = \text{id}_{\mathcal{A}}$. In this case, we suggest a relation to the torsions of the regular m -fold cyclic covering spaces E_L^m . Here the m -fold cyclic covering $p : E_L^m \rightarrow S^3 \setminus L$ is associated with the surjection $\pi_1(S^3 \setminus L) \rightarrow \mathbb{Z}/m$.

For the purpose, we begin reviewing a ring homomorphism in [Nos]. For $\ell \leq m$ and $a \in \mathcal{A}$, we let $\mathcal{D}_\ell(a)$ be a diagonal $(\ell \times \ell)$ -matrix of the form

$$\mathcal{D}_\ell(a) := \begin{pmatrix} \kappa^\ell(a) & 0 & \cdots & 0 \\ 0 & \kappa^{\ell-1}(a) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \kappa^1(a) \end{pmatrix},$$

and let $\mathbb{O}_{s,t}$ be the zero $(s \times t)$ -matrix. We define a square matrix

$$M_\ell(a) := \begin{pmatrix} \mathbb{O}_{m-\ell,\ell} & \mathcal{D}_{m-\ell}(\kappa^{\ell+1}(a)) \\ \mathcal{D}_\ell(a) & \mathbb{O}_{\ell,m-\ell} \end{pmatrix} \in \text{Mat}(m \times m; \mathcal{A}). \quad (7)$$

Consider the Laurent polynomial ring $\mathcal{A}_{\text{id}}((t))$, where t is a commutative indeterminate. Then, the paper [Nos] introduced a ring homomorphism Υ defined by setting

$$\Upsilon : \mathcal{A}_\kappa((\tau)) \longrightarrow \text{Mat}(m \times m; \mathcal{A}_{\text{id}}((t))); \quad \sum_{j=k} a_j \tau^j \longmapsto \sum_{j=k} M_j(a_j) t^j. \quad (8)$$

Hence, we can consider the pushforwards of the K_1 -classes: $\Upsilon_*(\mathcal{T}(S^3 \setminus L, \rho))$ and $\Upsilon_*(A_{F,W})$. Here, by the Morita invariance on K_1

$$\mathcal{I} : K_1(\text{Mat}(m \times m; \mathcal{A}_{\text{id}}((t)))) \cong K_1(\mathcal{A}_{\text{id}}((t))),$$

the K_1 -classes are regarded as quotient elements of $K_1(\mathcal{A}_{\text{id}}((t)))$.

Meanwhile, notice that the image of the composite $\rho \circ p_* : \mathbb{Z}[\pi_1(E_L^m)] \rightarrow \mathcal{A}_\kappa((\tau))$ is contained in $\mathcal{A}_\kappa((\tau^m)) = \mathcal{A}_{\text{id}}((t))$. Thus, we can define the torsion $\mathcal{T}(E_L^m, \rho \circ p_*)$ in $K_1(\mathcal{A}_{\text{id}}((t)))$.

Theorem 5.1. *Let L be a knot. Under the above situation, the torsion of E_L^m is equal to the pushforward of the torsion of $S^3 \setminus L$, that is,*

$$\mathcal{I} \circ \Upsilon_*(\mathcal{T}(S^3 \setminus L, \rho)) = \mathcal{T}(E_L^m, \rho \circ p_*) \in K_1(\mathcal{A}_{\text{id}}((t))) / \pm \rho \circ p_*(\pi_1(E_L^m)).$$

Proof. We first analyze $\pi_1(E_L^m)$. Recall from (1) the presentation

$$\pi_1(S^3 \setminus L) \cong \langle x_1, \dots, x_{2g}, \mathfrak{m} \mid r_i^F := \mathfrak{m}^{-1} y_i \mathfrak{m} z_i^{-1} \quad (1 \leq i \leq 2g) \rangle.$$

For $k \in \mathbb{Z}/m$, let $x_i^{(k)}$ be a copy of x_i , and $y_i^{(k)}$ be the word obtained by replacing x_i by $x_i^{(k)}$ in the word y_i . We similarly define the word $z_i^{(k)}$. Then, by a Reidemeister-Schreier method (see, e.g., [LS, Kab]), $\pi_1(E_L^m)$ is presented by

$$\langle x_1^{(k)}, \dots, x_{2g}^{(k)}, \bar{\mathbf{m}} \mid r_i^{(k)} := \bar{\mathbf{m}}^{-1} y_i^{(k)} \bar{\mathbf{m}} (z_i^{(k+1)})^{-1} \quad (1 \leq i \leq 2g, k \in \mathbb{Z}/m) \rangle,$$

and the injection $p_* : \pi_1(E_L^m) \rightarrow \pi_1(S^3 \setminus L)$ is represented by the correspondence $x_i^{(k)} \mapsto x_i$, $\bar{\mathbf{m}} \mapsto \mathbf{m}^m$.

Then, similarly to (6), we have the cellular chain complex of $C_*(E_L^m; \mathcal{A}_{\text{id}}((t)))$ as

$$0 \rightarrow \bigoplus_{j=1}^m \bigoplus_{k=1}^{2g} \mathcal{A}_{\text{id}}((t)) r_j^{(k)} \xrightarrow{\partial_2} \mathcal{A}_{\text{id}}((t)) \bar{\mathbf{m}} \oplus \bigoplus_{i=1}^m \bigoplus_{k=1}^{2g} \mathcal{A}_{\text{id}}((t)) x_i^{(k)} \xrightarrow{\partial_1} \mathcal{A}_{\text{id}}((t)) \rightarrow 0,$$

where the boundary maps are given by

$$\partial_2(r_j^{(k)}) = (1 - \rho(y_j^{(k)})) \bar{\mathbf{m}} + \sum_{i,k'} \rho\left(\frac{\partial r_j^{(k)}}{\partial x_i^{(k')}}\right) x_i^{(k')}, \quad \text{and} \quad \partial_1(x_i^{(k)}) = 1 - \rho(x_i^{(k)}), \quad \partial_1(\bar{\mathbf{m}}) = 1 - t.$$

Let \mathcal{J} be the $(2gm \times 2gm)$ -matrix $\{\rho(\partial r_i^{(k)} / \partial x_j^{(k')})\}_{i,j \leq g, k,k' \leq m}$. Then, similarly to the proof of Theorem 3.1, Proposition 4.1 implies

$$\mathcal{T}(E_L^m, \rho \circ p_*) = (1 - t^m)^{-1} \cdot \mathcal{J} \in K_1(\mathcal{A}_{\text{id}}((t))) / \pm \rho \circ p(\pi_1(E_L^m)). \quad (9)$$

On the other hand, by the definition of Υ , we can easily check

$$\Upsilon(\rho(\frac{\partial r_i^F}{\partial x_j})) = \left\{ \rho(\frac{\partial r_i^{(s)}}{\partial x_j^{(t)}}) \right\}_{1 \leq s, t \leq m} \in \text{Mat}(m \times m; \mathcal{A}_{\text{id}}((t))),$$

for any $i, j \leq m$. Therefore, as $(2gm \times 2gm)$ -matrices, $\Upsilon(A_{F,W}) = \mathcal{J}$, where $A_{F,W}$ is the matrix in (2). Notice that $\Upsilon(1 - \tau) = 1 - t^m$ in $K_1(\mathcal{A}_{\text{id}}((t)))$, and recall $\mathcal{T}(S^3 \setminus L, \rho) = (1 - \tau)^{-1} A_{F,W}$ by the proof of Theorem 3.1. Hence, combining those with (9) deduces the required equality. \square

6 Reciprocity of some K_1 -Alexander invariants

As is well-known, the (classical) Alexander polynomial of a knot has symmetry, i.e., it can be expanded as $\sum_{i=-m}^m a_i t^i$ such that $a_i \in \mathbb{Z}$ and $a_{-i} = a_i$. Such a symmetry is called *Reciprocity*. As a generalization, if we choose an appropriate representation, the twisted Alexander polynomial also has reciprocity in some sense; see, e.g., [FKK, FV, Ki1, Kitan]. Moreover, reciprocity of some Reidemeister torsions is generalized for K_1 -groups; see [Mil1, §10] or [Tur, Theorem 14.1]. Thus, it is reasonable to ask reciprocity on the K_1 -classes $\Delta_\rho^{K_1}$. Unfortunately, some papers on reciprocity require some conditions to show duality theorems. Similarly, the author could not show directly reciprocity for every $\Delta_\rho^{K_1}$; however, this paper will observe (Theorem 6.1) a reciprocity of the pushforward by the ring homomorphism Υ_* , under a certain situation. Here the point is to find a situation applicable to the theorem of Turaev.

The situation in this section is described as follows:

(\star) As in Example 2.1, we consider a group homomorphism $h : \pi_1(S^3 \setminus L) \rightarrow H \rtimes \mathbb{Z}$ such that H is a group of finite order, and $h(x_i) = (g_i, 1)$ for some $g_i \in H$. Let \mathcal{A} be the group ring $\mathbb{Q}[H]$ over \mathbb{Q} . Let $m \in \mathbb{Z}_{\geq 0}$ be the minimal such that $\kappa^m = \text{id}_{\mathcal{A}}$.

Starting from the situation (\star), we discuss some involutions and observe some K_1 . In this section, let \mathcal{A} be $\mathbb{Q}[H]$, and consider the (skew) Laurent polynomial ring $\mathcal{A}_\kappa[\tau^{\pm 1}]$, instead of Novikov rings. Since $\mathcal{A} = \mathbb{Q}[H]$ is semi-simple, the Wedderburn theorem immediately implies that there are division rings D_1, \dots, D_m over \mathbb{Q} and integers n_1, \dots, n_k which ensure the ring isomorphism

$$\mathbb{Q}[H] \cong \bigoplus_{i:1 \leq i \leq k} \text{Mat}(n_i \times n_i; D_i).$$

For a division ring D , let $D(t)$ be the fractional field of $D[t^{\pm 1}]$ with $\bar{t} = t^{-1}$ and $\iota_D : D[t^{\pm 1}] \rightarrow D(t)$ be the inclusion. Then, the direct sum $\oplus \iota_{D_i}$ gives rise to

$$(\oplus \iota_{D_i})_* : \text{Mat}(m \times m; \mathcal{A}_{\text{id}}[t^{\pm 1}]) \longrightarrow \bigoplus_{i:1 \leq i \leq k} \text{Mat}(n_i m \times n_i m; D_i(t)).$$

Formally, we will denote by $\mathbb{Q}[H](t)$ by the direct sum $\oplus_{i=1}^k D_i(t)$. Then, Morita invariance on K_1 -groups again implies the isomorphisms

$$K_1\left(\bigoplus_{i:1 \leq i \leq k} \text{Mat}(n_i m \times n_i m; D_i(t))\right) \cong \bigoplus_{i:1 \leq i \leq k} K_1(D_i(t)) \cong K_1(\mathbb{Q}[H](t)).$$

By the assumption (\star), $A_{F,W}$ in (2) is regarded as a matrix over the (uncompleted) Laurent polynomial ring $\mathbb{Q}[H]_\kappa[\tau^{\pm 1}]$. In the same way as (8), consider the ring homomorphism Υ defined by setting

$$\Upsilon : \mathcal{A}_\kappa[\tau^{\pm 1}] \longrightarrow \text{Mat}(m \times m; \mathcal{A}_{\text{id}}(t^{\pm 1})); \quad \sum_{i=k} a_i \tau^i \longmapsto \sum_{i=k} M_i(a) t^i,$$

where $M_j(a_j)$ is defined in (14). Then, the pushforward $(\oplus \iota_{D_i})_* \circ \Upsilon_*(A_{F,W})$ can be regarded over $\mathbb{Q}[H](t)$. Thus, $\Upsilon_*(A_{F,W})$ can be considered to be an element of the $K_1(\mathbb{Q}[H](t))$, where we omit writing $(\oplus \iota_{D_i})_*$.

Theorem 6.1. *Let $\mathfrak{l} \in \pi_1(S^3 \setminus L)$ be the preferred longitude of the knot L . Suppose the above situation (\star), and invertibility of the pushforward $\Upsilon_*(A_{F,W})$ over $\mathbb{Q}[H](t)$. In addition, we assume that either (a) $\Upsilon(1 - \rho(\mathfrak{l}))$ is an invertible matrix or (b) $\rho(\mathfrak{l}) = 1 \in \mathbb{Q}[H]$.*

Then, under the isomorphism $(\oplus \iota_{D_i})_$, the following equality holds:*

$$\overline{\Upsilon_*(A_{F,W})} = \Upsilon_*(A_{F,W}) \in \frac{K_1(\mathbb{Q}[H](t))}{\{K_1(\mathbb{Q}[H]), t^\ell\}_{\ell \in \mathbb{Z}}}.$$

Remark 6.2. As a result, we can observe the reciprocity on the K_1 -class $\Upsilon_*(\Delta_\rho^{K_1})$ in the K -group of a Novikov ring $\mathcal{Q}_{\mathbb{Q}[H](t), \text{id}}$ as well. Indeed, since $D_i(t)$ and $D_i((t))$ are division rings, the associated inclusions $j_{D_i} : D_i(t) \rightarrow D_i((t))$ induce $\iota : \mathbb{Q}[H](t) \rightarrow \mathbb{Q}[H]((t))$; thus, the reciprocity inherits on $\iota_*(\Upsilon_*(A_{F,W}))$ in $\mathcal{Q}_{\mathbb{Q}[H](t), \text{id}}$.

Furthermore, since the Dieudonné determinant over any division ring D induces an isomorphism $K_1(D) \cong (D^\times)_{\text{ab}}$, the K_1 -group $K_1(\mathbb{Q}[H](t))$ is identified with $(\mathbb{Q}[H](t)^\times)_{\text{ab}}$. In particular, if H is abelian, we can interpret Theorem 6.1 as the reciprocity of a polynomial.

Proof of Theorem 6.1. Let E_L^m be the cyclic covering space as above. For simplicity, let Υ_ρ denote the composite $\Upsilon \circ \rho$. By virtue of Theorem 5.1, we may show only the reciprocity in Reidemeister torsions of E_L^m .

We first give the proof in the case (a). Then, [Tur, Corollary 14.2] immediately says

$$\mathcal{T}(E_L^m, \partial E_L^m; \Upsilon_\rho) = \overline{\mathcal{T}(E_L^m; \Upsilon_\rho)} \in K_1(\mathbb{Q}[H](t)) / \{K_1(\mathbb{Q}[H]), \tau^\ell\}_{\ell \in \mathbb{Z}}. \quad (10)$$

Moreover, thanks to the multiplicity of torsions with respect to short exact sequences (see [Tur, Theorem 3.4]), we immediately have

$$\mathcal{T}(E_L^m, \partial E_L^m; \Upsilon_\rho) \mathcal{T}(\partial E_L^m; \Upsilon_\rho) = \mathcal{T}(E_L^m; \Upsilon_\rho) \in K_1(\mathbb{Q}[H](t)) / \{K_1(\mathbb{Q}[H]), t^\ell\}_{\ell \in \mathbb{Z}}. \quad (11)$$

By Lemma 6.3 below, the second term $\mathcal{T}(\partial E_L^m; \Upsilon_\rho)$ is equal to 1. Hence, the equalities (10) and (11) readily mean the required equality.

Next, we discuss the case (b). Let M_0 be the closed 3-manifold obtained by 0-surgery of E_L^m along L . Notice $\partial M_0 = \emptyset$. Let $p : E_L^m \rightarrow S^3 \setminus L$ be the covering. Since $\rho(\mathfrak{l}) = 1$, the composite $\rho \circ p_*$ induces $\mathbb{Z}[\pi_1(M_0)] \rightarrow \mathcal{A}_\kappa((\tau))$. Therefore, by [Tur, Corollary 14.2] again, we readily have the reciprocity $\mathcal{T}(M_0; \Upsilon_{\rho \circ p_*}) = \overline{\mathcal{T}(M_0; \Upsilon_{\rho \circ p_*})}$. Hence, it is enough for the proof to show $(1-t)\mathcal{T}(M_0; \Upsilon_{\rho \circ p_*}) = \mathcal{T}(E_L^m; \Upsilon_\rho)$.

To show this, consider the exact sequence appearing in a Mayer-Vietoris argument

$$0 \longrightarrow C_*(S^1 \times S^1) \xrightarrow{\iota_1 \oplus \iota_2} C_*(D^2 \times S^1) \oplus C_*(E_L^m) \longrightarrow C_*(M_0) \longrightarrow 0 \quad (\text{exact}).$$

Since $1 - \Upsilon_\rho(\mathfrak{l}) = 0$, the boundary maps in the second $C_*(D^2 \times S^1)$ are zero, and ι_1 has a splitting as a chain map. Hence, the sequence is reduced to

$$0 \longrightarrow C_*(S^1) \xrightarrow{\iota_2} C_*(E_L^m) \longrightarrow C_*(M_0) \longrightarrow 0 \quad (\text{exact}).$$

Since the first and second complexes are acyclic, so is the third. Notice $\mathcal{T}(C_*(S^1)) = (1-t)$ by definition. Hence, by [Tur, Theorem 3.4] again, we obtain $(1-t)\mathcal{T}(M_0; \Upsilon_{\rho \circ p_*}) = \mathcal{T}(E_L^m; \Upsilon_\rho)$ as required. \square

Lemma 6.3. *If $1 - \Upsilon_\rho(\mathfrak{l}) \in \mathbb{Q}[H]$ is invertible, then the torsion $\mathcal{T}(\partial E_L^m; \Upsilon_\rho)$ is 1.*

Proof. Since ∂E_L^m is torus, we can describe the cellular complex as

$$0 \longrightarrow \mathbb{Q}[H](t)c_2 \xrightarrow{(1-\Upsilon_\rho(\overline{\mathfrak{m}}), 1-\Upsilon_\rho(\mathfrak{l}))} \mathbb{Q}[H](t)c_1 \oplus \mathbb{Q}[H](t)c'_1 \xrightarrow{\begin{pmatrix} \Upsilon_\rho(\mathfrak{l}) - 1 \\ 1 - \Upsilon_\rho(\overline{\mathfrak{m}}) \end{pmatrix}} \mathbb{Q}[H](t)c_0 \longrightarrow 0,$$

which is acyclic by assumption. If we choose

$$b_2 = \{c_2\}, \quad b_1 = \{c_1\}, \quad b_0 = \{c_0\},$$

then

$$[b_2/c_2] = 1, \quad [b_1\partial_2 b_2/c_1] = \begin{pmatrix} 1 - \Upsilon_\rho(\mathfrak{l}) & 1 \\ 1 - \Upsilon_\rho(\overline{\mathfrak{m}}) & 0 \end{pmatrix}, \quad [b_0\partial_1 b_1/c_0] = 1 - \Upsilon_\rho(\overline{\mathfrak{m}}).$$

Hence, by the definition of \mathcal{T} , we get the conclusion. \square

7 Applications to m -fold metabelian Alexander polynomials.

The paper [Nos] introduces an (m -fold) metabelian Alexander polynomial. As applications of the previous sections, we will see properties of the polynomial.

Let us recall the polynomial. Let L be a knot, and H be the torsion subgroup $\text{Tor}(H_1(\tilde{E}_L^m; \mathbb{Z}))$. Suppose that $mH = H$, e.g., the case m is a prime power. Then, the commutator subgroup of $\pi_1(S^3 \setminus L)$ surjects on H via an abelianization. Thus, we canonically have a group epimorphism $\rho_m^{\text{meta}} : \pi_1(S^3 \setminus L) \rightarrow H \rtimes \mathbb{Z}$ satisfying the assumption (\dagger) . Notice $\mathcal{A} = \mathbb{Q}[H]$ is isomorphic to a direct sum of cyclotomic fields over \mathbb{Q} ; see, e.g., [Tur, Corollary 12.10]. Then, the metabelian Alexander polynomial is defined to be the determinant of the pushforward $\Upsilon_*(A_{F,W})$:

$$\Delta_{\rho_m^{\text{meta}}} := \det(\Upsilon_*(A_{F,W})) \in \mathbb{Q}[H](t)^\times / (\mathbb{Q}[H]^\times).$$

As seen in Appendix A, this polynomial can be also interpreted as a torsion obtained from a S^1 -valued Morse theory.

As a slight generalization of Lemma 4 of [CG], we will show the invertibility of $\Upsilon_*(A_{F,W})$.

Proposition 7.1 (cf. [CG]). *The matrix $\Upsilon_*(A_{F,W})$ in $\text{Mat}(2gm \times 2gm; \mathbb{Q}[H](t))$ is invertible.*

Proof. Let $S \subset H = \text{Tor}H_1(E_L^m; \mathbb{Z})$ be any direct summand isomorphic to \mathbb{Z}/p^n for some $n \in \mathbb{N}$ and prime p . For the proof, we may replace H by S , since $GL_n(\mathbb{Q}[H_1 \oplus H_2]) = GL_n(\mathbb{Q}[H_1]) \times GL_n(\mathbb{Q}[H_2])$ for any finite abelian groups H_i .

Let \tilde{E}_L be the infinite cyclic covering space of $S^3 \setminus L$. Let $\tilde{M} \rightarrow \tilde{E}_L$ be the abelian finite covering associated with the projection $\pi_1(\tilde{E}_L) \xrightarrow{\text{proj}} \text{Tor}H_1(E_L^m; \mathbb{Z}) \xrightarrow{\text{proj}} S$. According to Lemma 4 and its corollary of [CG], the rational homology $H_*(\tilde{M}; \mathbb{Q})$ is shown to be zero. Moreover, by Shapiro lemma (see, e.g., [KL]), the cellular complex $C_*(\tilde{M}; \mathbb{Q})$ is isomorphic to $C_*(E_L^m; \mathbb{Q}[S][t^{\pm 1}])$ of finite dimension. Thus, the tensored complex $C_*(E_L^m; \mathbb{Q}[S](t)) = C_*(E_L^m; \mathbb{Q}[S][t^{\pm 1}]) \otimes \mathbb{Q}[S](t)$ over $\mathbb{Q}[S](t)$ is acyclic. In usual, the first boundary ∂_1 is a splittable surjection; hence, the acyclicity of $C_*(E_L^m; \mathbb{Q}[S](t))$ implies the invertibility of ∂_2 . As seen in the proof of Theorem 3.1, ∂_2 is represented by the matrix $\Upsilon_*(A_{F,W})$. Hence, we complete the proof. \square

As a corollary, we will see the reciprocity of $\Delta_{\rho_m^{\text{meta}}}$. Here, notice that $\rho(\mathfrak{l}) = 0_H$ since the preferred longitude \mathfrak{l} is bounded by a Seifert surface in \tilde{E}_L . Therefore, our situation fulfills the conditions in Theorem 6.1; hence, we immediately have

Corollary 7.2. *The metabelian Alexander polynomial has reciprocity in the sense of*

$$\overline{\Delta_{\rho_m^{\text{meta}}}} = \Delta_{\rho_m^{\text{meta}}} \in \mathbb{Q}[H](t)^\times / \{\mathbb{Q}[H]^\times, t^{\pm 1}\}.$$

This situation is similar to the Casson-Gordon invariant [CG], which discusses obstructions of sliceness. Thus it is reasonable to consider applications for sliceness from our results.

Theorem 7.3 (cf. [KL, Theorem 6.2]). *Suppose that the knot L is topological slice, and $H_1(E_L^m; \mathbb{Z})$ is isomorphic to $\mathbb{Z} \oplus T$ for some torsion module T .*

Then, there is a subgroup $B \subset H = \text{Tor}H_1(E_L^m; \mathbb{Z})$ such that $|B|^2 = |H|$ and that there exists a polynomial $f(t) \in \mathbb{Q}[H/B](t)$ satisfying

$$\mathcal{P}_*(\Delta_{\rho_m^{\text{meta}}}) = at^n(1-t)f(t)\overline{f(t)} \in \mathbb{Q}[H/B](t), \quad (12)$$

for some $a \in \mathbb{Q}[H/B]^\times$, $n \in \mathbb{Z}$, where $\mathcal{P} : H \rightarrow H/B$ is the projection.

Remark 7.4. We give a comparison with the works [KL, HKL]. The papers suppose m to be a prime power, and consider a homomorphism $\chi : H_1(B_L^m; \mathbb{Z}) \rightarrow \mathbb{Z}/p^d\mathbb{Z}$ for some prime p . Choose a homomorphism $\lambda : \mathbb{Z}/p^d\mathbb{Z} \rightarrow GL_1(\mathbb{C})$. Then, [KL, Theorem 6.2] claims that the further pushforward $(\lambda \circ \chi)_* \circ \mathcal{P}_*(\Delta_{\rho_m^{\text{meta}}})$ is decomposed as (12). Thus, Theorem 7.3 is a slight generalization of [KL, Theorem 6.2], although the proof below is outlined on discussions in [CG, HKL, KL].

7.1 Proof of Theorem 7.3

For the proof, we review Reidemeister torsions suitable to non-acyclic cases, in terms of determinants. Let \mathbb{F} be a commutative field of characteristic zero. Let X be a finite connected CW-complex X . Choose homomorphisms $\rho : \pi_1(X) \rightarrow \mathbb{F}^\times$ and $\alpha : \pi_1(X) \rightarrow \mathbb{Z} = \{t^n\}_{n \in \mathbb{Z}}$. We have the tensor representation $\rho \otimes \alpha : \pi_1(X) \rightarrow \mathbb{F}[t^{\pm 1}]^\times$. Then, the cellular complex with local coefficients is defined to be

$$C_*(X; \mathbb{F}(t)) = \mathbb{F}(t) \otimes_{\mathbb{F}[\mathbb{Z}]} C_*(X; \mathbb{F}[\mathbb{Z}]) = \mathbb{F}(t) \otimes_{\rho \otimes \alpha} C_*(\tilde{X}).$$

If C_* is a based chain complex, c_i is a basis for C_i , b_i a basis for the boundaries B_i , h_i a basis for the homology H_i , then the Reidemeister torsion \mathcal{T}' of the based chain complex is defined by

$$\mathcal{T}(X, \rho, h_*)' = \frac{\prod_i \det[b_{2i} \tilde{h}_{2i} \tilde{b}_{2i} / c_{2i}]}{\prod_i \det[b_{2i-1} \tilde{h}_{2i-1} \tilde{b}_{2i-1} / c_{2i-1}]} \in \mathbb{F}(t)^\times / \{t^n, a \in \mathbb{F}^\times\}.$$

In this expression, \tilde{h}_i is a choice of lift of h_i to C_i , and \tilde{b}_i is a choice of b_i to C_{i+1} using the differential $\partial_{i+1} : C_{i+1} \rightarrow C_i$. It is known this $\mathcal{T}(X, \rho, h_*)'$ is independent of the choices of b_i and of the lifts. Notice that, if C_* is acyclic and $\dim X \leq 3$, then this $\mathcal{T}(X, \rho, \emptyset)'$ is recovered by the torsion in §4: precisely, $\det \mathcal{T}(X, \rho \otimes \alpha) = \mathcal{T}(X, \rho, \emptyset)'$ by definitions.

In order to start the proof of Theorem 7.3, following [Mil1, Mil2], let us notice the following theorem (which is also stated in [KL, Theorem 5.1]).

Theorem 7.5 (see [KL, Theorem 5.1]). *Let X be an oriented connected compact C^∞ -manifold with boundary $\partial(X)$. Choose a triangulation of X as a CW-complex. Suppose that $C_*(X; \mathbb{F}(t))$ is not acyclic, but $C_*(\partial X; \mathbb{F}(t))$ is acyclic, and choose an appropriate basis h_q for $H_q(X; \mathbb{F}(t))$ for each q . Then, with respect to these bases,*

$$\mathcal{T}(\partial X, \rho)' = \mathcal{T}(X, \rho, h_*)' \overline{\mathcal{T}(X, \rho, h_*)'}^{(-1)^{\dim(X)}}.$$

Proof of Theorem 7.3. We give some preparations. Let $D \subset B^4$ be the slice disk of L in the 4-ball, and $\tilde{B} \rightarrow B^4 \setminus D$ be the m -fold covering. Let M_0 (resp. X) be the oriented closed manifold obtained by 0-surgery of E_L^m along L (resp. of \tilde{B} along D). Then, we notice $\partial X = M_0$, and $\text{Tor} H_1(M_0; \mathbb{Z}) \cong H = \text{Tor} H_1(\tilde{E}_L^m; \mathbb{Z})$. Let $B \subset H$ be the subgroup consisting of metabolizers of the linking form on M_0 . Then, it is known (see [CG] or [KL, §6]) that $|B|^2 = |H|$ and the projection $\rho_m^{\text{meta}} : \pi_1(M_0) \rightarrow H$ extends to $\pi_1(X) \rightarrow H/B$. Recall that the group ring $\mathbb{Q}[H/B]$ is ring isomorphic to $\oplus_i \mathbb{F}_i$, where \mathbb{F}_i is a cyclotomic field over \mathbb{Q} . Let $q_i : \mathbb{Q}[H/B] \rightarrow \mathbb{F}_i$ be the projection.

Notice that we can find $f_i \in \mathbb{F}_i(t)$ satisfying

$$q_i \circ \mathcal{T}(M_0, \rho)' = a_i t^{n_i} f_i(t) \overline{f_i(t)} \in \mathbb{F}_i(t), \quad (13)$$

for some $a_i \in \mathbb{F}_i^\times, n_i \in \mathbb{Z}$. Indeed, we may only let ρ_i be $q_i \circ \rho_m^{\text{meta}}$, and $f_i(t)$ be $\mathcal{T}(X, \rho_i, h_*)$ in Theorem 7.5. Moreover, as in Theorem 6.1, each $\mathcal{T}(M_0, \rho_i)'$ has reciprocity, we may suppose $n_1 = n_2 = \dots$. Furthermore, by the end of the proof of Theorem 6.1, we notice $q_i \circ \mathcal{P}_*(\Delta_{\rho_m^{\text{meta}}}) = \mathcal{T}(E_L^m, \rho_i)' = (1-t)\mathcal{T}(M_0, \rho_i)'$. To conclude, since $\mathbb{Q}[H/B](t) \cong \oplus_i \mathbb{F}_i(t)$, the multiplications of (13) running over i implies the required equality (12). \square

A Appendix; relation to S^1 -valued Morse theoretic torsions

Fix a preferred longitude $\mathfrak{l} \in \pi_1(S^3 \setminus L)$. Under an assumption $\rho(\mathfrak{l}) = 1$, the K_1 -class $\Phi_{\rho, k}$ can be described from S^1 -valued Morse theory; however, this section essentially contains nothing new. In fact, this sections are analogous to [P, Ki2]. Here, we suppose notation in §§2 and 4.

Let us roughly review the (twisted) S^1 -valued Morse theory, and the main result of [P]; see [P] for the details. Let M be a connected closed C^∞ -manifold, and $f : M \rightarrow S^1$ be a Morse map. Suppose that $f_* : H_1(M) \rightarrow H_1(S^1) = \mathbb{Z}$ is surjective. As in Example 2.1, let G be a semidirect product $H \rtimes \mathbb{Z}$ with projection $\chi : G \rightarrow \mathbb{Z}$, and \mathcal{A} be $\mathbb{Q}[H]$. Then, we have the Novikov ring $\mathcal{A}_\kappa((\tau))$.

We study a logarithm from $K_1(\mathcal{A}_\kappa((\tau)))$. For $n \in \mathbb{Z}$, let Γ_n be the set of conjugacy classes contained in $\chi^{-1}(n)$. Take the \mathbb{Q} -vectors space $\mathbb{Q}\Gamma_n$ spanned by Γ_n , and define \mathfrak{G} by $\prod_{n \geq 0} \mathbb{Q}\Gamma_n$. Furthermore, consider the multiplicative subgroup, W , of $\mathcal{A}_\kappa((\tau))^\times$ consisting of elements of the form $1 + a_1\tau + a_2\tau^2 + \dots$. The image of W in $K_1(\mathcal{A}_\kappa((\tau)))$ will be denoted by \widehat{W} . The logarithm $\log : W \rightarrow \mathfrak{G}$ is defined by

$$\log(1 + \mu\tau) = \mu\tau - \frac{(\mu\tau)^2}{2} + \dots + (-1)^{n-1} \frac{(\mu\tau)^n}{n} + \dots \quad \text{with } \mu \in \mathcal{A}_\kappa[[\tau]].$$

Then, it is shown [P, Lemma 1.1] that this map induces a homomorphism $\mathcal{L} : \widehat{W} \rightarrow \mathfrak{G}$. The subgroup \widehat{W} is known to be a direct summand of $K_1(\mathcal{A}_\kappa((\tau)))$ [PR]; hence, \mathcal{L} is regarded as a homomorphism from $K_1(\mathcal{A}_\kappa((\tau)))$.

Next, we will review non-abelian eta functions. For a vector field v , which satisfies a “Kupka-Smale condition”, let $\text{Cl}(v)$ be the closed orbits of v . For a closed orbit $\gamma \in \text{Cl}(v)$, let $\epsilon(\gamma) \in \{\pm 1\}$ denote the index of the corresponding Poincaré map; let $m(\gamma)$ denote the multiplicity of γ . Here is the definition of *the non-abelian eta function* of $-v$:

$$\eta_L(-v) = \sum_{\gamma \in \text{Cl}(-v)} \frac{\epsilon(\gamma)}{m(\gamma)} \{\gamma\} \in \mathfrak{G}.$$

Let denote $\mathcal{G}(f)$ be the set of vector fields satisfying the Kupka-Smale condition.

The main theorem of [P] clarified the difference between torsions from the usual complex and from the Novikov complex. To be precise, there is an open dense subset $\mathcal{G}_0(f) \subset \mathcal{G}(f)$ with respect to a certain C^0 -topology such that, for every $v \in \mathcal{G}_0(f)$, there is a chain homotopy equivalence

$$\phi : C_*(v) \longrightarrow C_*(\widetilde{M}; \mathbb{Z}) \otimes_{\mathbb{Z}[H]} \mathcal{A}_\kappa((\tau))$$

such that

$$\mathcal{L}(\tau(\phi)) = \eta_L(-v).$$

Here $\tau(\phi)$ is the torsion of the acyclic complex of the cokernel $\text{Coker}(\phi)$.

We will give a conclusion from the assumption $(*)$ and suppose $\rho(\mathfrak{l}) = 1$ as above (For instance, the setting in §7 is satisfied). Let $M_{L,0}$ be the closed 3-manifold obtained by 0-surgery of $S^3 \setminus L$ along L . Then, ρ induces a homomorphism $\pi_1(M_{L,0}) \rightarrow \mathcal{A}_\kappa((\tau))$. As in the proof of Theorem 6.1 implies the torsion $\mathcal{T}(M_{L,0}, \rho)$ is equal to $(1-t)^{-1}\mathcal{T}(S^3 \setminus L, \rho)$. Then, by the proof Theorem 3.1, we have $[\Phi_{\rho,k}] = (1-t)\mathcal{T}(M_{L,0}, \rho)$. To conclude, the torsions $\mathcal{L}([\Phi_{\rho,k}])$ can be described by the torsion defined from S^1 -valued Morse complex and the eta function.

B Relation to the higher order Alexander polynomial

The previous papers [C, Har] suggested a generalization of the classical Alexander polynomial; see also [GS, Fri2] for the studies. The point is that the generalized polynomial is defined from a group G , which is locally indicable and amenable, not any group. The purpose of this section is to show Proposition B.1 below, which suggests a relation between the generalized polynomial and the K_1 -class $\Delta_\rho^{K_1}$. The facts and explanations in this section are essentially based on [Fri2]; there is almost nothing new in this appendix. This appendix supposes terminology in §§2–4.

For this, we start by reviewing the situation in [C, Har]. Let $\text{Ab} : \pi_1(S^3 \setminus L) \rightarrow \mathbb{Z}$ be an abelianization. Let G be a locally indicable and amenable group. It is known that the group ring $\mathbb{Z}[G]$ embeds in a (skew) fractional field. Fix an epimorphism $\phi : \pi_1(S^3 \setminus L) \rightarrow G$ such that there exists a group homomorphism $\varphi_G : G \rightarrow \mathbb{Z}$ satisfying $\text{Ab} = \varphi_G \circ \phi$. Such a pair (φ_G, ϕ) is called *an admissible pair* for $\pi_1(S^3 \setminus L)$, following [Har, Definition 1.4].

Let G' be the kernel $\text{Ker}(\varphi_G : G \rightarrow \mathbb{Z})$. Since G' is also locally indicable and amenable, $\mathbb{Z}[G']$ embeds in a fractional field $\mathbb{K}^{G'}$. Let $\mu \in G$ be $\phi(\mathfrak{m})$, and define $\gamma : \mathbb{K}^{G'} \rightarrow \mathbb{K}^{G'}$ to be the homomorphism induced by $\gamma(g) = \mu g \mu^{-1}$. Then, we obtain the skew polynomial ring $\mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]$, and a ring homomorphism

$$\nu : \mathbb{Z}[G] \longrightarrow \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]; \quad \sum_g n_g g \longmapsto \sum_g n_g g \mu^{-\phi(g)} \tau^{\phi(g)}.$$

Let \mathcal{A} be the division ring $\mathbb{K}^{G'}$, and κ be γ . Then, the following composite satisfies Assumption (\dagger) :

$$\rho_G : \mathbb{Z}[\pi_1(S^3 \setminus L)] \xrightarrow{\phi} \mathbb{Z}[G] \xrightarrow{\nu} \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}] \hookrightarrow \mathbb{K}_\gamma^{G'}((\tau)) \quad (14)$$

Next, we review the higher order Alexander polynomial. The ring $\mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]$ is known to be a principal ideal domain since $\mathbb{K}_\gamma^{G'}$ is a skew field. For a finitely generated right $\mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]$ -module H , the elementary divisor theorem claims an isomorphism

$$H \cong \bigoplus_{i: 1 \leq i \leq \ell} \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}] / p_i(t) \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]$$

for some $\ell \in \mathbb{N}$ and $p_i(t) \in \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]$ for $i \in \{1, \dots, \ell\}$. Following [C, Fri2], we define $\text{ord}(H)$ by the product $p_1(t) \cdots p_\ell(t)$. Although it is a subject to discuss which set this $\text{ord}(H)$ should be contained in, according to [Fri2], we regard $\text{ord}(H)$ as an element in $\mathbb{K}_\gamma^{G'}(\tau)_{\text{ab}}^\times \cup \{0\}$ up to

multiplication by an element of the form $k\tau^e$ with $k \in \mathbb{K}_\gamma^{G'}$ and $e \in \mathbb{Z}$. As is known, $\text{ord}(H)$ depends on only H ; see [Fri2, Theorem 3.1]. Then, the order of the i -th homology

$$\Delta_i^\psi := \text{ord} H_i(S^3 \setminus L; \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]) \in \mathbb{K}_\gamma^{G'}(\tau)_{\text{ab}}^\times / \{k\tau^e\}_{k \in \mathbb{K}_\gamma^{G'}, e \in \mathbb{Z}}$$

is called *the higher order Alexander polynomial*.

To state Proposition B.1, let us consider the natural inclusion λ from the fractional field $\mathbb{K}_\gamma^{G'}(\tau)$ into $\mathbb{K}_\gamma^{G'}((\tau))$, since $\mathbb{K}_\gamma^{G'}((\tau))$ is also a field. Then, by the isomorphism $K_1(\mathbb{K}_\gamma^{G'}((\tau))^\times) \cong (\mathbb{K}_\gamma^{G'}((\tau))^\times)_{\text{ab}}$, the inclusion yields the homomorphism

$$\lambda_* : \mathbb{K}_\gamma^{G'}(\tau)_{\text{ab}}^\times / \{k\tau^e\}_{k \in \mathbb{K}_\gamma^{G'}, e \in \mathbb{Z}} \longrightarrow \mathcal{Q}_{\mathcal{A}, \gamma} / \{\tau^e\}_{e \in \mathbb{Z}}.$$

Then, the pushforward of Δ_1^ψ turns out to be almost the K_1 -class $\Delta_{\rho_G}^{K_1}$; More precisely,

Proposition B.1. *Suppose that the matrix $A_{F,W}$ is invertible. Then, Δ_i^ψ is zero if $i > 1$, and $\Delta_0^\psi = 1 - \tau$. Furthermore, the following equality holds:*

$$\lambda_*(\Delta_1^\psi) = \Delta_{\rho_G}^{K_1} \in \mathcal{Q}_{\mathcal{A}, \gamma} / \{\tau^e\}_{e \in \mathbb{Z}}.$$

Proof. From the cellular complex (6), we see that the i -th homology is vanishes for $i > 1$, i.e., $\Delta_i^\psi = 0$, and if $i = 0$, we find

$$H_0(S^3 \setminus L; \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}]) \cong \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}] / (1 - \tau) \mathbb{K}_\gamma^{G'}[\tau^{\pm 1}].$$

Thus, $\Delta_0^\psi = 1 - \tau$.

Let us consider the case $i = 1$. Then, the main theorem of [Fri2],

$$\mathcal{T}(S^3 \setminus L, \rho_G) = \Delta_1^\psi / \Delta_0^\psi \in \mathbb{K}_\gamma^{G'}(\tau)_{\text{ab}}^\times / \{k\tau^e\}_{k \in \mathbb{K}_\gamma^{G'}, e \in \mathbb{Z}},$$

where ρ_G is the composite in (14). In the proof of Theorem 2.3, $\lambda_*(\mathcal{T}(S^3 \setminus L, \rho_G))$ was shown to be $[A_{F,W}] \cdot (1 - \tau)^{-1}$. Hence, we have the required equality. \square

However, the point is that, in general, the fractional field $\mathbb{K}_\gamma^{G'}$ is quite incomprehensible from quantitative viewpoint; in particular, it is difficult to compute the abelianization $\mathbb{K}_\gamma^{G'}(\tau)_{\text{ab}}^\times$ and to distinguish whether two elements in $\mathbb{K}_\gamma^{G'}(\tau)_{\text{ab}}^\times$ are equal or not. By the reasons, it is believed that there is no example of computing their Alexander polynomial with non-triviality even if L is an easy knot.

In contrast, in the definition of the K_1 -class $\Delta_{\rho}^{K_1}$, we considered no fraction. In [Nos, §§4–6], the author showed some non-trivial examples of $\Delta_{\rho}^{K_1}$ without using fraction, but using a logarithm and the homomorphism Υ_* . This is a reason why we employ Novikov rings instead of fractional fields and polynomial ring.

References

- [C] T. D. Cochran. *Noncommutative knot theory*. Algebr. Geom. Topol., **4**: 347–398, 2004.
- [CG] A Casson, C. M Gordon, *Cobordism of classical knots*, Progr. Math. 62, Birkhäuser, Boston (1986) 181–199.
- [Fox] R. H. Fox, *Free differential calculus. I. Derivation in the free group ring*, Ann. of Math. **57** (1953), no. 2, 547–560.
- [Fri1] S. Friedl, *Novikov homology and noncommutative Alexander polynomials*, Journal of Knot Theory and its Ramifications 26, 1740013 (2017)

- [Fri2] S. Friedl, *Reidemeister torsion, the Thurston norm and Harvey's invariants*, Pacific Journal of Mathematics, Vol. 230: 271–296 (2007)
- [FKK] S. Friedl, T. Kim, T. Kitayama *Poincare duality and degrees of twisted Alexander polynomials*, Indiana Univ. Math. J. 61 (2012) 147–192.
- [FV] S. Friedl, S. Vidussi, *A survey of twisted Alexander polynomials*, The mathematics of knots, 45–94, Contrib. Math. Comput. Sci., **1**, Springer, Heidelberg, 2011.
- [GS] H. Goda, T. Sakasai, *Factorization formulas and computations of higher-order Alexander invariants for homologically fibered knots*, Journal of Knot Theory and Its Ramifications 20 (2011), 1355–1380.
- [HT] R. Hakamata, M. Teragaito, *Left-orderable fundamental group and Dehn surgery on genus one two-bridge knots*, Algebraic and Geometric Topology **14** (2014), 4, 2125–2148.
- [Har] S. L. Harvey. *Monotonicity of degrees of generalized Alexander polynomials of groups and 3-manifolds*, Math. Proc. Cambridge Philos. Soc. 140 no. 3, 431–450, 2006.
- [HKL] C. Herald, P. Kirk, C. Livingston, *Metabelian representations, twisted Alexander polynomials, knot slicing, and mutation*, Math. Z. **265** (2010), no. 4, 925–949.
- [Kab] Y. Kabaya, *Cyclic branched coverings of knots and quandle homology*, Pacific Journal of Mathematics, **259** (2012), No. 2, 315–347.
- [KL] P. Kirk, C. Livingston. *Twisted Alexander invariants, Reidemeister torsion, and Casson-Gordon invariants*, Topology, **38** (1999), 635–661.
- [Kitan] T. Kitano, *Twisted Alexander polynomial and Reidemeister torsion*, Pacific J. Math. **174** (1996), no. 2, 431–442.
- [Ki1] T. Kitayama, *Normalization of twisted Alexander invariants*, Internat. J. Math. **26** (2015) 1550077
- [Ki2] T. Kitayama, *Non-commutative Reidemeister torsion and Morse-Novikov theory*, Proc. Amer. Math. Soc. **138** (2010) 3345–3360.
- [Lic] W.B. Lickorish, *An introduction to knot theory*, Springer-Verlag, Berlin - New York, 1974.
- [Lin] X.S. Lin, *Representations of knot groups and twisted Alexander polynomials*, Acta Math. Sin. (Engl. Ser.) **17** (2001), 361–380.
- [LS] R. C. Lyndon, P. E. Schupp, *Combinatorial group theory*, Springer, Berlin, 1977.
- [Nos] T. Nosaka, *Twisted Alexander invariants of knot group representations*, preprint, available at arXiv:2002.10192
- [Mil1] J. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc. 72 (1966), 358–426.
- [Mil2] J. Milnor, *A duality theorem for Reidemeister torsion*, Annals of Mathematics, 1962, **76** (1), 137–147.
- [P] A.V. Pajitnov, *Closed orbits of gradient flows and logarithms of non-abelian Witt vectors*, K-theory, vol. 21. No. 4, 2000, 301–324.
- [PR] A.V. Pajitnov and A. Ranicki, *The Whitehead group of the Novikov ring*, K-theory **21** (2000), 325–365.
- [Tro] H. F. Trotter, *Homology of group systems with applications to knot theory*, Ann. of Math. **76** (1962), 464–498.
- [Tur] V. Turaev, *Introduction to combinatorial torsions*, Lectures in Mathematics, ETH Zürich, Birkhäuser, Basel, 2001. MR 2001m:57042 Zbl 0970.57001
- [Wada] M. Wada, *Twisted Alexander polynomial for finitely presentable groups*, Topology **33** (1994), 241–256.
- [Wal] F. Waldhausen, *Algebraic K-theory of generalized free products. I, II*, Ann. of Math. (2) 108 (1978) 135–204
- [Wei] C. A. Weibel, *The K-book*, volume 145 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K-theory.

DEPARTMENT OF MATHEMATICS TOKYO INSTITUTE OF TECHNOLOGY 2-12-1 OOKAYAMA ,
MEGURO-KU TOKYO 152-8551 JAPAN