

MedDialog: Two Large-scale Medical Dialogue Datasets

Xuehai He, Shu Chen, Zeqian Ju, Xiangyu Dong, Hongchao Fang, Sicheng Wang, Yue Yang, Jiaqi Zeng, Ruisi Zhang, Ruoyu Zhang, Meng Zhou, Penghui Zhu, Pengtao Xie

UC San Diego

PENGTAOXIE2008@GMAIL.COM

Abstract

Medical dialogue systems are promising in assisting in telemedicine to increase access to healthcare services, improve the quality of patient care, and reduce medical costs. To facilitate the research and development of medical dialogue systems, we build two large-scale medical dialogue datasets: MedDialog-EN and MedDialog-CN. MedDialog-EN is an English dataset containing 0.3 million conversations between patients and doctors and 0.5 million utterances. MedDialog-CN is a Chinese dataset containing 1.1 million conversations and 4 million utterances. To our best knowledge, MedDialog-(EN,CN) are the largest medical dialogue datasets to date. The dataset is available at <https://github.com/UCSD-AI4H/Medical-Dialogue-System>

1. Introduction

Telemedicine refers to the practice of delivering patient care remotely, where doctors provide medical consultations to patients using HIPAA compliant video-conferencing tools. As an important complement to traditional face-to-face medicine practiced physically in hospitals and clinics, telemedicine has a number of advantages. First, it increases access to care. For people living in medically under-served communities (e.g., rural areas) that are in shortage of clinicians, telemedicine enables them to receive faster and cheaper care compared with traveling over a long distance to visit a clinician. Second, it reduces healthcare cost. In a study¹ by Jefferson Health, it is shown that diverting patients from emergency departments with telemedicine can save more than \$1,500 per visit. Third, telemedicine can improve quality of care. The study in (Pande and Morris, 2015) shows that telemedicine patients score lower for depression, anxiety, and stress, and have 38% fewer hospital admissions. Other advantages include improving patient engagement and satisfaction, improving provider satisfaction, etc. Please refer to (Wootton et al., 2017) for a more comprehensive review.

While telemedicine is promising, it has several limitations. First, it puts additional burden to physicians. In addition to practicing face-to-face medicine which already makes physicians highly occupied, physicians need to provide remote consultations in telemedicine, which further increases the risk of physician burnout. Second, different from in-hospital patients, the progression of whose medical conditions can be easily tracked by clinicians,

1. <https://www.healthleadersmedia.com/clinical-care/cost-savings-telemedicine-estimated-19-120-patient-visit>

remote patients are difficult to track and monitor. To address such problems, there has been increasing research interests in developing artificial intelligence (AI) methods to assist in telemedicine. In particular, medical dialogue systems are being developed to server as “virtual doctors”. These “virtual doctors” are aimed to interact with patients via natural dialogues, asking about the medical conditions and history of patients and providing clinical advice. They can also proactively reach out to patients to ask about the progression of patients’ conditions and provide timely interventions accordingly.

To build medical dialogue systems, a large collection of conversations between patients and doctors are needed as training data. Due to data privacy concerns, such data is very difficult to obtain. The existing medical dialogue datasets are limited in size or biased to certain diseases, which cannot adequately serve the purpose to train medical dialogue systems that can achieve doctor-level intelligence and cover all specialties in medicine.

To address the limitations of existing datasets, we build two large-scale medical dialogue datasets: MedDialog-EN in English and MedDialog-CN in Chinese. MedDialog-EN contains 0.3 million patient-doctor consultations and 0.5 million utterances. MedDialog-CN contains 1.1 million consultations and 4 million utterances. Dialogs in these two datasets cover almost all specialties in medicine, ranging from internal medicine to family medicine and covers a wide spectrum of diseases, including cancer, pneumonia, etc. To our best knowledge, they are the largest medical dialogue datasets to date. The data is open to the public.

2. Datasets

2.1. MedDialog-EN

The MedDialog-EN dataset contains 257,454 English consultations between patients and doctors. The total number of utterances is 514,908: 257,454 from doctors and 257,454 from patients. Each consultation consists of two parts: (1) description of patient’s medical conditions; (2) conversation between patient and doctor. Figure 1 shows an exemplar consultation. The data is crawled from icliniq.com² and healthcaremagic.com³, which are two online platforms of healthcare services, including symptom self-checker, video consultation, online chat with doctors, etc.

The consultations cover 51 categories of communities including diabetes, elderly problems, pain management, etc. and 96 specialties including andrology, cardiology, nephrology, pharmacology, etc. The consultations are conducted from 2008 to 2020.

2.2. MedDialog-CN

The MedDialog-CN dataset contains 1,145,231 Chinese consultations between patients and doctors. The total number of utterances is 3,959,333: 2,179,008 from doctors and 1,780,325 from patients. Each consultation consists of three parts: (1) description of patient’s medical condition and history; (2) conversation between patient and doctor; (3) (optional) diagnosis and treatment suggestions given by the doctor. In the description of patient’s medical condition and history, the following fields are included: present disease, detailed description of present disease, what help is needed from the doctor, how long the disease has been,

2. <https://www.icliniq.com/>

3. <https://www.healthcaremagic.com/>

Description

I get mild left-sided chest pain with low Hb and vitamin B12 levels. Please help.

Dialogue

Patient:

Hello doctor,

I am a 39-year-old woman. I have mild pain in the left side of the chest (below the neck and above the breast) and then sensation in the upper back for four days. It comes and goes. Sometimes it goes to the right side of the chest also.

I had my ECG and blood test 6 months ago. ECG and blood sugar were normal. No hypertension but hemoglobin was 10 and Vitamin B 12 was below average. What can I do as in lockdown it is not possible to see the doctor as a person. Please help.

Doctor:

Hello. I would like to ask you some more questions, do you have symptoms of acid peptic disease or GERD? Or any burning sensation in the epigastric region, the center of your chest (heartburn). If you do, it could also present as chest pain so I will guide you accordingly. I would also like to rule out any muscular pain, for which I will encourage you to take a muscle relaxant and see if it helps. Take tablet Muscoril (Thiocolchicoside 4 mg) one tablet once a day when you experience pain and let me know in the follow up in a couple of days to see if it relieves the pain. Like you said your Vitamin B12 levels are below normal, it could also be neuropathic pain for which I will only advise vitamin B12 supplements (tablet Methylcobalamin once a day for three months) or B complex supplements. You can also improve these levels with diet. Take a diet enriched with beef, liver, and chicken. Fish and shellfish such as trout, salmon, tuna fish, and clams. Fortified breakfast cereal. Low-fat milk, yogurt, cheese, and eggs. Since your Hb is also below the ideal levels, take iron-rich foods which are usually Vitamin B12 rich as well. So I will advise diet modification to incorporate these in your everyday routine or take supplements with iron and Vitamin B12 as well. Because anemia can also present with the said symptoms. Lastly, I will encourage you to reduce weight with exercise and changing your diet and switching to a low-fat diet with the addition of more fruits and vegetables to your diet. Try to start with at least 30-40 minutes of cardio workout every day and work it up from there according to your stamina and you will see the visible change that you feel. You will feel more active and fresh. Try to bring your BMI (body mass index) as close to the normal range as you can. Because obesity itself can cause countless problems as well. I hope this helps. ECG (electrocardiography). Do not lift heavyweights. Take a low fat, high fiber diet. After two days.

Figure 1: An exemplar consultation, which includes (1) description of medical conditions of the patient, (2) dialogue between doctor and patient.

medications, allergies, and past disease. Figure 2 shows an exemplar consultation. In the conversation, there are cases that multiple consecutive utterances are from the same person (either doctor or patient) and these utterances were posted at different time points. If we combine consecutive utterances from the same person into a single one, there are 3,209,660 utterances: 1,981,844 from doctors and 1,227,816 from patients. The data is crawled from haodf.com⁴, which is an online platform of healthcare services, including medical consultation, scheduling appointment with doctors, etc.

The consultations cover 29 broad categories of specialties including internal medicine, pediatrics, dentistry, etc. and 172 fine-grained specialties including cardiology, neurology, gastroenterology, urology, etc. The consultations are conducted from 2010 to 2020.

3. Advantages of our datasets

To our best knowledge, MedDialog-EN and MedDialog-CN are the largest English and Chinese medical dialog dataset respectively. They have the following advantages.

- **Large number of conversations and utterances.** MedDialog-EN has about 0.3 million conversations and 0.5 million utterances. MedDialog-CN has about 1.1 million conversations and 4 million utterances.
- **Broad coverage of medical specialties.** Consultations in MedDialog-EN are about 96 categories of specialties. Consultations in MedDialog-CN are about 29 broad categories of specialties and 172 fine-grained specialties.
- **Diversity of the patients.** The patients in MedDialog-EN are from all over the world, with different nationalities, ethics, age, gender, occupation, education, income, etc. The patients in MedDialog-CN are from 31 provincial-level administrative divisions in China, with different ethics, age, gender, occupation, education, income, etc. Such diversity greatly minimizes population biases in these two datasets.

4. Related Works

Table 1 shows a comparison of our dataset with several other medical dialogue datasets. The number of dialogs and diseases in our dataset are both much larger than those in other datasets.

Dataset	#dialogs	#diseases
Muzhi (Wei et al., 2018)	710	4
Dxy (Xu et al., 2019)	527	5
COVID-EN (Yang et al., 2020)	603	1
COVID-CN (Yang et al., 2020)	1,088	1
MedDialog-EN	257,454	96
MedDialog-CN	3,407,494	172

Table 1: Comparison with other datasets.

4. <https://www.haodf.com/>

5. Conclusions

To facilitate the research and development of medical dialogue systems that can potentially assist in telemedicine, we build two large-scale medical dialogue datasets. MedDialog-EN contains 0.3 million conversations between patients and doctors and 0.5 million utterances. MedDialog-CN contains 1.1 million conversations and 4 million utterances. The datasets are publicly available and are continuously growing.

References

- Reena L Pande and Michael Morris. Leveraging remote behavioral health interventions to improve medical outcomes and reduce costs. *Am J Manag Care*, 21(2):e000–e000, 2015.
- Zhongyu Wei, Qianlong Liu, Baolin Peng, Huaixiao Tou, Ting Chen, Xuan-Jing Huang, Kam-Fai Wong, and Xiang Dai. Task-oriented dialogue system for automatic diagnosis. In *Proceedings of the 56th Annual Meeting of the Association for Computational Linguistics (Volume 2: Short Papers)*, pages 201–207, 2018.
- Richard Wootton, John Craig, and Victor Patterson. *Introduction to telemedicine*. CRC Press, 2017.
- Lin Xu, Qixian Zhou, Ke Gong, Xiaodan Liang, Jianheng Tang, and Liang Lin. End-to-end knowledge-routed relational dialogue system for automatic diagnosis. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 33, pages 7346–7353, 2019.
- Wenmian Yang, Guangtao Zeng, Bowen Tan, Zeqian Ju, Subrato Chakravorty, Xuehai He, Shu Chen, Xingyi Yang, Qingyang Wu, Zhou Yu, et al. On the generation of medical dialogues for covid-19. *arXiv preprint arXiv:2005.05442*, 2020.

Description of medical conditions and history

疾病：宝宝眼角红红的，严重时轻微溃烂。

(Disease: The baby's eyes are red and slightly ulcerated when becoming severe.)

病情描述：宝宝眼角红红的，用手挠，严重时轻微溃烂，怎么回事。用了紫草膏很快消失过两天又出来了
(Description of medical condition: The baby's eyes are red and itchy, scratched with hand, and slightly ulcerated when becoming severe. After using Burt's bee Res-Q ointment, it disappeared quickly but came out after two days.)

希望获得的帮助：宝宝眼角红红怎么回事

(Help needed: What's wrong with baby's red eyes?)

患病多久：一月内

(Hong long the condition has been: Less than one month)

过敏史：无

(Allergies: No)

既往病史：无

(Past medical history: No)

Dialogue

医生：感谢您的信任，病情资料我已详细阅读。根据您现有的资料，建议：睑缘炎。图片不是很清楚。经常揉是吧？

(Doctor: Thank you for your trust. I have read the medical information in detail. Based on the existing information, the diagnosis is blepharitis. The picture is not very clear. Scratch it often, right?)

病人：出生到现在奶量一直很少，嘴唇老是干干的，也不像别的宝宝流口水

(Patient: Drinks little amount of milk since birth, and the baby's lips are always dry, and not drooling like other babies.)

医生：眼部是局部炎症

(Doctor: Eyes have local arthritis.)

病人：是的

(Patient: Yes)

医生：给予典必殊眼膏一天两次

(Doctor: Use Tobramycin and Dexamethasone eye ointment twice a day)

病人：这个怎么回事

(Patient: What's going on?)

医生：考虑睑腺炎或者睑缘炎

(Doctor: Consider blepharitis or blepharitis)

病人：严重吗

(Patient: is it severe?)

医生：目前看，不算严重。用几天药物看看

(Doctor: At present, it is not severe. Try to take the medications for a few days first.)

病人：哦哦

(Patient: OK)

医生：边治疗边交流吧

(Doctor: Let me know how it works.)

Diagnosis and suggestions

病情摘要及初步印象：睑腺炎

(Summary of the condition and initial impressions: Blepharitis)

总结建议：局部炎症，给予典必殊眼膏一天两次，观察恢复情况，必要时医院就诊。

(Summary of recommendations: For local inflammation, use Tobramycin and Dexamethasone eye ointment eye ointment twice a day, monitor the recovery, and go to the hospital if necessary.)

Figure 2: An exemplar consultation, which includes (1) description of medical conditions and history of the patient, (2) dialogue between doctor and patient, and (3) diagnosis and treatment suggestions given by the doctor.