One-generated nilpotent Novikov algebras <sup>1</sup>

## Luisa M. Camacho<sup>a</sup>, Iqboljon Karimjanov<sup>b</sup>, Ivan Kaygorodov<sup>c</sup> & Abror Khudoyberdiyev<sup>d</sup>

<sup>a</sup> University of Sevilla, Sevilla, Spain.

<sup>b</sup> Andijan State University, Andijan, Uzbekistan.

<sup>c</sup> CMCC, Universidade Federal do ABC, Santo André, Brazil.

<sup>d</sup> National University of Uzbekistan, Institute of Mathematics Academy of Sciences of Uzbekistan, Tashkent, Uzbekistan

E-mail addresses:

Luisa M. Camacho (lcamacho@us.es) Iqboljon Karimjanov (iqboli@gmail.com) Ivan Kaygorodov (kaygorodov.ivan@gmail.com) Abror Khudoyberdiyev (khabror@mail.ru)

**Abstract**: We give a classification of 5- and 6-dimensional complex one-generated nilpotent Novikov algebras.

**Keywords**: Novikov algebra, nilpotent algebra, algebraic classification, central extension.

MSC2010: 17D25, 17A30.

#### Introduction

The algebraic classification (up to isomorphism) of algebras of dimension n from a certain variety defined by some family of polynomial identities is a classic problem in the theory of non-associative algebras. There are many results related to the algebraic classification of low-dimensional algebras in the varieties of Jordan, Lie, Leibniz, Zinbiel and many other algebras [1, 11, 14–18, 27–29, 32, 41, 42]. Another interesting direction is a study of one-generated objects. The description of one-generated groups is well-known: there is only one one-generated group of order n. In the case of algebras, there are some similar results, such as the description of n-dimensional one-generated nilpotent associative [19], noncommutative Jordan [34], Leibniz and Zinbiel algebras [43]. It was proven that there is only one n-dimensional one-generated nilpotent algebra in these varieties. But on the other hand, as we can see in varieties of Novikov [36], assosymmetric [33], bicommutative [39], commutative [25], and terminal [38] algebras, there are more than one 4-dimensional one-generated nilpotent algebra from these varieties. It is easy to see that a nilpotent one-generated algebra is a central extension of a nilpotent one-generated algebra of a smaller dimension. In the present paper, we give an algebraic classification of 5- and 6-dimensional one-generated nilpotent Novikov algebras introduced by Novikov and Balinsky in 1985 [3].

The variety of Novikov algebras is defined by the following identities:

$$(xy)z = (xz)y,$$
  
$$(xy)z - x(yz) = (yx)z - y(xz).$$

It contains the commutative associative algebras as a subvariety. On the other hand, the variety of Novikov algebras is the intersection of the variety of right commutative algebras (defined by the first Novikov identity) and the variety of left symmetric (Pre-Lie) algebras (defined by the second Novikov identity). Also, any Novikov algebra under the commutator multiplication is a Lie algebra [8,9], and Novikov algebras are related with Tortken and Novikov-Poisson algebras [6,20,47]. The systematic study of Novikov algebras started after the paper by Zelmanov where all complex finite-dimensional simple Novikov algebras were found [49]. The first nontrivial examples of infinite-dimensional simple Novikov algebras were constructed in [26]. Also, the simple Novikov algebras were described in infinite-dimensional case and over fields of positive characteristic [44,46,48]. The

1

<sup>&</sup>lt;sup>1</sup> This work was supported by RFBR 18-31-20004; FAPESP 18/15712-0, 19/00192-3; Ministerio de Economía y Competitividad (Spain), grant MTM2016-79661-P (European FEDER support included, UE). The authors thank Farukh Mashurov and the referee of the paper for constructive comments.

algebraic classification of 3-dimensional Novikov algebras was given in [4], and for some classes of 4-dimensional algebras, it was given in [7]; the geometric classification of 3-dimensional Novikov algebras was given in [5] and of 4-dimensional nilpotent Novikov algebras in [35]. Many other pure algebraic properties were studied in a series of papers by Dzhumadildaev [21–24].

Our method to classify nilpotent Novikov algebras is based on the calculation of central extensions of smaller nilpotent algebras from the same variety. The algebraic study of central extensions of Lie and non-Lie algebras has been an important topic for years [2, 30, 31, 36, 40, 45, 50]. First, Skjelbred and Sund used central extensions of Lie algebras for a classification of nilpotent Lie algebras [45]. After that, using the method described by Skjelbred and Sund, all non-Lie central extensions of all 4-dimensional Malcev algebras were described [31], and also all non-associative central extensions of 3-dimensional Jordan algebras [30], all anticommutative central extensions of the 3-dimensional anticommutative algebras [10], and all central extensions of the 2-dimensional algebras [12]. Note that the method of central extensions is an important tool in the classification of nilpotent algebras, which was used to describe all 4-dimensional nilpotent associative algebras [18], all 5-dimensional nilpotent Jordan algebras [29], all 5-dimensional nilpotent restricted Lie algebras [16], all 5-dimensional nilpotent Malcev algebras [32], all 6-dimensional nilpotent anticommutative algebras [37], all 8-dimensional dual Mock Lie algebras [13], and some others.

#### 1. THE ALGEBRAIC CLASSIFICATION OF NILPOTENT NOVIKOV ALGEBRAS

1.1. **Method of classification of nilpotent algebras.** Throughout this paper, we use the notations and methods well written in [12, 30, 31], which we have adapted for the Novikov case with some modifications. Further in this section we give some important definitions.

Let  $(\mathbf{A}, \cdot)$  be a Novikov algebra over  $\mathbb{C}$  and  $\mathbb{V}$  a vector space over  $\mathbb{C}$ . The  $\mathbb{C}$ -linear space  $Z^2(\mathbf{A}, \mathbb{V})$  is defined as the set of all bilinear maps  $\theta \colon \mathbf{A} \times \mathbf{A} \longrightarrow \mathbb{V}$  such that

$$\theta(xy,z) = \theta(xz,y),$$
  
$$\theta(xy,z) - \theta(x,yz) = \theta(yx,z) - \theta(y,xz).$$

These elements will be called *cocycles*. For a linear map f from  $\mathbf{A}$  to  $\mathbb{V}$ , if we define  $\delta f : \mathbf{A} \times \mathbf{A} \longrightarrow \mathbb{V}$  by  $\delta f(x,y) = f(xy)$ , then  $\delta f \in \mathbf{Z}^2(\mathbf{A},\mathbb{V})$ . We define  $\mathbf{B}^2(\mathbf{A},\mathbb{V}) = \{\theta = \delta f : f \in \mathrm{Hom}(\mathbf{A},\mathbb{V})\}$ . We define the *second cohomology space*  $\mathbf{H}^2(\mathbf{A},\mathbb{V})$  as the quotient space  $\mathbf{Z}^2(\mathbf{A},\mathbb{V}) / \mathbf{B}^2(\mathbf{A},\mathbb{V})$ .

Let  $\operatorname{Aut}(\mathbf{A})$  be the automorphism group of  $\mathbf{A}$  and let  $\phi \in \operatorname{Aut}(\mathbf{A})$ . For  $\theta \in \operatorname{Z}^2(\mathbf{A}, \mathbb{V})$  define the action of the group  $\operatorname{Aut}(\mathbf{A})$  on  $\operatorname{H}^2(\mathbf{A}, \mathbb{V})$  by  $\phi\theta(x, y) = \theta(\phi(x), \phi(y))$ . It is easy to verify that  $\operatorname{B}^2(\mathbf{A}, \mathbb{V})$  is invariant under the action of  $\operatorname{Aut}(\mathbf{A})$ . So, we have an induced action of  $\operatorname{Aut}(\mathbf{A})$  on  $\operatorname{H}^2(\mathbf{A}, \mathbb{V})$ .

Let  $\mathbf{A}$  be a Novikov algebra of dimension m over  $\mathbb{C}$  and  $\mathbb{V}$  be a  $\mathbb{C}$ -vector space of dimension k. For  $\theta \in \mathbf{Z}^2$   $(\mathbf{A}, \mathbb{V})$ , define on the linear space  $\mathbf{A}_{\theta} = \mathbf{A} \oplus \mathbb{V}$  the bilinear product " $[-,-]_{\mathbf{A}_{\theta}}$ " by  $[x+x',y+y']_{\mathbf{A}_{\theta}} = xy + \theta(x,y)$  for all  $x,y \in \mathbf{A}, x',y' \in \mathbb{V}$ . The algebra  $\mathbf{A}_{\theta}$  is called an k-dimensional central extension of  $\mathbf{A}$  by  $\mathbb{V}$ . One can easily check that  $\mathbf{A}_{\theta}$  is a Novikov algebra if and only if  $\theta \in \mathbf{Z}^2(\mathbf{A}, \mathbb{V})$ .

Call the set  $Ann(\theta) = \{x \in \mathbf{A} : \theta(x, \mathbf{A}) + \theta(\mathbf{A}, x) = 0\}$  the *annihilator* of  $\theta$ . We recall that the *annihilator* of an algebra  $\mathbf{A}$  is defined as the ideal  $Ann(\mathbf{A}) = \{x \in \mathbf{A} : x\mathbf{A} + \mathbf{A}x = 0\}$ . Observe that  $Ann(\mathbf{A}_{\theta}) = (Ann(\theta) \cap Ann(\mathbf{A})) \oplus \mathbb{V}$ .

The following result shows that every algebra with a non-zero annihilator is a central extension of a smaller-dimensional algebra. As a consequence, it is easy to see that a one-generated nilpotent algebra is a central extension of a one-generated nilpotent algebra of a smaller dimension.

**Lemma 1.** Let **A** be an n-dimensional Novikov algebra such that  $\dim(\operatorname{Ann}(\mathbf{A})) = m \neq 0$ . Then there exists, up to isomorphism, a unique (n-m)-dimensional Novikov algebra  $\mathbf{A}'$  and a bilinear map  $\theta \in \mathrm{Z}^2(\mathbf{A}, \mathbb{V})$  with  $\operatorname{Ann}(\mathbf{A}) \cap \operatorname{Ann}(\theta) = 0$ , where  $\mathbb{V}$  is a vector space of dimension m, such that  $\mathbf{A} \cong \mathbf{A}'_{\theta}$  and  $\mathbf{A}/\operatorname{Ann}(\mathbf{A}) \cong \mathbf{A}'$ .

*Proof.* Let  $\mathbf{A}'$  be a linear complement of  $\mathrm{Ann}(\mathbf{A})$  in  $\mathbf{A}$ . Define a linear map  $P \colon \mathbf{A} \longrightarrow \mathbf{A}'$  by P(x+v) = x for  $x \in \mathbf{A}'$  and  $v \in \mathrm{Ann}(\mathbf{A})$ , and define a multiplication on  $\mathbf{A}'$  by  $[x,y]_{\mathbf{A}'} = P(xy)$  for  $x,y \in \mathbf{A}'$ . For  $x,y \in \mathbf{A}$ , we have

$$P(xy) = P((x - P(x) + P(x))(y - P(y) + P(y))) = P(P(x)P(y)) = [P(x), P(y)]_{\mathbf{A}'}.$$

Since P is a homomorphism  $P(\mathbf{A}) = \mathbf{A}'$  is a Novikov algebra and  $\mathbf{A}/\operatorname{Ann}(\mathbf{A}) \cong \mathbf{A}'$ , which gives us the uniqueness. Now, define the map  $\theta \colon \mathbf{A}' \times \mathbf{A}' \longrightarrow \operatorname{Ann}(\mathbf{A})$  by  $\theta(x,y) = xy - [x,y]_{\mathbf{A}'}$ . Thus,  $\mathbf{A}'_{\theta}$  is  $\mathbf{A}$  and therefore  $\theta \in \mathrm{Z}^2(\mathbf{A},\mathbb{V})$  and  $\operatorname{Ann}(\mathbf{A}) \cap \operatorname{Ann}(\theta) = 0$ .

**Definition 2.** Let **A** be an algebra and I be a subspace of Ann(**A**). If  $\mathbf{A} = \mathbf{A}_0 \oplus I$  then I is called an annihilator component of **A**.

**Definition 3.** A central extension of an algebra A without annihilator component is called a non-split central extension.

Our task is to find all central extensions of an algebra  $\mathbf{A}$  by a space  $\mathbb{V}$ . In order to solve the isomorphism problem we need to study the action of  $\operatorname{Aut}(\mathbf{A})$  on  $\operatorname{H}^2(\mathbf{A},\mathbb{V})$ . To do that, let us fix a basis  $e_1,\ldots,e_s$  of  $\mathbb{V}$ , and  $\theta\in\operatorname{Z}^2(\mathbf{A},\mathbb{V})$ . Then  $\theta$  can be uniquely written as  $\theta(x,y)=\sum_{i=1}^s\theta_i(x,y)\,e_i$ , where  $\theta_i\in\operatorname{Z}^2(\mathbf{A},\mathbb{C})$ . Moreover,  $\operatorname{Ann}(\theta)=\operatorname{Ann}(\theta_1)\cap\operatorname{Ann}(\theta_2)\cap\cdots\cap\operatorname{Ann}(\theta_s)$ . Furthermore,  $\theta\in\operatorname{B}^2(\mathbf{A},\mathbb{V})$  if and only if all  $\theta_i\in\operatorname{B}^2(\mathbf{A},\mathbb{C})$ . It is not difficult to prove (see [31, Lemma 13]) that given a Novikov algebra  $\mathbf{A}_\theta$ , if we write as above  $\theta(x,y)=\sum_{i=1}^s\theta_i(x,y)\,e_i\in\operatorname{Z}^2(\mathbf{A},\mathbb{V})$  and  $\operatorname{Ann}(\theta)\cap\operatorname{Ann}(\mathbf{A})=0$ , then  $\mathbf{A}_\theta$  has an annihilator component if and only if  $[\theta_1]$ ,  $[\theta_2]$ , ...,  $[\theta_s]$  are linearly dependent in  $\operatorname{H}^2(\mathbf{A},\mathbb{C})$ .

Let  $\mathbb{V}$  be a finite-dimensional vector space over  $\mathbb{C}$ . The  $Grassmannian\ G_k\ (\mathbb{V})$  is the set of all k-dimensional linear subspaces of  $\mathbb{V}$ . Let  $G_s\ (\mathbb{H}^2\ (\mathbf{A},\mathbb{C}))$  be the  $Grassmannian\ of\ subspaces\ of\ dimension\ s\ in\ \mathbb{H}^2\ (\mathbf{A},\mathbb{C})$ . There is a natural action of  $\operatorname{Aut}(\mathbf{A})$  on  $G_s\ (\mathbb{H}^2\ (\mathbf{A},\mathbb{C}))$ . Let  $\phi\in\operatorname{Aut}(\mathbf{A})$ . For  $W=\langle[\theta_1]\ , [\theta_2]\ ,\dots, [\theta_s]\rangle\in G_s\ (\mathbb{H}^2\ (\mathbf{A},\mathbb{C}))$  define  $\phi W=\langle[\phi\theta_1]\ , [\phi\theta_2]\ ,\dots, [\phi\theta_s]\rangle$ . We denote the orbit of  $W\in G_s\ (\mathbb{H}^2\ (\mathbf{A},\mathbb{C}))$  under the action of  $\operatorname{Aut}(\mathbf{A})$  by  $\operatorname{Orb}(W)$ . Given

$$W_1 = \langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle, W_2 = \langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle \in G_s \left( H^2 (\mathbf{A}, \mathbb{C}) \right),$$

we easily have that if  $W_1 = W_2$ , then  $\bigcap_{i=1}^s \operatorname{Ann}(\theta_i) \cap \operatorname{Ann}(\mathbf{A}) = \bigcap_{i=1}^s \operatorname{Ann}(\vartheta_i) \cap \operatorname{Ann}(\mathbf{A})$ , and therefore we can introduce the set

$$\mathbf{T}_{s}(\mathbf{A}) = \left\{ W = \langle \left[\theta_{1}\right], \left[\theta_{2}\right], \dots, \left[\theta_{s}\right] \rangle \in G_{s}\left(\mathbf{H}^{2}\left(\mathbf{A}, \mathbb{C}\right)\right) : \bigcap_{i=1}^{s} \operatorname{Ann}(\theta_{i}) \cap \operatorname{Ann}(\mathbf{A}) = 0 \right\},\,$$

which is stable under the action of  $Aut(\mathbf{A})$ .

Now, let  $\mathbb{V}$  be an s-dimensional linear space and let us denote by  $\mathbf{E}(\mathbf{A}, \mathbb{V})$  the set of all non-split s-dimensional central extensions of  $\mathbf{A}$  by  $\mathbb{V}$ . By above, we can write

$$\mathbf{E}\left(\mathbf{A}, \mathbb{V}\right) = \left\{\mathbf{A}_{\theta} : \theta\left(x, y\right) = \sum_{i=1}^{s} \theta_{i}\left(x, y\right) e_{i} \text{ and } \left\langle\left[\theta_{1}\right], \left[\theta_{2}\right], \dots, \left[\theta_{s}\right]\right\rangle \in \mathbf{T}_{s}(\mathbf{A})\right\}.$$

We also have the following result, which can be proved as in [31, Lemma 17].

**Lemma 4.** Let  $\mathbf{A}_{\theta}$ ,  $\mathbf{A}_{\vartheta} \in \mathbf{E}(\mathbf{A}, \mathbb{V})$ . Suppose that  $\theta(x, y) = \sum_{i=1}^{s} \theta_{i}(x, y) e_{i}$  and  $\vartheta(x, y) = \sum_{i=1}^{s} \vartheta_{i}(x, y) e_{i}$ . Then the Novikov algebras  $\mathbf{A}_{\theta}$  and  $\mathbf{A}_{\vartheta}$  are isomorphic if and only if

$$\operatorname{Orb} \langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle = \operatorname{Orb} \langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle.$$

This shows that there exists a one-to-one correspondence between the set of  $\operatorname{Aut}(\mathbf{A})$ -orbits on  $\mathbf{T}_s(\mathbf{A})$  and the set of isomorphism classes of  $\mathbf{E}(\mathbf{A}, \mathbb{V})$ . Consequently we have a procedure that allows us, given a Novikov algebra  $\mathbf{A}'$  of dimension n-s, to construct all non-split central extensions of  $\mathbf{A}'$ . This procedure is:

Procedure

- (1) For a given Novikov algebra  $\mathbf{A}'$  of dimension n-s, determine  $\mathrm{H}^2(\mathbf{A}',\mathbb{C})$ ,  $\mathrm{Ann}(\mathbf{A}')$  and  $\mathrm{Aut}(\mathbf{A}')$ .
- (2) Determine the set of  $Aut(\mathbf{A}')$ -orbits on  $T_s(\mathbf{A}')$ .
- (3) For each orbit, construct the Novikov algebra associated with a representative of it.
- 1.2. **Notations.** Let **A** be a Novikov algebra with a basis  $e_1, e_2, \ldots, e_n$ . Then by  $\Delta_{ij}$  we will denote the bilinear form  $\Delta_{ij} : \mathbf{A} \times \mathbf{A} \longrightarrow \mathbb{C}$  with  $\Delta_{ij} (e_l, e_m) = \delta_{il}\delta_{jm}$ . Then the set  $\{\Delta_{ij} : 1 \le i, j \le n\}$  is a basis for the linear space of the bilinear forms on **A**. Then every bilinear form  $\theta$  can be uniquely written as  $\theta = \sum_{1 \le i, j \le n} c_{ij} \Delta_{ij}$ , where  $c_{ij} \in \mathbb{C}$ . Let us fix the following notations:
  - $\mathcal{N}_{i}^{i}$  jth i-dimensional one-generated Novikov algebra.
- 1.3. **Low dimensional one-generated nilpotent Novikov algebras.** Thanks to [35] we have the algebraic classification of 2-, 3- and 4- dimensional and 5-dimensional with 2-dimensional annihilator one-generated nilpotent algebras. Let us give a list if 1-, 2-, 3- and 4-dimensional one-generated nilpotent Novikov algebras.

**Remark 5.** Note that a non-split central extension of a split algebra can not be a one-generated algebra. Hence, we will consider central extensions only of non-split one-generated nilpotent algebras.

### 2. Classification of 5-dimensional one-generated nilpotent Novikov algebras

All the necessary information about coboundaries, cocycles and second cohomology spaces of 5-dimensional one-generated nilpotent Novikov algebras is given in Table A (see Table A from Appendix).

From the results of [35], it is not difficult to see that there are two 5-dimensional one-generated Novikov algebras with 2-dimensional annihilator:

2.1. Central extensions of  $\mathcal{N}_{01}^4$ . Let us use the following notations

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{14}] - [\Delta_{41}], \quad \nabla_3 = [\Delta_{22}] + [\Delta_{31}] + 2[\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{01}^4$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & xy & x^3 & 0 \\ v & xy & 0 & x^3 \end{pmatrix}.$$

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 & \alpha_2 \\ 0 & \alpha_3 & 0 & 0 \\ \alpha_3 & 0 & 0 & 0 \\ 2\alpha_3 - \alpha_2 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha^{**} & \alpha_1^* & \alpha_2^* \\ \alpha^{***} & \alpha_3^* & 0 & 0 \\ \alpha_3^* & 0 & 0 & 0 \\ 2\alpha_3^* - \alpha_2^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}^4_{01})$  on the subspace  $\langle \sum\limits_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^4 \alpha_1, \quad \alpha_2^* = x^4 \alpha_2, \quad \alpha_3^* = x^4 \alpha_3.$$

- 2.1.1. *1-dimensional central extensions of*  $\mathcal{N}_{01}^4$ . Since we are interested only in new algebras, we have the following cases:
  - (1) if  $\alpha_3 \neq 0$ , then we have the representative  $\langle \lambda \nabla_1 + \mu \nabla_2 + \nabla_3 \rangle$ ;
  - (2) if  $\alpha_3 = 0$ , then  $(\alpha_1, \alpha_2) \neq (0, 0)$  and we have the representative  $\langle \nabla_1 + \lambda \nabla_2 \rangle$  with  $\lambda \neq 0$ .

Hence, we have the following new 5-dimensional algebras:

2.1.2. 2-dimensional central extensions of  $\mathbb{N}^4_{01}$ . Consider the vector space generated by the following two cocycles

$$\begin{array}{lcl} \theta_1 & = & \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3, \\ \theta_2 & = & \beta_1 \nabla_1 + \beta_2 \nabla_2. \end{array}$$

If  $\alpha_3 = 0$ , then we have the representative  $\langle \nabla_1, \nabla_2 \rangle$ . If  $\alpha_3 \neq 0$ , then

- (1)  $\beta_2 = 0$ , then we have the family of representatives  $\langle \nabla_1, \lambda \nabla_2 + \nabla_3 \rangle$ ;
- (2)  $\beta_2 \neq 0$ , then we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_2, \mu \nabla_1 + \nabla_3 \rangle$ .

Hence, we have the following 6-dimensional algebras:

2.2. Central extensions of  $\mathcal{N}_{02}^4$ . Let us use the following notations

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = 2[\Delta_{14}] - [\Delta_{23}] - [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{02}^4$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & 0 & 0 & x^4 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & 0 & 2\alpha_2 \\ 0 & 0 & -\alpha_2 & 0 \\ 0 & 0 & 0 & 0 \\ -\alpha_2 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha_1^* & 0 & 2\alpha_2^* \\ \alpha^{**} & 0 & -\alpha_2^* & 0 \\ 0 & 0 & 0 & 0 \\ -\alpha_2^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}^4_{02})$  on the subspace  $\langle \sum_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^3 \alpha_1, \quad \alpha_2^* = x^5 \alpha_2.$$

- 2.2.1. 1-dimensional central extensions of  $\mathbb{N}_{02}^4$ . Since we are interested only in new algebras, we have assume  $\alpha_2 \neq 0$ , and we have the following cases:
  - (1) if  $\alpha_1 = 0$ , then we have the representative  $\langle \nabla_2 \rangle$ ;
  - (2) if  $\alpha_1 \neq 0$ , then by choosing  $x = \sqrt{\alpha_1/\alpha_2}$ , we have the representative  $\langle \nabla_1 + \nabla_2 \rangle$ .

Thus, we have the following 5-dimensional algebras:

2.2.2. 2-dimensional central extensions of  $\mathbb{N}_{02}^4$ . Since dim  $(H^2(\mathbb{N}_{02}^4)) = 2$ , there is only one 2-dimensional central extension of  $\mathbb{N}_{02}^4$ . Thus, we have the following 6-dimensional algebra:

$$\mathcal{N}_{04}^6 \quad : \quad e_1e_1 = e_2 \quad e_1e_2 = e_5 \quad e_1e_3 = e_4 \quad e_1e_4 = 2e_6 \quad e_2e_1 = e_3 \quad e_2e_3 = -e_6 \quad e_3e_1 = -e_4 \quad e_4e_1 = -e_6 \quad e_3e_1 = -e_4 \quad e_4e_2 = -e_6 \quad e_3e_3 = -e_6 \quad e_3e_4 = -e_6 \quad e_3e_5 = -e_6 \quad e_5e_5 = -e_6$$

2.3. Central extensions of  $\mathcal{N}_{03}^4(\lambda)_{\lambda \notin \{0,1\}}$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = (3 - 2\lambda)[\Delta_{14}] + \lambda(2 - \lambda)[\Delta_{23}] + \lambda[\Delta_{32}] + \lambda[\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{03}^4(\lambda)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & 2xy & 0 & x^4 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & (3-2\lambda)\alpha_2 \\ \alpha_1 & 0 & \lambda(2-\lambda)\alpha_2 & 0 \\ 0 & \lambda\alpha_2 & 0 & 0 \\ \lambda\alpha_2 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \alpha^* & 0 & (3-2\lambda)\alpha_2^* \\ \alpha_1^* + \lambda\alpha^* & 0 & \lambda(2-\lambda)\alpha_2^* & 0 \\ 0 & \lambda\alpha_2^* & 0 & 0 \\ \lambda\alpha_2^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}^4_{03}(\lambda))$  on the subspace  $\langle \sum_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^2(x\alpha_1 + 2\lambda(\lambda - 1)y\alpha_2), \quad \alpha_2^* = x^5\alpha_2.$$

2.3.1. *1-dimensional central extensions of*  $\mathcal{N}_{03}^4(\lambda)_{\lambda\notin\{0,1\}}$ . Since we are interested only in new algebras, we have  $\alpha_2\neq 0$ . Then choosing  $y=\frac{x\alpha_1}{2\lambda(1-\lambda)\alpha_2}$ , we have the representative  $\langle \nabla_2 \rangle$ . Hence, we have the following 5-dimensional algebras:

$$\begin{array}{llll} \mathcal{N}^{5}_{07}(\lambda)_{\lambda\notin\{0,1\}} & : & e_{1}e_{1}=e_{2} & e_{1}e_{2}=e_{3} & e_{1}e_{3}=(2-\lambda)e_{4} & e_{1}e_{4}=(3-2\lambda)e_{5} & e_{2}e_{1}=\lambda e_{3} \\ & & e_{2}e_{2}=\lambda e_{4} & e_{2}e_{3}=\lambda(2-\lambda)e_{5} & e_{3}e_{1}=\lambda e_{4} & e_{3}e_{2}=\lambda e_{5} & e_{4}e_{1}=\lambda e_{5} \end{array}$$

2.3.2. 2-dimensional central extensions of  $\mathcal{N}_{03}^4(\lambda)_{\lambda\notin\{0,1\}}$ . Since dim  $\left(H^2(\mathcal{N}_{03}^4(\lambda)_{\lambda\neq0;1})\right)=2$ , there is only one 2-dimensional central extension of  $\mathcal{N}_{03}^4(\lambda)_{\lambda\notin\{0,1\}}$ . Hence, we have the following 6-dimensional algebras:

2.4. Central extensions of  $\mathcal{N}_{03}^4(0)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{14}], \quad \nabla_2 = [\Delta_{21}], \quad \nabla_3 = 2[\Delta_{23}] - 2[\Delta_{32}] + [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{03}^4(0)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & xy & x^3 & 0 \\ t & 2xz & 2x^2y & x^4 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & \alpha_1 \\ \alpha_2 & 0 & 2\alpha_3 & 0 \\ 0 & -2\alpha_3 & 0 & 0 \\ \alpha_3 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha^{**} & \alpha^{***} & \alpha_1^* \\ \alpha_2^* & 0 & 2\alpha_3^* & 0 \\ 0 & -2\alpha_3^* & 0 & 0 \\ \alpha_3^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{03}^4(0))$  on the subspace  $\langle \sum\limits_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^5 \alpha_1, \quad \alpha_2^* = x^3 \alpha_2 + (4x^2z - 2xy^2)\alpha_3, \quad \alpha_3^* = x^5 \alpha_3.$$

- 2.4.1. 1-dimensional central extensions of  $\mathcal{N}_{03}^4(0)$ . Since we are interested only in new algebras, we have  $(\alpha_1, \alpha_3) \neq (0, 0)$  and:
  - (1) if  $\alpha_3 = 0$ ,  $\alpha_2 = 0$ , then we have the representative  $\langle \nabla_1 \rangle$ ;
  - (2) if  $\alpha_3 = 0, \alpha_2 \neq 0$ , then choosing  $x = \sqrt{\alpha_2/\alpha_1}$ , we have the representative  $\langle \nabla_1 + \nabla_2 \rangle$ ;
  - (3) if  $\alpha_3 \neq 0$ , then choosing  $y = 0, z = -\frac{\alpha_2}{4\alpha_3 \sqrt[5]{\alpha_3}}$ , we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_3 \rangle$ .

Hence, we have the following 5-dimensional algebras:

2.4.2. 2-dimensional central extensions of  $\mathcal{N}_{03}^4(0)$ . Consider the vector space generated by the following two cocycles

$$\theta_1 = \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3$$
  
$$\theta_2 = \beta_1 \nabla_1 + \beta_2 \nabla_2.$$

If  $\alpha_3 = 0$ , then we have the representative  $\langle \nabla_1, \nabla_2 \rangle$ . If  $\alpha_3 \neq 0$ , then

- (1) if  $\beta_1 = 0, \beta_2 \neq 0$ , then we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_3, \nabla_2 \rangle$ ;
- (2) if  $\beta_1 \neq 0, \beta_2 \neq 0$ , then choosing  $x = \sqrt{\beta_2/\beta_1}, y = 0, z = \frac{x^3\alpha_1 x\alpha_2}{4\alpha_3}$ , we have the representative  $\langle \nabla_1 + \nabla_2, \nabla_3 \rangle$ ;
- (3) if  $\beta_1 \neq 0, \beta_2 = 0$ , then we have the representative  $\langle \nabla_1, \nabla_3 \rangle$ .

Hence we have the following 6-dimensional algebras:

2.5. Central extensions of  $\mathcal{N}_{03}^4(1)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = [\Delta_{14}] + [\Delta_{23}] + [\Delta_{32}] + [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{03}^4(1)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & 2xy & x^3 & 0 \\ t & 2xz + y^2 & 3x^2y & x^4 \end{pmatrix}.$$

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & \alpha_2 \\ \alpha_1 & 0 & \alpha_2 & 0 \\ 0 & \alpha_2 & 0 & 0 \\ \alpha_2 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha^{**} & \alpha^* & \alpha_2^* \\ \alpha_1^* + \alpha^{**} & \alpha^* & \alpha_2^* & 0 \\ \alpha^* & \alpha_2^* & 0 & 0 \\ \alpha_2^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{03}^4(1))$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^3 \alpha_1, \quad \alpha_2^* = x^5 \alpha_2.$$

- 2.5.1. *1-dimensional central extensions of*  $\mathcal{N}^4_{03}(1)$ . We may assume  $\alpha_2 \neq 0$ , so we have the following cases:
  - (1) if  $\alpha_1 = 0$ , then we have the representative  $\langle \nabla_2 \rangle$ ;
  - (2) if  $\alpha_1 \neq 0$ , then choosing  $x = \sqrt{\alpha_1/\alpha_2}$  we have the representative  $\langle \nabla_1 + \nabla_2 \rangle$ .

Hence, we have the following 5-dimensional algebras:

2.5.2. 2-dimensional central extensions of  $\mathcal{N}_{03}^4(1)$ . Since dim  $\left(H^2(\mathcal{N}_{03}^4(1))\right)=2$ , there is only one 2-dimensional central extension of  $\mathcal{N}_{03}^4(1)$ . Hence, we have the following 6-dimensional algebra:

2.6. Central extensions of  $\mathcal{N}_{04}^4$ . Let us use the following notations

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = 2[\Delta_{14}] + [\Delta_{22}] + [\Delta_{31}], \quad \nabla_3 = [\Delta_{14}] - 2[\Delta_{23}] + 2[\Delta_{32}] - [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{04}^4$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & x & 1 & 0 \\ z & x + 2y & 2x & 1 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 & 2\alpha_2 + \alpha_3 \\ 0 & \alpha_2 & -2\alpha_3 & 0 \\ \alpha_2 & 2\alpha_3 & 0 & 0 \\ -\alpha_3 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha^{**} & 2\alpha^* + \alpha_1^* & 2\alpha_2^* + \alpha_3^* \\ \alpha^* & \alpha_2^* & -2\alpha_3^* & 0 \\ \alpha_2^* & 2\alpha_3^* & 0 & 0 \\ -\alpha_3^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathcal{N}_{04}^4)$  on the subspace  $\langle \sum_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = \alpha_1 + 2(x - 2x^2 + 4y)\alpha_3, \quad \alpha_2^* = \alpha_2, \quad \alpha_3^* = \alpha_3.$$

- 2.6.1. *1-dimensional central extensions of*  $\mathcal{N}_{04}^4$ . Since we are interested only in new algebras, we assume  $(\alpha_2, \alpha_3) \neq (0, 0)$  and we have the following cases:
  - (1) if  $\alpha_2 \neq 0, \alpha_3 = 0$ , then we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_2 \rangle$ ;
  - (2) if  $\alpha_3 \neq 0$ , then by choosing  $y = -\frac{\alpha_1 + (2x 4x^2)\alpha_3}{8\alpha_3}$ , we have the family of representatives  $\langle \lambda \nabla_2 + \nabla_3 \rangle$ .

Hence, we have the following 5-dimensional algebras:

2.6.2. 2-dimensional central extensions of  $\mathbb{N}_{04}^4$ . Consider the vector space generated by the following two cocycles

$$\theta_1 = \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3, \theta_2 = \beta_1 \nabla_1 + \beta_2 \nabla_2.$$

If  $\alpha_3 = 0$ , then we have the representative  $\langle \nabla_1, \nabla_2 \rangle$ . If  $\alpha_3 \neq 0$ , then

- (1) if  $\beta_2 = 0$ , then we have the family of representatives  $\langle \nabla_1, \lambda \nabla_2 + \nabla_3 \rangle$ ;
- (2) if  $\beta_2 \neq 0$ , then we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_2, \nabla_3 \rangle$ .

Hence, we have the following 6-dimensional algebras:

2.7. Central extensions of  $\mathcal{N}_{05}^4$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = -2[\Delta_{13}] + [\Delta_{14}] + [\Delta_{23}] + [\Delta_{32}] + [\Delta_{41}].$$

The automorphism group of  $\mathbb{N}_{05}^4$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & 2x & 1 & 0 \\ z & x^2 + x + 2y & 3x & 1 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & -2\alpha_2 & \alpha_2 \\ \alpha_1 & 0 & \alpha_2 & 0 \\ 0 & \alpha_2 & 0 & 0 \\ \alpha_2 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha^{**} & -2\alpha_2^* + \alpha^* & \alpha_2^* \\ \alpha_1^* + \alpha^* + \alpha^{**} & \alpha^* & \alpha_2^* & 0 \\ \alpha^* & \alpha_2^* & 0 & 0 \\ \alpha_2^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}^4_{05})$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = \alpha_1, \quad \alpha_2^* = \alpha_2.$$

2.7.1. 1-dimensional central extensions of  $\mathcal{N}_{05}^4$ . Assuming  $\alpha_2 \neq 0$ , we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_2 \rangle$ . Hence, we have the following 5-dimensional algebras:

2.7.2. 2-dimensional central extensions of  $\mathcal{N}_{05}^4$ . Since dim  $(H^2(\mathcal{N}_{05}^4)) = 2$ , there is only one 2-dimensional central extension of  $\mathcal{N}_{05}^4$ . Hence, we have the following 6-dimensional algebra:

$$\mathcal{N}_{12}^6$$
:  $e_1e_1=e_2$   $e_1e_2=e_3$   $e_1e_3=e_4-2e_5$   $e_1e_4=e_5$   $e_2e_1=e_3+e_4+e_6$   
 $e_2e_2=e_4$   $e_2e_3=e_5$   $e_3e_1=e_4$   $e_3e_2=e_5$   $e_4e_1=e_5$ 

Summarizing the results of the present sections and using the information from Table A from Appendix, we have the following theorem.

**Theorem 6.** Let  $\mathbb{N}$  be a 5-dimensional complex one-generated nilpotent Novikov algebra. Then  $\mathbb{N}$  is isomorphic to one of the algebras  $\mathbb{N}_{j}^{5}$ ,  $j = 01, \ldots, 13$  listed in Table B presented in Appendix.

#### 3. CLASSIFICATION OF 6-DIMENSIONAL ONE-GENERATED NILPOTENT NOVIKOV ALGEBRAS

The multiplication tables of 5-dimenensional one-generated nilpotent Novikov algebras are given in Table B presented in Appendix. All the necessary information about coboundaries, cocycles and second cohomology spaces of 5-dimensional one-generated nilpotent Novikov algebras is given in Table C presented in Appendix.

### 3.1. Central extensions of $\mathcal{N}_{01}^5$ . Let us use the following notations

$$\nabla_1 = [\Delta_{14}], \quad \nabla_2 = 2[\Delta_{13}] + [\Delta_{22}] + [\Delta_{41}], \quad \nabla_3 = 2[\Delta_{15}] - [\Delta_{23}] - [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}^5_{01}$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 \\ z & xy & x^3 & 0 & 0 \\ v & xy & 0 & x^3 & 0 \\ w & 0 & -x^2y & x^2y & x^4 \end{pmatrix}.$$

Since

$$\phi^{T} \begin{pmatrix} 0 & 0 & 2\alpha_{2} & \alpha_{1} & 2\alpha_{3} \\ 0 & \alpha_{2} & -\alpha_{3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \alpha_{2} & 0 & 0 & 0 & 0 \\ -\alpha_{3} & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & 2\alpha_{2}^{*} + \alpha^{*} & \alpha_{1}^{*} & 2\alpha_{3}^{*} \\ \alpha^{**} & \alpha_{2}^{*} & -\alpha_{3}^{*} & 0 & 0 \\ -\alpha^{*} & 0 & 0 & 0 & 0 \\ \alpha_{2}^{*} & 0 & 0 & 0 & 0 \\ -\alpha_{3}^{*} & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}_{01}^5)$  on the subspace  $\langle \sum_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^4 \alpha_1 + 2x^3 y \alpha_3, \quad \alpha_2^* = x^4 \alpha_2 - x^3 y \alpha_3, \quad \alpha_3^* = x^5 \alpha_3.$$

We assume  $\alpha_3 \neq 0$ , since otherwise we obtain algebras with 2-dimensional annihilator. Then we have the following cases:

- (1) if  $\alpha_1 = -2\alpha_2$ , then choosing  $y = \frac{\alpha_2 x}{\alpha_3}$ , we have the representative  $\langle \nabla_3 \rangle$ , which gives an algebra with 2-dimensional annihilator;
- (2) if  $\alpha_1 \neq -2\alpha_2$ , then choosing  $y = \frac{\alpha_2 x}{\alpha_3}$ ,  $x = \frac{\alpha_1 + 2\alpha_2}{\alpha_3}$ , we have the representative  $\langle \nabla_1 + \nabla_3 \rangle$ . Hence, we have only one new algebra:

$$\mathcal{N}_{13}^6$$
:  $e_1e_1 = e_2$   $e_1e_2 = e_4$   $e_1e_3 = e_5$   $e_1e_4 = e_6$   $e_1e_5 = 2e_6$   $e_2e_1 = e_3$   $e_2e_3 = -e_6$   $e_3e_1 = -e_5$   $e_5e_1 = -e_6$ 

## 3.2. Central extensions of $\mathcal{N}_{02}^5(\lambda)_{\lambda\neq 0}$ . Let us use the following notations

$$\nabla_1 = [\Delta_{14}] - [\Delta_{41}], \quad \nabla_2 = [\Delta_{13}] - \lambda[\Delta_{41}], \quad \nabla_3 = (3 - 2\lambda)[\Delta_{15}] + \lambda(2 - \lambda)[\Delta_{23}] + \lambda[\Delta_{32}] + \lambda[\Delta_{51}].$$

The automorphism group of  $\mathcal{N}^5_{02}(\lambda)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 & 0 \\ z & (\lambda+1)xy & x^3 & 0 & 0 \\ v & xy & 0 & x^3 & 0 \\ w & \lambda y^2 + 2xz & (-\lambda^2 + 2\lambda + 2)x^2y & \lambda^2(\lambda-1)x^2y & x^4 \end{pmatrix}.$$

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_2 & \alpha_1 & (3-2\lambda)\alpha_3 \\ 0 & 0 & \lambda(2-\lambda)\alpha_3 & 0 & 0 \\ 0 & \lambda\alpha_3 & 0 & 0 & 0 \\ -\alpha_1 - \lambda\alpha_2 & 0 & 0 & 0 & 0 \\ \lambda\alpha_3 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha_2^* + (2-\lambda)\alpha^* & \alpha_1^* & (3-2\lambda)\alpha_3^* \\ \alpha^{**} & \lambda\alpha^* & \lambda(2-\lambda)\alpha_3^* & 0 & 0 \\ \lambda\alpha^* & \lambda\alpha_3^* & 0 & 0 & 0 \\ -\alpha_1^* - \lambda\alpha_2^* & 0 & 0 & 0 & 0 \\ \lambda\alpha_3^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathcal{N}_{02}^5(\lambda)_{\lambda\neq 0})$  on the subspace  $\langle \sum_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^4 \alpha_1 + x^3 y \lambda^2 (\lambda - 1)(3 - 2\lambda) \alpha_3, \quad \alpha_2^* = x^4 \alpha_2 + x^3 y \lambda (\lambda - 1)(\lambda - 3) \alpha_3, \quad \alpha_3^* = x^5 \alpha_3.$$

We assume  $\alpha_3 \neq 0$  and  $(\alpha_1, \alpha_2) \neq (0, 0)$ . Then we have the following cases

- (1)  $\lambda = 1$ . Then
- (a) α₂ ≠ 0, then choosing x = α₂/α₃, we have the family of representatives ⟨α∇₁ + ∇₂ + ∇₃⟩;
  (b) α₂ = 0, then choosing x = α₁/α₃, we have the representative ⟨∇₁ + ∇₃⟩.
  (2) λ = 3. If α₂ ≠ 0, then choosing x = α₂/α₃, y = xα₁/54α₃ we have the representative ⟨∇₂ + ∇₃⟩. If α₂ = 0, then we have an algebra with 2-dimensional annihilator.
- (3)  $\lambda \notin \{1,3\}$ . Then in the case of  $\alpha_1 \neq \frac{\alpha_2 \lambda (3-2\lambda)}{\lambda-3}$ , choosing  $x = \frac{\alpha_1(\lambda-3) \alpha_2 \lambda (3-2\lambda)}{\alpha_3(\lambda-3)}$ ,  $y = \frac{-x\alpha_2}{\alpha_3 \lambda(\lambda-1)(\lambda-3)}$ , we have the representative  $\langle \nabla_1 + \nabla_3 \rangle$ . If  $\alpha_1 = \frac{\alpha_2 \lambda (3-2\lambda)}{\lambda-3}$ , then we have an algebra with 2-dimensional annihilator.

Thus, we have three representatives of distinct orbits  $\langle \alpha \nabla_1 + \nabla_2 + \nabla_3 \rangle_{\lambda=1}$ ,  $\langle \nabla_2 + \nabla_3 \rangle_{\lambda=3}$ ,  $\langle \nabla_1 + \nabla_3 \rangle_{(\lambda \notin \{0;3\})}$ , which give the following algebras:

3.3. Central extensions of  $\mathcal{N}_{02}^5(0)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{14}] - [\Delta_{41}], \quad \nabla_2 = 2[\Delta_{14}] + [\Delta_{22}] + [\Delta_{31}], \quad \nabla_3 = [\Delta_{15}], \quad \nabla_4 = 2[\Delta_{23}] - 2[\Delta_{32}] + [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{02}^5(0)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 \\ z & xy & x^3 & 0 & 0 \\ v & xy & 0 & x^3 & 0 \\ w & 2xz & 2x^2y & 0 & x^4 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & \alpha_1 + 2\alpha_2 & \alpha_3 \\ 0 & \alpha_2 & 2\alpha_4 & 0 & 0 \\ \alpha_2 & -2\alpha_4 & 0 & 0 & 0 \\ -\alpha_1 & 0 & 0 & 0 & 0 \\ \alpha_4 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha^{**} & \alpha_1^* + 2\alpha_2^* & \alpha_3^* \\ \alpha^* & \alpha_2^* & 2\alpha_4^* & 0 & 0 \\ \alpha_2^* & -2\alpha_4^* & 0 & 0 & 0 \\ -\alpha_1^* & 0 & 0 & 0 & 0 \\ \alpha_4^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{02}^5(0))$  on the subspace  $\langle \sum_{i=1}^4 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^4 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^4 \alpha_1, \quad \alpha_2^* = x^4 \alpha_2, \quad \alpha_3^* = x^5 \alpha_3, \quad \alpha_4^* = x^5 \alpha_4.$$

We assume  $(\alpha_1, \alpha_2) \neq (0, 0)$  and  $(\alpha_3, \alpha_4) \neq (0, 0)$ . Then we have the following cases:

- (1) if  $\alpha_2 = 0$ ,  $\alpha_4 = 0$ , then  $\alpha_1 \neq 0$ ,  $\alpha_3 \neq 0$  and choosing  $x = \frac{\alpha_1}{\alpha_3}$ , we have the representative  $\langle \nabla_1 + \nabla_3 \rangle$ ;
- (2) if  $\alpha_2 = 0$ ,  $\alpha_4 \neq 0$ , then  $\alpha_1 \neq 0$  and choosing  $x = \frac{\alpha_1}{\alpha_4}$ , we have the family of representatives  $\langle \nabla_1 + \alpha \nabla_3 + \nabla_4 \rangle$ ;

- (3) if  $\alpha_2 \neq 0$ ,  $\alpha_4 = 0$ , then  $\alpha_3 \neq 0$  and choosing  $x = \frac{\alpha_2}{\alpha_3}$ , we have the family of representatives  $\langle \alpha \nabla_1 + \nabla_2 + \nabla_3 \rangle$ ;
- (4) if  $\alpha_2 \neq 0$ ,  $\alpha_4 \neq 0$ , then choosing  $x = \frac{\alpha_2}{\alpha_4}$ , we have the family of representatives  $\langle \alpha \nabla_1 + \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle$ . Hence, we have the following new algebras:

# 3.4. Central extensions of $\mathcal{N}_{03}^5(\lambda,\mu)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{14}] - [\Delta_{41}],$$

$$\nabla_3 = (2\mu - 1)[\Delta_{15}] + \lambda(2-\mu)[\Delta_{23}] + \mu(2-\mu)[\Delta_{24}] + \Big(3 - (\lambda+1)(2-\mu)\Big)[\Delta_{32}] + (2-\mu)[\Delta_{42}] + (2-\mu)[\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{03}^5(\lambda,\mu)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 & 0 \\ z & xy & x^3 & 0 & 0 \\ v & xy & 0 & x^3 & 0 \\ w & (\lambda+1)xz + 2xv + y^2 & (\lambda+\mu+1)x^2y & (4-\mu)x^2y & x^4 \end{pmatrix}.$$

Since

$$\phi^{T} \begin{pmatrix} 0 & 0 & \alpha_{1} & \alpha_{2} & (2\mu - 1)\alpha_{3} \\ 0 & 0 & \lambda(2 - \mu)\alpha_{3} & \mu(2 - \mu)\alpha_{3} & 0 \\ 0 & (3 - (\lambda + 1)(2 - \mu))\alpha_{3} & 0 & 0 & 0 \\ -\alpha_{2} & (2 - \mu)\alpha_{3} & 0 & 0 & 0 \\ (2 - \mu)\alpha_{3} & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha_{1}^{*} + \lambda\alpha^{*} & \alpha_{2}^{*} + \mu\alpha^{*} & (2\mu - 1)\alpha_{3}^{*} \\ \alpha^{**} & \alpha^{*} & \lambda(2 - \mu)\alpha_{3}^{*} & \mu(2 - \mu)\alpha_{3}^{*} & 0 \\ \alpha^{*} & (3 - (\lambda + 1)(2 - \mu))\alpha_{3}^{*} & 0 & 0 & 0 \\ (2 - \mu)\alpha^{*} - \alpha_{2}^{*} & (2 - \mu)\alpha_{3}^{*} & 0 & 0 & 0 \\ (2 - \mu)\alpha_{3}^{*} & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{03}^5(\lambda,\mu))$  on the subspace  $\langle \sum\limits_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^4 \alpha_1 + x^3 y \alpha_3(\mu + 1) \Big( 2\mu - 1 + \lambda(\mu - 2) \Big), \quad \alpha_2^* = x^4 \alpha_2 + x^3 y \alpha_3(\mu - 1)(\mu - 2)^2, \quad \alpha_3^* = x^5 \alpha_3.$$

Assuming  $\alpha_3 \neq 0$ , we have the following cases:

- (1) if  $\mu \notin \{1, 2\}$ , then choosing  $y = -\frac{x\alpha_2}{(\mu 1)(\mu 2)^2\alpha_3}$ , we get  $\alpha_2^* = 0$  and have the representatives  $\langle \nabla_3 \rangle$  and  $\langle \nabla_1 + \nabla_3 \rangle$ , depending on whether  $\alpha_1 \frac{(\mu + 1)(2\mu 1 + \lambda(\mu 2))}{(\mu 1)(\mu 2)^2}\alpha_2 = 0$  or not;
- (2) if  $\mu = 1$ , then we consider following subcases
  - (a) if  $\lambda \neq 1$ , then choosing  $y = \frac{x\alpha_1}{2(\lambda-1)\alpha_3}$ , we have the representatives  $\langle \nabla_3 \rangle$  and  $\langle \nabla_2 + \nabla_3 \rangle$ , depending on whether  $\alpha_2 = 0$  or not;

- (b) if  $\lambda=1$ , then we get  $\alpha_1^*=x^4\alpha_1,\ \alpha_2^*=x^4\alpha_2,\ \alpha_3^*=x^5\alpha_3$  and have the representatives  $\langle\nabla_3\rangle,\langle\nabla_1+\nabla_3\rangle$  and  $\langle \nu \nabla_1 + \nabla_2 + \nabla_3 \rangle;$
- (3) if  $\mu=2$ , then choosing  $y=-\frac{x\alpha_1}{9\alpha_3}$ , we have the representatives  $\langle \nabla_3 \rangle$  and  $\langle \nabla_2 + \nabla_3 \rangle$ , depending on whether  $\alpha_2=0$  or

Thus, we obtain the following representatives of distinct orbits

$$\langle \nabla_3 \rangle$$
,  $\langle \nabla_1 + \nabla_3 \rangle_{\mu \neq 2}$ ,  $\langle \nabla_2 + \nabla_3 \rangle_{\mu=1}$ ,  $\langle \nu \nabla_1 + \nabla_2 + \nabla_3 \rangle_{\lambda=1,\mu=1,\nu\neq 0}$ ,  $\langle \nabla_2 + \nabla_3 \rangle_{\mu=2}$ ,

Hence, we have the following new algebras:

3.5. Central extensions of  $\mathcal{N}_{04}^5(\lambda)_{\lambda\neq 0}$ . Let us use the following notations

$$\nabla_1 = [\Delta_{14}] - [\Delta_{41}], \quad \nabla_2 = [\Delta_{22}] + [\Delta_{31}] + 2[\Delta_{41}], \quad \nabla_3 = -2[\Delta_{15}] + [\Delta_{23}] + \lambda[\Delta_{24}] - [\Delta_{32}] + [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{04}^5(\lambda)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 \\ z & xy & x^3 & 0 & 0 \\ v & xy & 0 & x^3 & 0 \\ w & xz & (\lambda + 1)x^2y & -\lambda x^2y & x^4 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & \alpha_1 & -2\alpha_3 \\ 0 & \alpha_2 & \alpha_3 & \lambda\alpha_3 & 0 \\ \alpha_2 & -\alpha_3 & 0 & 0 & 0 \\ 2\alpha_2 - \alpha_1 & 0 & 0 & 0 & 0 \\ \alpha_3 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha^* & \alpha_1^* + \lambda\alpha^* & -2\alpha_3^* \\ \alpha^{**} & \alpha_2^* & \alpha_3^* & \lambda\alpha_3^* & 0 \\ \alpha_2^* & -\alpha_3^* & 0 & 0 & 0 \\ 2\alpha_2^* - \alpha_1^* - \lambda\alpha^* & 0 & 0 & 0 \\ \alpha_3^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{04}^5(\lambda))$  on the subspace  $\langle \sum_{i=1}^{3} \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^{3} \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^4 \alpha_1 + 2x^3 y \lambda(\lambda + 2) \alpha_3, \quad \alpha_2^* = x^4 \alpha_2 + x^3 y \lambda \alpha_3, \quad \alpha_3^* = x^5 \alpha_3.$$

Assuming  $\alpha_3 \neq 0$ , we have the following cases:

- (1) if  $\alpha_1=2(\lambda+2)\alpha_2$ , then choosing  $y=-\frac{x\alpha_2}{\lambda\alpha_3}$ , we have the representative  $\langle\nabla_3\rangle$ ; (2) if  $\alpha_1\neq 2(\lambda+2)\alpha_2$ , then choosing  $x=\frac{\alpha_1-2(\lambda+2)\alpha_2}{\alpha_3}$ ,  $y=-\frac{x\alpha_2}{\lambda\alpha_3}$ , we have the representative  $\langle\nabla_1+\nabla_3\rangle$ .

Hence, we have the following new algebras:

## 3.6. Central extensions of $\mathcal{N}_{05}^5$ . Let us use the following notations

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = 3[\Delta_{15}] - 2[\Delta_{24}] + [\Delta_{33}] - [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{05}^5$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 & 0 \\ y & 0 & x^3 & 0 & 0 \\ z & 0 & 0 & x^4 & 0 \\ v & xz & -x^2y & 0 & x^5 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & 0 & 0 & 3\alpha_2 \\ 0 & 0 & 0 & -2\alpha_2 & 0 \\ 0 & 0 & \alpha_2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -\alpha_2 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha_1^* & -\alpha^* & 0 & 3\alpha_2^* \\ \alpha^{**} & 0 & 0 & -2\alpha_2^* & 0 \\ \alpha^* & 0 & \alpha_2^* & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -\alpha_2^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}_{05}^5)$  on the subspace  $\langle \sum_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^3 \alpha_1 + 3x^2 z \alpha_2, \quad \alpha_2^* = x^6 \alpha_2.$$

It is easy to see that we have only one non-trivial orbit with the representative  $\langle \nabla_2 \rangle$  and obtain the algebra:

$$\mathcal{N}_{27}^6$$
:  $e_1e_1 = e_2$   $e_1e_3 = e_4$   $e_1e_4 = 2e_5$   $e_1e_5 = 3e_6$   $e_2e_1 = e_3$   $e_2e_3 = -e_5$   $e_2e_4 = -2e_6$   $e_3e_1 = -e_4$   $e_3e_3 = e_6$   $e_4e_1 = -e_5$   $e_5e_1 = -e_6$ 

# 3.7. Central extensions of $\mathcal{N}_{06}^5$ . Let us use the following notations

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = 3[\Delta_{15}] - [\Delta_{22}] - 2[\Delta_{24}] - 2[\Delta_{31}] + [\Delta_{33}] - [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{06}^5$  consists of invertible matrices of the form

$$\phi_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ x & 0 & 1 & 0 & 0 \\ y & 0 & 0 & 1 & 0 \\ z & y & -x & 0 & 1 \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ x & 0 & -1 & 0 & 0 \\ y & 0 & 0 & 1 & 0 \\ z & -y & -x & 0 & -1 \end{pmatrix}.$$

$$\phi_i^T \begin{pmatrix} 0 & \alpha_1 & 0 & 0 & 3\alpha_2 \\ 0 & -\alpha_2 & 0 & -2\alpha_2 & 0 \\ -2\alpha_2 & 0 & \alpha_2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -\alpha_2 & 0 & 0 & 0 & 0 \end{pmatrix} \phi_i = \begin{pmatrix} \alpha^{***} & \alpha_1^* & \alpha^* & 0 & 3\alpha_2^* \\ \alpha^{**} & -2\alpha_2^* & 0 & -2\alpha_2^* & 0 \\ -2\alpha_2^* - \alpha^* & 0 & \alpha_2^* & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -\alpha_2^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{06}^5)$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = (-1)^i \alpha_1 + 3y\alpha_2, \quad \alpha_2^* = \alpha_2.$$

It is easy to see that we have only one non-trivial orbit with the representative  $\langle \nabla_2 \rangle$  and obtain the algebra:

3.8. Central extensions of  $\mathcal{N}_{07}^5(\lambda)_{\lambda \notin \{0,1\}}$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = (4 - 3\lambda)[\Delta_{15}] + \lambda(3 - 2\lambda)[\Delta_{24}] + \lambda(2 - \lambda)[\Delta_{33}] + \lambda[\Delta_{42}] + \lambda[\Delta_{51}].$$

The automorphism group of  $\mathfrak{N}^5_{07}(\lambda)_{\lambda \neq \{0,1\}}$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 & 0 \\ 0 & 0 & x^3 & 0 & 0 \\ y & 0 & 0 & x^4 & 0 \\ z & (3-\lambda)xy & 0 & 0 & x^5 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & 0 & (4-3\lambda)\alpha_2 \\ \alpha_1 & 0 & 0 & \lambda(3-2\lambda)\alpha_2 & 0 \\ 0 & 0 & \lambda(2-\lambda)\alpha_2 & 0 & 0 \\ 0 & \lambda\alpha_2 & 0 & 0 & 0 \\ \lambda\alpha_2 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \\ = \begin{pmatrix} \alpha^{**} & \alpha^* & 0 & 0 & (4-3\lambda)\alpha_2^* \\ \alpha_1^* + \lambda\alpha^* & 0 & 0 & \lambda(3-2\lambda)\alpha_2^* & 0 \\ 0 & 0 & \lambda(2-\lambda)\alpha_2^* & 0 & 0 \\ 0 & \lambda\alpha_2^* & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}^5_{07}(\lambda)_{\lambda \not\in \{0,1\}})$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^3 \alpha_1 - 3x^2 y \lambda (\lambda - 1)(\lambda - 2)\alpha_2, \quad \alpha_2^* = x^6 \alpha_2.$$

Assuming  $\alpha_2 \neq 0$ , we have the following cases:

- (1) if  $\lambda \neq 2$ , then choosing  $y = \frac{x\alpha_1}{3\lambda(\lambda-1)(\lambda-2)\alpha_2}$ , we have the representative  $\langle \nabla_2 \rangle$ ;
- (2) if  $\lambda=2$ , then we have two representatives  $\langle \nabla_2 \rangle$  and  $\langle \nabla_1 + \nabla_2 \rangle$  depending on whether  $\alpha_1=0$  or not. Hence, we have the following new algebras:

3.9. Central extensions of  $\mathcal{N}_{07}^5(0)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{15}], \quad \nabla_2 = [\Delta_{21}], \quad \nabla_3 = 2[\Delta_{23}] - 2[\Delta_{32}] + [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{07}^5(0)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 \\ z & xy & x^3 & 0 & 0 \\ v & 2xz & 2x^2y & x^4 & 0 \\ w & 3xv & 6x^2z & 3x^3y & x^5 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & 0 & \alpha_1 \\ \alpha_2 & 0 & 2\alpha_3 & 0 & 0 \\ 0 & -2\alpha_3 & 0 & 0 & 0 \\ \alpha_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha^{**} & \alpha^* & \alpha_1^* \\ \alpha_2^* & 0 & 2\alpha_3^* & 0 & 0 \\ 0 & -2\alpha_3^* & 0 & 0 & 0 \\ \alpha_3^* & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{07}^5(0))$  on the subspace  $\langle \sum\limits_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^6 \alpha_1, \quad \alpha_2^* = x^3 \alpha_2 + 2x(2xz - y^2)\alpha_3, \quad \alpha_3^* = x^5 \alpha_3.$$

Assuming  $\alpha_1 \neq 0$ , we have the following cases:

- (1) if  $\alpha_3 \neq 0$ , then choosing  $x = \frac{\alpha_3}{\alpha_1}, y = 0, z = -\frac{x\alpha_2}{4\alpha_3}$ , we have the representative  $\langle \nabla_1 + \nabla_3 \rangle$ ;
- (2) if  $\alpha_3 = 0, \alpha_2 \neq 0$ , then choosing  $x = \sqrt[3]{\alpha_2/\alpha_1}$ , we have the representative  $\langle \nabla_1 + \nabla_2 \rangle$ ;
- (3) if  $\alpha_3 = \alpha_2 = 0$ , then we have the representative  $\langle \nabla_1 \rangle$ .

Hence, we have the following new algebras:

3.10. Central extensions of  $\mathcal{N}_{07}^5(1)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = [\Delta_{15}] + [\Delta_{24}] + [\Delta_{33}] + [\Delta_{42}] + [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{07}^5(1)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 & 0 \\ y & x^2 & 0 & 0 & 0 \\ z & 2xy & x^3 & 0 & 0 \\ v & 2xz + y^2 & 3x^2y & x^4 & 0 \\ w & 2xv + 2yz & 3x^2z + 3xy^2 & 4x^3y & x^5 \end{pmatrix}.$$

$$\phi^{T} \begin{pmatrix} 0 & 0 & 0 & 0 & \alpha_{2} \\ \alpha_{1} & 0 & 0 & \alpha_{2} & 0 \\ 0 & 0 & \alpha_{2} & 0 & 0 \\ 0 & \alpha_{2} & 0 & 0 & 0 \\ \alpha_{2} & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha^{**} & \alpha^{*} & \alpha_{2}^{*} \\ \alpha_{1}^{*} + \alpha^{***} & \alpha^{**} & \alpha^{*} & \alpha_{2}^{*} & 0 & 0 \\ \alpha^{**} & \alpha^{*} & \alpha_{2}^{*} & 0 & 0 & 0 \\ \alpha_{2}^{*} & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}_{07}^5(1))$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = x^3 \alpha_1, \quad \alpha_2^* = x^6 \alpha_2.$$

Assuming  $\alpha_2 \neq 0$ , we have the following cases:

- (1) if  $\alpha_1 = 0$ , then we have the representative  $\langle \nabla_2 \rangle$ ;
- (2) if  $\alpha_1 \neq 0$ , then choosing  $x = \sqrt[3]{\alpha_1/\alpha_2}$ , we have the representative  $\langle \nabla_1 + \nabla_2 \rangle$ . Hence, we have the algebras:

# 3.11. Central extensions of $\mathcal{N}_{08}^5$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = 2[\Delta_{15}] + [\Delta_{22}] + [\Delta_{31}], \quad \nabla_3 = [\Delta_{23}] - [\Delta_{32}] + [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{08}^5$  consists of invertible matrices of the form

$$\phi_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ x & 1 & 0 & 0 & 0 \\ y & x & 1 & 0 & 0 \\ z & y & x & 1 & 0 \\ v & x+z & y & x & 1 \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ x & 1 & 0 & 0 & 0 \\ y & -x & -1 & 0 & 0 \\ z & -y & x & 1 & 0 \\ v & -x-z & y & -x & -1 \end{pmatrix}.$$

Since

$$\phi_i^T \begin{pmatrix} 0 & 0 & 0 & 0 & 2\alpha_2 \\ \alpha_1 & \alpha_2 & \alpha_3 & 0 & 0 \\ \alpha_2 & -\alpha_3 & 0 & 0 & 0 \\ \alpha_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \phi_i = \begin{pmatrix} \alpha^{****} & \alpha^{***} & \alpha^{**} & \alpha^* & 2\alpha_2^* \\ \alpha_1^* + \alpha^* & \alpha_2^* & \alpha_3^* & 0 & 0 \\ \alpha_2^* & -\alpha_3^* & 0 & 0 & 0 \\ \alpha_3^* & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}^5_{08})$  on the subspace  $\langle \sum\limits_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\begin{array}{lll} \text{for} \ i=1: & \alpha_1^*=\alpha_1-(x^2-2y)\alpha_3, & \alpha_2^*=\alpha_2, & \alpha_3^*=\alpha_3, \\ \text{for} \ i=2: & \alpha_1^*=-\alpha_1+(x^2+2y)\alpha_3, & \alpha_2^*=\alpha_2, & \alpha_3^*=-\alpha_3. \end{array}$$

Assuming  $\alpha_2 \neq 0$  and using  $\phi_1$  we have the following cases:

- (1) if  $\alpha_3 = 0$ , then we have the family of representatives  $\langle \lambda \nabla_1 + \nabla_2 \rangle$ ;
- (2) if  $\alpha_3 \neq 0$ , then choosing  $y = \frac{x^2\alpha_3 \alpha_1}{2\alpha_3}$ , we have the family of representatives  $\langle \nabla_2 + \lambda \nabla_3 \rangle_{\lambda \neq 0}$ . Hence, we have the following new algebras:

Moreover, using the action of  $\phi_2$  we have that  $\mathcal{N}_{34}^6(\lambda) \cong \mathcal{N}_{34}^6(-\lambda)$  and  $\mathcal{N}_{35}^6(\lambda) \cong \mathcal{N}_{35}^6(-\lambda)$ .

3.12. Central extensions of  $\mathcal{N}_{10}^5$ . Let us use the following notations

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = -2[\Delta_{13}] + [\Delta_{15}] + [\Delta_{24}] + [\Delta_{33}] + [\Delta_{42}] + [\Delta_{51}].$$

The automorphism group of  $\mathcal{N}_{10}^5$  consists of invertible matrices of the form

$$\phi_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ x & 1 & 0 & 0 & 0 & 0 \\ y & 2x & 1 & 0 & 0 \\ z & x^2 + 2y & 3x & 1 & 0 \\ v & 2xy + x + 2z & 3x^2 + 3y & 4x & 1 \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ x & 1 & 0 & 0 & 0 & 0 \\ y & -2x & -1 & 0 & 0 & 0 \\ z & x^2 - 2y & 3x & 1 & 0 \\ v & 2xy - x - 2z & -3x^2 + 3y & -4x & -1 \end{pmatrix}.$$

Since

$$\phi_i^T \begin{pmatrix} 0 & \alpha_1 & -2\alpha_2 & 0 & \alpha_2 \\ 0 & 0 & 0 & \alpha_2 & 0 \\ 0 & 0 & \alpha_2 & 0 & 0 \\ 0 & \alpha_2 & 0 & 0 & 0 \\ \alpha_2 & 0 & 0 & 0 & 0 \end{pmatrix} \phi_i = \begin{pmatrix} \alpha^{****} & \alpha_1^* + \alpha^{***} & -2\alpha_2^* + \alpha^{**} & \alpha^* & \alpha_2^* \\ \alpha^{***} & \alpha^{**} & \alpha^{**} & \alpha_2^* & 0 \\ \alpha^{**} & \alpha_2^* & \alpha_2^* & 0 & 0 \\ \alpha_2^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathbb{N}^5_{10})$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = (-1)^i \alpha_1 - 4x\alpha_2, \quad \alpha_2^* = \alpha_2.$$

It is easy to see that we have only one non-trivial orbit with representative  $\langle \nabla_2 \rangle$ . The corresponding algebra is

$$\mathcal{N}_{36}^6$$
:  $e_1e_1 = e_2$   $e_1e_2 = e_3$   $e_1e_3 = e_4 - 2e_6$   $e_1e_4 = e_5$   $e_1e_5 = e_6$   $e_2e_1 = e_3 + e_5$   $e_2e_2 = e_4$   $e_2e_3 = e_5$   $e_2e_4 = e_6$   $e_3e_1 = e_4$   $e_3e_2 = e_5$   $e_3e_3 = e_6$   $e_4e_1 = e_5$   $e_4e_2 = e_6$   $e_5e_1 = e_6$ 

3.13. Central extensions of  $\mathcal{N}_{11}^5(\lambda)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{15}] + [\Delta_{32}], \quad \nabla_3 = [\Delta_{14}] - 2[\Delta_{23}] + 2[\Delta_{32}] - [\Delta_{41}].$$

The automorphism group of  $\mathcal{N}_{11}^5(\lambda)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ x & 1 & 0 & 0 & 0 & 0 \\ y & x & 1 & 0 & 0 \\ z & x + 2y & 2x & 1 & 0 \\ v & x^2 + (\lambda + 1)y + 2z & (\lambda + 3)x + 4y & 2x & 1 \end{pmatrix}.$$

Since

we have that the action of  $\operatorname{Aut}(\mathfrak{N}^5_{11}(\lambda))$  on the subspace  $\langle \sum\limits_{i=1}^3 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^3 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = \alpha_1 + (3x + 4y - 2x^2)\alpha_2 + (8y + 2x - 4x^2)\alpha_3, \quad \alpha_2^* = \alpha_2, \quad \alpha_3^* = \alpha_3.$$

Assuming  $\alpha_2 \neq 0$ , we have the follofing cases

(1) if  $\alpha_2 \neq -2\alpha_3$ , then choosing  $y = -\frac{\alpha_1 + x(3\alpha_2 + 2\alpha_3) - 2x^2(\alpha_2 + 2\alpha_3)}{4(\alpha_2 + 2\alpha_3)}$ , we have the family of orbits with the representatives  $\langle \nabla_2 + \mu \nabla_3 \rangle_{\mu \neq -\frac{1}{2}}$ ;

(2) if  $\alpha_2 = -2\alpha_3$ , then choosing  $x = -\frac{\alpha_1}{2\alpha_2}$ , we have the family of orbits with the representatives  $\langle \nabla_2 - \frac{1}{2} \nabla_3 \rangle$ . Thus, we obtain the family of algebras:

3.14. Central extensions of  $\mathcal{N}_{13}^5(\lambda)$ . Let us use the following notations

$$\nabla_1 = [\Delta_{21}], \quad \nabla_2 = (1 - 2\lambda)[\Delta_{13}] - 2[\Delta_{14}] + [\Delta_{15}] - [\Delta_{23}] + [\Delta_{24}] + [\Delta_{33}] + [\Delta_{41}] + [\Delta_{42}] + [\Delta_{51}].$$

The automorphism group of  $\mathfrak{N}_{13}^5(\lambda)$  consists of invertible matrices of the form

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ x & 1 & 0 & 0 & 0 & 0 \\ y & 2x & 1 & 0 & 0 & 0 \\ z & x^2 + x + 2y & 3x & 1 & 0 \\ v & \lambda x + 2xy - 2y + 2z & 3x^2 - 3x + 3y & 4x & 1 \end{pmatrix}.$$

Since

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & (1-2\lambda)\alpha_2 & -2\alpha_2 & \alpha_2 \\ 0 & 0 & -\alpha_2 & \alpha_2 & 0 \\ 0 & 0 & \alpha_2 & 0 & 0 \\ \alpha_2 & \alpha_2 & 0 & 0 & 0 \\ \alpha_2 & 0 & 0 & 0 & 0 \end{pmatrix} \phi = \\ \begin{pmatrix} \alpha^{****} & \alpha_1^* + \alpha^{***} & (1-2\lambda)\alpha_2^* + \alpha^{**} - 2\alpha^* & -2\alpha_2^* + \alpha^* & \alpha_2^* \\ \alpha^{***} + \alpha^{**} + \lambda\alpha^* & \alpha^{**} & -\alpha_2^* + \alpha^* & \alpha_2^* & 0 \\ \alpha^{**} & \alpha^* & \alpha_2^* & 0 & 0 \\ \alpha_2^* + \alpha^* & \alpha_2^* & 0 & 0 & 0 \\ \alpha_2^* & 0 & 0 & 0 & 0 \end{pmatrix},$$

we have that the action of  $\operatorname{Aut}(\mathfrak{N}^5_{13}(\lambda))$  on the subspace  $\langle \sum\limits_{i=1}^2 \alpha_i \nabla_i \rangle$  is given by  $\langle \sum\limits_{i=1}^2 \alpha_i^* \nabla_i \rangle$ , where

$$\alpha_1^* = \alpha_1 - (\lambda x - x^2 + x + y)\alpha_2, \quad \alpha_2^* = \alpha_2.$$

It is easy to see that we have only one non-trivial orbit with the representative  $\langle \nabla_2 \rangle$  and obtain the family of algebras:

Summarizing the results of the present section and using the information from Tables B and C from Appendix, we have the following theorem.

**Theorem 7.** Let  $\mathbb{N}$  be a 6-dimensional complex one-generated nilpotent Novikov algebra. Then  $\mathbb{N}$  is isomorphic to  $\mathbb{N}_j^6$ , where  $j = 01, \ldots, 38$  (see Table D presented in Appendix).

### APPENDIX

**Table A.** The list of cohomology spaces of 4-dimensional one-generated nilpotent Novikov algebras.

```
Z^2 \left( \mathcal{N}_{01}^4 \right)
                                                                    \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13}, \Delta_{14} - \Delta_{41}, \Delta_{22} + \Delta_{31} + 2\Delta_{41}
B^2 (\mathcal{N}_{01}^4)
                                                                   \Delta_{11}, \Delta_{12}, \Delta_{21}
                                                    =
\mathrm{H}^2\left(\mathcal{N}_{01}^4\right)
                                                                    [\Delta_{13}], [\Delta_{14}] - [\Delta_{41}], [\Delta_{22}] + [\Delta_{31}] + 2[\Delta_{41}]
Z^2 \left( \mathcal{N}_{02}^4 \right)
                                                                   \Delta_{11}, \Delta_{12}, \Delta_{13} - \Delta_{31}, \Delta_{21}, 2\Delta_{14} - \Delta_{23} - \Delta_{41}
                                                    =
B^2 (N_{02}^4)
                                                                   \Delta_{11}, \Delta_{13} - \Delta_{31}, \Delta_{21}
                                                    =
H^2 (N_{02}^4)
                                                                   [\Delta_{12}], 2[\Delta_{14}] - [\Delta_{23}] - [\Delta_{41}]
                                                                     \Delta_{11}, \Delta_{12}, \Delta_{21}, (2-\lambda)\Delta_{13} + \lambda\Delta_{22} + \lambda\Delta_{31},
Z^2 \left( \mathcal{N}_{03}^4(\lambda)_{\lambda \notin \{0,1\}} \right)
                                                                    (3-2\lambda)\Delta_{14}+(2-\lambda)\lambda\Delta_{23}+\lambda\Delta_{32}+\lambda\Delta_{41}
\mathrm{B}^2\left(\mathcal{N}_{03}^4(\lambda)_{\lambda \not\in \{0,1\}}\right)
                                                                   \Delta_{11}, \Delta_{12} + \lambda \Delta_{21}, (2-\lambda)\Delta_{13} + \lambda \Delta_{22} + \lambda \Delta_{31}
                                                    =
\mathrm{H}^2 \left( \mathcal{N}_{03}^4(\lambda)_{\lambda \notin \{0,1\}} \right)
                                                                    [\Delta_{21}], (3-2\lambda)\Delta_{14} + (2-\lambda)\lambda\Delta_{23} + \lambda\Delta_{32} + \lambda\Delta_{41}
Z^2\left(\mathcal{N}_{03}^4(0)\right)
                                                                   \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14}, \Delta_{21}, 2\Delta_{23} - 2\Delta_{32} + \Delta_{41}
B^{2}(\mathcal{N}_{03}^{4}(0))
                                                                    \Delta_{11}, \Delta_{12}, \Delta_{13}
                                                    =
H^2\left(\mathcal{N}_{03}^4(0)\right)
                                                                    [\Delta_{14}], [\Delta_{21}], 2[\Delta_{23}] - 2[\Delta_{32}] + [\Delta_{41}]
Z^2\left(\mathcal{N}_{03}^4(1)\right)
                                                                   \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13} + \Delta_{22} + \Delta_{31}, \Delta_{14} + \Delta_{23} + \Delta_{32} + \Delta_{41}
                                                    =
B^2 (\mathcal{N}_{03}^4(1))
                                                                   \Delta_{11}, \Delta_{12} + \Delta_{21}, \Delta_{13} + \Delta_{22} + \Delta_{31}
                                                    =
\mathrm{H}^2\left(\underline{\mathcal{N}_{03}^4(1)}\right)
                                                                    [\Delta_{21}], [\Delta_{14}] + [\Delta_{23}] + [\Delta_{32}] + [\Delta_{41}]
Z^2 \left( N_{04}^4 \right)
                                                                    \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{21}, 2\Delta_{14} + \Delta_{22} + \Delta_{31}, \Delta_{14} - 2\Delta_{23} + 2\Delta_{32} - \Delta_{41}
                                                    =
B^2 \left( \widetilde{\mathcal{N}_{04}^4} \right)
                                                                   \Delta_{11}, \Delta_{12}, 2\Delta_{13} + \Delta_{21}
                                                    =
H^2(N_{04}^4)
                                                    =
                                                                    [\Delta_{13}], 2[\Delta_{14}] + [\Delta_{22}] + [\Delta_{31}], [\Delta_{14}] - 2[\Delta_{23}] + 2[\Delta_{32}] - [\Delta_{41}]
Z^2 \left( \mathcal{N}_{05}^4 \right)
                                                                    \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13} + \Delta_{22} + \Delta_{31}, -2\Delta_{13} + \Delta_{14} + \Delta_{23} + \Delta_{32} + \Delta_{41}
                                                    =
B^2 \left( \widetilde{\mathcal{N}_{05}^4} \right)
                                                                    \Delta_{11}, \Delta_{12} + \Delta_{21}, \Delta_{13} + \Delta_{21} + \Delta_{22} + \Delta_{31}
\mathrm{H}^2\left(\mathcal{N}_{05}^4\right)
                                                                    [\Delta_{21}], -2[\Delta_{13}] + [\Delta_{14}] + [\Delta_{23}] + [\Delta_{32}] + [\Delta_{41}]
                                                    =
```

<b>Table B.</b> The list of 5-dimensional one-generated nilpotent Novikov algebras.							
$\mathcal{N}_{01}^{5}$	:	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_1e_3 = e_5$	$e_2e_1=e_3$	$e_3e_1 = -e_5$	
$\mathcal{N}^5_{02}(\lambda)$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = (2 - \lambda)e_5$			
		$e_2e_1 = \lambda e_3 + e_4$	$e_2e_2 = \lambda e_5$	$e_3e_1 = \lambda e_5$			
$\mathcal{N}_{03}^5(\lambda,\mu)$	:	$e_1e_1=e_2$	$e_1e_2=e_3$	$e_1e_3 = \lambda e_5$	$e_1e_4 = \mu e_5$		
		$e_2e_1=e_4$	$e_2e_2=e_5$	$e_3e_1=e_5$	$e_4e_1 = (2 - \mu)e_5$		
$\mathcal{N}_{04}^5(\lambda)_{\lambda \neq 0}$	:	$e_1e_1=e_2$	$e_1e_2=e_3$	$e_1e_3=e_5$	$e_1e_4 = \lambda e_5$		
		$e_2e_1=e_4$	$e_4e_1 = -\lambda e_5$				
$\mathcal{N}_{05}^{5}$	:	$e_1e_1=e_2$	$e_1e_3=e_4$	$e_1e_4 = 2e_5$	$e_2e_1=e_3$		
		$e_2e_3=-e_5$	$e_3e_1 = -e_4$	$e_4e_1 = -e_5$			
$\mathcal{N}_{06}^{5}$	:	$e_1e_1=e_2$	$e_1e_2 = e_5$	$e_1e_3=e_4$	$e_1e_4 = 2e_5$		
		$e_2e_1=e_3$	$e_2e_3=-e_5$	$e_3e_1 = -e_4$			
$\mathcal{N}^5_{07}(\lambda)$	:	$e_1e_1=e_2$	$e_1e_2 = e_3$			_	
		$e_2 e_2 = \lambda e_4$	$e_2e_3 = \lambda(2-\lambda)e_5$	$e_3e_1 = \lambda e_4$	$e_3e_2 = \lambda e_5$	$e_4 e_1 = \lambda e_5$	
$\mathcal{N}_{08}^{5}$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$e_1e_4 = e_5$	$e_2e_1=e_5$	
$\mathcal{N}_{09}^5(\lambda)$	:	$e_1e_1=e_2$	$e_1e_2=e_3$	$e_1e_3 = 2e_4$	$e_1e_4 = \lambda e_5$		
		$e_2e_3=2e_5$	$e_3e_2 = -2e_5$	$e_4e_1=e_5$			
$\mathcal{N}_{10}^5$	:	$e_1e_1=e_2$	$e_1e_2=e_3$	$e_1e_3=e_4$	$e_1e_4 = e_5$	$e_2e_1 = e_3 + e_5$	
		$e_2e_2=e_4$	$e_2e_3=e_5$	$e_3e_1=e_4$		$e_4e_1=e_5$	
$\mathcal{N}_{11}^5(\lambda)$	:	$e_1e_1=e_2$	$e_1e_2=e_3$	$e_1e_3 = 2e_4 + \lambda e_5$	$e_1e_4 = 2e_5$		
.,,		$e_2e_1=e_4$	$e_2e_2=e_5$	$e_3e_1=e_5$			
$\mathcal{N}_{12}^5(\lambda)$	:	$e_1e_1=e_2$	$e_1e_2=e_3$	$e_1e_3 = 2e_4$	$e_1 e_4 = (2\lambda + 1)e_5$	$e_2e_1=e_4$	

		$e_2e_2 = \lambda e_5$	$e_2e_3 = -2e_5$	$e_3e_1 = \lambda e_5$	$e_3e_2 = 2e_5$	$e_4e_1 = -e_5$
$\mathcal{N}_{13}^5(\lambda)$	:	$e_1e_1 = e_2$	$e_1e_2=e_3$	$e_1 e_3 = e_4 - 2e_5$	$e_1e_4 = e_5$	$e_2 e_1 = e_3 + e_4 + \lambda e_5$
		$e_2e_2=e_4$	$e_2 e_3 = e_5$	$e_3e_1 = e_4$	$e_3 e_2 = e_5$	$e_4 e_1 = e_5$

$$\mathcal{N}_{04}^5(0) \cong \mathcal{N}_{02}^5(0).$$

```
        Table C. The list of cohomology spaces of 5-dimensional one-generated nilpotent Novikov algebras.

\overline{\mathrm{Z}^2}\left(\mathbb{N}_{01}^5\right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13} - \Delta_{31}, \Delta_{14}, \Delta_{21}, 2\Delta_{13} + \Delta_{22} + \Delta_{41}, 2\Delta_{15} - \Delta_{23} - \Delta_{51}
B^{2}(N_{01}^{5})
                                                                                            \begin{array}{l} \Delta_{11}, \Delta_{12}, \Delta_{13} - \Delta_{31}, \Delta_{21} \\ [\Delta_{14}], 2[\Delta_{13}] + [\Delta_{22}] + [\Delta_{41}], 2[\Delta_{15}] - [\Delta_{23}] - [\Delta_{51}] \end{array} \rangle
                                                                    =
H^2(N_{01}^5)
                                                                                                \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{\overline{14} - \Delta_{41}, \Delta_{13} - \lambda \Delta_{41}, \Delta_{22} + \Delta_{31} + (2 - \lambda)\Delta_{41},
Z^2 \left( \mathcal{N}_{02}^5(\lambda)_{\lambda \neq 0} \right)
                                                                                               (3-2\lambda)\Delta_{15} + \lambda(2-\lambda)\Delta_{23} + \lambda\Delta_{32} + \lambda\Delta_{51}
B^2 \left( \mathcal{N}_{02}^5(\lambda)_{\lambda \neq 0} \right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{21}, (2-\lambda)\Delta_{13} + \lambda\Delta_{22} + \lambda\Delta_{31}
                                                                                             \frac{\left[\Delta_{14}\right] - \left[\Delta_{41}\right], \left[\Delta_{13}\right] - \lambda\left[\Delta_{41}\right], \left(3 - 2\lambda\right)\left[\Delta_{15}\right] + \lambda(2 - \lambda)\left[\Delta_{23}\right] + \lambda\left[\Delta_{32}\right] + \lambda\left[\Delta_{51}\right]}{\Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{14} - \Delta_{41}, \Delta_{13}, 2\Delta_{14} + \Delta_{22} + \Delta_{31}, \Delta_{15}, 2\Delta_{23} - 2\Delta_{32} + \Delta_{51}}
\mathrm{H}^2\left(\mathcal{N}_{02}^5(\lambda)_{\lambda\neq 0}\right)
Z^{2}\left(\mathcal{N}_{02}^{5}(0)\right)
B^2 \left( \mathcal{N}_{02}^{\overline{5}}(0) \right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13}
                                                                                             \frac{\left[\Delta_{14}\right] - \left[\Delta_{41}\right], 2\left[\Delta_{14}\right] + \left[\Delta_{22}\right] + \left[\Delta_{31}\right], \left[\Delta_{15}\right], 2\left[\Delta_{23}\right] - 2\left[\Delta_{32}\right] + \left[\Delta_{51}\right] \ \rangle}{\Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13}, \Delta_{14} - \Delta_{41}, \Delta_{22} + \Delta_{31} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2 - \mu)\Delta_{23} + 2\Delta_{41}, (2\mu - 1)\Delta_{15} + \lambda(2\mu -
B^{2}\left(\mathcal{N}_{02}^{5}(0)\right)
Z^2\left(\mathcal{N}_{03}^5(\lambda,\mu)\right)
                                                                                               \mu(2-\mu)\Delta_{24} + (3-(\lambda+1)(2-\mu))\Delta_{32} + (2-\mu)\Delta_{42} + (2-\mu)\Delta_{51}
B^2\left(\mathcal{N}_{03}^5(\lambda,\mu)\right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{21}, \lambda \dot{\Delta}_{13} + \mu \Delta_{14} + \Delta_{22} + \Delta_{31} + (2 - \mu) \Delta_{41}
                                                                                               [\Delta_{13}], [\Delta_{14}] - [\Delta_{41}], (2\mu - 1)[\Delta_{15}] + \lambda(2 - \mu)[\Delta_{23}] + \mu(2 - \mu)[\Delta_{24}] + \mu(2 - \mu)[\Delta_{13}]
\mathrm{H}^2\left(\mathcal{N}_{03}^5(\lambda,\mu)\right)
                                                                                                 (3-(\lambda+1)(2-\mu))[\Delta_{32}]+(2-\mu)[\Delta_{42}]+(2-\mu)[\Delta_{51}]
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{21}, \Delta_{14} - \Delta_{41}, \Delta_{22} + \Delta_{31} + 2\Delta_{41}, -2\Delta_{15} + \Delta_{23} + \lambda\Delta_{24} - \Delta_{32} + \Delta_{51}
Z^2 \left( \mathcal{N}_{04}^5(\lambda)_{\lambda \neq 0} \right)
B^2 \left( \mathcal{N}_{04}^5(\lambda)_{\lambda \neq 0} \right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13} + \lambda(\Delta_{14} - \Delta_{41})
                                                                                            \mathrm{H}^2\left(\mathcal{N}_{04}^5(\lambda)_{\lambda\neq 0}\right)
                                                                                             \Delta_{11}, \Delta_{12}, \overline{\Delta_{13}, \Delta_{15}, \Delta_{21}, \Delta_{14}} - \Delta_{41}, 2\Delta_{14} + \Delta_{22} + \Delta_{31}, 2\Delta_{23} - 2\Delta_{32} + \Delta_{51}
 Z^2\left(\mathcal{N}_{04}^5(0)\right)
B^2\left(\mathcal{N}_{04}^5(0)\right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{21}
H^2(N_{04}^5(0))
                                                                                             [\Delta_{15}], [\Delta_{14}] - [\Delta_{41}], 2[\Delta_{14}] + [\Delta_{22}] + [\Delta_{31}], 2[\Delta_{23}] - 2[\Delta_{32}] + [\Delta_{51}]
                                                                                            \Delta_{11}, \Delta_{12}, \Delta_{13} - \Delta_{31}, 2\Delta_{14} - \Delta_{23} - \Delta_{41}, \Delta_{21}, 3\Delta_{15} - 2\Delta_{24} + \Delta_{33} - \Delta_{51}
\overline{\mathrm{Z}^{2}}\left( \mathcal{N}_{05}^{5}\right)
B^2\left(\mathcal{N}_{05}^5\right)
                                                                    =
                                                                                             \Delta_{11}, \Delta_{21}, \Delta_{13} - \Delta_{31}, 2\Delta_{14} - \Delta_{23} - \Delta_{41}
H^2(N_{05}^5)
                                                                                             [\Delta_{12}], 3[\Delta_{15}] - 2[\Delta_{24}] + [\Delta_{33}] - [\Delta_{51}]
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13} - \Delta_{31}, \Delta_{21}, 2\Delta_{14} - \Delta_{23} - \Delta_{41}, 3\Delta_{15} - \Delta_{22} - 2\Delta_{24} - 2\Delta_{31} + \Delta_{33} - \Delta_{51}
Z^2 \left( \mathcal{N}_{06}^5 \right)
                                                                     =
B^2 \left( \mathcal{N}_{06}^5 \right)
                                                                                             \Delta_{11}, \Delta_{21}, \Delta_{13} - \Delta_{31}, \Delta_{12} + 2\Delta_{14} - \Delta_{23} - \Delta_{41}
                                                                    =
H^2 (N_{06}^5)
                                                                                             [\Delta_{12}], 3[\Delta_{15}] - [\Delta_{22}] - 2[\Delta_{24}] - 2[\Delta_{31}] + [\Delta_{33}] - [\Delta_{51}]
                                                                     =
                                                                                               \Delta_{11}, \Delta_{12}, \Delta_{21}, (2-\lambda)\Delta_{13} + \lambda\Delta_{22} + \lambda\Delta_{31}, (3-2\lambda)\Delta_{14} + \lambda(2-\lambda)\Delta_{23} + \lambda\Delta_{32} + \lambda\Delta_{41},
Z^2 \left( \mathcal{N}_{07}^5(\lambda)_{\lambda \neq 0} \right)
                                                                                               (4-3\lambda)\Delta_{15} + \lambda(3-2\lambda)\Delta_{24} + \lambda(2-\lambda)\Delta_{33} + \lambda\Delta_{42} + \lambda\Delta_{51}
B^2 \left( \mathcal{N}_{07}^5(\lambda)_{\lambda \neq 0} \right)
                                                                                             \Delta_{11}, \Delta_{12} + \lambda \Delta_{21}, (2-\lambda)\Delta_{13} + \lambda \Delta_{22} + \lambda \Delta_{31}, (3-2\lambda)\Delta_{14} + \lambda (2-\lambda)\Delta_{23} + \lambda \Delta_{32} + \lambda \Delta_{41}
\mathrm{H}^2\left(\mathcal{N}_{07}^5(\lambda)_{\lambda\neq0}\right)
                                                                                             [\Delta_{21}], (4-3\lambda)[\Delta_{15}] + \lambda(3-2\lambda)[\Delta_{24}] + \lambda(2-\lambda)[\Delta_{33}] + \lambda[\Delta_{42}] + \lambda[\Delta_{51}]
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14}, \Delta_{15}, \Delta_{21}, 2\Delta_{23} - 2\Delta_{32} + \Delta_{41}
Z^2\left(\mathcal{N}_{07}^5(0)\right)
B^2\left(\mathcal{N}_{07}^5(0)\right)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14}
                                                                     =
H^2(N_{07}^5(0))
                                                                                             [\Delta_{15}], [\Delta_{21}], 2[\Delta_{23}] - 2[\Delta_{32}] + [\Delta_{41}] \ \rangle
                                                                                             \Delta_{11}, \overline{\Delta_{12}, \Delta_{13}, \Delta_{14}, \Delta_{21}, 2\Delta_{15} + \Delta_{22} + \Delta_{31}, \Delta_{23} - \Delta_{32} + \Delta_{41}}
\overline{\mathrm{Z}^{2}\left( \mathcal{N}_{08}^{5}\right) }
                                                                     =
B^2 (N_{08}^5)
                                                                                             \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14} + \Delta_{21}
                                                                    =
H^2 (N_{08}^5)
                                                                                             [\Delta_{21}], 2[\Delta_{15}] + [\Delta_{22}] + [\Delta_{31}], [\Delta_{23}] - [\Delta_{32}] + [\Delta_{41}]
```

```
Z^2\left(\mathcal{N}_{09}^5(\lambda)\right)
                                                         \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14}, \Delta_{21}, 2\Delta_{23} - 2\Delta_{32} + \Delta_{41}
B^2(\mathcal{N}_{09}^5(\lambda))
                                                         \Delta_{11}, \Delta_{12}, \Delta_{13}, \lambda \Delta_{14} + 2\Delta_{23} - 2\Delta_{32} + \Delta_{41}
                                          =
\mathrm{H}^2\left(\mathcal{N}_{09}^5(\lambda)\right)
                                                         [\Delta_{14}], [\Delta_{21}]
                                                          \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13} + \Delta_{22} + \Delta_{31}, \Delta_{14} + \Delta_{23} + \Delta_{32} + \Delta_{41},
Z^{2}(N_{10}^{5})
                                                           -2\Delta_{13} + \Delta_{15} + \Delta_{24} + \Delta_{33} + \Delta_{42} + \Delta_{51}
B^2 (N_{10}^5)
                                                         \Delta_{11}, \Delta_{12} + \Delta_{21}, \Delta_{13} + \Delta_{22} + \Delta_{31}, \Delta_{14} + \Delta_{21} + \Delta_{23} + \Delta_{32} + \Delta_{41}
H^2(N_{10}^5)
                                                          [\Delta_{12}], -2[\Delta_{13}] + [\Delta_{15}] + [\Delta_{24}] + [\Delta_{33}] + [\Delta_{42}] + [\Delta_{51}]
                                                         \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{15} + \Delta_{32}, \Delta_{21}, 2\Delta_{14} + \Delta_{22} + \Delta_{31}, \Delta_{14} - 2\Delta_{23} + 2\Delta_{32} - \Delta_{41} 
Z^2\left(\mathcal{N}_{11}^5(\lambda)\right)
B^2\left(\mathcal{N}_{11}^5(\lambda)\right)
                                                          \Delta_{11}, \Delta_{12}, 2\Delta_{13} + \Delta_{21}, \lambda\Delta_{13} + 2\Delta_{14} + \Delta_{22} + \Delta_{31}
                                          =
\mathrm{H}^2\left(\mathcal{N}_{11}^5(\lambda)\right)
                                                         [\Delta_{13}], [\Delta_{15}] + [\Delta_{32}], [\Delta_{14}] - 2[\Delta_{23}] + 2[\Delta_{32}] - [\Delta_{41}]
\overline{\mathrm{Z}^2\left(\mathcal{N}_{12}^5(\lambda)\right)}
                                                         \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{21}, \Delta_{14} - 2\Delta_{23} + 2\Delta_{32} - \Delta_{41}, \Delta_{22} + \Delta_{31} + 2\Delta_{14}
                                          =
B^2(\mathcal{N}_{12}^5(\lambda))
                                                         \Delta_{11}, \Delta_{12}, 2\Delta_{13} + \Delta_{21}, (1+2\lambda)\Delta_{14} + \lambda\Delta_{22} - 2\Delta_{23} + \lambda\Delta_{31} + 2\Delta_{32} - \Delta_{41}
                                          =
\mathrm{H}^2\left(\mathcal{N}_{12}^5(\lambda)\right)
                                                         [\Delta_{13}], [\Delta_{14}] - 2[\Delta_{23}] + 2[\Delta_{32}] - [\Delta_{41}]
                                                          \Delta_{11}, \Delta_{12}, \Delta_{21}, \Delta_{13} + \Delta_{22} + \Delta_{31}, -2\Delta_{13} + \Delta_{14} + \Delta_{23} + \Delta_{32} + \Delta_{41},
Z^2\left(\mathcal{N}_{13}^5(\lambda)\right)
                                                          (1-2\lambda)\Delta_{13} - 2\Delta_{14} + \Delta_{15} - \Delta_{23} + \Delta_{24} + \Delta_{33} + \Delta_{41} + \Delta_{42} + \Delta_{51}
B^2\left(\mathcal{N}_{13}^5(\lambda)\right)
                                                         \Delta_{11}, \Delta_{12} + \Delta_{21}, \Delta_{13} + \Delta_{21} + \Delta_{22} + \Delta_{31}, -2\Delta_{13} + \Delta_{14} + \lambda \Delta_{21} + \Delta_{23} + \Delta_{32} + \Delta_{41}
\mathrm{H}^2\left(\mathcal{N}_{13}^5(\lambda)\right)
                                                         [\Delta_{12}], (1-2\lambda)[\Delta_{13}] - 2[\Delta_{14}] + [\Delta_{15}] - [\Delta_{23}] + [\Delta_{24}] + [\Delta_{33}] + [\Delta_{41}] + [\Delta_{42}] + [\Delta_{51}]
```

$N_{01}^{6}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = e_5$	$e_1e_4 = e_6$	
		$e_2e_1 = e_4$	$e_4 e_1 = -e_6$			
$\mathcal{N}_{02}^{6}(\lambda)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = e_5$	$e_1e_4 = \lambda e_6$	
		$e_2e_1 = e_4$	$e_2e_2 = e_6$	$e_3e_1 = e_6$	$e_4 e_1 = (2 - \lambda) e_6$	
$N_{03}^6(\lambda,\mu)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = \lambda e_5 + \mu e_6$	$e_1e_4 = e_5$	
		$e_2 e_1 = e_4$	$e_2 e_2 = e_6$	$e_3 e_1 = e_6$	$e_4e_1 = -e_5 + 2e_6$	
$N_{04}^{6}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_5$	$e_1e_3 = e_4$	$e_1e_4 = 2e_6$	
		$e_2e_1 = e_3$	$e_2e_3 = -e_6$	$e_3e_1 = -e_4$	$e_4e_1 = -e_6$	
$N_{05}^6(\lambda)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = (2 - \lambda)e_4$	$e_1e_4 = (3 - 2\lambda)e_5$	$e_2e_1 = \lambda e_3 + e_6$
		$e_2e_2 = \lambda e_4$	$e_2 e_3 = \lambda (2 - \lambda) e_5$	$e_3e_1 = \lambda e_4$	$e_3e_2 = \lambda e_5$	$e_4e_1 = \lambda e_5$
$\mathcal{N}_{06}^{6}(\lambda)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_4$	$e_1e_4 = \lambda e_5$	$e_2e_1 = e_6$
		$e_2e_3 = 2e_5$	$e_3e_2 = -2e_5$	$e_4e_1 = e_5$		
$N_{07}^{6}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_4$	$e_1 e_4 = e_5$	
		$e_2e_1 = e_5$	$e_2 e_3 = 2e_6$	$e_3 e_2 = -2e_6$	$e_4 e_1 = e_6$	
$N_{08}^{6}$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = 2e_4$	$e_1 e_4 = e_5$	
		$e_2e_3 = 2e_6$	$e_3e_2 = -2e_6$	$e_4e_1 = e_6$		
$N_{09}^{6}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_4 + e_6$	$e_1e_4 = 2e_5$	
		$e_2 e_1 = e_4$	$e_2 e_2 = e_5$	$e_3 e_1 = e_5$		
$\mathcal{N}_{10}^6(\lambda)$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_4 + e_6$	$e_1 e_4 = (2\lambda + 1)e_5$	$e_2e_1 = e_4$
		$e_2e_2 = \lambda e_5$	$e_2e_3 = -2e_5$	$e_3e_1 = \lambda e_5$	$e_3e_2 = 2e_5$	$e_4e_1 = -e_5$
$\mathcal{N}_{11}^6(\lambda)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_4 + \lambda e_6$	$e_1e_4 = e_5 + 2e_6$	$e_2e_1 = e_4$
		$e_2 e_2 = e_6$	$e_2 e_3 = -2e_5$	$e_3 e_1 = e_6$	$e_3 e_2 = 2e_5$	$e_4e_1 = -e_5$
$N_{12}^{6}$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = e_4 - 2e_5$	$e_1 e_4 = e_5$	$e_2e_1 = e_3 + e_4 + e_6$
		$e_2e_2 = e_4$	$e_2e_3 = e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	$e_4e_1 = e_5$
$N_{13}^{6}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_1e_3 = e_5$	$e_1e_4 = e_6$	$e_1e_5 = 2e_6$
		$e_2e_1 = e_3$	$e_2e_3 = -e_6$	$e_3e_1 = -e_5$	$e_5e_1 = -e_6$	
$\mathcal{N}_{14}^{6}(\alpha)$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_5 + e_6$	$e_1e_4 = \alpha e_6$	
		$e_1e_5 = e_6$	$e_2e_1 = e_3 + e_4$	$e_2e_2 = e_5$	$e_2e_3 = e_6$	
·		$e_3 e_1 = e_5$	$e_3 e_2 = e_6$	$e_4 e_1 = -(\alpha + 1)e_6$	$e_5 e_1 = e_6$	
$N_{15}^{6}$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = -e_5 + e_6$	$e_1e_5 = -3e_6$	
		$e_2e_1 = 3e_3 + e_4$	$e_2e_2 = 3e_5$	$e_2e_3 = -3e_6$	$e_3e_1 = 3e_5$	
26 (2)		$e_3e_2 = 3e_6$	$e_4e_1 = -3e_6$	$e_5e_1 = 3e_6$		
$\mathcal{N}_{16}^6(\lambda)_{\lambda \neq 3}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = (2 - \lambda)e_5$	$e_1 e_4 = e_6$	
		$e_1e_5 = (3 - 2\lambda)e_6$	$e_2e_1 = \lambda e_3 + e_4$	$e_2e_2 = \lambda e_5$	$e_2e_3 = \lambda(2-\lambda)e_6$	
2.6 ( )		$e_3e_1 = \lambda e_5$	$e_3e_2 = \lambda e_6$	$e_4e_1 = -e_6$	$e_5e_1 = \lambda e_6$	
$N_{17}^6(\alpha)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_5$	$e_1 e_4 = e_6$	
		$e_1e_5 = \alpha e_6$	$e_2e_1 = e_4$	$e_2e_3 = 2e_6$	$e_3e_1 = e_6$	
N6 (-)		$e_3e_2 = -2e_6$	$e_4e_1 = -e_6$	$e_5e_1 = e_6$	(- + 0)	
$\mathcal{N}_{18}^{6}(\alpha)$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = 2e_5$	$e_1e_4 = (\alpha + 2)e_6$	$e_1 e_5 = e_6$
26 ( 2)		$e_2e_1 = e_4$	$e_2 e_2 = e_6$	$e_3e_1 = e_6$	$e_4e_1 = -\alpha e_6$	
$\mathcal{N}_{19}^6(\alpha,\beta)$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = 2e_5$	$e_1e_4 = (\alpha + 2)e_6$	
		$e_1 e_5 = \beta e_6$ $e_3 e_1 = e_6$	$e_2 e_1 = e_4$ $e_3 e_2 = -2e_6$	$e_2 e_2 = e_6$ $e_4 e_1 = -\alpha e_6$	$e_2 e_3 = 2e_6$ $e_5 e_1 = e_6$	

$\mathcal{N}_{20}^6(\lambda,\mu)$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = \lambda e_5$	$e_1 e_4 = \mu e_5$	$e_1e_5 = (2\mu - 1)e_6$
Σ·20(Λ, μ)	•	$e_1e_1 = e_2$ $e_2e_1 = e_4$	$e_1e_2 - e_3$ $e_2e_2 = e_5$	$e_1e_3 = \lambda e_5$ $e_2e_3 = \lambda(2-\mu)e_6$	$e_1e_4 = \mu e_5$ $e_2e_4 = \mu(2-\mu)e_6$	ο <sub>1</sub> ο <sub>0</sub> = (2μ 1)e <sub>6</sub>
		$e_3e_1 = e_5$	$e_4e_1 = (2 - \mu)e_5$	$e_3e_2 = (3 - (\lambda + 1)(2 - \mu))e_6$	$e_4 e_2 = (2 - \mu)e_6$	$e_5e_1 = (2 - \mu)e_6$
$N_{21}^{6}(\lambda, \mu)_{\mu \neq 2}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = \lambda e_5 + e_6$	$e_1 e_4 = \mu e_5$	$e_1e_5 = (2\mu - 1)e_6$
21 / /		$e_2e_1 = e_4$	$e_2e_2 = e_5$	$e_2e_3 = \lambda(2-\mu)e_6$	$e_2e_4 = \mu(2-\mu)e_6$	
		$e_3e_1=e_5$	$e_4e_1 = (2 - \mu)e_5$	$e_3 e_2 = (3 - (\lambda + 1)(2 - \mu))e_6$	$e_4 e_2 = (2 - \mu)e_6$	$e_5e_1 = (2 - \mu)e_6$
$\mathcal{N}_{22}^{6}(\lambda)$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = \lambda e_5$	$e_1e_4 = e_5 + e_6$	$e_1e_5 = e_6$
		$e_2e_1 = e_4$	$e_2e_2 = e_5$	$e_2e_3=\lambda e_6$	$e_2e_4 = e_6$	
- 6 ( )		$e_3e_1 = e_5$	$e_4e_1 = e_5 - e_6$	$e_3e_2 = (2-\lambda)e_6$	$e_4 e_2 = e_6$	$e_5e_1 = e_6$
$N_{23}^6(\nu)_{\nu\neq 0}$	:	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_5 + \nu e_6$	$e_1 e_4 = e_5 + e_6$	$e_1 e_5 = e_6$
		$e_2e_1 = e_4$	$e_2e_2 = e_5$	$e_2e_3 = e_6$	$e_2e_4 = e_6$	0-0-
$\mathcal{N}_{24}^6(\lambda)$		$e_3e_1 = e_5$	$e_4e_1 = e_5 - e_6$	$e_3 e_2 = e_6$ $e_1 e_3 = \lambda e_5$	$e_4e_2 = e_6$ $e_1e_4 = 2e_5 + e_6$	$e_5e_1 = e_6$
24(^)	•	$e_1 e_1 = e_2$ $e_2 e_1 = e_4$	$e_1 e_2 = e_3$ $e_2 e_2 = e_5$	$e_{1}e_{3} = \lambda e_{5}$ $e_{3}e_{1} = e_{5}$	$e_3e_2 = e_6$	$e_1 e_5 = e_6$
$\mathcal{N}_{25}^{6}(\lambda)_{\lambda \neq 0}$		$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_5$	$e_1e_4 = \lambda e_5$	$e_1e_5 = -2e_6$
**25(**/ <i>X</i> ≠0	•	$e_2e_1 = e_4$	$e_2e_3 = e_6$	$e_2e_4 = \lambda e_6$	$e_3e_2 = -e_6$	0103 = 200
		$e_4e_1 = -\lambda e_5$	$e_5e_1 = e_6$	2 1	3 2 -0	
$\mathcal{N}_{26}^6(\lambda)_{\lambda \neq 0}$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_5$	$e_1e_4 = \lambda e_5 + e_6$	$e_1e_5 = -2e_6$
201 / //-0		$e_2e_1 = e_4$	$e_2e_3 = e_6$	$e_2e_4 = \lambda e_6$	$e_3e_2 = -e_6$	
		$e_4e_1 = -\lambda e_5 - e_6$	$e_5 e_1 = e_6$			
$N_{27}^{6}$	:	$e_1 e_1 = e_2$	$e_1 e_3 = e_4$	$e_1 e_4 = 2e_5$	$e_1e_5 = 3e_6$	
		$e_2e_1 = e_3$	$e_2e_3 = -e_5$	$e_2e_4 = -2e_6$	$e_3e_1 = -e_4$	
2.6		$e_3e_3 = e_6$	$e_4e_1 = -e_5$	$e_5e_1 = -e_6$	0	
$N_{28}^{6}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_5$	$e_1 e_3 = e_4$	$e_1e_4 = 2e_5$	$e_1e_5 = 3e_6$
		$e_2e_1 = e_3$ $e_3e_3 = e_6$	$e_2 e_2 = -e_6$ $e_4 e_1 = -e_5$	$e_2e_3 = -e_5$ $e_5e_1 = -e_6$	$e_2e_4 = -2e_6$	$e_3e_1 = -e_4 - 2e_6$
$\mathcal{N}_{29}^{6}(\lambda)$		$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1 e_3 = (2 - \lambda)e_4$	$e_1e_4 = (3 - 2\lambda)e_5$	$e_1e_5 = (4 - 3\lambda)e_6$
3129(N)	•	$e_2e_1 = \lambda e_3$	$e_1e_2 = e_3$ $e_2e_2 = \lambda e_4$	$e_2e_3 = \lambda(2-\lambda)e_5$	$e_2e_4 = \lambda(3 - 2\lambda)e_6$	$e_3e_1 = \lambda e_4$
		$e_3e_2 = \lambda e_5$	$e_3e_3 = \lambda(2-\lambda)e_6$	$e_4e_1 = \lambda e_5$	$e_4e_2 = \lambda e_6$	$e_5e_1 = \lambda e_6$
$N_{30}^{6}$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_4 = -e_5$	$e_1e_5 = -2e_6$	
		$e_2e_1 = 2e_3 + e_6$	$e_2e_2 = 2e_4$	$e_2e_4 = -e_6$	$e_3e_1 = 2e_4$	
E		$e_3e_2 = 2e_5$	$e_4e_1 = 2e_5$	$e_4 e_2 = 2e_6$	$e_5e_1 = 2e_6$	
$N_{31}^{6}$	:	$e_1e_1=e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 2e_4$	$e_1e_4 = 3e_5$	
2.6		$e_1 e_5 = e_6$	$e_2e_3 = 2e_6$	$e_3e_2 = -2e_6$	$e_4 e_1 = e_6$	
$\mathcal{N}_{32}^6$	:	$e_1 e_1 = e_2$ $e_1 e_4 = 3e_5$	$e_1 e_2 = e_3$ $e_1 e_5 = e_6$	$e_1 e_3 = 2e_4$ $e_2 e_1 = e_6$		
$N_{33}^{6}$	-	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$\rho_1 \rho_A = \rho_r$	P1 PT - PC
33	•	$e_1e_1 = e_2$ $e_2e_1 = e_3 + e_6$	$e_1e_2 = e_3$ $e_2e_2 = e_4$	$e_1e_3 = e_4$ $e_2e_3 = e_5$	$e_1 e_4 = e_5$ $e_2 e_4 = e_6$	$e_1 e_5 = e_6$ $e_3 e_1 = e_4$
		$e_3e_2 = e_5$	$e_3e_3 = e_6$	$e_4 e_1 = e_5$	$e_4 e_2 = e_6$	$e_5e_1 = e_6$
$\mathcal{N}_{34}^{6}(\lambda)$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$e_1e_4 = e_5$	
01		$e_1e_5 = 2e_6$	$e_2e_1 = e_5 + \lambda e_6$	$e_{2}e_{2} = e_{6}$	$e_3e_1 = e_6$	
$\mathcal{N}_{35}^6(\lambda)_{\lambda \neq 0}$	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_1e_4 = e_5$	
		$e_1e_5 = 2e_6$	$e_2e_1 = e_5$	$e_2e_2 = e_6$	$e_2e_3 = \lambda e_6$	
2.65		$e_3e_1 = e_6$	$e_3e_2 = -\lambda e_6$	$e_4e_1 = \lambda e_6$		
$N_{36}^6$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4 - 2e_6$	$e_1e_4 = e_5$	$e_1e_5 = e_6$
		$e_2e_1 = e_3 + e_5$	$e_2e_2 = e_4$	$e_2e_3 = e_5$ $e_4e_1 = e_5$	$e_2 e_4 = e_6$ $e_4 e_2 = e_6$	$e_3e_1 = e_4$ $e_5e_1 = e_6$
$N_{37}^6(\lambda,\mu)$		$e_3e_2 = e_5$	$e_3e_3 = e_6$	$e_1e_1 - e_5$ $e_1e_3 = 2e_4 + \lambda e_5$	$e_1e_4 = 2e_5 + \mu e_6$	c <sub>0</sub> c <sub>1</sub> = c <sub>6</sub>
37(1, 1)	•	$e_1 e_1 = e_2$ $e_1 e_5 = e_6$	$e_1 e_2 = e_3$ $e_2 e_1 = e_4$	$e_1e_3 = 2e_4 + \lambda e_5$ $e_2e_2 = e_5$	$e_1e_4 = 2e_5 + \mu e_6$ $e_2e_3 = -2\mu e_6$	
		$e_3e_1 = e_5$	$e_3 e_2 = (1 + 2\mu)e_6$	$e_4 e_1 = -\mu e_6$	- 2 - 3 =0	
$\mathcal{N}_{38}^{6}(\lambda)$	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4 - 2e_5 + (1 - 2\lambda)e_6$	$e_1e_4 = e_5 - 2e_6$	$e_1e_5 = e_6$
30 . ,		$e_2e_1 = e_3 + e_4 + \lambda e_5$	$e_2 e_2 = e_4$	$e_2e_3 = e_5 - e_6$	$e_2e_4 = e_6$	$e_3e_1 = e_4$
				$e_4e_1 = e_5 + e_6$		

$$\begin{array}{lll} \mathcal{N}^6_{16}(3) \cong \mathcal{N}^6_{05}(3), & \mathcal{N}^6_{21}(\lambda,2) \cong \mathcal{N}^6_{20}(\lambda,2), & \mathcal{N}^6_{23}(0) \cong \mathcal{N}^6_{22}(1), & \mathcal{N}^6_{25}(0) \cong \mathcal{N}^6_{06}(-2), \\ \mathcal{N}^6_{26}(0) \cong \mathcal{N}^6_{17}(-2), & \mathcal{N}^6_{35}(0) \cong \mathcal{N}^6_{34}(0), & \mathcal{N}^6_{34}(\lambda) \cong \mathcal{N}^6_{34}(-\lambda), & \mathcal{N}^6_{35}(\lambda) \cong \mathcal{N}^6_{35}(-\lambda). \end{array}$$

# REFERENCES

- [1] Abdelwahab H., Calderón A.J., Kaygorodov I., The algebraic and geometric classification of nilpotent binary Lie algebras, International Journal of Algebra and Computation, 29 (2019), 6, 1113–1129.
- [2] Adashev J., Camacho L., Omirov B., Central extensions of null-filiform and naturally graded filiform non-Lie Leibniz algebras, Journal of Algebra, 479 (2017), 461–486.
- [3] Balinsky A., Novikov S., Poisson brackets of hydrodynamic type, Frobenius algebras and Lie algebras, Soviet Mathematics Doklady, 32 (1985), 1, 228–231.
- [4] Bai C., Meng D., The classification of Novikov algebras in low dimensions, Journal of Physics A, 34 (2001), 8, 1581-1594.
- [5] Beneš T., Burde D., Classification of orbit closures in the variety of three-dimensional Novikov algebras, Journal of Algebra and its Applications, 13 (2014), 2, 1350081, 33 pp.
- [6] Bokut L., Chen Yu., Zhang Z., On free Gelfand-Dorfman-Novikov-Poisson algebras and a PBW theorem, Journal of Algebra, 500 (2018), 153-170.

- [7] Burde D., de Graaf W., Classification of Novikov algebras, Applicable Algebra in Engineering, Communication and Computing, 24 (2013), 1, 1–15.
- [8] Burde D., Dekimpe K., Novikov structures on solvable Lie algebras, Journal of Geometry and Physics, 56 (2006), 9, 1837–1855.
- [9] Burde D., Dekimpe K., Vercammen K., Novikov algebras and Novikov structures on Lie algebras, Linear Algebra and its Applications, 429 (2008), 1, 31-41
- [10] Calderón Martín A., Fernández Ouaridi A., Kaygorodov I., The classification of n-dimensional anticommutative algebras with (n-3)-dimensional annihilator, Communications in Algebra, 47 (2019), 1, 173–181.
- [11] Calderón Martín A., Fernández Ouaridi A., Kaygorodov I., The classification of 2-dimensional rigid algebras, Linear and Multilinear Algebra, 68 (2020), 4, 828–844.
- [12] Calderón Martín A., Fernández Ouaridi A., Kaygorodov I., On the classification of bilinear maps with radical of a fixed codimension, Linear and Multilinear Algebra, 2020, to appear, arXiv:1806.07009
- [13] Camacho L., Kaygorodov I., Lopatkin V., Salim M., The variety of dual Mock-Lie algebras, Communications in Mathematics, 2020, to appear, arXiv:1910.01484
- [14] Cañete E., Khudoyberdiyev A., The classification of 4-dimensional Leibniz algebras, Linear Algebra and its Applications, 439 (2013), 1, 273–288.
- [15] Cicalò S., De Graaf W., Schneider C., Six-dimensional nilpotent Lie algebras, Linear Algebra and its Applications, 436 (2012), 1, 163-189.
- [16] Darijani I., Usefi H., The classification of 5-dimensional p-nilpotent restricted Lie algebras over perfect fields, I., Journal of Algebra, 464 (2016), 97–140.
- [17] De Graaf W., Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2, Journal of Algebra, 309 (2007), 2, 640–653.
- [18] De Graaf W., Classification of nilpotent associative algebras of small dimension, International Journal of Algebra and Computation, 28 (2018), 1, 133–161.
- [19] Dekimpe K., Ongenae V., Filiform left-symmetric algebras, Geometriae Dedicata, 74 (1999), 2, 165–199.
- [20] Dzhumadildaev A., Novikov-Jordan algebras, Communications in Algebra, 30 (2002), 11, 5207-5240.
- [21] Dzhumadildaev A., Codimension growth and non-Koszulity of Novikov operad, Communications in Algebra, 39 (2011), 8, 2943-2952.
- [22] Dzhumadildaev A., Ismailov N.,  $S_n$  and  $GL_n$ -module structures on free Novikov algebras, Journal of Algebra, 416 (2014), 287–313.
- [23] Dzhumadildaev A., Löfwall C., Trees, free right-symmetric algebras, free Novikov algebras and identities, Homology, Homotopy and Applications, 4 (2002), 2, 165–190.
- [24] Dzhumadildaev A., Tulenbaev K., Engel theorem for Novikov algebras, Communications in Algebra, 34 (2006), 3, 883–888.
- [25] Fernández Ouaridi A., Kaygorodov I., Khrypchenko M., Volkov Yu., Degenerations of nilpotent algebras, arXiv:1905.05361
- [26] Filippov V., A class of simple nonassociative algebras, Mathematical Notes, 45 (1989), 1-2, 68-71.
- [27] Gorshkov I., Kaygorodov I., Kytmanov A., Salim M., The variety of nilpotent Tortkara algebras, Journal of Siberian Federal University. Mathematics & Physics, 12 (2019), 2, 173–184.
- [28] Gorshkov I., Kaygorodov I., Khrypchenko M., The algebraic classification of nilpotent Tortkara algebras, Communications in Algebra, 2020, DOI: 10.1080/00927872.2020.1742347
- [29] Hegazi A., Abdelwahab H., Classification of five-dimensional nilpotent Jordan algebras, Linear Algebra and its Applications, 494 (2016), 165-218.
- [30] Hegazi A., Abdelwahab H., The classification of n-dimensional non-associative Jordan algebras with (n-3)-dimensional annihilator, Communications in Algebra, 46 (2018), 2, 629–643.
- [31] Hegazi A., Abdelwahab H., Calderón Martín A., The classification of n-dimensional non-Lie Malcev algebras with (n-4)-dimensional annihilator, Linear Algebra and its Applications, 505 (2016), 32–56.
- [32] Hegazi A., Abdelwahab H., Calderón Martín A., Classification of nilpotent Malcev algebras of small dimensions over arbitrary fields of characteristic not 2, Algebras and Representation Theory, 21 (2018), 1, 19–45.
- [33] Ismailov N., Kaygorodov I., Mashurov F., The algebraic and geometric classification of nilpotent assosymmetric algebras, Algebras and Representation Theory, 2020, DOI: 10.1007/s10468-019-09935-y.
- [34] Jumaniyozov D., Kaygorodov I., Khudoyberdiyev A., The algebraic and geometric classification of nilpotent noncommutative Jordan algebras, Journal of Algebra and its Applications, to appear, arXiv:1912.02691
- [35] Karimjanov I., Kaygorodov I., Khudoyberdiyev A., The algebraic and geometric classification of nilpotent Novikov algebras, Journal of Geometry and Physics, 143 (2019), 11–21.
- [36] Karimjanov I., Kaygorodov I., Ladra M., Central extensions of filiform associative algebras, Linear and Multilinear Algebra, 2019, DOI: 10.1080/03081087.2019.1620674
- [37] Kaygorodov I., Khrypchenko M., Lopes S., The algebraic and geometric classification of nilpotent anticommutative algebras, Journal of Pure and Applied Algebra, 224(2020), 8, 106337.
- [38] Kaygorodov I., Khrypchenko M., Popov Yu., The algebraic and geometric classification of nilpotent terminal algebras, arXiv:1909.00358.
- [39] Kaygorodov I., Páez-Guillán P., Voronin V., The algebraic and geometric classification of nilpotent bicommutative algebras, Algebras and Representation Theory, 2020, DOI: 10.1007/s10468-019-09944-x.
- [40] Kaygorodov I., Rakhimov I., Said Husain Sh. K., The algebraic classification of nilpotent associative commutative algebras, Journal of Algebra and its Applications, 2019, DOI: 10.1142/S0219498820502205
- [41] Kaygorodov I., Volkov Yu., The variety of 2-dimensional algebras over an algebraically closed field, Canadian Journal of Mathematics, 71 (2019), 4, 819–842.
- [42] Khudoyberdiyev A., Rakhimov I., Said Husain Sh.K., On classification of 5-dimensional solvable Leibniz algebras, Linear Algebra and its Applications, 457 (2014), 428–454.
- [43] Masutova K., Omirov B., On some zero-filiform algebras, Ukrainian Mathematical Journal, 66 (2014), 4, 541-552.
- [44] Osborn J., Infinite-dimensional Novikov algebras of characteristic 0, Journal of Algebra, 167 (1994), 1, 146–167

- [45] Skjelbred T., Sund T., Sur la classification des algebres de Lie nilpotentes, C. R. Acad. Sci. Paris Ser. A-B, 286 (1978), 5, A241–A242.
- [46] Xu X., Classification of simple Novikov algebras and their irreducible modules of characteristic 0, Journal of Algebra, 246 (2001), 2, 673-707.
- [47] Xu X., Novikov–Poisson algebras, Journal of Algebra, 190 (1997), 2, 253–279.
- [48] Xu X., On simple Novikov algebras and their irreducible modules, Journal of Algebra, 185 (1996), 3, 905–934.
- [49] Zelmanov E., A class of local translation-invariant Lie algebras, Soviet Mathematics Doklady, 35 (1987), 6, 216–218.
- [50] Zusmanovich P., Central extensions of current algebras, Transactions of the American Mathematical Society, 334 (1992), 1, 143–152.