arXiv:2004.03896v1 [nlin.SI] 8 Apr 2020

GEOMETRIC STUDY OF GAS BEHAVIOR IN A ONE-DIMENSIONAL NOZZLE (THE CASE OF THE VAN DER WAALS GAS)

I.S. KRASIL′SHCHIK AND V.V. LYCHAGIN

Abstract. We construct a three-component system of PDEs describing dynamics of van der Walls gas in one-dimensional nozzle. The group of conservation laws for this system is described. We also compute the Lie algebras of point symmetries and present group classification. Examples of exact invariant solutions are given.

Introduction

Methods of differential geometry applied to the problems arising in analysis of partial differential equations (PDEs) allow one to construct a lot of important invariants, such as, e.g., symmetries and conservation laws (see [2], [4], [6]). In turn, these invariants may be used for qualitative study of equations, their classification, construction of exact solutions, etc.

In what follows, we use these methods to study gas behavior in in a one dimensional nozzle. In Section 1, we construct the model and derive the equations for the case the van der Waals gas. Section 2 deals with basic constructions from the geometry of PDEs necessary for the exposition. Conservation laws of the obtained system are described in Section 3, while Section 4 deals with the computation of symmetries and classification of the equations at hand based on their symmetry algebras. Finally, we discuss invariant solutions in Section 5.

1. The model

We consider a gas flow in a wind tunnel (nozzle) of a variable radius. Assume that the x-axes is the symmetry axis of the tunnel, $A(x)$ is an area of section at the point x and (y, z) components of the gas flow velocity are negligibly small in comparison with the x-component $u(x, t)$. Then the Euler equations of the momentum consrevation and the mass conservation laws take the form

$$(A\rho u)_t + (p + A\rho u^2)_x = 0, \quad (A\rho)_t + (A\rho u)_x = 0, \tag{1}$$

where ρ is the gas density and p is the pressure. We also assume that the gas flow is adiabatic, i.e.,

$$s_t + us_x = 0, \tag{2}$$

where $s(x, t)$ is the specific entropy.

The thermodynamic variables s, v, e, p ,and T satisfy in addition the state equations

$$p = RT\Pi_v, \quad e = RT^2\Pi_T, \quad s = R(\Pi + T\Pi_T),$$

where $v = \rho^{-1}$ is the specific volume, e being the specific inner energy, T temperature, R the universal gas constant, while $\Pi = \Pi(v, T)$ is the Massieu-Plank potential (see, for example, [3, 7]).

Let us introduce potential functions φ and ψ for Equations (1):

$$A\rho u = -\varphi_x, \qquad p + A\rho u^2 = \varphi_t,$$
$$A\rho = \psi_x, \qquad A\rho u = -\psi_t.$$

Then System (1) takes the form

$$\varphi_x - \psi_t = 0, \qquad p = \varphi_t + \psi_t^2\psi_x^{-1},$$

2010 *Mathematics Subject Classification.* 76N15, 37K05.

Key words and phrases. Gas dynamics, one-dimensional nozzle, symmetries, conservation laws, invariant solution.

Partially supported by Russian Foundation for Basic Research, Grant 18-29-10013.

$$\rho = \psi_x A^{-1}, \quad u = -\psi_t \psi_x^{-1}.$$

Note the function $\psi(x,t)$ is the first integral of the vector field $\partial_t + u\partial_x$ and rewrite Equation (2) in the form

$$s = S(\psi), \tag{3}$$

for some function S. In order to eliminate the unknown function (3), we use Equation (2) in the form

$$s_v(v_t + uv_x) + s_T(T_t + uT_x) = 0,$$

assuming that the function $s = s(v,T)$ is given by the state equations. Then our system takes the form

$$\varphi_t = p - \varphi_x^2\psi_x^{-1}, \quad \psi_t = \varphi_x, \quad T_t = -uT_x - \frac{s_v}{s_T}(v_t + uv_x),$$

where

$$v = A\psi_x^{-1}, \qquad u = -\varphi_x\psi_x^{-1}.$$

In the case of ideal gas flows we have the following state equations

$$p = Rv^{-1}T, \quad e = \frac{n}{2}RT, \quad s = \ln\left(T^{n/2}v\right) + \text{const},$$

where n is the degree of freedom. For the van der Waals gas the state equations read (see [5])

$$p = \frac{RTv^2 - a(v-b)}{v^2(v-b)}, \quad e = \frac{nRT}{2} - \frac{a}{b}, \quad s = \ln\left(T^{4n/3}(v-b)^{8/3}\right) + \text{const}\,.$$

The constant a takes into account the intermolecular forces and the constant b is the molecular volume, while n is the degree of freedom.

As the result, we arrive to the following system of evolutionary equations

$$\begin{aligned}
\varphi_t &= -\frac{\psi_x(aA\psi_x + ab\psi_x^2 + A^3\varphi_x^2 + bA^2\varphi_x^2\psi_x + R\tau A^2)}{A^2(A + b\psi_x)},\\
\psi_t &= -\varphi_x,\\
\tau_t &= -\frac{A(2\tau\psi_x\varphi_{xx} + n\varphi_x\tau_x\psi_x - 2\tau\varphi_x\psi_{xx}) + 2\tau A_x\varphi_x\psi_x + nb\varphi_x\tau_x\psi_x^2}{\psi_x^2 n(A + b\psi_x)},
\end{aligned} \tag{4}$$

where the temperature T is relabeled to τ and ψ to $-\psi$.

We study this system in the forthcoming sections. Note that the case $a,\, b = 0$ corresponds to the ideal gas.

2. Preliminary facts and notation

We consider the space $J^\infty(2,3)$ with the coordinates x, t, φ_{kl}, ψ_{kl}, τ_{kl}, $k,\, l = 0, 1, \ldots$, where the subscript corresponds to the partial derivative $\partial^{k+l}/\partial x^k \partial t^l$, and the subspace $\mathcal{E} \subset J^\infty(2,3)$ defined by Equations (4) and all their differential consequences. The functions x, t, $\varphi_k = \varphi_{k0}$, $\psi_k = \psi_{k0}$, and $\tau_k = \tau_{k0}$ may be chosen for internal coordinates in $\mathcal{E}$. Denote by r_φ and r_τ the right-hand sides of the first and third equations in (4), respectively. Then the total derivatives

$$\begin{aligned}
D_x &= \frac{\partial}{\partial x} + \sum_{k\geq 0}\left(\varphi_{k+1}\frac{\partial}{\partial\varphi_k} + \psi_{k+1}\frac{\partial}{\partial\psi_k} + \tau_{k+1}\frac{\partial}{\partial\tau_k}\right),\\
D_t &= \frac{\partial}{\partial t} - \sum_{k\geq 0}\left(D_x^k(r_\varphi)\frac{\partial}{\partial\varphi_k} + \varphi_{k+1}\frac{\partial}{\partial\psi_k} + D_x^k(r_\tau)\frac{\partial}{\partial\tau_k}\right)
\end{aligned}$$

are vector fields on $\mathcal{E}$. For any function Y on $\mathcal{E}$ define its linearization

$$\ell_Y(f,g,h) = \sum_{k\geq 0}\left(\frac{\partial Y}{\partial\varphi_k}D_x(f) + \frac{\partial Y}{\partial\psi_k}D_x(g) + \frac{\partial Y}{\partial\tau_k}D_x(h)\right).$$

A symmetry of $\mathcal{E}$ is a vector field that commutes with D_x and D_t. Any symmetry is of the form

$$\mathbf{E}_{\mathcal{S}} = \sum_{k\geq 0}\left(D_x^k(\mathcal{S}_\varphi)\frac{\partial}{\partial\varphi_k} + D_x^k(\mathcal{S}_\psi)\frac{\partial}{\partial\psi_k} + D_x^k(\mathcal{S}_\tau)\frac{\partial}{\partial\tau_k}\right),$$

where the vector-function $\mathcal{S} = (\Phi, \Psi, \Theta)$ must satisfy the linear system

$$D_t(\Phi) = \ell_{r_\varphi}(\Phi,\Psi,\Theta),\ D_t(\Psi) = -D_x(\Phi),\ D_t(\Theta) = \ell_{r_\varphi}(\Phi,\Psi,\Theta). \tag{5}$$

We identify symmetries with the corresponding functions $\mathcal{S}$.

A conservation law of $\mathcal{E}$ is a differential form $\omega = X\,dx + T\,dt$ such that $D_t(X) = D_x(T)$. It is called trivial if there exists a potential Z such that $X = D_x(Z)$, $T = D_t(Z)$. The generating function of ω is the triple $G = (G_\varphi, G_\psi, G_\tau)$, where

$$G_\varphi = \frac{\delta X}{\delta\varphi},\quad G_\psi = \frac{\delta X}{\delta\psi},\quad \mathcal{S}_\tau = \frac{\delta X}{\delta\tau} \tag{6}$$

and

$$\frac{\delta X}{\delta\varphi} = \sum_{k\geq 0}(-1)^k D_x^k\left(\frac{\partial X}{\partial\varphi_x}\right),$$

etc., are the variational derivatives. To find generating functions, one needs to solve the system

$$\begin{aligned}
D_t(G_\varphi) &= \sum_{k\geq 0}(-1)^{k+1}\left(D_x^k\left(\frac{\partial G_\varphi}{\partial\varphi_k}\right) + D_x^k\left(\frac{\partial G_\psi}{\partial\varphi_k}\right) + D_x^k\left(\frac{\partial G_\tau}{\partial\varphi_k}\right)\right),\\
D_t(G_\psi) &= \sum_{k\geq 0}(-1)^{k+1}\left(D_x^k\left(\frac{\partial G_\varphi}{\partial\psi_k}\right) + D_x^k\left(\frac{\partial G_\psi}{\partial\psi_k}\right) + D_x^k\left(\frac{\partial G_\tau}{\partial\psi_k}\right)\right),\\
D_t(G_\tau) &= \sum_{k\geq 0}(-1)^{k+1}\left(D_x^k\left(\frac{\partial G_\varphi}{\partial\tau_k}\right) + D_x^k\left(\frac{\partial G_\psi}{\partial\tau_k}\right) + D_x^k\left(\frac{\partial G_\tau}{\partial\tau_k}\right)\right)
\end{aligned}$$

adjoint to (5). A conservaton law is trivial if and only if its generating function vanishes.

3. Conservation laws

Solving this system in the case when G depends on x, t, φ_k, ψ_k, τ_k, $k \leq 2$, we get the solutions

$$G^0 = (0,1,0), \qquad G^\infty = (0, G_\psi, G_\tau),$$

where

$$\begin{aligned}
G_\psi = &-\frac{1}{n\tau\psi_x^2(b\psi_x + A)^2}\Big(2A^2\tau\Big(\psi_x H - \tau^{\frac{n}{2}}(b\psi_x + A)\frac{\partial H}{\partial y}\Big)\psi_{xx}\\
&+ \psi_x^2\Big(2\tau A\frac{\partial A}{\partial x} + n\tau_x(b\psi_x + A)^2\Big)H - 2A\tau\psi_x^3(b\psi_x + A)\frac{\partial H}{\partial\psi}\\
&- A\tau^{\frac{n}{2}}\psi_x(b\psi_x + A)\Big(2\tau\frac{\partial A}{\partial x} + n\tau_x(b\psi_x + A)\Big)\frac{\partial H}{\partial y}\Big),\\
G_\tau = &\frac{H\psi_x}{\tau}.
\end{aligned}$$

where $H = H(y,\psi)$ is an arbitrary smooth function and

$$y = \frac{b\psi_x + A}{\psi_x}\cdot\tau^{\frac{n}{2}}.$$

Using relations (6), one can reconstruct the corresponding conservation laws and obtain

$$\omega_0 = \psi\,dx + \varphi\,dt$$

and

$$\omega_H = \frac{2H\ln(y)}{n}\cdot(\psi_x\,dx - \varphi_x\,dt).$$

4. Symmetries

We compute here point symmetries, i.e., such that the function $\mathcal{S}$ depends on x, t and φ_k, ψ_k, τ_k for $k \leq 2$. Note first that the second equation in (5) means that the form $\Psi\,dx - \Phi\,dt$ is a conservation law. Due to the results of Section 3, we have

$$\Phi = -\lambda\varphi + \frac{2H\ln(y)}{n}\varphi_x - D_t(P), \qquad \Psi = \lambda\psi + \frac{2H\ln(y)}{n}\psi_x + D_x(P),$$

where $\lambda = \mathrm{const}$ and $P = P(x, t, \varphi, \psi, \tau, \varphi_x, \psi_x, \tau_x)$ is an arbitrary smooth function. Denote by $\mathfrak{g}$ the Lie algebra of symmetries. The structure of $\mathfrak{g}$ depends on the values of the parameters that enter the basic equations[1]. Note that in all the cases below the algebra $\mathfrak{g}$ contains the 3-dimensional Abelian ideal $\mathfrak{g}_0$ spanned by the symmetries

$$\mathcal{S}_1 = (1, 0, 0), \quad \mathcal{S}_2 = (0, 1, 0), \quad \mathcal{S}_3 = (\varphi_t, \psi_t, \tau_t).$$

Describing the algebras, we present only those generators that do not belong to $\mathfrak{g}_0$. In the description of Lie algebra structures we indicate nonzero commutators only.

4.1. $b = 0$, $a \neq 0$. There are eight subcases.

4.1.1. $A = p = \mathrm{const}$, $n = 2$. The algebra $\mathfrak{g}$ is 6-dimensional. Its generators are

$$\Phi_4 = \frac{\psi_x^2 at + pt(p\varphi_x^2 + R\tau)\psi_x - \varphi p^2}{p^2}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = -\frac{\tau t\varphi_x\psi_{xx}}{\psi_x^2} + \frac{\tau t\varphi_{xx}}{\psi_x} + \frac{t\varphi_x\tau_x}{\psi_x} - 2\tau - \frac{2a\psi_x}{pR};$$
$$\Phi_5 = \varphi_x, \quad \Psi_5 = \psi_x, \quad \Theta_5 = \tau_x;$$
$$\Phi_6 = x\varphi_x - 2\varphi, \quad \Psi_6 = x\psi_x\psi, \quad \Theta_6 = x\tau_x - 2\tau - \frac{2a\psi_x}{pR}.$$

with the commutators

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_1, \mathcal{S}_6] = -2\mathcal{S}_1,\ [\mathcal{S}_2, \mathcal{S}_6] = -\mathcal{S}_2,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3, [\mathcal{S}_5, \mathcal{S}_6] = -\mathcal{S}_5.$$

4.1.2. $A = p = \mathrm{const}$, $n \neq 2$. We have $\dim\mathfrak{g} = 5$ with the generators

$$\Phi_4 = \frac{\psi_x^2 at + pt(p\varphi_x^2 + R\tau)\psi_x - p^2(\varphi_x x - \varphi)}{p^2}, \quad \Psi_4 = t\varphi_x - \psi_x x + \psi,$$
$$\Theta_4 = \frac{-2\tau t\varphi_x\psi_{xx} + 2\tau t\varphi_{xx}\psi_x + (t\varphi_x - \psi_x x)\tau_x\psi_x n}{\psi_x^2 n};$$
$$\Phi_5 = \varphi_x, \quad \Psi_5 = \psi_x, \quad \Theta_5 = \tau_x.$$

The commutators are

$$[\mathcal{S}_2, \mathcal{S}_4] = \mathcal{S}_2,\ [\mathcal{S}_2, \mathcal{S}_5] = 0,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3,\ [\mathcal{S}_4, \mathcal{S}_5] = -\mathcal{S}_5.$$

4.1.3. $A = pe^{qx}$, p, $q \neq 0$, $n = 2$. One has $\dim\mathfrak{g} = 5$. The generators are

$$\Phi_4 = \frac{e^{-2qx}\psi_x^2 at + e^{-qx}\psi_x Rpt\tau + \psi_x\varphi_x^2 p^2 t - \varphi p^2}{p^2}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = \frac{\tau t\varphi_{xx}\psi_x pR - t\tau\varphi_x\psi_{xx}pR + \psi_x(-2\psi_x^2 ae^{-qx} + (-2\psi_x\tau + t\varphi_x(q\tau + \tau_x))pR)}{\psi_x^2 pR};$$
$$\Phi_5 = \varphi_x, \quad \Psi_5 = \psi_x, \quad \Theta_5 = \frac{-2e^{-qx}qa\psi_x - pR(q\tau - \tau_x)}{pR}.$$

The commutators read

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

[1]The full classification includes the case $n = -2$, which is physically senseless and we omit it. So, everywhere below $n \neq -2$

4.1.4. $A = pe^{qx}$, $p,\ q \neq 0$, $n \neq 2$. The algebra of symmetries is 4-dimensional with the generators

$$\Phi_4 = \frac{e^{-2qx}\psi_x^2 aqt + e^{-qx}\psi_x Rpqt\tau + \psi_x\varphi_x^2 p^2 qt - \varphi p^2 q - \varphi_x p^2}{(p^2 q)}, \quad \Psi_4 = t\varphi_x - \frac{\psi_x}{q},$$
$$\Theta_4 = \frac{2\tau t\varphi_{xx}\psi_x q - 2t\tau\varphi_x\psi_{xx}q + (-n(q\tau+\tau_x)\psi_x + qt\varphi_x(n\tau_x + 2q\tau))\psi_x}{n\psi_x^2 q}.$$

and commutators

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

4.1.5. $A = (px+r)^q$, $p,\ q \neq 0$, $n = 2$. The symmetry algebra is 5-dimensional and generated by

$$\Phi_4 = \psi_x^2(px+r)^{-q}at + \psi_x(px+r)^{-q}Rt\tau + \varphi_x^2\psi_x t - \varphi, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = \frac{-t\tau\varphi_x\psi_{xx}}{\psi_x^2} + \frac{t\tau\varphi_{xx}}{\psi_x} + \frac{t\varphi_x\tau_x}{\psi_x} + \frac{qpt\tau\varphi_x}{(px+r)\psi_x} - 2\tau - \frac{2(px+r)^{-q}a\psi_x}{R};$$
$$\Phi_5 = -2\varphi + \frac{(px+r)\varphi_x}{p}, \quad \Psi_5 = -\psi + \frac{(px+r)\psi_x}{p},$$
$$\Theta_5 = \frac{(-2apq\psi_x - 2\psi_x)(px+r)^{-q}ap + (\tau_x(px+r) - p\tau(q+2))R}{pR}$$

with the commutators

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_1, \mathcal{S}_5] = -2\mathcal{S}_1,\ [\mathcal{S}_2, \mathcal{S}_5] = -\mathcal{S}_2,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

4.1.6. $A = (px+r)^q$, $p,\ q \neq 0$, $n \neq 2$. Here $\dim \mathfrak{g} = 4$. Generators are

$$\Phi_4 = -2\varphi + (q+1)\varphi - (q+1)t\psi_x(R(px+r)^{-q}\tau + \psi_x(px+r)^{-2q}a + \varphi_x^2)$$
$$+ \frac{(px+r)\varphi_x}{p}, \qquad \Psi_4 = -\psi - (q+1)t\varphi_x + \frac{(px+r)\psi_x}{p},$$
$$\Theta_4 = \frac{1}{(px+r)\psi_x^2 npR}\Big(- 2\psi_x^3 anpq((px+r)^{-q+1} + 2(px+r)(px+r)^{-q})$$
$$- 2R(\tau\psi_x pt(q+1)(px+r)\varphi_{xx} - \tau\varphi_x pt(q+1)(px+r)\psi_{xx}$$
$$+ (-\frac{1}{2}((px+r)\tau_x + pq\tau)(px+r)n\psi_x + (\frac{1}{2}\tau_x n(px+r) + pq\tau)\varphi_x p(q+1)t)\psi_x)\Big)$$

while the brackets read

$$[\mathcal{S}_1, \mathcal{S}_4] = (q-1)\mathcal{S}_1,\ [\mathcal{S}_2, \mathcal{S}_4] = -\mathcal{S}_2,\ [\mathcal{S}_3, \mathcal{S}_4] = -(q+1)\mathcal{S}_3.$$

4.1.7. *A general,* $n = 2$. The algebra is 4-dimensional with the generators

$$\Phi_4 = \frac{(\psi_x^2 at + At(A\varphi_x^2 + R\tau)\psi_x - \varphi A^2)}{A^2}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = -\frac{t\tau\varphi_x\psi_{xx}}{\psi_x^2} + \frac{\tau t\varphi_{xx}}{\psi_x} + \frac{t\varphi_x\tau_x}{\psi_x} + \frac{A_x t\tau\varphi_x}{\psi_x A} - 2\tau - \frac{2a\psi_x}{AR}.$$

and the commutator

$$[\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

4.1.8. *A general,* $n \neq 2$. Here $\mathfrak{g}$ coincides with $\mathfrak{g}_0$.

4.2. $b = 0$, $a = 0$ **(the ideal gas).** We have four cases.

4.2.1. $A = p = \text{const}$. The 6-dimensional algebra is generated by

$$\Phi_4 = \frac{-\varphi p + t\psi_x(\varphi_x^2 p + R\tau)}{p}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = -\frac{2t\tau\varphi_x\psi_{xx}}{\psi_x^2 n} + \frac{2t\tau\varphi_{xx}}{\psi_x n} + \frac{t\varphi_x\tau_x}{\psi_x} - 2\tau;$$
$$\Phi_5 = \varphi_x, \quad \Psi_5 = \psi_x, \quad \Theta_5 = \tau_x;$$
$$\Phi_6 = x\varphi_x - 2\varphi, \quad \Psi_6 = x\psi_x - \psi, \quad \Theta_6 = x\tau_x - 2\tau.$$

The commutators are

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_1, \mathcal{S}_6] = -2\mathcal{S}_1,\ [\mathcal{S}_2, \mathcal{S}_6] = -2\mathcal{S}_2,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3,\ [\mathcal{S}_5, \mathcal{S}_6] = -\mathcal{S}_5.$$

4.2.2. $A = pe^{qx}$, $p,\ q \neq 0$. One has $\dim \mathfrak{g} = 5$. The algebra is generated by

$$\Phi_4 = \frac{\psi_x e^{-qx} R t\tau + p(\psi_x t\varphi_x^2 - \varphi)}{p}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = \frac{2\tau t\varphi_{xx}\psi_x - 2t\tau\varphi_x\psi_{xx} + \psi_x(-2\psi_x n\tau + t\varphi_x(n\tau_x + 2q\tau))}{n\psi_x^2};$$
$$\Phi_5 = \varphi_x, \quad \Psi_5 = \psi_x, \quad \Theta_5 = -q\tau + \tau_x.$$

The commutators are

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

4.2.3. $A = (px + r)^q$, $p,\ q \neq 0$. One has $\dim \mathfrak{g} = 5$ and the algebra is generated by

$$\Phi_4 = \psi_x(px + r)^{-q} R t\tau + \psi_x\varphi_x^2 t - \varphi, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = -\frac{2t\varphi_x\tau\psi_{xx}}{\psi_x^2 n} + \frac{2\tau t\varphi_{xx}}{\psi_x n} + \frac{t\varphi_x\tau_x}{\psi_x} + \frac{2qpt\tau\varphi_x}{(px + r)\psi_x n} - 2\tau;$$
$$\Phi_5 = -2\varphi + \frac{(px + r)\varphi_x}{p}, \quad \Psi_5 = -\psi + \frac{(px + r)\psi_x}{p}, \quad \Theta_5 = \frac{(px + r)\tau_x - p\tau(q + 2)}{p}.$$

The commutator relations are

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_1, \mathcal{S}_5] = -2\mathcal{S}_1,\ [\mathcal{S}_2, \mathcal{S}_5] = -\mathcal{S}_2,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

4.2.4. *A general.* The 4-dimensional algebra is generated by

$$\Phi_4 = \frac{t\psi_x(\varphi_x^2 A + R\tau) - \varphi A}{A}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = -\frac{2t\tau\varphi_x\psi_{xx}}{\psi_x^2 n} + \frac{2\tau t\varphi_{xx}}{\psi_x n} + \frac{t\varphi_x\tau_x}{\psi_x} + \frac{2tA_x\tau\varphi_x}{\psi_x A n} - 2\tau.$$

and the brackets are

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3.$$

4.3. $b \neq 0$**.** The four subcases here are as follows.

4.3.1. $a = 0$, $A = p = \text{const}$. We have $\dim \mathfrak{g} = 6$. The algebra is generated by

$$\Phi_4 = \frac{\varphi_x^2\psi_x^2 bt + (pt\varphi_x^2 + Rt\tau - \varphi b)\psi_x - p\varphi}{(b\psi_x + p}, \quad \Psi_4 = t\varphi_x,$$
$$\Theta_4 = \frac{2tp\tau\varphi_{xx}\psi_x - 2tp\tau\varphi_x\psi_{xx} - (2(b\psi_x + p))\psi_x(-\frac{1}{2}t\varphi_x\tau_x + \psi_x\tau)n}{\psi_x^2 n(b\psi_x + p)};$$
$$\Phi_5 = \varphi_x, \quad \Psi_5 = \psi_x, \quad \Theta_5 = \tau_x;$$
$$\Phi_6 = x\varphi_x - 2\varphi, \quad \Psi_6 = x\psi_x - \psi, \quad \Theta_6 = x\tau_x - 2\tau$$

with the commutators

$$[\mathcal{S}_1, \mathcal{S}_4] = -\mathcal{S}_1,\ [\mathcal{S}_1, \mathcal{S}_6] = -2\mathcal{S}_1,\ [\mathcal{S}_2, \mathcal{S}_6] = -\mathcal{S}_2,\ [\mathcal{S}_3, \mathcal{S}_4] = \mathcal{S}_3,\ [\mathcal{S}_5, \mathcal{S}_6] = -\mathcal{S}_5.$$

4.3.2. $a=0$, *A general.* The symmetry algebra has 4 generators

$$\Phi_4=\frac{\varphi_x^2\psi_x^2bt+(At\varphi_x^2+Rt\tau-\varphi b)\psi_x-A\varphi}{b\psi_x+A},\quad \Psi_4=\varphi_x t,$$
$$\Theta_4=\frac{1}{\psi_x^2n(b\psi_x+A)}\Big(2A\tau t\varphi_{xx}\psi_x-2tA\tau\varphi_x\psi_{xx}$$
$$+(-2bn\tau\psi_x^2+(\varphi_x\tau_xbnt-2An\tau)\psi_x+\varphi_xt(A\tau_xn+2A_x\tau))\psi_x\Big)$$

that enjoy the relations

$$[\mathcal{S}_1,\mathcal{S}_4]=-\mathcal{S}_1,\ [\mathcal{S}_3,\mathcal{S}_4]=\mathcal{S}_3.$$

4.3.3. $a\neq 0$, $A=p=\text{const}$. Dimension of $\mathfrak{g}$ is five here. The generators are

$$\Phi_4=\frac{\psi_x^3abt+pt(bp\varphi_x^2+a)\psi_x^2+p^2(pt\varphi_x^2+Rt\tau-bx\varphi_x+b\varphi)\psi_x-p^3(\varphi_x x-\varphi)}{p^2(b\psi_x+p)},$$
$$\Psi_4=t\varphi_x-\psi_x x+\psi,$$
$$\Theta_4=\frac{2p\tau t\varphi_{xx}\psi_x-2t\varphi_xp\tau\psi_{xx}-\psi_x\tau_xn(-t\varphi_x+\psi_x x)(b\psi_x+p)}{\psi_x^2n(b\psi_x+p)},$$
$$\Phi_5=\varphi_x,\quad \Psi_5=\psi_x,\quad \Theta_5=\tau_x.$$

They are subject to the relations

$$[\mathcal{S}_1,\mathcal{S}_4]=\mathcal{S}_1,\ [\mathcal{S}_2,\mathcal{S}_4]=\mathcal{S}_2,\ [\mathcal{S}_3,\mathcal{S}_4]=\mathcal{S}_3,\ [\mathcal{S}_4,\mathcal{S}_5]=-\mathcal{S}_5.$$

4.3.4. $a\neq 0$, *A general.* One has $\mathfrak{g}=\mathfrak{g}_0$.

5. Examples of exact solutions

Let us now describe two types of exact solutions

- stationary ones, i.e., independent of t,
- traveling waves in the case $A=\text{const}$.

5.1. **Stationary solutions.** Consider the stationary case, i.e., assume $\psi_t=\varphi_t=\tau_t=0$ in Equations (4). Then the second equation reads $\varphi_x=0$, i.e., $\varphi=\text{const}$, the entire system reduces to the sole equation

$$\psi_x(aA\psi_x+ab\psi_x^2+RA^2\tau)=0.$$

Under the assumption $ab\neq 0$, we obtain three solutions

$$\psi_x=-\frac{\big(a+\sqrt{a^2-4abR\tau}\big)A}{2ab},\quad \psi_x=\frac{\big(-a+\sqrt{a^2-4abR\tau}\big)A}{2ab},\quad \psi_x=0$$

of which only the first one has physical meaning, because by definition $\psi_x=-\rho A<0$ (recall that in the final system we relabeled ψ to $-\psi$). Thus, we obtain

$$\rho=\frac{a+\sqrt{a^2-4abR\tau}}{2ab},$$

or

$$\tau=\frac{a\rho(1-b\rho)}{R},$$

which gives a simple dependence between temperature and density.

5.2. **Traveling-wave solutions.** If $A = \text{const}$ then System (4) becomes invariant with respect to x-translation symmetry $\mathfrak{X} = (\varphi_x, \psi_x, \tau_x)$, and one can consider traveling-wave solutions, i.e., solutions invariant with respect to $\mathfrak{X} + \nu\mathcal{T}$, where $\mathcal{T} = (\varphi_t, \psi_t, \tau_t)$, ν being the velocity. Such solutions are functions of the variable $z = x - \nu t$ and one has

$$\frac{\partial}{\partial x} = \frac{d}{dz}, \qquad \frac{\partial}{\partial t} = -\nu\frac{d}{dz}.$$

Consequently, (4) reduces to the two equations $\dot{\varphi} = \nu\dot{\psi}$ and

$$\big(\nu^2A^2b\dot{\psi}^3 + (ab + \nu^2A^3)\dot{\psi}^2 + (aA - \nu^2A^2b)\dot{\psi} - \nu^2A^3 + R\tau A^2\big)\dot{\psi} = 0, \tag{7}$$

where 'dot' denotes the z-derivative, while the definitions of the potentials (see Section 1) take the form

$$\rho = -\frac{\dot{\psi}}{A}, \quad u = \nu, \quad p = -\nu(1+\nu)\dot{\psi}.$$

Now, solving (7) with respect to τ we obtain dependence of temperature

$$\tau = \frac{(b\rho - 1)(\nu^2A^3\rho^2 - a\rho - \nu^2A)}{R}$$

on density.

Acknowledgments

Computations were supported by the Jets software, [1].

References

[1] H. Baran, M. Marvan, *Jets. A software for differential calculus on jet spaces and diffeties*. http://jets.math.slu.cz.
[2] A.V. Bocharov et al., *Symmetries of Differential Equations in Mathematical Physics and Natural Sciences*, edited by A.M. Vinogradov and I.S. Krasil′shchik). Factorial Publ. House, 1997 (in Russian). English translation: Amer. Math. Soc., 1999.
[3] V.E. Fortov, *Equation of state*, Thermopedia, http://dx.doi.org/10.1615/AtoZ.e.equation_of_state.
[4] I.S. Krasil′shchik, V.V. Lychagin, A.M. Vinogradov, *Geometry of Jet Spaces and Nonlinear Partial Differential Equations*. Gordon and Breach, New York, 1986.
[5] V.V. Lychagin, M.D. Roop, *Critical phenomena in filtration processes of real gases*, Lobachevskii J. of Mathematics, **41** (2020) 3.
[6] L.V. Ovsiannikov, *Group Analysis of Differential Equations Paperback*, Academic Press, New York, 1982.
[7] M. Planck, Über das Gesetz der Energieverteilung im Normalspectrum, Ann. der Phys, **309** (1901) 3, 553–563.

Trapeznikov Institute of Control Sciences, 65 Profsoyuznaya street, Moscow 117997, Russia
E-mail address: josephkra@gmail.com

Trapeznikov Institute of Control Sciences, 65 Profsoyuznaya street, Moscow 117997, Russia
E-mail address: lychagin@math.uit.no