

Cartan Connection for Schrödinger equation. The nature of vacuum.

Radosław A. Kycia^{1,2,a}

¹Masaryk University
Department of Mathematics and Statistics
Kotlářská 267/2, 611 37 Brno, The Czech Republic

²Cracow University of Technology
Faculty of Materials Science and Physics
Warszawska 24, Kraków, 31-155, Poland

^akycia.radoslaw@gmail.com

Abstract

We reinterpret the Schrödinger equation as a continuity equation in the space with the Cartan connection given by scaling Lie-Bäcklund group on a specific jet space. In this space, the wave function and their gradient coordinates are treated as independent coordinates. This approach gives a full Cartan connection form a divergence-free condition. Once constructed, the connection makes it possible to investigate the geometry of the space on which this Schrödinger-Cartan connection is constructed. This is the idea that generalizes the concepts present in de Broglie-Bohm (pilot wave) theory in a geometric way. We also present this procedure for constructing (non-uniquely) torsion-free Cartan connections for general Partial Differential Equations.

Keywords: Cartan connection on jet space; Schrödinger equation; de Broglie-Bohm theory; pilot-wave theory; vacuum; geometric theory of differential equations; jet space;

1 Introduction

Construction of a connection for Schrödinger equation opens up interpretation of quantum mechanics in geometrical terms. Some approaches are using the quantum connection, e.g., [11, 10] or contact geometry [7, 8]. Moreover, the symmetry methods [13, 14, 16, 15] for Partial Differential Equations (PDEs) are with success applicable to Schrödinger equations including scale transformations, e.g., [6, 23].

There are various notions of connection. The most general is the Ehresmann connection [12]. The Cartan connection in physics usually appears in the context of the Einstein-Cartan theory [19], and for reductive geometries, it is equivalent to the Ehresmann connection. However, the Cartan connection carries full geometric information on geometry, especially for embedded submanifolds [18], [1], [12].

These results suggest the different approaches to the construction of connection for the Schrödinger equation. We will develop a different approach that merges the

scaling group and the Cartan connection in a (larger) jet space [13, 14, 16, 15]. As an entry point, we reinterpret Schrödinger equation as a divergence equation in a jet space, where divergence operator is given by some Cartan connection induced by a scaling group on this jet space. Then matching the general divergence equation and the Schrödinger equation fixes some of the Cartan connection coefficients. Unfortunately, some of them are undefined, and therefore, some additional constraint, as torsion freeness must be imposed to limit some freedom.

The idea of some sort of scaling of the length of the object during its movement was introduced in the General Relativity context by Hermann Weyl [21], [22]. It was rejected as obviously non-physical but gave rise to the gauge principle as the scaling of phase and is base of all modern fundamental Yang-Mills type theories. In this paper, we interpret this scaling/gauge idea in the context of a change of scale of the wave function and its gradient components. This gives rise to the Cartan connection for the scaling group, which in turn, can be associated with this background, which we will call the vacuum. We will show that all these ideas can be connected in the procedure that allows us to construct a Cartan connection for Schrödinger equation.

We sketch the basic ingredient of this procedure. Consider flat Euclidean space with 'time-space' coordinates (t, x) . Then we have the Schrödinger equation (we set the Planck constant $\hbar = 1$, $i = \sqrt{-1}$) for a complex-valued L^2 function $u = u(x, t)$

$$i\partial_t u + \partial_x^2 u = Vu, \quad (1)$$

where $V = V(x, t)$ is a real-valued function called potential, which by physical assumptions is bounded from below, and it can be set¹ $V > 0$. Usually $V = V(x)$, but in general $V = V(x, t)$. The additional normalization condition $\int |u(x, t)|^2 dx = 1$ is imposed for probabilistic interpretation of $|u(x, t)|^2 dx$ as a probability measure for finding particle at (t, x) . This normalization, which is projection from $\mathbb{C} \setminus \{(0, 0)\} \rightarrow \mathbb{C}P^1 \cong S^2$ can be imposed in the end, so therefore we will be omitting it in what follows.

As it was pointed out in [17], the equation (1) can be written as a conservation law

$$d\lambda = 0, \quad (2)$$

for

$$\lambda = in(x)u(x, t)dx - \left(n(x)\partial_x u(x, t) - \frac{dn(x)}{dx} \right) dt, \quad (3)$$

with the condition

$$\frac{d^2 n(x)}{dx^2} = V(x)n(x). \quad (4)$$

The condition (4) is an elliptic problem and for suitable boundary conditions the solution can be assumed to be positive², or generally $n : \mathbb{R}^2 \rightarrow \mathbb{C} \setminus \{(0, 0)\}$.

This suggest the following ansatz $u(x, t) \rightarrow n(x)u(x, t)$ in (1), that yields

$$n(i\partial_t u + \partial_x^2 u) = -2\frac{dn}{dx}\partial_x u + u \left(Vn - \frac{d^2 n}{dx^2} \right), \quad (5)$$

¹For bounded from below V one can set $u \rightarrow e^{iAt}$ for $A > \min_{\{t, x\}}(V(x, t))$, which results in $V \rightarrow V + A > 0$.

²We can write a nontivial weak solution (so also a smooth one) of this equation as an unique minimizer of functional $F(n) = \int (\frac{1}{2}|\nabla n|^2 + \frac{1}{2}Vn^2) dx > 0$ for $V > 0$, i.e., $F(u) \leq F(u_+)$, for $u_+ := \max\{0, u\}$.

which can be rewritten as the system

$$\begin{cases} i\partial_t u + \partial_x^2 u + \frac{2}{n} \frac{dn}{dx} \partial_x u = 0 \\ \frac{1}{n} \frac{d^2 n(x)}{dx^2} = V(x). \end{cases} \quad (6)$$

This decomposition, which works for linear Schrödinger equation, has interesting interpretation. The second (elliptic) equation couples potential with scale n . This scale factor then couples, in the first (evolutionary) equation, to the wave function of the object under interest by the gradient term $\frac{2}{n} \frac{dn}{dx} \partial_x u$. Therefore, this system of equations can be interpreted as evolution of u on 'vacuum' background n , and thus we will be calling the scale n the vacuum. In addition, due to elliptic character of equation for n the background is non-casual - it is fixed once V is given or, loosely speaking, 'the perturbation propagates with infinite speed'. It is worth mentioning that similar behaviour was restored in hydrodynamics by small droplet jumping on the surface of water [2], [3], [4]. The droplet corresponds to isolated quantum system and the surface of water is the background on which evolution of this system happens. In the next sections we provide suitable geometric interpretation of these equations.

In order to make the link between (3) and the Cartan connection consider the divergence-free condition or continuity equation [20], [5]

$$\nabla_a T^{ab} = 0, \quad b = 1 \dots n, \quad (7)$$

for energy-momentum tensor T^{ab} in a n -dimensional manifold M with coordinates (x^1, \dots, x^n) with some connection ∇ . In field theory it is assumed that the connection ∇ arises from metric structure on M . However, we can reinterpret (7) as a general divergence-free equation in a space where connection is not a Riemannian connection, but arises from a Lie scaling group. We elaborate this idea further. For fixed b , the vector $u^a := T^{ab}$ can be used to construct the 'flux-density' form [17]

$$\lambda = u_b \star dx^b, \quad (8)$$

where \star is the Hodge star induced by metric. Then (7) can be written in the form

$$d^\nabla \lambda = 0, \quad (9)$$

where d^∇ is covariant exterior derivative [12], which in flat space is usual exterior derivative in Riemannian case. This is exactly (2).

Now, in general case of connection for arbitrary Lie group, the equation (7) can be written in full form as

$$\nabla_i u^i = \partial_{x^i} u^i + \omega_j^i(\partial_{x^i}) u^j = 0, \quad (10)$$

for connection one-forms ω_b^a valued in some Lie algebra used as a gauge group. We want to stress again that in general relativity is is the Lorentz group, however we want to consider scaling group on jet space.

We are interested in recovering connection from (10) comparing it with (5), however, it is difficult task to find out connection (one-forms) ω knowing only the equation (5). Therefore, the procedure will be non-unique in general.

The phenomenon of interaction between particle and background is explicitly present in de Broglie-Bohm/pilot-wave approach to quantum mechanics [9]. The basic

idea is to assume that $u(x, t) = \sqrt{\rho(x, t)}e^{iS(x, t)}$ for two real functions ρ and S interpreted as probability and phase of u . Upon substituting into (1) and comparing real and imaginary parts we get coupled system

$$\begin{cases} \partial_t \rho + \partial_x(2\rho \partial_x S) = 0 \\ -\partial_t S = (\partial_x S)^2 + V + Q, \end{cases} \quad (11)$$

where $Q = -\frac{\partial_x^2 \sqrt{\rho}}{\sqrt{\rho}}$ is called the quantum potential. In this substitution the first equation describes conservation of the flux of probability ρ that flows with the velocity $2\rho \partial_x S$. The second is the Hamilton-Jacobi equation with 'quantum piloting' correction Q with addition to classical potential V . The Hamilton-Jacobi equations can be further converted to equations of motion for particle in effective $V + Q$ potential. The splitting in (6) is similar - n is a background and u represents the 'particle' moving on this background.

The paper is organized as follows: In the next section, we present the general procedure of extracting connection (or at least its part) from the arbitrary differential equation treating the equation as a divergence-free equation with connection given by a scaling group. This general procedure is then illustrated on the example of Schrödinger equation. The paper ends with a short discussion of the results, possible implications, and philosophical questions on the nature of quantum theory.

2 General theory

In this section, the general idea of recovering a Cartan connection from the partial differential equation (PDE) will be presented. This procedure is non-unique, as we will see. The next section describes the construction of specific connections for Schrödinger equation as an illustration. Therefore it is advisable to read this section along with one of the examples of the next section.

Consider the linear differential equation, where the part of equation that contains highest order derivatives of each variable is normalized and diagonalized. We will focus on operator of second order in x for simplicity, however the procedure can be applied to equations of any order. We have

$$\sum_{i=1}^n a_i \partial_i^2 u + A(u, \partial_i u) = 0, \quad (12)$$

where A is a linear function in u and $\partial_i u$ and $a_i = \pm 1$. We will treat this equation as a continuity equation (10) where the connection form is determined by some Lie group.

Since continuity equation (10) does not contain each coefficient, so from our PDE we will not be able to extract the connection in unique way, however we can set all not present coefficients to zero for simplicity. However, once we set up connection for one gauge and find the gauge group H then we get a geometric object that can be studied independently of PDE we started from. Here are detailed steps of the construction:

Step 1. Introduce the jet space. For treating (12) as (10) we have to treat all derivatives up to order 1 as variables, so we introduce the first jet space [17], [13], [14], [16], [15] on M , where e.g., $M = R^n$. For function u treated as a section of $E \rightarrow M$, where, e.g., $E = M \times \mathbb{R}$, we construct $J^1(M)$ where local coordinates are (x^i, v_0, u_i) , $i = 1 \dots n$. On this space we have prolongation of u to $J^1(M)$ given in

local coordinates by $j^1 u = (x^i, u(x), \partial_i u(x))$, so $v_0(j^1 u) = u$ and $v_i(j^1 u) = \partial_i u$. Then the equation (12) is

$$\sum_{i=1}^n a_i \partial_i v_i + A(v_i) = 0, \quad (13)$$

In general, in this step, by introducing suitable jet space, we 'flatten' r th order PDE to the first order PDE on jet space. The number of v_i coordinates also coincides with the number of initial/boundary data that we have to impose for the equation.

Step 2. Select a group G_0 . Due to motivation in previous section we introduce a Lie-Bäcklund group[17] on $J^1(M)$ that scales the differential term of (13) by some function $a \neq 0$. We take the commutative matrix group G with elements $g \in G_0$ of the form

$$g = \begin{bmatrix} a & 0 & \dots & \dots & 0 \\ b_1 & a & \ddots & \dots & 0 \\ b_2 & 0 & a & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ b_n & 0 & \dots & \dots & a \end{bmatrix}. \quad (14)$$

In general, G_0 does not have to be prolongation of the multiplication group $u \rightarrow au$, that is, we does not have to fix $b_i = \partial_i a$.

Applying the group to the equation (13) we get

$$a \sum_{i=1}^n a_i \partial_i v_i + A'(g, v, v_i) = 0, \quad (15)$$

where A' is still linear in v, v_i and depends on elements of matrix g and their derivatives.

Step 3. Consistency condition for connection. The next step is to calculate all possible coefficients of connection one-form. First, introduce metric $\eta = \text{diag}(a_1, \dots, a_n)$ and rewrite (15) as

$$a \eta^{ij} \partial_i v_j + A'(h, v, v_i) = 0. \quad (16)$$

Then the A' term is compared with (non-differential) Cartan connection part

$$\frac{1}{a} A'(h, v, v_i) = \omega_j^i (\partial_{x^i}) v_j. \quad (17)$$

Solution for ω_j^i is non-unique since the system is not determined. However, we can set, from practical reasons, all non-occurring in (17) coefficients of ω to zero. Some additional conditions will appear later from demanding torsion-freeness of the curvature of this connection.

In this step we can also embed the connection ω and the group G_0 inside the Weyl group G , which is a bookkeeping device for tracing the scale a . The extension is

$$g_0 \hookrightarrow \begin{bmatrix} a & 0 \\ 0 & g_0 \end{bmatrix}, \quad (18)$$

where a is the diagonal element of $g_0 \in G_0$ and 0 are zero vectors of dimension $n + 1$. Then the connection ω is then embedded in Weyl connection

$$\omega_W = \begin{bmatrix} \epsilon & 0 \\ \theta & \omega \end{bmatrix}, \quad (19)$$

where θ is a vector of dual elements to tangent base in $J^1(M)$, that is, $\theta = [dv_0, \dots, dv_n]^T$. In order to check that this is in fact the Weyl structure [18] we have to show that under commutative scaling group by b (that is a subgroup of G) the scaling part gauges as $\bar{\epsilon} = \epsilon + d \ln(b)$. This will be shown in the next step.

Step 4. Determination of the gauge group H . We want to find such subgroup H of G that the differential equation (15) transforms under $h \in H$ as connection [18], [1]

$$\omega'_W = Ad(h^{-1})\omega_W + h^*\omega_{MC}. \quad (20)$$

$h^*\omega_{MC}$ is a pullback of the Maurer-Cartan form [18] for H along h , which for matrix group is $h^{-1}dh$. Observe that the linear transformation is associated with Ad -transformed part and the Maurer-Cartan part involves derivatives of h . Therefore, the natural condition is that the part of (13) containing derivatives will transform under h as continuity equation for Maurer-Cartan connection, that is,

$$\eta^{ij} \partial_i(hv_j) = \lambda \eta^{ij} \partial_i(v_j) + B(h, v_j) = \lambda \eta^{ij} \partial_i(v_j) + \sum_i h^{-1} dh(\partial_i)u_i, \quad (21)$$

where λ is some arbitrary constant scale³, and B is linear in v_i . This is determining additional equation for matrix elements of h , so defines subgroup H of G . Note also that the Maurer-Cartan form is determined up to scale.

To this end, for checking consistency of procedure, we have to also show that Ad -part also transforms in the same way as $A'(h.v_i)$. We have for

$$h = \begin{bmatrix} e & 0 \\ 0 & A \end{bmatrix}, \quad (22)$$

that

$$Ad(h^{-1})\omega_W = \begin{bmatrix} \frac{1}{e} & 0 \\ 0 & A^{-1} \end{bmatrix} \begin{bmatrix} \epsilon & 0 \\ \theta & \omega \end{bmatrix} \begin{bmatrix} e & 0 \\ 0 & A \end{bmatrix} = \begin{bmatrix} \epsilon & 0 \\ sA^{-1}\theta & Ad(A^{-1})\omega \end{bmatrix}, \quad (23)$$

which under change of base $\bar{\theta} = sA^{-1}\theta$ is

$$Ad(h^{-1})\omega_W = \begin{bmatrix} \epsilon & 0 \\ \bar{\theta} & (Ad(A^{-1})\omega)A\frac{1}{s} \end{bmatrix} \quad (24)$$

Since group G is commutative so $Ad(A^{-1})\omega = \omega$ and we are left with lower-right block of the form $\frac{1}{s}\omega A$, which is exactly the same operation that is made on $A'(v_i)$ of (13) under transformation $v_i \rightarrow hv_i$. This shows that the gauge transformations on constructed connection are consistent with the same operations on differential equation.

Note also that the (1,1) block of (23) is $\bar{\epsilon} = \epsilon + d \ln(e)$, therefore, the defining condition for existence of Weyl structure is indeed fulfilled.

Step 5. Torsion-free condition. The curvature of (19) is [18], [1], [12]

$$\Omega_W := d\omega_W + \omega_W \wedge \omega_W = \begin{bmatrix} d\epsilon & 0 \\ d\theta + \theta \wedge \epsilon + \omega \wedge \theta & d\omega + \omega \wedge \omega \end{bmatrix}, \quad (25)$$

and the vanishing torsion condition is

$$d\theta + \theta \wedge \epsilon + \omega \wedge \theta = 0. \quad (26)$$

³We have a freedom to take arbitrary scale which we use here. We did not use it in the previous step for simplicity.

This condition imposes additional constraints on connection coefficients.

From the physical point of view, the vanishing torsion is a reasonable condition, which means that we require commutativity of parallel transport in jet space of solution. Since the initial vector in jet space is determined by initial data for the equation, so the torsion-free condition forces that the initial data propagates in the same way along any path on M .

Please also note that the torsion-freeness can be used to determine Weyl structure ϵ , since it is a 1-form.

3 Application to Schrödinger equation

In this section, we construct a Cartan connection using the method presented above. First, for clarity, we present the construction in $1 + 1$ dimension and then generalize construction to $1 + n$ case.

3.1 $1 + 1$ dimensional case

We will consider the following one-dimensional Schrödinger equation

$$i\partial_t u + \partial_x^2 u - V u = 0, \quad (27)$$

for $V = V(x, t)$ bounded from below. Underlying manifold is $M = \mathbb{R}^2$ and u can be considered as a section of $M \times \mathbb{C}^2 \rightarrow M$. We will follow the steps described in the previous section.

Step 1. The order of derivatives is $(1, 2)$ in (t, x) . Therefore, it is sufficient to introduce $J^1(M)$ with coordinates (t, x, v, v_1) , which on holonomic sections $j^1 u$ are $v(j^1 u) = u$, and $v_1(j^1 u) = \partial_x u$. On $J^1(M)$ there is also a coordinate associated with $\partial_t u$, however we do not need it and we omit it. This can be seen as a projection from J^1 to subspace where there is no 'time-derivative' coordinate. Note also that we can leave i factor next to derivative, which is associated with change $t \rightarrow t/i$, or we can take iv variable instead of v , however this introduces i in v_1 .

The equation in these coordinates is

$$i\partial_t v + \partial_x v_1 - V v = 0, \quad (28)$$

which is of first order.

Step 2. The group G_0 consists of 2×2 matrices g of the form

$$g = \begin{bmatrix} a & 0 \\ b & a \end{bmatrix}. \quad (29)$$

Under this element the equation transforms to

$$i\partial_t v + \partial_x v + v \frac{1}{a} (i\partial_0 a + \partial_x b - V a) + v_1 \frac{1}{a} (b + \partial_x a) = 0. \quad (30)$$

Step 3. The connection associated with G_0 has the form

$$\omega = \begin{bmatrix} \alpha & 0 \\ \beta & \alpha \end{bmatrix}, \quad (31)$$

where $\alpha = \alpha_t dt + \alpha_x dx$ and $\beta = \beta_t dt + \beta_x dx$. The continuity equation is

$$i\partial_t v + \partial_x v + v(\alpha_t + \beta_x) + v_1 \alpha_x = 0. \quad (32)$$

Comparing (30) and (32) we have system of equations for connection coefficients

$$\begin{cases} \alpha_t + \beta_x = \frac{1}{a}(\partial_x b - Va) \\ \alpha_x = \frac{1}{a}(b + \partial_x a) \end{cases} \quad (33)$$

Note indeterminism of this system - we have the following symmetry

$$\begin{cases} \alpha_t \rightarrow \alpha_t + f(x, t)dt \\ \beta_x \rightarrow \beta_x - f(x, t)dx \\ \beta_t = g(x, t)dt, \end{cases} \quad (34)$$

for some arbitrary functions f and g . We can choose, e.g. $\alpha_t = \frac{1}{a}i\partial_0 a dt$ and $\beta_t = 0$. In this way we have determined non-uniquely a connection matrix ω .

We can also embed it in Weyl structure. The Weyl connection is

$$\omega_W = \begin{bmatrix} \epsilon & 0 & 0 \\ dv & \frac{1}{a}(b + \partial_x a)dx + \frac{1}{a}i\partial_0 a dt & 0 \\ dv_1 & \frac{1}{a}(\partial_x b - Va)dx & \frac{1}{a}(b + \partial_x a)dx + \frac{1}{a}i\partial_0 a dt \end{bmatrix}, \quad (35)$$

for some ϵ to be determined later.

Step 4. We will now construct a subgroup H_0 of G_0 with element h of the form

$$h = \begin{bmatrix} e & 0 \\ f & e \end{bmatrix}. \quad (36)$$

The Maurer-Cartan form for this element is

$$h^* \omega_{MC} = h^{-1} dh = \begin{bmatrix} d \ln(e) & 0 \\ d \frac{f}{e} & d \ln(e) \end{bmatrix} \quad (37)$$

We want to associate in part containing the highest order derivatives in the equation with (37). From (28) we get after using h

$$i\partial_t(hv) + \partial_x(hv_1) = i\partial_t v + \partial_x v_1 + v \frac{1}{e}(i\partial_t e + \partial_x f) + v_1 \frac{1}{e}(f + \partial_x e) = 0, \quad (38)$$

and from the continuity equation for connection given by (37) we obtain

$$i\partial_t v + \partial_x v_1 + v(\frac{1}{e}\partial_t e + \partial_x(f/e)) + v_1 \frac{1}{e}\partial_x e = 0. \quad (39)$$

The coefficients at v and v_1 should be proportional⁴ and therefore we get the system

$$\begin{cases} \frac{1}{e}(i\partial_t e + \partial_x f) = \lambda(\frac{1}{e}\partial_t e + \partial_x(f/e)) \\ \frac{1}{e}(f + \partial_x e) = \lambda \frac{1}{e}\partial_x e \end{cases} \quad (40)$$

These system can be rewritten as

$$\begin{cases} f = (\lambda - 1)\partial_x e \\ (i - \lambda)\partial_t \ln(e) - (\lambda - 1)(\partial_x \ln(e))^2 = \partial_x^2 \ln(e) \end{cases} \quad (41)$$

which is a constraint for group manifold H . Note that for $\lambda = 1$ we get $f = 0$ and H_0 consist of diagonal matrices.

⁴The linear equation has symmetry of scaling by nonzero constant factor λ , which is a special case of group we consider.

Step 5. Starting from (35) and using the fact that we use holonomic coordinate system $\theta = [dv, dv_1]^T$, that is $d\theta = 0$, we get vanishing torsion condition in the form

$$0 = \theta \wedge \epsilon + \omega \wedge \theta = \begin{bmatrix} \epsilon \\ \epsilon \end{bmatrix} + \begin{bmatrix} \alpha & 0 \\ \beta & \alpha \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}. \quad (42)$$

This gives $\epsilon = -\alpha = -(\partial_x \ln(a)dx + \frac{1}{a}i\partial_0adt)$ and $\beta = 0$. Therefore, torsion-free condition is

$$0 = \frac{1}{a}(\partial_x b - Va). \quad (43)$$

It is understand that once we choose a , then this equation determines b , or vice versa.

We can consider special cases, when the group G_0 is a prolongation to $J^1(M)$ of the multiplication group $u \rightarrow au$, i.e., $b = \partial_x a$. We get then

$$\frac{1}{a}\partial_x^2 a - V = 0, \quad (44)$$

which is exactly the condition from (6) for $a = n$. This suggest that the form of (6) has fundamental geometric meaning. However this condition changes, when we choose differently α_t and β_x to fulfil the equation (33).

3.2 $1 + n$ dimensional case

The $1 + n$ ($n > 0$) case is a straightforward generalization of previous subsection.

Step 1. We have $M = \mathbb{R}^{1+n}$ with coordinates (t, x^1, \dots, x^n) . Then for the functions $u : M \rightarrow \mathbb{C}$, which can be seen as sections of $M \times \mathbb{C} \rightarrow M$, we construct jet bundle $J^1(M)$ with coordinates $(t, x, v, v_1, \dots, v_n)$, where we have $v_i(j^1u) = \partial_{x^i}u$. As before, the coordinate associated with $\partial_t u$ is excluded by the projection.

The Schrödinger equation is

$$i\partial_t v + \sum_{i=1}^n \partial_i v_i - Vv = 0. \quad (45)$$

Step 2. The group G_0 contains matrices of the form

$$g = \begin{bmatrix} a & 0 & \dots & \dots & 0 \\ b_1 & a & \ddots & \dots & 0 \\ b_2 & 0 & a & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ b_n & 0 & \dots & \dots & a \end{bmatrix}, \quad (46)$$

with connection one-form

$$\omega = \begin{bmatrix} \alpha & 0 & \dots & \dots & 0 \\ \beta_1 & \alpha & \ddots & \dots & 0 \\ \beta_2 & 0 & \alpha & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ \beta_n & 0 & \dots & \dots & \alpha \end{bmatrix}, \quad (47)$$

with $\alpha = \alpha_t dt + \sum_{i=1}^n \alpha_i dx^i$ and $b_i = \beta_i dt + \sum_{j=1}^n \beta_{ij} dx^j$.

Step 3. The equation after transformation by g is

$$i\partial_t v + \sum_{i=1}^n \partial_i v_i + v \frac{1}{a} (i\partial_t a + \sum_{i=1}^n \partial_i b_i - Va) + \sum_{i=1}^n v_i \frac{1}{a} (b_i + \partial_i a) = 0, \quad (48)$$

and the continuity equation for connection (47) is

$$i\partial_t v + \sum_{i=1}^n \partial_i v_i + v \left(\alpha_t + \sum_{i=1}^n \beta_{ii} \right) + \sum_{i=1}^n v_i \alpha_i = 0. \quad (49)$$

Comparing coefficients at v and v_i we obtain an undetermined system of equations for connection

$$\begin{cases} (\alpha_t + \sum_{i=1}^n \beta_{ii}) = \frac{1}{a} (i\partial_t a + \sum_{i=1}^n \partial_i b_i - Va) \\ \alpha_i = \frac{1}{a} (b_i + \partial_i a). \end{cases} \quad (50)$$

Here the choice of connection coefficient is also non-unique. Therefore, we will do it in analogical way as for 1 + 1 case, and symmetrize potential V among β coefficients, i.e.,

$$\begin{cases} \alpha = \frac{1}{a} i\partial_t a dt + \sum_{i=1}^n \frac{1}{a} (b_i + \partial_i a) dx^i \\ \beta_i = \frac{1}{a} (\partial_i b_i - \frac{1}{n} Va) dx^i \quad (\text{no summation}). \end{cases} \quad (51)$$

The embedding of connection into the Weyl connection is as before

$$\omega_W = \begin{bmatrix} \epsilon & 0 \\ \theta & \omega \end{bmatrix}, \quad (52)$$

where $\theta = [dv, dv_1, \dots, dv_n]^T$, and the Weyl structure ϵ will be determined later by torsion-free condition.

Step 4. We will now look for H_0 subgroup of G_0 that we match with by differential part of the equation with the Maurer-Cartan form for H_0 . Element $h \in H_0$ is of the form

$$h = \begin{bmatrix} e & 0 & \dots & \dots & 0 \\ f_1 & e & \ddots & \dots & 0 \\ f_2 & 0 & e & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ f_n & 0 & \dots & \dots & e \end{bmatrix}, \quad (53)$$

and the Maurer-Cartan form for h is

$$h^* \omega_{MC} = h^{-1} dh = \begin{bmatrix} d\ln(e) & 0 & \dots & \dots & 0 \\ d\frac{f_1}{e} & d\ln(e) & \ddots & \dots & 0 \\ d\frac{f_2}{e} & 0 & d\ln(e) & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ d\frac{f_n}{e} & 0 & \dots & \dots & d\ln(e) \end{bmatrix}. \quad (54)$$

The differential part of (45) under h is changed to

$$i\partial_t v + \sum_{i=1}^n \partial_i v_i + v \frac{1}{e} (i\partial_t e + \sum_{i=1}^n \partial_i f_i) + \sum_{i=1}^n v_i \frac{1}{e} (f_i + \partial_i e) = 0, \quad (55)$$

which we compare with continuity equation for the Maurer-Cartan form of H_0 , that is,

$$i\partial_t v + \sum_{i=1}^n \partial_i v_i + v \left(\partial_t \ln(e) + \sum_{i=1}^n \partial_i (f_i/e) \right) + \sum_{i=1}^n v_i \partial_i \ln(e) = 0. \quad (56)$$

The coefficients at v , v_i of both equations should be proportional by some constant λ , and therefore we have matching conditions

$$\begin{cases} f_i = (\lambda - 1)\partial_i e \\ (i - \lambda)\partial_i \ln(e) - \sum_{i=1}^n (\partial_i \ln(e))^2 = (\lambda - 1) \sum_{i=1}^n \partial_i^2 \ln(e), \end{cases} \quad (57)$$

which gives conditions defining H_0 .

Step 5. The final step is to impose torsion-free condition. Since we are using holonomic coordinates, so $d\theta = 0$, and the condition is

$$0 = \theta \wedge \epsilon + \omega \wedge \theta = \begin{bmatrix} \epsilon \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \epsilon \end{bmatrix} + \begin{bmatrix} \alpha & 0 & \dots & \dots & 0 \\ \beta_1 & \alpha & \ddots & \dots & 0 \\ \beta_2 & 0 & \alpha & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ \beta_n & 0 & \dots & \dots & \alpha \end{bmatrix} \begin{bmatrix} 1 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ 1 \end{bmatrix}. \quad (58)$$

That gives $\epsilon = -\alpha$ and $\beta_i = 0$. This means that for our specific choice of connection we have

$$\partial_i b_i - \frac{1}{n} V a = 0, \quad i = 1, \dots, n, \quad (59)$$

or

$$\sum_{i=1}^n \partial_i b_i = V a. \quad (60)$$

This can be treated as equation for b when a is some selected function or vice versa.

Again, for G_0 that is lift of multiplication group on u , that is $b_i = \partial_i a$, we get condition in the form

$$\triangle a = V a. \quad (61)$$

This is n -dimensional version of (4). Again, different choices for α and β results in change of this condition.

3.3 Other equations

Concluding this section, we also comment on other PDEs.

The results of this section easily extend to the diffusion equation by changing $i \rightarrow 1$. The Schrödinger equation is vital due to motivation and physical context, however, the same procedure can be applied to other PDEs.

4 Conclusions

The above construction of the Cartan connection for the scaling group of wave functions shows that the Schrödinger equation is of geometric type. Since it is associated with the continuity equation, therefore it defines only some of the connection components. The other components must be fixed otherwise. This shows that the

Schrödinger equation has big freedom of selecting the connection one-form, and therefore, the freedom to be geometrized. The torsion-free condition for the curvature of the connection imposes additional conditions on connection and makes the connection to belonging to the diagonal subgroup of scalings on jet space.

The above geometric approach is present in the first jet space where components of the vector are the wave function and its gradient. However, knowing the Cartan connection, we can also ask how arbitrary vector in jet space, that apart of the wave function has other components that are not necessary gradient components of the wave function, evolves. Therefore, by the construction described in the paper, we significantly extended degrees of freedom of quantum mechanics.

As it was mentioned in the Introduction, the scaling function (sometimes non-normalizable) appears in many places in quantum mechanics. The approach presented in this paper assigns to this scaling a geometric object that can be associated with the background on which quantum evolution takes place. We believe that it is the first step to a better understanding of the nature of quantum vacuum and construct powerful non-perturbative geometric methods for solving quantum-mechanical problems.

The procedure of constructing the Cartan connection from Schrödinger equation is non-unique because the equation is treated as a continuity equation, which is some kind of 'divergence-free' condition of some object. Therefore, this approach suggests that some additional information must be imposed if quantum mechanics from this viewpoint should be considered as a geometric theory of Cartan connection. This raises a serious question on completeness of quantum theory described by Schrödinger equation. In this approach, the same problem appears if we consider general PDE and try to treat it as a continuity equation. From a philosophical point of view, this lack of additional input perhaps may be attributed to the fact that the continuity equation describes how we can manipulate of a given quantity of 'material' to preserve it, however, it does not describe what kind of 'material' on which we manipulate. Therefore to describe of nature of this 'material' additional set of laws has to be provided.

From a mathematical point of view, the construction of the Cartan connection for PDE was provided. The procedure is non-unique, however, once performed, it can give a geometric interpretation of the equation.

Acknowledgments

These ideas in the paper grew for many years during discussions. I would like to thanks Andrew Waldron for inspiring discussion on gauge theories and quantum mechanics during 38th Winter School of Geometry and Physics in Srni, to Josef Silhan and Jan Slovák for support and for showing me a beautiful world of differential geometry of Cartan connection, and to Jarek Duda for discussion on nature of space and time. Last but not least, I would like to thanks Anatoliy K. Prykarpatsky for continuous motivation, and Rikard von Unge and Klaus Bering Larsen for discussion on the nature of the quantum world.

The author was supported by the GACR grant GA19-06357S, the grant 8J20DE004 of Ministry of Education, Youth and Sports of the CR, and Masaryk University grant MUNI/A/0885/2019. I would like to acknowledge COST CA18223 action for support.

References

- [1] A. Čap, Jan Slovák, *Parabolic Geometries: Background and General Theory*, (AMS, 2009)
- [2] Y. Couder, A. Boudaoud, S. Protière, E. Fort, *Walking droplets, a form of wave-particle duality at macroscopic scale?*, Europhysics News 41, 1, (2010) 14–18; DOI: <https://doi.org/10.1051/epn/2010101>
- [3] Y. Couder, S. Protière, E. Fort, A. Boudaoud, *Walking and orbiting droplets*, Nature 437, 208 (2005); DOI: <https://doi.org/10.1038/437208a>
- [4] Y. Couder, E. Fort, C.-H. Gautier, A. Boudaoud, *From Bouncing to Floating: Noncoalescence of Drops on a Fluid Bath*, Phys. Rev. Lett. 94, 177801 (2005); DOI: <https://doi.org/10.1103/PhysRevLett.94.177801>
- [5] T. Frankel, *Gravitational Curvature*, (Dover Publications, 2011)
- [6] C.R. Hagen, *Scale and conformal transformations in galilean-covariant field theory*, Phys.Rev.D 5 377–388 (1972); DOI: 10.1103/PhysRevD.5.377
- [7] G. Herczeg, A. Waldron, *Contact Geometry and Quantum Mechanics*, Physics Letters B, Volume 781, p. 312-315, 2018; DOI: 10.1016/j.physletb.2018.04.008
- [8] G. Herczeg, E. Latini, A. Waldron, *Contact Quantization: Quantum Mechanics = Parallel transport*, arXiv:1805.11731 [hep-th]
- [9] B.J. Hiley, D. Bohm, *The Undivided Universe: An Ontological Interpretation of Quantum Theory*, (Routledge, 1995)
- [10] A. Jadczyk, J. Janyška, M. Modugno, *Galilei general relativistic quantum mechanics revisited*, in "Geometria, Física-Matemática e outros Ensaios", Homenagem a António Ribeiro Gomes, A. S.Alves, F. J. Craveiro de Carvalho and J. A. Pereira da Silva Eds., Coimbra 1998, 253–313
- [11] J. Janyška, M. Modugno, *Covariant Schrödinger operator*, Jour. Phys. A.: Math. Gen.,35, 1–28, (2002)
- [12] I. Kolář, P.W. Michor, J. Slovák, *Natural Operations in Differential Geometry*, (Springer, 1993)
- [13] I.S. Krasil'shchik, A.M.Vinogradov (Eds.), *Symmetries and Conservation Laws for Differential Equations*, (AMS, 1999)
- [14] A. Kushner, V. Lychagin, V. Rubtsov, *Contact Geometry and Nonlinear Differential Equations*, (Cambridge University Press, 2007)
- [15] R.A. Kycia, M. Ulan, E. Schneider (Eds.), *Nonlinear PDEs, Their Geometry, and Applications*, (Birkhäuser Basel, 2019)
- [16] P.J.F. Olver, *Applications Of Lie Groups To Differential Equations*, 2nd edition (Springer, 2000)
- [17] W.-H. Steeb, *Continuous Symmetries, Lie Algebras, Differential Equations and Computer Algebra*, 2nd edition, (World Scientific, 2007)
- [18] R.W.F. Sharpe, *Differential Geometry: Cartan's Generalization of Klein's Erlangen Program*, (Springer, 2000)
- [19] A. Trautman, *Einstein-Cartan Theory*, Encyclopedia of Mathematical Physics, edited by J.-P. Francoise, G.L. Naber and Tsou S.T. Oxford: Elsevier, vol. 2, pages 189–195, 2006

- [20] R.M. Wald, *General Relativity*, (University of Chicago Press, 1984)
- [21] H. Weyl, *Eine neue Erweiterung der Relativitätstheorie*, Ann. der Physik 364, 10, 101–133, (1919)
- [22] H. Weyl, *Space, Time, Matter*, (Dover Publications, 1952)
- [23] P.-M. Zhang, M. Elbistan, P.A. Horvathy, P. Kosinski, *A generalized Noether theorem for scaling symmetry*, Eur.Phys.J.Plus 135, 2, 223 (2020); DOI: 10.1140/epjp/s13360-020-00247-5