

# On the Problem of Steady Bifurcation of a Falling Sphere in a Navier-Stokes Liquid

Giovanni P Galdi

Department of Mechanical Engineering and Materials Science  
University of Pittsburgh, USA

## Abstract

We study steady bifurcation for the coupled system body-liquid consisting of a sphere freely falling in a Navier-Stokes liquid under the action of gravity. In particular we show that, under the assumption that for the bifurcating solution the translational velocity of the sphere is parallel to the gravity, bifurcation takes place provided 1 is a simple eigenvalue of a suitable linear operator and the transversality property holds. Moreover, we also give sufficient conditions for symmetry breaking.

## Introduction

Bifurcation –steady and time-periodic– is a common phenomenon in fluid mechanics that, over the past several decades, has been the object of numerous rigorous investigations. Typical and significant examples are the so called Taylor–Couette and Bénard–Rayleigh problems, for which one can provide a rather complete picture of steady-state bifurcation; see, e.g., [2], [16], [25, Sections 72.7–72.9] and the reference therein.

The fundamental issue that one has to address when investigating this kind of questions is to find the right *functional setting* that allows for the use of general abstract results. In this regard, it should be pointed out that the steady bifurcation problems previously mentioned, as many of those collected in classical fluid mechanics literature, regard flow occurring in a *bounded* spatial region. In such a case, the problem is naturally formulated as a nonlinear equation in a suitable Banach space, with the relevant operator being a compact perturbation of the identity. As a consequence, the corresponding linearization possesses a purely discrete spectrum that enables one to provide bifurcation criteria in terms of the spectral properties of the 0 eigenvalue, like simplicity and transversality [22, Chapter IV].

However, if the liquid occupies a spatial region that is *unbounded* in all directions, as in the flow past an obstacle, the above functional setting is no longer available because of lack of compactness of the nonlinear operator and, in addition, the linearized operator has a large essential spectrum containing 0 [4]. The general problem of bifurcation from the essential spectrum has been addressed by several authors; see [23] and the references therein. Though of great interest, the theory there developed, however, does not apply to the flow of a liquid past a body, because it requires a self-adjointness property that is not satisfied in such a case.

Recently, we have undertaken a systematic study of bifurcation, in both steady and time-periodic cases, of a Navier–Stokes flow past a body [13], [8], [10], [11], [12]. In the particular case of steady bifurcation, in order to overcome the issue of the essential spectrum, we found

it more appropriate to formulate the problem in the “natural” function class where the steady-state Navier–Stokes problem is well set. This class consists of “graded” (homogeneous) Sobolev spaces, characterized by requiring different summability properties of the various derivatives involved [8], [10]. In doing so, we have been able to furnish necessary as well as sufficient conditions for the occurrence of steady bifurcation from a (steady-state) Navier–Stokes flow past a rigid body. It should be emphasized that all the results established in [8], [10] are obtained under the assumption that the motion of the rigid body is *prescribed*.

Objective of the present paper is to present a steady bifurcation theory in the case when the motion of the body is *not* prescribed, thus becoming a further unknown. To our knowledge, this is the first contribution to the rigorous study of bifurcation in a liquid-solid interaction problem. More specifically, we shall study the problem of a sphere of constant density that is steadily and freely falling in a Navier–Stokes liquid under the action of gravity,  $\mathbf{g}$ . Here, the driving mechanism is represented by the dimensionless buoyancy,  $\lambda$  (*Galilei number*), that we take to be positive, namely, the sphere is falling, not rising. The basic motion  $\mathbf{s}_0$ , of the coupled system consists then of a constant translation of the sphere with velocity  $\boldsymbol{\xi}$  parallel to  $\mathbf{g}$ , while the liquid executes a corresponding flow of steady-state nature, when referred to a frame translating with velocity  $\boldsymbol{\xi}$  as well. Such a flow is rotationally symmetric around the direction of  $\boldsymbol{\xi}$ . There is an abundant literature about this problem from both experimental and numerical viewpoints (see [24, 20, 17, 18, 6, 3] and the references therein) showing that there exists a critical value  $\lambda_0$  of Galilei number at which the flow of the liquid is no longer rotationally symmetric, and a new flow sets in that only possesses planar symmetry. Our goal is to find a suitable functional setting that allows us to furnish necessary and sufficient conditions for the existence of such  $\lambda_0$ . Precisely, under the assumption that the bifurcating solution also translates in the direction of  $\mathbf{g}$ , we show that the problem can be correctly formulated in a suitable Hilbert space,  $\mathcal{H}$ , constituted by functions having a finite Dirichlet integral and reducing to a rigid motion on the boundary; see (1.5). The equation for the bifurcating branch is then written as a nonlinear equation in  $\mathcal{H}$ , where the relevant operator,  $F$ , suitably defined on a dense set of  $\mathcal{H}$ , can be written as the sum of the identity plus a nonlinear operator that, however, unlike the case of a bounded domain, is *not* compact; see (1.21). Nevertheless, we show that the derivative of  $F$  at criticality is Fredholm of index 0 (Lemma 2.4) and this allows us to apply general bifurcation results. In this way we prove that a necessary condition for the occurrence of bifurcation at a certain  $\lambda_0$ , is that 1 is an eigenvalue for a suitable linear operator  $\lambda M = \lambda M(\lambda)$  evaluated at  $\lambda = \lambda_0$ ; see (2.17) and Remark 3.2. In order to make this assumption meaningful, we show that, in fact, the intersection of the spectrum of  $M(\lambda)$  with the positive real semi-line is constituted only by eigenvalues with finite multiplicity, for each fixed  $\lambda > 0$ ; see Lemma 2.6. We then prove that the above condition is also sufficient, provided 1 is simple and the transversality condition holds; see Theorem 3.1. It seems quite remarkable that, *formally*, we can formulate bifurcation conditions very similar to those given for flow in a bounded region. Finally, we address the question of symmetry breaking. As suggested by numerical test [6], this certainly happens if, at the critical value  $\lambda = \lambda_0$ , the sphere start spinning transversally. Based on this considerations, we furnish a sufficient condition for the transversal component of the angular velocity to be non-zero at criticality, which guarantees symmetry breaking; see Theorem 3.2.

We end this introductory section by pointing out a significant open question. It concerns the assumption that in the bifurcating solution the sphere translates parallel to the gravity. We need this hypothesis for merely technical reasons and we believe that it can be removed. However, to date, the proof of the latter seems to be away from our reach.

The plan of the paper is as follows. In Section 1, after formulating the problem, we show that the bifurcation problem can be written as a nonlinear operator equation in an appropriate Hilbert space. Successively, in Section 2, we study the relevant properties of the operators involved in the equation. Finally, in Section 3, we give a proof of the main results.

## 1 Formulation of the Problem and its Functional Setting

Consider a sphere,  $\mathcal{S}$ , of constant density  $\rho_{\mathcal{S}}$  and radius  $R$ , freely falling under the action of gravity in an otherwise quiescent Navier–Stokes liquid,  $\mathcal{L}$ . Let  $\mathcal{F} = \{O, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  be the frame with the origin at the center of  $\mathcal{S}$  and the axis  $\mathbf{e}_1$  oriented along the acceleration of gravity  $\mathbf{g}$ . The steady-state motions of the coupled system  $\mathcal{S} \cup \mathcal{L}$  in  $\mathcal{F}$  are then governed by the following set of nondimensional equations [7, Section 4]

$$\left. \begin{aligned} \Delta \mathbf{v} + \lambda \boldsymbol{\xi} \cdot \nabla \mathbf{v} &= \lambda \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p \\ \operatorname{div} \mathbf{v} &= 0 \end{aligned} \right\} \quad \text{in } \Omega$$

$$\mathbf{v} = \boldsymbol{\xi} + \boldsymbol{\omega} \times \mathbf{x} \quad \text{at } \partial\Omega, \quad \lim_{|\mathbf{x}| \rightarrow \infty} \mathbf{v}(\mathbf{x}) = \mathbf{0}, \quad (1.1)$$

$$\int_{\partial\Omega} \mathbb{T}(\mathbf{v}, p) \cdot \mathbf{n} = \lambda \mathbf{e}_1, \quad \int_{\partial\Omega} \mathbf{x} \times \mathbb{T}(\mathbf{v}, p) \cdot \mathbf{n} = 0.$$

Here  $\mathbf{v}, p$  are (non-dimensional) velocity and pressure fields of  $\mathcal{L}$ , while  $\boldsymbol{\xi}, \boldsymbol{\omega}$  represent (non-dimensional) translational and angular velocities of  $\mathcal{S}$ . Moreover,  $\lambda = \sqrt{\alpha g R^3} / \nu$  ( $> 0$ ) is the (dimensionless) Galilei number, where  $\alpha = (\rho_{\mathcal{S}} / \rho_{\mathcal{L}} - 1)$  with  $\rho_{\mathcal{L}}$  density of the liquid, and  $\nu$  its kinematic viscosity. We are assuming that the sphere has a negative buoyancy, so that  $\alpha > 0$ . Finally,  $\Omega$  is the region occupied by the liquid (the exterior of  $\mathcal{S}$ ), and

$$\mathbb{T}(\mathbf{v}, p) = -p \mathbb{1} + 2 \mathbb{D}(\mathbf{v}), \quad \mathbb{D}(\mathbf{v}) := \frac{1}{2} (\nabla \mathbf{v} + (\nabla \mathbf{v})^\top),$$

is the Cauchy tensor with  $\mathbb{1}$  identity tensor and  $\top$  denoting transpose.

On physical ground, it is expected that, for any (positive) value of the Galilei number, (1.1) always admits a solution where  $\mathcal{S}$  moves with no spin and translational velocity directed along the gravity. In this regard, in [7, Theorem 4.7] –completed with the results of [9, Section X.6]– the following theorem is proved.

**Theorem 1.1** *For any given  $\lambda > 0$ , problem (1.1) has at least one solution  $\mathbf{s}_0 = \mathbf{s}_0(\lambda) := (\mathbf{v}_0, p_0, \boldsymbol{\xi}_0, \boldsymbol{\omega}_0)(\lambda)$  such that<sup>1</sup>*

$$\begin{aligned} \mathbf{v}_0 &\in D^{2,s}(\Omega) \cap D^{1,r}(\Omega) \cap L^q(\Omega) \cap C^\infty(\Omega), ; \\ p_0 &\in D^{1,s}(\Omega) \cap L^{s_1}(\Omega) \cap C^\infty(\Omega), \\ \boldsymbol{\xi}_0 &= \xi_0 \mathbf{e}_1, \quad \xi_0 > 0; \quad \boldsymbol{\omega}_0 = \mathbf{0}, \end{aligned} \quad (1.2)$$

all  $s > 1, r > 4/3, q > 2, s_1 > 3/2$ , and satisfying the “energy equality”

$$\|\mathbb{D}(\mathbf{v}_0)\|_2^2 = \lambda \xi_0. \quad (1.3)$$

*This solution is rotationally symmetric around the  $x_1$ -axis.*

---

<sup>1</sup>We use standard notation:  $L^q$  is the Lebesgue space with norm  $\|\cdot\|_q$ , and  $W^{1,2}$  is Sobolev space. Furthermore,  $D^{k,t}$  represent homogeneous Sobolev spaces with semi-norm  $\sum_{|l|=k} \|D^l u\|_t$ .

The main question we want to address is to establish necessary and sufficient conditions for the occurrence of steady bifurcation for the branch  $\mathbf{s}_0 = \mathbf{s}_0(\lambda)$ , at some Galilei number  $\lambda = \lambda_0$ , when the translational velocity remains parallel to the gravity direction  $\mathbf{e}_1$ .

In order to accomplish this goal, we proceed as follows. In the first place, we will equivalently rewrite (1.1) as an operator equation in an appropriate function space. Successively, we will show certain fundamental characteristics of the involved operator, notably, its Fredholm property. In this way, we shall finally be able to apply classical abstract bifurcation theorems and obtain the desired results.

We begin to introduce some function spaces. Let

$$\begin{aligned}\mathcal{C} &= \mathcal{C}(\Omega) = \{\boldsymbol{\varphi} \in C_0^\infty(\overline{\Omega}) : \operatorname{div} \boldsymbol{\varphi} = 0 \text{ in } \Omega; \\ &\quad \boldsymbol{\varphi}(x) = \xi_{\boldsymbol{\varphi}} \mathbf{e}_1 + \boldsymbol{\omega}_{\boldsymbol{\varphi}} \times \mathbf{x}, \quad \xi_{\boldsymbol{\varphi}} \in \mathbb{R}, \quad \boldsymbol{\omega}_{\boldsymbol{\varphi}} \in \mathbb{R}^3, \text{ in a neighborhood of } \partial\Omega\}, \\ \mathcal{C}_0 &= \mathcal{C}_0(\Omega) := \{\boldsymbol{\varphi} \in \mathcal{C} : \boldsymbol{\varphi} = \mathbf{0} \text{ at } \partial\Omega\},\end{aligned}$$

and define

$$\begin{aligned}\mathcal{H} &= \mathcal{H}(\Omega) \equiv \{\text{completion of } \mathcal{C}(\Omega) \text{ in the norm } \|\mathbb{D}(\cdot)\|_2\}, \\ \mathcal{H}_0 &= \mathcal{H}_0(\Omega) \equiv \{\text{completion of } \mathcal{C}_0(\Omega) \text{ in the norm } \|\mathbb{D}(\cdot)\|_2\}\end{aligned}$$

The proof of the next lemma is given in [7, Lemmas 9–11].

**Lemma 1.1**  *$\mathcal{H}$  is a Hilbert space endowed with the scalar product*

$$[\mathbf{u}, \mathbf{w}] = \int_{\Omega} \mathbb{D}(\mathbf{u}) : \mathbb{D}(\mathbf{w}), \quad \mathbf{u}, \mathbf{w} \in \mathcal{H}, \quad (1.4)$$

and the following characterizations hold

$$\mathcal{H} = \{\mathbf{u} \in W_{\text{loc}}^{1,2}(\overline{\Omega}) : \mathbf{u} \in L^6(\Omega), \mathbb{D}(\mathbf{u}) \in L^2(\Omega); \operatorname{div} \mathbf{u} = 0 \text{ in } \Omega; \mathbf{u}(y) = \xi_{\mathbf{u}} \mathbf{e}_1 + \boldsymbol{\omega}_{\mathbf{u}} \times \mathbf{y}, y \in \partial\Omega\}. \quad (1.5)$$

and

$$\mathcal{H}_0 = \{\mathbf{u} \in \mathcal{H} : \mathbf{u} = \mathbf{0} \text{ at } \partial\Omega\}.$$

Moreover, we have

$$\|\nabla \mathbf{u}\|_2 \leq \sqrt{2} \|\mathbb{D}(\mathbf{u})\|_2 \leq 2 \|\nabla \mathbf{u}\|_2, \quad (1.6)$$

and

$$\|\mathbf{u}\|_6 \leq c_0 \|\mathbb{D}(\mathbf{u})\|_2, \quad \mathbf{u} \in \mathcal{H}, \quad (1.7)$$

for some numerical constant  $c_1 > 0$ .<sup>2</sup> Finally, there is another positive numerical constant  $c_0$  such that

$$|\xi_{\mathbf{u}}| + |\boldsymbol{\omega}_{\mathbf{u}}| \leq c_1 \|\mathbb{D}(\mathbf{u})\|_2. \quad (1.8)$$

Let  $\mathcal{H}^{-1}$  be the (strong) dual of  $\mathcal{H}$ , and denote by  $\langle \cdot, \cdot \rangle$  and  $\|\cdot\|_{-1}$  the corresponding duality pair and associated norm, respectively. We then introduce the following space that will play a fundamental role in our work:<sup>3</sup>

$$\mathcal{X}(\Omega) := \{\mathbf{u} \in \mathcal{H}(\Omega) : \partial_1 \mathbf{u} \in \mathcal{H}^{-1}\}$$

---

<sup>2</sup>Recall that, in our non-dimensionalization, the sphere has radius 1.

<sup>3</sup> $\partial_1 \equiv \partial/\partial x_1$ .

where  $\partial_1 \mathbf{u} \in \mathcal{H}^{-1}$  means that there is  $C = C(\mathbf{u}) > 0$  such that

$$|(\partial_1 \mathbf{u}, \varphi)| \leq C \|\mathbb{D}(\varphi)\|_2, \quad \text{for all } \varphi \in \mathcal{C}. \quad (1.9)$$

In fact, if (1.9) holds, since  $\mathcal{C}$  is dense in  $\mathcal{H}$ , by the Hahn–Banach theorem  $\partial_1 \mathbf{u}$  can be uniquely extended to a bounded linear functional on the whole of  $\mathcal{H}$ , with

$$\|\partial_1 \mathbf{u}\|_{-1} := \sup_{\varphi \in \mathcal{C}; \|\mathbb{D}(\varphi)\|_2 = 1} |(\partial_1 \mathbf{u}, \varphi)|.$$

It is readily shown that the functional

$$\|\mathbf{u}\|_{\mathcal{X}} := \|\mathbb{D}(\mathbf{u})\|_2 + \|\partial_1 \mathbf{u}\|_{-1},$$

defines a norm in  $\mathcal{X}(\Omega)$ . We have the following property whose proof is entirely analogous to [10, Proposition 65] and therefore it will be omitted.

**Lemma 1.2** *The space  $\mathcal{X}(\Omega)$  endowed with the norm  $\|\cdot\|_{\mathcal{X}}$  is a reflexive, separable Banach space.*

We also have.

**Lemma 1.3** *The space  $\mathcal{X}(\Omega)$  is continuously embedded in  $L^4(\Omega)$ . Moreover, there is  $c > 0$  such that*

$$\|\mathbf{u}\|_4 \leq c (\|\partial_1 \mathbf{u}\|_{-1}^{\frac{1}{4}} \|\mathbb{D}(\mathbf{u})\|_2^{\frac{3}{4}} + \|\mathbb{D}(\mathbf{u})\|_2).$$

*Proof.* For a given  $\mathbf{u} \in \mathcal{X}(\Omega)$ , we set

$$\mathbf{V} = -\frac{1}{2} \text{curl} [\text{curl} (\zeta \xi_{\mathbf{u}} \mathbf{e}_1 x_2^2) + \zeta |\mathbf{x}|^2 \boldsymbol{\omega}_{\mathbf{u}}],$$

where  $\zeta$  is a function in  $C_0^\infty(\overline{\Omega})$  that is 1 near  $\partial\Omega$ . Clearly,  $\mathbf{V}$  is smooth with bounded support and

$$\text{div } \mathbf{V} = 0 \text{ in } \Omega, \quad \mathbf{V}(x) = \xi_{\mathbf{u}} + \boldsymbol{\omega}_{\mathbf{u}} \times \mathbf{x} \text{ at } \partial\Omega,$$

so that  $\mathbf{V} \in \mathcal{C}$ . Moreover, by a straightforward calculation and with the help of Lemma 1.1, one shows

$$\|\mathbf{V}\|_4 + \|\nabla \mathbf{V}\|_2 + \|\partial_1 \mathbf{V}\|_{-1} \leq C (|\xi_{\mathbf{u}}| + |\boldsymbol{\omega}_{\mathbf{u}}|) \leq C_1 \|\mathbb{D}(\mathbf{u})\|_2. \quad (1.10)$$

We next write

$$\mathbf{u} = \mathbf{u} - \mathbf{V} + \mathbf{V} := \mathbf{w} + \mathbf{V}. \quad (1.11)$$

and observe that, also by the above properties of  $\mathbf{V}$ , we have

$$\mathbf{w} \in \mathcal{H}_0. \quad (1.12)$$

Furthermore, from the identity  $\Delta \varphi = 2 \text{div } \mathbb{D}(\varphi)$ ,  $\varphi \in \mathcal{C}_0$ , and the density of  $\mathcal{C}_0$  in  $\mathcal{H}_0$  (see Lemma 1.1), we easily obtain

$$\|\nabla \mathbf{w}\|_2 = \sqrt{2} \|\mathbb{D}(\mathbf{w})\|_2, \quad \mathbf{w} \in \mathcal{H}_0. \quad (1.13)$$

Thus, in particular, with the help of (1.10), (1.11), we get

$$\begin{aligned} |\partial_1 \mathbf{w}|_{-1} &:= \sup_{\varphi \in \mathcal{C}_0; \|\nabla \varphi\|_2 = 1} |(\partial_1 \mathbf{w}, \varphi)| \leq \frac{1}{\sqrt{2}} \sup_{\varphi \in \mathcal{C}_0; \|\mathbb{D}(\varphi)\|_2 = 1} |(\partial_1 \mathbf{w}, \varphi)| \\ &\leq \frac{1}{\sqrt{2}} \|\partial_1 \mathbf{w}\|_{-1} \leq C (\|\partial_1 \mathbf{u}\|_{-1} + \|\mathbb{D}(\mathbf{u})\|_2). \end{aligned} \quad (1.14)$$

From (1.12)–(1.14) and [8, Proposition 1.1], we then conclude,  $\mathbf{w} \in L^4(\Omega)$  which, in turn, by (1.10) and (1.11) implies  $\mathbf{u} \in L^4(\Omega)$ . Furthermore, again by [8, Proposition 1.1] we have

$$\|\mathbf{w}\|_4 \leq c_1 \left( |\partial_1 \mathbf{w}|_{-1}^{\frac{1}{4}} \|\nabla \mathbf{w}\|_2^{\frac{3}{4}} + \|\nabla \mathbf{w}\|_2 \right),$$

that once combined with (1.10), (1.11), (1.13) and (1.14) completes the proof of the lemma.  $\square$

With the above two lemmas in hand, we are now able to write (1.1) as an operator equation in the space  $\mathcal{H}^{-1}$ . To do this, we introduce the following *notation*:

For  $\varphi \in \mathcal{H}$ , we denote by  $\overline{\varphi} := \xi_{\mathbf{u}} \mathbf{e}_1 + \boldsymbol{\omega}_{\mathbf{u}} \times \mathbf{x}$  its trace at  $\partial\Omega$ .

Consider now the operators

$$\begin{aligned} \mathbf{S} : \mathbf{v} \in \mathcal{X}(\Omega) &\mapsto \mathbf{S}\mathbf{v} \in \mathcal{H}^{-1}, \\ \mathbf{g} : \mathbf{v} \in \mathcal{X}(\Omega) &\mapsto \mathbf{g} \in \mathcal{H}^{-1}, \\ \mathbf{N} : \mathbf{v} \in \mathcal{X}(\Omega) &\mapsto \mathbf{N}(\mathbf{v}) \in \mathcal{H}^{-1}, \end{aligned}$$

where

$$\begin{aligned} \langle \mathbf{S}\mathbf{v}, \varphi \rangle &:= (\mathbb{D}(\mathbf{v}), \mathbb{D}(\varphi)); \\ \langle \mathbf{g}, \varphi \rangle &:= \xi_{\varphi}; \\ \langle \mathbf{N}(\mathbf{v}), \varphi \rangle &:= (\mathbf{v} \cdot \mathbb{D}(\varphi), \mathbf{v}) + \xi_{\mathbf{v}} \langle \partial_1 \mathbf{v}, \varphi \rangle; \end{aligned} \quad \varphi \in \mathcal{H}. \quad (1.15)$$

By using Lemma 1.1, Lemma 1.3 and Schwarz inequality, it is at once established that all the operators introduced above are well defined. Furthermore, it is readily checked that problem (1.1) can be reformulated as the following operator equation

$$\mathbf{S}\mathbf{v} - \lambda \mathbf{g} - \lambda \mathbf{N}(\mathbf{v}) = 0 \quad \text{in } \mathcal{H}^{-1}. \quad (1.16)$$

Actually, if we dot-multiply by  $\varphi \in \mathcal{C}$  both sides of (1.1)<sub>1</sub>, integrate by parts over  $\Omega$  and take into account (1.1)<sub>2,3,5,6</sub> we get

$$(\mathbb{D}(\mathbf{v}), \mathbb{D}(\varphi)) - \lambda \xi_{\varphi} = \lambda (\mathbf{v} \cdot \mathbb{D}(\varphi), \mathbf{v}) - \lambda \int_{\partial\Omega} \overline{\mathbf{v}} \cdot \mathbf{n} (\overline{\mathbf{v}} \cdot \overline{\varphi}) + \lambda \xi_{\mathbf{v}} (\partial_1 \mathbf{v}, \varphi), \quad \varphi \in \mathcal{C}. \quad (1.17)$$

Denote by  $\mathcal{I}$  the surface integral in (1.17). Observing that, for  $x \in \partial\Omega$ ,  $\mathbf{x}$  and  $\mathbf{n}$  are parallel vectors, and  $\int_{\partial\Omega} \mathbf{n} = \mathbf{0}$ , we get

$$\mathcal{I} = \int_{\partial\Omega} [\xi_{\mathbf{v}}^2 \mathbf{e}_1 \cdot \mathbf{n} \mathbf{e}_1 \cdot (\boldsymbol{\omega}_{\varphi} \times \mathbf{x}) + \xi_{\varphi} \xi_{\mathbf{v}} \mathbf{e}_1 \cdot \mathbf{n} \mathbf{e}_1 \cdot (\boldsymbol{\omega}_{\mathbf{v}} \times \mathbf{x}) + \xi_{\mathbf{v}} \mathbf{e}_1 \cdot \mathbf{n} (\boldsymbol{\omega}_{\varphi} \times \mathbf{x}) \cdot (\boldsymbol{\omega}_{\mathbf{v}} \times \mathbf{x})] := \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3.$$

Recalling that  $\mathbb{R}^3 - \overline{\Omega} = \Omega_0$ , by Gauss theorem it follows that

$$\mathcal{I}_1 = \xi_v^2 \int_{\Omega_0} \mathbf{e}_1 \cdot \nabla(\boldsymbol{\omega}_\varphi \times \mathbf{x}) \cdot \mathbf{e}_1 = \xi_v^2 \int_{\Omega_0} \mathbf{e}_1 \cdot \mathbb{D}(\boldsymbol{\omega}_\varphi \times \mathbf{x}) \cdot \mathbf{e}_1 = 0$$

and, analogously,

$$\mathcal{I}_2 = 0.$$

Moreover, again by Gauss theorem,

$$\mathcal{I}_3 = \xi_v \int_{\Omega_0} [\partial_1(\boldsymbol{\omega}_v \times \mathbf{x}) \cdot (\boldsymbol{\omega}_\varphi \times \mathbf{x}) + \partial_1(\boldsymbol{\omega}_\varphi \times \mathbf{x}) \cdot (\boldsymbol{\omega}_v \times \mathbf{x})] = 0.$$

Thus,

$$\int_{\partial\Omega} \bar{\mathbf{v}} \cdot \mathbf{n}(\bar{\mathbf{v}} \cdot \bar{\boldsymbol{\varphi}}) = 0 \quad (1.18)$$

and, as a result, (1.17), by density, gives (1.16). Conversely, (1.16) implies (1.17), and so choosing in (1.17)  $\varphi \in \mathcal{C}_0$  we obtain

$$(\mathbb{D}(\mathbf{v}), \mathbb{D}(\varphi)) = \lambda(\mathbf{v} \cdot \mathbb{D}(\varphi), \mathbf{v}) + \lambda \xi_v (\partial_1 \mathbf{v}, \varphi), \quad \varphi \in \mathcal{C},$$

which, after integrating by parts and using the arbitrariness of  $\varphi \in \mathcal{C}_0$  shows that there exists a suitable pressure field  $p$  such that  $(\mathbf{v}, p, \xi_v \mathbf{e}_1, \boldsymbol{\omega}_v)$  satisfies (1.1)<sub>1,2,3</sub>. If we integrate by parts the first and the third term in (1.17), the latter furnishes

$$(\operatorname{div} \mathbb{T}(\mathbf{v}, p) + \lambda \partial_1 \mathbf{v} - \mathbf{v} \cdot \nabla \mathbf{v}, \varphi) = \int_{\partial\Omega} \bar{\boldsymbol{\varphi}} \cdot \mathbb{T}(\mathbf{v}, p) \cdot \mathbf{n} - \lambda \xi_\varphi, \quad \text{for all } \varphi \in \mathcal{C}.$$

However, the left-hand side of this equation vanishes because  $\mathbf{v}, p$  and  $\xi_v$  satisfy (1.1)<sub>1</sub>, so that the right-hand side vanishes as well and, by the arbitrariness of  $\varphi$ , this implies that also (1.1)<sub>4,5</sub> are satisfied.

Our main goal is to investigate bifurcation of the branch  $\mathbf{s}_0(\lambda) := (\mathbf{v}_0, p_0, \boldsymbol{\xi}_0, \mathbf{0})(\lambda)$  obtained in Theorem 1.1. In this regard, we premise the following lemma.

**Lemma 1.4** *Let  $\mathbf{s}_0(\lambda) := (\mathbf{v}_0, p_0, \boldsymbol{\xi}_0, \mathbf{0})(\lambda)$  be the solution in Theorem 1.1 corresponding to  $\lambda > 0$ . Then  $\mathbf{v}_0 \in \mathcal{X}(\Omega)$ .*

*Proof.* We only need to show

$$\|\partial_1 \mathbf{v}_0\|_{-1} < \infty. \quad (1.19)$$

We recall that  $\mathbf{s}_0$  solves the following problem

$$\left. \begin{aligned} \Delta \mathbf{v}_0 + \lambda \xi_0 \partial_1 \mathbf{v}_0 &= \lambda \mathbf{v}_0 \cdot \nabla \mathbf{v}_0 + \nabla p_0 \\ \operatorname{div} \mathbf{v}_0 &= 0 \end{aligned} \right\} \quad \text{in } \Omega$$

$$\mathbf{v}_0 = \xi_0 \mathbf{e}_1 \quad \text{at } \partial\Omega, \quad \lim_{|\mathbf{x}| \rightarrow \infty} \mathbf{v}_0(\mathbf{x}) = \mathbf{0}, \quad (1.20)$$

$$\int_{\partial\Omega} \mathbb{T}(\mathbf{v}_0, p_0) \cdot \mathbf{n} = \lambda \mathbf{e}_1, \quad \int_{\partial\Omega} \mathbf{x} \times \mathbb{T}(\mathbf{v}_0, p_0) \cdot \mathbf{n} = 0.$$

If we dot-multiply both sides of (1.20)<sub>1</sub> by  $\varphi \in \mathcal{C}$ , integrate by parts over  $\Omega$  and use (1.20)<sub>2,3,5,6</sub> we get

$$\lambda \xi_0 (\partial_1 \mathbf{v}_0, \varphi) = (\mathbb{D}(\mathbf{v}_0), \mathbb{D}(\varphi)) - \lambda \xi_\varphi - \lambda (\mathbf{v}_0 \cdot \mathbb{D}(\varphi), \mathbf{v}_0).$$

Thus, using in the latter Schwarz inequality and Lemma 1.1, we infer

$$\lambda \xi_0 |(\partial_1 \mathbf{v}_0, \boldsymbol{\varphi})| \leq c \left( \|\mathbb{D}(\mathbf{v}_0)\|_2 + \lambda + \lambda \|\mathbf{v}_0\|_4^2 \right) \|\mathbb{D}(\boldsymbol{\varphi})\|_2,$$

and since, by Theorem 1.1,  $\mathbf{v}_0 \in L^4(\Omega)$ , the proof of the lemma is completed.  $\square$

We thus write in (1.16)  $\mathbf{v} = \mathbf{v}_0 + \mathbf{u}$  to obtain the following equation for  $\mathbf{u} \in \mathcal{X}(\Omega)$

$$\mathbf{L}_\lambda \mathbf{u} - \lambda \mathbf{K}_{\mathbf{v}_0} \mathbf{u} - \lambda \mathbf{N}(\mathbf{u}) = 0 \quad \text{in } \mathcal{H}^{-1}, \quad (1.21)$$

where, for all  $\boldsymbol{\varphi} \in \mathcal{H}$ ,

$$\begin{aligned} \langle \mathbf{L}_\lambda \mathbf{u}, \boldsymbol{\varphi} \rangle &:= (\mathbf{S} \mathbf{u}, \boldsymbol{\varphi}) - \lambda \xi_0 \langle \partial_1 \mathbf{u}, \boldsymbol{\varphi} \rangle \\ \langle \mathbf{K}_{\mathbf{v}_0} \mathbf{u}, \boldsymbol{\varphi} \rangle &:= 2(\mathbf{u} \cdot \mathbb{D}(\boldsymbol{\varphi}), \mathbf{v}_0) + \xi_{\mathbf{u}} \langle \partial_1 \mathbf{v}_0, \boldsymbol{\varphi} \rangle. \end{aligned} \quad (1.22)$$

**Remark 1.1** Since, in general  $\mathbf{v}_0$  is a function of the Galilei number  $\lambda$ , so is  $\mathbf{K}_{\mathbf{v}_0}$ . Whenever needed, we shall emphasize this property by writing  $\mathbf{K}_{\mathbf{v}_0}(\lambda)$ .  $\triangle$

It is plain that  $(\lambda_0, \mathbf{v}_0(\lambda_0))$  is a bifurcation point for (1.15) if and only if  $(\lambda_0, \mathbf{0})$  is a bifurcation point for (1.21). Therefore, our problem reduces to find a branch of non-trivial solutions  $(\lambda, \mathbf{u}(\lambda))$  to (1.21) in a neighborhood of  $(\lambda_0, \mathbf{0})$ . To reach this goal, we need several preparatory results concerning the relevant functional properties of the operators defined in (1.21).

## 2 Preparatory Results

We begin with the following.

**Lemma 2.1** *For any  $\mathbf{u} \in \mathcal{X}(\Omega)$  we have*

$$\langle \partial_1 \mathbf{u}, \mathbf{u} \rangle = 0.$$

*Proof.* From the splitting (1.11) and (1.10) we get  $\mathbf{w}, \mathbf{V} \in \mathcal{X}(\Omega)$  and

$$\langle \partial_1 \mathbf{u}, \mathbf{u} \rangle = \langle \partial_1 \mathbf{w}, \mathbf{w} \rangle + (\partial_1 \mathbf{w}, \mathbf{V}) + (\partial_1 \mathbf{V}, \mathbf{w}) + (\partial_1 \mathbf{V}, \mathbf{V}). \quad (2.1)$$

By integrating by parts and recalling that  $\mathbf{w} = 0$  at  $\partial\Omega$  we show

$$(\partial_1 \mathbf{w}, \mathbf{V}) + (\partial_1 \mathbf{V}, \mathbf{w}) = 0. \quad (2.2)$$

Also, recalling that  $\mathbb{R}^3 - \overline{\Omega} = \Omega_0$ , by another integration by parts we obtain

$$(\partial_1 \mathbf{V}, \mathbf{V}) = \frac{1}{2} \int_{\partial\Omega} \overline{\mathbf{u}} \cdot \overline{\mathbf{u}} \overline{\mathbf{u}} \cdot \mathbf{n} = \int_{\Omega_0} \overline{\mathbf{u}} \cdot \mathbb{D}(\overline{\mathbf{u}}) \cdot \overline{\mathbf{u}} = 0. \quad (2.3)$$

Thus, from (2.1)–(2.3) it follows that

$$\langle \partial_1 \mathbf{u}, \mathbf{u} \rangle = \langle \partial_1 \mathbf{w}, \mathbf{w} \rangle.$$

However,  $\mathbf{w} \in \mathcal{H}_0$  with  $|\partial_1 \mathbf{w}|_{-1} < \infty$  (see (1.14)). Therefore, from [8, Proposition 1.2] we infer  $\langle \partial_1 \mathbf{w}, \mathbf{w} \rangle = 0$ , which concludes the proof.  $\square$



**Lemma 2.2** *For any fixed  $\lambda > 0$ , the operator  $K_{v_0} = K_{v_0}(\lambda)$  is compact.*

*Proof.* For sake of simplicity, in what follows we will not distinguish between a given sequence and its subsequences. Thus, suppose  $\{\mathbf{u}_n\} \subset \mathcal{X}(\Omega)$  with  $\|\mathbf{u}_n\|_{\mathcal{X}} \leq M$  and  $M$  independent of  $n$ . By Lemma 1.3 this implies, in particular,

$$\|\mathbf{u}_n\|_4 + \|\mathbb{D}(\mathbf{u}_n)\|_2 \leq M_1, \quad (2.4)$$

for another  $M_1$  independent of  $n$ . Therefore, by Lemma 1.3, there exists  $\mathbf{u} \in \mathcal{X}(\Omega)$  such that

$$\mathbf{u}_n \rightharpoonup \mathbf{u} \text{ in } \mathcal{X}(\Omega).$$

Also, from (1.6), (2.4) and classical compact embedding theorems, we deduce

$$\mathbf{u}_n \rightarrow \mathbf{u} \text{ in } L^4(\Omega_R), \text{ for all } R > R_*, \quad (2.5)$$

By (2.4) and (1.8) there exist  $\xi_0 \in \mathbb{R}$ ,  $\omega_0 \in \mathbb{R}^3$  such that

$$\xi_{\mathbf{u}_n} \rightarrow \xi_*, \quad \omega_{\mathbf{u}_n} \rightarrow \omega_* \text{ in } \mathbb{R}. \quad (2.6)$$

However, by trace theorems (e.g. [9, Theorem II.4.1]) and (1.6) we also have

$$\int_{\partial\Omega} |\xi_{\mathbf{u}_n} - \xi_{\mathbf{u}} + (\omega_{\mathbf{u}_n} - \omega_{\mathbf{u}}) \times \mathbf{x}|^2 \leq c(\|\mathbf{u}_n - \mathbf{u}\|_{4,\Omega_R}^2 + \|\mathbf{u}_n - \mathbf{u}\|_{4,\Omega_R} \|\mathbb{D}(\mathbf{u}_n - \mathbf{u})\|_2). \quad (2.7)$$

Thus, passing to the limit  $n \rightarrow \infty$  in (2.7) and employing (2.4)–(2.6) we infer

$$\xi_{\mathbf{u}_n} \rightarrow \xi_{\mathbf{u}}, \quad \omega_{\mathbf{u}_n} \rightarrow \omega_{\mathbf{u}} \text{ in } \mathbb{R}. \quad (2.8)$$

Consequently, setting  $\mathbf{w}_n := \mathbf{u}_n - \mathbf{u}$ ,  $\mu_n := \xi_{\mathbf{u}_n} - \xi_{\mathbf{u}}$ , from (1.22) with the help of Schwarz inequality we prove

$$\|K_{v_0} \mathbf{w}_n\|_{-1} \leq \|\mathbf{w}_n\|_{4,\Omega_R} \|\mathbf{v}_0\|_4 + \|\mathbf{w}_n\|_4 \|\mathbf{v}_0\|_{4,\Omega^R} + |\mu_n| \|\partial_1 \mathbf{v}_0\|_{-1}.$$

If we let  $n \rightarrow \infty$  into this relation and use (2.4), (2.5) and (2.8) we get

$$\limsup_{n \rightarrow \infty} \|K_{v_0} \mathbf{w}_n\|_{-1} \leq M_1 \|\mathbf{v}\|_{4,\Omega^R},$$

which, in turn, by setting  $R \rightarrow \infty$  and using the absolute continuity of Lebesgue integral, proves the lemma. □

**Lemma 2.3** *The operator  $S \equiv L_0$  defined in (1.22)<sub>1</sub> is a homeomorphism of  $\mathcal{H}$  onto  $\mathcal{H}^{-1}$ , whereas if  $\lambda \neq 0$ ,  $L_\lambda$  is a homeomorphism of  $\mathcal{X}(\Omega)$  onto  $\mathcal{H}^{-1}$ .*

*Proof.* By Riesz theorem, for any  $\mathbf{f} \in \mathcal{H}^{-1}$  there is a unique  $\mathbf{u} \in \mathcal{H}$  such that

$$[\mathbf{u}, \boldsymbol{\varphi}] = \langle \mathbf{f}, \boldsymbol{\varphi} \rangle, \quad \text{all } \boldsymbol{\varphi} \in \mathcal{H},$$

with  $[\cdot, \cdot]$  defined in (1.4), which proves the first part of the lemma. To show the second part, it is enough to show that for any  $\mathbf{f} \in \mathcal{H}^{-1}$  there is a unique  $\mathbf{u} \in \mathcal{H}$  verifying

$$(\mathbb{D}(\mathbf{u}), \mathbb{D}(\boldsymbol{\varphi})) - \lambda \xi_0 (\partial_1 \mathbf{u}, \boldsymbol{\varphi}) = \langle \mathbf{f}, \boldsymbol{\varphi} \rangle, \quad \text{for all } \boldsymbol{\varphi} \in \mathcal{C} \quad (2.9)$$

and such that

$$\|\mathbb{D}(\mathbf{u})\|_2 \leq \|\mathbf{f}\|_{-1}. \quad (2.10)$$

In fact, from (2.9), (2.10) it also follows

$$\lambda \xi_0 |(\partial_1 \mathbf{u}, \varphi)| \leq 2\|\mathbf{f}\|_{-1} \|\mathbb{D}(\varphi)\|_2, \quad \text{for all } \varphi \in \mathcal{C},$$

which proves  $\mathbf{u} \in \mathcal{X}(\Omega)$ . Let  $\{\varphi_k\} \subset \mathcal{C}$  be a basis in  $\mathcal{H}$  with  $[\varphi_k, \varphi_{k'}] = \delta_{kk'}$ . In view of (1.7), we have that for any given  $\varphi \in \mathcal{H}$ , there is  $\{\gamma_k\} \subset \mathbb{R}$  such that setting  $\Phi_N = \sum_{k=1}^N \gamma_k \varphi_k$ , it follows

$$\|\mathbb{D}(\Phi_N - \varphi)\|_2 + \|\Phi_N - \varphi\|_6 \rightarrow 0 \text{ as } N \rightarrow \infty. \quad (2.11)$$

We look for an “approximating” solution to (2.9) of the type

$$\mathbf{u}_m = \sum_{\ell=1}^m c_{\ell m} \varphi_\ell$$

where the coefficients  $c_{\ell m}$  are solutions to the system

$$(\mathbb{D}(\mathbf{u}_m), \mathbb{D}(\varphi_k)) - \lambda \xi_0 (\partial_1 \mathbf{u}_m, \varphi_k) = \langle \mathbf{f}, \varphi_k \rangle, \quad k = 1, \dots, m, \quad (2.12)$$

or, equivalently,

$$\sum_{k=1}^n (\delta_{\ell k} - \lambda \xi_0 \mathcal{A}_{\ell k}) c_{\ell m} = \langle \mathbf{f}, \varphi_k \rangle, \quad k = 1, \dots, m, \quad (2.13)$$

where

$$\mathcal{A}_{\ell k} = (\partial_1 \varphi_\ell, \varphi_k).$$

Integrating by parts, we have

$$\mathcal{A}_{\ell k} = -\mathcal{A}_{k\ell} + \int_{\partial\Omega} \mathbf{e}_1 \cdot \mathbf{n} \overline{\varphi_\ell} \cdot \overline{\varphi_k}$$

However, arguing as in the proof of (1.18), we show

$$\int_{\partial\Omega} \mathbf{e}_1 \cdot \mathbf{n} \overline{\varphi_\ell} \cdot \overline{\varphi_k} = 0, \quad (2.14)$$

which implies that  $\mathcal{A}_{\ell k}$  is skew-symmetric. As a result [9, Lemma IX.3.1], (2.13) has a unique solution  $c_{\ell k}$ ,  $k = 1, \dots, m$ , or, equivalently, (2.12) has a unique solution  $\mathbf{u}_m$  for each  $m \in \mathbb{N}$ . Multiplying both sides of (2.12) by  $c_{\ell k}$ , summing over  $k$  from 1 to  $m$  and using (2.14) entails

$$\|\mathbb{D}(\mathbf{u}_m)\|_2^2 = \langle \mathbf{f}, \mathbf{u}_m \rangle$$

from which we easily deduce

$$\|\mathbb{D}(\mathbf{u}_m)\|_2 \leq \|\mathbf{f}\|_{-1}. \quad (2.15)$$

Thus, there is  $\mathbf{u} \in \mathcal{H}$  such that  $\mathbf{u}_m \rightharpoonup \mathbf{u}$  in  $\mathcal{H}$  and, moreover,  $\mathbf{u}$  satisfies (2.10). Also, passing to the limit  $m \rightarrow \infty$  in (2.12) we show

$$(\mathbb{D}(\mathbf{u}), \mathbb{D}(\varphi_k)) - \lambda \xi_0 (\partial_1 \mathbf{u}, \varphi_k) = \langle \mathbf{f}, \varphi_k \rangle, \quad \text{for all } k \in \mathbb{N}. \quad (2.16)$$

We can now replace in (2.16)  $\varphi_k$  with linear combinations of the type  $\Phi_N$  given in (2.11). Therefore, taking the limit  $N \rightarrow \infty$  we obtain that actually  $\mathbf{u}$  satisfies (2.9), which completes the proof of existence. Suppose, next, that

$$\langle \mathbb{L}_\lambda \mathbf{u}, \varphi \rangle \equiv (\mathbb{D}(\mathbf{u}), \mathbb{D}(\varphi)) - \lambda \xi_0 \langle \partial_1 \mathbf{u}, \varphi \rangle = 0, \text{ for all } \varphi \in \mathcal{H}.$$

If we set  $\varphi = \mathbf{u}$  into this relation and use Lemma 2.1 we get  $\|\mathbb{D}(\mathbf{u})\|_2 = 0$ , namely,  $\mathbf{u} = 0$ , and uniqueness follows.  $\square$

From Lemma 2.2 and Lemma 2.3 we deduce, in particular, the following property.

**Lemma 2.4** *For any fixed  $\rho \neq 0$ , and  $\lambda > 0$  the operator  $\mathbb{L}_\rho - \rho \mathbb{K}_{\mathbf{v}_0}(\lambda)$  is Fredholm of index 0.*

We next recall that, by Lemma 2.3,  $\mathbb{S}$  is an homeomorphism from  $\mathcal{H}$  onto  $\mathcal{H}^{-1}$ . We may then introduce the operator

$$\mathbb{M} : \mathbf{u} \in D(\mathbb{M}) \equiv \mathcal{X}(\Omega) \subset \mathcal{H} \mapsto \mathbb{S}^{-1}[\xi_0 \partial_1 \mathbf{u} - \mathbb{K}_{\mathbf{v}_0} \mathbf{u}] \in \mathcal{H}. \quad (2.17)$$

We recall that, in general, the operator  $\mathbb{M}$  is a function of the Galilei number  $\lambda$ ; see Remark 1.1. In the next lemmas we shall show some relevant properties of  $\mathbb{M}$ .

**Lemma 2.5** *For each fixed  $\lambda > 0$ , the operator  $\mathbb{M} = \mathbb{M}(\lambda)$  is densely defined and closed.*

*Proof.* Since  $\mathcal{C} \subset \mathcal{X}(\Omega) \subset \mathcal{H}$ , the density property is obvious. Let  $\{\mathbf{v}_k\} \subset \mathcal{X}(\Omega)$ ,  $\mathbf{m}_k := \mathbb{M}\mathbf{v}_k$ , with  $\mathbf{v}_k \rightarrow \mathbf{v}$  and  $\mathbf{m}_k \rightarrow \mathbf{u}$  in  $\mathcal{H}$ , for some  $\mathbf{v}, \mathbf{u} \in \mathcal{H}$ . Let us prove that, in fact,  $\mathbf{v}_k \rightarrow \mathbf{v}$  in  $\mathcal{X}(\Omega)$ . Recalling the definition of  $\mathbb{S}$ , we obtain that the relation  $\mathbb{M}(\mathbf{v}_k - \mathbf{v}_{k'}) = \mathbf{m}_k - \mathbf{m}_{k'}$  is equivalent to

$$\xi_0 \langle \partial_1(\mathbf{v}_k - \mathbf{v}_{k'}), \varphi \rangle - 2(\mathbf{v}_0, \mathbb{D}(\varphi), (\mathbf{v}_k - \mathbf{v}_{k'})) - (\xi_k - \xi_{k'}) \langle \partial_1 \mathbf{v}_0, \varphi \rangle = (\mathbb{D}(\mathbf{m}_k - \mathbf{m}_{k'}), \mathbb{D}(\varphi)). \quad (2.18)$$

for all  $\varphi \in \mathcal{H}$ . Employing Hölder inequality on the right hand side of (2.18), we show

$$\xi_0 \|\partial_1(\mathbf{v}_k - \mathbf{v}_{k'})\|_{-1} \leq 2\|\mathbf{v}_0\|_3 \|\mathbf{v}_k - \mathbf{v}_{k'}\|_6 + |\xi_k - \xi_{k'}| \|\partial \mathbf{v}_0\|_{-1} + \|\mathbb{D}(\mathbf{m}_k - \mathbf{m}_{k'})\|_2. \quad (2.19)$$

Since  $(\mathbf{v}_k - \mathbf{v}_{k'}) \rightarrow 0$  in  $\mathcal{H}$ , by Lemma 1.1 we know that

$$(\mathbf{v}_k - \mathbf{v}_{k'}) \rightarrow 0 \text{ in } L^6(\Omega); \quad (\xi_k - \xi_{k'}) \rightarrow 0 \text{ in } \mathbb{R},$$

and also, from Theorem 1.1 and Lemma 1.4,  $\|\mathbf{v}_0\|_3 + \|\partial \mathbf{v}_0\|_{-1} < \infty$ . Therefore, letting  $k, k' \rightarrow \infty$  it follows that  $\mathbf{v} \in \mathcal{X}(\Omega) \equiv D(\mathbb{M})$ , and  $\mathbf{v}_k \rightarrow \mathbf{v}$  in  $\mathcal{X}(\Omega)$ . In view of the latter, by formally setting in (1.10)  $\mathbf{v}_{k'} \equiv \mathbf{v}$ ,  $\mathbf{m}_{k'} \equiv \mathbf{u}$  and then passing to the limit  $k \rightarrow \infty$ , we conclude  $\mathbb{M}\mathbf{v} = \mathbf{u}$  which completes the proof that  $\mathbb{M}$  is closed.  $\square$

**Lemma 2.6** *For any fixed  $\lambda > 0$  and  $\mu \neq 0$  the operator*

$$\mathbb{H}_\mu = \mu \mathbb{I} - \lambda \mathbb{M}(\lambda)$$

*is Fredholm of index 0. Furthermore, denoting by  $\mathcal{H}_c$  the complexification of  $\mathcal{H}$ , by  $\mathbb{M}_c$  the natural extension of  $\mathbb{M}$  to  $\mathcal{H}_c$  and by  $\sigma(\mathbb{M}_c)$  the spectrum of  $\mathbb{M}_c$ , we have that  $\sigma(\mathbb{M}_c) \cap (0, \infty)$  consists at most of a countable number of eigenvalues of finite algebraic multiplicity that can only cluster at 0.*

*Proof.* Since

$$\mathbf{H}_\mu = \mu \mathbf{S}^{-1} (\mathbf{L}_\mu - \frac{1}{\mu} \mathbf{K} \mathbf{v}_0) := \mathbf{S}^{-1} \mathbf{T}_\mu, \quad (2.20)$$

and, by Lemma 2.4,  $\mathbf{T}_\mu$  is Fredholm of index 0, we have

$$\dim N[\mathbf{H}_\mu] = \dim N[\mathbf{T}_\mu] = m < \infty.$$

Moreover, from

$$\mathcal{H}^{-1} = R(\mathbf{T}_\mu) \oplus S_m$$

with  $S_m$   $m$ -dimensional subspace, we deduce that for every  $\mathbf{w} \in \mathcal{H}$ , it is  $\mathbf{S}\mathbf{w} = \mathbf{w}_1 + \mathbf{w}_2$ ,  $\mathbf{w}_1 \in R(\mathbf{T}_\mu)$ ,  $\mathbf{w}_2 \in S_m$ . Therefore,  $\mathbf{w} = \mathbf{S}^{-1}\mathbf{w}_1 + \mathbf{S}^{-1}\mathbf{w}_2$ , with  $\mathbf{S}^{-1}\mathbf{w}_1 \in R(\mathbf{H}_\mu)$ , and  $\mathbf{S}^{-1}\mathbf{w}_2 \in \mathbf{S}^{-1}S_m$ , which completes the proof. Clearly, the natural complexification  $\mathbf{H}_{\mu c}$ , of  $\mathbf{H}_\mu$  is also Fredholm of index 0, for all  $\mu > 0$ . It then follows that the essential spectrum  $\sigma_{\text{ess}}(\mathbf{M}_c)$ , of  $\mathbf{M}_c$  defined as the set of  $\mu$  where  $\mathbf{H}_{\mu c}$  is not Fredholm has empty intersection with  $(0, \infty)$ . We shall next show that the resolvent set  $P(\mathbf{M}_c)$  of  $\mathbf{M}_c$  has a non-empty intersection with  $(0, \infty)$ . Since  $\mathbf{H}_\mu$  is Fredholm of index 0, it is enough to show that, for sufficiently large  $\mu > 0$ , it is  $N[\mathbf{H}_\mu] = \{0\}$ . From (2.20), we see that the latter is equivalent to

$$\mu (\mathbb{D}(\mathbf{u}), \mathbb{D}(\boldsymbol{\varphi})) - \xi_0 \langle \partial_1 \mathbf{u}, \boldsymbol{\varphi} \rangle - 2(\mathbf{u} \cdot \mathbb{D}(\boldsymbol{\varphi}), \mathbf{v}_0) - \xi_{\mathbf{u}} \langle \partial_1 \mathbf{v}_0, \boldsymbol{\varphi} \rangle = 0, \quad \boldsymbol{\varphi} \in \mathcal{H}.$$

Choosing  $\boldsymbol{\varphi} = \mathbf{u}$  in this relation and using Lemma 2.1, (1.7) and (1.8) along with Hölder inequality furnishes

$$\mu \|\mathbb{D}(\mathbf{u})\|_2^2 \leq C (\|\mathbf{v}_0\|_3 + \|\partial_1 \mathbf{v}_0\|_{-1}) \|\mathbb{D}(\mathbf{u})\|_2^2.$$

Thus, if  $\mu > C (\|\mathbf{v}_0\|_3 + \|\partial_1 \mathbf{v}_0\|_{-1}) \equiv \bar{\mu}$ , we conclude  $\mathbf{u} \equiv \mathbf{0}$ , namely,  $P(\mathbf{M}_c) \cap (\bar{\mu}, \infty) \neq \emptyset$ . Summarizing, we have shown that  $\sigma_{\text{ess}}(\mathbf{M}_c) \cap (0, \infty) = \emptyset$  while  $P(\mathbf{M}_c) \cap (\bar{\mu}, \infty) \neq \emptyset$ . Therefore, the stated property about eigenvalues is a consequence of classical results in spectral theory [14, Theorem XVII.2]. □

We conclude this section by observing that, in view of the homomorphism property of  $\mathbf{S}$ , the equation (1.21) is equivalent to the following one

$$\mathbf{F}(\lambda, \mathbf{u}) := \mathbf{u} - \lambda \mathbf{M}(\lambda) \mathbf{u} - \lambda \mathbf{B}(\mathbf{u}) = 0 \quad \text{in } \mathcal{H}. \quad (2.21)$$

with  $\mathbf{B} := \mathbf{S}^{-1} \mathbf{N}$ .

### 3 Necessary and Sufficient Conditions for Steady Bifurcation.

The main objective of this section is to investigate steady bifurcation of the flow branch determined in Theorem 1.1 around Galilei number  $\lambda_0$ . This leads us to study, in a neighborhood of  $\lambda_0$ , the existence of a nontrivial branch of solutions  $\mathbf{u} = \mathbf{u}(\lambda)$  to equation (2.21). Since  $\mathbf{B}$  is quadratic in  $\mathbf{u}$ , we deduce that  $\mathbf{F}$  is analytic with respect to the  $\mathbf{u}$ -variable and, in particular, its Frechet derivative at  $(\lambda = \lambda_0, \mathbf{u} = \mathbf{0})$  is given by

$$D_{\mathbf{u}} \mathbf{F}(\lambda_0, \mathbf{0}) \mathbf{w} = \mathbf{w} - \lambda_0 \mathbf{M}(\lambda_0) \mathbf{w}. \quad (3.1)$$

As for the regularity in  $\lambda$ , we make the following assumption,

There is a neighborhood  $U_0$  of  $\lambda_0$  such that the map  $\lambda \in U_0 \mapsto \mathbf{v}_0(\lambda)$  is of class  $C^2$  (H)

which implies that  $F$  is of class  $C^2$  in  $U_0 \times \mathcal{X}(\Omega)$ . We next observe that, by Lemma 2.5,  $\mu I - \lambda M(\lambda)$  is Fredholm of index 0 for all fixed  $\lambda > 0$  and all  $\mu \neq 0$ . Thus, we may define a simple eigenvalue of  $M$  as follows [25, Definition 79.14]. The number  $\mu \neq 0$  is a *simple eigenvalue* if

$$\begin{aligned} \dim N[\mu I - \lambda M(\lambda)] &= 1; \\ N[\mu I - \lambda M(\lambda)] \cap R[I - \lambda M(\lambda)] &= \{0\}. \end{aligned} \quad (3.1)$$

It is very well known that the second condition can be reformulated in an equivalent way in terms of an eigenvector of the adjoint operator,  $M^*$ , of  $M$ . Actually, from (3.1)<sub>1</sub> and the Fredholm property we deduce that  $\dim N[\mu I - \lambda M^*(\lambda)] = \text{codim } R[\mu I - \lambda M(\lambda)] = 1$ . Thus, denoting by  $w_1 \in \mathcal{H}$  and  $w_1^* \in \mathcal{H}^{-1}$  non-zero elements of  $N[\mu I - \lambda M(\lambda)]$  and  $N[\mu I - \lambda M^*(\lambda)]$ , respectively, (3.1)<sub>2</sub> is shown to be equivalent (after suitable normalization) to

$$\langle w_1^*, w_1 \rangle = 1. \quad (3.2)$$

The following result is a consequence of (2.21) and [25, Corollary 79.16].

**Lemma 3.1** *Suppose there exists  $\lambda_0 > 0$  such that 1 is a simple eigenvalue of the operator  $\lambda_0 M(\lambda_0)$  and that (H) holds. Then, there is  $\overline{U}_0 \subseteq U_0$  such that the eigenvalue  $\mu = \mu(\lambda)$  of  $\lambda M(\lambda)$ ,  $\lambda \in \overline{U}_0$ , is still simple and of class  $C^2$ . Moreover, we have*

$$\mu'(\lambda_0) = -\langle w_1^*, (M(\lambda_0) + \lambda_0 M'(\lambda_0))w_1 \rangle,$$

where the prime denotes differentiation with respect to  $\lambda$ .

We are now in a position to prove our main bifurcation result.

**Theorem 3.1** *A necessary condition for  $(\lambda_0, \mathbf{0})$  to be a bifurcation point of (2.21) is that  $\dim N[I - \lambda_0 M(\lambda_0)] \geq 1$ , namely, the equation*

$$w - \lambda_0 M(\lambda_0)w = \mathbf{0} \quad (3.3)$$

*has at least one non-trivial solution  $w_1$ . Conversely, suppose that 1 is a simple eigenvalue of  $\lambda_0 M(\lambda_0)$ , namely, (2.21) holds with  $\mu = 1$ . Then, if  $\mu'(\lambda_0) \neq 0$  (transversality condition), in a suitable neighborhood of  $(\lambda_0, \mathbf{0})$  there exists exactly one continuous curve of nontrivial solutions to (2.21),  $(\lambda, u(\lambda))$ , with  $(\lambda_0, u(\lambda_0)) = (\lambda_0, \mathbf{0})$ .*

*Proof.* The necessary condition for bifurcation at  $(\lambda_0, \mathbf{0})$  is that the derivative  $D_u F(\lambda_0, \mathbf{0})$  is singular. By (3.1) and Lemma 2.6 this derivative is Fredholm of index 0, and, therefore, it is singular if and only if (3.3) has a nontrivial solution  $w_1$ . Conversely, since  $D_u F(\lambda_0, \mathbf{0})$  is Fredholm of index 0, if  $\dim N[I - \lambda_0 M(\lambda_0)] = 1$ , a classical bifurcation result [1, Theorem 4.1.12] ensures the stated sufficient property provided

$$D_{\lambda u}^2 F(\lambda_0, \mathbf{0}) w_1 \notin R[D_u F(\lambda_0, \mathbf{0})],$$

or, equivalently,

$$\langle w_1^*, D_{\lambda u}^2 F(\lambda_0, \mathbf{0}) w_1 \rangle \neq 0. \quad (3.4)$$

By a straightforward computation, from (2.21) we show that

$$D_{\lambda u}^2 F(\lambda_0, \mathbf{0}) w_1 = M(\lambda_0)w_1 + \lambda_0 M'(\lambda_0)w_1,$$

so that, if 1 is a simple eigenvalue, by Lemma 3.1, condition (3.4) is equivalent to  $\mu'(\lambda_0) \neq 0$ , which concludes the proof of the theorem.  $\square$

We would like to make several comments regarding Theorem 3.1 that we shall collect in as many remarks.

**Remark 3.1** If the branch  $\mathbf{v}_0(\lambda)$  is constant around  $\lambda_0$ , the hypothesis  $\mu'(\lambda_0) \neq 0$  is equivalent to the request that 1 is a simple eigenvalue and, therefore, the latter can be omitted.  $\triangle$

**Remark 3.2** Taking into account the definition of the operator  $\mathbf{M}$  given in (2.17), we show that (3.3) is equivalent to the following one

$$\frac{1}{\lambda_0}(\mathbb{D}(\mathbf{w}), \mathbb{D}(\boldsymbol{\varphi})) - \xi_0 \langle \partial_1 \mathbf{w}, \boldsymbol{\varphi} \rangle - 2(\mathbf{w} \cdot \mathbb{D}(\boldsymbol{\varphi}), \mathbf{v}_0) - \xi_{\mathbf{w}} \langle \partial_1 \mathbf{v}_0(\lambda_0), \boldsymbol{\varphi} \rangle = 0, \quad \text{for all } \boldsymbol{\varphi} \in \mathcal{H}, \quad (3.5)$$

where  $\mathbf{w} \in \mathcal{X}(\Omega)$ . Thus, by classical regularity results [9, Section VII.1] and Theorem 1.1 it follows that, on the one hand,  $\mathbf{w} \in C^\infty(\Omega)$  and, on the other hand, there is a pressure field  $p \in C^\infty(\Omega)$  such that equation (3.5) is equivalent to the following ones

$$\left. \begin{aligned} \Delta \mathbf{w} + \lambda_0 \xi_0 \partial_1 \mathbf{w} - \lambda_0 [\mathbf{v}_0(\lambda_0) \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_0(\lambda_0) - \xi_{\mathbf{w}} \partial_1 \mathbf{v}_0(\lambda_0)] &= \nabla p \\ \operatorname{div} \mathbf{w} &= 0 \end{aligned} \right\} \quad \text{in } \Omega$$

$$\mathbf{w} = \xi_{\mathbf{w}} \mathbf{e}_1 + \boldsymbol{\omega}_{\mathbf{w}} \times \mathbf{x} \quad \text{at } \partial\Omega, \quad (3.6)$$

$$\int_{\partial\Omega} \mathbb{T}(\mathbf{w}, p) \cdot \mathbf{n} = \mathbf{0}, \quad \int_{\partial\Omega} \mathbf{x} \times \mathbb{T}(\mathbf{w}, p) \cdot \mathbf{n} = \mathbf{0}.$$

We omit the proof of the latter, since it is obtained from (3.5) by an argument entirely analogous to show that (1.17) is equivalent to (1.1). In view of the summability properties of  $\mathbf{v}_0$  stated in Theorem 1.1 and the fact that  $\mathbf{w} \in L^4(\Omega)$ , it can be shown that the quantity in bracket in (3.5)<sub>1</sub> is in  $L^t(\Omega)$ , for all  $t > 1$ . As a consequence, from known results on the Oseen problem [9, Section VII.7] it follows that  $(\mathbf{w}, p)$  belongs to the same functional class as  $(\mathbf{v}_0, p_0)$  given in (1.2).  $\triangle$

**Remark 3.3** The bifurcation result given in Theorem 3.1 is of *local* nature, namely, the existence of the bifurcating branch is ensured only in a neighborhood of  $(\lambda_0, \mathbf{0})$ . As a matter of fact, thanks to [5, Theorem 7.2], the same assumptions as those of Theorem 3.1 lead to the following result of *global* nature (see also [13, Corollary 6.1]).

**Theorem 2.1'** *Let the assumptions of Theorem 3.1 be satisfied and take in (H)  $U_0 \equiv (a, b)$ . Moreover, denote by  $\mathcal{C}$  the connected component of the closure (in  $\mathbb{R}^3 \times \mathcal{X}(\Omega)$ ) of  $F^{-1}(0) \setminus (a, b) \times \{\mathbf{0}\}$  that contains  $(\lambda_0, \mathbf{0})$ . Then, one of the following three conditions hold:*

- (i)  $\mathcal{C}$  contains a point  $(a, \mathbf{u})$  or  $(b, \mathbf{u})$  for some  $\mathbf{u} \in \mathcal{X}(\Omega)$ ;
- (ii)  $\mathcal{C}$  is not compact;
- (iii)  $\mathcal{C}$  contains a point  $(\lambda_*, \mathbf{0})$  with  $\lambda_* \neq \lambda_0$ .

It should be observed that, for the validity of the above result, the  $C^2$  assumption in (H) can be replaced by requiring only  $C^1$  regularity [21, Theorem 6.1]. We also notice that the statement in (ii) could be replaced by “ $\mathcal{C}$  is unbounded,” on condition that  $\mathbf{F}$  is proper on the closed bounded sets of  $[a + \varepsilon, b - \varepsilon] \times \mathcal{X}(\Omega)$ , for small  $\varepsilon > 0$ . However, the truthfulness of this property does not seem obvious and it is yet to be ascertained.

△

As already noticed in the introductory section, lab and numerical tests show that steady bifurcation occurs by breaking the rotational symmetry of the flow while, however, still keeping planar symmetry along the direction of fall. In particular, the detailed numerical investigation carried out in [6, Section 3] evidences that symmetry breaking is induced by transversal rotation of the sphere, namely, along a direction perpendicular to the translational velocity. Conversely, it is clear that a non-zero value of such rotation is incompatible with rotational symmetry. Our next objective is then to provide sufficient conditions for the non-vanishing of the transversal component of the angular velocity. Of course, it is enough to furnish such conditions on the solution to (3.1) or, equivalently, (3.6). To this end, let  $(\mathbf{H}, P)$  solve the following Stokes problem:

$$\left. \begin{aligned} \operatorname{div} \mathbb{T}(\mathbf{H}, P) &= \mathbf{0} \\ \operatorname{div} \mathbf{H} &= 0 \end{aligned} \right\} \quad \text{in } \Omega \quad (3.7)$$

$$\mathbf{H}(y) = \mathbf{e}_3 \times \mathbf{y}, \quad y \in \partial\Omega.$$

The fields  $\mathbf{H}$  and  $P$  are well known, and given by [19, § 334]

$$\mathbf{H} = \mathbf{e}_3 \times \frac{\mathbf{x}}{|\mathbf{x}|^3}, \quad P = \text{const.} \quad (3.8)$$

Clearly,  $\mathbf{H} \in \mathcal{H}$ , and so we can choose  $\boldsymbol{\varphi} = \mathbf{H}$  in (3.5) to obtain

$$\frac{1}{\lambda_0} (\mathbb{D}(\mathbf{w}), \mathbb{D}(\mathbf{H})) - \xi_0 (\partial_1 \mathbf{w}, \mathbf{H}) - 2(\mathbf{w} \cdot \mathbb{D}(\mathbf{H}), \mathbf{v}_0) - \xi_{\mathbf{w}} (\partial_1 \mathbf{v}_0(\lambda_0), \mathbf{H}) = 0. \quad (3.9)$$

Notice that we have changed  $\langle \cdot, \cdot \rangle$  into  $(\cdot, \cdot)$  for the terms involving  $\partial_1$ , because now they become meaningful in view of the summability properties of  $\mathbf{w}$  (see Remark 3.2),  $\mathbf{v}_0$  (see Theorem 1.1) and  $\mathbf{H}$  (see (3.8)). Also, by dot-multiplying both sides of (3.7)<sub>1</sub> by  $\mathbf{w}$ , integrating by parts over  $\Omega$  and taking into account (3.6)<sub>3</sub>, we deduce

$$\boldsymbol{\omega}_{\mathbf{w}} \cdot \int_{\partial\Omega} \mathbf{x} \times \mathbb{T}(\mathbf{H}, P) \cdot \mathbf{n} = (\mathbb{D}(\mathbf{w}), \mathbb{D}(\mathbf{H})), \quad (3.10)$$

where we have used the fact that, as is well known [15, p. 187] and easily checked,

$$\int_{\partial\Omega} \mathbf{e}_1 \cdot \mathbb{T}(\mathbf{H}, P) \cdot \mathbf{n} = 0.$$

Furthermore,

$$\int_{\partial\Omega} \mathbf{x} \times \mathbb{T}(\mathbf{H}, P) \cdot \mathbf{n} = -8\pi \mathbf{e}_3. \quad (3.11)$$

Collecting (3.9)–(3.11), we conclude

$$-\frac{8}{\pi} \lambda_0 \boldsymbol{\omega}_{\mathbf{w}} \cdot \mathbf{e}_3 = \xi_0 (\partial_1 \mathbf{w}, \mathbf{H}) + 2(\mathbf{w} \cdot \mathbb{D}(\mathbf{H}), \mathbf{v}_0) + \xi_{\mathbf{w}} (\partial_1 \mathbf{v}_0(\lambda_0), \mathbf{H}),$$

from which we deduce the following.

**Theorem 3.2** *Let  $(\mathbf{w}, p, \xi_{\mathbf{w}} \mathbf{e}_1, \omega_{\mathbf{w}})$  be a solution to the “criticality” equation (3.6). A sufficient condition for symmetry breaking bifurcation to occur is that*

$$\xi_0(\partial_1 \mathbf{w}, \mathbf{H}) + 2(\mathbf{w} \cdot \mathbb{D}(\mathbf{H}), \mathbf{v}_0) + \xi_{\mathbf{w}}(\partial_1 \mathbf{v}_0(\lambda_0), \mathbf{H}) \neq 0,$$

where  $\mathbf{H}$  is given in (3.8).

## References

- [1] Berger, M.S., *Nonlinearity and Functional Analysis* Lectures on Nonlinear Problems in Mathematical Analysis, Academic Press, 1977
- [2] Chossat, P. and Iooss, G., *The Couette–Taylor problem*. Applied Mathematical Sciences, **102**. Springer-Verlag, New York, 1994
- [3] Dušek, J., Path instabilities of axisymmetric bodies falling or rising under the action of gravity and hydrodynamic forces in a Newtonian fluid. *Particles in flows*, 397–451, Adv. Math. Fluid Mech., Birkhäuser/Springer, Cham, 2017
- [4] Farwig, R., Neustupa, J. On the spectrum of a Stokes-type operator arising from flow around a rotating body. *Manuscripta Mathematica* **122** (2007) 419–437
- [5] Fitzpatrick, P. M., Pejsachowicz, J., and Rabier, P. J. The degree of proper  $C^2$  Fredholm mappings. I. *J. Reine Angew. Math.* **427** (1992), 133
- [6] Giacobello, Ooi, A. and S. Balachandar, Wake structure of a transversely rotating sphere at moderate Galilei numbers, *J. Fluid Mech.*, **621** (2009) 103–130.
- [7] Galdi, G.P. On the motion of a rigid body in a viscous liquid: a mathematical analysis with applications. *Handbook of mathematical fluid dynamics*, Vol. I, 653–791, North-Holland, Amsterdam, 2002
- [8] Galdi, G.P., Further properties of steady-state solutions to the Navier–Stokes problem past a three-dimensional obstacle. *J. Math. Phys.* **48** (2007), no. 6, 065207, 43 pp.
- [9] Galdi, G.P., *An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems*, Second edition. Springer Monographs in Mathematics, Springer, New York (2011)
- [10] Galdi, G.P., Steady-state Navier–Stokes problem past a rotating body: geometric–functional properties and related questions. *Topics in mathematical fluid mechanics*, 109197, Lecture Notes in Math., 2073, Fond. CIME/CIME Found. Subser., Springer, Heidelberg, 2013.
- [11] Galdi, G.P. A timeperiodic bifurcation theorem and its applications to Navier–Stokes flow past an obstacle, *Mathematical Analysis of Viscous Incompressible Flow*, Ed. Tashiaki Hishida, RIMS Kokyuroku, Kyoto University, Japan (2015) 1–27
- [12] Galdi, G.P., On bifurcating time-periodic flow of a Navier–Stokes liquid past a cylinder. *Arch. Ration. Mech. Anal.* **222** (2016), no. 1, 285–315



- [13] Galdi, G.P. and Rabier, P.J., Functional properties of the Navier-Stokes operator and bifurcation of stationary solutions: planar exterior domains. *Topics in nonlinear analysis*, 273–303, Progr. Nonlinear Differential Equations Appl., **35**, Birkhuser, Basel, 1999
- [14] Gohberg, I., Goldberg, S. and Kaashoek, M.A., *Classes of linear operators:I. Operator Theory*, Advances and Applications, Vol.49, Birkhäuser Verlag, Basel, 1990
- [15] Happel, J. and Brenner, H., *Low Galilei number hydrodynamics with special applications to particulate media*. Prentice–Hall, Inc., Englewood Cliffs, N.J. 1965
- [16] Haragus, M. and Iooss, G. *Local bifurcations, center manifolds, and normal forms in infinite-dimensional dynamical systems*. Universitext. Springer–Verlag London, Ltd., London; EDP Sciences, Les Ulis, 2011
- [17] Jenny, M., Bouchet, G., Dušek, J., Nonvertical ascension or fall of a free sphere in a Newtonian fluid. *Phys. Fluids* **15** (2003) 9–12
- [18] Jenny, M., Dušek, J., Bouchet, G., Instabilities and transition of a sphere falling or ascending freely in a Newtonian fluid. *J. Fluid Mech.* **508** (2004) 201–239.
- [19] Lamb, H., *Hydrodynamics*, Cambridge Univ. Press (1932)
- [20] Nakamura, I., Steady wake behind a sphere, *Physics of Fluids* **19** (1976) 5–8
- [21] Pejsachowicz, J. and Rabier, P.J., Degree theory for  $C^1$  Fredholm mappings of index 0. *J. Anal. Math.* **76** (1998), 289–319
- [22] Sattinger, D.H., *Topics in stability and bifurcation theory*. Lecture Notes in Mathematics, Vol. 309. Springer–Verlag, Berlin–New York, 1973
- [23] Stuart, C. A., Bifurcation from the essential spectrum. Topological nonlinear analysis, II (Frascati, 1995), 397–443, Progr. Nonlinear Differential Equations Appl., **27**, Birkhäuser Boston, Boston, MA, 1997
- [24] Taneda, S., Experimental investigation of the wake behind a sphere at low Galilei numbers. *J. Phys. Soc. Japan* **11**(1956) 1104–1108
- [25] Zeidler, E., *Nonlinear Functional Analysis and Applications*, Vol.4, Application to Mathematical Physics, Springer-Verlag, New York, 1988