arXiv:2004.06776v1 [math.DS] 14 Apr 2020

THE CIRCUMBILLIARD: ANY TRIANGLE CAN BE A 3-PERIODIC

DAN REZNIK AND RONALDO GARCIA

Abstract. A Circumconic passes through a triangle's vertices. We define the Circumbilliard, a circumellipse to a generic triangle for which the latter is a 3-periodic. We study its properties and associated loci.

Keywords elliptic billiard, periodic trajectories, triangle center, circumconic, circumellipse, circumhyperbola, conservation, invariance, invariant.

MSC 51M04 and 37D50 and 51N20 and 51N35 and 68T20

1. Introduction

Given a triangle, a *circumconic* passes through its three vertices and satifies two additional constraints, e.g., center or perspector[1]. We study properties and invariants of such conics derived from a 1d family of triangles: 3-periodics in an Elliptic Billiard (EB): these are triangles whose bisectors coincide with normals to the boundary (bounces are elastic), see Figure 1.

Amongst all planar curves, the EB is uniquely integrable [8]. It can be regarded as a special case of Poncelet's Porism [3]. These two propeties imply two classic invariances: N-periodics have constant perimeter and envelop a confocal Caustic. The seminal work is [17] and more recent treatments include [11, 16].

[1]Where reference and polar triangles are perspective [19].




Figure 1. 3-periodics (blue) in the Elliptic Billiard (EB, black): normals to the the boundary at vertices (black arrows) are bisectors. The family is constant-perimeter and envelopes a confocal Caustic (brown). This family conserves the ratio inradius-to-circumradius and has a stationary Mittenpunkt at the EB center. **Video**: [13, PL#01].

We have shown 3-periodics also conserve the Inradius-to-Circumradius[2] ratio which implies an invariant sum of cosines, and that their *Mittenpunkt*[3] is stationary at the EB center [14]. Indeed many such invariants have been effectively generalized for $N > 3$ [1, 2].

We have also studied the loci of 3-periodic Triangle Centers over the family: out of the first 100 listed in [9], 29 sweep out ellipses (a remarkable fact on its own) with the remainder sweeping out higher-order curves [6]. Related is the study of loci described by the Triangle Centers of the Poristic Triangle family [12].

Summary of the paper: given a generic triangle T we define its *Circumbilliard* CB: a Circumellipse to T for which the latter is a 3-periodic. We then analyze the dynamic of geometry of Circumbilliards for triangles derived from the 3-periodic family such as the Excentral, Anticomplementary, Medial, and Orthic, as well as the loci swept by their centers. Additional results include:

- Proposition 5 in Section 3 describes regions of the EB which produce acute, right-triangle, and obtuse 3-periodics.
- Theorem 1 in Section 3: The aspect ratio of Circumbilliards of the Poristic Triangle Family [4] is invariant. This is a family of triangle with fixed Incircle and Circumcircle.

A reference table with all Triangle Centers, Lines, and Symbols appears in Appendix B. Videos of many of the experiments are assembled on Table 2 in Section 4.

2. The Circumbilliard

Let the boundary of the EB satisfy:

$$f(x,y) = \left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1. \tag{1}$$

Where $a > b > 0$ denote the EB semi-axes, and $c = a^2 - b^2$ throughout the paper. Below we use *aspect ratio* as the ratio of an ellipse's semi-axes. When referring to Triangle Centers we adopt Kimberling X_i notation [9], e.g., X_1 for the Incenter, X_2 for the Barycenter, etc., see Table 3 in Appendix B.

The following five-parameter equation is assumed for all circumconics not passing through $(0,0)$.

$$1 + c_1 x + c_2 y + c_3 xy + c_4 x^2 + c_5 y^2 = 0 \tag{2}$$

Proposition 1. *Any triangle $T = (P_1, P_2, P_3)$ is associated with a unique ellipse E_9 for which T is a billiard 3-periodic. The center of E_9 is T's Mittenpunkt.*

Proof. If T is a 3-periodic of E_9, by Poncelet's Porism, T is but an element of a 1d family of 3-periodics, all sharing the same confocal Caustic[4]. This family will share a common Mittenpunkt X_9 located at the center of E_9 [14].

[2] As does the Poristic Family [4].

[3] Where lines drawn from each Excenter thru sides' midpoints meet.

[4] This turns out to be the Mandart Inellipse I_9 of the family [19].

Appendix A shows how to obtain the parameters for (2) such that it passes through P_1, P_2, P_3 and is centered on X_9: this yields a 5×5 linear system. Solving it its is obtained a quadratic equation with positive discriminant, hence the conic is an ellipse. □

E_9 is called the Circumbilliard (CB) of T. Figure 2 shows examples of CBs for two sample triangles.




FIGURE 2. Two random triangles are shown as well as their Circumbilliards (CBs). Notice their axes in general are not horizontal/vertical. An algorithm for computing the CB is given in Appendix A. **Video:** [13, PL#02]

3. Circumbilliards of Derived Triangles

Figure 3 shows CBs for the Excentral, Anticomplementary (ACT), and Medial Triangles, derived from 3-periodics.

3.1. Excentral Triangle.

The locus of the Excenters is shown in Figure 3 (left). It is an ellipse similar to the 90°-rotated locus of X_1 and its axes a_e, b_e are given by [5, 6]:

$$a_e = \frac{b^2+\delta}{a}, \quad b_e = \frac{a^2+\delta}{b}$$

Where $\delta = \sqrt{a^4 - a^2b^2 + b^4}$.

Proposition 2. *The locus of the Excenters the stationary MacBeath Circumellipse E_6' [19] of the Excentral Triangles.*

Proof. The center of E_6' is the Symmedian Point X_6 [19, MacBeath Circumconic]. The Excentral Triangle's X_6 coincides with the Mittenpunkt X_9 of the reference [9]. Since over the 3-periodics the vertices of the Excentral lie on an ellipse and its center is stationary, the result follows. □

Proposition 3. *The Excentral CB is centered on X_{168}, whose trilinears are irrational, and whose locus is non-elliptic.*

Proof. X_{168} is the Mittenpunkt of the Excentral Triangle [9] and its trilinears are irrational[5] on the sidelengths. To determine if its locus is an ellipse we use the algebro-numeric techniques described in [6]. Namely, a least-squares fit of a zero-centered, axis-aligned ellipse to a sample of X_{168} positions of the 3-periodic family produces finite error, therefore it cannot be an ellipse. □

[5]No Triangle Center whose trilinears are irrational on sidelengths has yet been found whose locus under the 3-periodic family is an ellipse [6].




FIGURE 3. Draw in black in each picture is an $a/b = \varphi \simeq 1.618$ EB and a 3-periodic at $t = 7.0°$. **Left**: the CB of the Excentral Triangle (solid green) centered on the latter's Mittenpunkt is X_{168} [9]. Its locus (red) is non-elliptic. Also shown (dashed green) is the elliptic locus of the Excenters (the MacBeath Circumellipse E_6' of the Excentrals [19]), whose center is X_9 [6]; **Top Right**: the CB of the Anticomplementary Triangle (ACT) (blue), axis-aligned with the EB. Its center is the Gergonne Point X_7, whose locus (red) is elliptic and similar to the EB [6]. The locus of the ACT vertices is not elliptic (dashed blue); **Bottom Right**: the CB of the Medial Triangle (teal), also axis-aligned with the EB, is centered on X_{142}, whose locus (red) is also elliptic and similar to the EB, since it is the midpoint of X_9X_7 [9]. The locus of the medial vertices is a dumb-bell shaped curve (dashed teal). **Video:** [13, PL#03]

This had been observed in [6] for several irrational centers such as X_i, i =13–18, as well as many others. Notice a center may be rational but produce a non-elliptic locus, the emblematic case being X_6, whose locus is a convex quartic. Other examples include X_j, j =19, 22–27, etc.

3.2. **Anticomplementary Triangle (ACT).** The ACT is shown in Figure 3 (top right). The locus of its vertices is clearly not an ellipse.

The ACT is perspective with the reference triangle (3-periodic) at X_2 and all of its triangle centers correspond to the anticomplement[6] of corresponding reference ones [19]. The center of the CB of the ACT is therefore X_7, the anticomplement of X_9. We have shown the locus of X_7 to be an ellipse similar to the EB with axes [6]:

$$(a_7, b_7) = k\,(a, b)\,, \quad \text{with: } k = \frac{2\delta - a^2 - b^2}{c^2}$$

Remark 1. *The axes of the ACT CB are parallel to the EB and of fixed length.*

This stems from the fact the ACT is homothetic to the 3-periodic.

[6]Anticomplement: a 1:2 reflection about X_2.

3.3. **Medial Triangle.** The locus of its vertices is the dumbbell-shaped curve, which at larger a/b is self-intersecting, and therefore clearly not an ellipse, Figure 3 (bottom right).

Like the ACT, the Medial is perspective with the reference triangle (3-periodic) at X_2. All of its triangle centers correspond to the complement[7] of corresponding reference ones [19]. The center of the CB of the Medial is therefore X_{142}, the complement of X_9. This point is known to sit midway between X_9 and X_7.

Remark 2. *The locus of X_{142} is an ellipse similar to the EB.*

This stems from the fact X_9 is stationary and the locus of X_7 is an ellipse similar to the EB (above). Therefore its axes will be given by:

$$(a_{142}, b_{142}) = (a_7, b_7)/2$$

Stemming from homothety of 3-periodic and its Medial:

Remark 3. *The axes of the Medial CB are parallel to the EB and of fixed length.*

3.4. **Superposition of ACT and Medial.**

Proposition 4. *The Intouchpoints of the ACT (resp. 3-periodic) are on the EB (resp. on the CB of the Medial)*

Proof. The first part was proved in [15, Thm. 2]. Because the 3-periodic can be regarded as the ACT of the Medial, the result follows. □

This phenomenon is shown in Figure 4. Also shown is the fact that X_i, i =7, 142, 2, 9, 144 are all collinear and their intermediate intervals are related as $3:1:2:6$. In [10] this line is known as $L(X_2, X_7)$ or $\mathcal{L}_{663}$. X_{144} is the perspector of the ACT and its Intouch Triangle (not shown).

3.5. **Orthic Triangle.** Let $\alpha_4 = \sqrt{2\sqrt{2}-1} \simeq 1.352$. In [15, Thm. 1] we show that if $a/b > \alpha_4$, the 3-periodic family will contain obtuse triangles.

Proposition 5. *If $a/b > \alpha_4$, the 3-periodic is a right triangle when one of its vertices is at four symmetric points $P_i^{\perp}$, $i = 1, 2, 3, 4$ given by $(\pm x^{\perp}, \pm y^{\perp})$ with:*

$$x^{\perp} = \frac{a^2\sqrt{a^4+3b^4-4b^2\delta}}{c^3}, \quad y^{\perp} = \frac{b^2\sqrt{-b^4-3a^4+4a^2\delta}}{c^3} \tag{3}$$

Proof. Let the coordinates of the 3-periodic vertices be $P_1 = (x_1, y_1), P_2 = (x_2, y_2), P_3 = (x_3, y_3)$ as derived in [5].

Computing the equation $\langle P_2 - P_1, P_3 - P_1\rangle = 0$, after careful algebraic manipulations, it follows that x_1 satisfies the quartic equation

$$c^8 x_1^4 - 2a^4c^2(a^4+3b^4)x_1^2 + a^8(a^4+2a^2b^2-7b^4) = 0.$$

For $a/b > \sqrt{2\sqrt{2}-1}$ the only positive root in the interval $(0, a)$ is given by

[7] Complement: a 2:1 reflection about X_2.




FIGURE 4. Construction for both ACT and Medial CBs, centered on X_7 and X_{142}, respectively. The Incircle of the ACT (resp. 3-periodic) is shown blue (resp. green). The former touches the ACT at the EB and the latter touches the 3-periodic sides at the Medial CB. Also shown is line $L(2,7) = L_{663}$ which cointains X_i, $i = 7, 142, 2, 9, 144$. Their consecutive distances are proportional to $3:1:2:6$. X_{144} was included since it is the perspector of the ACT and its Intouch Triangle (not shown) [19]. **Video**: [13, PL#04,05]

$$x^{\perp} = \frac{a^2\sqrt{a^4+3b^4-4b^2\delta}}{c^3}. \tag{4}$$

With $y^{\perp}$ obtainable from (1). □

Equivalently, a 3-periodic will be obtuse iff one of its vertices lies on top or bottom halves of the EB between the $P_i^{\perp}$, see Figure 5.

Consider the elliptic arc along the EB between $(\pm x^{\perp}, y^{\perp})$. When a vertex of the 3-periodic lies within (resp. outside) this interval, the 3-periodic is obtuse (resp. acute).

Proposition 6. *When $a/b > \alpha_4$, the locus of the center of the Orthic CB has four pieces: 2 for when the 3-periodic is acute (equal to the X_6 locus), and 2 when it is obtuse (equal to the locus of X_6 of $T'' = P_2P_3X_4$.*

Proof. It is well-known that [9] an acute triangle T has an Orthic whose vertices lie on the sidelines. Furthermore the Orthic's Mittenpunkt coincides with the Symmedian X_6 of T. Also known is the fact that:

Remark 4. *Let triangle $T' = P_1P_2P_3$ be obtuse on P_1. Its Orthic has one vertex on P_2P_3 and two others exterior to T'. Its Orthocenter X_4 is also*




FIGURE 5. Two 3-periodics are shown: one acute (solid blue) and one obtuse (dashed blue) inscribed into an $a/b = 1.618$ EB. Red arcs along the top and bottom halves of the EB indicate that when a 3-periodic vertex is there, the 3-periodic is obtuse. These only exist when $a/b > \alpha_4 \simeq 1.352$.

exterior. Furthermore, the Orthic's Mittenpunkt is the Symmedian Point X_6 of acute triangle $T'' = P_2P_3X_4$.

To see this, notice the Orthic of T'' is also[8] T'. T'' must be acute since its Orthocenter is P_2. □

The CB of the orthic is shown in Figures 6 for four 3-periodic configurations in an EB whose $a/b > \alpha_4$.

Proposition 7. *The coordinates $(\pm x^*, \pm y^*)$ where the locus of the center of the Orthic's CB transitions from one curve to the other are given by:*

$$x^* = \frac{x^\perp}{c^6}\left(a^6 + 2a^2b^4 - b^2\delta(3a^2 + b^2) + b^6\right)$$
$$y^* = -\frac{y^\perp}{c^6}\left(b^6 + 2a^4b^2 - a^2\delta(3b^2 + a^2)\delta + a^6\right)$$

Proof. Let $P_1 = (x_1, y_1)$ be the right-triangle vertex of a 3-periodic, given by $(x^\perp, y^\perp)$ as in (4). Using [5], obtain $P_2 = (p_{2x}/q_2, p_{2y}/q_2)$ and $P_3 = (p_{3x}/q_3, p_{3y}/q_3)$, with:

$$\begin{aligned}
p_{2x} &= b^4c^2x_1^3 - 2a^4b^2x_1^2y_1 + a^4c^2x_1y_1^2 - 2a^6y_1^3\\
p_{2y} &= 2b^6x_1^3 - b^4c^2x_1^2y_1 + 2a^2b^4x_1y_1^2 - a^4c^2y_1^3\\
q_2 &= b^4(a^2+b^2)x_1^2 - 2a^2b^2c^2x_1y_1 + a^4(a^2+b^2)y_1^2\\
p_{3x} &= b^4c^2x_1^3 + 2a^4b^2x_1^2y_1 + a^4c^2x_1y_1^2 + a^6y_1^3\\
p_{3y} &= -2b^6x_1^3 - b^4c^2x_1^2y_1 - 2a^2b^4x_1y_1^2 - a^4c^2y_1^3\\
q_3 &= b^4(a^2+b^2)x_1^2 + 2a^2b^2c^2x_1y_1 + a^4(a^2+b^2)y_1^2
\end{aligned}$$

It can be shown the Symmedian point X_6 of a right-triangle is the midpoint of its right-angle vertex altitude. Computing X_6 using this property leads to the result. □

[8] The anti-orthic pre-images of T' are both the 3-periodic and T''.

Let $\alpha_{eq} = \sqrt{4\sqrt{3}-3} \simeq 1.982$ be the only positive root of $x^4 + 6x^2 - 39$. It can be shown, see Figure 7:

Proposition 8. *At $a/b = \alpha_{eq}$, the locus of the Orthic CB is tangent to EB's top and bottom vertices. If a 3-periodic vertex is there, the Orthic is equilateral.*

Proof. Let T be an equilateral with side s_{eq} and center C. Let h be the distance from any vertex of T to C. It can be easily shown that $h/s_{eq} = \sqrt{3}/3$. Let T' be the Excentral Triangle of T: its sides are $2s_{eq}$. Now consider the upside down equilateral in Figure 7, which is the Orthic of an upright isosceles 3-periodic. h is clearly the 3-periodic's height and $2s_{eq}$ is its base. The height and width of the upright isosceles are obtained from explicit expressions for the vertices [5]:

$$s_{eq} = \frac{\alpha^2}{\alpha^2-1}\sqrt{2\delta-\alpha^2-1}, \quad h = \frac{\alpha^2+\delta+1}{\alpha^2+\delta}$$

where $\alpha = a/b$. Setting $h/s_{eq} = \sqrt{3}/3$ and solving for α yields the required result for α_{eq}. □

3.6. **Summary.** Table 1 summarizes the CBs discussed above, their centers, and their loci.

Triangle	Center	Elliptic Locus
3-Periodic	X_9	n/a
Excentral	X_{168}	No
ACT	X_7	Yes
Medial	X_{142}	Yes
Orthic	X_6^*	No

TABLE 1. CBs mentioned in this Section, their Centers and loci types.

3.7. **Circumbilliard of the Poristic Family.** The Poristic Triangle Family is a set of triangles (blue) with fixed Incircle and Circumcircle [4]. It is a cousing of the 3-periodic family in that by definition its Inradius-to-Circumradius r/R ratio is constant.

Weaver [18] proved the Antiorthic Axis[9] of this family is stationary. Odehnal showed the locus of the Excenters is a circle centered on X_{40} and of radius $2R$ [12]. He also showed that over the family, the locus of the Mittenpunkt X_9 is a circle whose radius is $2d^2(4R+r)$ and center is $X_1 + (X_1 - X_3)(2R - r)/(4R + r)$, where $d = |X_1X_3| = \sqrt{R(R-2r)}$ [12, page 17].

Let $\rho = r/R$ and a_9, b_9 be the semi-axis lengths of the Circumbilliard a poristic triangle. As shown in Figure 8:

Theorem 1. *The ratio a_9/b_9 is invariant over the family and is given by:*

[9]The line passing through the intersections of reference and Excentral sidelines [19].




FIGURE 6. Orthic CB for an EB with $a/b = 1.5 > \alpha_4$, i.e., containing obtuse 3-periodics, which occur when a 3-periodic vertex lies on the top or bottom areas of the EB between the $P^\perp$. **Top left**: 3-periodic is sideways isosceles and acute (vertices outside $P^\perp$, so 3 orthic vertices lie on sidelines. The Orthic CB centers is simply the mittenpunkt of the Orthic, i.e, X_6 of the 3-periodic (blue curve: a convex quartic [6]). **Top right**: The position when a vertex is at a $P^\perp$ and the 3-periodic is a right triangle: its Orthic and CB degenerate to a segment. Here the CB center is at a first (of four) transition points shown in the other insets as Q_i, $i = 1, 2, 3, 4$. **Bottom left**: The 3-periodic is obtuse, the Orthic has two exterior vertices, and the center of the CB switches to the Symmedian of $T'' = P_1P_2X_4$ (red portion of locus). **Bottom right:**. The 3-periodic is an upright isosceles, still obtuse, the center of the Orthic CB reaches its highest point along its locus (red). **Video**: [13, PL#06].

$$\frac{a_9}{b_9} = \sqrt{\frac{\rho^2 + 2(\rho+1)\sqrt{1-2\rho} + 2}{\rho(\rho+4)}}$$

where $\rho = r/R$.

Proof. The following expression for r/R has been derived for the 3-periodic family of an a, b EB [7, Equation 7]:

$$\rho = \frac{r}{R} = \frac{2(\delta - b^2)(a^2 - \delta)}{c^4} \tag{5}$$

Solving the above for a/b yields the result. □

4. Conclusion

Videos mentioned above have been placed on a playlist [13]. Table 2 contains quick-reference links to all videos mentioned, with column "PL#" providing video number within the playlist.




FIGURE 7. At $a/b = \alpha_{eq} \simeq 1.982$, the Orthic is an equilateral triangle when a 3-periodic vertex lies on a top or bottom vertex of the EB. Therefore its CB is a circle.




FIGURE 8. Two configurations (left and right) of the Poristic Triangle Family (blue), whose Incircle (green) and Circumcircle (purple) are fixed. Here $R = 1, r = 0.3625$. Over the family, the Circumbilliard (black) has invariant aspect ratio, in this case $a_9/b_9 \simeq 1.5$. Also shown is the circular locus of X_9 [12, page 17]. **Video**: [13, PL#07].

PL#	Title	Section
01	Mittenpunkt stationary at EB center	1
02	Circumbilliards (CB) of Various Triangles	2
03	CBs of Derived Triangles and Loci of Centers	3
04	CBs of ACT and Medial (separate)	3
05	CBs of ACT and Medial (superposed)	3
06	CB of Orthic and Locus of its Mittenpunkt	3
07	Invariant Aspect Ratio of Circumbilliard of Poristic Family	3

TABLE 2. Videos mentioned in the paper. Column "PL#" indicates the entry within the playlist [13]

ACKNOWLEDGMENTS

We would like to thank Peter Moses and Clark Kimberling, for their prompt help with dozens of questions. We would like to thank Boris Odehnal for his help with some proofs. A warm thanks goes out to Profs. Jair Koiller and Daniel Jaud who provided critical editorial help.

The second author is fellow of CNPq and coordinator of Project PRONEX/ CNPq/ FAPEG 2017 10 26 7000 508.

REFERENCES

[1] Akopyan, A., Schwartz, R., Tabachnikov, S.: Billiards in ellipses revisited (2020). URL https://arxiv.org/abs/2001.02934 2

[2] Bialy, M., Tabachnikov, S.: Dan Reznik's identities and more (2020). URL https://arxiv.org/abs/2001.08469 2

[3] Dragović, V., Radnović, M.: Poncelet Porisms and Beyond: Integrable Billiards, Hyperelliptic Jacobians and Pencils of Quadrics. Frontiers in Mathematics. Springer, Basel (2011). URL https://books.google.com.br/books?id=QcOmDAEACAAJ 1

[4] Gallatly, W.: The modern geometry of the triangle. Francis Hodgson (1914) 2, 8

[5] Garcia, R.: Elliptic billiards and ellipses associated to the 3-periodic orbits. American Mathematical Monthly **126**(06), 491–504 (2019). URL https://doi.org/10.1080/00029890.2019.1593087 3, 5, 7, 8

[6] Garcia, R., Reznik, D., Koiller, J.: Loci of 3-periodics in an elliptic billiard: why so many ellipses? (2020). URL https://arxiv.org/abs/2001.08041 2, 3, 4, 9, 13

[7] Garcia, R., Reznik, D., Koiller, J.: New properties of triangular orbits in elliptic billiards (2020). URL https://arxiv.org/abs/2001.08054 9

[8] Kaloshin, V., Sorrentino, A.: On the integrability of Birkhoff billiards. Phil. Trans. R. Soc. **A**(376) (2018). DOI https://doi.org/10.1098/rsta.2017.0419 1

[9] Kimberling, C.: Encyclopedia of triangle centers (2019). URL https://faculty.evansville.edu/ck6/encyclopedia/ETC.html 2, 3, 4, 6, 12

[10] Kimberling, C.: Central lines of triangle centers (2020). URL https://faculty.evansville.edu/ck6/encyclopedia/CentralLines.html 5, 13

[11] Lynch, P.: Integrable elliptic billiards and ballyards. European Journal of Physics **41**(1), 015,005 (2019). DOI 10.1088/1361-6404/ab4748. URL https://arxiv.org/abs/1907.09295 1

[12] Odehnal, B.: Poristic loci of triangle centers. J. Geom. Graph. **15**(1), 45–67 (2011) 2, 8, 10

[13] Reznik, D.: Playlist for "Circumphenomena of 3-Periodics in Elliptic Billiards" (2020). URL https://bit.ly/2WDPeVk 1, 3, 4, 6, 9, 10, 11

[14] Reznik, D., Garcia, R., Koiller, J.: Can the elliptic billiard still surprise us? Math Intelligencer **42** (2019). DOI 10.1007/s00283-019-09951-2. URL https://rdcu.be/b2cg1 2
[15] Reznik, D., Garcia, R., Koiller, J.: The ballet of triangle centers on the elliptic billiard (2020). URL https://arxiv.org/abs/2002.00001 5
[16] Rozikov, U.A.: An Introduction To Mathematical Billiards. World Scientific Publishing Company (2018) 1
[17] Tabachnikov, S.: Geometry and Billiards, *Student Mathematical Library*, vol. 30. American Mathematical Society, Providence, RI (2005). DOI 10.1090/stml/030. URL http://www.personal.psu.edu/sot2/books/billiardsgeometry.pdf. Mathematics Advanced Study Semesters, University Park, PA 1
[18] Weaver, J.H.: Invariants of a poristic system of triangles. Bull. Amer. Math. Soc. **33**(2), 235–240 (1927). DOI 10.1090/S0002-9904-1927-04367-1. URL https://doi.org/10.1090/S0002-9904-1927-04367-1 8
[19] Weisstein, E.: Mathworld (2019). URL http://mathworld.wolfram.com 1, 2, 3, 4, 5, 6, 8

Appendix A. Computing a Circumconic

Let a Circumconic have center $M = (x_m, y_m)$ Equation (2) is subject to the following 5 constraints[10]: it must be satisfied for vertices P_1, P_2, P_3, and its gradient must vanish at M:

$$f(P_i) = 0, \quad i = 1, 2, 3$$
$$\frac{dg}{dx}(x_m, y_m) = c_1 + c_3 y_m + 2c_4 x_m = 0$$
$$\frac{dg}{dy}(x_m, y_m) = c_2 + c_3 x_m + 2c_5 y_m = 0$$

Written as a linear system:

$$\begin{bmatrix} x_1 & y_1 & x_1 y_1 & x_1^2 & y_1^2 \\ x_2 & y_2 & x_2 y_2 & x_2^2 & y_2^2 \\ x_3 & y_3 & x_3 y_3 & x_3^2 & y_3^2 \\ 1 & 0 & y_m & 2\,x_m & 0 \\ 0 & 1 & x_m & 0 & 2\,y_m \end{bmatrix} \cdot \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \\ c_5 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ -1 \\ 0 \\ 0 \end{bmatrix}$$

Given sidelenghts s_1, s_2, s_3, the coordinates of $X_9 = (x_m, y_m)$ can be obtained by converting its Trilinears $(s_2 + s_3 - s_1 :: ...)$ to Cartesians [9].

Principal axes' directions are given by the eigenvectors of the Hessian matrix H (the jacobian of the gradient), whose entries only depend on c_3, c_4, and c_5:

$$H = J(\nabla g) = \begin{bmatrix} 2\,c_4 & c_3 \\ c_3 & 2\,c_5 \end{bmatrix} \tag{6}$$

The ratio of semiaxes' lengths is given by the square root of the ratio of H's eigenvalues:

$$a/b = \sqrt{\lambda_2/\lambda_1} \tag{7}$$

[10] If M is set to X_9 one obtains the Circumbilliard.

Let $U = (x_u, y_u)$ be an eigenvector of H. The length of the semiaxis along u is given by the distance t which satisfies:

$$g(M + t\,U) = 0$$

This yields a two-parameter quadratic $d_0 + d_2 t^2$, where:

$$\begin{aligned} d_0 &= 1 + c_1 x_m + c_4 x_m^2 + c_2 y_m + c_3 x_m y_m + c_5 y_m^2 \\ d_2 &= c_4 x_u^2 + c_3 x_u y_u + c_5 y_u^2 \end{aligned}$$

The length of the semi-axis associated with U is then $t = \sqrt{-d_0/d_2}$. The other axis can be computed via (7).

The eigenvectors (axes of the conic) of H are given by the zeros of the quadratic form

$$q(x, y) = c_3(y^2 - x^2) + 2(c_2 - c_5)xy$$

Appendix B. Table of Symbols

Tables 3 and 4 lists most Triangle Centers and symbols mentioned in the paper.

Center	Meaning	Note
X_1	Incenter	Locus is Ellipse
X_2	Barycenter	Perspector of Steiner Circum/Inellipses
X_3	Circumcenter	Locus is Ellipse, Perspector of M
X_4	Orthocenter	Exterior to EB when 3-periodic is obtuse
X_5	Center of the 9-Point Circle	
X_6	Symmedian Point	Locus is Quartic [6]
X_6^*	X_9 of Orthic	Detached from X_6 locus for obtuse triangles
X_7	Gergonne Point	Perspector of Incircle
X_8	Nagel Point	Perspector of I_9, X_1 of ACT Incircle
X_9	Mittenpunkt	Center of (Circum)billiard
X_{10}	Spieker Point	Incenter of Medial
X_{11}	Feuerbach Point	on confocal Caustic
X_{40}	Bevan Point	X_3 of Excentral
X_{69}	X_6 of the ACT	Perspector of I_3
X_{100}	Anticomplement of X_{11}	On Circumcircle and EB, J_{exc} center
X_{142}	X_9 of Medial	Midpoint of X_9X_7, lies on $L(2,7)$
X_{144}	Anticomplement of X_7	Perspector of ACT and its Intouch Triangle
X_{168}	X_9 of the Excentral Triangle	Non-elliptic Locus
$L(2,7)$	ACT-Medial Mittenpunkt Axis	Line $\mathcal{L}_{663}$ [10]

Table 3. Kimberling Centers and Central Lines mentioned in the paper.

Dan Reznik, Data Science Consulting,, Rio de Janeiro, RJ, Brazil
E-mail address: dan@dat-sci.com

Ronaldo Garcia, Inst. de MatemÃątica e EstatÃdstica,, Univ. Federal de GoiÃąs,, GoiÃĉnia, GO, Brazil
E-mail address: ragarcia@ufg.br

Symbol	Meaning	Note
a, b	EB semi-axes	$a > b > 0$
P_i, s_i	Vertices and sidelengths of 3-periodic	invariant $\sum s_i$
P_i'	Vertices of the Excentral Triangle	
a_9, b_9	Semi-axes of Poristic Circumbilliard	
r, R, ρ	Inradius, Circumradius, r/R	ρ is invariant
δ	Oft-used constant	$\sqrt{a^4 - a^2b^2 + b^4}$
d	Distance $\lvert X_1X_3\rvert$	$\sqrt{R(R-2r)}$
α	EB aspect ratio	a/b
α_4	a/b threshold for obtuse 3-Periodics	$\sqrt{2\sqrt{2}-1}$
α_{eq}	a/b for equilateral Orthic	$\sqrt{4\sqrt{3}-3}$
$P^\perp$	Obtuse 3-periodic limits on EB	
x^*, y^*	where X_6^* detaches from X_6 locus	Occurs when some P_i is at $P^\perp$

TABLE 4. Symbols mentioned in the paper.