

On Reductions of Hintikka Sets for Higher-Order Logic

Alexander Steen¹, Christoph Benzmüller²

¹ University of Luxembourg, FSTM, alexander.steen@uni.lu

² Freie Universität Berlin, FB Mathematik und Informatik, c.benzmueller@fu-berlin.de

January 31, 2022

Abstract

Steen’s (2018) Hintikka set properties for Church’s type theory based on primitive equality are reduced to the Hintikka set properties of Brown (2007). Using this reduction, a model existence theorem for Steen’s properties is derived.

1 Introduction and Preliminaries

Abstract consistency properties and Hintikka sets play an important role in the study (e.g., of Henkin-completeness) of proof calculi for Church’s type theory [7, 2], aka. classical higher-order logic (HOL). Technically quite different definitions of these terms have been used in the literature, since they depend on the primitive logical connectives assumed in each case. The definitions of Benzmüller, Brown and Kohlhasse [3, 4], for example, are based on negation, disjunction and universal quantification, while Steen [8], in the tradition of Andrews [1], works with primitive equality only. Despite their conceptual relationship, important semantical corollaries that are implied by these syntax related Hintikka properties (such as model existence theorems) can hence not be directly transferred between formalisms.

Brown [6, 5] presents generalised abstract consistency properties in which the primitive logical connectives can vary. In this paper we show that the properties of Steen can be reduced to those of Brown.¹ Theorem 1 that we establish in this paper paves way for the convenient reuse of (e.g., model existence) results from the work of Brown [6, 5] in the context of Steen’s setting.

Paper structure. In the remainder of this introduction we recapitulate some relevant (syntactic) notions on HOL, mainly to clarify our notation; for further details on HOL as relevant for this paper. §2 we present the Hintikka properties as used by Brown, and in §3 we give the related properties as used by Steen. In §3 we then show various lemmas that are implied in Steen’s setting, and it are (some of) those lemmas which prepare the main reduction result of this paper (Theorem 1), which is given in §4. A model existence theorem for Hintikka sets as defined by Steen is then derived in §5.

Equality conventions. Different notions of equality will be used in the following: If a concept is defined (as an abbreviation), the symbol $:=$ is used. Primitive equality, written $=^\tau$, refers to a logical constant symbol from the HOL language such that $s_\tau =^\tau t_\tau$ is a term of the logic (assuming s_τ and t_τ are, where τ is a type annotation), cf. details further below. Leibniz-equality, written \doteq , is a defined term; usually it stands for $\lambda X_\tau. \lambda Y_\tau. \forall P_{o\tau}. (P\ X) \Rightarrow (P\ Y)$, where \Rightarrow is a (primitive) logical connective. Meta equality \equiv denotes set-theoretic identity between objects. Finally, \equiv_\star , for $\star \subseteq \{\beta, \eta\}$ is used for syntactic equality modulo β -, η - and $\beta\eta$ -conversion, respectively (as in the related work, α -conversion is taken as implicit).

Syntax of HOL. The set \mathcal{T} of simple types is freely generated from the base types o and ι by juxtaposition. The types o and ι represent the type of Booleans and individuals, respectively. A type $\nu\tau$ represents the type of a total function from objects of type τ to objects of type ν .

¹In an earlier reduction attempt we tried to reduce Steen’s [8] abstract consistency properties to those of Benzmüller, Brown and Kohlhasse [3]. This attempt had gaps that could not easily be closed (as was pointed out by an unknown reviewer). The mapping that we applied in this initial reduction attempt between the two formalisms replaced primitive equations (in Steen’s Hintikka sets) by Leibniz equations (to obtain Hintikka sets in the style of Benzmüller, Brown and Kohlhasse). It thereby introduced additional universally quantified formulas which in turn triggered the applicability of abstract consistency conditions whose validity could not be ensured by referring to those of Steen. In this work we therefore instead map to the structurally better suited, generalised conditions of Brown [6, 5], which enables us to circumvent the problems induced by our previous reduction attempt to the work of Benzmüller, Brown and Kohlhasse.

Let Σ_τ be a set of constant symbols of type $\tau \in \mathcal{T}$ and let $\Sigma := \bigcup_{\tau \in \mathcal{T}} \Sigma_\tau$ be the union of all typed symbols, called a *signature*. Let further \mathcal{V} denote a set of (typed) variable symbols. From these the terms of HOL are constructed by the following abstract syntax ($\tau, \nu \in \mathcal{T}$):

$$s, t ::= c_\tau \in \Sigma \mid X_\tau \in \mathcal{V} \mid (\lambda X_\tau. s_\nu)_{\nu\tau} \mid (s_{\nu\tau} t_\tau)_\nu$$

The terms are called *constants*, *variables*, *abstractions* and *applications*, respectively. The set of all terms of type τ over a signature Σ is denoted $\Lambda_\tau(\Sigma)$, and $\Lambda_\tau^c(\Sigma)$ is used for closed terms, respectively. The notion of free and bound variables are defined as usual, and a term t is called *closed* if t does not contain any free variables. It is assumed that the set \mathcal{V} contains countably infinitely many variable symbols for each type $\tau \in \mathcal{T}$.

The type of a term is written as subscript but may be dropped by convention if clear from the context (or if not important). Also, parentheses are omitted whenever possible, and application is assumed to be left-associative. Furthermore, the scope of an λ -abstraction's body reaches as far to the right as is consistent with the remaining brackets. Nested applications $s t^1 \dots t^n$ may also be written in vector notation $s \overline{t^n}$.

Variants of HOL often differ regarding the choice of the primitive logical connectives in the signature Σ . In any case, $s \neq t$ is used in the remainder as an abbreviation for $\neg(s = t)$. Also, for simplicity, binary logical connectives may be written in infix notation; e.g., the term $p_o \vee q_o$ formally represents the application $(\vee_{ooo} p_o q_o)$. Furthermore, binder notation is used for universal and existential quantification: The term $\forall X_\tau. s_o$ is used as a short-hand for $\Pi^\tau (\lambda X_\tau. s_o)$, where Π^τ is a constant symbol. Finally, Leibniz-equality, denoted \doteq , is defined as $\doteq := \lambda X_\tau. \lambda Y_\tau. \forall P_{o\tau}. (P X) \Rightarrow (P Y)$. A Σ -*formula* s_o is a term from $s_o \in \Lambda_o(\Sigma)$ of type o and a Σ -*sentence* if it is a closed Σ -formula. The reference to Σ may be omitted if clear from the context.

In the following, variables are denoted by capital letters such as X_τ, Y_τ, Z_τ , and, more specifically, the variable symbols P_o, Q_o and $F_{\nu\tau}, G_{\nu\tau}$ are used for predicate or Boolean variables and variables of functional type, respectively. Analogously, lower case letters s_τ, t_τ, u_τ denote general terms and $f_{\nu\tau}, g_{\nu\tau}$ are used for terms of functional type.

Semantics of HOL. The semantics of HOL, including the notions of Σ -models and Σ -Henkin models, is not discussed here; cf. [3, 8, 6, 5] for details.

2 Hintikka sets as defined by Brown [6]

In the formulation of HOL as employed by Brown [6], the set of primitive logical connectives is not fixed and may be chosen arbitrarily from the set $\{\top_o, \perp_o, \neg_{oo}, \wedge_{ooo}, \vee_{ooo}, \supset_{ooo}, \equiv_{ooo}\} \cup \{\Pi_{o(o\tau)} \mid \tau \in \mathcal{T}\} \cup \{\Sigma_{o(o\tau)} \mid \tau \in \mathcal{T}\} \cup \{=_\tau \mid \tau \in \mathcal{T}\}$. All remaining constant symbols from Σ are called parameters. In this presentation, the notation from Brown is adapted and we apply the conventions from above. For example, we denote general terms with lower case symbols s and t instead of upper case letters as used by Brown, and instead of $wff_\tau(\Sigma)$, which Brown uses to denote the set of internal terms of type τ , we use $\Lambda_\tau(\Sigma)$.

Brown distinguishes between internal terms, the elements of $\Lambda_\tau(\Sigma)$, and external propositions (meta-level propositions) in a set $prop(\Sigma)$, which are defined as follows [6, Def. 2.1.20] (we use the color **red** to visually highlight Brown's meta-level connectives and their associated Hintikka set properties in the remainder of this paper):

- If $s \in \Lambda_o(\Sigma)$, then $s \in prop(\Sigma)$.
- If $\alpha \in \mathcal{T}$ and $s, t \in \Lambda_\alpha(\Sigma)$, then $[s \doteq t] \in prop(\Sigma)$.
- $\top \in prop(\Sigma)$.
- If $s \in prop(\Sigma)$, then $[\neg s] \in prop(\Sigma)$.
- If $s, t \in prop(\Sigma)$, then $[s \vee t] \in prop(\Sigma)$.
- If $s \in prop(\Sigma)$, then $[\forall X_\alpha s] \in prop(\Sigma)$.

Closed propositions $s \in prop(\Sigma)$ are also called sentences; we then write $s \in sent(\Sigma)$.

Brown introduces the following Hintikka set properties for external propositions.

Properties of extensional Hintikka sets \mathcal{H} [6, Def. 5.5.4]:

- $\vec{\nabla}_c : s \notin \mathcal{H} \text{ or } \neg s \notin \mathcal{H}.$
- $\vec{\nabla}_{\beta\eta} : \text{If } s \in \mathcal{H}, \text{ then } s^\downarrow \in \mathcal{H}.$
- $\vec{\nabla}_\perp : \neg \top \notin \mathcal{H}.$

$\vec{\nabla}_{\neg}$: If $\neg\neg s \in \mathcal{H}$, then $s \in \mathcal{H}$.
 $\vec{\nabla}_{\vee}$: If $s \vee t \in \mathcal{H}$, then $s \in \mathcal{H}$ or $t \in \mathcal{H}$.
 $\vec{\nabla}_{\wedge}$: If $\neg(s \vee t) \in \mathcal{H}$, then $\neg s \in \mathcal{H}$ and $\neg t \in \mathcal{H}$.
 $\vec{\nabla}_{\forall}$: If $\forall X_{\tau} s \in \mathcal{H}$, then $[t/x]s \in \mathcal{H}$ for every closed term $t \in \Lambda_{\tau}^c(\Sigma)$.
 $\vec{\nabla}_{\exists}$: If $\neg(\forall X_{\tau} s) \in \mathcal{H}$, then there is a parameter $p_{\tau} \in \Sigma_{\tau}$ such that $\neg([p/X]s) \in \mathcal{H}$.
 $\vec{\nabla}^{\sharp}$: If $s \in \mathcal{H}$, then $s^{\sharp} \in \mathcal{H}$. Also, if $\neg s \in \mathcal{H}$, then $\neg s^{\sharp} \in \mathcal{H}$.
 $\vec{\nabla}_m$: If $\neg(h \overline{s^n}) \in \mathcal{H}$ and $(h \overline{t^n}) \in \mathcal{H}$, then there is an i with $1 \leq i \leq n$ such that $\neg(s^i \doteq t^i) \in \mathcal{H}$.
 $\vec{\nabla}_{dec}$: If p is a parameter and $\neg((p \overline{s^n}) \doteq^{\iota} (p \overline{t^n})) \in \mathcal{H}$, then there is an i , $1 \leq i \leq n$, s.t. $\neg(s^i \doteq t^i) \in \mathcal{H}$.
 $\vec{\nabla}_b$: If $\neg(s \doteq^o t) \in \mathcal{H}$, then $\{s, \neg t\} \subseteq \mathcal{H}$ or $\{\neg s, t\} \subseteq \mathcal{H}$.
 $\vec{\nabla}_f$: If $\neg(f \doteq^{\nu\tau} g) \in \mathcal{H}$, then there is a parameter $p_{\tau} \in \Sigma_{\tau}$ such that $\neg(f p \doteq^{\nu} g p) \in \mathcal{H}$.
 $\vec{\nabla}_{\doteq^o}$: If $s \doteq^o t \in \mathcal{H}$, then $\{s, t\} \subseteq \mathcal{H}$ or $\{\neg s, \neg t\} \subseteq \mathcal{H}$.
 $\vec{\nabla}_{\doteq^{\nu\tau}}$: If $f \doteq^{\nu\tau} g \in \mathcal{H}$, then $(f u \doteq^{\nu} g u) \in \mathcal{H}$ for every closed term $u \in \Lambda_{\tau}^c(\Sigma)$.
 $\vec{\nabla}^r$: $\neg(s \doteq^i s) \notin \mathcal{H}$.
 $\vec{\nabla}^u$: Suppose $(s \doteq^i t) \in \mathcal{H}$ and $\neg(u \doteq^i v) \in \mathcal{H}$. Then $\neg(s \doteq^i u) \in \mathcal{H}$ or $\neg(t \doteq^i v) \in \mathcal{H}$. Also, $\neg(s \doteq^i v) \in \mathcal{H}$ or $\neg(t \doteq^i u) \in \mathcal{H}$.

The collection of all sets of sentences satisfying all these properties is called $\mathfrak{Hint}_{\beta\mathfrak{fb}}(\Sigma)$.

3 Hintikka sets as defined by Steen [8]

In the formulation of HOL as employed by Steen [8], the equality predicate, denoted \doteq^{τ} , for each type τ , is assumed to be the only logical connective present in the signature Σ , i.e., $\{=\tau \mid \tau \in \mathcal{T}\} \subseteq \Sigma$. All (potentially) remaining constant symbols from Σ are called parameters. Such signatures are also referred to as Σ^{τ} . A formulation of HOL based on equality as sole logical connective originates from Andrew's system \mathcal{Q}_0 , cf. [1] and the references therein. The usual logical connectives are defined as follows (technically our formulation is a modification of the one used by Andrews [1], since the order of terms in defining equations is swapped in many cases):

$$\begin{aligned}
\top_o &:= \doteq_{ooo}^o =_{o(ooo)(ooo)}^{ooo} \doteq_{ooo}^o \\
\perp_o &:= (\lambda P_o. P) \doteq^{oo} (\lambda P_o. \top) \\
\neg_{oo} &:= \lambda P_o. P \doteq^o \perp \\
\wedge_{ooo} &:= \lambda P_o. \lambda Q_o. (\lambda F_{ooo}. F \top \top) \doteq^{o(ooo)} (\lambda F_{ooo}. F P Q) \\
\vee_{ooo} &:= \lambda P_o. \lambda Q_o. \neg(\neg P \wedge \neg Q) \\
\Rightarrow_{ooo} &:= \lambda P_o. \lambda Q_o. \neg P \vee Q \\
\Leftrightarrow_{ooo} &:= \lambda P_o. \lambda Q_o. P \doteq^o Q \\
\Pi_{o(o\tau)}^{\tau} &:= \lambda P_{o\tau}. P \doteq^{o\tau} \lambda X_{\tau}. \top
\end{aligned}$$

The (primitive and defined) connectives of this formulation of HOL are written in blue in the following, as are the properties below.

Properties for acceptable Hintikka sets [8, Def. 3.15]

$\vec{\nabla}_c$: $s \notin \mathcal{H}$ or $\neg s \notin \mathcal{H}$.
 $\vec{\nabla}_{\beta\eta}$: If $s \equiv_{\beta\eta} t$ and $s \in \mathcal{H}$, then $t \in \mathcal{H}$.
 $\vec{\nabla}^r$: $(s \neq s) \notin \mathcal{H}$.
 $\vec{\nabla}^s$: If $u[s]_p \in \mathcal{H}$ and $s \doteq t \in \mathcal{H}$ then $u[t]_p \in \mathcal{H}$.
 $\vec{\nabla}_b^+$: If $s \doteq t \in \mathcal{H}$, then $\{s, t\} \subseteq \mathcal{H}$ or $\{\neg s, \neg t\} \subseteq \mathcal{H}$.
 $\vec{\nabla}_b^-$: If $s \neq t \in \mathcal{H}$, then $\{s, \neg t\} \subseteq \mathcal{H}$ or $\{\neg s, t\} \subseteq \mathcal{H}$.
 $\vec{\nabla}_f^+$: If $f_{\nu\tau} \doteq_{\nu\tau} g_{\nu\tau} \in \mathcal{H}$, then $f s \doteq g s \in \mathcal{H}$ for any closed term $s \in \Lambda_{\tau}^c(\Sigma)$.
 $\vec{\nabla}_f^-$: If $f_{\nu\tau} \neq_{\nu\tau} g_{\nu\tau} \in \mathcal{H}$, then $f w \neq g w \in \mathcal{H}$ for some parameter $w \in \Sigma_{\tau}$.
 $\vec{\nabla}_m$: If s, t are atomic and $s, \neg t \in \mathcal{H}$, then $s \neq t \in \mathcal{H}$.
 $\vec{\nabla}_d$: If $h \overline{s^n} \neq h \overline{t^n} \in \mathcal{H}$, then there is an i with $1 \leq i \leq n$ such that $s^i \neq t^i \in \mathcal{H}$.

The collection of all sets satisfying these properties is called \mathfrak{H} . Every element $\mathcal{H} \in \mathfrak{H}$ is called acceptable.

Definition 1. A set \mathcal{H} of formulas is called saturated iff $s \in \mathcal{H}$ or $\neg s \in \mathcal{H}$ for every closed formula s .

Derived properties

Lemma 1 (Basic properties). *Let $\mathcal{H} \in \mathfrak{H}$. Then it holds that*

- (a) $\perp \notin \mathcal{H}$
- (b) $\neg\top \notin \mathcal{H}$
- (c) $\top=\perp \notin \mathcal{H}$
- (d) If $s_o=\top \in \mathcal{H}$ or $\top=s_o \in \mathcal{H}$ ($s_o \neq \perp \in \mathcal{H}$ or $\perp \neq s_o \in \mathcal{H}$), then $\{s_o, \top\} \subseteq \mathcal{H}$ ($\{s_o, \neg\perp\} \subseteq \mathcal{H}$)
- (e) If $s_o=\perp \in \mathcal{H}$ or $\perp=s_o \in \mathcal{H}$ ($s_o \neq \top \in \mathcal{H}$ or $\top \neq s_o \in \mathcal{H}$), then $\{\neg s_o, \neg\perp\} \subseteq \mathcal{H}$ ($\{\neg s_o, \top\} \subseteq \mathcal{H}$)
- (f) If $\neg\perp \in \mathcal{H}$, then $\top \in \mathcal{H}$
- (g) If $\top \in \mathcal{H}$, then $\neg\perp \in \mathcal{H}$
- (h) If $s=t \in \mathcal{H}$ and $t=u \in \mathcal{H}$, then $s=u \in \mathcal{H}$.

Proof. Let $\mathcal{H} \in \mathfrak{H}$ be an acceptable Hintikka set.

- (a) Assume $\perp \in \mathcal{H}$. By definition of \perp it holds $(\lambda P. P) = (\lambda P. \top) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$, it follows that $w=\top \in \mathcal{H}$ for any closed term w . Taking $w \equiv \neg\top$ we obtain $\neg\top=\top \in \mathcal{H}$. But then, $\vec{\nabla}_b^+$ gives us a contradiction to $\vec{\nabla}_c$. Hence $\perp \notin \mathcal{H}$.
- (b) Assume $\neg\top \in \mathcal{H}$. By definition of \perp it holds $(=\neq) \in \mathcal{H}$, which contradicts $\vec{\nabla}_=^r$. Hence, $\neg\top \notin \mathcal{H}$.
- (c) Assume $\top=\perp \in \mathcal{H}$. Applying $\vec{\nabla}_b^+$ gives us that either $\{\top, \perp\} \subseteq \mathcal{H}$ or $\{\neg\top, \neg\perp\} \subseteq \mathcal{H}$. Either case is impossible by either (a) or (b) of this lemma. Hence, $\top=\perp \notin \mathcal{H}$.
- (d) Let $s_o=\top \in \mathcal{H}$ or $\top=s_o \in \mathcal{H}$. In both cases it follows by $\vec{\nabla}_b^+$ that either $\{s, \top\} \subseteq \mathcal{H}$ or $\{\neg s, \neg\top\} \subseteq \mathcal{H}$. Since the latter case contradicts (b) from above, it follows that $\{s, \top\} \subseteq \mathcal{H}$. The negative cases are analogous using $\vec{\nabla}_b^-$.
- (e) Let $s_o=\perp \in \mathcal{H}$ or $\perp=s_o \in \mathcal{H}$. In both cases it follows by $\vec{\nabla}_b^+$ that either $\{s, \perp\} \subseteq \mathcal{H}$ or $\{\neg s, \neg\perp\} \subseteq \mathcal{H}$. Since the former case contradicts (a) from above, it follows that $\{\neg s, \neg\perp\} \subseteq \mathcal{H}$. The negative case is analogous using $\vec{\nabla}_b^-$.
- (f) Let $\neg\perp \in \mathcal{H}$. Then, by definition of \perp , $((\lambda P. P) \neq (\lambda P. \top)) \in \mathcal{H}$. By $\vec{\nabla}_f^-$ and $\vec{\nabla}_{\beta\eta}$ it holds that $p \neq \top \in \mathcal{H}$ for some parameter p . By $\vec{\nabla}_b^-$ it follows that either $\{p, \neg\top\} \subseteq \mathcal{H}$ or $\{\neg p, \top\} \subseteq \mathcal{H}$. Since the former case is ruled out by (b) from above, the latter case yields the desired result.
- (g) Let $\top \in \mathcal{H}$, that is, $=^o =^{ooo} =^o \in \mathcal{H}$. By $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$ it follows that $(s_o=t_o) = (s_o=t_o) \in \mathcal{H}$ for every two closed formulas s, t . For $s \equiv t \equiv \neg\perp$ it follows that $(\neg\perp=\neg\perp) = (\neg\perp=\neg\perp) \in \mathcal{H}$, and hence, by $\vec{\nabla}_b^+$, either $\neg\perp=\neg\perp \in \mathcal{H}$ or $\neg\perp \neq \neg\perp \in \mathcal{H}$. Since the latter case is ruled out by $\vec{\nabla}_=^r$, it follows that $\neg\perp=\neg\perp \in \mathcal{H}$. Again, by $\vec{\nabla}_b^+$, it follows that either $\neg\perp \in \mathcal{H}$ or $\neg(\neg\perp) \in \mathcal{H}$. The latter case is impossible by $\vec{\nabla}_=^r$ since $\neg(\neg\perp) \equiv (\perp \neq \perp)$ and hence $\neg\perp \in \mathcal{H}$.
- (h) Let $s=t \in \mathcal{H}$ and $t=u \in \mathcal{H}$. By $\vec{\nabla}_=^s$ it follows directly that $s=u \in \mathcal{H}$.

□

Lemma 2 (Properties of usual connectives). *Let $\mathcal{H} \in \mathfrak{H}$. Then it holds that*

- (a) If $\neg\neg s_o \in \mathcal{H}$, then $s \in \mathcal{H}$
- (b) If $(s_o \vee t_o) \in \mathcal{H}$, then $s \in \mathcal{H}$ or $t \in \mathcal{H}$.
- (c) If $(s_o \wedge t_o) \in \mathcal{H}$, then $s \in \mathcal{H}$ and $t \in \mathcal{H}$.
- (d) If $\Pi^\alpha F \in \mathcal{H}$, then $F s \in \mathcal{H}$ for every closed term s .
- (e) If $\neg\Pi^\alpha F \in \mathcal{H}$, then $\neg(F w) \in \mathcal{H}$ for some parameter $w \in \Sigma$.

Proof. Let $\mathcal{H} \in \mathfrak{H}$ be an acceptable Hintikka set.

- (a) Let $\neg\neg s_o \in \mathcal{H}$. By definition of \neg and $\vec{\nabla}_{\beta\eta}$ it holds $(s \neq \perp) \in \mathcal{H}$. Hence, by $\vec{\nabla}_6^-$, either $\{s, \neg\perp\} \subseteq \mathcal{H}$, or $\{\neg s, \perp\} \subseteq \mathcal{H}$. As the latter case is impossible by Lemma 1(a), it follows that $\{s, \neg\perp\} \subseteq \mathcal{H}$ and, in particular, that $s \in \mathcal{H}$.
- (b) Let $(s_o \vee t_o) \in \mathcal{H}$. By definition of \vee , \neg and $\vec{\nabla}_{\beta\eta}$ it holds $((\lambda P. P \top \top) \neq \lambda P. P (\neg s) (\neg t)) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^-$ and $\vec{\nabla}_{\beta\eta}$, it follows that $(p \top \top) \neq (p (\neg s) (\neg t))$ for some parameter $p \in \Sigma$. By $\vec{\nabla}_d$ either (i) $\top \neq \neg s \in \mathcal{H}$ or (ii) $\top \neq \neg t \in \mathcal{H}$. Hence, by $\vec{\nabla}_6^-$, applied to both cases, it holds that either (i) $\neg\neg s \in \mathcal{H}$, or (ii) $\neg\neg t \in \mathcal{H}$ (because $\neg\top \notin \mathcal{H}$ by Lemma 1(b)). It follows that $s \in \mathcal{H}$ or $t \in \mathcal{H}$ by (a) of this lemma.
- (c) Let $(s_o \wedge t_o) \in \mathcal{H}$. By definition of \wedge and $\vec{\nabla}_{\beta\eta}$ it holds $(\lambda P. P \top \top) = (\lambda P. P s t) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$, it follows that $(w \top \top) = (w s t) \in \mathcal{H}$ for every closed term w . By $\vec{\nabla}_{\beta\eta}$, using $w \equiv \lambda x. \lambda y. x$ and $w \equiv \lambda x. \lambda y. y$, it holds $\top = s \in \mathcal{H}$ and $\top = t \in \mathcal{H}$, respectively. Application of Lemma 1(d) yields the desired result.
- (d) Let $\Pi^\alpha s \in \mathcal{H}$. By definition of Π^α and $\vec{\nabla}_{\beta\eta}$ it holds $(s = \lambda x. \top) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$, it follows that $s t = \top \in \mathcal{H}$ for every closed term t . Application of Lemma 1(d) yields the desired result.
- (e) Let $\neg\Pi^\alpha s \in \mathcal{H}$. By definition of Π^α and $\vec{\nabla}_{\beta\eta}$ it holds that $(s \neq (\lambda x. \top)) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^-$ and $\vec{\nabla}_{\beta\eta}$, it follows that $(s p) \neq \top \in \mathcal{H}$ for some parameter p . Application of Lemma 1(e) yields the desired result.

□

Lemma 3 (Properties of Leibniz equality). *Let $\mathcal{H} \in \mathfrak{H}$. Then it holds that*

- (a) *If $s \doteq t \in \mathcal{H}$, then $s = t \in \mathcal{H}$.*
- (b) *If $\neg(s \doteq t) \in \mathcal{H}$, then $s \neq t \in \mathcal{H}$.*
- (c) *$\neg(s \doteq s) \notin \mathcal{H}$.*
- (d) *If $u[s]_p \in \mathcal{H}$ and $s \doteq t \in \mathcal{H}$, then $u[t]_p \in \mathcal{H}$.*
- (e) *If $s \doteq t \in \mathcal{H}$, then $t \doteq s \in \mathcal{H}$.*
- (f) *If $s \doteq t \in \mathcal{H}$ and $t \doteq u \in \mathcal{H}$, then $s \doteq u \in \mathcal{H}$.*

Proof. Let $\mathcal{H} \in \mathfrak{H}$ be an acceptable Hintikka set.

- (a) Let $(s \doteq t) \in \mathcal{H}$. By definition of \doteq and $\vec{\nabla}_{\beta\eta}$ we have $(\lambda P. (P s) \Rightarrow (P t)) = (\lambda P. \top) \in \mathcal{H}$, and hence, by $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$, it holds that $((w s) \Rightarrow (w t)) = \top \in \mathcal{H}$ for every closed term w . Then, $(w s) \Rightarrow (w t) \in \mathcal{H}$ by Lemma 1(d). By definition of \Rightarrow and $\vec{\nabla}_{\beta\eta}$ it holds that $\neg(w s) \vee (w t) \in \mathcal{H}$ and hence, by Lemma 2(b), that $\neg(w s) \in \mathcal{H}$ or $(w t) \in \mathcal{H}$. For $w \equiv (\lambda X. s = X)$ it follows by $\vec{\nabla}_{\beta\eta}$ that $\neg(s = s) \in \mathcal{H}$ or $(s = t) \in \mathcal{H}$. Since the former case contradicts $\vec{\nabla}_=^r$, it follows that $(s = t) \in \mathcal{H}$.
- (b) Let $\neg(s \doteq t) \in \mathcal{H}$. By definition of \doteq and $\vec{\nabla}_{\beta\eta}$ we have $(\lambda P. (P s) \Rightarrow (P t)) \neq (\lambda P. \top) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^-$ and $\vec{\nabla}_{\beta\eta}$, it holds that $((p s) \Rightarrow (p t)) \neq \top \in \mathcal{H}$ for some parameter p . By Lemma 1(e) it follows that $\neg((p s) \Rightarrow (p t)) \in \mathcal{H}$. Then, by Lemma 2(a) and 2(c), it follows that $\neg\neg(p s) \in \mathcal{H}$ and $\neg(p t) \in \mathcal{H}$. Moreover, $(p s) \in \mathcal{H}$ by Lemma 2(a). By $\vec{\nabla}_m$ it then follows that $(p s) \neq (p t) \in \mathcal{H}$, and finally, by $\vec{\nabla}_d$, that $s \neq t \in \mathcal{H}$.
- (c) Assume $\neg(s \doteq s) \in \mathcal{H}$. By (b) above it follows that $s \neq s \in \mathcal{H}$ which contradicts $\vec{\nabla}_=^r$. Hence, $\neg(s \doteq s) \notin \mathcal{H}$.
- (d) Let $u[s]_p \in \mathcal{H}$ and $s \doteq t \in \mathcal{H}$. By (a) above it holds that $s = t \in \mathcal{H}$ and thus by $\vec{\nabla}_=^s$ it follows that $u[t]_p \in \mathcal{H}$.
- (e) Let $(s \doteq t) \in \mathcal{H}$. By definition of \doteq and $\vec{\nabla}_{\beta\eta}$ we have $(\lambda P. (P s) \Rightarrow (P t)) = (\lambda P. \top) \in \mathcal{H}$. Hence, by $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$, it holds that $((w s) \Rightarrow (w t)) = \top \in \mathcal{H}$ for every closed term w . Then, $(w s) \Rightarrow (w t) \in \mathcal{H}$ by Lemma 1(d). By definition of \Rightarrow and $\vec{\nabla}_{\beta\eta}$ it holds that $\neg(w s) \vee (w t) \in \mathcal{H}$ and hence by Lemma 2(b) that $\neg(w s) \in \mathcal{H}$ or $(w t) \in \mathcal{H}$. For $w \equiv (\lambda X. t \doteq s)$, it follows by $\vec{\nabla}_{\beta\eta}$ that $\neg(t \doteq s) \in \mathcal{H}$ or $(t \doteq s) \in \mathcal{H}$. Assume $\neg(t \doteq s) \in \mathcal{H}$. Since by Lemma 3(a) it holds that $s = t \in \mathcal{H}$, it follows by $\vec{\nabla}_=^s$ that $\neg(t \doteq t) \in \mathcal{H}$, which contradicts (c) above. Hence, $(t \doteq s) \in \mathcal{H}$.
- (f) Let $(s \doteq t) \in \mathcal{H}$ and $(t \doteq u) \in \mathcal{H}$. By (d) above it follows that $(s \doteq u) \in \mathcal{H}$.

□

Lemma 4 (Sufficient conditions for saturatedness). *Let $\mathcal{H} \in \mathfrak{H}$. Then it holds that*

- (a) *If $\top \in \mathcal{H}$, then \mathcal{H} is saturated.*
- (b) *If $\neg s \in \mathcal{H}$ for some closed term s , then \mathcal{H} is saturated.*
- (c) *If $s \vee t \in \mathcal{H}$ for some closed terms s, t , then \mathcal{H} is saturated.*
- (d) *If $s \wedge t \in \mathcal{H}$ for some closed terms s, t , then \mathcal{H} is saturated.*
- (e) *If $\Pi^\tau P \in \mathcal{H}$ for some closed term $P_{\tau \rightarrow o}$, then \mathcal{H} is saturated.*
- (f) *If $s \doteq t \in \mathcal{H}$ for some closed terms s, t , then \mathcal{H} is saturated.*

Proof. Let $\mathcal{H} \in \mathfrak{H}$ be an acceptable Hintikka set.

- (a) Let $\top \in \mathcal{H}$, that is, $=^o =^{ooo} =^o \in \mathcal{H}$. By $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$ it follows that $(s_o = t_o) = (s_o = t_o) \in \mathcal{H}$ for every two closed formulas s, t . For $s \equiv t \equiv c$ for some closed term c , it follows that $(c = c) = (c = c) \in \mathcal{H}$ and thus, by $\vec{\nabla}_b^+$ and $\vec{\nabla}_=^r$, it holds that $c = c \in \mathcal{H}$. By $\vec{\nabla}_b^+$ it follows that $c \in \mathcal{H}$ or $\neg c \in \mathcal{H}$. Hence, \mathcal{H} is saturated.
- (b) If $\neg s \in \mathcal{H}$ for some closed term s , then $s = \perp \in \mathcal{H}$. By $\vec{\nabla}_b^+$, it follows that either $\{s, \perp\} \subseteq \mathcal{H}$ or $\{\neg s, \neg \perp\} \subseteq \mathcal{H}$. Since the former case is ruled out by Lemma 1(a), it follows that $\neg \perp \in \mathcal{H}$. By Lemma 1(f) it follows that $\top \in \mathcal{H}$ and by (a) above it follows that \mathcal{H} is saturated.
- (c) If $s \vee t \in \mathcal{H}$ for some closed terms s, t , then by definition of \vee and $\vec{\nabla}_{\beta\eta}$ we have $\neg(\neg s \wedge \neg t) \in \mathcal{H}$. An application of (b) yields the desired result.
- (d) If $s \wedge t \in \mathcal{H}$ for some closed terms s, t , then by definition of \wedge and $\vec{\nabla}_{\beta\eta}$ it holds $(\lambda g. g \ s \ t) = (\lambda g. g \ \top \ \top) \in \mathcal{H}$. By $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$ it follows that $s = \top \in \mathcal{H}$ (take $\lambda x. \lambda y. x$). By $\vec{\nabla}_b^+$, it follows that either $\{s, \top\} \subseteq \mathcal{H}$ or $\{\neg s, \neg \top\} \subseteq \mathcal{H}$. Since the latter case is ruled out by Lemma 1(b), it follows that $\top \in \mathcal{H}$. An application of (a) above yields the desired result.
- (e) If $\Pi^\tau s \in \mathcal{H}$ for some closed terms s , then by definition of Π^τ and $\vec{\nabla}_{\beta\eta}$ it holds that $s = (\lambda x. \top) \in \mathcal{H}$. By $\vec{\nabla}_f^+$ and $\vec{\nabla}_{\beta\eta}$ it follows that $(s \ w) = \top \in \mathcal{H}$ for every closed term w . By $\vec{\nabla}_b^+$, it follows that either $\{(s \ w), \top\} \subseteq \mathcal{H}$ or $\{\neg(s \ w), \neg \top\} \subseteq \mathcal{H}$. Since the latter case is ruled out by Lemma 1(b), it follows that $\top \in \mathcal{H}$. An application of (a) above yields the desired result.
- (f) Let $s \doteq t \in \mathcal{H}$. By definition of Π^τ and $\vec{\nabla}_{\beta\eta}$ it holds that $\Pi(\lambda P. (P \ s) \Rightarrow (P \ t)) \in \mathcal{H}$. An application of (e) yields the desired result.

□

Corollary 1. *Let $\mathcal{H} \in \mathfrak{H}$ and let $s \neq t \in \mathcal{H}$ or $\neg(s \doteq t) \in \mathcal{H}$ for some closed terms s, t . Then, \mathcal{H} is saturated.*

Proof. As $(s \neq t) \equiv \neg(s = t)$, both cases are a special instance of Lemma 4(b). □

Lemma 5 (Saturated sets properties). *Let $\mathcal{H} \in \mathfrak{H}$ and let \mathcal{H} be saturated. Then it holds that*

- (a) *If $s = t \in \mathcal{H}$, then $s \doteq t \in \mathcal{H}$.*
- (b) *If $s = t \in \mathcal{H}$ then $t = s \in \mathcal{H}$.*
- (c) *$s = s \in \mathcal{H}$ for every closed term s .*
- (d) *$s \doteq s \in \mathcal{H}$ for every closed term s .*

Proof. Let $\mathcal{H} \in \mathfrak{H}$ and let \mathcal{H} be saturated.

- (a) Let $s = t \in \mathcal{H}$ and assume $s \doteq t \notin \mathcal{H}$. Since \mathcal{H} is saturated we have $\neg(s \doteq t) \in \mathcal{H}$. Then, by Lemma 3(b), it follows that $s \neq t \in \mathcal{H}$, and thus $\{s = t, s \neq t\} \subseteq \mathcal{H}$, which contradicts $\vec{\nabla}_c$. Hence, $s \doteq t \in \mathcal{H}$.
- (b) Let $s = t \in \mathcal{H}$ and assume $t = s \notin \mathcal{H}$. Since \mathcal{H} is saturated we have $t \neq s \in \mathcal{H}$. Then, by $\vec{\nabla}_=^s$, it follows that $t \neq t \in \mathcal{H}$ which contradicts $\vec{\nabla}_=^r$. Hence, $t = s \in \mathcal{H}$.
- (c) Let s be a closed term of some type and assume that $s = s \notin \mathcal{H}$. Since \mathcal{H} is saturated we have that $s \neq s \in \mathcal{H}$. Since this contradicts $\vec{\nabla}_=^r$ it follows that $s = s \in \mathcal{H}$.
- (d) Let s be a closed term of some type and assume that $s \doteq s \notin \mathcal{H}$. Since \mathcal{H} is saturated we have that $\neg(s \neq s) \in \mathcal{H}$. Since this is impossible by Lemma 3(c) it follows that $s \doteq s \in \mathcal{H}$.

□

Lemma 6 (Properties of negated equalities). *Let $\mathcal{H} \in \mathfrak{H}$. It holds that*

- (a) *If $s \neq t \in \mathcal{H}$, then $t \neq s \in \mathcal{H}$.*
- (b) *If $\neg(s \doteq t) \in \mathcal{H}$, then $\neg(t \doteq s) \in \mathcal{H}$*
- (c) *If $s \neq t \in \mathcal{H}$, then $\neg(s \doteq t) \in \mathcal{H}$.*

Proof. Let $\mathcal{H} \in \mathfrak{H}$.

- (a) Let $s \neq t \in \mathcal{H}$ and assume that $t \neq s \notin \mathcal{H}$. By Corollary 1 it follows that \mathcal{H} is saturated and hence we have that $t = s \in \mathcal{H}$. By saturation and Lemma 5(b) it follows that $s = t \in \mathcal{H}$, and thus $\{s \neq t, s = t\} \subseteq \mathcal{H}$, which contradicts $\vec{\nabla}_c$. Hence, $t \neq s \in \mathcal{H}$.
- (b) Let $\neg(s \doteq t) \in \mathcal{H}$ and assume $\neg(t \doteq s) \notin \mathcal{H}$. By Corollary 1 it follows that \mathcal{H} is saturated and hence we have that $t \doteq s \in \mathcal{H}$. Then, by Lemma 3(d), it follows that $s \doteq t \in \mathcal{H}$, and thus $\{\neg(s \doteq t), s \doteq t\} \subseteq \mathcal{H}$, which contradicts $\vec{\nabla}_c$. Hence, $\neg(t \doteq s) \in \mathcal{H}$.
- (c) Let $s \neq t \in \mathcal{H}$ and assume $\neg(s \doteq t) \notin \mathcal{H}$. By Corollary 1 it follows that \mathcal{H} is saturated and hence we have that $(s \doteq t) \in \mathcal{H}$. Then, by Lemma 3(a), it follows that $s = t \in \mathcal{H}$, and thus $\{s = t, s \neq t\} \subseteq \mathcal{H}$, which contradicts $\vec{\nabla}_c$. Hence, $\neg(s \doteq t) \in \mathcal{H}$.

□

Definition 2 (Leibniz-free). *Let S be a set of formulas. S is called Leibniz-free iff $s \doteq t \notin S$ for any terms s, t .*

Corollary 2 (Impredicativity Gap). *Let $\mathcal{H} \in \mathfrak{H}$. \mathcal{H} is saturated or Leibniz-free.*

Proof. Assume that \mathcal{H} is not Leibniz-free. Then, there exists some formula $s \doteq t \in \mathcal{H}$. An application of Lemma 4(f) yields the desired result. □

Summary of properties of equality and Leibniz-equality. The following table contains an overview of the implied properties of $=$ and \doteq , respectively. A property that holds unconditionally is marked with \checkmark , a property that holds for saturated Hintikka sets is marked with **sat.**.

Property	$\star \equiv =$	$\star \equiv \doteq$
$s \star s \in \mathcal{H}$	sat.	sat.
$\neg(s \star s) \notin \mathcal{H}$	\checkmark	\checkmark
If $s \star t \in \mathcal{H}$ and $t \star u \in \mathcal{H}$, then $s \star u \in \mathcal{H}$	\checkmark	\checkmark
If $s \star t \in \mathcal{H}$, then $t \star s \in \mathcal{H}$	sat.	\checkmark
If $u[s]_p \in \mathcal{H}$ and $s \star t \in \mathcal{H}$, then $u[t]_p \in \mathcal{H}$	\checkmark	\checkmark

4 Reduction of \mathfrak{H} (Steen) to $\mathfrak{H}\text{int}_{\beta\text{fb}}$ (Brown)

We reduce the notion of Hintikka sets of Steen to the notion of Hintikka sets by Brown. Conceptually, every formula from a set $\mathcal{H} \in \mathfrak{H}$ is first translated to its red equivalent under a signature $\Sigma \subseteq \{\neg\} \cup \{=\tau \mid \tau \in \mathcal{T}\}$ containing no further primitive logical connectives. Note that this involves mapping a primitive connective to a primitive connective (i.e. $=$ to $=$) as well as mapping a defined connective to a primitive connective (i.e. \neg to \neg). In a second step, the red formulas are translated into their meta-counterparts, i.e. into external propositions.

Formally, we define the mapping as follows: Let Σ be given as introduced above. We define $\mathcal{H}_\triangleright := \{(s[\neg\backslash\neg])(=\backslash=) \mid s \in \mathfrak{H}\}$, where $s[l\backslash r]$ denotes that term in $\Lambda^c(\Sigma)$ that is obtained by replacing all occurrences of l in s by r . Obviously, if $s \in \Lambda_o^c(\Sigma)$ then $(s[\neg\backslash\neg])(=\backslash=) \in \Lambda_o^c(\Sigma)$, and thus $\mathcal{H}_\triangleright \subseteq \Lambda_o^c(\Sigma)$.

As the Hintikka properties of Brown are defined in terms of external propositions, we proceed by constructing a set $\mathcal{H}_\triangleright^\# \subseteq \text{prop}(\Sigma)$ from $\mathcal{H}_\triangleright$ by enriching $\mathcal{H}_\triangleright$ with its external counterparts: To that end, for any term $s_o \in \Lambda_o(\Sigma)$, we denote by $s^\# \in \text{prop}(\Sigma)$ the term that is constructed by replacing the primitive connective at head position by its external equivalent², i.e. we have that $(\neg s)^\#$ is equal to $\neg s$ and that $(s =^o t)^\#$ is equal to $s =^o t$. If there is no connective at head position, the term is left unchanged, i.e., $(p \overline{s^n})^\#$ equals $p \overline{s^n}$ and $(X \overline{s^n})^\#$ equals $X \overline{s^n}$ whenever $p \in \Sigma$ is a parameter and X is a variable.

Then, $\mathcal{H}_\triangleright^\#$ is defined inductively as follows. For $\mathcal{H} \in \mathfrak{H}$, let $\mathcal{H}_\triangleright^\#$ be the smallest set of external propositions over Σ such that

²This corresponds to the original definition of $s^\#$ by Brown [6, Def. 2.1.38].

- (1) if $s_o \in \mathcal{H}$, then $s \in \mathcal{H}_\triangleright^\sharp$, and
- (2) if $s_o \in \mathcal{H}$, then $s^\sharp \in \mathcal{H}_\triangleright^\sharp$, and
- (3) if $\neg s \in \mathcal{H}_\triangleright^\sharp$, then $\neg s^\sharp \in \mathcal{H}_\triangleright^\sharp$.

Intuitively, this translation adopts the translated object-level terms of \mathcal{H} (clause (1)), and additionally augments the set corresponding meta-level terms by replacing connectives at head positions by its meta-connective equivalent (clause (2)). Also, connectives directly under a (meta-)negation are considered (clause (3)).³ Deep translations are then implicitly provided by the Hintikka closure properties.

We provide simple examples of unsaturated Hintikka sets \mathcal{H} , $\mathcal{H}_\triangleright$ and $\mathcal{H}_\triangleright^\sharp$:

- Let $\mathcal{H} := \{s | s \equiv_{\beta\eta} t \text{ for } t \in \{a =^t b, a =^t a, b =^t b, b =^t a\}\}$. Then $\mathcal{H}_\triangleright = \{s | s \equiv_{\beta\eta} t \text{ for } t \in \{a =^t b, a =^t a, b =^t b, b =^t a\}\}$ and $\mathcal{H}_\triangleright^\sharp = \mathcal{H}_\triangleright \cup \{s | s \equiv_{\beta\eta} t \text{ for } t \in \{a \neq^t b, a \neq^t a, b \neq^t b, b \neq^t a\}\}$.
- Let $\mathcal{H} := \{s | s \equiv_{\beta\eta} t \text{ for } t \in \{a =^t p(\neg), a =^t a, p(\neg) =^t p(\neg), p(\neg) =^t a\}\}$. Then $\mathcal{H}_\triangleright = \{s | s \equiv_{\beta\eta} t \text{ for } t \in \{a =^t p(\neg), a =^t a, p(\neg) =^t p(\neg), p(\neg) =^t a\}\}$ and $\mathcal{H}_\triangleright^\sharp = \mathcal{H}_\triangleright \cup \{s | s \equiv_{\beta\eta} t \text{ for } t \in \{a \neq^t p(\neg), a \neq^t a, p(\neg) \neq^t p(\neg), p(\neg) \neq^t a\}\}$.

We now show that if $\mathcal{H} \in \mathfrak{H}$ then $\mathcal{H}_\triangleright^\sharp \in \mathfrak{H}\text{int}_{\beta\text{fb}}$ (i.e., $\mathcal{H}_\triangleright^\sharp$ fulfils all $\vec{\nabla}$ from §2).

Theorem 1 (Reduction of \mathfrak{H} to $\mathfrak{H}\text{int}_{\beta\text{fb}}$). *If $\mathcal{H} \in \mathfrak{H}$, then there exists an extensional Hintikka set $\mathcal{H}' \in \mathfrak{H}\text{int}_{\beta\text{fb}}$ such that $\mathcal{H}_\triangleright^\sharp \equiv \mathcal{H}'$.*

Proof. Let $\mathcal{H} \in \mathfrak{H}$ be a Hintikka set according §3, i.e., fulfilling all $\vec{\nabla}$ properties. We verify every $\vec{\nabla}$ -property for $\mathcal{H}_\triangleright^\sharp$:

- $\vec{\nabla}_c$: Assume that both $s \in \mathcal{H}_\triangleright^\sharp$ and $\neg s \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $s \in \mathcal{H}$ and $\neg s \in \mathcal{H}$. As this contradicts $\vec{\nabla}_c$, it follows that $s \notin \mathcal{H}_\triangleright^\sharp$ or $\neg s \notin \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_{\beta\eta}$: Let $s \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $s \in \mathcal{H}$. Since $s \equiv_{\beta\eta} s^\downarrow$ it follows by $\vec{\nabla}_{\beta\eta}$ that $s^\downarrow \in \mathcal{H}$. By definition, we then have $s^\downarrow \in \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_\perp$: $\neg \top \notin \mathcal{H}_\triangleright^\sharp$ is vacuously true, as \top is never the result of any term translation.
- $\vec{\nabla}_\neg$: Let $\neg \neg s \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $\neg \neg s \in \mathcal{H}$. By Lemma 2(a) it follows that $s \in \mathcal{H}$ and hence $s \in \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_\vee$: Vacuously true, as \vee is never the result of any term translation.
- $\vec{\nabla}_\wedge$: Vacuously true, as \wedge is never the result of any term translation.
- $\vec{\nabla}_\forall$: Vacuously true, as \forall is never the result of any term translation.
- $\vec{\nabla}_\exists$: Vacuously true, as \exists is never the result of any term translation.
- $\vec{\nabla}^\sharp$: Holds by definition of $\mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_m$: Let $\neg(p \overline{s^n}) \in \mathcal{H}_\triangleright^\sharp$ and let $(p \overline{t^n}) \in \mathcal{H}_\triangleright^\sharp$ for some parameter p . Then, by definition, $\neg(p \overline{s^n}) \in \mathcal{H}$ and $(p \overline{t^n}) \in \mathcal{H}$. By $\vec{\nabla}_m$ it follows that $(p \overline{t^n}) \neq (p \overline{s^n}) \in \mathcal{H}$. By $\vec{\nabla}_d$ it then follows that there is some i , $1 \leq i \leq n$, such that $t^i \neq s^i \in \mathcal{H}$. By Lemma 6(a) it follows that $s^i \neq t^i \in \mathcal{H}$ and hence $\neg(s^i \neq t^i) \in \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_{dec}$: Let p be a parameter and let $\neg((p \overline{s^n}) \neq (p \overline{t^n})) \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $\neg((p \overline{s^n}) \neq (p \overline{t^n})) \in \mathcal{H}$. By $\vec{\nabla}_d$ it follows that there is some i , $1 \leq i \leq n$, such that $\neg(s^i = t^i) \in \mathcal{H}$ and hence $\neg(s^i \neq t^i) \in \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_b$: Let $\neg(s \neq^o t) \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $\neg(s \neq^o t) \in \mathcal{H}$. By $\vec{\nabla}_b$ it follows that $\{s, \neg t\} \subseteq \mathcal{H}$ or $\{\neg s, t\} \subseteq \mathcal{H}$. Hence we have that $\{s, \neg t\} \subseteq \mathcal{H}_\triangleright^\sharp$ or $\{\neg s, t\} \subseteq \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_f$: Let $\neg(f \neq^{\nu\tau} g) \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $\neg(f \neq^{\nu\tau} g) \in \mathcal{H}$. By $\vec{\nabla}_f$ it follows that $\neg(f p \neq^{\nu} g p) \in \mathcal{H}$ for some parameter p_τ . Hence we have that $\neg(f p \neq^{\nu} g p) \in \mathcal{H}_\triangleright^\sharp$.
- $\vec{\nabla}_=$: Let $s \neq^o t \in \mathcal{H}_\triangleright^\sharp$. Then, by definition, $s \neq^o t \in \mathcal{H}$. By $\vec{\nabla}_=$ it follows that $\{s, t\} \subseteq \mathcal{H}$ or $\{\neg s, \neg t\} \subseteq \mathcal{H}$. Hence we have that $\{s, t\} \subseteq \mathcal{H}_\triangleright^\sharp$ or $\{\neg s, \neg t\} \subseteq \mathcal{H}_\triangleright^\sharp$.

³Clauses (2) and (3) intuitively reflect property $\vec{\nabla}^\sharp$ of Hintikka sets by Brown [6, Def. 5.5.1].

$\vec{\nabla}_{\rightarrow}^{\neg}$: Let $f \stackrel{\neg}{=}^{\nu\tau} g \in \mathcal{H}_{\triangleright}^{\#}$. Then, by definition, $f \stackrel{\neg}{=}^{\nu\tau} g \in \mathcal{H}$. By $\vec{\nabla}_{\rightarrow}^{\neg}$ it follows that $f s \stackrel{\neg}{=}^{\nu} g s \in \mathcal{H}$ for every closed term $s_{\tau} \in \Lambda_{\tau}^c(\Sigma)$. Hence we have that $f s \stackrel{\neg}{=}^{\nu} g s \in \mathcal{H}_{\triangleright}^{\#}$ for every closed $s_{\tau} \in \Lambda_{\tau}^c(\Sigma)$.

$\vec{\nabla}_{=}^r$: Assume $\neg(s \stackrel{\neg}{=}^i s) \in \mathcal{H}_{\triangleright}^{\#}$. Then, by definition, $\neg(s \stackrel{\neg}{=}^i s) \in \mathcal{H}$, contradicting $\vec{\nabla}_{=}^r$. Hence we have that $\neg(s \stackrel{\neg}{=}^i s) \notin \mathcal{H}_{\triangleright}^{\#}$.

$\vec{\nabla}_{=}^u$: Let $(s \stackrel{\neg}{=}^i t) \in \mathcal{H}_{\triangleright}^{\#}$ and $\neg(u \stackrel{\neg}{=}^i v) \in \mathcal{H}_{\triangleright}^{\#}$. Then, by definition, $(s \stackrel{\neg}{=}^i t) \in \mathcal{H}$ and $\neg(u \stackrel{\neg}{=}^i v) \in \mathcal{H}$. By $\vec{\nabla}_m$ it follows that $(s \stackrel{\neg}{=}^i t) \neq^o (u \stackrel{\neg}{=}^i v) \in \mathcal{H}$. By $\vec{\nabla}_d$ we then have $s \neq^i u \in \mathcal{H}$ or $t \neq^i v \in \mathcal{H}$. Hence, $\neg(s \stackrel{\neg}{=}^i u) \in \mathcal{H}_{\triangleright}^{\#}$ or $\neg(t \stackrel{\neg}{=}^i v) \in \mathcal{H}_{\triangleright}^{\#}$. For the second half, we know by Corollary 1 that \mathcal{H} is saturated and hence by Lemma 5(b) it holds that $(t \stackrel{\neg}{=}^i s) \in \mathcal{H}$. Applying $\vec{\nabla}_{=}^s$ yields $t \neq^i u \in \mathcal{H}$ or $s \neq^i v \in \mathcal{H}$, and thus $\neg(t \stackrel{\neg}{=}^i u) \in \mathcal{H}_{\triangleright}^{\#}$ or $\neg(s \stackrel{\neg}{=}^i v) \in \mathcal{H}_{\triangleright}^{\#}$.

□

We claim that this result analogously holds if the original definitions of Andrews for the logical connectives are assumed instead of the slightly modified ones introduced in §3 and used in [8]; a technical proof remains future work.

5 Use Case: Bridging Model Existence

In this section, we apply the above reduction to derive a model existence theorem for Steen's Hintikka sets. Informally, we proceed as follows: There exists an extensional $\{=, \neg\}$ -model $\mathcal{M} \in \mathfrak{M}_{\beta\text{bf}}$ such that $\mathcal{M} \models \mathcal{H}_{\triangleright}^{\#}$. Because we constructed $\mathcal{H}_{\triangleright}^{\#}$ as an extensional Hintikka set based on negation and equality, we know that the domain of Booleans in \mathcal{M} is bivalent. In order to get a model solely based on equality (as required in the notion of Steen), we subsequently restrict \mathcal{M} to terms over $\{=\tau \mid \tau \in \mathcal{T}\}$. Finally, we find an extensional model over frames isomorphic to it.⁴

First, we summarize important results by Brown [6] used in this reduction.

Theorem 2 (Model Existence for Extensional Hintikka Sets [6, Theorem 5.7.17]). *Let \mathcal{H} be an extensional Σ -Hintikka set (i.e., $\mathcal{H} \in \mathfrak{Hint}_{\beta\text{bf}}(\Sigma)$). There is an extensional Σ -model \mathcal{M} (i.e., $\mathcal{M} \in \mathfrak{M}_{\beta\text{bf}}(\Sigma)$) such that $\mathcal{M} \models \mathcal{H}$.*

Theorem 3 (Property **b** [6, Theorem 3.3.7]). *Let Σ be a signature and \mathcal{M} be an Σ -model. Suppose either $\top, \perp \in \Sigma$ or $\neg \in \Sigma$. Then \mathcal{M} satisfies **b** iff \mathcal{D}_o has two elements.*

Theorem 4 (Isomorphic Models over Frames [6, Theorem 3.5.6]). *Let $\mathcal{M} = (\mathcal{D}, @, \mathcal{E}, v) \in \mathfrak{M}_{\beta\text{bf}}(\Sigma)$ be an extensional Σ -model such that \mathcal{D}_o has two elements. There is an isomorphic Σ -model $\mathcal{M}^h = (\mathcal{D}^h, @^h, \mathcal{E}^h, v^h)$ over frames, in particular $\mathcal{D}_o^h = \{T, F\}$ and v^h is the identity.*

Now we infer a model existence theorem for Steen's Hintikka properties by bridging to those of Brown: Let $\mathcal{H} \in \mathfrak{H}$ be a Hintikka set (according to Steen) over a signature $\Sigma^=$. Let $\mathcal{H}_{\triangleright}^{\#}$ be the translated set according to §4. By Theorem 1 it follows that $\mathcal{H}_{\triangleright}^{\#}$ is an extensional Σ -Hintikka set, for $\Sigma := \{\neg\} \cup \{=\tau \mid \tau \in \mathcal{T}\} \cup \{p \mid p \in \Sigma^= \text{ is parameter}\}$. By Theorem 2 it follows that there is an extensional model $\mathcal{M} = (\mathcal{D}, @, \mathcal{E}, v) \in \mathfrak{M}_{\beta\text{bf}}(\Sigma)$ such that $\mathcal{M} \models \mathcal{H}_{\triangleright}^{\#}$. Since $\neg \in \Sigma$ it follows by Theorem 3 that \mathcal{D}_o is bivalent.⁵

Now we eliminate \neg from the signature Σ to get an extensional model over $\{=\tau \mid \tau \in \mathcal{T}\}$; we refer to this signature as $\Sigma^=$, i.e., let $\Sigma^= := \Sigma \setminus \{\neg\}$. To that end, let $\mathcal{M}' := (\mathcal{D}, @, \mathcal{E}|_{\Lambda(\Sigma^=)}, v)$. \mathcal{M}' is an extensional $\Sigma^=$ -model, in particular \mathcal{D}_o is bivalent [6, Theorem 3.3.15]. Now by Theorem 4 it follows that there is an $\Sigma^=$ -model over frames $\mathcal{M}^h = (\mathcal{D}^h, @^h, \mathcal{E}^h, v^h)$ isomorphic to \mathcal{M}' .

We now construct our desired Henkin model $\mathcal{M}_{\triangleleft}$ for \mathcal{H} over $\Sigma^=$ as follows: $\mathcal{M}_{\triangleleft} := (\mathcal{D}_{\triangleleft}, \mathcal{I}_{\triangleleft})$, where

- $\mathcal{D}_{\triangleleft} := \mathcal{D}^h$, and
- $\mathcal{I}_{\triangleleft} := c \mapsto \mathcal{E}^h(c)$ for all $c \in \Sigma^=$.

It is easy to see that $\mathcal{M}_{\triangleleft}$ is a Henkin model; in particular $\mathcal{I}(=\tau)(a, b) = T$ iff $a \equiv b$ for every $a, b \in \mathcal{D}_{\triangleleft}^h$, $\tau \in \mathcal{T}$, by property $\mathcal{L}_{=\tau}(q)$ of \mathcal{M}^h for $q \equiv \mathcal{E}(=\tau) \in \mathcal{D}_{o\tau\tau}^h$. Finally we need to verify that $\mathcal{M}_{\triangleleft} \models \mathcal{H}$ indeed holds. Let $s_o \in \mathcal{H}$. By definition $\mathcal{M}_{\triangleleft} \models s$ if and only if $\|s\|^{\mathcal{M}_{\triangleleft}, g} \equiv T$ for every g . An induction over the structure of s yields the desired result: If s is an equality of the form $(l =^{\tau} r)$, then $\|l =^{\tau} r\|^{\mathcal{M}_{\triangleleft}, g} \equiv T$ iff

⁴We take a slight indirection here: We first assume negation is part of the red signature, to make sure there exists an element in $\mathbf{n} \in \mathcal{D}_{oo}$ that is the interpretation of negation. Sadly, it seems there is currently no easier way to enforce its existence; a more convenient way would be to show that there is an extensional model that satisfies $\mathcal{L}_{\neg}(\mathbf{n})$ without having negation in the signature, cf. [6, 3] for details on these semantic \mathcal{L} -properties.

⁵The fact that \mathcal{M} satisfies property **b** follows directly from the fact that $\mathcal{M} \in \mathfrak{M}_{\beta\text{bf}}(\Sigma)$.

$\|l\|^{\mathcal{M}_{\triangleleft},g} \equiv \|r\|^{\mathcal{M}_{\triangleleft},g}$. Since $l =^{\tau} r \in \mathcal{H}_{\triangleright}^{\sharp}$, we know $\mathcal{M} \models l =^{\tau} r$ and, consequently, that $\mathcal{E}_{\varphi}(l) \equiv \mathcal{E}_{\varphi}(r)$ for every assignment φ . It follows that $\mathcal{E}_{\varphi}^h(l[\backslash=]) \equiv \mathcal{E}_{\varphi}^h(r[\backslash=])$ and hence, by definition of $\mathcal{M}_{\triangleleft}$ and the induction hypothesis, $\mathcal{I}_{\triangleleft}(=^{\tau})(\|r\|^{\mathcal{M}_{\triangleleft},g}, \|r\|^{\mathcal{M}_{\triangleleft},g}) = T$ and thus $\|l =^{\tau} r\|^{\mathcal{M}_{\triangleleft},g} \equiv T$, for every g . For parameters $p \in \Sigma^=$, the proposition follows directly. For complex formulas of the form $(s \overline{s^n})$ and for abstractions, we apply the induction hypotheses to every sub-formula. This construction yields:

Theorem 5 (Bridged Model Existence). *Let $\mathcal{H} \in \mathfrak{H}$ be a $\Sigma^=$ -Hintikka set. Then there exists a $\Sigma^=$ -Henkin model \mathcal{M} such that $\mathcal{M} \models \mathcal{H}$.*

Acknowledgements.

We thank an anonymous reviewer for very valuable feedback to this work (cf. footnote 1).

References

- [1] Peter B. Andrews. *An Introduction to Mathematical Logic and Type Theory*. Applied Logic Series. Springer, 2002.
- [2] Christoph Benzmüller and Peter Andrews. Church’s type theory. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*, pages pp. 1–62 (in pdf version). Metaphysics Research Lab, Stanford University, summer 2019 edition, 2019.
- [3] Christoph Benzmüller, Chad Brown, and Michael Kohlhase. Higher-order semantics and extensionality. *Journal of Symbolic Logic*, 69(4):1027–1088, 2004. Preprint: <http://christoph-benzmueller.de/papers/J6.pdf>.
- [4] Christoph Benzmüller, Chad Brown, and Michael Kohlhase. Semantic techniques for cut-elimination in higher-order logics. SEKI Report SR-2004-07, Saarland University, Saarbrücken, Germany, 2004. Preprint: <http://christoph-benzmueller.de/papers/R37.pdf>.
- [5] Chad E. Brown. *Set Comprehension in Church’s Type Theory*. PhD thesis, Department of Mathematical Sciences, Carnegie Mellon University, 2004.
- [6] Chad E. Brown. *Automated Reasoning in Higher-Order Logic. Set Comprehension and Extensionality in Church’s Type Theory*, volume 10 of *Studies in Logic*. College Publications, 2007.
- [7] Alonzo Church. A formulation of the simple theory of types. *J. Symb. Log.*, 5(2):56–68, 1940.
- [8] Alexander Steen. *Extensional paramodulation for higher-order logic and its effective implementation Leo-III*. PhD thesis, Freie Universität Berlin, Berlin-Dahlem, Germany, 2018. Published as DISKI – Dissertations in Artificial Intelligence, Vol. 345, AKA Verlag and IOS Press, 2018. ISBN 978-1-61499-919-5.