

THE FUNDAMENTAL GROUPS OF PRESYMPLECTIC HAMILTONIAN G -MANIFOLDS

HUI LI

ABSTRACT. We consider presymplectic manifolds equipped with Hamiltonian G -actions, G being a connected compact Lie group. A presymplectic manifold is foliated by the integral submanifolds of the kernel of the presymplectic form. For a presymplectic Hamiltonian G -manifold, Lin and Sjamaar propose a condition under which they show that the moment map image has the same “convex and polyhedral” property as the moment map image of a symplectic Hamiltonian G -manifold, a result proved independently by Atiyah, Guillemin-Sternberg, and Kirwan. In this paper, under the condition Lin and Sjamaar proposed on presymplectic Hamiltonian G -manifolds, we study the fundamental groups of such manifolds, comparing with earlier results on the fundamental groups of symplectic Hamiltonian G -manifolds. We observe that the results on the symplectic case are special cases of the results on the presymplectic case.

1. INTRODUCTION

Let M be a smooth manifold, ω be a closed 2-form on M with constant rank. If $\ker \omega = 0$, then (M, ω) is a symplectic manifold, otherwise, (M, ω) is called a **presymplectic manifold**, and ω is called a presymplectic form. Symplectic and contact manifolds are special cases of presymplectic manifolds. Let G be a connected compact Lie group acting on (M, ω) preserving ω . If (M, ω) is symplectic and the G -action is Hamiltonian with a *proper* moment map ϕ , by Atiyah, Guillemin-Sternberg, and Kirwan’s theorems, $\phi(M) \cap \mathfrak{t}_+^*$ is a closed convex polyhedral set ([2, 8, 13]), where $\mathfrak{t}_+^* \subset \mathfrak{t}^*$ is a closed positive Weyl chamber, \mathfrak{t}^* being the dual Lie algebra of a maximal torus of G . If (M, ω) is a presymplectic G -manifold, we can similarly define Hamiltonian G -actions and moment maps (see Sec. 2). If (M, ω) is presymplectic and the G -action on M is Hamiltonian with a proper moment map ϕ , then $\phi(M) \cap \mathfrak{t}_+^*$ may not be convex, or be a polyhedral set (see [20]). In recent years, Lin and Sjamaar propose a condition, called **cleanness** of the G -action, and show that under this condition, $\phi(M) \cap \mathfrak{t}_+^*$ is a closed convex polyhedral set; this set is a convex polytope if M is compact ([20]). See also [22] for a presymplectic convexity result for torus actions by Ratiu and Zung.

In [16, 17, 18], the author studies the fundamental groups of symplectic Hamiltonian G -manifolds, respectively for circle actions, for G -actions on compact and noncompact manifolds. The proof relies in a certain degree on the polyhedral property of the image $\phi(M) \cap \mathfrak{t}_+^*$ above. In this paper, when the presymplectic Hamiltonian G -manifold has this same property, we study the fundamental groups of such manifolds.

Key words and phrases. presymplectic manifold, Hamiltonian G -action, moment map, presymplectic quotient, fundamental group.

2010 classification. Primary : 53D05, 53D10, 53D20; Secondary : 55Q05.

Throughout this paper, G always denotes a connected compact Lie group, T denotes a maximal torus of G , \mathfrak{g} and \mathfrak{t} are respectively the Lie algebras of G and T , \mathfrak{g}^* and \mathfrak{t}^* are respectively the dual Lie algebras, and \mathfrak{t}_+^* denotes a closed positive Weyl chamber in \mathfrak{t}^* .

If (M, ω) is a compact connected symplectic manifold with a Hamiltonian G -action and moment map ϕ , for any $a \in \phi(M)$, we have the symplectic quotient space M_a . Due to the existence of different kinds of singular values on $\phi(M) \cap \mathfrak{t}_+^*$, the M_a 's may have many different topological types. In previous works, we show that the quotient map $M \rightarrow M/G$ induces an isomorphism $\pi_1(M) \cong \pi_1(M/G)$, and there are natural isomorphisms $\pi_1(M/G) \cong \pi_1(M_a)$ for all the symplectic quotients M_a 's. These results may no longer be true for presymplectic Hamiltonian G -manifolds. For example, $M = S^1$ is a presymplectic manifold with trivial presymplectic form, let the circle group S^1 act on M freely, then $\pi_1(M) \not\cong \pi_1(M/S^1)$. For another example, by [19], the fundamental group of a connected compact contact toric manifold of Reeb type is a finite cyclic group, while the quotient space is contractible (see also Example 4.5). For presymplectic Hamiltonian G -manifolds, we can similarly define presymplectic quotients, see Definition 1.1; Examples 8.6 and 8.7 provide examples for $\pi_1(M/G) \not\cong \pi_1(M_a)$ in the presymplectic case.

In this paper, assuming (M, ω) is a connected presymplectic Hamiltonian G -manifold, we look at the relations between $\pi_1(M)$, $\pi_1(M/G)$ and the $\pi_1(M_a)$'s. In Section 2 we will explain the terminologies null subgroup, cleanness of the action, leafwise nontangency and leafwise transitivity of the action, where the latter two are special cases of the cleanness of the action. Under the condition of the cleanness of the G action, we obtain good results. In the following Theorems 1, 2 and 3, we concentrate on compact manifolds. Since G is connected, the quotient map always induces a surjection $\pi_1(M) \twoheadrightarrow \pi_1(M/G)$. This map may have a nontrivial kernel in the presymplectic case. The kernel comes from the images of the fundamental groups of the orbits of the null group action — the group action whose orbits are everywhere tangent to the leaves of the foliation given by $\ker(\omega)$.

Theorem 1. *Let (M, ω) be a connected compact presymplectic Hamiltonian G -manifold. If the G -action is leafwise nontangent everywhere, then the quotient map induces an isomorphism $\pi_1(M) \cong \pi_1(M/G)$.*

In Theorem 1, if (M, ω) is symplectic, then the assumption on the nontangency of the action is automatically satisfied.

Suppose G acts on a manifold M . Let $M_{(H)}$ be the set of points in M with stabilizer groups conjugate to $H \subset G$, it is called the **(H) -orbit type**. It is clear that the G -orbits in $M_{(H)}$ are diffeomorphic to each other. If $M_{(H)}$ is closed, it is called a **closed orbit type**. For more general cases than that in Theorem 1, Theorem 2 gives a description on the kernel of the map $\pi_1(M) \rightarrow \pi_1(M/G)$.

Theorem 2. *Let (M, ω) be a connected compact presymplectic Hamiltonian G -manifold with a clean G -action. Let N be the null subgroup. From each closed orbit type of the N -action, take an N -orbit \mathcal{O} . Let $\langle \text{im}(\pi_1(\mathcal{O})) \rangle$ be the normal subgroup of $\pi_1(M)$ generated by the images of the $\pi_1(\mathcal{O})$'s in $\pi_1(M)$. Then the quotient map induces an isomorphism $\pi_1(M)/\langle \text{im}(\pi_1(\mathcal{O})) \rangle \cong \pi_1(M/G)$.*

Theorem 2 recovers the case of symplectic manifolds, where N is trivial, see [17].

If (M, α) is a contact manifold, where α is a contact 1-form, if we take $\omega = d\alpha$, then it is a presymplectic form. The leaves of the null foliation are the Reeb orbits of α . For a contact G -action on M , there is automatically a contact moment map $\phi: M \rightarrow \mathfrak{g}^*$ given by $\langle \phi, \xi \rangle = \phi^\xi = \alpha(\xi_M)$ for any $\xi \in \mathfrak{g}$, where ξ_M is the vector field on M generated by ξ . This moment map is G -equivariant (see e.g. [6]). It is easy to check that up to a sign, this moment map is the same as the one defined in Sec. 2 for the presymplectic G -manifold $(M, d\alpha)$.

Let (M, α) be a connected compact contact G -manifold. The G -action is called of **Reeb type** if there is a Lie algebra element $\xi \in \mathfrak{g}$ which generates the Reeb vector field of α . We can perturb α to a G -invariant contact form α' so that $\ker(\alpha) = \ker(\alpha')$, and the Reeb vector field of α' is generated by a rational element $\xi' \in \mathfrak{g}$, which is close to ξ , so ξ' is the generator of the Lie algebra of a circle subgroup of G (see for example [5]). Clearly, $\pi_1(M)$ and $\pi_1(M/G)$ are independent of the chosen invariant contact form. For contact G -manifolds, we specialize the result as follows. See Example 4.5 for an application.

Theorem 3. *Let (M, α) be a connected compact contact manifold with a clean G -action. Then the action is either trivial or is of Reeb type. When the action is of Reeb type, perturb the contact form to an invariant contact form so that its Reeb orbits are generated by a circle subgroup $S^1 \subset G$, and let $m = \text{lcm}\{k \mid \mathbb{Z}_k \subset S^1 \text{ is a stabilizer group}\}$, then S^1/\mathbb{Z}_m corresponds to a loop \mathcal{O} in M . Let $\langle \text{im}(\pi_1(\mathcal{O})) \rangle$ be the normal subgroup of $\pi_1(M)$ generated by the image of $\pi_1(\mathcal{O})$ in $\pi_1(M)$. Then the quotient map induces an isomorphism $\pi_1(M)/\langle \text{im}(\pi_1(\mathcal{O})) \rangle \cong \pi_1(M/G)$.*

For a presymplectic Hamiltonian G -manifold, we define the quotient spaces as follows, which is similar to the symplectic quotient spaces.

Definition 1.1. Let (M, ω) be a presymplectic Hamiltonian G -manifold, and let $\phi: M \rightarrow \mathfrak{g}^*$ be the G -equivariant moment map. For any $a \in \phi(M)$, define the presymplectic quotient space at a to be $M_a = \phi^{-1}(a)/G_a = \phi^{-1}(G \cdot a)/G$, where G_a is the stabilizer group of a under the coadjoint action.

Theorem 5 in Sec. 2 proved by Lin and Sjamaar states that when we have a clean Hamiltonian G -action on a connected presymplectic manifold M with a proper moment map ϕ , then $\phi(M) \cap \mathfrak{t}_+^*$ is a closed convex polyhedral set. If this set is not a single point, then there are infinitely many presymplectic quotient spaces; and like in the symplectic case, the existence of possibly different kinds of singular values infers that the quotient spaces may have many different topological types. The quotient space M_a in Definition 1.1 may not be presymplectic or smooth, similar to the symplectic case. Here we are not looking for a condition to make a certain M_a to be a smooth presymplectic manifold. We consider the family $\{M_a \mid a \in \phi(M)\}$ as topological spaces, and study their fundamental groups as topological spaces.

Theorem 4. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Then we have natural isomorphisms $\pi_1(M/G) \cong \pi_1(M_a)$ for all $a \in \phi(M)$.*

This theorem includes the symplectic case as a special case [18, Theorem 1.5].

In Example 8.1, we give an application of Theorems 2 and 4.

The paper is organized as follows. In Section 2, we introduce clean G -actions, and the notions appeared in the theorems. We also introduce equivariant moment

maps on presymplectic G -manifolds. In Section 3, We prove Theorems 1 and 2. In Section 4, we study compact contact G -manifolds and prove Theorem 3. In Section 5, we consider presymplectic Hamiltonian G -manifolds when the action is leafwise transitive everywhere. In this case, we can use the symplectization of the presymplectic G -manifold and previous results on symplectic manifolds to prove Theorem 4. In Section 6, for the proof of Theorem 4 in the general case, we state the local normal form theorem in the presymplectic case, which is established in [20]; we address the cross section theorem and a convergence theorem in the presymplectic case. In Section 7, we prove Theorem 4. In Section 8, we give an application of Theorems 2 and 4, and we give two counter examples of the theorems for non-clean G -actions.

Acknowledgement. I would like to thank Reyer Sjamaar and Yi Lin for some helpful discussions. This work is supported by the NSFC grant K110712116.

2. CLEAN G -ACTIONS AND MOMENT MAPS ON PRESYMPLECTIC G -MANIFOLDS

In this section, we explain the terminologies occurring in the theorems in the Introduction, and we set up the materials needed for the next sections.

Let (M, ω) be a presymplectic manifold. The distribution

$$\ker(\omega) = \{v \in T_m M, m \in M \mid \omega(v, \cdot) = 0\}$$

is involutive [4], hence by Frobenius' theorem, integrates to a regular foliation \mathcal{F} of M , called the **null foliation**. The leaves of \mathcal{F} may not be closed.

In [20], Lin and Sjamaar introduce the concept *clean G -actions*, unifying earlier considerations of *nontangent G -actions* by He ([11]) and *leafwise transitive torus actions* by Ishida ([12]). Here we introduce this concept, one may refer to [20] for more detail. Let a connected compact Lie group G act on a presymplectic manifold (M, ω) preserving ω . We call the action **leafwise nontangent everywhere** if

$$T_m(G \cdot m) \cap T_m \mathcal{F} = 0 \text{ for each } m \in M.$$

Let $\mathfrak{n} \subset \mathfrak{g}$ be the Lie subalgebra consisting of all elements $\xi \in \mathfrak{g}$ such that the induced vector field ξ_M is tangent everywhere to \mathcal{F} , and let N be the connected immersed Lie subgroup with Lie algebra \mathfrak{n} . It can be shown that the adjoint action of G on \mathfrak{g} preserves \mathfrak{n} , so \mathfrak{n} is an ideal and hence N is a normal subgroup ([20]). The ideal \mathfrak{n} is called the **null ideal**, and the normal subgroup N is called the **null subgroup**. We call the G -action **leafwise transitive everywhere** if

$$N \cdot m = \mathcal{F}(m) \text{ for each } m \in M, \text{ where } \mathcal{F}(m) \text{ is the leaf through } m.$$

In [20], Lin and Sjamaar defined cleanness of the G -action at any point and show that for connected manifold M , if the G -action is clean at each point on M , then the following equation holds:

$$(2.1) \quad T_m(G \cdot m) \cap T_m \mathcal{F} = T_m(N \cdot m) \text{ for each } m \in M.$$

In this paper, we only use the notion of cleanness of the action on the whole manifold, so we use this equation as the definition of cleanness of the action on the manifold, i.e., the G -action is called **clean on M** if (2.1) holds.

Next, we introduce Hamiltonian G -actions on presymplectic manifolds. The presymplectic G -action on (M, ω) is called **Hamiltonian** if there exists a **moment map** $\phi: M \rightarrow \mathfrak{g}^*$ such that

- $i(\xi_M)\omega = d\langle\phi, \xi\rangle$ for each $\xi \in \mathfrak{g}$, where ξ_M is the vector field generated by the ξ -action, and
- $\phi(g \cdot m) = \text{Ad}_g^*(\phi(m))$ for all $g \in G$ and all $m \in M$. This property is called the equivariance of ϕ .

In the case of Hamiltonian G -actions, the null ideal \mathfrak{n} satisfies

$$\mathfrak{n} = \{\xi \in \mathfrak{g} \mid \xi_M \in T\mathcal{F}\} = \{\xi \in \mathfrak{g} \mid \phi^\xi = \langle\phi, \xi\rangle \text{ is locally constant on } M\}.$$

We have the following claim on the image of the moment map for a Hamiltonian G -action. Note that this claim does not need the action to be clean.

Proposition 2.2. [20] *Let (M, ω) be a connected presymplectic G -manifold with moment map ϕ . Let \mathfrak{n} be the null ideal. Then the affine span of the image $\phi(M)$ is of the form $\lambda + \mathfrak{n}^\circ$, where λ is a fixed point in \mathfrak{g}^* for the coadjoint action, and \mathfrak{n}° is the annihilator of \mathfrak{n} .*

Under the cleanness assumption of the action, Lin and Sjamaar obtain the following theorem.

Theorem 5. [20] *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Then the fibers of ϕ are connected, $\phi(M) \cap \mathfrak{t}_+^*$ is a closed convex polyhedral set, and $\phi(M) \cap \mathfrak{t}_+^*$ is rational if and only if the null subgroup N is closed. If M is compact, then $\phi(M) \cap \mathfrak{t}_+^*$ is a convex polytope.*

Let us clarify some terms occurring in Theorem 5. A convex polyhedral set in a finite-dimensional real vector space is the intersection of closed half-spaces that is locally finite. A convex polyhedron is the intersection of a finite number of closed half-spaces. A convex polytope is a bounded convex polyhedron.

3. PRESYMPLECTIC G -MANIFOLDS AND THE PROOF OF THEOREMS 1 AND 2

In this section, we prove Theorems 1 and 2.

Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with moment map ϕ . Let g be a G -invariant Riemannian metric on M compatible with ω , which means that g is an invariant metric such that on the symplectic subbundle $(T\mathcal{F})^\perp$ perpendicular to $T\mathcal{F}$, g is compatible with $\omega|_{(T\mathcal{F})^\perp}$, i.e., for any $X, Y \in (T\mathcal{F})^\perp$, $g(X, Y) = \omega(JX, Y)$ for a G -invariant almost complex structure J on $(T\mathcal{F})^\perp$ determined by g . Let $\xi \in \mathfrak{g}$, and let $\phi^\xi = \langle\phi, \xi\rangle$ be a moment map component. Let $\text{grad}(\phi^\xi)$ be the gradient vector field of ϕ^ξ , i.e., for any $X \in TM$, $g(\text{grad}(\phi^\xi), X) = d\phi^\xi(X)$. Hence It is easy to check that

$$\text{grad}(\phi^\xi) = -J\bar{\xi}_M,$$

where $\bar{\xi}_M$ is the projected vector field of ξ_M to $(T\mathcal{F})^\perp$.

Proof of Theorem 1. First, since G is connected, by the path lifting property (see [3, Theorem 6.2 or Corollary 6.3]), the quotient map induces a surjection

$$\pi_1(M) \twoheadrightarrow \pi_1(M/G).$$

To show that the quotient map induces an injection, we need to show that if a loop in M projects to a homotopically trivial loop in M/G , then the loop in M is homotopically trivial. By the homotopy lifting property [3, Theorem 7.3], we only need to show that the lifting of any constant loop (i.e., a point) in M/G , which is

a loop in a G -orbit in M , is homotopically trivial in M . Let $G \cdot x$ be any G -orbit and α be a loop in this orbit. Identify $G \cdot x$ with G/H , where H is the stabilizer group of x . Then α lifts to a path in G from the identity to some $h \in H$. Let T be the maximal torus of G containing the identity and h . We can deform the path in G , fixing the endpoints, to a path in T , and then deform the path in T to the form $\exp(t\xi)$, where $t \in [0, 1]$, $\xi \in \mathfrak{t}$, and $\exp \xi = h$. Then α is homotopic to the loop $\exp(t\xi) \cdot x$. If this loop is not a constant, then along the flow of $\text{grad}(\phi^\xi)$ or $-\text{grad}(\phi^\xi)$ with respect to some G -invariant metric, it can be deformed to a loop $\exp(t\xi) \cdot x_0$ in the minimum set of ϕ^ξ . Since for any point $m \in \exp(t\xi) \cdot x_0$, $d\phi^\xi(m) = 0$, so $T_m(\exp(t\xi) \cdot x_0) \in T_m\mathcal{F}$. Since the action is nontangent everywhere, $T_m(\exp(t\xi) \cdot x_0) = 0$, hence $\exp(t\xi) \cdot x_0$ is a constant loop. \square

Remark 3.1. Here are easy examples of leafwise nontangent actions. Let (M_1, ω_1) be a connected compact symplectic manifold with a nontrivial symplectic (or Hamiltonian) action of G , and let M_2 be any connected compact smooth manifold with a trivial G -action. Let $M = M_1 \times M_2$ with $\omega = \pi^*\omega_1$, where $\pi: M \rightarrow M_1$ is the projection. Then (M, ω) is a presymplectic manifold with a *nontrivial* action of G which is leafwise nontangent everywhere. (Compare these examples with Proposition 4.2.)

Recall that if M is a G -manifold, $M_{(H)}$ denotes the (H) -orbit type, and all the G -orbits in $M_{(H)}$ are diffeomorphic to each other. If $M_{(K)}$ is in the closure of $M_{(H)}$, then $(H) \subset (K)$, i.e., H is conjugate to a subgroup of K . Let $O \subset M_{(K)}$ be a G -orbit, and suppose that a tubular neighborhood of O intersects $M_{(H)}$. Then by the slice theorem, we can see that a tubular neighborhood N_O of O deformation retracts to O , so a loop α' in a G -orbit O' in $M_{(H)} \cap N_O$ can be deformed to a loop α in the orbit O , so that $[\alpha'] = [\alpha] \in \pi_1(M)$.

Proof of Theorem 2. As before, since G is connected, the map

$$q_*: \pi_1(M) \rightarrow \pi_1(M/G)$$

induced by the quotient map is surjective. Since for each \mathcal{O} , $\text{im}(\pi_1(\mathcal{O}))$ maps to the trivial element by q_* , $\langle \text{im}(\pi_1(\mathcal{O})) \rangle$ maps to the trivial element by q_* , so the map induced by the quotient

$$\pi_1(M)/\langle \text{im}(\pi_1(\mathcal{O})) \rangle \rightarrow \pi_1(M/G)$$

is well defined and surjective. This shows that $\langle \text{im}(\pi_1(\mathcal{O})) \rangle \subset \ker(q_*)$. To show $\ker(q_*) \subset \langle \text{im}(\pi_1(\mathcal{O})) \rangle$, by the homotopy lifting property [3, Theorem 7.3], we only need to show that any loop in any G -orbit is homotopic to a loop in some \mathcal{O} . Let α be a loop at x in the G -orbit $G \cdot x$. As in the proof of Theorem 1, we can deform α to a loop at x of the form $\exp(t\xi) \cdot x$, where $t \in [0, 1]$, and $\xi \in \mathfrak{t}$; moreover, if $\exp(t\xi) \cdot x$ is not a constant, it can be further deformed to a loop $\exp(t\xi) \cdot x_0$, either a constant or in the minimum set of ϕ^ξ . Since for each point $m \in \exp(t\xi) \cdot x_0$, $d\phi^\xi(m) = 0$, we have $\xi_{M,m} \in T_m(N \cdot m) \subset T_m\mathcal{F}$ by cleanness of the action. So the loop $\exp(t\xi) \cdot x_0$ is in an N -orbit (which is in some orbit type of the N -action). By the paragraph above, the loop $\exp(t\xi) \cdot x_0$ can be further deformed to a loop in an N -orbit \mathcal{O} lying in a closed orbit type of the N -action. \square

For the special case of compact symplectic manifolds, these arguments shortens those in [17].

For the case of noncompact connected presymplectic Hamiltonian G -manifolds, in certain cases, one can modify the statements from the compact case.

4. CONTACT G -MANIFOLDS AND THE PROOF OF THEOREM 3

In this section, we study contact G -manifolds and prove Theorem 3.

Lemma 4.1. *Let (M, α) be a connected contact G -manifold. Then for each $\xi \in \mathfrak{n}$, either $\xi_{M,m} = 0$ for all $m \in M$ or $\xi_{M,m} \neq 0$ for all $m \in M$. Here, \mathfrak{n} is the null ideal, and ξ_M is the vector field on M generated by ξ .*

Proof. If $\mathfrak{n} = 0$, the claim is trivial. Now assume $\mathfrak{n} \neq 0$, and let $\xi \in \mathfrak{n}$. Then by the definition of \mathfrak{n} , ξ_M is tangent to the Reeb orbit (the leaf of $\ker(\omega = d\alpha)$) at any point, so ϕ^ξ is a constant on M (M is connected). While for any point $m \in M$, $\phi^\xi(m) = \alpha_m(\xi_{M,m})$, so either $\xi_{M,m} = 0$ or $\xi_{M,m} \neq 0$ for all m . \square

The next result says that on compact contact G -manifolds, if the action is leafwise nontangent everywhere, then the action is trivial.

Proposition 4.2. *Let (M, α) be a connected compact contact G -manifold. If the action is leafwise nontangent everywhere, then the action is trivial.*

Proof. Let ϕ be the moment map. By Theorem 5, $\phi(M) \cap \mathfrak{t}_+^*$ is a convex polytope. Choose a G -invariant metric on \mathfrak{g}^* . Let v be a vertex on the polytope furthest from the origin. Let ϕ_T be the moment map for the T -action. Then the image of ϕ_T is the orthogonal projection to \mathfrak{t}^* of the image of ϕ . Then v is an extremal value of ϕ_T and $\phi^{-1}(v) = \phi_T^{-1}(v)$. For each $\xi \in \mathfrak{t}$, and each $m \in \phi_T^{-1}(v)$, $d\phi^\xi(m) = 0$, so $\xi_M(m) \in T_m\mathcal{F}$, which implies $\xi_M(m) = 0$ since the action is nontangent everywhere. Then $\phi^\xi(m) = \alpha(\xi_M)(m) = 0$ for all $\xi \in \mathfrak{t}$ and all $m \in \phi_T^{-1}(v)$, which means that $\phi_T(\phi_T^{-1}(v)) = v = 0$. By the choice of v , we have $\phi(M) = 0$. Since each point is fixed by the T -action, each point is fixed by the G -action. \square

Proposition 4.3. *Let (M, α) be a connected compact contact G -manifold with a clean G -action. Then the action is either trivial or is of Reeb type.*

Proof. Since the action is clean, $T_m(G \cdot m) \cap T_m\mathcal{F} = T_m(N \cdot m)$ at each $m \in M$. By Lemma 4.1, the action is either leafwise nontangent everywhere or is of Reeb type (leafwise transitive everywhere). By Proposition 4.2, the action is trivial when it is leafwise nontangent everywhere. \square

Remark 4.4. Note that when a G -action on a contact manifold is of Reeb type, the null subgroup (which is non trivial) is a normal subgroup of G . For example, $SU(2)$ (or $SO(3)$) has no nontrivial connected normal subgroups, so $SU(2)$ (or $SO(3)$) cannot act on a contact manifold which is of Reeb type. By Proposition 4.3, there is no nontrivial clean $SU(2)$ ($SO(3)$) action on compact contact manifolds.

Now we prove Theorem 3.

Proof of Theorem 3. By Proposition 4.3, the action is either trivial or is of Reeb type. Now assume the action is of Reeb type, and the contact form is perturbed as stated in the theorem, we will show the rest of the claims.

As before, since G is connected, the quotient map $q: M \rightarrow M/G$ induces a surjection

$$q_*: \pi_1(M) \twoheadrightarrow \pi_1(M/G).$$

If the null subgroup S^1 acts freely, then $m = 1$, and S^1 corresponds to a Reeb orbit which is a loop \mathcal{O} . Clearly, $\text{im}(\pi_1(\mathcal{O})) \subset \ker(q_*)$. Since $\ker(q_*)$ is a normal subgroup, $\langle \text{im}(\pi_1(\mathcal{O})) \rangle \subset \ker(q_*)$. Now assume S^1 acts locally freely. If \mathbb{Z}_k is a stabilizer group for the S^1 -action, then S^1/\mathbb{Z}_k represents a Reeb orbit. Since M is compact, we have finitely many distinct stabilizer groups \mathbb{Z}_{k_i} 's for the Reeb orbits, where $i = 0, 1, \dots, l$. Then $\frac{m}{k_i}$ are relatively prime, so there are integers a_0, \dots, a_l , such that

$$\sum_{i=0}^l a_i \frac{1}{k_i} = \frac{1}{m}.$$

For $i = 0, \dots, l$, let $\mathcal{O}_i \approx S^1/\mathbb{Z}_{k_i}$ be a Reeb orbit, and let $x_i \in \mathcal{O}_i$ be a point. Let β_i be a path from x_0 to x_i . Let

$$\mathcal{O} = (\mathcal{O}_0^{a_0}) \cdot (\beta_1 \cdot \mathcal{O}_1^{a_1} \cdot \beta_1^{-1}) \cdot (\beta_2 \cdot \mathcal{O}_2^{a_2} \cdot \beta_2^{-1}) \cdots (\beta_l \cdot \mathcal{O}_l^{a_l} \cdot \beta_l^{-1}).$$

Then \mathcal{O} is a loop at x_0 , it is parametrized by S^1/\mathbb{Z}_m . Clearly, $q_*((\pi_1(\mathcal{O}_i))) = 0$. So $q_*((\pi_1(\mathcal{O}))) = 0$, hence $\text{im}(\pi_1(\mathcal{O})) \subset \ker(q_*)$. Hence $\langle \text{im}(\pi_1(\mathcal{O})) \rangle \subset \ker(q_*)$ since $\ker(q_*)$ is normal. Therefore, the quotient map induces a surjective homomorphism

$$\pi_1(M)/\langle \text{im}(\pi_1(\mathcal{O})) \rangle \rightarrow \pi_1(M/G).$$

To show that $\ker(q_*) \subset \langle \text{im}(\pi_1(\mathcal{O})) \rangle$, by the homotopy lifting property [3, Theorem 7.3], we need to show that any loop in any G -orbit is homotopic to a loop in \mathcal{O} . By Theorem 2, the loop in the G -orbit is homotopic to a loop in some \mathcal{O}_i . We claim that the generator of $\pi_1(\mathcal{O}_i)$ is a multiple of the generator of $\pi_1(\mathcal{O})$, hence $\ker(q_*) \subset \langle \text{im}(\pi_1(\mathcal{O})) \rangle$ follows. The claim can be shown in the following way. We consider the map

$$(q_1)_*: \pi_1(M) \rightarrow \pi_1(M/\mathbb{Z}_m)$$

induced by the quotient $q_1: M \rightarrow M/\mathbb{Z}_m$ (not a covering!). This map is injective. To see this, by the homotopy lifting property, we only need to show that the lift of any constant loop in M/\mathbb{Z}_m is homotopically trivial in M . This is clear since the lift is a point in a \mathbb{Z}_m -orbit. The group S^1/\mathbb{Z}_m acts freely on M/\mathbb{Z}_m , any Reeb orbit in M/\mathbb{Z}_m is parametrized by S^1/\mathbb{Z}_m . It is clear that $(q_1)_*([\mathcal{O}])$ is a Reeb orbit in M/\mathbb{Z}_m . Since $(q_1)_*([\mathcal{O}_i])$ is a multiple of $(q_1)_*([\mathcal{O}])$, $[\mathcal{O}_i]$ is the same multiple of $[\mathcal{O}]$. \square

Recall that a **contact toric** manifold is a contact manifold of dimension $2n+1$ with an effective T^{n+1} -action. Contact toric manifolds are the odd dimensional analog of symplectic toric manifolds.

Example 4.5. In [19], we have shown that a connected compact contact toric manifold M of Reeb type has finite cyclic fundamental group \mathbb{Z}_k for some $k \geq 1$. Suppose that the contact form is chosen so that its Reeb orbit is generated by an S^1 -action. For simplicity, we assume that S^1 acts freely. Then we have an S^1 principal bundle $M \rightarrow M/S^1$, and the long exact sequence

$$\cdots \rightarrow \pi_2(M/S^1) \xrightarrow{\partial} \pi_1(S^1) \rightarrow \pi_1(M) \rightarrow \pi_1(M/S^1) \rightarrow \pi_0(S^1) = 0 \rightarrow \cdots.$$

We see that $\pi_1(M)/\text{im}(\pi_1(S^1)) = \pi_1(M/S^1)$. Since M/S^1 is a compact connected symplectic toric manifold, $\pi_1(M/S^1) = 0$. Since M/T is contractible, $\pi_1(M/T) = 0$. So we have

$$\pi_1(M)/\text{im}(\pi_1(S^1)) = \pi_1(M/T) = 0.$$

Here, $\text{im}(\pi_1(S^1))$ corresponds to the image of $\pi_1(\mathcal{O})$ in $\pi_1(M)$, where \mathcal{O} is a Reeb orbit, parametrized by S^1 . Since $\pi_1(M) = \mathbb{Z}_k$, we have

$$\text{im}(\pi_1(S^1)) = \mathbb{Z}_k.$$

For this case when the Reeb orbits are free S^1 -orbits, we can also see in the following way that $\pi_1(M)$ is finite cyclic. By the long exact sequence above, we have

$$\pi_1(M) = \pi_1(S^1)/\text{im}(\partial(\pi_2(M/S^1))),$$

where $\pi_2(M/S^1) \cong H_2(M/S^1; \mathbb{Z})$ since $\pi_1(M/S^1) = 0$. Since M/S^1 is a compact symplectic manifold, $H_2(M/S^1; \mathbb{Z})$ has dimension at least 1; $H_2(M/S^1; \mathbb{Z})$ has no torsion since M/S^1 is toric. If the Euler class of the S^1 bundle $M \rightarrow M/S^1$ is $e = a_1 t_1 + \cdots + a_l t_l$, where $a_i \in \mathbb{Z}$ for all i , and $\{t_1, \dots, t_l\}$ is a basis of $H_2(M/S^1; \mathbb{Z})$, then $\text{im}(\partial(\pi_2(M/S^1))) = k\mathbb{Z}$, where $k = \gcd(a_1, \dots, a_l)$. So $\pi_1(M) = \mathbb{Z}_k$.

5. PROOF OF THEOREM 4 WHEN THE G -ACTION IS LEAFWISE TRANSITIVE

We observe that when the G -action is leafwise transitive everywhere, we can use the symplectization of the presymplectic G -manifold and previous results on symplectic manifolds to prove Theorem 4. The reason is that when the Hamiltonian G -action is leafwise transitive everywhere on the presymplectic manifold with a proper moment map, the moment map on its symplectization is proper onto its image. In general case, however, I do not know how to do it this way.

We now introduce the symplectization of a presymplectic G -manifold (M, ω) . Let $T^*\mathcal{F}$ be the dual bundle of $T\mathcal{F}$ and let $\pi: T^*\mathcal{F} \rightarrow M$ be the projection. We choose a G -invariant Riemannian metric on M . Let $TM \rightarrow T\mathcal{F}$ be the orthogonal projection, and let $j: T^*\mathcal{F} \rightarrow T^*M$ be the dual embedding. Let ω_0 be the standard symplectic form on T^*M , and let

$$\Omega = \pi^*\omega + j^*\omega_0.$$

Then Ω is symplectic in a neighborhood of M in $T^*\mathcal{F}$. The G -action lifts to $T^*\mathcal{F}$ and to T^*M . The germ at M of the G -manifold $T^*\mathcal{F}$ is called the **symplectization of M** , denoted X . The lifted G -action on T^*M is Hamiltonian with moment map $\phi_0: T^*M \rightarrow \mathfrak{g}^*$ given by

$$(5.1) \quad \langle \phi_0(m, y), \xi \rangle = \langle y, \xi_M(m) \rangle,$$

where $m \in M$, $y \in T_m^*M$, $\xi \in \mathfrak{g}$, and ξ_M is the vector field on M generated by ξ . Moreover, if the G -action on M is Hamiltonian with moment map ϕ , then the lifted G -action on X is Hamiltonian, with moment map

$$(5.2) \quad \psi = \pi^*\phi + j^*\phi_0.$$

Lemma 5.3. *Let M be a connected presymplectic manifold with a clean Hamiltonian G -action and moment map ϕ . Let X be the symplectization of M , and ψ be the moment map for the lifted G -action on X . Then for any $(m, y) \in X$ with $m \in M$ and $y \in T_m^*\mathcal{F}$, we have*

$$\psi(m, y) - \phi(m) = \Lambda_m(y),$$

where $\Lambda_m: T_m^*\mathcal{F} \rightarrow \mathfrak{g}^*$ is a linear map such that

- $\ker(\Lambda_m) = \{y \in T_m^*\mathcal{F} \mid y|_{T_m(N \cdot m)} = 0\}$, and
- $\text{im}(\Lambda_m) \cap \mathfrak{n}^\circ = 0$.

Here, N is the null subgroup and \mathfrak{n} the null ideal.

Proof. By (5.2) and (5.1), $\psi^\xi(m, y) - \phi^\xi(m) = \langle y, \xi_M(m) \rangle$ for any $\xi \in \mathfrak{g}$. We write

$$\langle y, \xi_M(m) \rangle = \Lambda_m^\xi(y) = \langle \Lambda_m(y), \xi \rangle,$$

then $\Lambda_m(y)$ is linear on y . Then $y \in T_m^* \mathcal{F}$ is in $\ker(\Lambda_m)$ if and only if $\langle y, \xi_M(m) \rangle = 0$ for all $\xi \in \mathfrak{g}$, and if and only if $y \in (T_m(N \cdot m))^\circ$ since the action is clean. Now suppose $\Lambda_m(y) \in \text{im}(\Lambda_m) \cap \mathfrak{n}^\circ$. Then $\Lambda_m^\xi(y) = \langle y, \xi_M(m) \rangle = 0$ for all $\xi \in \mathfrak{n}$. Since $y \in T_m^* \mathcal{F}$, $\langle y, \xi_M(m) \rangle = 0$ for all $\xi \in \mathfrak{g}$, hence $\Lambda_m(y) = 0$. \square

Lemma 5.4. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with proper moment map ϕ . Suppose that the G -action is leafwise transitive everywhere. Then the moment map of the G action on the symplectization X of M is proper onto its image.*

Proof. By Lemma 5.3 and 2.2, the moment map of the G -action on X is

$$\psi(m, y) = \phi(m) + \Lambda_m(y): X \rightarrow \mathfrak{n}^\circ \oplus \mathfrak{n}^*.$$

By assumption, ϕ is proper. The action is leafwise transitive implies that $\ker(\Lambda_m) = 0$, so Λ_m is proper onto its image. So $\psi: X \rightarrow \mathfrak{g}^*$ is proper onto its image. \square

A special case for Theorem 4 is as follows.

Theorem 6. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with proper moment map ϕ . Suppose that the G -action is leafwise transitive everywhere. Then $\pi_1(M/G) = \pi_1(M_a)$ for all $a \in \phi(M)$.*

Proof. By Lemma 5.4, the moment map of the G -action on X is proper onto its image. By the symplectic case [18, Theorem 1.5 and Remark 1.7],

$$\pi_1(X/G) = \pi_1(X_a) \quad \text{for any } a \in \psi(X).$$

It is clear that X/G is homotopy equivalent to M/G . Hence

$$\pi_1(X/G) = \pi_1(M/G).$$

If $a \in \phi(M)$, then $a \in \psi(X)$, and $\psi^{-1}(a) = \phi^{-1}(a)$, so $X_a = M_a$. Therefore

$$\pi_1(X_a) = \pi_1(M_a) \quad \text{for any } a \in \phi(M).$$

The claim follows. \square

Remark 5.5. In the proof above, the symplectic manifold X is not compact, we may not have $\pi_1(X) = \pi_1(X/G)$.

6. THE LOCAL NORMAL FORM THEOREM, THE CROSS SECTION THEOREM, AND A CONVERGENCE THEOREM

For the purpose of proving Theorem 4, in this section, we address three important theorems for presymplectic Hamiltonian G -actions.

First let us describe the local normal form theorem for a presymplectic Hamiltonian G -manifold (M, ω) with a clean G -action and moment map ϕ . Let $x \in M$, H be the stabilizer group of x , and $G_{\phi(x)}$ be the stabilizer group of $\phi(x)$ under the coadjoint action. By the equivariance of ϕ , $H \subseteq G_{\phi(x)}$. Let $\mathfrak{h} = \text{Lie}(H)$, and $\mathfrak{g}_{\phi(x)} = \text{Lie}(G_{\phi(x)})$ be the Lie algebras. Let $\mathfrak{m} = \mathfrak{g}_{\phi(x)}/\mathfrak{h}$. Let $\mathfrak{p} = \mathfrak{n}/(\mathfrak{n} \cap \mathfrak{h})$, where \mathfrak{n} is the null ideal. By Proposition 2.2, $\mathfrak{n} \subset \mathfrak{g}_{\phi(x)}$, so we may take $\mathfrak{q} = \mathfrak{m}/\mathfrak{p}$, let \mathfrak{q}^* be the dual of \mathfrak{q} . Using these notations, we can describe the local normal form theorem as follows, the theorem is established in [20, Appendix C].

Theorem 7. (*The local normal form*) Let (M, ω) be a presymplectic Hamiltonian G -manifold with a clean G -action and moment map ϕ . Let $x \in M$, with stabilizer group H . Then a G -invariant neighborhood of the orbit $G \cdot x$ in M is isomorphic as a presymplectic Hamiltonian G -manifold to

$$A = G \times_H (\mathfrak{q}^* \times S \times V),$$

where S is the “symplectic slice” on which H acts symplectically, V is the “null slice”, which is a linear H -invariant subspace of $T_x \mathcal{F}$. The moment map on A is $\phi_A([g, a, s, v]) = \text{Ad}(g)^*(\phi(x) + a + \psi(s))$, where ψ is the moment map for the H -action on S .

Note that the “null slice” V has no contribution to the moment map image.

In the local normal form theorem for a symplectic Hamiltonian G -manifold, there is no “null slice” V , and in place of \mathfrak{q}^* above, it is \mathfrak{m}^* , the dual space of the above \mathfrak{m} . See [9, 21].

Next, similar to the symplectic Hamiltonian G -action case, we establish the cross section theorem for presymplectic Hamiltonian G -actions, where G is nonabelian.

Suppose that a Lie group G acts on a manifold M . Given a point m in M with stabilizer group G_m , a submanifold $U \subset M$ containing m is called a **slice at \mathfrak{m}** if U is G_m -invariant, $G \cdot U$ is a neighborhood of m , and the map

$$G \times_{G_m} U \longrightarrow G \cdot U, \text{ with } [g, u] \longrightarrow g \cdot u,$$

is an isomorphism. For instance, consider the coadjoint action of G on \mathfrak{g}^* . Let $a \in \mathfrak{t}_+^*$. Let τ be the open face of \mathfrak{t}_+^* containing a and let G_a be the stabilizer group of a . Since all the points on τ have the same stabilizer group, we also use G_τ to denote G_a . Then the natural slice at a is $U = G_a \cdot \{b \in \mathfrak{t}_+^* \mid G_b \subset G_a\} = G_a \cdot \bigcup_{\tau' \subset \tau} \tau'$, and it is an open subset of $\mathfrak{g}_\tau^* = \mathfrak{g}_a^*$.

We have the following cross section theorem. The cross section theorem in the symplectic case is due to Guillemin and Sternberg [10, Theorem 26.7], also see [7, Cor. 2.3.6] and [15, Theorem 3.8].

Theorem 8. (*The cross section*) Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and moment map ϕ . Let $a \in \text{im}(\phi) \cap \mathfrak{t}_+^*$, let U be the natural slice at a , and G_a be the stabilizer group of a . Then the **cross section** $R = \phi^{-1}(U)$ is a G_a -invariant presymplectic submanifold of M , and every leaf of R is a leaf of M . Furthermore, the restriction $\phi|_R$ is a moment map for the action of G_a on R .

Proof. We can similarly prove the theorem as in the symplectic case with slightly more care, we refer to [15, Theorem 3.8] for more detail if necessary. Here we outline the main points of the proof. First, since the coadjoint orbits intersect U transversely, and ϕ is equivariant, hence ϕ is also transverse to U , so $R = \phi^{-1}(U)$ is a submanifold. Since U is G_a -invariant and ϕ is equivariant, R is G_a -invariant. We need to show that for any point $x \in R$, $T_x R$ is presymplectic in $T_x M$ such that $\ker(\omega_x|_{T_x R}) = \ker(\omega_x)$. Let \mathfrak{f} be the orthogonal complement of \mathfrak{g}_a in \mathfrak{g} with respect to some G -invariant metric, so $\mathfrak{g} = \mathfrak{g}_a \oplus \mathfrak{f}$. Let $\mathfrak{f}_{M,x} = \{\xi_{M,x} \mid \xi \in \mathfrak{f}\}$, the subspace tangent to the orbit $G \cdot x$ generated by \mathfrak{f} . We can check the following two things:

- (1) $T_x R$ is symplectically perpendicular to $\mathfrak{f}_{M,x}$ in $T_x M$, and
- (2) $\mathfrak{f}_{M,x}$ is a symplectic vector space of $T_x M$.

Then due to the fact $T_x M = T_x R \oplus \mathfrak{f}_{M,x}$, the theorem follows. (1) can be checked directly. Now we say some words about the proof of (2). Since the G -action is clean, $T_x(G \cdot x) \cap T_x \mathcal{F} \cong T_x(N \cdot x)$, where N is the null group. Since $G \cdot a \subset \lambda + \mathfrak{n}^\circ$ by Proposition 2.2, $\mathfrak{n} \subset \mathfrak{g}_a$, hence $\mathfrak{n} \cap \mathfrak{f} = \emptyset$. So

$$(6.1) \quad \mathfrak{f}_{M,x} \cap T_x \mathcal{F} = 0,$$

which means that $\mathfrak{f}_{M,x}$ is transverse to $\ker(\omega_x)$. Note that for any $\xi, \eta \in \mathfrak{g}$, we have

$$(6.2) \quad \omega_x(\xi_{M,x}, \eta_{M,x}) = \langle \xi, d\phi_x(\eta_{M,x}) \rangle = \langle \xi, ad^*(\eta) \cdot \phi(x) \rangle = -\langle [\xi, \eta], \phi(x) \rangle.$$

(6.1) and (6.2) imply that $\mathfrak{f}_{M,x}$ is symplectic if and only if $ad^*(\mathfrak{f})(\phi(x))$ is symplectic in $T_{\phi(x)}(G \cdot \phi(x))$. We have the splitting $T_{\phi(x)}(G \cdot \phi(x)) = T_{\phi(x)}(G_a \cdot \phi(x)) \oplus ad^*(\mathfrak{f})(\phi(x))$, and $T_{\phi(x)}(G_a \cdot \phi(x))$ is a symplectic subspace. Moreover, the subspaces $T_{\phi(x)}(G_a \cdot \phi(x))$ and $ad^*(\mathfrak{f})(\phi(x))$ are symplectically perpendicular in $T_{\phi(x)}(G \cdot \phi(x))$: let $\xi \in \mathfrak{g}_a$, $\eta \in \mathfrak{f}$, then

$$\langle [\xi, \eta], \phi(x) \rangle = \langle \eta, ad^*(\xi)\phi(x) \rangle = 0$$

since \mathfrak{f} annihilates $T_{\phi(x)}U$. Hence $ad^*(\mathfrak{f})(\phi(x))$ is a symplectic subspace. \square

The highest dimensional face τ^P of \mathfrak{t}_+^* which intersects $\phi(M)$ is called the **principal face**. If U^P is the slice at τ^P , then the cross section $R^P = \phi^{-1}(U^P) = \phi^{-1}(\tau^P)$ is called the **principal cross section**, on which only the maximal torus of G acts.

In the rest part of this section, we establish a convergence theorem for presymplectic clean Hamiltonian G -actions with proper moment maps.

Theorem 9. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Assume $0 \in \phi(M)$. Then a small open G -invariant neighborhood of $\phi^{-1}(0)$ equivariantly deformation retracts to $\phi^{-1}(0)$.*

Proof. The case when $\ker(\omega) = 0$, i.e., the symplectic Hamiltonian G -action case, is proved in [14, 24]. The same idea of proof applies here. Here we follow the proof in [14]. The main point is as follows: the action is clean equips us with the local normal form theorem (Theorem 7), which gives us local real analytic coordinates and allows us to write $\|\phi\|^2$ locally as a real analytic function, then we can use the properties of local real analytic functions and the same arguments in [14] to prove the claim. \square

Corollary 6.3. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Assume $a \in \phi(M)$. Then a small open G -invariant neighborhood of $\phi^{-1}(G \cdot a)$ equivariantly deformation retracts to $\phi^{-1}(G \cdot a)$.*

Proof. Let U be the slice at a , and let $R = \phi^{-1}(U)$ be the G_a -invariant cross section. Since a is a fixed point of G_a , by a shift of $\phi|_R$, we may think of a as the value 0 of $\phi|_R$ on R . By Theorem 8, the G -action on M is clean implies that the G_a -action on R is clean, i.e., $T_x(G_a \cdot x) \cap T_x \mathcal{F} = T_x(N \cdot x)$ for all $x \in R$. By Theorem 9, a small open G_a -invariant neighborhood of $\phi^{-1}(a)$ in R G_a -equivariantly deformation retracts to $\phi^{-1}(a)$. By the equivariance of ϕ , the claim follows. \square

7. THE FUNDAMENTAL GROUPS OF THE QUOTIENTS AND THE PROOF OF THEOREM 4

In this section, we prove Theorem 4. The method is similar to that of the proof of the theorem when $\ker(\omega) = 0$, i.e., the symplectic case ([18]). For the current presymplectic case, we need to take a deeper look at the structure of connected compact Lie groups and their Lie algebras. We use mainly two operations: removing strata from stratified spaces, and doing local deformation retractions using Theorem 9 and Corollary 6.3. For the removing process to work, we need to prove that the links of the removed strata are connected and simply connected. For the local deformation retractions, we need the (local) properness of the moment map.

A **stratified space** X is a Hausdorff and paracompact topological space defined recursively as follows: X can be decomposed into a disjoint union of (locally finite) connected pieces, called strata, which are manifolds, such that given any point x in a (connected) stratum S , there exist an open neighborhood U of x , an open ball B around x in S , a compact stratified space L , called the **link of x** , and a homeomorphism

$$B \times \overset{\circ}{C}L \longrightarrow U$$

that preserves the decompositions. Here, $\overset{\circ}{C}L$ is a cone over the link L , i.e., $(L \times [0, \infty))/L \times \{0\}$. We also call the link of x the **link of S** .

We will use the following two useful results. Lemma 7.1 can be checked using the Van-Kampen Theorem.

Lemma 7.1. [17] *Let X be a connected stratified space. If X_0 is a closed stratum in X with connected and simply connected link, then $\pi_1(X) \cong \pi_1(X - X_0)$.*

Theorem 10. [1] *Let K be a compact Lie group acting on a compact path connected and simply connected metric space X . Let H be the smallest normal subgroup of K which contains the identity component of K and all those elements of K which have fixed points. Then $\pi_1(X/K) \cong K/H$.*

Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Recall that by Theorem 5, $\phi(M) \cap \mathfrak{t}_+^*$ is a closed convex polyhedral set. We call a value a of ϕ **generic** if $\phi^{-1}(G \cdot a)$ consists of points with the smallest dimensional stabilizer groups on M . A connected set of generic values on the principal face of \mathfrak{t}_+^* is called a **chamber** of $\phi(M) \cap \mathfrak{t}_+^*$.

7.1. When $G = T$ is a torus.

In this part, we prove Theorem 4 for $G = T$, a torus. In this case, let us keep in mind that $\phi(M) \cap \mathfrak{t}_+^* = \phi(M)$, and it is a locally finite closed convex polyhedral set. Let \mathcal{H} be the set of closed half spaces involved in $\phi(M)$. Each closed half space in \mathcal{H} has an interior and a boundary, we call them **faces** of the closed half space. We call the intersections of the faces of the half spaces in \mathcal{H} **faces** of the polyhedral set $\phi(M)$. A face can consists of interior points of the polyhedral set $\phi(M)$, or it can be lower dimensional internal or external set of intersection points. The different dimensional faces of $\phi(M)$ are caused by different dimensional stabilizer groups of the action. Lower dimensional faces other than the chambers defined above are called **non-generic faces**.

Lemma 7.2. *Let (M, ω) be a connected presymplectic Hamiltonian T -manifold with a clean T -action and proper moment map ϕ . Then for any values a and b in the same chamber of $\phi(M)$, we have $\pi_1(M_a) = \pi_1(M_b)$.*

Proof. Assume a and b are in the same chamber U . Then $\phi: \phi^{-1}(U) \rightarrow U$ is a proper (equivariant) submersion. By Ehresmann's lemma, $\phi^{-1}(a)$ and $\phi^{-1}(b)$ are (equivariantly) diffeomorphic. So the claim follows. \square

Proposition 7.3. *Let (M, ω) be a connected presymplectic Hamiltonian T -manifold with a clean T -action and proper moment map ϕ . Let \mathbf{F} be a non-generic face of $\phi(M)$. Suppose M_H is an orbit type such that $M_H \cap \phi^{-1}(\mathbf{F}) \neq \emptyset$. Let \overline{U} be the closure of one chamber U such that $\mathbf{F} \subset \overline{U}$. Then the link L_H of $(M_H \cap \phi^{-1}(\mathbf{F}))/T$ in $\phi^{-1}(\overline{U})/T$ is always connected and simply connected.*

Proof. By Theorem 7, a neighborhood of an orbit with stabilizer group H is isomorphic as a presymplectic Hamiltonian T -manifold to

$$A = T \times_H (\mathfrak{q}^* \times S \times V),$$

where \mathfrak{q}^* , S and V are as stated in the theorem. Split $\mathfrak{q}^* \cong \mathbb{R}^l \times \mathbb{R}^m$, where \mathbb{R}^m is the subspace which is mapped to \mathbf{F} , split $S = S^H \times S'$ and $V = V^H \times V'$, where S^H and V^H are the subspaces fixed by H . Either of these subspaces can be 0 or the whole space. Then

$$A_H \cap \phi^{-1}(\mathbf{F}) = T \times_H (\mathbb{R}^m \times S^H \times V^H).$$

So

$$(A_H \cap \phi^{-1}(\mathbf{F}))/T = (\mathbb{R}^m \times S^H \times V^H)/H = \mathbb{R}^m \times S^H \times V^H.$$

We have

$$A \cap \phi^{-1}(\overline{U}) = T \times_H (\mathbb{R}^m \times S^H \times V^H \times (\mathbb{R}^+)^l \times (S' \cap \psi^{-1}(\overline{U})) \times V'),$$

where \mathbb{R}^+ denotes the nonnegative half space of \mathbb{R} . Then

$$(A \cap \phi^{-1}(\overline{U}))/T = (\mathbb{R}^m \times S^H \times V^H) \times ((\mathbb{R}^+)^l \times (S' \cap \psi^{-1}(\overline{U})) \times V')/H.$$

So the link of $(A_H \cap \phi^{-1}(\mathbf{F}))/T$ in $(A \cap \phi^{-1}(\overline{U}))/T$ is

$$L_H = \mathcal{S}((\mathbb{R}^+)^l \times (S' \cap \psi^{-1}(\overline{U})) \times V')/H,$$

where $\mathcal{S}(\cdot)$ denotes the sphere of the corresponding space. This is the same as the link of $(M_H \cap \phi^{-1}(\mathbf{F}))/T$ in $\phi^{-1}(\overline{U})/T$. Since by Theorem 7, V' has no contribution to $\text{im}(\phi)$, and \mathbf{F} is not a generic face, we have

$$(\mathbb{R}^+)^l \times (S' \cap \psi^{-1}(\overline{U})) \neq 0.$$

(1) First assume $S' \cap \psi^{-1}(\overline{U}) = S'$. (This happens when a neighborhood of \mathbf{F} meets only one chamber or when ψ is trivial on S' .) Then

$$L_H = \mathcal{S}((\mathbb{R}^+)^l \times S' \times V')/H.$$

If $(\mathbb{R}^+)^l \neq 0$, then $\mathcal{S}((\mathbb{R}^+)^l \times S' \times V')$ is always connected and simply connected no matter what the vector spaces S' and V' are, and H fixes $(\mathbb{R}^+)^l$, by Theorem 10, L_H is connected and simply connected. Next assume $(\mathbb{R}^+)^l = 0$, then S' needs to be a nontrivial symplectic H -representation with a nontrivial moment map (in order to have the chamber). We must have $\dim H > 0$. If $S' \cong \mathbb{C}$, and $V' = 0$, then L_H must be a point, hence is connected and simply connected. If $\dim(S' \times V') \geq 3$, then

$\mathcal{S}(S' \times V')$ is connected and simply connected. By Theorem 10, $\mathcal{S}(S' \times V')/H^0$ is connected and simply connected, where H^0 is the identity component of H . Since each element of $\Gamma = H/H^0$ acts on S' as an element of a circle, it must have a nonzero fixed point in $\mathcal{S}(S')/H^0$, by Theorem 10 again, $\mathcal{S}(S' \times V')/H^0/\Gamma = L_H$ is connected and simply connected.

(2) Next assume that $S' \cap \psi^{-1}(\overline{U}) \subsetneq S'$.

(2a) Suppose first that $(\mathbb{R}^+)^l = 0$. Then $S' \cong \mathbb{C}^k$, with $k \geq 1$, is a nontrivial symplectic H -representation with nontrivial moment map ψ . With no loss of generality, we assume the H -action on S' is effective, so $\dim(H) \leq k$. Since the H action is linear and the moment map ψ is homogeneous, to prove $\mathcal{S}((S' \cap \psi^{-1}(\overline{U})) \times V')/H$ is connected and simply connected, we only need to prove $((\mathbb{C}^k \times V' - \{0\}) \cap \psi^{-1}(\overline{U}))/H$ is connected and simply connected. The moment map for the closed subtorus H action on the symplectic vector space S'

$$\psi: S' \cong \mathbb{C}^k \longrightarrow \mathfrak{h}^*$$

is of the form $\psi(z_1, \dots, z_k) = \sum_{i=1}^k |z_i|^2 \alpha_i$, where the α_i 's are weight vectors in \mathfrak{h}^* . With no loss of generality, we may assume that the cone $\text{im}(\psi) \cap \overline{U}$ is spanned by the first certain number of α_i 's, and denote the index set of these i 's by J , where $|J| \geq \dim(H)$. By writing the rest of the α_i 's as linear combinations of the first linearly independent $\dim(H)$ number of α_i 's, we have that the map

$$\psi: S' \cong \mathbb{C}^k \longrightarrow \overline{U}$$

is given by $\psi(z_1, \dots, z_k) = \sum_{i \in J} f_i \alpha_i$, where f_i is obtained in the way above, and is of the form

$$f_i(z) = \sum_{j=1}^k a_{ij} |z_j|^2 \geq 0.$$

Let $A_i = \{(z, x) \in \mathbb{C}^k \times V' \mid f_i(z) > 0, f_j(z) \geq 0 \text{ for } j \neq i\}$. Then

$$((\mathbb{C}^k \times V' - \{0\}) \cap \psi^{-1}(\overline{U}))/H = \bigcup_{i \in J} A_i/H.$$

We may argue that each A_i/H is connected and simply connected (as in (1), we may argue that A_i/H^0 is connected and simply connected, and then argue that $A_i/H^0/\Gamma = A_i/H$ is connected and simply connected), and $(A_i/H) \cap (A_j/H)$ is connected when $i \neq j$. Then by the Van-Kampen theorem, the above union set, hence L_H is connected and simply connected. We leave this as an exercise, or we refer to the proof of [17, Lemma 3.9].

(2b) Suppose next that $(\mathbb{R}^+)^l \neq 0$. The moment map

$$\phi: (\mathbb{R}^+)^l \times S' \times V' \longrightarrow \overline{U}$$

is given by $\phi(r, z, x) = r + \psi(z)$, where $r = (r_1, \dots, r_l) \in (\mathbb{R}^+)^l$, and the moment map ψ on S' is of the form $\psi = \sum_{i \in J} f_i \alpha_i$ similar to that in (2a). For $i \in J$, let $A'_i = \{(r, z, x) \in (\mathbb{R}^+)^l \times S' \times V' \mid r_j \geq 0 \text{ for all } j = 1, \dots, l, f_i > 0 \text{ and } f_j \geq 0 \text{ for all } j \neq i\}$. For $i = 1, \dots, l$, let $B_i = \{(r, z, x) \in (\mathbb{R}^+)^l \times S' \times V' \mid r_i > 0, r_j \geq 0 \text{ for all } j \neq i\}$.

0 for $j \neq i$, and $f_j \geq 0$ for all $j \in J$. Then

$$((\mathbb{R}^+)^l \times S' \times V' - \{0\}) \cap \phi^{-1}(\overline{U}) / H = \left(\bigcup_{i \in J} A'_i / H \right) \bigcup \left(\bigcup_{i=1}^l B_i / H \right).$$

Similar to the last case, we can show that this set, hence the link L_H is connected and simply connected. \square

The proofs of the following Lemma 7.4 and Proposition 7.5 are the same as in [17]. For clarity and display of the ideas, we include the proofs here.

Lemma 7.4. *Let (M, ω) be a connected presymplectic Hamiltonian T -manifold with a clean T -action and proper moment map ϕ . Let c be a non-generic value, and a be a generic value very near c . Then $\pi_1(M_c) = \pi_1(M_a)$.*

Proof. Let O be a small open neighborhood of c containing a , and U be the chamber containing a . Let $V = O \cap U$, and let \overline{V} be the closure of V in O . By Theorem 9, $\phi^{-1}(O)$, hence $\phi^{-1}(\overline{V})$ -equivariantly deformation retracts to $\phi^{-1}(c)$, hence

$$\pi_1(\phi^{-1}(\overline{V})/T) \cong \pi_1(M_c).$$

Let B be the set of values in $\overline{V} - V$. Using Lemma 7.1 and Proposition 7.3 repeatedly in the right order, we get

$$\pi_1(\phi^{-1}(\overline{V})/T) \cong \pi_1(\phi^{-1}(\overline{V})/T - \phi^{-1}(B)/T).$$

Since $\phi^{-1}(\overline{V})/T - \phi^{-1}(B)/T$ deformation retracts to $\phi^{-1}(a)/T = M_a$, the claim follows. \square

Proposition 7.5. *Let (M, ω) be a connected presymplectic Hamiltonian T -manifold with a clean T -action and proper moment map ϕ . Then $\pi_1(M/T) = \pi_1(M_a)$ for some value a .*

Proof. Using deforming (Theorem 9 and Corollary 6.3) and removing (Lemma 7.1 and Proposition 7.3) alternately, we can achieve the proof. There can be different processes and different choices of the values a 's. Follow the same arguments as in the proof of [17, Theorem 1.6] for $G = T$. \square

Theorem 4 for the case $G = T$ follows from Lemmas 7.2, 7.4, and Proposition 7.5.

7.2. When G is nonabelian.

In this part, we prove Theorem 4 for the case when G is nonabelian. In this subsection, without specification, G always denotes a connected compact *nonabelian* Lie group.

We first prove the following facts about Lie groups which will be important to us in the sequel.

Proposition 7.6. *Let G be a connected compact semisimple nonabelian Lie group. Let $H \subset G$ be a closed subgroup with Lie algebra \mathfrak{h} , let $\mathfrak{f} = \mathfrak{g}/\mathfrak{h}$, and we may view \mathfrak{f} as a direct summand of \mathfrak{g} complementary to \mathfrak{h} . Let \mathfrak{a} be an ideal of \mathfrak{g} , let $\mathfrak{p} = \mathfrak{a}/\mathfrak{a} \cap \mathfrak{h}$, and let $\mathfrak{q} = \mathfrak{f}/\mathfrak{p}$.*

- (1) *If $\mathfrak{q} = 0$, then either $\mathfrak{a} = \mathfrak{g}$ or $H \subseteq G$ is a nonabelian subgroup.*

- (2) If $\mathfrak{q} \neq 0$, then $\dim(\mathfrak{q}) \geq 2$, and for the adjoint action of H on \mathfrak{q} , the smallest normal subgroup of H containing the identity component of H and those elements which have nonzero fixed points is H itself. If $\dim(\mathfrak{q}) = 2$, then $\mathcal{S}(\mathfrak{q})/H$ is a point, where $\mathcal{S}(\mathfrak{q})$ denotes the sphere in \mathfrak{q} .

Proof. Since G is semisimple, we can split

$$\mathfrak{g} = \bigoplus_{i=1}^k \mathfrak{s}_i,$$

where each \mathfrak{s}_i is a simple ideal with $[\mathfrak{s}_i, \mathfrak{s}_j] = 0$ for $i \neq j$, and $\text{span}[\mathfrak{s}_i, \mathfrak{s}_i] = \mathfrak{s}_i$ (see [23, Theorem 5.18]). Since \mathfrak{a} is an ideal, it is a direct sum of some factors of \mathfrak{g} , with no loss of generality, we assume

$$\mathfrak{a} = \bigoplus_{i=1}^l \mathfrak{s}_i \quad \text{with } l \leq k.$$

Since $H \subset G$ is a closed subgroup,

$$\mathfrak{h} = \bigoplus_{i=1}^k \mathfrak{h}_i,$$

where $\mathfrak{h}_i \subseteq \mathfrak{s}_i$ is a subalgebra for each i . Then $\mathfrak{a} \cap \mathfrak{h} = \bigoplus_{i=1}^l \mathfrak{h}_i$, so

$$\mathfrak{p} = \mathfrak{a}/\mathfrak{a} \cap \mathfrak{h} = \bigoplus_{i=1}^l (\mathfrak{s}_i/\mathfrak{h}_i).$$

While

$$\mathfrak{f} = \mathfrak{g}/\mathfrak{h} = \bigoplus_{i=1}^k (\mathfrak{s}_i/\mathfrak{h}_i).$$

So

$$\mathfrak{q} = \mathfrak{f}/\mathfrak{p} = \bigoplus_{i=l+1}^k (\mathfrak{s}_i/\mathfrak{h}_i).$$

(1) Assume that $\mathfrak{q} = 0$. Then either $\mathfrak{f} = 0$ which means that $H = G$ hence H is nonabelian, or $l = k$ which means that $\mathfrak{a} = \mathfrak{g}$, or $l < k$ and $\mathfrak{s}_i = \mathfrak{h}_i$ for all $l+1 \leq i \leq k$, which implies that H is nonabelian (since \mathfrak{s}_i is nonabelian).

(2) Assume that $\mathfrak{q} \neq 0$. Then $l < k$, and there is at least one i with $l+1 \leq i \leq k$ so that $\mathfrak{h}_i \subsetneq \mathfrak{s}_i$. For each i with $\mathfrak{h}_i \subsetneq \mathfrak{s}_i$, since \mathfrak{s}_i is nonabelian and \mathfrak{h}_i is a subalgebra, $\dim(\mathfrak{s}_i/\mathfrak{h}_i) \geq 2$, so $\dim(\mathfrak{q}) \geq 2$.

Let $S_i = \exp \mathfrak{s}_i$. Then $G \cong (S_1 \times \cdots \times S_k)/F$, where F is a finite central subgroup of G ([23, Theorem 5.22]). So up to dividing a finite central subgroup, $H = H_1 \times \cdots \times H_k$ with $H_i \subset S_i$ a subgroup for $1 \leq i \leq k$. Let $H^0 = H_1^0 \times \cdots \times H_k^0$ be the identity component of H . There are finitely many elements g 's of the form $g = (g_1, \dots, g_k) \in H/H^0$, where for each j , g_j is either 1 or $g_j \notin H_j^0$. Now consider a fixed i above with $l+1 \leq i \leq k$ and $\mathfrak{h}_i \subsetneq \mathfrak{s}_i$. If $g_i = 1$, then $\text{Ad}(g_i)X = X$ for all $X \in \mathfrak{s}_i/\mathfrak{h}_i$, hence $\text{Ad}(g)X = \text{Ad}(g_1, \dots, g_k)X = \text{Ad}(g_1) \cdots \text{Ad}(g_k)X = X$ since $\text{Ad}(g_j)X = X$ for $j \neq i$ (due to the fact $[\mathfrak{s}_i, \mathfrak{s}_j] = 0$ when $i \neq j$). If $g_i \notin H_i^0$, then H_i^0 is nontrivial, and g_i is in a maximal torus T_i of S_i but not in a maximal torus of H_i^0 , and $\text{Ad}(g_i)Y = Y$ for all $0 \neq Y \in \mathfrak{t}_i = \text{Lie}(T_i)$. So $\text{Ad}(g_i)Y' = Y'$, where $Y' \neq 0$ is the component of Y in $\mathfrak{s}_i/\mathfrak{h}_i$, then similar to the above, $\text{Ad}(g)Y' = Y'$. We have

shown that any $g \in H/H^0$ has a nonzero fixed point in \mathfrak{q} . So the smallest normal subgroup of H containing the identity component of H and all those elements which have nonzero fixed points is H itself.

Now assume $\dim(\mathfrak{q}) = 2$. Then there is exactly one i with $l+1 \leq i \leq k$ such that $\mathfrak{h}_i \subsetneq \mathfrak{s}_i$, and $\mathfrak{q} = \mathfrak{s}_i/\mathfrak{h}_i$, with $\mathfrak{h}_i \neq 0$. If we consider the real root space decomposition of respectively \mathfrak{h}_i and \mathfrak{s}_i , we can see that the Cartan subalgebra of \mathfrak{h}_i and of \mathfrak{s}_i must be the same, and the space \mathfrak{q} can be identified with a 2-dimensional (nonzero) root space of \mathfrak{s}_i . Let X and Y be the two eigenvectors in this 2-dimensional root space. Then there is a nonzero element Z in the Cartan subalgebra of \mathfrak{s}_i so that X , Y , and Z generate a Lie algebra isomorphic to that of $SU(2)$ (or $SO(3)$). The one parameter subgroup of H generated by Z acts on $\mathcal{S}(\mathfrak{q})$ transitively, hence $\mathcal{S}(\mathfrak{q})/H$ is a point. \square

Now we proceed with the steps of the proof of Theorem 4.

Proposition 7.7. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Let \mathcal{C} be the (closed) central face of \mathfrak{t}_+^* , and assume that $\mathcal{C} \cap \phi(M) \neq \emptyset$ and that \mathcal{C} is not the only face of \mathfrak{t}_+^* which intersects $\phi(M)$. For each orbit type $M_{(H)}$ such that $M_{(H)} \cap \phi^{-1}(\mathcal{C}) \neq \emptyset$, let L_H be the link of $(M_{(H)} \cap \phi^{-1}(\mathcal{C}))/G$ in M/G . Then L_H is connected and simply connected.*

Proof. We write $G = (G_1 \times T_c)/F$, where G_1 is a connected compact semisimple Lie group, T_c is a connected (central) torus, and F is a finite central subgroup ([23, Theorem 5.22]). Let \mathfrak{g}_1 and \mathfrak{t}_c be respectively the Lie algebras of G_1 and T_c , and \mathfrak{g}_1^* and \mathfrak{t}_c^* be their dual Lie algebras.

By Theorem 7, a neighborhood of a G -orbit in $\phi^{-1}(\mathcal{C})$ with stabilizer group (H) is isomorphic to

$$A = G \times_H (\mathfrak{q}^* \times S \times V),$$

where \mathfrak{q}^* , S and V are as explained in Theorem 7 (and the paragraph above it). Up to dividing a finite *central* subgroup, $H = H_1 \times T_1$, where $H_1 \subset G_1$ and $T_1 \subset T_c$ are closed subgroups (since H is closed). Let \mathfrak{h} , \mathfrak{h}_1 and \mathfrak{t}_1 be respectively the Lie algebras of H , H_1 and T_1 . Under the splitting of G above, let the null ideal

$$\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2, \text{ where } \mathfrak{n}_1 \subseteq \mathfrak{g}_1 \text{ and } \mathfrak{n}_2 \subseteq \mathfrak{t}_c \text{ are ideals.}$$

Then

$$\mathfrak{p} = \mathfrak{n}/\mathfrak{n} \cap \mathfrak{h} = (\mathfrak{n}_1/(\mathfrak{n}_1 \cap \mathfrak{h}_1)) \oplus (\mathfrak{n}_2/(\mathfrak{n}_2 \cap \mathfrak{t}_1)) \triangleq: \mathfrak{p}_1 \oplus \mathfrak{p}_2.$$

The stabilizer group under the coadjoint action of each point on the central face \mathcal{C} is G , so

$$\mathfrak{m} = \mathfrak{g}/\mathfrak{h} = (\mathfrak{g}_1/\mathfrak{h}_1) \oplus (\mathfrak{t}_c/\mathfrak{t}_1) \triangleq: \mathfrak{m}_1 \oplus \mathfrak{m}_2.$$

Then

$$\mathfrak{q} = \mathfrak{m}/\mathfrak{p} = (\mathfrak{m}_1/\mathfrak{p}_1) \oplus (\mathfrak{m}_2/\mathfrak{p}_2) \triangleq: \mathfrak{q}_1 \oplus \mathfrak{q}_2.$$

So we can write

$$A = G \times_H ((\mathfrak{q}_1^* \times \mathfrak{q}_2^*) \times S \times V).$$

By the moment map description on A , we have

$$A_{(H)} \cap \phi^{-1}(\mathcal{C}) = G \times_H (\mathfrak{q}_2^* \times S^H \times V^H),$$

where S^H and V^H are respectively the subspaces of S and V fixed by H . So

$$(A_{(H)} \cap \phi^{-1}(\mathcal{C}))/G = (\mathfrak{q}_2^* \times S^H \times V^H)/H = \mathfrak{q}_2^* \times S^H \times V^H.$$

While

$$A/G = (\mathfrak{q}_2^* \times S^H \times V^H) \times ((\mathfrak{q}_1^* \times S' \times V')/H),$$

where S' and V' are respectively the complementary subspaces of S^H in S and V^H in V . The link of $(A_{(H)} \cap \phi^{-1}(\mathcal{C}))/G$ in A/G is

$$L_H = \mathcal{S}(\mathfrak{q}_1^* \times S' \times V')/H,$$

where $\mathcal{S}(\cdot)$ denotes the sphere in the corresponding space. This is the same as the link of $(M_{(H)} \cap \phi^{-1}(\mathcal{C}))/G$ in M/G . By assumption, $\phi(M)$ intersects at least another higher dimensional face of \mathfrak{t}_+^* . Let $a \in \phi(M)$ be on this higher dimensional face, then $G_a \cap G_1 \subsetneq G_1$ (otherwise, $G_1 \subseteq G_a$, then $G = G_a$, so a is in \mathcal{C} , a contradiction). Since the coadjoint orbit $G \cdot a$ lies on the affine space spanned by \mathfrak{n}° (by Proposition 2.2), we have $\mathfrak{n} \subseteq \mathfrak{g}_a = \text{Lie}(G_a)$, hence $\mathfrak{n}_1 \subsetneq \mathfrak{g}_1$. Then by Proposition 7.6 applied for the semisimple group G_1 , we have 2 possibilities:

- (1) $\mathfrak{q}_1^* = 0$ and H is nonabelian, and
- (2) $\dim(\mathfrak{q}_1^*) \geq 2$ and we have the claims in part (2) of Proposition 7.6.

First assume we are in case (1). Then S' must be a nontrivial symplectic H -representation with a nontrivial moment map, hence is of dimension at least 4. So $\mathcal{S}(S' \times V')$ is connected and simply connected. By Theorem 10, $\mathcal{S}(S' \times V')/H^0$ is connected and simply connected, where H^0 is the identity component of H . Since each element in $\Gamma = H/H^0$ acts on $S' \cong \mathbb{C} \times \cdots \times \mathbb{C}$ as an element of a circle, each element in Γ has a fixed point in $\mathcal{S}(S')/H^0$. By Theorem 10 again, $(\mathcal{S}(S' \times V')/H^0)/\Gamma = \mathcal{S}(S' \times V')/H = L_H$ is connected and simply connected. Now assume we are in case (2). If $\dim(\mathfrak{q}_1^* \times S' \times V') > 2$, then $\mathcal{S}(\mathfrak{q}_1^* \times S' \times V')$ is connected and simply connected. Note that the central component T_1 of H fixes \mathfrak{q}_1^* . By Proposition 7.6 (2) and Theorem 10, L_H is connected and simply connected. If $\dim(\mathfrak{q}_1^* \times S' \times V') = 2$, i.e., $S' = V' = 0$ and $\dim(\mathfrak{q}_1^*) = 2$, then by Proposition 7.6 (2), L_H is a point hence is connected and simply connected. \square

The proofs of the following Lemmas 7.8, 7.9 and 7.10 are the same as in [17] and [18]. In the proof of Lemma 7.8, we use the cross section theorem Theorem 8. Here we omit their proofs.

Lemma 7.8. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Let $c \in \tau \cap \phi(M)$, where $\tau \neq \tau^P$ is a face of \mathfrak{t}_+^* , τ^P being the principal face, and let a be a generic value on τ^P very near c . Let O be a small open invariant neighborhood of c in \mathfrak{g}^* containing a . Let B be the set of values in $O \cap \mathfrak{t}_+^*$ other than those on the chamber of generic values containing a on τ^P . Then*

$$\pi_1(\phi^{-1}(O)/G) \cong \pi_1(\phi^{-1}(O)/G - \phi^{-1}(G \cdot B)/G).$$

Lemma 7.9. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Let $c \in \tau \cap \phi(M)$, where $\tau \neq \tau^P$ is a face of \mathfrak{t}_+^* , τ^P being the principal face, and let a be a generic value on τ^P very near c . Then $\pi_1(M_c) \cong \pi_1(M_a)$.*

Lemma 7.10. *Let (M, ω) be a connected presymplectic Hamiltonian G -manifold with a clean G -action and proper moment map ϕ . Let $\tau^P \subset \mathfrak{t}_+^* \cap \phi(M)$ be the principal face. Then $\pi_1(\phi^{-1}(G \cdot \tau^P)/G) \cong \pi_1(M_a)$ for all $a \in \tau^P$.*

Now we can finish the proof of Theorem 4:

Proof of Theorem 4 for nonabelian G . Similar to the proof of Lemma 7.8, by going to the cross sections, using Proposition 7.7 in the cross sections, and by equivariance of the action and of the moment map, we can inductively remove $\phi^{-1}(G \cdot \tau)/Gs'$ from M/G for all the nonprincipal faces $\tau s'$ of \mathfrak{t}_+^* (note that the links are local). Now assume we have achieved that

$$\pi_1(M/G) \cong \pi_1\left(M/G - \bigcup_{\tau \neq \tau^P} \phi^{-1}(G \cdot \tau)/G\right) \cong \pi_1(\phi^{-1}(G \cdot \tau^P)/G).$$

Then the theorem follows from Lemmas 7.9 and 7.10. \square

8. AN APPLICATION OF THEOREMS 2 AND 4 AND TWO COUNTER EXAMPLES

First we look at an example of Theorems 2 and 4.

Example 8.1. Let (M, ω) be a compact connected symplectic manifold of dimension $2n$ equipped with a Hamiltonian G -action and moment map ϕ . Choose a Riemannian metric on M compatible with ω . Let Y be the unitary frame bundle of M , $\pi: Y \rightarrow M$, then $(Y, \pi^*(\omega))$ is a compact connected presymplectic manifold whose leaves are the $U(n)$ -orbits. The G -action on M naturally lifts to a G -action on Y , which commutes with the $U(n)$ action. The $G \times U(n)$ -action is leafwise transitive, hence clean on Y , with null subgroup $N \supseteq U(n)$. The action of $G \times U(n)$ is Hamiltonian on Y with moment map

$$\Phi = \phi \times 0: Y \longrightarrow \mathfrak{g}^* \times \mathfrak{u}(n)^*.$$

Let \mathcal{O} be an N -orbit, which is diffeomorphic to $U(n)$. By Theorem 2,

$$(8.2) \quad \pi_1(Y)/\langle \text{im}(\pi_1(\mathcal{O})) \rangle = \pi_1(Y/(G \times U(n))) = \pi_1(M/G).$$

By Theorem 4,

$$\pi_1(Y/(G \times U(n))) = \pi_1(Y_a) \quad \text{for any } a \in \Phi(Y).$$

If $a \in \phi(M)$, then $a \in \Phi(Y)$, and $Y_a = M_a$. So

$$(8.3) \quad \pi_1(Y/(G \times U(n))) = \pi_1(M/G) = \pi_1(M_a) \quad \text{for any } a \in \phi(M).$$

On the other hand, by the long exact sequence of the $U(n)$ -fibration $\pi: Y \rightarrow M$,

$$\cdots \rightarrow \pi_1(U(n)) \rightarrow \pi_1(Y) \rightarrow \pi_1(M) \rightarrow \pi_0(U(n)) = 0 \rightarrow \cdots,$$

we get

$$(8.4) \quad \pi_1(Y)/\text{im}(\pi_1(U(n))) = \pi_1(M).$$

Since the N -orbit \mathcal{O} is diffeomorphic to $U(n)$, we have

$$(8.5) \quad \pi_1(Y)/\langle \text{im}(\pi_1(\mathcal{O})) \rangle = \pi_1(Y)/\text{im}(\pi_1(U(n))).$$

Since M is a compact connected Hamiltonian G -manifold, by [17],

$$\pi_1(M) = \pi_1(M/G).$$

This shows the consistence of (8.2) and (8.4). Equations (8.2), (8.3), (8.4) and (8.5) recover the claim for the compact symplectic manifold M ([17]):

$$\pi_1(M) = \pi_1(M/G) = \pi_1(M_a) \quad \text{for any } a \in \phi(M).$$

Now we look at two counter examples of the theorems.

Example 8.6. Let $M = S^1 \times T^2$, and $\alpha = \cos t d\theta_1 + \sin t d\theta_2$, where t is the coordinate on the first factor and (θ_1, θ_2) are the coordinates on the second factor. Then (M, α) is a contact manifold, and $(M, d\alpha)$ is presymplectic. The Reeb vector field is $R = \cos t \frac{\partial}{\partial \theta_1} + \sin t \frac{\partial}{\partial \theta_2}$. The null foliation on M is given by the orbits of the flow of R .

Let T^2 act on M by acting freely on the second factor and acting trivially on the first factor. This T^2 -action is *not clean*, one can check it by definition. Let us look at the moment map image. The moment map for the T^2 -action is $\phi(t, \theta_1, \theta_2) = (\cos t, \sin t)$, $\phi(M)$ is a circle, not a convex polytope. For any $a \in \phi(M)$, $M_a = (\text{pt} \times T^2)/T^2 = \text{pt}$, so $\pi_1(M_a) = 0$. But $M/T^2 = S^1$, so $\pi_1(M/T^2) = \mathbb{Z}$.

Example 8.7. Consider the contact manifold in Example 8.6. Let $S^1 \subset T^2$ act on M by acting freely on the first coordinate of T^2 . This S^1 -action is *not clean*, checking by definition. The moment map of this S^1 -action is $\phi(t, \theta_1, \theta_2) = \cos t$, so $\phi(M) = [-1, 1]$. We have that

$$\begin{aligned} M/S^1 &= S^1 \times S^1, \\ \phi^{-1}(0) &= 2 \text{ points} \times T^2, \text{ so } M_0 \approx 2 \text{ points} \times S^1, \text{ and} \\ \phi^{-1}(1) &= 1 \text{ point} \times T^2, \text{ so } M_1 \approx S^1. \end{aligned}$$

REFERENCES

- [1] M. A. Armstrong, *Calculating the fundamental group of an orbit space*, Proc. Amer. Math. Soc., **84** (1982), no. 2, 267-271.
- [2] M. Atiyah, *Convexity and commuting Hamiltonians*, Bull. Lond. Math. Soc. **14** (1982), 1-15.
- [3] G. Bredon, *Introduction to compact transformation groups*, Elsevier, Vol. **46** in Pure and Applied mathematics, 1972.
- [4] J. Block and E. Getzler, *Quantization of foliations*, Proceedings of the XXth International Conference on Differential Geometric Methods in Theoretical Physics (New York, 1991) (S. Catto and A. Rocha, eds.), World Scientific Publishing Co., River Edge, NJ, 1992, pp. 471-487.
- [5] C. P. Boyer and K. Galicki, *A note on toric contact geometry*, J. Geom. Phys. **35** (2000), 288-298.
- [6] H. Geiges, *Constructions of contact manifolds*, Math. Proc. Camb. Phil. Soc. (1997), 121, 455-464.
- [7] V. Guillemin, E. Lerman, and S. Sternberg, *Symplectic fibrations and multiplicity diagrams*, Cambridge University Press, Cambridge, 1996.
- [8] V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), no. 3, 491-513.
- [9] V. Guillemin and S. Sternberg, *A normal form for the moment map*, Differential geometric methods in mathematical physics, (S. Sternberg, Ed.) Reidel, Dordrecht, Holland, 1984.
- [10] V. Guillemin and S. Sternberg, *Symplectic techniques in physics*, Cambridge University Press, Cambridge, 1990.
- [11] Z. He, *Odd-dimensional symplectic manifolds*, Ph.D. thesis, Massachusetts Institute of Technology, (2010), <http://hdl.handle.net/1721.1/60189>.
- [12] H. Ishida, *Torus invariant transverse Kähler foliations*, Trans. Amer. Math. Soc. **369** (2017), no. 7, 5137-5155.
- [13] F. C. Kirwan, *Convexity properties of the moment mapping III*, Invent. Math. **77** (1984), 547-552.
- [14] E. Lerman, *Gradient flow of the norm squared of a moment map*, Enseign. Math., **51** (2005), no. 1-2, 117-127.
- [15] E. Lerman, E. Meinrenken, S. Tolman, and C. Woodward, *Nonabelian convexity by symplectic cuts*, Topology **37** (1998), no. 2, 245-259.
- [16] H. Li, π_1 of Hamiltonian S^1 -manifolds, Proc. Amer. Math. Soc. **131** (2003), no. 11, 3579-3582.

- [17] H. Li, *The fundamental group of symplectic manifolds with Hamiltonian Lie group actions*, J. of Symp. Geom., **4**, no. 3, 345-372, 2007.
- [18] H. Li, *The fundamental group of G-manifolds*, Communications in Contemporary Mathematics, **15**, no. 3, (2013) 1250056.
- [19] H. Li, *The fundamental groups of contact toric manifolds*, J. of Symplectic Geom., **18**, no. 3, 2020.
- [20] Y. Lin and R. Sjamaar, *Convexity properties of presymplectic moment maps*, J. of Symplectic Geom., **17** (2019), no. 4, 1159-1200.
- [21] C. M. Marle, *Le voisinage d'une orbite d'une action hamiltonienne d'un group de Lie*, In: Séminaire sud-rhodanien de géométrie, II (Lyon, 1983), 19-35, Travaux en cours, Hermann, Paris, 1984.
- [22] T. Ratiu and N. T. Zung, *Presymplectic convexity and (ir)rational polytopes*, J. of Symplectic Geom., **17** (2019), no. 5, 1479-1511.
- [23] M. Sepanski, *Compact Lie groups*, Springer-Verlag Berlin Heidelberg, 2007.
- [24] C. Woodward, *The Yang-Mills heat flow on the moduli space of framed bundles on a surface*, Amer. J. Math. **128** (2006), no. 2, 311-359.

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA.

Email address: hui.li@suda.edu.cn