

STABLE PAIR INVARIANTS OF LOCAL CALABI-YAU 4-FOLDS

YALONG CAO, MARTIJN KOOL, AND SERGEJ MONAVARI

ABSTRACT. In 2008, Klemm-Pandharipande defined Gopakumar-Vafa type invariants of a Calabi-Yau 4-fold X using Gromov-Witten theory. Recently, Cao-Maulik-Toda proposed a conjectural description of these invariants in terms of stable pair theory.

When X is the total space of the sum of two line bundles over a surface S , and all stable pairs are scheme theoretically supported on the zero section, we express stable pair invariants in terms of intersection numbers on Hilbert schemes of points on S . As an application, we obtain new verifications of the Cao-Maulik-Toda conjectures for low degree curve classes and find connections to Carlsson-Okounkov numbers. Some of our verifications involve genus zero Gopakumar-Vafa type invariants recently determined in the context of the log-local principle by Bousseau-Brini-van Garrel.

Finally, using the vertex formalism, we provide a few more verifications of the Cao-Maulik-Toda conjectures when thickened curves contribute and also for the case of local \mathbb{P}^3 .

1. INTRODUCTION

1.1. GW/GV invariants of Calabi-Yau 4-folds. Gromov-Witten invariants are rational numbers, which are virtual counts of stable maps from curves to a fixed algebraic variety. Due to multiple cover contributions, they are in general not integers. For Calabi-Yau 4-folds, Klemm-Pandharipande [KP] defined Gopakumar-Vafa type invariants using Gromov-Witten theory and conjectured their integrality. More specifically, let X be a smooth projective Calabi-Yau 4-fold. Gromov-Witten invariants vanish for genus $g \geq 2$ for dimensional reasons and one only needs to consider the genus zero and one cases.

The genus zero Gromov-Witten invariants of X for class $\beta \in H_2(X, \mathbb{Z})$ are defined using an insertion. Consider the evaluation map $\text{ev}: \overline{M}_{0,1}(X, \beta) \rightarrow X$. For $\gamma \in H^4(X, \mathbb{Z})$, one defines

$$\text{GW}_{0,\beta}(\gamma) = \int_{[\overline{M}_{0,1}(X, \beta)]^{\text{vir}}} \text{ev}^*(\gamma).$$

The genus zero Gopakumar-Vafa type invariants

$$(1.1) \quad n_{0,\beta}(\gamma) \in \mathbb{Q}$$

are defined in [KP] by the identity

$$\sum_{\beta > 0} \text{GW}_{0,\beta}(\gamma) q^\beta = \sum_{\beta > 0} n_{0,\beta}(\gamma) \sum_{d=1}^{\infty} d^{-2} q^{d\beta},$$

where the sum is over all non-zero effective classes in $H_2(X, \mathbb{Z})$. For the genus one case, the virtual dimension of $\overline{M}_{1,0}(X, \beta)$ is zero and one defines

$$\text{GW}_{1,\beta} = \int_{[\overline{M}_{1,0}(X, \beta)]^{\text{vir}}} 1 \in \mathbb{Q}.$$

The genus one Gopakumar-Vafa type invariants

$$(1.2) \quad n_{1,\beta} \in \mathbb{Q}$$

are defined in [KP] by the identity

$$\begin{aligned} \sum_{\beta > 0} \text{GW}_{1,\beta} q^\beta &= \sum_{\beta > 0} n_{1,\beta} \sum_{d=1}^{\infty} \frac{\sigma(d)}{d} q^{d\beta} + \frac{1}{24} \sum_{\beta > 0} n_{0,\beta} (c_2(X)) \log(1 - q^\beta) \\ &\quad - \frac{1}{24} \sum_{\beta_1, \beta_2} m_{\beta_1, \beta_2} \log(1 - q^{\beta_1 + \beta_2}), \end{aligned}$$

where $\sigma(d) = \sum_{i|d} i$ and $m_{\beta_1, \beta_2} \in \mathbb{Z}$ are called meeting invariants, which can be inductively determined by the genus zero Gromov-Witten invariants of X . In [KP], both of the invariants (1.1), (1.2) are conjectured to be integers. Using localization techniques and mirror symmetry, they calculate the Gromov-Witten invariants of X in numerous examples in support of their

integrality conjecture. The genus zero integrality conjecture has been proved by Ionel-Parker using symplectic geometry [IP, Thm. 9.2].

1.2. Stable pair invariants of Calabi-Yau 4-folds. Stable pairs were introduced in general by Le Potier [LeP] and used by Pandharipande-Thomas to define virtual invariants of smooth projective threefolds [PT1, PT2, PT3]. Stable pair invariants of threefolds are related to Gromov-Witten invariants by the celebrated GW/PT correspondence [MNOP, PT1], which has been proved in many cases by Pandharipande-Pixton [PP1, PP2].

In [CMT2], Cao-Maulik-Toda studied stable pair theory of a smooth projective Calabi-Yau 4-fold X . They used stable pair invariants of X to give a sheaf theoretical interpretation of the Gopakumar-Vafa type invariants (1.1) and (1.2)¹.

Let $P_n(X, \beta)$ be the moduli space of stable pairs $\{s : \mathcal{O}_X \rightarrow F\}$ with $\text{ch}(F) = (0, 0, 0, \beta, n)$. There exists a virtual class

$$(1.3) \quad [P_n(X, \beta)]^{\text{vir}} \in H_{2n}(P_n(X, \beta), \mathbb{Z}),$$

in the sense of Borisov-Joyce [BJ], which depends on the choice of an orientation of a certain (real) line bundle over $P_n(X, \beta)$ [CGJ]. For $\gamma \in H^4(X, \mathbb{Z})$, we define primary insertions

$$\tau : H^4(X, \mathbb{Z}) \rightarrow H^2(P_n(X, \beta), \mathbb{Z}), \quad \tau(\gamma) = \pi_{P*}(\pi_X^* \gamma \cup \text{ch}_3(\mathbb{F})),$$

where π_X, π_P are projections from $X \times P_n(X, \beta)$ to the corresponding factors and

$$\mathbb{I}^\bullet = \{\mathcal{O} \rightarrow \mathbb{F}\}$$

is the universal stable pair on $X \times P_n(X, \beta)$. Note that $\text{ch}_3(\mathbb{F})$ is Poincaré dual to the fundamental cycle of \mathbb{F} . The stable pair invariants of X with primary insertions are defined by

$$(1.4) \quad P_{n, \beta}(\gamma) := \int_{[P_n(X, \beta)]^{\text{vir}}} \tau(\gamma)^n.$$

When $n = 0$, we simply denote this invariant by $P_{0, \beta}$. We set $P_{0, 0} := 1$ and $n_{0, 0}(\gamma) := 0$.

Conjecture 1.1. ([CMT2]) *Let X be a smooth projective Calabi-Yau 4-fold, $\beta \in H_2(X, \mathbb{Z})$, $\gamma \in H^4(X, \mathbb{Z})$, and $n \geq 1$. Then there exist choices of orientations such that*

$$P_{n, \beta}(\gamma) = \sum_{\substack{\beta_0 + \beta_1 + \dots + \beta_n = \beta \\ \beta_0, \beta_1, \dots, \beta_n \geq 0}} P_{0, \beta_0} \cdot \prod_{i=1}^n n_{0, \beta_i}(\gamma),$$

where the sum is over all effective decompositions of β .

Conjecture 1.2. ([CMT2]) *Let X be a smooth projective Calabi-Yau 4-fold. Then there exist choices of orientations such that*

$$\sum_{\beta \geq 0} P_{0, \beta} q^\beta = \prod_{\beta > 0} M(q^\beta)^{n_{1, \beta}},$$

where $M(q) = \prod_{k \geq 1} (1 - q^k)^{-k}$ denotes the MacMahon function.

Conjecture 1.1 can be interpreted as a wall-crossing formula in the category of D0-D2-D8 bound states in Calabi-Yau 4-folds [CT1], while Conjecture 1.2 seems to be more mysterious. In [CMT2], these conjectures were verified in the following cases (modulo some minor assumptions in some of the cases). In each case, Conjecture 1.1 was only verified for $n = 1$.

- X is a general sextic and $\beta = [\ell], 2[\ell]$, where $\ell \subseteq X$ is a line.
- X is a Weierstrass elliptic fibration and $\beta = r[F]$, where $[F]$ is the fibre class and $r > 0$ (in the case of Conjecture 1.1 only for $r = 1$).
- $X = Y \times E$, where Y is a smooth projective Calabi-Yau threefold, E is an elliptic curve, and β is the push-forward of an irreducible class on $Y \times \{\text{pt}\}$.
- $X = Y \times E$, where Y is a smooth projective Calabi-Yau threefold, E is an elliptic curve, and $\beta = r[E]$, where $[E]$ is the fibre class and $r > 0$ (Conjecture 1.2 only).

When X is either the total space of a smooth projective Fano threefold, or the total space of $\mathcal{O}(-1) \oplus \mathcal{O}(-2)$ on \mathbb{P}^2 , or $\mathcal{O}(-1, -1) \oplus \mathcal{O}(-1, -1)$ on $\mathbb{P}^1 \times \mathbb{P}^1$, the moduli spaces $P_n(X, \beta)$ are projective and it makes sense to consider Conjectures 1.1 and 1.2. In this setting, the conjectures were verified in some cases for irreducible curve classes in [CMT2].

One of the main goals of this paper is to provide more verifications for these local geometries for more general low degree curve classes.

¹In [CMT1, CT2], the authors also proposed a sheaf theoretical interpretation of (1.1), (1.2) using Donaldson-Thomas type counting invariants of one dimensional stable sheaves on X .

1.3. Stable pair invariants of local surfaces. Let S be a smooth projective surface and let L_1, L_2 be two line bundles on S satisfying $L_1 \otimes L_2 \cong K_S$. Then the total space X of $L_1 \oplus L_2$ over S is a non-proper Calabi-Yau 4-fold, which we refer to as a *local surface*. Consider the moduli space $P_n(X, \beta)$ of stable pairs (F, s) with $\chi(F) = n$ and such that F has proper scheme theoretic support in class $\beta \in H_2(X, \mathbb{Z})$. Although $P_n(X, \beta)$ is in general non-proper, it can be proper in several interesting cases (Propositions 3.1, 3.8). Then we can define virtual classes (1.3) and corresponding stable pair invariants (1.4).

Example 1.3. For $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$ and $(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, the moduli space $P_n(X, \beta)$ is projective for all n, β (see Proposition 3.1).

Example 1.4. For $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, $P_n(X, \beta)$ is in general non-proper. E.g. let $H_1 := \{\text{pt}\} \times \mathbb{P}^1$, take $\beta = [H_1]$, and $n = \chi(\mathcal{O}_{H_1}) = 1$. Then $N_{H_1/X} \cong \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-2)$ has sections in the first fibre direction, so $H_1 \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \subseteq X$ can move off the zero section $\mathbb{P}^1 \times \mathbb{P}^1 \subseteq X$ and $P_1(X, [H_1])$ is non-proper. On the other hand, for $H_2 := \mathbb{P}^1 \times \{\text{pt}\}$ and $\beta = [H_2]$, we have $\beta \cdot L_1 < 0$ and $\beta \cdot L_2 < 0$, so $P_1(X, [H_2])$ is projective by Proposition 3.1.

When S is toric, the local surface X is toric and the vertex formalism for calculating stable pair invariants of X has been developed in [CK2, CKM] in analogy with [PT2]. Let $T \subseteq (\mathbb{C}^*)^4$ denote the 3-dimensional subtorus preserving the Calabi-Yau volume form, then the fixed locus $P_n(X, \beta)^T$ consists of finitely many isolated reduced points [CK2, Sect. 2.2], though the number of fixed points is typically very large making calculations using the vertex formalism cumbersome.

Although we perform a few new calculations using the vertex formalism as well, we mainly focus on another approach, where we use the global geometry of S . We consider the case when all stable pairs on X are scheme theoretically supported on the zero section $\iota : S \hookrightarrow X$, i.e. we have an isomorphism

$$\iota_* : P_n(S, \beta) \cong P_n(X, \beta).$$

Under this isomorphism, we have (Proposition 4.2)

$$(1.5) \quad [P_n(X, \beta)]^{\text{vir}} = (-1)^{\beta \cdot L_2 + n} \cdot e(-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)) \cdot [P_n(S, \beta)]^{\text{vir}},$$

where $[P_n(S, \beta)]^{\text{vir}}$ is the virtual class of the pairs obstruction theory on S , $e(\cdot)$ denotes Euler class, $\pi_{P_S} : S \times P_n(S, \beta) \rightarrow P_n(S, \beta)$ is the projection, $\mathbf{R}\mathcal{H}om_{\pi_{P_S}} = \mathbf{R}\pi_{P_S*} \circ \mathbf{R}\mathcal{H}om$, and \mathbb{F} is the universal 1-dimensional sheaf on $S \times P_n(S, \beta)$. The sign $(-1)^{\beta \cdot L_2 + n} = (-1)^{\beta \cdot c_1(Y) + n}$, where $Y = \text{Tot}_S(L_1)$, comes from a preferred choice of orientation on $P_n(X, \beta)$ which was discussed in a similar situation in [Cao].

In order to use (1.5) for calculations, we need the fact that $P_n(S, \beta)$ is isomorphic to a relative Hilbert scheme. More precisely, assume $b_1(S) = 0$ and denote by $|\beta|$ the linear system determined by β . Denote by $\mathcal{C} \rightarrow |\beta|$ the universal curve, then [PT3, Prop. B.8] gives

$$P_n(S, \beta) \cong \text{Hilb}^m(\mathcal{C}/|\beta|),$$

where $\text{Hilb}^m(\mathcal{C}/|\beta|)$ denotes the relative Hilbert scheme of m points on the fibres of $\mathcal{C} \rightarrow |\beta|$ and

$$m = n + g(\beta) - 1 = n + \frac{1}{2}\beta(\beta + K_S).$$

This isomorphism was exploited in order to determine the surface contribution to stable pair invariants of local surfaces $\text{Tot}_S(K_S)$ in [KT2]. The relative Hilbert scheme $\text{Hilb}^m(\mathcal{C}/|\beta|)$ is an incidence locus in a smooth ambient space

$$\text{Hilb}^m(\mathcal{C}/|\beta|) \subseteq S^{[m]} \times |\beta|,$$

where $S^{[m]}$ denotes the Hilbert scheme of m points on S . More precisely, $\text{Hilb}^m(\mathcal{C}/|\beta|)$ is cut out tautologically by a section of a vector bundle on $S^{[m]} \times |\beta|$ as we recall in Section 4.1. This allows us to express the stable pair invariants of X in terms of intersection numbers on $S^{[m]} \times |\beta|$, or more precisely, on the “virtual” ambient space $S^{[m]} \times \mathbb{P}^{\chi(\beta)-1}$, where

$$\chi(\beta) := \chi(\mathcal{O}_S(\beta)).$$

In what follows, $\mathcal{Z} \subseteq S \times S^{[m]}$ denotes the universal subscheme and \mathcal{I} is the corresponding ideal sheaf. For any line bundle \mathcal{L} on S , the corresponding tautological bundle is defined by

$$\mathcal{L}^{[m]} := p_* q^* \mathcal{L},$$

where $p : \mathcal{Z} \rightarrow S^{[m]}$ and $q : \mathcal{Z} \rightarrow S$ are projections. Moreover, we consider the “twisted tangent bundle” [CO12]

$$(1.6) \quad T_{S^{[m]}}(\mathcal{L}) := \mathbf{R}\Gamma(S, \mathcal{L}) \otimes \mathcal{O} - \mathbf{R}\mathcal{H}om_{\pi}(\mathcal{I}, \mathcal{I} \boxtimes \mathcal{L}),$$

where $\pi : S \times S^{[m]} \rightarrow S^{[m]}$ denotes projection. Finally, we denote the total Chern class by c and the tautological line bundle on $\mathbb{P}^{\chi(\beta)-1}$ by $\mathcal{O}(1)$. We prove the following result (Theorem 4.4).

Theorem 1.5. *Let S be a smooth projective surface with $b_1(S) = p_g(S) = 0$ and $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$. Suppose $\beta \in H_2(S, \mathbb{Z})$ and $n \geq 0$ are chosen such that $P_n(X, \beta) \cong P_n(S, \beta)$ for $X = \text{Tot}_S(L_1 \oplus L_2)$. Denote by $[\text{pt}] \in H^4(X, \mathbb{Z})$ the pull-back of the Poincaré dual of the point class on S . Let $P_n(X, \beta)$ be endowed with the orientation as in (1.5). Then*

$$P_{n,\beta}([\text{pt}]) = (-1)^{\beta \cdot L_2 + n} \int_{S^{[m]} \times \mathbb{P}^{X(\beta)-1}} c_m(\mathcal{O}_S(\beta)^{[m]}(1)) \frac{h^n(1+h)^{X(L_1(\beta))}(1-h)^{X(L_2(\beta))} c(T_{S^{[m]}}(L_1))}{c(L_1(\beta)^{[m]}(1)) \cdot c((L_2(\beta)^{[m]}(1))^{\vee})},$$

when $\beta^2 \geq 0$. Here $m := n + g(\beta) - 1$ and $h := c_1(\mathcal{O}(1))$. Moreover, $P_{n,\beta}([\text{pt}]) = 0$ when $\beta^2 < 0$.

The main assumption in this theorem is $P_n(X, \beta) \cong P_n(S, \beta)$. For (S, L_1, L_2) with S minimal and toric, $L_1 \otimes L_2 \cong K_S$ with L_1^{-1}, L_2^{-1} non-trivial and nef, we classify all cases for which $n \geq 0$, $P_n(X, \beta) \cong P_n(S, \beta)$, and $P_n(S, \beta)$ is non-empty (Proposition 3.9, Remark 3.10). Note that $P_n(X, \beta) \cong P_n(S, \beta)$ more or less forces $p_g(S) = 0$, because as soon as L_1 or L_2 has non-zero sections this isomorphism does not hold. See Remark 4.5 for an extension to the case $b_1(S) > 0$.

1.4. Verifications. In this paper, we apply Theorem 1.5 to examples for which S is in addition toric². Then the integrals on $S^{[m]}$ of Theorem 1.5 can be calculated using Atiyah-Bott localization for the lift of the 2-dimensional torus action from S to $S^{[m]}$ as described in Section 4.4. This leads to the tables for stable pair invariants in Appendix A.

Denote by $[H] \in H_2(\mathbb{P}^2, \mathbb{Z})$ the class of a line and let $[H_1], [H_2] \in H_2(\mathbb{P}^1 \times \mathbb{P}^1, \mathbb{Z})$ be as in Example 1.4. In [KP, Sect. 3], Klemm-Pandharipande determined the Gromov-Witten invariants of $X = \text{Tot}_S(L_1 \oplus L_2)$ for $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$ and $(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$. They tabulated the corresponding values of the Gopakumar-Vafa type invariants for $\beta = d[H]$ with $d \leq 10$ resp. $\beta = d_1[H_1] + d_2[H_2]$ with $d_1, d_2 \leq 6$. Combining their calculations and the tables in Appendix A, we deduce the following:

Corollary 1.6. *In the following cases, Conjectures 1.1 and 1.2 are true for $X = \text{Tot}_S(L_1 \oplus L_2)$.*

- $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $d = 1$, and any $n \geq 0$.
- $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $d = 2, 3, 4$, and $n = 0, 1$.
- $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $d = 2, 3$, and $n = 2$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (1, 0), (0, 1), (1, 1)$, any $n \geq 0$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (0, d), (d, 0)$ with $d \geq 2$, and $0 \leq n \leq d$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (1, d), (d, 1)$ with $d \geq 2$, and $n = 0, 1, 2$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (2, 2), (2, 3), (3, 2), (2, 4), (4, 2), (3, 3)$, and $n = 0$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (2, 2), (2, 3), (3, 2)$, and $n = 1$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (2, 2)$, and $n = 2$.

Remark 1.7. In all these cases $P_n(X, \beta) \cong P_n(S, \beta)$. In fact, these are *all* (S, L_1, L_2) with $L_1 \otimes L_2 \cong K_S$ for which L_1^{-1}, L_2^{-1} are ample, $n \geq 0$, and $P_n(X, \beta) \cong P_n(S, \beta)$ by Propositions 3.2 and 3.9. Calculations based on Theorem 1.5 are often more efficient than the vertex formalism [CK2, CKM]. For instance, for $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (2, 4)$ and $n = 0$, $P_n(X, \beta)$ has 182 T -fixed points, whereas Theorem 1.5 only involves an integral over $S^{[2]} \times \mathbb{P}^{14}$.

Bousseau-Brini-van Garrel [BBG] recently determined the genus zero Gromov-Witten (and hence Gopakumar-Vafa type) invariants of several local surfaces for their verifications of the log-local principle conjectured in general in [GGR]. Combining their numbers with the tables for stable pair invariants in Appendix A allows us to provide some further verifications of Conjecture 1.1 as we will now describe. For any $a \geq 1$, consider the Hirzebruch surface

$$\mathbb{F}_a = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(a)).$$

We denote by $[F]$ the class of a fibre and by $[B]$ the class of the unique section satisfying $B^2 = -a$. We write $\mathcal{O}(m, n) := \mathcal{O}(mB + nF)$ and consider curve classes $\beta := d_1[B] + d_2[F]$, $d_1, d_2 \geq 0$.

Corollary 1.8. *In the following cases, Conjecture 1.1 is true for $X = \text{Tot}_S(L_1 \oplus L_2)$.*

- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 1)$, and any $n \geq 1$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, d)$ with $d \geq 2$, and $n = d$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (2, 2), (2, 3), (1, d), (d, 1)$ with $d \geq 1$, and $n = 1$.

²To our knowledge, all local surfaces which are Calabi-Yau 4-folds and for which Gopakumar-Vafa type invariants have been calculated so far are toric.

- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (1, d)$ with $d \geq 2$, and $n = 2$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 1)$, and any $n \geq 1$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, d)$ with $d \geq 2$, and $n = d$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (2, 2), (2, 3), (2, 4), (1, d)$ with $d \geq 1$, and $n = 1$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (1, d)$ with $d \geq 2$, and $n = 2$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(0, -1), \mathcal{O}(-2, -2))$, $(d_1, d_2) = (2, 2), (2, 3), (1, d)$ with $d \geq 1$, $n = 1$.
- $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 1)$, and any $n \geq 1$.
- $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, d)$ with $d \geq 2$, and $n = d$.
- $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (2, 3), (2, 4), (2, 5), (1, d)$ with $d \geq 1$, and $n = 1$.
- $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (1, d)$ with $d \geq 2$, and $n = 2$.

In all these cases $P_n(X, \beta) \cong P_n(S, \beta)$ and $n > 0$. Since Bousseau-Brini-van Garrel only determined the *genus zero* Gopakumar-Vafa type invariants for the above geometries, we can only verify Conjecture 1.1 in these cases. In fact, in Proposition 3.9 and Remark 3.10, we classify *all* cases (S, L_1, L_2) such that S is minimal toric, $L_1 \otimes L_2 \cong K_S$, L_1^{-1}, L_2^{-1} are non-trivial and nef, $n \geq 0$, $P_n(X, \beta) \cong P_n(S, \beta)$, and $P_n(S, \beta)$ is non-empty. Using Theorem 1.5, we determined the stable pair invariants in all these cases, *including* the $n = 0$ case (see Appendix A).

Remark 1.9. For all calculations done in Appendix A for which $P_n(X, \beta) \cong P_n(S, \beta)$ and the invariant is non-zero, we have

$$(1.7) \quad P_{n, \beta}([\text{pt}]) = \pm \int_{S^{[m]}} e(T_{S^{[m]}}(L_1)).$$

These numbers were calculated by Carlsson-Okounkov [CO12] and are determined by the formula

$$\sum_{m=0}^{\infty} q^m \int_{S^{[m]}} e(T_{S^{[m]}}(L_1)) = \prod_{m=1}^{\infty} (1 - q^m)^{-c_2(T_S \otimes L_1)},$$

where $c_2(T_S \otimes L_1) = c_2(S) - L_1 L_2$. We do not know whether (1.7) is a mere coincidence.

1.5. Vertex calculations. Although most calculations in this paper are based on Theorem 1.5, we also did some computations using the vertex formalism.

Proposition 1.10. *For the following cases, Conjectures 1.1 and 1.2 are true.*

- $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$, $d = 1$, and any $n \geq 0$.
- $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$, $d = 2, 3$, and $n = 0, 1$.
- $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$, $d = 2$, and $n = 2$.

For the following cases Conjecture 1.1 is true for $X = \text{Tot}_S(L_1 \oplus L_2)$.

- $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $d = 2$, and $n = 3$.
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (0, 2), (2, 0), (1, 2), (2, 1)$, and $n = 3$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 2)$, and $n = 3$.

The invariants in this proposition are defined by localization on the fixed locus [CK2, CKM]. In the cases above where $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$, we have $P_n(X, \beta) \cong P_n(\mathbb{P}^3, \beta)$ and

$$[P_n(X, \beta)]^{\text{vir}} = (-1)^{\beta \cdot c_1(\mathbb{P}^3) + n} \cdot [P_n(\mathbb{P}^3, \beta)]_{\text{pair}}^{\text{vir}},$$

where $[P_n(\mathbb{P}^3, \beta)]_{\text{pair}}^{\text{vir}}$ is the virtual class of the pairs perfect obstruction theory on \mathbb{P}^3 (discussed in (4.9), see also [CMT2, Lem. 3.1] in a similar setting). The sign in this formula is a preferred choice of orientation on $P_n(X, \beta)$ similar to (1.5). Then the Graber-Pandharipande virtual localization formula [GP] can be applied to the right hand side to show that the local invariants of Proposition 1.10 are equal to the global invariants (1.4). The same method works for the local surface case $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $d = 2$, $n = 3$, because then all stable pairs are scheme theoretically supported in the threefold $\text{Tot}_S(L_1)$.³ See Remark A.2 for more details.

We remark that most stable pair invariants of local surfaces calculated in this paper are small (see Section A.1). For $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$, the numbers are rather big:

$$P_{0,3[\ell]} = 11200, \quad P_{1,2[\ell]}([\ell]) = -820, \quad P_{1,3[\ell]}([\ell]) = -68060, \quad P_{2,2[\ell]}([\ell]) = 400,$$

where $[\ell] \in H_2(\mathbb{P}^3, \mathbb{Z}) \cong H_2(X, \mathbb{Z})$ denotes the class of a line $\ell \subseteq \mathbb{P}^3$ and we also write $[\ell] \in H^4(X, \mathbb{Z})$ for the pull-back of its Poincaré dual from \mathbb{P}^3 to X . This provides further good evidence for Conjectures 1.1 and 1.2.

³For the other local surfaces cases of Proposition 1.10, equating our invariants to the global invariants requires a more general virtual localization formula. Recently, Oh-Thomas announced such a formula [OT].

1.6. Acknowledgements. We warmly thank Pierrick Bousseau, Andrea Brini, and Michel van Garrel for providing their values of the genus zero Gopakumar-Vafa type invariants for the geometries listed in Corollary 1.8 and for useful discussions. This allowed us to use our calculations of stable pair invariants for verifying Conjecture 1.1 in more cases. We are grateful to Daveshe Maulik and Yukinobu Toda for various helpful discussions and communications. We also thank the anonymous referee for several helpful comments, which improved the exposition of the paper. Y.C. is partially supported by the World Premier International Research Center Initiative (WPI), MEXT, Japan, the JSPS KAKENHI Grant Number JP19K23397, and Royal Society Newton International Fellowships Alumni 2019. M.K. is supported by NWO grant VI.Vidi.192.012. S.M. is supported by NWO grant TOP2.17.004.

2. BACKGROUND

2.1. DT invariants of Calabi-Yau 4-folds. Let X be a smooth projective Calabi-Yau 4-fold with ample divisor ω and take a cohomology class $v \in H^*(X, \mathbb{Q})$. The coarse moduli space $M_\omega(v)$ of ω -Gieseker semistable sheaves E on X with $\text{ch}(E) = v$ is a projective scheme. We always assume that $M_\omega(v)$ is a fine moduli space, i.e. any point $[E] \in M_\omega(v)$ is stable and there exists a universal family \mathcal{E} on $X \times M_\omega(v)$ flat over $M_\omega(v)$. For instance, the moduli space of 1-dimensional stable sheaves E with $[E] = \beta$, $\chi(E) = 1$ and Hilbert schemes of closed subschemes satisfy this assumption [CK1, CK2, CMT1, CT2].

Borisov-Joyce [BJ] (in general) and Cao-Leung [CL1] (in special cases) constructed a virtual fundamental class

$$(2.1) \quad [M_\omega(v)]^{\text{vir}} \in H_{2-\chi(v,v)}(M_\omega(v), \mathbb{Z}),$$

where $\chi(\cdot, \cdot)$ denotes the Euler pairing. In order to construct the above virtual class (2.1) with coefficients in \mathbb{Z} (instead of \mathbb{Z}_2), we need an orientability result for $M_\omega(v)$, which can be stated as follows. Let

$$\mathcal{L} := \det(\mathbf{R}\mathcal{H}om_{\pi_M}(\mathcal{E}, \mathcal{E})) \in \text{Pic}(M_\omega(v)), \quad \pi_M: X \times M_\omega(v) \rightarrow M_\omega(v)$$

be the determinant line bundle of $M_\omega(v)$, which is equipped with the nondegenerate symmetric pairing Q induced by Serre duality. An *orientation* of (\mathcal{L}, Q) is a reduction of its structure group from $\text{O}(1, \mathbb{C})$ to $\text{SO}(1, \mathbb{C}) = \{1\}$. In other words, we require a choice of square root of the isomorphism

$$(2.2) \quad Q: \mathcal{L} \otimes \mathcal{L} \rightarrow \mathcal{O}_{M_\omega(v)}.$$

Existence of orientations was first proved when the Calabi-Yau 4-fold X satisfies $\text{Hol}(X) = \text{SU}(4)$ and $H^{\text{odd}}(X, \mathbb{Z}) = 0$ in [CL2], and was recently generalized to arbitrary Calabi-Yau 4-folds in [CGJ, Cor. 1.17]. Notice that the collection of orientations forms a torsor for $H^0(M_\omega(v), \mathbb{Z}_2)$. The virtual class (2.1) depends on the choice of orientation, but we suppress it from the notation.

Roughly speaking, in order to construct (2.1), one chooses at every point $[E] \in M_\omega(v)$, a half-dimensional real subspace

$$\text{Ext}_+^2(E, E) \subseteq \text{Ext}^2(E, E)$$

of the usual obstruction space $\text{Ext}^2(E, E)$, on which the quadratic form Q defined by Serre duality is real and positive definite. Then one glues local Kuranishi-type models of the form

$$\kappa_+ = \pi_+ \circ \kappa: \text{Ext}^1(E, E) \rightarrow \text{Ext}_+^2(E, E),$$

where κ is the Kuranishi map for $M_\omega(v)$ at $[E]$ and π_+ denotes projection on the first factor of the decomposition $\text{Ext}^2(E, E) = \text{Ext}_+^2(E, E) \oplus \sqrt{-1} \cdot \text{Ext}_+^2(E, E)$.

In [CL1], local models are glued in three special cases:

- (1) when $M_\omega(v)$ consists of locally free sheaves only,
- (2) when $M_\omega(v)$ is smooth,
- (3) when $M_\omega(v)$ is a shifted cotangent bundle of a quasi-smooth derived scheme.

In each case, the corresponding virtual classes are constructed using either gauge theory or algebro-geometric perfect obstruction theory.

The general gluing construction, due to Borisov-Joyce [BJ], is based on Pantev-Töen-Vaquié-Vezzosi's theory of shifted symplectic geometry [PTVV] and Joyce's theory of derived C^∞ -geometry. The corresponding virtual class is constructed using Joyce's D-manifold theory (a machinery similar to Fukaya-Oh-Ohta-Ono's theory of Kuranishi space structures used for defining Lagrangian Floer theory).

Examples computed in this paper only involve virtual class constructions in (2) and (3) mentioned above. We briefly review them:

- When $M_\omega(v)$ is smooth, the obstruction sheaf $\text{Ob} \rightarrow M_\omega(v)$ is a vector bundle endowed with a quadratic form Q via Serre duality. Then the virtual class is given by

$$[M_\omega(v)]^{\text{vir}} = \text{PD}(e(\text{Ob}, Q)),$$

where $\text{PD}(\cdot)$ denotes Poincaré dual and $e(\text{Ob}, Q)$ is the half-Euler class of (Ob, Q) , i.e. the Euler class of its real form Ob_+ . In this case, a choice of orientation (2.2) is equivalent to a choice of orientation of Ob_+ . The half-Euler class satisfies

$$\begin{aligned} e(\text{Ob}, Q)^2 &= (-1)^{\frac{\text{rk}(\text{Ob})}{2}} e(\text{Ob}), & \text{if } \text{rk}(\text{Ob}) \text{ is even,} \\ e(\text{Ob}, Q) &= 0, & \text{if } \text{rk}(\text{Ob}) \text{ is odd.} \end{aligned}$$

- Suppose $M_\omega(v)$ is the classical truncation of the shifted cotangent bundle of a quasi-smooth derived scheme. Roughly speaking, this means that at any closed point $[E] \in M_\omega(v)$, we have a Kuranishi map of the form

$$\kappa: \text{Ext}^1(E, E) \rightarrow \text{Ext}^2(E, E) = V_E \oplus V_E^*,$$

where κ factors through a maximal isotropic subspace V_E of $(\text{Ext}^2(E, E), Q)$. Then the virtual class of $M_\omega(v)$ is essentially the virtual class of the perfect obstruction theory formed by $\{V_E\}_{[E] \in M_\omega(v)}$. When $M_\omega(v)$ is furthermore smooth as a scheme, then it is simply the Euler class of the vector bundle $\{V_E\}_{[E] \in M_\omega(v)}$ over $M_\omega(v)$.

2.2. Stable pair invariants of Calabi-Yau 4-folds. As in [PT1], a stable pair (F, s) on a smooth projective Calabi-Yau 4-fold X consists of

- a pure dimension 1 sheaf F on X ,
- a section $s \in H^0(X, F)$ with 0-dimensional or trivial cokernel.

For $\beta \in H_2(X, \mathbb{Z})$ and $n \in \mathbb{Z}$, denote by $P_n(X, \beta)$ be the moduli space of stable pairs (F, s) on X such that F has scheme theoretic support with class β and $\chi(F) = n$. By [PT1], it can alternatively be seen as the moduli space parametrizing 2-term complexes

$$I^\bullet = \{\mathcal{O}_X \xrightarrow{s} F\} \in D^b(\text{Coh}(X))$$

in the bounded derived category of coherent sheaves on X . This viewpoint produces an obstruction theory on $P_n(X, \beta)$, which is however not perfect because $\text{Ext}^2(I^\bullet, I^\bullet)_0$ is in general non-vanishing. Nonetheless, using the methods of Borisov-Joyce [BJ], one can construct a virtual class (see [CMT2, Thm. 1.4])

$$[P_n(X, \beta)]^{\text{vir}} \in H_{2n}(P_n(X, \beta), \mathbb{Z})$$

depending on a choice of orientation. Existence of orientations was proved in [CGJ, Cor. 1.17].

3. MODULI SPACES

3.1. Compactness I. In the previous section, we assumed X is a smooth *projective* Calabi-Yau 4-fold. As we will discuss in more detail in Section 4.2, the previous section also applies to certain cases where X is a smooth quasi-projective Calabi-Yau 4-fold and $P_n(X, \beta)$ is proper.

Suppose S is a smooth projective surface and $L_1, L_2 \in \text{Pic}(S)$ satisfy

$$L_1 \otimes L_2 \cong K_S.$$

Then $X = \text{Tot}_S(L_1 \oplus L_2)$ is a smooth quasi-projective Calabi-Yau 4-fold, which we refer to as a *local surface*. One way to ensure the properness of $P_n(X, \beta)$ is as follows.

Proposition 3.1. *Suppose S is a smooth projective surface with $L_1, L_2 \in \text{Pic}(S)$ satisfying $L_1 \otimes L_2 \cong K_S$ and let $X = \text{Tot}_S(L_1 \oplus L_2)$. Let $\beta \in H_2(S, \mathbb{Z})$ and suppose for any $0 \neq \beta' \leq \beta$,⁴ we have $\beta' \cdot L_i < 0$ for $i = 1, 2$. Then $P_n(X, \beta)$ is projective for any $n \in \mathbb{Z}$.*

Proof. Let $[(F, s)] \in P_n(X, \beta)$. We first show that F is set theoretically supported on the zero section $S \subseteq X$. Let D be an irreducible component of the scheme theoretic support of F , then we want to show $D_{\text{red}} \subseteq S$. Let $Y = \text{Tot}_S(L_1)$ and consider the projection $p: X = \text{Tot}_Y(L_2) \rightarrow Y$ (here and below, we suppress the pull-back of L_2 along the projection $Y \rightarrow S$). Since D_{red} is a proper irreducible reduced curve, $\mathcal{O}_{D_{\text{red}}}$ is stable. By the spectral construction, it corresponds to a stable Higgs pair $(p_*\mathcal{O}_{D_{\text{red}}}, \phi)$, where

$$\phi: p_*\mathcal{O}_{D_{\text{red}}} \rightarrow p_*\mathcal{O}_{D_{\text{red}}} \otimes L_2.$$

Denote the curve class of the scheme theoretic support of $p_*\mathcal{O}_{D_{\text{red}}}$ by $\beta' \in H_2(Y, \mathbb{Z}) \cong H_2(S, \mathbb{Z})$. Then $0 \neq \beta' \leq \beta$, so $\beta' \cdot L_2 < 0$. Combined with stability of the Higgs pairs $(p_*\mathcal{O}_{D_{\text{red}}}, \phi)$ and

⁴The notation $\beta' \leq \beta$ means that there exist effective curve classes $\beta', \beta'' \in H_2(S, \mathbb{Z})$ such that $\beta = \beta' + \beta''$.

$(p_*\mathcal{O}_{D_{\text{red}}} \otimes L_2, \phi \otimes \text{id}_{L_2})$, this implies $\phi = 0$ so $D_{\text{red}} \subseteq Y = \text{Tot}_S(L_1)$ (see [TT, Prop. 7.4] for a similar argument). Reversing the roles of L_1, L_2 , we deduce $D_{\text{red}} \subseteq Y = \text{Tot}_S(L_2)$, so $D_{\text{red}} \subseteq S$.

Since each element of $P_n(X, \beta)$ is set theoretically supported on S , we conclude that $P_n(X, \beta)$ is projective. Indeed, there is a $d \gg 0$ such that every element of $P_n(X, \beta)$ is scheme theoretically supported in dS , where dS denotes the d times thickening of the zero section $S \subseteq X$, i.e. the closed subscheme of X defined by $I^d \subseteq \mathcal{O}_X$, where $I \subseteq \mathcal{O}_X$ denotes the ideal of the zero section. Therefore $P_n(X, \beta) \cong P_n(dS, \beta)$, \square

Suppose L_1^{-1} and L_2^{-1} are ample. Then K_S^{-1} is ample, i.e. S is del Pezzo, and $P_n(X, \beta)$ is projective for all β, n by Proposition 3.1. As noted in [CMT2, Sect. 4.2], there are only two possibilities:

Proposition 3.2. *Let S be a smooth projective surface and $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$. Suppose L_1^{-1} and L_2^{-1} are ample. Then, up to permutating L_1, L_2 , we only have $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$ or $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$.*

Proof. Suppose S contains a (-1) -curve C . Then the Nakai criterion and adjunction imply

$$-2 \geq \deg(L_1|_C) + \deg(L_2|_C) = \deg(K_S|_C) = -1,$$

so S does not contain (-1) -curves. The classification of del Pezzo surfaces yields the result. \square

For both geometries of this proposition, the Gromov-Witten (and hence Gopakumar-Vafa type) invariants were determined in [KP, Sect. 3].

Let us go back to an arbitrary smooth projective surface S with $L_1, L_2 \in \text{Pic}(S)$ satisfying $L_1 \otimes L_2 \cong K_S$. Consider the moduli space $P_n(S, \beta)$ of stable pairs (F, s) on S with $\chi(F) = n$ and scheme theoretic support of F in class $\beta \in H_2(S, \mathbb{Z})$. Any stable pair $I^\bullet = \{\mathcal{O}_S \rightarrow F\}$ gives rise to a stable pair

$$\{\mathcal{O}_X \rightarrow \iota_*\mathcal{O}_S \rightarrow \iota_*F\}$$

on $X = \text{Tot}_S(L_1 \oplus L_2)$, where $\iota : S \hookrightarrow X$ denotes inclusion of the zero section. This gives a closed embedding

$$(3.1) \quad P_n(S, \beta) \hookrightarrow P_n(X, \beta).$$

We refer to elements of $P_n(X, \beta)$ in the image as “stable pairs which are scheme theoretically supported on S ”. Requiring $P_n(X, \beta)$ to be proper poses restrictions on n, β . The following result is very useful for finding “candidates” for proper moduli spaces $P_n(X, \beta)$ (as we will see later in this section in Proposition 3.8).

Proposition 3.3. *Let S be a smooth projective surface, $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$ and let $X = \text{Tot}_S(L_1 \oplus L_2)$. Let $\beta \in H_2(S, \mathbb{Z})$ and $n \in \mathbb{Z}$ such that $P_n(X, \beta)$ is proper and $P_n(S, \beta) \neq \emptyset$. Suppose $C_1, C_2 \subseteq S$ are effective divisors satisfying*

- $C_1 \cong \mathbb{P}^1$ and $[C_1 + C_2] = \beta$,
- $L_i \cdot C_1 = 0$ for $i = 1$ or $i = 2$.

Then

$$-\frac{1}{2}\beta(\beta + K_S) \leq n \leq -\frac{1}{2}C_2(C_2 + K_S).$$

Proof. Suppose $P_n(S, \beta) \neq \emptyset$, $P_n(X, \beta)$ is proper, and let $C_1, C_2 \subseteq S$ be as stated. Then for any element $[(F, s)] \in P_n(S, \beta)$ with underlying scheme theoretic support C , we have

$$n = \chi(F) \geq \chi(\mathcal{O}_C) = -\frac{1}{2}\beta(\beta + K_S).$$

Suppose

$$n \geq 1 - \frac{1}{2}C_2(C_2 + K_S).$$

Since $C_1 \cong \mathbb{P}^1$ and $\deg(L_i|_{C_1}) = L_i \cdot C_1 = 0$, for $i = 1$ or $i = 2$, the line bundle $L_i|_{C_1}$ is trivial. Hence we can take a nowhere vanishing section D_1 of the line bundle $L_i|_{C_1} \cong \mathbb{P}^1 \times \mathbb{C}$. In particular, D_1 and C_2 are disjoint. Therefore

$$\begin{aligned} \chi(\mathcal{O}_{D_1 \sqcup C_2}) &= \chi(\mathcal{O}_{D_1}) + \chi(\mathcal{O}_{C_2}) \\ &= 1 - \frac{1}{2}C_2(C_2 + K_S). \end{aligned}$$

Twisting $\mathcal{O}_{D_1 \sqcup C_2}$ by an effective divisor of appropriate length, we obtain a stable pair $[(F, s)] \in P_n(X, \beta) \setminus P_n(S, \beta)$ with underlying scheme theoretic support $D_1 \sqcup C_2$. Since D_1 does not lie in the zero-section, using the \mathbb{C}^* -scaling action on L_i , we get a family of stable pairs with part of the support (i.e. D_1) moving off to infinity, contradicting properness of $P_n(X, \beta)$. \square

We want to apply this proposition to smooth projective surfaces S with $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$ and L_1^{-1}, L_2^{-1} non-trivial and nef. These surfaces were recently studied in the context of the log-local principle by Bousseau-Brini-van Garrel [BBG]. In particular, they determined the genus zero Gromov-Witten (and hence Gopakumar-Vafa type) invariants of $\text{Tot}_S(L_1 \oplus L_2)$ in many new cases.

Smooth projective surfaces S with K_S^{-1} nef and big are called weak del Pezzo surfaces. The weak toric del Pezzo surfaces are: \mathbb{P}^2 , $\mathbb{P}^1 \times \mathbb{P}^1$, \mathbb{F}_1 , \mathbb{F}_2 , or certain repeated toric blow-ups of \mathbb{P}^2 in at most 6 points as specified in [Sat]. In this paper, we only consider the *minimal* cases, i.e. the first four cases. Using the notation for Hirzebruch surfaces from the introduction, the only possibilities for $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$ with L_1^{-1}, L_2^{-1} non-trivial and nef are (up to permutations of L_1, L_2):

- $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$,
- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$ or $(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$,
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$ or $(\mathbb{F}_1, \mathcal{O}(0, -1), \mathcal{O}(-2, -2))$,
- $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$.

Example 3.4. Suppose $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$. Let $H_1 = \{\text{pt}\} \times \mathbb{P}^1$ and $H_2 = \mathbb{P}^1 \times \{\text{pt}\}$. We define $(d_1, d_2) := d_1 H_1 + d_2 H_2$. For all $d_1, d_2 \in \mathbb{Z}$, (d_1, d_2) is effective if and only if $d_1, d_2 \geq 0$. For $(0, d)$ with $d \geq 1$, the moduli space $P_n(X, (0, d))$ is projective for all n by Proposition 3.1. Now suppose $d_1 > 0$, $d_2 \geq 0$, and $n \geq 0$. Let $C_1 \in |H_1|$ and $C_2 \in |(d_1 - 1)H_1 + d_2 H_2|$. Then $L_1 \cdot C_1 = 0$ and the inequalities of Proposition 3.3 reduce to

$$d_1 + d_2 - d_1 d_2 \leq n \leq d_1 + 2d_2 - d_1 d_2 - 1.$$

These inequalities have the following solutions:

- $(d_1, d_2) = (3, 2), (2, d)$ with $d \geq 2$ and $n = 0$,
- $(d_1, d_2) = (d, 1), (1, d)$ with $d \geq 1$ and $n = 1$, or $(d_1, d_2) = (2, d)$ with $d \geq 1$ and $n = 1$,
- $(1, d)$ with $2 \leq n \leq d$.

Example 3.5. Suppose $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$ and use the notation for Hirzebruch surfaces from the introduction, so $(d_1, d_2) := d_1 B + d_2 F$ for all $d_1, d_2 \in \mathbb{Z}$. Then (d_1, d_2) is effective if and only if $d_1, d_2 \geq 0$ (this holds for all Hirzebruch surfaces). For $(0, d)$ with $d \geq 1$, the moduli space $P_n(X, (0, d))$ is projective for all n by Proposition 3.1. Suppose $d_1 > 0$, $d_2 \geq 0$, and $n \geq 0$. Let $C_1 \in |B|$ and $C_2 \in |(d_1 - 1)B + d_2 F|$. Then $L_1 \cdot C_1 = 0$ and the inequalities of Proposition 3.3 reduce to

$$\frac{1}{2}d_1(d_1 + 1) - d_2(d_1 - 1) \leq n \leq \frac{1}{2}d_1(d_1 - 1) - d_2(d_1 - 2).$$

These inequalities have the following solutions:

- $(d_1, d_2) = (3, 3), (2, d)$ with $d \geq 3$ and $n = 0$,
- $(d_1, d_2) = (2, d)$ with $d \geq 2$ and $n = 1$,
- $(1, d)$ with $1 \leq n \leq d$.

Example 3.6. Suppose $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(0, -1), \mathcal{O}(-2, -2))$. Suppose $d_1 > 0$, $d_2 \geq 0$, and $n \geq 0$. Taking C_1, C_2 as in Example 3.5 leads to the same list. Additionally, we can take $d_1 \geq 0$, $d_2 > 0$, $n \geq 0$, $C_1 \in |F|$ and $C_2 \in |d_1 B + (d_2 - 1)F|$. Then $L_1 \cdot C_1 = 0$ and the inequalities of Proposition 3.3 reduce to

$$\frac{1}{2}d_1(d_1 + 1) - d_2(d_1 - 1) \leq n \leq \frac{1}{2}d_1(d_1 + 1) - (d_2 - 1)(d_1 - 1).$$

The solutions to these inequalities *and* the ones from Example 3.5 are:

- $(d_1, d_2) = (2, 3), (2, 4), (3, 3)$ and $n = 0$,
- $(d_1, d_2) = (2, 2), (2, 3), (1, d)$ with $d \geq 1$ and $n = 1$.

Example 3.7. Suppose $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$. For $(0, d)$ with $d \geq 1$, the moduli space $P_n(X, (0, d))$ is projective for all n by Proposition 3.1. Suppose $d_1 > 0$, $d_2 \geq 0$, and $n \geq 0$. Let $C_1 \in |B|$ and $C_2 \in |(d_1 - 1)B + d_2 F|$. Then $L_1 \cdot C_1 = 0$ and the inequalities of Proposition 3.3 reduce to

$$d_1^2 - d_2(d_1 - 1) \leq n \leq (d_1 - 1)^2 - d_2(d_1 - 2).$$

These inequalities have the following solutions:

- $(d_1, d_2) = (2, d)$, $d \geq 4$, and $n = 0$,
- $(d_1, d_2) = (2, d)$, $d \geq 3$, and $n = 1$,
- $(1, d)$ with $1 \leq n \leq d$.

In these examples we listed, for given (S, L_1, L_2) , all the cases for which $n \geq 0$ and *potentially* $P_n(X, \beta)$ is proper and $P_n(S, \beta) \neq \emptyset$ (Proposition 3.3). For $\beta = (0, d)$ with $d \geq 1$, $P_n(S, \beta) \neq \emptyset$ if and only if $n \geq d$, and $P_n(X, \beta)$ is proper by Proposition 3.1. For all other cases listed, $P_n(S, \beta)$ is also non-empty since $|\beta| \neq \emptyset$ and $n \geq \chi(\mathcal{O}_C)$ for any $C \in |\beta|$. Indeed adding sufficiently many points to C one obtains a stable pair (F, s) on S with $\chi(F) = n$. We now prove that in each of the cases listed, $P_n(X, \beta)$ is indeed proper.

Proposition 3.8. *In each of the cases listed in Examples 3.4–3.7, $P_n(X, \beta)$ is projective.*

Proof. We write out the proof for Example 3.4. The other cases are analogous. Recall that $H_1 := \{\text{pt}\} \times \mathbb{P}^1$, $H_2 := \mathbb{P}^1 \times \{\text{pt}\}$, $L_1 := \mathcal{O}(-H_1)$, $L_2 := \mathcal{O}(-H_1 - 2H_2)$, and $\beta := d_1 H_1 + d_2 H_2$. Suppose n, β are as listed in Example 3.4. As in Proposition 3.1, it is enough to show that all elements of $P_n(X, \beta)$ are set theoretically supported on S . Suppose $[(F, s)] \in P_n(X, \beta)$ has scheme theoretic support C and let D be an irreducible component of C which is *not* set theoretically supported on S . Then we claim D_{red} is a proper irreducible reduced curve with class $[D_{\text{red}}] \in H_2(X, \mathbb{Z}) \cong H_2(S, \mathbb{Z})$ satisfying $[D_{\text{red}}] \cdot L_1 \geq 0$ or $[D_{\text{red}}] \cdot L_2 \geq 0$. Indeed suppose $[D_{\text{red}}] \cdot L_1 < 0$ and $[D_{\text{red}}] \cdot L_2 < 0$. Using the spectral construction as in the proof of Proposition 3.1, stability of $\mathcal{O}_{D_{\text{red}}}$, then implies $D_{\text{red}} \subseteq S$ contrary to our assumption.

The only non-zero effective curve classes β' on S such that $\beta' \cdot L_1 \geq 0$ or $\beta' \cdot L_2 \geq 0$ are $\beta' = mH_1$ for some $m > 0$. Hence there exists a $\Sigma \in |H_1|$ such that

$$p|_{D_{\text{red}}} : D_{\text{red}} \rightarrow \Sigma \subseteq S,$$

where $p : X \rightarrow S$ denotes the projection. Note that $L_1|_{\Sigma} \cong \mathcal{O}$ and $L_2|_{\Sigma} \cong \mathcal{O}(-2)$. Since $[D_{\text{red}}] \cdot L_2 < 0$, a similar argument as above shows that $D_{\text{red}} \subseteq \text{Tot}_{\Sigma}(L_1) \cong \mathbb{P}^1 \times \mathbb{C}$. Therefore D_{red} is a non-zero section of $\text{Tot}_{\Sigma}(L_1) \cong \mathbb{P}^1 \times \mathbb{C}$.

Denote the irreducible components of C which are *not* set theoretically supported on S by D_1, \dots, D_{ℓ} and let D' be the union of the remaining components. Above, we showed each $D_{i,\text{red}} \cong \mathbb{P}^1$ and $D_{i,\text{red}}$ is a non-zero section of $\text{Tot}_{\Sigma_i}(L_1) \cong \mathbb{P}^1 \times \mathbb{C}$ for some $\Sigma_i \in |H_1|$. It follows that $D_{1,\text{red}}, \dots, D_{\ell,\text{red}}, D'_{\text{red}}$ are mutually disjoint. Denote the multiplicity of D_i at $D_{i,\text{red}}$ by $\delta_i \geq 1$. Consider the classes $p_*[D_i], p_*[D'] \in H_2(S, \mathbb{Z})$, where $p : X \rightarrow S$ is the projection. Then

$$p_*[D_i] := \delta_i H_1, \quad p_*[D'] := \beta - \delta H_1,$$

where $\delta := \sum_{i=1}^{\ell} \delta_i$. We claim

$$(3.2) \quad \chi(\mathcal{O}_{D_i}) \geq 1, \quad \chi(\mathcal{O}_{D'}) \geq 1 - g(p_*[D']) = -\frac{1}{2}(\beta - \delta H_1)(\beta - \delta H_1 + K_S),$$

for all $i = 1, \dots, \ell$, where the last equality is by the Riemann-Roch formula. When D_i (resp. D') are reduced, these inequalities are equalities. In general, since $N_{D_{i,\text{red}}/X} \cong \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-2)$ and $N_{S/X} \cong L_1 \oplus L_2$ with L_1^{-1}, L_2^{-1} nef, we have inequalities as stated⁵. From (3.2) and the fact that D_1, \dots, D_{ℓ}, D' are mutually disjoint, we deduce

$$\begin{aligned} n = \chi(F) &\geq \chi(\mathcal{O}_C) = \sum_{i=1}^{\ell} \chi(\mathcal{O}_{D_i}) + \chi(\mathcal{O}_{D'}) \\ &\geq \delta - \frac{1}{2}(\beta - \delta H_1)(\beta - \delta H_1 + K_S) \\ &= \delta - \frac{1}{2}(d_1 - \delta)(d_2 - 2) - \frac{1}{2}d_2(d_1 - \delta - 2). \end{aligned}$$

However, for each of the cases listed in Example 3.4, it is easy to see that $n \leq \delta - 1 - \frac{1}{2}(d_1 - \delta)(d_2 - 2) - \frac{1}{2}d_2(d_1 - \delta - 2)$ for all $1 \leq \delta \leq d_1$ by explicit calculation. We have reached a contradiction. \square

Conclusion. For any (S, L_1, L_2) with $L_1 \otimes L_2 \cong K_S$, L_1^{-1}, L_2^{-1} non-trivial and nef, S minimal and toric, we classified *all* $n \geq 0$, $\beta \in H_2(S, \mathbb{Z})$ such that $P_n(X, \beta)$ is proper and $P_n(S, \beta) \neq \emptyset$.

3.2. Compactness II. In the previous section, we studied properness of $P_n(X, \beta)$ for local surfaces. In particular, for $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$ or $(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, the moduli space $P_n(X, \beta)$ is always proper (Proposition 3.1). We are now interested in the cases where $P_n(X, \beta) \cong P_n(S, \beta)$, i.e. the embedding (3.1) is an isomorphism. For the 3-fold $\text{Tot}(K_{\mathbb{P}^2})$, this question was considered by Choi-Katz-Klemm in [CKK, Prop. 2]. In the proof of the following proposition, we use some of their techniques (adapted to the 4-fold setting).

⁵One way to see this is by using filtrations by thickenings of $D_{i,\text{red}} \subseteq X$ and $S \subseteq X$ as in the proof of Proposition 3.9 below.

Proposition 3.9. *Let $X = \text{Tot}_{\mathbb{P}^2}(\mathcal{O}(-1) \oplus \mathcal{O}(-2))$, $\beta = d[H]$ with $d \geq 1$, and $n \geq 0$. Then*

$$P_n(X, \beta) \cong P_n(\mathbb{P}^2, \beta)$$

if and only if

- (1) $d = 1$ and any $n \geq 0$, or
- (2) $d = 2, 3, 4$ and $n = 0, 1$, or
- (3) $d = 2, 3$ and $n = 2$.

Let $X = \text{Tot}_{\mathbb{P}^1 \times \mathbb{P}^1}(\mathcal{O}(-1, -1) \oplus \mathcal{O}(-1, -1))$, $\beta = d_1[H_1] + d_2[H_2] \neq 0$ with $d_1, d_2 \geq 0$, and $n \geq 0$. Then

$$P_n(X, \beta) \cong P_n(\mathbb{P}^1 \times \mathbb{P}^1, \beta),$$

if and only if

- (1) $(d_1, d_2) = (1, 0), (0, 1), (1, 1)$ and any $n \geq 0$, or
- (2) $(d_1, d_2) = (0, d), (d, 0)$ with $d \geq 2$ and $0 \leq n \leq d$, or
- (3) $(d_1, d_2) = (1, d), (d, 1)$ with $d \geq 2$ and $n = 0, 1, 2$, or
- (4) $(d_1, d_2) = (2, 2), (2, 3), (3, 2), (2, 4), (4, 2), (3, 3)$ and $n = 0$, or
- (5) $(d_1, d_2) = (2, 2), (2, 3), (3, 2)$ and $n = 1$, or
- (6) $(d_1, d_2) = (2, 2)$ and $n = 2$.

Proof. Let $[(F, s)] \in P_n(X, \beta)$ be a stable pair with scheme theoretic support $C := \text{supp}(F)$. The stable pair (F, s) is set theoretically supported on the zero section $S \subseteq X$ by Proposition 3.1. Let $Y_i = \text{Tot}_S(L_i)$ for $i = 1, 2$. We consider the ideals of $C \subseteq X$ and $Y_1 \subseteq X$:

$$J := I_{C \subseteq X} \subseteq \mathcal{O}_X, \quad I_2 := I_{Y_1 \subseteq X} \subseteq \mathcal{O}_X.$$

Note that I_2 is a line bundle on X . Since (F, s) is set theoretically supported on $S \subseteq X$ (and therefore $Y_1 \subseteq X$), there exists an $\ell \geq 0$ such that $J + I_2^{\ell+1} = J$ and we have

$$(3.3) \quad \chi(\mathcal{O}_C) = \sum_{j=0}^{\ell} \chi\left(\frac{J + I_2^j}{J + I_2^{j+1}}\right).$$

For each j , we have a surjective map

$$p^* L_2^{-j} \cong \frac{I_2^j}{I_2^{j+1}} \rightarrow \frac{J + I_2^j}{J + I_2^{j+1}},$$

where $p : Y_1 \rightarrow S$ denotes projection. Hence $\frac{J + I_2^j}{J + I_2^{j+1}} \cong \mathcal{O}_{C_j} \otimes p^* L_2^{-j}$ for some closed subscheme $C_j \subseteq Y_1$ of dimension ≤ 1 . Moreover, we have $C_j \supseteq C_{j+1}$ for all j .⁶ From the fact that C is Cohen-Macaulay, it also follows that, when non-empty, C_j is not 0-dimensional.

For a fixed j , we consider the ideals of $C_j \subseteq Y_1$ and $S \subseteq Y_1$:

$$J_j := I_{C_j \subseteq Y_1} \subseteq \mathcal{O}_{Y_1}, \quad I_1 := I_{S \subseteq Y_1} \subseteq \mathcal{O}_{Y_1}.$$

Note that I_1 is a line bundle on Y_1 . As above, there exists an $\ell_j \geq 0$ such that $J_j + I_1^{\ell_j+1} = J_j$ and we have

$$\chi(\mathcal{O}_{C_j}) = \sum_{i=0}^{\ell_j} \chi\left(\frac{J_j + I_1^i}{J_j + I_1^{i+1}}\right).$$

As above, for all i , we have $\frac{J_j + I_1^i}{J_j + I_1^{i+1}} \cong \mathcal{O}_{C_{ij}} \otimes L_1^{-i}$ for some closed subscheme $C_{ij} \subseteq S$ of dimension ≤ 1 . As above, we also have $C_{ij} \supseteq C_{i+1,j}$ for all i . This time, we leave open the possibility that C_{ij} is 0-dimensional, because C_j need not be Cohen-Macaulay. Nonetheless, denoting $\beta_{ij} := [C_{ij}]$, we have

$$\beta = \sum_{j=0}^{\ell} \sum_{i=0}^{\ell_j} \beta_{ij} \in H_2(S, \mathbb{Z}).$$

Consider the torsion filtration

$$0 \rightarrow \mathcal{T}_0 \rightarrow \mathcal{O}_{C_{ij}} \rightarrow \mathcal{O}_{C_{ij}}^{\text{pure}} \rightarrow 0$$

and the exact sequence

$$0 \rightarrow \mathcal{O}_S(-C_{ij}^{\text{pure}}) \rightarrow \mathcal{O}_S \rightarrow \mathcal{O}_{C_{ij}}^{\text{pure}} \rightarrow 0.$$

⁶This follows from the natural surjection $\mathcal{O}_{C_j} \otimes p^* L_2^{-j-1} \cong \frac{J + I_2^j}{J + I_2^{j+1}} \otimes \frac{I_2}{I_2^2} \rightarrow \frac{J + I_2^{j+1}}{J + I_2^{j+2}} \cong \mathcal{O}_{C_{j+1}} \otimes p^* L_2^{-j-1}$.

The support of \mathcal{T}_0 is 0-dimensional. Applying the Hirzebruch-Riemann-Roch formula gives

$$\begin{aligned} \chi\left(\frac{J_j + I_1^i}{J_j + I_1^{i+1}} \otimes p^* L_2^{-j}\right) &= \chi(\mathcal{O}_{C_{ij}} \otimes L_1^{-i} \otimes L_2^{-j}) \geq \chi(\mathcal{O}_{C_{ij}^{\text{pure}}} \otimes L_1^{-i} \otimes L_2^{-j}) \\ &= -\frac{1}{2}\beta_{ij}(\beta_{ij} + K_S) - (iL_1 + jL_2)\beta_{ij}. \end{aligned}$$

Combining with (3.3), we obtain

$$\begin{aligned} \chi(F) &\geq \chi(\mathcal{O}_C) \geq -\sum_{j=0}^{\ell} \sum_{i=0}^{\ell_j} \left(\frac{1}{2}\beta_{ij}(\beta_{ij} + K_S) + (iL_1 + jL_2)\beta_{ij}\right) \\ (3.4) \quad &\geq -\frac{1}{2}\beta(\beta + K_S) - \beta(L_1 + L_2) \\ &\quad + \frac{1}{2} \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} \beta_{ij}\beta_{i'j'} + \beta_{00}(L_1 + L_2) + L_1 \sum_{j=1}^{\ell} \beta_{0j} + L_2 \sum_{i=1}^{\ell_0} \beta_{i0}, \end{aligned}$$

where we used that L_1 and L_2 are nef line bundles.

Case 1. Let $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $\beta = d[H]$ with $d \geq 1$, and $n \geq 0$. Suppose there exists an element $[(F, s)] \in P_n(X, \beta) \setminus P_n(S, \beta)$. We use the notation above for its scheme theoretic support C and the associated schemes C_{ij} . Let $\beta_{ij} = d_{ij}[H]$, then (3.4) gives

$$\begin{aligned} \chi(F) &\geq -\frac{1}{2}d^2 + \frac{9}{2}d - 3d_{00} - \sum_{j=1}^{\ell} d_{0j} - 2 \sum_{i=1}^{\ell_0} d_{i0} + \frac{1}{2} \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} d_{ij}d_{i'j'} \\ &\geq -\frac{1}{2}d^2 + \frac{5}{2}d - d_{00} + \frac{1}{2} \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} d_{ij}d_{i'j'} \\ &\geq -\frac{1}{2}d^2 + \frac{5}{2}d, \end{aligned}$$

where the last inequality uses that there exists an $(i, j) \neq (0, 0)$ with $d_{ij} \geq 1$, because we assumed C is not scheme theoretically supported in the zero section $S \subseteq X$. Hence $n \leq -\frac{1}{2}d^2 + \frac{5}{2}d - 1$ implies $P_n(S, \beta) \cong P_n(X, \beta)$. In particular, we find that for the cases (1)–(3) we have $P_n(S, \beta) \cong P_n(X, \beta)$. For case (1) this is obvious from Proposition 3.1 and the fact that β is irreducible.

For β, n other than (1)–(3), it is easy to construct a $(\mathbb{C}^*)^4$ -fixed stable pair in

$$P_n(X, \beta) \setminus P_n(S, \beta)$$

using the combinatorial description of stable pairs in [PT2, CK2], where $(\mathbb{C}^*)^4$ denotes the torus of the toric Calabi-Yau 4-fold X .

Case 2. Let $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $\beta = d_1[H_1] + d_2[H_2]$ for some $d_1, d_2 \geq 0$ not both zero, and $n \geq 0$. Suppose there exists an element $[(F, s)] \in P_n(X, \beta) \setminus P_n(S, \beta)$. We use the notation above for its scheme theoretic support C and the associated schemes C_{ij} . Let $\beta_{ij} = d_{1,ij}[H_1] + d_{2,ij}[H_2]$, then (3.4) gives

$$\begin{aligned} \chi(F) &\geq 3d_1 + 3d_2 - d_1d_2 - 2d_{1,00} - 2d_{2,00} - \sum_{j=1}^{\ell} (d_{1,0j} + d_{2,0j}) - \sum_{i=1}^{\ell_0} (d_{1,i0} + d_{2,i0}) \\ (3.5) \quad &\quad + \frac{1}{2} \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} (d_{1,ij}d_{2,i'j'} + d_{1,i'j'}d_{2,ij}) \\ &\geq 2d_1 + 2d_2 - d_1d_2 - d_{1,00} - d_{2,00} + \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} d_{1,ij}d_{2,i'j'} \\ &\geq d_1 + d_2 - d_1d_2 + 1 + \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} d_{1,ij}d_{2,i'j'}, \end{aligned}$$

where the last inequality uses that C does not lie scheme theoretically in S . Suppose $d_1, d_2 \geq 2$, then

$$(3.6) \quad \sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} d_{1,ij}d_{2,i'j'} \geq 2.$$

Therefore $n \leq d_1 + d_2 - d_1 d_2 + 2$ implies $P_n(S, \beta) \cong P_n(X, \beta)$. In particular, for β, n as in (4)–(6), *except* for $(d_1, d_2) = (3, 3)$ and $n = 0$ (!), we deduce that $P_n(S, \beta) \cong P_n(X, \beta)$. Cases (1)–(3) can be found from (3.5) by a similar reasoning.

For $(d_1, d_2) = (3, 3)$, we still use (3.5), but we need to sharpen (3.6). Recall that the schemes C_j and C_{ij} constructed in the first part of the proof are nested. This implies $d_{1,ij} \geq d_{1,i+1j}$ and $d_{2,ij} \geq d_{2,i+1j}$ for all i, j . Using these inequalities for $(d_1, d_2) = (3, 3)$, one can show that

$$\sum_{\substack{((i,j),(i',j')) \\ (i,j) \neq (i',j')}} d_{1,ij} d_{2,i'j'} \geq 3.$$

It follows that for $[(F, s)] \in P_n(X, (3, 3)) \setminus P_n(S, (3, 3))$, we have $\chi(F) \geq 1$. Hence $P_0(S, (3, 3)) \cong P_0(X, (3, 3))$.

For β, n other than (1)–(6), it is easy to construct a $(\mathbb{C}^*)^4$ -fixed stable pair in $P_n(X, \beta) \setminus P_n(S, \beta)$ using the combinatorial description of stable pairs in [PT2, CK2]. \square

Remark 3.10. For (S, L_1, L_2) as in Examples 3.4–3.7, we found *all* cases for which $n \geq 0$, $P_n(X, \beta)$ is proper, and $P_n(S, \beta) \neq \emptyset$ (Propositions 3.3 and 3.8). A similar reasoning as in the proof of Proposition 3.9 (using (3.4)) can be applied to find out when $P_n(X, \beta) \cong P_n(S, \beta)$. In the following cases, we have $n \geq 0$, $P_n(X, \beta) \cong P_n(S, \beta)$:

- $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 1)$ and any $n \geq 0$, or $(d_1, d_2) = (0, d)$ for any $d \geq 2$ and $n = d$, or $(d_1, d_2) = (2, 2), (2, 3), (3, 2), (2, 4)$ and $n = 0$, or $(d_1, d_2) = (2, 2), (2, 3), (1, d), (d, 1)$ for any $d \geq 1$ and $n = 1$, or $(d_1, d_2) = (1, d)$ for any $d \geq 2$ and $n = 2$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 1)$ and any $n \geq 0$, or $(d_1, d_2) = (0, d)$ for any $d \geq 2$ and $n = d$, or $(d_1, d_2) = (2, 3), (2, 4), (3, 3), (2, 5)$ and $n = 0$, or $(d_1, d_2) = (2, 2), (2, 3), (2, 4)$ and $n = 1$, or $(d_1, d_2) = (1, d)$ for $n = 1, 2$ and any $d \geq n$.
- $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(0, -1), \mathcal{O}(-2, -2))$, $(d_1, d_2) = (2, 3), (2, 4), (3, 3)$ and $n = 0$, or $(d_1, d_2) = (2, 2), (2, 3), (1, d)$ with $d \geq 1$ and $n = 1$.
- $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$, $(d_1, d_2) = (0, 1)$ and any $n \geq 0$, or $(d_1, d_2) = (0, d)$ for any $d \geq 2$ and $n = d$, or $(d_1, d_2) = (2, 4), (2, 5), (2, 6)$ and $n = 0$, or $(d_1, d_2) = (2, 3), (2, 4), (2, 5)$ and $n = 1$, or $(d_1, d_2) = (1, d)$ for $n = 1, 2$ and any $d \geq n$.

Furthermore, in all cases listed in Examples 3.4–3.7 but *not* in the above list, one can easily construct a $(\mathbb{C}^*)^4$ -fixed stable pair in $P_n(X, \beta) \setminus P_n(S, \beta)$ using the combinatorial description of stable pairs on toric varieties [PT2, CK2].

Conclusion. For any (S, L_1, L_2) with $L_1 \otimes L_2 \cong K_S$, L_1^{-1}, L_2^{-1} non-trivial and nef, and S minimal and toric, we have classified *all* $\beta \in H_2(S, \mathbb{Z})$ and $n \geq 0$ such that $P_n(X, \beta) \cong P_n(S, \beta) \neq \emptyset$.

In the next section, we develop a method to determine the stable pair invariants $P_{n,\beta}([\text{pt}])$ in all of these cases (tabulated in Appendix A).

4. INVARIANTS

4.1. Virtual classes of relative Hilbert schemes. For S a smooth projective surface, $\beta \in H_2(S, \mathbb{Z})$, and $n \in \mathbb{Z}$, the moduli space $P_n(S, \beta)$ has a nice description in terms of relative Hilbert schemes due to Pandharipande-Thomas [PT3]. Given a stable pair $[(F, s)] \in P_n(S, \beta)$, one has a short exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow F \rightarrow Q \rightarrow 0,$$

where C is the scheme theoretic support of F . Dualizing on C yields a short exact sequence

$$0 \rightarrow F^* \rightarrow \mathcal{O}_C \rightarrow \mathcal{E}xt^1(Q, \mathcal{O}_C) \rightarrow 0,$$

where we used $\mathcal{E}xt^1(F, \mathcal{O}_C) = 0$ by [PT3, Lem. B.2]. Hence $\mathcal{E}xt^1(Q, \mathcal{O}_C) \cong \mathcal{O}_Z$ for some 0-dimensional subscheme $Z \subseteq C$ of length

$$m = n + g(\beta) - 1 = n + \frac{1}{2}\beta(\beta + K_S).$$

As shown in [PT3, Prop. B.8.], the family version of this argument gives an isomorphism

$$(4.1) \quad P_n(S, \beta) \cong \text{Hilb}^m(\mathcal{C}/H_\beta),$$

where $\text{Hilb}^m(\mathcal{C}/H_\beta)$ denotes the relative Hilbert scheme of m points on the fibres of the universal curve $\mathcal{C} \rightarrow H_\beta$ and H_β denotes the Hilbert scheme of effective divisors on S in class β . The description in terms of relative Hilbert schemes helps to establish smoothness. Although we do not need it for this paper, we include the following observation.

Proposition 4.1. *In all the cases listed in Proposition 3.9 and Remark 3.10, $P_n(S, \beta)$ is smooth.*

Proof. The method in this proof was also used in [KST]. Let $S = \mathbb{P}^2$, then for any $\beta = d[H]$ with $d \geq 1$, and any $n \in \mathbb{Z}$, we have a morphism

$$(4.2) \quad P_n(S, \beta) \cong \text{Hilb}^m(\mathcal{C}/|\mathcal{O}(d)|) \rightarrow S^{[m]},$$

where $m = n + \frac{1}{2}d(d-3)$. The fibre over $Z \in S^{[m]}$ is the projectivization of the kernel of the evaluation map $H^0(\mathbb{P}^2, \mathcal{O}(d)) \rightarrow H^0(Z, \mathcal{O}(d)|_Z)$. It suffices to show that for n and $d \neq 1$ as in Proposition 3.9, this map is surjective. Then it follows that the fibres of (4.2) are equidimensional projective spaces and $P_n(S, \beta)$ is smooth, because $S^{[m]}$ is also smooth. Surjectivity of the evaluation map for all $Z \in S^{[m]}$ is equivalent to $(m-1)$ -very ampleness of $\mathcal{O}(d)$ (by definition, [BS]). Beltrametti-Sommese showed that $\mathcal{O}(d)$ is $(m-1)$ -very ample if and only if $m-1 \leq d$, i.e.

$$n \leq d - \frac{1}{2}d(d-3) + 1.$$

This inequality is satisfied for all n and $d \neq 1$ in Cases (2), (3) of Proposition 3.9. Smoothness of $P_n(S, \beta)$ for $d = 1$ and any n is clear, because in this case the fibres of (4.2) are $\text{Sym}^m(\mathbb{P}^1) \cong \mathbb{P}^m$.

For $S = \mathbb{P}^1 \times \mathbb{P}^1$, $\mathcal{O}(d_1H_1 + d_2H_2)$ is k -very ample if and only if $k \leq \min\{d_1, d_2\}$ [BS]. For $S = \mathbb{F}_a$ (for any $a \geq 1$), $\mathcal{O}(d_1B + d_2F)$ is k -very ample if and only if $k \leq \min\{d_1, d_2 - ad_1\}$ [BS]. The proof in the remaining cases of Proposition 3.9 and Remark 3.10 then follows similarly. \square

Let $\mathcal{Z} \subseteq S \times S^{[m]}$ be the universal subscheme and denote the pull-back of \mathcal{Z} to $S \times S^{[m]} \times H_\beta$ by the same symbol (and similarly for $\mathcal{C} \subseteq S \times H_\beta$). Consider the rank m vector bundle

$$\mathcal{O}(\mathcal{C})^{[m]} := \pi_*(\mathcal{O}(\mathcal{C})|_{\mathcal{Z}})$$

on $S^{[m]} \times H_\beta$, where $\pi : S \times S^{[m]} \times H_\beta \rightarrow S^{[m]} \times H_\beta$ is the projection. By [KT1, App. A], there exists a tautological section s of $\mathcal{O}(\mathcal{C})^{[m]}$ cutting out $\text{Hilb}^m(\mathcal{C}/H_\beta)$ from its ambient space

$$\begin{array}{c} \mathcal{O}(\mathcal{C})^{[m]} \\ \downarrow \Bigg) s \\ s^{-1}(0) \cong \text{Hilb}^m(\mathcal{C}/H_\beta) \hookrightarrow S^{[m]} \times H_\beta. \end{array}$$

In general H_β is not smooth (or smooth but not of expected dimension). Therefore, this construction only provides a *relative* perfect obstruction theory on $\text{Hilb}^m(\mathcal{C}/H_\beta) \rightarrow H_\beta$. The Hilbert scheme of divisors H_β has a natural perfect obstruction theory

$$(\mathbf{R}p_*\mathcal{O}_{\mathcal{C}}(\mathcal{C}))^\vee \rightarrow \mathbb{L}_{H_\beta},$$

where $p : S \times H_\beta \rightarrow H_\beta$ denotes projection. This is the perfect obstruction theory used to define the Poincaré/Seiberg-Witten invariants of S in [DKO, CK]. Taken together, these provide an *absolute* perfect obstruction theory on $P_n(S, \beta) \cong \text{Hilb}^m(\mathcal{C}/H_\beta)$ by [KT1, App. A.3]. The virtual tangent bundle of this absolute perfect obstruction theory is

$$\mathbf{R}\mathcal{H}om_{\pi_S}(\mathbb{I}_S^\bullet, \mathbb{F})$$

where $\mathbb{I}_S^\bullet = \{\mathcal{O} \rightarrow \mathbb{F}\}$ denotes the universal stable pair on $S \times P_n(S, \beta)$ and $\pi_S : S \times P_n(S, \beta) \rightarrow P_n(S, \beta)$ is the projection. By [Koo, Prop. 2.1], the resulting virtual class satisfies

$$(4.3) \quad j_*[\text{Hilb}^m(\mathcal{C}/H_\beta)]^{\text{vir}} = (S^{[m]} \times [H_\beta])^{\text{vir}} \cdot e(\mathcal{O}(\mathcal{C})^{[m]}).$$

The corresponding virtual class on $P_n(S, \beta)$, via the isomorphism (4.1), is denoted by $[P_n(S, \beta)]^{\text{vir}}$.

4.2. Comparison of virtual classes. Let S be a smooth projective surface and $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$. We consider the local surface $X = \text{Tot}_S(L_1 \oplus L_2)$, which is a Calabi-Yau 4-fold. Fix $n \in \mathbb{Z}$ and $\beta \in H_2(S, \mathbb{Z})$ such that $P_n(X, \beta)$ is proper. Then it has a virtual class

$$(4.4) \quad [P_n(X, \beta)]^{\text{vir}} \in H_{2n}(P_n(X, \beta), \mathbb{Z}),$$

in the sense of Borisov-Joyce [BJ], which depends on a choice of orientation on $P_n(X, \beta)$. In order to apply [BJ] to our case, we need the existence of a (-2) -shifted symplectic structure on $P_n(X, \beta)$ and an orientability result as reviewed in Section 2.1. The existence of a (-2) -shifted symplectic structure on $P_n(X, \beta)$ was shown in [Bus, Thm. 7.3.2] and [Pre, Thm. 4.0.8]. By [CMT2, Thm. 5.3], the existence of orientations on $P_n(X, \beta)$ can be reduced to the existence of orientations on the moduli stack $\mathfrak{M}_n(X, \beta)$ of coherent sheaves F on X with 1-dimensional proper support of class β and $\chi(F) = n$. Taking $Y = \text{Tot}_S(L_1)$, we have $X = \text{Tot}_Y(K_Y)$. By considering the derived enhancement of $\mathfrak{M}_n(X, \beta)$, it is the (-2) -shifted cotangent bundle of a

derived moduli stack of sheaves on Y . Therefore $\mathfrak{M}_n(X, \beta)$ has an orientation (see e.g. [Tod, Lem. 4.3] for a similar argument in the case of Calabi-Yau threefolds).

We denote by $[\text{pt}] \in H^4(X, \mathbb{Z})$ the pull-back along $\pi : X \rightarrow S$ of the Poincaré dual of the point class on S . Using the same notation as in Section 1.2, we define stable pair invariants

$$(4.5) \quad P_{n,\beta}([\text{pt}]) := \int_{[P_n(X,\beta)]^{\text{vir}}} \tau([\text{pt}])^n \in \mathbb{Z}.$$

When $n = 0$, we simply write $P_{0,\beta} := P_{0,\beta}([\text{pt}])$.

Assuming $P_n(X, \beta) \cong P_n(S, \beta)$, we can compare the virtual class (4.4) to the virtual class on the relative Hilbert scheme (4.3) studied in [KT1, KT2]. In Proposition 3.9 and Remark 3.10 we gave a list of examples where this assumption is satisfied.

Proposition 4.2. *Let S be a smooth projective surface, $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$ and let $X = \text{Tot}_S(L_1 \oplus L_2)$. Suppose $\beta \in H_2(S, \mathbb{Z})$ and $n \geq 0$ are chosen such that $P_n(X, \beta) \cong P_n(S, \beta)$. Then there exists a choice of orientation such that*

$$[P_n(X, \beta)]^{\text{vir}} = (-1)^{\beta \cdot L_2 + n} \cdot e(-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)) \cdot [P_n(S, \beta)]^{\text{vir}}.$$

Here $[P_n(S, \beta)]^{\text{vir}}$ is the virtual class induced from the relative Hilbert scheme (Section 4.1), $\mathbb{I}_S^\bullet = \{\mathcal{O} \rightarrow \mathbb{F}\}$ denotes the universal stable pair on $S \times P_n(S, \beta)$, and $\pi_{P_S} : S \times P_n(S, \beta) \rightarrow P_n(S, \beta)$ is the projection. The sign results from a preferred choice of orientation.

Proof. Let $Y = \text{Tot}_S(L_1)$. Then $X = \text{Tot}_Y(K_Y)$ is the total space of the canonical bundle of Y . By the assumption, we have isomorphisms of moduli spaces

$$(4.6) \quad P_n(S, \beta) \cong P_n(Y, \beta) \cong P_n(X, \beta).$$

Let $\iota : S \hookrightarrow Y$ denote the zero section. A stable pair $I_S^\bullet = \{s : \mathcal{O}_S \rightarrow F\} \in P_n(S, \beta)$ on S induces a stable pair

$$I_Y^\bullet = \{\mathcal{O}_Y \rightarrow \iota_* \mathcal{O}_S \xrightarrow{\iota^* s} \iota_* F\}$$

on Y . Consider the distinguished triangle

$$(4.7) \quad I_Y^\bullet \rightarrow \mathcal{O}_Y \rightarrow \iota_* F.$$

Applying $\mathbf{R}\mathcal{H}om_Y(I_Y^\bullet, \cdot)$ and taking out trace gives a distinguished triangle

$$\mathbf{R}\mathcal{H}om_Y(I_Y^\bullet, \iota_* F) \rightarrow \mathbf{R}\mathcal{H}om_Y(I_Y^\bullet, I_Y^\bullet)_0[1] \rightarrow \mathbf{R}\mathcal{H}om_Y(\iota_* F, \mathcal{O}_Y)[2].$$

Applying adjunction and the isomorphism

$$(4.8) \quad \mathbf{L}\iota^* I_Y^\bullet \cong I_S^\bullet \oplus F \otimes L_1^{-1}$$

gives a long exact sequence

$$\cdots \rightarrow \text{Ext}_S^i(I_S^\bullet, F) \oplus \text{Ext}_S^i(F, F \otimes L_1) \rightarrow \text{Ext}_Y^{i+1}(I_Y^\bullet, I_Y^\bullet)_0 \rightarrow \text{Ext}_Y^{i+2}(\iota_* F, \mathcal{O}_Y) \rightarrow \cdots.$$

Note that $\text{Ext}_Y^1(\iota_* F, \mathcal{O}_Y) \cong \text{Ext}_Y^2(\mathcal{O}_Y, \iota_* F \otimes K_Y)^\vee = 0$. Furthermore, the isomorphism (4.6) induces an isomorphism on Zariski tangent spaces

$$\text{Ext}_S^0(I_S^\bullet, F) \cong \text{Ext}_Y^1(I_Y^\bullet, I_Y^\bullet)_0.$$

Therefore, we deduce $\text{Hom}_S(F, F \otimes L_1) = 0$ (similarly $\text{Hom}_S(F, F \otimes L_2) = 0$). This vanishing allows us to conclude that the natural (Le Potier) pair obstruction theory

$$(4.9) \quad (\mathbf{R}\mathcal{H}om_{\pi_{P_Y}}(\mathbb{I}_Y^\bullet, \iota_{P_Y*} \mathbb{F}))^\vee \rightarrow \mathbb{L}_{P_n(Y, \beta)}$$

is *perfect*, i.e. 2-term, as we will now show⁷. Here $\mathbb{I}_Y^\bullet = \{\mathcal{O} \rightarrow \iota_{P_Y*} \mathbb{F}\}$ denotes the universal stable pair on $Y \times P_n(Y, \beta)$, $\iota_{P_Y} : S \times P_n(Y, \beta) \hookrightarrow Y \times P_n(Y, \beta)$ is the base change of the zero section, and $\pi_{P_Y} : Y \times P_n(Y, \beta) \rightarrow P_n(Y, \beta)$ denotes the projection.

From the distinguished triangle

$$(4.10) \quad \mathbf{R}\mathcal{H}om_Y(\iota_* F, \iota_* F) \rightarrow \mathbf{R}\mathcal{H}om_Y(\mathcal{O}_Y, \iota_* F) \rightarrow \mathbf{R}\mathcal{H}om_Y(I_Y^\bullet, \iota_* F),$$

we obtain an exact sequence

$$0 = H^2(Y, \iota_* F) \rightarrow \text{Ext}_Y^2(I_Y^\bullet, \iota_* F) \rightarrow \text{Ext}_Y^3(\iota_* F, \iota_* F) \rightarrow 0 \rightarrow \text{Ext}_Y^3(I_Y^\bullet, \iota_* F) \rightarrow 0.$$

Moreover, by adjunction and $\mathbf{L}\iota^* F \cong F \oplus F \otimes L_1^{-1}[1]$, we have

$$\text{Ext}_Y^3(\iota_* F, \iota_* F) \cong \text{Ext}_S^3(F, F) \oplus \text{Ext}_S^2(F, F \otimes L_1) \cong \text{Hom}_S(F, F \otimes L_2)^\vee = 0.$$

Hence $\text{Ext}_Y^2(I_Y^\bullet, \iota_* F) \cong \text{Ext}_Y^3(\iota_* F, \iota_* F) = 0$. Also note that $\text{Hom}_Y(\iota_* F, \iota_* F) \rightarrow \text{Hom}_Y(\mathcal{O}_Y, \iota_* F)$ is injective. Therefore $\text{Ext}_Y^i(I_Y^\bullet, \iota_* F) = 0$ unless $i = 0, 1$ and the complex (4.9) is 2-term. We denote the corresponding virtual class by $[P_n(Y, \beta)]_{\text{pair}}^{\text{vir}}$.

⁷This was proved for irreducible β in [CMT2, Lem. 3.1].

We can now use the argument of [CMT2, Prop. 4.3] to deduce that the 4-fold virtual class $[P_n(X, \beta)]^{\text{vir}}$ of (4.4) equals the pairs virtual class $[P_n(Y, \beta)]_{\text{pair}}^{\text{vir}}$. For completeness, we repeat the argument. Just like pushing forward from S to Y gives (4.7) and (4.8), pushing forward further to X gives

$$\begin{aligned} \mathbf{R}\text{Hom}_X(I_X^\bullet, j_* \iota_* F) &\rightarrow \mathbf{R}\text{Hom}_X(I_X^\bullet, I_X^\bullet)_0[1] \rightarrow \mathbf{R}\text{Hom}_X(j_* \iota_* F, \mathcal{O}_X)[2], \\ \mathbf{L}j^* I_X^\bullet &\cong I_Y^\bullet \oplus \iota_* F \otimes K_Y^{-1}, \end{aligned}$$

where

$$I_X^\bullet = \{\mathcal{O}_X \rightarrow j_* \iota_* \mathcal{O}_S \xrightarrow{j_* \iota_* s} j_* \iota_* F\}$$

and we denote the zero section by $j: Y \hookrightarrow X$. Let T be the cone of the composition

$$\mathbf{R}\text{Hom}_Y(I_Y^\bullet, \iota_* F) \rightarrow \mathbf{R}\text{Hom}_X(I_X^\bullet, j_* \iota_* F) \rightarrow \mathbf{R}\text{Hom}_X(I_X^\bullet, I_X^\bullet)_0[1].$$

Then T fits in the distinguished triangles

$$(4.11) \quad \begin{aligned} \mathbf{R}\text{Hom}_Y(I_Y^\bullet, \iota_* F) &\rightarrow \mathbf{R}\text{Hom}_X(I_X^\bullet, I_X^\bullet)_0[1] \rightarrow T, \\ \mathbf{R}\text{Hom}_Y(\iota_* F, \iota_* F \otimes K_Y) &\rightarrow T \rightarrow \mathbf{R}\text{Hom}_X(j_* \iota_* F, \mathcal{O}_X)[2]. \end{aligned}$$

Applying Serre duality to the first and third term of the second distinguished triangle, dualizing, and shifting gives the following distinguished triangle

$$\mathbf{R}\text{Hom}_Y(\iota_* F, \iota_* F)[2] \rightarrow \mathbf{R}\text{Hom}_X(\mathcal{O}_X, j_* \iota_* F)[2] \rightarrow T^\vee.$$

Comparing to (4.10), we obtain $T \cong \mathbf{R}\text{Hom}_Y(I_Y^\bullet, \iota_* F)^\vee[-2]$. Hence from (4.11) we get a short exact sequence

$$0 \rightarrow \text{Ext}_Y^1(I_Y^\bullet, \iota_* F) \rightarrow \text{Ext}_X^2(I_X^\bullet, I_X^\bullet)_0 \rightarrow \text{Ext}_Y^1(I_Y^\bullet, \iota_* F)^\vee \rightarrow 0,$$

where we crucially used $\text{Ext}_Y^2(I_Y^\bullet, \iota_* F) = 0$ which was shown above. This way, we obtain a half-dimensional subspace $\text{Ext}_Y^1(I_Y^\bullet, \iota_* F)$ of $\text{Ext}_X^2(I_X^\bullet, I_X^\bullet)_0$. One can show that it is isotropic by the exact same argument as in the proof of [CMT2, Prop. 3.3, Prop. 2.11]. From this, it is concluded in loc. cit. that

$$[P_n(X, \beta)]^{\text{vir}} = (-1)^{\beta \cdot L_2 + n} \cdot [P_n(Y, \beta)]_{\text{pair}}^{\text{vir}}.$$

Here the sign comes from a choice of preferred orientation discussed in a similar setting in [Cao].

Finally, we express the pairs virtual class on Y in terms of the pairs virtual class on S . By adjunction, we have

$$\begin{aligned} \mathbf{R}\text{Hom}_{\pi_{P_Y}}(\mathbb{I}_Y^\bullet, \iota_{P_Y}^* \mathbb{F}) &\cong \mathbf{R}\text{Hom}_{\pi_{P_S}}(\mathbf{L}\iota_{P_Y}^* \mathbb{I}_Y^\bullet, \mathbb{F}) \\ &\cong \mathbf{R}\text{Hom}_{\pi_{P_S}}(\mathbb{I}_S^\bullet, \mathbb{F}) \oplus \mathbf{R}\text{Hom}_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1), \end{aligned}$$

where $\pi_{P_S}: S \times P_n(S, \beta) \rightarrow P_n(S, \beta) \cong P_n(Y, \beta)$ denotes the projection. From the vanishing $\text{Hom}_S(F, F \otimes L_1) = \text{Hom}_S(F, F \otimes L_2) = 0$, for all $[(F, s)] \in P_n(S, \beta)$, we deduce that

$$-\mathbf{R}\text{Hom}_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1) \cong \mathcal{E}xt_{\pi_{P_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L_1)$$

is locally free on $P_n(Y, \beta) \cong P_n(S, \beta)$. Hence the two virtual tangent bundles on $P_n(Y, \beta) \cong P_n(S, \beta)$ differ by a locally free sheaf (in degree 1). Therefore, by [Sie, Thm. 4.6], we have

$$[P_n(Y, \beta)]_{\text{pair}}^{\text{vir}} = e(-\mathbf{R}\text{Hom}_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)) \cdot [P_n(S, \beta)]^{\text{vir}}. \quad \square$$

Remark 4.3. Let S be a smooth projective surface satisfying $b_1(S) = p_g(S) = 0$. Let $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$ and $X = \text{Tot}_S(L_1 \oplus L_2)$. Suppose $P_n(X, \beta)$ is proper. Once the virtual localization formula for stable pair theory on Calabi-Yau 4-folds is established⁸, we expect that it will induce a virtual class on each of the connected components of $P_n(X, \beta)^{\mathbb{C}^*}$, where \mathbb{C}^* is the 1-dimensional subtorus, preserving the Calabi-Yau volume form, inside the torus $\mathbb{C}^* \times \mathbb{C}^*$ acting on the fibres of X . When $P_n(S, \beta) \subseteq P_n(X, \beta)^{\mathbb{C}^*}$ is open and closed, this gives a virtual class, which we expect to be given by (for an appropriate choice of orientation)

$$(4.12) \quad (-1)^{\beta \cdot L_2 + n} \cdot e(-\mathbf{R}\text{Hom}_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1) \otimes \mathbf{t}_1) \cdot [P_n(S, \beta)]^{\text{vir}},$$

where $[P_n(S, \beta)]^{\text{vir}}$ is the virtual class induced from the relative Hilbert scheme (Section 4.1), \mathbf{t}_1 is a primitive character corresponding to the first component of the action of $\mathbb{C}^* \times \mathbb{C}^*$, and $e(\cdot)$ denotes equivariant Euler class. When $P_n(X, \beta)$ is non-proper, one could define the contribution of $P_n(S, \beta)$ to the stable pair invariants of X by (4.12) (capped with appropriate insertions).

⁸In the special case S is moreover toric and $P_n(S, \beta) \cong P_n(X, \beta)$ is smooth, a virtual localization formula was proved in [CK2, Thm. A.1]. A general virtual localization formula has been announced by Oh-Thomas [OT].

4.3. Main theorem. We are now ready to prove the theorem of the introduction. Recall from (1.6) that we denote by $T_{S^{[m]}}(\mathcal{L})$ the twisted (by \mathcal{L}) tangent bundle of $S^{[m]}$.

Theorem 4.4. *Let S be a smooth projective surface with $b_1(S) = p_g(S) = 0$ and $L_1, L_2 \in \text{Pic}(S)$ such that $L_1 \otimes L_2 \cong K_S$. Suppose $\beta \in H_2(S, \mathbb{Z})$ and $n \geq 0$ are chosen such that $P_n(X, \beta) \cong P_n(S, \beta)$ for $X = \text{Tot}_S(L_1 \oplus L_2)$. Denote by $[\text{pt}] \in H^4(X, \mathbb{Z})$ the pull-back of the Poincaré dual of the point class on S . Let $P_n(X, \beta)$ be endowed with the orientation as in (1.5). Then*

$$P_{n,\beta}([\text{pt}]) = (-1)^{\beta \cdot L_2 + n} \int_{S^{[m]} \times \mathbb{P}^{\chi(\beta)-1}} c_m(\mathcal{O}_S(\beta)^{[m]}(1)) \frac{h^n(1+h)^{\chi(L_1(\beta))}(1-h)^{\chi(L_2(\beta))} c(T_{S^{[m]}}(L_1))}{c(L_1(\beta)^{[m]}(1)) \cdot c((L_2(\beta)^{[m]}(1))^\vee)},$$

when $\beta^2 \geq 0$. Here $m := n + g(\beta) - 1$ and $h := c_1(\mathcal{O}(1))$. Moreover, $P_{n,\beta}([\text{pt}]) = 0$ when $\beta^2 < 0$.

Proof. Suppose β is an effective divisor and $m \geq 0$, otherwise $P_n(S, \beta) \cong \text{Hilb}^m(\mathcal{C}/H_\beta) = \emptyset$ and $P_{n,\beta}([\text{pt}]) = 0$. Consider the closed embedding

$$j : \text{Hilb}^m(\mathcal{C}/H_\beta) \hookrightarrow S^{[m]} \times H_\beta,$$

as in Section 4.1. Below, we will show that there exists a class $\psi \in K_0(S^{[m]} \times H_\beta)$ restricting to $e(-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)) \cdot \tau([\text{pt}])^n$ on $\text{Hilb}^m(\mathcal{C}/H_\beta)$. By Proposition 4.2, it follows that

$$(4.13) \quad P_{n,\beta}([\text{pt}]) = \int_{S^{[m]} \times [H_\beta]^{\text{vir}}} c_m(\mathcal{O}(\mathcal{C})^{[m]}) \cdot \psi.$$

Since $b_1(S) = p_g(S) = 0$, we have [DKO, KT2]

$$[H_\beta]^{\text{vir}} = |\beta|^{\text{vir}} = h^{h^1(\mathcal{O}(\beta))} \cap |\beta| \in H_{2\chi(\beta)-2}(|\beta|),$$

where h denotes the class of the hyperplane on $H_\beta = |\beta|$. Furthermore, we have

$$\mathcal{O}(\mathcal{C})^{[m]} := \pi_*(\mathcal{O}(\mathcal{C})|_{\mathcal{Z}}) \cong \mathcal{O}(\beta)^{[m]}(1),$$

which follows from the isomorphism $\mathcal{O}(\mathcal{C}) \cong \mathcal{O}_S(\beta) \boxtimes \mathcal{O}(1)$ on $S \times |\beta|$. Therefore $P_{n,\beta}([\text{pt}]) = 0$, unless $\chi(\beta) \geq 1$, which we assume from now on.

Recall from the proof of Proposition 4.2 that $-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1) \cong \mathcal{E}xt_{\pi_{P_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L_1)$ is locally free on $P_n(S, \beta)$ and its rank is β^2 . Therefore $P_{n,\beta}([\text{pt}]) = 0$ unless $\beta^2 \geq 0$, which we assume from now on. Next, we extend the complex $-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)$ from $P_n(S, \beta)$ to $S^{[m]} \times |\beta|$. In K -theory, we have $\mathbb{I}_S^\bullet = \mathcal{O} - \mathbb{F}$ and

$$\begin{aligned} -\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1) &= -\chi(L_1) \otimes \mathcal{O} + \mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathcal{O}, \mathbb{I}_S^\bullet \boxtimes L_1) \\ &\quad + \mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{I}_S^\bullet, L_1) - \mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{I}_S^\bullet, \mathbb{I}_S^\bullet \boxtimes L_1) \\ &= -\chi(L_1) \otimes \mathcal{O} + \mathbf{R}\mathcal{H}om_{\pi_{P_S}}((\mathbb{I}_S^\bullet)^\vee, L_1) \\ &\quad + \mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathcal{O}, (\mathbb{I}_S^\bullet)^\vee \boxtimes L_1) - \mathbf{R}\mathcal{H}om_{\pi_{P_S}}((\mathbb{I}_S^\bullet)^\vee, (\mathbb{I}_S^\bullet)^\vee \boxtimes L_1), \end{aligned}$$

where we suppressed some obvious pull-backs. On $S \times |\beta| \times S^{[m]}$, we have the sheaf $\mathcal{I} \boxtimes \mathcal{O}_S(\beta) \boxtimes \mathcal{O}(1)$, where we use the notation from the introduction. By [KT1, Lem. A.4], we have

$$(\mathbb{I}_S^\bullet)^\vee \cong \mathcal{I} \boxtimes \mathcal{O}_S(\beta) \boxtimes \mathcal{O}(1)|_{\text{Hilb}^m(\mathcal{C}/|\beta|) \times S}.$$

Next we can replace \mathcal{I} by $\mathcal{O} - \mathcal{O}_{\mathcal{Z}}$ in K -theory. Then $-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)$ is the restriction of the following element in the K -group of $S^{[m]} \times |\beta|$

$$\begin{aligned} &-\chi(L_1) \otimes \mathcal{O} + \mathbf{R}\mathcal{H}om_{\pi}((\mathcal{O} - \mathcal{O}_{\mathcal{Z}}) \boxtimes (\mathcal{O}_S(\beta) \otimes L_1^{-1}) \boxtimes \mathcal{O}(1), \mathcal{O}) \\ &\quad + \mathbf{R}\mathcal{H}om_{\pi}(\mathcal{O}, (\mathcal{O} - \mathcal{O}_{\mathcal{Z}}) \boxtimes (\mathcal{O}_S(\beta) \otimes L_1) \boxtimes \mathcal{O}(1)) - \mathbf{R}\mathcal{H}om_{\pi}(\mathcal{I}, \mathcal{I} \boxtimes L_1) \\ (4.14) \quad &= -\chi(L_1) \otimes \mathcal{O} + \chi(L_1(\beta)) \otimes \mathcal{O}(1) + \chi(L_2(\beta)) \otimes \mathcal{O}(-1) \\ &\quad - (L_1(\beta))^{[m]} \boxtimes \mathcal{O}(1) - ((L_2(\beta))^{[m]})^\vee \boxtimes \mathcal{O}(-1) - \mathbf{R}\mathcal{H}om_{\pi}(\mathcal{I}, \mathcal{I} \boxtimes L_1), \end{aligned}$$

where $\pi : S \times S^{[m]} \times |\beta| \rightarrow S^{[m]} \times |\beta|$ denotes the projection and we used Serre duality, $L_1 \otimes L_2 \cong K_S$, and $L_i(\beta) := \mathcal{O}_S(\beta) \otimes L_i$.

Finally, we consider primary insertions

$$\tau : H^4(X, \mathbb{Z}) \rightarrow H^2(P_n(X, \beta), \mathbb{Z}), \quad \tau(\gamma) = \pi_{P*}(\pi_X^* \gamma \cup \text{ch}_3(\iota_* \mathbb{F})),$$

where π_X, π_P are projections from $X \times P_n(X, \beta)$ to corresponding factors, and $\iota : S \times P_n(S, \beta) \hookrightarrow X \times P_n(S, \beta) \cong X \times P_n(X, \beta)$ is the base change of the inclusion of the zero section. Note that $\text{ch}_3(\iota_* \mathbb{F})$ is Poincaré dual to the fundamental class of the scheme theoretic support of $\iota_* \mathbb{F}$, which we denote by $[\iota_* \mathbb{F}]$. The fundamental class of the scheme theoretic support of \mathbb{F} , which we

denote by $[\mathbb{F}]$, equals $j^*[\mathcal{C}]$ where $\mathcal{C} \subseteq S \times S^{[m]} \times |\beta|$ is the pullback of the universal curve over $|\beta|$. Consider the commutative diagram

$$\begin{array}{ccccc}
 & & \xrightarrow{\quad \iota \quad} & & \\
 X & \xleftarrow{\quad p \quad} & S & \xleftarrow{\quad \pi_S \quad} & S \times S^{[m]} \times |\beta| \\
 \uparrow \pi_X & & \uparrow \pi_S & & \\
 X \times P_n(X, \beta) & \xrightarrow{\quad p \quad} & S \times P_n(S, \beta) & \xrightarrow{\quad j \quad} & S \times S^{[m]} \times |\beta| \\
 \downarrow \pi_P & & \downarrow \pi_P & & \downarrow \pi \\
 P_n(X, \beta) & \xrightarrow{\quad \cong \quad} & P_n(S, \beta) & \xrightarrow{\quad j \quad} & S^{[m]} \times |\beta|
 \end{array}$$

For $\gamma := p^*[\text{pt}] \in H^4(X, \mathbb{Z})$, where $[\text{pt}]$ denotes the Poincaré dual of the point class on S , we work our way through the diagram

$$\begin{aligned}
 \tau(\gamma) &= \pi_{P*}(\pi_X^* \gamma \cup [\iota_* \mathbb{F}]) \\
 &= \pi_{P*} p_*(p^* \pi_S^* [\text{pt}] \cup \iota_* [\mathbb{F}]) \\
 &= \pi_{P*}(\pi_S^* [\text{pt}] \cup [\mathbb{F}]) \\
 &= j^*(\pi_*(\pi_S^* [\text{pt}] \cup [\mathcal{C}])).
 \end{aligned}
 \tag{4.15}$$

Using, once more, that on $S \times |\beta|$ we have $\mathcal{O}(\mathcal{C}) \cong \mathcal{O}_S(\beta) \boxtimes \mathcal{O}(1)$, we conclude

$$\pi_*(\pi_S^* [\text{pt}] \cup [\mathcal{C}]) = h \int_S \text{pt} = h.$$

Therefore $\tau([\text{pt}])^n$ is simply the restriction of the class h^n on $S^{[m]} \times |\beta|$.

Since $\text{rk}(-\mathbf{R}\mathcal{H}om_{\pi_{P_S}}(\mathbb{F}, \mathbb{F} \boxtimes L_1)) = \beta^2 = 2m + \chi(\beta) - 1 - m - n$, we can replace Euler class by total Chern class. The result now follows from (4.13), (4.14), and (4.15). \square

Remark 4.5. For surfaces with $p_g(S) = 0$ and $b_1(S) > 0$, we can still use the formula for the virtual class from Proposition 4.2. Suppose $h^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$. Then the virtual class $[H_\beta]^{\text{vir}}$ can be calculated by fixing a sufficiently ample effective divisor A on S and considering the embedding $H_\beta \hookrightarrow H_{[A]+\beta}$ as in [DKO] (see also [KT1, Prop. A.2], [KT2]). Therefore the invariant can be expressed as an integral over $S^{[m]} \times H_{[A]+\beta}$, where $H_{[A]+\beta}$ is a projective bundle over $\text{Pic}_{[A]+\beta}(S)$ via the Abel-Jacobi map. Pushing forward along the Abel-Jacobi map, the invariant can be expressed as an integral over $S^{[m]} \times \text{Pic}_{A+\beta}(S)$.

4.4. Atiyah-Bott localization. In Corollary 3.9, Remark 3.10, we gave examples of (S, L_1, L_2) for which the assumptions of Theorem 4.4 are satisfied. In all of these cases, S is a toric surface. As a consequence, $X = \text{Tot}_S(L_1 \oplus L_2)$ is also toric, so in principle one could calculate the invariant $P_{n,\beta}([\text{pt}])$ using the vertex formalism for stable pair invariants on toric Calabi-Yau 4-folds developed in [CK2, CKM]⁹. However, the number of $(\mathbb{C}^*)^4$ -fixed points is typically very large. For instance, for $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$, $(d_1, d_2) = (2, 4)$, and $n = 0$, we have 182 fixed points, whereas Theorem 4.4 only involves an integral over $S^{[2]} \times \mathbb{P}^{14}$.

The calculation of intersection numbers on Hilbert schemes of points on toric surfaces is a classical subject (see e.g. [ES]). Let S be a smooth projective toric surface with torus $T = (\mathbb{C}^*)^2$. The action of T on S lifts to an action of T on $S^{[m]}$ for any m . Let P be a polynomial expression in Chern classes of

$$(4.16) \quad \mathbf{R}\Gamma(\mathcal{L}) \otimes \mathcal{O} - \mathbf{R}\mathcal{H}om_\pi(\mathcal{I}, \mathcal{I} \boxtimes \mathcal{L}), \quad \mathcal{L}^{[m]},$$

for various choices of T -equivariant line bundles \mathcal{L} on S and where $\pi : S \times S^{[m]} \rightarrow S^{[m]}$ denotes the projection. Note that this includes Chern classes of the tangent bundle, which can be expressed as $-\mathbf{R}\mathcal{H}om_\pi(\mathcal{I}, \mathcal{I})_0$ (since $\text{Ext}^i(I_Z, I_Z)_0 = 0$ for $Z \in S^{[m]}$ and $i \neq 1$). Suppose also that the degree of P , as a class in the Chow ring $A^*(S^{[m]})$, equals $\dim S^{[m]} = 2m$. By the Atiyah-Bott localization formula [AB], we have

$$\int_{S^{[m]}} P = \int_{(S^{[m]})^T} \frac{P|_{(S^{[m]})^T}}{e(N_{(S^{[m]})^T/S^{[m]}})},$$

where $e(\cdot)$ denotes the T -equivariant Euler class and $N_{(S^{[m]})^T/S^{[m]}}$ is the normal bundle of the fixed point locus $(S^{[m]})^T \subseteq S^{[m]}$. Furthermore, in this formula one has to choose a T -equivariant

⁹In loc. cit. it is assumed that the fixed locus $P_n(X, \beta)^{(\mathbb{C}^*)^4}$ is at most 0-dimensional. This is the case for all local Calabi-Yau 4-fold surfaces.

lift of P . More precisely, one can choose a T -equivariant structure on all (complexes of) sheaves appearing in P and replace all Chern classes appearing in P by T -equivariant Chern classes.

The fixed point locus consists of isolated reduced points, which can be described combinatorially. Consider a cover by maximal T -invariant affine open subsets:

$$\{U_\sigma \cong \operatorname{Spec} \mathbb{C}[x_\sigma, y_\sigma]\}_{\sigma=1}^{e(S)}.$$

Then the fixed locus $(S^{[m]})^T$ precisely consists of the closed subschemes of S defined by collections of monomial ideals

$$\{I_\sigma \subseteq \mathbb{C}[x_\sigma, y_\sigma]\}_{\sigma=1}^{e(S)}$$

of total colength m . The monomial ideals of finite colength in $\mathbb{C}[x, y]$ are in bijective correspondence with partitions. Explicitly, $\lambda = (\lambda_1 \geq \dots \geq \lambda_\ell)$ corresponds to the ideal

$$(y^{\lambda_1}, xy^{\lambda_2}, \dots, x^{\ell-1}y^{\lambda_\ell}, x^\ell),$$

where $\ell(\lambda) = \ell$ is the length of λ . Hence we can index the points of the fixed locus $(S^{[m]})^T$ by collections of partitions

$$\lambda = \{\lambda^{(\sigma)}\}_{\sigma=1}^{e(S)}$$

of total size

$$\sum_{\sigma=1}^{e(S)} |\lambda^{(\sigma)}| = \sum_{\sigma=1}^{e(S)} \sum_{i=1}^{\ell(\lambda^{(\sigma)})} \lambda_i^{(\sigma)} = m.$$

Denote the closed subscheme corresponding to λ by Z_λ .

In order to calculate integrals such as the one in Theorem 4.4 by Atiyah-Bott localization, we need to consider Chern classes of

(4.17)

$$\begin{aligned} \mathcal{L}^{[m]}|_{Z_\lambda} &= H^0(\mathcal{L}|_{Z_\lambda}) \in K_0^T(\operatorname{pt}) = \mathbb{Z}[t_1^{\pm 1}, t_2^{\pm 1}], \\ \left(\mathbf{R}\Gamma(\mathcal{L}) \otimes \mathcal{O} - \mathbf{R}\mathcal{H}om_\pi(\mathcal{I}, \mathcal{I} \boxtimes \mathcal{L}) \right)|_{Z_\lambda} &= \mathbf{R}\Gamma(\mathcal{L}|_{Z_\lambda}) - \mathbf{R}\mathcal{H}om_S(I_{Z_\lambda}, I_{Z_\lambda} \otimes \mathcal{L}) \in K_0^T(\operatorname{pt}), \end{aligned}$$

where t_1, t_2 are the equivariant parameters of T .

Suppose Z_λ is a 0-dimensional T -equivariant subscheme supported entirely on a maximal T -invariant affine open subset U_σ and set $\lambda := \lambda^{(\sigma)}$. Suppose we choose coordinates such that $U_\sigma = \operatorname{Spec} \mathbb{C}[x, y]$ and the torus action (on coordinate functions) is given by $(t_1, t_2) \cdot (x, y) = (t_1 x, t_2 y)$. Denote the character corresponding to $\mathcal{L}|_{U_\sigma}$ by $\chi(t_1, t_2)$. Then

$$\begin{aligned} H^0(\mathcal{L}|_{Z_\lambda}) &= \chi(t_1, t_2) \cdot Z_\lambda, \\ \text{where } Z_\lambda &:= \sum_{i=1}^{\ell(\lambda)} \sum_{j=1}^{\lambda_i} t_1^{i-1} t_2^{j-1}. \end{aligned} \quad (4.18)$$

Now suppose W_μ is a second 0-dimensional T -equivariant subscheme supported entirely on U_σ and write $\mu := \mu^{(\sigma)}$. The following formula can be deduced from a well-known calculation using Čech cohomology (e.g. see [GK, Prop. 4.1]):

$$\begin{aligned} \mathbf{R}\mathcal{H}om_S(\mathcal{O}_{W_\mu}, \mathcal{O}_{Z_\lambda} \otimes \mathcal{L}) &= \chi(t_1, t_2) W_\mu^* Z_\lambda \frac{(1-t_1)(1-t_2)}{t_1 t_2} \in K_0^T(\operatorname{pt}), \\ \text{where } W_\mu &:= \sum_{i=1}^{\ell(\mu)} \sum_{j=1}^{\mu_i} t_1^{i-1} t_2^{j-1}. \end{aligned} \quad (4.19)$$

Here $(\cdot)^*$ is the involution defined by dualizing.

For arbitrary Z_λ , the K -group classes of (4.17) can be determined from (4.18) and (4.19) by using the following equalities in K -theory

$$\begin{aligned} \mathcal{O}_{Z_\lambda} &= \sum_{\sigma=1}^{e(S)} \mathcal{O}_{Z_{\lambda^{(\sigma)}}}, \\ I_{\mathcal{O}_{Z_\lambda}} &= \mathcal{O}_S - \mathcal{O}_{Z_\lambda}, \end{aligned}$$

where $Z_{\lambda^{(\sigma)}}$ denotes the 0-dimensional closed subscheme supported on U_σ determined by $\lambda^{(\sigma)}$.

Consider Theorem 4.4 for the examples of (S, L_1, L_2) listed in Proposition 3.9 and Remark 3.10. In each case, we calculated the invariant $P_{n,\beta}([\operatorname{pt}])$ by first integrating out the linear system $\mathbb{P}^{X(\beta)-1}$. This amounts to expanding the integrand in powers of $h = c_1(\mathcal{O}(1))$ and taking the coefficient of $h^{X(\beta)-1}$. This gives a polynomial expression P in Chern classes of complexes of the

form (4.16). The integral $\int_{S[m]} P$ is then calculated by Atiyah-Bott localization as described. The resulting stable pair invariants are tabulated in Appendix A.

With the numbers of Appendix A, we are able to do various new checks of the Cao-Maulik-Toda conjectures (Conjectures 1.1 and 1.2). Combining our tables in Appendix A with the tables for $n_{0,\beta}([\text{pt}])$, $n_{1,\beta}$ in [KP, Sect. 3] gives Corollary 1.6 of the introduction.

Bousseau-Brini-van Garrel [BBG] determined all the genus zero Gromov-Witten (and therefore Gopakumar-Vafa type) invariants of $X = \text{Tot}_S(L_1 \oplus L_2)$ with (S, L_1, L_2) as in Remark 3.10 (as well as for other cases). Note that the method of Bousseau-Brini-van Garrel does not produce genus one Gopakumar-Vafa type invariants, so we can only do verifications of Conjecture 1.1 in these cases. Combining the tables in Appendix A with the values for genus zero Gopakumar-Vafa type invariants provided to us by Bousseau-Brini-van Garrel gives Corollary 1.8 of the introduction.

Recall that for (S, L_1, L_2) with $L_1 \otimes L_2 \cong K_S$, L_1^{-1} , L_2^{-1} non-trivial and nef, and S minimal and toric, we classified all values of $n \geq 0$ for which $P_n(S, \beta) \cong P_n(X, \beta)$, and $P_n(S, \beta) \neq \emptyset$ (Remark 3.10). In these cases we therefore calculated *all* stable pair invariants $P_{n,\beta}([\text{pt}])$.

APPENDIX A. TABLES

A.1. Local surfaces. In this section, we list the stable pair invariants $P_{n,\beta}([\text{pt}])$ of $X = \text{Tot}_S(L_1 \oplus L_2)$ for the cases mentioned in Proposition 3.9 and Remark 3.10, and for a few additional cases. We use the following conventions and notation:

- $P_{0,0} := 1$ and $P_{n,0}([\text{pt}]) = 0$ for all $n > 0$.
- Entries decorated with \star were defined by a virtual localization formula on the fixed locus and have been calculated using the vertex formalism as discussed in [CK2, CKM]. In these cases $P_n(X, \beta) \setminus P_n(S, \beta) \neq \emptyset$. See Remark A.2 for a comparison to the globally defined invariants. All other entries have been computed using Theorem 4.4,
- Zeroes decorated with \dagger have non-empty underlying moduli space $P_n(X, \beta)$. In this sense, they are “non-trivial” zeroes.

For $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, we calculated the following values for $P_{0,d} := P_{0,d[H]}$ and $P_{n,d}([\text{pt}]) := P_{n,d[H]}([\text{pt}])$ (for $n > 0$).

$d \setminus n$	0	1	2	3	4
1	0	-1	0^\dagger	0^\dagger	0^\dagger
2	0	1	1	$0^{\star,\dagger}$	
3	-1	-1	-2		
4	2	3			

$$P_{n,1}([\text{pt}]) = 0^\dagger, \quad \forall n \geq 2.$$

For $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -1))$ and $P_{0,(d_1,d_2)} := P_{0,d_1[H_1]+d_2[H_2]}$ and $P_{n,(d_1,d_2)}([\text{pt}]) := P_{n,d_1[H_1]+d_2[H_2]}([\text{pt}])$ (for $n > 0$), where H_1, H_2 are defined in Example 1.4, we have

$P_{0,(d_1,d_2)}$	0	1	2	3	4
0	1	0	0	0	0
1	0	0	0	0	0
2	0	0	1	2	5
3	0	0	2	10	
4	0	0	5		

$P_{1,(d_1,d_2)}([\text{pt}])$	0	1	2	3	4
0	0	1	0	0	0
1	1	1	1	1	1
2	0	1	2	5	
3	0	1	5		
4	0	1			

$P_{2,(d_1,d_2)}([\text{pt}])$	0	1	2	3	4
0	0	0^\dagger	1	0	0
1	0^\dagger	2	2	2	2
2	1	2	5		
3	0	2			
4	0	2			

$P_{3,(d_1,d_2)}([\text{pt}])$	0	1	2	3	4
0	0	0^\dagger	$0^{\star,\dagger}$	1	0
1	0^\dagger	0^\dagger	3^\star		
2	$0^{\star,\dagger}$	3^\star			
3	1				
4	0				

$$\begin{aligned}
P_{0,(1,d)} &= P_{0,(d,1)} = 0, \quad \forall d \geq 0 \\
P_{1,(1,d)}([\text{pt}]) &= P_{1,(d,1)}([\text{pt}]) = 1, \quad \forall d \geq 0 \\
P_{2,(1,d)}([\text{pt}]) &= P_{2,(d,1)}([\text{pt}]) = 2, \quad \forall d \geq 1 \\
P_{n,(0,d)}([\text{pt}]) &= P_{n,(d,0)}([\text{pt}]) = \delta_{n,d}, \quad \forall 0 \leq n \leq d \\
P_{n,(1,1)}([\text{pt}]) &= 0^\dagger, \quad \forall n \geq 3, \\
P_{n,(0,1)}([\text{pt}]) &= P_{n,(1,0)}([\text{pt}]) = 0^\dagger, \quad \forall n \geq 2.
\end{aligned}$$

For $(S, L_1, L_2) = (\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(-1, 0), \mathcal{O}(-1, -2))$, we have:

$$\begin{aligned}
P_{0,(2,2)} &= 1, \quad P_{0,(2,3)} = 2, \quad P_{0,(2,4)} = 5, \quad P_{0,(3,2)} = 2, \\
P_{1,(2,2)}([\text{pt}]) &= 2, \quad P_{1,(2,3)}([\text{pt}]) = 5, \\
P_{1,(d,1)}([\text{pt}]) &= P_{1,(1,d)}([\text{pt}]) = 1, \quad \forall d \geq 1 \\
P_{2,(1,d)}([\text{pt}]) &= 2, \quad \forall d \geq 2, \\
P_{n,(0,1)}([\text{pt}]) &= 0^\dagger, \quad \forall n \geq 2 \\
P_{n,(0,n)}([\text{pt}]) &= 1, \quad \forall n \geq 1.
\end{aligned}$$

Recall the notation for Hirzebruch surfaces from the introduction. Consider $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(-1, -1), \mathcal{O}(-1, -2))$. We write $P_{0,(d_1,d_2)} := P_{0,d_1[B]+d_2[F]}$ and, for $n > 0$, $P_{n,(d_1,d_2)}([\text{pt}]) := P_{n,d_1[B]+d_2[F]}([\text{pt}])$. In the tables below, the rows are for d_1 and the columns for d_2 .

$P_{0,(d_1,d_2)}$	0	1	2	3	4	5	6	7
0	1	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0
2	0	0	0	1	2	5		
3	0	0	0	-1				

$P_{1,(d_1,d_2)}([\text{pt}])$	0	1	2	3	4	5	6	7
0	0	1	0	0	0	0	0	0
1		-1	-1	-1	-1	-1	-1	-1
2	0	0	1	2	5			
3	0	0	0					

$$\begin{aligned}
P_{1,(1,d)}([\text{pt}]) &= -1, \quad \forall d \geq 1 \\
P_{2,(1,d)}([\text{pt}]) &= -2, \quad \forall d \geq 2 \\
P_{3,(0,2)}([\text{pt}]) &= 0^{\star,\dagger}, \\
P_{n,(0,1)}([\text{pt}]) &= 0^\dagger, \quad \forall n \geq 2 \\
P_{n,(0,n)}([\text{pt}]) &= 1, \quad \forall n \geq 1.
\end{aligned}$$

Remark A.1. Denoting the exceptional curve of \mathbb{F}_1 by B , we have $N_{B/X} = \mathcal{O}(-1) \oplus \mathcal{O} \oplus \mathcal{O}(-1)$, which has sections in the direction of L_1 . Therefore $P_1(X, [B])$ is non-proper, which explains the gap in the table for $P_{1,(1,0)}([\text{pt}])$.

For $(S, L_1, L_2) = (\mathbb{F}_1, \mathcal{O}(0, -1), \mathcal{O}(-2, -2))$, we have

$P_{0,(d_1,d_2)}$	0	1	2	3	4
0	1	0	0	0	0
1	0	0	0	0	0
2	0	0	0	1	2
3	0	0	0	-1	

$P_{1,(d_1,d_2)}([\text{pt}])$	0	1	2	3	4
0	0		0	0	0
1		-1	-1	-1	-1
2	0	0	1	2	
3	0	0	0		

$$P_{1,(1,d)}([\text{pt}]) = -1, \quad \forall d \geq 1$$

For $(S, L_1, L_2) = (\mathbb{F}_2, \mathcal{O}(-1, -2), \mathcal{O}(-1, -2))$, we have

$P_{0,(d_1,d_2)}$	0	1	2	3	4	5	6	7
0	1	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0
2	0	0	0	0	1	2	5	

$P_{1,(d_1,d_2)}([\text{pt}])$	0	1	2	3	4	5	6	7
0	0	1	0	0	0	0	0	0
1		1	1	1	1	1	1	1
2	0	0	0	1	2	5		

$$\begin{aligned}
P_{1,(1,d)}([\text{pt}]) &= 1, \quad \forall d \geq 1 \\
P_{2,(1,d)}([\text{pt}]) &= 2, \quad \forall d \geq 2 \\
P_{n,(0,1)}([\text{pt}]) &= 0^\dagger, \quad \forall n \geq 2 \\
P_{n,(0,n)}([\text{pt}]) &= 1, \quad \forall n \geq 1.
\end{aligned}$$

A.2. Local \mathbb{P}^3 . Consider $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$. Let $P_{0,d} := P_{0,d[\ell]}$ and $P_{n,d}([\ell]) := P_{n,d[\ell]}([\ell])$ (for $n > 0$), where $[\ell] \in H_2(\mathbb{P}^3, \mathbb{Z}) \cong H_2(X, \mathbb{Z})$ denotes the class of a line $\ell \subseteq \mathbb{P}^3$ and we also write $[\ell] \in H^4(X, \mathbb{Z})$ for the pull-back of its Poincaré dual from \mathbb{P}^3 to X . Obviously, $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$ is not a local surface so Theorem 4.4 does not apply. All stable pair invariants in this section have been calculated using the vertex formalism of [CK2, CKM] (this is stressed by decorating the invariants with \star). We determined the following values of $P_{0,d}$ and $P_{n,d}([\ell])$.

$d \setminus n$	0	1	2	3	4
1	0^\star	-20^\star	$0^{\star,\dagger}$	$0^{\star,\dagger}$	$0^{\star,\dagger}$
2	0^\star	-820^\star	400^\star		
3	11200^\star	-68060^\star			

$$P_{n,1}([\ell]) = 0, \quad \forall n \geq 2.$$

Remark A.2. For $X = \text{Tot}_{\mathbb{P}^3}(K_{\mathbb{P}^3})$ and all the cases in this table, we have $P_n(\mathbb{P}^3, \beta) \cong P_n(X, \beta)$. This can be deduced from a filtration argument similar to Proposition 3.9 combined with the fact that all degree 2 Cohen-Macaulay curves C on \mathbb{P}^3 satisfy $\chi(\mathcal{O}_C) \geq 1$ (see also [CT2, p. 20]). Therefore, the reasoning of Proposition 4.2 yields

$$(A.1) \quad [P_n(X, \beta)]^{\text{vir}} = (-1)^{\beta \cdot c_1(\mathbb{P}^3) + n} \cdot [P_n(\mathbb{P}^3, \beta)]_{\text{pair}}^{\text{vir}},$$

where $[P_n(\mathbb{P}^3, \beta)]_{\text{pair}}^{\text{vir}}$ is the virtual class of the pair perfect obstruction theory (4.9) on \mathbb{P}^3 (see also [CMT2, Lem. 3.1] in a similar setting). The sign in this formula is a preferred choice of orientation on $P_n(X, \beta)$ as for (1.5). Now we are in the world of “ordinary” perfect obstruction theories and the torus action on \mathbb{P}^3 can be used to apply the Graber-Pandharipande virtual localization formula [GP] to the right hand side of (A.1). Similar to the argument for [CK2, Thm. A.1], it then follows that the invariants in this table (defined by localization on the fixed locus [CK2, CKM]) are equal to the global invariants (4.5). This reasoning also works for the local surface case $(S, L_1, L_2) = (\mathbb{P}^2, \mathcal{O}(-1), \mathcal{O}(-2))$, $d = 2$, $n = 3$, because then all stable pairs are scheme theoretically supported in the threefold $Y = \text{Tot}_S(L_1)$.

REFERENCES

- [AB] M. F. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology 23 no. 1 (1984) 1–28.
- [BS] M. Beltrametti and A. J. Sommese, *On k -spannedness for projective surfaces*, in 1988 L’Aquila Proceedings: Hyperplane sections, Lecture Notes in Math., Vol. 1417 24–51, Springer-Verlag (1990).
- [BJ] D. Borisov and D. Joyce, *Virtual fundamental classes for moduli spaces of sheaves on Calabi-Yau four-folds*, Geom. Topol. 21 (2017) 3231–3311.
- [BBG] P. Bousseau, A. Brini, and M. van Garrel, *Stable maps to Looijenga pairs*, arXiv:2011.08830.
- [Bus] V. Bussi, *Derived symplectic structures in generalized Donaldson-Thomas theory and categorification*, PhD thesis (2014), University of Oxford.
- [Cao] Y. Cao, *Genus zero Gopakumar-Vafa type invariants for Calabi-Yau 4-folds II: Fano 3-folds*, Commun. Contemp. Math. 22 (2020) no. 7 1950060.
- [CGJ] Y. Cao, J. Gross and D. Joyce, *Orientability of moduli spaces of $\text{Spin}(7)$ -instantons and coherent sheaves on Calabi-Yau 4-folds*, Adv. Math. 368 (2020) 107134.
- [CK1] Y. Cao and M. Kool, *Zero-dimensional Donaldson-Thomas invariants of Calabi-Yau 4-folds*, Adv. Math. 338 (2018) 601–648.
- [CK2] Y. Cao and M. Kool, *Curve counting and DT/PT correspondence for Calabi-Yau 4-folds*, Adv. Math. 375 (2020) 107371.
- [CKM] Y. Cao, M. Kool, and S. Monavari, *K-theoretic DT/PT correspondence for toric Calabi-Yau 4-folds*, arXiv:1906.07856.
- [CL1] Y. Cao and N. C. Leung, *Donaldson-Thomas theory for Calabi-Yau 4-folds*, arXiv:1407.7659.
- [CL2] Y. Cao and N. C. Leung, *Orientability for gauge theories on Calabi-Yau manifolds*, Adv. Math. 314 (2017) 48–70.
- [CMT1] Y. Cao, D. Maulik, and Y. Toda, *Genus zero Gopakumar-Vafa type invariants for Calabi-Yau 4-folds*, Adv. Math. 338 (2018) 41–92.

- [CMT2] Y. Cao, D. Maulik, and Y. Toda, *Stable pairs and Gopakumar-Vafa type invariants for Calabi-Yau 4-folds*, arXiv:1902.00003. To appear in JEMS.
- [CT1] Y. Cao and Y. Toda, *Curve counting via stable objects in derived categories of Calabi-Yau 4-folds*, arXiv:1909.04897.
- [CT2] Y. Cao and Y. Toda, *Gopakumar-Vafa type invariants on Calabi-Yau 4-folds via descendent insertions*, Comm. Math. Phys. (2020), <https://doi.org/10.1007/s00220-020-03897-9>
- [CO12] E. Carlsson and A. Okounkov, *Exts and vertex operators*, Duke Math. J. 161 (2012) 1797–1815.
- [CK] H.-L. Chang and Y.-H. Kiem, *Poincaré invariants are Seiberg-Witten invariants*, Geom. Topol. 17 (2013) 1149–1163.
- [CKK] J. Choi, S. Katz, and A. Klemm, *The refined BPS index from stable pair invariants*, Comm. Math. Phys. 328 (2014) 903–954.
- [DKO] M. Dürr, A. Kabanov, and Ch. Okonek, *Poincaré invariants*, Topology 46 (2007) 225–294.
- [ES] G. Ellingsrud and S. A. Strømme, *On a cell decomposition of the Hilbert scheme of points in the plane*, Invent. Math. 91 (1988) 365–370.
- [GGR] M. van Garrel, T. Graber, and H. Ruddat, *Local Gromov-Witten invariants are log invariants*, Adv. Math. 350 (2019) 860–876.
- [GK] L. Göttsche and M. Kool, *Virtual refinements of the Vafa-Witten formula*, Comm. Math. Phys. 376 (2020) 1–49.
- [GP] T. Graber and R. Pandharipande, *Localization of virtual classes*, Invent. Math. 135 (1999) 487–518.
- [IP] E. N. Ionel and T. Parker, *The Gopakumar-Vafa formula for symplectic manifolds*, Ann. of Math. 187 (2018) 1–64.
- [KP] A. Klemm and R. Pandharipande, *Enumerative geometry of Calabi-Yau 4-folds*, Comm. Math. Phys. 281 (2008) 621–653.
- [Koo] M. Kool, *Stable pair invariants of surfaces and Seiberg-Witten invariants*, Quart. Jour. Math. 67 (2016) 365–386.
- [KST] M. Kool, V. Shende, and R. P. Thomas, *A short proof of the Göttsche conjecture*, Geom. Topol. 15 (2011) 397–406.
- [KT1] M. Kool and R. P. Thomas, *Reduced classes and curve counting on surfaces I: theory*, Alg. Geom. 1 (2014) 334–383.
- [KT2] M. Kool and R. P. Thomas, *Reduced classes and curve counting on surfaces II: calculations*, Alg. Geom. 1 (2014) 384–399.
- [MNOP] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory, I*, Compos. Math. 142 (2006) 1263–1285.
- [OT] J. Oh and R. P. Thomas, *Counting sheaves on Calabi-Yau 4-folds, I*, arXiv:2009.05542.
- [PP1] R. Pandharipande and A. Pixton, *Gromov-Witten/pairs descendent correspondence for toric 3-folds*, Geom. Topol. 18 (2014) 2747–2821.
- [PP2] R. Pandharipande and A. Pixton, *Gromov-Witten/Pairs correspondence for the quintic 3-fold*, JAMS 30 (2017) 389–449.
- [PT1] R. Pandharipande and R. P. Thomas, *Curve counting via stable pairs in the derived category*, Invent. Math. 178 (2009) 407–447.
- [PT2] R. Pandharipande and R. P. Thomas, *The 3-fold vertex via stable pairs*, Geom. Topol. 13 (2009) 1835–1876.
- [PT3] R. Pandharipande and R. P. Thomas, *Stable pairs and BPS invariants*, JAMS 23 (2010) 267–297.
- [PTVV] T. Pantev, B. Töen, M. Vaquié, and G. Vezzosi, *Shifted symplectic structures*, Publ. Math. I.H.E.S. 117 (2013) 271–328.
- [LeP] J. Le Potier, *Systèmes cohérents et structures de niveau*, Astérisque 214 (1993).
- [Pre] A. Preygel, *Some remarks on shifted symplectic structures on non-compact mapping spaces*, preprint, see webpage of A. Preygel <http://www.tolypreygel.com/index-research.html>
- [Sat] H. Sato, *The classification of smooth toric weakened Fano 3-folds*, Manuscripta Math. 109 (2002) 73–84.
- [Sie] B. Siebert, *Virtual fundamental classes, global normal cones and Fulton’s canonical classes*, in: Frobenius manifolds, Aspects Math. 36 (2004) 341–358, Vieweg.
- [TT] Y. Tanaka and R. P. Thomas, *Vafa-Witten invariants for projective surfaces I: stable case*, Jour. Alg. Geom. 29 (2020) 603–668.
- [Tod] Y. Toda, *On categorical Donaldson-Thomas theory for local surfaces*, arXiv:1907.09076.

KAVLI INSTITUTE FOR THE PHYSICS AND MATHEMATICS OF THE UNIVERSE (WPI), THE UNIVERSITY OF TOKYO
 INSTITUTES FOR ADVANCED STUDY, THE UNIVERSITY OF TOKYO, KASHIWA, CHIBA 277-8583, JAPAN
Email address: yalong.cao@ipmu.jp

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY, P.O. BOX 80010 3508 TA UTRECHT, THE NETHERLANDS
Email address: m.kool1@uu.nl

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY, P.O. BOX 80010 3508 TA UTRECHT, THE NETHERLANDS
Email address: s.monavari@uu.nl