PRIME NUMBER THEOREM FOR REGULAR TOEPLITZ SUBSHIFTS

KRZYSZTOF FRACZEK, ADAM KANIGOWSKI, AND MARIUSZ LEMAŃCZYK

ABSTRACT. We prove that neither a prime nor an *l*-almost prime number theorem hold in the class of regular Toeplitz subshifts. But, when a quantitative strengthening of the regularity with respect to the periodic structure involving Euler's totient function is assumed, then the two theorems hold.

1. Introduction

Given a topological dynamical system (X,T), where T is a homeomorphism of a compact metric space X, one says that a prime number theorem (PNT) holds for (X,T) if the limit

(1)
$$\lim_{N \to \infty} \frac{1}{\pi(N)} \sum_{p < N} f(T^p x)$$

(p stands always for a prime number) exists for each $x \in X$, an arbitrary $f \in C(X)$ and $\pi(N)$ denotes the number of primes up to N. Then, via the Riesz theorem, for all $f \in C(X)$, we have

(2)
$$\lim_{N \to \infty} \frac{1}{\pi(N)} \sum_{n < N} f(T^p x) = \int_X f \, d\nu_x$$

for a Borel probability measure ν_x on X, where ν_x depends only on $x \in X$.

Let us first consider the cyclic case $X = \mathbb{Z}/k\mathbb{Z}$ and Tx = x + 1. Fix $x \in X$ and notice that (1) indeed holds by the classical prime number theorem in arithmetic progressions, where ν_x is the uniform probability measure on the "coset" $\{a < k : (a, k) = 1\} + x$. Hence, a PNT holds in cyclic (and therefore also in finite) systems.

Consider now the procyclic case, that is, assume we are given an odometer system (H,T) with

$$H = \lim_{t\to\infty} \mathbb{Z}/n_t\mathbb{Z}, \ Tx = x + (1, 1, \ldots)$$

(here $n_t|n_{t+1}$ for $t \ge 0$). In this case, a PNT still holds. Indeed, the space H has a sequence of natural partitions $D^t = (D_0^t, \ldots, D_{n_{t-1}}^t)$, $t \ge 0$, consisting of clopen sets and such that $TD_i^t = D_{i+1 \bmod n_t}^t$. It follows that the sets D_i^t , $i \le n_t - 1$, have the same diameter which goes to 0 as $t \to \infty$. Moreover, it is not hard to see that each character of the group H is constant on the levels of the towers D^t for t sufficiently large. Hence, each $f \in C(H)$ can be approximated **uniformly** by functions which are constant on the levels of the towers D^t and a PNT holds because it does in the finite case.

Our main results concern prime number theorems for extensions of odometers. Recall that odometers are zero entropy topological systems which are minimal (all

²⁰⁰⁰ Mathematics Subject Classification. 37B10, 37A45, 11N05, 11N13.

Key words and phrases. prime number theorem, Toeplitz systems, almost prime numbers, polynomial ergodic theorems.

Research supported by Narodowe Centrum Nauki grant 2019/33/B/ST1/00364.

T-orbits are dense) and uniquely ergodic (there is only one T-invariant measure -Haar measure in this case). Before we describe our results, let us discuss a PNT in the class of uniquely ergodic systems. First, recall that for all such systems (1) holds a.e. with respect to the unique invariant measure [3], [24]. On the other hand, one can easily construct a counterexample to the validity of (1) for all $x \in X$. Indeed, denote by \mathbb{P} the set of prime numbers and consider the left shift S on $\{0,1\}^{\mathbb{Z}}$ and the subshift $(X_{\mathbf{1}_{\mathbb{P}\cup-\mathbb{P}}},S)$ obtained by the orbit closure of the characteristic function $\mathbf{1}_{\mathbb{P}_{\mathcal{V}}-\mathbb{P}}$ of the "symmetrized" primes. It has a unique invariant measure of zero entropy (which is the Dirac measure at the fixed point ... 0.00...) and a PNT fails in it (see e.g. [8] for details). Now, this particular uniquely ergodic model of the one-point system implies paradoxically that each ergodic dynamical system has a uniquely ergodic model $(X,T)^{-1}$ in which a PNT does not hold. To see this, take any uniquely ergodic model (Y, ν, R) of the given measure-theoretic dynamical system. Since the one-point system is (Furstenberg) disjoint with any other system, the product system $(X_{\mathbf{1}_{P\cup P}} \times Y, S \times R)$ is still uniquely ergodic, with the unique invariant measure $\delta_{\dots 0.00\dots} \otimes \nu$. It is not hard to see that the product system is still measure-theoretically isomorphic to the original system. Since the new system has $(X_{\mathbf{1}_{\mathbb{P}\cup-\mathbb{P}}},S)$ as its **topological** factor, a PNT does not hold in $(X_{\mathbf{1}_{\mathbb{P}\cup-\mathbb{P}}}\times Y,S\times R)$. Hence, if we think about a necessary condition for a PNT to hold, it looks reasonable to add the minimality assumption to avoid a problem of "exotic" orbits on which PNT does not hold (we also recall that a uniquely ergodic system has a unique subsystem which is strictly ergodic). However, in this class one can still produce counterexamples to a PNT, see [19] for the first symbolic counterexamples (although their entropy is not determined in [19]), or [13] for non-symbolic counterexamples. On the other hand, we have quite a few classes in which a PNT holds, including systems of algebraic origin [11], [23], symbolic systems [4], [10], [16], [17] or recently [13] in the category of smooth systems, where a PNT has been proved in the class of analytic Anzai skew products. Finding a sufficient dynamical condition for a PNT to hold, postulated a few years ago by P. Sarnak [21] seems to be an important and difficult task in dynamics, however we rather expect the following:

Working Conjecture: Each ergodic and aperiodic³ measure-theoretic dynamical system has a strictly ergodic model in which a PNT fails.

If true, this makes Sarnak's postulate even harder to realize. The present paper should be viewed as introductory steps in trying to understand the conjecture.

A PNT can be reformulated as the existence of a limit of $\frac{1}{N} \sum_{n < N} f(T^n x) \Lambda(n)$, where Λ stands for the von Mangoldt function: $\Lambda(p^{\ell}) = \log p$ for $\ell \ge 1$ and 0 otherwise. Proving dynamical prime number theorems for zero entropy systems is

¹Recall that the Jewett-Kreiger theorem says the following: Suppose (Z, κ, R) is an ergodic measure-theoretic dynamical system. Then there exists a uniquely ergodic (even strictly ergodic, that is, additionally minimal) topological system (Y, S) with the unique invariant measure ν such that (Z, κ, R) and (Y, ν, S) are **measure-theoretically** isomorphic.

²To illustrate this, consider an irrational rotation R_{α} on \mathbb{T} for which a PNT holds because of Vinogradov's theorem (prime "orbits" are equidistributed). However, our observation shows that there is a uniquely ergodic model of R_{α} in which the eigenfunctions are still continuous but a PNT fails, that is, some of the prime "orbits" are not equidistributed.

³The set of periodic points has measure zero.

closely related to Sarnak's Möbius disjointness conjecture [20]:

(3)
$$\lim_{N \to \infty} \frac{1}{N} \sum_{n < N} f(T^n x) \boldsymbol{\mu}(n) = 0$$

for each $x \in X$, $f \in C(X)$ in each **zero entropy** dynamical system (X,T) (μ stands for the Möbius function: $\mu(1) = 1$, $\mu(p_1 \cdot \ldots \cdot p_k) = (-1)^k$ for different primes p_1, \ldots, p_k , and $\mu(n) = 0$ for the remaining $n \in \mathbb{N}$). Here, the class of systems for which we expect the positive answer is precisely defined. In fact, in quite a few cases (see [4], [5], [9]-[11], [16] and [17]) one can observe the following principle: once we can prove Sarnak's conjecture for (X,T) with a "sufficient" speed of convergence to zero in (3) then a PNT holds in (X,T).

With all the above in mind we come back to extensions of odometers that we intend to study. We stay in the **zero entropy** category of systems and we assume **minimality**. Further, we assume that the systems are **almost 1-1 extensions** of odometers.⁴ We also assume that our systems are **symbolic**.⁵ All these natural assumptions determine however a very precise class of topological systems, namely Toeplitz subshifts (X_x, S) , where x is a Toeplitz sequence over a finite alphabet \mathcal{A} , see Section 7 in Downarowicz's survey [7]. That is, $x \in \mathcal{A}^{\mathbb{Z}}$ has the property that for every $a \in \mathbb{Z}$ there is $\ell \in \mathbb{N}$ such that $x(a) = x(a + k\ell)$ for each $k \in \mathbb{Z}$, and K_x is the set of all $k \in \mathbb{Z}$ with the property that all subblocks of $k \in \mathbb{Z}$ also appear in $k \in \mathbb{Z}$. One shows then that there is a sequence $k \in \mathbb{Z}$ then

(4)
$$\bigcup_{t\geq 0} \operatorname{Per}_{n_t}(x) = \mathbb{Z}.$$

Moreover, there is a natural continuous factor map $\pi: X_x \to H$, where H stands for the odometer determined by (n_t) . In fact, we will restrict our attention to so called regular Toeplitz subshifts, whose formal definition is that the density of $\bigcup_{t=0}^M \operatorname{Per}_{n_t}(x)$ goes to 1. Regular Toeplitz subshifts are zero entropy strictly ergodic systems, and measure-theoretically isomorphic to the rotation given by their maximal equicontinuous factors. Although in [7] there are four other equivalent conditions for regularity (see Theorem 13.1 in [7]), we will choose a different path. Since $\pi: X_x \to H$ is a continuous and equivariant surjection,

$$E^t := \pi^{-1}(D^t) = (E_0^t, \dots, E_{n_t-1}^t) \text{ with } E_j^t = \pi^{-1}(D_j^t)$$

is an S-tower of height n_t whose levels are closed (hence clopen). By the minimality of (X_x, S) there is a unique tower with clopen levels and of fixed height. Let us consider a metric on $\mathcal{A}^{\mathbb{Z}}$ inducing the product topology given by

$$d(x,y) = 2^{-\inf\{|n|:x(n)\neq y(n)\}}.$$

The diameters of the levels of towers E^t do not converge to zero, unless x is periodic. Moreover, the diameters of different levels are in general different as the shift S is

⁴If (H,T) is a factor of (X,S) via $\pi: X \to H$, then (X,S) is called an almost 1-1 extension of (H,T) if there is a point $h \in H$ such that $|\pi^{-1}(h)| = 1$; in fact, in this case the set of points with singleton fibers is G_{δ} and dense.

⁵We recall that each zero entropy system has an extension which is symbolic [2], and clearly if a PNT holds for a system, it does for a factor.

not an isometry. Let us consider the diameter of the tower E^t given by:

$$\delta(E^t) := \sum_{0 \le j < n_t} \operatorname{diam}(E_j^t).$$

It is not hard to see (see Appendix A) that the regularity of a Toeplitz sequence is equivalent to

$$\lim_{t \to \infty} \frac{\delta(E^t)}{n_t} = 0.$$

It is also not hard to see that this property does not depend on the choice of (n_t) satisfying (4). We recall that the Möbius disjointness of subshifts given by regular Toeplitz sequences has been proved in [1]. Here are two first results of the paper proved in Section 2 and Section 4, respectively:

Theorem A. A PNT does not hold in the class of minimal almost 1-1 symbolic extensions of odometers satisfying (5). That is, a PNT need not hold in a strictly ergodic subshift determined by a **regular** Toeplitz sequence.

Theorem B. A PNT holds in the class of minimal almost 1-1 symbolic extensions of odometers in which (5) holds with a speed

(6)
$$\lim_{t \to \infty} \frac{\delta(E^t)}{\varphi(n_t)} = 0,$$

where φ denotes the Euler totient function.

As for all Toeplitz dynamical systems constructed in the proof of Theorem A, we have

$$0 < \liminf_{t \to \infty} \frac{\delta(E^t)}{\varphi(n_t)} \leqslant \limsup_{t \to \infty} \frac{\delta(E^t)}{\varphi(n_t)} < +\infty,$$

which shows that the condition (6) in Theorem B is optimal to have a PNT. The systems in Theorem B are strictly ergodic and since they all have non-trivial cyclic factors, the measures ν_y , $y \in X_x$, in (2) are never S-invariant.⁶

We then turn our attention to an l-almost prime number theorem (P_lNT) which is much less explored than the PNT case and which, for the first time in dynamics, is studied in [14] (for some smooth Anzai skew products). Recall that for any $l \ge 1$ a natural number is called an l-almost prime if it is a product of l primes. We denote the set of l-almost prime numbers by \mathbb{P}_l . By \mathbb{P}_l^N we denote the set of l-almost prime numbers $\le N$ and we let $\pi_l(N)$ stand for the cardinality of \mathbb{P}_l^N . A classical result of Landau asserts that

(7)
$$\lim_{N \to \infty} \frac{\pi_l(N)}{\frac{N}{\log N} \frac{(\log \log N)^{l-1}}{(l-1)!}} = 1,$$

see $\S 56$ in [15].

Analogously to the PNT, we say that a topological dynamical system (X, T) satisfies a P_lNT if the limit

$$\lim_{N \to \infty} \frac{1}{\pi_l(N)} \sum_{n \in \mathbb{P}_l^N} f(T^n x)$$

exists for each $x \in X$ and each $f \in C(X)$.

⁶To be compared with the case of Sturmian systems, see Theorem B.1, in which ν_y , $y \in X_{\alpha,\beta}$, are equal to the unique S-invariant measure.

In Section 3 and Section 5 we provide sketches of proofs of the exact analogues of Theorems A and B for a P_lNT for regular Toeplitz subshifts.

In Section 6.1 we prove a new polynomial ergodic theorem:

$$\lim_{N \to \infty} \frac{1}{N} \sum_{n \le N} f(S^{P(n)}x) \text{ exists}$$

for monic polynomials P with positive integer coefficients for all symbolic minimal almost 1-1 extensions of odometers with a modified condition (6). In Section 6.2 we provide a regular Toeplitz subshift which does not satisfy the polynomial ergodic theorem for squares but it satisfies a PNT. We refer again to [19] for the first examples of strictly ergodic systems (of low complexity), where the Birkhoff ergodic averages along squares do not converge.

While Theorem A confirms the Working Conjecture for a subclass of odometers, we have been unable to confirm it for the whole class of odometers. Confirming Working Conjecture for the class of automorphisms with discrete spectrum seems to be the first step toward a possible general statement. In Appendix B, we provide a simple argument showing that a PNT holds for all symbolic models of irrational rotations given by Sturmian sequences. The Sturmian systems are strictly ergodic and are almost 1-1 extensions of irrational rotations.

2. Regular Toeplitz subshifts which do not satisfy PNT (proof of Theorem A)

For all $K, n \in \mathbb{N}$ and $a \in \mathbb{Z}$ let

$$\pi(K; n, a) = \{1 \le p \le K : p \in \mathbb{P}, p = a \mod n\}.$$

Theorem 2.1 (PNT in arithmetic progressions, see [22]). For any natural n and any integer a with (a, n) = 1 we have

$$\lim_{K \to \infty} \frac{\pi(K; n, a)}{\pi(K)} = \frac{1}{\varphi(n)}.$$

We construct a Toeplitz sequence $x \in \{0,1\}^{\mathbb{Z}}$ with the period structure (n_t) :

(8)
$$n_{t+1} = k_{t+1}n_t, (k_{t+1}, n_t) = 1$$

for each $t \ge 1$. We will show that for this x:

$$\lim_{t \to \infty} \frac{1}{\pi(n_t)} \sum_{p < n_t} F(S^p x) \text{ does not exist},$$

where $F(y) = (-1)^{y(0)}$. At stage t, x is approximated by the infinite concatenation of $x_t[0, n_t - 1] \in \{0, 1, ?\}^{n_t}$ (that is, we see a periodic sequence of 0, 1, ? with period n_t). Successive "?" will be filled in the next steps of construction of x. We require that:

(9)
$$\frac{\varphi(n_t)}{n_t} \leqslant \frac{1}{2^t},$$

$$\{0 \leqslant i < n_t : x_t(i) = ?\} \subset \{0 \leqslant j < n_t : (j, n_t) = 1\},\$$

(11)
$$\#\{0 \leqslant i < n_t : x_t(i) = ?\} \geqslant \left(1 - \sum_{l=1}^t \frac{1}{100^l}\right) \varphi(n_t),$$

(12)
$$\#\{p < n_t : x_t(p) = ?\} \geqslant \frac{1}{2}\pi(n_t).$$

We choose k_{t+1} satisfying (8) and:

(13)
$$\frac{\varphi(k_{t+1})}{k_{t+1}} \leqslant \frac{1}{2}, \ ^{7}$$

(15)
$$8\log n_{t+1} \leqslant \pi(n_{t+1}), \quad 8\pi(n_t) \leqslant \pi(n_{t+1})$$

and for each $0 < a < n_t$, $(a, n_t) = 1$, we have

(16)
$$\#(\{a+jn_t: j=0,\ldots,k_{t+1}\} \cap \mathbb{P}) = \pi(n_{t+1};n_t,a) \leq 2\frac{\pi(n_{t+1})}{\varphi(n_t)}.$$

The latter we obtain from Theorem 2.1 (remembering that n_t is fixed, so the number of a is known, we can obtain the accuracy as good as we want by taking k_{t+1} sufficiently large).

We need two simple observations:

(17)
$$\{0 \leqslant k < n_{t+1} : (k, n_{t+1}) = 1\} \subset \bigcup_{\substack{0 \leqslant a < n_t \\ (a, n_t) = 1}} \{a + jn_t : j = 0, \dots, k_{t+1} - 1\}.$$

Lemma 2.2. For every $0 \le a < n_t$ with $(a, n_t) = 1$, we have

$$\#\{0 \le j < k_{t+1} : (a + jn_t, n_{t+1}) = 1\} = \varphi(k_{t+1}).$$

Proof. First note that $(a+jn_t, n_{t+1}) = 1$ iff $(a+jn_t, k_{t+1}) = 1$. Indeed, assume that $(a+jn_t, k_{t+1}) = 1$. If for some prime p we have $p|(a+jn_t)$ and $p|n_{t+1} = n_t k_{t+1}$, then $p|k_{t+1}$. Otherwise, we have $p|n_t$, so p|a. As $(a, n_t) = 1$, this gives a contradiction. Thus $(a+jn_t, k_{t+1}) = 1$ implies $(a+jn_t, n_{t+1}) = 1$. The opposite implication is obvious. Thus

$$\{0 \le j < k_{t+1} : (a+jn_t, n_{t+1}) = 1\} = \{0 \le j < k_{t+1} : (a+jn_t, k_{t+1}) = 1\}.$$

Let us consider the affine map

$$\mathbb{Z}/k_{t+1}\mathbb{Z}\ni j\stackrel{A}{\mapsto}a+jn_t\in\mathbb{Z}/k_{t+1}\mathbb{Z}.$$

If
$$J := \{0 \le \ell < k_{t+1} : (\ell, k_{t+1}) = 1\}$$
 then

$$\{0 \le j < k_{t+1} : (a + jn_t, k_{t+1}) = 1\} = A^{-1}(J).$$

Since $(n_t, k_{t+1}) = 1$, the map A is a bijection. It follows that

$$\#\{0 \le j < k_{t+1} : (a+jn_t, k_{t+1}) = 1\} = \#\{0 \le \ell < k_{t+1} : (\ell, k_{t+1}) = 1\}$$
$$= \varphi(k_{t+1}),$$

which completes the proof. \Box

We need to describe now which and how we fill "?" in $x_{t+1}[0, n_{t+1}-1]$. This block is divided into k_{t+1} subblocks

$$\underbrace{x_t[0, n_t - 1]x_t[0, n_t - 1] \dots x_t[0, n_t - 1]}_{k_{t+1}}.$$

⁷Note that if p_i stand for the *i*-th prime then $\sum_{j\geqslant i}1/p_j=+\infty$, whence remembering that $\varphi(p_ip_{i+1}\dots p_{i+s})=p_ip_{i+1}\dots p_{i+s}\prod_{j=0}^s(1-1/p_{i+j})$, we have $\prod_{j=0}^s(1-1/p_{i+j})\to 0$, and therefore $\prod_{j=0}^s(1-1/p_{i+j})<1/2$ for s large enough.

We fill in **all** "?" in the first block $x_t[0, n_t - 1]$ in such a way to "destroy" a PNT for the time n_t , namely

$$\frac{1}{\pi(n_t)} \sum_{p < n_t} F(S^p x) = \frac{1}{\pi(n_t)} \sum_{\substack{p < n_t \\ p \mid n_t}} (-1)^{x(p)} + \frac{1}{\pi(n_t)} \left(\sum_{\substack{p < n_t \\ (p, n_t) = 1 \\ x_t(p) = 0}} 1 - \sum_{\substack{p < n_t \\ (p, n_t) = 1 \\ x_t(p) = 1}} 1 + \sum_{\substack{p < n_t \\ (p, n_t) = 1 \\ x_t(p) = ?}} (-1)^{x(p)} \right).$$

As the number of the primes dividing n_t is bounded by $\log n_t$, it is negligible compared to $\pi(n_t) = n_t/\log n_t$. It follows that

$$\left| \frac{1}{\pi(n_t)} \sum_{\substack{p < n_t \\ p \mid n_t}} (-1)^{x(p)} \right| \le \frac{\log n_t}{\pi(n_t)} = o(1),$$

so the first summand does not affect the asymptotic of the averages in (1). Since the number of p in the last summand is at least $\frac{1}{2}\pi(n_t)$ in view of (12), we can fill in x_{t+1} at places $\{p < n_t : (p, n_t) = 1, x_t(p) = ?\}$ to obtain the sum completely different that the known number which we had from stage t-1.

We fill in (in an arbitrary way) all remaining places between 0 and $n_t - 1$ and all places $a + jn_t$ for $0 \le j < k_{t+1}$ such that this number is not coprime with n_{t+1} , so that (10) will be satisfied at stage t + 1. We must remember that for certain $0 < a < n_t$ coprime to n_t , $x_t(a)$ was already defined at previous stages, so along the corresponding arithmetic progressions $a + jn_t$, $0 \le j < k_{t+1}$, these places are also filled in previously. On the other hand, if $x_{t+1}(a+jn_t) \ne ?$ (that is, $x_{t+1}(a+jn_t) = 0$ or $x_{t+1}(a+jn_t) = 1$) and $(a+jn_t, n_{t+1}) = 1$ for some $0 < j < k_{t+1}$ then $x_t(a) \ne ?$. In view of (17), it gives

$$\{0 \leqslant i < n_{t+1} : (i, n_{t+1}) = 1, x_{t+1}(i) \neq ?\}$$

$$\subset \{0 < a < n_t : (a, n_t) = 1, x_{t+1}(a) \neq ?\}$$

$$\cup \bigcup_{\substack{0 \leqslant a < n_t \\ (a, n_t) = 1 \\ x_t(a) \neq ?}} \{a + jn_t : 0 < j < k_{t+1}, (a + jn_t, n_{t+1}) = 1\}.$$

By (10), Lemma 2.2, (11) and (14), it follows that

$$\#\{0 \leqslant i < n_{t+1} : (i, n_{t+1}) = 1, x_{t+1}(i) \neq ?\}$$

$$\leqslant \varphi(n_t) + \#\{0 \leqslant a < n_t : (a, n_t) = 1, x_t(a) \neq ?\} \varphi(k_{t+1})$$

$$\leqslant \varphi(n_t) + \left(\sum_{k=1}^t \frac{1}{100^k}\right) \varphi(n_t) \varphi(k_{t+1}) = \left(\frac{1}{\varphi(k_{t+1})} + \sum_{k=1}^t \frac{1}{100^k}\right) \varphi(n_{t+1})$$

$$\leqslant \sum_{k=1}^{t+1} \frac{1}{100^k} \varphi(n_{t+1}) \leqslant \frac{1}{99} \varphi(n_{t+1}).$$

In particular, at stage t + 1, also (11) is satisfied.

Similar arguments show that

$$\{p < n_{t+1} : x_{t+1}(p) \neq ?\} \subset \{p < n_{t+1} : p | n_{t+1}\} \cup \{p < n_t : x_{t+1}(p) \neq ?\}$$

$$\cup \bigcup_{\substack{0 \le a < n_t \\ (a,n_t) = 1 \\ x_t(a) \neq ?}} \{a + jn_t : 0 < j < k_{t+1}, \ a + jn_t \in \mathbb{P}\}.$$

In view of (16), (11) and (15), it follows that

$$\#\{p < n_{t+1} : x_{t+1}(p) \neq ?\}$$

$$\leq \log n_{t+1} + \pi(n_t) + 2\#\{0 \leq a < n_t : (a, n_t) = 1, x_t(a) \neq ?\} \frac{\pi(n_{t+1})}{\varphi(n_t)}$$

$$\leq \log n_{t+1} + \pi(n_t) + \frac{2}{99}\varphi(n_t) \frac{\pi(n_{t+1})}{\varphi(n_t)} \leq \frac{1}{2}\pi(n_{t+1}).$$

Therefore, at stage t + 1, also (12) is satisfied.

Finally, note that

$$\frac{\varphi(n_{t+1})}{n_{t+1}} = \frac{\varphi(n_t)}{n_t} \frac{\varphi(k_{t+1})}{k_{t+1}} \leqslant \frac{\varphi(n_t)}{n_t} \frac{1}{2},$$

so (9) holds and the resulting Toeplitz sequence is regular.

3. Toeplitz subshifts for which a P_tNT does not hold

We now intend to give an example of a (regular) Toeplitz sequence x such that a P_lNT does not hold for the corresponding subshift. In fact,

$$\lim_{t \to \infty} \frac{1}{\pi_l(n_t)} \sum_{p^{(l)} \in \mathbb{P}_l^{n_t}} F(S^{p^{(l)}}x) \text{ does not exist.}$$

For any natural m and $0 \le a < m$, let

$$\pi_l(N; m, a) := \#(\mathbb{P}_l^N \cap (a + m\mathbb{Z})).$$

Lemma 3.1. *If* (a, m) > 1 *then*

(18)
$$\pi_l(N; m, a) = o(\pi_l(N)).$$

If (a, m) = 1 then

(19)
$$\lim_{N \to \infty} \frac{\pi_l(N; m, a)}{\pi_l(N)} = \frac{1}{\varphi(m)}.$$

Proof. The proof is by induction on l. If l = 1 and (a, m) > 1 then $\pi_l(N; m, a) \leq 1$, so (18) holds. If (a, m) = 1 then (19) is given by Theorem 2.1.

Suppose that (18) and (19) are satisfied for all parameters less than some natural number $l \ge 2$. Assume that $(a, m) \in \mathbb{P}_j$ for some $j \ge 1$. If j > l then $\pi_l(N; m, a) = 0$. If $(a, m) \in \mathbb{P}_l$ then $\pi_l(N; m, a) \le 1$, so (18) holds. If $p^{(j)} := (a, m) \in \mathbb{P}_j$ for some $1 \le j < l$ then,

$$\pi_{l}(N; m, a) \leq \pi_{l-j}([N/p^{(j)}]; m/p^{(j)}, a/p^{(j)}) = O(\pi_{l-j}(N))$$

$$= O\left(\frac{N(\log\log N)^{l-j-1}}{\log N}\right) = o\left(\frac{N(\log\log N)^{l-1}}{\log N}\right) = o(\pi_{l}(N)).$$

Now, suppose that (a, m) = 1. Assume that $p_1 \leq p_2 \leq \ldots \leq p_l$ are prime numbers such that $p^{(l)} = p_1 \cdots p_l \leq N$, $p^{(l)} = a \mod m$. Then $p_1 \leq \sqrt[l]{N}$. Since

 $(p_1, m) = 1$, there exists a unique $0 \le a(p_1) < m$ such that $p_1 \cdot a(p_1) = a \mod m$ and $(a(p_1), m) = 1$. Then

$$\pi_l(N; m, a) = \sum_{p_1 \leq \sqrt[l]{N}} \pi_{l-1}([N/p_1]; m, a(p_1)).$$

As $p_1 \leqslant \sqrt[l]{N}$ implies $N/p_1 \geqslant N^{1-1/l}$, by assumption, for every $\varepsilon > 0$ there exists N_{ε} such that for all $N \geqslant N_{\varepsilon}$ and $p_1 \leqslant \sqrt[l]{N}$ with $(p_1, m) = 1$, we have

$$(1-\varepsilon)\frac{\pi_{l-1}([N/p_1])}{\varphi(m)} < \pi_{l-1}([N/p_1]; m, a(p_1)) < (1+\varepsilon)\frac{\pi_{l-1}([N/p_1])}{\varphi(m)}.$$

Since $\pi_l(N) = \sum_{p_1 \leqslant \sqrt[l]{N}} \pi_{l-1}([N/p_1])$, it follows that

$$(1-\varepsilon)\frac{\pi_l(N)}{\varphi(m)} < \pi_l(N; m, a) < (1+\varepsilon)\frac{\pi_l(N)}{\varphi(m)}$$

for every $N \ge N_{\varepsilon}$, so we have (19). \square

Lemma 3.2. For every $l \ge 2$, we have

(20)
$$\#\{p^{(l)} \in \mathbb{P}_l^N : (p^{(l)}, N) > 1\} = o(\pi_l(N)).$$

Proof. Notice that $\#\{p^{(l)} \in \mathbb{P}_l^N : (p^{(l)}, N) > 1\} \leqslant \sum_{p|N} \pi_{l-1}(\frac{N}{p})$. Therefore, using (7),

$$\begin{aligned} \#\{p^{(l)} \in \mathbb{P}_{l}^{N} : (p^{(l)}, N) > 1\} &= O\Big(\sum_{p \mid N} \frac{N/p}{\log(N/p)} \frac{(\log\log(N/p))^{l-2}}{(l-2)!}\Big) \\ &= O\Big(\frac{N}{\log N} \frac{(\log\log(N))^{l-1}}{(l-1)!} \frac{(l-1)}{\log\log N} \sum_{p \mid N} \frac{\log N}{p \log(N/p)}\Big). \end{aligned}$$

So again, by (7), the result will follow by showing that

$$\frac{1}{\log\log N} \sum_{p|N} \frac{\log N}{p\log(N/p)} = o(1).$$

Note that

$$\sum_{p|N} \frac{\log N}{p \log(N/p)} = \sum_{\substack{p|N \\ p \leqslant N^{1/2}}} \frac{\log N}{p \log(N/p)} + \sum_{\substack{p|N \\ p > N^{1/2}}} \frac{\log N}{p \log(N/p)},$$

and that the second term contains at most one prime p. Moreover, as $l \ge 2$ and $(p^{(l)}, N) > 1$, the number N is not prime, so $N/p \ge 2$. Using this, we get

$$\sum_{p|N} \frac{\log N}{p \log(N/p)} \le 2 \sum_{p|N} \frac{1}{p} + \frac{1}{\log 2} \frac{\log N}{N^{1/2}} = O(\log \log \log N),$$

as $\sum_{p|N} \frac{1}{p} = O(\log \log \log N)$, see e.g. [12]. This finishes the proof. \square

Now, we repeat the scheme of the construction from Section 2 almost word for word, although we have to take care how to choose k_{t+1} .

First of all, we require that k_{t+1} is large enough so that

(21)
$$\pi_l(n_{t+1}; n_t, a) \leqslant 2 \frac{\pi_l(n_{t+1})}{\varphi(n_t)} \text{ for every } 0 \leqslant a < n_t \text{ with } (a, n_t) = 1,$$

(22)
$$\sum_{0 \leq a < n_t, (a, n_t) > 1} \pi_l(n_{t+1}; n_t, a) \leq \frac{\varepsilon}{8} \pi_l(n_{t+1}),$$

(23)
$$\#\{p^{(l)} \in \mathbb{P}_l^{n_{t+1}}, (p^{(l)}, n_{t+1}) > 1\} = o(\pi_l(n_{t+1})).$$

The existence of such k_{t+1} is guaranteed by Lemmas 3.1 and 3.2. Next, we replace (12) by

$$\#\{p^{(l)} \in \mathbb{P}_l^{n_t} : x_t(p^{(l)}) = ?\} \geqslant \frac{1}{2}\pi_l(n_t)$$

and requiring (instead of (16)) that for $(a, n_t) = 1$, we have

$$\#(\{a+jn_t: 0 \le j < k_{t+1}\} \cap \mathbb{P}_l) \le 2\frac{\pi_l(n_{t+1})}{\varphi(n_t)},$$

cf. (21). Furthermore, we replace (15) by the requirement that

$$\#\{p^{(l)} \in \mathbb{P}_l^{n_t} : p^{(l)} \equiv a \bmod n_t \text{ with } (a, n_t) > 1\} \leqslant \frac{1}{8} \pi_l(n_{t+1}),$$

cf. (22). To carry over the previous proof, it remains to show that

$$\frac{1}{\pi_l(n_t)} \sum_{p^{(l)} \in \mathbb{P}_l^{n_t}, (p^{(l)}, n_t) > 1} (-1)^{x(p^{(l)})} = o(1).$$

This follows from (23) applied in the previous step of the construction.

4. Regular Toeplitz subshifts which satisfy a PNT (proof of Theorem B)

Let $x \in \mathcal{A}^{\mathbb{Z}}$ be a regular Toeplitz sequence. Then, for every $k \in \mathbb{N}$, there is an n_k -periodic sequence $x_k \in (\mathcal{A} \cup \{?\})^{\mathbb{Z}}$ so that

$$x_k(j) \neq ?$$
 implies $x(j) = x_k(j) = x_l(j)$ for all $l \geqslant k$

and

$$?_k = ?_k(x) := \#\{0 \le j < n_k : x_k(j) = ?\} = o(n_k).$$

For every Toeplitz sequence $x \in \mathcal{A}^{\mathbb{Z}}$ and natural m let us consider a new Toeplitz sequence $x^{(m)} \in (\mathcal{A}^{2m+1})^{\mathbb{Z}}$ given by

$$x^{(m)}(j) = (x(j-m), \dots, x(j+m))$$
 for every $j \in \mathbb{Z}$.

If $(n_t)_{t\geq 1}$ is a periodic structure of x, then it is also a periodic structure of $x^{(m)}$. Moreover,

(24)
$$?_k(x^{(m)}) \le (2m+1)?_k(x) \text{ for every } k \ge 1.$$

Hence, the regularity of x implies the regularity of $x^{(m)}$.

Theorem B follows directly from Lemma A.1 and the following result.

Theorem 4.1. Suppose that (X_x, S) is a Toeplitz system such that

$$?_k = o(\varphi(n_k)).$$

Then (X_x, S) satisfies a PNT.

Proof. To show a PNT for (X_x, S) , it suffices to show that for every continuous $F: X_x \to \mathbb{C}$ and every $\varepsilon > 0$ there exists N_{ε} so that for every $N, M \geqslant N_{\varepsilon}$ and every $r \in \mathbb{Z}$, we have

(25)
$$\left| \frac{1}{\pi(N)} \sum_{p \le N} F(S^{p+r}x) - \frac{1}{\pi(M)} \sum_{p \le M} F(S^{p+r}x) \right| < \varepsilon.$$

Note that the above is stronger than what is needed as it shows that the convergence in (1) is uniform in $y \in X_x$. We first assume that $F: X_x \to \mathbb{R}$ depends only on the zero coordinate, i.e. F(y) = f(y(0)) for some $f: A \to \mathbb{R}$.

Fix $\varepsilon > 0$. Fix also $k \ge 1$ so that

Next choose N_{ε} such that for every $N \geq N_{\varepsilon}$, we have

(27)
$$\left| \pi(N; n_k, a) - \frac{\pi(N)}{\varphi(n_k)} \right| < \frac{\varepsilon}{8} \frac{\pi(N)}{\varphi(n_k)} \text{ for all } a \in \mathbb{Z} \text{ with } (a, n_k) = 1,$$

(28)
$$\#\{p \leqslant N : p|n_k\} \leqslant \log n_k < \frac{\varepsilon}{8}\pi(N).$$

We will show that for all $N \geqslant N_{\varepsilon}$ and $r \in \mathbb{Z}$ we have

(29)
$$\left| \frac{1}{\pi(N)} \sum_{p \leqslant N} F(S^{p+r}x) - \frac{1}{\varphi(n_k)} \sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k) = 1 \\ x_k(a) \neq ?}} F(S^ax) \right| \leqslant \varepsilon ||F||_{\sup},$$

which implies (25).

Recall that $x_k \in (\mathcal{A} \cup \{?\})^{\mathbb{Z}}$ is an n_k -periodic sequence (used to construct x at stage k). If for some $a \in \mathbb{Z}$ we have

$$x_k(a) \neq ?$$

then

$$x(a+j\cdot n_k)=x_k(a)$$
 for every $j\in\mathbb{Z}$.

This implies that if $p \leq N$ and $x_k(p+r \mod n_k) \neq ?$, then

$$(30) F(S^{p+r}x) = F(S^{p+r \mod n_k}x).$$

Note that

$$\#\{p \leqslant N : x_k(p+r \mod n_k) = ?\}$$

$$\leqslant \sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k)=1 \\ x_k(a)=?}} \#\{p \leqslant N : p = a-r \mod n_k\}$$

$$+ \sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k)>1}} \#\{p \leqslant N : p = a-r \mod n_k\}.$$

Assume that $N \ge N_{\varepsilon}$. By (27) and (28), for every integer v with $(v, n_k) = 1$ we have

$$\#\{p \leqslant N : p = v \mod n_k\} = \pi(N; n_k, v) \leqslant (1 + \varepsilon/8) \frac{\pi(N)}{\varphi(n_k)}$$

and

(31)
$$\sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k) > 1}} \#\{p \leqslant N : p = a - r \mod n_k\} \leqslant \#\{p \leqslant N : p | n_k\} < \frac{\varepsilon}{8}\pi(N),$$

where left inequality follows from the fact that if $(a - r, n_k) > 1$ and $p_a = a - r$ mod n_k for a prime p_a , then $(a - r, n_k) = p_a$ and

$$\{p \le N : p = a - r \mod n_k\} = \{p_a\}.$$

It follows that (use also (26))

$$\#\{p \leqslant N : x_k(p+r \mod n_k) = ?\}$$

$$\leqslant \#\{0 \leqslant a < n_k : (a-r,n_k) = 1, x_k(a) = ?\}(1+\varepsilon/8)\frac{\pi(N)}{\varphi(n_k)} + \frac{\varepsilon}{8}\pi(N)$$

$$\leqslant ?_k(1+\varepsilon/8)\frac{\pi(N)}{\varphi(n_k)} + \frac{\varepsilon}{8}\pi(N) \leqslant \frac{\varepsilon}{2}\pi(N).$$

Let

$$P_N := \{ p \leqslant N : x_k(p + r \mod n_k) \neq ? \}.$$

Then by the above, for every $N \ge N_{\varepsilon}$,

(32)
$$\left| \frac{1}{\pi(N)} \sum_{p \leqslant N} F(S^{p+r}x) - \frac{1}{\pi(N)} \sum_{p \in P_N} F(S^{p+r}x) \right| \leqslant \frac{\varepsilon}{2} \|F\|_{\sup}.$$

But by (30).

$$\sum_{p \in P_N} F(S^{p+r}x) = \sum_{\substack{0 \le a < n_k \\ x_k(a) \ne ?}} \sum_{\substack{p \le N \\ p \equiv a - r \mod n_k}} F(S^a x)$$

$$= \sum_{\substack{0 \le a < n_k \\ x_k(a) \ne ?}} F(S^a x) \# \{ p \le N, p = a - r \mod n_k \}.$$

If $(a-r, n_k) = 1$, then again by (27), we have

$$\left| \#\{p \leqslant N, p = a - r \mod n_k\} - \frac{\pi(N)}{\varphi(n_k)} \right| = \left| \pi(N; n_k, a - r) - \frac{\pi(N)}{\varphi(n_k)} \right| < \frac{\varepsilon}{8} \frac{\pi(N)}{\varphi(n_k)}.$$

In view of (31), it follows that

$$\begin{split} & \left| \frac{1}{\pi(N)} \sum_{p \in P_N} F(S^{p+r}x) - \frac{1}{\varphi(n_k)} \sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k) = 1 \\ x_k(a) \neq ?}} F(S^ax) \right| \\ & = \left| \sum_{\substack{0 \leqslant a < n_k \\ x_k(a) \neq ?}} F(S^ax) \frac{\pi(N; n_k, a-r)}{\pi(N)} - \frac{1}{\varphi(n_k)} \sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k) = 1 \\ x_k(a) \neq ?}} F(S^ax) \right| \\ & \leqslant \frac{1}{\pi(N)} \sum_{\substack{0 \leqslant a < n_k \\ (a-r,n_k) = 1 \\ x_k(a) \neq ?}} |F(S^ax)| \left| \pi(N; n_k, a-r) - \frac{\pi(N)}{\varphi(n_k)} \right| + \frac{\varepsilon}{8} \|F\|_{\sup} \\ & \leqslant \|F\|_{\sup} \left(\frac{\varepsilon}{8} \frac{\#\{0 \leqslant a < n_k : x_k(a) \neq ?, (a-r,n_k) = 1\}}{\varphi(n_k)} + \frac{\varepsilon}{8} \right) \leqslant \|F\|_{\sup} \frac{\varepsilon}{2}. \end{split}$$

Together with (32), this gives (29), which completes the proof in the case of F depending only on the zero coordinate.

Now suppose that $F: X_x \to \mathbb{C}$ depends only on finitely many coordinates. Then there exists natural m and $f: \mathcal{A}^{2m+1} \to \mathbb{C}$ such that $F(y) = f(y(-m), \dots, y(m))$ for every $y = (y(k))_{k \in \mathbb{Z}} \in X_x$. Denote by $X_{x^{(m)}} \subset (\mathcal{A}^{2m+1})^{\mathbb{Z}}$ the orbit closure of $x^{(m)} \in (\mathcal{A}^{2m+1})^{\mathbb{Z}}$. Then every $y^{(m)} \in X_{x^{(m)}}$ is of the form $y^{(m)}(k) = (y(k-m), \dots, y(k+m))$ for some $y = (y(k))_{k \in \mathbb{Z}} \in X_x$.

In view of (24), $(X_{x^{(m)}}, S)$ is a regular Toeplitz shift with $?_k(x^{(m)}) = o(\varphi(n_k))$. Let us consider $\bar{F}: X_{x^{(m)}} \to \mathbb{C}$ given by $\bar{F}(y^{(m)}) = f(y^{(m)}(0)) = f(y(-m), \dots, y(m))$ for $y^{(m)} \in X_{x^{(m)}}$. Since \bar{F} depends only on the zero coordinate, by (25) applied to $x^{(m)}$ and the map \bar{F} , for every $\varepsilon > 0$ there exists N_{ε} such that for $N, M \geq N_{\varepsilon}$, we have

$$\left| \frac{1}{\pi(N)} \sum_{p \leqslant N} F(S^{p+r}x) - \frac{1}{\pi(M)} \sum_{p \leqslant M} F(S^{p+r}x) \right|$$

$$= \left| \frac{1}{\pi(N)} \sum_{p \leqslant N} \bar{F}(S^{p+r}x^{(m)}) - \frac{1}{\pi(M)} \sum_{p \leqslant M} \bar{F}(S^{p+r}x^{(m)}) \right| < \varepsilon.$$

Thus (25) holds for every $F: X_x \to \mathbb{C}$ depending only on finitely many coordinates. As the set of such functions is dense in $C(X_x)$, (25) also holds for every $F \in C(X_x)$, which completes the proof. \square

As $\varphi(n) \to \infty$ when $n \to \infty$, we obtain the following result.

Corollary 4.2. If x is Toeplitz for which the sequence $(?_k)$ is bounded then (X_x, S) satisfies a PNT.

5. Toeplitz subshifts for which a P_lNT holds

Theorem 5.1. Suppose that (X_x, S) is a Toeplitz system such that

$$?_k = o(\varphi(n_k)).$$

Then, for every $F \in C(X_x)$ and $y \in X_x$, the limit

$$\lim_{N \to \infty} \frac{1}{\pi_l(N)} \sum_{p^{(l)} \in \mathbb{P}_l^N} F(S^{p^{(l)}}y) \text{ exists.}$$

Proof. The proof proceeds along the same lines as the proof of Theorem 4.1. It relies on the following analogue of (29): for every $\varepsilon > 0$ there exists a natural N_{ε} such that for all $N \geq N_{\varepsilon}$ and $r \in \mathbb{Z}$, we have

(33)
$$\left| \frac{1}{\pi_l(N)} \sum_{p^{(l)} \in \mathbb{P}_l^N} F(S^{p^{(l)} + r} x) - \frac{1}{\varphi(n_k)} \sum_{\substack{0 \le a < n_k \\ (a - r, n_k) = 1 \\ x_k(a) \neq ?}} F(S^a x) \right| \le \varepsilon ||F||_{\sup}.$$

In turn, the proof of (29) is based on only two elements: (27) and (31). Their l-almost prime counterparts follow directly from (19) and (18), respectively. Now, we repeat the arguments of the proof of (29) almost word for word, replacing (27) and (31) by their l-almost prime counterparts. \square

Remark 5.2. In view of (29) and (33), under the assumption $?_k = o(\varphi(n_k))$, we have

$$\lim_{N \to \infty} \frac{1}{\pi_l(N)} \sum_{p^{(l)} \in \mathbb{P}_l^N} F(S^{p^{(l)}} y) = \lim_{N \to \infty} \frac{1}{\pi(N)} \sum_{p < N} F(S^p y)$$

for every $F \in C(X_x)$ and $y \in X_x$, so a PNT and a P_l NT fully coincide for this class of regular Toeplitz systems.

6. Ergodic averages along polynomial times

Let P be a monic polynomial⁸ of degree d > 1 with non-negative integer coefficients. Note that, under these assumptions, $P(\cdot)$ is a strictly increasing function on \mathbb{N} . For every $n \in \mathbb{N}$, let

 $R_n^P := \{0 \leqslant a < n : a = P(m) \mod n \text{ for some } m \in \mathbb{N}\} \text{ and } \psi^P(n) := \#R_n^P.$

For all $N, n \in \mathbb{N}$ and $a \in \mathbb{R}_n^P$, let

$$\rho^{P}(N; n, a) = \#\{1 \le m \le N : P(m) = a \mod n\}.$$

and

$$\rho^{P}(n, a) := \rho^{P}(n; n, a), \quad \rho^{P}(n) := \max_{a \in R_{r}^{P}} \rho^{P}(n; n, a).$$

Lemma 6.1. The function ψ^P is multiplicative, i.e. $\psi^P(n_1n_2) = \psi^P(n_1)\psi^P(n_2)$ if $(n_1, n_2) = 1$. If $a \in \mathbb{Z}/n\mathbb{Z}$, n_1, \ldots, n_k are pairwise coprime and $n = n_1 \cdots n_k$ then $a \in R_n^P$ iff $a_i \in R_{n_i}^P$ for $i = 1, \ldots, k$, where $0 \le a_i < n_i$ is the remainder of a when divided by n_i (that is, $0 \le a_i < n_i$ and $a_i = a \mod n_i$). Moreover,

(34)
$$\rho^{P}(n,a) = \prod_{i=1}^{k} \rho^{P}(n_{i}, a_{i}).$$

Proof. Note that the multiplicativity of ψ^P follows from the second part of the lemma.

Moreover, note that $a \in R_n^P$ iff $a = P(m) \mod n$ for some $0 \le m < n$. Indeed, if $a = P(m) \mod n$ for some $m \in \mathbb{N}$, then $a = P(m') \mod n$, where $0 \le m' < n$ is the remainder of m when divided by n.

If $a \in R_{n_1 \cdots n_k}^P$, i.e. $a = P(m) \mod n_1 \cdots n_k$ for some $0 \le m < n$, then $a_i = a = P(m) = P(m_i) \mod n_i$ for every $i = 1, \dots, k$, where $0 \le m_i < n_i$ is the remainder of m when divided by n_i .

Now, suppose $a \in \mathbb{Z}/n\mathbb{Z}$, $a_i = a \mod n_i$ and $a_i \in R_{n_i}^P$ for $i = 1, \ldots, k$. Then, for every $i = 1, \ldots, k$, there exists $0 \le m_i < n_i$ such that $a_i = P(m_i) \mod n_i$. By the Chinese Remainder Theorem, there exists a unique $0 \le m < n$ such that $m = m_i \mod n_i$ for $i = 1, \ldots, k$. It follows that

$$P(m) = P(m_i) = a_i = a \mod n_i \text{ for all } i = 1, \dots, k.$$

This yields $a = P(m) \mod n_1 \cdots n_k$ and $a \in \mathbb{R}_n^P$.

The argument above also shows (34). \square

Remark 6.2. Note that in the argument above we used the fact that the a_i 's determine a as by the ChRT there exists only one $0 \le a < n$ such that $a = a_i \mod n_i$ for each $i = 1, \ldots, k$.

For any natural n denote by $\omega(n)$ the number of its prime divisors (counted without multiplicities) and by p(n) the product of its prime divisors.

Corollary 6.3. The arithmetic function ρ_P is multiplicative and $\rho^P(n) \leqslant \frac{d^{\omega(n)}}{p(n)} n$.

⁸The leading coefficient of P equals 1. This assumption is only for simplicity. In fact, Theorem 6.8 below is true whenever the set of (non-zero) coefficients of P - P(0) is coprime, see the proof of Corollary 6.3 and the assumptions of Albis theorem in [18].

Proof. The multiplicativity of ρ^P follows directly from (34). By Albis theorem (see Corollary 3 of Theorem 1.23 in [18] ⁹), for any prime number we have $\rho^P(p^n) \leq dp^{n-1}$. This result combined with the multiplicativity of ρ^P gives the required bound of $\rho^P(n)$. \square

Lemma 6.4. For all $n \in \mathbb{N}$, $a \in \mathbb{R}_n^P$ and $N \geqslant P(n)$, we have

$$\rho^{P}(n,a) \left(\frac{P^{-1}(N)}{n} - 1 \right) \le \#\{m \in \mathbb{N} : 1 \le P(m) \le N, P(m) = a \mod n\}$$

$$\le \rho^{P}(n,a) \left(\frac{P^{-1}(N)}{n} + 1 \right).$$

Proof. Let $s := \rho^P(n, a)$ and let $1 \le m_1 < \ldots < m_s \le n$ be all numbers such that $P(m_i) = a \mod n$. Note that a natural number m satisfies $P(m) \le N$ and $P(m) = a \mod n$ iff m = jn + r with $0 \le j \le (P^{-1}(N) - r)/n$ and $0 < r \le n$ satisfies $P(r) = a \mod n$. Thus, $r = m_i$ for some $i = 1, \ldots, s$. It follows that

$$\rho := \#\{m \in \mathbb{N} : 1 \leqslant P(m) \leqslant N, P(m) = a \mod n\}$$
$$= \sum_{i=1}^{s} \left(\left[\frac{P^{-1}(N) - m_i}{n} \right] + 1 \right).$$

Since

$$\frac{P^{-1}(N)}{n} - 1 \leqslant \frac{P^{-1}(N) - m_i}{n} < \left[\frac{P^{-1}(N) - m_i}{n}\right] + 1$$
$$\leqslant \frac{P^{-1}(N) - m_i}{n} + 1 < \frac{P^{-1}(N)}{n} + 1,$$

by summing up, this gives

$$s\left(\frac{P^{-1}(N)}{n} - 1\right) \leqslant \rho \leqslant s\left(\frac{P^{-1}(N)}{n} + 1\right).$$

Remark 6.5. As P is an increasing function, we can apply the above inequalities to P(N) instead of N (as $P(N) \ge N$). Then $P(m) \le P(N)$ iff $m \le N$, and the result of the lemma implies

$$\rho^{P}(n,a)\left(\frac{N}{n}-1\right) \leqslant \rho^{P}(N;n,a) \leqslant \rho^{P}(n,a)\left(\frac{N}{n}+1\right).$$

We now focus on the simplest case when $P(n) = n^2$. We continue to write R for R^P , ψ for ψ^P and ρ for ρ^P . In view of Theorems 1.27 and 1.30 in [18], we have the following result.

Proposition 6.6. For every prime number p > 2, for every $a \in R_{p^N}$, where N = 2n or 2n + 1, we have

$$\rho(p^N, a) = \begin{cases} 2 & \text{if } a = a' \mod p \text{ for } a' \in R_p \setminus \{0\} \\ 2p^r & \text{if } a = p^{2r}a' \text{ and } a' = a'' \mod p \text{ for } a'' \in R_p \setminus \{0\} \\ p^n & \text{if } a = 0. \end{cases}$$

$$\rho^{P}(n; n, a) = \lambda_{P-a}(n), \ \rho^{P}(n) = \max_{a \in R_{-}^{P}} \lambda_{P-a}(n);$$

the estimate on λ_P in [18] depends **only** on the degree of the polynomial.

⁹Note that compared to notation from [18], we have:

Moreover, we have

$$\psi(p^{2n+1}) = \frac{p^{2n+2} + 2p + 1}{2(p+1)} \text{ and } \psi(p^{2n}) = \frac{p^{2n+1} + p + 2}{2(p+1)}.$$

Furthermore, If p=2 then

$$\rho(2, a) = 1$$
 for all $a \in R_2$, $\rho(4, a) = 2$ for all $a \in R_4$

and for any $N \ge 3$, where N = 2n or 2n + 1, for every $a \in R_{2N}$, we have

$$\rho(2^N, a) = \begin{cases} 4 & \text{if } a = 1 \mod 8 \\ 4 \cdot 2^r & \text{if } a = 2^{2r}a', \ 2r \leqslant N - 3, \ a' = 1 \mod 8 \\ 2 \cdot 2^r & \text{if } a = 2^{2r}a', \ 2r = N - 2, \ a' = 1 \mod 4 \\ 2^r & \text{if } a = 2^{2r}a', \ 2r = N - 1, \ a' = 1 \mod 2 \\ 2^n & \text{if } a = 0. \end{cases}$$

Moreover.

$$\psi(2^{2n}) = \frac{2^{2n-1}+4}{3}$$
 and $\psi(2^{2n+1}) = \frac{2^{2n}+5}{3}$.

Corollary 6.7. For every natural $n \ge 2$, we have $\rho(n) \le 4\sqrt{n}$. Moreover, if n is square-free, then $\rho(n) \leq 2^{\omega(n)}$.

Proof. By a direct inspection of the formulas in Proposition 6.6, we obtain:

$$\rho(2^N) \le 2\sqrt{2^N}, \ \rho(3^N) \le 2\sqrt{3^N}$$

but for all $p \ge 5$, we have

$$\rho(p^N) \leqslant \sqrt{p^N}$$

Indeed, for the cases $a = a' \mod p$ (for $a' \in R_p \setminus \{0\}$) and a = 0, it is direct. For the case $\rho(p^N, a) = 2p^r$, we have $a = p^{2r}a' < p^N$, so $2r \le N - 1$ and then indeed $2p^r \leqslant p^{N/2}$.

The second inequality follows directly from $\rho(p) \leq 2$.

For some future purposes, we are interested in cases (in Proposition 6.6) which gives possibly smallest values for the function ρ , hence, for every prime number p and any natural n, let

$$\widetilde{R}_{p^n} := \begin{cases} \{0 \leqslant a < p^n : a = a' \mod p \text{ for } a' \in R_p \setminus \{0\}\} & \text{if } p > 2 \\ R_2 & \text{if } p^n = 2 \\ R_4 & \text{if } p^n = 4 \\ \{0 \leqslant a < 2^n : a = 1 \mod 8\} & \text{if } n \geqslant 3. \end{cases}$$

By Proposition 6.6, $\widetilde{R}_{p^n} \subset R_{p^n}$. Let $n = p_1^{m_1} p_2^{m_2} \cdots p_k^{m_k}$ be the canonical representation of n. Let

$$\Phi: \mathbb{Z}/n\mathbb{Z} \to \mathbb{Z}/p_1^{m_1}\mathbb{Z} \times \ldots \times \mathbb{Z}/p_k^{m_k}\mathbb{Z}$$

$$\psi(p^{2n}) = \frac{p-1}{2}p^{2n-1} + \sum_{r=1}^{n-1} \frac{p-1}{2}p^{2n-2r-1} + 1 = 1 + p\frac{p-1}{2}\sum_{r=0}^{n-1} p^{2(n-r-1)} = 1 + \frac{p(p-1)}{2}\frac{(p^2)^n - 1}{p^2 - 1} = \frac{p^{2n+1} + p + 2}{2(p+1)}.$$

 $^{^{10}}$ We obtain these formulas by using the formulas for the values of σ and counting the number of the possibilites in each row, so for N=2n, we have:

be the canonical ring isomorphism. Recall (cf. Lemma 6.1 and Remark 6.2) that Φ establishes a one-to-one correspondence between R_n and $R_{p_1^{m_1}} \times \ldots \times R_{p_k^{m_k}}$. Set

$$\widetilde{R}_n := \Phi^{-1}(\widetilde{R}_{p_1^{m_1}} \times \ldots \times \widetilde{R}_{p_k^{m_k}})$$

and

$$\widetilde{\psi}(n) := \#\widetilde{R}_n.$$

Then, clearly, $\widetilde{\psi}$ is a multiplicative function. Moreover, by Proposition 6.6, for each $a \in \widetilde{R}_{p^N}$, we have

$$\rho(p^{N}, a) = \begin{cases} 1 & \text{if } p^{N} = 2\\ 2 & \text{if } p^{N} = 2 \text{ or } p > 2\\ 4 & \text{if } p = 2 \text{ and } N \geqslant 3. \end{cases}$$

Hence, in view of (34), for every $a \in \widetilde{R}_n$, we have

(35)
$$\frac{1}{2} \cdot 2^{\omega(n)} \leqslant \rho(n, a) \leqslant 2 \cdot 2^{\omega(n)}.$$

Moreover, by definition,

$$\widetilde{\psi}(p^n) := \begin{cases} p^{n-1} \frac{p-1}{2} & \text{if } p > 2\\ 2 & \text{if } p^n = 2\\ 2 & \text{if } p^n = 4\\ 2^{n-3} & \text{if } p = 2 \text{ and } n \geqslant 3. \end{cases}$$

It follows that

(36)
$$\frac{1}{2} \prod_{p|n} \left(1 - \frac{1}{p} \right) \leqslant \frac{2^{\omega(n)} \widetilde{\psi}(n)}{n} \leqslant 4 \prod_{p|n} \left(1 - \frac{1}{p} \right).$$

(To obtain these inequalities, for $n = p_1^{m_1} p_2^{m_2} \cdots p_k^{m_k}$, write $\frac{2^{\omega(n)} \widetilde{\psi}(n)}{n} = \prod_{i=1}^k \frac{2\widetilde{\psi}(p_i^{m_i})}{p_i^{m_i}}$ and apply the formula above.)

6.1. Polynomial ergodic theorem. In the result below P is a monic polynomial of degree d > 1 with non-negative integer coefficients.

Theorem 6.8. Suppose that (X_x, S) is a Toeplitz system such that

(37)
$$?_k = o(n_k/\rho^P(n_k)).$$

Then, for every continuous map $F: X_x \to \mathbb{C}$ and $y \in X_x$, the limit

(38)
$$\lim_{N \to \infty} \frac{1}{N} \sum_{m \leqslant N} F(S^{P(m)}y)$$

exists.

Proof. To show (38), we need to prove that for every $\varepsilon > 0$ there exists N_{ε} so that for every $N, M \ge N_{\varepsilon}$ and every $r \in \mathbb{Z}$, we have

(39)
$$\left| \frac{1}{N} \sum_{m \le N} F(S^{P(m)+r}x) - \frac{1}{M} \sum_{m \le M} F(S^{P(m)+r}x) \right| < \varepsilon.$$

We first assume that $F: X_x \to \mathbb{R}$ depends only on the zero coordinate, i.e. F(y) = f(y(0)) for some $f: \mathcal{A} \to \mathbb{R}$.

Fix $\varepsilon > 0$. Choose $k \ge 1$ so that

Next, choose $N_{\varepsilon} \geq 8n_k^2/\varepsilon$. Then, in view of Remark 6.5 (and the choice of N_{ε}), for every $N \geq N_{\varepsilon}$ and $a \in R_{n_k}^P$, we have

(41)
$$\left| \rho^P(N; n_k, a) - \rho^P(n_k, a) \frac{N}{n_k} \right| < \rho^P(n_k) \leqslant n_k \leqslant \frac{\varepsilon}{8} \frac{N}{n_k}.$$

From now on, we write that an integer number v belongs to $R_{n_k}^P$ if there exists $0 \le v' < n_k$ such that $v' = v \mod n_k$ and $v' \in R_{n_k}^P$. We will show that for all $N \ge N_{\varepsilon}$ and $r \in \mathbb{Z}$, we have

$$\left| \frac{1}{N} \sum_{m \leqslant N} F(S^{P(m)+r}x) - \frac{1}{n_k} \sum_{\substack{0 \leqslant a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} \rho^P(n_k, a - r) F(S^a x) \right| \leqslant \varepsilon ||F||_{\sup},$$

and this implies (39).

Recall that $x_k \in (\mathcal{A} \cup \{?\})^{\mathbb{Z}}$ is an n_k -periodic sequence (used to construct x at stage k). Note that for every $a \in \mathbb{Z}$, we have

$$x_k(a) \neq ? \Rightarrow x(a+j \cdot n_k) = x_k(a)$$
 for every $j \in \mathbb{Z}$.

This implies that if $m \leq N$ and $x_k(P(m) + r \mod n_k) \neq ?$, then

(43)
$$F(S^{P(m)+r}x) = F(S^{P(m)+r \mod n_k}x).$$

Therefore,

$$\#\{m \leqslant N : x_k(P(m) + r \mod n_k) = ?\}
= \sum_{\substack{0 \leqslant a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) = ?}} \#\{m \leqslant N : P(m) = a - r \mod n_k\} = \sum_{\substack{0 \leqslant a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) = ?}} \rho^P(N; n_k, a - r).$$

Assume that $N \ge N_{\varepsilon}$. By (41), for every integer $v \in R_{n_k}^P$, we have

$$\rho^P(N; n_k, v) \leqslant 2\rho^P(n_k) \frac{N}{n_k}.$$

In view of (40), it follows that

$$\#\{m \le N : x_k(P(m) + r \mod n_k) = ?\}$$

$$\le \#\{0 \le a < n_k : a - r \in R_{n_k}^P, x_k(a) = ?\} 2\rho^P(n_k) \frac{N}{n_k}$$

$$\le 2?_k \rho^P(n_k) \frac{N}{n_k} \le \frac{\varepsilon}{4} N.$$

Let

$$U_N := \{ m \leqslant N : x_k(P(m) + r \mod n_k) \neq ? \}.$$

Then by the above, for every $N \geqslant N_{\varepsilon}$,

$$\left| \frac{1}{N} \sum_{m \leq N} F(S^{P(m)+r}x) - \frac{1}{N} \sum_{m \in U_N} F(S^{P(m)+r}x) \right| \leqslant \frac{\varepsilon}{4} ||F||_{\sup}.$$

But by (43),

$$\sum_{m \in U_N} F(S^{P(m)+r}x) = \sum_{\substack{0 \le a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} \sum_{\substack{m \le N \\ p(m) = a - r \mod n_k}} F(S^a x)$$

$$= \sum_{\substack{0 \le a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} F(S^a x) \#\{m \le N : P(m) = a - r \mod n_k\}$$

$$= \sum_{\substack{0 \le a < n_k \\ x_k(a) \neq ? \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} F(S^a x) \rho^P(N; n_k, a - r).$$

By (41), we have

$$\left| \rho^P(N; n_k, a - r) - \rho^P(n_k, a - r) \frac{N}{n_k} \right| < \frac{\varepsilon}{8} \frac{N}{n_k}.$$

It follows that

$$\begin{split} & \left| \frac{1}{N} \sum_{m \in U_N} F(S^{P(m)+r}x) - \frac{1}{n_k} \sum_{\substack{0 \leq a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} \rho^P(n_k, a - r) F(S^a x) \right| \\ & = \left| \frac{1}{N} \sum_{\substack{0 \leq a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} F(S^a x) \rho^P(N; n_k, a - r) - \frac{1}{n_k} \sum_{\substack{0 \leq a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} \rho^P(n_k, a - r) F(S^a x) \right| \\ & \leq \frac{1}{N} \sum_{\substack{0 \leq a < n_k \\ a - r \in R_{n_k}^P \\ x_k(a) \neq ?}} |F(S^a x)| \left| \rho^P(N; n_k, a - r) - \rho^P(n_k, a - r) \frac{N}{n_k} \right| \\ & \leq \|F\|_{\sup} \frac{\varepsilon}{8} \frac{\#\{0 \leq a < n_k : x_k(a) \neq ?, a - r \in R_{n_k}^P\}}{n_k} \leq \|F\|_{\sup} \frac{\varepsilon}{8}. \end{split}$$

Together with (44), this gives (42), which completes the proof in the case of F depending only on the zero coordinate. The rest of the proof runs as in the proof of Theorem 4.1, this is by passing to the Toeplitz sequences $x^{(m)} \in (\mathcal{A}^{2m+1})^{\mathbb{Z}}$ for $m \ge 1$. \square

Remark 6.9. Denote by $\mathbb{P}_{(n_t)}$ the set of all prime divisors of elements of the sequence $(n_t)_{t\geq 1}$. In view of Corollary 6.3, $?_t = o(p(n_t)/d^{\omega(n_t)})$ implies (37). Unfortunately, if $\mathbb{P}_{(n_t)}$ is finite then the sequence $(p(n_t)/d^{\omega(n_t)})_{t\geq 1}$ is bounded, so Theorem 6.8, in the way, is not applicable. Fortunately, if $\mathbb{P}_{(n_t)}$ is infinite then $p(n_t)/d^{\omega(n_t)} \to +\infty$ as $t \to +\infty$, so Theorem 6.8 applies to a non-trivial class of regular Toeplitz shifts, in particular, it applies when the periodic sequences x_t defining x have a bounded number of "?".

However, Theorem 6.8 applies to a much wider class of regular Toeplitz shifts when $P(n) = n^2$. Then, by Corollary 6.7, $?_t = o(\sqrt{n_t})$ implies (37). Here the finiteness or the infinity of the set $\mathbb{P}_{(n_t)}$ does not matter.

The assumption (37) about the growth of the sequence $(?_t)_{t\geqslant 1}$ is the least restrictive when all n_t are square-free. Then, by the second part of Corollary 6.7, $?_t = o(n_t/2^{\omega(n_t)})$ implies (37). Therefore, $?_t = O(n_t^{1-\frac{1}{\log_2\log_2 n_t}})$ also implies (37). Indeed, it suffices to show that $2^{\omega(n)} = o(n^{\frac{1}{\log_2\log_2 n}})$ for square-free numbers $n \to +\infty$. Suppose that $\omega(n) = k$ and denote by $(p_l)_{l\geqslant 1}$ the increasing sequence of all prime numbers. Since

$$\ln n \geqslant \sum_{l=1}^{k} \ln p_l \geqslant k \ln k,$$

we have

$$\frac{2^{\omega(n)}}{n^{\frac{1}{\log_2\log_2 n}}} = \frac{2^k}{2^{\frac{\log_2 n}{\log_2\log_2 n}}} \leqslant \frac{2^k}{2^{\frac{k\log_2 k}{\log_2(k\log_2 k)}}} = \frac{1}{2^{\frac{k\log_2\log_2 k}{\log_2 k + \log_2\log_2 k}}}.$$

As $\frac{k \log_2 \log_2 k}{\log_2 k + \log_2 \log_2 k} \to +\infty$ when $k \to +\infty$, this gives $2^{\omega(n)} = o(n^{\frac{1}{\log_2 \log_2 n}})$.

6.2. Counter-examples. We will show that there exists a regular Toeplitz sequence $x \in \{0, 1\}^{\mathbb{Z}}$ with the period structure $(n_t)_{t \ge 1}$ satisfying

(45)
$$n_{t+1} = k_{t+1}n_k \text{ with } (k_{t+1}, n_t) = 1, \ n_{t+1} \ge 2^4 n_t^2 \text{ and } \sum_{p \in \mathbb{P}_{(n_t)}} \frac{1}{p} < +\infty$$

and such that

$$\lim_{t \to \infty} \frac{1}{\sqrt{n_t}} \sum_{0 \le m < \sqrt{n_t}} F(S^{m^2} x) \text{ does not exist,}$$

where $F(y) = (-1)^{y(0)}$. Let

$$0<\beta:=\frac{1}{16}\prod_{p\in\mathbb{P}_{(n_t)}}\frac{p-1}{p}.$$

By (36), for every $t \ge 1$, we have

(46)
$$\frac{2^{\omega(n_t)}\widetilde{\psi}(n_t)}{n_t} \geqslant 8\beta.$$

Passing to a subsequence of $(n_t)_{t\geq 1}$ (and remembering that $\widetilde{\psi}(m) \to \infty$ when $m \to \infty$), we can assume that

$$\sum_{t>1} \frac{1}{\widetilde{\psi}(k_t)} \leqslant \frac{1}{2}.$$

Set

$$\gamma_t := \sum_{l=1}^t \frac{1}{\widetilde{\psi}(k_l)} \ (\leqslant \frac{1}{2}).$$

At stage t, x is approximated by the infinite concatenation of $x_t[0, n_t-1] \in \{0, 1, ?\}^{n_t}$ (that is, we see a periodic sequence of 0, 1, ? with period n_t). Successive "?" will be filled in the next steps of construction of x. We require that:

$$\{0 \le i < n_t : x_t(i) = ?\} \subset R_{n_t};$$

(48)
$$\#\{a \in \widetilde{R}_{n_t} : x_t(a) = ?\} \geqslant (1 - \gamma_t)\widetilde{\psi}(n_t);$$

(49)
$$\#\{0 \le m < \sqrt{n_t} : x_t(m^2) = ?\} \ge \beta \sqrt{n_t}.$$

Recall that, in view of Lemma 6.4 (remembering that $P^{-1}(n_{t+1}) = \sqrt{n_{t+1}}$), (35) and (45), for each $a \in \widetilde{R}_{n_t}$, we have¹¹

$$\#(\{a+jn_{t}: 0 < j < k_{t+1}\} \cap \mathbb{N}^{2}) \geqslant \#(\{m^{2} = a \mod n_{t}: m^{2} < n_{t+1}\}) - 1$$

$$\geqslant \left(\frac{\sqrt{n_{t+1}}}{n_{t}} - 1\right)\rho(n_{t}, a) - 1 \geqslant \left(\frac{\sqrt{n_{t+1}}}{n_{t}} - 1\right)\frac{1}{2}2^{\omega(n_{t})} - 1$$

$$\geqslant \frac{1}{2}2^{\omega(n_{t})}\left(\frac{\sqrt{n_{t+1}}}{n_{t}} - 2\right) \geqslant \frac{1}{4}2^{\omega(n_{t})}\frac{\sqrt{n_{t+1}}}{n_{t}},$$

SO

(50)
$$\#(\{a+jn_t: 0 < j < k_{t+1}\} \cap \mathbb{N}^2) \geqslant \frac{2^{\omega(n_t)}}{4} \frac{\sqrt{n_{t+1}}}{n_t}.$$

By the definition of the sets R_n and \widetilde{R}_n , we have

(51)
$$R_{n_{t+1}} \subset \bigcup_{a \in R_{n_t}} \{a + jn_t : 0 \le j < k_{t+1}\},$$

(52)
$$\widetilde{R}_{n_{t+1}} \subset \bigcup_{a \in \widetilde{R}_{n_t}} \{a + jn_t : 0 \leqslant j < k_{t+1}\}.$$

Moreover, by Lemma 6.1, for every $a \in \widetilde{R}_{n_t}$, we have

(53)
$$\#\{i \in \widetilde{R}_{n_{t+1}} : i = a \mod n_t\} = \#\widetilde{R}_{k_{t+1}} = \widetilde{\psi}(k_{t+1}).$$

We need to describe now which and how we fill "?" in $x_{t+1}[0, n_{t+1}-1]$. This block is divided into k_{t+1} subblocks

$$\underbrace{x_t[0, n_t - 1]x_t[0, n_t - 1] \dots x_t[0, n_t - 1]}_{k_{t+1}}$$
.

We fill in **all** "?" in the first block $x_t[0, n_t - 1]$ in such a way to "destroy" the convergence of averages in (45) for the time n_t , namely

$$\frac{1}{\sqrt{n_t}} \sum_{0 \leqslant m < \sqrt{n_t}} F(S^{m^2} x) = \frac{1}{\sqrt{n_t}} \left(\sum_{\substack{m < \sqrt{n_t} \\ x_t(m^2) = 0}} 1 - \sum_{\substack{m < \sqrt{n_t} \\ x_t(m^2) = 1}} 1 + \sum_{\substack{m < \sqrt{n_t} \\ x_t(m^2) = ?}} (-1)^{x(m^2)} \right).$$

And, since the number of m in the last summand is at least $\beta\sqrt{n_t}$ in view of (49), we can fill in these places at stage t+1 to obtain the sum completely different that the known number which we had from stage t-1. We also fill in (in an arbitrary way) the remaining places in $\{0, \ldots, n_t-1\}$.

We fill in (in an arbitrary way) all places in $\{n_t, \ldots, n_{t+1} - 1\} \setminus R_{n_{t+1}}$ and only these places, so that (47) will be satisfied at stage t + 1.

We must remember that for any $a \in R_{n_t}$ if $x_t(a) \neq ?$ then for every $0 \leq j < k_{t+1}$, we have $x_{t+1}(a+jn_t) = x_t(a+jn_t) = x_t(a) \neq ?$. Moreover, for any $a \in \widetilde{R}_{n_t}$ if $x_t(a) = ?$ then for every $0 < j < k_{t+1}$ with $a + jn_t \in \widetilde{R}_{n_{t+1}}$ we have $x_{t+1}(a+jn_t) = ?$. In view of (52), this gives

$$\#\{i \in \widetilde{R}_{n_{t+1}} : x_{t+1}(i) \neq ?\}$$

$$\leq \widetilde{\psi}(n_t) + \sum_{a \in \widetilde{R}_{n_t} : x_t(a) \neq ?} \#\{a + jn_t \in \widetilde{R}_{n_{t+1}} : 0 < j < k_{t+1}\}.$$

¹¹ \mathbb{N}^2 stands for $\{m^2 : m \ge 0\}$.

In view of (53) and (48), it follows that

$$\#\{i \in \widetilde{R}_{n_{t+1}} : x_{t+1}(i) \neq ?\} \leqslant \widetilde{\psi}(n_t) + (\widetilde{\psi}(k_{t+1}) - 1) \#\{a \in \widetilde{R}_{n_t} : x_t(a) \neq ?\}$$

$$\leqslant \widetilde{\psi}(n_t) + (\widetilde{\psi}(k_{t+1}) - 1) \gamma_t \widetilde{\psi}(n_t) = \left(\gamma_t + \frac{1 - \gamma_t}{\widetilde{\psi}(k_{t+1})}\right) \widetilde{\psi}(n_{t+1}) \leqslant \gamma_{t+1} \widetilde{\psi}(n_{t+1}).$$

Therefore, at stage t + 1, also (48) is satisfied.

A similar argument combined with (50), (48) and (46) shows that

$$\#\{0 \leqslant m^{2} < n_{t+1} : x_{t+1}(m^{2}) =?\} = \#\{i \in R_{n_{t+1}} \cap \mathbb{N}^{2} : x_{t+1}(i) =?\}$$

$$\geqslant \sum_{a \in R_{n_{t}} : x_{t}(a) =?} \#\{a + jn_{t} \in R_{n_{t+1}} \cap \mathbb{N}^{2} : 0 < j < k_{t+1}\}$$

$$\geqslant \sum_{a \in \widetilde{R}_{n_{t}} : x_{t}(a) =?} \frac{2^{\omega(n_{t})}}{4} \frac{\sqrt{n_{t+1}}}{n_{t}} = \frac{\sqrt{n_{t+1}}}{4n_{t}} 2^{\omega(n_{t})} \#\{a \in \widetilde{R}_{n_{t}} : x_{t}(a) =?\}$$

$$= (1 - \gamma_{t}) \frac{\sqrt{n_{t+1}}}{4n_{t}} 2^{\omega(n_{t})} \widetilde{\psi}(n_{t}) \geqslant \beta \sqrt{n_{t+1}}.$$

Therefore, at stage t + 1, also (49) is satisfied. This completes the construction.

Remark 6.10. In view of (47), in the constructed example of Toeplitz system (X_x, S) we have $?_t \le \psi(n_t)$. Moreover, $\psi(n_t) = o(\varphi(n_t))$. Indeed, by Proposition 6.6, for every prime number p we have $\psi(p^n) \le p^{n-1}\frac{p+2}{2}$. It follows that

$$\frac{\psi(p^n)}{\varphi(p^n)} \leqslant \frac{1}{2} \cdot \frac{p+2}{p-1} \leqslant \frac{3}{4}$$

for all prime $p \ge 7$. It follows that

$$\frac{\psi(n_t)}{\varphi(n_t)} = O\left(\left(\frac{3}{4}\right)^{\omega(n_t)}\right) = o(1).$$

Consequently, we have $?_t = o(\varphi(n_t))$. Therefore, in view of Theorem 4.1, (X_x, S) satisfies a PNT.

APPENDIX A. THE DIAMETER OF A TOWER

Let $x \in \mathcal{A}^{\mathbb{Z}}$ be a Toeplitz sequence with the periodic structure given by $(n_t)_{t \geq 1}$. Recall that

$$\operatorname{Per}_{n_t}(x) = \{ a \in \mathbb{Z} : x(a + jn_t) = x(a) \text{ for all } j \in \mathbb{Z} \}.$$

Let $\operatorname{Aper}_{n_t}(x) := \mathbb{Z} \backslash \operatorname{Per}_{n_t}(x)$. Then, we define the periodic sequence $x_t \in (\mathcal{A} \cup \{?\})^{\mathbb{Z}}$ by: $x_t(k) = x(k)$ if $k \in \operatorname{Per}_{n_t}(x)$ and $x_t(k) = ?$ if $k \in \operatorname{Aper}_{n_t}(x)$. Note that the density of the set $\operatorname{Aper}_{n_t}(x)$ is equal to $\frac{?_t}{n_t}$, where

$$?_t = \#\{0 \le k < n_t : x_t(k) = ?\} = \#(\operatorname{Aper}_{n_t}(x) \cap \{0, 1, \dots, n_t - 1\}).$$

It follows that the regularity of (X_x, S) is equivalent to $?_t = o(n_t)$.

Lemma A.1. For any Toeplitz sequence $x \in A^{\mathbb{Z}}$ we have

$$?_t \leq \delta(E^t) \leq 3?_t \text{ for every } t \geq 1.$$

Proof. First note that for every $0 \le j < n_t$ we have

$$E_j^t = \{ y \in X_x : y(k-j) = x(k) = x_t(k) \text{ for all } k \in Per_{n_t} \}.$$

Moreover, if $k \in \operatorname{Aper}_{n_t}(x)$ then we can find $y, z \in E_j^t$, so that $y(k-j) \neq z(k-j)$. It follows that

$$diam(E_i^t) = 2^{-\inf\{|n|: n \in Aper_{n_t}(x) - \{j\}\}}.$$

Suppose that

$$\operatorname{Aper}_{n_t}(x) \cap \{0, 1, \dots, n_t - 1\} = \{l_1, l_2, \dots, l_s\}$$

with $1 \leq l_1 \leq \ldots \leq l_s \leq n_t$ and $s = ?_t$. Thus, $\operatorname{diam}(E_{l_i}^t) = 1$ and if $l_{i-1} < j < l_i$ $(l_0 = l_s - n_t \text{ and } l_{s+1} = l_1 + n_t)$ then $\operatorname{diam}(E_j^t) = 2^{-\min\{j - l_{i-1}, l_i - j\}}$. Therefore,

$$\delta(E^t) = \sum_{0 \le j < n_t} \operatorname{diam}(E_j^t) \geqslant \sum_{i=1}^s \operatorname{diam}(E_{l_i}^t) = s$$

and

$$\delta(E^t) = \sum_{0 \le j < n_t} \operatorname{diam}(E_j^t) = \sum_{i=1}^s \sum_{\frac{l_{i-1} + l_i}{2} \le j < \frac{l_i + l_{i+1}}{2}} \operatorname{diam}(E_j^t)$$
$$= \sum_{i=1}^s \left(1 + \sum_{1 \le j < \frac{l_{i+1} - l_i}{2}} 2^{-j} + \sum_{1 \le j \le \frac{l_i - l_{i-1}}{2}} 2^{-j} \right) \le 3s,$$

which completes the proof. \Box

As the regularity of x is equivalent to $?_t = o(n_t)$, we have the following conclusion.

Corollary A.2. A Toeplitz sequence is regular if and only if $\delta(E^t) = o(n_t)$.

APPENDIX B. STURMIAN DYNAMICAL SYSTEMS SATISFY A PNT

Let $T: \mathbb{T} \to \mathbb{T}$ ($\mathbb{T} := \mathbb{R}/\mathbb{Z}$) be an irrational rotation on \mathbb{T} by α . For every non-zero $\beta \in \mathbb{T}$ let $\{A_0, A_1\}$ be the partition given by the intervals $A_0 = [0, \beta)$ and $A_1 = [\beta, 1)$. For every $x \in \mathbb{T}$ denote by $\bar{x} \in \{0, 1\}^{\mathbb{Z}}$ the code of x defined by $\bar{x}(k) = i$ if and only if $T^k x \in A_i$. Finally, denote by $X_{\alpha,\beta} \subset \{0, 1\}^{\mathbb{Z}}$ the closure of the set $\{\bar{x} \in \{0, 1\}^{\mathbb{Z}} : x \in \mathbb{T}\}$. Since $X_{\alpha,\beta}$ is an invariant subset for the left shift S on $\{0, 1\}^{\mathbb{Z}}$, we can focus the topological dynamical system $S: X_{\alpha,\beta} \to X_{\alpha,\beta}$.

Theorem B.1. For the topological dynamical system $S: X_{\alpha,\beta} \to X_{\alpha,\beta}$ a PNT holds.

Proof. For every $y=(y(n))_{n\in\mathbb{Z}}\in X_{\alpha,\beta}$ the set $\bigcap_{n\in\mathbb{Z}}\overline{A}_{y(n)}\subset\mathbb{T}$ has exactly one element $\pi(y)\in\mathbb{T}$. Moreover, $\pi:X_{\alpha,\beta}\to\mathbb{T}$ is a continuous map intertwining S and T and there exists a unique S-invariant probability measure μ on $X_{\alpha,\beta}$. The π -image of μ coincides with Lebesgue measure on \mathbb{T} .

By Vinogradov's theorem, for any character $f(x) = e^{2\pi i n x}$, $n \in \mathbb{Z}$, we have

(54)
$$\lim_{N \to \infty} \frac{1}{\pi(N)} \sum_{p < N} f(T^p x) = \int_{\mathbb{T}} f(x) \, dx \text{ for every } x \in \mathbb{T}.$$

Since every continuous function $f: \mathbb{T} \to \mathbb{C}$ is uniformly approximated by trigonometric polynomials, (54) holds also for any continuous f. Moreover, (54) holds for any Riemann integrable $f: \mathbb{T} \to \mathbb{R}$. Indeed, for every $\varepsilon > 0$ there are two continuous functions $f_-, f_+: \mathbb{T} \to \mathbb{R}$ such that $f_-(x) \leq f(x) \leq f_+(x)$ for every $x \in \mathbb{T}$ and

 $\int_{\mathbb{T}} (f_{+}(x) - f_{-}(x)) dx < \varepsilon$. It follows that

$$\limsup_{N \to \infty} \frac{1}{\pi(N)} \sum_{p < N} f(T^p x) \leqslant \lim_{N \to \infty} \frac{1}{\pi(N)} \sum_{p < N} f_+(T^p x)$$
$$= \int_{\mathbb{T}} f_+(x) \, dx < \int_{\mathbb{T}} f(x) \, dx + \varepsilon$$

and

$$\lim_{N \to \infty} \inf \frac{1}{\pi(N)} \sum_{p < N} f(T^p x) \geqslant \lim_{N \to \infty} \frac{1}{\pi(N)} \sum_{p < N} f_-(T^p x)$$

$$= \int_{\mathbb{T}} f_-(x) \, dx > \int_{\mathbb{T}} f(x) \, dx - \varepsilon.$$

As $\varepsilon > 0$ can be chosen freely, this gives (54).

Suppose that $f: X_{\alpha,\beta} \to \mathbb{R}$ depends only on finitely many coordinates. More precisely, assume that $f(y) = g(y(-n), \dots, y(n))$ for some $g: \{0, 1\}^{2n+1} \to \mathbb{R}$. Then there exists $F: \mathbb{T} \to \mathbb{R}$ such that $f = F \circ \pi$ and F is constant on the atoms of the partition $\bigvee_{i=-n}^n T^{-i}\{A_0, A_1\}$ (for example, if n = 0 and f is the characteristic function of $\{y \in X_{\alpha,\beta}: y(0) = 0\}$ then F is $\mathbf{1}_{A_0}$). It follows that F is Riemann integrable. Therefore, for every $y \in X_{\alpha,\beta}$, we have

$$\frac{1}{\pi(N)} \sum_{p < N} f(S^p y) = \frac{1}{\pi(N)} \sum_{p < N} F(T^p \pi(y)) \to \int_{\mathbb{T}} F(x) dx = \int_{X_{\alpha, \beta}} f \, d\mu.$$

Since every continuous function $f: X_{\alpha,\beta} \to \mathbb{R}$ is uniformly approximated by functions depending on finitely many coordinates,

$$\frac{1}{\pi(N)} \sum_{p < N} f(S^p y) \to \int_{X_{\alpha,\beta}} f \, d\mu \text{ for any } y \in X_{\alpha,\beta}$$

holds for every continuous f. \square

Acknowledgements. We would like to thank the anonymous referee for suggestions to improve the paper.

References

- [1] H. El Abdalaoui, M. Lemańczyk, S. Kasjan, 0-1 sequences of the Thue-Morse type and Sarnak's conjecture, Proc. Amer. Math. Soc. 144 (2016), 161-176.
- [2] M. Boyle, D. Fiebig, U. Fiebig, Residual entropy, conditional entropy and subshift covers, Forum Math. 14 (2002), 713-757.
- [3] J. Bourgain, An approach to pointwise ergodic theorems, In Geometric aspects of functional analysis (1986/87), volume 1317 of Lecture Notes in Math., pages 204-223. Springer, Berlin, 1988.
- [4] J. Bourgain, Möbius-Walsh correlation bounds and an estimate of Mauduit and Rivat, J. Anal. Math. 119 (2013), 147-163.
- [5] J. Bourgain, On the correlation of the Möbius function with rank-one systems, J. Anal. Math. 120 (2013), 105–130.
- [6] P.G.L. Dirichlet, Lectures on number theory. Supplements by R. Dedekind. Translated from the 1863 German original and with an introduction by John Stillwell. History of Mathematics, 16. American Mathematical Society, Providence, RI; London Mathematical Society, London, 1999. xx+275 pp.
- [7] T. Downarowicz, Survey of odometers and Toeplitz flows. Algebraic and topological dynamics, 7–37, Contemp. Math., 385, Amer. Math. Soc., Providence, RI, 2005.

- [8] S. Ferenczi, J. Kułaga-Przymus, M. Lemańczyk, Sarnak's Conjecture what's new, in: Ergodic Theory and Dynamical Systems in their Interactions with Arithmetics and Combinatorics, CIRM Jean-Morlet Chair, Fall 2016, Editors: S. Ferenczi, J. Kułaga-Przymus, M. Lemańczyk, Lecture Notes in Mathematics 2213, Springer International Publishing, pp. 418.
- [9] S. Ferenczi, C. Mauduit, On Sarnak's conjecture and Veech's question for interval exchanges,
 J. Anal. Math. 134 (2018), 545–573.
- [10] B. Green, On (not) computing the Möbius function using bounded depth circuits, Combin. Probab. Comput. 21 (2012), 942-951.
- [11] B. Green, T. Tao, The Möbius function is strongly orthogonal to nilsequences, Annals of Math. (2), 175 (2012), 541-566.
- [12] A. Kanigowski, M. Lemańczyk, M. Radziwiłł, Rigidity in dynamics and Möbius disjointness, arXiv:1905.13256 (submitted).
- [13] A. Kanigowski, M. Lemańczyk, M. Radziwiłł, *Prime number theorem for analytic skew products*, arXiv:2004.01125 (submitted).
- [14] A. Kanigowski, M. Lemańczyk, M. Radziwiłł, Semiprime number theorem for smooth Anzai skew products, in preparation.
- [15] E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen.* 2 Bände. (German) 2d ed. With an appendix by Paul T. Bateman. Chelsea Publishing Co., New York, 1953.
- [16] C. Mauduit, J. Rivat, Prime numbers along Rudin-Shapiro sequences, J. Eur. Math. Soc. 17 (2015), 2595–2642.
- [17] C. Müllner, Automatic sequences fulfill the Sarnak conjecture, Duke Math. J. 166 (2017), 3219-3290.
- [18] W. Narkiewicz, *Number theory*. Translated from the Polish by S. Kanemitsu. World Scientific Publishing Co., Singapore; distributed by Heyden & Son, Inc., Philadelphia, PA, 1983. xii+371 pp.
- [19] R. Pavlov, Some counterexamples in topological dynamics, Ergodic Theory Dynam. Systems 28 (2008), 1291-1322.
- [20] P. Sarnak, Three lectures on the Möbius function, randomness and dynamics, http://publications.ias.edu/sarnak/.
- [21] P. Sarnak, Möbius randomness and Dynamics six years later at CIRM at 1h 08 minute https://library.cirm-math.fr/Record.htm?idlist=1&record=19282918124910001909
- [22] A. Selberg, An elementary proof of the prime-number theorem for arithmetic progressions, Canad. J. Math. 2 (1950), 66-78.
- [23] I.M. Vinogradov, The method of trigonometrical sums in the theory of numbers, (Russian) Trav. Inst. Math. Stekloff 23, (1947). 109 pp.
- [24] M. Wierdl, Pointwise ergodic theorem along the prime numbers, Israel J. Math. **64** (1989), 315-336.

Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, ul. Chopina $12/18,\,87\text{-}100$ Toruń, Poland

Email address: fraczek@mat.umk.pl

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, 4176 CAMPUS DRIVE, WILLIAM E. KIRWAN HALL, COLLEGE PARK, MD 20742-4015, USA

 $Email\ address \hbox{: akanigow@umd.edu}$

Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, ul. Chopina $12/18,\,87\text{-}100$ Toruń, Poland

Email address: mlem@mat.umk.pl