

ON TWO CONJECTURES REGARDING GENERALIZED SEQUENCE OF DERANGEMENTS

ERYK LIPKA AND PIOTR MISKA

ABSTRACT. In [1] the second author studied arithmetic properties of a class of sequences that generalize the sequence of derangements. The aim of the following paper is to disprove two conjectures stated in [1]. The first conjecture regards the set of prime divisors of their terms. The latter one is devoted to the order of magnitude of considered sequences.

1. INTRODUCTION

We denote the sets of non-negative integers, positive integers, prime numbers, integers and rational numbers by $\mathbb{N}$, $\mathbb{N}_+$, $\mathbb{P}$, $\mathbb{Z}$ and $\mathbb{Q}$, respectively.

If there exists $n_0 \in \mathbb{N}$ such that some property holds for each integer $n \geq n_0$, then we will write, that this property holds for each $n \gg 0$.

We call that a sequence $(a_n)_{n \in \mathbb{N}}$ is an ultimately geometric progression with ratio b if there are c, b belonging to some ring such that $a_n = cb^n$ for each $n \gg 0$.

Having a given sequence $\mathbf{a} = (a_n)_{n \in \mathbb{N}}$ we define the set

$$\mathcal{P}_{\mathbf{a}} = \{p \in \mathbb{P} : \exists n \in \mathbb{N} \quad p \mid a_n\}.$$

A derangement of a set with n elements is a permutation of this set without any fixed points. We denote the number of derangements in S_n by D_n . The sequence of numbers of derangements (or the sequence of derangements, for short) $(D_n)_{n \in \mathbb{N}}$ satisfies the following recurrence relation

$$(1) \quad D_0 = 1, \quad D_n = nD_{n-1} + (-1)^n, \quad n > 0.$$

Let us consider a class $\mathcal{R}$ of generalizations of this sequence. For given $f, g, h \in \mathbb{Z}[X]$ we define the sequence $\mathbf{a}(f, g, h) = (a_n)_{n \in \mathbb{N}}$ by the recurrence

$$(2) \quad a_0 = g(0), \quad a_n = f(n)a_{n-1} + g(n)(h(n))^n, \quad n > 0.$$

Let us notice that the class $\mathcal{R}$ contains many well-known sequences:

- if $f, h = 1, g = c \in \mathbb{Z}$, then $(a_n)_{n \in \mathbb{N}} = (c(n+1))_{n \in \mathbb{N}}$ is an arithmetic progression;
- if $f = q \in \mathbb{Z}, g = c \in \mathbb{Z}, h = 0$, then $(a_n)_{n \in \mathbb{N}} = (cq^n)_{n \in \mathbb{N}}$ is a geometric progression;
- if $f = 1, g = c \in \mathbb{Z}, h = q \in \mathbb{Z}$, then $(a_n)_{n \in \mathbb{N}} = (\sum_{j=0}^n cq^j)_{n \in \mathbb{N}}$ is the sequence of partial sums of a geometric progression;
- if $f = X, g = 1, h = 0$, then $(a_n)_{n \in \mathbb{N}} = (n!)_{n \in \mathbb{N}}$ is the sequence of factorials;
- if $f = 2X + l, l \in \{0, 1\}, g = 1, h = 0$, then $(a_n)_{n \in \mathbb{N}} = ((2n+l)!)_{n \in \mathbb{N}}$ is the sequence of double factorials.

The second author investigated arithmetic properties of this class of sequences in [1], and he stated the following two conjectures.

Conjecture 1 (Conjecture 5.1. from [1]). *If $\mathbf{a} = (a_n)_{n \in \mathbb{N}} \in \mathcal{R}$ is not a geometric progression, then the set $\mathcal{P}_{\mathbf{a}}$ is infinite.*

2020 *Mathematics Subject Classification.* 11B83, 11B99.

Key words and phrases. derangement, geometric progression, prime divisor, recurrence.

Conjecture 2 (Conjecture 5.2. from [1]). *If $\mathbf{a} = (a_n)_{n \in \mathbb{N}} \in \mathcal{R}$ and there are $b, c \in \mathbb{Z}$ such that $a_n = cb^n$ for $n \gg 0$, then $a_n = cb^n$ for each $n \in \mathbb{N}$.*

2. ULTIMATELY GEOMETRIC SEQUENCES

While attempting to prove the mentioned conjectures, we found unrecoverable error in proof, which led us to a rather simple counterexamples.

Theorem 1. *Let $f, g, h \in \mathbb{Z}[X]$. If $(a_n)_{n \in \mathbb{N}} = \mathbf{a}(f, g, h)$ and there are $b, c \in \mathbb{Z}$ such that $a_n = cb^n$ for $n \gg 0$, then $h \equiv b$ or $gh \equiv 0$.*

Proof. If $c = 0$, then $a_n = 0$ for each sufficiently large positive integer n . This yields $gh \equiv 0$. Hence, we assume that $c, g, h \neq 0$.

If h is not constant, then we have $|h(n)| > |b|$ for $n \gg 0$. Thus

$$|a_n| = |f(n)cb^{n-1} + g(n)h(n)^n| \geq |b+1|^n > |cb^n|, \quad n \gg 0.$$

Hence, $a_n \neq cb^n$ for sufficiently large n . If $h(n)$ is constant and $|h(n)| > |b|$, then again using the same argument we conclude that $|a_n| > |cb^n|$ for $n \gg 0$. This means that $a_n \neq cb^n$ for sufficiently large n . We get that h is constant and $|h| \leq |b|$.

If $b = 0$, then $h \equiv 0$, so from now on we assume that $b \neq 0$. If $|b| = 1$, then of course $|h| = |b|$. If $|b| > 1$, then we consider the prime decomposition

$$|b| = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots, \quad p_i \in \mathbb{P}, \alpha_i \in \mathbb{N}_+.$$

The divisibility $p_i^{n\alpha_i} | a_n$ combined with the recurrence for $\mathbf{a}(f, g, h)$ implies

$$p_i^{(n-1)\alpha_i} | f(n)cb^{n-1} + g(n)h^n$$

and $p_i^{(n-1)\alpha_i} | g(n)h^n$ for $n \gg 0$. If $p_i^{\alpha_i} \nmid h$, then we have $p_i^{n-\alpha_i} | g(n)$, which is a contradiction for arbitrary large n as g is a nonzero polynomial. As a result, $p_i^{\alpha_i} | h$ for every i , hence $b | h$. Finally, $|h| \geq |b|$, which leads us to equality $|h| = |b|$.

If $h = -b$, then using the recurrence for $\mathbf{a}(f, g, h)$ and $n \gg 0$ we have the following chain of equalities.

$$\begin{aligned} cb^n &= f(n)cb^{n-1} + g(n)b^n(-1)^n \\ cb &= f(n)c + g(n)b(-1)^n \\ f(n) &= b + g(n)\frac{b}{c}(-1)^{n-1} \end{aligned}$$

If g is not constant, then the value of $f(n)$ changes sign infinitely many times, which is impossible for a polynomial. If g is a nonzero constant then $f(n)$ would attain two values alternately which is also impossible for a polynomial. Finally we get $h = b$. $\square$

A formula for a sequence $(a_n)_{n \in \mathbb{N}} = \mathbf{a}(f, g, h)$ with $gh \equiv 0$ can be rewritten as

$$(3) \quad a_n = g(0) \prod_{i=1}^n f(i), \quad n \in \mathbb{N}.$$

If such a sequence is an ultimately geometric progression, then f is constant and the sequence is a geometric progression. However, if $gh \neq 0$, then it is possible that $\mathbf{a}(f, g, h)$ is an ultimately geometric progression but not a geometric progression.

Theorem 2. *Let $f, g, h \in \mathbb{Z}[X]$. If $(a_n)_{n \in \mathbb{N}} = \mathbf{a}(f, g, h)$, $gh \neq 0$ and there are $b, c \in \mathbb{Z}$ such that $a_n = cb^n$ for $n \gg 0$, then*

$$g = c - \frac{c}{b}f.$$

Moreover, if $n_0 = \min \{n \in \mathbb{N} : f(n) = 0\}$, then $a_n = cb^n$ if and only if $n \geq n_0$.

Proof. It is easy to see that $bc = 0$ would imply $gh \equiv 0$. We already know from previous theorem that $h = b$, so for $n \gg 0$ we have

$$\begin{aligned} cb^n &= f(n)cb^{n-1} + g(n)b^n \\ cb &= f(n)c + g(n)b \\ g(n) &= c - f(n)\frac{c}{b} \end{aligned}$$

If two polynomials are equal for infinitely many values, then they are equal, so $g = c - \frac{c}{b}f$.

Now, if $a_n = cb^n$ and $a_{n-1} \neq cb^{n-1}$, then

$$cb^n = a_n = f(n)a_{n-1} + g(n)b^n = f(n)a_{n-1} + cb^n - cb^{n-1}f(n),$$

which implies

$$0 = f(n)(a_{n-1} - cb^{n-1}).$$

Thus, $f(n)$ must be zero. On the other hand, if $a_{n-1} = cb^{n-1}$, then

$$a_n = f(n)cb^{n-1} + g(n)b^n = f(n)cb^{n-1} + cb^n - f(n)cb^{n-1} = cb^n.$$

If $f(n) = 0$, then

$$a_n = f(n)a_{n-1} + g(n)b^n = f(n)(a_{n-1} - cb^{n-1}) + cb^n = cb^n.$$

From this we can see, that if $a_n = cb^n$ for $n \geq n_0$, then n_0 is the least non-negative integer zero of the polynomial f . $\square$

3. COUNTEREXAMPLES

The following are examples of sequences $\mathbf{a} \in \mathcal{R}$ being ultimately geometric progressions but not geometric progressions and such that the set $\mathcal{P}_{\mathbf{a}}$ is finite. The presented sequences are counterexamples for both Conjectures 1 and 2.

Example 1. The following example shows that for each pair (b, c) of integers there exists a sequence $\mathbf{a} = (a_n)_{n \in \mathbb{N}} \in \mathcal{R}$ such that it is not a geometric progression but $a_n = cb^n$ for $n \gg 0$. Let $f = b(2 - X)$, $g = c(X - 1)$, $h = b$ and $(a_n)_{n \in \mathbb{N}} = \mathbf{a}(f, g, h)$. Then, we have

$$\begin{aligned} a_0 &= -c, \\ a_1 &= b \cdot (-c) + 0 \cdot b = -cb, \\ a_2 &= 0 \cdot (-bc) + c \cdot b^2 = cb^2, \\ a_n &= b(2 - n) \cdot cb^{n-1} + c(n - 1) \cdot b^n = cb^n \text{ for } n > 2. \end{aligned}$$

So, we see that this sequence is not a geometric progression despite being ultimately geometric. Because $\mathcal{P}_{\mathbf{a}} = \{p \in \mathbb{P} : p \mid bc\}$, we thus see that $\mathcal{P}_{\mathbf{a}}$ is finite on condition that $bc \neq 0$.

Example 2. This example shows that for each $n_0 \in \mathbb{N}$ there exists a sequence $\mathbf{a} = (a_n)_{n \in \mathbb{N}} \in \mathcal{R}$ without zero terms such that $(a_n)_{n \geq n_0}$ is a geometric progression but $(a_n)_{n \geq n_0 - 1}$ is not. Let us take some $b', c' \in \mathbb{Z}$ and put $f = 2b'n_0! - 2b' \prod_{j=0}^{n_0-1} (X - j)$, $g = -c'n_0! + 2c' \prod_{j=0}^{n_0-1} (X - j)$ and $h = b'n_0!$. Then, the terms of the sequence $\mathbf{a}(f, g, h)$ are equal to

$$\begin{aligned} a_n &= (1 - 2^{n+1})c'b'^n n_0!^{n+1}, \text{ for } n < n_0, \\ a_n &= c'b'^n n_0!^{n+1}, \text{ for } n \geq n_0. \end{aligned}$$

As we can see, $a_n = cb^n$ for $n \geq n_0$, where $b = b'dn_0!$ and $c = c'dn_0!$. On the other hand $a_n \neq cb^n$ for any $n \in \{0, \dots, n_0 - 1\}$. If $bc \neq 0$, then the set $\mathcal{P}_{\mathbf{a}}$ is finite.

Example 3. This example also shows that for each $n_0 \in \mathbb{N}$ there exists a sequence $\mathbf{a} = (a_n)_{n \in \mathbb{N}} \in \mathcal{R}$ without zero terms such that $(a_n)_{n \geq n_0}$ is a geometric progression but $(a_n)_{n \geq n_0-1}$ is not. To the opposition of the previous example, the terms of the sequence $\mathbf{a}(f, g, h)$ have the same sign if $b > 0$. Let us take some $b', c', d \in \mathbb{Z}$ and put $f = b'n_0! - b' \prod_{j=0}^{n_0-1} (X - j)$, $g = c'(d-1)n_0! + c' \prod_{j=0}^{n_0-1} (X - j)$ and $h = b'dn_0!$. Then, the terms of the sequence $\mathbf{a}(f, g, h)$ are equal to

$$\begin{aligned} a_n &= (d^{n+1} - 1)c'(b')^n(n_0!)^{n+1}, \text{ for } n < n_0, \\ a_n &= c'(b')^n d^{n+1}(n_0!)^{n+1}, \text{ for } n \geq n_0. \end{aligned}$$

As we can see, $a_n = cb^n$ for $n \geq n_0$, where $b = b'dn_0!$ and $c = c'dn_0!$. On the other hand $a_n \neq cb^n$ for any $n \in \{0, \dots, n_0 - 1\}$. If $bc \neq 0$, then the set $\mathcal{P}_{\mathbf{a}}$ is finite.

Remark 1. If we allow to consider polynomials f, g, h belonging to the ring $\mathbb{Q}_{\mathbb{Z}}[X] := \{P \in \mathbb{Q}[X] : P(\mathbb{Z}) \subset \mathbb{Z}\}$, then putting $f = 2b - \frac{2b}{n_0!} \prod_{j=0}^{n_0-1} (X - j)$, $g = -c + \frac{2c}{n_0!} \prod_{j=0}^{n_0-1} (X - j)$ and $h = b$ we obtain for any pair (b, c) of integers and $n_0 \in \mathbb{N}$ a sequence of the form $\mathbf{a}(f, g, h) = (a_n)_{n \in \mathbb{N}}$ such that $a_n = cb^n$ if and only if $n \geq n_0$. Just as in Example 2 we obtain $a_n = (1 - 2^{n+1})cb^n$ for $n < n_0$.

Similarly we can modify Example 3. Taking $f = b' - \frac{b'}{n_0!} \prod_{j=0}^{n_0-1} (X - j)$, $g = c'(d-1) + \frac{c'}{n_0!} \prod_{j=0}^{n_0-1} (X - j)$ and $h = b'd$, we get

$$\begin{aligned} a_n &= (d^{n+1} - 1)c'(b')^n, \text{ for } n < n_0, \\ a_n &= cb^n, \text{ for } n \geq n_0, \end{aligned}$$

where $b = b'd$ and $c = c'd$.

4. FURTHER STUDY

Despite the fact that Conjectures 1 and 2 turned out to be false, there remains an open problem of classification of the sequences $\mathbf{a} \in \mathcal{R}$ such that the set $\mathcal{P}_{\mathbf{a}}$ is finite. We know, that $\mathbf{a} \in \mathcal{R}$ may be bounded but if this is the case, then $\mathbf{a}$ is an ultimately geometric progression with ratio being -1 , 0 or 1 (see [2]). Hence, at this moment, the only known sequences of the form $\mathbf{a} \in \mathcal{R}$ with finite set $\mathcal{P}_{\mathbf{a}}$ are ultimately geometric progressions. This is why we state the revised version of Conjecture 1 as follows.

Conjecture 3. If $\mathbf{a} \in \mathcal{R}$ is not an ultimately geometric progression, then the set $\mathcal{P}_{\mathbf{a}}$ is infinite.

REFERENCES

- [1] Piotr Miska, *On a generalization of the recurrence defining the number of derangements*, Colloq. Math., published on-line first 05.03.2020, DOI: 10.4064/cm7510-4-2019.
- [2] Piotr Miska, *When a constant subsequence implies ultimate periodicity*, Bulletin Of The Polish Academy Of Sciences Mathematics, vol. 67, no. 1 (2019), 41–52.

INSTITUTE OF MATHEMATICS, PEDAGOGICAL UNIVERSITY OF CRACOW, UL. PODCHORAŻYCH 2, 30-348 KRAKÓW, POLAND,
E-mail address: erylkipka@gmail.com

INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, JAGIELLONIAN UNIVERSITY IN CRACOW, UL. ŁOJASIEWICZA 6, 30-084 KRAKÓW, POLAND,
E-mail address: piotr.miska@uj.edu.pl