

RING ISOMORPHISMS OF CLOSED STATE SPACES VIA HOMOLOGICAL MIRROR SYMMETRY

SANGWOOK LEE

ABSTRACT. We investigate how ring isomorphisms between mirror closed states of compact toric Fano manifolds come from homological mirror symmetry. A natural closed-open relation on B -side is also discussed.

1. INTRODUCTION

1.1. Mirror symmetry. Mirror symmetry, which is one of central concepts in string theory, has also enabled mathematicians to predict a number of striking coincidences of structures which come from different origins. It was first recognized as a powerful tool in mathematics when it successfully predicted curve countings in Calabi-Yau manifolds([4]) in terms of period integrals. If we consider a Fano manifold X instead of Calabi-Yau manifolds, then a version of (closed string) mirror symmetry conjecture translates into the following ring isomorphism:

$$(1.1) \quad QH^*(X) \cong \text{Jac}(W)$$

where the holomorphic function W is the *Landau-Ginzburg mirror* of X . We will call these algebras as closed state spaces(quantum cohomologies are A -models, while Jacobian rings are B -models). Obviously the isomorphism is surprising: it compares a symplectic geometry data(counting of rational curves) on one hand and a deformation theoretic invariant of a holomorphic function on the other hand. There have been several works concerning this statement, for example [3, 18].

Later Kontsevich introduced the mirror conjecture for open strings in mathematical language, so-called the *homological mirror symmetry*. For Calabi-Yau mirror pairs (X, \check{X}) , it compares two A_∞ -categories, one is the Fukaya category(A -model) $Fu(X)$ and the other is $D_\infty^b \text{Coh}(\check{X})$, the (dg-enhancement of) derived category of coherent sheaves(B -model) on \check{X} . For a Fano manifold, if the A -model is its Fukaya category, then the B -model is a category of matrix factorizations $MF(W)$ of the Landau-Ginzburg mirror W .

We focus on compact toric manifolds. In this case, Fukaya-Oh-Ohta-Ono[14] constructed the mirror isomorphism (1.1) in terms of disc countings. They show that every ambient cycle can be used as a “bulk parameter”, so that the obstruction m_0^b deforms to a new multiple of the unit $m_0^{b,b}$ if b is a weak bounding cochain. Furthermore any derivative of $m_0^{b,b}$ with respect to a direction in ambient cycles is still a multiple of the unit, and the coefficients are used to define the isomorphism.

For toric Fano manifolds, we have homological mirror symmetry. In [6] they construct a functor from the Fukaya category of a toric Fano manifold to the category of matrix factorizations. The functor sends a full subcategory \mathcal{C} generated by strongly balanced torus fibers to the whole category, because image objects are

easily seen to be generators. On the other hand, in [10] Evans-Lekili proved a generation result for monotone Hamiltonian G -manifolds: in particular we deduce from their result that the subcategory \mathcal{C} generates the Fukaya category, so we obtain the homological mirror symmetry for Fano case. A more general theorem of the generations of Fukaya categories of compact toric manifolds has been announced by [1].

1.2. Closed-open relations. It has been widely believed that the homological mirror symmetry “induces” the closed string mirror symmetry. In particular, if we concentrate on the ring structure of mirror closed state spaces, then we expect that they are isomorphic to *Hochschild cohomology algebras* of mirror categories. If a holomorphic function W has isolated singularities, then we indeed have a ring isomorphism

$$\gamma : HH^*(MF(W)) \cong Jac(W)$$

which can be seen as a “closed-open map” on the Landau-Ginzburg B -model. We will construct γ explicitly and justify why it deserves the name.

For Fukaya categories, it is in general hard to prove that their Hochschild cohomologies are isomorphic to quantum cohomologies. By [10] again, for monotone Hamiltonian G -manifolds we have isomorphisms between them, by a corollary of the generation result.

1.3. Result of this paper. We study how \mathfrak{ts} can be realized as a ring isomorphism from homological mirror symmetry. More precisely, we prove the following.

Theorem 1.1. *Let X be a compact toric Fano manifold and $\mathfrak{P}\mathfrak{D}$ be its LG superpotential. Then there is a $Fu(X)$ - $\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D})$ -bimodule \mathcal{M} such that the following diagram commutes:*

$$\begin{array}{ccc} QH^*(X) & \xrightarrow{c\mathcal{O}} & HH^*(Fu(X)) \\ \downarrow \mathfrak{ts} & & \searrow \cong \\ & & L_{\mathcal{M}}^1 \\ & & \nearrow \cong \\ \bigoplus_{\eta \in Crit(\mathfrak{P}\mathfrak{D})} Jac(\mathfrak{P}\mathfrak{D}; \eta) & \xrightarrow{\gamma} & HH^*(\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D})) \\ & & \nearrow R_{\mathcal{M}}^1 \\ & & Hom(\mathcal{M}, \mathcal{M}). \end{array}$$

We will construct \mathcal{M} explicitly from the localized mirror functor between $Fu(X)$ and $\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D})$. The maps $L_{\mathcal{M}}^1$ and $R_{\mathcal{M}}^1$ from Hochschild cohomologies to the endomorphism space of \mathcal{M} are described in [22] and will be reviewed. The main content of the proof begins with an explicit description of the map γ in terms of the bar resolution. Then the proof is a usual check of A_∞ -relations together with their differentiations.

We organize the paper as follows: in Section 2 we recall algebraic preliminaries concerning A_∞ -categories. Then we recollect relevant definitions and facts of Lagrangian Floer theory on compact toric manifolds in Section 3. Section 4 deals with categories of matrix factorizations. In Section 5 we study mirror symmetries, both closed and open string ones. Finally, we state and prove our main theorem in Section 6.

Acknowledgements. The author is grateful to Cheol-hyun Cho for his encouragement and interest in this work. He also thanks Bumsig Kim who invited him to KIAS where he began to work on this subject. The author was supported by NRF-2007-0093859.

2. A_∞ -CATEGORIES AND A_∞ -BIMODULES

2.1. Basic definitions. We refer readers to [15, 21, 22] for more detailed explanations and proofs of the results discussed in this section.

Definition 2.1. The Novikov field is $\Lambda := \left\{ \sum_{i \geq 0} a_i T^{\lambda_i} \mid a_i \in \mathbb{C}, \lambda_i \in \mathbb{R}, \lambda_i \rightarrow \infty \text{ as } i \rightarrow \infty \right\}$.

A filtration $F^\bullet \Lambda$ of Λ is given by

$$F^\lambda \Lambda := \left\{ \sum_{i \geq 0} a_i T^{\lambda_i} \mid \lambda_i \geq \lambda \text{ for all } i \right\} \subset \Lambda.$$

We write

$$F^+ \Lambda := \left\{ \sum_{i \geq 0} a_i T^{\lambda_i} \mid \lambda_i > 0 \text{ for all } i \right\}.$$

The Novikov ring Λ_0 is $F^0 \Lambda$.

Definition 2.2. A filtered A_∞ -category \mathcal{C} over Λ consists of a class of objects $Ob(\mathcal{C})$ and the set of morphisms $hom_{\mathcal{C}}(A, B)$ for each pair of objects A and B (we write $\mathcal{C}(A, B)$ instead of $hom_{\mathcal{C}}(A, B)$ for simplicity), with the following conditions:

- (1) $\mathcal{C}(A, B)$ is a filtered $\mathbb{Z}/2$ -graded Λ -vector space for any $A, B \in Ob(\mathcal{C})$,
- (2) for $k \geq 0$ there are multilinear maps of degree 1

$$m_k : \mathcal{C}(A_0, A_1)[1] \otimes \mathcal{C}(A_1, A_2)[1] \otimes \cdots \otimes \mathcal{C}(A_{k-1}, A_k)[1] \rightarrow \mathcal{C}(A_0, A_k)[1]$$

such that they preserve the filtration and satisfy the A_∞ -relation

$$(2.1) \quad \sum_{k_1+k_2=n+1} \sum_{i=1}^{k_1} (-1)^\epsilon m_{k_1}(x_1, \dots, x_{i-1}, m_{k_2}(x_i, \dots, x_{i+k_2-1}), x_{i+k_2}, \dots, x_n) = 0$$

where $\epsilon = \sum_{j=1}^{i-1} (|x_j| + 1)$.

The meaning of m_0 is that for each object A we have $m_0^A \in hom^1(A, A)[1] = hom^2(A, A)$.

For convenience, we write $|x_j|' := |x_j| + 1$.

We can also define A_∞ -operations by coderivations. Let

$$BC := \bigoplus_{X_0, \dots, X_k \in Ob(\mathcal{C})} \mathcal{C}(X_0, X_1)[1] \otimes \cdots \otimes \mathcal{C}(X_{k-1}, X_k)[1].$$

Then define

$$\hat{m} : BC \rightarrow BC,$$

$$\hat{m}(x_0, \dots, x_{k-1}) = \sum_{l,j} (-1)^{\sum_{i=0}^l |x_i|'} x_0 \otimes \cdots \otimes x_l \otimes m_j(x_{l+1}, \dots, x_{l+j}), x_{l+j+1} \otimes \cdots \otimes x_{k-1}.$$

Now (2.1) is equivalent to $\hat{m} \circ \hat{m} = 0$.

Definition 2.3. Let \mathcal{C} and \mathcal{C}' be A_∞ -categories. An A_∞ -functor between \mathcal{C} and \mathcal{C}' is a collection $\mathcal{F} = \{\mathcal{F}_i\}_{i \geq 0}$ consisting of

- $\mathcal{F}_0 : Ob(\mathcal{C}) \rightarrow Ob(\mathcal{C}')$,
- $\mathcal{F}_k : \mathcal{C}(A_0, A_1)[1] \otimes \cdots \otimes \mathcal{C}(A_{k-1}, A_k)[1] \rightarrow \mathcal{C}'(\mathcal{F}_0(A_0), \mathcal{F}_0(A_k))[1]$, *multilinear maps of degree 0*

which are subject to the following condition:

$$\begin{aligned} & \sum_{i,j} (-1)^{|x_1|' + \cdots + |x_{i-1}|'} \mathcal{F}_{i-j+k}(x_1, \dots, x_{i-1}, m_{j-i+1}^{\mathcal{C}}(x_i, \dots, x_j), x_{j+1}, \dots, x_k) \\ &= \sum_l m_{l+1}^{\mathcal{C}'}(\mathcal{F}_{i_1-1}(x_1, \dots, x_{i_1}), \mathcal{F}_{i_2-i_1}(x_{i_1+1}, \dots, x_{i_2}), \dots, \mathcal{F}_{k-i_l}(x_{i_l+1}, \dots, x_k)). \end{aligned}$$

Again, we can define an A_∞ -functor \mathcal{F} as an equation

$$\mathcal{F} \circ \hat{m} = m \circ \hat{\mathcal{F}}$$

where

$$\begin{aligned} & \hat{\mathcal{F}} : BC \rightarrow BC, \\ & \hat{\mathcal{F}}(x_1, \dots, x_k) = \sum_{i_1, \dots, i_l} \mathcal{F}_{i_1}(x_1, \dots, x_{i_1}) \otimes \cdots \otimes \mathcal{F}_{i_l}(x_{k-i_l+1}, \dots, x_k). \end{aligned}$$

For $x \in hom(A, B)$, we have

$$(2.2) \quad m_1^2(x) + m_2(m_0^A, x) + (-1)^{|x|+1} m_2(x, m_0^B) = 0.$$

Hence if $m_0 = 0$, then $m_1^2 = 0$, but if m_0 is nonzero then m_1 does not have to be a differential.

To achieve $m_1^2 = 0$ so that we can think of m_1 -homologies, we need to deform the original A_∞ -category. We recall more definitions.

Definition 2.4. For an object A in an A_∞ -category \mathcal{C} , $e_A \in \mathcal{C}(A, A)$ is called a unit if it satisfies

- (1) $m_2(e_A, x) = x$, $m_2(y, e_A) = (-1)^{|y|} y$ for any $x \in \mathcal{C}(A, B)$, $y \in \mathcal{C}(B, A)$,
- (2) $m_{k+1}(x_1, \dots, e_A, \dots, x_k) = 0$ for any $k \neq 1$.

Definition 2.5. An element $b \in F^+ \mathcal{C}^1(A, A)$ is called a weak bounding cochain of A if it is a solution of the weak Maurer-Cartan equation

$$(2.3) \quad m(e^b) := m_0^A + m_1(b) + m_2(b, b) + \cdots = \mathfrak{P}\mathfrak{D}(A, b) \cdot e_A$$

for some $PO(A, b) \in \Lambda$. If such a solution exists, then A is called weakly unobstructed. If there exists a solution b such that $\mathfrak{P}\mathfrak{D}(A, b) = 0$, then b is called a bounding cochain and A is called unobstructed. $\mathfrak{P}\mathfrak{D}(A, b)$ is called the Landau-Ginzburg superpotential of b .

We denote $\mathcal{M}_{weak}(A)$ be the set of weak bounding cochains of A . Then $\mathfrak{P}\mathfrak{D}(A, \cdot)$ is a function on $\mathcal{M}_{weak}(A)$. We also define

$$\mathcal{M}_{weak}^\lambda(A) := \{b \in \mathcal{M}_{weak}(A) \mid \mathfrak{P}\mathfrak{D}(A, b) = \lambda\}.$$

Given an A_∞ -category \mathcal{C} , under the assumption $\mathcal{M}_{weak}^\lambda(A)$ is nonempty for some objects, we define a new A_∞ -category \mathcal{C}_λ as

$$Ob(\mathcal{C}_\lambda) = \bigcup_{A \in Ob(\mathcal{C})} \{A\} \times \mathcal{M}_{weak}^\lambda(A),$$

$$hom_{\mathcal{C}_\lambda}((A_1, b_1), (A_2, b_2)) = hom_{\mathcal{C}}(A_1, A_2)$$

with the following A_∞ -structure maps

$$m_k^{b_0, \dots, b_k} : \mathcal{C}_\lambda((A_0, b_0), (A_1, b_1)) \otimes \cdots \otimes \mathcal{C}_\lambda((A_{k-1}, b_{k-1}), (A_k, b_k)) \rightarrow \mathcal{C}_\lambda((A_0, b_0), (A_k, b_k)),$$

$$m_k^{b_0, \dots, b_k}(x_1, \dots, x_k) := \sum_{l_0, \dots, l_k} m_{k+l_0+\dots+l_k}(b_0^{l_0}, x_1, b_1^{l_1}, \dots, b_{k-1}^{l_{k-1}}, x_k, b_k^{l_k})$$

where $x_i \in \mathcal{C}_\lambda((A_i, b_i), (A_{i+1}, b_{i+1}))$. A_∞ -relation is induced by the weak Maurer-Cartan equation (2.3).

Theorem 2.6. *Let $(A_0, b_0), (A_1, b_1) \in \text{Ob}(\mathcal{C}_\lambda)$. Then $(m_1^{b_0, b_1})^2 = 0$.*

Proof. Let $x \in \mathcal{C}_\lambda((A_0, b_0), (A_1, b_1))$. Then the A_∞ -equation is

$$(m_1^{b_0, b_1})^2 + m_2(m(e^{b_0}), x) + (-1)^{|x|+1} m_2(x, m(e^{b_1})) = 0.$$

By $m(e^{b_0}) = \lambda \cdot e_{A_0}$, $m(e^{b_1}) = \lambda \cdot e_{A_1}$ and by definition of units,

$$m_2(m(e^{b_0}), x) + (-1)^{|x|+1} m_2(x, m(e^{b_1})) = 0,$$

so $(m_1^{b_0, b_1})^2 = 0$. □

Hence, if there exist weak Maurer-Cartan solutions, we get a new A_∞ -category with $m_1^2 = 0$ by restricting to objects sharing certain value of a Landau-Ginzburg (LG for short) superpotential.

Definition 2.7. *Let \mathcal{C} and \mathcal{D} be A_∞ -categories. A \mathcal{C} - \mathcal{D} -bimodule \mathcal{M} is the following data:*

- For $V \in \text{Ob}(\mathcal{C})$ and $V' \in \text{Ob}(\mathcal{D})$, $\mathcal{M}(V, V')$ is a graded k -vector space,
- degree 1 $-r-s$ multilinear maps

$$\begin{aligned} \mu^{r|1|s} : \mathcal{C}(V_0, V_1) \otimes \dots \otimes \mathcal{C}(V_{r-1}, V_r) \otimes \mathcal{M}(V_r, W_0) \otimes \mathcal{D}(W_0, W_1) \otimes \dots \otimes \mathcal{D}(W_{s-1}, W_s) \\ \rightarrow \mathcal{M}(V_0, W_s) \end{aligned}$$

for any $V_i \in \text{Ob}(\mathcal{C})$ and $W_j \in \text{Ob}(\mathcal{D})$ satisfying

$$\begin{aligned} & \sum (-1)^\epsilon \mu^{i+1|1|s-j-1}(v_0, \dots, v_i, \mu^{r-i-1|1|j+1}(v_{i+1}, \dots, v_{r-1}, \underline{m}, w_0, \dots, w_j), w_{j+1}, \dots, w_{s-1}) \\ & + \sum (-1)^\epsilon \mu^{i+r-j+1|1|s}(v_0, \dots, v_i, m_{j-i}^{\mathcal{C}}(v_{i+1}, \dots, v_j), v_{j+1}, \dots, v_{r-1}, \underline{m}, w_0, \dots, w_{s-1}) \\ & + \sum (-1)^{\epsilon'} \mu^{r|1|i+s-l+1}(v_0, \dots, v_{r-1}, \underline{m}, w_0, \dots, w_i, m_{l-i}^{\mathcal{D}}(w_{i+1}, \dots, w_l), w_{l+1}, \dots, w_{s-1}) \\ & = 0 \end{aligned}$$

where

$$\epsilon = |v_0|' + \dots + |v_i|', \quad \epsilon' = |v_0|' + \dots + |v_{r-1}|' + |m| + |w_0|' + \dots + |w_i|'.$$

Definition 2.8. *A pre-morphism of \mathcal{C} - \mathcal{D} -bimodules $\mathcal{F} : \mathcal{M} \rightarrow \mathcal{M}'$ of degree k is a collection of multilinear maps*

$$\begin{aligned} \mathcal{F}^{r|1|s} : \mathcal{C}(V_0, V_1) \otimes \dots \otimes \mathcal{C}(V_{r-1}, V_r) \otimes \mathcal{M}(V_r, W_0) \otimes \mathcal{D}(W_0, W_1) \otimes \dots \otimes \mathcal{D}(W_{s-1}, W_s) \\ \rightarrow \mathcal{M}'(V_0, W_s) \end{aligned}$$

of degree $k-r-s$, and the composition $\mathcal{F}' \circ \mathcal{F}$ is defined by

$$\begin{aligned} & (\mathcal{F}' \circ \mathcal{F})(v_1, \dots, v_r, \underline{m}, w_1, \dots, w_s) \\ & := \sum (-1)^{|\mathcal{F}|(|v_1|' + \dots + |v_i|')} \mathcal{F}'^{i|1|s-j}(v_1, \dots, v_i, \mathcal{F}^{r-i|1|j}(v_{i+1}, \dots, \underline{m}, w_1, \dots, w_j), w_{j+1}, \dots, w_s). \end{aligned}$$

The differential δ on premorphisms is defined by

$$\begin{aligned} & (\delta\mathcal{F})(v_1, \dots, v_r, \underline{m}, w_1, \dots, w_s) \\ &:= \sum (-1)^{\epsilon_1} \mu_{\mathcal{M}'}^{i|1|s-j} (v_1, \dots, v_i, \mathcal{F}^{r-i|1|j} (v_{i+1}, \dots, v_r, \underline{m}, w_1, \dots, w_j), w_{j+1}, \dots, w_s) \\ & \quad - \sum (-1)^{\epsilon_2} \mathcal{F}^{i|1|s-j} (v_1, \dots, v_i, \mu^{r-i|1|s-j} (v_{i+1}, \dots, v_r, \underline{m}, w_1, \dots, w_j), w_{j+1}, \dots, w_s) \\ & \quad - \sum \mathcal{F}^{*|1|s} (\hat{m}^{\mathcal{C}}(v_1, \dots, v_r), \underline{m}, w_1, \dots, w_s) \\ & \quad - \sum (-1)^{\epsilon_3} \mathcal{F}^{r|1|*} (v_1, \dots, v_r, \underline{m}, \hat{m}^{\mathcal{D}}(w_1, \dots, w_s)) \end{aligned}$$

where

$$\epsilon_1 = |\mathcal{F}|(|v_1|' + \dots + |v_i|'), \quad \epsilon_2 = |v_1|' + \dots + |v_i|', \quad \epsilon_3 = |v_1|' + \dots + |v_r|' + |m|.$$

More succinctly,

$$\delta\mathcal{F} = \mu_{\mathcal{M}'} \circ \hat{\mathcal{F}} - (-1)^{|\mathcal{F}|} \mathcal{F} \circ \hat{\mu}_{\mathcal{M}}.$$

Remark 2.9. The definition of the degree of a pre-morphism of bimodules is motivated by the fact that the degree k pre-morphism is indeed given by multilinear maps

$$\mathcal{C}[1]^{\otimes r} \otimes \mathcal{M} \otimes \mathcal{D}[1]^{\otimes s} \rightarrow \mathcal{M}$$

of degree k . Once we accept such a perspective on degrees of maps, all signs obey only Koszul conventions and so we sometimes just write $(-1)^{\text{Koszul}}$ for signs when it is sufficient.

The readers can easily check that \mathcal{C} - \mathcal{D} -bimodules together with premorphisms form a dg category. If $\mathcal{F} : \mathcal{M} \rightarrow \mathcal{N}$ is a premorphism such that $\delta\mathcal{F} = 0$ and its cohomology level map $[\mathcal{F}^{0|1|0}]$ is an isomorphism, then \mathcal{F} is called a *quasi-isomorphism* and write $\mathcal{M} \cong \mathcal{N}$.

Example 2.10. For an A_∞ -category \mathcal{C} , the diagonal bimodule \mathcal{C}_Δ is a \mathcal{C} - \mathcal{C} -bimodule defined by

$$\mathcal{C}_\Delta(X, Y) := \mathcal{C}(X, Y)$$

with module structure maps

$$\mu^{r|1|s} := m_{r+s+1}^{\mathcal{C}}.$$

We recall some operations on bimodules.

Definition 2.11 (Base change). Let $\mathcal{F} : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ and $\mathcal{G} : \mathcal{D}_1 \rightarrow \mathcal{D}_2$ be A_∞ -functors. Suppose that \mathcal{M} is a \mathcal{C}_2 - \mathcal{D}_2 -bimodule. Then a \mathcal{C}_1 - \mathcal{D}_1 -bimodule $(\mathcal{F} \otimes \mathcal{G})^* \mathcal{M}$ is defined on objects by

$$(\mathcal{F} \otimes \mathcal{G})^* \mathcal{M}(X, Y) := \mathcal{M}(\mathcal{F}(X), \mathcal{G}(Y))$$

for $X \in \text{Ob}(\mathcal{C}_1)$, $Y \in \text{Ob}(\mathcal{D}_1)$, and the structure maps are given by

$$\begin{aligned} & \mu_{(\mathcal{F} \otimes \mathcal{G})^* \mathcal{M}}^{r|1|s} (v_1, \dots, v_r, \underline{m}, w_1, \dots, w_s) \\ &:= \sum_{k,l} \mu_{\mathcal{M}}^{k|1|l} (\mathcal{F}_{i_1}(v_1, \dots), \dots, \mathcal{F}_{i_k}(\dots, v_r), \underline{m}, \mathcal{G}_{j_1}(w_1, \dots), \dots, \mathcal{G}_{j_l}(\dots, w_s)). \end{aligned}$$

Definition 2.12 (Tensor product). Let \mathcal{M} be a \mathcal{C} - \mathcal{D} -bimodule and \mathcal{N} be a \mathcal{D} - \mathcal{E} -bimodule for A_∞ -categories \mathcal{C} , \mathcal{D} and \mathcal{E} . Then $\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N}$ is a \mathcal{C} - \mathcal{E} -bimodule such that

$$(\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N})(C, E) := \bigoplus_{D_1, \dots, D_k \in \text{Ob}(\mathcal{D})} \mathcal{M}(C, D_1) \otimes \mathcal{D}(D_1, D_2) \otimes \dots \otimes \mathcal{D}(D_{k-1}, D_k) \otimes \mathcal{N}(D_k, E)$$

for $C \in \text{Ob}(\mathcal{C})$ and $E \in \text{Ob}(\mathcal{E})$, and the structure maps are given as follows:

$$\begin{aligned} & \mu_{\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N}}^{0|1|0}(\underline{m} \otimes d_1 \otimes \cdots \otimes d_k \otimes \underline{n}) \\ &:= \sum \mu_{\mathcal{M}}^{0|1|i}(\underline{m}, d_1, \dots, d_i) \otimes d_{i+1} \otimes \cdots \otimes d_k \otimes \underline{n} \\ & \quad + \sum (-1)^{\text{Koszul}} \underline{m} \otimes d_1 \otimes \cdots \otimes d_i \otimes m_{j-i}^{\mathcal{D}}(d_{i+1}, \dots, d_j) \otimes d_{j+1} \otimes \cdots \otimes d_k \otimes \underline{n} \\ & \quad + \sum (-1)^{\text{Koszul}} \underline{m} \otimes d_1 \otimes \cdots \otimes d_i \otimes \mu_{\mathcal{N}}^{k-i|1|0}(d_{i+1}, \dots, d_k, \underline{n}), \end{aligned}$$

$$\begin{aligned} & \mu_{\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N}}^{r|1|0}(c_1, \dots, c_r, \underline{m} \otimes d_1 \otimes \cdots \otimes d_k \otimes \underline{n}) \\ &:= \sum \mu_{\mathcal{M}}^{r|1|p}(c_1, \dots, c_r, \underline{m}, d_1, \dots, d_p) \otimes d_{p+1} \otimes \cdots \otimes d_k \otimes \underline{n}, \end{aligned}$$

$$\begin{aligned} & \mu_{\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N}}^{0|1|s}(\underline{m} \otimes d_1 \otimes \cdots \otimes d_k \otimes \underline{n}, e_1, \dots, e_s) \\ &:= \sum (-1)^{\text{Koszul}} \underline{m} \otimes d_1 \otimes \cdots \otimes d_p \otimes \mu_{\mathcal{N}}^{k-p|1|s}(d_{p+1}, \dots, d_k, \underline{n}, e_1, \dots, e_s) \end{aligned}$$

and $\mu_{\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N}}^{r|1|s} = 0$ if r and s are both nonzero.

Definition 2.13. Two A_{∞} -categories \mathcal{C} and \mathcal{D} are Morita equivalent if there is a \mathcal{C} - \mathcal{D} -bimodule \mathcal{M} and a \mathcal{D} - \mathcal{C} -bimodule \mathcal{N} such that

$$\mathcal{M} \otimes_{\mathcal{D}} \mathcal{N} \cong \mathcal{C}_{\Delta}, \quad \mathcal{N} \otimes_{\mathcal{C}} \mathcal{M} \cong \mathcal{D}_{\Delta}.$$

In this case, we call \mathcal{M} and \mathcal{N} be Morita bimodules.

2.2. Hochschild cohomology.

Definition 2.14. Let \mathcal{M} be an A_{∞} -bimodule over \mathcal{C} . We define the Hochschild cochain complex of \mathcal{M}

$$CH^*(\mathcal{C}, \mathcal{M}) := \prod_{X_0, \dots, X_k \in \text{Ob}(\mathcal{C})} \text{hom}^{\bullet}(\mathcal{C}(X_0, X_1)[1] \otimes \cdots \otimes \mathcal{C}(X_{k-1}, X_k)[1], \mathcal{M}(X_0, X_k))[-1]$$

with differential b^* defined by

$$\begin{aligned} & b^* \phi(x_0, \dots, x_{k-1}) \\ &:= \sum \phi(\hat{m}(x_0, \dots, x_{k-1})) \\ & \quad + \sum (-1)^{|\phi|'(|x_0|' + \cdots + |x_i|')} \mu_{\mathcal{M}}^{i+1|1|k-l-1}(x_0, \dots, x_i, \phi(x_{i+1}, \dots, x_l), x_{l+1}, \dots, x_{k-1}). \end{aligned}$$

Its cohomology of b^* is called the Hochschild cohomology of \mathcal{C} with coefficient in \mathcal{M} . If $\mathcal{M} = \mathcal{C}_{\Delta}$, then we write $CH^*(\mathcal{C}) := CH^*(\mathcal{C}, \mathcal{C}_{\Delta})$.

Proposition 2.15 ([17]). $CH^*(\mathcal{C})$ is an A_{∞} -algebra with A_{∞} -operations given by

$$\begin{aligned} & M^k(\phi_1, \dots, \phi_k)(x_1, \dots, x_n) \\ &:= \sum (-1)^{\epsilon} m_*(\vec{x}_{i_1}, \phi_1(\vec{x}_{j_1}), \vec{x}_{i_2}, \phi_2(\vec{x}_{j_2}), \dots, \vec{x}_{i_k}, \phi_k(\vec{x}_{j_k}), \vec{x}_{i_{k+1}}). \end{aligned}$$

with $\vec{x}_{i_1} \otimes \vec{x}_{j_1} \otimes \cdots \otimes \vec{x}_{i_{k+1}} = x_1 \otimes \cdots \otimes x_n$, $M^0 = 0$, $M^1 = b^*$, and

$$\epsilon = \sum_{l=1}^k |\phi_l|'(|\vec{x}_{i_1}|' + |\vec{x}_{j_1}|' + \cdots + |\vec{x}_{j_{l-1}}|' + |\vec{x}_{i_l}|').$$

In particular, the binary product M^2 induces the *Yoneda product* \cup on the cohomology $HH^*(\mathcal{C})$ by

$$\phi \cup \psi := (-1)^{|\phi|} M^2(\phi, \psi).$$

Then the Yoneda product is associative.

2.3. Morita invariance of Hochschild cohomology. Let \mathcal{A} and \mathcal{B} be Morita equivalent with Morita bimodules \mathcal{M} and \mathcal{N} which are over $\mathcal{A}\text{-}\mathcal{B}$ and $\mathcal{B}\text{-}\mathcal{A}$ respectively. Recall the following:

Lemma 2.16 ([22]). *The following are A_∞ quasi-isomorphisms*

$$L_{\mathcal{M}} : CH^*(\mathcal{A}) \rightarrow \text{hom}_{\mathcal{A}\text{-}\mathcal{B}}^*(\mathcal{M}, \mathcal{M}),$$

$$R_{\mathcal{M}} : CH^*(\mathcal{B})^{op} \rightarrow \text{hom}_{\mathcal{A}\text{-}\mathcal{B}}^*(\mathcal{M}, \mathcal{M})$$

which are defined as follows:

$$\begin{aligned} & L_{\mathcal{M}}^p(\phi_1, \dots, \phi_p)(a_1, \dots, a_r, \underline{m}, b_1, \dots, b_s) \\ &:= \sum (-1)^{\text{Koszul}} \mu_{\mathcal{M}}^{*|1|s}(\vec{a}_1, \phi_1(\vec{a}_2), \vec{a}_3, \dots, \phi_p(\vec{a}_{2p}), \vec{a}_{2p+1}, \underline{m}, b_1, \dots, b_s), \\ & R_{\mathcal{M}}^p(\phi_1, \dots, \phi_p)(a_1, \dots, a_r, \underline{m}, b_1, \dots, b_s) \\ &:= \sum (-1)^{\text{Koszul}} \mu_{\mathcal{M}}^{r|1|*}(\vec{a}_1, \dots, a_r, \underline{m}, \vec{b}_1, \phi_p(\vec{b}_2), \vec{b}_3, \dots, \phi_1(\vec{b}_{2p}), \vec{b}_{2p+1}). \end{aligned}$$

In particular, $L_{\mathcal{M}}^1$ and $R_{\mathcal{M}}^1$ induce ring isomorphisms on cohomology level.

Later we will examine the maps $L_{\mathcal{M}}^1$ and $R_{\mathcal{M}}^1$ carefully for the Morita bimodule between mirror pairs $Fu(X)$ and $\overline{MF}_{A_\infty}(W)$.

3. LAGRANGIAN FLOER THEORY ON COMPACT TORIC MANIFOLDS

3.1. Review on Fukaya categories. Let (X, ω) be a symplectic manifold and L_0, \dots, L_k be its Lagrangian submanifolds. Pick a compatible almost complex structure J_0 on (X, ω) . For a while we only assume an extreme case of intersecting patterns of those Lagrangians: there are only transverse intersections among L_0, \dots, L_k . But we will also deal with the other extreme case $L_0 = \dots = L_k$ later.

Let $CF(L_i, L_{i+1}) := \bigoplus_{p \in L_i \cap L_{i+1}} \Lambda \cdot p$ be a $\mathbb{Z}/2$ -graded vector space over Λ . The degree is defined by the Maslov index of each intersection point. Let $\beta \in \pi_2(X, L_0 \cup \dots \cup L_k)$. Define a moduli space

$$\begin{aligned} & \widehat{\mathcal{M}}(p_0, \dots, p_{k-1}; q; \beta; J_0) \\ &:= \{u : (D^2, z_0, \dots, z_{k-1}, z_k) \xrightarrow{J_0\text{-hol}} (X, p_0, \dots, p_{k-1}, q) \mid \\ & \quad u(\widehat{z_i z_{i+1}}) \subset L_{i+1}, u(\widehat{z z_0}) \subset L_0, [u] = \beta\}. \end{aligned}$$

and $\mathcal{M}(p_0, \dots, p_{k-1}; q; \beta; J_0)$ be its stable map compactification. In general such moduli spaces may not be transversal, so we apply Kuranishi perturbations and achieve moduli spaces of correct dimensions. We denote such ‘correct’ moduli space by $\mathcal{M}(p_0, \dots, p_{k-1}; q; \beta)$, omitting J_0 . Suppose that we can take suitable Kuranishi perturbations so that boundary components of the moduli spaces are expressed as fiber products of moduli spaces of the same kind with respect to evaluation maps at boundary marked points (these technical assumptions can be indeed achieved for compact toric manifolds, see [12, 13, 14]).

Now, define

$$m_k(p_0, \dots, p_{k-1}) := \sum_{\substack{\beta \in \pi_2(X, L_0 \cup \dots \cup L_k) \\ q \in L_k \cap L_0}} \# \mathcal{M}(p_0, \dots, p_{k-1}; q; \beta) \cdot T^{\omega(\beta)} \cdot q.$$

We only count the moduli space when the correct virtual dimension is zero. The above technical assumptions enables us to say that these operations satisfy a filtered A_∞ -relation (2.1).

Definition 3.1. *The Fukaya category $Fu(X)$ is an A_∞ -category whose objects are Lagrangian submanifolds and morphism spaces are $CF(L, L')$. The A_∞ -structure is given by above operations $\{m_k\}_{k \geq 0}$. The derived Fukaya category $D^\pi Fu(X)$ is the homotopy category of the triangulated envelope of $Fu(X)$ followed by Karoubi completion.*

In general, there might be nonzero m_0 , which comes from holomorphic discs whose boundaries are on Lagrangian submanifolds. More precisely, for $\beta \in \pi_2(X, L)$, let $\mathcal{M}_1^{main}(\beta)$ be the (Kuranishi perturbed) moduli space of holomorphic discs of homotopy classes β with a boundary marked point. Define $m_{0,\beta}^L := PD((ev_0)_* \mathcal{M}_1(\beta))$ which is a cocycle in L , and let $m_0^L := \sum_{\beta \in \pi_2(X, L)} m_{0,\beta}^L$. Higher A_∞ -algebra operations on a single Lagrangian are defined as follows. Let $\mathcal{M}_{k+1}^{main}(\beta)$ be the moduli space of holomorphic discs of class β with $k+1$ boundary marked points which respect the counterclockwise cyclic order. Let

$$ev_0^\beta, \dots, ev_k^\beta : \mathcal{M}_{k+1}^{main}(\beta) \rightarrow L$$

be evaluation maps at marked points (z_0, \dots, z_k) of each disc. Let p_1, \dots, p_k be cycles of L . Then

$$m_{k,\beta}^L : H^*(L; \Lambda)^{\otimes k} \rightarrow H^*(L; \Lambda)$$

is defined by

$$(PD(p_1), \dots, PD(p_k)) \mapsto PD[(ev_0^\beta)_! (ev_1^\beta \times \dots \times ev_k^\beta)^* (p_1 \times \dots \times p_k)].$$

Define

$$m_k^L := \sum_{\beta} T^{\omega(\beta)} \cdot m_{k,\beta}^L.$$

We refer to [11, 12] for the well-definedness of above operations on the de Rham model of each single Lagrangian submanifold. If we take Kuranishi perturbations successfully, then the fundamental class $[L]$ plays the role of the A_∞ -unit, so the following definition makes sense (see Definition 2.5).

Definition 3.2. *We say L is weakly unobstructed if there is a chain $b \in C^*(L; \Lambda)$ such that for some constant $\mathfrak{P}\mathfrak{D}(L, b) \in \Lambda$,*

$$(m_0^L)^b := m_0^L + m_1^L(b) + m_2^L(b, b) + \dots = c \cdot [L].$$

The constant $\mathfrak{P}\mathfrak{D}(L, b)$ is called the superpotential of (L, b) .

3.2. Floer theory on compact toric manifolds. Let X be a compact toric manifold and $\pi : X \rightarrow P$ be the moment map. For $\mathbf{u} \in \text{Int}(P)$, let $L(\mathbf{u}) := \pi^{-1}(\mathbf{u})$, i.e. a Lagrangian torus fiber over \mathbf{u} . Recall that P can be expressed as

$$P = \{\mathbf{u} \in \mathbb{R}^n \mid l_j(\mathbf{u}) \geq 0, j = 1, \dots, m\}$$

for certain affine functionals $l_j : \mathbb{R}^n \rightarrow \mathbb{R}$. Let \vec{v}_j be the primitive normal vector to the facet $\partial_i P = \{\mathbf{u} \in \mathbb{R}^n \mid l_j(\mathbf{u}) = 0\}$. Let

$$\vec{v}_j = (v_{j,1}, \dots, v_{j,n}).$$

Based on the classification of holomorphic discs bounded by torus fibers [8] and the transversality result [12], we have the following substantial result.

Theorem 3.3 ([12]). *• Every torus fiber $L(\mathbf{u})$ of a compact toric manifold X is weakly unobstructed and $H^1(L(\mathbf{u}); \Lambda_0)$ is contained in the weak Maurer-Cartan solution space.*

- Let $\{\mathbf{e}_i\}_{i=1, \dots, n}$ be the basis of $L(\mathbf{u})$, consisting of elements representing dt_i . Let $\{x_i\}_{i=1, \dots, n}$ be the dual basis and $b = \sum_i x_i \mathbf{e}_i$. Then the potential $\mathfrak{PD}(L(\mathbf{u}), b)$ has the expression

$$\mathfrak{PD}(L(\mathbf{u}), b) = z_1 + \dots + z_m + \sum_{k=1}^N P_k(z_1, \dots, z_m)$$

where $z_i := T^{l_j(\mathbf{u})}(y_1^{\mathbf{u}})^{v_{j,1}} \dots (y_n^{\mathbf{u}})^{v_{j,n}}$ and $y_1^{\mathbf{u}} := e^{x_i}$. If X is furthermore Fano, then P_k are all zero.

- The above expression $\mathfrak{PD}(L(\mathbf{u}), b)$ only depends on X (and independent of \mathbf{u}), so we denote the above $\mathfrak{PD}(L(\mathbf{u}), b)$ just by $\mathfrak{PD}(x_1, \dots, x_n)$, or \mathfrak{PD} for short.

We are particularly interested in $\text{Crit}(\mathfrak{PD})$ because of the following fact:

Theorem 3.4. [8, 12] *If $b = \sum a_i \mathbf{e}_i \in H^1(L(\mathbf{u}), \Lambda_0)$ then the following are equivalent:*

- $\frac{\partial \mathfrak{PD}}{\partial x_i}(b) = 0$ for all $i = 1, \dots, n$.
- $HF((L(\mathbf{u}), b), (L(\mathbf{u}), b); \Lambda) \cong H^*(T^n; \Lambda)$.
- $HF((L(\mathbf{u}), b), (L(\mathbf{u}), b)) \neq 0$.

Finally we recall the following generation result.

Theorem 3.5 ([1, 10]). *For a compact toric Fano manifold X , we have a decomposition*

$$D^\pi Fu(X) = \bigoplus_{\eta \in \text{Crit}(\mathfrak{PD})} D^\pi Fu_\eta(X)$$

and every torus fiber with a weak bounding cochain in $\text{Crit}(\mathfrak{PD})$ generates $D^\pi Fu_\eta(X)$.

3.3. Closed-open maps. Let $\mathcal{M}_{k+1,l}^{\text{main}}(\beta)$ be the moduli space of holomorphic discs of class β with $k+1$ boundary marked points respecting the cyclic order and l interior marked points. Define the evaluation map

$$ev = (ev_1^+, \dots, ev_l^+, ev_0, \dots, ev_k) : \mathcal{M}_{k+1,l}^{\text{main}}(\beta) \rightarrow X^l \times L^{k+1}$$

and consider the map [11, 13]:

$$\mathfrak{q}_{l,k;\beta} : E_l(\mathcal{A}) \otimes H^*(L; \Lambda)^{\otimes k} \rightarrow H^*(L; \Lambda),$$

$$\begin{aligned} & \mathfrak{q}_{l,k;\beta}([\alpha_1 \otimes \cdots \otimes \alpha_l], p_1, \dots, p_k) \\ & := (ev_0)_*(ev_1^+ \times \cdots \times ev_l^+ \times ev_0 \otimes \cdots \otimes ev_k)^*([\alpha_1 \otimes \cdots \otimes \alpha_l], p_1 \otimes \cdots \otimes p_l). \end{aligned}$$

Here $E_l(\mathcal{A})$ means the subspace of $\mathcal{A}^{\otimes l}$, consisting of S_l -invariant elements and

$$[\alpha_1 \otimes \cdots \otimes \alpha_l] = \sum_{\sigma \in S_k} \frac{1}{l!} \alpha_{\sigma(1)} \otimes \cdots \otimes \alpha_{\sigma(l)}.$$

We similarly define

$$\mathfrak{q}_{l,k;\beta}([\alpha_1 \otimes \cdots \otimes \alpha_l], p_1, \dots, p_k)$$

for p_1, \dots, p_k being transverse intersections among distinct Lagrangians L_0, \dots, L_k , using moduli spaces of holomorphic polygons with faces on L_0, \dots, L_k together with interior marked points. We also define (both for a single Lagrangian or a transversal collection of Lagrangians)

$$\mathfrak{q}_{l,k}([\alpha_1 \otimes \cdots \otimes \alpha_l], p_1, \dots, p_k) := \sum_{\beta \in \pi_2(M, \vec{L})} \mathfrak{q}_{l,k;\beta}([\alpha_1 \otimes \cdots \otimes \alpha_l], p_1, \dots, p_k) \cdot T^{\omega(\beta)}.$$

Theorem 3.6 ([13]). *Let X be a compact toric manifold. Then for any $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$ and $b \in H^1(L(\mathfrak{u}); \Lambda)$ the following always holds:*

$$\sum_{l=0}^{\infty} \sum_{k=0}^{\infty} \mathfrak{q}_{l,k}(\mathfrak{b}^l; b^k) \equiv 0 \pmod{\Lambda_+ \mathbf{e}}.$$

We denote the left hand side by $\mathfrak{P}\mathfrak{D}^{\mathfrak{b}}(b) \cdot \mathbf{e}$ and call the coefficient $\mathfrak{P}\mathfrak{D}^{\mathfrak{b}}(b)$ a bulk-deformed potential.

Now we define the following.

Definition 3.7. *Let X be a compact toric manifold and $\mathfrak{b} \in \mathcal{A}(\Lambda_+)$. Let b_0, \dots, b_k be weak bounding cochains. Then the following operations define a new A_{∞} -structure:*

$$m_k^{\mathfrak{b}, \vec{b}}(x_1, \dots, x_k) := \sum_{l=0}^{\infty} \mathfrak{q}_{l,*}(\mathfrak{b}^l; e^{b_0}, x_1, e^{b_1}, x_2, \dots, e^{b_{k-1}}, x_k, e^{b_k}).$$

Let $\mathfrak{b} = t\alpha$ where $\alpha \in \mathcal{A}$ and t is a formal variable, and $b = \sum x_i \mathbf{e}_i \in H^1(L(\mathfrak{u}), \Lambda_0)$ for some $\mathfrak{u} \in \text{Int}(P)$. Then $\mathfrak{P}\mathfrak{D}^{\mathfrak{b}}(b)$ is a formal power series in t . We observe the following:

$$\widetilde{\mathfrak{f}\mathfrak{s}}(\alpha) := \frac{\partial \mathfrak{P}\mathfrak{D}^{\mathfrak{b}}(b)}{\partial t} \Big|_{t=0} = \sum_{k \geq 0} \mathfrak{q}_{1,k}(\alpha; b^k).$$

By above theorem, $\widetilde{\mathfrak{f}\mathfrak{s}}(\alpha)$ is a multiple of the unit \mathbf{e} . Again, the coefficient is a power series in x_1, \dots, x_n , or a Laurent polynomial in y_1, \dots, y_n . We will later employ $\widetilde{\mathfrak{f}\mathfrak{s}}$ for the statement of the closed string mirror symmetry for toric Fano manifolds.

We can also think of the t -derivative of A_{∞} -equations of $\{m_k^{t\alpha, \vec{b}}\}$ at $t = 0$ as follows (we omit the decoration e^{b_i} for simplicity).

$$\begin{aligned} (3.1) \quad 0 &= \sum (-1)^{|x_1|' + \cdots + |x_i|'} m_{i+k-j+1}(x_1, \dots, x_i, \mathfrak{q}(\alpha; x_{i+1}, \dots, x_j), x_{j+1}, \dots, x_k) \\ &\quad + \sum (-1)^{|x_1|' + \cdots + |x_i|'} \mathfrak{q}(\alpha; x_1, \dots, x_i, m_{j-i}(x_{i+1}, \dots, x_j), x_{j+1}, \dots, x_k). \end{aligned}$$

This equation will be crucially used in Section 6. Remark that it is related to the degeneration of holomorphic discs with one interior constraint (see Figure 1).




FIGURE 1. A degeneration of holomorphic discs with interior constraint α . In broken discs, components without interior constraints correspond to $m_k(\vec{p})$ and components with interior constraints correspond to $\mathfrak{q}_{1,k}(\alpha; \vec{p})$.

4. CATEGORY OF MATRIX FACTORIZATIONS

Let R be a commutative algebra and $W \in R$ be a non-zero-divisor. A *matrix factorization* of W is a $\mathbb{Z}/2$ -graded R -module $E = E^0 \oplus E^1$ with a degree 1 endomorphism

$$Q = \begin{pmatrix} 0 & Q_{01} \\ Q_{10} & 0 \end{pmatrix}, \text{ where } Q_{ij} \in \text{Hom}(E^j, E^i),$$

satisfying $Q^2 = W \cdot \text{Id}$. We denote the above data by (E, Q) for short.

Matrix factorizations of W form a differential $\mathbb{Z}/2$ -graded category $MF(R, W)$ as follows: given two matrix factorizations $(E, Q), (F, Q')$, $\mathbb{Z}/2$ -graded morphisms from (E, Q) to (F, Q') are given by homomorphisms

$$\Phi = \begin{pmatrix} \Phi_{00} & \Phi_{01} \\ \Phi_{10} & \Phi_{11} \end{pmatrix}, \text{ where } \Phi_{ij} \in \text{Hom}(E^j, F^i).$$

Compositions of morphisms are defined in the obvious way. The differential on a morphism is defined as

$$\delta\Phi := Q\Phi - (-1)^{|\Phi|}\Phi Q$$

for morphisms of homogeneous degrees.

We often write as $MF(W)$ instead of $MF(R, W)$. The cohomology category $[MF(W)] := H^0(MF(W))$ with respect to δ is a triangulated category. The following theorem illustrates the geometric relevance of categories of matrix factorizations.

Theorem 4.1 ([19]). $[MF(W)] \simeq D_{sg}(\text{Spec}(R/W))$.

$D_{sg}(X)$ is defined for a variety X by the quotient of $D^b(X)$ by $\text{Perf}(X)$ which is a thick subcategory consisting of complexes which are quasi-isomorphic to bounded complexes of locally free sheaves (in affine case, projective modules). If X is smooth

then $D_{sg}(X)$ is trivial, hence the category roughly measures how the variety is singular.

We are particularly interested in the case $W \in R = \Lambda[x_1^\pm, \dots, x_n^\pm]$ with isolated critical points. According to the above theorem of Orlov, $[MF(W - \lambda)]$ is nontrivial if and only if λ is a critical value of W . For a critical point η of W , let

$$\hat{R}_\eta := \Lambda[[x_1 - \eta_1, \dots, x_n - \eta_n]].$$

If there are finitely many critical points, we take a direct sum of categories over all critical points as

$$\overline{MF}(W) := \bigoplus_{\eta \in \text{Crit}(W)} MF(\hat{R}_\eta, W - W(\eta)).$$

The direct sum of categories means that there are no nontrivial morphisms between objects in different summands.

5. MIRROR SYMMETRY

5.1. Closed string mirror symmetry. Let X be compact toric Fano. One of the simplest version of closed string mirror symmetry is the following:

Conjecture 5.1. *There is a ring homomorphism*

$$QH^*(X) \cong \text{Jac}(\mathfrak{P}\mathfrak{D})$$

where W is the Landau-Ginzburg mirror.

There were some related results due to [3, 18]. While these approaches are explicit and given by algebraic methods, the following Fukaya-Oh-Ohta-Ono's construction of the isomorphism is rather geometric.

Theorem 5.2 ([14]). *For a compact toric manifold X (not necessarily Fano), the ring isomorphism above can be given by*

$$\begin{aligned} \mathfrak{ts} : QH^*(X) &\rightarrow \text{Jac}(\mathfrak{P}\mathfrak{D}), \\ \alpha &\mapsto [\widetilde{\mathfrak{ts}}(\alpha)] \end{aligned}$$

where $\mathfrak{P}\mathfrak{D}$ is the leading order potential and

$$\text{Jac}(\mathfrak{P}\mathfrak{D}) = \frac{\Lambda[y_1^\pm, \dots, y_n^\pm]}{\left(y_1 \frac{\partial \mathfrak{P}\mathfrak{D}}{\partial y_1}, \dots, y_n \frac{\partial \mathfrak{P}\mathfrak{D}}{\partial y_n}\right)}.$$

Recall the definition of $\widetilde{\mathfrak{ts}}$ from Section 3.3: we counted holomorphic discs with interior constraints.

5.2. Localized mirror functor. Given a potential function $W \in R$, we modify the convention of the dg category $(MF(W), \delta, \circ)$ to obtain an A_∞ -category $(MF_{A_\infty}(W), m_1^{MF}, m_2^{MF})$. Objects are still matrix factorizations of W , and morphism spaces are

$$\text{Hom}_{MF_{A_\infty}(W)}((E, Q_E), (F, Q_F)) := \text{Hom}_R(F, E).$$

m_1^{MF} and m_2^{MF} are defined as

$$\begin{aligned} m_1^{MF}(\Phi) &:= \delta(\Phi) = Q_E \circ \Phi - (-1)^{|\Phi|} \Phi \circ Q_F, \\ m_2^{MF}(\Phi, \Psi) &:= (-1)^{|\Phi|} \Phi \circ \Psi. \end{aligned}$$

Then m_1^{MF} and m_2^{MF} satisfy A_∞ -relations (2.1). We also define

$$\overline{MF_{A_\infty}}(W) := \bigoplus_{\eta \in \text{Crit}(W)} MF_{A_\infty}(\hat{R}_\eta, W - W(\eta)).$$

Let (\mathbb{L}, b) be a weakly unobstructed Lagrangian where $b = x_1 X_1 + \cdots + x_n X_n$, and let $W = m_0^b$ be a function on the space of weak Maurer-Cartan solutions. For any other weakly unobstructed Lagrangian L with potential value λ (i.e. there exists a Maurer-Cartan element b_0 such that $m_0^{b_0} = \lambda$), the A_∞ -equation gives the following matrix factorization identity

$$(m_1^{b_0, b})^2 = W - \lambda.$$

Theorem 5.3 ([5]). *We have an A_∞ -functor $\mathcal{LM}^\mathbb{L}$ from Fukaya A_∞ -category to $A_\infty(MF(W))$. On the level of objects, it sends a weakly unobstructed Lagrangian L with Maurer-Cartan element b_0 to the matrix factorization (E, Q) , which is given by*

$$E := \text{Hom}((L, b_0), (\mathbb{L}, b)), \quad Q := -m_1^{b_0, b}$$

On the level of morphisms, $\mathcal{LM}_k^\mathbb{L}$ is defined as

$$(5.1) \quad \mathcal{LM}_k^\mathbb{L}(x_1, \dots, x_k)(\bullet) := m_{k+1}(x_1, \dots, x_k, \bullet).$$

Then these data define an A_∞ -functor.

Remark 5.4. In [5, 6], they considered $CF((\mathbb{L}, b), (L, b_0))$ instead. In the current convention, we do not have any sign in (5.1).

Proof. To avoid confusion, let $\{m_k^{Fu}\}$ be A_∞ -operations of the Fukaya category, while m_1^{MF} and m_2^{MF} are A_∞ -operations of $MF_{A_\infty}(W)$. We need to show that

$$(5.2) \quad \begin{aligned} & \mathcal{LM}^\mathbb{L}(\hat{m}^{Fu}(x_1, \dots, x_k)) \\ &= \sum_{\vec{x}_1 \otimes \vec{x}_2 = x_1 \otimes \cdots \otimes x_k} m_2^{MF}(\mathcal{LM}^\mathbb{L}(\vec{x}_1), \mathcal{LM}^\mathbb{L}(\vec{x}_2)) + m_1^{MF}(\mathcal{LM}^\mathbb{L}(x_1, \dots, x_k)). \end{aligned}$$

The left hand side can be written as

$$\sum_{\vec{x}_1 \otimes \vec{x}_2 \otimes \vec{x}_3 = x_1 \otimes \cdots \otimes x_k} (-1)^{|\vec{x}_1|'} m^{Fu}(\vec{x}_1 \otimes m^{Fu}(\vec{x}_2) \otimes \vec{x}_3, \bullet).$$

By definition of m_2^{MF} ,

$$\begin{aligned} & \sum_{\vec{x}_1 \otimes \vec{x}_2 = x_1 \otimes \cdots \otimes x_k} m_2^{MF}(\mathcal{LM}^\mathbb{L}(\vec{x}_1), \mathcal{LM}^\mathbb{L}(\vec{x}_2)) \\ &= \sum_{\vec{x}_1 \otimes \vec{x}_2 = x_1 \otimes \cdots \otimes x_k} (-1)^{|\vec{x}_1|'+1} \mathcal{LM}^\mathbb{L}(\vec{x}_1) \circ \mathcal{LM}^\mathbb{L}(\vec{x}_2) \\ &= \sum_{\vec{x}_1 \otimes \vec{x}_2 = x_1 \otimes \cdots \otimes x_k} (-1)^{|\vec{x}|'+1} m^{Fu}(\vec{x}_1, m^{Fu}(\vec{x}_2, \bullet)). \end{aligned}$$

The last term equals to

$$-m_1^{Fu} m_{k+1}^{Fu}(x_1, \dots, x_k, \bullet) - (-1)^{|x_1|'+\cdots+|x_k|'+1} m_{k+1}^{Fu}(x_1, \dots, x_k, -m_1^{Fu}(\bullet)).$$

Summarizing, the equation (5.2) is nothing but an A_∞ -relation of the Fukaya category, and we are done. \square

For compact toric Fano manifolds, the above construction indeed gives homological mirror symmetry.

Theorem 5.5 ([6]). *For any compact toric Fano manifold X , the localized mirror functor*

$$\mathcal{LM}^{\mathbb{L}} : Fu(X) \rightarrow \overline{MF_{A_\infty}}(\mathfrak{P}\mathfrak{D})$$

is an A_∞ -equivalence for the monotone fiber \mathbb{L} with $b = x_1 e_1 + \cdots + x_n e_n$, where $\{e_i\}$ is a basis of $H^1(\mathbb{L}, \Lambda)$ and x_i are formal variables.

6. MAIN RESULT

Recall the closed-open map $\mathcal{CO} : QH^*(X) \rightarrow HH^*(Fu(X))$. We give a mirror counterpart $\gamma : Jac(\mathfrak{P}\mathfrak{D}) \rightarrow HH^*(\overline{MF_{A_\infty}}(\mathfrak{P}\mathfrak{D}))$ of the map \mathcal{CO} by the following lemma. Recall that the notation $\overline{MF_{A_\infty}}(\mathfrak{P}\mathfrak{D})$ means a direct sum of categories

$$\overline{MF_{A_\infty}}(\mathfrak{P}\mathfrak{D}) = \bigoplus_{\eta \in Crit(\mathfrak{P}\mathfrak{D})} MF_{A_\infty}(\hat{R}_\eta, \mathfrak{P}\mathfrak{D} - \mathfrak{P}\mathfrak{D}(\eta)).$$

Also recall that if $\mathfrak{P}\mathfrak{D}$ has isolated critical points we have a decomposition of $Jac(\mathfrak{P}\mathfrak{D})$ into local rings

$$Jac(\mathfrak{P}\mathfrak{D}) = \bigoplus_{\eta \in Crit(\mathfrak{P}\mathfrak{D})} Jac(\mathfrak{P}\mathfrak{D}; \eta).$$

For simplicity let $\mathcal{A} := Fu(X)$, $\mathcal{B} := \overline{MF_{A_\infty}}(\mathfrak{P}\mathfrak{D})$ and $\mathcal{B}_\eta := MF_{A_\infty}(\hat{R}_\eta, \mathfrak{P}\mathfrak{D} - \mathfrak{P}\mathfrak{D}(\eta))$.

Lemma 6.1. *The following map is a well-defined ring isomorphism:*

$$\gamma : Jac(\mathfrak{P}\mathfrak{D}) \rightarrow \bigoplus_{\eta \in Crit(\mathfrak{P}\mathfrak{D})} HH^*(\mathcal{B}_\eta), [r] \mapsto \sum_{\eta \in Crit(\mathfrak{P}\mathfrak{D})} [\phi_{r_\eta}]$$

where

$$[r] = \sum_{\eta \in Crit(\mathfrak{P}\mathfrak{D})} [r_\eta]$$

for $[r_\eta] \in Jac(\mathfrak{P}\mathfrak{D}; \eta)$, and

$$\phi_{r_\eta} = \bigoplus_{X \in Ob(\mathcal{B}_\eta)} r_\eta \cdot \text{id}_X \in CH^*(\mathcal{B}_\eta)$$

which is a Hochschild cocycle with length zero part only.

Proof. Let $\mathcal{C} = MF_{A_\infty}(\hat{R}, \mathfrak{P}\mathfrak{D})$, where \hat{R} is the completion of R at the ideal \mathfrak{m} which supports a critical point of $\mathfrak{P}\mathfrak{D}$. Recall from [9] that the isomorphism $\gamma : Jac(\mathfrak{P}\mathfrak{D}) \rightarrow HH^*(\mathcal{C})$ is induced by the map

$$Jac(\mathfrak{P}\mathfrak{D}) \rightarrow \mathbb{R}Hom_{\mathcal{C}-\mathcal{C}}(\mathcal{C}_\Delta, \mathcal{C}_\Delta), [r] \mapsto (\mathcal{C}_\Delta \xrightarrow{r} \mathcal{C}_\Delta).$$

Replacing with the bar resolution $B(\mathcal{C}_\Delta)$, the map $r : \mathcal{C}_\Delta \rightarrow \mathcal{C}_\Delta$ lifts to the map of complexes of \mathcal{C} -bimodules

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \bigoplus_{B_0, B_1 \in \mathcal{C}} \mathcal{C}(-, B_0) \otimes \mathcal{C}(B_0, B_1) \otimes \mathcal{C}(B_1, -) & \longrightarrow & \bigoplus_{B \in \mathcal{C}} \mathcal{C}(-, B) \otimes \mathcal{C}(B, -) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow r \cdot m_2^{\mathcal{C}} & & \\ 0 & \longrightarrow & 0 & \longrightarrow & \mathcal{C}_\Delta & \longrightarrow & 0 \end{array}$$

and by the following identification

$$\begin{aligned} \text{hom}_{\mathcal{C}-\mathcal{C}}(\mathcal{C}(-, B_0) \otimes \mathcal{C}(B_0, B_1) \otimes \cdots \otimes \mathcal{C}(B_{p-1}, B_p) \otimes \mathcal{C}(B_p, -), X) \\ \simeq \text{hom}_k(\mathcal{C}(B_0, B_1) \otimes \cdots \otimes \mathcal{C}(B_{p-1}, B_p), X(B_0, B_p)) \end{aligned}$$

the above map of complexes changes to

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \bigoplus_{B_0, B_1 \in \mathcal{C}} \mathcal{C}(B_0, B_1) & \longrightarrow & \bigoplus_{B \in \mathcal{C}} k & \longrightarrow & 0 \\ & & \downarrow & & \downarrow r & & \\ 0 & \longrightarrow & 0 & \longrightarrow & \bigoplus_{B \in \mathcal{C}} \mathcal{C}(B, B) & \longrightarrow & 0 \end{array}$$

where the map r sends 1 to $r \cdot \text{id}_B$ for each B . \square

We justify that the map γ is indeed the closed-open map on the B -model as follows: given the open-closed map $\mathcal{CO} : HH_*(Fu(X)) \rightarrow QH^*(X)$ on the A -model, let $\sigma \in HH_*(Fu(X))$ be the preimage of the unit $1 \in QH^*(X)$. For $\psi = a_0 \otimes a_1 \otimes \cdots \otimes a_n \in HH_*(\mathcal{C})$ with \mathcal{C} being an A_∞ -category, recall the *cap product*

$$- \cap \psi : HH^*(\mathcal{C}) \rightarrow HH_*(\mathcal{C}),$$

$$\phi \cap \psi := \sum (-1)^* m_*^{\mathcal{C}}(a_{l+1}, \dots, a_i, \phi(a_{i+1}, \dots, a_j) \otimes a_{j+1} \otimes \cdots \otimes a_n \otimes \underline{a_0} \otimes \cdots \otimes a_k) \otimes a_{k+1} \otimes \cdots \otimes a_l$$

where

$$\begin{aligned} \star &= |\phi|'(|a_0|' + |a_1|' + \cdots + |a_i|') \\ &\quad + (|a_{l+1}|' + \cdots + |\phi(a_{i+1}, \dots, a_j)|' + \cdots + |a_n|')(|a_0|' + |a_1|' + \cdots + |a_l|'). \end{aligned}$$

Via the cap product, $HH_*(\mathcal{C})$ is equipped with a module structure over $HH^*(\mathcal{C})$. Then we have the following important fact:

$$\mathcal{CO} \circ (\cap \sigma) \circ \mathcal{CO} = \text{id}.$$

In other words, $\mathcal{CO} = (\cap \sigma)^{-1} \circ \mathcal{CO}^{-1}$ as a linear map.

Now consider the “open-closed map” on the B -model

$$\xi_\eta : HH_*(\mathcal{B}_\eta) \rightarrow Jac(\mathfrak{P}\mathfrak{D}; \eta)$$

which is explained in [7] (for original descriptions see [20, 23]). Since it is an isomorphism, let $\psi := \xi_\eta^{-1}(1) \in HH_*(\mathcal{B}_\eta)$. Let $\psi = \underline{a_0} \otimes a_1 \otimes \cdots \otimes a_n$. Pick $r_\eta \in Jac(\mathfrak{P}\mathfrak{D}; \eta)$. Since $\gamma(r_\eta) = \bigoplus_{X \in Ob(\mathcal{B}_\eta)} r_\eta \cdot \text{id}_X$, we have the following computation of cap products:

$$\gamma(r_\eta) \cap \psi = m_2(r_\eta \cdot \text{id}, \underline{a_0}) \otimes a_1 \otimes \cdots \otimes a_n = r_\eta \cdot \underline{a_0} \otimes a_1 \otimes \cdots \otimes a_n,$$

so the following is straightforward:

$$(6.1) \quad \xi_\eta \circ (\cap \psi) \circ \gamma = \text{id}.$$

We can summarize the meaning of the above arguments as follows: since $HH_*(\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D}))$ is a module over $HH^*(\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D}))$ which is isomorphic to $Jac(\mathfrak{P}\mathfrak{D})$ via γ , $HH_*(\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D}))$ is also equipped with a $Jac(\mathfrak{P}\mathfrak{D})$ -module structure. On the other hand, there is also a canonical $Jac(\mathfrak{P}\mathfrak{D})$ -module structure on $HH_*(\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D}))$ by multiplication. The above condition (6.1) tells us that these two $Jac(\mathfrak{P}\mathfrak{D})$ -module structures coincide.

Theorem 6.2. *Let X be a compact toric Fano manifold and $\mathfrak{P}\mathfrak{D}$ be its LG superpotential. Then the following diagram commutes:*

$$\begin{array}{ccc}
 QH^*(X) & \xrightarrow{\mathcal{CO}} & HH^*(Fu(X)) \\
 \downarrow \mathfrak{fs} & & \searrow \xrightarrow{[L_{\mathcal{M}}^1]} \\
 & & Hom_{\mathcal{A}-\mathcal{B}-bimod}(\mathcal{M}, \mathcal{M}) \\
 & & \nearrow \xrightarrow{[R_{\mathcal{M}}^1]} \\
 Jac(\mathfrak{P}\mathfrak{D}) & \xrightarrow{\gamma} & HH^*(\overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D}))
 \end{array}$$

so \mathfrak{fs} becomes a ring isomorphism.

Proof. Let $\mathcal{LM}^{\mathbb{L}} : Fu(X) \rightarrow \overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D})$ be the localized mirror functor whose reference \mathbb{L} is the monotone torus fiber. Recall our notations $\mathcal{A} = Fu(X)$ and $\mathcal{B} = \overline{MF}_{A_\infty}(\mathfrak{P}\mathfrak{D})$. For a compact toric Fano manifold X , $\mathcal{LM}^{\mathbb{L}}$ is actually an A_∞ -quasiequivalence by the result of [6], and the Morita bimodules are given by

$$\mathcal{M} = (\mathcal{LM}^{\mathbb{L}} \otimes 1)^* \mathcal{B}_\Delta, \quad \mathcal{N} = (1 \otimes \mathcal{LM}^{\mathbb{L}})^* \mathcal{B}_\Delta$$

so that $\mathcal{M} \otimes_{\mathcal{B}} \mathcal{N} \simeq \mathcal{A}_\Delta$ and $\mathcal{N} \otimes_{\mathcal{A}} \mathcal{M} \simeq \mathcal{B}_\Delta$.

Let $\mathcal{F}_\alpha := (L_{\mathcal{M}}^1 \circ \mathcal{CO})(\alpha) \in hom_{\mathcal{A}-\mathcal{B}}^\bullet(\mathcal{M}, \mathcal{M})$ for $\alpha \in QH^*(X)$. Let $(a_i : L_i \rightarrow L_{i+1})_{i=1, \dots, r}$ be a tuple of morphisms in $Fu_\lambda(X)$, fix a reference Lagrangian \mathbb{L} which has a critical value λ and let

$$\underline{m} \in \mathcal{M}(L_{r+1}, P) = \mathcal{B}_\Delta(\mathcal{LM}^{\mathbb{L}}(L_{r+1}), P) = \mathcal{B}(\mathcal{LM}^{\mathbb{L}}(L_{r+1}), P)$$

for some $P \in Ob(\mathcal{B})$.

$$\begin{aligned}
 & \mathcal{F}_\alpha^{r|1|0}(a_1, \dots, a_r, \underline{m}) \\
 &= L_{\mathcal{M}}^1(\mathcal{CO}(\alpha))(a_1, \dots, a_r, \underline{m}) \\
 &= \sum (-1)^{|a_1|' + \dots + |a_i|'} \mu_{\mathcal{M}}^{i+r-j+1|1|0}(a_1, \dots, a_i, \mathbf{q}(\alpha; a_{i+1}, \dots, a_j), a_{j+1}, \dots, a_r, \underline{m}) \\
 (6.2) \quad &= \sum (-1)^{|a_1|' + \dots + |a_i|'} m_2^{\mathcal{B}} \left(\mathcal{LM}^{\mathbb{L}}(a_1, \dots, a_i, \mathbf{q}(\alpha; a_{i+1}, \dots, a_j), a_{j+1}, \dots, a_r), \underline{m} \right)
 \end{aligned}$$

and $\mathcal{F}_\alpha^{r|1|s} = 0$ if r and s are both nonzero, since \mathcal{B} has A_∞ -operations only up to m_2 .

Also, for $\mathcal{G}_\alpha := (R_{\mathcal{M}}^1 \circ \gamma \circ \mathfrak{fs})(\alpha) \in hom_{\mathcal{A}-\mathcal{B}}^\bullet(\mathcal{M}, \mathcal{M})$,

$$\mathcal{G}_\alpha^{0|1|0}(\underline{m}) = R_{\mathcal{M}}^1(\gamma(\mathfrak{fs}(\alpha)))(\underline{m}) = (-1)^{|\underline{m}|} m_2^{\mathcal{B}}(\underline{m}, \mathfrak{fs}(\alpha) \cdot \text{id}_P)$$

and $\mathcal{G}_\alpha^{r|1|s} = 0$ if $r \neq 0$ or $s \neq 0$. The sign $(-1)^{|\underline{m}|}$ is due to the definition of $R_{\mathcal{M}}^p$ in Lemma 2.16.

We show that

$$\mathcal{G}_\alpha - \mathcal{F}_\alpha = \delta \xi_\alpha$$

for some $\xi_\alpha \in hom_{\mathcal{A}-\mathcal{B}}^\bullet(\mathcal{M}, \mathcal{M})$, so $[\mathcal{F}_\alpha] = [\mathcal{G}_\alpha]$ in $Hom_{\mathcal{A}-\mathcal{B}}(\mathcal{M}, \mathcal{M})$. For any $r \geq 0$, let

$$\xi_\alpha^{r|1|0}(a_1, \dots, a_r, \underline{m}) := m_2^{\mathcal{B}}(\mathbf{q}(\alpha; a_1, \dots, a_r, \bullet), \underline{m})$$

and $\xi_\alpha^{r|1|s} = 0$ if $s \neq 0$. Then $|\xi_\alpha| = 0$. Here, $\mathbf{q}(\alpha; a_1, \dots, a_r, \bullet)$ is a morphism in \mathcal{B} from $CF(L_1, \mathbb{L})$ to $CF(L_{r+1}, \mathbb{L})$, i.e. the bullet means an input in $CF(L_{r+1}, \mathbb{L})$.

First let $r \geq 1$. Then $(\mathcal{F}_\alpha - \mathcal{G}_\alpha)^{r|1|0} = \mathcal{F}_\alpha^{r|1|0}$, and continuing from (6.2),

$$\begin{aligned} & \mathcal{F}_\alpha^{r|1|0}(a_1, \dots, a_r, \underline{m}) \\ &= \sum (-1)^{|a_1|' + \dots + |a_i|'} m_2^{\mathcal{B}}(\mathcal{LM}^{\mathbb{L}}(a_1, \dots, a_i, \mathbf{q}(\alpha; a_{i+1}, \dots, a_j), a_{j+1}, \dots, a_r), \underline{m}) \\ &= \sum (-1)^{|a_1|' + \dots + |a_i|'} m_2^{\mathcal{B}}(m_{i+r-j+1}^{\mathcal{A}}(a_1, \dots, a_i, \mathbf{q}(\alpha; a_{i+1}, \dots, a_j), \dots, a_r, \bullet), \underline{m}) \end{aligned} \quad (6.3)$$

$$= \sum m_2^{\mathcal{B}} \left((-1)^{|\vec{a}_1|' + 1} \mathbf{q}(\alpha; \vec{a}_1, m^{\mathcal{A}}(\vec{a}_2, \bullet)) + (-1)^{|\vec{a}'_1|' + 1} \mathbf{q}(\alpha; \vec{a}'_1, m^{\mathcal{A}}(\vec{a}'_2), \vec{a}'_3, \bullet), \underline{m} \right) \quad (6.4)$$

$$+ \sum (-1)^{|\vec{a}_1|' + 1} m_2^{\mathcal{B}} \left(m^{\mathcal{A}}(\vec{a}_1, \mathbf{q}(\alpha; \vec{a}_2, \bullet)), \underline{m} \right) \quad (6.5)$$

$$\pm \sum m_2^{\mathcal{B}} \left(\mathbf{q}(\alpha; a_1, \dots, a_r, \bullet, m_0^{\mathbb{L}}(1)), \underline{m} \right) \quad (6.6)$$

$$\pm \sum m_2^{\mathcal{B}} \left(m_{r+2}^{\mathcal{A}}(a_1, \dots, a_r, \bullet, \mathbf{q}_0^{\mathbb{L}}(\alpha)), \underline{m} \right).$$

Recall that the identity at (6.3) is given by the formula (3.1). Also observe that

$$(6.3) = -(-1)^{|\xi_\alpha|} (\xi_\alpha \circ \hat{\mu}_{\mathcal{M}})(a_1, \dots, a_r, \underline{m}).$$

On the other hand, the following also holds

$$(6.4) = -(\mu_{\mathcal{M}} \circ \hat{\xi}_\alpha)(a_1, \dots, a_r, \underline{m}),$$

by computations below:

$$\begin{aligned} & \sum m_2^{\mathcal{B}} \left((-1)^{|\vec{a}_1|' + 1} m^{\mathcal{A}}(\vec{a}_1, \mathbf{q}(\alpha; \vec{a}_2, \bullet)), \underline{m} \right) \\ &= \sum m_2^{\mathcal{B}} \left((-1)^{|\vec{a}_1|' + 1} m^{\mathcal{A}}(\vec{a}_1, \bullet) \circ \mathbf{q}(\alpha; \vec{a}_2, \bullet), \underline{m} \right) \\ &= \sum m_2^{\mathcal{B}} \left(m_2^{\mathcal{B}}(m^{\mathcal{A}}(\vec{a}_1, \bullet), \mathbf{q}(\alpha; \vec{a}_2, \bullet)), \underline{m} \right) \\ &= \sum (-1)^{|\vec{a}_1|'} m_2^{\mathcal{B}} \left(m^{\mathcal{A}}(\vec{a}_1, \bullet), m_2^{\mathcal{B}}(\mathbf{q}(\alpha; \vec{a}_2, \bullet), \underline{m}) \right) \\ &= \sum (-1)^{|\vec{a}_1|' + |\vec{a}_1|' + 1} m_2^{\mathcal{B}} \left(m^{\mathcal{A}}(\vec{a}_1, m_2^{\mathcal{B}}(\mathbf{q}(\alpha; \vec{a}_2, \bullet), \underline{m})) \right) \\ &= -(\mu_{\mathcal{M}} \circ \hat{\xi}_\alpha)(a_1, \dots, a_r, \underline{m}). \end{aligned}$$

Furthermore, (6.5) and (6.6) vanish since $m_0^{\mathbb{L}}(1)$ and $\mathbf{q}_0^{\mathbb{L}}(\alpha)$ are both constant multiples of A_∞ -unit. Hence,

$$(\mathcal{G}_\alpha - \mathcal{F}_\alpha)^{r|1|0} = (\delta \xi_\alpha)^{r|1|0} \text{ for } r > 0.$$

If $\underline{m} \in \mathcal{B}(\mathcal{LM}^{\mathbb{L}}(L), P)$, then we have

$$\begin{aligned} & (\mathcal{F}_\alpha - \mathcal{G}_\alpha)^{0|1|0}(\underline{m}) \\ &= m_2^{\mathcal{B}}(\mathcal{LM}^{\mathbb{L}}(\mathbf{q}_0^{\mathbb{L}}(\alpha)), \underline{m}) - (-1)^{|\underline{m}|} m_2^{\mathcal{B}}(\underline{m}, \mathbf{ks}^\lambda(\alpha) \cdot \text{id}_P) \\ &= m_2^{\mathcal{B}}(\mathcal{LM}^{\mathbb{L}}(\mathbf{q}_0^{\mathbb{L}}(\alpha)), \underline{m}) - \underline{m} \circ (\mathbf{ks}^\lambda(\alpha) \cdot \text{id}_P) \\ &= m_2^{\mathcal{B}} \left(m_2^{\mathcal{A}}(\mathbf{q}_0^{\mathbb{L}}(\alpha), \bullet) - (-1)^{|\bullet|} m_2^{\mathcal{A}}(\bullet, \mathbf{q}_0^{\mathbb{L}}(\alpha)), \underline{m} \right) \end{aligned} \quad (6.7)$$

and by (3.1) again,

$$(6.7)$$

$$(6.8) \quad = -m_2^{\mathcal{B}}\left(\mathbf{q}(\alpha; m_1^{\mathcal{A}}(\bullet)) + m_1^{\mathcal{A}}(\mathbf{q}(\alpha; \bullet)), \underline{m}\right)$$

$$(6.9) \quad -m_2^{\mathcal{B}}\left(\mathbf{q}(\alpha; m_0^{\mathcal{L}}(1), \bullet) + (-1)^{|\bullet|'} \mathbf{q}(\alpha; \bullet, m_0^{\mathcal{L}}(1)), \underline{m}\right)$$

and since $m_0^{\mathcal{L}}(1)$ and $m_0^{\mathcal{L}}(1)$ are both (multiples of) A_{∞} -units, (6.9) vanish. On the other hand,

$$(6.10) \quad \begin{aligned} & (\delta\xi_{\alpha})^{0|1|0}(\underline{m}) \\ &= \xi_{\alpha}(m_1^{\mathcal{B}}(\underline{m})) + m_1^{\mathcal{B}}(\xi_{\alpha}(\underline{m})) \\ &= m_2^{\mathcal{B}}\left(\mathbf{q}(\alpha; \bullet), m_1^{\mathcal{B}}(\underline{m})\right) + m_1^{\mathcal{B}}\left(m_2^{\mathcal{B}}(\mathbf{q}(\alpha; \bullet), \underline{m})\right) \end{aligned}$$

and we observe that

$$\mathbf{q}(\alpha; -m_1^{\mathcal{A}}(\bullet)) - m_1^{\mathcal{A}}(\mathbf{q}(\alpha; \bullet)) = m_1^{\mathcal{B}}(\bullet \mapsto \mathbf{q}(\alpha; \bullet)),$$

but by A_{∞} -relation on \mathcal{B} we have

$$m_2^{\mathcal{B}}\left(m_1^{\mathcal{B}}(\mathbf{q}(\alpha; \bullet)), \underline{m}\right) + m_2^{\mathcal{B}}\left(\mathbf{q}(\alpha; \bullet), m_1^{\mathcal{B}}(\underline{m})\right) + m_1^{\mathcal{B}}\left(m_2^{\mathcal{B}}(\mathbf{q}(\alpha; \bullet), \underline{m})\right) = 0,$$

hence

$$(6.8) = -(6.10) = -(\delta\xi_{\alpha})^{0|1|0}(\underline{m}).$$

Finally, we show $(\delta\xi_{\alpha})^{r|1|s} = 0$ for $s \neq 0$, so that in this case

$$(\mathcal{G}_{\alpha} - \mathcal{F}_{\alpha})^{r|1|s} = (\delta\xi_{\alpha})^{r|1|s}$$

where the left hand side is automatically zero by definition of \mathcal{G}_{α} and \mathcal{F}_{α} .

Since \mathcal{B} has no A_{∞} -operations $m_{\geq 3}$, we only need to compute $(\delta\xi_{\alpha})^{r|1|1}$.

$$(6.11) \quad \begin{aligned} & \delta\xi_{\alpha}(a_1, \dots, a_r, \underline{m}, b) \\ &= (-1)^{|a_1|' + \dots + |a_r|'} \xi_{\alpha}(a_1, \dots, a_r, m_2^{\mathcal{B}}(\underline{m}, b)) \\ &+ \xi_{\alpha}(\hat{m}(a_1, \dots, a_r), \underline{m}, b) \end{aligned}$$

$$(6.12) \quad \begin{aligned} &+ \sum (-1)^{|a_1|' + \dots + |a_i|'} \xi_{\alpha}(a_1, \dots, a_i, \mu_{\mathcal{M}}(a_{i+1}, \dots, a_r, \underline{m}), b) \\ &+ m_2^{\mathcal{B}}(\xi_{\alpha}(a_1, \dots, a_r, \underline{m}), b). \end{aligned}$$

By property $\xi_{\alpha}^{r|1|s} = 0$ for $s \neq 0$, (6.11) and (6.12) are zero, and

$$(-1)^{|a_1|' + \dots + |a_r|'} \xi_{\alpha}(a_1, \dots, a_r, m_2^{\mathcal{B}}(\underline{m}, b)) = (-1)^{|a_1|' + \dots + |a_r|'} m_2^{\mathcal{B}}\left(\mathbf{q}(\alpha; a_1, \dots, a_r, \bullet), m_2^{\mathcal{B}}(\underline{m}, b)\right),$$

$$m_2^{\mathcal{B}}\left(\xi_{\alpha}(a_1, \dots, a_r, \underline{m}), b\right) = m_2^{\mathcal{B}}\left(m_2^{\mathcal{B}}(\mathbf{q}(\alpha; a_1, \dots, a_r, \bullet), \underline{m}), b\right).$$

The sum of above two terms is zero due to the A_{∞} -relation of $m_2^{\mathcal{B}}$, hence

$$\delta\xi_{\alpha}(a_1, \dots, a_r, \underline{m}, b) = 0.$$

Summarizing all above arguments, we conclude that

$$(\mathcal{G}_{\alpha} - \mathcal{F}_{\alpha})^{r|1|s} = (\delta\xi_{\alpha})^{r|1|s}$$

for any r and s , so on the cohomology level,

$$[\mathcal{F}_{\alpha}] = [\mathcal{G}_{\alpha}].$$

□

REFERENCES

- [1] M. Abouzaid, K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, in preparation.
- [2] D. Aranha, *On Hochschild cohomology, Koszul duality and DG categories*, Mathematisch-Naturwissenschaftliche Fakultät der Rheinischen Friedrich-Wilhelms-Universität Bonn.
- [3] V. Batyrev, *Quantum cohomology rings of toric manifolds*, Journées de Géométrie Algébrique d'Orsay, 1992, Astérisque 218, 9-34.
- [4] P. Candelas, X. de la Ossa, P. Green and L. Parkes, *A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, Nucl. Phys., B359(1991), pp. 21-74.
- [5] C.-H. Cho, H. Hong and S.-C. Lau, *Localized mirror functor for Lagrangian immersions, and homological mirror symmetry for $\mathbb{P}_{a,b,c}^1$* , J. Differential Geom. 106 (2017), no. 1, 45-126.
- [6] C.-H. Cho, H. Hong and S.-C. Lau, *Localized mirror functor constructed from a Lagrangian torus*, arXiv:1406.4597.
- [7] C.-H. Cho, S. Lee and H.-S. Shin, *Open string pairing and homological mirror symmetry*, in progress.
- [8] C.-H. Cho and Y.-G. Oh, *Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds*, Asian J. Math. 10 (2006), no. 4, 773-814.
- [9] T. Dyckerhoff, *Compact generators in categories of matrix factorizations*, Duke Math. J. 159 (2011), no. 2, 223-274.
- [10] J. Evans and Y. Lekili, *Generating the Fukaya categories of Hamiltonian G -manifolds*, arXiv:1507.05842.
- [11] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian intersection Floer theory: anomaly and obstruction. Part I*, AMS/IP Studies in Advanced Mathematics, vol. 46, American Mathematical Society, Providence, RI, 2009.
- [12] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory on compact toric manifolds I*, Duke Math. J. 151 (2010), no. 1, 23-174.
- [13] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory for compact toric manifolds II: bulk deformations*, Selecta Math. (N.S.) 17 (2011), no. 3, 609-711.
- [14] K. Fukaya, Y.-G. Oh, H. Ohta and K. Ono, *Lagrangian Floer theory and mirror symmetry on compact toric manifolds*, Astérisque No. 376 (2016), vi+340 pp.
- [15] S. Ganatra, *Symplectic cohomology and duality for the wrapped Fukaya category*, Thesis (Ph.D.)-Massachusetts Institute of Technology, 2012.
- [16] S. Ganatra, T. Perutz and N. Sheridan, *Mirror symmetry: from categories to curve counts*, arXiv:1510.03839.
- [17] E. Getzler, *Cartan homotopy formulas and the Gauss-Manin connection in cyclic homology*, Israel Math. Conf. Proc. 7 (1993), 1-12.
- [18] A. Givental, *Homological geometry and mirror symmetry*, Proceeding of the ICM 1994, Zürich, Basel, 1995, 472-480.
- [19] D. Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, Proc. Steklov Inst. Math. 2004, no. 3(246), 227-248.
- [20] E. Segal, *The closed state space of affine Landau-Ginzburg B-models*, J. Noncommut. Geom. 7 (2013), no. 3, 857-883.
- [21] P. Seidel, *Fukaya category and Picard-Lefschetz theory*, Zürich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [22] N. Sheridan, *Formulae in noncommutative Hodge theory*, arXiv:1510.30795.
- [23] D. Shklyarov, *Matrix factorizations and higher residue pairings*, Adv. Math. 292 (2016), 181-209.

KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGI-RO DONGDAEMUN-GU, SEOUL 02455, KOREA, SWLEE@KIAS.RE.KR