UNIQUENESS OF ANCIENT SOLUTIONS TO GAUSS CURVATURE FLOW ASYMPTOTIC TO A CYLINDER

BEOMJUN CHOI, KYEONGSU CHOI, AND PANAGIOTA DASKALOPOULOS

ABSTRACT. We address the classification of ancient solutions to the Gauss curvature flow under the assumption that the solutions are contained in a cylinder of bounded cross-section. For each cylinder of convex bounded cross-section, we show that there are only two ancient solutions which are asymptotic to this cylinder: the non-compact translating soliton and the compact oval solution obtained by gluing two translating solitons approaching each other from time $-\infty$ from two opposite ends.

Contents

1.	Introduction	1
2.	Preliminaries	4
3.	Convergence of solution around tip	9
4.	Uniqueness of non-compact ancient solution	13
5.	Existence of compact ancient solution	13
6.	Uniqueness of compact ancient solution	17
References		24

1. INTRODUCTION

A one-parameter family $\Sigma_t := F(M^n, t)$ of complete convex embedded hypersurfaces defined by $F: M^n \times [0, T) \to \mathbb{R}^{n+1}$ is a solution of the *Gauss* curvature flow (GCF in abbreviation) if F(p, t) satisfies

(1.1)
$$\frac{\partial}{\partial t}F(p,t) = -K(p,t)\nu(p,t),$$

where K(p,t) is the Gauss curvature of Σ_t at F(p,t), and $\nu(p,t)$ is the unit normal vector of Σ_t at F(p,t) pointing outward of the convex hull of Σ_t .

The GCF was first introduced by W. Firey [31] in 1974 as a model that describes the deformation of a compact convex body $\hat{\Sigma}_0$ embedded in \mathbb{R}^{n+1} which is subject to wear under impact from any random angle. An example can be a stone on a beach impacted by the sea. The probability of impact at any point P on the surface $\Sigma_t = \partial \hat{\Sigma}_t$ is proportional to the Gauss curvature K of Σ at P. W. Firey showed, assuming that a solution exists, that the GCF shrinks smooth, compact, strictly convex and centrally symmetric

2 BEOMJUN CHOI, KYEONGSU CHOI, AND PANAGIOTA DASKALOPOULOS

hypersurfaces embedded in \mathbb{R}^3 to round points. The existence of solutions in any dimension was established in 1985 by Tso [39]. Tso showed that under the assumption that the initial surface Σ_0 is smooth, compact and strictly convex the Gauss curvature flow admits a unique solution Σ_t which shrinks to a point at the exact time $T^* := V/4\pi$, where V is the volume enclosed by the initial surface Σ_0 . Around the same time Chow [23] proved that, under certain restrictions on the second fundamental form of the initial surface, the Gauss curvature flow shrinks smooth compact strictly convex hypersurfaces to round points. Later, in [3] Andrews showed that the Gauss Curvature flow shrinks any compact convex hypersurface in \mathbb{R}^3 to a round point. For higher dimensions $n \geq 3$, P. Guan and L. Ni in [32] obtained the convergence of the flow after rescaling to a self-shrinking soliton. K. Choi and P. Daskalopoulos [19] have recently shown that the sphere is the unique self-shrinking soliton which combined with the result in [32] shows that the only finite time singularities in the n-dimensional GCF are the spheres.

In this work we will study the ancient solutions to the GCF, that is solutions which exist for all time $t \in (-\infty, T)$, for some $T \in (\infty, \infty]$. Shrinking and translating solitons are typical important models of ancient solutions. A shrinking soliton refers a solution which homothetically shrinks to a point. A shrinking soliton which shrinks to at spatial origin at time t = 0, is of the form $\Sigma_t = (-t)^{\frac{1}{1+n}} \Sigma_{-1}$ and Σ_t satisfies $K = \frac{1}{(n+1)(-t)} \langle F, \nu \rangle$. A translating soliton refers to a solution which moves by translation $\Sigma_t = \Sigma_0 + \omega t$, along a fixed direction $\omega \in \mathbb{R}^{n+1}$, it is defined for all $t \in (-\infty, \infty)$, and satisfies $K = \langle -\nu, \omega \rangle$.

In one dimension, the GCF coincides with the *Curve Shortening Flow* (CSF in abbreviation) for curves embedded in \mathbb{R}^2 . In this case, there is only one translating soliton (up to isometries and rescaling). It is called the *Grim Reaper solution* and is given by the graphical representation $y = t - \ln \cos x$, for $(x, t) \in (-\pi/2, \pi/2) \times (-\infty, \infty)$.

The result of P. Daskalopoulos, R. Hamilton and N. Sesum [25] reveals that shrinking and translating solitons are major building blocks of ancient solutions. It was shown in [25] that the only compact convex ancient solutions to CSF are the shrinking round sphere or the Angenent oval. The latter, looks as if it is constructed by the gluing of two Grim Reapers coming from opposite ends. It is given by the implicit equation $\cos x = e^t \cosh y$ and, as $t \to -\infty$, it is approximately the intersection of the two Grim Reapers $y = (t - \ln 2) - \ln \cos x$ and $y = -(t - \ln 2) + \ln \cos x$. Moreover recently, T. Bourni, M. Langford, and G. Tinaglia [10] completed this classification by showing that the shrinking circle, the Angenent oval, the Grim Reaper, and the stationary line are the only convex ancient solutions to the curve shortening flow.

A similar classification holds true in the two-dimensional *Ricci flow*. It was shown by P. Daskalopoulos, R. Hamilton, and N. Sesum in [26], that the shrinking sphere and the King solution, are the only ancient solutions

defined on \mathbb{S}^2 . For the *mean curvature flow* and the *Ricci flow* in higher dimensions, the classifications are done under a non-collapsing along with convexity, low entropy or certain other conditions. See [13], [8], [12], [29].

In this paper, we prove the classification of ancient solutions to the Gauss curvature flow under the assumption that the solution is contained in a cylinder of bounded cross-section. The relevance of this assumption is found in the classification of translating solitons to the GCF given by J. Urbas [40, 41] where each translator is shown to be a graph on a convex bounded domain $\Omega \subset \mathbb{R}^n$. Note also that the Grim Reaper and the Angenent oval are solutions to the curve shortening flow contained in a strip, and the rotationally symmetric steady cigar soliton and the King solution are ancient solutions of the 2-dim Ricci flow asymptotic to a fixed round cylinder. However, a significant difference between those previous results and ours in this work, is that a translator exists for each $\Omega \subset \mathbb{R}^n$ (see J. Urbas [40, 41]) and hence there are *infinitely many ancient solutions*. We will show the *uniqueness* of ancient solutions having asymptotic cylinder $\Omega \times \mathbb{R}$, according to the definition below.

Definition 1.1 (Asymptotic cylinder of an ancient solution). Assume that $\Sigma_t, t \in (-\infty, T), T \in (-\infty, +\infty]$ is a complete ancient GCF solution such that $\cup_{t \in (-\infty,T)} \Sigma_t \subset \operatorname{cl}(\Omega) \times \mathbb{R}$, for a some open bounded domain $\Omega \subset \mathbb{R}^n$, and that $\cup_{t \in (-\infty,T)} \Sigma_t$ is not contained in any smaller cylinder $\operatorname{cl}(\Omega') \times \mathbb{R}$. We will then refer to the cylinder $\Omega \times \mathbb{R}$ as the *asymptotic cylinder* of the ancient solution $\Sigma_t, t \in (-\infty,T)$.

Similarly, we define the asymptotic cylinder of a time slice $\Sigma_{t'}$ by the smallest cylinder containing $\Sigma_{t'}$.

Our first result given below settles the *uniqueness* of *non-compact* ancient solutions.

Theorem 1.2 (Uniqueness of non-compact ancient solutions). Given a convex bounded $\Omega \subset \mathbb{R}^n$, the translating soliton asymptotic to $\Omega \times \mathbb{R}$ is the unique non-compact ancient solution asymptotic to $\Omega \times \mathbb{R}$. This uniqueness holds up to translations along the e_{n+1} direction and reflection about $\{x_{n+1} = 0\}$.

Regarding *compact ancient solutions*, our next result shows the existence of *ancient oval solutions* which are the analogue of the Angenent oval solution for *curve shortening flow*.

Theorem 1.3 (Existence of ancient oval solutions). Given a convex bounded $\Omega \subset \mathbb{R}^n$ with $C^{1,1}$ boundary, there exists a compact ancient solution $\Gamma_t \subset \mathbb{R}^{n+1}$ to Gauss curvature flow which is defined for all $t \in (-\infty, T)$, it becomes extinct at $T := -\frac{2V_\Omega}{\omega_n}$, and has asymptotic cylinder $\Omega \times \mathbb{R}$. Here V_Ω is the volume under the graph of the translating soliton asymptotic to $\Omega \times \mathbb{R}$ which is finite according to Lemma 5.1. Furthermore, the solution Γ_t satisfies the properties in Propositions 5.3 and 5.4.

Our last result shows that Γ_t in Theorem 1.3 is unique compact ancient solution asymptotic to $\Omega \times \mathbb{R}$.

Theorem 1.4. Given a convex bounded $\Omega \subset \mathbb{R}^n$ with $C^{1,1}$ boundary, let $\Sigma_t, t \in (-\infty, T)$ be a compact ancient solution to the Gauss curvature flow in \mathbb{R}^{n+1} which is asymptotic to $\Omega \times \mathbb{R}$. Assuming that the solution becomes extinct at time $T := -\frac{2V_{\Omega}}{\omega_n}$, there is $v \in \mathbb{R}$ such that $\Sigma_t + ve_{n+1} = \Gamma_t$, for all $t \in (-\infty, T)$, where Γ_t is the solution constructed in Theorem 1.3.

Remark 1.5. In the proofs of Theorem 1.3 and Theorem 1.4, the only use of the $C^{1,1}$ assumption on Ω is to ensure that V_{Ω} is finite. In other words, if one can show Lemma 5.1 without such an assumption, this condition can be removed from the theorems. In two theorems, the extinction time $T = -\frac{2V_{\Omega}}{\omega_n}$ is chosen so that the maximum height of Γ_t , $h(t) := \max_{x \in \Gamma_t} |x_{n+1}|$, satisfies $h(t) = \lambda |t| + o(1)$ as $t \to -\infty$. Here $\lambda := \frac{\omega_n}{2|\Omega|}$ is the speed of the translating soliton asymptotic to $\Omega \times \mathbb{R}$.

The organization of this paper is as follows: In Section 2, we give some preliminary results and define an appropriate notion of weak solution. This is needed as translating solitons defined on non-strictly convex domains are not necessarily smooth and the corresponding compact ancient solutions may not be smooth as well. Theorem 1.2-1.4 will be shown in later sections with this notion of weak solution.

In Section 3, we show Theorem 3.3, the asymptotic convergence of an ancient solution to a translating soliton, as $t \to -\infty$, if one translates the solution so that its tip is fixed. In their recent recent work [17, 18] the authors established the forward in time convergence of non-compact solutions asymptotic to a cylinder to the corresponding translating soliton. Theorem 3.3 is obtained as an application of this result, and hence we will refer to the results in [18] when they are needed. Theorem 1.2, the uniqueness of non-compact ancient solutions, will be shown in Section 4 as a consequence of Theorem 3.3 and [18].

In Section 5, we show Theorem 1.3, the existence of a compact ancient solution asymptotic to a given cylinder, by showing Propositions 5.3 and 5.4. These two propositions also establish additional properties of the constructed solution Γ_t which will be used when we show the uniqueness Theorem 1.4 by comparing Γ_t with an arbitrary ancient solution Σ_t .

In Section 6, we show Theorem 1.4, the uniqueness of a compact ancient solution asymptotic to a given cylinder. Part of our proof is inspired by the recent significant works of Bourni-Langford-Tinaglia [9, 10] where they use the rate change in the enclosed the volume as a function of time to estimate the location of the tips.

2. Preliminaries

In this section we collect some preliminary results. Throughout this paper, h_{ij} denotes the second fundamental form. For a strictly convex solution,

one may consider the inverse b^{ij} of the second fundamental form h_{ij} , which satisfies $b^{ik}h_{kj} = \delta^i_j$. Let us recall the unique existence of translating solitons by J. Urbas and denote them as follows.

Definition 2.1 (Theorem of J.Urbas [40, 41]). Given a convex bounded domain $\Omega \subset \mathbb{R}^n$, we define $u_\Omega : \Omega \to \mathbb{R}$ by the graph function of the unique translating soliton which is asymptotic to $\Omega \times \mathbb{R}$, moves in the positive e_{n+1} direction, and satisfies inf $u_\Omega(\cdot) = 0$. In other words, the hypersurface given by $\partial\{(x, x_{n+1}) \in \mathbb{R}^{n+1} : x_{n+1} > u(x)\}$ defines the translating soliton. The existence and the uniqueness is shown in [40, 41].

Remark 2.2. In the case where Ω is not a strictly convex domain, it is possible that $\limsup_{x\to x_0} u_{\Omega}(x) < \infty$, for some $x_0 \in \partial\Omega$, hence the hypersurface $\{x_{n+1} = u_{\Omega}(x)\}$ may not be always complete. This is the reason why we denote the translating soliton by $\partial\{x_{n+1} > u(x)\}$. Urbas [41] showed the existence of such solitons and their uniqueness among solutions realized in certain generalized sense. To be more specific, Urbas [41] showed if a convex function u(x) defined on Ω satisfies the translating soliton equation

(2.1)
$$\det D^2 u = \beta \left(1 + |Du|^2\right)^{\frac{n+1}{2}}$$

for some $\beta > 0$ in the sense of Alexandrov, and $|\mathbb{R}^n - Du(\Omega)| = 0$, then $u = u_{\Omega} + C$, for some constant C. We will use this characterization of soliton in the proofs of Theorem 3.3.

Definition 2.3. For given convex bounded domain $\Omega \subset \mathbb{R}^n$, let us note the speed of associated translating soliton by

(2.2)
$$\lambda := \frac{1}{|\Omega|} \left[\int_{\mathbb{R}^n} \frac{1}{(\sqrt{1+|p|^2})^{n+1}} dp \right] = \frac{\omega_n}{2|\Omega|}$$

where $\omega_n = |\mathbb{S}^n|$. One can find a derivation of this λ in [41] or the equality case of (3.4).

In [20] it was shown that a translating soliton may be weakly convex with flat sides in which case it fails to be smooth at the boundary $\partial \Omega \times \mathbb{R}$ of its asymptotic cylinder. This requires a suitable notion of weak solutions. In Definition 2.4, we define a weak solution in such a way that it satisfies global comparisons with smooth classical solutions. The existence and the uniqueness of ancient solutions will then be shown in this class of weak solutions. Here and the remaining sections, an ancient solution is assumed to be a *weak solution to the GCF* in the sense of Definition 2.4 unless otherwise stated. Also, throughout the paper, we will use $\hat{\Sigma}$ to denote the closed region which is bounded by Σ .

Definition 2.4 (Definition 2.6 in [18]). Suppose that $\hat{\Sigma}_t \subset \mathbb{R}^{n+1}$ for $t \in [T_0, T_1]$ is a one-parameter family of closed convex sets with non-empty interior. $\Sigma_t = \partial \hat{\Sigma}_t \subset \mathbb{R}^{n+1}$ is a *weak subsolution* to the GCF if the following holds: for given a smooth strictly convex solution to the GCF $\Sigma'_t = \partial \hat{\Sigma}'_t$

defined for $t \in [a,b] \subset [T_0,T_1]$ with initial data $\hat{\Sigma}'_a \subset \hat{\Sigma}_a$, $\hat{\Sigma}'_t \subset \hat{\Sigma}_t$ holds for all $t \in [a,b]$. We define a *weak supersolution* in a similar way with the opposite inclusion. $\Sigma_t = \partial \hat{\Sigma}_t$ is a *weak solution* if it is both a weak sub- and super-solution. $\Sigma_t = \hat{\Sigma}_t$ for $t \in (-\infty,T)$ is a *weak ancient solution* if Σ_t , $t \in [a,b]$, is a weak solution for all $-\infty < a \le b < T$.

The following result shows the existence and uniqueness of a weak solution starting at any convex hypersurface $\Sigma_0 = \partial \hat{\Sigma}_0 \subset \mathbb{R}^{n+1}$ which is compact or non-compact and asymptotic to a cylinder.

Theorem 2.5 (Theorem 2.7 in [18]). Let $\hat{\Sigma}_0 \subset \mathbb{R}^{n+1}$ be a convex set with non-empty interior. If $\Sigma_0 = \partial \hat{\Sigma}_0$ is compact then there is a unique weak solution Σ_t to the GCF running from Σ_0 and defined over $t \in [0,T)$ for some $T < +\infty$. If $\Sigma_0 = \partial \hat{\Sigma}_0$ is non compact and asymptotic to a cylinder $\Omega \times \mathbb{R}$, then there is a unique weak solution Σ_t to the GCF running from Σ_0 defined for all $t \in [0, +\infty)$. Moreover, each time slice Σ_t is non-compact and asymptotic to $\Omega \times \mathbb{R}$ for all $t \in [0, \infty)$.

Proof. Let us only give an outline of the proof and refer the reader to [18] for details. Let $\hat{\Sigma}_{i,0}$ be a sequence of convex sets with smooth strictly convex boundaries which strictly increases to $\hat{\Sigma}_0$. We denote by $\Sigma_{i,t}$ the unique GCF solution with initial data $\Sigma_{i,0} = \partial \hat{\Sigma}_{i,0}$ and we simply define Σ_t to be the limit of $\Sigma_{i,t}$. Since each $\Sigma_{i,t}$ is smooth it can be compared with any smooth solution or any weak solution. The existence and the uniqueness of a weak solution, as stated in the theorem, follow from this comparison principle.

Lemma 2.6. The following hold:

- (i) The limit of any monotone sequence of weak solutions is a weak solution, provided that the limit is compact.
- (ii) The comparison principle between weak solutions holds.
- (iii) $\partial_t Vol(\hat{\Sigma}_t) = -|\mathbb{S}^n| = -\omega_n$, for any compact weak solution Σ_t .

Proof. Let us begin by showing (i). Suppose $\partial \hat{\Sigma}_{i,t}$ is an increasing sequence of weak solutions and $\hat{\Sigma}_t = \operatorname{cl}(\cup_i \hat{\Sigma}_{i,t})$. Then $\Sigma_t := \partial \hat{\Sigma}_t$ is a supersolution. To show that Σ_t is a subsolution, suppose that $\hat{\Sigma}'_a \subset \hat{\Sigma}_a$ and assume, without loss of generality, that $0 \in \operatorname{int}(\hat{\Sigma}'_a)$. For $\lambda < 1$, $\lambda \Sigma'_{a+\lambda^{-(n+1)}(t-a)}$ is a smaller solution and thus $\lambda \hat{\Sigma}'_a \subset \hat{\Sigma}_{i,a}$ for large $i > i_{\lambda}$ (here we use the compactness of Σ_t). This shows that $\lambda \hat{\Sigma}'_{a+\lambda^{-(n+1)}(t-a)} \subset \hat{\Sigma}_t$. By letting $\lambda \uparrow 1$ we obtain $\hat{\Sigma}'_t \subset \hat{\Sigma}_t$. A similar argument shows the same for decreasing sequences. (ii) and (iii) follow by approximation with smooth solutions as discussed in the proof of Theorem 2.5.

Lemma 2.6, in particular, implies that in the case that Σ_0 is compact, the maximal time of existence in Theorem 2.5 is given by $T = \operatorname{Vol}(\hat{\Sigma}_0)/\omega_n$. The following is the Harnack inequality for the GCF and its consequence to graphical solutions.

Theorem 2.7 (B. Chow [24], Proposition 3.2 [18]). Let Σ_t , $t \ge 0$, be a smooth compact strictly convex solution to the GCF. Then,

(2.3)
$$\frac{1}{K}(\partial_t K - b^{ij}\nabla_i K \nabla_j K) \ge -\frac{n}{1+n}\frac{1}{t}$$

Let $x_{n+1} = u(x', t)$, be a smooth strictly convex graphical solution to the GCF which could be possibly incomplete. If the solution satisfies (2.3), then

(2.4)
$$u_{tt} \ge -\frac{n}{1+n}\frac{u_t}{t}$$

and hence, for $t_2 \ge t_1 > 0$,

(2.5)
$$u_t(\cdot, t_2) \ge \left(\frac{t_1}{t_2}\right)^{\frac{n}{1+n}} u_t(\cdot, t_1).$$

We finish this section with the following regularity result, which roughly says that a weak GCF becomes smooth on region which are away from the initial surface. A similar property holds for other degenerate equations, such as the porous medium equation. While this property is known to hold true for GCF as well, its proof doesn't seem to exist in the literature (see in [29] for a related result). We include it here for completeness. The proof uses some of main results shown in [18]. We suggest to the reader to skip the proof of this proposition at their first reading.

Proposition 2.8. Assume that $\Sigma_0 = \partial \hat{\Sigma}_0$ is a convex hypersurface which is either compact or non-compact asymptotic to a cylinder $\Omega \times \mathbb{R}$ of bounded cross-section. Let Σ_t be a weak GCF solution starting at Σ_0 . If a point $p \in \Sigma_{t'}$, for some t' > 0, is away from Σ_0 and $\hat{\Sigma}_{t'}$ has non empty interior, then the solution Σ_t is strictly convex and smooth around p in spacetime, i.e. there is $B_r(p) \subset \mathbb{R}^{n+1}$ such that $B_r(p) \cap \Sigma_T$ is smooth for $t \in (0, t']$. Moreover, the Harnack inequality (2.3) holds on smooth part of Σ_t .

Proof. Consider a sequence smooth strictly convex compact solutions $\Sigma_{i,t}$ which approximates Σ_t from the inside in the sense that $\operatorname{cl}(\cup_i \hat{\Sigma}_{i,t}) = \hat{\Sigma}_t$. See the proof of Theorem 2.5, which is Theorem 2.7 in [18], for the construction of such an approximation. Let us denote the graphical representations of lower part of $\Sigma_{i,t}$ by $x_{n+1} = u_i(x,t)$. Since $\Sigma_{i,t}$ are compact solutions, $\Sigma_{i,t}$ and $u_i(x,t)$ satisfy Theorem 2.7.

Let p be a point in $\Sigma_{t'} \setminus \Sigma_0$ for some t' > 0. After a translation we may assume that $p = (0,0) \in \mathbb{R}^{n+1}$. Next, since a convex hypersurface is locally a convex graph, after a rotation and renaming the index *i*, we may find positive constants r', δ , *L* and *M* with the following significance: all $\Sigma_{i,t'}$ enclose the sphere of radius r' centered at $(0, L) \in \mathbb{R}^{n+1}$ and the graphical representation $u_i(x, t)$ defined on $D_{r'}(0) = \{x \in \mathbb{R}^n : |x| \le r'\}$ for $t \in [0, t']$. Moreover on $D_{r'}(0)$, $-M \leq u_i(x,0) \leq -2\delta$ and $-\delta \leq u_i(x,t')$. Our goal is to find some $\epsilon > 0$, C_1 , C_2 such that

(2.6)
$$0 < C_2 \le \partial_t u_i(x,t) \le C_1$$
, on $(x,t) \in D_{r'/2}(0) \times [t'-\epsilon,t']$.

Once we have these graphical speed bounds. Proposition 4.3, Theorem 4.4, and the argument in Corollary 4.5 in [18] can be applied to $x_{n+1} = u_i(x,t)$ on $D_{r'/4}(0) \times [t' - \frac{\epsilon}{2}, t']$ showing positive upper and lower bound on the curvature and gradient estimate (uniform in *i*). This would give a uniform smooth estimate and would show the spacetime C^{∞} convergence of $\Sigma_{i,t}$ to Σ_t around *p*.

The main tool in showing (2.6) is the Harnack estimate. Let us fix a point $x' \in D_{r'/2}(0)$. First, we are going to show the upper bound for $\partial_t u_i(x', t')$. Using the spherical solution starting from the sphere of radius r' centered at $(0, L) \in \mathbb{R}^{n+1}$ at time t = t' as a barrier, we may find some t'' > t' such that $u_i(x', t) \leq L$ on $x' \in D_{r'/2}(0)$ and $t \leq t''$. Inequality (2.5) of Theorem 2.7, yields that for any 0 < t < t'',

$$M + L \ge u_i(x', t'') - u_i(x', t)$$

= $\int_t^{t''} \partial_t u_i(x', s) ds \ge \partial_t u_i(x', t) \int_t^{t''} \left(\frac{t}{s}\right)^{\frac{n}{1+n}} ds,$

and it gives

$$\partial_t u_i(x',t) \le \frac{M+L}{(1+n)t^{\frac{n}{1+n}}[(t'')^{\frac{1}{1+n}} - t^{\frac{1}{1+n}}]}$$

In particular, for $t \in [t'/2, t']$, we have

$$\partial_t u_i(x',t) \le C_1$$
, for some $C_1 = C_1(M+L,t',t'',n)$

proving the upper bound in (2.6)

Let us now show the lower bound of $\partial_t u_i(x', t)$ in (2.6). For $0 \le \tau \le t'/2$, the previous upper bound implies

$$u_i(x',t'-\tau) \ge u_i(x',t') - \tau C_1$$

= $u_i(x',0) + (u_i(x',t') - u_i(x',0)) - \tau C_1 \ge u(x',0) + \delta - \tau C_1.$

Hence, for $\tau \leq \min(\frac{t'}{2}, \frac{\delta}{2C_1}) =: \epsilon$, by integrating inequality (2.5) of Theorem 2.7, we obtain

$$\frac{\delta}{2} \le u(x', t' - \tau) - u(x', 0) = \int_0^{t' - \tau} \partial_t u(x', s) ds \le (1 + n)(t' - \tau) \partial_t u(x', t' - \tau)$$

which readily shows the lower bound

$$\partial_t u_i(x',t) \ge \frac{\delta}{2(1+n)t'} =: C_2, \quad \text{for } t \in [t'-\epsilon,t'].$$

This proves the desired estimate which implies the smooth convergence of $\Sigma_{i,t}$ to Σ_t around the point p.



FIGURE 1. Definition 3.1

3. Convergence of solution around tip

Throughout this section we will assume that Σ_t , $t \in (-\infty, T)$ is a weak ancient complete solution to the GCF which is asymptotic to the cylinder $\Omega \times \mathbb{R}$, as $t \to -\infty$ (see Definition 1.1). The goal is to show if we translate the solution and observe our solution around the tip region, then as $t \to -\infty$ it converges to the unique translating soliton asymptotic to $\Omega \times \mathbb{R}$.

As we mentioned earlier, an ancient solution may touch the boundary of its asymptotic cylinder $\Omega \times \mathbb{R}$ (c.f. in [20]). For this reason, some of the results in this section are written and shown in terms of $\operatorname{int}(\hat{\Sigma}_t)$.

Definition 3.1. For an ancient convex GCF solution Σ_t , $t \in (-\infty, T)$, asymptotic to the cylinder $\Omega \times \mathbb{R}$, we define

$$h^+(t) := \sup_{x \in \Sigma_t} \langle x, e_{n+1} \rangle$$
 and $h^-(t) := \inf_{x \in \Sigma_t} \langle x, e_{n+1} \rangle$

to be the maximum and minimum heights, respectively. They are both finite if Σ_t is compact. For the non-compact case, after reflection, we will assume that $-\infty < h^-(t) < +\infty$ and $h^+(t) = \infty$.

We also define $p^+(t)$ and $p^-(t) \in \Sigma_t$ to be the tips of Σ_t by the condition

(3.1)
$$\langle p^+, e_{n+1} \rangle = h^+$$
 and $\langle p^-, e_{n+1} \rangle = h^-$.

In the non-compact case we only have one tip $p^{-}(t)$.

Definition 3.2. Let $\Sigma_t = \partial \hat{\Sigma}_t$, $t \in (-\infty, T)$ be an ancient convex GCF solution, asymptotic to the cylinder $\Omega \times \mathbb{R}$. For each $t \in (-\infty, T)$, $\operatorname{int}(\hat{\Sigma}_t)$ can be represented as the region between two graphs $u^+(\cdot, t)$ and $u^-(\cdot, t)$ defined on some domain $\Omega_t \subset \Omega$ as follows:

 $\operatorname{int}(\hat{\Sigma}_t) = \{ (x', x_{n+1}) \in \mathbb{R}^{n+1} : u^-(x', t) < x_{n+1} < u^+(x', t) \text{ and } x' \in \Omega_t \}.$

Here, Ω_t is the image of the projection of $int(\hat{\Sigma}_t)$ to the hyperplane $\{x_{n+1} = 0\}$, which is an open bounded convex set. For non-compact case, we set $u^+ = \infty$.

Note that since $\Omega \times \mathbb{R}$ is the smallest open cylinder containing $\cup_t \operatorname{int}(\hat{\Sigma}_t)$, the domains Ω_t increase to Ω , as $t \to -\infty$. For the non-compact case, the last assertion in Theorem 2.5 implies $\Omega_t = \Omega$ for all t. The functions $u^+(\cdot, t)$ and $u^{-}(\cdot, t)$ are graphical solutions to the GCF, which are defined on the domain Ω_t . Before proceeding to the next theorem, recall the definition of translating soliton u_{Ω} in Definition 2.1 and the speed λ in Definition 2.3.

Theorem 3.3. Let Σ_t , $t \in (-\infty, T)$, be a complete ancient convex weak solution of GCF which is asymptotic to the cylinder $\Omega \times \mathbb{R}$. Then, as $t \to -\infty$, $\Sigma_t - h^-(t)e_{n+1}$ converges locally smoothly to the unique translating soliton $\{x_{n+1} = u_{\Omega}(x)\}$. In the case that Σ_t is compact, $\Sigma_t - h^+(t)e_{n+1}$ also converges locally smoothly to the translating soliton $\{x_{n+1} = -u_{\Omega}(x)\}$. More precisely, we have

 $u^{-}(x,t) - h^{-}(t) \rightarrow u_{\Omega}(x)$ and $u^{+}(x,t) - h^{+}(t) \rightarrow -u_{\Omega}(x)$ in $C^{\infty}_{loc}(\Omega)$ as $t \rightarrow -\infty$.

We need several lemmas before giving the proof of this theorem.

Lemma 3.4. We have $h^-(t) \to -\infty$ as $t \to -\infty$. If Σ_t is compact, then $h^+(t) \to \infty$ as $t \to -\infty$ as well. Furthermore, in both cases we have $\cup_t \operatorname{int}(\hat{\Sigma}_t) = \Omega \times \mathbb{R}$.

Proof. We give the proof assuming that Σ_t is compact. The proof in the non-compact case is similar. We first show that $h^-(t) \to -\infty$ as $t \to -\infty$ by a contradiction argument. Suppose that $h_- \geq -C$, for some $C < \infty$ for all time. This implies $\cup_t \operatorname{int}(\hat{\Sigma}_t) \subset \Omega \times [-C, \infty)$. Then we may find a translating soliton which is asymptotic to a slightly but strictly larger cylinder while containing $\Omega \times [-C, \infty)$. By comparing this soliton with our solution Σ_t , starting at large negative times $t_0 \ll -1$, we conclude that Σ_t has to be empty for each t. This is a contradiction and hence $\lim_{t\to -\infty} h^- = -\infty$. Similarly, $\lim_{t\to -\infty} h^+ = \infty$.

Let us now see that $\cup_t \operatorname{int}(\hat{\Sigma}_t) = \Omega \times \mathbb{R}$. Since $\cup_t \operatorname{int}(\hat{\Sigma}_t)$ is a convex set and its boundary contains $p^+(t)$ and $p^-(t)$ which move to opposite infinities as $t \to -\infty$, it easy to see that the sections $\cup_t \operatorname{int}(\hat{\Sigma}_t) \cap \{e_{n+1} = l\}$ have to be identical and thus $\cup_t \operatorname{int}(\hat{\Sigma}_t)$ is a convex open cylinder. Since, by assumption, $\cup_t \operatorname{int}(\hat{\Sigma}_t)$ is contained in no smaller cylinder than $\Omega \times \mathbb{R}$, we obtain the conclusion. \Box

From now on, we will concentrate on the convergence of $u^-(x,t) - h^-(t)$. The convergence of $u^+(x,t) - h^+(t)$ follows by a similar arguments. Lemma 3.4 and Proposition 2.8 imply the following regularity lemma.

Lemma 3.5 (c.f. Theorem 2.7 or [24]). An ancient weak solution Σ_t , $t \in (-\infty, T)$, satisfying the assumptions of Theorem 3.3 is smooth and strictly convex away from $\partial\Omega \times \mathbb{R}$ provided that $\hat{\Sigma}_t$ has non empty interior. Moreover, the Harnack inequality

(3.2)
$$\frac{1}{K}(\partial_t K - b^{ij}\nabla_i K \nabla_j K) \ge 0$$

holds on all points where Σ_t is smooth. As a consequence, we have

- (i) $\partial_t u^-(x,t) = \frac{K}{\langle -\nu, e_{n+1} \rangle}$ satisfies $\partial_{tt}^2 u^-(x,t) \ge 0$ on $(x,t) \in \bigcup_t (\Omega_t \times \{t\}).$
- (ii) Let $K(\nu, t)$ be the Gaussian curvature of a point on Σ_t parametrized by its outer unit normal ν . Then $\partial_t K(\nu_0, t) \ge 0$ whenever Σ_t is smooth around the point p_0 with $\nu(p_0, t) = \nu_0$.

In the following steps, we are interested in establishing lower bounds on the Gaussian curvature K for any weak ancient solution Σ_t satisfying the assumptions of the Theorem 3.3.

Lemma 3.6. Let

(3.3)
$$\beta := \lim_{t \to -\infty} \partial_t h^- = \lim_{t \to -\infty} K(-e_{n+1}, t)$$

which exists by Lemma 3.5. Then, we have $\beta \geq \lambda$, where λ is the speed of the translating soliton in Definition 2.3.

Proof. We argue by contradiction. By translating the solution in time if necessary, we may assume $\hat{\Sigma}_0$ is not empty. Suppose that $\beta < \lambda - 2\epsilon$, for some $\epsilon > 0$. Consider a strictly larger cylinder containing $\Omega \times \mathbb{R}$ whose corresponding translating soliton has the speed $\lambda - \epsilon$. Let us denote by $\bar{\Sigma}^-$ to be such a soliton moving in positive e_{n+1} direction and having $\inf_{x \in \bar{\Sigma}^-} \langle x, e_{n+1} \rangle = 0$ (namely its tip is the point $(0,0) \in \mathbb{R}^{n+1}$). Then there is $C_1 > 0$ such that $\bar{\Sigma}^- - C_1 e_{n+1}$ encloses $\Omega \times [0, \infty)$. Therefore, $\bar{\Sigma}^- + (-C_1 + h^-(t)) e_{n+1}$ encloses Σ_t for t < 0. Then, the comparison principle implies that the surface $\bar{\Sigma}^- + (-C_1 + h^-(t) + (\lambda - \epsilon)\tau) e_{n+1}$ encloses $\Sigma_{t+\tau}$ for $\tau \ge 0$. By letting $\tau = -t$, we conclude that $\bar{\Sigma}^- + (-C_1 + h^-(t) - (\lambda - \epsilon)t) e_{n+1}$ encloses Σ_0 for all t < 0. On the other hand, as $t \to -\infty$, $h^-(t) \ge (\lambda - 2\epsilon)t + o(t)$ by (3.3). Thus

$$-C_1 + h^-(t) - (\lambda - \epsilon) t \ge -C_1 - \epsilon t + o(t) \to \infty, \quad \text{as } t \to -\infty,$$

which contradicts the assumption $\hat{\Sigma}_0$ is non-empty.

Proposition 3.7. Let Σ_t , $t \in (-\infty, T)$ be an ancient solution satisfying the assumptions of the Theorem 3.3. Given any $\Omega' \subset \subset \Omega$, there is $t_0 < 0$ and c > 0 such that

$$c \le \partial_t u^- \le c^{-1}, \qquad for \ t \le t_0.$$

Proof. Let $\epsilon > 0$ be such that dist $(\Omega', \partial \Omega) = 2\epsilon > 0$. By Lemmas 3.4 and 3.6, we may choose $t_0 \ll -1$ so that the following hold for $t \leq t_0$:

- (i) $\beta(t_0 t) \le h^-(t_0) h^-(t) \le 2\beta(t_0 t).$
- (ii) If $\Omega_{t,0}$ is the cross-section of $\hat{\Sigma}_t$ at $x_{n+1} = 0$, namely we have $(\hat{\Sigma}_t \cap \{\langle x, e_{n+1} \rangle = 0\} =: \Omega_{t,0} \times \{0\})$, then $\Omega' \subset \Omega_{t,0}$ and dist $(\Omega', \partial \Omega_{t,0}) \ge \epsilon$.

From now on, assume $t \leq t_0$ and let $x' \in \Omega'$ be an arbitrary point. By the monotonicity of $\partial_t u^-(\cdot, t)$ in t which follows from 2.4, we have $\partial_t u^-(x', t) \leq \partial_t u^-(x', t_0) \leq \sup_{\Omega'} \partial_t u(\cdot, t_0) < \infty$, which proves the upper bound.

We next show the lower bound. Since Σ_t is convex, it has to enclose a cone generated by the base $\Omega_{t,0} \times \{0\}$ and the vertex $p^-(t)$. Together with property (ii) above, this implies the bound $u^-(x',t) \leq \frac{e}{C}h^-(t)$, where $C = \operatorname{diam} \Omega$ (recall that both u^- and h^- are negative). Using also that $h^-(t) \leq u^-(x',t)$, we conclude that for any $\tau_1 < \tau_2 \leq t_0$, we have

$$u(x', \tau_2) \ge h^-(\tau_2) \ge 2\beta (\tau_2 - t_0) + h^-(t_0)$$

and

$$u(x', \tau_1) \le \frac{\epsilon}{C} h^-(\tau_1) \le \frac{\epsilon}{C} (\beta (\tau_1 - t_0) + h^-(t_0)).$$

Subtracting these two inequalities and using $t_0 < 0$ and $h^-(t_0) < 0$ yields

$$u(x',\tau_2) - u(x',\tau_1) \ge \beta \left(2\tau_2 - \frac{\epsilon}{C}\tau_1\right) - \beta \left(2 - \frac{\epsilon}{C}\right)t_0 + \left(1 - \frac{\epsilon}{C}\right)h^-(t_0)$$
$$\ge \beta \left(2\tau_2 - \frac{\epsilon}{C}\tau_1\right) + h^-(t_0).$$

If we choose $\tau_1 = \frac{C(2+L)}{\epsilon}\tau_2$ for L > 0, the monotonicity of u_t^- implies $u_t^-(x',\tau_2)\left(\tau_2 - \frac{C(2+L)}{\epsilon}\tau_2\right) \ge u(x',\tau_2) - u(x',\frac{C(2+L)}{\epsilon}\tau_2) \ge L\beta(-\tau_2) + h^-(t_0)$ which gives

$$u_t^-(x',\tau_2) \ge \frac{L\beta}{\frac{C(2+L)}{\epsilon} - 1} + \frac{h^-(t_0)}{\tau_2 - \frac{C(2+L)}{\epsilon}\tau_2}.$$

Finally, taking $L \to \infty$, we obtain the desired lower bound $u_t^-(x', \tau_2) \ge \frac{\epsilon\beta}{C}$.

Proposition 3.8. Let Σ_t , $t \in (-\infty, T)$, be an ancient solution satisfying the assumptions of the Theorem 3.3. For any given sequence $\tau_i \to -\infty$, passing to a subsequence if necessary, $u_i^-(x,t) := u^-(x,t+\tau_i) - h^-(\tau_i)$ converges to $u_{\infty}(x,t)$ in $C_{loc}^{\infty}(\Omega \times \mathbb{R})$. Moreover, the limiting graphical solution $x_{n+1} = u_{\infty}(x,t)$ satisfies $\partial_t u_{\infty} \equiv \beta$, where β is as in (3.3). The solution u_{∞} represents a translating soliton which may possibly be incomplete.

Proof. Since we have the bounds on the graphical speed $u_t = K \langle -\nu, e_{n+1} \rangle^{-1}$ from Proposition 3.7, we can apply Proposition 4.3 and Theorem 4.4 in [18] as they are applied in Corollary 4.5 in [18] and obtain the following: for any given $\Omega' \subset \subset \Omega$, there is $t_0 < 0$ and C > 0 such that

$$|Du^{-}|, \lambda_{\min}^{-1}, \lambda_{\max} \leq C \text{ on } \Omega' \times (-\infty, t_0].$$

The equation of graphical GCF becomes uniformly parabolic provided we have the gradient bound, positive upper and lower curvature bounds. Thus we may pass to a limit u_i^- , $i \to \infty$, by the standard regularity theory of parabolic equations, obtaining a graphical eternal solution $u_{\infty}(x,t)$. In view of (i) in Lemma 3.5, $\partial_t u_{\infty}(x,t)$ must be independent of t, that is $\partial_t u_{\infty}(x,t) = \beta_{\infty}(x)$. Furthermore, the fact that $|Du_{\infty}(\cdot,t)|$ is bounded, globally in time, on every compact set $\Omega' \subset \subset \Omega$, implies that $\beta_{\infty}(\cdot)$ has to be a constant. From Lemma 3.6, we conclude that $\partial_t u_{\infty} \equiv \beta$.

Proof of Theorem 3.3. By Proposition 3.8, $u_{\infty}(x,t) = u_{\infty,0}(x) + \beta t$ where $u_{\infty,0}(\cdot) := u(\cdot,0)$. It remains to prove that $u_{\infty,0}(\cdot) = u_{\Omega}(\cdot)$. By the

characterization of u_{Ω} given after Definition 2.1, it suffices to show $|\mathbb{R}^n - Du_{\infty,0}(\Omega)| = 0$. Note that

$$u_t \equiv \beta = (1 + |Du_{\infty,0}|^2)^{\frac{1}{2}} \left[\frac{\det D^2 u_{\infty,0}}{(1 + |Du_{\infty,0}|^2)^{\frac{n+2}{2}}} \right] \text{ on } \Omega.$$

This implies

(3.4)
$$\beta|\Omega| = \int_{\Omega} \frac{\det D^2 u_{\infty,0}}{(1+|Du_{\infty,0}|^2)^{\frac{n+2}{2}-\frac{1}{2}}} = \int_{Du_{\infty,0}(\Omega)} \frac{1}{(\sqrt{1+|p|^2})^{n+1}} \\ \leq \int_{\mathbb{R}^n} \frac{1}{(\sqrt{1+|p|^2})^{n+1}} = \lambda |\Omega|.$$

We have shown $\beta \geq \lambda$ in Lemma 3.6 and therefore the equality must hold in (3.4). In particular, this shows $\beta = \lambda$ and $|\mathbb{R}^n - Du_{\infty,0}(\Omega)| = 0$.

4. UNIQUENESS OF NON-COMPACT ANCIENT SOLUTION

We are ready to give a proof of the uniqueness of non-compact ancient solution.

Proof of Theorem 1.2. Let Σ_t be an ancient solution as in the statement of the theorem. By Theorem 2.5, the solution exist for all $t \in (-\infty, \infty)$ and Σ_t is asymptotic to $\Omega \times \mathbb{R}$ for each time slice. i.e. the domain of graphical representation of Σ_t does not change over time. Therefore, we may represent Σ_t by a graph $\partial\{(x, x_{n+1}) \in \mathbb{R}^{n+1} : x_{n+1} \geq u^-(x, t)\}$ for $(x, t) \in \Omega \times (-\infty, \infty)$.

The main result, Theorem 1.1, in [18] proves the forward-in-time convergence to the soliton $u_{\Omega}(x)$, namely

$$u^{-}(x,t) - h^{-}(t) \to u_{\Omega}(x)$$
 in $C^{\infty}_{loc}(\Omega)$, as $t \to \infty$.

Let us fix an arbitrary $x' \in \Omega$. Together with Theorem 3.3,

$$\lim_{t\to\infty} u^-_t(x',t) = \lim_{t\to-\infty} u^-_t(x',t) = \lambda$$

where λ is the speed of the translating soliton u_{Ω} . By (i) Lemma 3.5, $u_t^- \equiv \lambda$ on $(x,t) \in \Omega \times \mathbb{R}$ showing Σ_t is a translating soliton with the speed λ . We may repeat the same argument in (3.4), while β replaced by λ , to conclude $|\mathbb{R}^n - Du^-(\cdot, t)(\Omega)| = 0$ and hence $u^-(x,t) = u_{\Omega}(x) + \lambda t + C$, for some constant C.

5. EXISTENCE OF COMPACT ANCIENT SOLUTION

Let $\Omega \subset \mathbb{R}^n$ is a bounded convex open domain. In this section we will construct an ancient compact solution of GCF which has asymptotic cylinder $\Omega \times \mathbb{R}$, as $t \to -\infty$; (see Definition 1.1).

14 BEOMJUN CHOI, KYEONGSU CHOI, AND PANAGIOTA DASKALOPOULOS

We recall that u_{Ω} denotes the translator associated with the domain Ω satisfying $\inf_{x\in\Omega} u_{\Omega}(x) = 0$. For the construction of compact ancient solutions we need to show that the volume under the translating soliton $V_{\Omega} := \int_{\Omega} u_{\Omega}(x) dx$ is finite. Although this is expected to hold for any compact convex domain Ω with no further regularity assumptions on $\partial\Omega$, we could show this under $C^{1,1}$ boundary condition.

Lemma 5.1. Assume that $\Omega \subset \mathbb{R}^n$ be a convex bounded open domain with $C^{1,1}$ boundary. Let $x_{n+1} = u_{\Omega}(x)$, $x \in \Omega$, be the translating soliton associated with the domain Ω and having $\inf_{x \in \Omega} u_{\Omega}(x) = 0$. Then, the volume under the translating soliton is finite, i.e. we have

(5.1)
$$V_{\Omega} := \int_{\Omega} u_{\Omega}(x) dx < \infty.$$

Remark 5.2. If this lemma is shown without $C^{1,1}$ assumption, it also proves Theorem 1.3 and Theorem 1.4 without $C^{1,1}$ assumption on $\partial\Omega$.

Proof of Lemma 5.1. The basic strategy is to find a supersolution of the graphical translating soliton equation

$$\frac{\det D^2 \phi}{(1+|D\phi|^2)^{\frac{n+1}{2}}} \le \lambda = \frac{\omega_n}{2|\Omega|}$$

which is integrable near the boundary of Ω .

Assume first that $\partial\Omega$ is smooth. Before going into the details, let us recall some properties of the distance function d(x) from any point $x \in \Omega$ to $\partial\Omega$. The function d(x) is well defined on $\{y \in \Omega : d(y,\partial\Omega) < \lambda_0\}$, $\lambda_0 := \sup_{y \in \partial\Omega} \lambda_{\max}(y)$, where at each $y \in \Omega$, $\lambda_{\max}(y) := \max_{i=1,\dots,n-1} \lambda_i(y)$ denotes the maximum of the principal curvatures of $\partial\Omega$ at y. Furthermore, d(x) is a smooth function in this tubular neighborhood.

Let $x \in \Omega$ be a point in this neighborhood and $\pi(x) \in \partial\Omega$ be the point such that $|\pi(x) - x| = \text{dist}(x, \partial\Omega)$. If we denote by λ_i , $i = 1, \ldots, n-1$ the principal curvatures of the hypersurface $\partial\Omega \subset \mathbb{R}^n$ at $\pi(x)$, then with respect to the orthonormal basis $\{(e_i)_{i=1}^{n-1}, -\frac{\pi(x)-x}{|\pi(x)-x|}\}$ of \mathbb{R}^n , we have

$$Dd(x) = (0, \dots, 0, 1)$$

and

$$D^{2}d(x) = \left[\begin{array}{c|c} \left[\operatorname{diag}\left(-\frac{\lambda_{i}}{1-\lambda_{i}d}\right)\right]_{n-1\times n-1} & \left[0\right]_{n-1\times 1}\\ \hline \left[0\right]_{1\times n-1} & 0\end{array}\right].$$

Define $\phi(x) = -L \log d(x)$ as our test function. Then in this neighborhood we have

(5.2)
$$\frac{\det D^2 \phi}{(1+|D\phi|^2)^{\frac{n+1}{2}}} = \frac{1}{(1+\frac{L^2}{d^2})^{\frac{n+1}{2}}} \frac{L^n}{d^{n+1}} \prod_{n=1}^{n-1} \frac{\lambda_i(\pi(x))}{1-\lambda(\pi(x)) d(x)} \le \frac{1}{L} \prod_{n=1}^{n-1} \frac{\lambda_i(\pi(x))}{1-\lambda_i(\pi(x)) d(x)}.$$

Assume next that $\partial\Omega$ is in $C^{1,1}$ and take a strictly monotone increasing sequence $\{\Omega_m\}$ of convex domains which approximates Ω from the inside in such a way that each $\partial\Omega_m$ is smooth and $\sup_{y\in\partial\Omega_m}\lambda_{\max}(y) < 2\lambda_0$. Here strictly monotone means $\Omega_m \subset \subset \Omega_{m+1}$. We may also assume that dist $(x,\partial\Omega) < \frac{1}{4\lambda_0}$ for all $x \in \partial\Omega_1$. Set $M := \sup_{x\in\Omega_1} u(x) > 0$ and define the functions

$$\phi_m(x) = -\frac{(4\lambda_0)^{n-1}}{\lambda} \log \frac{(\operatorname{dist}(x, \partial \Omega_m))}{\operatorname{diam} \Omega} + M, \quad \text{for } x \in \Omega_m \setminus \Omega_1.$$

Then, by our choice of Ω_m , each ϕ_m is smooth in the interior of $\Omega_m \setminus \Omega_1$. Furthermore, from (5.2) we have

$$\frac{\det D^2 \phi_m}{(1+|D\phi_m|^2)^{\frac{n+1}{2}}} \le \frac{\lambda}{(4\lambda_0)^{n-1}} \left(\frac{2\lambda_0}{1-\frac{2\lambda_0}{4\lambda_0}}\right)^{n-1} \le \lambda.$$

We will next compare ϕ_n with u_Ω to conclude that $V_\Omega < \infty$. Since $\phi_n \ge u$ on $\partial \Omega_1$ and it becomes infinite on $\partial \Omega_m$, the comparison principle implies $\phi_m \ge u$ in the interior of $\Omega_m \setminus \Omega_1$. Note that ϕ_m converges locally uniformly to $\phi := -\frac{(4\lambda_0)^{n-1}}{\lambda} \log(\operatorname{dist}(x,\partial\Omega)) + M$ on $\Omega \setminus \Omega_1$, which implies $\phi(x) \ge u(x)$ in this region. Since ϕ is integrable on $\Omega \setminus \Omega_1$, this implies $\int_{\Omega} u = V_\Omega$ is finite.

Proof of Theorem 1.3. Recall that the speed of the translator u_{Ω} defined on the domain Ω is given by $\lambda = \frac{\omega_n}{2|\Omega|}$. Theorem 1.3 is implied by two propositions below.

Proposition 5.3. Let $\Omega \subset \mathbb{R}^n$ be a convex bounded domain with $C^{1,1}$ boundary. Then, there is a compact weak ancient solution Γ_t of the Gauss curvature flow, defined on $t \in (-\infty, T)$ with $T := -\frac{2V_\Omega}{\omega_n}$, such that

- (i) Vol $(\hat{\Gamma}_t) = -\omega_n t 2V_{\Omega}$, where V_{Ω} is given by (5.1);
- (ii) Γ_t has reflection symmetry with respect to $x_{n+1} = 0$;
- (iii) Γ_t is contained in $\Omega \times \mathbb{R}$, but not in a smaller cylinder, i.e. Γ_t is asymptotic to $\Omega \times \mathbb{R}$;
- (iv) Γ_t is smooth in the interior of $\Omega \times \mathbb{R}$, for t < T and satisfies the differential Harnack inequality.

Proof. For our given bounded convex domain Ω , denote by u_{Ω} the graph of translating soliton corresponding to the domain Ω having speed $\lambda := \frac{\omega_n}{2|\Omega|}$ and satisfying $\inf_{\Omega} u_{\Omega}(x) = 0$ (see Definition 2.1). To simplify the notation, from now on we will denote $u_{\Omega}(x)$ simply by u(x).

The graphs $x_{n+1} = u(x) + \lambda t$ and $x_{n+1} = -u(x) - \lambda t$, $t \in \mathbb{R}$, define translating solitons, moving in opposite directions and having tips at a distance $2\lambda |t|$ from each other. The basic idea here is to construct our solution Γ_t as limit of hypersufaces which for $t \ll -1$ are approximated by the boundary of the region $\{x_{n+1} > u(x) + \lambda t\} \cap \{x_{n+1} < -u(x) - \lambda t\}$.

To make this rigorous, for any s < 0 define $\Gamma_{s,0}$ to be the convex region which is bounded between the hypersurfaces $x_{n+1} = u(x) - \lambda |s|$ and $x_{n+1} = -u(x) + \lambda |s|$. By Lemma 5.1, we can deduce that

(5.3)
$$(\operatorname{Vol}(\widehat{\Gamma}_{s,0}) - \omega_n |s|) \uparrow -2V_{\Omega}, \quad \text{as } s \to -\infty.$$

Note that $\partial \hat{\Gamma}_{\tau,0}$, viewed as a one-parameter family of hypersurfaces in τ , is a weak subsolution to the GCF.

Let $\Gamma_{s,t}$, $t \in [0, \tau_s)$ with $\tau_s = \operatorname{Vol}(\widehat{\Gamma}_{s,0})/\omega_n$, be the weak solution to the GCF starting from $\Gamma_{s,0}$. Consider the time translated solutions $\Gamma_{s,t-s}$. $t \in [s, \tau_s + s)$. For each fixed t, we claim that $\Gamma_{s,t-s}$, $s \leq t$, are monotone decreasing as $s \to -\infty$: for $s_1 < s_2 < 0$, by the comparison principle between $\Gamma_{s_1,t}$ and $\Gamma_{s_1+t,0}$, Γ_{s_1,s_2-s_1} is contained in $\Gamma_{s_2,0}$. Again by the comparison principle between $\Gamma_{s_1,s_2-s_1+\tau}$ and $\Gamma_{s_2,\tau}$, we conclude $\Gamma_{s_1,t-s_1}$ is contained in $\Gamma_{s_2,t-s_2}$. In view of (1) Lemma 2.6, weak solutions $\Gamma_{s,t-s}$ monotonically converge to a weak solution Γ_t as $s \to -\infty$.

(5.4)
$$\Gamma_{s,t-s} \downarrow \Gamma_t, \quad \text{as } s \to -\infty.$$

Since $\lim_{s\to-\infty} \operatorname{Vol}(\hat{\Gamma}_{s,t-s}) = -t \,\omega_n - 2V_{\Omega}$, $\hat{\Gamma}_t$ has non empty interior for $t < -\frac{2V_{\Omega}}{\omega_n}$ and has empty interior for $t > -\frac{2V_{\Omega}}{\omega_n}$. This defines the ancient solution Γ_t for $t \in (-\infty, -\frac{2V_{\Omega}}{\omega_n})$.

We will now see that Γ_t satisfies properties (i)-(iv) in the statement of our theorem. Properties (i) and the reflection symmetry property (ii) clearly hold by construction. Furthermore, property (iv) is just a consequence of Lemma 3.5. It remains to show property (iii). By construction, $\hat{\Gamma}_t$ is contained in $\hat{\Gamma}_{t,0}$ which is contained in $\Omega \times \mathbb{R}$. Hence, $\hat{\Gamma}_t$ is contained in $\Omega \times \mathbb{R}$. Suppose there is a smaller $\Omega' \subset \Omega$ such that $\hat{\Gamma}_t$ is contained in $\Omega' \times \mathbb{R}$ for all t. Since $\hat{\Gamma}_t \subset \hat{\Gamma}_{t,0}$,

$$\sup_{\hat{\Gamma}_t} x_{n+1} - \inf_{\hat{\Gamma}_t} x_{n+1} \le \sup_{\hat{\Gamma}_{t,0}} x_{n+1} - \inf_{\hat{\Gamma}_{t,0}} x_{n+1} = 2\lambda(-t).$$

Therefore, $\operatorname{Vol}(\hat{\Gamma}_t) \leq 2\lambda(-t)|\Omega'| = \frac{|\Omega'|}{|\Omega|}(-t\omega_n)$. On the other hand, we know that $\operatorname{Vol}(\hat{\Gamma}_t) = -t\omega_n - 2V_{\Omega}$. If $|\Omega'| < |\Omega|$, we have a contradiction by taking $t \to -\infty$ in the above inequality. This shows there is no such smaller Ω' .

We next provide some extra properties of the solution Γ_t constructed above. Those properties will be used in the proof of our uniqueness Theorem 1.4 in the next section.

Proposition 5.4. The constructed ancient solution Γ_t , $t \in (-\infty, T)$, with $T := -\frac{2V_{\Omega}}{\omega_n}$, satisfies $\left(\sup_{\Gamma_t} |x_{n+1}| - \lambda |t|\right) \uparrow 0$ as $t \to -\infty$.

Proof. The Harnack inequality implies that the speed of each tip of Γ_t is greater than λ , hence the quantity $\sup_{\Gamma_t} |x_{n+1}| - \lambda |t|$ decreases, as t increases. In addition, the solution $\hat{\Gamma}_t$ is contained in $\hat{\Gamma}_{t,0}$, by construction, where $\hat{\Gamma}_{t,0}$

is given in the proof of Proposition 5.3. Hence $\sup_{\Gamma_t} |x_{n+1}| \leq \lambda |t|$, i.e. $\lim_{t \to -\infty} (\sup_{\Gamma_t} |x_{n+1}| - \lambda |t|) := L \leq 0$. If L < 0, this implies that the solution $\hat{\Gamma}_t$ is contained in $\Omega \times [-(L + \lambda |t|), L + \lambda |t|]$. By the convergence of solution to the translating soliton around the tips, we have that

(5.5)
$$\limsup_{t \to -\infty} \operatorname{Vol}(\hat{\Gamma}_t) - 2[(L+\lambda|t|)|\Omega| - V_{\Omega}] \le 0.$$

Since $2[(L+\lambda|t|)|\Omega|-V_{\Omega}] = \omega_n|t|+2L|\Omega|-2V_{\Omega}$, this contradicts Proposition 5.3 (i). This proves the assertion.

6. Uniqueness of compact ancient solution

Given a bounded domain $\Omega \in \mathbb{R}^n$ we recall that u_Ω denotes the translator associated with the domain Ω satisfying $\inf_{x \in \Omega} u_\Omega(x) = 0$. Let us recall that if Ω is $C^{1,1}$

$$V_{\Omega} := \int_{\Omega} u_{\Omega}(x) \, dx < \infty$$

by Lemma 5.1. We have shown in the previous section that there exists a compact ancient solution Γ_t , $t \in (-\infty, -\frac{2V_{\Omega}}{\omega_n})$ which is asymptotic to the cylinder $\Omega \times \mathbb{R}$ and which becomes extinct at time $T := -\frac{2V_{\Omega}}{\omega_n}$. We will next show that Γ_t is unique up to translations in space along the axis e^{n+1} and translations in time.

Let us briefly *outline the proof* of the Theorem which will be given below. As we stated in Theorem 1.4, our goal is to show that any given compact ancient solution Σ_t asymptotic to $\Omega \times \mathbb{R}$ which becomes extinct at time $t = -\frac{2V_{\Omega}}{\omega_n}$ is same as Γ_t , the solution constructed in the previous section, up to a translation in e_{n+1} direction. For now, let us set aside to deal with this translation. The main step in our proof is to show the inclusion

$$\hat{\Gamma}_t \subset \hat{\Sigma}_t$$
, for all $t < T$.

Recall that Γ_t was obtained as the limit of $\Gamma_{s,t-s}$ as $s \to -\infty$, where $\Gamma_{s,\tau}$ is the GCF running from $\Gamma_{s,0}$ and $\Gamma_{s,0}$ is the compact surface obtained from the gluing of two translators so that the distances from each tip to the origin is equal to $|s| \lambda$. Thus, it would have been sufficient to show that $\hat{\Gamma}_{s,0} \subset \hat{\Sigma}_s$, for all $s \ll -1$. However, this is unlikely to hold in general. Instead, it suffices to find a family of convex sets $\hat{K}_s \subset \hat{\Sigma}_s$ satisfying $\hat{K}_s \subset \hat{\Gamma}_{s,0}$ and $\operatorname{Vol}(\hat{\Gamma}_{s,0} \setminus \hat{K}_s) \to 0$, as $s \to -\infty$. If $\hat{K}_{s,\tau}$ is the GCF from K_s , then $\hat{K}_{s,t-s} \subset \hat{\Sigma}_t$ for all $s \ll -1$. Meanwhile, $\hat{K}_{s,t-s} \subset \hat{\Gamma}_{s,t-s}$ and $\operatorname{Vol}(\hat{K}_{s,t-s}) \operatorname{Vol}(\hat{\Gamma}_{s,t-s}) = \operatorname{Vol}(\hat{K}_{s,0}) - \operatorname{Vol}(\hat{\Gamma}_{s,0}) \to 0$ as $s \to -\infty$, showing that $\hat{K}_{s,t-s} \to$ $\hat{\Gamma}_t$ as $s \to -\infty$. This proves $\hat{\Gamma}_t \subset \hat{\Sigma}_t$. In this argument, we used the following two properties in a strong way:

(i) $\partial_t(\operatorname{Vol}(\hat{\Sigma}_t)) = -\omega_n$, for any GCF solution $\Sigma_t = \partial \hat{\Sigma}_t$, and

18 BEOMJUN CHOI, KYEONGSU CHOI, AND PANAGIOTA DASKALOPOULOS

(ii) if two convex sets M_1, M_2 satisfy $\hat{M}_1 \subset \hat{M}_2$ and $\operatorname{Vol}(\hat{M}_1) = \operatorname{Vol}(\hat{M}_2)$, then $\hat{M}_1 = \hat{M}_2$.

Let us next describe how we find such a family \hat{K}_s . Instead of the translator u_{Ω} in the domain Ω , we will consider a hypersurface $x_{n+1} = u_{\epsilon}(x)$ on $(1 + \epsilon)^{-\frac{1}{n}}\Omega$ which is the translator of the same speed λ on the domain $(1 + \epsilon)^{-\frac{1}{n}}\Omega \setminus B_{\epsilon}(0)$ (see in Lemmas 6.1 and 6.2 below). When the domain shrinks from Ω to $(1 + \epsilon)^{-\frac{1}{n}}\Omega$, the associated translator speed larger that λ , but we can adjust the speed to be equal to λ by subtracting a small ball $B_{\epsilon}(0)$ from $(1 + \epsilon)^{-\frac{1}{n}}\Omega$. If we glue two such hypersurfaces at distance $|s| \lambda$, then the convergence of tip regions to the translator and the comparison principle from $-\infty$ time imply that Σ_s contains such a hypersurface as $s \ll -1$ (see in Lemma 6.3). Let $\hat{K}_s^{\epsilon_s}$ be the best possible (meaning the smallest ϵ_s) convex set which can be inserted in $\hat{\Sigma}_s$ by the argument above. We want Vol $(\hat{\Gamma}_{s,0} \setminus \hat{K}_s^{\epsilon_s}) \to 0$. Roughly,

$$\operatorname{Vol}(\widehat{\Gamma}_{s,0}) \approx \operatorname{Vol}(\Omega \times [-|s|\lambda, |s|\lambda]) - 2V_{\Omega} = 2\lambda |s| \operatorname{Vol}(\Omega) - 2V_{\Omega}$$

and

$$\operatorname{Vol}(\hat{K}_{s}^{\epsilon_{s}}) \approx \operatorname{Vol}((1+\epsilon_{s})^{-\frac{1}{n}}\Omega \times [-|s|\lambda, |s|\lambda]) - 2V_{\epsilon_{s}}$$
$$= 2\lambda |s| \operatorname{Vol}((1+\epsilon_{s})^{-\frac{1}{n}}\Omega) - 2V_{\epsilon_{s}}.$$

Here, V_{ϵ} denotes the volume under the surface $x_{n+1} = u_{\epsilon}(x)$ and it converges to V_{Ω} , as $\epsilon \to 0$ (see in Lemma 6.1). Since $\operatorname{Vol}((1+\epsilon_s)^{-\frac{1}{n}}\Omega) \approx (1-\epsilon_s)\operatorname{Vol}(\Omega)$ for small ϵ_s , we need $\epsilon_s = o(|s|^{-1})$ to approximate the volume of $\hat{\Gamma}_{s,0}$ by $K_s^{\epsilon_s}$ as $s \to -\infty$. A stronger statement of this assertion will be shown in Proposition 6.6.

We will now give the detailed proof of Theorem 1.4. Without loss of generality we may assume that $0 \in \Omega$ and that $\inf u_{\Omega}(\cdot) = u_{\Omega}(0) = 0$. Let us fix $r_0 > 0, R_0 > 0$ such that $B_{2r_0}(0) \subset \Omega \subset B_{R_0}(0)$. We begin with a few preliminary results, where η denotes a standard cut off function supported in $B_1 \subset \mathbb{R}^n$ such that $\int \eta \, dx = 1$.

Lemma 6.1. Let $\epsilon_0 := \min(\frac{r_0}{2}, 1)$ and $\lambda = \frac{\omega_n}{2|\Omega|}$. Given $\epsilon \in (0, \epsilon_0)$, there is a unique convex solution $u_{\epsilon} : \Omega_{\epsilon} := (1+\epsilon)^{-\frac{1}{n}} \Omega \to \mathbb{R}$ to the equation

(6.1)
$$\sqrt{1+|Du_{\epsilon}(x)|^2} K(u_{\epsilon},x) = \lambda \left(1+\epsilon^{1+n}\eta(\epsilon^{-1}x)\right),$$

satisfying the conditions

 $\inf_{\Omega_{\epsilon}} u_{\epsilon} = 0 \qquad and \qquad Du_{\epsilon}((1+\epsilon)^{-\frac{1}{n}}\Omega) = \mathbb{R}^{n}.$

Moreover, $V_{\epsilon} := \int_{(1+\epsilon)^{-\frac{1}{n}\Omega}} u_{\epsilon}(x) dx \to V_{\Omega}$, as $\epsilon \to 0$.

Proof. The result of Urbas in [40, 41] guarantee the existence of a unique solution of equation (6.1) satisfying the required conditions. In addition, standard regularity estimates for equations of Monge-Ampère type imply

that as $\epsilon \to 0$, $u_{\epsilon}(x)$ converges to the translator $u_{\Omega}(x)$ having $\inf_{\Omega} u(x) = 0$ and the convergence is in the C_{loc}^{∞} sense. The convergence of $V_{\epsilon} \to V_{\Omega}$ easily follows, since the proof of Lemma 5.1 can be applied uniformly to the solutions u_{ϵ} and gives

$$\sup_{\epsilon < \min(\frac{r_0}{2}, 1)} \int_{\{x \in (1+\epsilon)^{-\frac{1}{n}}\Omega : \operatorname{dist}(x, \partial(1+\epsilon)^{-\frac{1}{n}}\Omega) \le \delta\}} u_{\epsilon}(x) dx = o(1) \text{ as } \delta \to 0.$$

Since Σ_t converges to the translating soliton near tip regions, there is $\tau_0 \ll -1$ and M > 0 such that

(6.2)
$$|Du^+(x,t)|, |Du^-(x,t)| \le M, \quad \text{on } B_{\epsilon_0}(0) \times (-\infty, \tau_0].$$

In particular, this implies $|Du_{\Omega}(x)| \leq M$ on $x \in B_{\epsilon_0}(0)$. We will use τ_0 and M in the remaining of this section. Also recall that we have assumed $u_{\Omega}(0) = \inf u_{\Omega}(\cdot) = 0$ in this section.

Lemma 6.2. For $\epsilon \in (0, \epsilon_0)$, u_{ϵ} defined in Lemma 6.1 satisfies

 $u_{\epsilon}(x) + M\epsilon \ge u_{\Omega}(x) \quad \text{for all } x \in (1+\epsilon)^{-\frac{1}{n}}\Omega.$

Proof. Note $u_{\epsilon}(x) + M\epsilon \geq M\epsilon \geq u(x)$ on B_{ϵ} and it becomes infinity at $\partial(1+\epsilon)^{-\frac{1}{n}}\Omega$. They are both solutions to the translating soliton equation of speed λ in the domain $(1+\epsilon)^{-\frac{1}{n}}\Omega \setminus B_{\epsilon}$. Hence, the comparison principle implies the lemma.

For the next lemma let us define $d(t) = \min(|u^+(0,t)|, |u^-(0,t)|)$. $u^+(0,t)$ and $u^-(0,t)$ are very similar to $h^+(t)$ and $h^-(t)$, respectively (recall Definition 3.1) in the following sense: since $u^-(x,t) - h^-(t)$ and $u^+(x,t) - h^+(t)$ converges to $u_{\Omega}(x)$ and $-u_{\Omega}(x)$ as $t \to -\infty$, respectively, and $\inf u_{\Omega} =$ u(0) = 0, we have $|u^-(0,t) - h^-(t)| = o(1)$ and $|u^+(0,t) - h^+(t)| = o(1)$ as $t \to -\infty$. Moreover, $\partial_t u^-(0,t) \ge \lambda$ and $\partial_t u^+(0,t) \le -\lambda$.

Lemma 6.3. Let $\epsilon \in (0, \epsilon_0)$ be a fixed given number. For a solution Σ_t satisfying the assumptions of Theorem 1.4, set $d(t) = \min(|u^+(0,t)|, |u^-(0,t)|)$. Then, the solution Σ_t encloses the convex body

$$\hat{K}_{t,\epsilon} := \{ (x', x'_{n+1}) \in \mathbb{R}^{n+1} : |x'_{n+1}| \le -u_{\epsilon}(x') - M\epsilon + d(t) \}$$

for all $t \leq \min(t_{\epsilon}, \tau_0)$, where τ_0 is a time satisfying (6.2) and

$$t_{\epsilon} := \sup\{t : ((1+\epsilon)^{-\frac{1}{n}}\Omega) \times \{0\} \subset \hat{\Sigma}_t \cap \{x_{n+1} = 0\}\}.$$

Proof. We apply again the comparison principle. Since $\partial_t u^-(0,t) \geq \lambda$ and $\partial_t u^+(0,t) \leq -\lambda$ (this is due to the convergence to the soliton shown in Theorem 3.3 and the Harnack inequality in Lemma 3.5 (i)), we have that $K_{t,\epsilon} := \partial \hat{K}_{t,\epsilon}$ is a supersolution of the Gauss curvature flow except the crosssection $\Sigma_t \cap \{x_{n+1} = 0\}$ and the two tip regions which are components of $(B_{\epsilon}(0) \times \mathbb{R}) \cap K_{t,\epsilon}$. From the choice of τ_0 and d(t), we have $u_{\epsilon}(x) + M\epsilon - d(t) \geq 0$

 $u^{-}(x,t)$ and $-u_{\epsilon}(x) - M\epsilon + d(t) \leq u^{+}(x,t)$ on $(x,t) \in B_{\epsilon} \times (-\infty, \tau_{0}]$. Thus if $t \leq \min(t_{\epsilon}, \tau_{0})$, then $K_{t,\epsilon}$ does not touch to Σ_{t} on these three regions.

Moreover, the convergence to the translating soliton around tips and Lemma 6.2 imply Σ_t contains $K_{t,\epsilon}$ for large negative times. By the comparison principle, we have $\hat{K}_{t,\epsilon} \subset \hat{\Sigma}_t$ for $t < \min(t_{\epsilon}, \tau_0)$.

A crucial ingredient in the proof Theorem 1.4 is Proposition 6.6 which is an estimate on the rate that shows how fast Σ_t becomes asymptotic to $\Omega \times \mathbb{R}$. We achieve this by a barrier argument. A one-parameter family of rotationally symmetric convex hypersurfaces in \mathbb{R}^{n+1} represented by

$$\Sigma_t = \{(x, x_{n+1}) \in \mathbb{R}^{n+1} : |x| = r(x_{n+1}, t)\}$$

is a solution to the GCF if $r_{xx} < 0$ and

$$r_t = r_{xx}(1 + r_x^2)^{-\frac{n+1}{2}}r^{-(n-1)}.$$

Hence, Σ_t plays the role of an inner barrier if

(6.3)
$$r_t \le r_{xx}(1+r_x^2)^{-\frac{n+1}{2}}r^{-(n-1)}$$

Lemma 6.4 (Ancient entasis). For given $\epsilon > 0$ and L > 1, consider 1parameter family of (incomplete) hypersurfaces $\{\Sigma_t^e\}_{t\leq 0}$ defined by

$$\Sigma_t^e = \{ (x_1, \dots, x_n, x_{n+1}) : |(x_1, \dots, x_n)| = r(x_{n+1}, t), |x_{n+1}| \le -t \},\$$

where r(x,t) on $|x| \leq -t$ is defined by

$$r(x,t) = 2\epsilon \left(2 - \exp \frac{t}{L} \cosh \frac{x}{L}\right),$$

where $L = (2\epsilon)^{-(n-1)}$. Then $\{\Sigma_t^e\}_{t \leq 0}$ is an (incomplete) inner barrier to the GCF. Moreover, on its boundary $|x_{n+1}| = |t|$, there holds

(6.4)
$$2\epsilon \le r(|t|, t) = r(-|t|, t) \le 3\epsilon.$$

Remark 6.5. Note that Σ_t^e has rotationally symmetry about x_{n+1} -axis and has reflection symmetry about $\{x_{n+1} = 0\}$.

Proof. Since $r_{xx} \leq 0$, $(1 + r_x^2) \geq 1$, and $r \geq 2\epsilon$, we have

$$-r_{xx}(1+r_x^2)^{-\frac{n+1}{2}}r^{-(n-1)} \le -(2\epsilon)^{-(n-1)}r_{xx}$$

Therefore, combining with $L = (2\epsilon)^{-(n-1)}$ yields (6.3). In addition, we can obtain (6.4) by observing $\frac{1}{2} \leq e^{\frac{t}{L}} \cosh(\frac{t}{L}) \leq 1$.

Proposition 6.6. Let Ω be a bounded convex domain with $C^{1,1}$ boundary and let $\Sigma_t = \partial \hat{\Sigma}_t$, for $t \in (-\infty, T)$, be a compact ancient solution which is asymptotic to $\Omega \times \mathbb{R}$. Then there exist large positive C, $L < \infty$ such that

$$\sup_{x \in \partial \Omega \times \{0\}} d(x, \Sigma_t \cap \{x_{n+1} = 0\}) \le Ce^{t/L}.$$

Here L depends only on Ω , and C depends only on Ω , Σ_t .

Proof. After a rescaling, we may assume $|\Omega| = \frac{\omega_n}{4}$ so that the translator asymptotic to $\Omega \times \mathbb{R}$ has the speed $\lambda = 2$. Then, by the Harnack Lemma 3.5 (i), we have

$$(6.5) |\partial_t u^+(x,t)| \ge 2, |\partial_t u^-(x,t)| \ge 2,$$

where $u^{\pm}(x,t)$ are the maximum and minimum height functions of Σ_t from Definition 3.2.

Next, due to the $C^{1,1}$ boundary assumption, there is some constant $\epsilon > 0$ such that at each point $p \in \partial \Omega$, we have $B_{6\epsilon}(p+6\epsilon n_p) \subset \Omega$, where n_p denotes the unit inward pointing normal vector to $\partial \Omega$ at p. Namely, given $p \in \partial \Omega$, Ω has an inscribed ball of radius 6ϵ tangent at p.

We denote the 0-level set of $\hat{\Sigma}_t$ by $\tilde{\Omega}_t := \{x \in \mathbb{R}^n : (x, 0) \in \hat{\Sigma}_t\}$. Since $\tilde{\Omega}_t$ increases to Ω as $t \to -\infty$, given $\delta \in (0, \epsilon]$ there is some $t_\delta \ll T$ such that

(6.6)
$$\sup_{x \in \partial \Omega} d(x, \tilde{\Omega}_{t_{\delta}}) \le \delta.$$

Now, we claim that given $p \in \partial\Omega$ and $\delta \in (0, \epsilon]$, Σ_t encloses $\Sigma_{t-t_{\epsilon}}^e + (p_{\delta}, 0)$ for $t \leq t_{\epsilon}$, where $p_{\delta} := p + (4\epsilon + \delta)n_p$ and Σ_t^e is the ancient entasis defined in Lemma 6.4. To prove the claim, we will apply the comparison principle by showing that the Dirichlet and initial boundaries of $\Sigma_{t-t_{\epsilon}}^e + (p_{\delta}, 0)$ are enclosed by Σ_t .

We first consider the Dirichlet condition. By (6.4), we have

$$\partial \Sigma_{t-t_{\epsilon}}^{e} + (p_{\delta}, 0) \subset B_{3\epsilon}(p_{\delta}) \times \{ \pm |t - t_{\epsilon}| \}.$$

Therefore, it is enough to show

(6.7)
$$\overline{B_{3\epsilon}(p_{\delta})} \times [-|t - t_{\epsilon}|, |t - t_{\epsilon}|] \subset \hat{\Sigma}_{t}$$

for $t \leq t_{\epsilon}$. Indeed, (6.6) implies $\overline{B_{3\epsilon}(p_{\delta})} \subset \tilde{\Omega}_{t_{\epsilon}}$, and thus we have $u^+(x, t_{\epsilon}) \geq 0$ and $u^-(x, t_{\epsilon}) \leq 0$ for $|x-p_{\delta}| \leq 3\epsilon$. Therefore, (6.5) yields $u^+(x, t) \geq 2|t-t_{\epsilon}|$ and $u^-(x, t_{\epsilon}) \leq -2|t-t_{\epsilon}|$ in $\overline{B_{3\epsilon}(p_{\delta})}$ for $t \leq t_{\epsilon}$. This gives us (6.7).

To deal with the initial condition, we notice that

$$\Sigma_{t-t_{\epsilon}}^{e} + (p_{\delta}, 0) \subset B_{4\epsilon}(p_{\delta}) \times [-|t - t_{\epsilon}|, |t - t_{\epsilon}|].$$

On the other hand, (6.6) yields $B_{4\epsilon}(p_{\delta}) \subset \tilde{\Omega}_{t_{\delta}}$ so that we obtain $u^+(x, t_{\delta}) \geq 0$ and $u^-(x, t_{\delta}) \leq 0$ in $B_{4\epsilon}(p_{\delta})$. Hence, (6.5) again leads to $u^+(x, t) \geq 2|t - t_{\delta}|$ and $u^-(x, t_{\epsilon}) \leq -2|t - t_{\delta}|$ in $\overline{B_{3\epsilon}(p_{\delta})} \times (-\infty, t_{\delta}]$. Therefore, there is some $\bar{t} \ll t_{\epsilon}$ depending only on t_{ϵ}, t_{δ} such that

$$B_{4\epsilon}(p_{\delta}) \times [-|t - t_{\epsilon}|, |t - t_{\epsilon}|] \subset \hat{\Sigma}_t$$

holds for $t \leq \bar{t}$.

Since $\Sigma_{t-t_{\epsilon}}^{e} + (p_{\delta}, 0)$ satisfies the boundary conditions as an inner barrier of Σ_{t} , the comparison principle guarantees $\Sigma_{t-t_{\epsilon}}^{e} + (p_{\delta}, 0) \subset \hat{\Sigma}_{t}$ for $t \leq t_{\epsilon}$.

Passing δ to 0 and interpreting the containment among zero level sets, we conclude

$$d(p, \tilde{\Omega}_t) \le d(p, \pi(\Sigma_{t-t_{\epsilon}}^e) + p + 4\epsilon n_p) = 4\epsilon - r(0, t - t_{\epsilon}) = 2\epsilon e^{\frac{t-t_{\epsilon}}{L}}$$

where $\pi(\Sigma_{t-t_{\epsilon}}^{e}) = \{x : (x,0) \in \Sigma_{t-t_{\epsilon}}^{e}\}$. This completes the proof, since ϵ , $L = (2\epsilon)^{-(n-1)}$, and t_{ϵ} are independent on p.

Remark 6.7. The exponential decay in Proposition 6.6 is sharp in the sense that the non-compact translators satisfies similar bounds (as in the proof of Lemma 5.1) and the Grim Reaper in \mathbb{R}^2

$$\sin y = e^t \cosh x$$

has no better decay.

Remembering $0 \in \Omega$, Proposition 6.6 implies that there exists large positive L such that

(6.8)
$$\epsilon_t := \inf\{\epsilon > 0 : (1+\epsilon)^{-\frac{1}{n}} \Omega \subset \Omega_t\} = O(e^{t/L}), \quad \text{as } t \to -\infty.$$

Instead of this exponential decay, $\epsilon_t = o(|t|^{-1})$ will be sufficient to conclude the proof of Theorem 1.4.

Proof of Theorem 1.4. We begin by choosing a number $t_1 \ll -1$ such that the cross-section $\tilde{\Omega}_{t_1} := \{x \in \mathbb{R}^n : (x,0) \in \hat{\Sigma}_{t_1}\}$ contains the origin, and hence ϵ_t defined in (6.8) is a finite number for each $t \leq t_1$. Recall our notation $\Gamma_t = \partial \hat{\Gamma}_t$ and $\Sigma_t = \partial \hat{\Sigma}_t$, where $\hat{\Gamma}_t$ is given in Proposition 5.3. As the key step, we claim

(6.9)
$$\hat{\Gamma}_{t-t_1} \subset \hat{\Sigma}_t, \quad \text{for all } t < t_1 - 2\omega_n^{-1}V_\Omega$$

Note first that, by Lemma 6.3 and the definition of ϵ_t , we have that

(6.10)
$$K_{t,\epsilon_t} \subset \Sigma_t$$

for $t \leq \tau_1$ where $\tau_1 := \min(t_1, \tau_0, \sup\{t : \epsilon_t < \epsilon_0)\})$. Here, τ_0 and ϵ_0 are defined in (6.2) and Lemma 6.1, respectively. Furthermore since $\partial_t u^-(0,t) \geq \lambda$ and $\partial_t u^+(0,t) \leq -\lambda$ by the Harnack, we obtain

(6.11)
$$\hat{K}'_t := \{ (x', x'_{n+1}) : |x'_{n+1}| \le -u_{\epsilon_t}(x') - M\epsilon_t + \lambda |t - t_1| \} \subset \hat{K}_{t,\epsilon_t},$$

for all $t \leq \tau_1$. Next, recalling $\Gamma_{s,0}$ from the proof of Proposition 5.3, we have

(6.12)
$$K'_t \subset \Gamma_{t-t_1,0}$$

for $t \leq \tau_1$ by Lemma 6.2. Moreover, we have

(6.13)
$$\operatorname{Vol}(\hat{\Gamma}_{t-t_1,0}) - \operatorname{Vol}(\hat{K}'_t) \to 0, \quad \text{as } t \to -\infty.$$

Indeed, points in $\hat{\Gamma}_{t-t_1,0} \setminus \hat{K}'_t$ belong to one of two cylinders either $\Omega \setminus \Omega_{\epsilon_t} \times [-\lambda|t-t_1|, \lambda|t-t_1|]$ or $\Omega_{\epsilon_t} \times [-\lambda|t-t_1|, \lambda|t-t_1|]$. As a consequence of (6.8) (which is a consequence of Proposition 6.6), the volume of the former cylinder converges to zero. Moreover, the volume of $\hat{\Gamma}_{t-t_1,0} \setminus \hat{K}'_t$ inside of

the later cylinder converges to zero since $\lim_{t\to-\infty} V_{\epsilon_t} = V_{\Omega}$ by Lemma 6.1. Now (6.13) follows by combining these two.

On the other hand, we denote the GCFs running from $K'_{t,0} = \partial \hat{K}'_t$ and $\Gamma_{s,0} = \partial \hat{\Gamma}_{s,0}$ by $K'_{t,s} = \partial \hat{K}'_{t,s}$ and $\Gamma_{s,t} = \partial \hat{\Gamma}_{s,t}$, respectively. Since (6.10) and (6.11) gave us $\hat{K}'_t \subset \hat{K}_{t,\epsilon_t} \subset \hat{\Sigma}_t$, the comparison principle implies

$$\hat{K}'_{t,t_2-t} \subset \hat{\Sigma}_{t_2}, \quad \text{for all } t_2 \ge t.$$

We remember (6.12) and apply (6.13) to obtain

$$\operatorname{Vol}(\hat{\Gamma}_{t-t_1,t_2-t}) - \operatorname{Vol}(\hat{K}'_{t,t_2-t}) = \operatorname{Vol}(\hat{\Gamma}_{t-t_1,0}) - \operatorname{Vol}(\hat{K}'_t) \to 0$$

as $t \to -\infty$. In addition, we have $\Gamma_{t-t_1,t_2-t} \downarrow \Gamma_{t_2-t_1}$ as $t \to -\infty$ by (5.4). Therefore, we conclude that

$$(6.14) \qquad \qquad \hat{\Gamma}_{t_2-t_1} \subset \hat{\Sigma}_{t_2}$$

for all $t_2 < t_1 - \frac{2V_{\Omega}}{\omega_n}$, which proves our claim (6.9).

We will now use the inclusion (6.9) to conclude the proof of our uniqueness theorem. First, since $\hat{\Gamma}_{t-t_1} \subset \hat{\Sigma}_t$ and $\partial_t \operatorname{Vol}(\hat{\Sigma}_t) = \partial_t \operatorname{Vol}(\hat{\Gamma}_{t-t_1}) = -\omega_n$, we obtain

(6.15)
$$\operatorname{Vol}(\hat{\Sigma}_t \setminus \hat{\Gamma}_{t-t_1})$$
 is constant in time.

We recall the definitions of heights h^{\pm} of Σ_t given in Definition 3.1. Since $\hat{\Gamma}_{t-t_1} \subset \hat{\Sigma}_t$, $\partial_t h^+(t) \leq -\lambda$ and $\partial_t h^-(t) \geq \lambda$, we obtain that, as $t \to -\infty$, $h^+(t) - \lambda |t|$ increases and $h^-(t) + \lambda |t|$ decreases. In addition, by Theorem 3.3, Proposition 5.4, and (6.15) we have

(6.16)
$$\lim_{t \to -\infty} h^+(t) - \lambda |t| = C^+, \qquad \lim_{t \to -\infty} h^-(t) + \lambda |t| = C^-,$$

for some constants C^+ and C^- . Next, recall $\hat{\Gamma}_{t-t_1} \subset \hat{\Gamma}_{t-t_1,0}$ by (5.4) and $\operatorname{Vol}(\hat{\Gamma}_{t-t_1,0} \setminus \hat{\Gamma}_{t-t_1})$ converges to 0 as $t \to -\infty$ by (5.3) and Proposition 5.3 (i). Thus $\operatorname{Vol}(\hat{\Sigma}_t \setminus \hat{\Gamma}_{t-t_1}) = \operatorname{Vol}(\hat{\Sigma}_t \setminus \hat{\Gamma}_{t-t_1,0}) + o(1)$ as $t \to -\infty$. Observe

$$\operatorname{Vol}(\hat{\Sigma}_{t} \setminus \hat{\Gamma}_{t-t_{1},0}) = \int [(u_{\Omega}(x) - \lambda | t - t_{1} |) - u^{-}(x,t)]_{+} dx + \int [u^{+}(x,t) + (u_{\Omega}(x) - \lambda | t - t_{1} |)]_{+} dx$$

where we interpret $u^- = \infty$ and $u^+ = -\infty$ outside of their domain. By the dominated convergence theorem, the two integrals above converge to $(-C^- - \lambda t_1)|\Omega|$ and $(C^+ - \lambda t_1)|\Omega|$, respectively, as $t \to -\infty$. Indeed, by (6.16), the first integrand is bounded by $u_{\Omega}(x) - C^- - \lambda t_1 \in L^1(\Omega)$ by Lemma 5.1 and the integrand converges locally uniformly to the constant $-C^- - \lambda t_1$ by Theorem 3.3. The same argument works for the second integral. This proves

$$\operatorname{Vol}(\hat{\Sigma}_t \setminus \hat{\Gamma}_{t-t_1,0}) = (C^+ - C^- - 2\lambda t_1)|\Omega| + o(1),$$

and we conclude $\operatorname{Vol}(\hat{\Sigma}_t \setminus \hat{\Gamma}_{t-t_1}) = (C^+ - C^- - 2\lambda t_1)|\Omega|$ as this is constant. Since both of Σ_t and Γ_t become extinct at $T = -\frac{2V_\Omega}{\omega_n}$, we have

$$(C^+ - C^- - 2\lambda t_1)|\Omega| = \operatorname{Vol}(\hat{\Sigma}_{T+t_1}) = -\omega_n t_1,$$

namely $C^+ = C^- =: C$.

It remains to show that $\Sigma_t = \Gamma_t + C e_{n+1}$ and the proof follows from what we have already done. We may summarize the first part of the proof as follows: if there exist constants t_0 , τ_1 , and a decreasing function $\epsilon_t = (|t|^{-1})$ as $t \to -\infty$, such that $\hat{\Sigma}_t$ contains

$$\hat{K}_{t,\epsilon_t,t_0} := \{ (x, x_{n+1}) : |x_{n+1}| \le -u_{\epsilon_t}(x) - M\epsilon_t + \lambda |t - t_0| \}$$

for all $t < \tau_1$, then $\hat{\Gamma}_{t-t_0} \subset \hat{\Sigma}_t$. By looking at the intersection between Σ_t and $\{x_{n+1} = C\}$, we may define ϵ'_t in the same way as ϵ_t is defined in (6.8), that is

$$\epsilon'_t := \inf\{\epsilon > 0 : (1+\epsilon)^{-\frac{1}{n}} \Omega \subset \Omega'_t\}$$

where Ω'_t is the cross-section of $\hat{\Sigma}_t$ by $\{x_{n+1} = C\}$. Similarly to Proposition 6.6, we have $\epsilon'_t = o(|t|^{-1})$. Next, for each small $\delta > 0$, Lemma 6.3 and (6.16) imply that there is $\tau_{\delta} \ll -1$ such that $\hat{K}_{t,\epsilon'_t,-\delta} + C\epsilon_{n+1} \subset \hat{\Sigma}_t$, for all $t < \tau_{\delta}$. Then, as we obtained (6.14), we can derive $\hat{\Gamma}_{t+\delta} + Ce_{n+1} \subset \hat{\Sigma}_t$ by repeating the argument from (6.11) to (6.14) after replacing t_1 by $-\delta$. Taking $\delta \to 0$, we get $\hat{\Gamma}_t + Ce_{n+1} \subset \hat{\Sigma}_t$. Since $\operatorname{Vol}(\hat{\Gamma}_t) = \operatorname{Vol}(\hat{\Sigma}_t) = -\omega_n t - 2V_\Omega$, we finally conclude that $\Gamma_t + Ce_{n+1} = \Sigma_t$.

References

- B. Andrews. Entropy inequalities for evolving hypersurfaces. Communications in Analysis and Geometry, 2(1): 53–64, 1994.
- B. Andrews. Contraction of convex hypersurfaces by their affine normal. Journal of Differential Geometry, 43(2): 207–230, 1996.
- B. Andrews. Gauss curvature flow: the fate of the rolling stones. *Inventiones mathe-maticae*, 138(1): 151–161, 1999.
- B. Andrews. Motion of hypersurfaces by Gauss curvature. Pacific Journal of Mathematics, 195(1): 1–34, 2000.
- [5] B. Andrews. Classification of limiting shapes for isotropic curve flows. Journal of the American Mathematical Society, 16(2): 443–459, 2003.
- [6] B. Andrews and X. Chen. Surfaces moving by powers of Gauss curvature. Pure and Applied Mathematics Quarterly, 8(4): 825–834, 2012.
- [7] B. Andrews, P. Guan, and L. Ni. Flow by powers of the Gauss curvature. Advances in Mathematics, 299: 174–201, 2016.
- [8] S.B. Angenent, P. Daskalopoulos, and N. Sesum. Uniqueness of two-convex closed ancient solutions to the mean curvature flow. Ann. Math., 192.2 (2020): 353-436.
- [9] T. Bourni, M. Langford, and G. Tinaglia. A collapsing ancient solution of mean curvature flow in ℝ³. To appear in Journal of Differential Geometry (arXiv:1705.06981 (2017)).
- [10] T. Bourni, M. Langford, and G. Tinaglia. Convex ancient solutions to curve shortening flow. Calc. Var. PDE., 59.4 (2020): 1-15.

- [11] S. Brendle, K. Choi and P. Daskalopoulos. Asymptotic behavior of flows by powers of the Gaussian curvature. Acta Mathematica, 219(1): 1–16, 2017.
- [12] S. Brendle. Ancient solutions to the Ricci flow in dimension 3. Acta Math., 225.1 (2020): 1-102.
- [13] S. Brendle and K. Choi. Uniqueness of convex ancient solutions to mean curvature flow in R³. Invent. Math., 217.1 (2019): 35-76.
- [14] E. Calabi. Improper affine hyperspheres of convex type and a generalization of a theorem by K.Jörgens. *Michigan Math. J.* 5: 105–126, 1958.
- [15] E. Calabi. Complete affine hyperspheres. I. Symposia Mathematica Vol. X 19–38 Academic Press, London, 1972.
- [16] S.Y. Cheng and S.T. Yau. Complete affine hypersurfaces. I. The completeness of affine metrics. Communications on Pure and Applied Mathematics, 39(6): 839–866, 1986.
- [17] B. Choi, K. Choi and P. Daskalopoulos. Convergence of curve shortening flow to translating solitons Amer. J. Math., 143.4 (2021): 1043-1077.
- [18] B. Choi, K. Choi and P. Daskalopoulos. Convergence of Gauss curvature flows to translating solitons. Adv. Math., 397 (2022): 108207.
- [19] K. Choi and P. Daskalopoulos. Uniqueness of closed self-similar solutions to the Gauss curvature flow. arXiv:1609.05487, 2016.
- [20] K. Choi, P. Daskalopoulos, and K.A. Lee. Translating solutions to the Gauss curvature flow with flat sides. Analysis & PDE 14.2 (2021): 595-616.
- [21] K. Choi, P. Daskalopoulos, L. Kim, and K.A. Lee. The evolution of complete noncompact graphs by powers of Gauss curvature. *Journal für die reine und angewandte Mathematik*, DOI: 10.1515/crelle- 2017–0032, 2017.
- [22] K. Choi, R. Hashofer and O. Hershkovits. Ancient low entropy flows, mean convex neighborhoods, and uniqueness. *To appear in Acta Math.*, arXiv:1810.08467 (2018).
- [23] B. Chow. Deforming convex hypersurfaces by the nth root of the Gaussian curvature. Journal of Differential Geometry 22(1): 117–138, 1985.
- [24] B. Chow. On Harnack's inequality and entropy for the Gaussian curvature flow. Communications on Pure and Applied Mathematics, 44(4): 469–483, 1991.
- [25] P. Daskalopoulos, R. Hamilton and N. Sesum. Classification of compact ancient solutions to the curve shortening flow. *Journal of Differential Geometry*, 84.3 (2010): 455-464.
- [26] P. Daskalopoulos, R. Hamilton and N. Sesum. Classification of ancient compact solutions to the Ricci flow on surfaces. *Journal of Differential Geometry*, 91.2 (2012): 171-214.
- [27] P. Daskalopoulos and R. Hamilton. The free boundary in the Gauss curvature flow with flat sides. Journal f
 ür die reine und angewandte Mathematik, 510: 187–227, 1999.
- [28] P. Daskalopoulos and K.A. Lee. Worn stones with flat sides all time regularity of the interface. *Invent. Math.*, 156(3): 445–493, 2004.
- [29] P. Daskalopoulos and O. Savin. C^{1,α} regularity of solutions to parabolic Monge-Ampère equations. Amer. J. Math, 134(4): 1051–1087, 2012.
- [30] P. Daskalopoulos and N. Sesum. Uniqueness of ancient compact non-collapsed solutions to the 3-dimensional Ricci flow. arXiv preprint arXiv:1907.01928 (2019).
- [31] W. Firey. Shapes of worn stones. *Mathematika*, 21(1): 1–11, 1974.
- [32] P. Guan and L. Ni. Entropy and a convergence theorem for Gauss curvature flow in high dimension. *Journal of European Mathematical Society*, 19(12): 3735–3761, 2017.
- [33] R. Hamilton. Worn stones with flat sides. Discourses Math. Appl, 3: 69–78, 1994.
- [34] R. Hamilton. Harnack estimate for the mean curvature flow. Journal of Differential Geometry, 41(1): 215–226, 1995.
- [35] H. Jian and X.J. Wang. Existence of entire solutions to the Monge-Ampère equation. Amer. J. Math., 136(4): 1093–1106, 2014.

26 BEOMJUN CHOI, KYEONGSU CHOI, AND PANAGIOTA DASKALOPOULOS

- [36] K. Jörgens. Über die Lösungen der Differentialgleichung $rt s^2 = 1$. Mathematische Annalen, 127: 130–134, 1954.
- [37] J.C.C. Nitsche. Elementary proof of Bernstein's theorem on minimal surfaces. Annals of Mathematics, 66(2): 543–544, 1957.
- [38] A.V. Pogorelov. On the improper convex affine hyperspheres. *Geometriae Dedicata*, 1(1): 33—46, 1972.
- [39] K. Tso. Deforming a hypersurface by its Gauss-Kronecker curvature. Communications on Pure and Applied Mathematics, 38(6): 867–882, 1985.
- [40] J. Urbas. Global Hölder estimates for equations of Monge-Ampère type. Invent. Math., 91(1): 1–29, 1988.
- [41] J. Urbas. Complete noncompact self-similar solutions of Gauss curvature flows I. Positive powers. *Mathematische Annalen*, 311(2): 251–274, 1998.

Acknowledgements

B. Choi has been partially supported by NSF grant DMS-1600658 and the National Research Foundation of Korea grant NRF-2022R1C1C1013511. B. Choi also thanks Columbia University and University of Toronto where the research was initiated and then developed.

K. Choi has been partially supported by KIAS Individual Grant MG078901 and TJ Park Science Fellowship.

P. Daskalopoulos has been partially supported by NSF grant DMS-1600658 and DMS-1900702.

Beomjun Choi: DEPARTMENT OF MATHEMATICS, POSTECH, 77 CHEONGAM-RO, NAM-GU, POHANG, GYEONGBUK 37673, REPUBLIC OF KOREA. *Email address*: bchoi@postech.ac.kr

Kyeongsu Choi: Korea Institute for Advanced Study, 85 Hoegiro, Dongdaemungu, Seoul 02455, Republic of Korea.

Email address: choiks@kias.re.kr

Panagiota Daskalopoulos: DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVER-SITY, 2990 BROADWAY, NEW YORK, NY 10027, USA. *Email address*: pdaskalo@math.columbia.edu